# Physical properties and the maximum compactness bound of a class of compact stars in  $f(Q)$  gravity

R. Sharma,\* A. Ghosh,<sup>[†](#page-1-1)</sup> and A. Paul<sup>[‡](#page-1-2)</sup>

Cooch Behar Panchanan Barma University, Vivekananda Street, Cooch Behar 736101, India. (Dated: September 10, 2024)

# Abstract

We investigate the physical behaviour of a stellar configuration by developing a compact stellar model within the framework of  $f(Q)$  gravity. We study the mass-radius  $(M - R)$  relationship and obtain the maximum compactness bound of the resultant stellar configuration by assuming the modification to be linear in non-metricity Q, i.e.  $f(Q) = \alpha Q + \beta$ . The maximum compactness bound proposed in  $f(Q)$  gravity is analogous to the Buchdahl bound in general relativity. We note that the compactness bound increases in  $f(Q)$  gravity. In the general relativistic limit ( $\alpha = -1$ ), our approach regains the Buchdahl bound for an incompressible star. Our observation might be relevant in the context of a recent observation with the MeerKAT observatory, which indicates the existence of high mass non-black hole compact objects which cannot be modelled by using the conventional neutron star equation of state (EoS).

## I. INTRODUCTION

The General Theory of Relativity (GTR) developed by Einstein remains the most modern theory of gravity to date, as the theory has successfully predicted many gravity tests, including the most recent discovery of gravitational waves. Despite its remarkable success, GTR faces many challenges on small and large scales. Among many other issues, one of the biggest challenges that GTR faces is its limitation in explaining the current accelerated expansion of the universe vis-a-vis the hidden source of the extra repulsive pressure. To explore the possible sources of late time cosmic expansion, different modified theories of gravity have so far been developed [\[1](#page-16-0)[–4\]](#page-16-1). Such modifications are mostly done in the matter segment of the Einstein–Hilbert action as well as in the geometry of the associated spacetime. Amongst many other alternative theories, one of the most popular proposals is the  $f(R)$  theory of gravity in which the Ricci scalar  $(R)$  is replaced by an arbitrary function  $f(R)$ in the Einstein–Hilbert (E–H) action [\[5\]](#page-16-2). GTR is based on the Riemannian geometry, where one considers the Ricci curvature  $R$  as a fundamental property of spacetime. However, this prescription does not include torsion or non-metricity, in general. A non-Riemannian geometry can incorporate torsion and non-metricity as additional geometrical properties of the

<span id="page-1-1"></span><span id="page-1-0"></span><sup>∗</sup> Corresponding author; [rsharma@associates.iucaa.in](mailto:rsharma@associates.iucaa.in)

<span id="page-1-2"></span><sup>†</sup> [arpitaghosh92727@gmail.com](mailto:arpitaghosh92727@gmail.com)

<sup>‡</sup> [paulabhishek.res@gmail.com](mailto:paulabhishek.res@gmail.com)

spacetime. Teleparallel gravity is an alternative to GTR, where the gravitational forces arise from the torsion T instead of the curvature R. As an extension of the symmetric teleparallel gravity, Harko *et al* [\[6\]](#page-16-3) proposed the modified  $f(Q)$  gravity. The consideration of the non-metricity  $(Q)$  serves as the mediator of gravitational interactions resulting in the  $f(Q)$ gravity [\[7\]](#page-16-4). It turns out that the choice of non-zero metricity can be an alternative approach to explain the late cosmic expansion [\[8\]](#page-16-5). Consequently,  $f(Q)$  gravity has gained significance in understanding its other cosmological and astrophysical implications. Hohmann et al [\[9\]](#page-16-6) examined the propagation velocity and potential polarization of gravitational waves in  $f(Q)$ gravity. Soudi et al. [\[10\]](#page-16-7) studied the strong field behaviour by analyzing the gravitational wave polarization. Various investigations have been carried out to explore the implications of  $f(Q)$  gravity in constraining observational data [\[11,](#page-16-8) [12\]](#page-16-9), energy conditions [\[13\]](#page-16-10), cosmography  $|14|$ , bouncing scenarios  $|15, 16|$  and studies of black holes  $|17|$ . For a comprehensive study of  $f(Q)$  gravity models, we refer to [\[18–](#page-17-4)[23\]](#page-17-5) and references therein.

Even though  $f(Q)$  gravity is primarily motivated by cosmological observations, the theory has gained widespread attention in astrophysics in the recent past. In particular, such a modification provides a platform to study relativistic, highly compact stars. Assuming Buchdahl metric potential [\[24\]](#page-17-6) for a strange star, Sokoliuk et al. [\[25\]](#page-17-7) analyzed the physical features of the star in both linear and non-linear forms of modified symmetric teleparallel gravity. Maurya et al [\[26\]](#page-17-8) investigated the maximum mass limit of a compact anisotropic star composed of strange quark matter in  $f(Q)$  gravity where  $f(Q)$  is linear in Q. The investigation predicts higher mass compact stars consistent with recent GW 190814 event observations. Lin and Zhai [\[27\]](#page-17-9) analyzed the implications of  $f(Q)$  gravity in the case of a spherically symmetric configuration. In their paper, with  $f(Q) = Q + \alpha Q^2$ , for a polytropic star, they have shown that a negative modification  $(\alpha < 0)$  could provide more stellar masses. In contrast, a positive value could reduce the amount of matter within the star. Making use of the Tolamn-Kuchowicz ansatz [\[28\]](#page-17-10) for a hybrid star composed of strange quark matter together with baryonic matter, Bhar *et al*  $[29]$  analyzed the maximum massradius relationship in  $f(Q)$  gravity. In a recent article, Araujo and Fortes [\[30\]](#page-17-12) investigated the maximum mass limit of a compact star admitting a polytropic EOS in  $f(Q) = Q + \xi Q^2$ gravity. The dynamical behaviour of compact sources has been studied by many investigators in  $f(Q)$  gravity [\[31–](#page-18-0)[33\]](#page-18-1).

It should be stressed here that physical features of compact stars have also been probed

in other modified theories of gravity which include the works in  $f(R)$  gravity [\[34,](#page-18-2) [35\]](#page-18-3),  $f(T)$ gravity [\[36,](#page-18-4) [37\]](#page-18-5),  $f(G)$  gravity [\[38,](#page-18-6) [39\]](#page-18-7),  $f(G, T)$  gravity [\[40,](#page-18-8) [41\]](#page-18-9), Rastall gravity [\[42\]](#page-18-10),  $f(Q, T)$ gravity [\[43\]](#page-18-11) and Teleparallel Palatini theory [\[44\]](#page-18-12). Using the minimal geometric deformation technique, Pradhan *et al* 45 studied geometrically deformed compact objects in  $f(Q, T)$ gravity in the presence of an electric field. Gul et al [\[46\]](#page-18-14) explored the viability and stability of compact stellar objects characterized by anisotropic matter within the framework of  $f(Q, T)$ gravity.

In this paper, we plan to analyze the implications of  $f(Q)$  gravity on the gross physical behaviour of a compact stellar configuration and its compatibility with recent observational data. In particular, we are interested in studying its effect on the maximum compactness bound. Note that in GTR, the maximum compactness bound is constrained by the Buchdahl bound  $M/R \leq 4/9$ . As far as mass is concerned, the known neutron star EOS provides stellar configurations within the mass range  $1 - 2$  M<sub>☉</sub>. Recently, Barr *et al* [\[47\]](#page-19-0) noted the masses of some of the pulsars in the mass gap region of known neutron stars (NSs) and low-mass black holes (BHs). The internal structure of such compact objects is unknown and demands further probe. We intend to study a stellar configuration in  $f(Q)$  gravity and explore its possible application in explaining such mass gap objects.

To achieve our goal, we first develop an internal solution for an anisotropic stellar configuration in  $f(Q)$  gravity whose exterior spacetime is described by the Schwarzschild-de Sitter metric. A relaxation in the pressure isotropy condition is relevant in the studies of compact stars, as pointed out by Herrea [\[48\]](#page-19-1). Various factors like a magnetic field, phase transition, viscosity, rotation and the mixture of two fluids, among others, can be potential sources for such as anisotropy [\[49–](#page-19-2)[52\]](#page-19-3). Amongst many others, anisotropic stellar distribution has been investigated by Sharma and Maharaj [\[53\]](#page-19-4) and Maurya et al [\[54,](#page-19-5) [55\]](#page-19-6).

Our paper is organized as follows: Section [II](#page-4-0) provides the mathematical framework of  $f(Q)$  gravity. Corresponding to a static and spherically symmetric anisotropic stellar configuration, we lay down the subsequent field equations in this section. In Section [III,](#page-7-0) making use of the Karmarker condition [\[56](#page-19-7)[–58\]](#page-19-8) together with a particular ansatz for one of the metric potentials, we provide an exact solution to the system. Using the relevant matching conditions in section [IV,](#page-8-0) we fix the values of the model parameters. Gross physical properties of the resultant stellar configuration are studied in section [V.](#page-9-0) In section [VI,](#page-11-0) we determine the maximum compactness bound in our developed model, which might be treated as  $f(Q)$  analogue of the Buchdal bound in GTR. The mass-radius relationship of the resultant configuration is studied and its relevance in the context of some recent observations is analyzed. Some concluding remarks are made in section [VII.](#page-13-0)

# <span id="page-4-0"></span>II. FORMALISM OF  $f(Q)$  GRAVITY

# A. General formalism

One formulates the  $f(Q)$  gravity model by setting the curvature R and the torsion T to be zero in the non-Riemannian geometry, assuming that the geometric information is encoded in the non-metricity  $Q$  only [\[59\]](#page-19-9). For the action

<span id="page-4-1"></span>
$$
S = \int \sqrt{-g} \, d^4x \left[ \frac{1}{2} f(Q) + \lambda_a^{bmn} R_{bmn}^a + \lambda_a^{mn} T_{mn}^a + L_m \right], \tag{1}
$$

the non-metricity  $Q$  is defined in terms of the affine connections as

$$
Q_{aun} \equiv \nabla_a g_{un} = \partial_a g_{un} - \Gamma^{\varrho}_{au} g_{\varrho n} - \Gamma^{\varrho}_{an} g_{u\varrho}, \tag{2}
$$

where

$$
\Gamma_{un}^{\varrho} = \{ \,_{un}^{\varrho} \} + K_{un}^{\varrho} + L_{un}^{\varrho}, \tag{3}
$$

$$
\{_{un}^{\varrho}\} = \frac{1}{2}g^{\varrho b} \left( \partial_u g_{bn} + \partial_n g_{bu} - \partial_b g_{un} \right), \tag{4}
$$

$$
K_{un}^{\varrho} = \frac{1}{2} T_{un}^{\varrho} + T_{(u \ n)}^{\ \varrho},\tag{5}
$$

$$
T_{un}^{\varrho} \equiv 2\Gamma_{[un]}^{\varrho},\tag{6}
$$

$$
L_{un}^{\varrho} = \frac{1}{2} Q_{un}^{\varrho} - Q_{(u \ n)}^{\ \varrho}.
$$
\n(7)

In equation [\(1\)](#page-4-1),  $\lambda_a^{bmn}$  corresponds to multipliers of the Lagrangian and  $L_m$  is the density of the Lagrangian matter. Expressing the non-metricity conjugate as

$$
P_{un}^{a} = -\frac{1}{4}Q_{un}^{a} + \frac{1}{2}Q_{(u}^{a}{}_{n)} + \frac{1}{4}\left(Q^{a} - \bar{Q}^{a}\right)g_{un} - \frac{1}{4}\delta^{a}{}_{(u}Q_{n)},\tag{8}
$$

we have

<span id="page-4-2"></span>
$$
Q = -Q_{aun}P^{aun},\tag{9}
$$

where,  $\overline{Q}_a \equiv Q_{au}^u$ ,  $Q_a \equiv Q_{a\,u}^{\,u}$ .

The subsequent field equations are obtained by varying the action [\(1\)](#page-4-1), which yields

<span id="page-5-2"></span>
$$
-T_{un} = \frac{2}{\sqrt{-g}} \nabla_a \left( \sqrt{-g} f_Q P_{un}^a \right) + \frac{1}{2} g_{un} f + f_Q \left( P_{uab} Q_n^{ab} - 2 Q_{abu} P_n^{ab} \right), \tag{10}
$$

where  $f_Q = \partial_Q f(Q)$ . Further, varying equation [\(1\)](#page-4-1) with respect to the affine connections, one obtains

<span id="page-5-0"></span>
$$
\nabla_{\rho} \varrho_a^{nu\rho} + \varrho_a^{un} = \sqrt{-g} f_Q p_a^{un} + H_a^{un}, \qquad (11)
$$

where the energy-momentum tensor has the form

$$
T_{un} = -\frac{2}{\sqrt{-g}} \frac{\partial \left(\sqrt{-g} \ L_{matter}\right)}{\partial \ g^{un}}.
$$
\n(12)

The density for the hyper-momentum tensor has the form

$$
H_a^{un} = -\frac{1}{2} \frac{\delta \ L_{matter}}{\delta \Gamma_{un}^a}.
$$
\n(13)

Now, using the asymmetric property of n and u, equation  $(11)$  can be written as

<span id="page-5-1"></span>
$$
\nabla_u \nabla_n \left( \sqrt{-g} f_Q P_a^{un} + H_a^{un} \right) = 0. \tag{14}
$$

Using the relation  $\nabla_u \nabla_n H_a^{un} = 0$ , we write equation [\(14\)](#page-5-1) as

$$
\nabla_u \nabla_n \left( \sqrt{-g} \ f_Q \ P^{un} \right) = 0. \tag{15}
$$

In a particular coordinate system, in the case of coincident gauge, the non-metricity is reduced to the following expression

$$
Q_{aun} = \partial_a \ g_{un},\tag{16}
$$

where, in the absence of curvature or torsion, the affine connections are given by

$$
\Gamma_{un}^a = \left(\frac{\partial x^a}{\partial \xi^{\varrho}}\right) \partial_u \partial_n \xi^{\varrho}.\tag{17}
$$

In the following sub-section, for a given line element, we obtain the non-metricity scalar and subsequent field equations.

## B. The line element

To understand the physical behaviour of a spherically symmetric, static and compact stellar object, we assume that the following line element describe the internal geometry of the star

<span id="page-6-0"></span>
$$
ds^{2} = -e^{\nu(r)} dt^{2} + e^{\lambda(r)} dr^{2} + r^{2} (\sin^{2} \theta d\phi^{2} + d\theta^{2}), \qquad (18)
$$

where  $\nu(r)$  and  $\lambda(r)$  are the undetermined functions of the radial coordinate r. Substituting equation [\(18\)](#page-6-0) into [\(9\)](#page-4-2), we obtain the non-metricity scalar  $Q$  as

$$
Q = -\frac{(2 e^{-\lambda(r)}) \left(\nu'(r) + \frac{1}{r}\right)}{r},\tag{19}
$$

where a prime denotes a derivative with respect to  $r$ . The energy-momentum tensor for an anisotropic matter is assumed to be of the form

<span id="page-6-1"></span>
$$
T_{ij} = (\rho + p_t) \zeta_i \zeta_j - p_t g_{ij} + (p_r - p_t) \xi_i \zeta_j, \tag{20}
$$

where  $\rho, p_r$  and  $p_t$  are the energy density, radial pressure and tangential pressure, respectively.  $\zeta_i$  is the four-velocity and  $\xi_i$  corresponds to a radial four-vector obeying the following relations

$$
\zeta^{a} = e^{-\frac{\nu}{2}} \delta^{a}_{0}, \ \zeta^{a} \zeta_{a} = 1, \ \xi^{a} = e^{-\frac{\lambda}{2}} \delta^{a}_{1}, \ \xi^{a} \xi_{a} = -1.
$$

For the metric [\(18\)](#page-6-0) and the energy-momentum tensor [\(20\)](#page-6-1), using [\(10\)](#page-5-2), we obtain the independent set of the field equations as

$$
\rho = -f_Q \left[ Q + \frac{1}{r^2} + \frac{e^{-\lambda(r)} \left( \lambda'(r) + \nu'(r) \right)}{r} \right] + \frac{f}{2},\tag{21}
$$

$$
p_r = f_Q\left(Q + \frac{1}{r^2}\right) - \frac{f}{2},\tag{22}
$$

$$
p_t = f_Q \left[ \frac{Q}{2} - e^{-\lambda(r)} \left[ \left( \frac{\nu'(r)}{4} + \frac{1}{2r} \right) (\nu'(r) - \lambda'(r)) + \frac{\nu''(r)}{2} \right] \right] - \frac{f}{2},\tag{23}
$$

$$
0 = \frac{\cot \theta}{2} Q' f_{QQ}.
$$
\n(24)

To close the system, we first assume a linear form of  $f(Q)$ 

$$
f(Q) = \alpha Q + \beta,\tag{25}
$$

where  $\alpha$  and  $\beta$  are constants.

We now apply the Karmakar condition [\[56\]](#page-19-7)

$$
R_{0101} R_{2323} = R_{0202} R_{1313} - R_{1202} R_{1303}.
$$
 (26)

and by determining the non-vanishing components of the Riemann tensor in Karmakar's condition, we obtain

<span id="page-7-1"></span>
$$
[\lambda'(r) - \nu'(r)] \nu'(r) e^{\lambda(r)} + 2 (1 - e^{\lambda(r)}) \nu''(r) + \nu'^2(r) = 0.
$$
 (27)

Integration of [\(27\)](#page-7-1) yields

<span id="page-7-3"></span>
$$
e^{\nu(r)} = \left(C + D \int \sqrt{e^{\lambda(r)} - 1} \, dr\right)^2,\tag{28}
$$

where C and D are constants of integration. Consequently the line element  $(18)$  is obtained in the form

$$
ds^{2} = -\left(C + D \int \sqrt{e^{\lambda(r)} - 1} \, dr\right)^{2} dt^{2} + e^{\lambda(r)} dr^{2} + r^{2} \left(d\theta^{2} + \sin^{2}\theta \, d\phi^{2}\right). \tag{29}
$$

Note that that the system can now be closed by choosing a physically meaningful form of  $\lambda(r)$ . This will be taken up in the following section.

#### <span id="page-7-0"></span>III. EXACT SOLUTION

In this section, we assume a particular form of the metric potential  $g_{rr}$ . We consider the Vaidya-and-Tikekar (VT) metric ansatz [\[60\]](#page-19-10), which has been found to be a useful geometric approach that can provide realistic stellar models. The VT ansatz is given by

<span id="page-7-2"></span>
$$
e^{\lambda(r)} = \frac{1 + K\frac{r^2}{L^2}}{1 - \frac{r^2}{L^2}}.\tag{30}
$$

The  $t =$ constant hyper-surface of the associated spacetime, when embedded in a 4–D Euclidean spacetime, turns out to be spheroidal rather than spherical. The spheroidal geometry is governed by the dimensionless curvature parameters  $K$  and another parameter  $L$  having the dimension of a  $length$ . Substituting [\(30\)](#page-7-2) in [\(28\)](#page-7-3) and integrating, we obtain

$$
e^{\nu(r)} = [C - D\sqrt{(1+K)(L^2 - r^2)}]^2.
$$
\n(31)

The constants C and D can be determined from the boundary conditions.

Subsequently, the physical variables are obtained in the form

<span id="page-8-1"></span>
$$
\rho = \frac{\beta}{2} - \frac{(K+1)(3L^2 + Kr^2)\alpha}{(L^2 + Kr^2)^2},
$$
\n
$$
p_r = \frac{1}{2(L^2 + Kr^2)(D(1+K) - C\sqrt{\frac{1+K}{L^2-r^2}}}\left(C\sqrt{\frac{1+K}{L^2-r^2}}\right)
$$
\n
$$
(-2\alpha(1+K) + \beta(L^2 + Kr^2)) - D(1+K)(-2\alpha(3+K) + \beta(L^2 + Kr^2))\right),
$$
\n(33)\n
$$
p_t = \frac{1}{2(L^2 + Kr^2)^2(D(1+K) - C\sqrt{\frac{1+K}{L^2-r^2}}}\left(C\sqrt{\frac{1+K}{L^2-r^2}}\right)
$$
\n
$$
(-2\alpha L^2(1+K) + \beta(L^2 + Kr^2)^2) - D(1+K)(-2\alpha(3+K) + \beta(L^2 + Kr^2))\right),
$$
\n(34)\n
$$
\Delta = p_t - p_r.
$$
\n(35)

The mass contained within a radius  $r$  is obtained as

$$
m(r) = \left(\frac{r^3}{12}\right) \left(\beta - \frac{6\alpha(K+1)}{L^2 + Kr^2}\right). \tag{36}
$$

# <span id="page-8-0"></span>IV. EXTERIOR SPACETIME AND BOUNDARY CONDITIONS

The exterior spacetime of the star under consideration is assumed to be described by the Schwartzchild de-Sitter metric

$$
ds^{2} = -\left(1 - \frac{2M}{r} - \frac{\Lambda}{3}r^{2}\right)dt^{2} + \left(1 - \frac{2M}{r} - \frac{\Lambda}{3}r^{2}\right)^{-1}dt^{2} + r^{2}\left(d\theta^{2} + \sin^{2}\theta \, d\phi^{2}\right), (37)
$$

where  $\Lambda = \frac{\beta}{2\alpha}$  is the cosmological constant and M is the total mass of the star. Matching the interior solution to the exterior solution across the physical boundary  $r = R$  of the star and demanding that the radial pressure should vanish at the boundary, we obtain the junction conditions as

$$
1 - \frac{2M}{R} - \frac{\Lambda}{3}R^2 = e^{\nu(R)} = e^{-\lambda(R)},
$$
\n(38)

$$
p_r|_{r=R} = 0. \tag{39}
$$

Substituting the values of  $e^{\nu(r)}$  and  $e^{\lambda(r)}$  in the above boundary conditions and utilizing

equation [\(33\)](#page-8-1), we determine the constants of the model as

$$
C = \sqrt{1 - \frac{2M}{R}} \left( \frac{3 + K}{2} - \frac{\beta}{4\alpha} (L^2 + KR^2) \right),
$$
\n(40)

$$
D = \frac{1}{\sqrt{(1+K)(L^2+KR^2)}} \left( \frac{1+K}{2} - \frac{\beta}{4\alpha} (L^2+KR^2) \right),\tag{41}
$$

$$
L = \frac{R}{\sqrt{2M}}\sqrt{R(1+K) - 2KM}.\tag{42}
$$

It should be stressed that the value of the cosmological constant is  $\sim 10^{-54} / km^2$ , and hence we take  $\beta \approx 0$  in our construction. Consequently, the constants can be fixed for different choices of  $\alpha$ , which will be taken up in the following section.

## <span id="page-9-0"></span>V. PHYSICAL ANALYSIS

In order to analyze the physical features of the resultant stellar configuration, we consider the pulsar  $4U1608-52$  whose mass and radius are estimated to be  $M = 1.74 \pm 0.14$   $M_{\odot}$  and  $R = 9.52 \pm 0.15 \; km$ , respectively [\[61\]](#page-19-11). For the given mass and radius, in Table [I,](#page-9-1) we show numerical values of the constants for different choices of  $\alpha$ .

$\alpha$	L	C	Ð
		$-1.0 63.5691 17.9892 0.0386$	
		$-1.5 63.5691 17.9892 0.0386$	
		$-2.0 63.5691 17.9892 0.0386$	

<span id="page-9-1"></span>TABLE I. Values of the constants for different choices of  $\alpha$  with  $\beta = 0$ . For numerical calculation, we have considered the estimated mass and radius of the pulsar  $4U1608 - 52$ .

In figure  $1(a)$ , we show the behaviour of the metric potentials at the interior of the star, which is regular throughout the stellar configuration. Interestingly, the metric potentials remain unaltered for different values of  $\alpha$ . To examine the  $f(Q)$  modification on other physical quantities, we plot the behaviour of physically meaningful quantities for different choices of  $\alpha$ . We note that the mass function  $m(r)$  increases for any departure from the GR limit  $(\alpha = -1)$  as shown in figure [1\(](#page-10-0)b).

We note that the density and both the radial and the tangential pressures increase as  $\alpha$ become more negative (figure  $2(a)$  and figure  $2(b)$ ). Note that the anisotropic parameter



<span id="page-10-0"></span>FIG. 1. (a) Metric potentials  $e^{\lambda(r)}$  and  $e^{\nu(r)}$  plotted against r. (b) Plot of mass function  $m(r)$ against radial distance r for different values of  $\alpha$ .



<span id="page-10-1"></span>FIG. 2. (a) Plot of radial pressure  $p_r$  and transverse pressure  $p_t$  against radial coordinate r for different values of  $\alpha$ . (b) Density( $\rho$ ) vs radius(r) plot for different values of  $\alpha$ .

 $\Delta > 0$  increases as  $\alpha$  departs from its GR limiting value (figure [3\(](#page-11-1)a)). The thermodynamic relation between the energy density and the radial pressure provides the equation of state (EOS) of the matter composition as shown in figure [3\(](#page-11-1)b). The EOS is almost linear and remains so for different values of  $\alpha$ . It is noteworthy that for  $\alpha > 0$ , we get unrealistic density and pressure profiles and hence we restrict our analysis for negative values of  $\alpha$  only.



<span id="page-11-1"></span>FIG. 3. (a) Anisotropic factor  $\Delta$  plotted against radial distance r for different values of  $\alpha$ . (b) Equation of state (EOS) for different values of  $\alpha$ .

## <span id="page-11-0"></span>VI. MAXIMUM COMPACTNESS BOUND IN  $f(Q)$  GRAVITY

#### A. Compactness bound

In [\(33\)](#page-8-1), by imposing the condition that the central pressure must not diverge, we get

<span id="page-11-2"></span>
$$
LD\sqrt{1+K} - C \ge 0 \quad (L \ne 0). \tag{43}
$$

We use this condition to determine the maximum compactness bound of the stellar configuration in  $f(Q)$  gravity. Substituting the values of C, D and L in equation [\(43\)](#page-11-2), we obtain a general compactness bound in  $f(Q)$  gravity as

<span id="page-11-3"></span>
$$
\frac{M}{R} \le \frac{2(3\alpha(1+K) + \alpha^2(2K+1))}{4K\alpha^2 - 9(1+K)}.
$$
\n(44)

In equation [\(44\)](#page-11-3), by setting  $\alpha = -1$  and  $K = 0$ , we regain the Buchdahl bound  $\frac{M}{R} \leq \frac{4}{9}$  $\frac{4}{9}$ . Note that for  $K = 0$  and  $\alpha = -1$ , the associated geometry is spherical and the model corresponds to an interior fluid sphere solution obtained by Schwarzschild. For  $K \neq 0$  with  $\alpha = -1$ , we regain the maximum compactness bound

$$
\frac{M}{R} \le \frac{2(K+2)}{(5K+9)},
$$
\n(45)

obtained earlier by Sharma et al  $[62]$ . The parameter K in this model was identified as the measure of anisotropic pressure. Thus, equation [\(44\)](#page-11-3) turns out to be a generalization of the compactness bound for an anisotropic star within the framework of  $f(Q)$  gravity. The bound is obtained simply by demanding that the central pressure must not diverge.

In our model, for a departure from the general relativistic limiting value of  $\alpha = -1$ , the compactness can go beyond the Buchdahl bound, as shown in figure [4.](#page-12-0) The bound, however, does not exceed the black hole limit 0.5. Moreover, for an arbitrary choice of  $K \neq 0$ ,  $\alpha$  cannot take values  $\alpha = \pm \frac{3}{2}$ 2  $\sqrt{(K+1)}$  $\frac{N^{(+)}}{K}$  as the compactness bound becomes singular for such choices. In figure [5,](#page-12-1) we show the maximum compactness bound for different values of K and  $\alpha$ .



<span id="page-12-0"></span>FIG. 4. Compactness bound  $\frac{M}{R}$  for different values of  $\alpha$  with  $K = 0$ .



<span id="page-12-1"></span>FIG. 5. Compactness bound  $M/R$  for different values of K and  $\alpha$ .

#### B. Mass-radius  $(M - R)$  relationship

To obtain the mass and radius  $(M - R)$  relationship, using equation [\(32\)](#page-8-1), we first obtain the surface density of the star

$$
\rho_s = \frac{\beta}{2} - \frac{(K+1)(3L^2 + KR^2)\alpha}{(L^2 + KR^2)^2}.
$$
\n(46)

For a given surface density  $\rho_s$  and curvature parameter K, we obtain the  $M-R$  relationship as shown in figure [\(6\)](#page-13-1). In our calculation, we assume the surface density to be 350 MeV fm<sup>-3</sup> and  $K = 50$ . From the figure, we note that for  $\alpha -1.0, -1.5, -2.0$ , the respective values of the maximum masses are obtained as 3.2479  $M_{\odot}$ , 3.9778  $M_{\odot}$  and 4.5932  $M_{\odot}$ , respectively. Thus, stellar models in  $f(Q)$  gravity can generate larger masses, which is consistent with some of the earlier observations. It should, however, be pointed out that due to a restriction on  $\alpha$ , one cannot arbitrarily increase the mass by this method.



<span id="page-13-1"></span>FIG. 6. Mass-radius relationship for different values of  $\alpha$ .

In table [II,](#page-14-0) we show the evaluated values of masses for the pulsar  $4U1608 - 52$  for its estimated radius  $R = 9.52$  km. While in the GR limit  $(\alpha = -1)$ , the calculated mass is 1.74  $M_{\odot}$ , it turns out that the mass increases in  $f(Q)$  gravity.

## <span id="page-13-0"></span>VII. CONCLUDING REMARKS

In this paper, we have developed a static anisotropic stellar model within the  $f(Q)$ gravity framework. We have developed a technique to obtain the maximum compactness

R(km)	$\alpha$	$M_{\odot}$
		$-1.0$   1.74
9.52		$-1.5 2.61$
		$-2.0 3.48$

<span id="page-14-0"></span>TABLE II. Table showing estimated mass of the pulsar  $4U1608-52$  for different values of  $\alpha$  where the radius fixed at  $R = 9.52 \pm 0.15$  km.

bound of the resultant configuration. The Buchdahl bound is regained in the GR limit for an incompressible fluid sphere. We have also obtained the mass-radius relationship, which clearly shows that  $f(Q)$  gravity has the potential of providing larger masses. Such an observation is interesting in the context of some recent observations which predicts the existence of binary pulsar systems possessing masses greater than the conventional highmass neutron stars or low-mass black holes. In particular, a recent deep survey of pulsars made with the radio telescope array MeerKAT [\[64,](#page-20-1) [65\]](#page-20-2) observed 13 millisecond pulsars in the globular cluster NGC 1851 [\[63\]](#page-20-3). The observation indicates the existence of three massive binary systems  $PSRJ0514 - 4002A$  [\[66\]](#page-20-4),  $PSRJ0514 - 4002D$  and  $PSRJ0514 4002E$  [\[67\]](#page-20-5). The latter has a spin period of 5.6 ms, an orbital period of 7.44 days and an orbital eccentricity of 0.71 [\[67\]](#page-20-5). An investigation carried out by Barr *et al* [\[47\]](#page-19-0) shows that  $PSRJ0514-4002E$  is a binary system of compact objects possessing masses 3.887 $\pm$ 0.004  $M_{\odot}$ and 2.09  $-$  2.71  $M_{\odot}$ . The large mass of the companion star is not akin to conventional neutron star models. In our model, we obtain the total mass  $2.09M_{\odot}$  and  $2.71M_{\odot}$  for respective values of  $\alpha = -1.07$  and  $\alpha = -1.8$ , where we have assumed  $K = 100$  and the surface density 874.6 MeV fm<sup>-3</sup>. The corresponding radii are estimated to be 6.1 km and 7.92 km, respectively. This is shown in figure [\(7\)](#page-15-0). Thus, our procedure in  $f(Q)$  gravity may provide a possible explanation for the large companion mass of the observed binary system  $PSRJ0514 - 4002E$ .

Notably, for a given value of  $K$ , our model shows that the Buchdahl bound can be exceeded by decreasing the value of  $\alpha$ . For example, assuming  $K = 50$ , we note that compactness goes beyond the Buchdahl bound for  $\alpha \leq -2.8$  as shown in figure [8\)](#page-15-1). However, a discontinuity in compactness is observed at  $\alpha = -1.514$  for the assumed value of  $K = 50$ . This, in a way, restricts the choice of  $\alpha$  in our analysis.

Some of the areas that remain unexplored in our analysis are as follows: Firstly, in our



<span id="page-15-0"></span>FIG. 7. Maximum mass vs radius curve for a pulsar  $PSRJ0514 - 4002E$ . For  $\alpha = -1.07$  and  $\alpha = -1.8$  we have have obtained masses  $2.09 M_\odot$  and  $2.71 M_\odot$  respectively.



<span id="page-15-1"></span>FIG. 8. Compactness  $(M/R)$  plotted against  $\alpha$  for a given value of K. The dotted straight line indicates the Buchdahl limit.

construction, we have assumed a modification which is linear  $(f(Q) = \alpha Q + \beta)$ . Moreover, due to the smallness of the cosmological constant, we have set  $\beta \sim 0$  in our analysis. It remains to be seen whether other forms of  $f(Q)$  can provide similar results, particularly in the context of the total mass of the resultant configuration. Secondly, we have used the Karmakar condition and a particular metric potential ansatz to close the system. Different techniques may be employed to generate a new class of solutions, and subsequently, the solutions may be used for similar studies.

## ACKNOWLEDGMENTS

RS gratefully acknowledges support from the Inter-University Centre for Astronomy and Astrophysics (IUCAA), Pune, India, under its Visiting Research Associateship Programme.

- <span id="page-16-0"></span>[1] R. Ferraro and F. Fiorini, *Born-Infeld gravity in Weitzenböck spacetime, Phys. Rev. D.* **78** (2008) 124019.
- [2] C.-Q. Geng et al, "Teleparallel" Dark Energy, Physics Letters B. 704 (2011) 384.
- <span id="page-16-1"></span>[3] Y.-F. Cai et al,  $f(T)$  teleparallel gravity and cosmology, Rep. Prog. Phys. **79** (2016) 106901.
- $[4]$  L. Järv and A. Toporensky, *General relativity as an attractor for scalar-torsion cosmology*, Phys. Rev. D. 93 (2016) 024051.
- <span id="page-16-2"></span>[5] S. Nojiri and S.D. Odintsov, Introduction to Modified Gravity and Gravitational Alternative for Dark Energy, Int. J. Geom. Meth. Mod. Phys. 4 (2007) 115.
- <span id="page-16-3"></span>[6] T. Harko, T. S. Koivisto, F. S. N. Lobo, G. J. Olmo and D. R. -Garcia, Coupling matter in modified Q gravity, Phys. Rev. D. 98 (2018) 084043.
- <span id="page-16-4"></span>[7] J. B. Jiménez, L. Heisenberg and T. Koivisto, *Coincident general relativity. Phys. Rev. D* 98 (2018) 044048.
- <span id="page-16-6"></span><span id="page-16-5"></span>[8] I. Kirsch, *Higgs mechanism for gravity*, *Phys. Rev. D* **72** (2005) 024001.
- [9] M. Hohmann, C. Pfeifer, U. Ualikhanova, and J. L. Said, Propagation of gravitational waves in symmetric teleparallel gravity theories, Phys. Rev. D. 99 (2019) 024009.
- <span id="page-16-7"></span>[10] I. Soudi, G. Farrugia, J. L. Said, V. Gakis and E. N. Saridakis, Polarization of gravitational waves in symmetric teleparallel theories of gravity and their modifications, Phys. Rev. D. 100 (2019) 044008.
- <span id="page-16-8"></span>[11] R. Lazkoz, F. S. N. Lobo, M. O. Baños and V. Salzano, *Observational constraints of*  $f(Q)$ gravity, Phys. Rev. D. 100 (2019) 104027.
- <span id="page-16-9"></span>[12] I. Ayuso, R. Lazkoz and V. Salzano, Observational constraints on cosmological solutions of  $f(Q)$  theories, Phys. Rev. D. 103 (2021) 063505.
- <span id="page-16-10"></span>[13] S. Mandal, P. K. Sahoo and J. R. L. Santos, Energy conditions in f(Q) gravity, Phys. Rev. D. 102 (2020) 024057.
- <span id="page-17-0"></span>[14] S. Mandal, D. Wang and P. K. Sahoo, *Cosmography in*  $f(Q)$  gravity, Phys. Rev. D. **102** (2020) 024057.
- <span id="page-17-1"></span>[15] S. Mandal, N. Myrzakulov, P. K. Sahoo and R. Myrzakulov, Cosmological bouncing scenarios in symmetric teleparallel gravity, Eur. Phys. J. Plus. 136 (2021) 760.
- <span id="page-17-2"></span>[16] F. Bajardi, D. Vernieri and S. Capozziello, Bouncing cosmology in f(Q) symmetric teleparallel gravity, Eur. Phys. J. Plus. 135 (2020) 912.
- <span id="page-17-3"></span>[17] F D'Ambrosio, S. D. B. Fell, L. Heisenberg, and S. Kuhn, *Black holes in*  $f(Q)$  *gravity, Phys.* Rev. D. 105 (2022) 024042.
- <span id="page-17-4"></span>[18] D. V. Nguyen, D. Sarnaaik, K. K. Boddy, E. O. Nadler and V. Gluscevic, Observational constraints on dark matter scattering with electrons, Phys. Rev. D. 104 (2021) 103521.
- [19] F. Esposito, S. Carloni, R. Cianci and S. Vignolo, Reconstructing isotropic and anisotropic  $f(Q)$  cosmologies, Phys. Rev. D. 105 (2022) 084061.
- [20] D. Zhao, *Covariant formulation of*  $f(Q)$  theory, Eur. Phys. J. C. 82 (2022) 303.
- [21] G. Mustafa, Z. Hassan, P.K. Sahoo, Traversable wormhole inspired by non-commutative geometries in  $f(Q)$  gravity with conformal symmetry, Annals of Physics. **437** (2021) 168751.
- [22] R. Kimura, A. Naruko and D. Yamauchi, On Lorentz-invariant bispin-2 theories, Phys. Rev. D. 104 (2021) 044021.
- <span id="page-17-5"></span>[23] N. Dimakis et al, Quantum cosmology in  $f(Q)$  theory, Class. Quantum Grav.. **38** (2021) 225003.
- <span id="page-17-7"></span><span id="page-17-6"></span>[24] H. A. Buchdahl, General Relativistic Fluid Spheres, Phys. Rev. 116 (1959) 1027.
- [25] O. Sokoliuk, S. Pradhan, P. K. Sahoo, A. Baransky, *Buchdahl quark stars within*  $f(Q)$  *theory*, Eur. Phys. J. Plus. 137 (2022) 1077.
- <span id="page-17-9"></span><span id="page-17-8"></span>[26] S. K. Maurya, G. Mustafa, M. Govender and K. N. Singh, , JCAP. 10 (2022) 003.
- [27] R-H Lin and X-H Zhai, Spherically symmetric configuration in  $f(Q)$  gravity, Phys. Rev. D. 103 (2022) 124001.
- <span id="page-17-11"></span><span id="page-17-10"></span>[28] R. C. Tolman, Phys. Rev, 55 (1939) 364; B. Kuchowicz, Acta. Phys. Pol. 33 (1968) 541.
- [29] P. Bhar, S. Pradhan, A. Malik, P. K. Sahoo, Physical characteristics and maximum allowable mass of hybrid star in the context of  $f(Q)$  gravity, Eur. Phys. J. C. 83 (2023) 646.
- <span id="page-17-12"></span>[30] J. C. N de Araujo, H. G. M. Fortes, Compact stars in  $f(Q) = Q + \xi Q^2$ , arXiv:2407.08884  $|gr\text{-}qc$ .
- <span id="page-18-0"></span>[31] S. Mandal, G. Mustafa, Z. Hassan, P.K. Sahoo, A study of anisotropic spheres in  $f(Q)$  gravity with quintessence field, Physics of dark Universe. **35** (2022) 100934.
- [32] A. Errehymy et al, Anisotropic electrically charged stars in  $f(Q)$  symmetric teleparallel gravity, Eur. Phys. J. Plus. 137 (2022) 1311.
- <span id="page-18-2"></span><span id="page-18-1"></span>[33] M. Z. Gul et al, Viable and stable compact stars in  $f(Q)$  theory, Eur. Phys. J. C. 84 (2024) 8.
- [34] T. Kobayashi and K.-I. Maeda, *Relativistic stars in*  $f(R)$  gravity, and absence thereof, *Phys.* Rev. D. 78 (2008) 064019.
- <span id="page-18-3"></span>[35] A. Upadhye, W.e Hu, *The existence of relativistic stars in*  $f(R)$  gravity, *Phys.Rev.D.* 80 (2009) 064002.
- <span id="page-18-4"></span>[36] J. Aftergood and A. DeBenedictis, Matter conditions for regular black holes in  $f(T)$  gravity, Phys. Rev. D. 90 (2014) 124006.
- <span id="page-18-5"></span>[37] A. DeBenedictis and S. Ilijić, Spherically symmetric vacuum in covariant  $f(T) = T + \frac{\alpha}{2}$  $\frac{\alpha}{2}T^2 +$  $OT^{\gamma}$  gravity theory, Phys. Rev. D. 94 (2016) 124025.
- <span id="page-18-6"></span>[38] G. Abbas et al, Anisotropic compact stars in f(G) gravity, Astrophys Space Sci. 357 (2015) 158.
- <span id="page-18-7"></span>[39] A. Malik, M. Ahmad, B. A. Alwan and Z. Naeem, *Bardeen compact stars in modified*  $f(G)$ gravity, Can. J. Phys. 100 (2022) 10.
- <span id="page-18-8"></span>[40] M. F. Shamir and M. Ahmad, *Emerging anisotropic compact stars in*  $f(G,T)$  gravity, *Eur*. Phys. J. C. 77 (2017) 674.
- <span id="page-18-9"></span>[41] M. A. Sadiq, R. Manzoor, and S. Mumtaz, *Collapsing axial system and*  $f(G,T)$  gravity, Int. J. Mod. Phys. D. 31 06 (2022) 2250044.
- <span id="page-18-10"></span>[42] P. Bhar et al, Study on anisotropic stars in the framework of Rastall gravity, Astrophys. Space Sci. **365**(2020) 145.
- <span id="page-18-11"></span>[43] K. P. Das and U. Debnath, Spherically symmetric anisotropic charged neutron stars in  $f(Q, T)$ gravity, Eur. Phys. J. C. 84 (2024) 513.
- <span id="page-18-13"></span><span id="page-18-12"></span>[44] J. B. Jiménez et al, Teleparallel Palatini theories, JCAP 08 (2018) 039.
- [45] S. Pradhan et al, Geometrically deformed charged anisotropic models in  $f(Q,T)$  gravity, arXiv:2408.02051 [gr-qc].
- <span id="page-18-14"></span>[46] M. Z. Gul et al, Physical Analysis of Spherically Stellar Structure in  $f(Q, T)$  Theory, Gen. Relativ. Gravit., 56 (2024) 45.
- <span id="page-19-0"></span>[47] E. D. Barret al, A pulsar in a binary with a compact object in the mass gap between neutron stars and black holes, Science. 383 (2024) 275.
- <span id="page-19-2"></span><span id="page-19-1"></span>[48] L. Herrera, Stability of the isotropic pressure condition, Phys. Rev. D 101 (2020) 104024.
- [49] M. Ruderman, Pulsars: Structure and Dynamics, ANNUAL REVIEW OF ASTRONOMY AND ASTROPHYSICS. 10 (1972) 427.
- [50] R. L. Bower et al, Anisotropic Spheres in General Relativity,Astrophysical Journal 188 (1974) 657.
- [51] H. Bondi,Anisotropic spheres in general relativity, Royal Astronomical Society. 259 (1992) 365-368.
- <span id="page-19-3"></span>[52] L. Herrera, N.O. Santos, Local anisotropy in self-gravitating systems, Physics Reports. 286 (1997) 53.
- <span id="page-19-4"></span>[53] R. Sharma, S. D. Maharaj, A class of relativistic stars with a linear equation of state, Monthly Notices of the Royal Astronomical Society. 375 (2007) 1265–1268.
- <span id="page-19-5"></span>[54] S. K. Maurya, Y. K. Gupta, S. Ray and B. Dayanandan, Anisotropic models for compact stars. Eur. Phys. J. C 75 (2015) 225.
- <span id="page-19-6"></span>[55] S. K. Maurya, A. Banerjee, and S. Hansraj, Role of pressure anisotropy on relativistic compact stars, Phys. Rev. D. 97 (2018) 044022.
- <span id="page-19-7"></span>[56] K. R. Karmarkar, Gravitational metrics of spherically symmetry and class one Proc. Ind. Acad. Sci. A **27** (1948) 56.
- [57] G. Abbas and H. Nazar, Stellar shear-free gravitational collapse with Karmarkar condition in  $f(R)$  gravity, International Journal of Modern Physics A. 34 33 (2019) 950220.
- <span id="page-19-8"></span>[58] M. Sharif and S. Naz, *Gravastars with Karmarkar condition in*  $f(R, T^2)$  gravity, International Journal of Modern Physics D. 31 11 (2022) 2240008.
- <span id="page-19-9"></span>[59] G. Mustafa et al, Study on physical properties and maximum mass limit of Finch-Skea anisotropic model under Karmarkar condition in  $f(Q)$ -gravity,  $arXiv:2312.16866v1$  [gr-qc] 28 Dec 2023.
- <span id="page-19-10"></span>[60] P. C. Vaidya and R. Tikekar, Exact Relativistic Model for a Superdense Star, J. Astrophys. Astr., 3 (1982) 325.
- <span id="page-19-11"></span>[61] T. Gangopadhyay et al, Strange star equation of state fits the refined mass measurement of 12 pulsars and predicts their radii, Mon. Not. R. Astron., 431 (2013) 3216.
- <span id="page-20-0"></span>[62] R. Sharma, A. Ghosh, S. Bhattacharya and S. Das, Anisotropic generalization of Buchdahl bound for specific stellar models, Eur. Phys. J. C. 81 (2021) 527.
- <span id="page-20-3"></span>[63] A. Ridolfi et al, Eight new millisecond pulsars from the first MeerKAT globular cluster census, Mon. Not. R. Astron. Soc. 504 (2021) 1407.
- <span id="page-20-1"></span>[64] J. L. Jonas et al, The MeerKAT Radio Telescope, MeerKAT Science: On the Pathway to the SKA. 001 (PoS 2016).
- <span id="page-20-2"></span>[65] Fernando Camilo, African star joins the radio astronomy firmament, Nat. Astron. 2 (2018) 594.
- <span id="page-20-4"></span>[66] A. Ridolfi, P. C. C. Freire, Y. Gupta and S. M. Ransom, Upgraded Giant Metrewave Radio Telescope timing of NGC 1851A: a possible millisecond pulsar - neutron star system, Mon. Not. R. Astron. Soc.. 490(3) (2019) 3860.
- <span id="page-20-5"></span>[67] A. Ridolfi et al,TRAPUM discovery of 13 new pulsars in NGC 1851 using MeerKAT, Astron. Astrophys. A27 (2022) 664.