

Jackknife Empirical Likelihood Ratio Test for Cauchy Distribution

Avhad Ganesh Vishnu^a, Ananya Lahiri^a, Sudheesh K. Kattumannil^b

^a*Department of Mathematics and Statistics, Indian Institute of Technology, Tirupati, India,*

^b*Statistical Sciences Division, Indian Statistical Institute, Chennai, Tamil Nadu, India.*

Abstract

Heavy-tailed distributions, such as the Cauchy distribution, are acknowledged for providing more accurate models for financial returns, as the normal distribution is deemed insufficient for capturing the significant fluctuations observed in real-world assets. Data sets characterized by outlier sensitivity are critically important in diverse areas, including finance, economics, telecommunications, and signal processing. This article addresses a goodness-of-fit test for the Cauchy distribution. The proposed test utilizes empirical likelihood methods, including the jackknife empirical likelihood (JEL) and adjusted jackknife empirical likelihood (AJEL). Extensive Monte Carlo simulation studies are conducted to evaluate the finite sample performance of the proposed test. The application of the proposed test is illustrated through the analysing two real data sets.

Keywords: Cauchy distribution · Cryptocurrency · U-statistics · Jackknife empirical likelihood · Wilk's theorem.

1. Introduction

The Cauchy distribution is widely used in statistical analysis, particularly within finance and economics, due to its relevance in modeling financial returns and economic variables that exhibit extreme values or outliers. Its heavy-tailed nature makes it a valuable distribution, but the lack of a finite mean and variance introduces specific challenges in statistical modeling and hypothesis testing. Numerous applications of the Cauchy distribution are documented in the literature; for instance, it has been utilized across various fields, such as modeling the behavior of polar and non-polar liquids in porous glasses within the mechanical and electrical theory (Stapf et al. (1996)), addressing physical anthropology and measurement problems (Johnson et al. (1995)), and modeling value-at-risk in risk and financial analysis (Liu et al. (2012)). Thus, a goodness-of-fit test for assessing whether the underlying distribution conforms to a Cauchy form would be extremely useful in practical applications. Recently, based on various characterizations, several authors, including Mahdizadeh and Zamanzade (2017), Mahdizadeh and Zamanzade (2019), Villaseñor and González-Estrada (2021), and Pekgör (2023), have proposed goodness-of-fit tests for the Cauchy distribution.

Email address: avhadgv@gmail.com (Avhad Ganesh Vishnu)

In recent years, empirical likelihood methods have gained popularity in non-parametric statistics for their flexibility and minimal reliance on strong parametric assumptions. The empirical likelihood ratio test has been particularly valued for constructing confidence regions and hypothesis tests. However, it has been observed that traditional empirical likelihood ratio tests may suffer from reduced accuracy and power, especially when applied to small sample sizes or when the underlying distribution deviates from the assumption of normality. To overcome these limitations, Jing et al. (2009) proposed the JEL ratio test as a more robust alternative. The jackknife method, known for reducing bias and variance in statistical estimates, has been employed to enhance the empirical likelihood framework. This enhancement has resulted in improved accuracy of the empirical likelihood ratio test, particularly when dealing with small samples and heavy-tailed distributions such as the Cauchy distribution. In this paper, we propose both a JEL ratio test and an AJEL ratio test for Cauchy distribution.

The test statistic and empirical likelihood methods, JEL and AJEL are introduced in Section 2. The power properties of these tests are evaluated using Monte Carlo simulations, and the results are presented in Section 3. Section 4 illustrates the proposed methods by analysing real data sets. Finally, Section 5 concludes with a summary.

2. Test statistics

A random variable X is said to have a standard Cauchy distribution, denoted by $X \sim \mathcal{C}(0, 1)$, and has a probability density function (p.d.f.)

$$f(x) = \frac{1}{\pi(1+x^2)}, \quad x \in \mathbb{R}, \quad (1)$$

and cumulative distribution function (c.d.f.)

$$F(x) = \frac{1}{2} + \frac{1}{\pi} \tan^{-1}(x), \quad x \in \mathbb{R}. \quad (2)$$

A sequence of independent and identically distributed (*i.i.d.*) continuous realizations X_1, X_2, \dots, X_n of a random variable X with an unknown distribution function (*d.f.*) F . Based on this sample, we are interested in testing the composite null hypothesis as follows:

$$\begin{aligned} H_0 &: \text{the distribution of } X \text{ is } \mathcal{C}(0, 1) \\ H_1 &: \text{the distribution of } X \text{ does not follow } \mathcal{C}(0, 1) \end{aligned} \quad (3)$$

with the support \mathbb{R} .

A new class of tests is introduced, derived from the following characterization of the Cauchy distribution.

Theorem 1. (*Arnold (1979)*) *Let X, X_1 and X_2 be absolutely continuous random variables. Let us define $Y = (X_1 - X_2^{-1})/2$. We recall X has d.f. F and let Y has d.f. G . X and $Y = (X_1 - X_2^{-1})/2$ are identically distributed if, and only if, $X \sim \mathcal{C}(0, 1)$.*

Proof. If X follows a standard Cauchy distribution, the transformation $Y = (X_1 - X_2^{-1})/2$, also following a standard Cauchy distribution. This result follows from the stability properties of the Cauchy distribution under linear transformations, where Y preserves the distributional shape characterized by heavy tails and symmetry about zero. Therefore, Y and X have the same distribution if, and only if, X is standard Cauchy distributed, as discussed in Arnold (1979). \square

This characterization is classified within the category of distributional characterizations based on equality. Constructing tests based on such characterizations has become popular in goodness-of-fit testing because they ensure equivalence across distribution functions, density functions, characteristic functions, and related analytical properties.

To test the hypothesis defined in (3), a measure of departure, denoted as Δ , is defined as follows:

$$\begin{aligned}\Delta &= \int_{-\infty}^{\infty} \left(P\left(\frac{X_1 X_2 - 1}{2X_2} \leq t\right) - P(X_3 < t) \right) dF(t) \\ &= \int_{-\infty}^{\infty} (G(t) - F(t)) dF(t).\end{aligned}$$

The measure Δ represents the integral of the difference between two cumulative distribution functions over the entire real line. For X_1 , X_2 , and X_3 ,

$$\begin{aligned}\int_{-\infty}^{\infty} G(t) dF(t) &= \int_{-\infty}^{\infty} P\left(\frac{X_1 X_2 - 1}{2X_2} \leq t\right) dF(t) \\ &= P\left(\frac{X_1 X_2 - 1}{2X_2} \leq X_3\right) \\ &= E\left(\mathbf{I}\left\{\left(\frac{X_1 X_2 - 1}{2X_2}\right) \leq X_3\right\}\right),\end{aligned}$$

and if F is a continuous d.f. then $\int_{-\infty}^{\infty} F(t) dF(t) = \frac{1}{2}$. Hence

$$\Delta = E\left(\mathbf{I}\left\{\left(\frac{X_1 X_2 - 1}{2X_2}\right) \leq X_3\right\}\right) - \frac{1}{2} = \Delta' - \frac{1}{2} \quad (\text{say}). \quad (4)$$

In view of the characterization, the following empirical U-statistics based test statistic is considered:

$$\Delta_n = \int_{-\infty}^{\infty} (G_n(t) - F_n(t)) dF_n(t), \quad (5)$$

where

$$G_n(t) = \binom{n}{2}^{-1} \sum_{i=1}^n \sum_{j=1; j < i}^n \mathbf{I}\left\{\left(\frac{X_i X_j - 1}{2X_j}\right) \leq t\right\} \quad \text{and} \quad F_n(t) = \frac{1}{n} \sum_{i=1}^n \mathbf{I}\{X_i \leq t\}$$

are U-empirical d.f.'s. Here, $G_n(t)$ represents an empirical distribution function based on pairs of observations, while $F_n(t)$ is the empirical distribution function based on

individual observations. These U-empirical d.f.'s are used to approximate the underlying distribution functions that appear in the definition of Δ . The difference between $G_n(t)$ and $F_n(t)$ quantifies how the data diverge from the hypothesized model, and the test statistic Δ_n is constructed as their weighted integral over the entire range of t .

After integration, an asymptotic equivalence test statistic of Δ_n becomes

$$\widehat{\Delta}_n^* = \binom{n}{3}^{-1} \sum_{i=1}^n \sum_{j=1; j < i}^n \sum_{k=1; k < j}^n \mathbb{I}\left\{\left(\frac{X_i X_j - 1}{2X_j}\right) \leq X_k\right\} - \frac{1}{2}. \quad (6)$$

The estimator $\widehat{\Delta}_n^*$ is unbiased for Δ . The symmetric kernel of $\widehat{\Delta}_n^*$ is given by

$$\begin{aligned} & h(X_1, X_2, X_3) \\ &= \frac{1}{3} \left(\mathbb{I}\left(\frac{X_1 X_2 - 1}{2X_2} \leq X_3\right) + \mathbb{I}\left(\frac{X_1 X_3 - 1}{2X_3} \leq X_2\right) + \mathbb{I}\left(\frac{X_2 X_3 - 1}{2X_3} \leq X_1\right) \right) - \frac{1}{2}. \end{aligned}$$

Next, the asymptotic properties of the test statistics are studied.

Theorem 2. *Let X, X_1, X_2, \dots, X_n be i.i.d. random variables with d.f. F . Assume $E[h^2(X_1, X_2, X_3)|X_1] < \infty$, then as $n \rightarrow \infty$, $\sqrt{n}(\widehat{\Delta}_n^* - \Delta)$ converges in distribution to a normal with mean zero and variance $9\sigma^2$, where σ^2 is given by*

$$\sigma^2 = \text{Var}(K(X)) \quad \text{where, } K(x) = \frac{2}{3}P\left(\frac{xX_2 - 1}{2X_2} \leq X_3\right) + \frac{1}{3}P\left(\frac{X_2 X_3 - 1}{2X_3} \leq x\right). \quad (7)$$

Proof. Let $\widehat{\Delta}_n^* = \widehat{\Delta}'_n - \frac{1}{2}$, where

$$\widehat{\Delta}'_n = \binom{n}{3}^{-1} \sum_{i=1}^n \sum_{j=1; j < i}^n \sum_{k=1; k < j}^n \mathbb{I}\left\{\frac{X_i X_j - 1}{2X_j} \leq X_k\right\}. \quad (8)$$

The estimator $\widehat{\Delta}'_n$ is an unbiased estimator of Δ' . Hence the asymptotic distribution of $\sqrt{n}(\widehat{\Delta}_n^* - \Delta)$ and $\sqrt{n}(\widehat{\Delta}'_n - \Delta')$ are the same. Note that $\widehat{\Delta}'_n$ is a U-statistic with kernel function $h^*(X_1, X_2, X_3) = \mathbb{I}\left(\frac{X_1 X_2 - 1}{2X_2} \leq X_3\right)$ of degree three. Symmetric version of kernel $h^*(\cdot)$ is given by

$$\begin{aligned} & h_1(X_1, X_2, X_3) \\ &= \frac{1}{3} \left(\mathbb{I}\left(\frac{X_1 X_2 - 1}{2X_2} \leq X_3\right) + \mathbb{I}\left(\frac{X_1 X_3 - 1}{2X_3} \leq X_2\right) + \mathbb{I}\left(\frac{X_2 X_3 - 1}{2X_3} \leq X_1\right) \right). \quad (9) \end{aligned}$$

Using the central limit theorem for U-statistics (see Lee (2019), Theorem 1, Page 76), as $n \rightarrow \infty$, $\sqrt{n}(\widehat{\Delta}'_n - \Delta')$ converges to normal with mean zero and variance $9\sigma^2$, where σ^2 is the asymptotic variance of $\widehat{\Delta}'_n$ and is given by

$$\sigma^2 = \text{Var}(E(h_1(X_1, X_2, X_3)|X_1)).$$

Using (9), we find the conditional expectation as

$$K(x) = E(h_1(X_1, X_2, X_3)|X_1 = x) = \frac{2}{3}P\left(\frac{xX_2 - 1}{2X_2} \leq X_3\right) + \frac{1}{3}P\left(\frac{X_2X_3 - 1}{2X_3} \leq x\right).$$

Hence, we obtain the variance expression as specified in the theorem. \square

The following result shows that the asymptotic distribution of the null distribution follows a normal distribution. Note that under H_0 , $\Delta = 0$.

Corollary 1. *Let X be a random variable with a distribution function specified in (2). As $n \rightarrow \infty$, $\sqrt{n}\widehat{\Delta}_n^*$ converges in distribution to a normal with mean zero and variance $9\sigma_0^2$, where σ_0^2 is given by*

$$\sigma_0^2 = \text{Var}(K^*(X)), \quad \text{where } K^*(x) = \frac{2}{3}P\left(\frac{xX_2 - 1}{2X_2} \leq X_3\right) + \frac{1}{3}P(X_1 \leq x). \quad (10)$$

Proof. Under H_0 , using the characterization of Cauchy distribution given in Theorem 1, we obtain

$$P\left(\frac{X_1X_2 - 1}{2X_2} \leq x\right) = P(X_1 \leq x).$$

Now, we know that $\sigma^2 = \text{Var}(E(h(X_1, X_2, X_3)|X_1))$. Hence, from (7), we have

$$K^*(x) = E(h_1(X_1, X_2, X_3)|X_1 = x) = \frac{2}{3}P\left(\frac{xX_2 - 1}{2X_2} \leq X_3\right) + \frac{1}{3}P(X_1 \leq x).$$

from where follows (10). \square

Thus, given Corollary 1, we obtain a test based on normal approximation, and we reject the null hypothesis H_0 against the alternative hypothesis H_1 if

$$\frac{\sqrt{n}|\widehat{\Delta}_n^*|}{\hat{\sigma}_0} \geq Z_{\alpha/2},$$

where $\hat{\sigma}_0$ is a consistent estimator of σ_0 and Z_α is the upper α -percentile point of standard normal distribution. As a result, implementing the test using normal approximation is challenging due to the difficulty in finding a consistent estimator of σ_0^2 . This challenge motivates the development of a nonparametric empirical likelihood-based test. Next, we discuss the JEL and AJEL-based tests for testing Cauchy distribution.

2.1. JEL ratio test

In this subsection, we construct a JEL test based on jackknife pseudo values that is constructed using (6). The jackknife pseudo-values are given by

$$\widehat{J}_i = n\widehat{\Delta}_n^* - (n-1)\widehat{\Delta}_{n-1}^{*(-i)}, \quad i = 1, 2, \dots, n, \quad (11)$$

where the value $\widehat{\Delta}_{n-1}^{*(-i)}$ is the computed using $\widehat{\Delta}_n^*$ based on the $(n-1)$ observations $X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n$. Hence, it can be easily shown that $\widehat{\Delta}_n^* = \frac{1}{n} \sum_{i=1}^n \widehat{J}_i$. Although \widehat{J}_i are typically dependent random variables, they are asymptotically independent under mild conditions (see Shi (1984)). This allows us to use the empirical likelihood method using the Jackknife pseudo value. Let p_i be the probability assigned to each \widehat{J}_i . The JEL ratio evaluated at Δ , becomes

$$\mathcal{R}(\Delta) = \max \left\{ \prod_{i=1}^n np_i : \sum_{i=1}^n p_i = 1, \sum_{i=1}^n p_i \widehat{J}_i = 0 \right\}.$$

By using the standard Lagrange multiplier method, the above maximization is attained at

$$\widehat{p}_i = \frac{1}{n} \frac{1}{(1 + \widehat{\lambda}_1 \widehat{J}_i)}$$

where $\widehat{\lambda}_1$ satisfies

$$\frac{1}{n} \sum_{i=1}^n \frac{\widehat{J}_i}{(1 + \widehat{\lambda}_1 \widehat{J}_i)} = 0.$$

Hence, we obtain

$$\log \mathcal{R}(\Delta) = - \sum_{i=1}^n \log (1 + \widehat{\lambda}_1 \widehat{J}_i).$$

Next, the asymptotic null distribution of the JEL ratio test is determined to construct a critical region for the JEL based test. Following Theorem 2 of Jing et al. (2009), the well known Wilks' theorem regarding the asymptotic null distribution of the likelihood ratio statistic is applied.

Theorem 3. *Assume that*

1. $E[h^2(X_1, X_2, X_3)] < \infty$, and
2. $\sigma_1^2 = \text{Var}[E\{h(X_1, X_2, X_3)|X_1\}] > 0$.

Then, as $n \rightarrow \infty$, $-2 \log \mathcal{R}(\Delta)$ converges in distribution to a chi-squared distribution with one degree of freedom.

In view of Theorem 3, the null hypothesis H_0 is rejected in favor of the alternative hypothesis H_1 at the α level of significance, if

$$-2 \log \mathcal{R}(\Delta) > \chi_{1,\alpha}^2,$$

where $\chi_{1,\alpha}^2$ is the upper α -percentile point of a χ^2 distribution with one degree of freedom.

2.2. AJEL ratio test

The convex hull of $\widehat{J}_i - E(\widehat{J}_i)$ may not contain zero when the sample size is very small. To overcome this difficulty, Chen and Ning (2016) combined the idea of jackknife and adjusted jackknife empirical likelihood and proposed an adjusted jackknife

empirical likelihood (AJEL) ratio test. In the AJEL method, Chen and Ning (2016) proposed incorporating an additional jackknife pseudo-value alongside the existing jackknife pseudo-values given in (11) and defined as

$$\widehat{J}_{n+1} = \frac{-k_n}{n} \sum_{i=1}^n \widehat{J}_i.$$

In the context of AJEL, Chen et al. (2008) suggest the most popular choice of k_n as $\max\{1, \log(n)/2\}$. Hence we consider the same value for k_n . The AJEL is defined as follows,

$$\mathcal{R}(\Delta) = \max \left\{ \prod_{i=1}^{n+1} ((n+1)p_i) : \sum_{i=1}^{n+1} p_i = 1, p_i \geq 0, \sum_{i=1}^{n+1} p_i \widehat{W}_i = 0 \right\},$$

where

$$\widehat{W}_i = \begin{cases} \widehat{J}_i & \text{for } i = 1, 2, \dots, n. \\ -\max\left(1, \frac{1}{2} \log(n)\right) \frac{1}{n} \sum_{i=1}^n \widehat{J}_i & \text{for } i = n+1. \end{cases} \quad (12)$$

Thus, the next theorem established that the asymptotic distribution of the empirical likelihood ratio test statistic follows a chi-squared distribution, analogous to Wilks' theorem.

Theorem 4. *Under the assumptions outlined in Theorem 3, under H_0 , $-2 \log \mathcal{R}(\Delta)$ converges in distribution to χ^2 with one degree of freedom where the log empirical likelihood ratio is obtained as*

$$\log \mathcal{R}(\Delta) = - \sum_{i=1}^{n+1} \log \left(1 + \widehat{\lambda}_2 \widehat{W}_i \right)$$

where $\widehat{\lambda}_2$ satisfies

$$\frac{1}{n+1} \sum_{i=1}^{n+1} \frac{\widehat{W}_i}{1 + \widehat{\lambda}_2 \widehat{W}_i} = 0, \quad \text{and} \quad \widehat{p}_i = \frac{1}{(n+1)} \frac{1}{(1 + \widehat{\lambda}_2 \widehat{W}_i)}.$$

Proof. Using the jackknife pseudo value defined in (12), we obtain $Z_n = \frac{1}{n} \sum_{i=1}^n \widehat{J}_i$. Define $S^2 = \frac{1}{n} \sum_{i=1}^n (\widehat{J}_i)^2$, hence by strong law of large number, we have $S^2 = \sigma^2 + O_p(1)$.

We have $|\widehat{\lambda}_2| = \mathcal{O}_p(1/\sqrt{n})$, when $k_n = O_p(n)$. Consider

$$\begin{aligned} -2 \log \mathcal{R}(\Delta) &= \sum_{i=1}^{n+1} 2 \log(1 + \widehat{\lambda}_2 \widehat{W}_i) \\ &= 2 \sum_{i=1}^{n+1} (\widehat{\lambda}_2 \widehat{W}_i - (\widehat{\lambda}_2 \widehat{W}_i)^2/2) + O_p(1) \\ &= 2n\widehat{\lambda}_2 Z_n - nS\widehat{\lambda}_2^2 + O_p(1) \\ &= \frac{nZ_n^2}{S^2} + O_p(1) \end{aligned}$$

where the preceding identity is established because the $(n+1)$ th term in the summation is $k_n O_p(n^{-3/2}) = O_p(n) O_p(n^{-3/2}) = O_p(1)$. Hence by Slutsky's theorem, as $n \rightarrow \infty$,

$$-2 \log \mathcal{R}(\Delta) = \frac{nZ_n^2}{S^2} \xrightarrow{d} \chi_1^2.$$

□

Thus in view of Theorem 4, we reject the null hypothesis H_0 against the alternative hypothesis H_1 at a α level of significance if

$$-2 \log \mathcal{R}(\Delta) > \chi_{1,\alpha}^2.$$

3. Simulation study

In this section, a comprehensive simulation study is presented to evaluate the empirical power of the new tests relative to established methods. All computations for this and the subsequent sections were carried out using the statistical software R. The simulations were conducted at significance level $\alpha = 0.05$ and for sample sizes $n = 20, 40, 60, 80$, and 100.

To evaluate the empirical power of the newly proposed tests in comparison with the following established tests for the Cauchy distribution:

- The classical Kolmogorov-Smirnov (KS_n) test:

$$KS_n = \max \left[\max_{1 \leq i \leq n} \left\{ \frac{i}{n} - F(x), F(x) - \frac{i-1}{n} \right\} \right],$$

where $F(x)$ is the c.d.f. defined in (2).

- Cramér-von Mises (CM_n) test:

$$CM_n = \sum_{i=1}^n \left(F(X_{(i)}) - \frac{i-0.5}{n} \right)^2 + \frac{1}{12n}.$$

- Modified Anderson-Darling (MA_n) test:

$$MA_n = \frac{-2}{n} \sum_{i=1}^n \left[(i - 0.5) \log\{F(X_{(i)})\} + (n - i + 0.5) \log\{1 - F(X_{(i)})\} \right] - n,$$

where $X_{(\cdot)}$ denotes the order statistics.

- Likelihood ratio based, Zhang (2002) tests:

$$\begin{aligned} ZA_n &= - \sum_{i=1}^n \left[\frac{\log\{F(X_{(i)})\}}{n - i + 0.5} + \frac{\log\{1 - F(X_{(i)})\}}{i - 0.5} \right], \\ ZB_n &= \sum_{i=1}^n \left[\log \left\{ \frac{1/F(X_{(i)}) - 1}{(n - 0.5)/(i - 0.75) - 1} \right\} \right]^2, \\ ZC_n &= \max_{1 \leq i \leq n} \left[(i - 0.5) \log \left\{ \frac{i - 0.5}{nF(X_{(i)})} \right\} \right. \\ &\quad \left. + (n - i + 0.5) \log \left\{ \frac{n - i + 0.5}{n(1 - F(X_{(i)}))} \right\} \right]. \end{aligned}$$

- Gürtler and Henze (2000) empirical characteristic function based test:

$$D_{n,\lambda} = \frac{2}{n} \sum_{i,j=1}^n \frac{\lambda}{\lambda^2 + (X_i - X_j)^2} - 4 \sum_{i=1}^n \frac{1 + \lambda}{(1 + \lambda)^2 + X_i^2} + \frac{2n}{2 + \lambda},$$

where $\lambda > 0$ be the weighting parameter. Based on the Gürtler and Henze (2000) simulation results, using $\lambda = 5$ proves to be effective in practice, leading to a robust test.

- Kullback (1997) distance (KL_n) test: The Kullback-Leibler (KL) distance is the most widely used information criterion for evaluating model discrepancies. Mahdizadeh and Zamanzade (2017) suggested a test statistic

$$KL_n = \exp \left\{ - HV_{n,m} - \frac{1}{n} \sum_{i=1}^n \log(f(X_i)) \right\},$$

where $m = n/2$ is the optimal window size mentioned in Mahdizadeh and Zamanzade (2019), $f(x)$ be the p.d.f defined in (1) and

$$HV_{n,m} = \frac{1}{n} \sum_{i=1}^n \log \left\{ \frac{n}{2m} (X_{(i+m)} - X_{(i-m)}) \right\}.$$

- Entropy estimator based test: Recently Mahdizadeh and Zamanzade (2019) proposed six tests based on entropy measure. In our simulation study, we consider the test which gives higher power than other tests. Hence we consider the test

given by

$$HE_{n,m} = \frac{1}{n} \sum_{i=1}^n \log \left\{ \frac{n}{c_i m} (Y_{(i+m)} - Y_{(i-m)}) \right\},$$

where

$$c_i = \begin{cases} 1 + \frac{1+i}{m} - \frac{i}{m^2} & 1 \leq i \leq m, \\ 2 & m+1 \leq i \leq n-m, \\ 1 + \frac{n-i}{m+1} & n-m+1 \leq i \leq n \end{cases}$$

and the $Y_{(i)}$'s are

$$\begin{cases} Y_{(i-m)} = p + \frac{i-1}{m} (X_{(1)} - p) & 1 \leq i \leq m, \\ Y_{(i)} = X_{(1)} & m+1 \leq i \leq n-m, \\ Y_{(i+m)} = q + \frac{n-i}{m} (q - X_{(n)}) & n-m+1 \leq i \leq n \end{cases}$$

where the constants p and q are to be determined such that $P(p \leq X \leq q) \approx 1$ and $m = n/2$ is the optimal window size.

First, we find empirical Type-I error. For this purpose we generate observations from standard Cauchy distribution. The empirical Type-I error rates for the tests based on JEL and AJEL are computed using 10,000 replications. The proportion of times the null hypothesis is rejected at the 5% significance level is assessed. The results, presented in Table 1, show that the empirical Type-I error rates converge to the specified significance level α as the sample size increases. Next, the empirical power of the proposed JEL and AJEL tests is assessed by examining their performance against the alternative distributions considered below.

- Student's t distribution with r degrees of freedom: t_r
- Normal distribution with parameters μ and σ^2 : $N(\mu, \sigma^2)$
- Gamma distribution with parameters α and β : $G(\alpha, \beta)$
- Laplace distribution with parameters μ and σ^2 : $L(\mu, \sigma^2)$
- Beta distribution with parameters α and β : $B(\alpha, \beta)$
- Uniform distribution with parameters α and β : $U(\alpha, \beta)$

The alternative distributions considered are t_3 , $N(0, 1)$, $G(2, 1)$, $L(0, 1)$, $B(2, 2)$, and $U(0, 1)$. Samples of sizes $n = 20, 40, 60, 80$, and 100 are generated from each of these distributions. The empirical power of the tests based on JEL and AJEL is calculated using 10,000 replications. The proportion of times the null hypothesis is rejected at the 5% significance level is computed. The results are presented in Tables 2. From Tables

2, we observed that the proposed test has very good power, even for a relatively small sample size.

Compared to classical tests, the JEL and AJEL tests show better performance. classical tests often struggle with small sample sizes or specific types of data, which can make them less reliable in some cases. However, the JEL and AJEL tests are more dependable and effective across a wide range of situations. They handle different sample sizes and various data distributions well, making them a versatile and robust choice for many different scenarios. In real-world situations, where conditions can vary greatly, the JEL and AJEL tests provide a more stable and dependable method for detecting significant differences.

Table 1: Empirical sizes at 0.05 level of significance

n	JEL	AJEL	KS_n	CM_n	MA_n	ZA_n	ZB_n	ZC_n	$D_{n,\lambda}$	KL_n	$HE_{n,m}$
20	0.044	0.030	0.057	0.048	0.047	0.048	0.045	0.039	0.053	0.035	0.035
40	0.032	0.029	0.044	0.051	0.046	0.061	0.060	0.059	0.045	0.054	0.040
60	0.047	0.035	0.069	0.078	0.073	0.047	0.047	0.043	0.058	0.044	0.042
80	0.049	0.047	0.067	0.061	0.062	0.064	0.039	0.041	0.055	0.064	0.044
100	0.051	0.049	0.078	0.069	0.091	0.058	0.059	0.068	0.058	0.056	0.038

4. Data analysis

Next, we analyze two real-world data sets employing all the proposed methodologies.

I. German Stock Index Data

Stock market return distributions frequently exhibit heavy tails, a feature not commonly associated with the normal distribution. Consequently, it is more appropriate to analyze stock market return data using Cauchy distributions, which are known for their ability to model data with heavy tails. In this study, we examine the returns of the closing prices of the German Stock Index using the Cauchy distribution. The DAX (Deutscher Aktienindex) is a stock market index that includes the 30 largest blue-chip companies listed on the Frankfurt Stock Exchange, representing a significant segment of the German equity market.

In this study, we apply goodness-of-fit tests to a dataset consisting of 30 returns of closing prices from the DAX (Deutscher Aktienindex). These data points were collected daily, starting from January 1, 1991, while excluding weekends and public holidays. The dataset, rounded to seven decimal places, is provided in Table 3 and sourced from the `datasets` package in the R statistical software. Figure 1 displays the Cauchy Q-Q plot, accompanied by a histogram with an overlaid Cauchy density function.

The values of JEL test statistic yield 0.3206 with a p-value 0.5711 and for AJEL test statistics is 0.3078 with a p-value 0.5613, whereas all other statistics are computed and presented in Table 9 of Mahdizadeh and Zamanzade (2019). For each test performed, the null hypothesis that the data are drawn from a Cauchy distribution cannot be rejected at the 0.05 significance level.

Table 2: Empirical power at 0.05 level of significance

Alt.	n	JEL	AJEL	KS_n	CM_n	MA_n	ZA_n	ZB_n	ZC_n	$D_{n,\lambda}$	KL_n	$HE_{n,m}$
t_3	20	0.553	0.482	0.028	0.035	0.019	0.006	0.001	0.018	0.006	0.099	0.043
	40	0.750	0.717	0.002	0.001	0.002	0.046	0.019	0.169	0.033	0.527	0.473
	60	0.865	0.852	0.003	0.004	0.007	0.424	0.198	0.398	0.246	0.741	0.813
	80	0.953	0.944	0.001	0.001	0.006	0.594	0.445	0.600	0.384	0.887	0.893
	100	0.968	0.965	0.006	0.002	0.019	0.862	0.705	0.589	0.562	0.909	0.911
N(0,1)	20	0.512	0.447	0.002	0.001	0.003	0.008	0.008	0.030	0.005	0.193	0.134
	40	0.686	0.644	0.001	0.002	0.003	0.549	0.299	0.509	0.398	0.798	0.686
	60	0.836	0.817	0.001	0.004	0.027	0.799	0.853	0.776	0.841	0.887	0.764
	80	0.923	0.916	0.005	0.010	0.039	0.830	0.910	0.807	0.908	0.908	0.884
	100	0.964	0.960	0.002	0.005	0.076	0.919	0.948	0.923	0.938	0.934	0.901
G(2,1)	20	0.718	0.484	0.010	0.016	0.015	0.077	0.021	0.009	0.023	0.168	0.091
	40	1.000	1.000	0.054	0.247	0.124	1.000	0.535	0.042	0.055	0.917	0.910
	60	1.000	1.000	0.193	0.600	0.383	1.000	0.960	0.277	0.322	0.990	0.995
	80	1.000	1.000	0.341	0.809	0.592	1.000	0.999	0.663	0.456	1.000	1.000
	100	1.000	1.000	0.476	0.955	0.794	1.000	1.000	0.878	0.679	1.000	1.000
L(0,1)	20	0.540	0.477	0.022	0.011	0.009	0.005	0.006	0.009	0.006	0.066	0.023
	40	0.742	0.720	0.006	0.002	0.007	0.051	0.052	0.052	0.073	0.681	0.402
	60	0.885	0.871	0.002	0.002	0.004	0.336	0.196	0.346	0.244	0.729	0.836
	80	0.932	0.924	0.002	0.004	0.009	0.800	0.390	0.781	0.400	0.887	0.902
	100	0.974	0.970	0.001	0.002	0.013	0.942	0.684	0.879	0.660	0.948	0.967
B(2,2)	20	1.000	1.000	0.020	0.033	0.050	0.229	0.170	0.013	0.013	0.514	0.224
	40	1.000	1.000	0.149	0.110	0.216	0.749	0.971	0.809	0.194	0.879	0.768
	60	1.000	1.000	0.706	0.668	0.794	0.939	1.000	0.990	0.964	0.959	0.961
	80	1.000	1.000	1.000	0.959	0.991	1.000	1.000	1.000	1.000	1.000	1.000
	100	1.000	1.000	1.000	0.997	1.000	1.000	1.000	1.000	1.000	1.000	1.000
U(0,1)	20	1.000	1.000	0.006	0.012	0.014	0.228	0.132	0.018	0.006	0.918	0.635
	40	1.000	1.000	0.809	0.226	0.246	0.820	0.910	0.801	0.615	1.000	0.920
	60	1.000	1.000	1.000	0.739	0.882	1.000	1.000	1.000	0.998	1.000	1.000
	80	1.000	1.000	1.000	0.912	0.962	1.000	1.000	1.000	1.000	1.000	1.000
	100	1.000	1.000	1.000	0.998	0.998	1.000	1.000	1.000	1.000	1.000	1.000

II. Cryptocurrency Data

Now, we utilize tests for the Cauchy distribution to analyze the log returns of EOS cryptocurrency. Cauchy distribution is found to be a comparably good model for such data sets, see Ebner et al. (2024). The dataset analyzed in this study comprises daily closing prices of EOS in USD, spanning from December 25, 2020, to July 07, 2024.

Table 3: Closing prices of DAX

0.0011848	-0.0057591	-0.0051393	-0.0051781	0.0020043	0.0017787
0.0026787	-0.0066238	-0.0047866	-0.0052497	0.0004985	0.0068006
0.0016206	0.0007411	-0.0005060	0.0020992	-0.0056005	0.0110844
-0.0009192	0.0019014	-0.0042364	0.0146814	-0.0002242	0.0024545
-0.0003083	-0.0917876	0.0149552	0.0520705	0.0117482	0.0087458




Figure 1: Q-Q plot and histogram for German Stock Index (DAX) Data

The data were accessible from CryptoDataDownload (www.cryptodatadownload.com/data). For the analysis, we utilize the daily returns $r_d = (P_d - P_{d-1})/P_{d-1}$ of the closing prices, where P_d be the closing price of the d -th day. Figure 2 displays the histogram of the dataset along with the fitted Cauchy distribution densities. Visually, the Cauchy model appears to be a reasonable approximation. The calculated value of the JEL statistic is 0.2084 with a corresponding p-value of 0.6479, while the AJEL test statistic value is 0.1192 with a p-value 0.6464. Based on JEL and AJEL values, we fail to reject the null hypothesis at the 0.05 significance level.

5. Summary

In this paper, jackknife and adjusted jackknife likelihood ratio tests are proposed, utilizing U-statistic theory and based on a straightforward characterization of the Cauchy distribution. Drawing inspiration from the work of Jing et al. (2009), the paper derives the asymptotic distribution of the empirical likelihood ratio test statistic based on the JEL method, analogously to Wilks' theorem. This result is then extended to the AJEL method. A simulation study is conducted to evaluate the efficacy of the proposed test




Figure 2: Q-Q plot and histogram for daily closing prices of Bitcoin data

procedures. Additionally, the methodologies are applied to real data sets, specifically the daily closing prices of the German stock index and EOS cryptocurrency. The data analysis shows that both data sets exhibit heavy-tailed behavior, suggesting that the Cauchy distribution is suitable for both data sets.

References

- Arnold, B. C. (1979). Some characterizations of the Cauchy distribution. *Australian Journal of Statistics*, 21(2):166–169.
- Chen, J., Variyath, A. M., and Abraham, B. (2008). Adjusted empirical likelihood and its properties. *Journal of Computational and Graphical Statistics*, 17(2):426–443.
- Chen, Y.-J. and Ning, W. (2016). Adjusted jackknife empirical likelihood. *arXiv preprint arXiv:1603.04093*.
- Ebner, B., Eid, L., and Klar, B. (2024). Cauchy or not Cauchy? new goodness-of-fit tests for the Cauchy distribution. *Statistical Papers*, 65(1):45–78.
- Gürtler, N. and Henze, N. (2000). Goodness-of-fit tests for the Cauchy distribution based on the empirical characteristic function. *Annals of the Institute of Statistical Mathematics*, 52(2):267–286.
- Jing, B., Yuan, J., and Zhou, W. (2009). Jackknife empirical likelihood. *Journal of the American Statistical Association*, 104(487):1224–1232.

- Johnson, N. L., Kotz, S., and Balakrishnan, N. (1995). *Continuous univariate distributions, Volume 2*. John Wiley & Sons.
- Kullback, S. (1997). *Information theory and statistics*. Courier Corporation.
- Lee, A. J. (2019). *U-statistics: Theory and Practice*. Routledge.
- Liu, T., Zhang, P., Dai, W.-S., and Xie, M. (2012). An intermediate distribution between Gaussian and Cauchy distributions. *Physica A: Statistical Mechanics and its Applications*, 391(22):5411–5421.
- Mahdizadeh, M. and Zamanzade, E. (2017). New goodness of fit tests for the Cauchy distribution. *Journal of Applied Statistics*, 44(6):1106–1121.
- Mahdizadeh, M. and Zamanzade, E. (2019). Goodness-of-fit testing for the Cauchy distribution with application to financial modeling. *Journal of King Saud University-Science*, 31(4):1167–1174.
- Pekgör, A. (2023). A novel goodness-of-fit test for Cauchy distribution. *Journal of Mathematics*, 2023(1):9200213.
- Shi, X. (1984). The approximate independence of jackknife pseudo-values and the bootstrap methods. *Journal of Wuhan Institute Hydra-Electric Engineering*, 2:83–90.
- Stapf, S., Kimmich, R., Seitter, R.-O., Maklakov, A., and Skirda, V. (1996). Proton and deuteron field-cycling nmr relaxometry of liquids confined in porous glasses. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 115:107–114.
- Villaseñor, J. and González-Estrada, E. (2021). Goodness-of-fit tests for Cauchy distributions using data transformations. *Advances in Statistics-Theory and Applications: Honoring the Contributions of Barry C. Arnold in Statistical Science*, pages 271–282.
- Zhang, J. (2002). Powerful goodness-of-fit tests based on the likelihood ratio. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 64(2):281–294.