# Lipschitz inextendibility of weak null singularities from curvature blow-up

Jan Sbierski\*

September 30, 2024

#### Abstract

We prove the  $C_{loc}^{0,1}$ -inextendibility of weak null singularities without any symmetry assumptions. The proof introduces a new strategy to infer  $C_{loc}^{0,1}$ -inextendibility from the blow-up of curvature. The assumed blow-up is expected to be satisfied for weak null singularities in the interior of generic rotating black holes. Thus, we expect the result presented here to directly contribute to the resolution of the  $C_{loc}^{0,1}$ -formulation of the strong cosmic censorship conjecture in a neighbourhood of subextremal Kerr.

### Contents

1	Introduction		
	1.1	Preliminaries	3
<b>2</b>	The spacetimes considered and gauge choice		
	2.1	The class of spacetimes	4
	2.2	Frame and Ricci coefficients	4
3	The	$C_{ m loc}^{0,1}$ -inextendibility of weak null singularities	5
	3.1	Auxiliary results	5
	3.2	The main theorem	8
Re	References		

## 1 Introduction

It is known from the seminal work of Dafermos and Luk [4] that dynamical vacuum black holes, which settle down to a subextremal Kerr black hole in the exterior, also have a Cauchy horizon in the black hole interior to which the Lorentzian metric extends continuously, see Figure 1. Results for the linearised Einstein equations [14], [23], [10] indicate that generically curvature blows up at the Cauchy horizon, turning it into a so-called *weak null singularity*, terminating the classical time-evolution. This scenario is a special case of the much broader strong cosmic censorship conjecture, according to which the maximal globally hyperbolic development of generic asymptotically flat initial data is inextendible as a suitably regular Lorentzian manifold. The main

<sup>\*</sup>School of Mathematics and Maxwell Institute for Mathematical Sciences, University of Edinburgh, James Clerk Maxwell Building, Peter Guthrie Tait Road, Edinburgh, EH9 3FD, United Kingdom

result of this paper, Theorem 3.13, shows, that the dynamical black hole interiors of [4], supplemented by a curvature blow-up assumption, are locally Lipschitz inextendible across the weak null singularity. The assumed blow-up of curvature is expected to be generically satisfied, see the discussion in Remark 3.16. Thus, we expect the main theorem to provide one of the ingredients for the resolution of the  $C_{loc}^{0,1}$ -formulation of the strong cosmic censorship conjecture in a neighbourhood of subextremal Kerr initial data.

The result presented in this paper is the first low-regularity<sup>1</sup> inextendibility result for singular spacetimes which are expected to arise from an open set of initial data. Existing results are either restricted to spacetimes with symmetries [21], [5], [22], [24], [12], [15], with warped product structure [22], or are conditional results  $[5], [3], [8]^2$ 



Figure 1: A Penrose-style diagram of a dynamical vacuum black hole settling down to a sub-extremal Kerr black hole in the exterior I. Very little is known about the singularity structure in the interior II beyond the weak null singularity.

We give a rough outline of the idea of the proof: assuming a  $C_{\text{loc}}^{0,1}$ -extension across a weak null singularity exists. The first step is to use upper bounds on the connection coefficients (in the gauge used in [4]) to show that the  $C^1$ -structure of the  $C^{0,1}_{loc}$ -extension has to be locally equivalent to the  $C^1$ -structure of the  $C^0$ -extension constructed in [4]. Indeed, this local uniqueness problem for  $C_{\rm loc}^{0,1}$ -extensions was studied in the precursor [25] to this paper and here, we verify that the assumptions of [25] are met so that we can conclude the local equivalence of  $C^1$ -structures. An important consequence is that a vector field which extends continuously in the  $C^0$ -extension of [4], also extends continuously in the assumed  $C^{0,1}_{loc}$ -extension.

In the second step we use the assumed blow-up of curvature: contract curvature with four vector fields which extend continuously in the  $C^0$ -extension of [4] and integrate over a sequence of compact regions asymptotically touching the weak null singularity. We assume that this sequence of integrals diverges. On the other hand, using the coordinates of the  $C_{loc}^{0,1}$ -extension, in which the Christoffel symbols remain uniformly bounded, we rewrite these integrals over curvature in terms of these Christoffel symbols. By virtue of the equivalence of the  $C^1$ -structures, the vector fields in this integral are also regular with respect to those coordinates.<sup>3</sup> One can now use Stokes' theorem to convert the integral over first derivatives of Christoffel symboles, which appears in curvature, into a boundary integral over Christoffel symbols. The expression obtained is then manifestly finite and uniformly bounded along the sequence, thus giving the desired contradiction.

<sup>&</sup>lt;sup>1</sup>By this we mean below  $C^2$  (or  $C_{loc}^{1,1}$ ). <sup>2</sup>An exception are the  $C^0$ -inextendibility results obtained in [6], [16] (see also [7]) which are based on geodesic completeness, do not require any symmetries, and do apply for example to the spacetimes arising from small perturbations of Minkowski initial data [2]. However, these inextendibility results do not concern singularities.

<sup>&</sup>lt;sup>3</sup>Indeed, we need that these vector fields are  $C^1$ , but the equivalence of the  $C^1$ -structures only gives that the vector fields are continuous. This is the reason for an additional smoothing procedure, which, as a consequence, requires that the integrals over curvature also diverge if we insert vector fields which are  $\varepsilon$ -close to the original ones.

The proof given here establishes for the first time a connection between curvature blow-up and lowregularity inextendibility.<sup>4</sup> The subtleties of such a connection were demonstrated in [17]. On the other hand, the  $C_{loc}^{0,1}$ -inextendibility of *spherically symmetric* weak null singularities was shown in [22] using a method based on direct computation of holonomy. While this method is in principle also applicable to weak null singularities without symmetry assumptions as considered in this paper, it would require, in addition to the control of the  $C^1$ -structure, (pointwise) lower bounds on one of the connection coefficients (namely  $\chi$ , see Section 2.2). The new approach presented here is more parsimonious in the sense that only an integrated lower bound on curvature is needed, which is easier to establish analytically.

Finally, let us remark that although the paper demonstrates a method to capture geometrically the *blow-up* of the Levi-Civita connection, the problem of how to capture geometrically the *rate of the blow up*, i.e., an inextendibility statement in  $C^0 \cap W_{\text{loc}}^{1,p}$ , 1 , remains a challenging open problem (but see Remark 3.30).

#### Acknowledgements

I acknowledge support through the Royal Society University Research Fellowship URF\R1\211216.

#### **1.1** Preliminaries

We begin by recalling two fundamental definitions and the result which constitutes the starting point for most low-regularity inextendibility results.

**Definition 1.1.** Let (M, g) be a Lorentzian manifold<sup>5</sup> and let  $\Gamma$  be a regularity class, for example  $\Gamma = C^k$ with  $k \in \mathbb{N} \cup \{\infty\}$  or  $\Gamma = C_{\text{loc}}^{0,1}$ . A  $\Gamma$ -extension of (M, g) consists of a smooth isometric embedding  $\iota : M \hookrightarrow \tilde{M}$ of M into a Lorentzian manifold  $(\tilde{M}, \tilde{g})$  of the same dimension as M where  $\tilde{g}$  is  $\Gamma$ -regular and such that  $\partial \iota(M) \subseteq \tilde{M}$  is non-empty.

If (M,g) admits a  $\Gamma$ -extension, then we say that (M,g) is  $\Gamma$ -extendible, otherwise we say (M,g) is  $\Gamma$ -inextendible.

**Definition 1.2.** Let (M,g) be a time-oriented Lorentzian manifold with  $g \in C^0$  and let  $\iota : M \hookrightarrow \tilde{M}$  be a  $C^0$ -extension of M. The future boundary of M is the set  $\partial^+\iota(M)$  consisting of all points  $\tilde{p} \in \tilde{M}$  such that there exists a smooth timelike curve  $\tilde{\gamma} : [-1,0] \to \tilde{M}$  such that  $\operatorname{Im}(\tilde{\gamma}|_{[-1,0]}) \subseteq \iota(M), \ \tilde{\gamma}(0) = \tilde{p} \in \partial\iota(M)$ , and  $\iota^{-1} \circ \tilde{\gamma}|_{[-1,0]}$  is future directed in M.

The next proposition is found in [20], Proposition 2.2.

**Proposition 1.3.** Let  $\iota : M \hookrightarrow \tilde{M}$  be a  $C^0$ -extension of a time-oriented globally hyperbolic Lorentzian manifold (M, g) with  $g \in C^0$  and with Cauchy hypersurface  $\Sigma$  – and let  $\tilde{p} \in \partial^+ \iota(M)$ . For every  $\delta > 0$  there exists a chart  $\tilde{\varphi} : \tilde{U} \to (-\varepsilon_0, \varepsilon_0) \times (-\varepsilon_1, \varepsilon_1)^d$ ,  $\varepsilon_0, \varepsilon_1 > 0$  with the following properties

- i)  $\tilde{p} \in \tilde{U}$  and  $\tilde{\varphi}(p) = (0, \dots, 0)$
- *ii*)  $|\tilde{g}_{\mu\nu} \eta_{\mu\nu}| < \delta$ , where  $\eta_{\mu\nu} = \text{diag}(-1, 1, ..., 1)$
- iii) There exists a Lipschitz continuous function  $f: (-\varepsilon_1, \varepsilon_1)^d \to (-\varepsilon_0, \varepsilon_0)$  with the following property:

$$\{(x_0,\underline{x}) \in (-\varepsilon_0,\varepsilon_0) \times (-\varepsilon_1,\varepsilon_1)^d \mid x_0 < f(\underline{x})\} \subseteq \tilde{\varphi}\big(\iota\big(I^+(\Sigma,M)\big) \cap \tilde{U}\big)$$
(1.4)

<sup>&</sup>lt;sup>4</sup>For the reverse direction, namely to prove extendibility under curvature bounds, we refer the reader to [18] and references therein. For synthetic curvature blow-up *in* extensions, see [9].

<sup>&</sup>lt;sup>5</sup>All differentiable manifolds are assumed to be smooth. The metric g in this definition is at least  $\Gamma$ -regular.

and

$$\{(x_0,\underline{x}) \in (-\varepsilon_0,\varepsilon_0) \times (-\varepsilon_1,\varepsilon_1)^d \mid x_0 = f(\underline{x})\} \subseteq \tilde{\varphi}\big(\partial^+\iota(M) \cap \tilde{U}\big) . \tag{1.5}$$

Moreover, the set on the left hand side of (1.5), i.e. the graph of f, is achronal<sup>6</sup> in  $(-\varepsilon_0, \varepsilon_0) \times (-\varepsilon_1, \varepsilon_1)^d$ .

Given a future boundary point  $\tilde{p}$ , we call a chart as above a *future boundary chart*. We also denote by  $\tilde{U}_{\leq}$  the (image under  $\tilde{\varphi}^{-1}$  of) left hand side of (1.4). The region  $\{x_0 \leq f(\underline{x})\}$  including the graph is denoted by  $\tilde{U}_{\leq}$ .

# 2 The spacetimes considered and gauge choice

#### 2.1 The class of spacetimes

We consider Lorentzian manifolds (M, g) in a double null gauge: assume  $M = (-1, 1) \times (-1, 0) \times \mathbb{S}^2$  with coordinates  $(u, \underline{u})$  on the first two factors and denote with  $(\theta^1, \theta^2) = \theta^A$ , A = 1, 2, an arbitrary set of smooth coordinates on  $\mathbb{S}^2$ . Let  $\gamma(u, \underline{u})$  be a Riemannian metric on  $\mathbb{S}^2$  for each  $(u, \underline{u}) \in (-1, 1) \times (-1, 0)$  such that  $\gamma_{AB}(u, \underline{u}, \theta^1, \theta^2)$  are at least  $C^2$ -regular functions of  $(u, \underline{u}, \theta^1, \theta^2)$  for each  $A, B \in \{1, 2\}$ . Moreover, let  $b(u, \underline{u})$ be a vector field on  $\mathbb{S}^2$  for each  $(u, \underline{u}) \in (-1, 1) \times (-1, 0)$  such that  $b^A(u, \underline{u}, \theta^1, \theta^2)$  are at least  $C^2$ -regular functions of  $(u, \underline{u}, \theta^1, \theta^2)$  for A = 1, 2. Finally, let  $\Omega : (-1, 1) \times (-1, 0) \times \mathbb{S}^2 \to (0, \infty)$  be a positive function that is at least  $C^2$ -regular. The metric g is then assumed to be given in local coordinates  $(u, \underline{u}, \theta^A)$  by

$$g = -2\Omega^2 (du \otimes d\underline{u} + d\underline{u} \otimes du) + \gamma_{AB} (d\theta^A - b^A d\underline{u}) \otimes (d\theta^B - b^B d\underline{u}) .$$

$$(2.1)$$

By assumption we have  $g \in C^2$ . Moreover, we define a time-orientation on (M, g) by stipulating that  $\partial_u$  is future-directed null. Note that (M, g) is globally hyperbolic. Indeed,  $t := u + 2\underline{u}$  is a time function and  $\{t = -1\}$  is a Cauchy hypersurface.

The double null gauge used here is the one used in [4]. However, here we only consider a finite chunk (M, g) of the interior of the dynamical black hole constructed in [4], cf. Figure 1. This is sufficient for our purposes, since the problem of inextendibility reduces to local problems, see Remark 3.17, and each of those finite chunks (M, g) of the weak null singularity is expected to satisfy the assumptions of Theorem 3.13.

#### 2.2 Frame and Ricci coefficients

The only non-vanishing components of the inverse metric  $g^{-1}$  in the  $(u, \underline{u}, \theta^A)$  coordinates are

$$(g^{-1})^{AB} = (\gamma^{-1})^{AB}$$
,  $(g^{-1})^{u\underline{u}} = -\frac{1}{2\Omega^2}$ ,  $(g^{-1})^{uA} = -\frac{1}{2\Omega^2}b^A$ .

Note that u and  $\underline{u}$  satisfy  $g^{-1}(du, du) = 0 = g^{-1}(d\underline{u}, d\underline{u})$ .

We define two null vectors by

$$e_3 := \frac{\partial}{\partial u}$$
 and  $e_4 := \frac{1}{\Omega^2} \left( \frac{\partial}{\partial \underline{u}} + b^A \frac{\partial}{\partial \theta^A} \right)$ .

Note that we have  $g(e_3, e_4) = -2$  and that  $e_3 = -2\Omega^2(d\underline{u})^{\sharp}$  and  $e_4 = -2(du)^{\sharp}$ , where  $\sharp$  denotes the raising of an index with  $g^{-1}$ .

<sup>&</sup>lt;sup>6</sup>With respect to *smooth* timelike curves.

We introduce the following Ricci coefficients with respect to the double null frame  $e_3, e_4, e_1, e_2$ , where  $e_A := \frac{\partial}{\partial \theta^A}, A = 1, 2$ :

$$\begin{split} \chi_{AB} &:= g(\nabla_{e_A} e_4, e_B) \;, & & \underline{\chi}_{AB} &:= g(\nabla_{e_A} e_3, e_B) \;, \\ \eta_A &:= -\frac{1}{2} g(\nabla_{e_3} e_A, e_4) \;, & & \underline{\eta}_A &:= -\frac{1}{2} g(\nabla_{e_4} e_A, e_3) \;, \\ & & \underline{\omega} &:= -\frac{1}{4} g(\nabla_{e_3} e_4, e_3) \;. \end{split}$$

Note that we have

$$4\omega := -g(\nabla_{e_4}e_3, e_4) = g(e_3, \nabla_{e_4}e_4) = 0$$
(2.2)

and

$$2\zeta_A := g(\nabla_{e_A} e_4, e_3) = -g(e_4, \nabla_{e_A} e_3) = -g(e_4, \nabla_{e_3} e_A) = 2\eta_A$$
(2.3)

by our choice of double null frame.

Moreover, we define the pointwise norms  $|\chi|_{\gamma} = \sqrt{(\gamma^{-1})^{AB}(\gamma^{-1})^{CD}\chi_{AC}\chi_{BD}}$  and  $|\eta|_{\gamma} = \sqrt{(\gamma^{-1})^{AB}\eta_{A}\eta_{B}}$ ; and similarly for  $\chi$  and  $\eta$ .

We also introduce the vector field  $L := \Omega^2 e_4 = \frac{\partial}{\partial \underline{u}} + b^A \frac{\partial}{\partial \theta^A}$  and denote its integral curve starting at  $(u_0, \underline{u}_0, \theta_0) \in M$  by  $s \mapsto \sigma_{(u_0, \underline{u}_0, \theta_0)}(s) = (u_0, \underline{u} + s, \sigma^{\theta}_{(u_0, \underline{u}_0, \theta_0)}(s))$ . The corresponding flow starting from the hypersurface  $\{\underline{u} = -\frac{1}{2}\}$  we denote by

$$\Phi: M \cap \{\underline{u} = -\frac{1}{2}\} \times [0, \frac{1}{2}) \to M \cap \{\underline{u} \ge -\frac{1}{2}\}, \qquad \Phi\left((u, \theta), s\right) = \sigma_{(u, -\frac{1}{2}, \theta)}(s).$$
(2.4)

Clearly,  $\Phi$  is a diffeomorphism.

Furthermore, we denote with h the auxiliary Riemannian metric on M given by  $h := du^2 + d\underline{u}^2 + \gamma(u, \underline{u})$ which measures deviation from the  $(u, \underline{u}, \theta)$ -differentiable structure.

And finally, our convention for the Riemann curvature tensor is

$$R(X,Y,Z,W) := -g(\nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z, W) = -g(\nabla_X (\nabla_Y Z) - \nabla_Y (\nabla_X Z) - \nabla_{[X,Y]} Z, W),$$

where X, Y, Z, W are smooth vector fields on (M, g).

# 3 The $C_{loc}^{0,1}$ -inextendibility of weak null singularities

#### 3.1 Auxiliary results

**Lemma 3.1.** Let (M,g) be a Lorentzian manifold with  $g \in C^0$ , let  $p \in M$  and let  $(b_0, b_1, \ldots, b_d)$  and  $(f_0, f_1, \ldots, f_d)$  be two orthonormal bases for  $T_pM$  with  $b_0$  and  $f_0$  timelike. Let L denote the Lorentz transformation relating the two bases, i.e.,  $f_{\alpha} = L_{\alpha\beta}b_{\beta}$ .<sup>7</sup> Then  $||L||_{\max} := \max_{\alpha\beta} |L_{\alpha\beta}| \le |g(b_0, f_0)|$ .

In other words, the Lorentz transformation relating two orthonormal bases can be uniformly bounded by the inner product of the two timelike basis vectors.

*Proof.* The metric g at p expressed in the orthonormal bases takes on the Minkowski form  $\eta = \text{diag}(-1, 1, \dots, 1)$ . It follows that

$$\eta_{\alpha\beta} = g(f_{\alpha}, f_{\beta}) = g(L_{\alpha\gamma}b_{\gamma}, L_{\beta\delta}b_{\delta}) = L_{\alpha\gamma}L_{\beta\delta}\eta_{\gamma\delta} ,$$

<sup>&</sup>lt;sup>7</sup>We use here the Einstein summation convention, i.e., there is an implicit sum over  $\beta$ .

which, in matrix notation, reads  $L\eta L^T = \eta$ . Thus we get  $L^{-1} = \eta L^T \eta$ , where we have used  $\eta^{-1} = \eta$ . Here  $L^{-1}$  is the inverse Lorentz transformation  $b_{\alpha} = (L^{-1})_{\alpha\beta} f_{\beta}$ . Thus we have

$$(L^{-1})_{00} = L_{00}$$
 and  $(L^{-1})_{0i} = -(L^T)_{0i} = -L_{i0}$ , (3.2)

where  $i = 1, \ldots, d$ . Consider

$$f_0 = L_{00}b_0 + L_{0i}b_i \tag{3.3}$$

and taking the inner product of both sides with  $b_0$  gives  $L_{00} = -g(f_0, b_0)$ . Taking the inner product of each side of (3.3) with itself gives

$$(L_{00})^2 = 1 + \sum_i (L_{0i})^2 , \qquad (3.4)$$

which provides the desired bound for  $L_{0i}$ . To obtain the bound for  $L_{i0}$ , we use (3.2) together with the equivalent of (3.4) for  $L^{-1}$ :

$$|L_{i0}|^2 = |(L^{-1})_{0i}|^2 \le |(L^{-1})_{00}|^2 - 1 = |L_{00}|^2 - 1.$$
(3.5)

Finally, taking the inner product of each side of  $f_i = L_{i0}b_0 + L_{ij}b_j$  with itself gives  $\sum_j (L_{ij})^2 = 1 + (L_{i0})^2 \leq (L_{00})^2$ , where we have used (3.5). This concludes the proof.

**Proposition 3.6.** Let (M, g) be a Lorentzian manifold in a double null gauge as in Section 2.1. Assume that the metric (2.1) in  $(u, \underline{u}, \theta^A)$ -coordinates extends continuously as a Lorentzian metric to the manifold with boundary<sup>8</sup>  $\overline{M} := (-1, 1) \times (-1, 0] \times \mathbb{S}^2$ . Moreover, assume that the following bounds on the Ricci coefficients (with respect to the double null frame from Section 2.2) are satisfied for some  $0 < C < \infty$ :

$$\sup_{M} \{ |\underline{\omega}| + |\eta|_{\gamma} + |\underline{\eta}|_{\gamma} + |\underline{\chi}|_{\gamma} \} \le C \quad \text{and} \quad \int_{-1}^{0} \left( \sup_{(u,\theta) \in (-1,1) \times \mathbb{S}^{2}} |\chi|_{\gamma}(u,\underline{u},\theta) \right) d\underline{u} \le C.$$
(3.7)

Let  $\tau : [0, t_0) \to M$  be an affinely parameterised future directed timelike geodesic with  $\lim_{s \to t_0} \underline{u}(\tau(s)) = 0$ and  $\lim_{s \to t_0} u(\tau(s)) < 1$ . Then  $\lim_{s \to t_0} \tau(s) \in \overline{M}$  exists,  $\tau$  has finite length  $0 < t_0 < \infty$ , and  $h(\dot{\tau}(s), \dot{\tau}(s)) \leq C < \infty$  for all  $s \in [0, t_0)$  for some constant C > 0.

This proposition establishes that the affine velocity of timelike geodesics approaching the weak null singularity remains regular with respect to the differentiable structure of  $\overline{M}$ , i.e., with respect to the differentiable structure of the  $C^0$ -extension constructed in [4].

Before we prove the proposition we briefly remark that there indeed exist future directed timelike geodesics in M that approach a point in  $\overline{M} \cap \{\underline{u} = 0\}$ . To see this we pick a point on  $\overline{M} \cap \{\underline{u} = 0\}$  and another one in its timelike past in M. Maximising the Lorentzian length of timelike curves connecting these two points (see [19], [6]) we find a future directed timelike geodesic which starts in M and approaches a point in  $\overline{M} \cap \{\underline{u} = 0\}$ .

Proof. We first show that  $\lim_{s\to t_0} \tau(s) \in \overline{M}$  exists and that the Lorentzian length of  $\tau$  is finite. Recall the time function  $t = u + 2\underline{u}$ . Its range on M is between -3 and 1. With slight abuse of notation we reparameterise  $\tau$  by t to obtain  $\tau : [t_1, t_2) \to M$  with  $-3 < t_1 < t_2 \leq 1$ . Thus we have  $1 = \dot{\tau}^u + 2\dot{\tau}^{\underline{u}}$ . Since both terms on the right and side are positive, they are uniformly bounded. Note that by the continuous extension of g to  $\overline{M}$  we in particular obtain that, along  $\tau$ , all metric components in (2.1) are uniformly bounded and that  $\gamma$  is a uniformly non-degenerate Riemannian metric on  $\mathbb{S}^2$ . This, together with

$$0 > g(\dot{\tau}, \dot{\tau}) = -4\Omega^2 \dot{\tau}^u \dot{\tau}^{\underline{u}} + \gamma_{AB} (\dot{\tau}^A - b^A \dot{\tau}^{\underline{u}}) (\dot{\tau}^B - b^B \dot{\tau}^{\underline{u}})$$
(3.8)

<sup>&</sup>lt;sup>8</sup>This means that  $\Omega$  extends as a continuous positive function to  $(-1,1) \times (-1,0] \times \mathbb{S}^2$ , the vector field  $b(u,\underline{u})$  on  $\mathbb{S}^2$  extends to  $(-1,1) \times (-1,0]$  such that  $b^A(u,\underline{u},\theta^1,\theta^2)$  are continuous up to and including  $\underline{u} = -1$ , and  $\gamma(u,\underline{u})$  extends as a Riemannian-metric-on- $\mathbb{S}^2$ -valued function to  $(-1,1) \times (-1,0]$  such that  $\gamma_{AB}(u,\underline{u},\theta^1,\theta^2)$  are continuous up to and including  $\underline{u} = -1$ .

and Cauchy-Schwarz gives the uniform boundedness of  $\gamma_{AB}\dot{\tau}^A\dot{\tau}^B$  and hence also that of  $\dot{\tau}^A$ . Firstly, this gives that  $\lim_{s\to t_2} \tau(s) \in \overline{M}$  exists – and secondly, using again the uniform boundedness of the metric components along  $\tau$  it is now straightforward that  $L(\tau) = \int_{t_1}^{t_2} \sqrt{-g(\dot{\tau}(s), \dot{\tau}(s))} \, ds$  is finite.

We return now to the affine parameterisation  $\tau : [0, t_0) \to M$  of  $\tau$  and show that its affine velocity vector remains bounded with respect to the  $(u, \underline{u}, \theta)$  differentiable structure.

We complement  $e_3$  and  $e_4$  with a choice of  $e_1$  and  $e_2$  along  $\tau$  to obtain a normalised double null frame along  $\tau$ , i.e.,  $g(e_{\mu}, e_A) = 0$  for  $\mu = 3, 4$  and A = 1, 2 and  $g(e_A, e_B) = \delta_{AB}$ . Expanding  $\dot{\tau}$  with respect to this frame we obtain  $\dot{\tau} = \dot{\tau}^4 e_4 + \dot{\tau}^3 e_3 + \dot{\tau}^2 e_2 + \dot{\tau}^1 e_1$  with  $\dot{\tau}^4 = \Omega^2 \dot{\tau}^{\underline{u}}$  and  $\dot{\tau}^3 = \dot{\tau}^u$ . We introduce the future directed timelike vector field  $T := e_3 + e_4$  on M and define the energy

$$E := -g(T, \dot{\tau}) = 2\dot{\tau}^4 + 2\dot{\tau}^3 = 2\Omega^2 \dot{\tau}^{\underline{u}} + 2\dot{\tau}^u .$$
(3.9)

Note that  $\dot{\tau}^3$  and  $\dot{\tau}^4$  are positive and thus are controlled by *E*. Moreover, the affine parameterisation of  $\tau$  implies  $-1 = g(\dot{\tau}, \dot{\tau}) = -4\dot{\tau}^3\dot{\tau}^4 + (\dot{\tau}^1)^2 + (\dot{\tau}^2)^2$  and thus

$$(\dot{\tau}^1)^2 + (\dot{\tau}^2)^2 \le 4\dot{\tau}^3 \dot{\tau}^4 . \tag{3.10}$$

Hence, we obtain

$$|\dot{\tau}^{\alpha}| \le E \qquad \text{for all } \alpha \in \{1, 2, 3, 4\} . \tag{3.11}$$

We compute with the convention that  $A, B \in \{1, 2\}$ 

$$\frac{d}{ds}E = -g(\nabla_{\dot{\tau}}T,\dot{\tau}) = -\sum_{\alpha,\beta=1}^{4} \dot{\tau}^{\alpha}\dot{\tau}^{\beta} \Big[g(\nabla_{e_{\alpha}}e_{3},e_{\beta}) + g(\nabla_{e_{\alpha}}e_{4},e_{\beta})\Big] 
= -\Big(\dot{\tau}^{A}\dot{\tau}^{B}\underline{\chi}_{AB} + \dot{\tau}^{A}\dot{\tau}^{4}(-2\zeta_{A}) + \dot{\tau}^{3}\dot{\tau}^{4}4\underline{\omega} + \dot{\tau}^{4}\dot{\tau}^{A}2\underline{\eta}_{A}\Big) 
- \Big(\dot{\tau}^{A}\dot{\tau}^{B}\chi_{AB} + \dot{\tau}^{A}\dot{\tau}^{3}2\zeta_{A} + \dot{\tau}^{3}\dot{\tau}^{A}2\eta_{A} + \dot{\tau}^{3}\dot{\tau}^{3}(-4\underline{\omega})\Big)$$
(3.12)

We now use (3.11), the first part of (3.7), and Cauchy Schwarz together with (3.10) for the  $\chi_{AB}$ -term to obtain<sup>10</sup>

$$\frac{d}{ds}E \le C(E^2 + E\dot{\tau}^4|\chi|_{\gamma})$$

Dividing by E and using the boundedness of  $\Omega^2$  along  $\tau$  we obtain

$$\frac{d}{ds}\log E \le C\Big(E + \dot{\tau}^{\underline{u}} \sup_{(u,\theta)\in(-1,1)\times\mathbb{S}^2} |\chi|_{\gamma} \big(u,\tau^{\underline{u}}(s),\theta\big)\Big),$$

We can now integrate over  $0 \le s < t_1 < t_0$  using (3.9) and the second part of (3.7) to get

$$\log E(t_1) \le \log E(0) + C \int_0^{t_1} [\dot{\tau}^{\underline{u}}(s) + \dot{\tau}^u(s)] \, ds + C \int_{\tau^{\underline{u}}(0)}^{\tau^{\underline{u}}(t_1)} \sup_{(u,\theta) \in (-1,1) \times \mathbb{S}^2} |\chi|_{\gamma} (u,\underline{u},\theta) \, d\underline{u} \le C$$

for all  $0 < t_1 < t_0$ . This shows the uniform boundedness of  $\dot{\tau}^u(s)$  and  $\dot{\tau}^{\underline{u}}(s)$ . The uniform boundedness of  $\gamma(\dot{\tau}(s), \dot{\tau}(s))$  follows as before from the right of (3.8) being equal to -1. This concludes the proof.

Note that the important structure of the proof was that the Ricci coefficient in (3.12) which is only integrable in  $\underline{u}$ , i.e.,  $\chi_{AB}$ , comes with a coefficient which is upper bounded in absolute value by  $E \cdot \dot{\tau}^4$ .

<sup>&</sup>lt;sup>9</sup>Note that this choice is different to the one made for  $e_1$  and  $e_2$  in Section 2.2. We will only use this new choice here in this proof.

 $<sup>^{10}</sup>$ The symbol C > 0 does not stand for a particular numerical value, but the constant may change value from line to line.

#### 3.2 The main theorem

**Theorem 3.13.** Consider the setting of Proposition 3.6 and assume in addition that there is a set of smooth coordinates  $(V_{(i)} \subseteq \mathbb{S}^2, \theta_{(i)})$  for  $\mathbb{S}^2$ , i = 1, 2, with  $V_{(1)} \cup V_{(2)} = \mathbb{S}^2$  and such that

$$\int_{-1}^{0} \sup_{(u,\theta_{(i)})\in(-1,1)\times V_{(i)}} \left|\frac{\partial}{\partial\theta^{B}_{(i)}} b^{A}_{(i)}(u,\underline{u},\theta_{i})\right| d\underline{u} < \infty$$
(3.14)

for i = 1, 2. Moreover, assume that for every  $\overline{p} \in \partial \overline{M}$  and any neighbourhood  $\overline{W} \subseteq \overline{M}$  of  $\overline{p}$  there exists  $\overline{q} \in \partial \overline{M} \cap \overline{W}$  and a compact neighbourhood  $\overline{V} \subseteq \overline{W}$  of  $\overline{q}$  such that for all  $\underline{u}' \leq 0$  close enough to 0 the set  $\overline{V} \cap \{\underline{u} \leq \underline{u}'\}$  is also a compact smooth manifold with corners<sup>11</sup>, and which, moreover, satisfies the following property: there exist continuous vector fields  $\overline{X}_i$  on  $\overline{V}$ ,  $i = 1, \ldots, 4$ , and an  $\varepsilon_0 > 0$  such that for any continuous vector fields  $\hat{X}_i \text{ on } \overline{V}$  with  $\sup_{\overline{V}} ||\hat{X}_i - \overline{X}_i||_h < \varepsilon_0$  we have

$$\left| \int_{\overline{V} \cap \{\underline{u} < \underline{u}_k\}} R(\hat{X}_1, \hat{X}_2, \hat{X}_3, \hat{X}_4) \operatorname{vol} \right| \to \infty$$
(3.15)

along a sequence  $\underline{u}_k \to 0$ .

Then there is no  $C^{0,1}_{\text{loc}}$ -extension  $\hat{\iota}: M \hookrightarrow \hat{M}$  with the property that there is an affinely parameterised, future directed and future inextendible timelike geodesic  $\tau: (-t_0, 0) \to M$  with  $\lim_{s\to 0} \underline{u}(\tau(s)) = 0$ ,  $\lim_{s\to 0} u(\tau(s)) < 1$ , and such that  $\lim_{s\to 0} (\hat{\iota} \circ \tau)(s) \in \hat{M}$  exists. Here,  $t_0 > 0$ .

Let us remark that the theorem only requires bounds on a subset of the connection coefficients. Bounding a connection strongly relies on the chosen frame – here we use the frame introduced in Section 2.2. We also briefly comment on the structure of the assumptions in the theorem: assuming the existence of a Lipschitz extension through a point  $\overline{p} \in \partial \overline{M}$  in the sense that a timelike geodesic leaves through this point, the first neighbourhood  $\overline{W} \subseteq \overline{M}$  arises just as a neighbourhood of  $\overline{p}$  which must also be contained in the extension. And while curvature may be bounded in the direction of isolated points on  $\partial \overline{M} \cap \overline{W}$ , the next statement is that one can always find an open set of boundary points towards which curvature blows up.

**Remark 3.16** (Discussion of the assumptions). All of the assumptions made in the setting of Proposition 3.6 are satisfied for each such finite chunk (M,g) of the Cauchy horizon by the work [4] of Dafermos and Luk. The bound (3.14) is not contained as a formal statement in [4] and requires some more work to establish. For the vector fields  $\overline{X}_i$  the expectation is that for example the choice  $\overline{X}_1 = \overline{X}_3 = e_4$  and  $\overline{X}_2 = e_A$ ,  $\overline{X}_4 = e_B$ gives (3.15). Indeed, unpublished work<sup>12</sup> by the author shows that the result of [23] can be slightly improved so that the version of (3.15) for linearised curvature is satisfied. Alternatively, this follows from the recent work [10].

**Remark 3.17** (Relation to global inextendibility result). The above theorem serves as one piece in a global low-regularity inextendibility statement of generic rotating vacuum black hole spacetimes. Such statements are proven by contradiction. Assuming the existence of a  $C_{loc}^{0,1}$ -extension, one knows that there is an inextendible timelike geodesic in the original spacetime that has a limit point in the extension (Theorem 2 in [6], Theorem 3.2 in [20], see also [16]). One then leads this statement to a contradiction by showing that none of the inextendible timelike geodesics can leave into a  $C_{loc}^{0,1}$ -extension. Here, one has to take the different geometries along those geodesics into account. Theorem 3.13 deals with those timelike geodesics approaching the weak null singularity, e.g.  $\tau_2$  in Figure 1. For timelike geodesics lying outside the black hole, for example  $\tau_1$  in

 $<sup>^{11}</sup>$ In the sense of [11].

 $<sup>^{12}\</sup>mathrm{It}$  may appear as part of forthcoming work.

Figure 1, one uses their completeness to derive a contradiction [6]. The geometry in the interior of the black hole to the future of the weak null singularity is still not well-understood.

**Remark 3.18** (The role of assumption (3.14)). The assumption (3.14) is used in two ways: on the one hand it is used to control the Ricci-coefficient  $g(\nabla_{e_4}e_A, e_B)$ . On the other hand it guarantees that the flow  $\Phi$  extends as a homeomorphism to  $\Phi: M \cap \{\underline{u} = -\frac{1}{2}\} \times [0, \frac{1}{2}] \to \overline{M} \cap \{\underline{u} \ge -\frac{1}{2}\}$ . To see the latter we first observe that by the assumption that g extends continuously to  $\overline{M}$  we have that the vector field L extends continuously to  $\underline{u} = 0$ – and thus also its integral curves. By a standard continuous dependence argument the flow map  $\Phi$  from (2.4) then extends continuously to  $M \cap \{\underline{u} = -\frac{1}{2}\} \times [0, \frac{1}{2}] \to \overline{M} \cap \{\underline{u} \ge -\frac{1}{2}\}$ . The surjectivity of this map follows from Peano's existence theorem. For the injectivity we consider two integral curves  $\sigma_j(s) = (u, s, \theta_j(s))$ , j = 1, 2, with  $\theta_1(-\frac{1}{2}) \neq \theta_2(-\frac{1}{2})$ . We want to show that  $\lim_{s\to 0} \theta_1(s) \neq \lim_{s\to 0} \theta_2(s)$ . For this we can assume that we can choose  $-1 < s_0 < 0$  close enough to 0 such that for  $s_0 \le s < 0$  the projections of the two integral curves onto the spheres lie in  $\mathring{V}_{(i)} \subseteq V_{(i)}$  for either i = 1 or i = 2, where  $\theta_{(i)}^A(\mathring{V}_{(i)}) \subseteq \mathbb{R}^2$  is convex. We now drop the index from the angular chart for the rest of the argument and compute in coordinates

$$\frac{d}{ds}||\theta_1^A(s) - \theta_2^A(s)||_{\mathbb{R}^2} \le ||b^A(\sigma_1(s)) - b^A(\sigma_2(s))||_{\mathbb{R}^2} \le \sup_{\substack{\theta \in V_{(i)}\\B,C \in \{1,2\}}} |\partial_{\theta^B} b^C| \cdot ||\theta_1^A(s) - \theta_2^A(s)||_{\mathbb{R}^2} \le ||\partial_{\theta^B} b^C| \cdot ||\theta_1^A(s) - \theta_2^A(s)||_{\mathbb{R}^2} \le ||\partial_{\theta^B} b^C| \cdot ||\partial_{\theta^$$

By (3.14) and a Gronwall backwards in s we obtain

$$||\theta_1^A(s_0) - \theta_2^A(s_0)||_{\mathbb{R}^2} \le C||\theta_1^A(s) - \theta_2^A(s)||_{\mathbb{R}^2}$$

for some C > 0 and for all  $s_0 \leq s < 0$ . This establishes the injectivity.<sup>13</sup> The continuity of the inverse then follows from the compactness of the domain (for constant u).

The statement of the theorem is adapted to the double null gauge chosen in [4], since presently this is the main application of this result. However, note that a double null gauge as in [13] would slightly change the statement of Theorem 3.13; in particular no assumption of the form (3.14) would be needed.

Proof. We begin by extending the continuous boundary extension  $\overline{M}$  to a continuous extension  $\tilde{\iota}: M \hookrightarrow \tilde{M}$ of (M, g) in the sense of Definition 1.1: we set  $\tilde{M} := (-1, 1) \times (-1, 1) \times \mathbb{S}^2$  and consider the trivial inclusion  $\tilde{\iota}: (-1, 1) \times (-1, 0) \times \mathbb{S}^2 \hookrightarrow (-1, 1) \times (-1, 1) \times \mathbb{S}^2$ . By pushing forward g via  $\tilde{\iota}$  we obtain a continuous metric on  $\overline{M} \subseteq \tilde{M}$  and extend it to  $\underline{u} > 0$  by setting, with respect to the local  $(u, \underline{u}, \theta)$ -coordinates,  $\tilde{g}_{\mu\nu}(u, \underline{u}, \theta) :=$  $\tilde{g}_{\mu\nu}(u, 0, \theta)$  for  $\underline{u} > 0$ .

To prove the theorem, we now assume that there is a  $C^{0,1}_{\text{loc}}$ -extension  $\hat{\iota}: M \hookrightarrow \hat{M}$  and a future directed and future inextendible timelike geodesic  $\tau: (-1,0) \to M$  with  $\lim_{s\to 0} \underline{u}(\tau(s)) = 0$ ,  $\lim_{s\to 0} u(\tau(s)) < 1$ , and  $\lim_{s\to 0} (\hat{\iota} \circ \tau)(s) =: \hat{p} \in \partial \hat{\iota}(M)$ . For the proof it is convenient to go over from the timelike geodesic  $\tau$  to a null geodesic (an integral curve of L) which also leaves through  $\hat{p}$ . In order to arrange for this, let us first observe that by Proposition 3.6 the limit  $\lim_{s\to 0} (\tilde{\iota} \circ \tau)(s) =: \tilde{p} \in \partial \tilde{\iota}(M)$  exists. We denote with  $\sigma: [-\frac{1}{2}, 0) \to M$ the integral curve of L (null geodesic), starting at parameter value  $s = -\frac{1}{2}$  at  $(u_0, -\frac{1}{2}, \theta_0) \in M$ , such that  $\lim_{s\to 0} (\tilde{\iota} \circ \sigma)(s) = \tilde{p}$ . There is exactly one such integral curve by Remark 3.18, i.e., since  $\Phi$  extends as a homeomorphism to  $M \cap \{\underline{u} = -\frac{1}{2}\} \times [0, \frac{1}{2}]$ .

**Step 1:** We show that  $\lim_{s\to 0} (\hat{\iota} \circ \sigma)(s) = \hat{p}$ .

We want to apply Proposition 5.1 from [25] with  $\gamma_1 = \tau$  and  $\gamma_2 = \sigma$ .<sup>14</sup> We start by also parameterising the timelike geodesic  $\tau$  by  $\underline{u}$  so that, by slight abuse of notation, we obtain

 $\tau(s) = \left(\tau^u(s), s, \tau^\theta(s)\right) \qquad \text{and} \qquad \sigma(s) = \left(\tilde{p}^u, s, \sigma^\theta(s)\right).$ 

 $<sup>^{13}</sup>$ For our application to weak null singularities inside generic rotating vacuum black holes this also follows from the estimate above (16.49) in [4].

<sup>&</sup>lt;sup>14</sup>Also note that the place of  $\tilde{M}$  in Proposition 5.1 from [25] is taken by  $\hat{M}$ .

We define the connecting curves  $\rho_n$  as follows: first consider the Riemannian metric  $\overline{h} := du^2 + \mathring{\gamma}_2$  on  $\mathbb{R} \times \mathbb{S}^2$ , where  $\mathring{\gamma}_2$  denotes the standard round metric on  $\mathbb{S}^2$ . Since  $\tau$  and  $\sigma$  have the same limit point  $\tilde{p}$ , we have

$$\ell_n := d_{\mathbb{R} \times \mathbb{S}^2} \left( \left[ \tau^u(-1/n), \tau^\theta(-1/n) \right], \left[ \tilde{p}^u, \sigma^\theta(-1/n) \right] \right) \to 0 \qquad \text{for } n \to \infty$$

Let now  $\overline{\rho}_n : [0, \ell_n] \to \mathbb{R} \times \mathbb{S}^2$  be the shortest curve in  $(\mathbb{R} \times \mathbb{S}^2, \overline{h})$  connecting  $[\tau^u(-1/n), \tau^\theta(-1/n)]$  with  $[\tilde{p}^u, \sigma^\theta(-1/n)]$ , parameterised by arclength. For *n* large enough, there is a unique such curve and thus  $\overline{\rho}_n$  is well-defined. We then define  $\rho_n : [0, \ell_n] \to M$  by

$$\rho_n(s) = \begin{cases} \left(\overline{\rho}_n^u(s), -\frac{1}{n} - s, \overline{\rho}_n^\theta(s)\right) & \text{for } 0 \le s \le \frac{1}{2}\ell_n \\ \left(\overline{\rho}_n^u(s), -\frac{1}{n} - \ell_n + s, \overline{\rho}_n^\theta(s)\right) & \text{for } \frac{1}{2}\ell_n \le s \le \ell_n . \end{cases}$$

Note that  $\rho_n$  connects  $\tau(-1/n)$  with  $\sigma(-1/n)$ . As will become clear later, it is important that the connecting curves  $\rho_n$  move also a little backwards and forwards in  $\underline{u}$  – otherwise their gap-length would not be finite in general.

We now introduce a reference orthonormal basis  $f_{\alpha}$  given by  $f_0 := \frac{1}{2}(e_3 + e_4)$ ,  $f_1 := \frac{1}{2}(e_3 - e_4)$ , and a smooth choice of  $f_2$  and  $f_3$ . The exact choice of  $f_2$  and  $f_3$  is not important and we only need this frame on the union of the curves  $\tau$ ,  $\sigma$ , and  $\rho_n$ , so that  $f_2$  and  $f_3$  can be smoothly defined.

Since  $\overline{h}(\dot{\overline{\rho}}(s), \dot{\overline{\rho}}(s)) = 1$ , it in particular follows that there is a constant C > 0, uniform in  $n \in \mathbb{N}$  and  $s \in [0, \ell_n]$ , such that

$$|g(\dot{\rho}_n(s), f_\alpha)| \le C . \tag{3.19}$$

In particular, if we expand  $\dot{\rho}_n(s) = \dot{\rho}_n^{\alpha} f_{\alpha}$ , then the components  $\dot{\rho}_n^{\alpha}$  with respect to the  $f_{\alpha}$  frame are uniformly bounded.

One of the assumptions of Proposition 5.1 in [25] is that the gap-length of  $\sigma$  is finite. This follows from  $\nabla_{e_4}e_4 = 0$ ,  $\dot{\sigma} = L = \Omega^2 e_4$ , and the uniform boundedness of  $\Omega^2$  along  $\sigma$ .

Next, we introduce a parallely propagated frame  $d_{\alpha}$  along  $\sigma$  and show that it is related to the  $f_{\alpha}$  frame by uniformly bounded Lorentz transformations.<sup>15</sup> We fix  $d_{\alpha}$  by demanding  $d_{\alpha}(\sigma(-\frac{1}{2})) = f_{\alpha}(\sigma(-\frac{1}{2}))$ . We claim that we have

$$|g(d_0(s), f_0)| \le C \tag{3.20}$$

for some C > 0 uniform along  $\sigma$ . By Lemma 3.1 this implies that the two frames are related by uniformly bounded Lorentz transformations.

To prove (3.20), we write  $d_0 = d_0^0 f_0 + d_0^i f_i$ . Since parallel transport preserves the spacetime inner product, we have

$$-1 = -(d_0^0)^2 + \sum_i (d_0^i)^2$$
(3.21)

and also  $d_0^0(s) > 0$  for all  $s \in [-\frac{1}{2}, 0)$ . By definition, we get

$$0 = g(\nabla_{\dot{\sigma}} d_0, f_0) = -\frac{d}{ds} d_0^0 + d_0^i \cdot g(\nabla_{\dot{\sigma}} f_i, f_0) .$$

Using (3.21) to bound  $d_0^i$  in terms of  $d_0^0$  and dividing by  $d_0^0$  we get

$$\frac{d}{ds} \log d_0^0 \le \sum_i |g(\nabla_{\sigma} f_i, f_0)| .$$

<sup>&</sup>lt;sup>15</sup>Indeed, this can also be more easily inferred just from  $\nabla_{e_4} e_4 = 0$ . However, the following method of proof will reappear later.

Note that  $\dot{\sigma} = L = \Omega^2 e_4$ ; and since  $\Omega^2$  is bounded, all the connection coefficients appearing on the right hand side are bounded<sup>16</sup> by (3.7). This proves the claim.

Next we consider a parallely propagated orthonormal frame  $b_{\beta}$  along  $\tau$  such that  $b_0 = \dot{\tau}$ . By Proposition 3.6 we have

$$|g(b_0, f_0)| \le C \tag{3.22}$$

for some constant C > 0 which is uniform along  $\tau$ .

We now claim that  $L_{\text{gap},b_{\beta}}(\rho_n) \to 0$  for  $n \to \infty$ , which is another assumption in Proposition 5.1 in [25]. To prove this, we first establish that for the parallel transport of  $b_{\beta}$  along  $\rho_n$  we have  $|g(b_0, f_0)| \leq C$  uniformly in  $s \in [0, \ell_n]$  and  $n \in \mathbb{N}$ .

We proceed as before. Writing  $b_0 = b_0^0 f_0 + b_0^i f_i$  it follows from  $0 = g(\nabla_{\dot{\rho}_n} b_0, f_0)$  in the same way as before that  $\frac{d}{ds} \log b_0^0 \leq \sum_i |g(\nabla_{\dot{\rho}_n} f_i, f_0)|$ . Using (3.19) this gives

$$\frac{d}{ds}\log b_0^0 \le C\sum_{i,\alpha} |g(\nabla_{f_\alpha} f_i, f_0)|$$

Integrating yields

$$\left|\log(b_0^0)|_{\rho_n(s)} - \log(b_0^0)|_{\tau(-1/n)}\right| \le C \left(\ell_n + 2 \int_{-1/n-\ell_n/2}^{-1/n} \sup_{u,\theta} |\hat{\chi}|_{\gamma}(\underline{u}) \, d\underline{u}\right) \to 0$$

for all  $s \in [0, \ell_n]$  and  $n \to \infty$ . Since  $0 < b_0^0|_{\tau(-1/n)} \leq C$  uniformly in  $n \in \mathbb{N}$  by (3.22), this gives

$$-g(b_0, f_0)\big|_{\rho_n(s)} = b_0^0|_{\rho_n(s)} \le C$$
(3.23)

uniformly in n and s.

By Lemma 3.1 the Lorentz transformations relating  $b_{\beta}$  and  $f_{\alpha}$  along  $\rho_n$  are uniformly bounded. Since by (3.19) the components of  $\dot{\rho}_n = \dot{\rho}_n^{\alpha} f_{\alpha}$  with respect to the frame  $f_{\alpha}$  are uniformly bounded, we also obtain that the components of  $\dot{\rho}_n = \dot{\rho}_n^{\beta} b_{\beta}$  with respect to the frame  $b_{\beta}$  are uniformly bounded in n and s. But this gives  $L_{\text{gap},b_{\beta}}(\rho_n) = \int_0^{\ell_n} \sqrt{\sum_{\beta} |\dot{\rho}_n^{\beta}(s)|^2} \, ds \to 0$  for  $n \to \infty$  since  $\ell_n \to 0$ .

Finally, note that by Lemma 3.1 and (3.20) together with (3.23) the Lorentz transformations relating the orthonormal bases  $b_{\beta}$  with  $f_{\alpha}$  are uniformly bounded along  $\sigma(-1/n)$  as well as the Lorentz transformations relating  $f_{\alpha}$  with  $d_{\alpha}$ . Thus the Lorentz transformations relating  $b_{\beta}$  with  $d_{\alpha}$  on  $\sigma(-1/n)$  are uniformly bounded in n. This verifies the assumptions of Proposition 5.1 in [25] and we infer  $\lim_{s\to 0} (\iota \circ \sigma)(s) = \hat{p}$ .

Step 2: Let  $\operatorname{id} := \hat{\iota} \circ \tilde{\iota}^{-1} : \tilde{\iota}(M) \to \hat{\iota}(M)$  denote the identification map. We show that there exists a future boundary chart  $\hat{\varphi} : \hat{U} \to (-\varepsilon_2, \varepsilon_2) \times (-\varepsilon_3, \varepsilon_3)^3$  around  $\hat{p} \subseteq \hat{M}$  and neighbourhoods  $\tilde{W} \subseteq \tilde{M}$  of  $\tilde{p}$  and  $\hat{W} \subseteq \hat{U}$  of  $\hat{p}$  such that  $\operatorname{id}|_{\tilde{W}_{\leq}} : \tilde{W}_{\leq} \to \hat{W}_{\leq}$  is a diffeomorphism and extends as a  $C^1$ -regular isometric diffeomorphism to  $\operatorname{id}|_{\tilde{W}_{\leq}} : \tilde{W}_{\leq} \to \hat{W}_{\leq}$ . Here we have  $\operatorname{set}^{17} \tilde{W}_{\leq} := \tilde{W} \cap \{\underline{u} < 0\}, \tilde{W}_{\leq} := \tilde{W} \cap \{\underline{u} \le 0\}$  and  $\hat{W}_{\leq} := \hat{W} \cap \hat{U}_{<}, \hat{W}_{\leq} := \hat{W} \cap \hat{U}_{<}$ .

We show the claim by verifying the assumptions of Proposition 5.10 in [25]. The setting of Proposition 5.10 is in terms of a future boundary chart  $\tilde{\varphi}: \tilde{U} \to (-\varepsilon_0, \varepsilon_0) \times (-\varepsilon_1, \varepsilon_1)^3$  around  $\tilde{p} \in \tilde{M}$ . Such a chart can be easily constructed by first making an affine coordinate transformation from the  $(u, \underline{u}, \theta^A)$  coordinates to new  $x^{\mu}$  coordinates in which  $\tilde{g}_{\mu\nu}|_{\tilde{p}} = \eta_{\mu\nu}$ , where  $\eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$ , and such that  $\frac{\partial}{\partial x^0}|_{\tilde{p}}$  is future directed – and then making  $\varepsilon_0, \varepsilon_1 > 0$  sufficiently small. Note that we have  $\tilde{U}_{\leq} = \tilde{U} \cap \{\underline{u} < 0\}$ .

<sup>&</sup>lt;sup>16</sup>Indeed,  $\chi$  does not appear. But even if it did, it would be fine.

<sup>&</sup>lt;sup>17</sup>See the definition of  $\hat{U}_{\leq}$  and  $\hat{U}_{\leq}$  below Proposition 1.3.

Recall that  $\Phi((u_0, \theta_0), \frac{1}{2}) = \tilde{p}$ . We now take a coordinate chart  $(V_{(i)}, \theta_{(i)})$  in which  $\theta_0$  is contained and we drop the index of this chart. Consider a coordinate ball  $B_{\rho}((u_0, \theta_0)) \subseteq \mathbb{R}^3$  around  $(u_0, \theta_0)$  with respect to  $(u, \theta^A)$  coordinates and then make  $1 - |u_0| > \rho > 0$  so small and choose  $0 \leq s_0 < \frac{1}{2}$  close enough to  $\frac{1}{2}$  such that  $\Phi(B_{\rho}((u_0, \theta_0)), [s_0, \frac{1}{2}]) \subseteq \tilde{U}$ . This uses the fact that  $\Phi$  extends continuously to  $M \cap \{\underline{u} = -\frac{1}{2}\} \times [0, \frac{1}{2}]$ . We thus have that  $\Phi : B_{\rho}((u_0, \theta_0)) \times [s_0, \frac{1}{2}] \to \tilde{U}_{\leq}$  onto its image, and that  $\partial_s x^{\mu}(\Phi(u, \theta, s)) = L^{\mu}$ is uniformly bounded.

After making  $\rho > 0$  smaller and choosing  $0 \le s_0 < \frac{1}{2}$  closer to  $\frac{1}{2}$  if necessary, we can assume that the closure of the projection of  $\Phi\left(B_{\rho}\left((u_0, \theta_0)\right) \times [s_0, \frac{1}{2})\right)$  onto  $\mathbb{S}^2$  is contained in one of the charts  $(V_{(i)}, \theta_{(i)})$ , of which we again drop the index in the following. We complement  $f_0$  and  $f_1$  to get a smooth orthonormal basis  $f_{\alpha}$  on  $\Phi\left(B_{\rho}\left((u_0, \theta_0)\right) \times \{s_0\}\right)$  and we define an orthonormal basis  $d_{\alpha}$  on  $\Phi\left(B_{\rho}\left((u_0, \theta_0)\right) \times [s_0, \frac{1}{2})\right)$  by parallely propagating  $d_{\alpha}$  along L with initial condition  $d_{\alpha} = f_{\alpha}$ .

<u>Claim</u>: The orthonormal basis  $d_{\alpha}$  extends continuously to  $\Phi\left(B_{\rho}\left((u_0,\theta_0)\right)\times[s_0,\frac{1}{2}]\right)$ .

To prove the claim, we use the smooth frame field  $e_{\alpha}$  from Section 2.2 on the domain under consideration, where  $e_A = \partial_{\theta^A}$ , for A = 1, 2. The dual basis is given by

$$\omega^{A} = g((d\theta^{A})^{\sharp_{\gamma}}, \cdot), \quad \omega^{3} = -\frac{1}{2}g(e_{4}, \cdot), \quad \omega^{4} = -\frac{1}{2}g(e_{3}, \cdot),$$

where  $\sharp_{\gamma}$  denotes the raising of an index with respect to the metric  $\gamma$  on the spheres. We then expand  $d_{\beta} = d^{\alpha}_{\ \beta} e_{\alpha}$  and contracting  $0 = \nabla_L d_{\beta} = \frac{d}{ds} d^{\alpha}_{\ \beta} \cdot e_{\alpha} + d^{\alpha}_{\ \beta} \cdot \nabla_L e_{\alpha}$  with  $\omega^{\rho}$  gives

$$\frac{d}{ds}d^{\rho}_{\ \beta} = -d^{\alpha}_{\ \beta}\omega^{\rho} \left(\nabla_{L}e_{\alpha}\right) =: -\mathbb{A}^{\rho}_{\ \alpha}d^{\alpha}_{\ \beta}$$

with

$$\begin{split} \mathbb{A}^4_{\ \alpha} &= -\frac{1}{2}g(e_3, \nabla_L e_\alpha) = \Omega^2 \frac{1}{2}g(\nabla_{e_4} e_3, e_\alpha) = \begin{cases} \Omega^2 \underline{\eta}_A & \text{for } \alpha = A \\ 0 & \text{for } \alpha \in \{3, 4\} \end{cases} \\ \mathbb{A}^3_{\ \alpha} &= -\frac{1}{2}g(e_4, \nabla_L e_\alpha) = 0 & \text{for } \alpha \in \{1, 2, 3, 4\} \end{cases} \\ \mathbb{A}^A_{\ \alpha} &= \gamma^{AB}g(e_B, \nabla_L e_\alpha) = \begin{cases} \gamma^{AB}g(e_B, \nabla_{e_C} L) + \gamma^{AB}g(e_B, [L, e_C]) = \Omega^2 \gamma^{AB} \chi_{BC} - \partial_{\theta^C} b^A & \text{for } \alpha = C \\ \Omega^2 \gamma^{AB}g(e_B, \nabla_{e_4} e_3) = 2\Omega^2 \gamma^{AB} \underline{\eta}_B & \text{for } \alpha = 3 \\ 0 & \text{for } \alpha = 4 \,. \end{cases} \end{split}$$

Assumptions (3.7) and (3.14), the continuity of  $\Omega^2$  together with  $1 - |u_0| > \rho > 0$ , as well as the continuity of  $\gamma_{AB}$  together with the closure of the projection of  $\Phi\left(B_\rho\left((u_0, \theta_0)\right) \times [s_0, \frac{1}{2})\right)$  onto  $\mathbb{S}^2$  being contained in one angular chart, give

$$\int_{-\frac{1}{2}}^{0} \sup_{(u,\theta)\in B_{\rho}\left((u_{0},\theta_{0})\right)} \left|\left|\mathbb{A}_{\alpha}^{\rho}\right|_{\Phi\left((u,\theta),s\right)}\right|\right| ds \leq C$$

$$(3.24)$$

for some C > 0. Hence, Gronwall gives

$$||d_{\beta}^{\rho}|_{\Phi((u,\theta),s)}|| \le C \qquad \text{uniformly for } (u,\theta) \in B_{\rho}((u_0,\theta_0)) \text{ and } s \in [-\frac{1}{2},0) .$$
(3.25)

Integrating the parallel transport equation we obtain for all  $s_0 \leq s_1 < s_2 < 0$ 

$$d^{\rho}_{\ \beta}|_{\Phi((u,\theta),s_2)} = d^{\rho}_{\ \beta}|_{\Phi((u,\theta),s_1)} - \int_{s_1}^{s_2} (\mathbb{A}^{\rho}_{\ \alpha} d^{\alpha}_{\ \beta})|_{\Phi((u,\theta),s)} \, ds$$

and combining this with (3.24) and (3.25) yields easily that  $d^{\rho}_{\beta}|_{\Phi((u,\theta),s)}$  extends continuously to  $(u,\theta,s) \in B_{\rho}((u_{0},\theta_{0})) \times [-\frac{1}{2},0]$ . Finally, since  $\Phi$  extends as a homeomorphism to this set and since the frame field  $e_{\alpha}$  is continuous on  $\Phi(B_{\rho}((u_{0},\theta_{0})) \times [s_{0},\frac{1}{2}])$ , so is  $d_{\beta} = d^{\alpha}_{\beta}e_{\alpha}$ . This proves the claim.

is continuous on  $\Phi\left(B_{\rho}((u_{0},\theta_{0}))\times[s_{0},\frac{1}{2}]\right)$ , so is  $d_{\beta}=d_{\beta}^{\alpha}e_{\alpha}$ . This proves the claim. Finally note that  $\partial_{s}\Phi^{\mu}((u,\theta),s)=L^{\mu}|_{\Phi\left((u,\theta),s\right)}$  is uniformly bounded on  $(u,\theta,s)\in B_{\rho}((u_{0},\theta_{0}))\times[-\frac{1}{2},0)$  in the  $x^{\mu}$  (or  $(u,\underline{u},\theta)$ ) coordinates. This verifies the last assumption in Proposition 5.10 in [25] so that we can conclude Step 2.

**Step 3:** Using the Lipschitz extension we construct vector fields  $\hat{X}_i$ , satisfying the assumptions of the theorem, such that (3.15) is uniformly bounded – which gives the contradiction.

First recall that  $\operatorname{id}|_{\tilde{W}_{<}}$  extending as a  $C^{1}$  isometric diffeomorphism to  $\operatorname{id}|_{\tilde{W}_{\leq}} : \tilde{W}_{\leq} \to \hat{W}_{\leq}$  means that one can extend the isometry  $\operatorname{id}|_{\tilde{W}_{<}}$  as a  $C^{1}$  diffeomorphism to some open subset of  $\tilde{W}$  containing  $\tilde{W}_{\leq}$ . So after making  $\tilde{W}$  and  $\hat{W}$  smaller if necessary we can assume that  $\operatorname{id}|_{\tilde{W}_{<}}$  extends as a  $C^{1}$ -diffeomorphism to  $\operatorname{id}|_{\tilde{W}} : \tilde{W} \to \hat{W}$ .

Let us denote the coordinates induced by  $\hat{\varphi} : \hat{U} \to (-\varepsilon_2, \varepsilon_2) \times (-\varepsilon_3, \varepsilon_3)^3$  by  $y^{\mu}$ . After making  $\tilde{W}$  and  $\hat{W}$  smaller if necessary we may assume that

$$|\hat{g}_{\mu\nu} - \eta_{\mu\nu}| \le \frac{1}{10}$$
 and  $|\partial_{\kappa}\hat{g}_{\mu\nu}| \le \Lambda$  on  $\hat{W}_{<}$  (3.26)

for some  $0 < \Lambda < \infty$ . As a consequence, if  $\hat{Y}, \hat{Z}$  are smooth vector fields on  $\hat{W}$  such that their components and the partial derivatives of those components with respect to the  $y^{\mu}$  coordinates are uniformly bounded, then the components

$$(\hat{\nabla}_{\hat{Y}}\hat{Z})^{\mu} = \hat{Y}^{\kappa} \frac{\partial}{\partial y^{\kappa}} \hat{Z}^{\mu} + \hat{\Gamma}^{\mu}_{\kappa\rho} \hat{Y}^{\kappa} \hat{Z}^{\rho} \le C$$
(3.27)

are uniformly bounded on  $W_{\leq}$ .

By the assumptions of the theorem there now exists  $\tilde{q} \in \tilde{W} \cap \{\underline{u} = 0\}$  and a compact neighbourhood  $\tilde{V} \subseteq \tilde{W}_{\leq}$  of  $\tilde{q}$  such that for all  $\underline{u}' \leq 0$  close enough to 0 the set  $\tilde{V} \cap \{\underline{u} \leq \underline{u}'\}$  is a smooth manifold with corners, and, moreover, there exist continuous vector fields  $\tilde{X}_i$  on  $\tilde{V}$ ,  $i = 1, \ldots, 4$ , and an  $\varepsilon_0 > 0$  such that for any continuous vector fields  $\hat{X}_i$  on  $\tilde{V}$  with<sup>18</sup> sup<sub> $\tilde{V}$ </sub>  $||\hat{X}_i - \tilde{X}_i||_h < \varepsilon_0$  we have  $|\int_{\tilde{V} \cap \{\underline{u} \leq \underline{u}_k\}} R(\hat{X}_1, \hat{X}_2, \hat{X}_3, \hat{X}_4) \operatorname{vol}| \to \infty$ 

along a sequence  $\underline{u}_k \to 0$ .



Figure 2: The  $C^1$ -identification of  $\hat{W}_{\leq}$  with  $\hat{W}_{\leq}$ .

Using the Tietze extension theorem in coordinates, we first extend the vector fields  $\tilde{X}_i$  to continuous vector fields on all of  $\tilde{W}$ , and, by abuse of notation, denote them by  $\tilde{X}_i$  as well. We then consider the vector fields  $(\mathrm{id}|_{\tilde{W}})_*\tilde{X}_i$  on  $\hat{W}$ , which are continuous as well, since  $(\mathrm{id}|_{\tilde{W}})_*$  is  $C^1$ . Also  $(\mathrm{id}|_{\tilde{W}})_*h$  is a continuous Riemannian metric on  $\hat{W}$ . Smoothing out  $(\mathrm{id}|_{\tilde{W}})_*\tilde{X}_i$  on  $\hat{W}$  with respect to the smooth structure of  $\hat{M}$  (note

 $<sup>^{18}\</sup>mathrm{We}$  extend the auxiliary Riemannian metric in the obvious way to  $\tilde{M}.$ 

that we have shown in Step 2 that the  $C^1$ -structures on  $\tilde{W}$  and  $\hat{W}$  are equivalent, but the  $C^2$ -structures are in general inequivalent) we can find smooth vector fields  $\hat{Y}_i$  on  $\hat{W}$  with

$$\sup_{\hat{V}} ||\hat{Y}_i - \mathrm{id}_* \tilde{X}_i||_{(\mathrm{id}|_{\tilde{W}})_* h} < \varepsilon_0 .$$

We now pull back the vector fields  $\hat{Y}_i$  to  $\tilde{W}$  to obtain continuous vector fields  $\hat{X}_i := (\mathrm{id}|_{\tilde{W}}^{-1})_* \hat{Y}_i$  on  $\tilde{W}$  satisfying

$$\sup_{\tilde{V}} ||\hat{X}_i - \tilde{X}_i||_h < \varepsilon_0 \; .$$

Thus, along a sequence  $\underline{u}_k \to 0$  we have

$$\frac{\left|\int\limits_{\tilde{V}\cap\{\underline{u}<\underline{u}_k\}} R(\hat{X}_1, \hat{X}_2, \hat{X}_3, \hat{X}_4) \operatorname{vol}\right| \to \infty}{=:I_k}$$
(3.28)

On the other hand we estimate

$$\begin{split} I_{k} &= \Big| \int_{\mathrm{id}(\tilde{V} \cap \{\underline{u} < \underline{u}_{k}\})} \hat{R}(\hat{Y}_{1}, \hat{Y}_{2}, \hat{Y}_{3}, \hat{Y}_{4}) \, \mathrm{vol} \Big| \\ &= \Big| \int_{\mathrm{id}(\tilde{V} \cap \{\underline{u} < \underline{u}_{k}\})} \hat{g}\Big(\hat{\nabla}_{\hat{Y}_{1}}(\hat{\nabla}_{\hat{Y}_{2}}\hat{Y}_{3}) - \hat{\nabla}_{\hat{Y}_{2}}(\hat{\nabla}_{\hat{Y}_{1}}\hat{Y}_{3}) - \hat{\nabla}_{[\hat{Y}_{1},\hat{Y}_{2}]}\hat{Y}_{3}, \hat{Y}_{4}\Big) \, \mathrm{vol} \Big| \\ &\leq C + \Big| \int_{\mathrm{id}(\tilde{V} \cap \{\underline{u} < \underline{u}_{k}\})} \Big[ \hat{Y}_{1}(\hat{g}(\hat{\nabla}_{\hat{Y}_{2}}\hat{Y}_{3}, \hat{Y}_{4})) - \hat{g}(\hat{\nabla}_{\hat{Y}_{2}}\hat{Y}_{3}, \hat{\nabla}_{\hat{Y}_{1}}\hat{Y}_{4}) - \hat{Y}_{2}(\hat{g}(\hat{\nabla}_{\hat{Y}_{1}}\hat{Y}_{3}, \hat{Y}_{4})) + \hat{g}(\hat{\nabla}_{\hat{Y}_{1}}\hat{Y}_{3}, \hat{\nabla}_{\hat{Y}_{2}}\hat{Y}_{4}) \Big] \, \mathrm{vol} \Big| \\ &\leq C + \Big| \int_{\mathrm{id}(\tilde{V} \cap \{\underline{u} < \underline{u}_{k}\})} d\Big(\hat{g}(\hat{\nabla}_{\hat{Y}_{2}}\hat{Y}_{3}, \hat{Y}_{4}) \cdot \hat{Y}_{1} \cup \mathrm{vol}\Big) - \hat{g}(\hat{\nabla}_{\hat{Y}_{2}}\hat{Y}_{3}, \hat{Y}_{4}) \cdot \mathrm{div}\hat{Y}_{1} \cdot \mathrm{vol} \Big| \\ &+ \Big| \int_{\mathrm{id}(\tilde{V} \cap \{\underline{u} < \underline{u}_{k}\})} d\Big(\hat{g}(\hat{\nabla}_{\hat{Y}_{1}}\hat{Y}_{3}, \hat{Y}_{4}) \cdot \hat{Y}_{2} \cup \mathrm{vol}\Big) - \hat{g}(\hat{\nabla}_{\hat{Y}_{1}}\hat{Y}_{3}, \hat{Y}_{4}) \cdot \mathrm{div}\hat{Y}_{2} \cdot \mathrm{vol} \Big| \\ &\leq C + \Big| \int_{\mathrm{id}(\tilde{V} \cap \{\underline{u} \le \underline{u}_{k}\})} \hat{g}(\hat{\nabla}_{\hat{Y}_{2}}\hat{Y}_{3}, \hat{Y}_{4}) \cdot \hat{Y}_{1} \cup \mathrm{vol}\Big| + \Big| \int_{\mathrm{id}(\tilde{V} \cap \{\underline{u} \le \underline{u}_{k}\})} \hat{g}(\hat{\nabla}_{\hat{Y}_{1}}\hat{Y}_{3}, \hat{Y}_{4}) \cdot \hat{Y}_{2} \cup \mathrm{vol}\Big| \\ &\leq C + C \sum_{i=1,2} \Big( \int_{\mathrm{id}(\tilde{V} \cap \{\underline{u} = \underline{u}_{k}\})} |\hat{Y}_{i} \cup \mathrm{vol}| + \int_{\mathrm{id}(\tilde{V})} |\hat{Y}_{i} \cup \mathrm{vol}|\Big) \\ &\leq C . \end{split}$$

$$(3.29)$$

Here, using the smoothness of the vector fields  $\hat{Y}_i$  we have just used the definition of the curvature tensor in the second step. In the third step we used the product rule on the first two terms and bounded the last term using the compactness of  $\hat{V}$ , the smoothness of  $[\hat{Y}_1, \hat{Y}_2]$  on  $\hat{W}$ , and (3.26) and (3.27). The fourth step is just Cartan's formula, and in the fifth step we use Stokes theorem for manifolds with corners and bound again the first order terms as before. In the sixth step we bound again the first order terms as before, estimate the integral of the three-form over the boundary by the corresponding density, and also split the boundary integral into a k-dependent part and estimate the remaining part by the integral over all of  $\partial id(\tilde{V})$ . Finally, the last integral is an integral of a continuous 3-density over a compact piecewise  $C^1$  boundary and thus finite, while to see the uniform boundedness of the first integral we may pull it back via id to obtain

$$\int_{\tilde{V} \cap \{\underline{u} = \underline{u}_k\}} |\tilde{X}_i \lrcorner \tilde{\mathrm{vol}}| = \int_{\tilde{V} \cap \{\underline{u} = \underline{u}_k\}} \sqrt{-\mathrm{det}\tilde{g}} |\tilde{X}_i^{\underline{u}}| \, du d\theta^1 d\theta^2$$

Since the integrand and the domain of integration are uniformly bounded in k, this integral is also uniformly bounded. This gives the contradiction to (3.28) and thus concludes the proof of the main theorem.

**Remark 3.30.** The method introduced in Step 3 of the proof also gives a possible approach for obtaining an inextendibility statement with  $\tilde{g} \in C^0$  and  $\partial \tilde{g} \in L^2_{loc}$ : if the extension is assumed to only have this regularity, then the boundary term in (3.29) is not necessarily bounded. With a minor modification, taking care of the boundary terms in the angular and the u direction, one can now square the whole estimate up to this point and integrate again in  $\underline{u}_k$  to get an expression which has to be finite for  $\tilde{g}$  in the above regularity class. If on the other hand the corresponding doubly integrated (and squared) curvature expression is assumed infinite, one again obtains the desired contradiction. The main problem with implementing this idea of proof is that the analogue of Step 2 is not known to hold; below Lipschitz regularity the methods of [25] break down. Indeed, in [1] it is shown that weak null singularities as considered here admit continuous extensions for which the identification map id does not extend as a  $C^1$ -regular diffeomorphism to the boundary!

# References

- [1] CAMERON, P., AND SBIERSKI, J. On the uniqueness of continuous spacetime extensions in 1 + 1 dimensions with applications to weak null singularities. *Forthcoming* (2024).
- [2] CHRISTODOULOU, D., AND KLAINERMAN, S. The Global Nonlinear Stability of the Minkowski Space. Princeton University Press, 1993.
- [3] CHRUŚCIEL, P., AND KLINGER, P. The annoying null boundaries. J. Phys. Conf. Ser. 968 (2018).
- [4] DAFERMOS, M., AND LUK, J. The interior of dynamical vacuum black holes I: The C<sup>0</sup>-stability of the Kerr Cauchy horizon. arXiv:1710.01772 (2017).
- [5] GALLOWAY, G., AND LING, E. Some Remarks on the C<sup>0</sup>-(in)extendibility of Spacetimes. Ann. Henri Poincaré 18, 10 (2017), 3427–3477.
- [6] GALLOWAY, G., LING, E., AND SBIERSKI, J. Timelike completeness as an obstruction to C<sup>0</sup>-extensions. Comm. Math. Phys. 359, 3 (2018), 937–949.
- [7] GRAF, M., AND LING, E. Maximizers in Lipschitz spacetimes are either timelike or null. Class. Quantum Grav. 35, 8 (2018).
- [8] GRAF, M., AND VAN DEN BELD-SERRANO, M. C<sup>0</sup>-inextendibility of FLRW spacetimes within a subclass of axisymmetric spacetimes. arXiv:2409.13799 (2024).
- [9] GRANT, J., KUNZINGER, M., AND SÄMANN, C. Inextendibility of spacetimes and Lorentzian length spaces. Ann. Glob. Anal. Geom. 55 (2019), 133–147.
- [10] GURRIARAN, S. Precise asymptotics of the spin +2 Teukolsky field in the Kerr black hole interior. arXiv:2409.0267 (2024).
- [11] LEE, J. Introduction to Smooth Manifolds. Springer, 2002.
- [12] LING, E. The  $C^0$ -inextendibility of some spatially flat FLRW spacetimes. arXiv:2404.08257v1 (2024).
- [13] LUK, J. Weak null singularities in general relativity. Journal of the American mathematical society 31, 1 (2018), 1–63.

- [14] MCNAMARA, J. Instability of black hole inner horizons. Proc. R. Soc. Lond. A 358 (1978), 499–517.
- [15] MIETHKE, B. C<sup>0</sup>-inextendibility of the Kasner spacetime. Master thesis, University of Vienna (also arXiv:2408.05257) (2024).
- [16] MINGUZZI, E., AND SUHR, S. Some regularity results for Lorentz-Finsler spaces. Ann. Glob. Anal. Geom. 56 (2019), 597–611.
- [17] ORI, A. Strength of curvature singularities. *Phys. Rev. D* 61 (2000).
- [18] RÁCZ, I. Spacetime extensions ii. Classical and Quantum Gravity 27, 15 (2010), 155007.
- [19] SÄMANN, C. Global Hyperbolicity for Spacetimes with Continuous Metrics. Annales Henri Poincaré 17, 6 (2016), 1429–1455.
- [20] SBIERSKI, J. On the proof of the C<sup>0</sup>-inextendibility of the Schwarzschild spacetime. J. Phys. Conf. Ser. 968 (2018).
- [21] SBIERSKI, J. The C<sup>0</sup>-inextendibility of the Schwarzschild spacetime and the spacelike diameter in Lorentzian geometry. J. Diff. Geom. 108, 2 (2018), 319–378.
- [22] SBIERSKI, J. On holonomy singularities in general relativity and the  $C_{loc}^{0,1}$ -inextendibility of space-times. Duke Mathematical Journal 171, 14 (2022), 2881 – 2942.
- [23] SBIERSKI, J. Instability of the Kerr Cauchy Horizon Under Linearised Gravitational Perturbations. Annals of PDE 9, 7 (2023).
- [24] SBIERSKI, J. The  $C^0$ -inextendibility of a class of FLRW spacetimes. arXiv:2312.07443 (2023).
- [25] SBIERSKI, J. Uniqueness and non-uniqueness results for spacetime extensions. International Mathematics Research Notices, rnae194 (2024).