# Null hypersurfaces in Einstein spacetimes as Carrollian manifolds

Samuel Blitz

Department of Mathematics and Statistics, Masaryk University Building 08, Kotlářská 2, Brno, CZ 61137 blitz@math.muni.cz

> David McNutt Center for Theoretical Physics, Polish Academy of Sciences, Warsaw mcnuttdd@gmail.com

> Pawel Nurowski Center for Theoretical Physics, Polish Academy of Sciences, Warsaw nurowski@fuw.edu.pl (Dated: October 1, 2024)

We explicitly determine all shear-free null hypersurfaces embedded in an Einstein spacetime as oriented 3-dimensional manifolds where each is equipped with a coframe basis, a structure group and a connection. Such manifolds are known as null hypersurface structures (NHSs). The coframe and connection one-forms for an NHS arise as solutions to an exterior differential system arising from the Cartan structure equations in the spacetime projected onto the null hypersurface. We then show that each NHS corresponds to a Carrollian structure equipped with a unique pair of Ehresmann connection and affine connection.

## I. INTRODUCTION

Within general relativity (GR) and more generally in Lorentzian geometry, null hypersurfaces are defined as hypersurfaces whose normal vector-field are non-zero but with zero magnitude with respect to the metric. Unlike timelike or spacelike hypersurfaces, where the normal vector-field has positive or negative magnitude, the induced metric for null hypersurfaces is necessarily degenerate [\[1\]](#page-11-0). Furthermore, the null normal vector-field lies in the tangent space of the hypersurface.

Despite this degeneracy, null hypersurfaces play an important role in GR and other gravity theories. For black hole solutions, the event horizon of the black hole is a null hypersurface [\[2,](#page-11-1) [3\]](#page-11-2). In the study of asymptotically flat Einstein spacetimes, Newman and Unti showed that on a foliation of null hypersurfaces, the Einstein field equations reduce to a set of first-order linear inhomogeneous differential equations [\[4\]](#page-11-3). Furthermore, asymptotic null infinity can, unsurprisingly, be treated as a null hypersurface using conformal methods [\[5](#page-12-0)]. Finally, when gluing solutions together using junction conditions, null hypersurfaces are the natural choice for black hole and wormhole solutions [\[6\]](#page-12-1).

Due to the degeneracy of the induced metric on null hypersurfaces, they cannot be realized as Riemannian geometries [\[7](#page-12-2)[–11\]](#page-12-3). Instead, the geometric structure is called a Carrollian geometry, whose structure group is the Carroll group  $[12]$ . This group is a dual contraction of the Poincaré group where the velocity of light is set to zero. As these are physically relevant hypersurfaces in spacetime, there is a growing interest in applications of Carrollian geometries in black hole physics [\[13](#page-12-5)[–19\]](#page-12-6) and field theories [\[20](#page-12-7)[–35\]](#page-12-8).

Mathematically, a Carrollian geometry is a  $(d-1)$ -dimensional differentiable manifold N equipped with a rank- $(d-2)$  semi-definite symmetric bilinear form h and a *fundamental vector field* k that spans the kernel of h. In this paper, we consider those Carrollian geometries induced by null hypersurfaces in four dimensional spacetimes.

To discuss such structures concretely, more data is required. With the addition of an Ehresmann connection  $\bf{K}$  that is dual to k, the tangent space TN is decomposed into a horizontal subbundle  $H\mathcal{N}$  and the span of k. Furthermore, when K is integrable, the horizontal subspace at a point can be integrated into a spacelike submanifold  $\mathcal C$  whose tangent space  $T_pC$  at a point  $p \in C$  is the horizontal subspace through that point  $H_p\mathcal{N}$ . Indeed, this picture explains why a Carrollian geometry paired with an Ehresmann connection is called *ruled* [\[19\]](#page-12-6), as it defines a foliation of the Carrollian geometry.

Within Carrollian geometry, there is substantial debate on the choice of affine connection used to construct differential geometric invariants. Following from arguments using Spencer cohomology, there is no unique torsion-free connection for Carrollian manifolds constructed only from the intrinsic geometry of the manifold[\[8\]](#page-12-9), unlike for (pseudo)Riemannian manifolds. As such, different geometric investigations may warrant different choices of additional geometric constraints that determine an affine connection.

In the literature, two choices seem to appear: an affine connection that is torsion-free but non-metric [\[18,](#page-12-10) [19,](#page-12-6) [36](#page-12-11)– [39\]](#page-12-12) and an affine connection that is metric but has torsion [\[8,](#page-12-9) [10](#page-12-13), [11,](#page-12-3) [21](#page-12-14), [40](#page-13-0)[–49\]](#page-13-1). While the latter is perhaps more well-motivated mathematically, the former seems to be more useful in physical contexts.

It is not clear, however, that these two options completely exhaust the natural geometric construction of affine connections on null hypersurfaces. To that end, we construct an affine connection for the Carrollian geometry in a new way. In [\[50\]](#page-13-2), Nurowski and Robinson used Cartan's moving frame approach to study the invariants of general null hypersurfaces, called null hypersurface structures (NHS), which included an examination of the bulk connection forms pulled back to the hypersurface. Inspired by their work, in this paper we examine the induced connection (found by gauge-fixing the freedom in the frame) on null hypersurfaces embedded in bulk spacetimes that are shear-free solutions to the vacuum Einstein equations. While somewhat restrictive, this family of hypersurfaces accounts for many solutions of physical interest (as, for example, many black hole spacetimes solve the vacuum Einstein equations). In doing so, we are able to produce closed form expressions for the connection coefficients and curvature components for all null hypersurfaces that can possibly be embedded in such spacetimes. The resulting induced connection can then be compared to the two connections described above. Furthermore, the intrinsic and extrinsic first-order invariants of a Carrollian geometry can be fully characterized for all null hypersurfaces satisfying the constraints we have imposed.

## <span id="page-1-3"></span>II. CARTAN STRUCTURE EQUATIONS AND THEIR PULLBACK TO A NULL HYPERSURFACE

We will consider the Newman-Penrose (NP) formalism as presented in [\[50\]](#page-13-2) for a four-dimensional spacetime  $(M, g)$ with a metric **g** and signature  $(+++-)$ . We define  $(e_1, e_2, e_3, e_4) = (\mathbf{m}, \bar{\mathbf{m}}, \ell, \mathbf{k})$  with the dual coframe  $(\theta^1, \theta^2, \theta^3, \theta^4) =$  $(M, M, L, K)$  so that

$$
\mathbf{M} = \mathbf{g}(\bar{\mathbf{m}}, \cdot), \quad \bar{\mathbf{M}} = \mathbf{g}(\mathbf{m}, \cdot), \quad \mathbf{L} = \mathbf{g}(\mathbf{k}, \cdot), \quad \mathbf{K} = \mathbf{g}(\ell, \cdot).
$$

With this choice the Lorentz frame transformations that preserve the null direction **k** can be written as

<span id="page-1-0"></span>
$$
\mathbf{M}' = e^{i\phi} [\mathbf{M} + z\mathbf{L}],
$$
  
\n
$$
\mathbf{L}' = A^{-1} \mathbf{L},
$$
  
\n
$$
\mathbf{K}' = A[\mathbf{K} + z\overline{z}\mathbf{L} + z\overline{\mathbf{M}} + \overline{z}\mathbf{M}].
$$
\n(1)

The first Cartan structure equations are

<span id="page-1-1"></span>
$$
d\mathbf{M} = -\mathbf{\Gamma}_{21} \wedge \mathbf{M} - \mathbf{\Gamma}_{23} \wedge \mathbf{L} - \mathbf{\Gamma}_{24} \wedge \mathbf{K},
$$
  
\n
$$
d\mathbf{L} = \mathbf{\Gamma}_{41} \wedge \mathbf{M} + \mathbf{\Gamma}_{42} \wedge \bar{\mathbf{M}} + \mathbf{\Gamma}_{43} \wedge \mathbf{L},
$$
  
\n
$$
d\mathbf{K} = \mathbf{\Gamma}_{31} \wedge \mathbf{M} + \mathbf{\Gamma}_{32} \wedge \bar{\mathbf{M}} + \mathbf{\Gamma}_{34} \wedge \mathbf{K}.
$$
\n(2)

where

<span id="page-1-2"></span>
$$
\Gamma_{41} = \sigma \mathbf{M} + \rho \bar{\mathbf{M}} + \tau \mathbf{L} + \kappa \mathbf{K},
$$
  
\n
$$
\Gamma_{23} = \mu \mathbf{M} + \lambda \bar{\mathbf{M}} + \nu \mathbf{L} + \pi \mathbf{K},
$$
  
\n
$$
\frac{1}{2} (\Gamma_{12} + \Gamma_{34}) = -\beta \mathbf{M} - \alpha \bar{\mathbf{M}} - \gamma \mathbf{L} - \epsilon \mathbf{K}.
$$
\n(3)

It is important to note here that traditionally the connection one-forms are written as  $\Gamma^a_{\ b} = \Gamma^a_{\ bc} \theta^c$ . However, the index a may be lowered to exploit the metric compatibility condition  $\Gamma_{ab} = -\Gamma_{ba}$ . For example,  $\Gamma_{3}^{1}$  is related to  $\Gamma_{2}^{4}$ since both can be expressed as  $\Gamma_{23}$ .

Under the transformations in equation [\(1\)](#page-1-0), these tranform as:

$$
\Gamma'_{41} = A^{-1}e^{-i\phi}\Gamma_{41},
$$
\n
$$
\Gamma'_{23} = Ae^{i\phi}(\Gamma_{23} + z[\Gamma_{12} + \Gamma_{34}] - z^2\Gamma_{14} - dz),
$$
\n
$$
[\Gamma_{12} + \Gamma_{34}]' = \Gamma_{12} + \Gamma_{34} + 2z\Gamma_{41} + d(\ln A) + i d\phi.
$$
\n(4)

Denoting the Weyl spinor scalars as  $\Psi_{\mu}$ ,  $\mu = 0, \ldots, 4$ , the Ricci scalar as R, and the traceless Ricci tensor as  $S_{ij} = R_{ij} - g_{ij}R/4$  i, j = 1, ..., 4, the second Cartan structure equations are [\[51](#page-13-3)]:

<span id="page-2-1"></span>
$$
d\Gamma_{23} = (\Gamma_{12} + \Gamma_{34}) \wedge \Gamma_{23} + \Psi_4 \bar{M} \wedge L + \Psi_3 (K \wedge L - M \wedge \bar{M})
$$
  
+  $(\Psi_2 + \frac{1}{12}R)K \wedge M + \frac{1}{2}S_{33}M \wedge L + \frac{1}{2}S_{32}(K \wedge L + M \wedge \bar{M}) + \frac{1}{2}S_{22}K \wedge \bar{M},$   
 $d\Gamma_{14} = \Gamma_{14} \wedge (\Gamma_{12} + \Gamma_{34}) + (-\Psi_2 - \frac{1}{12}R)\bar{M} \wedge L - \Psi_1(K \wedge L - M \wedge \bar{M})$   
 $- \Psi_0 K \wedge M - \frac{1}{2}S_{11}M \wedge L - \frac{1}{2}S_{41}(K \wedge L + M \wedge \bar{M}) - \frac{1}{2}S_{44}K \wedge \bar{M},$   
 $\frac{1}{2}(d\Gamma_{12} + d\Gamma_{34}) = \Gamma_{23} \wedge \Gamma_{14} - \Psi_3 \bar{M} \wedge L - (\Psi_2 - \frac{1}{24}R)(K \wedge L - M \wedge \bar{M})$   
 $- \Psi_1 K \wedge M - \frac{1}{2}S_{31}M \wedge L - \frac{1}{4}(S_{12} + S_{34})(K \wedge L + M \wedge \bar{M}) - \frac{1}{2}S_{42}K \wedge \bar{M}.$  (5)

Without loss of generality, a degenerate 3-manifold  $C$  is realized as a null hypersurface embedded in the spacetime as  $\mathcal{N} = \varphi(C)$  in the spacetime M, via the embedding  $\varphi : C \to M$  such that  $\varphi^* \mathbf{L} = 0$ . Similarly, using the natural projection operator to TN induced by the choice of frame,  $\top : TM|_{\mathcal{N}} \to TN$ , we have that  $\top \ell = 0$ . Thus using index notation, for any tensorial quantity, components with an index equal to 3 must be zero when viewed as objects on the hypersurface. This observation is key to deriving the intrinsic classification of null hypersurfaces as Carrollian manifolds. Then the Carrolian structure **h**, **k** on C is given by  $h = \phi^* g = 2M\overline{M}$  and **k** being the degenerate direction for h, and the structure equations on C are the pulbacks via  $\phi$  of Cartan's first and second structures equations for the spacetime.

For example, by combining equations [\(2\)](#page-1-1) with [\(3\)](#page-1-2), we find that

$$
d\mathbf{L} = (\bar{\rho} - \rho)\mathbf{M} \wedge \bar{\mathbf{M}} + (\bar{\alpha} + \beta - \tau)\mathbf{M} \wedge \mathbf{L} + (\alpha + \bar{\beta} - \bar{\tau})\bar{\mathbf{M}} \wedge \mathbf{L} + \kappa \mathbf{K} \wedge \mathbf{M} + \bar{\kappa}\mathbf{K} \wedge \bar{\mathbf{M}} + (\epsilon + \bar{\epsilon})\mathbf{K} \wedge \mathbf{L}.
$$
 (6)

Pulling back this equation to the null hypersurface, the left-hand side must vanish, whereas on the right-hand side only select terms vanish. Thus we find that

$$
Im(\rho)|_{\mathcal{N}} = 0, \quad \kappa|_{\mathcal{N}} = 0,\tag{7}
$$

implying that k is hypersurface-orthogonal and geodesic. As an abuse of notation in what follows, when obvious from context, we will omit  $\vert_{\mathcal{N}}$  and work with the same symbols when working with pulled back quantities.

Instead of describing a NHS as a three-dimensional oriented manifold  $\mathcal N$  equipped with a degenerate metric, **h** of signature  $(++)$ ), we may instead describe it as an oriented manifold N equipped with a basis of coframe fields. Recalling that in the spacetime, we may adapt our null coframe to the null hypersurface and the resulting metric takes the form

$$
\mathbf{g} = -2\mathbf{K}\mathbf{L} + 2\mathbf{M}\bar{\mathbf{M}},\tag{8}
$$

then the degenerate metric on  $\mathcal N$  is

<span id="page-2-0"></span>
$$
\mathbf{h} = 2\mathbf{M}\bar{\mathbf{M}} \tag{9}
$$

where M is a complex-valued one-form. On N we may always include a real-valued K so that  $M \wedge \overline{M} \wedge K \neq 0$ . Pulling back the Lorentz frame transformations in equation [\(1\)](#page-1-0) yields the following transformations

$$
\mathbf{M}' = e^{i\phi}\mathbf{M}, \n\mathbf{K}' = A[K + z\bar{\mathbf{M}} + \bar{z}\mathbf{M}].
$$
\n(10)

This is just a 1-dimensional extension of the Carroll group. The original definition of a NHS can be restated as [\[50\]](#page-13-2):

**Definition 1.** A null hypersurface structure (NHS) is a 3-dimensional manifold  $N$  equipped with an equivalence class of one-forms  $(M, K) \equiv (M, \overline{M}, K)$  such that

- $M$  is complex-valued and  $K$  is real-valued.
- $\mathbf{M} \wedge \mathbf{M} \wedge \mathbf{K} \neq 0$  at every point of N.
- Given  $(M, K)$  and  $(M', K')$ , they are equivalent if and only if they are related by the transformations [\(10\)](#page-2-0).

4

This definition is somewhat more general than a pre-Carrollian structure [\[11](#page-12-3)] due to the transformation group admitting a real-valued parameter, A. However, by setting this parameter to one and choosing a fundamental vector field k, we recover a Carrollian structure. In a similar manner as Carrollian manifolds, we may also choose a connection for the NHS:

## Definition 2. A null hypersurface manifold is a NHS equipped with an affine connection.

As mentioned in the introduction, we are interested in the characterization of embedded null hypersurfaces in spacetime. Following the classification in [\[50\]](#page-13-2), we could consider the invariants arising from the exterior differential systems that arise by pulling back to  $N$ . Pulling back to the null hypersurface, the first Cartan structure equations become:

<span id="page-3-0"></span>
$$
d\mathbf{M} = -\mathbf{\Gamma}_{21} \wedge \mathbf{M} - \mathbf{\Gamma}_{24} \wedge \mathbf{K},
$$
  
\n
$$
0 = \mathbf{\Gamma}_{41} \wedge \mathbf{M} + \mathbf{\Gamma}_{42} \wedge \bar{\mathbf{M}},
$$
  
\n
$$
d\mathbf{K} = \mathbf{\Gamma}_{31} \wedge \mathbf{M} + \mathbf{\Gamma}_{32} \wedge \bar{\mathbf{M}} + \mathbf{\Gamma}_{34} \wedge \mathbf{K}.
$$
\n(11)

A similar pullback of the second Cartan structure equations is

$$
d\Gamma_{23} = (\Gamma_{12} + \Gamma_{34}) \wedge \Gamma_{23} - \Psi_3 - M \wedge \bar{M} + (\Psi_2 + \frac{1}{12}R)K \wedge M + \frac{1}{2}S_{32}M \wedge \bar{M} + \frac{1}{2}S_{22}K \wedge \bar{M},
$$
  
\n
$$
d\Gamma_{14} = \Gamma_{14} \wedge (\Gamma_{12} + \Gamma_{34}) + \Psi_1 M \wedge \bar{M} - \Psi_0 K \wedge M - \frac{1}{2}S_{41}M \wedge \bar{M} - \frac{1}{2}S_{44}K \wedge \bar{M},
$$
\n(12)

<span id="page-3-1"></span>
$$
\frac{1}{2}(d\Gamma_{12}+d\Gamma_{34})=\Gamma_{23}\wedge\Gamma_{14}+(\Psi_2-\frac{1}{24}R){\bf{M}}\wedge\bar{{\bf{M}}}-\Psi_1{\bf{K}}\wedge{\bf{M}}-\frac{1}{4}(S_{12}+S_{34}){\bf{M}}\wedge\bar{{\bf{M}}}-\frac{1}{2}S_{42}{\bf{K}}\wedge\bar{{\bf{M}}}.
$$

This approach is independent of the choice of connection on  $\mathcal N$  and explicitly classifies null hypersurfaces in terms of invariantly-defined quantities. In particular, the quantities  $\rho$  and  $\sigma$  can be used to broadly classify null hypersurfaces into four distinct cases [\[50\]](#page-13-2). We will see that this approach picks out a preferred connection for a given hypersurface embedded in the spacetime.

We note that the equations in [\(5\)](#page-2-1) could be expanded to give the NP field equations. Similarly, by exterior differentiating [\(5\)](#page-2-1) we may recover the Bianchi identities. An advantage of deriving the equations in this manner is that one may explicitly derive identities that are helpful when pulled back to the null hypersurface.

#### III. FIELD EQUATIONS FOR NULL HYPERSURFACES IN EINSTEIN SOLUTIONS

In the mathematics literature, a spacetime is said to be Einstein when it satisfies the vacuum Einstein field equations. Explicitly, this means that

$$
R_{ij} - \frac{1}{2}g_{ij}R + \Lambda g_{ij} = 0,
$$

for some constant Λ. Taking the trace of this expression we find that  $R = 4\Lambda$ —a constant. This family of spacetimes includes many of physical interest, including but not limited to the prototypical black hole solutions.

Recall from Section [II](#page-1-3) that the pullback of the first Cartan structure equations [\(11\)](#page-3-0) reduce to two equations and a constraint equation. Pulling back the definition of the connection coefficients in equation [\(3\)](#page-1-2), we have that

<span id="page-3-2"></span>
$$
\Gamma_{41} = \sigma \mathbf{M} + \rho \bar{\mathbf{M}} + \kappa \mathbf{K},
$$
  
\n
$$
\Gamma_{23} = \mu \mathbf{M} + \lambda \bar{\mathbf{M}} + \pi \mathbf{K},
$$
  
\n
$$
\frac{1}{2} (\Gamma_{12} + \Gamma_{34}) = -\beta \mathbf{M} - \alpha \bar{\mathbf{M}} - \epsilon \mathbf{K}.
$$
\n(13)

Similarly, imposing the Einstein condition on the pullback of the second Cartan structure equations [\(12\)](#page-3-1) yields

<span id="page-3-3"></span>
$$
d\Gamma_{23} = (\Gamma_{12} + \Gamma_{34}) \wedge \Gamma_{23} - \Psi_3 - M \wedge \bar{M} + (\Psi_2 + \frac{1}{12}R)K \wedge M,
$$
  
\n
$$
d\Gamma_{14} = \Gamma_{14} \wedge (\Gamma_{12} + \Gamma_{34}) + \Psi_1 M \wedge \bar{M} - \Psi_0 K \wedge M
$$
  
\n
$$
\frac{1}{2}(d\Gamma_{12} + d\Gamma_{34}) = \Gamma_{23} \wedge \Gamma_{14} + (\Psi_2 - \frac{1}{24}R)M \wedge \bar{M} - \Psi_1 K \wedge M.
$$
\n(14)

<span id="page-4-0"></span>5

We remind the reader that we have chosen to write the pulled back connection 1-forms with indices lowered. In particular, we have the following relationships

$$
\Gamma_{23} = -\Gamma_{2}^{4}, \n\Gamma_{11} = -\Gamma_{4}^{2}, \n\Gamma_{12} = \Gamma_{2}^{2}, \n\Gamma_{34} = \Gamma_{4}^{4}.
$$
\n(15)

along with their complex conjugates. The metric compatibility condition in the spacetimes implies that  $\Gamma_{11a} = \Gamma_{22a}$  $0, a = 1, 2, 4$ , which ensures that the above connection one-forms on the null hypersurface contain all the information necessary to specify a connection.

Combining equations [\(11\)](#page-3-0), [\(13\)](#page-3-2), and the second equation in [\(14\)](#page-3-3) we may write  $\Psi_0$  as

$$
\Psi_0 = \mathbf{k}(\sigma) + (\bar{\epsilon} - 3\epsilon - \rho)\sigma \tag{16}
$$

From the coarse classification of NHSs in [\[50\]](#page-13-2) using  $\rho$  and  $\sigma$  there are four cases:

1.  $\sigma = \rho = 0$ , 2.  $\sigma = 0, \rho \neq 0,$ 3.  $\sigma \neq 0, \rho = 0,$ 4.  $\sigma \neq 0, \rho \neq 0.$ 

For Einstein spacetimes, the third case is necessarily excluded due to the null Raychaudhuri equation [\[51](#page-13-3), Chapter 5]:

$$
-\mathbf{k}(\rho) + \rho^2 + |\sigma|^2 = -\frac{1}{2}S_{44}.\tag{17}
$$

In the case of an Einstein spacetime with  $\rho = 0$ , this equation implies that  $\sigma = 0$ . In this paper we will focus on the null hypersurfaces where  $\sigma = 0$  as these hypersurfaces have physical relevance as shear-free hypersurfaces. Thus, imposing that  $\sigma = 0$  in equation [\(16\)](#page-4-0), it follows that  $\Psi_0 = 0$ .

We may treat the pullbacks of  $\Psi_i$ ,  $i = 1, 2, 3$  onto the hypersurfaces as extrinsic data that give information about the interaction of the hypersurface with the bulk spacetime. Following the analysis by Szekeres [\[52](#page-13-4)],  $\Psi_2$  describes the Coulomb-like part of the gravitational field while  $\Psi_1$  and  $\Psi_3$  describe incoming or outgoing longitudinal gravitation waves in the spacetime. The pullback of these quantities give insight into dynamics of the hypersurface.

We may explicitly integrate the field equations for any shear-free NHS in an Einstein spacetime to determine the coframe and connection one-forms. In what follows, we we will employ index notation for the connection coefficients instead of the quasi-NP formalism used in [\[50](#page-13-2)]. The following index convention will be used for covariant differentiation of basis one-forms:

$$
\nabla_a \theta^b = \Gamma^b_{\phantom{b}ca} \theta^c, \quad a, b, c = 1, 2, 4. \tag{18}
$$

This ordering is consistent with the ordering used in the Cartan structure equations for the original spacetime. We present the solutions to the two cases for shear-free NHSs as two propositions and include the proofs in the following subsections.

<span id="page-4-1"></span>**Proposition 1.** Any solution to the Einstein structure equations [\(14\)](#page-3-3) on a null hypersurface N satisfying  $\rho = \sigma = 0$ is gauge equivalent to

$$
\mathbf{M} = e^w \mathrm{d}\zeta, \quad \mathbf{K} = \mathrm{d}r + (X + r\bar{w}_{,\zeta})\mathrm{d}\zeta + (\bar{X} + rw_{,\bar{\zeta}})\mathrm{d}\bar{\zeta},\tag{19}
$$

with the following connection coefficients and their complex conjugates:

$$
\Gamma_{41}^{2} = 0, \ \Gamma_{42}^{2} = 0, \Gamma_{44}^{2} = 0,
$$
\n
$$
\Gamma_{21}^{2} = -\bar{w}_{,\zeta}e^{-w}, \quad \Gamma_{22}^{2} = w_{,\bar{\zeta}}e^{-\bar{w}}, \quad \Gamma_{24}^{2} = 0,
$$
\n
$$
\Gamma_{41}^{4} = -\bar{w}_{,\zeta}e^{-w}, \quad \Gamma_{42}^{4} = -w_{,\bar{\zeta}}e^{-\bar{w}}, \quad \Gamma_{44}^{4} = 0,
$$
\n
$$
\Gamma_{21}^{4} = -\left[\left(\frac{R}{8}e^{w} + \bar{w}_{,\zeta,\bar{\zeta}}e^{-\bar{w}}\right)r + \left(S + \frac{1}{2}\left(X_{,\bar{\zeta}} - \bar{X}_{,\zeta} + w_{,\bar{\zeta}}X - \bar{w}_{,\zeta}\bar{X}\right)\right)e^{-\bar{w}}\right]e^{-w},
$$
\n
$$
\Gamma_{22}^{4} = 0, \quad \Gamma_{24}^{4} = 0.
$$
\n(20)

Here,  $w = w(\zeta, \bar{\zeta}), X = X(\zeta, \bar{\zeta})$  are arbitrary complex-valued functions, and  $S = S(\zeta, \bar{\zeta})$  is an arbitrary real-valued function.

The pulled back NP Weyl scalars are

$$
\begin{aligned} \Psi_0 &= 0 \\ \Psi_1 &= 0 \\ \Psi_2 &= \frac{R}{24} + \bar{w}_{,\zeta\bar{\zeta}}e^{-(w+\bar{w})} \\ \Psi_3 &= e^{-(w+\bar{w})}\left[ \left( \left( \frac{R}{8}e^w + \bar{w}_{,\zeta\bar{\zeta}} \right) r + \frac{1}{2}e^{-\bar{w}}(2S+X_{,\bar{\zeta}}-\bar{X}_\cdot\zeta + w_{,\bar{\zeta}}X - \bar{w}_\cdot\zeta\bar{X})\right)_{,\bar{\zeta}} - \left( \frac{R}{8}e^w + \bar{w}_{,\zeta\bar{\zeta}}e^{-\bar{w}} \right)(\bar{X} + rw_{,\bar{\zeta}}) \right]. \end{aligned}
$$

<span id="page-5-0"></span>**Proposition 2.** Any solution to the Einstein structure equations [\(14\)](#page-3-3) in the on a null hypersurface  $\mathcal{N}$ , satisfying  $\rho \neq 0$  and  $\sigma = 0$  is gauge equivalent to

$$
\mathbf{M} = \frac{d\zeta}{\rho}, \quad \mathbf{K} = -\mathrm{d}\left(\frac{1}{\rho}\right) + \frac{1}{2}\rho(x\mathrm{d}\zeta + \bar{x}d\bar{\zeta}),\tag{21}
$$

with the following connection coefficients and their complex conjugates:

$$
\begin{split} \Gamma^2{}_{41} &= 0, \quad \Gamma^2{}_{42} = -\rho, \quad \Gamma^2{}_{44} = 0, \\ \Gamma^2{}_{21} &= -\frac{1}{2}x\rho^3, \quad \Gamma^2{}_{22} = \frac{1}{2}\bar{x}\rho^3, \quad \Gamma^2{}_{24} = 0, \\ \Gamma^4{}_{41} &= -\frac{1}{2}x\rho^3, \quad \Gamma^4{}_{42} = -\frac{1}{2}\bar{x}\rho^3, \quad \Gamma^4{}_{44} = 0, \\ \Gamma^4{}_{21} &= -\rho\left[-\frac{1}{6}\rho^4|x|^2 + \frac{1}{2}\rho^2x_{,\bar{\zeta}} + \rho a - \frac{R}{24\rho^2}\right], \\ \Gamma^4{}_{22} &= -\rho q, \Gamma^4{}_{24} = 0. \end{split}
$$

Here,  $x = x(\zeta, \bar{\zeta})$ ,  $q = q(\zeta, \bar{\zeta})$  are arbitrary complex-valued functions, and  $a = a(\zeta, \bar{\zeta})$  is an arbitrary real-valued function.

The pulled back NP Weyl scalars are

$$
\Psi_0 = 0,\n\Psi_1 = x\rho^4,\n\Psi_2 = a\rho^3 - \frac{2}{3}\rho^6 x \bar{x} + \rho^4 x_{,\bar{\zeta}}\n\Psi_3 = \frac{1}{3}\rho^8 x \bar{x}^2 - \frac{1}{3}\rho^6 \left(2\bar{x}x_{,\bar{\zeta}} - \frac{1}{2}x\bar{x}_{,\bar{\zeta}}\right) - \frac{1}{2}\rho^5 \bar{x}a + \rho^4 \left(\frac{1}{2}x_{,\zeta\bar{\zeta}} - qx\right) + \rho^4 a_{,\bar{\zeta}} - \rho^2 q_{,\zeta} - \frac{R}{24}\bar{x}.
$$
\n(22)

#### A. Proof of Proposition [1](#page-4-1)

Setting  $\rho = \sigma = 0$  in  $\Gamma_{41}$  in [\(13\)](#page-3-2), this implies that  $\Gamma_{41} = 0$ . Imposing this condition, the second equation in [\(11\)](#page-3-0) vanishes automatically, while the first equation in [\(11\)](#page-3-0), implies that

$$
d\mathbf{M} \wedge \mathbf{M} = -\mathbf{\Gamma}_{21} \wedge \mathbf{M} \wedge \mathbf{M} = 0 \tag{23}
$$

and so there exists (see for example [\[53\]](#page-13-5)) two complex functions w and  $\zeta$  such that

$$
\mathbf{M} = e^w \mathrm{d}\zeta \,. \tag{24}
$$

Since  $\mathbf{M} \wedge \bar{\mathbf{M}} \neq 0$  it follows that  $e^{2Re(w)}d\zeta \wedge d\bar{\zeta} \neq 0$ .

Imposing  $\Gamma_{41} = 0$  on the equations in [\(14\)](#page-3-3), the second equation implies that  $\Psi_1 = 0$  while the third equation implies

$$
d(\Gamma_{12} + \Gamma_{34}) \wedge M = d(\Gamma_{12} + \Gamma_{34}) \wedge d\zeta = 0.
$$
\n(25)

This implies the existence of two complex functions  $\eta$  and h such that  $\Gamma_{12} + \Gamma_{34} = d\eta + h d\zeta$  [\[53\]](#page-13-5). Using the frame freedom [\(10\)](#page-2-0) we can fix the parameters A and  $\phi$  so that

<span id="page-6-0"></span>
$$
\Gamma_{12} + \Gamma_{34} = h \mathrm{d}\zeta \,. \tag{26}
$$

Now inserting this 1-form into the third equation in [\(14\)](#page-3-3), we find

$$
dh \wedge d\zeta = 2\left(\Psi_2 - \frac{R}{24}\right) d\zeta \wedge d\bar{\zeta}.
$$
\n(27)

It follows that h is a function of  $\zeta$  and  $\overline{\zeta}$  only, and further that

$$
h_{,\bar{\zeta}} = 2\left(\frac{R}{24} - \Psi_2\right)e^{w+\bar{w}}.
$$
\n(28)

From [\(26\)](#page-6-0), it follows that  $\Gamma_{12} = \frac{1}{2} (h d \zeta - \bar{h} d \bar{\zeta})$ . We may plug this into the first equation of [\(11\)](#page-3-0) to find that  $w = w(\zeta, \bar{\zeta})$ only and

<span id="page-6-1"></span>
$$
h = -2\bar{w}_{,\zeta}.\tag{29}
$$

From [\(28\)](#page-6-1), and the above identity, we may solve for  $\Psi_2$ 

$$
\Psi_2 = \frac{R}{24} + \bar{w}_{,\zeta\bar{\zeta}}e^{-(w+\bar{w})}.
$$
\n(30)

Furthermore, the first equation in [\(14\)](#page-3-3) gives

$$
d\Gamma_{23} = -2\bar{w}_{,\zeta}d\zeta \wedge \Gamma_{23} - \Psi_3 e^{w+\bar{w}}d\zeta \wedge d\bar{\zeta} + \left(\frac{R}{8} + \bar{w}_{,\zeta\bar{\zeta}}e^{-(w+\bar{w})}\right)e^w \mathbf{K} \wedge d\zeta.
$$
 (31)

Thus, we find that  $d\Gamma_{23} \wedge d\zeta = 0$ , and there must exist two complex-valued functions s and q such that  $\Gamma_{23} = ds + qd\zeta$ [\[53\]](#page-13-5). Fixing the z-parameter in [\(10\)](#page-2-0), this can be set to be

<span id="page-6-5"></span><span id="page-6-4"></span><span id="page-6-3"></span><span id="page-6-2"></span>
$$
\Gamma_{23} = qd\zeta. \tag{32}
$$

We note that all of the frame transformation parameters have been fixed now.

Inserting equation [\(32\)](#page-6-2) into equation [\(31\)](#page-6-3) we get

$$
\mathrm{d}q = \Psi_3 e^{w + \bar{w}} \mathrm{d}\bar{\zeta} + \left(\frac{R}{8} + \bar{w}_{,\zeta\bar{\zeta}} e^{-(w + \bar{w})}\right) e^w \mathbf{K} + s \mathrm{d}\zeta,\tag{33}
$$

where  $s$  is an arbitrary complex-valued function. Using this in the third equation in  $(11)$ , we have

$$
d\mathbf{K} = (\bar{q}e^{w} - qe^{w})d\zeta \wedge d\bar{\zeta} + \mathbf{K} \wedge [\bar{w}_{,\zeta}d\zeta + w_{,\bar{\zeta}}d\bar{\zeta}]).
$$
\n(34)

Supplementing  $\zeta$  and  $\bar{\zeta}$  by a real variable r to produce a coordinate system  $(r, \zeta, \bar{\zeta})$  on N such that  $\mathbf{k} = \partial_r$ , we may express  $K$  as

$$
\mathbf{K} = \mathrm{d}r + Y\mathrm{d}\zeta + \bar{Y}\mathrm{d}\bar{\zeta},\tag{35}
$$

where Y is a complex-valued function of  $(r, \zeta, \overline{\zeta})$ . Plugging this expression for **K** into equation [\(34\)](#page-6-4) we find

$$
dY = q e^{\bar{w}} d\bar{\zeta} + \bar{w}_{,\zeta} [dr + \bar{Y} d\bar{\zeta}] + u d\zeta + t d\bar{\zeta},\tag{36}
$$

where  $t$  and  $u$  are real-valued and complex-valued functions, respectively.

This equation is equivalent to a system of partial differential equations (PDEs):

$$
Y_{,r} = \bar{w}_{,\zeta},
$$
  
\n
$$
Y_{,\bar{\zeta}} = \bar{w}_{,\zeta}\bar{Y} + qe^{\bar{w}} + t,
$$
  
\n
$$
Y_{,\zeta} = u.
$$
\n(37)

Integrating the first equation of [\(37\)](#page-7-0), we have

<span id="page-7-2"></span><span id="page-7-1"></span><span id="page-7-0"></span>
$$
Y = r\bar{w}_{,\zeta} + X,\tag{38}
$$

where  $X = X(\zeta, \bar{\zeta})$  is an arbitrary function. Using this expression for Y in the second equation of [\(37\)](#page-7-0) and taking the imaginary part gives

$$
r(\bar{w}_{,\zeta\bar{\zeta}} - w_{,\zeta\bar{\zeta}}) + X_{,\bar{\zeta}} - \bar{X}_{,\zeta} = \bar{w}_{,\zeta}\bar{X} - w_{,\bar{\zeta}}X + qe^{\bar{w}} - \bar{q}e^w.
$$
\n(39)

If we transcribe equation [\(33\)](#page-6-5) into a system of PDEs, we have

$$
q_{,r} = \frac{R}{8}e^w + \bar{w}_{,\zeta\bar{\zeta}}e^{-\bar{w}},
$$
  
\n
$$
q_{,\bar{\zeta}} = \Psi_3 e^{w+\bar{w}} + \left(\frac{R}{8}e^w + \bar{w}_{,\zeta\bar{\zeta}}e^{-\bar{w}}\right)\bar{Y}.
$$
\n(40)

Integrating the first equation in [\(40\)](#page-7-1) we find

$$
q = \left[\frac{R}{8}e^w + \bar{w}_{,\zeta\bar{\zeta}}e^{-\bar{w}}\right]r + Qe^{-\bar{w}}
$$
\n(41)

where  $Q = Q(\zeta, \bar{\zeta})$  is an arbitrary function. Note this integration required that R is constant, which followed from the Einstein condition. Inserting this into equation [\(39\)](#page-7-2) we find a constraint on the imaginary part of Q:

$$
Q - \bar{Q} = X_{,\bar{\zeta}} - \bar{X}_{,\zeta} - \bar{w}_{,\zeta}\bar{X} + w_{,\bar{\zeta}}X.
$$
\n
$$
(42)
$$

Writing  $Q + \bar{Q} = S$ , this is an arbitrary function of  $\zeta$  and its conjugate. Using the second equation in [\(40\)](#page-7-1) we may solve for  $\Psi_3$ . We have now fully integrated out the differential conditions for **M, K** and the connection 1-forms on the null hypersurface  $N$ , the results of which are summarized in Proposition [1.](#page-4-1)

#### B. Proof of Proposition [2](#page-5-0)

Setting  $\sigma = 0$  in  $\Gamma_{41}$  in [\(13\)](#page-3-2), this implies that

$$
\Gamma_{41} = \rho \bar{M}, \quad \bar{\rho} = \rho. \tag{43}
$$

Furthermore, from equation [\(16\)](#page-4-0), it follows that

<span id="page-7-4"></span><span id="page-7-3"></span>
$$
\Psi_0 = 0.\tag{44}
$$

The second equation in [\(11\)](#page-3-0) with equation[\(43\)](#page-7-3) implies that  $d\Gamma_{41} \wedge \Gamma_{41} = 0$  which in turn implies the existence of two complex-valued functions f and  $\zeta$  such that  $\Gamma_{14} = \bar{f}d\bar{\zeta}$ . Using the A and  $\phi$  parameters in the frame transformation group [\(10\)](#page-2-0) we may set  $\bar{f} = -1$  giving

$$
\Gamma_{41} = -d\bar{\zeta}.\tag{45}
$$

Then equating [\(43\)](#page-7-3) with [\(45\)](#page-7-4) gives

<span id="page-7-5"></span>
$$
\mathbf{M} = \frac{\mathrm{d}\zeta}{\rho}.\tag{46}
$$

Using  $(45)$  in the second equation in  $(14)$  we can express another connection 1-form:

$$
\Gamma_{12} + \Gamma_{34} = -\frac{\Psi_1}{\rho^2} \mathrm{d}\zeta + h \Gamma_{14},\tag{47}
$$

where h is an arbitrary complex-valued function. We now may fix the remaining  $z$  parameter in the frame transfor-mation group [\(10\)](#page-2-0) to set  $h = 0$  and yield

<span id="page-8-0"></span>
$$
\Gamma_{12} + \Gamma_{34} = -\frac{\Psi_1}{\rho^2} d\zeta.
$$
\n(48)

Taking M,  $\Gamma_{12}$  and  $\Gamma_{24}$  in equation [\(46\)](#page-7-5), the real part of equation [\(48\)](#page-8-0), and the complex conjugate of [\(43\)](#page-7-3), respectively, then the first equation in [\(11\)](#page-3-0) we can solve explicitely for the remaining coframe element:

$$
\mathbf{K} = -\mathrm{d}\left(\frac{1}{\rho}\right) + \frac{1}{2}\frac{\Psi_1}{\rho^3}\mathrm{d}\zeta + \frac{1}{2}\frac{\bar{\Psi}_1}{\rho^3}\mathrm{d}\bar{\zeta}.\tag{49}
$$

In addition, since  $\rho$  is real-valued, we can adapt  $(\rho, \zeta, \overline{\zeta})$  as a coordinate system on N.

Substituting  $\Gamma_{12} + \Gamma_{34}$  in [\(48\)](#page-8-0) into the third equation of [\(14\)](#page-3-3), it follows that there exists three complex-valued functions  $p, q$  and s such that

<span id="page-8-2"></span>
$$
\Gamma_{23} = \frac{p}{\rho^2} d\zeta + q d\bar{\zeta},\tag{50}
$$

along with

$$
\Psi_1 d\left(\frac{\Psi_1}{\rho^4}\right) + \left(2\rho + \frac{R}{12} - 2\Psi_2 - \frac{\Psi_1\bar{\Psi}_1}{\rho^2}\right) d\bar{\zeta} + 2s d\zeta = 0
$$
\n(51)

Notice that this second equation implies that

$$
\Psi_1 d\left(\frac{\Psi_1}{\rho^4}\right) \wedge d\zeta \wedge d\bar{\zeta} = 0.
$$
\n(52)

There are two cases, depending on whether  $\Psi_1$  vanishes or not. We will consider the case when  $\Psi_1 \neq 0$  first. Then equation [\(52\)](#page-8-1) implies that there is a function  $x = x(\zeta, \zeta)$  such that

<span id="page-8-3"></span><span id="page-8-1"></span>
$$
\Psi_1 = \rho^4 x. \tag{53}
$$

Setting this into  $(51)$  allows for p to be solved for algebraically

$$
p = \Psi_2 - \frac{R}{24} + \frac{1}{2}\rho^4(-x_{,\bar{\zeta}} + \rho^2 x\bar{x}).
$$
\n(54)

With this function set into the third equation in [\(11\)](#page-3-0), it can only be satisfied if

$$
\bar{\Psi}_2 - \Psi_2 = \rho^4 (\bar{x}_{,\zeta} - x_{,\bar{\zeta}}). \tag{55}
$$

The remaining equation which must be examined is the first equation of [\(14\)](#page-3-3). Computing the d $\rho \wedge d\zeta$  and  $d\rho \wedge d\zeta$ terms:

$$
q_{,r} = 0, \n\Psi_2 = a(\zeta, \bar{\zeta})\rho^3 - \frac{2}{6}\rho^6 x\bar{x} + \rho^4 x_{,\bar{\zeta}},
$$
\n(56)

where a is an arbitrary function of  $\zeta$  and  $\bar{\zeta}$ . The first equation implies that q must be independent of  $\rho$ , i.e.,  $q = q(\zeta, \bar{\zeta})$ . The second equation is compatible with equation [\(55\)](#page-8-3) if and only if  $a = \overline{a}$ , i.e., a is a real-valued function.

Finally, the d $\zeta \wedge d\zeta$  term in the first equation in [\(14\)](#page-3-3) gives an algebraic expression for  $\Psi_3$ 

$$
\Psi_3 = \frac{1}{3}\rho^8 x \bar{x}^2 - \frac{1}{3}\rho^6 \left(2\bar{x}x_{,\bar{\zeta}} - \frac{1}{2}x\bar{x}_{,\bar{\zeta}}\right) - \frac{1}{2}\rho^5 \bar{x}a + \rho^4 \left(\frac{1}{2}x_{,\zeta\bar{\zeta}} - qx\right) + \rho^4 a_{,\bar{\zeta}} - \rho^2 q_{,\zeta} - \frac{R}{24}\bar{x}.\tag{57}
$$

This solves the equations [\(11\)](#page-3-0) and [\(14\)](#page-3-3) in the case that  $\Psi_1 \neq 0$ . We note that solutions to these equations when  $\Psi_1 = 0$  can be formally obtained by setting  $x = 0$ . The results of this analysis are summarized in Proposition [2.](#page-5-0)

### IV. SHEAR-FREE NULL HYPERSURFACE STRUCTURES AS CARROLLIAN MANIFOLDS

For N an NHS embedded in a shear-free Einstein spacetime, there is automatically a degenerate metric,  $h = 2MM$ . Following from Proposition [1](#page-4-1) and [2,](#page-5-0) where the coframe basis has been determined invariantly, we may view  $\bf{K}$  as an Ehresmann connection. Then, by computing the dual of the coframe,  $(m, \bar{m}, k)$ , we may recover a vector-field that exists in the tangent space of  $\mathcal N$  that we use as the fundamental vector field:

Proposition 1(
$$
\sigma = \rho = 0
$$
):  $\mathbf{m} = -(X + r\bar{w}_{,\zeta})\partial_r + e^w \partial_{\zeta}$ ,  $\bar{\mathbf{m}} = -(\bar{X} + rw_{,\bar{\zeta}})\partial_r + e^{\bar{w}} \partial_{\bar{\zeta}}$ ,  $\mathbf{k} = \partial_r$ 

Proposition 2 
$$
(\sigma = 0, \rho \neq 0)
$$
:  $\mathbf{m} = -\frac{x\rho^4}{2}\partial_{\rho} + \rho\partial_{\zeta}$ ,  $\bar{\mathbf{m}} = -\frac{\bar{x}\rho^4}{2}\partial_{\rho} + \rho\partial_{\bar{\zeta}}$ ,  $\mathbf{k} = \rho^2\partial_{\rho}$ . (58)

Thus, the NHS may be treated as a Carrollian structure  $(N, \mathbf{h}, \mathbf{k})$  through a restriction of the frame transformation group to the Carrollian group by fixing the A parameter in equation [\(1\)](#page-1-0). Furthermore, this Carrollian structure is equipped with a distinguished affine connection, and so it is actually a Carrollian manifold.

We have seen that both the coframe and the connection for a NHS can be determined by solving the exterior differential system arising from the pullback of the Cartan structure equations. While we have required that the connection is torsion-free and metric-compatible in the bulk, these conditions do not necessarily hold for the induced connection on the NHS.

Taking the degenerate metric, it is a straightforward calculation to determine the nonmetricity tensor,  $\mathbf{Q} = \nabla \mathbf{h}$ ,

$$
\sigma = \rho = 0: \mathbf{Q} = 0,\n\sigma = 0, \rho \neq 0: \mathbf{Q} = \rho(\mathbf{M} \otimes \mathbf{K} \otimes \bar{\mathbf{M}} + \bar{\mathbf{M}} \otimes \mathbf{K} \otimes \mathbf{M} + \mathbf{K} \otimes \mathbf{h})
$$
\n(59)

Similarly, computing the intrinsic torsion of the connection for arbitrary vector-fields,  $X$  and  $Y$ , in the tangent space of  $\mathcal{N}$ :

$$
\mathbf{T}(\mathbf{X}, \mathbf{Y}) = \nabla_{\mathbf{X}} \mathbf{Y} - \nabla_{\mathbf{Y}} \mathbf{X} - [\mathbf{X}, \mathbf{Y}],
$$
\n(60)

we find that in the case that  $\sigma = \rho = 0$  for N, the torsion tensor is then

$$
\mathbf{T} = 2e^{w-\bar{w}} \left[ (\bar{w}_{,\zeta\bar{\zeta}} - w_{,\zeta\bar{\zeta}})r + Xw_{,\bar{\zeta}} - \bar{X}\bar{w}_{,\zeta} - \bar{X}_{,\zeta} + X_{,\bar{\zeta}} \right] \mathbf{k} \otimes (\mathbf{M} \wedge \bar{\mathbf{M}}) + 2\bar{w}_{\zeta}e^{-w} \mathbf{k} \otimes (\mathbf{K} \wedge \bar{\mathbf{M}}) + 2w_{\bar{\zeta}}e^{-\bar{w}} \mathbf{k} \otimes (\mathbf{K} \wedge \mathbf{M}).
$$
\n(61)

while in the case where  $\sigma = 0$  and  $\rho \neq 0$  the torsion tensor is

$$
\mathbf{T} = \frac{\rho^3}{2} (\bar{x}_{,\zeta} - x_{,\bar{\zeta}}) \mathbf{k} \otimes (\mathbf{M} \wedge \bar{\mathbf{M}}) + 2\rho^3 \bar{x} \mathbf{k} \otimes (\mathbf{K} \wedge \bar{\mathbf{M}}) + \rho^3 x \mathbf{k} \otimes (\mathbf{K} \wedge \mathbf{M}).
$$
 (62)

With the data contained in Propositions [1](#page-4-1) and [2,](#page-5-0) we may also compute the curvature tensor for these Carrollian geometries. The arbitrary functions that are contained in the connection coefficients but not the frame or coframe appear within the components of the curvature tensor. We note further that while expressions similar to  $\Psi_1$  and  $\Psi_2$ appear in the components of the curvature tensor,  $\Psi_3$  and  $\Psi_4$  will not. To understand this, observe that a similar phenomenon occurs even for non-null hypersurfaces. Indeed, for any signature hypersurface and a geometricallydetermined (co)frame along it, differentiation tangent to the hypersurface can only extract components of the bulk curvature tensor contracted with at most one copy of the vector field transverse to the hypersurface. As the constructed intrinsic covariant derivative on the Carrollian manifold is equal to the tangential Levi-Civita action modified by an extrinsic invariant that is first-order in derivatives on the bulk metric, the Carrollian curvature can only pick up components of the bulk curvature containing at most one copy of the transverse vector field. Because both  $\Psi_3$  and  $\Psi_4$  are curvature components involving two copies of the transverse vector field, the fact that we do not see these components in the Carrollian curvature is expected.

The connections we have found for the shear-free NHS embedded in Einstein spacetimes are new. Since  $\nabla \mathbf{h} \neq 0$ these connections are not metric compatible [\[11](#page-12-3)]. Of course, if  $\mathbf{Q} \neq \mathbf{0}$  one could introduce a shift tensor,  $\Delta \mathbf{\Gamma}$ ,

$$
\Delta \Gamma_Q = \rho (\mathbf{m} \otimes \mathbf{k} \otimes \mathbf{M} + \bar{\mathbf{M}} \otimes \mathbf{k} \otimes \bar{\mathbf{M}})
$$
(63)

so that  $\Gamma' = \Gamma + \Delta \Gamma$  is a metric compatible connection. However, for this new connection it is not possible to require that  $\nabla$ **k** = 0 or that  $\nabla$ **K**' = 0 without imposing conditions on the arbitrary functions, as would be required if the connection constructed here could be modified to produce the connection in [\[11\]](#page-12-3). This is expected as the connection discussed there is intended to describe the intrinsic properties of embedded null hypersurfaces as Carrollian structures.

However, using the idea of a shift tensor we may construct a new torsion-free connection on  $\mathcal N$  that is not metric compatible. We remark that this shift tensor is not quite a contorsion tensor as this would require raising and lowering of indices which is not permitted with a degenerate metric. For any NHS where  $\sigma = \rho = 0$ , we may use

<span id="page-10-0"></span>
$$
\Delta\Gamma_{T} = -2e^{w-\bar{w}} \left[ (\bar{w}_{,\zeta\bar{\zeta}} - w_{,\zeta\bar{\zeta}})r + Xw_{,\bar{\zeta}} - \bar{X}\bar{w}_{,\zeta} - \bar{X}_{,\zeta} + X_{,\bar{\zeta}} \right] \mathbf{k} \otimes \mathbf{M} \otimes \bar{\mathbf{M}} - 2\bar{w}_{,\zeta}e^{-w} \mathbf{k} \otimes \mathbf{M} \otimes \mathbf{K} - 2w_{,\bar{\zeta}}e^{-\bar{w}} \mathbf{k} \otimes \bar{\mathbf{M}} \otimes \mathbf{K}.
$$
\n(64)

while for the case where  $\sigma = 0$  but  $\rho \neq 0$  we may use

$$
\Delta\Gamma_T = -\frac{\rho^3}{2}(\bar{x}_{,\zeta} - x_{,\bar{\zeta}}) \mathbf{k} \otimes \mathbf{M} \otimes \bar{\mathbf{M}} - 2\rho^3 \bar{x} \mathbf{k} \otimes \mathbf{K} \otimes \bar{\mathbf{M}} - \rho^3 x \mathbf{k} \otimes \mathbf{K} \otimes \mathbf{M}.
$$
 (65)

Using  $\Gamma' = \Gamma + \Delta \Gamma_T$  we now have a torsion-free connection that still contains the arbitrary functions that arise from extrinsic information. This torsion-free connection is reminiscent of another choice of connection that is increasingly popular. Indeed, in the physics literature, this particular choice (summarized well in [\[18\]](#page-12-10)) is also a connection that both contains extrinsic embedding data and is torsion-free (but is not metric compatible).

In that context, this choice is made because it allows one to encode the bulk gravitational effects on the null hypersurface as aspects of the intrinsic geometry of the hypersurface, such as incoming gravitational radiation. The desired connection is determined by several constraints. First, one demands that the identity operator on the tangent bundle of the ruled Carrollian geometry is preserved under covariant differentiation and that the connection is symmetric. Then, to uniquely fix the connection, one uses the rigging connection [\[37,](#page-12-15) [38\]](#page-12-16) to determine the action of the connection on the Ehresmann connection and the fundamental vector field. The pullback and projection of the resulting tensors then complete the specification of the connection. In this way, extrinsic embedding data is built in to the connection at the cost of no longer being adapted to the Carrollian structure.

It is thus worthwhile to consider when the connection constructed in this manuscript can be shifted to the connection built from a rigging connection. By definition a symmetric connection is torsion-free so this condition is already satisfied. Taking the connection in Propositions [1](#page-4-1) and [2](#page-5-0) and shifting them by the shift tensor in equation [\(64\)](#page-10-0) and [\(65\)](#page-10-1) respectively, we may verify that the identity operator,

<span id="page-10-1"></span>
$$
\mathbf{q} = \mathbf{m} \otimes \mathbf{M} + \bar{\mathbf{m}} \otimes \bar{\mathbf{M}} + \mathbf{k} \otimes \mathbf{K},\tag{66}
$$

satisfies  $\nabla \mathbf{q} = 0$ . Thus, the shifted connections can be treated as a rigging connection and we may specify the action of the torsion-free connections on the fundamental vector-field and Ehresmann connection.

Corollary 1. For any null hypersurface embedded in an Einstein spacetime with  $\sigma = \rho = 0$  we may always construct a Carrollian geometry admitting a rigging connection where the fundamental vector-field has vanishing inaffinity,  $\mathbf{K}_b\mathbf{k}^a\nabla_a\mathbf{k}^b$  .

Proof. Using the torsion-free connection  $\Gamma' = \Gamma + \Delta \Gamma$  with  $\Gamma$  defined in Proposition [1](#page-4-1) and  $\Delta \Gamma$  defined in equation [\(64\)](#page-10-0). Computing the covariant derivative of the fundamental vector field,  $\bf{k}$ , and the Ehresmann connection,  $\bf{K}$ , we find:

$$
\nabla \mathbf{k} = \mathbf{k} \otimes (\bar{w}_{,\zeta} e^{-w} \mathbf{M} + w_{,\bar{\zeta}} e^{-\bar{w}} \bar{\mathbf{M}}),
$$
  
\n
$$
\nabla \mathbf{K} = \Gamma^3_{(12)} \mathbf{h} + (\Gamma^3_{[12]} + [\Delta \Gamma_T]^3_{[12]})(\mathbf{M} \wedge \bar{\mathbf{M}})
$$
  
\n
$$
+ 2 (\bar{w}_{,\zeta} e^{-w} \mathbf{M} + w_{,\bar{\zeta}} e^{-\bar{w}} \bar{\mathbf{M}}) \otimes \mathbf{K} + \mathbf{K} \otimes (\bar{w}_{,\zeta} e^{-w} \mathbf{M} + w_{,\bar{\zeta}} e^{-\bar{w}} \bar{\mathbf{M}}).
$$
\n(67)

**Corollary 2.** For any null hypersurface embedded in an Einstein spacetime with  $\sigma = 0$  and  $\rho \neq 0$  we may always construct a Carrollian geometry admitting a rigging connection where the fundamental vector-field has vanishing inaffinity,  $\mathbf{K}_b \mathbf{k}^a \nabla_a \mathbf{k}^b$ .

Proof. Using the torsion-free connection  $\Gamma' = \Gamma + \Delta \Gamma$  with  $\Gamma$  defined in Proposition [2](#page-5-0) and  $\Delta \Gamma$  defined in equation [\(65\)](#page-10-1). Computing the covariant derivative of the fundamental vector field,  $\bf{k}$ , and the Ehresmann connection,  $\bf{K}$ , we find:

$$
\nabla \mathbf{k} = -\rho (\mathbf{q} - \mathbf{k} \otimes \mathbf{K}) - \mathbf{k} \otimes \left( \frac{\rho^3 x}{2} \mathbf{M} + \frac{\rho^3 \bar{x}}{2} \bar{\mathbf{M}} \right),
$$
  
\n
$$
\nabla \mathbf{K} = \Gamma^3_{(12)} \mathbf{h} + (\Gamma^3_{[12]} + [\Delta \Gamma_T]^3_{[12]})(\mathbf{M} \wedge \bar{\mathbf{M}}) + \rho q \mathbf{M} \otimes \mathbf{M} + \rho \bar{q} \bar{\mathbf{M}} \otimes \bar{\mathbf{M}}
$$
  
\n
$$
+ \mathbf{k} \otimes \left( \frac{\rho^3 x}{2} \mathbf{M} + \frac{\rho^3 \bar{x}}{2} \bar{\mathbf{M}} \right) + (\rho^3 x \mathbf{M} + \rho^3 \bar{x} \bar{\mathbf{M}}) \otimes \mathbf{k}.
$$
\n(68)

 $\Box$ 

If we permit a scaling of **k** and **K** to preserve  $\mathbf{K}(\mathbf{k}) = 1$ , then employing the Leibnitz rule for covariant differentiation gives the next result.

Corollary 3. For any shear-free NHS embedded in an Einstein spacetime, we may construct a Carrollian geometry with non-vanishing inaffinity by setting  $\mathbf{k}' = A^{-1}\mathbf{k}$  and  $\mathbf{K}' = A\mathbf{K}$  where A is a function satisfying  $\mathbf{k}(A) \neq 0$ .

## V. CONCLUSIONS

By explicitly solving the exterior differential system that arises from the Cartan structure equations projected onto a null hypersurface, we have determined all possible shear-free null hypersurfaces in Einstein spacetimes as null hypersurface structures (NHS) equipped with a unique pairing of coframe basis and connection. The structure group for NHSs contains the Carrollian group as a subgroup and can be recovered by fixing one of the group parameters. Of particular note, all non-expanding horizons (NEHs) and hence weakly isolated horizons (WIH) [\[2\]](#page-11-1) in Einstein spacetimes are determined explicitly in this paper as NHSs.

Motivated by this observation, we then proved that for such hypersurfaces there is a uniquely determined Carrolian manifold with a unique pair of Ehresmann connection and affine connection. As such, this resolves a significant roadblock in discussing Carrollian geometry: how one geometrically fixes the Ehresmann connection. Indeed, previous work by Ashtekar et. al. [\[2\]](#page-11-1) showed that an Ehresmann connection can be canonically chosen for non-extremal WIHs. However, the method provided here provides a canonical choice regardless of whether the WIH is extremal.

As our mapping from a NHS to a Carrollian structure fixes the A parameter in the NHS structure group to unity, one must no longer make an arbitrary choice of representative in the generating null direction for the fundamental vector field. Instead, our construction chooses the unique representative with vanishing inaffinity. By permitting the A parameter to be a fixed function, it is possible to construct related Carrollian manifolds with non-vanishing inaffinity. In this setting, the geometry constructed here is equivalent to that of [\[18\]](#page-12-10).

In principle we have determined all WIHs embedded in Einstein spacetimes and our results could be used to study the potential uniqueness of the horizon for the extremal Kerr black hole [\[54](#page-13-6), [55\]](#page-13-7). However, it is not at all clear how extremality can be investigated for the current NHSs and their corresponding Carrollian geometries. If we naively relate inaffinity to surface gravity, then the canonical choice for Carrollian geometries always picks out a fundamental vector field with zero surface gravity.

While the construction provided in this manuscript has focused on Einstein spacetimes, the results can potentially be generalized to study null hypersurfaces in far more general settings. We note that if instead one considers a spacetime with a cosmological constant which also admits a null electromagnetic field whose null direction is proportional to K the resulting field equations in [\(11\)](#page-3-0) and [\(14\)](#page-3-3) would still be applicable. This implies that the shear-free NHSs in this paper would also describe NHS in these slightly more general spacetimes. As a more ambitious step, one may consider NHSs in asymptotically flat spacetimes with non-null EM fields as a source, such as the outer event horizon in Kerr-Newman spacetimes. Given that [\[16\]](#page-12-17) employs a connection that lacks extrinsic data, more could be said about such null hypersurfaces if the appropriate frame and connection were determined.

## ACKNOWLEDGMENTS

The authors would like to thank David Robinson who made considerable stylistic improvements to the initial draft of this paper during their collaboration with PN in the late nineties. SB is supported by the Operational Programme Research Development and Education Project No. CZ.02.01.01/00/ 22-010/0007541. DM and PN were supported by the Norwegian Financial Mechanism 2014-2021 (project registration number 2019/34/H/ST1/00636).

<span id="page-11-0"></span><sup>[1]</sup> R Penrose. The geometry of impulsive gravitational waves 1972 General Relativity, Papers in Honour of JL Synge, edited by L. O'Raifeartaigh, 1972.

<span id="page-11-1"></span><sup>[2]</sup> A. Ashtekar, C. Beetle, and J. Lewandowski. Geometry of generic isolated horizons. Classical and Quantum Gravity, 19(6):1195, 2002.

<span id="page-11-2"></span><sup>[3]</sup> R. M. Wald. General relativity. University of Chicago press, 2010.

<span id="page-11-3"></span><sup>[4]</sup> E. T. Newman and T. W. J. Unti. Behavior of asymptotically flat empty spaces. Journal of Mathematical Physics, 3(5):891–901, 1962.

- <span id="page-12-0"></span>[5] Y. Herfray. Carrollian manifolds and null infinity: a view from cartan geometry. Classical and Quantum Gravity, 39(21):215005, sep 2022.
- <span id="page-12-1"></span>[6] M. Mars and J. M. M. Senovilla. Geometry of general hypersurfaces in spacetime: junction conditions. Classical and Quantum Gravity, 10(9):1865, 1993.
- <span id="page-12-2"></span>[7] M. Henneaux. Geometry of Zero Signature Space-times. Bull. Soc. Math. Belg., 31:47–63, 1979.
- <span id="page-12-9"></span>[8] J. Figueroa-O'Farrill. On the intrinsic torsion of spacetime structures. arXiv preprint arXiv:2009.01948, 2020.
- [9] N. Pinzani-Fokeeva and A. Yarom. Enstrophy without boost symmetry. SciPost Physics, 12(4):136, April 2022.
- <span id="page-12-13"></span>[10] E. Have, K. Nguyen, S. Prohazka, and J. Salzer. Massive carrollian fields at timelike infinity. preprint, 2024.
- <span id="page-12-3"></span>[11] Samuel Blitz and David McNutt. Horizons that gyre and gimble: a differential characterization of null hypersurfaces. The European Physical Journal C, 84(6):561, 2024.
- <span id="page-12-4"></span>[12] J.M. L´evy-Leblond. Une nouvelle limite non-relativiste du groupe de poincar´e. Annales De L Institut Henri Poincare-physique Theorique, 3:1–12, 1965.
- <span id="page-12-5"></span>[13] R. F. Penna. BMS invariance and the membrane paradigm. Journal of High Energy Physics, 2016(3):23, March 2016.
- [14] L. Donnay and C. Marteau. Carrollian physics at the black hole horizon. Classical and Quantum Gravity, 36(16):165002, jul 2019.
- [15] J. Redondo-Yuste and L. Lehner. Non-linear black hole dynamics and Carrollian fluids. Journal of High Energy Physics, 2023(2):240, February 2023.
- <span id="page-12-17"></span>[16] F. Gray, D. Kubizňák, T. R. Perche, and J. Redondo-Yuste. Carrollian motion in magnetized black hole horizons. Physical Review D, 107(6):064009, 2023.
- [17] L. Ciambelli and L. Lehner. Fluid-gravity correspondence and causal first-order relativistic viscous hydrodynamics. Physical Review D, 108(12):126019, December 2023.
- <span id="page-12-10"></span>[18] L. Freidel and P. Jai-akson. Carrollian hydrodynamics and symplectic structure on stretched horizons. Journal of High Energy Physics, 2024(5):135, May 2024.
- <span id="page-12-6"></span>[19] L. Ciambelli, L. Freidel, and R.G. Leigh. Null Raychaudhuri: canonical structure and the dressing time. Journal of High Energy Physics, 2024(1):166, January 2024.
- <span id="page-12-7"></span>[20] A. Bagchi, R. Basu, A. Kakkar, and A. Mehra. Flat holography: aspects of the dual field theory. Journal of High Energy Physics, 2016(12):147, December 2016.
- <span id="page-12-14"></span>[21] L. Ciambelli, C. Marteau, A. C. Petkou, P. M. Petropoulos, and K. Siampos. Flat holography and carrollian fluids. Journal of High Energy Physics, 2018(7), jul 2018.
- [22] A. Bagchi, A. Mehra, and P. Nandi. Field theories with conformal carrollian symmetry. Journal of High Energy Physics, 2019(5), may 2019.
- [23] A. Bagchi, R. Basu, A. Mehra, and P. Nandi. Field theories on null manifolds. Journal of High Energy Physics, 2020(2), feb 2020.
- [24] N. Gupta and N. V. Suryanarayana. Constructing Carrollian CFTs. Journal of High Energy Physics, 2021(3):194, March 2021.
- [25] K. Banerjee, R. Basu, A. Mehra, A. Mohan, and A. Sharma. Interacting conformal carrollian theories: Cues from electrodynamics. Physical Review D, 103(10), may 2021.
- [26] D. Rivera-Betancour and M. Vilatte. Revisiting the carrollian scalar field. Physical Review D, 106(8):085004, 2022.
- [27] L. Donnay, A. Fiorucci, Y. Herfray, and R. Ruzziconi. Carrollian Perspective on Celestial Holography. Physical Review Letters, 129(7):071602, August 2022.
- [28] A. Bagchi, S. Banerjee, S. Basu, and S. Dutta. Scattering amplitudes: Celestial and carrollian. Physical Review Letters, 128(24), jun 2022.
- [29] B. Chen, R. Liu, and Y. Zheng. On higher-dimensional Carrollian and Galilean conformal field theories. SciPost Physics, 14(5):088, May 2023.
- [30] A. Bagchi, A. Banerjee, S. Dutta, K. S. Kolekar, and P. Sharma. Carroll covariant scalar fields in two dimensions. Journal of High Energy Physics, 2023(1):72, January 2023.
- [31] S. Baiguera, G. Oling, W. Sybesma, and B. T. Søgaard. Conformal carroll scalars with boosts. SciPost Physics, 14(4):086, 2023.
- [32] B. Chen, R. Liu, H. Sun, and Y. Zheng. Constructing Carrollian field theories from null reduction. Journal of High Energy Physics, 2023(11):170, November 2023.
- [33] A. Mehra and A. Sharma. Toward Carrollian quantization: Renormalization of Carrollian electrodynamics. Physical Review D, 108(4):046019, August 2023.
- [34] J. Salzer. An embedding space approach to Carrollian CFT correlators for flat space holography. Journal of High Energy Physics, 2023(10):84, October 2023.
- <span id="page-12-8"></span>[35] K. Banerjee, R. Basu, B. Krishnan, S. Maulik, Ad. Mehra, and A. Ray. One-loop quantum effects in Carroll scalars. Physical Review D, 108(8):085022, October 2023.
- <span id="page-12-11"></span>[36] V. Chandrasekaran, É. É. Flanagan, I. Shehzad, and A.J. Speranza. Brown-York charges at null boundaries. Journal of High Energy Physics, 2022(1), jan 2022.
- <span id="page-12-15"></span>[37] M. Mars and G. Sánchez-Pérez. Double null data and the characteristic problem in general relativity. Journal of Physics A: Mathematical and Theoretical, 56(3):035203, January 2023.
- <span id="page-12-16"></span>[38] M. Mars and G. Sánchez-Pérez. Covariant definition of double null data and geometric uniqueness of the characteristic initial value problem. Journal of Physics A: Mathematical and Theoretical, 56(25):255203, June 2023.
- <span id="page-12-12"></span>[39] L. Freidel and A. Riello. Renormalization of conformal infinity as a stretched horizon, 2024.
- <span id="page-13-0"></span>[40] J. Hartong. Gauging the carroll algebra and ultra-relativistic gravity. Journal of High Energy Physics, 2015(8), aug 2015.
- [41] E. Bergshoeff, J. Gomis, B. Rollier, J. Rosseel, and T. Ter Veldhuis. Carroll versus Galilei gravity. Journal of High Energy Physics, 2017(3):165, March 2017.
- [42] X. Bekaert and K. Morand. Connections and dynamical trajectories in generalised newton-cartan gravity. ii. an ambient perspective. Journal of Mathematical Physics, 59(7), 2018.
- [43] L Ciambelli, C. Marteau, A. C. Petkou, P. M. Petropoulos, and K. Siampos. Covariant galilean versus carrollian hydrodynamics from relativistic fluids. Classical and Quantum Gravity, 35(16):165001, jul 2018.
- [44] L. Ciambelli, R. G. Leigh, C. Marteau, and P. M. Petropoulos. Carroll structures, null geometry, and conformal isometries. Physical Review D, 100(4), aug 2019.
- [45] A. D. Gallegos, U. Gürsoy, S. Verma, and N. Zinnato. Non-Riemannian gravity actions from double field theory. Journal of High Energy Physics, 2021(6):173, June 2021.
- [46] J. Figueroa-O'Farrill, E. Have, S. Prohazka, and J. Salzer. The gauging procedure and carrollian gravity. Journal of High Energy Physics, 2022(9):243, September 2022.
- [47] O. Mišković, R. Olea, P. M. Petropoulos, D. Rivera-Betancour, and K. Siampos. Chern-Simons action and the Carrollian Cotton tensors. Journal of High Energy Physics, 2023(12):130, December 2023.
- [48] V. E. Marotta and R. J. Szabo. Godbillon-Vey invariants of Non-Lorentzian spacetimes and Aristotelian hydrodynamics. Journal of Physics A: Mathematical and Theoretical, 56(45):455201, November 2023.
- <span id="page-13-1"></span>[49] S. Baiguera, G. Oling, W. Sybesma, and B. T. Søgaard. Conformal Carroll scalars with boosts. SciPost Physics, 14(4):086, April 2023.
- <span id="page-13-2"></span>[50] P. Nurowski and D. C. Robinson. Intrinsic geometry of a null hypersurface. Classical and Quantum Gravity, 17(19):4065, 2000.
- <span id="page-13-3"></span>[51] H. Stephani, D Kramer, M. MacCallum, C. Hoenselaers, and E. Herlt. Exact Solutions of Einstein's Field Equations. Cambridge University Press, 2009.
- <span id="page-13-4"></span>[52] P Szekeres. The gravitational compass. Journal of Mathematical Physics, 6(9):1387–1391, 1965.
- <span id="page-13-5"></span>[53] Pawel Nurowski. General relativity, a one year course, lecture 15. https://www.fuw.edu.pl/~nurowski/teaching/gr1\_sum\_2011/gr5.p 2011.
- <span id="page-13-6"></span>[54] P. Nurowski and M. Randall. Generalized ricci solitons. The Journal of Geometric Analysis, 26:1280-1345, 2016.
- <span id="page-13-7"></span>[55] M. Dunajski and J. Lucietti. Intrinsic rigidity of extremal horizons. arXiv preprint arXiv:2306.17512, 2023.