EQUIVALENCE OF SOBOLEV NORMS FOR KOLMOGOROV OPERATORS WITH SCALING-CRITICAL DRIFT

THE ANH BUI, XUAN THINH DUONG, AND KONSTANTIN MERZ

Abstract. We consider the ordinary or fractional Laplacian plus a homogeneous, scalingcritical drift term. This operator is non-symmetric but homogeneous, and generates scales of L^p -Sobolev spaces which we compare with the ordinary homogeneous Sobolev spaces. Unlike in previous studies concerning Hardy operators, i.e., ordinary or fractional Laplacians plus scaling-critical scalar perturbations, handling the drift term requires an additional, possibly technical, restriction on the range of comparable Sobolev spaces, which is related to the unavailability of gradient bounds for the associated semigroup.

CONTENTS

²⁰²⁰ Mathematics Subject Classification. Primary: 46E36; Secondary: 26D10.

Key words and phrases. Fractional Laplacian, Kolmogorov operator, Hardy inequality, Hardy drift, Sobolev space.

1. Introduction and main result

1.1. Introduction. We consider Kolmogorov operators, i.e., ordinary or fractional Laplacians plus gradient perturbations, given by $(-\Delta)^{\alpha/2} + b(x) \cdot \nabla$, acting on functions in \mathbb{R}^d with $d \in \mathbb{N} := \{1, 2, \ldots\}.$ Here, $\alpha \in (0, 2]$ and $b : \mathbb{R}^d \to \mathbb{R}^d$ is a vector field. Their study is motivated, among others, by probability theory, where they arise as generators of Brownian motion or α -stable processes with drift. Kolmogorov operators are also important in physics, biology, finance, and further applied sciences, where the inclusion of gradient perturbations allows for modeling additional forces or influences acting on the system, thereby providing a more comprehensive description of the underlying processes. For example, in turbulent fluids, Kolmogorov operators describe anomalous diffusion, i.e., particles spreading at a rate different from what Brownian motion or α -stable processes predict [\[MK00\]](#page-44-0). In biology, gradient perturbations can represent directed movement or taxis [\[KRS08\]](#page-44-1), and in finance, they can represent drifts in asset price dynamics [\[CT04\]](#page-43-0). For further examples and references, we refer, e.g., to the review [\[Sti19\]](#page-44-2).

In this paper, we consider the following Kolmogorov operator, formally given by

$$
\Lambda_{\kappa} := (-\Delta)^{\alpha/2} + \frac{\kappa}{|x|^{\alpha}} x \cdot \nabla,
$$
\n(1.1)

for $d \in \mathbb{N}$ and $\alpha \in (1, 2]$ with $\alpha < (d+2)/2$ and a coupling constant $\kappa \in \mathbb{R}$. We call the gradient perturbation "attractive" for $\kappa > 0$ and "repulsive" for $\kappa < 0$, following terminology in [\[KS23,](#page-44-3) [KMS24\]](#page-43-1), and motivated by considering the action of $x \cdot \nabla$ on positive functions decaying at infinity. The assumption $\alpha > 1$ is necessary for the heat kernel of Λ_{κ} —which will be a key technical tool in our paper—to have a chance to be comparable with that of $Λ_0$. We defer a more detailed explanation to Remark [2.6.](#page-15-0) Besides, $α \geq 1$ is used in technical integral estimates, especially in Section [4,](#page-24-0) below. The assumption $\alpha < (d+2)/2$ is crucial as it ensures that Λ_{κ} generates a holomorphic semigroup and poses only an additional restriction when $d = 1$ or $d = 2$. We will review the precise definition of Λ_{κ} in Subsection [1.2](#page-2-0) below.

Importantly, Λ_{κ} is homogeneous of degree $-\alpha$. Therefore, one expects Λ_{κ} and Λ_{0} to be comparable to each other in some sense. That this is indeed the case, is the main result of the present paper. In Theorem [1.2](#page-4-0) below, we compare the $L^p(\mathbb{R}^d)$ -Sobolev spaces generated by powers of Λ_{κ} with the ordinary, homogeneous Sobolev spaces. Our result joins a line of recent research [\[KMV](#page-43-2)⁺18, [FMS21,](#page-43-3) [Mer21,](#page-44-4) [BD23\]](#page-42-2) on the so-called Hardy operator,

$$
L_{\kappa} := (-\Delta)^{\alpha/2} + \frac{\kappa}{|x|^{\alpha}} \quad \text{in } L^{2}(\mathbb{R}^{d}), \tag{1.2}
$$

i.e., the fractional Laplacian plus the scalar-valued, so-called, Hardy potential $\kappa/|x|^{\alpha}$. We also refer to [\[FM23,](#page-43-4) [BM23\]](#page-43-5) for studies concerning the regional fractional Laplacian [\[BBC03\]](#page-42-3) on the half-space \mathbb{R}^d_+ with Hardy potential κ/x_d^{α} depending on the distance to the bounding halfplane. As $(-\Delta)^{\alpha/2}$ and $|x|^{-\alpha}$ share the same scaling behavior, these operators compete with each on every length scale, thereby leading to the emergence of a critical coupling constant. More precisely, by the sharp Hardy–Kato–Herbst inequality [\[Har19,](#page-43-6) [Har20,](#page-43-7) [Kat66,](#page-43-8) [Her77\]](#page-43-9)^{[1](#page-1-2)}

$$
\left\|(-\Delta)^{\alpha/4} f\right\|_{L^{2}(\mathbb{R}^{d})}^{2} \geq \kappa_{*} \left\| |x|^{-\alpha/2} f \right\|_{L^{2}(\mathbb{R}^{d})}^{2}
$$

for all $\alpha \in (0, 2] \cap (0, d)$ and $f \in C_{c}^{\infty}(\mathbb{R}^{d})$ (1.3)

¹See also [\[KPS81,](#page-44-5) [Yaf99,](#page-44-6) [FLS08,](#page-43-10) [FS08\]](#page-43-11) for other proofs of [\(1.3\)](#page-1-3) with the optimal constant κ_* . Formula (1.3) is often simply called Hardy inequality.

with

$$
\kappa_* = \kappa_*(d, \alpha) := 2^{\alpha} \frac{\Gamma\left(\frac{d+\alpha}{4}\right)^2}{\Gamma\left(\frac{d-\alpha}{4}\right)^2},\tag{1.4}
$$

the quadratic form corresponding to L_{κ} is bounded from below if and only if $\kappa \geq -\kappa_*$. Moreover, if $\kappa \geq -\kappa_*$, then this form is nonnegative and L_{κ} can be realized as a self-adjoint operator by a theorem of Friedrichs. While L_{κ} has no eigenvalues, there is a strictly monotonously increasing function

$$
(-\infty, \kappa_*] \ni \kappa \mapsto \delta_\kappa \in (-M, (d-\alpha)/2], \tag{1.5}
$$

where

$$
M := \begin{cases} \alpha & \text{if } \alpha < 2, \\ \infty & \text{if } \alpha = 2, \end{cases}
$$
 (1.6)

such that $L_{\kappa}|x|^{-\delta_{\kappa}} = 0$ holds pointwise almost everywhere. As $|x|^{-\delta_{\kappa}}$ is, however, not squareintegrable, this function is sometimes called generalized eigenfunction or generalized ground state, and 0 is called the corresponding generalized eigenvalue of L_{κ} . We refer, e.g., to [\[FMS21,](#page-43-3) p. 2286] for an expression of δ_{κ} and merely record $\delta_{-\infty} = -M$, $\delta_0 = 0$, and $\delta_{\kappa*} = (d - \alpha)/2$.

1.2. Main result. To state our main result, Theorem [1.2,](#page-4-0) we introduce notation related to the homogeneity of Λ_{κ} and, in fact, two, corresponding critical coupling constants. In the following, we parameterize κ via

$$
\Psi(\beta) := \begin{cases}\n\frac{2^{\alpha}}{\beta - \alpha} \cdot \frac{\Gamma(\frac{\beta}{2})\Gamma(\frac{d}{2} - \frac{\beta - \alpha}{2})}{\Gamma(\frac{d}{2} - \frac{\beta}{2})\Gamma(\frac{\beta - \alpha}{2})} & \text{for } \alpha \in (1, 2) \text{ and } \beta \in (\alpha, d + \alpha), \\
d - \beta & \text{for } \alpha = 2 \text{ and } \beta \in \mathbb{R}.\n\end{cases}
$$
\n(1.7)

The parameter β enters in the generalized eigenvalue equation $\Lambda^*_{\kappa}|x|^{\beta-d} = 0$ with $\Lambda^*_{\kappa} = \Lambda_0 \kappa \nabla \cdot (|x|^{-\alpha}x)$ being the formal adjoint of Λ_{κ} , sometimes called Fokker–Planck operator. As δ^κ plays an important role in the analysis of Lκ, we anticipate that the parameter β will be important in the present study. We note that the fraction of the four Gamma functions in [\(1.7\)](#page-2-2) becomes maximal for $\beta = (d+\alpha)/2$, i.e., the midpoint of the interval (α, d) ; cf. [\[FMS21\]](#page-43-3). Thus, we denote the corresponding value of κ by

$$
\kappa_{\rm c} := \Psi\left(\frac{d+\alpha}{2}\right) = \begin{cases} \frac{2^{\alpha+1}}{d-\alpha} \cdot \frac{\Gamma\left(\frac{d+\alpha}{4}\right)^2}{\Gamma\left(\frac{d-\alpha}{4}\right)^2} & \text{if } \alpha < 2, \\ \frac{d-2}{2} & \text{if } \alpha = 2, \end{cases}
$$
(1.8)

which turns out to be one of the critical coupling constants as we will discuss shortly. However, $\Psi(\beta)$ is not symmetric around the midpoint $\beta = (d+\alpha)/2$, see, e.g., [\[KSS21,](#page-44-7) p. 1868] and [\[KS23,](#page-44-3) p. 347] for plots. Note further that $\lim_{\beta \nearrow d} \Psi(\beta) = \lim_{\beta \searrow d} \Psi(\beta) = 0$, $\lim_{\beta \nearrow d + \alpha} \Psi(\beta) = 0$ $-\infty$, and

$$
\kappa_{\rm c} = \frac{2}{d - \alpha} \kappa_*(d, \alpha). \tag{1.9}
$$

In Appendix [A.1](#page-36-1) we show that $\Psi(\beta)$ is decreasing for $\beta > (d+\alpha)/2$.

Lemma 1.1. Let $\alpha \in (1,2]$ and $M = \alpha$ if $\alpha < 2$ and $M = \infty$ if $\alpha = 2$. Then, the map $((d + \alpha)/2, d + M) \ni \beta \mapsto \Psi(\beta)$, given by [\(1.7\)](#page-2-2) is strictly monotonously decreasing.

We now explain how the scaling criticality of the gradient perturbation $\kappa |x|^{-\alpha} x \cdot \nabla$, called Hardy drift in [\[KSS21\]](#page-44-7), manifests itself in the emergence of two critical coupling constants. (1) The first, and for us more important, critical coupling constant is κ_c . For $\kappa < \kappa_c$ the Hardy drift is a Rellich perturbation of $(-\Delta)^{\alpha/2}$, i.e.,

$$
\|\kappa|x|^{-\alpha}x \cdot \nabla (\zeta + (-\Delta)^{\alpha/2})^{-1} \|_{L^2(\mathbb{R}^d) \to L^2(\mathbb{R}^d)} < 1.
$$
 (1.10)

This follows from Hardy's inequality [\(1.3\)](#page-1-3) in the form

$$
\|\kappa |x|^{1-\alpha} (-\Delta)^{-(\alpha-1)/2} \|_{L^2(\mathbb{R}^d) \to L^2(\mathbb{R}^d)} < 1,
$$
\n(1.11)

whenever $\kappa < \kappa_*(d, 2(\alpha - 1)) = \kappa_c$ and $\alpha < (d + 2)/2$. Consequently, Λ_{κ} can be constructed as the algebraic sum $(-\Delta)^{\alpha/2} + \kappa |x|^{-\alpha} x \cdot \nabla$ in $L^2(\mathbb{R}^d)$ with domain being the Sobolev space $H^{\alpha}(\mathbb{R}^d)$, and Λ_{κ} generates a holomorphic semigroup, denoted by $e^{-t\Lambda_{\kappa}}$, in $L^2(\mathbb{R}^d)$ whenever $\kappa < \kappa_c$; see also [\[KSS21,](#page-44-7) Proposition 8], [\[KS23,](#page-44-3) §8], and [\[KS20,](#page-44-8) Theorem 4.[2](#page-3-0)]². Since the holomorphic semigroup corresponding to Λ_{κ} plays an important technical role in the following, we restrict our attention from now on to $\kappa < \kappa_c$, or, equivalently, to

$$
\beta \in \left(\frac{d+\alpha}{2}, d+M\right) \quad \text{with } d \in \mathbb{N} \text{ and } \alpha \in (1,2] \cap (1, (d+2)/2) \tag{1.12}
$$

in view of [\(1.8\)](#page-2-3) and Lemma [1.1.](#page-2-1) Moreover, Kinzebulatov, Semenov, and Szczypkowski [\[KS20,](#page-44-8) [KSS21,](#page-44-7) [KS23\]](#page-44-3) proved that $e^{-t\Lambda_{\kappa}}$ is an L^{∞} contraction and extends by continuity to a C_0 semigroup on $L^r(\mathbb{R}^d)$ for all $r \in [2,\infty)$ whenever $\kappa < \kappa_c$. Furthermore, for $\alpha < 2$, they proved that the kernel $e^{-t\Lambda_{\kappa}}(x, y)$ is bounded from above and below by positive constants times the heat kernel of Λ_0 times a weight, depending on only one of the spatial variables. The reason that this weight depends only on one of the variables is due to the non-symmetry of Λ_{κ} . See [\(2.12\)](#page-14-1) below for these bounds. Analogous bounds for $\alpha = 2$ do not appear to be available yet. However, in view of the factorization of the bounds for $e^{-t\Lambda_{\kappa}}$ with $\alpha < 2$, we expect them to be of the form [\(2.14\)](#page-14-2).

(2) The second critical coupling constant is $2\kappa_c$. While the Hardy drift is not a Rellich perturbation anymore for $\kappa \geq \kappa_c$, the operators Λ_{κ} and $e^{-t\Lambda_{\kappa}}$ can still be defined for all $\kappa \in [\kappa_c, 2\kappa_c]$. More precisely, for $\alpha < 2$, [\[KSS21\]](#page-44-7) define $e^{-t\Lambda_{\kappa}}$ as the limit of the heat kernels associated to $(-\Delta)^{\alpha/2} + \kappa(|x|^2 + \varepsilon)^{-\alpha/2}x \cdot \nabla + \alpha \kappa \varepsilon(|x|^2 + \varepsilon)^{-\alpha/2-1}$, defined on the Sobolev space $(1 - \Delta)^{-\alpha/2} L^p(\mathbb{R}^d)$ for all $p \in [1, \infty)$, as $\varepsilon \searrow 0$ in $L^r(\mathbb{R}^d)$ for all $r \in (r_{\kappa}, \infty)$ with an explicit $r_{\kappa} \geq 2$. Correspondingly, for $\kappa \in [\kappa_c, 2\kappa_c)$, Λ_{κ} is defined to be the generator of the so-constructed heat kernel $e^{-t\Lambda_{\kappa}}$. We refer to [\[KSS21,](#page-44-7) Section 3] and [\[KS20,](#page-44-8) Section 4] for the precise procedures for $\alpha < 2$ and $\alpha = 2$, respectively. For $\alpha < 2$, the work [\[KSS21\]](#page-44-7) shows the heat kernel estimates for $\kappa < \kappa_c$ discussed above extend to all $\kappa < 2\kappa_c$. One of the reasons why $2\kappa_c$ also deserves to be called a critical constant is that for $\alpha = 2$ and $\kappa > 2\kappa_c$, appropriately defined weak solutions to the corresponding parabolic equation cease to be unique [\[KS20,](#page-44-8) p. 1588. Moreover, for $d = 3$ and $\kappa = 2\kappa_c$, the properties of the corresponding semigroup are drastically different from the properties of $e^{t\Delta}$ and $e^{-t\Lambda_{\kappa}}$ with $\kappa < 2\kappa_c$, see, e.g., [\[FL19\]](#page-43-12). Noteworthy, for $\alpha \in (1, 2)$, an optimal analog of Hardy's inequality (1.3) in L^p in [\[BJLPP22\]](#page-43-13) allows to extend certain estimates used in [\[KSS21\]](#page-44-7) to construct $e^{-t\Lambda_{\kappa}}$ and Λ_{κ} even for $\kappa > 2\kappa_c$; see [\[KSS21,](#page-44-7) Remark 6].

²While [\[KS20,](#page-44-8) [KSS21,](#page-44-7) [KS23\]](#page-44-3) state their results only for $d \in \{3, 4, ...\}$ and $\alpha \in (1, 2]$, an inspection of their proofs, taking [\(1.11\)](#page-3-1) into account, show that all their results actually hold for all $d \in \mathbb{N}$ and $\alpha \in (1, 2]$ with $\alpha < (d+2)/2$.

We now state our main result using the notation

$$
d_{\beta} := \frac{d}{(d-\beta) \vee 0} \quad \text{and} \quad (d_{\beta})' := 1 \vee \frac{d}{\beta} \tag{1.13}
$$

for $\beta > 0$ with $A \vee B := \max\{A, B\}$ and the convention $1/0 = \infty$. It says that the Sobolev spaces generated by powers of Λ_{κ} and Λ_0 are comparable with each other when the coupling constant is not too large in a quantitative sense.

Theorem 1.2 (Equivalence of Sobolev norms). Let $d \in \mathbb{N}$, $\alpha \in (1,2]$ with $\alpha < (d+2)/2$, $s \in (0,1], \beta \in ((d+\alpha)/2, d+M)$, and $\kappa = \Psi(\beta)$ be defined by [\(1.7\)](#page-2-2). If $\alpha = 2$, assume that the upper heat kernel bound [\(2.14\)](#page-14-2) for $e^{-tA_{\kappa}}$ holds. If

$$
(d_{\beta})' < p < \frac{d}{\alpha s} \wedge d_{\beta},\tag{1.14}
$$

then the following statements hold.

(1) Assume furthermore that $\alpha s < \alpha - 1$. Then,

$$
\|(\Lambda_0)^s f\|_{L^p(\mathbb{R}^d)} \lesssim_{d,\alpha,s,\beta,p} \|(\Lambda_\kappa)^s f\|_{L^p(\mathbb{R}^d)} \quad \text{for all } f \in C_c^\infty(\mathbb{R}^d). \tag{1.15}
$$

(2) We have

$$
\|(\Lambda_{\kappa})^s f\|_{L^p(\mathbb{R}^d)} \lesssim_{d,\alpha,s,\beta,p} \|(\Lambda_0)^s f\|_{L^p(\mathbb{R}^d)} \quad \text{for all } f \in C_c^{\infty}(\mathbb{R}^d). \tag{1.16}
$$

Here and in the following, we write $A \wedge B := \min\{A, B\}$. Moreover, for $A, B \ge 0$ and a parameter τ , we write $A \leq_{\tau} B$ whenever there is a τ -dependent constant $c_{\tau} > 0$ such that $A \leq c_{\tau}B$. The notation $A \sim_{\tau} B$ means $B \lesssim_{\tau} A \lesssim_{\tau} B$. The dependence on fixed parameters like d, α, s, β, p is usually omitted. Generic (real positive) constants will often be denoted by c or C. For brevity, we sometimes write $||f||_p$ instead of $||f||_{L^p(\mathbb{R}^d)}$.

The rest of this introduction is structured as follows. First, we state auxiliary tools, socalled reversed and generalized Hardy inequalities, and use them to prove Theorem [1.2.](#page-4-0) These inequalities are also of independent interest. Afterwards, we compare Theorem [1.2](#page-4-0) to earlier, related results and outline implications. Finally, we outline the rest of the paper.

1.3. Proof of Theorem [1.2.](#page-4-0) The ideas to prove Theorem [1.2](#page-4-0) are similar to those in [\[KMV](#page-43-2)⁺18, [FMS21,](#page-43-3) [Mer21,](#page-44-4) [BD23,](#page-42-2) [FM23,](#page-43-4) [BM23\]](#page-43-5), where heat kernel bounds were paramount. As the heat kernel bounds for Λ_{κ} and are tighter for $\alpha = 2$ than those for $\alpha < 2$ (compare [\(2.12\)](#page-14-1) and [\(2.14\)](#page-14-2) below), we will prove Theorem [1.2](#page-4-0) and the following auxiliary statements involved in its proof only for $\alpha \in (1, 2)$, and omit the extension of the proofs to the case $\alpha = 2$.

Compared to the above-mentioned studies, our proof of Theorem [1.2](#page-4-0) requires slightly more technical effort. First, as opposed to [\[FMS21,](#page-43-3) [FM23\]](#page-43-4), due to the non-symmetry of Λ_{κ} , establishing the equivalence of L^2 -Sobolev norms via the spectral theorem is impossible. Therefore, we will resort to the continuous square function estimates established in [\[BD23\]](#page-42-2) and again ap-plied in [\[BM23\]](#page-43-5). Due to the non-symmetry of Λ_{κ} , we need to investigate the boundedness of the following two square functions: The first is associated with Λ_{κ} and the latter is associated with its adjoint Λ_{κ}^* . For $\gamma > 0$, they are defined as

$$
S_{\Lambda_{\kappa},\gamma}f(x) = \left(\int_0^\infty |(t\Lambda_{\kappa})^\gamma e^{-t\Lambda_{\kappa}}f|^2 \frac{dt}{t}\right)^{1/2} \tag{1.17}
$$

and

$$
S_{\Lambda^*_{\kappa},\gamma}f(x) = \left(\int_0^\infty |(t\Lambda^*_{\kappa})^\gamma e^{-t\Lambda^*_{\kappa}}f|^2 \frac{dt}{t}\right)^{1/2}.\tag{1.18}
$$

As Λ_{κ} generates a holomorphic semigroup [\[KSS21\]](#page-44-7), the functional calculus in [\[McI86\]](#page-44-9) implies that the square functions $S_{\Lambda_{\kappa},\gamma}$ and $S_{\Lambda_{\kappa}^*,\gamma}$ are bounded on $L^2(\mathbb{R}^d)$. In the following theorem, which we prove in Section [3,](#page-18-0) we show that $S_{\Lambda_{\kappa},\gamma}$ and $S_{\Lambda_{\kappa}^*,\gamma}$ are also bounded on $L^p(\mathbb{R}^d)$ with $p \neq 2$ and, in particular, obtain a square function representation of $\|\Lambda_{\kappa}^{s}f\|_{p}$.

Theorem 1.3. Let $\gamma \in (0,1)$, $\alpha \in (1,2]$ with $\alpha < (d+2)/2$, $\beta \in ((d+\alpha)/2, d+M)$, and $\kappa = \Psi(\beta)$ be defined by [\(1.7\)](#page-2-2). If $\alpha = 2$, assume that the upper heat kernel bound [\(2.14\)](#page-14-2) holds. Then,

$$
||S_{\Lambda_{\kappa},\gamma}f||_{L^{p}(\mathbb{R}^{d})} \lesssim ||f||_{L^{p}(\mathbb{R}^{d})} \quad \text{for all } (d_{\beta})' < p < \infty \tag{1.19}
$$

and

$$
||S_{\Lambda^*_{\kappa},\gamma}f||_{L^p(\mathbb{R}^d)} \lesssim ||f||_{L^p(\mathbb{R}^d)} \quad \text{for all } 1 < p < d_{\beta}. \tag{1.20}
$$

Consequently, for all $s \in (0,1)$ and $(d_{\beta})' < p < d_{\beta}$,

$$
\left\| \left(\int_0^\infty t^{-2s} |t \Lambda_\kappa e^{-t \Lambda_\kappa} f|^2 \frac{dt}{t} \right)^{1/2} \right\|_{L^p(\mathbb{R}^d)} \sim \| \Lambda_\kappa^s f \|_{L^p(\mathbb{R}^d)}.
$$
\n(1.21)

Thus, to prove Theorem [1.2,](#page-4-0) it suffices to compare the square functions associated to Λ_{κ}^{s} and $(-\Delta)^{\alpha s/2}$ with each other. To that end, we prove the following two statements, called reversed and generalized Hardy inequalities. We first state the reversed Hardy inequalities, expressed in terms of the square functions $S_{\Lambda_{\kappa},\gamma}$. These inequalities give an upper bound for the difference of the square functions associated to Λ_{κ} and Λ_0 in terms of the scalar Hardy potential. While one would expect upper bounds in terms of the gradient perturbation, the bounds below are sufficient for our purposes. In particular, the advantage to incorporate the Hardy potential is that it is sign-definite and a mere multiplication operator. In turn, these properties enable us to straightforwardly prove that powers of the Hardy potential are bounded by powers of Λ_{κ} , leading to the generalized Hardy inequality in Theorem [1.5](#page-6-0) below.

Theorem 1.4 (Reversed Hardy inequality). Let $\alpha \in (1,2]$ with $\alpha < (d+2)/2$, $\beta \in ((d+2)/2)$ α /2, $d + M$), $s \in (0, 1)$, and $\kappa = \Psi(\beta)$ be defined by [\(1.7\)](#page-2-2). If $\alpha = 2$, assume that the upper heat kernel bound [\(2.14\)](#page-14-2) holds. Then the following statements hold for all $p \in ((d_{\beta})', \infty)$.

(1) If
$$
\alpha s < \alpha - 1
$$
, then
\n
$$
\left\| \left(\int_0^\infty t^{-2s} \left| \left(t \Lambda_\kappa e^{-t \Lambda_\kappa} - t \Lambda_0 e^{-t \Lambda_0} \right) f \right|^2 \frac{dt}{t} \right)^{1/2} \right\|_{L^p(\mathbb{R}^d)} \lesssim \left\| \frac{f}{|x|^{\alpha s}} \right\|_{L^p(\mathbb{R}^d)}.
$$
\n(1.22)

(2) For any
$$
\gamma \in (0, 1)
$$
 such that $1 - \gamma \leq \alpha s < \alpha - \gamma$, we have

$$
\left\| \left(\int_0^\infty t^{-2s} \left| \left(t\Lambda_\kappa e^{-t\Lambda_\kappa} - t\Lambda_0 e^{-t\Lambda_0} \right) f \right|^2 \frac{dt}{t} \right)^{1/2} \right\|_{L^p(\mathbb{R}^d)} \lesssim \left\| \frac{\left| \Lambda_0^{\frac{1-\gamma}{\alpha}} f(x) \right|}{|x|^{\alpha s + \gamma - 1}} \right\|_{L^p(\mathbb{R}^d)} + \left\| \frac{f}{|x|^{\alpha s}} \right\|_{L^p(\mathbb{R}^d)}.
$$
\n(1.23)

The proof of [\(1.22\)](#page-5-2) uses ideas in [\[FMS21,](#page-43-3) [Mer21,](#page-44-4) [BD23\]](#page-42-2) and relies on pointwise estimates for the difference of the heat kernels associated to Λ_{κ} and Λ_{0} , as well as their time derivatives. These bounds are summarized in Proposition [4.5](#page-28-1) below. To obtain these estimates, we use Duhamel's formula

$$
e^{-t\Lambda_{\kappa}}(x,y) - e^{-t\Lambda_{0}}(x,y) = -\kappa \int_{0}^{t} ds \int_{\mathbb{R}^{d}} dz \, e^{-(t-s)\Lambda_{\kappa}}(x,z) |z|^{-\alpha} z \cdot \nabla_{z} e^{-s\Lambda_{0}}(z,y).
$$

Due to the gradient perturbation, we only obtain suitable bounds if $\alpha s < \alpha - 1$. In Appendix [B.2,](#page-38-0) we discuss the necessity of this assumption and argue that our pointwise kernel bounds in Proposition [4.5](#page-28-1) do not hold if $\alpha s \geq \alpha - 1$. On the other hand, [\(1.23\)](#page-5-3) also covers the case $\alpha s \geq \alpha - 1$ as this bound also involves the operator Λ_0 on the right-hand side of the estimate. The proof of this inequality is technically more involved and does not rely on kernel bounds. Instead, we estimate $|(t\Lambda_\kappa e^{-t\Lambda_\kappa}-t\Lambda_0 e^{-t\Lambda_0})f(x)|$ pointwise. While one might suspect that there is a bound similar to [\(1.23\)](#page-5-3) involving $|||x|^{1-\gamma-\alpha s}|\Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}}f||_{p}$ on the right-hand side, we are, unfortunately, not able to prove such a bound yet. One idea to obtain such a bound is to rewrite the above Duhamel formula as

$$
e^{-t\Lambda_{\kappa}}(x,y) - e^{-t\Lambda_{0}}(x,y) = \kappa \int_{0}^{t} ds \int_{\mathbb{R}^{d}} dz \, e^{-(t-s)\Lambda_{0}}(x,z) |z|^{-\alpha} z \cdot \nabla_{z} e^{-s\Lambda_{\kappa}}(z,y).
$$

This approach requires, however, gradient bounds for $e^{-t\Lambda_{\kappa}}$, e.g., those stated in [\(B.3\)](#page-38-1), see also Appendix [B.3](#page-40-0) for a more detailed discussion. As such bounds are currently unavailable and are likely difficult to obtain, we believe it is an interesting problem to derive sharp bounds for $|\nabla_x e^{-t\Lambda_\kappa}(x, y)|$.

We proceed and now present generalized Hardy inequalities giving upper bounds for the scalar-valued Hardy potential in terms of Λ_{κ} .

Theorem 1.5 (Generalized Hardy inequality). Let $\alpha \in (1, 2]$ with $\alpha < (d+2)/2$, $\beta \in ((d+2)/2)$ α)/2, $d + M$), $s \in (0, 1]$, and $\kappa = \Psi(\beta)$ be defined by [\(1.7\)](#page-2-2). If $\alpha = 2$, assume that the upper heat kernel bound [\(2.14\)](#page-14-2) holds. If $p \in ((d_{\beta})', d/(\alpha s))$, then

$$
\| |x|^{-\alpha s} g \|_{L^p(\mathbb{R}^d)} \lesssim \| (\Lambda_\kappa)^s g \|_{L^p(\mathbb{R}^d)} \quad \text{for all } g \in C_c^\infty(\mathbb{R}^d)
$$
 (1.24)

and, for all $1 > \gamma \geq 1 - \alpha s \geq 0$,

$$
\| |x|^{\alpha(\frac{1-\gamma}{\alpha}-s)} |\Lambda_\kappa^{\frac{1-\gamma}{\alpha}} g| \|_{L^p(\mathbb{R}^d)} \lesssim \| (\Lambda_\kappa)^s g \|_{L^p(\mathbb{R}^d)} \quad \text{for all } g \in C_c^\infty(\mathbb{R}^d). \tag{1.25}
$$

To prove [\(1.15\)](#page-4-2) for $s = 1$ without using the above-outlined square function representation, the estimate $|||x|^{-\alpha}x \cdot \nabla f||_p \lesssim ||\Lambda_{\kappa}g||_p$ would be beneficial. As our proof of Theorem [1.5](#page-6-0) relies on Riesz and heat kernel bounds for $e^{-t\Lambda_{\kappa}}$, this would require gradient estimates for $e^{-t\Lambda_{\kappa}}$, too. In Proposition [C.1,](#page-42-4) we prove a new generalized Hardy inequality for $|x|^{-\alpha}x \cdot \nabla$ assuming that suitable bounds for the gradient of $e^{-t\Lambda_{\kappa}}$ are available.

With Theorems [1.3,](#page-5-0) [1.4,](#page-5-1) and [1.5](#page-6-0) at hand, we now give the

Proof of Theorem [1.2.](#page-4-0) For $s \in (0,1)$, the square function estimates in [\(1.21\)](#page-5-4), the triangle inequality, the reversed and generalized Hardy inequalities (Theorems [1.4](#page-5-1)[–1.5\)](#page-6-0) yield, whenever $\alpha s < \alpha - 1$,

$$
\|(\Lambda_0)^s u\|_{L^p(\mathbb{R}^d)} \sim \left\| \left(\int_0^\infty t^{-2s} |t\Lambda_0 e^{-t\Lambda_0} f|^2 \frac{dt}{t} \right)^{1/2} \right\|_{L^p(\mathbb{R}^d)}
$$

\n
$$
\leq \left\| \left(\int_0^\infty t^{-2s} |t\Lambda_\kappa e^{-t\Lambda_\kappa} f|^2 \frac{dt}{t} \right)^{1/2} \right\|_{L^p(\mathbb{R}^d)}
$$

\n
$$
+ \left\| \left(\int_0^\infty t^{-2s} |(t\Lambda_0 e^{-t\Lambda_0} - t\Lambda_\kappa e^{-t\Lambda_\kappa}) f|^2 \frac{dt}{t} \right)^{\frac{1}{2}} \right\|_{L^p(\mathbb{R}^d)}
$$

\n
$$
\lesssim \|(\Lambda_\kappa)^s u\|_{L^p(\mathbb{R}^d)} + \| |x|^{-\alpha s} f \|_{L^p(\mathbb{R}^d)} \lesssim \|(\Lambda_\kappa)^s u\|_{L^p(\mathbb{R}^d)}.
$$
\n(1.26)

This proves [\(1.15\)](#page-4-2). Similarly, for any $\gamma \in (0,1)$ with $1 - \gamma \leq \alpha s < \alpha - \gamma$ if $\alpha s \geq \alpha - 1$,

$$
\begin{split}\n\|(\Lambda_{\kappa})^{s}u\|_{L^{p}(\mathbb{R}^{d})} &\sim \left\| \left(\int_{0}^{\infty} t^{-2s} |t\Lambda_{\kappa}e^{-t\Lambda_{\kappa}}f|^{2} \frac{dt}{t} \right)^{1/2} \right\|_{L^{p}(\mathbb{R}^{d})} \\
&\leq \left\| \left(\int_{0}^{\infty} t^{-2s} |t\Lambda_{0}e^{-t\Lambda_{0}}f|^{2} \frac{dt}{t} \right)^{1/2} \right\|_{L^{p}(\mathbb{R}^{d})} \\
&\quad + \left\| \left(\int_{0}^{\infty} t^{-2s} |(t\Lambda_{0}e^{-t\Lambda_{0}} - t\Lambda_{\kappa}e^{-t\Lambda_{\kappa}})f|^{2} \frac{dt}{t} \right)^{\frac{1}{2}} \right\|_{L^{p}(\mathbb{R}^{d})} \\
&\lesssim \|(\Lambda_{0})^{s}u\|_{L^{p}(\mathbb{R}^{d})} + \left\| |x|^{-\alpha s}f \right\|_{L^{p}(\mathbb{R}^{d})} + \left\| \frac{|\Lambda_{0}^{\frac{1-\gamma}{\alpha}}f|}{|x|^{\alpha s + \gamma - 1}} \right\|_{L^{p}(\mathbb{R}^{d})} \\
&\lesssim \|(\Lambda_{0})^{s}u\|_{L^{p}(\mathbb{R}^{d})},\n\end{split} \tag{1.27}
$$

which proves [\(1.16\)](#page-4-3) for all $s \in (0, 1)$. For $s = 1$, the square function estimates are not necessary. We merely use the triangle inequality and Hardy's inequality (see [\(1.25\)](#page-6-1) with $\kappa = 0$, $s = 1$, and $\gamma = 0$) to get the desired estimate.

Remark 1.6. In view of the above discussion, it is an open problem to decide whether [\(1.15\)](#page-4-2) holds for a wider range of parameters. If the gradient bound in $(B.3)$ was true for some $\gamma > 2\alpha - d$, then, in view of the above proof, [\(B.17\)](#page-42-5), Proposition [C.1,](#page-42-4) and [\(1.25\)](#page-6-1), Formula [\(1.15\)](#page-4-2) would hold for all $s \leq 1$ and $\beta > d - 1$.

1.4. Discussion of Theorem [1.2.](#page-4-0) In this subsection, we put Theorem [1.2](#page-4-0) into context and compare it with earlier, related results.

1.4.1. Comparison with earlier results. We compare Theorem [1.2](#page-4-0) with the results in [\[KMV](#page-43-2)⁺18, [FMS21,](#page-43-3) [Mer21,](#page-44-4) BD23, where the L^p -Sobolev spaces generated by the Hardy operator L_{κ} = $(-\Delta)^{\alpha/2} + \kappa/|x|^{\alpha}$ with $\kappa \geq -\kappa_*$ were considered^{[3](#page-7-1)}. Using the parameterization of κ via [\(1.5\)](#page-2-4), the following statements hold for $s \in (0,1]$ and $1 < p < \infty$:

- (1) If $\frac{d}{d-(\delta_{\kappa}\vee 0)} < p < \frac{d}{(\delta_{\kappa}+\alpha s)\vee 0}$, then $||(L_0)^s f||_{L^p(\mathbb{R}^d)} \lesssim ||(L_{\kappa})^s f||_{L^p(\mathbb{R}^d)}$ for all $f \in C_c^{\infty}(\mathbb{R}^d)$ (1.28)
- (2) If $\frac{d}{d-(\delta_{\kappa}\vee 0)} < p < \frac{d}{\alpha s\vee \delta_{\kappa}},$ then $||(L_{\kappa})^s f||_{L^p(\mathbb{R}^d)} \lesssim ||(L_0)^s f||_{L^p(\mathbb{R}^d)} \text{ for all } f \in C_c^{\infty}(\mathbb{R}^d)$ (1.29)

Let us compare the assumptions on p in these statements for L_{κ} with the conditions in Theo-rem [1.2.](#page-4-0) Keeping in mind that β has a similar interpretation as δ_{κ} , we note two differences.

- (1) The lower bound $p > \frac{d}{d (\delta_{\kappa} \vee 0)}$ in [\(1.28\)](#page-7-2)–[\(1.29\)](#page-7-3) is replaced with $p > 1 \vee d/\beta$ in [\(1.15\)](#page-4-2) $(1.16).$ $(1.16).$
- (2) The upper bounds on p in [\(1.28\)](#page-7-2)–[\(1.29\)](#page-7-3) are replaced with $p < d/(\alpha s) \wedge d/((d \beta) \vee 0)$ in (1.15) – (1.16) .

The lower bounds on p in (1.28) – (1.29) in [\[Mer21,](#page-44-4) [BD23\]](#page-42-2) and in (1.15) – (1.16) in Theorem [1.2](#page-4-0) arise in the reversed Hardy inequality in [\[Mer21,](#page-44-4) Proposition 3] or [\[BD23,](#page-42-2) Theorem 4.2] and the generalized Hardy inequality [\[Mer21,](#page-44-4) Proposition 1] involving L_{κ} , and their analogs in Theorems [1.4](#page-5-1) and [1.5](#page-6-0) involving Λ_{κ} .

The different upper bounds on p in (1.28) – (1.29) come from the reversed Hardy inequality requiring $p < d/\max\{\delta_{\kappa}, 0\}$ and the generalized Hardy inequality requiring $p < d/\max\{\alpha s +$

³We point out two misprints in [\[BD23,](#page-42-2) (3)]: There, the operators $(-\Delta)^{\alpha s/2}$ and $(\mathcal{L}_a)^{\alpha s/4}$ should be replaced with $(-\Delta)^{\alpha s/4}$ and $(\mathcal{L}_a)^{s/2}$, respectively.

 δ_{κ} , 0. In the present situation, the reversed Hardy inequality holds for all $p < \infty$ and the generalized Hardy inequality holds for $p < d/(\alpha s)$. In particular, these two upper bounds are independent of β , which is due to the non-symmetry of Λ_{κ} and the resulting sharp bounds for the associated semigroup in [\(2.12\)](#page-14-1). Additionally, due to our use of square function estimates (Theorem [1.3\)](#page-5-0), we also assume $p < d/((d - \beta) \vee 0)$ for the validity of [\(1.15\)](#page-4-2)–[\(1.16\)](#page-4-3).

Another novelty compared to the previous results on equivalence of Sobolev norms is the additional restriction $\alpha s < \alpha - 1$ for the validity of [\(1.15\)](#page-4-2), which is due to the gradient in the perturbation $\kappa |x|^{-\alpha} x \cdot \nabla$ as we discussed in the proof of Theorem [1.2](#page-4-0) above.

1.4.2. Impact of Theorem [1.2.](#page-4-0) Homogeneous operators like Λ_{κ} or L_{κ} frequently arise as model operators or scaling limits in more complicated problems. Hence, their analysis is paramount to advance studies of the more complicated problems they originate from. In many applications, one has to deal with functions, e.g., powers greater than one, of these operators, which are, however, difficult to handle. The equivalence of Sobolev norm results in [\[KMV](#page-43-2)+18, [FMS21,](#page-43-3) [Mer21,](#page-44-4) [BD23,](#page-42-2) [FM23,](#page-43-4) [BM23\]](#page-43-5) and Theorem [1.2](#page-4-0) here are useful to overcome this obstacle as they allow to replace the usually difficult to handle operator functions of Λ_{κ} or L_{κ} with easier to handle operator functions of $(-\Delta)^{\alpha/2}$. For instance, the equivalence of Sobolev norms for L_{κ} established in [\[KMV](#page-43-2)+18] for $\alpha = 2$ was paramount for Killip, Miao, Murphy, Visan, Zhang, and Zheng [\[KMV](#page-43-14)⁺17, [KMVZ17\]](#page-43-15) to study global well-posedness and scattering of nonlinear Schrödinger equations with the inverse-square potentials $\kappa/|x|^2$. In mathematical physics, the equivalence of Sobolev norms for L_{κ} with $\alpha = 1$ in [\[FMS21\]](#page-43-3) was crucial, e.g., to study the electron density of large, relativistically described atoms close to the nucleus, and, in particular, to prove Lieb's strong Scott conjecture [\[Lie81,](#page-44-10) [Sim84\]](#page-44-11) in [\[FMSS20,](#page-43-16) [MS22\]](#page-44-12); see also [\[FMS23b\]](#page-43-17) for a shorter proof and the recent review [\[FMS23a\]](#page-43-18) for an introduction to the Scott conjectures. Therefore, we believe that the results in the present paper will be useful to study nonlinear, dispersive equations involving Λ_{κ} or to advance the spectral theory of many-particle Schrödinger operators based on Λ_{κ} .

Organization. In Section [2,](#page-8-1) we recall kernel bounds for $e^{-t\Lambda_{\kappa}}$ and use them, together with a Phragmén–Lindelöf argument, to prove bounds for complex-time heat kernels. We use these bounds to prove novel bounds for the kernels of $(t\Lambda_{\kappa})^k e^{-t\Lambda_{\kappa}}$ and corresponding $L^p \to L^q$ estimates. We apply the latter bounds in Section [3](#page-18-0) to prove the square function estimates for $(\Lambda_{\kappa})^s$ in Theorem [1.3.](#page-5-0) In Section [4,](#page-24-0) we prove bounds for the kernel of the difference $t\Lambda_0e^{-t\Lambda_0}$ – $t\Lambda_{\kappa}e^{-t\Lambda_{\kappa}}$. These bounds are crucial to prove the reversed Hardy inequality (Theorem [1.4\)](#page-5-1) in Section [5.](#page-32-0) In Section [6,](#page-34-0) we prove the generalized Hardy inequality (Theorem [1.5\)](#page-6-0). In the appendices, we prove auxiliary results and discuss the necessity of $\alpha s < \alpha - 1$ for the validity of [\(1.15\)](#page-4-2).

2. ESTIMATES INVOLVING THE HEAT KERNEL OF Λ_{κ}

In this section, we collect known pointwise estimates for the heat kernel $e^{-t\Lambda_{\kappa}}$ and derive novel bounds for the complex-time heat kernel and $(t\Lambda_{\kappa})^k e^{-t\Lambda_{\kappa}}$ with $k \in \mathbb{N}$.

To begin, we introduce some notation. We denote the average of a measurable function f over a measurable set E with $0 < |E| < \infty$ by

$$
\oint_{E} f(x)dx = \frac{1}{|E|} \int_{E} f(x)dx.
$$

Given a ball B, we associate annuli $S_j(B) := 2^j B \setminus 2^{j-1}B$ for $j = 1, 2, 3, \ldots$ For $j = 0$, we write $S_0(B) = B$. For $p > 0$, the Hardy-Littlewood maximal function \mathcal{M}_p is defined as

$$
\mathcal{M}_p f(x) := \sup_{B \ni x} \left(\frac{1}{|B|} \int_B |f(y)|^p \, dy \right)^{1/p}, \ x \in \mathbb{R}^d,
$$

where the supremum is taken over all balls B containing x. When $r = 1$, we write M instead of \mathcal{M}_1 .

2.1. Spatially averaged estimates. In this section, we prove spatially weighted estimates for a family of operators whose kernels behave similarly to the heat kernel bounds of Λ_{κ} , presented in [\(2.12\)](#page-14-1) below. We will use them to prove bounds for time-derivatives of the heat kernel of Λ_{κ} in Proposition [2.9](#page-17-0) and the square function estimates in Section [3](#page-18-0) below.

We start with some auxiliary estimates, whose manual proofs we omit.

Lemma 2.1. Let $d \in \mathbb{N}$ and $\alpha > 0$. Then the following estimates hold.

(a) For all $\varepsilon \in (0,d)$ there exists $C = C_{\varepsilon} > 0$ so that for all $t > 0$,

$$
\int_{B(0,t)} \left(\frac{t}{|x|}\right)^{d-\varepsilon} dx \le Ct^d.
$$

(b) For all $\varepsilon > 0$, there exists $C = C_{\varepsilon} > 0$ such that

$$
\int_{\mathbb{R}^d} \frac{1}{t^{d/\alpha}} \Big(\frac{t^{1/\alpha} + |x - y|}{t^{1/\alpha}} \Big)^{-d-\varepsilon} dy \le C
$$

for all $x \in \mathbb{R}^d$.

(c) For all $\varepsilon > 0$, there exists $C = C_{\varepsilon} > 0$ such that

$$
\int_{\mathbb{R}^d} \frac{1}{t^{d/\alpha}} \Big(\frac{t^{1/\alpha} + |x - y|}{t^{1/\alpha}}\Big)^{-d-\varepsilon} f(y) dy \le C\mathcal{M}f(x)
$$

for all $t > 0$, $x \in \mathbb{R}^d$, and $f \in L^1_{loc}(\mathbb{R}^d)$.

Using these estimates, we prove the following estimates.

Theorem 2.2. Let $d \in \mathbb{N}$, $\alpha > 0$, and $\beta > 0$. Let $\{T_t\}_{t>0}$ be a family of linear operators on $L^2(\mathbb{R}^d)$ with their associated kernels $T_t(x, y)$. Assume that there exist $C > 0$ and $\theta \in (-\infty, d)$ such that for all $t > 0$ and $x, y \in \mathbb{R}^d \setminus \{0\},\$

$$
|T_t(x,y)| \le Ct^{-d/\alpha} \Big(\frac{t^{1/\alpha} + |x-y|}{t^{1/\alpha}}\Big)^{-d-\beta} \Big(1 + \frac{t^{1/\alpha}}{|y|}\Big)^{\theta}.
$$
 (2.1)

Assume that $\frac{d}{d-\theta} \vee 1 < p \le q < \infty$, where d_{θ} is defined as in [\(1.13\)](#page-4-4). Then, for any ball B, every $t > 0$, and $j \geq 2$, we have

$$
\left(\int_{S_j(B)} |T_t f|^q\right)^{1/q} \lesssim \max\left\{\left(\frac{r_B}{t^{1/\alpha}}\right)^{d/p}, \left(\frac{r_B}{t^{1/\alpha}}\right)^d\right\} \left(1 + \frac{2^j r_B}{t^{1/\alpha}}\right)^{-d-\beta} \left(\int_B |f|^p\right)^{1/p} \tag{2.2}
$$

for all $f \in L^p(\mathbb{R}^d)$ supported in B, and

$$
\left(\int_{B} |T_t f|^q\right)^{1/q} \lesssim \max\left\{ \left(\frac{2^j r_B}{t^{1/\alpha}}\right)^d, \left(\frac{2^j r_B}{t^{1/\alpha}}\right)^{d/p} \right\} \left(1 + \frac{2^j r_B}{t^{1/\alpha}}\right)^{-d-\beta} \left(\int_{S_j(B)} |f|^p\right)^{1/p} \tag{2.3}
$$

for all $f \in L^p(\mathbb{R}^d)$ supported in $S_j(B)$.

Proof. Note that the estimate (2.1) boils down to

$$
|T_t(x,y)| \le Ct^{-d/\alpha} \Big(\frac{t^{1/\alpha}+|x-y|}{t^{1/\alpha}}\Big)^{-d-\beta},
$$

as long as $\theta \in (-\infty, 0)$.

Hence, it suffices to prove the theorem for $\theta \in [0, d)$. In this case, $\frac{d}{d-\theta} \vee 1 = \frac{d}{d-\theta}$. Since $|x - y| \sim 2^{j} r_B$ whenever $x \in S_j(B)$ and $y \in B$,

$$
||T_t f||_{L^q(S_j(B))} \le \left\{ \int_{S_j(B)} \left[\int_B t^{-d/\alpha} \left(\frac{t^{1/\alpha} + |x - y|}{t^{1/\alpha}} \right)^{-d-\beta} \left(1 + \frac{t^{1/\alpha}}{|y|} \right)^{\theta} |f(y)| dy \right]^q dx \right\}^{1/q}
$$

$$
\le t^{-d/\alpha} \left(1 + \frac{2^j r_B}{t^{1/\alpha}} \right)^{-d-\beta} \left\{ \int_{S_j(B)} \left[\int_B \left(1 + \frac{t^{1/\alpha}}{|y|} \right)^{\theta} |f(y)| dy \right]^q dx \right\}^{1/q}
$$

$$
\le t^{-d/\alpha} |2^j B|^{1/q} \left(1 + \frac{2^j r_B}{t^{1/\alpha}} \right)^{-d-\beta} \int_B \left(1 + \frac{t^{1/\alpha}}{|y|} \right)^{\theta} |f(y)| dy. \tag{2.4}
$$

We now estimate the integral on the right hand side of [\(2.4\)](#page-10-0). We have

$$
\int_{B} \left(1 + \frac{t^{1/\alpha}}{|y|}\right)^{\theta} |f(y)| dy
$$
\n
$$
= \int_{B \cap B(0, t^{1/\alpha})} \left(1 + \frac{t^{1/\alpha}}{|y|}\right)^{\theta} |f(y)| dy + \int_{B \cap B(0, t^{1/\alpha})^c} \left(1 + \frac{t^{1/\alpha}}{|y|}\right)^{\theta} |f(y)| dy
$$
\n
$$
=: I_1 + I_2.
$$

By Hölder's inequality and Lemma [2.1](#page-9-2) (since $\theta p' < d$),

$$
I_1 \leq \Big[\int_{B(0,t^{1/\alpha})} \Big(1+\frac{t^{1/\alpha}}{|x|}\Big)^{\theta p'}dy\Big]^{1/p'}\|f\|_{L^p(B)} \leq Ct^{\frac{d}{\alpha p'}}|B|^{1/p}\Big(\int_B|f|^p\Big)^{1/p}.
$$

We proceed to consider I_2 . Here, we have $\left(1+\frac{t^{1/\alpha}}{|u|}\right)$ $|y|$ $\Big)^\theta \sim 1.$ Hence,

$$
I_2 \lesssim ||f||_{L^1(B)} \lesssim |B| \left(\int_B |f|^p \right)^{1/p}.
$$

Consequently,

$$
\int_{B} \left(1 + \frac{t^{1/\alpha}}{|y|}\right)^{\theta} |f(y)| dy \lesssim \left[t^{\frac{d}{\alpha p'}} |B|^{1/p} + |B|\right] \left(\int_{B} |f|^p\right)^{1/p}.
$$

Inserting this into [\(2.4\)](#page-10-0) gives

$$
||T_t f||_{L^q(S_j(B))} \leq t^{-d/\alpha} |2^j B|^{1/q} \left(1 + \frac{2^j r_B}{t^{1/\alpha}}\right)^{-d-\beta} \left[t^{\frac{d}{\alpha p'}} |B|^{1/p} + |B| \right] \left(\int_B |f|^p \right)^{1/p}
$$

$$
\leq |2^j B|^{1/q} \left(1 + \frac{2^j r_B}{t^{1/\alpha}}\right)^{-d-\beta} \left[t^{-\frac{d}{\alpha p}} |B|^{1/p} + t^{-d/\alpha} |B| \right] \left(\int_B |f|^p \right)^{1/p}
$$

$$
\leq |2^j B|^{1/q} \left(1 + \frac{2^j r_B}{t^{1/\alpha}}\right)^{-d-\beta} \max \left\{ \left(\frac{r_B}{t^{1/\alpha}} \right)^{d/p}, \left(\frac{r_B}{t^{1/\alpha}} \right)^d \right\} \left(\int_B |f|^p \right)^{1/p}.
$$

Similarly, we obtain

$$
||T_t f||_{L^q(B)} \le \left\{ \int_B \Big[\int_{S_j(B)} t^{-d/\alpha} \Big(\frac{t^{1/\alpha} + |x - y|}{t^{1/\alpha}} \Big)^{-d-\beta} \Big(1 + \frac{t^{1/\alpha}}{|y|} \Big)^\theta |f(y)| dy \Big]^q dx \right\}^{1/q}
$$

$$
\le t^{-d/\alpha} \Big(1 + \frac{2^j r_B}{t^{1/\alpha}} \Big)^{-d-\beta} \left\{ \int_B \Big[\int_{S_j(B)} \Big(1 + \frac{t^{1/\alpha}}{|y|} \Big)^\theta |f(y)| dy \Big]^q dx \right\}^{1/q}
$$

$$
\le t^{-d/\alpha} |B|^{1/q} \Big(1 + \frac{2^j r_B}{t^{1/\alpha}} \Big)^{-d-\beta} \int_{S_j(B)} \Big(1 + \frac{t^{1/\alpha}}{|y|} \Big)^\theta |f(y)| dy. \tag{2.5}
$$

We now write

$$
\int_{S_j(B)} \left(1 + \frac{t^{1/\alpha}}{|y|} \right)^{\theta} |f(y)| dy
$$
\n
$$
= \int_{S_j(B) \cap B(0, t^{1/\alpha})} \left(1 + \frac{t^{1/\alpha}}{|y|} \right)^{\theta} |f(y)| dy + \int_{S_j(B) \cap B(0, t^{1/\alpha})^c} \left(1 + \frac{t^{1/\alpha}}{|y|} \right)^{\theta} |f(y)| dy
$$
\n
$$
=: II_1 + II_2.
$$

Similarly to the term I_1 , by Hölder's inequality and Lemma [2.1,](#page-9-2)

$$
II_1 \leq \Big[\int_{B(0,t^{1/\alpha})} \Big(1 + \frac{t^{1/\alpha}}{|x|} \Big)^{\theta p'} dy \Big]^{1/p'} \|f\|_{L^p(S_j(B))}
$$

$$
\leq C t^{\frac{d}{\alpha p'}} |2^j B|^{1/p} \Big(\int_{S_j(B)} |f|^p \Big)^{1/p}.
$$

We proceed to consider II_2 . Here, $\left(1 + \frac{t^{1/\alpha}}{|u|}\right)$ $|y|$ $\int^{\theta} \sim 1$. Therefore,

$$
II_2 \lesssim ||f||_{L^1(S_j(B))} \lesssim |2^j B| \Big(\int_{S_j(B)} |f|^p \Big)^{1/p}.
$$

The estimates for II_1 and II_2 yield

$$
\int_{S_j(B)} \left(1 + \frac{t^{1/\alpha}}{|y|}\right)^{\theta} |f(y)| dy \lesssim \left[t^{\frac{d}{\alpha p'}} |2^j B|^{1/p} + |2^j B| \right] \left(\oint_{2^j B} |f|^p \right)^{1/p}.
$$

Inserting this into [\(2.5\)](#page-11-0) gives

$$
||T_t f||_{L^q(B)} \le t^{-d/\alpha} |B|^{1/q} \left(1 + \frac{2^j r_B}{t^{1/\alpha}}\right)^{-d-\beta} \left[t^{\frac{d}{\alpha p'}} |2^j B|^{1/p} + |2^j B|\right] \left(\int_{S_j(B)} |f|^p\right)^{1/p}
$$

\n
$$
\le |B|^{1/q} \left(1 + \frac{2^j r_B}{t^{1/\alpha}}\right)^{-d-\beta} \left[t^{-\frac{d}{\alpha p}} |2^j B|^{1/p} + t^{-d/\alpha} |2^j B|\right] \left(\int_{S_j(B)} |f|^p\right)^{1/p}
$$

\n
$$
\le |B|^{1/q} \left(1 + \frac{2^j r_B}{t^{1/\alpha}}\right)^{-d-\beta} \max\left\{\left(\frac{2^j r_B}{t^{1/\alpha}}\right)^{d/p}, \left(\frac{2^j r_B}{t^{1/\alpha}}\right)^d\right\} \left(\int_{S_j(B)} |f|^p\right)^{1/p}.
$$

\nThis concludes the proof.

We now prove similar estimates for kernels of operators which are formally adjoint to those with integral kernels obeying the assumptions in Theorem [2.2.](#page-9-3)

Theorem 2.3. Let $d \in \mathbb{N}$, $\alpha > 0$, and $\beta > 0$. Let $\{S_t\}_{t>0}$ be a family of linear operators on $L^2(\mathbb{R}^d)$ with their associated kernels $S_t(x, y)$. Assume that there exist $C > 0$ and $\theta \in (-\infty, d)$ such that for all $t > 0$ and $x, y \in \mathbb{R}^d \setminus \{0\},\$

$$
|S_t(x,y)| \le Ct^{-d/\alpha} \Big(\frac{t^{1/\alpha} + |x-y|}{t^{1/\alpha}}\Big)^{-d-\beta} \Big(1 + \frac{t^{1/\alpha}}{|x|}\Big)^{\theta}.
$$
 (2.6)

Assume that $1 < p \le q < \frac{d}{\theta \vee 0}$ (with the convention $d/0 = \infty$). Then, for any ball B, every $t > 0$, and $j \geq 2$, we have

$$
\left(\int_{S_j(B)} |S_t f|^q\right)^{1/q} \lesssim \left(\frac{r_B}{t^{1/\alpha}}\right)^d \left(1 + \frac{t^{1/\alpha}}{2^j r_B}\right)^{d/q} \left(1 + \frac{2^j r_B}{t^{1/\alpha}}\right)^{-d-\beta} \left(\int_B |f|^p\right)^{1/p} \tag{2.7}
$$

for all $f \in L^p(\mathbb{R}^d)$ supported in B, and

$$
\left(\int_{B} |S_t f|^q\right)^{1/q} \lesssim \left(\frac{2^j r_B}{t^{1/\alpha}}\right)^d \left(1 + \frac{t^{1/\alpha}}{r_B}\right)^{d/q} \left(1 + \frac{2^j r_B}{t^{1/\alpha}}\right)^{-d-\beta} \left(\int_{S_j(B)} |f|^p\right)^{1/p} \tag{2.8}
$$

for all $f \in L^p(\mathbb{R}^d)$ supported in $S_j(B)$.

Proof. Note that the estimate [\(2.6\)](#page-11-1) boils down to

$$
|S_t(x,y)| \le Ct^{-d/\alpha} \Big(\frac{t^{1/\alpha} + |x-y|}{t^{1/\alpha}}\Big)^{-d-\beta}
$$

as long as $\theta \in (-\infty, 0)$. Hence, it suffices to prove the theorem for $\theta \in [0, d)$. In this case, $\frac{d}{\theta \vee 0} = \frac{d}{\theta}$ with the convention $d/0 = \infty$.

Since $|x - y| \sim 2^{j} r_B$ whenever $x \in S_j(B)$ and $y \in B$,

$$
||S_{t}f||_{L^{q}(S_{j}(B))} \leq \left\{ \int_{S_{j}(B)} \left[\int_{B} t^{-d/\alpha} \left(\frac{t^{1/\alpha} + |x - y|}{t^{1/\alpha}} \right)^{-d-\beta} \left(1 + \frac{t^{1/\alpha}}{|x|} \right)^{\theta} |f(y)| dy \right]^{q} dx \right\}^{1/q}
$$

$$
\leq t^{-d/\alpha} \left(1 + \frac{2^{j} r_{B}}{t^{1/\alpha}} \right)^{-d-\beta} \left\{ \int_{S_{j}(B)} \left[\int_{B} \left(1 + \frac{t^{1/\alpha}}{|x|} \right)^{\theta} |f(y)| dy \right]^{q} dx \right\}^{1/q}
$$

$$
\leq t^{-d/\alpha} \left(1 + \frac{2^{j} r_{B}}{t^{1/\alpha}} \right)^{-d-\beta} \left[\int_{S_{j}(B)} \left(1 + \frac{t^{1/\alpha}}{|x|} \right)^{\theta q} dx \right]^{1/q} ||f||_{L^{1}(B)}.
$$

(2.9)

On the other hand, by Lemma [2.1,](#page-9-2)

$$
\int_{S_j(B)} \left(1 + \frac{t^{\frac{1}{\alpha}}}{|x|}\right)^{\theta q} dx = \int_{S_j(B) \cap B(0, t^{\frac{1}{\alpha}})} \left(1 + \frac{t^{\frac{1}{\alpha}}}{|x|}\right)^{\theta q} dx + \int_{S_j(B) \cap B(0, t^{1/\alpha})^c} \left(1 + \frac{t^{\frac{1}{\alpha}}}{|x|}\right)^{\theta q} dx
$$
\n
$$
\lesssim \int_{B(0, t^{1/\alpha})} \left(1 + \frac{t^{1/\alpha}}{|x|}\right)^{\theta q} dx + \int_{S_j(B)} dx
$$
\n
$$
\lesssim t^{d/\alpha} + |2^j B| \sim |2^j B| \left(1 + \frac{t^{1/\alpha}}{2^j r_B}\right)^d.
$$
\n(2.10)

Inserting this into [\(2.9\)](#page-12-0), we get

$$
||S_t f||_{L^q(S_j(B))} \lesssim t^{-d/\alpha} |2^j B|^{1/q} \left(1 + \frac{2^j r_B}{t^{1/\alpha}}\right)^{-d-\beta} \left(1 + \frac{t^{1/\alpha}}{2^j r_B}\right)^{d/q} ||f||_{L^1(B)}
$$

$$
\lesssim t^{-d/\alpha} |2^j B| \left(1 + \frac{2^j r_B}{t^{1/\alpha}}\right)^{-d-\beta} \left(1 + \frac{t^{1/\alpha}}{2^j r_B}\right)^{d/q} |B| \left(\int_B |f|^p\right)^{1/p}
$$

$$
\lesssim |2^j B|^{1/q} \left(\frac{r_B}{t^{1/\alpha}}\right)^d \left(1 + \frac{t^{1/\alpha}}{2^j r_B}\right)^{d/q} \left(1 + \frac{2^j r_B}{t^{1/\alpha}}\right)^{-d-\beta} \left(\int_B |f|^p\right)^{1/p},
$$

where we used Hölder's inequality in the second inequality. This proves (2.7) .

Similarly,

$$
||T_t f||_{L^q(B)} \le \left\{ \int_B \Big[\int_{S_j(B)} t^{-d/\alpha} \Big(\frac{t^{1/\alpha} + |x - y|}{t^{1/\alpha}} \Big)^{-d-\beta} \Big(1 + \frac{t^{1/\alpha}}{|x|} \Big)^\theta |f(y)| dy \Big]^q dx \right\}^{1/q}
$$

$$
\le t^{-d/\alpha} \Big(1 + \frac{2^j r_B}{t^{1/\alpha}} \Big)^{-d-\beta} \left\{ \int_B \Big[\int_{S_j(B)} \Big(1 + \frac{t^{1/\alpha}}{|y|} \Big)^\theta |f(y)| dy \Big]^q dx \right\}^{1/q}
$$

$$
\le t^{-d/\alpha} \Big(1 + \frac{2^j r_B}{t^{1/\alpha}} \Big)^{-d-\beta} \Big[\int_B \Big(1 + \frac{t^{1/\alpha}}{|x|} \Big)^\theta q dx \Big]^{1/q} ||f||_{L^1(S_j(B))}.
$$
 (2.11)

Similarly to [\(2.10\)](#page-12-2),

$$
\int_{B} \left(1 + \frac{t^{1/\alpha}}{|x|}\right)^{\theta q} dx \lesssim |B| + t^{\frac{d}{\alpha}} \sim |B| \left(1 + \frac{t^{1/\alpha}}{r_B}\right)^d.
$$

Therefore,

$$
||T_t f||_{L^q(B)} \lesssim t^{-d/\alpha} |B|^{1/q} \Big(1 + \frac{t^{1/\alpha}}{r_B} \Big)^{d/q} \Big(1 + \frac{2^j r_B}{t^{1/\alpha}} \Big)^{-d-\beta} ||f||_{L^1(S_j(B))}
$$

$$
\lesssim t^{-d/\alpha} |B|^{1/q} \Big(1 + \frac{t^{1/\alpha}}{r_B} \Big)^{d/q} \Big(1 + \frac{2^j r_B}{t^{1/\alpha}} \Big)^{-d-\beta} |2^j B| \Big(\int_{S_j(B)} |f|^p \Big)^{1/p}
$$

$$
\lesssim |B|^{1/q} \Big(\frac{2^j r_B}{t^{1/\alpha}} \Big)^d \Big(1 + \frac{t^{1/\alpha}}{r_B} \Big)^{d/q} \Big(1 + \frac{2^j r_B}{t^{1/\alpha}} \Big)^{-d-\beta} \Big(\int_{S_j(B)} |f|^p \Big)^{1/p}.
$$

This concludes the proof. \Box

Theorem 2.4. Let $d \in \mathbb{N}$, $\alpha > 0$, and $\beta > 0$.

(a) Let $\{T_t\}_{t>0}$ be a family of linear operators as in Theorem [2.2](#page-9-3) satisfying [\(2.1\)](#page-9-1). Then, for any $\frac{d}{d-\theta} < p \le q \le \infty$, we have

$$
||T_t||_{p \to q} \lesssim t^{-\frac{d}{\alpha}(\frac{1}{p} - \frac{1}{q})}.
$$

(b) Let $\{S_t\}_{t>0}$ be a family of linear operators as in Theorem [2.3](#page-11-2) satisfying [\(2.6\)](#page-11-1). Then, for any $1 \leq p \leq q < \frac{d}{\theta}$, we have

$$
||S_t||_{p \to q} \lesssim t^{-\frac{d}{\alpha}(\frac{1}{p} - \frac{1}{q})}.
$$

Proof. The proof of (b) is similar to that of (a) with similar modifications as in the proof of Theorem [2.3.](#page-11-2) Hence, we only provide the proof of (a). For $f \in L^p(\mathbb{R}^d)$, we have

$$
||T_t f||_{L^q(\mathbb{R}^d)} \leq \left\{ \int_{\mathbb{R}^d} \Big[\int_{\mathbb{R}^d} t^{-d/\alpha} \Big(\frac{t^{1/\alpha} + |x - y|}{t^{1/\alpha}} \Big)^{-d-\beta} \Big(1 + \frac{t^{1/\alpha}}{|y|} \Big)^{\theta} |f(y)| dy \Big]^q dx \right\}^{1/q}
$$

\n
$$
\leq \left\{ \int_{\mathbb{R}^d} \Big[\int_{B(0,t^{1/\alpha})} t^{-d/\alpha} \Big(\frac{t^{1/\alpha} + |x - y|}{t^{1/\alpha}} \Big)^{-d-\beta} \Big(1 + \frac{t^{1/\alpha}}{|y|} \Big)^{\theta} |f(y)| dy \Big]^q dx \right\}^{1/q}
$$

\n
$$
+ \left\{ \int_{\mathbb{R}^d} \Big[\int_{B(0,t^{1/\alpha})^c} t^{-\frac{d}{\alpha}} \Big(\frac{t^{\frac{1}{\alpha}} + |x - y|}{t^{\frac{1}{\alpha}}} \Big)^{-d-\beta} \Big(1 + \frac{t^{1/\alpha}}{|y|} \Big)^{\theta} |f(y)| dy \Big]^q dx \right\}^{1/q}
$$

\n
$$
=: I_1 + I_2.
$$

We first consider I_1 . By Minkowski's inequality,

$$
I_1 \lesssim \int_{B(0,t^{1/\alpha})^c} \left\{ \int_{\mathbb{R}^d} \left[t^{-d/\alpha} \left(\frac{t^{1/\alpha} + |x-y|}{t^{1/\alpha}} \right)^{-d-\beta} \right]^q dx \right\}^{1/q} \left(1 + \frac{t^{1/\alpha}}{|y|} \right)^{\theta} |f(y)| dy
$$

$$
\lesssim t^{\frac{d}{\alpha q} - \frac{d}{\alpha}} \int_{B(0,t^{1/\alpha})^c} \left(1 + \frac{t^{1/\alpha}}{|y|} \right)^{\theta} |f(y)| dy.
$$

Together with Hölder's inequality and Lemma [2.1](#page-9-2) (since $\theta p' < d$), this estimate yields

$$
I_1 \lesssim t^{\frac{d}{\alpha q} - \frac{d}{\alpha}} \Big[\int_{B(0, t^{1/\alpha})} \left(1 + \frac{t^{1/\alpha}}{|x|} \right)^{\theta p'} dy \Big]^{1/p'} \|f\|_p
$$

$$
\lesssim t^{\frac{d}{\alpha q} - \frac{d}{\alpha}} t^{\frac{d}{\alpha p'}} \|f\|_p \lesssim t^{-\frac{d}{\alpha}(\frac{1}{p} - \frac{1}{q})} \|f\|_p.
$$

We proceed to consider I_2 . Here, we have $\left(1+\frac{t^{1/\alpha}}{|u|}\right)$ $|y|$ \int_{0}^{θ} ~ 1 whenever $y \in B(0, t^{1/\alpha})^c$. Hence,

$$
I_2 \lesssim \left\{ \int_{\mathbb{R}^d} \left[\int_{\mathbb{R}^d} t^{-d/\alpha} \left(\frac{t^{1/\alpha} + |x - y|}{t^{1/\alpha}} \right)^{-d-\beta} |f(y)| dy \right]^q dx \right\}^{1/q}
$$

$$
\sim ||e^{-t(-\Delta)^{\alpha/2}} |f||_q \lesssim t^{-\frac{d}{\alpha}(\frac{1}{p} - \frac{1}{q})} ||f||_p.
$$

This concludes the proof. \Box

2.2. Pointwise estimates. In the following, $p_t(x, y)$ and $\tilde{p}_t(x, y)$ shall denote the kernels of $e^{-t\Lambda_{\kappa}}$ and $e^{-t\Lambda_{0}},$ respectively, i.e.,

$$
p_t(x, y) := e^{-t\Lambda_{\kappa}}(x, y)
$$
 and $\tilde{p}_t(x, y) := e^{-t\Lambda_0}(x, y)$.

As indicated in Section [1,](#page-1-0) we only consider $\kappa < \kappa_c$, i.e., $\beta > (d+\alpha)/2$, since in that situation, $e^{-tL_{\kappa}}$ is a holomorphic semigroup in $L^2(\mathbb{R}^d)$ and a C_0 semigroup on $L^r(\mathbb{R}^d)$ for all $r \in [2,\infty)$. As in [\[KMV](#page-43-2)⁺18, [FMS21,](#page-43-3) [Mer21,](#page-44-4) [BD23,](#page-42-2) [FM23,](#page-43-4) [BM23\]](#page-43-5), a paramount role to prove the reversed and generalized Hardy inequalities is played by bounds for $p_t(x, y)$. According to [\[KSS21,](#page-44-7) [KS23\]](#page-44-3), we have,

$$
p_t(x, y) \sim \tilde{p}_t(x, y) \cdot \left(1 \wedge \frac{|y|}{t^{1/\alpha}}\right)^{\beta - d} \quad \text{for all } \alpha \in (1, 2), t > 0, \text{ and } x, y \in \mathbb{R}^d \tag{2.12}
$$

with

$$
\tilde{p}_t(x,y) \sim t^{-d/\alpha} \left(\frac{t^{1/\alpha}}{t^{1/\alpha} + |x-y|}\right)^{d+\alpha} \quad \text{for all } \alpha \in (0,2), t > 0, \text{ and } x, y \in \mathbb{R}^d,
$$
 (2.13)

see, e.g., [\[BG60\]](#page-42-6). When $\alpha = 2$, corresponding estimates for $e^{-t\Lambda_{\kappa}}$ are unavailable to the best of our knowledge. However, in view of the factorization of the bounds in [\(2.12\)](#page-14-1) as product of the heat kernel with $\kappa = 0$ times a singular weight, we expect

$$
t^{-d/2} \left(1 \wedge \frac{|y|}{t^{1/2}}\right)^{\beta-d} e^{-c_1|x-y|^2/t} \lesssim e^{-t\Lambda_\kappa}(x, y) \lesssim t^{-d/2} \left(1 \wedge \frac{|y|}{t^{1/2}}\right)^{\beta-d} e^{-c_2|x-y|^2/t}
$$
\nfor $\alpha = 2$, all $t > 0$, and all $x, y \in \mathbb{R}^d$,\n
$$
(2.14)
$$

and certain $c_1 \geq c_2 > 0$, depending only on d and β .

Remark 2.5. Metafune, Negro, Sobajima, and Spina [\[MSS17,](#page-44-13) [MNS18\]](#page-44-14) proved matching upper and lower bounds for the heat kernel associated to

$$
-\Delta + (a-1)\sum_{j,\ell=1}^d \frac{x_j x_\ell}{|x|^2} \frac{\partial}{\partial x_j} \frac{\partial}{\partial x_\ell} + \kappa |x|^{-2} x \cdot \nabla + \frac{b}{|x|^2} \quad \text{in } L^2(\mathbb{R}^d, |x|^{(d-1-\kappa)/a-(d-1)} dx). \tag{2.15}
$$

In particular, in this weighted L^2 -space, the operator can be realized as a self-adjoint operator. Thus, unlike (2.12) or (2.14) , the corresponding heat kernel bounds are symmetric in x and y. We refrain from attempting to prove an analog of Theorem [1.2](#page-4-0) for the operator in [\(2.15\)](#page-14-3) to keep the paper at a reasonable length. Such a proof likely requires a combination of techniques in $KMV+18$ and the present ones, together with more technical work due to the necessity to work in weighted Lebesgue spaces. Additionally, new ideas will be needed to deal with the case $a \neq 1$.

Remark 2.6. In the critical case $\alpha = 1$ and the supercritical case $\alpha \in (0,1)$, Kinzebulatov, Madou, and Semenov [\[KMS24,](#page-43-1) Theorem 1] showed $e^{-t\Lambda_{\kappa}}(x,y) \lesssim t^{-d/\alpha}(1+\vert y\vert/t^{1/\alpha})^{\beta-d}$ for $t \leq 1, \ \kappa < 0, \ d \geq 3$, and $\kappa = \Psi(\beta)$ defined in $(1.7)^4$ $(1.7)^4$. This bound is insufficient for our purposes since it does not decay at all for $|x - y| \gtrsim t$ and is only stated for $t \lesssim 1$. The decay in $|x-y|/t^{1/\alpha}$ is, however, crucial to estimate several integrals discussed below. We do not know if the estimate in [\[KMS24\]](#page-43-1) can be improved in this regard. Moreover, it remains to be explored if their estimate holds for $t \geq 1$, and if a corresponding lower bound holds. For Kato-class gradient perturbations, this possibility was ruled out in [\[BJ07,](#page-43-19) p. 181] by noting that the heat kernel of $(-\Delta)^{\alpha/2} + |x|^{-1}x \cdot \nabla$ in $d = 1$ equals $e^{-t\Lambda_0}(y - (x + t))$ for all $x, y \in \mathbb{R}$, $t > 0$, and $\alpha \in (0, 2)$. Thus, taking $x = y$ and considering $t \searrow 0$ and $\alpha \in (0, 1]$, this kernel is bounded from above and below by $t^{-\alpha}$, whereas the estimate in [\[KMS24,](#page-43-1) Theorem 1], if it was also true for $d = 1$, $\kappa > 0$, and $t \gtrsim 1$, would predict an upper bound of the order $t^{-1/\alpha}$.

Remark 2.7. The heat kernels of $(-\Delta)^{\alpha/2} + b(x) \cdot \nabla$ with less singular, decaying drifts $b : \mathbb{R}^d \to$ \mathbb{R}^d are bounded from above and below by multiples of $e^{-t\Lambda_0}(x, y)$, at least for small times; see [\[BJ07\]](#page-43-19) for $\alpha \in (1, 2)$ and, for instance, [\[GL90,](#page-43-20) [Zha95,](#page-44-15) [Zha97,](#page-44-16) [KS06\]](#page-44-17) for $\alpha = 2$.

Given the above bounds for $e^{-t\Lambda_{\kappa}}$ and the estimates in Subsection [2.1,](#page-9-0) we now prove further estimates for $e^{-t\Lambda_{\kappa}}$. First, by Theorem [2.4,](#page-13-0) we see that $e^{-t\Lambda_{\kappa}}$ is $L^p(\mathbb{R}^d)$ -bounded whenever $p \in (d/\beta, \infty]$. We now prove bounds for time derivatives of $e^{-t\Lambda_{\kappa}}$. To that end, we use that Λ_{κ} generates, according to [\[KSS21,](#page-44-7) Proposition 8], a holomorphic semigroup in $L^2(\mathbb{R}^d)$, and prepare the following bounds for the holomorphic extension of $e^{-t\Lambda_{\kappa}}$ to the right complex half-plane. In the following, we write $\mathbb{C}_{\theta} := \{z \in \mathbb{C} : |\arg z| < \theta\}$ for $\theta \in [0, \pi/2]$.

Proposition 2.8. Let $\alpha \in (1, 2 \wedge (d+2)/2)$, $\beta \in ((d+\alpha)/2, d+\alpha)$, and $\kappa = \Psi(\beta)$ be defined by [\(1.7\)](#page-2-2). Then for any $\varepsilon \in (0,1)$, there exists a constant C such that

$$
|p_z(x,y)| \le C|z|^{-d/\alpha} \left(\frac{|z|^{1/\alpha}}{|z|^{1/\alpha} + |x-y|}\right)^{(d+\alpha)(1-\varepsilon)} \left(1 + \frac{|z|^{1/\alpha}}{|y|}\right)^{d-\beta} \tag{2.16}
$$

for all $x, y \in \mathbb{R}^d$ and $z \in \mathbb{C}_{\varepsilon \pi/4}$, where $p_z(x, y)$ is the kernel associated to the semigroup $e^{-z\Lambda_\kappa}$.

Proof. We use Davies' method [\[Dav90\]](#page-43-21), which relies on a Phragmén–Lindelöf argument. For $z \in \mathbb{C}$, set

$$
w_z(x) = \left(1 + \frac{t^{1/\alpha}}{|x|}\right)^{-(d-\beta)}, \quad x \in \mathbb{R}^d.
$$

We first claim that there exists $C > 0$ such that

$$
|p_z(x, y)w_z(y)| \le \frac{C}{|z|^{d/\alpha}}\tag{2.17}
$$

⁴The terminologies "supercritical" and "critical" are motivated by the fact that a gradient perturbation $b \cdot \nabla$ is of the same weight as $\sqrt{-\Delta}$, while if $\alpha < 1$, then, formally, $b \cdot \nabla$ dominates $(-\Delta)^{\alpha/2}$.

for all $x, y \in \mathbb{R}^d$ and $z \in \mathbb{C}_{\pi/4}$. To that end, let w be a nonnegative locally integrable function on \mathbb{R}^d . We define

$$
L_w^1(\mathbb{R}^d) = \left\{ f : |f|_{L_w^1(\mathbb{R}^d)} := \int_{\mathbb{R}^d} |f(x)| w(x) dx < \infty \right\}.
$$

Hence, the inequality [\(2.17\)](#page-15-2) is equivalent to

$$
\|e^{-z\Lambda_{\kappa}}\|_{L^1_{w_z^{-1}}(\mathbb{R}^d)\to L^\infty(\mathbb{R}^d)}\leq \frac{C}{|z|^{d/\alpha}}.
$$

For $z \in \mathbb{C}_{\pi/4}$ we can write $z = 2t + is$ for some $t \geq 0$ and $s \in \mathbb{R}$ such that $t \sim |z|$. Then,

$$
||e^{-z\Lambda_{\kappa}}||_{L_{w_z^{-1}}^1(\mathbb{R}^d)\to L^{\infty}(\mathbb{R}^d)}\leq ||e^{-t\Lambda_{\kappa}}||_{L^2(\mathbb{R}^d)\to L^{\infty}(\mathbb{R}^d)}||e^{-is\Lambda_{\kappa}}||_{L^2(\mathbb{R}^d)\to L^2(\mathbb{R}^d)}||e^{-t\Lambda_{\kappa}}||_{L_{w_z^{-1}}^1(\mathbb{R}^d)\to L^2(\mathbb{R}^d)}.
$$

This, together with Theorem [2.4](#page-13-0) and the fact $||e^{-is\Lambda_{\kappa}}||_{L^2(\mathbb{R}^d)\to L^2(\mathbb{R}^d)} \lesssim 1$ (since Λ_{κ} has a bounded functional calculus on $L^2(\mathbb{R}^d)$ for $\beta > (d+\alpha)/2$, see [\[KSS21,](#page-44-7) [KS23\]](#page-44-3)), implies

$$
\left\|e^{-z\Lambda_{\kappa}}\right\|_{L_{w_z^{-1}}^1(\mathbb{R}^d)\to L^{\infty}(\mathbb{R}^d)} \lesssim t^{-\frac{d}{2\alpha}}\left\|e^{-t\Lambda_{\kappa}}\right\|_{L_{w_z^{-1}}^1(\mathbb{R}^d)\to L^2(\mathbb{R}^d)}.\tag{2.18}
$$

It remains to estimate $||e^{-t\Lambda_{\kappa}}||_{L^1_{w_z^{-1}}(\mathbb{R}^d)\to L^2(\mathbb{R}^d)}$. For $f \in L^1_u$ $_{w_z^{-1}}^{1}(\mathbb{R}^d)$, we have, by (2.12) ,

$$
\|\mathrm{e}^{-t\Lambda_\kappa}f\|_{L^2(\mathbb{R}^d)}\lesssim \Big[\int_{\mathbb{R}^d}\Big|\int_{\mathbb{R}^d}t^{-d/\alpha}\Big(\frac{t^{1/\alpha}+|x-y|}{t^{1/\alpha}}\Big)^{-d-\alpha}w_z(y)^{-1}|f(y)|dy\Big|^2dx\Big]^{1/2}.
$$

By Minkowski's inequality,

$$
\begin{split} \| {\rm e}^{-t\Lambda_\kappa} f\|_{L^2(\mathbb{R}^d)} \lesssim & \int_{\mathbb{R}^d} \Big[\int_{\mathbb{R}^d} \Big| t^{-d/\alpha} \Big(\frac{t^{1/\alpha}+|x-y|}{t^{1/\alpha}}\Big)^{-d-\alpha} \Big|^2 dx \Big]^{1/2} w_z(y)^{-1} |f(y)| dy \\ \lesssim & \sup_{y\in \mathbb{R}^d} \Big[\int_{\mathbb{R}^d} \Big| t^{-d/\alpha} \Big(\frac{t^{1/\alpha}+|x-y|}{t^{1/\alpha}}\Big)^{-d-\alpha} \Big|^2 dx \Big]^{1/2} \|f\|_{L^1_{w_z^{-1}}} . \end{split}
$$

Using Lemma [2.1,](#page-9-2) we get

$$
\|e^{-t\Lambda_{\kappa}}f\|_{L^{2}(\mathbb{R}^{d})}\lesssim t^{-\frac{d}{2\alpha}}\|f\|_{L^{1}_{w_{x}^{-1}}},
$$

which yields

$$
\|e^{-t\Lambda_{\kappa}}\|_{L^1_{w_z^{-1}}(\mathbb{R}^d)\to L^2(\mathbb{R}^d)}\lesssim t^{-\frac{d}{2\alpha}}.
$$

Inserting this into [\(2.18\)](#page-16-0) gives

$$
\|e^{-z\Lambda_{\kappa}}\|_{L^1_{w_z^{-1}}(\mathbb{R}^d)\to L^\infty(\mathbb{R}^d)} \leq t^{-\frac{d}{\alpha}} \sim \frac{1}{|z|^{d/\alpha}},
$$

which implies (2.17) , i.e.,

$$
|p_z(x,y)| \lesssim \frac{1}{|z|^{d/\alpha}} \Big(1 + \frac{|z|^{1/\alpha}}{|y|}\Big)^{d-\beta}
$$

for all $x, y \in \mathbb{R}^d$ and $z \in \mathbb{C}_{\pi/4}$. By the Phragmén–Lindelöf argument in [\[DR96,](#page-43-22) Proposition 3.3] or [\[Mer22,](#page-44-18) Theorem 2.1], we get [\(2.16\)](#page-15-3).

This completes our proof.

For each $k \in \mathbb{N}$, we denote by $p_{t,k}(x, y)$ the kernel of $\Lambda_{\kappa}^{k}e^{-t\Lambda_{\kappa}}$. As a consequence of Proposition [2.8](#page-15-4) and Cauchy's formula, we obtain the following result.

Proposition 2.9. Let $\alpha \in (1, 2 \wedge (d+2)/2)$, $\beta \in ((d+\alpha)/2, d+\alpha)$, and $\kappa = \Psi(\beta)$ be defined by [\(1.7\)](#page-2-2). Then for any $\varepsilon > 0$ and $k \in \mathbb{N}$,

$$
|p_{t,k}(x,y)| \lesssim_{k,\varepsilon} t^{-(k+d/\alpha)} \left(\frac{t^{1/\alpha} + |x-y|}{t^{1/\alpha}}\right)^{-d-\alpha+\varepsilon} \left(1 + \frac{t^{1/\alpha}}{|y|}\right)^{d-\beta}
$$

for all $x, y \in \mathbb{R}^d$ and $t > 0$.

Proof. Applying Cauchy's formula, we get for every $t > 0$ and $k \in \mathbb{N}$,

$$
\Lambda_{\kappa}^{k} e^{-t\Lambda_{\kappa}} = \frac{(-1)^{k} k!}{2\pi i} \int_{|\xi - t| = \eta t} e^{-\xi} \frac{d\xi}{(\xi - t)^{k}},
$$

where $\eta > 0$ is small enough so that $\{\xi : |\xi - t| = \eta t\} \subset \mathbb{C}_{\tilde{\epsilon}\pi/4}$ with $\tilde{\epsilon} = \frac{\varepsilon}{d+1}$ $\frac{\varepsilon}{d+\alpha}$, and the integral does not depend on the choice of η . We now apply Proposition [2.8](#page-15-4) and the fact that $|\xi| \sim |\xi - t| \sim t$ to deduce

$$
|p_{t,k}(x,y)| \le C_{k,\tilde{\varepsilon}} t^{-(k+d/\alpha)} \left(\frac{t^{1/\alpha} + |x-y|}{t^{1/\alpha}}\right)^{-(d+\alpha)(1-\tilde{\varepsilon})} \left(1 + \frac{t^{1/\alpha}}{|y|}\right)^{d-\beta}
$$

$$
\le C_{k,\varepsilon} t^{-(k+d/\alpha)} \left(\frac{t^{1/\alpha} + |x-y|}{t^{1/\alpha}}\right)^{-(d+\alpha-\varepsilon)} \left(1 + \frac{t^{1/\alpha}}{|y|}\right)^{d-\beta}
$$

for all $x, y \in \mathbb{R}^d$ and $t > 0$.

We also need estimates for $|\nabla_x \partial_t e^{-t\Lambda_0}(x, y)|$. To that end, we first record the following bounds for spatial derivatives, essentially contained in [\[BJ07,](#page-43-19) Lemma 5]; see also [\[KSS21,](#page-44-7) (10), Lemma 1] for further estimates.

Lemma 2.10 ([\[BJ07,](#page-43-19) [KSS21\]](#page-44-7)). Let $d \in \mathbb{N}$ and $\alpha \in (0, 2)$. Then, for all $t > 0$ and $x, y \in \mathbb{R}^d$,

$$
|\nabla_x e^{-t\Lambda_0}(x, y)| \sim \frac{|x - y|}{t^{1/\alpha}} \cdot t^{-\frac{d+1}{\alpha}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha} + |x - y|}\right)^{d+2+\alpha}
$$

$$
\lesssim t^{-\frac{d+1}{\alpha}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha} + |x - y|}\right)^{d+1+\alpha}.
$$
 (2.19)

Lemma 2.11. Let $d \in \mathbb{N}$ and $\alpha \in (0, 2)$. Then, for any $\varepsilon > 0$,

$$
|\nabla_x \partial_t e^{-t\Lambda_0}(x, y)| \lesssim_{\varepsilon} t^{-\frac{d+\alpha+1}{\alpha}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha} + |x - y|}\right)^{d+\alpha+1-\varepsilon}.\tag{2.20}
$$

Proof. Note that $\nabla_x \partial_t e^{-t\Lambda_0} = \partial_t \nabla_x e^{-t\Lambda_0}$. Hence, the proof is similar to that of Proposition [2.9](#page-17-0) by using Lemma [2.10](#page-17-1) and hence we omit the details.

Lemma 2.12. Let $d \in \mathbb{N}$, $\alpha \in (0, 2)$, and $0 \leq \gamma \leq (d+1)/\alpha$. Then,

$$
|\nabla \Lambda_0^{-\gamma} e^{-t\Lambda_0}(x, y)| \lesssim_{\gamma} t^{-\frac{d+1-\gamma\alpha}{\alpha}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha} + |x - y|}\right)^{d+1-\alpha\gamma}.
$$
 (2.21)

Moreover, for $\alpha \in (1, 2 \wedge (d+2)/2)$, $0 \leq \gamma < d/\alpha$, $\beta \in ((d+\alpha)/2, d+\alpha)$, and $\kappa = \Psi(\beta)$ defined by (1.7) , we have

$$
|\Lambda_{\kappa}^{-\gamma} e^{-t\Lambda_{\kappa}}(x,y)| \lesssim_{\gamma} t^{-\frac{d-\gamma\alpha}{\alpha}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha}+|x-y|}\right)^{d-\alpha\gamma} \left(1 \wedge \frac{|y|}{t^{1/\alpha}+|x-y|}\right)^{\beta-d}.\tag{2.22}
$$

Note that the statements in this lemma are consistent their limits as $t \to 0$ in view of the Riesz kernel bounds for Λ_{κ} in Lemma [6.1](#page-34-1) below.

Proof. We first prove [\(2.21\)](#page-17-2). Using the formula

$$
\Lambda_0^{-\gamma} = c_\gamma \int_0^\infty s^\gamma e^{-s\Lambda_0} \frac{ds}{s},
$$

we have

$$
\nabla \Lambda_0^{-\gamma} e^{-t\Lambda_0}(x, y) = c_\gamma \int_0^\infty s^\gamma \nabla e^{-(s+t)\Lambda_0} \frac{ds}{s}
$$

.

.

.

This, together with Lemma [2.10,](#page-17-1) implies

$$
|\nabla \Lambda_0^{-\gamma} e^{-t\Lambda_0}(x, y)| \lesssim \int_0^\infty \frac{s^\gamma}{(s+t)^{\frac{d+1}{\alpha}}} \left(\frac{(s+t)^{1/\alpha}}{(s+t)^{1/\alpha}+|x-y|}\right)^{d+1+\alpha} \frac{ds}{s}
$$

$$
= \int_0^{t+|x-y|^\alpha} \dots + \int_{t+|x-y|^\alpha}^\infty \dots =: I_1 + I_2.
$$

We have

$$
I_1 \sim \int_0^{t+|x-y|^{\alpha}} s^{\gamma}(t+s) \left(\frac{1}{(s+t)^{1/\alpha}+|x-y|}\right)^{d+1+\alpha} \frac{ds}{s}
$$

$$
\lesssim (t^{1/\alpha}+|x-y|)^{\alpha(\gamma+1)} \left(\frac{1}{t^{1/\alpha}+|x-y|}\right)^{d+1+\alpha}
$$

$$
\sim \left(\frac{1}{t^{1/\alpha}+|x-y|}\right)^{d+1-\alpha\gamma} = \frac{1}{t^{\frac{d+1-\alpha\gamma}{\alpha}}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha}+|x-y|}\right)^{d+1-\alpha\gamma}
$$

Similarly, since $\alpha \gamma < d+1$, we have

$$
I_2 \sim \int_{t+|x-y|^{\alpha}}^{\infty} \frac{s^{\gamma}}{s^{\frac{d+1}{\alpha}}} \frac{ds}{s}
$$

$$
\sim \left(\frac{1}{t^{1/\alpha} + |x-y|}\right)^{d+1-\alpha\gamma} = \frac{1}{t^{\frac{d+1-\alpha\gamma}{\alpha}}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha} + |x-y|}\right)^{d+1-\alpha\gamma}
$$

This completes the proof of (2.21) . As the proof of (2.22) is similar, we omit the details. \Box

3. Proof of square function estimates (Theorem [1.3\)](#page-5-0)

To prove Theorem [1.3,](#page-5-0) we recall two criteria for singular integrals to be bounded on Lebesgue spaces, which will play an important role in the proof of the boundedness of the square functions. The first theorem gives a criterion on the boundedness on $L^p(\mathbb{R}^d)$ spaces with $p \in (1, 2)$, while the second one covers the range $p > 2$.

Theorem 3.1. Let $1 \leq p_0 < 2$ and let T be a sublinear operator which is bounded on $L^2(\mathbb{R}^d)$. Assume that there exists a family of operators $\{\mathcal{A}_t\}_{t>0}$ satisfying that for every $j \geq 2$ and every ball B

$$
\left(\int_{S_j(B)} |T(I - A_{r_B})f|^2\right)^{1/2} \le \alpha(j) \left(\int_B |f|^{p_0}\right)^{1/p_0} \tag{3.1}
$$

and

$$
\left(\int_{S_j(B)} |\mathcal{A}_{r_B} f|^2\right)^{1/2} \le \alpha(j) \left(\int_B |f|^{p_0}\right)^{1/p_0} \tag{3.2}
$$

for all f supported in B. If $\sum_j \alpha(j)2^{jd} < \infty$, then T is bounded on $L^p(\mathbb{R}^d)$ for all $p \in (p_0, 2)$.

Theorem 3.2. Let $2 < q_0 \leq \infty$. Let T be a bounded sublinear operator on $L^2(\mathbb{R}^d)$. Assume that there exists a family of operators $\{\mathcal{A}_t\}_{t>0}$ satisfying

$$
\left(\int_{B} \left|T(I - \mathcal{A}_{r_B})f\right|^2 dx\right)^{1/2} \leq C \mathcal{M}_2(f)(x) \tag{3.3}
$$

and

$$
\left(\int_{B} \left| T\mathcal{A}_{r_{B}}f \right|^{q_{0}} dx\right)^{1/q_{0}} \leq C\mathcal{M}_{2}(Tf)(x)
$$
\n(3.4)

for all balls B with radius r_B , all $f \in C_c^{\infty}(\mathbb{R}^d)$ and all $x \in B$. Then T is bounded on $L^p(\mathbb{R}^d)$ for all $2 < p < q_0$.

For the proof of Theorems [3.1](#page-18-1)[–3.2,](#page-18-2) see [\[Aus07\]](#page-42-7).

Proof of Theorem [1.3.](#page-5-0) As detailed below, the proof of the theorem only relies on the kernel estimates in Proposition [2.9.](#page-17-0) It suffices to prove the theorem for $\beta \in ((d+\alpha)/2, d]$ since the proof for the case $\beta \in (d, d + M)$ is similar to the case $\beta = d$. Since the proof of [\(1.19\)](#page-5-5) is similar to that of (1.20) (in fact, even easier), we only prove (1.20) .

Fix $\delta \in (0, \alpha)$ such that $\delta/\alpha > \gamma$. According to [\[KSS21,](#page-44-7) Proposition 8], Λ_{κ} generates a holomorphic semigroup in L^2 . Therefore, Λ_{κ} and Λ_{κ}^* have a bounded functional calculus on $L^2(\mathbb{R}^d)$ and $S_{\Lambda^*_{\kappa},\gamma}$ is bounded on $L^2(\mathbb{R}^d)$ (see [\[McI86\]](#page-44-9)). We now prove the boundedness of $S_{\Lambda^*_{\kappa},\gamma}$ and distinguish between whether $p < 2$ or $p > 2$.

Step 1: Proof of the L^p -boundedness for $1 < p < 2$

Fix $1 < p \leq 2$. Due to Theorem [3.1,](#page-18-1) it suffices to prove

$$
\left(\int_{S_j(B)} |S_{\Lambda^*_\kappa,\gamma}(I - A_{r_B})f(x)|^2 dx\right)^{1/2} \lesssim 2^{-(d+\delta)j} \left(\int_B |f(x)|^p dx\right)^{1/p} \tag{3.5}
$$

and

$$
\left(\int_{S_j(B)} |\mathcal{A}_{r_B} f(x)|^2 dx\right)^{1/2} \lesssim 2^{-(d+\delta)j} \left(\int_B |f(x)|^p dx\right)^{1/p} \tag{3.6}
$$

for all $j \geq 2$ and for every function f supported in B, where

$$
\mathcal{A}_{r_B} = I - (I - e^{-r_B^{\alpha} \Lambda_{\kappa}^*})^m, \quad m > \frac{d}{\alpha p'} + \frac{\delta}{\alpha} + 1.
$$

Since

$$
\mathcal{A}_{r_B} = \sum_{k=1}^m C_k^m e^{-kr_B^{\alpha} \Lambda_{\kappa}^*},
$$

the estimate [\(3.6\)](#page-19-0) follows directly from [\(2.12\)](#page-14-1) and Theorem [2.3.](#page-11-2)

It remains to prove [\(3.5\)](#page-19-1). To that end, we write

$$
\left(\int_{S_j(B)} |S_{\Lambda^*_{\kappa},\gamma}(I - e^{-r_B^{\alpha}\Lambda^*_{\kappa}})^m f|^2 dx\right)^{1/2} \n\leq \left(\int_0^{r_B^{\alpha}} \|(t\Lambda^*_{\kappa})^{\gamma}e^{-t\Lambda^*_{\kappa}}(I - e^{-r_B^{\alpha}\Lambda^*_{\kappa}})^m f\|_{L^2(S_j(B),\frac{dx}{|S_j(B)|})}^2 \frac{dt}{t}\right)^{1/2} \n+ \left(\int_{r_B^{\alpha}}^{\infty} \|(t\Lambda^*_{\kappa})^{\gamma}e^{-t\Lambda^*_{\kappa}}(I - e^{-r_B^{\alpha}\Lambda^*_{\kappa}})^m f\|_{L^2(S_j(B),\frac{dx}{|S_j(B)|})}^2 \frac{dt}{t}\right)^{1/2} \n:= E_1 + E_2.
$$
\n(3.7)

We first take care of I_1 . Note that

$$
(\Lambda_{\kappa}^*)^{\gamma} = \frac{1}{\Gamma(1-\gamma)} \int_0^{\infty} u^{1-\gamma} \Lambda_{\kappa}^* e^{-u\Lambda_{\kappa}^*} \frac{du}{u}.
$$
 (3.8)

Using this and Minkowski's inequality, we have

$$
E_1 \lesssim \left(\int_0^{r_B^{\alpha}} \left[\int_0^{r_B^{\alpha}} \left(\frac{u}{t}\right)^{1-\gamma} \left\|t\Lambda_{\kappa}^* e^{-(t+u)\Lambda_{\kappa}} (I - e^{-r_B^{\alpha}\Lambda_{\kappa}^*})^m f\right\|_{L^2(S_j(B), \frac{dx}{|S_j(B)|})} \frac{du}{u}\right]^2 \frac{dt}{t}\right)^{1/2} + \left(\int_0^{r_B^{\alpha}} \left[\int_{r_B^{\alpha}}^{\infty} \left(\frac{u}{t}\right)^{1-\gamma} \left\|t\Lambda_{\kappa}^* e^{-(t+u)\Lambda_{\kappa}^*} (I - e^{-r_B^{\alpha}\Lambda_{\kappa}^*})^m f\right\|_{L^2(S_j(B), \frac{dx}{|S_j(B)|})} \frac{du}{u}\right]^2 \frac{dt}{t}\right)^{1/2} := E_{11} + E_{12}.
$$

Using the identity

$$
(I - e^{-r_B^{\alpha} \Lambda_{\kappa}^*})^m = \sum_{k=0}^m (-1)^k C_k^m e^{-kr_B^{\alpha} \Lambda_{\kappa}^*}
$$

for certain numerical values C_k^m , we have

$$
E_{11} \lesssim \Big(\int_0^{r_B^{\alpha}} \Big[\int_0^{r_B^{\alpha}} \Big(\frac{u}{t}\Big)^{1-\gamma} \frac{t}{t+u} \Big\|(t+u)\Lambda_{\kappa}^* e^{-(t+u)\Lambda_{\kappa}^*} f\Big\|_{L^2(S_j(B),\frac{dx}{|S_j(B)|})} \frac{du}{u}\Big]^2 \frac{dt}{t}\Big)^{1/2} + \sum_{k=1}^m \Big(\int_0^{r_B^{\alpha}} \Big[\int_0^{r_B^{\alpha}} \Big(\frac{u}{t}\Big)^{1-\gamma} \frac{t}{t+u+kr_B^{\alpha}}\Big\|(t+u+kr_B^{\alpha})\Lambda_{\kappa}^* e^{-(t+u+kr_B^{\alpha})\Lambda_{\kappa}^*} f\Big\|_{L^2(S_j(B),\frac{dx}{|S_j(B)|})} \frac{du}{u}\Big]^2 \frac{dt}{t}\Big)^{1/2}.
$$

This, in combination with Proposition [2.9,](#page-17-0) Theorem [2.3,](#page-11-2) and the facts that $t + u \lesssim r_B^{\alpha}$ and $(t+u+kr_B^{\alpha})^{1/\alpha} \sim r_B$ for $u, t \in (0, r_B^{\alpha}]$ and $k \ge 1$, yields

$$
E_{11} \lesssim \Big(\int_0^{r_B^{\alpha}} \Big[\int_0^{r_B^{\alpha}} \Big(\frac{u}{t} \Big)^{1-\gamma} \frac{t}{t+u} \Big(\frac{r_B}{(t+u)^{1/\alpha}} \Big)^d \Big(\frac{2^j r_B}{(t+u)^{1/\alpha}} \Big)^{-d-\delta} \|f\|_{L^p(B, \frac{dx}{|B|})} \frac{du}{u} \Big]^2 \frac{dt}{t} \Big)^{\frac{1}{2}}
$$

+
$$
\sum_{k=1}^m \Big(\int_0^{r_B^{\alpha}} \Big[\int_0^{r_B^{\alpha}} \Big(\frac{u}{t} \Big)^{1-\gamma} \frac{t}{r_B^{\alpha}} 2^{-j(d+\delta)} \|f\|_{L^p(B, \frac{dx}{|B|})} \frac{du}{u} \Big]^2 \frac{dt}{t} \Big)^{1/2}.
$$

Hence,

$$
E_{11} \lesssim 2^{-j(d+\delta)} \Big(\int_B |f|^p dx\Big)^{1/p}.
$$

To estimate E_{12} , we use

$$
(I - e^{-r_B^{\alpha} \Lambda_{\kappa}^*})^m = \int_0^{r_B^{\alpha}} \dots \int_0^{r_B^{\alpha}} (\Lambda_{\kappa}^*)^m e^{-(s_1 + \dots + s_m)\Lambda_{\kappa}^*} d\vec{s}, \tag{3.9}
$$

where $d\vec{s} := ds_1 \dots ds_m$, to write

$$
E_{12} \lesssim \Big(\int_0^{r_B^{\alpha}} \Big[\int_{[0,r_B^{\alpha}]^m} \int_{r_B^{\alpha}}^{\infty} \Big(\frac{u}{t}\Big)^{1-\gamma} \Big\| t(\Lambda_{\kappa}^*)^{m+1} e^{-(t+u+s_1+\ldots+s_m)\Lambda_{\kappa}^*} f \Big\|_{L^2(S_j(B),\frac{dx}{|S_j(B)|})} \frac{du}{u} d\vec{s}\Big]^2 \frac{dt}{t}\Big)^{1/2}.
$$

In this case, $u \sim t + u + s_1 + \ldots + s_m \geq r_B^{\alpha}$. Hence, by Proposition [2.9](#page-17-0) and Theorem [2.3,](#page-11-2) we obtain

$$
E_{12} \lesssim \left[\int_0^{r_B^{\alpha}} \left(\int_{[0,r_B^{\alpha}]^m} \int_{r_B^{\alpha}}^{\infty} \left(\frac{u}{t} \right)^{1-\gamma} \frac{t}{u^{m+1}} \left(\frac{r_B}{u^{1/\alpha}} \right)^d \left(1 + \frac{u^{1/\alpha}}{2^j r_B} \right)^{d/2} \right] \tag{3.10}
$$

$$
\left(1 + \frac{2^j r_B}{u^{1/\alpha}} \right)^{-(d+\delta)} \|f\|_{L^p(B, \frac{dx}{|B|})} \frac{du}{u} d\vec{s} \right)^2 \frac{dt}{t} \Big]^{1/2}.
$$

We see that

$$
\int_{r_B^{\alpha}}^{\infty} \left(\frac{u}{t}\right)^{1-\gamma} \frac{t}{u^{m+1}} \left(\frac{r_B}{u^{1/\alpha}}\right)^d \left(1 + \frac{u^{1/\alpha}}{2^j r_B}\right)^{d/2} \left(1 + \frac{2^j r_B}{u^{1/\alpha}}\right)^{-(d+\delta)} \frac{du}{u}
$$
\n
$$
= \int_{r_B^{\alpha}}^{(2^j r_B)^{\alpha}} \dots + \int_{(2^j r_B)^{\alpha}}^{\infty} \dots
$$
\n
$$
\sim \int_{r_B^{\alpha}}^{(2^j r_B)^{\alpha}} \left(\frac{u}{t}\right)^{1-\gamma} \frac{t}{u^{m+1}} \left(\frac{r_B}{u^{1/\alpha}}\right)^d \left(\frac{2^j r_B}{u^{1/\alpha}}\right)^{-(d+\delta)} \frac{du}{u}
$$
\n
$$
+ \int_{(2^j r_B)^{\alpha}}^{\infty} \left(\frac{u}{t}\right)^{1-\gamma} \frac{t}{u^{m+1}} \left(\frac{r_B}{u^{1/\alpha}}\right)^d \left(\frac{u^{1/\alpha}}{2^j r_B}\right)^{d/2} \frac{du}{u}
$$
\n
$$
\lesssim 2^{-j(d+\delta)} \left(\frac{r_B^{\alpha}}{t}\right)^{1-\gamma} \frac{t}{r_B^{\alpha(m+1)}} + 2^{-jd} \left(\frac{(2^j r_B)^{\alpha}}{t}\right)^{1-\gamma} \frac{t}{(2^j r_B)^{\alpha(m+1)}}
$$
\n
$$
\lesssim 2^{-j(d+\delta)} \left(\frac{r_B^{\alpha}}{t}\right)^{1-\gamma} \frac{t}{r_B^{\alpha(m+1)}},
$$

for all $m \geq 2$. Inserting this into [\(3.10\)](#page-20-0) yields

$$
E_{12} \lesssim 2^{-j(d+\delta)} \|f\|_{L^p(B, \frac{dx}{|B|})} \Big(\int_0^{r_B^{\alpha}} \Big[\int_{[0,r_B^{\alpha}]^m} \Big(\frac{r_B^{\alpha}}{t} \Big)^{1-\gamma} \frac{t}{r_B^{\alpha(m+1)}} d\vec{s} \Big]^2 \frac{dt}{t} \Big)^{1/2}
$$

$$
\lesssim 2^{-j(d+\delta)} \Big(\int_B |f|^p dx \Big)^{1/p}.
$$

Collecting the estimates for E_{11} and E_{12} gives

$$
E_1 \lesssim 2^{-j(d+\delta)} \Big(\int_B |f|^p dx\Big)^{1/p}.
$$

We now take care of E_2 . To that end, we write

$$
E_2 \lesssim \Big(\int_{r_B^{\alpha}}^{\infty} \Big[\int_0^{r_B^{\alpha}} \Big(\frac{u}{t} \Big)^{1-\gamma} \Big\| t \Lambda_{\kappa}^* e^{-(t+u)\Lambda_{\kappa}^*} (I - e^{-r_B^{\alpha}\Lambda_{\kappa}^*})^m f \Big\|_{L^2(S_j(B), \frac{d}{|S_j(B)|})} \frac{du}{u} \Big]^2 \frac{dt}{t} \Big)^{1/2} + \Big(\int_{r_B^{\alpha}}^{\infty} \Big[\int_{r_B^{\alpha}}^{\infty} \Big(\frac{u}{t} \Big)^{1-\gamma} \Big\| t \Lambda_{\kappa}^* e^{-(t+u)\Lambda_{\kappa}^*} (I - e^{-r_B^{\alpha}\Lambda_{\kappa}^*})^m f \Big\|_{L^2(S_j(B), \frac{dx}{|S_j(B)|})} \frac{du}{u} \Big]^2 \frac{dt}{t} \Big)^{1/2} =: E_{21} + E_{22}.
$$

Similarly to [\(3.10\)](#page-20-0),

$$
E_{12} \lesssim \Big(\int_{r_B^{\alpha}}^{\infty} \Big[\int_{[0,r_B^{\alpha}]^m} \int_0^{r_B^{\alpha}} \Big(\frac{u}{t} \Big)^{1-\gamma} \frac{t}{t^{m+1}} \Big(\frac{r_B}{t^{1/\alpha}} \Big)^d \Big(1 + \frac{t^{1/\alpha}}{2^j r_B} \Big)^{d/2} \Big(1 + \frac{2^j r_B}{t^{1/\alpha}} \Big)^{-(d+\delta)} \| f \|_{L^p(B,\frac{dx}{|B|})} \frac{du}{u} d\vec{s} \Big]^2 \frac{dt}{t} \Big)^{\frac{1}{2}}
$$

$$
\lesssim \Big(\int_{r_B^{\alpha}}^{\infty} \Big[\Big(\frac{r_B^{\alpha}}{t} \Big)^{1-\gamma} \frac{r_B^{\alpha m}}{t^m} \Big(\frac{r_B}{t^{1/\alpha}} \Big)^d \Big(1 + \frac{t^{1/\alpha}}{2^j r_B} \Big)^{d/2} \Big(1 + \frac{2^j r_B}{t^{1/\alpha}} \Big)^{-(d+\delta)} \| f \|_{L^p(B,\frac{dx}{|B|})} \Big]^2 \frac{dt}{t} \Big)^{1/2}
$$

$$
\lesssim 2^{-j(d+\delta)} \Big(\int_B |f|^p dx \Big)^{1/p},
$$

for all $m \geq 2$. Similarly as before,

$$
\begin{split} &E_{22} \lesssim \Big(\int_{r_B^\alpha}^\infty \Big[\int_{[0,r_B^\alpha]^m} \int_{r_B^\alpha}^\infty \Big(\frac{u}{t}\Big)^{1-\gamma} \frac{t}{(t+u)^{m+1}} \Big(\frac{r_B}{(t+u)^{1/\alpha}} \Big)^d \Big(1+ \frac{(t+u)^{1/\alpha}}{2^j r_B} \Big)^{d/2} \\ &\times \Big(1+ \frac{2^j r_B}{(t+u)^{1/\alpha}} \Big)^{-(d+\delta)} \| f \|_{L^p(B,\frac{dx}{|B|})} \frac{du}{u} d\vec{s} \Big]^2 \frac{dt}{t} \Big)^{1/2} \\ \lesssim \Big(\int_{r_B^\alpha}^\infty \Big[\int_{[0,r_B^\alpha]^m} \int_{r_B^\alpha}^\infty \Big(\frac{u}{t}\Big)^{1-\gamma} \frac{1}{t^{m-1} u} \Big(\frac{r_B}{t^{1/\alpha}} \Big)^d \Big(1+ \frac{t^{1/\alpha}}{2^j r_B} \Big)^{d/2} \Big(1+ \frac{2^j r_B}{t^{1/\alpha}} \Big)^{-(d+\delta)} \| f \|_{L^p(B,\frac{dx}{|B|})} \frac{du}{u} d\vec{s} \Big]^2 \frac{dt}{t} \Big)^{\frac{1}{2}} \\ \lesssim \Big(\int_{r_B^\alpha}^\infty \Big[\Big(\frac{r_B^\alpha}{t} \Big)^{1-\gamma} \frac{r_B^{\alpha m}}{t^{m-1} r_B^\alpha} \Big(\frac{r_B}{t^{1/\alpha}} \Big)^d \Big(1+ \frac{t^{1/\alpha}}{2^j r_B} \Big)^{d/2} \Big(\frac{2^j r_B}{t^{1/\alpha}} \Big)^{-(d+\delta)} \| f \|_{L^p(B,\frac{dx}{|B|})} \Big]^2 \frac{dt}{t} \Big)^{1/2} \\ \lesssim 2^{-j(d+\delta)} \Big(\int_B |f|^p dx \Big)^{1/p} , \end{split}
$$

for all $m \geq 2$. Taking all the estimates of E_{21}, E_{22} and E_1 into account, we conclude that

$$
\Big(\int_{S_j(B)}|S_{\Lambda^*_\kappa,\gamma}(I-\mathrm{e}^{-r_B^\alpha\Lambda^*_\kappa})^m f|^2dx\Big)^{1/2}\lesssim 2^{-j(d+\delta)}|2^jB|^{1/2}\Big(\int_B|f|^2dx\Big)^{1/2}
$$

This completes the proof of [\(3.5\)](#page-19-1).

$\textbf{Step 2: Proof of the } L^p\textbf{-boundedness for } 2 < p < \frac{d}{d-\beta}$

By Theorem [3.2,](#page-18-2) for any $q \in (2, \frac{d}{d-1})$ $\frac{d}{d-\beta}$) it suffices to prove that

$$
\left(\int_{B} \left| S_{\Lambda_{\kappa}^*,\gamma}(I - \mathcal{A}_{r_B})f \right|^2 dx \right)^{1/2} \leq C \mathcal{M}_2(f)(x),\tag{3.11}
$$

and

$$
\left(\int_{B} \left|S_{\Lambda_{\kappa}^*,\gamma} \mathcal{A}_{r_B} f\right|^q dx\right)^{1/q} \leq C \mathcal{M}_2(|S_{\Lambda_{\kappa}^*,\gamma} f|)(x) \tag{3.12}
$$

for all balls B with radius r_B , all $f \in C_c^{\infty}(\mathbb{R}^d)$ and all $x \in B$ with $\mathcal{A}_{r_B} = I - (I - e^{-r_B^{\alpha} \Lambda_{\kappa}^*})^m$, and $m > 2$.

To prove [\(3.11\)](#page-22-0), we write

$$
\left(\int_B |S_{\Lambda^*_\kappa,\gamma}(I-\mathrm{e}^{-r_B^\alpha\Lambda^*_\kappa})^m f|^2 dx\right)^{1/2} \leq \sum_{j\geq 0} \left(\int_B |S_{\Lambda^*_\kappa,\gamma}(I-\mathrm{e}^{-r_B^\alpha\Lambda^*_\kappa})^m f_j|^2 dx\right)^{1/2} =: \sum_{j=0}^\infty F_j,
$$

where $f_j = f \chi_{S_j(B)}$.

For $j = 0, 1$, using the L²-boundedness of $S_{\Lambda_{\kappa}, \gamma}$ and \mathcal{A}_{r_B} , we have

$$
F_j \lesssim \mathcal{M}_2(f)(x).
$$

Hence, it suffices to prove that

$$
F_j \lesssim 2^{-j\beta} \Big(\int_{S_j(B)} |f|^2 dx \Big)^{1/2} \tag{3.13}
$$

for all $j \geq 2$. To do this, for $j \geq 2$, we write

$$
\left(\int_{B} |S_{\Lambda_{\kappa}^*,\gamma}(I - e^{-r_B^{\alpha}\Lambda_{\kappa}^*})^m f_j|^2 dx\right)^{1/2} \le \left(\int_0^{r_B^{\alpha}} \left\|(t\Lambda_{\kappa}^*)^{\gamma}e^{-t\Lambda_{\kappa}^*}(I - e^{-r_B^{\alpha}\Lambda_{\kappa}^*})^m f_j\right\|_{L^2(B,\frac{dx}{|B|})}^2 \frac{dt}{t}\right)^{1/2} + \left(\int_{r_B^{\alpha}}^{\infty} \left\|(t\Lambda_{\kappa}^*)^{\gamma}e^{-t\Lambda_{\kappa}^*}(I - e^{-r_B^{\alpha}\Lambda_{\kappa}^*})^m f_j\right\|_{L^2(B,\frac{dx}{|B|})}^2 \frac{dt}{t}\right)^{1/2}.
$$

.

At this stage, we can argue as in the proof of [\(3.5\)](#page-19-1) in Step 1. However, in this case, we will utilize (2.8) instead of (2.7) . By doing so, we arrive at the expression (3.13) . As the proof follows a similar structure, we omit the details.

It remains to prove [\(3.12\)](#page-22-2). We first write

$$
\left(\int_{B} |S_{\Lambda_{\kappa}^*,\gamma} \mathcal{A}_{r_B} f(x)|^q dx\right)^{\frac{1}{q}} = \left[\int_{B} \left(\int_0^{\infty} \Big| \sum_{k=1}^m C_k^m e^{-kr_B^{\alpha} \Lambda_{\kappa}^*} (t \Lambda_{\kappa}^*)^{\gamma} e^{-t \Lambda_{\kappa}^*} f(x) \Big|^{2} \frac{dt}{t}\right)^{q/2} dx\right]^{\frac{1}{q}}
$$

$$
\lesssim \sum_{j\geq 0} \left[\int_{B} \left(\int_0^{\infty} \Big| \sum_{k=1}^m C_k^m e^{-kr_B^{\alpha} \Lambda_{\kappa}^*} [(t \Lambda_{\kappa}^*)^{\gamma} e^{-t \Lambda_{\kappa}^*} f \chi_{S_j(B)}](x) \Big|^{2} \frac{dt}{t}\right)^{q/2} dx\right]^{1/q}
$$

which, along with Minkowski's inequality, Proposition [2.9,](#page-17-0) and (2.8) in Theorem [2.3,](#page-11-2) gives

$$
\left(\int_{B} |S_{\Lambda_{\kappa}^*,\gamma} A_{r_B} f(x)|^q dx\right)^{1/q} \n\lesssim \sum_{j\geq 0} \left(\int_0^{\infty} \Big\|\sum_{k=1}^m e^{-kr_B^{\alpha}\Lambda_{\kappa}^*} [(t\Lambda_{\kappa}^*)^{\gamma}e^{-t\Lambda_{\kappa}^*} f\chi_{S_j(B)}]\Big\|_{L^q(B,\frac{dx}{|B|})}^2 \frac{dt}{t}\right)^{1/2} \n\lesssim \sum_{j\geq 0} 2^{-j\beta} \left(\int_0^{\infty} \|(t\Lambda_{\kappa}^*)^{\gamma}e^{-t\Lambda_{\kappa}^*} f\|_{L^2(S_j(B),\frac{dx}{|S_j(B)|})}^2 \frac{dt}{t}\right)^{1/2} \n\lesssim \sum_{j\geq 0} 2^{-j\beta} \left(\int_{2^jB} |S_{\Lambda_{\kappa}^*,\gamma} f(x)|^2 dx\right)^{1/2}.
$$

This implies [\(3.12\)](#page-22-2). Hence the proof of Step 2 is completed.

Thus, we proved that the square function $S_{\Lambda^*_{\kappa},\gamma}$ is bounded on $L^p(\mathbb{R}^d)$ for all $1 < p < \infty$, i.e.,

$$
||S_{\Lambda_{\kappa}^*,\gamma}f||_{L^p(\mathbb{R}^d)} \lesssim ||f||_{L^p(\mathbb{R}^d)}.
$$

As we remarked at the beginning of this proof, the L^p -boundedness of $S_{\Lambda_{\kappa},\gamma}$ is proved analogously. We new prove the reversed square function inequalities using duality. For the sake of brevity, we only prove $||f||_p \lesssim ||S_{\Lambda_{\kappa}},\gamma f||_p$; the proof for Λ_{κ}^* is again similar.

By functional calculus, for any $g \in L^{p'}(\mathbb{R}^d)$, we have

$$
\int_{\mathbb{R}^d} f(x)g(x)dx = c(\gamma) \int_{\mathbb{R}^d} \int_0^\infty (t\Lambda_\kappa)^{2\gamma} e^{-2t\Lambda_\kappa} f(x)g(x) \frac{dt}{t} dx,
$$

where $c(\gamma) = \int_0^\infty t^{2\gamma} e^{-2t} \frac{dt}{t}$ $\frac{dt}{t}$. Using Hölder's inequality, we obtain

$$
\int_{\mathbb{R}^d} f(x)g(x)dx = c(\gamma) \int_{\mathbb{R}^d} \int_0^\infty (t\Lambda_\kappa)^\gamma e^{-t\Lambda_\kappa} f(x) (t\Lambda_\kappa^*)^\gamma e^{-t\Lambda_\kappa^*} g(x) \frac{dt}{t} dx
$$

$$
\lesssim \int_{\mathbb{R}^d} S_{\Lambda_\kappa,\gamma} f(x) S_{\Lambda_\kappa^*,\gamma} g(x) dx
$$

$$
\lesssim ||S_{\Lambda_\kappa,\gamma} f||_{L^p(\mathbb{R}^d)} ||S_{\Lambda_\kappa^*,\gamma} g||_{L^{p'}(\mathbb{R}^d)}.
$$

By (1.20) , i.e., $||S_{\Lambda^*_{\kappa},\gamma}g||_{L^{p'}(\mathbb{R}^d)} \lesssim ||g||_{L^{p'}(\mathbb{R}^d)}$, we get

$$
\int_{\mathbb{R}^d} f(x)g(x)dx \lesssim ||S_{\Lambda_{\kappa},\gamma}f||_{L^p(\mathbb{R}^d)}||g||_{L^{p'}(\mathbb{R}^d)}.
$$

As a consequence,

 $||f||_{L^p(\mathbb{R}^d)} \lesssim ||S_{\Lambda_{\kappa},\gamma}f||_{L^p(\mathbb{R}^d)},$

which completes the proof. \Box

4. Bounds for differences of kernels

The goal of this section is to prove pointwise estimates for the difference

$$
Q_t(x,y) := t\Lambda_0 e^{-t\Lambda_0}(x,y) - t\Lambda_\kappa e^{-t\Lambda_\kappa}(x,y)
$$
\n(4.1)

and $|(Q_t f)(x)|$ for any $f \in L^p(\mathbb{R}^d)$, $p > 1 \vee d/\beta$. These will be instrumental to prove the reversed Hardy inequality (Theorem [1.4\)](#page-5-1) via Schur tests in the ensuing section. For the sake of concreteness, we always assume $\alpha \in (1, 2)$, and $\beta \in ((d + \alpha)/2, d + \alpha)$ from now on.

To that end, we use Duhamel's formula and integral bounds involving, for $t > 0$, $x, y \in \mathbb{R}^d$, and $\gamma, \delta > 0$ the functions

$$
T_t^{\gamma}(x,y) := \left(1 \wedge \frac{|y|}{t^{1/\alpha}}\right)^{\beta - d} t^{-d/\alpha} \left(\frac{t^{1/\alpha}}{t^{1/\alpha} + |x - y|}\right)^{d + \gamma}
$$
(4.2)

and

$$
H_t^{\gamma,\delta}(x,y) := t^{-\frac{d+\delta}{\alpha}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha} + |x-y|}\right)^{d+\gamma}.
$$
 (4.3)

For $\delta = 1$ and appropriate values of γ , these functions arise from estimating $t | \partial_t e^{-t\Lambda_{\kappa}} |$ and $t|\nabla_x \partial_t e^{-t\Lambda_0}(x, y)|$, respectively; see Propositions [2.9](#page-17-0) and [2.11.](#page-17-4) We will formulate our bounds on $|Q_t(x, y)|$ and $|(Q_t f)(x)|$ in terms of the functions

$$
L_t^{\gamma,\delta}(x,y) := \mathbf{1}_{\{|y| \le t^{1/\alpha}\}} \left(\frac{|y|}{t^{1/\alpha}}\right)^{\beta - d - (\alpha - \gamma)} t^{-\frac{d}{\alpha}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha} + |x - y|}\right)^{d+\gamma} \cdot t^{(1-\delta)/\alpha}
$$

+
$$
\mathbf{1}_{\{|y| \le t^{1/\alpha}, |x| \sim |y|\}} \left(\frac{|y|}{t^{1/\alpha}}\right)^{1+\beta - d - \alpha} t^{-\frac{d}{\alpha}} \cdot t^{(1-\delta)/\alpha}
$$

+
$$
\mathbf{1}_{\{|y| \ge t^{1/\alpha}\}} \mathbf{1}_{|x-y| \ge (|x|\wedge |y|)/2} t^{-\frac{d}{\alpha}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha} + |x - y|}\right)^{d+\gamma} |y|^{1-\delta}
$$
(4.4)

and

$$
M_t^{\gamma,\delta}(x,y) := \mathbf{1}_{|y| \ge t^{1/\alpha}} \mathbf{1}_{|x-y| \le (|x|\wedge|y|)/2} \cdot \left(\frac{|x| \vee |y|}{t^{1/\alpha}}\right)^{\delta-\alpha} \frac{1}{t^{d/\alpha}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha} + |x-y|}\right)^{d+\gamma}.\tag{4.5}
$$

These two functions are similar to those in [\[FMS21,](#page-43-3) Lemma 3.1]. One major difference between the function $M_t^{\gamma,\delta}(x, y)$ here and its analog in [\[FMS21,](#page-43-3) p. 2295] is that the power $\delta - \alpha$ of the factor $|y|/t^{1/\alpha}$ is only $-\alpha$ in [\[FMS21\]](#page-43-3). In parts of our proofs below we will have $\delta = 0$, while in other parts $\delta > 0$ is necessary, which is essentially due to the gradient perturbation.

In the following subsection, we estimate integrals involving the functions T_t^{γ} $t^{\gamma}(x, y)$ and $H_t^{\gamma, \delta}$ t in terms of the functions L_t^{γ} $t^{\gamma}(x, y)$ and $M_t^{\gamma, \delta}(x, y)$. In the ensuing two subsections, we prove pointwise bounds for $|Q_t(x, y)|$ and, afterwards, for $|(Q_t f)(x)|$. The former bounds will be used to prove part (1), while the latter will be used to prove part (2) in Theorem [1.4.](#page-5-1)

4.1. Integral bounds. In this section, we estimate integrals involving the functions T_t^{γ} $\mathfrak{f}^\gamma_t(x,y)$ and $H_t^{\gamma,\delta}$ $\hat{t}^{\gamma,0}$. The estimates for $\delta = 1$ will be crucial in Subsection [4.2,](#page-28-0) while those for $\delta = \gamma$ will be important in Subsection [4.3.](#page-30-0) Our techniques are similar to those in [\[FM23,](#page-43-4) [BM23\]](#page-43-5).

The following lemma, whose proof we defer to Appendix [A.2,](#page-36-2) will be useful for our endeavors.

Lemma 4.1. Let $d \in \mathbb{N}$. Then for all $N > 0$, $0 < s < t$, and $x, y \in \mathbb{R}^d$, we have

$$
\int_{\mathbb{R}^d} dz \,(t-s)^{-d} \left(\frac{(t-s)}{(t-s)+|x-z|}\right)^{d+N} s^{-d} \left(\frac{s}{s+|z-y|}\right)^{d+N} \sim \frac{1}{t^d} \left(\frac{t}{t+|x-y|}\right)^{d+N}.\tag{4.6}
$$

In the following lemma, we consider integrals arising when studying Q_t in the region $|x-y|$ $(|x| \wedge |y|)/2$, where we expect cancellation effects.

Lemma 4.2. Let $\alpha \in (1,2], \beta \in ((d+\alpha)/2, d+\alpha)$, and $\gamma, \delta \in (0,\alpha), t > 0$, and $x, y \in \mathbb{R}^d$. Then,

$$
\mathbf{1}_{|y| \ge t^{1/\alpha}} \mathbf{1}_{|x-y| \le (|x|\wedge|y|)/2} \int_{\mathbb{R}^d} \int_0^t T_{t-s}^{\gamma}(x,z) |z|^{-\alpha+1} H_s^{\gamma,\delta}(z,y) \, ds \, dz \lesssim |y|^{1-\delta} M_t^{\gamma,\delta}(x,y). \tag{4.7}
$$

Despite the cancellations introduced by taking the gradient of the heat kernel, we do not expect that [\(4.7\)](#page-25-0) holds with $H_s^{\gamma,1}(z, y)$ and $M_t^{\gamma,0}$ instead of $H_s^{\gamma,\delta}(z, y)$ and $M_t^{\gamma,\delta}$. See Appendix [B.2](#page-38-0) for an argument.

Proof. By a scaling argument, it suffices to consider $t = 1$. Set

$$
S := \{(x, y) \in \mathbb{R}^d \times \mathbb{R}^d : |y| \ge 1, |x - y| \le (|x| \wedge |y|)/2 \}.
$$

We write

$$
\begin{split} \mathbf{1}_{|y|\geq 1} \mathbf{1}_{|x-y|\leq (|x|\wedge |y|)/2} & \int_{|x|/16\leq |z|\leq 4|x|} \int_{0}^{1} T_{t-s}^{\gamma}(x,z) |z|^{-\alpha+1} H_{s}^{\gamma,\delta}(z,y) \, ds \, dz \\ & = \mathbf{1}_{S}(x,y) \int_{\mathbb{R}^{d}} \int_{0}^{1} T_{1-s}^{\gamma}(x,z) |z|^{1-\alpha} H_{s}^{\gamma,\delta}(z,y) \, ds \, dz \\ & + \mathbf{1}_{S}(x,y) \int_{\mathbb{R}^{d}} \int_{0}^{1} T_{1-s}^{\gamma}(x,z) |z|^{1-\alpha} H_{s}^{\gamma,\delta}(z,y) \, ds \, dz \\ & =: F_{1} + F_{2}. \end{split}
$$

To bound F_2 , we use $|z| \ge |y|/8$ and $|y| \ge 1$, Lemma [4.1,](#page-24-2) and obtain

$$
F_2 \lesssim \frac{1}{|y|^{\alpha-1}} \int_0^1 \int_{\mathbb{R}^d} (1-s)^{-\frac{d}{\alpha}} \left(\frac{(1-s)^{1/\alpha}}{(1-s)^{1/\alpha} + |x-z|} \right)^{d+\gamma} s^{-\frac{d+\delta}{\alpha}} \left(\frac{s^{1/\alpha}}{s^{1/\alpha} + |z-y|} \right)^{d+\gamma} dz ds
$$

$$
\lesssim \frac{1}{|y|^{\alpha-1}} \left(\frac{1}{1+|x-y|} \right)^{d+\gamma} \lesssim |y|^{1-\delta} \left(\frac{1}{|y|} \right)^{\alpha-\delta} \left(\frac{1}{1+|x-y|} \right)^{d+\gamma} \lesssim |y|^{1-\delta} M_1^{\gamma,\delta}(x,y).
$$
\n(4.8)

We now consider F_1 . In this situation, we have $|x| \sim |y|, |x - z| \sim |x|$ and $|y - z| \sim |y|$ as $|z| \le |y|/8$ and $|x - y| \le (|x| \wedge |y|)/2$. Hence,

$$
F_1 \sim \int_{B(0,|y|/2)} \int_0^1 s^{-\frac{d+\delta}{\alpha}} (1-s)^{-\frac{d}{\alpha}} |z|^{1-\alpha} \left(\frac{|x|}{(1-s)^{1/\alpha}}\right)^{-d-\gamma} \left(\frac{|y|}{s^{1/\alpha}}\right)^{-d-\gamma} \left(1+\frac{s^{1/\alpha}}{|z|}\right)^{d-\beta} ds dz
$$

$$
\sim \frac{1}{|x|^{d+\gamma}|y|^{d+\gamma}} \int_0^1 (1-s)^{\gamma/\alpha} s^{(\gamma-\delta)/\alpha} \int_{B(0,|y|/8)} |z|^{1-\alpha} \left(1+\frac{s^{1/\alpha}}{|z|}\right)^{d-\beta} dz ds.
$$

By Lemma [2.1,](#page-9-2)

$$
\int_{B(0,|y|/8)} |z|^{-\alpha+1} \left(1 + \frac{s^{1/\alpha}}{|z|}\right)^{d-\beta} dz \le \int_{B(0,1)} |z|^{-\alpha+1} \left(1 + \frac{1}{|z|}\right)^{d-\beta} dz \n+ \int_{1 \le |z| \le |y|/8} |z|^{-\alpha+1} \left(1 + \frac{1}{|z|}\right)^{d-\beta} dz \n\sim \int_{B(0,1)} \frac{1}{|z|^{d-\beta+\alpha-1}} dz + \int_{1 \le |z| \le |y|/8} |z|^{-\alpha+1} dz \n\lesssim 1 + |y|^{d-\alpha+1} \sim |y|^{d-\alpha+1},
$$

where in the last inequality we used $|y| \gtrsim 1$. Plugging this into the bound of F_1 and using $|x| \sim |y| \gtrsim 1$, we obtain

$$
F_1 \lesssim \frac{|y|^{d-\alpha+1}}{|x|^{d+\gamma}|y|^{d+\gamma}} \int_0^1 (1-s)^{\gamma/\alpha} s^{(\gamma-\delta)/\alpha} ds \lesssim |y|^{1-\delta} \cdot |y|^{\delta-\gamma-\alpha} \cdot \frac{1}{|x|^{d+\gamma}} \lesssim |y|^{1-\delta} M_1^{\gamma,\delta-\gamma}(x,y).
$$

Since $M_1^{\gamma,\delta-\gamma}(x,y) \leq M_1^{\gamma,\delta}(x,y)$, this completes the proof of [\(4.7\)](#page-25-0).

Remark 4.3. Note that in [\(4.8\)](#page-25-1), the integral over $|z| \in [(1 - \varepsilon_1)|y|, (1 + \varepsilon_2)|y|]$ ^c for arbitrary but fixed $\varepsilon_1, \varepsilon_2 > 0$ is actually bounded by $M_1^{\gamma,0}(x, y)$ since in that case we can use

$$
|z|s^{-\frac{d+1}{\alpha}}\left(\frac{s^{1/\alpha}}{s^{\frac{1}{\alpha}}+|z-y|}\right)^{d+\gamma} \lesssim s^{-\frac{d}{\alpha}}\left(\frac{s^{1/\alpha}}{s^{\frac{1}{\alpha}}+|z-y|}\right)^{d+\gamma-1} \text{ for } |z| \in [(1-\varepsilon_1)|y|, (1+\varepsilon_2)|y|]^{c}.
$$

The remaining integrals are convergent whenever $\gamma \in (1, \alpha)$.

The following lemma concerns the region where no cancellation effects are expected anymore. It will be important in Subsection [4.3.](#page-30-0)

Lemma 4.4. Let
$$
\alpha \in (1, 2]
$$
, $\beta \in ((d + \alpha)/2, d + \alpha)$, $\gamma \in (0, 1)$, $t > 0$, and $x, y \in \mathbb{R}^d$. Then,

$$
\mathbf{1}_{|y| \le t^{1/\alpha}} \int_{|x|/16 \le |z| \le 4|x|} \int_0^t T_{t-s}^{\gamma}(x, z)|z|^{-\alpha+1} H_s^{\gamma, \delta}(z, y) ds dz
$$
\n
$$
+ \mathbf{1}_{|y| \ge t^{1/\alpha}} \mathbf{1}_{|x-y| \ge (|x|\wedge|y|)/2} \int_{|x|/16 \le |z| \le 4|x|} \int_0^t T_{t-s}^{\gamma}(x, z)|z|^{-\alpha+1} H_s^{\gamma, \delta}(z, y) ds dz \tag{4.9}
$$
\n
$$
\lesssim L_t^{\gamma, \delta}(x, y).
$$

Proof. By a scaling argument, it suffices to consider $t = 1$. We first prove

$$
{\bf 1}_{|y|\ge 1} {\bf 1}_{|x-y|\ge (|x|\wedge |y|)/2} \int\limits_{|x|/16\le |z|\le 4|x|} \int_0^1 T_{1-s}^\gamma(x,z) |z|^{-\alpha +1} H^{\gamma,\delta}_s(z,y)\, ds\, dz \lesssim L_1^{\gamma,\delta}(x,y).
$$

To do this, we break the integral with respect to dz as

$$
\int_{|x|/16\leq|z|\leq 4|x|}\dots dz = \int_{\substack{|x|/16\leq|z|\leq 4|x|\\|z|\geq |y|/100}}\dots dz + \int_{\substack{|x|/16\leq|z|\leq 4|x|\\|z|<|y|/100}}\dots dz.
$$

Subcase 1.1: $|x| \sim |z| \ge |y|/100$. By Lemma [4.1](#page-24-2) and $|z|^{-\alpha+1} \lesssim |y|^{-\alpha+1} \lesssim |y|^{1-\gamma}$, we have

$$
\begin{split} &\mathbf{1}_{|y|\geq 1}\mathbf{1}_{|x-y|\geq (|x|\wedge |y|)/2}\int\limits_{|x|/16\leq |z|\leq 4|x|}\int_{0}^{1}T_{1-s}^{\gamma}(x,z)|z|^{-\alpha+1}H_{s}^{\gamma,\delta}(z,y)\,ds\,dz\\ &\lesssim |y|^{1-\alpha}\Big(\frac{1}{1+|x-y|}\Big)^{d+\gamma}\int_{0}^{1}s^{-\delta/\alpha}\,ds\\ &\lesssim |y|^{1-\delta}\Big(\frac{1}{1+|x-y|}\Big)^{d+\gamma}\\ &\lesssim L_{1}^{\gamma,\delta}(x,y). \end{split}
$$

Subcase 1.2: $|x| ∼ |z| < |y|/100$. In this case $|x - y| ∼ |z - y| ∼ |y|$. Hence,

$$
\begin{array}{lcl}\n\mathbf{1}_{|y|\geq 1}\mathbf{1}_{|x-y|\geq (|x|\wedge |y|)/2} & \int\limits_{|x|/16\leq |z|\leq 4|x| \atop |z|<|y|/100}\int_0^1 T^\gamma_{1-s}(x,z)|z|^{-\alpha+1}H^{\gamma,\delta}_s(z,y)\,ds\,dz\\
\lesssim & \int\limits_{|x|/16\leq |z|\leq 4|x| \atop |z|<|y|/100}\int_0^1 T^\gamma_{1-s}(x,z)|z|^{-\alpha+1}\frac{1}{s^{\frac{d+\delta}{\alpha}}}\Big(\frac{s^{1/\alpha}}{|y|}\Big)^{d+\gamma}ds\,dz.\n\end{array}
$$

By Lemma [2.1,](#page-9-2)

$$
\int_{\mathbb{R}^d} T_{1-s}^{\gamma}(x,z) |z|^{-\alpha+1} dz \lesssim \int_{|z| \le (1-s)^{1/\alpha}} \frac{1}{(1-s)^{d/\alpha}} \left(\frac{(1-s)^{1/\alpha}}{|z|}\right)^{d-\beta} |z|^{1-\alpha} dz + \int_{|z| > (1-s)^{1/\alpha}} \frac{1}{(1-s)^{d/\alpha}} \left(\frac{(1-s)^{1/\alpha}}{(1-s)^{1/\alpha} + |x-z|}\right)^{d+\gamma} |z|^{1-\alpha} dz \tag{4.10}
$$

$$
\lesssim (1-s)^{-1+1/\alpha}.
$$

Hence,

$$
\begin{split} \mathbf{1}_{|y|\geq 1} \mathbf{1}_{|x-y|\geq (|x|\wedge |y|)/2} & \int_{\substack{|x|/16 \leq |z|\leq 4|x| \\ |z| < |y|/100}} \int_{0}^{1} T_{1-s}^{\gamma}(x, z) |z|^{-\alpha+1} H_{s}^{\gamma, \delta}(z, y) \, ds \, dz \\ &\lesssim \int_{0}^{1} (1-s)^{-1+1/\alpha} \frac{s^{(\gamma-1)/\alpha}}{|y|^{d+\gamma}} ds \lesssim \frac{1}{|y|^{d+\gamma}} \lesssim L_{1}^{\gamma, \delta}(x, y). \end{split}
$$

We have proved that

$$
\mathbf{1}_{|y|\geq 1}\mathbf{1}_{|x-y|\geq (|x|\wedge |y|)/2}\int\limits_{|x|/16\leq |z|\leq 4|x|}\int_0^1 T_{1-s}^\gamma(x,z)|z|^{-\alpha+1}H_s^{\gamma,\delta}(z,y)\,ds\,dz\lesssim L_1^{\gamma,\delta}(x,y).
$$

It remains to prove

$$
{\bf 1}_{|y|<1}\int\limits_{|x|/16\le |z|\le 4|x|}\int_0^1 T_{1-s}^\gamma(x,z)|z|^{-\alpha+1}H_s^{\gamma,\delta}(z,y)\,ds\,dz\lesssim L_1^{\gamma,\delta}(x,y).
$$

We now break the integral with respect to dz into four integrals as

$$
\int_{|x|/16 \le |z| \le 4|x|} \dots dz = \int_{\substack{|x|/16 \le |z| \le 4|x| \\ |y|/100 < |z| < 2|y|}} \dots dz + \int_{\substack{|x|/16 \le |z| \le 4|x| \\ |z| \ge t^{1/\alpha}}} \dots dz + \int_{\substack{|x|/16 \le |z| \le 4|x| \\ |y| < |z| \le |z|x|}} \dots dz + \int_{\substack{|x|/16 \le |z| \le 4|x| \\ |z| \le |y|/100}} \dots dz.
$$

Subcase 2.1: $|y|/100 < |z| < 2|y|$. In this case, by Lemma [4.1](#page-24-2) and $|x| ∼ |z| ∼ |y|$, we have

$$
\mathbf{1}_{|y|<1} \int\limits_{\substack{|x|/16\leq |z|\leq 4|x\\ |y|/100<|z|<2|y|}} \int_0^1 T^\gamma_{1-s}(x,z) |z|^{1-\alpha} H^{\gamma,\delta}_s(z,y)\, ds\, dz\lesssim L^{\gamma,\delta}_1(x,y).
$$

Subcase 2.2: $|z| \ge 2|y|$. Similarly to Subcase 2.1, we have

$$
\begin{split} \mathbf{1}_{|y|<1}&\int\limits_{|x|/16\leq |z|\leq 4|x|}\int_0^1 T^\gamma_{1-s}(x,z)|z|^{-\alpha+1}H^{\gamma,\delta}_s(z,y)\,ds\,dz\\ &\lesssim |y|^{-\alpha+1}\Big(\frac{1}{1+|x-y|}\Big)^{d+1+\gamma}\lesssim L^{\gamma,\delta}_1(x,y). \end{split}
$$

Subcase 2.3: $|z| \le |y|/100$. In this case, $|z - y| \sim |y| \ge |x - y|$. Hence,

$$
\begin{split}\n\mathbf{1}_{|y|<1} \int_{|x|/16 \leq |z| \leq 4|x| \atop |z| \leq |y|/100} \int_{0}^{1} T_{1-s}^{\gamma}(x,z) |z|^{-\alpha+1} H_{s}^{\gamma,\delta}(z,y) \, ds \, dz \\
&\lesssim \int_{|x|/16 \leq |z| \leq 4|x| \atop |z| \leq |y|/100} \int_{0}^{1} T_{1-s}^{\gamma}(x,z) |z|^{-\alpha+1} \frac{1}{s^{\frac{d+\delta}{\alpha}}} \left(\frac{s^{1/\alpha}}{s^{1/\alpha} + |y|} \right)^{d+\gamma} \left(\mathbf{1}_{s \in (0,1/2)} + \mathbf{1}_{s \in (1/2,1)} \right) ds dz \\
&=: E_{1} + E_{2}.\n\end{split}
$$

To bound E_2 , we use (4.10) , and obtain

$$
E_2 \lesssim \left(\frac{1}{1+|x-y|}\right)^{d+\gamma} \int_{1/2}^1 (1-s)^{-1+1/\alpha} ds \lesssim L^{\gamma,\delta}(x,y).
$$

For the term E_1 , we have

$$
E_1 \lesssim \int_0^{1/2} \int_{|x|/16 \le |z| \le 4|x|} \left(\frac{1}{|z|}\right)^{d-\beta} |z|^{-\alpha+1} \frac{1}{s^{\frac{d+\delta}{\alpha}}} \left(\frac{s^{1/\alpha}}{s^{1/\alpha} + |y|}\right)^{d+\gamma} dz ds
$$

$$
\lesssim \int_0^{1/2} \left(\frac{1}{|x|}\right)^{d-\beta} |x|^{d-\alpha+1} \frac{1}{s^{\frac{d+\delta}{\alpha}}} \left(\frac{s^{1/\alpha}}{s^{1/\alpha} + |y|}\right)^{d+\gamma} dz ds
$$

$$
\lesssim |x|^{\beta+1-\alpha} \left[\int_0^{|y|^\alpha} \dots + \int_{|y|^\alpha}^{1/2} \dots\right]
$$

$$
\lesssim |y|^{\beta+1-d-\alpha} \lesssim L_1^{\gamma,\delta}(x,y).
$$

Therefore,

$$
\mathbf{1}_{|y|<1} \int\limits_{|x|/16\leq |z|\leq 4|x|} \int_0^1 T^\gamma_{1-s}(x,z) |z|^{-\alpha+1} H^{\gamma,\delta}_s(z,y)\, ds\, dz \lesssim L^{\gamma,\delta}_1(x,y).
$$

This completes our proof.

4.2. An estimate for $|Q_t(x, y)|$. The estimate in this subsection will be used to prove part (1) in Theorem [1.4](#page-5-1) for $\alpha s < \alpha - 1$.

Proposition 4.5. Let $\alpha \in (1, 2)$, $\beta \in ((d + \alpha)/2, d + \alpha)$, and $\kappa = \Psi(\beta)$ be defined by [\(1.7\)](#page-2-2). Then, for any $\gamma \in (0, \alpha)$, $x, y \in \mathbb{R}^d$, and $t > 0$,

$$
|Q_t(x,y)| \lesssim L_t^{\gamma,1}(x,y) + M_t^{\gamma,1}(x,y). \tag{4.11}
$$

Proof. By Proposition [2.9,](#page-17-0)

$$
|\widetilde{p}_t(x, y)| + t |\widetilde{p}_{t,1}(x, y)| \lesssim H_t^{\gamma,1}(x, y) \text{ and}
$$

$$
|p_t(x, y)| + t |p_{t,1}(x, y)| \lesssim T_t^{\gamma}(x, y)
$$

for all $t > 0$ and $x, y \in \mathbb{R}^d$. We now consider two cases.

Case 1: $[|y| \le t^{1/\alpha}]$ OR $[|y| \ge t^{1/\alpha}$ and $|x - y| \ge (|x| \wedge |y|)/2]$. Since in this case

$$
H_t^{\gamma,1}(x,y) + T_t^{\gamma}(x,y) \lesssim L_t^{\gamma,1}(x,y),
$$

we get $|Q_t(x, y)| \lesssim L_t^{\gamma, 1}$ $t^{\gamma,1}(x,y)$ as desired.

Case 2: $|y| \ge t^{1/\alpha}$ and $|x - y| < (|x| \wedge |y|)/2$.

By Duhamel's formula,

$$
\tilde{p}_t(x, y) - p_t(x, y) = \kappa \int_0^t \int_{\mathbb{R}^d} p_{t-s}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_s(z, y) dz ds
$$

\n
$$
= \kappa \int_0^{t/2} \int_{\mathbb{R}^d} p_{t-s}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_s(z, y) dz ds
$$
(4.12)
\n
$$
+ \kappa \int_0^{t/2} \int_{\mathbb{R}^d} p_s(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_{t-s}(z, y) dz ds.
$$

Differentiating both sides with respect to t and multiplying by t gives

$$
Q_t(x, y) = \kappa t \int_{\mathbb{R}^d} p_{t/2}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_{t/2}(z, y) dz + \kappa t \int_0^{t/2} \int_{\mathbb{R}^d} p_{t-s, 1}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_s(z, y) dz ds + \kappa t \int_{t/2}^t \int_{\mathbb{R}^d} p_{t-s}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_{s, 1}(z, y) dz ds =: I_1 + I_2 + I_3.
$$
 (4.13)

Without loss of generality, we now assume $t = 1$. The term I_1 can be written as

$$
\kappa \int_{\mathbb{R}^d} p_{1/2}(x,z)|z|^{-\alpha}z \cdot \nabla_z \tilde{p}_{1/2}(z,y)dz = 6\kappa \int_{1/3}^{1/2} \int_{\mathbb{R}^d} p_{1/2}(x,z)|z|^{-\alpha}z \cdot \nabla_z \tilde{p}_{1/2}(z,y)dzds.
$$

By Lemma [4.2](#page-25-2) and [\(2.19\)](#page-17-5), we have $H_{1-\alpha}^{\gamma,1}$ $T_{1-s}^{\gamma,1}(z,y) \sim H_{1/2}^{\gamma,1}$ $T_{1/2}^{\gamma,1}(z,y) \gtrsim |\nabla_z \tilde{p}_{1/2}(z,y)|$ and $T_s^{\gamma}(\cdot,\cdot) \sim$ T_1^{γ} $T_{1/2}^{\gamma}(\cdot,\cdot) \gtrsim p_{1/2}(\cdot,\cdot)$ for $s \in [1/3,1/2]$. Thus,

$$
I_1 \lesssim \int_{1/3}^{1/2} \int_{\mathbb{R}^d} T_{1-s}^{\gamma}(x, z) |z|^{1-\alpha} H_s^{\gamma, 1}(z, y) dz ds \lesssim \int_0^1 \int_{\mathbb{R}^d} T_{1-s}^{\gamma}(x, z) |z|^{1-\alpha} H_s^{\gamma, 1}(z, y) dz ds
$$

$$
\lesssim M_1^{\gamma, 1}(x, y).
$$

We now estimate I_2 . Since $1 - s \sim t$ for $s \in (0, 1/2)$,

$$
|p_{1-s,1}(x,z)| \sim (1-s)|p_{1-s,1}(x,z)| \lesssim T_{1-s}^{\gamma}(x,z)
$$

and

$$
|\nabla_z \tilde{p}_s(z, y)| \lesssim H_s^{\gamma,1}(z, y)
$$

due to [\(2.19\)](#page-17-5). Therefore, by Lemma [4.2,](#page-25-2)

$$
I_2 \lesssim \int_0^{1/2} \int_{\mathbb{R}^d} T_{1-s}^{\gamma}(x, z) |z|^{1-\alpha} H_s^{\gamma, 1}(z, y) dz ds \lesssim M_1^{\gamma, 1}(x, y).
$$

For the last term I_3 , by Proposition [2.11,](#page-17-4)

$$
|\nabla_z \tilde{p}_{s,1}(z,y)| \lesssim s^{-1} H_s^{\gamma,1}(z,y) \sim H_s^{\gamma,1}(z,y)
$$

for $s \in [1/2, 1]$. This, together with $|p_{1-s}(x, z)| \lesssim T_1^{\gamma}$. $_{1-s}^{\gamma}(x, z)$ and (4.7) in Lemma [4.2,](#page-25-2) implies

$$
I_3 \lesssim \int_{1/2}^1 \int_{\mathbb{R}^d} T_{1-s}^{\gamma}(x, z) |z|^{-\alpha+1} H_s^{\gamma, 1}(z, y) \, dz \, ds \lesssim M_1^{\gamma, 1}(x, y).
$$

This completes our proof. \Box

4.3. An estimate for $|(Q_t f)(x)|$. In this subsection, we prove an estimate, which we use to show part (2) in Theorem [1.4.](#page-5-1)

Proposition 4.6. Let $\alpha \in (1, 2)$, $\beta \in ((d + \alpha)/2, d + \alpha)$, and $\kappa = \Psi(\beta)$ be defined by [\(1.7\)](#page-2-2). Then, for any $\gamma \in (0,1)$, $x \in \mathbb{R}^d$, $t > 0$, and $f \in L^p(\mathbb{R}^d)$ with $p > 1 \vee d/\beta$,

$$
\left| (Q_t f)(x) \right| \lesssim \int_{\mathbb{R}^d} \left[L_t^{\gamma,1}(x, y) + M_t^{\gamma,0}(x, y) \right] |f(y)| dy
$$

+
$$
\int_{\mathbb{R}^d} L_t^{\gamma, \gamma}(x, y) \left| (\Lambda_0^{\frac{1-\gamma}{\alpha}} f)(y) \right| dy
$$

+
$$
\int_{\mathbb{R}^d} M_t^{\gamma, \gamma}(x, y) ||y|^{1-\gamma} (\Lambda_0^{\frac{1-\gamma}{\alpha}} f)(y) | dy.
$$
 (4.14)

Proof. By Duhamel's formula,

$$
\tilde{p}_t(x, y) - p_t(x, y) = \kappa \int_0^t \int_{\mathbb{R}^d} p_{t-s}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_s(z, y) dz ds
$$

\n
$$
= \kappa \int_0^{t/2} \int_{\mathbb{R}^d} p_{t-s}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_s(z, y) dz ds
$$

\n
$$
+ \kappa \int_0^{t/2} \int_{\mathbb{R}^d} p_s(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_{t-s}(z, y) dz ds.
$$

Differentiating both sides with respect to t and multiplying by t gives

$$
Q_t(x,y) = \kappa t \int_{\mathbb{R}^d} p_{t/2}(x,z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_{t/2}(z,y) dz
$$

+ $\kappa t \int_0^{t/2} \int_{\mathbb{R}^d} p_{t-s,1}(x,z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_s(z,y) dz ds$
+ $\kappa t \int_{t/2}^t \int_{\mathbb{R}^d} p_{t-s}(x,z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_{s,1}(z,y) dz ds,$

which implies

$$
Q_t f(x) = \kappa t \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} p_{t/2}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_{t/2}(z, y) f(y) dz dy
$$

+ $\kappa t \int_{\mathbb{R}^d} \int_0^{t/2} \int_{\mathbb{R}^d} p_{t-s,1}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_s(z, y) f(y) dz ds dy$
+ $\kappa t \int_{\mathbb{R}^d} \int_{t/2}^t \int_{\mathbb{R}^d} p_{t-s}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_{s,1}(z, y) f(y) dz ds dy.$

Without loss of generality, we now assume $t = 1$. Set $S_{x,1} = \{z : |x|/16 < |z| < 4|x|\},$ $S_{x,2} = \mathbb{R}^d \setminus S_{x,1}, R_x = \{y : |y| \ge 1, |x - y| < (|x| \wedge |y|)/2 \}.$ Then we can write

$$
(Q_1f)(x) = \int_{\mathbb{R}^d \setminus R_x} Q_1(x, y) f(y) dy + \int_{R_x} Q_1(x, y) f(y) dy.
$$

The first term can be done similarly to Case 1 in the proof of Proposition [4.5.](#page-28-1) We have

$$
\left| \int_{\mathbb{R}^d \setminus R_x} Q_1(x, y) f(y) dy \right| \lesssim \int_{\mathbb{R}^d} L_1^{\gamma, 1}(x, y) |f(y)| dy.
$$

For the second term, we write

$$
\int_{R_x} Q_1(x, y) f(y) dy = \kappa \sum_{i=1,2} \int_{R_x} \int_{S_{x,i}} p_{1/2}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_{1/2}(z, y) f(y) dz dy \n+ \kappa \sum_{i=1,2} \int_{R_x} \int_0^{1/2} \int_{S_{x,i}} p_{1-s,1}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_s(z, y) f(y) dz ds dy \n+ \kappa \sum_{i=1,2} \int_{R_x} \int_{1/2}^1 \int_{S_{x,i}} p_{1-s}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_{s,1}(z, y) f(y) dz ds dy \n=: Q_{1,1} f(x) + Q_{1,2} f(x).
$$

For $Q_{1,2}f$, by the kernel bounds in Proposition [2.9,](#page-17-0) Lemmas [2.10](#page-17-1) and [2.11](#page-17-4) and Lemma [4.2,](#page-25-2) (using $H_t^{\gamma+1,\gamma} \leq H_t^{\gamma,\gamma}$ $\binom{\gamma,\gamma}{t}$

$$
|Q_{1,2}f| \lesssim \int_{\mathbb{R}^d} L_1^{\gamma,1}(x,y) |f(y)| dy + \int_{\mathbb{R}^d} M^{\gamma,0}(x,y) |f(y)| dy.
$$

To study $Q_{1,1}f$, we use $\nabla e^{-\Lambda_0} = \nabla \Lambda_0^{-\frac{1-\gamma}{\alpha}} e^{-\Lambda_0} \Lambda_0^{\frac{1-\gamma}{\alpha}}$. Thus,

$$
Q_{1,1}f = \kappa \int_{\mathbb{R}^d} \int_{S_{x,1}} p_{1/2}(x, z) |z|^{-\alpha} z \cdot \nabla_z \Lambda_0^{-\frac{1-\gamma}{\alpha}} e^{-\frac{1}{2}\Lambda_0}(z, y) \Lambda_0^{\frac{1-\gamma}{\alpha}} f(y) dz dy + \kappa \int_{\mathbb{R}^d} \int_0^{1/2} \int_{S_{x,1}} p_{1-s,1}(x, z) |z|^{-\alpha} z \cdot \nabla_z \Lambda_0^{-\frac{1-\gamma}{\alpha}} e^{-s\Lambda_0}(z, y) \Lambda_0^{\frac{1-\gamma}{\alpha}} f(y) dz ds dy + \kappa \int_{\mathbb{R}^d} \int_{1/2}^1 \int_{S_{x,1}} p_{1-s}(x, z) |z|^{-\alpha} z \cdot \nabla_z \Lambda_0^{-\frac{1-\gamma}{\alpha}+1} e^{-s\Lambda_0}(z, y) \Lambda_0^{\frac{1-\gamma}{\alpha}} f(y) dz ds dy - \kappa \int_{\mathbb{R}^d \setminus R_x} \int_{S_{x,1}} p_{1/2}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_{1/2}(z, y) f(y) dz dy - \kappa \int_{\mathbb{R}^d \setminus R_x} \int_0^{1/2} \int_{S_{x,1}} p_{1-s,1}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_s(z, y) f(y) dz ds dy - \kappa \int_{\mathbb{R}^d \setminus R_x} \int_0^1 \int_{S_{x,1}} p_{1-s}(x, z) |z|^{-\alpha} z \cdot \nabla_z \tilde{p}_{s,1}(z, y) f(y) dz ds dy.
$$
\n(4.15)

We now set

$$
Q_{1,1}^{1}f(x) = \kappa \int_{\mathbb{R}^{d}} \int_{S_{x,1}} p_{1/2}(x, z)|z|^{-\alpha} z \cdot \nabla_{z} \Lambda_{0}^{-\frac{1-\gamma}{\alpha}} e^{-\frac{1}{2}\Lambda_{0}}(z, y) \Lambda_{0}^{\frac{1-\gamma}{\alpha}} f(y) dz dy
$$

+ $\kappa \int_{\mathbb{R}^{d}} \int_{0}^{1/2} \int_{S_{x,1}} p_{1-s,1}(x, z)|z|^{-\alpha} z \cdot \nabla_{z} \Lambda_{0}^{-\frac{1-\gamma}{\alpha}} e^{-s\Lambda_{0}}(z, y) \Lambda_{0}^{\frac{1-\gamma}{\alpha}} f(y) dz ds dy$ (4.16)
+ $\kappa \int_{\mathbb{R}^{d}} \int_{1/2}^{1} \int_{S_{x,1}} p_{1-s}(x, z)|z|^{-\alpha} z \cdot \nabla_{z} \Lambda_{0}^{-\frac{1-\gamma}{\alpha}+1} e^{-s\Lambda_{0}}(z, y) \Lambda_{0}^{\frac{1-\gamma}{\alpha}} f(y) dz ds dy$

and

$$
Q_{1,1}^{2}f(x) = -\kappa \int_{\mathbb{R}^{d} \setminus R_{x}} \int_{S_{x,1}} p_{1/2}(x, z)|z|^{-\alpha} z \cdot \nabla_{z} \tilde{p}_{1/2}(z, y) f(y) dz dy
$$

$$
- \kappa \int_{\mathbb{R}^{d} \setminus R_{x}} \int_{0}^{1/2} \int_{S_{x,1}} p_{1-s,1}(x, z)|z|^{-\alpha} z \cdot \nabla_{z} \tilde{p}_{s}(z, y) f(y) dz ds dy
$$
(4.17)

$$
- \kappa \int_{\mathbb{R}^{d} \setminus R_{x}} \int_{1/2}^{1} \int_{S_{x,1}} p_{1-s}(x, z)|z|^{-\alpha} z \cdot \nabla_{z} \tilde{p}_{s,1}(z, y) f(y) dz ds dy.
$$

From the kernel bounds in Proposition [2.9](#page-17-0) and Lemma [2.12,](#page-17-6) by using Lemmas [4.4](#page-26-0) and [4.2,](#page-25-2) we obtain

$$
Q_{1,1}^1 f(x) \lesssim \int_{\mathbb{R}^d} L_1^{\gamma,\gamma}(x,y) \big| (\Lambda_0^{\frac{1-\gamma}{\alpha}} f)(y) \big| dy + \int_{\mathbb{R}^d} M_1^{\gamma,\gamma}(x,y) \big| |y|^{1-\gamma} (\Lambda_0^{\frac{1-\gamma}{\alpha}} f)(y) \big| dy. \tag{4.18}
$$

To bound $Q_{1,1}^2 f(x)$, we use the kernel bounds in Proposition [2.9,](#page-17-0) Lemmas [2.10](#page-17-1) and [2.11,](#page-17-4) and Lemma [4.4.](#page-26-0) We obtain

$$
|Q_{1,1}^{2}f(x)| \lesssim \int_{\mathbb{R}^{d}} L_{1}^{\gamma,1}(x,y)|f(y)|dy.
$$

This completes our proof.

5. Proof of the reversed Hardy inequality (Theorem [1.4\)](#page-5-1)

We now use the previous bounds for the difference of kernels to prove Theorem [1.4,](#page-5-1) i.e., the reversed Hardy inequality, expressed in terms of our square functions.

Proof of part (1) in Theorem [1.4.](#page-5-1) Let $\alpha s < \alpha - 1$. By Proposition [4.5,](#page-28-1)

$$
\left(\int_{0}^{\infty} t^{-2s} \left| \left(t\Lambda_{0}e^{-t\Lambda_{0}} - t\Lambda_{\kappa}e^{-t\Lambda_{\kappa}}\right) f(x)\right|^{2} \frac{dt}{t}\right)^{1/2} \n= \left[\sum_{j\in\mathbb{Z}} \int_{2^{\alpha j}}^{2^{\alpha(j+1)}} t^{-2s} \left| \left(t\Lambda_{0}e^{-t\Lambda_{0}} - t\Lambda_{\kappa}e^{-t\Lambda_{\kappa}}\right) f(x)\right|^{2} \frac{dt}{t}\right]^{1/2} \n\lesssim \left[\sum_{j\in\mathbb{Z}} \int_{2^{\alpha j}}^{2^{\alpha(j+1)}} t^{-2s} \left(\int_{\mathbb{R}^{d}} [L_{t}^{\gamma,1}(x,y) + M_{t}^{\gamma,1}(x,y)] |f(y)| dy\right)^{2} \frac{dt}{t}\right]^{1/2} \n\lesssim \sum_{j\in\mathbb{Z}} 2^{-js\alpha} \int_{\mathbb{R}^{d}} \left[L_{2^{\alpha j}}^{\gamma,1}(x,y) + M_{2^{j\alpha}}^{\gamma,1}(x,y)\right] |y|^{\alpha s} \frac{|f(y)|}{|y|^{\alpha s}} dy
$$
\n(5.1)

where in the last inequality we used the embedding $\ell_1 \hookrightarrow \ell_2$. Thus, it suffices to show the $L^p(\mathbb{R}^d)$ -boundedness of the operator with kernel

$$
\sum_{j\in\mathbb{Z}} 2^{-j\alpha s} \left[L_{2^{\alpha j}}^{\gamma,1}(x,y) + M_{2^{j\alpha}}^{\gamma,1}(x,y) \right] |y|^{\alpha s}.
$$
 (5.2)

To that end, we use Schur tests similar to those in [\[Mer21\]](#page-44-4). For the sake of completeness, we give the details. We begin with the Schur test involving $M_{2^{j\alpha}}^{\gamma,1}$. In particular, we will see that these Schur tests require $\alpha s < \alpha - 1$. In the following, let $N := 2^{-j} \in 2^{\mathbb{Z}}$. Using $1_{x\in\mathbb{R}^d}1_{|y|>t^{1/\alpha}}\leq 1_{|x|\vee|y|>t^{1/\alpha}}$ and noting that on the support of $M_{N^{-\alpha}}^{\gamma,1}(x,y)$, we have $|y|/2\leq$ $|x| \leq 2|y|$, we may replace the kernel with a symmetric kernel, i.e., it suffices to carry out a single Schur test. We estimate

$$
\sup_{y \in \mathbb{R}^d} \int_{\mathbb{R}^d} dx \sum_{N \in 2^{\mathbb{Z}}} N^{\alpha s} M_{N^{-\alpha}}^{\gamma,1}(x,y) (|x||y|)^{\frac{\alpha s}{2}} \n\sim \sup_{y \in \mathbb{R}^d} \int_{\frac{1}{2}|y| \le |x| \le 2|y|} dx \sum_{N \ge (|x| \vee |y|)^{-1}} N^{\alpha s} \frac{N^{1-\alpha+d}}{(|x| \vee |y|)^{\alpha-1}} \left(1 + \frac{N^{-\gamma-d}}{|x-y|^{d+\gamma}}\right) (|x||y|)^{\frac{\alpha s}{2}} \n\lesssim \sup_{y \in \mathbb{R}^d} |y|^{\alpha s - \alpha + 1} \int_{\frac{1}{2}|y| \le |x| \le 2|y|} dx \sum_{N \ge (2|y|)^{-1}} N^{\alpha s + 1 - \alpha + d} \left(1 + \frac{N^{-\gamma-d}}{|x-y|^{d+\gamma}}\right).
$$
\n(5.3)

Interchanging the order of integration and summation shows that the right-hand side is bounded by

$$
\sup_{y \in \mathbb{R}^d} |y|^{\alpha s + 1 - \alpha} \sum_{N \ge (2|y|)^{-1}} N^{\alpha s + 1 - \alpha + d} \int_{\frac{1}{2}|y| \le |x| \le 2|y|} dx \left(1 \wedge \frac{N^{-\gamma - d}}{|x - y|^{d + \gamma}} \right)
$$

$$
\le \sup_{y \in \mathbb{R}^d} |y|^{\alpha s + 1 - \alpha} \sum_{N \ge (2|y|)^{-1}} N^{\alpha s + 1 - \alpha + d} \int_{\mathbb{R}^d} dx \left(1 \wedge \frac{N^{-\gamma - d}}{|x - y|^{d + \gamma}} \right)
$$

$$
\sim \sup_{y \in \mathbb{R}^d} |y|^{\alpha s + 1 - \alpha} \sum_{N \ge (2|y|)^{-1}} N^{\alpha s + 1 - \alpha} \sim 1
$$
 (5.4)

where we used $\gamma > 0$ and $\alpha s < \alpha - 1$. This concludes the Schur test involving the kernel $M_{2^{j\alpha}}^{\gamma,1}$.

It remains to carry out the Schur tests involving $L_{2\alpha j}^{\gamma,1}$. The L^p -boundedness of the second summand of $L_t^{\gamma,1}$ $\tilde{t}^{\gamma,1}$ follows from

$$
\int_0^\infty \frac{dt}{t} \int_{\mathbb{R}^d} \mathbf{1}_{\{|y| \le t^{1/\alpha}, |x| \sim |y|\}} \frac{|y|^{\alpha s}}{t^s} \left(\frac{|y|}{t^{1/\alpha}}\right)^{1+\beta-d-\alpha} t^{-\frac{d}{\alpha}} |g(y)| dy \lesssim (\mathcal{M}_1 g)(x)
$$

and the L^p -boundedness of the Hardy–Littlewood maximal operator. To treat the other two summands, let $N := 2^{-j} \in 2^{\mathbb{Z}}$ as before. The tests for the regions $|x| \vee |y| < t^{1/\alpha}$ and $|x| \wedge |y| >$ $t^{1/\alpha}$ are similar to those in [\[Mer21\]](#page-44-4); however, unlike in [Mer21], where $p \in (d/\beta, d/(d - \beta))$ was required, we only need $p > d/\beta$ here because of the absent singular weight $(|x|/t^{1/\alpha})^{\beta-d}$ in the region $|x| \vee |y| < t^{1/\alpha}$, which is due to the non-symmetry of Λ_{κ} . Let us now give the details. First, we bound

$$
|y|^{\beta - d - (\alpha - \gamma)} \sum_{N \le |y|^{-1}} N^{\alpha s + \beta - (\alpha - \gamma)} \left(\frac{N^{-1}}{N^{-1} + |x - y|} \right)^{d + \gamma} + \sum_{N \ge |y|^{-1}} N^{\alpha s + d} \left(\frac{N^{-1}}{N^{-1} + |x - y|} \right)^{d + \gamma} \mathbf{1}_{|x - y| > (|x| \wedge |y|)/2} \lesssim |y|^{\beta - d} |x - y|^{-\alpha s - \beta} + |x - y|^{-d - \alpha s} \mathbf{1}_{|x - y| > \frac{|x| \wedge |y|}{2}},
$$
\n
$$
(5.5)
$$

where the summability relied on $\alpha s+\beta>0$ (which follows from $\beta\geq (d+\alpha)/2$) and $\alpha s-\gamma<0$ (which follow from $s < 1$ and that $\gamma < \alpha$ may be chosen arbitrarily close to α). Thus,

$$
\left\| \int_{\mathbb{R}^d} dy \sum_{N \in 2^{\mathbb{Z}}} N^{\alpha s} L_{N^{-\alpha}}^{\gamma,1}(x,y) |y|^{\alpha s} g(y) \right\|_{L^p(\mathbb{R}^d)}
$$

\n
$$
\lesssim \|g\|_{L^p(\mathbb{R}^d)} + \left\| \int_{\mathbb{R}^d} dy |y|^{\alpha s + \beta - d} |x - y|^{-\alpha s - \beta} g(y) \right\|_{L^p(\mathbb{R}^d)}
$$

\n
$$
+ \left\| \int_{\mathbb{R}^d} dy \frac{\mathbf{1}_{|x-y| > (|x| \wedge |y|)/2}}{|x - y|^{d + \alpha s}} |y|^{\alpha s} g(y) \right\|_{L^p(\mathbb{R}^d)}.
$$
\n
$$
(5.6)
$$

Now, we estimate the last two summands by multiples of $||g||_p$ using Schur tests with weights being powers of $|x|/|y|$. By

$$
\int\limits_{|x-y| > (|x|\wedge |y|)/2} \frac{dy}{|x-y|^{d+\alpha s}} |y|^{\alpha s} \left(\frac{|x|}{|y|}\right)^{\delta/p} + \int\limits_{|x-y| > (|x|\wedge |y|)/2} \frac{dx}{|x-y|^{d+\alpha s}} |y|^{\alpha s} \left(\frac{|y|}{|x|}\right)^{\delta/p'} < \infty
$$

with $0 < \delta/p < d + \alpha s$ and $-\alpha s < \delta/p' < d$ (such δ exist for all $1 < p < \infty$), the second summand on the right-hand side of [\(5.6\)](#page-33-0) is bounded by $||g||_{L^p(\mathbb{R}^d)}$. We now consider the first summand on the right-hand side of [\(5.6\)](#page-33-0). We have

$$
\int_{\mathbb{R}^d} |y|^{\alpha s+\beta-d} |x-y|^{-\alpha s-\beta} \left(\frac{|x|}{|y|}\right)^{\delta/p} \, dy + \int_{\mathbb{R}^d} |y|^{\alpha s+\beta-d} |x-y|^{-\alpha s-\beta} \left(\frac{|y|}{|x|}\right)^{\delta/p'} \, dx < \infty
$$

if $0 < \delta/p < \alpha s + \beta$ and $d - \alpha s - \beta < \delta/p' < d$, or, equivalently, $p > \delta/(\alpha s + \beta)$. Since $p > d/\beta$ and $d/\beta > \delta/(\alpha s + \beta)$ (since $d(\alpha s + \beta) > \alpha d > \delta \beta$ because $d > \beta > \alpha$ and $\delta < \alpha$), there are δ such that the condition $\delta/p < \alpha s + \beta$ is fulfilled. Thus, the first summand on the right-hand side of [\(5.6\)](#page-33-0) is bounded by $||g||_{L^p(\mathbb{R}^d)}$, too. This concludes the proof of part (1) in Theorem [1.4.](#page-5-1)

Proof of part (2) in Theorem [1.4](#page-5-1) We argue as before, but use Proposition [4.6](#page-30-1) instead of Proposition [4.5.](#page-28-1) More precisely, fix $s \in (0,1)$ and take $\gamma \in (0,1)$ such that $1-\gamma \leq \alpha s < \alpha - \gamma$. Then, by Proposition [4.6,](#page-30-1)

$$
\begin{split}\n&\left(\int_{0}^{\infty} t^{-2s} \left| \left(t\Lambda_{0}e^{-t\Lambda_{0}} - t\Lambda_{\kappa}e^{-t\Lambda_{\kappa}}\right) f(x)\right|^{2} \frac{dt}{t}\right)^{1/2} \\
&= \left[\sum_{j\in\mathbb{Z}} \int_{2^{\alpha j}}^{2^{\alpha(j+1)}} t^{-2s} \left| \left(t\Lambda_{0}e^{-t\Lambda_{0}} - t\Lambda_{\kappa}e^{-t\Lambda_{\kappa}}\right) f(x)\right|^{2} \frac{dt}{t}\right]^{1/2} \\
&\leq \sum_{j\in\mathbb{Z}} 2^{-js\alpha} \int_{\mathbb{R}^{d}} \left[L_{2^{\alpha(j+1)}}^{\gamma,\gamma}(x,y)|y|^{\gamma-1} + M_{2^{j\alpha}}^{\gamma,\gamma}(x,y)\right] |y|^{\alpha s} \frac{|\Lambda_{0}^{\frac{1-\gamma}{\alpha}} f(y)|}{|y|^{\alpha s+\gamma-1}} dy \\
&+ \sum_{j\in\mathbb{Z}} 2^{-js\alpha} \int_{\mathbb{R}^{d}} \left[L_{2^{\alpha(j+1)}}^{\gamma,1}(x,y) + M_{2^{j\alpha}}^{\gamma,0}(x,y)\right] |y|^{\alpha s} \frac{|f(y)|}{|y|^{\alpha s}} dy.\n\end{split} \tag{5.7}
$$

Since $s\alpha < \alpha - \gamma$, we have, using similar Schur tests as above,

$$
\left\| \left(\int_0^\infty t^{-2s} \left| \left(t \Lambda_0 e^{-t \Lambda_0} - t \Lambda_\kappa e^{-t \Lambda_\kappa} \right) f(x) \right|^2 \frac{dt}{t} \right)^{1/2} \right\|_{L^p(\mathbb{R}^d)}
$$
\n
$$
\lesssim \left\| \frac{\left| \Lambda_0^{\frac{1-\gamma}{\alpha}} f(x) \right|}{|x|^{\alpha s + \gamma - 1}} \right\|_{L^p(\mathbb{R}^d)} + \| |x|^{-\alpha s} f \|_{L^p(\mathbb{R}^d)}.
$$
\n(5.8)

This concludes also the proof of part (2) in Theorem [1.4.](#page-5-1)

Remark 5.1. Note that the power $d + \gamma$ in the Schur test for $M_{2^{j\alpha}}^{\gamma,1}$ (see the factor $\left(1 \wedge \frac{N^{-\gamma-d}}{|x-y|^{d+1}}\right)$ $\frac{N^{-\gamma-d}}{|x-y|^{d+\gamma}}$ in the second to last line of [\(5.4\)](#page-33-1)) does not affect the range of admissible s, as long as $\gamma > 0$.

6. Proof of the generalized Hardy inequality (Theorem [1.5\)](#page-6-0)

In this section, we prove the generalized Hardy inequality (Theorem [1.5\)](#page-6-0). To that end, we prove the following Riesz kernel bounds, which we obtain by integrating the heat kernel bounds [\(2.12\)](#page-14-1) against monomials in time.

Lemma 6.1. Let $\alpha \in (1, 2 \wedge (d+2)/2)$, $\beta \in ((d+\alpha)/2, d+\alpha)$, $s \in (0,1]$, and $\kappa = \Psi(\beta)$ be defined by (1.7) . Then,

$$
\Lambda_{\kappa}^{-s}(x,y) \sim |x-y|^{\alpha s - d} \left(1 \wedge \frac{|y|}{|x-y|} \right)^{\beta - d}.
$$
\n
$$
(6.1)
$$

Proof. By the functional calculus, we have

$$
\Lambda_{\kappa}^{-s}(x,y) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{dt}{t} t^s e^{-t\Lambda_{\kappa}}(x,y).
$$

By scaling and the heat kernel bounds [\(2.12\)](#page-14-1),

$$
\Lambda_{\kappa}^{-s}(x,y) \sim |x-y|^{\alpha s-d} \int_0^{\infty} \frac{dt}{t} t^s \frac{t}{(t^{1/\alpha}+1)^{d+\alpha}} \cdot \left(1 \wedge \frac{|y|/|x-y|}{t^{1/\alpha}}\right)^{\beta-d}
$$

In the following, we distinguish between $|x - y| > |y|/2$ and $|x - y| < |y|/2$. Case $|x - y| > |y|/2$. We get

$$
\Lambda_{\kappa}^{-s}(x,y) \sim |x-y|^{\alpha s-d} \left[\int_0^{(|y|/|x-y|)^\alpha} \frac{dt}{t} t^{s+1} + \int_{(|y|/|x-y|)^\alpha}^1 \frac{dt}{t} t^{s+1} \cdot \left(\frac{|y|/|x-y|}{t^{1/\alpha}} \right)^{\beta-d} \right]
$$

+
$$
\int_1^\infty \frac{dt}{t} t^{s+1-1-d/\alpha} \left(\frac{|y|/|x-y|}{t^{1/\alpha}} \right)^{\beta-d} \right]
$$

$$
\sim |x-y|^{\alpha s-d} \left[\left(\frac{|y|}{|x-y|} \right)^{(s+1)\alpha} + \left(\frac{|y|}{|x-y|} \right)^{\beta-d} \right]
$$

$$
\sim |x-y|^{\alpha s-d} \cdot \left(\frac{|y|}{|x-y|} \right)^{\beta-d}.
$$

Case $|x-y| < |y|/2$. We get

$$
\Lambda_{\kappa}^{-s}(x,y) \sim |x-y|^{\alpha s-d} \left[\int_0^1 \frac{dt}{t} t^{s+1} + \int_1^{(|y|/|x-y|)^\alpha} \frac{dt}{t} t^{s+1-1-d/\alpha} + \int_{(|y|/|x-y|)^\alpha}^{\infty} \frac{dt}{t} t^{s+1-1-d/\alpha} \left(\frac{|y|/|x-y|}{t^{1/\alpha}} \right)^{\beta-d} \right]
$$

$$
\sim |x-y|^{\alpha s-d} \left[1 + \left(\frac{|y|}{|x-y|} \right)^{\alpha s-d} \right] \sim |x-y|^{\alpha s-d}.
$$

Combining the above two estimates concludes the proof. \Box

We are now ready to prove the generalized Hardy inequality.

Proof of Theorem [1.5.](#page-6-0) It suffices to show the L^p -boundedness of the operator with integral kernel $|x|^{-\alpha s} \Lambda_{\kappa}^{-s}(x, y)$. To that end, we use weighted Schur tests and distinguish between the following regions.

Case $|x - y| < 4(|x| \wedge |y|)$. In that case, $|x| \sim |y|$ and the integral kernel in question is bounded from above and below by constants times $|x|^{-\alpha s}|x - y|^{\alpha s - d}$. By a Schur test,

$$
\int_{|x-y|<4|x|} |x|^{-\alpha s}|x-y|^{\alpha s-d} dy + \int_{|x-y|<4|y|} |x|^{-\alpha s}|x-y|^{\alpha s-d} dx \lesssim 1.
$$

Case $4|x| < |x - y| < 4|y|$. In that case, $|x - y| \sim |y| \ge |x|$ and the integral kernel in question is bounded from above and below by constants times $|x|^{-\alpha s}|y|^{\alpha s-d}$. A Schur test with weight being a power of $|x|/|y|$ gives

$$
\int_{|x|<|y|} |x|^{-\alpha s}|y|^{\alpha s-d} (|y|/|x|)^{\delta/p'} dx + \int_{|x|<|y|} |x|^{-\alpha s}|y|^{\alpha s-d} (|x|/|y|)^{\delta/p} dy \lesssim 1,
$$

whenever $p\alpha s < \delta < p'(d-\alpha s)$. Such δ exist since $\alpha s - \frac{\alpha s}{p}$ $\frac{\alpha s}{p} < \frac{d - \alpha s}{p}$ $\frac{-\alpha s}{p}$ is true under the assumption $p < d/(\alpha s)$.

Case $4|y| < |x - y| < 4|x|$. In that case, $|x - y| \sim |x| > |y|$ and the integral kernel in question is bounded from above and below by constants times

$$
|x|^{-\alpha s}|y|^{\beta-d}|x-y|^{\alpha s-\beta} \sim |x|^{-\beta}|y|^{\beta-d}.
$$

.

the assumption $p > d/\beta$.

Finally, (1.25) follows from (1.24) since

$$
\||y|^{\alpha(\frac{1-\gamma}{\alpha}-s)}|\Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}}f|\|_{L^{p}(\mathbb{R}^d)}=\||y|^{\alpha(\frac{1-\gamma}{\alpha}-s)}\Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}}f\|_{L^{p}(\mathbb{R}^d)}\lesssim \|\Lambda_{\kappa}^{s-\frac{1-\gamma}{\alpha}}\Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}}f\|_{L^{p}(\mathbb{R}^d)}=\|\Lambda_{\kappa}^sf\|_{L^{p}(\mathbb{R}^d)}.
$$

This concludes the proof. \Box

Appendix A. Proofs of auxiliary statements

A.1. **Proof of Lemma [1.1.](#page-2-1)** For $\alpha = 2$, the claim is obvious. Thus, let $\alpha \in (1, 2)$ and $\beta \in ((d+\alpha)/2, d+\alpha)$. We compute

$$
\frac{d\Psi(\beta)}{d\beta} = \frac{2^{\alpha-1}\Gamma\left(\frac{\beta}{2}\right)\Gamma\left(\frac{1}{2}(d+\alpha-\beta)\right)}{(\alpha-\beta)^2\Gamma\left(\frac{\beta-\alpha}{2}\right)\Gamma\left(\frac{d-\beta}{2}\right)} \times \left[(\beta-\alpha)\left(\psi\left(\frac{\beta}{2}\right)-\psi\left(\frac{\beta-\alpha}{2}\right)+\psi\left(\frac{d-\beta}{2}\right)-\psi\left(\frac{1}{2}(d+\alpha-\beta)\right)\right)-2\right]
$$

with the Digamma function $\psi(z) = \Gamma'(z)/\Gamma(z)$. Thus, the claim would follow from

$$
\psi\left(\frac{d-\beta}{2}\right)-\psi\left(\frac{d-\beta+\alpha}{2}\right)+\psi\left(\frac{\beta}{2}\right)-\psi\left(\frac{\beta-\alpha}{2}\right)<0,\quad \beta\in((d+\alpha)/2,d+\alpha).
$$

To show this inequality, we use [\[DLMF23,](#page-43-23) (5.7.6)], i.e.,

$$
\psi(z) = -\gamma_{\rm E} + \sum_{k \ge 0} \left(\frac{1}{k+1} - \frac{1}{k+z} \right)
$$

with the Euler–Mascheroni constant $\gamma_{\rm E}$. Thus,

$$
\psi\left(\frac{d-\beta}{2}\right) - \psi\left(\frac{d-\beta+\alpha}{2}\right) + \psi\left(\frac{\beta}{2}\right) - \psi\left(\frac{\beta-\alpha}{2}\right)
$$

$$
= \frac{\alpha}{2} \sum_{k\geq 0} \left[\frac{1}{(k+\beta/2)(k+(\beta-\alpha)/2)} - \frac{1}{(k+(d-\beta)/2)(k+(d+\alpha-\beta)/2)} \right]
$$

which is negative for all $\beta > (d + \alpha)/2$. Hence, the strict monotonicity of $\Psi(\beta)$ for $\beta \in ((d + \alpha)/2, d + \alpha)$ follows. $((d + \alpha)/2, d + \alpha)$ follows.

A.2. Proof of Lemma [4.1.](#page-24-2) The upper bound was proved in [\[FM23,](#page-43-4) Lemma 22]. The lower bound for all $N \in (0, 2]$ was proved in [\[BM23,](#page-43-5) Lemma 4.1]. We now give another proof of the lower bound which covers all $N > 0$. We consider two cases $s \in (0, t/2)$ and $s \in (t/2, t)$. Since these two cases are similar, we only give the proof for the first case $s \in (0, t/2)$. In this situation $t - s \sim t$, and hence

$$
\int_{\mathbb{R}^d} dz \, (t-s)^{-d} \left(\frac{(t-s)}{(t-s) + |x-z|} \right)^{d+N} s^{-d} \left(\frac{s}{s + |z-y|} \right)^{d+N}
$$

$$
\sim \int_{\mathbb{R}^d} dz \, t^{-d} \left(\frac{t}{t + |x-z|} \right)^{d+N} s^{-d} \left(\frac{s}{s + |z-y|} \right)^{d+N}.
$$

If $|x-y| < 2t$, then

$$
\int_{\mathbb{R}^d} dz \, t^{-d} \left(\frac{t}{t + |x - z|} \right)^{d+N} s^{-d} \left(\frac{s}{s + |z - y|} \right)^{d+N}
$$
\n
$$
\geq \int_{B(x, 4t)} dz \, t^{-d} s^{-d} \left(\frac{s}{s + |z - y|} \right)^{d+N} \geq \int_{B(y, s)} dz \, t^{-d} s^{-d} \left(\frac{s}{s + |z - y|} \right)^{d+N}
$$
\n
$$
\geq t^{-d} \sim \frac{1}{t^d} \left(\frac{t}{t + |x - y|} \right)^{d+N}.
$$

If $|x - y| \ge 2t$, then we have $|x - y| \sim |x - y|$ for $z \in B(y, |x - y|/2)$. Hence,

$$
\int_{\mathbb{R}^d} dz \, t^{-d} \left(\frac{t}{t + |x - z|} \right)^{d+N} s^{-d} \left(\frac{s}{s + |z - y|} \right)^{d+N}
$$
\n
$$
\geq \int_{B(y, |x - y|/2)} dz \, t^{-d} \left(\frac{t}{t + |x - y|} \right)^{d+N} s^{-d} \left(\frac{s}{s + |z - y|} \right)^{d+N}
$$
\n
$$
\geq t^{-d} \left(\frac{t}{t + |x - y|} \right)^{d+N} \int_{B(y, s)} dz \, s^{-d} \left(\frac{s}{s + |z - y|} \right)^{d+N} \gtrsim \frac{1}{t^d} \left(\frac{t}{t + |x - y|} \right)^{d+N}.
$$

This concludes the proof of Lemma [4.1.](#page-24-2)

APPENDIX B. ON THE NECESSITY OF $\alpha s < \alpha - 1$

In this appendix, we discuss the restriction $\alpha s < \alpha - 1$ for [\(1.15\)](#page-4-2) in our main result, Theorem [1.2,](#page-4-0) and the reversed Hardy inequality [\(1.22\)](#page-5-2) in Theorem [1.4.](#page-5-1) Moreover, we discuss whether one can expect a variant of Proposition [4.6](#page-30-1) where $\Lambda_0^{\frac{1-\gamma}{\alpha}}$ is replaced with $\Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}}$.

B.1. On the restriction on s in [\(1.15\)](#page-4-2). Let us discuss the restriction $\alpha s < \alpha - 1$ in [\(1.15\)](#page-4-2). To prove

$$
\|\Lambda_0^s \Lambda_\kappa^{-s} f\|_{L^p(\mathbb{R}^d)} \lesssim \|f\|_{L^p(\mathbb{R}^d)}
$$

it suffices, by the identity

$$
\Lambda_{\kappa}^{-s} = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{dt}{t} t^s e^{-t\Lambda_{\kappa}},
$$

to estimate the kernel of $\Lambda_0^s e^{-t\Lambda_\kappa}$. By Duhamel's formula

$$
e^{-t\Lambda_{\kappa}}(x,y) = e^{-t\Lambda_0}(x,y) - \kappa \int_0^t ds \int_{\mathbb{R}^d} dz \, e^{-(t-s)\Lambda_0}(x,z) |z|^{-\alpha} z \cdot \nabla_z e^{-s\Lambda_{\kappa}}(z,y),
$$

and integration by parts (since gradient bounds for $e^{-t\Lambda_{\kappa}}$ are unavailable at the time of this writing and likely difficult to obtain), this requires to treat, among others,

$$
\int_0^t \int_{\mathbb{R}^d} \nabla_z (\Lambda_0^s e^{-(t-\tau)\Lambda_0}(x,z)) \frac{z}{|z|^\alpha} e^{-\tau \Lambda_\kappa}(z,y) dz d\tau.
$$
 (B.1)

By scaling,

$$
|\nabla \Lambda_0^s e^{-(t-\tau)\Lambda_0}(x,z)| \lesssim (t-\tau)^{-\frac{1}{\alpha}-s} \frac{1}{(t-\tau)^{d/\alpha}} \left(\frac{(t-\tau)^{1/\alpha}}{(t-\tau)^{1/\alpha}+|x-z|}\right)^{d+\gamma}
$$
(B.2)

for some $\gamma > 0$. (One can deduce this bound, e.g., using [\(3.8\)](#page-19-2) and Lemma [2.10.](#page-17-1)) By scaling, the time-integral in [\(B.1\)](#page-37-2) converges at $\tau = t$, if and only if $\alpha s < \alpha - 1$.

The above argument indicates that [\(1.15\)](#page-4-2) may only hold for $\alpha s < \alpha - 1$. However, if the following gradient bound

$$
|\nabla_x e^{-t\Lambda_\kappa}(x,y)| \lesssim t^{-\frac{d+1}{\alpha}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha} + |x-y|}\right)^{d+\gamma} \left(1 \wedge \frac{|y|}{t^{1/\alpha}}\right)^{\beta - d} \tag{B.3}
$$

held for some $\gamma > 0$, then we may circumvent the restriction $\alpha s < \alpha - 1$. Indeed, given [\(B.3\)](#page-38-1), we would not need to integrate by parts in the above Duhamel formula. In particular, in that case, the factor $(t - \tau)^{-1/\alpha}$ appearing in [\(B.2\)](#page-37-3) would be replaced by a factor $s^{-1/\alpha}$, which, in turn, would make the τ -integration converge for $s < 1$. In view of scaling and Lemma [2.10](#page-17-1) one may wonder if [\(B.3\)](#page-38-1) holds.

Let us also remark that if the gradient perturbation $|z|^{-\alpha}z \cdot \nabla$ was replaced with a scalar perturbation, we would have to consider

$$
\Lambda_0^s e^{-(t-\tau)\Lambda_0}(x,z) \lesssim (t-\tau)^{-s} \frac{1}{(t-\tau)^{d/\alpha}} \left(\frac{(t-\tau)^{1/\alpha}}{(t-\tau)^{1/\alpha}+|x-z|}\right)^{d+\gamma}
$$
(B.4)

instead of $(B.2)$. In this case, the remaining time-integral would converge whenever $s < 1$.

B.2. On the restriction on s in (1.22) . Here, we argue that one can prove (1.22) in Theorem [1.4](#page-5-1) using only pointwise bounds for the kernel $Q_t(x, y)$ (defined in [\(4.1\)](#page-24-3)) only if $\alpha s < \alpha - 1$. This restriction arose when performing the Schur tests [\(5.3\)](#page-32-1)–[\(5.4\)](#page-33-1) involving the function $M_t^{\gamma,1}(x,y)$, defined in [\(4.5\)](#page-24-4). As we noted in Remark [5.1,](#page-34-2) the reason why the Schur test involving $M_t^{\gamma,1}$ is positive only for $\alpha s < \alpha - 1$ is due to the exponent $1 - \alpha$ of the factor $\int |x|\vee|y|$ $\left[\frac{x|y|y|}{t^{1/\alpha}}\right]^{1-\alpha}$, not because of the decay of the term depending on $|x-y|$ in $M_t^{\gamma,1}(x,y)$. The proof of Proposition [4.5](#page-28-1) reveals that the function $M_t^{\gamma,1}$ arose when estimating the right-hand side of [\(4.13\)](#page-29-0) on the set $\{(x, y) \in \mathbb{R}^{2d} : |y| \ge t^{1/\alpha}, |x - y| \le (|x| \wedge |y|)/2 \}$. These estimates were carried out in Lemma [4.2](#page-25-2) and were based on Duhamel's formula. In particular, the factor $\int |x|\vee|y|$ $\left[\frac{c|v|y|}{t^{1/\alpha}}\right]^{1-\alpha}$ only appears in the integral over $|z| \in [(1-\varepsilon_1)|y|, (1+\varepsilon_2)|y|]$ for arbitrary but fixed $\varepsilon_1, \varepsilon_2 > 0$; see also Remark [4.3.](#page-26-1)

In the following, we argue that the Duhamel integrals in [\(4.13\)](#page-29-0) are bounded from below by a constant times

$$
\left(\frac{|y|}{t^{1/\alpha}}\right)^{1-\alpha} t^{-d/\alpha} \left(\frac{t^{1/\alpha}}{t^{1/\alpha}+|x-y|}\right)^{d+\gamma}
$$

for some $\gamma > 0$ and $x, y \in \mathbb{R}^d$ such that $|y| > t^{1/\alpha}$ and $|x - y| < (|x| \wedge |y|)/2$. A strong argument in that favor would be the lower bound

$$
-\int_0^t ds \int_{|z| \in [(1-\varepsilon_1)|y|, (1+\varepsilon_2)|y|]} dz (t-s)^{-d/\alpha} \left(\frac{(t-s)^{1/\alpha}}{(t-s)^{1/\alpha}+|x-z|}\right)^{d+\gamma_1} |z|^{-\alpha}
$$

$$
\times z \cdot \frac{(z-y)/|z-y|}{s^{1/\alpha}} s^{-d/\alpha} \left(\frac{s^{1/\alpha}}{s^{1/\alpha}+|z-y|}\right)^{d+\gamma_2}
$$

\n
$$
\geq \left(\frac{|y|}{t^{1/\alpha}}\right)^{1-\alpha} t^{-d/\alpha} \left(\frac{t^{1/\alpha}}{t^{1/\alpha}+|x-y|}\right)^{d+\gamma}
$$
\n(B.5)

for some $\gamma_1, \gamma_2 > 0$, arbitrary but fixed $\varepsilon_1, \varepsilon_2 > 0$, and some $\gamma > 0$ and $x, y \in \mathbb{R}^d$ such that $|y| > t^{1/\alpha}$ and $|x-y| \leq (|x| \wedge |y|)/2$. Let us motivate the left-hand side of [\(B.5\)](#page-38-2). The term

$$
(t-s)^{-d/\alpha} \left(\frac{(t-s)^{1/\alpha}}{(t-s)^{1/\alpha} + |x-z|} \right)^{d+\gamma_1}
$$

comes from the bounds for $e^{-t\Lambda_{\kappa}}(x, z)$ and its time-derivative (see [\(2.12\)](#page-14-1) and Proposition [2.9\)](#page-17-0) when $|y| > t^{1/\alpha}$, i.e., when the singular weight $(1 \wedge |y|/t^{1/\alpha})^{\beta-d} \sim 1$. The term

$$
-\frac{(z-y)/|z-y|}{s^{1/\alpha}}s^{-d/\alpha}\left(\frac{s^{1/\alpha}}{s^{1/\alpha}+|z-y|}\right)^{d+\gamma_2}
$$

comes from computing the spatial, and possibly additional temporal, derivatives of the righthand side of the bounds for $e^{-t\Lambda_0}$ in [\(2.13\)](#page-14-4); see Propositions [2.10](#page-17-1) and [2.11.](#page-17-4) Here, the prefactor $s^{-1/\alpha}(z-y)/|z-y|$ comes from the fact that $t^{d/\alpha}e^{-t\Lambda_0}(x,y)$ is a function depending only on $|x-y|/t^{1/\alpha}.$

In the following, we establish the lower bound [\(B.5\)](#page-38-2) in the technically simpler (but artificial) one-dimensional case.

Proposition B.1. Let $d = 1$, $\alpha \in (0, 2)$, $\gamma_1, \gamma_2 > 0$, and $t > 0$. Then there are $\gamma > 0$, $y > t^{1/\alpha}$, and $x > 0$ such that $|x - y| \le (x \wedge y)/2$ and [\(B.5\)](#page-38-2) hold.

Proof. By a scaling argument, it suffices to consider $t = 1$. Let $\varepsilon_1 < 1/6$ and $x = (1-2\varepsilon_1)y < y$. Then, $|x - y| = 2\varepsilon_1 y \le (1/2 - \varepsilon_1)y = x/2$. Thus, it suffices to accomplish the following two tasks for some $0 < \varepsilon_2 < 1$.

(a) Show

$$
\int_0^1 ds \int_y^{(1+\varepsilon_2)y} dz \, (1-s)^{-\frac{d}{\alpha}} \left(\frac{(1-s)^{1/\alpha}}{(1-s)^{1/\alpha} + |x-z|} \right)^{d+\gamma_1} z^{1-\alpha} s^{-\frac{d+1}{\alpha}} \left(\frac{s^{1/\alpha}}{s^{1/\alpha} + |z-y|} \right)^{d+\gamma_2}
$$

$$
\leq \int_0^1 ds \int_{(1-\varepsilon_1)y}^y dz \, (1-s)^{-\frac{d}{\alpha}} \left(\frac{(1-s)^{1/\alpha}}{(1-s)^{1/\alpha} + |x-z|} \right)^{d+\gamma_1} z^{1-\alpha} s^{-\frac{d+1}{\alpha}} \left(\frac{s^{1/\alpha}}{s^{1/\alpha} + |z-y|} \right)^{d+\gamma_2} . \tag{B.6}
$$

(b) Show

$$
\int_{0}^{1} ds \int_{(1-\varepsilon_{1})y}^{y} dz (1-s)^{-\frac{d}{\alpha}} \left(\frac{(1-s)^{1/\alpha}}{(1-s)^{1/\alpha} + |x-z|} \right)^{d+\gamma_{1}} z^{1-\alpha} s^{-\frac{d+1}{\alpha}} \left(\frac{s^{1/\alpha}}{s^{1/\alpha} + |z-y|} \right)^{d+\gamma_{2}}
$$

$$
\gtrsim y^{1-\alpha} \left(\frac{1}{1+|x-y|} \right)^{d+\gamma} . \tag{B.7}
$$

We now accomplish these two tasks:

(a) We show that [\(B.6\)](#page-39-0) even holds pointwise for all $s \in [0,1]$. To see this, we use that

$$
z^{1-\alpha} \left(\frac{(1-s)^{1/\alpha}}{(1-s)^{1/\alpha} + |x-z|} \right)^{d+\gamma_1} \Big|_{z \in [y,(1+\varepsilon_2)y]} \n\leq y^{1-\alpha} \left(\frac{(1-s)^{1/\alpha}}{(1-s)^{1/\alpha} + y - x} \right)^{d+\gamma_1} \n\leq z^{1-\alpha} \left(\frac{(1-s)^{1/\alpha}}{(1-s)^{1/\alpha} + |x-z|} \right)^{d+\gamma_1} \Big|_{z \in [(1-\varepsilon_1)y,y]},
$$

where we used $|x-z| = z - x$ when $z \in [(1-\varepsilon_1)y, y]$ since $x = (1-2\varepsilon_1)y$. Thus, it suffices to show

$$
\int_{y}^{(1+\varepsilon_{2})y} dz \left(\frac{s^{1/\alpha}}{s^{1/\alpha}+z-y}\right)^{d+\gamma_{2}} \le \int_{(1-\varepsilon_{1})y}^{y} dz \left(\frac{s^{1/\alpha}}{s^{1/\alpha}+y-z}\right)^{d+\gamma_{2}}.\tag{B.8}
$$

But this just follows from shifting $z \mapsto z+y$ in both integrals and replacing $z \mapsto -z$ in one of the integrals. In fact, we could get a strict inequality by taking $\varepsilon_2 < \varepsilon_1$. This concludes the first task.

(b) We now show [\(B.7\)](#page-39-1). As $z \sim y$, it suffices to show

$$
y^{1-\alpha} \int_0^1 ds \int_{(1-\varepsilon_1)y}^{y} dz \left(1-s\right)^{-\frac{d}{\alpha}} \left(\frac{(1-s)^{1/\alpha}}{(1-s)^{1/\alpha}+|x-z|}\right)^{d+\gamma_1} s^{-\frac{d+1}{\alpha}} \left(\frac{s^{1/\alpha}}{s^{1/\alpha}+|z-y|}\right)^{d+\gamma_2} \approx y^{1-\alpha} \left(\frac{1}{1+|x-y|}\right)^{d+\gamma}.
$$
\n(B.9)

Let

$$
F(s, x, y, z) := y^{1-\alpha}(1-s)^{-\frac{d}{\alpha}} \left(\frac{(1-s)^{1/\alpha}}{(1-s)^{1/\alpha} + |x-z|} \right)^{d+\gamma_1} s^{-\frac{d+1}{\alpha}} \left(\frac{s^{1/\alpha}}{s^{1/\alpha} + |z-y|} \right)^{d+\gamma_2}
$$

By part (s) we bound

By part (a), we bound

$$
\int_0^1 ds \int_{(1-\varepsilon_1)y}^y dz F(s, x, y, z) \gtrsim \int_0^1 ds \int_{(1-\varepsilon_1)y}^{(1+\varepsilon_1)y} dz F(s, x, y, z)
$$
\n
$$
= \int_0^1 ds \int_{\mathbb{R}} dz F(s, x, y, z) - \int_0^1 ds \int_{\mathbb{R}\setminus[(1-\varepsilon_1)y, (1+\varepsilon_1)y]} dz F(s, x, y, z).
$$
\n(B.10)

Since

$$
\int_0^1 ds \int_{\mathbb{R}\setminus[(1-\varepsilon_1)y,(1+\varepsilon_1)y]} dz F(s,x,y,z) \lesssim y^{-\alpha} \left(\frac{1}{1+|x-y|}\right)^{d+(\gamma_1\wedge\gamma_2)} \tag{B.11}
$$

by Remark [4.3](#page-26-1) and $y^{-\alpha} \lesssim y^{1-\alpha}$ for $y > 1$, it suffices to estimates the first summand on the right-hand side of [\(B.10\)](#page-40-1). By [\(4.6\)](#page-24-5), we have, for $\gamma = \gamma_1 \vee \gamma_2$,

$$
y^{1-\alpha} \int_0^1 ds \int_{\mathbb{R}^d} dz \, (1-s)^{-\frac{d}{\alpha}} \left(\frac{(1-s)^{1/\alpha}}{(1-s)^{1/\alpha} + |x-z|} \right)^{d+\gamma_1} s^{-\frac{d+1}{\alpha}} \left(\frac{s^{1/\alpha}}{s^{1/\alpha} + |z-y|} \right)^{d+\gamma_2}
$$

\n
$$
\gtrsim y^{1-\alpha} \int_0^1 ds \int_{\mathbb{R}^d} dz \, (1-s)^{-\frac{d}{\alpha}} \left(\frac{(1-s)^{1/\alpha}}{(1-s)^{1/\alpha} + |x-z|} \right)^{d+\gamma} s^{-\frac{d}{\alpha}} \left(\frac{s^{1/\alpha}}{s^{1/\alpha} + |z-y|} \right)^{d+\gamma} \quad (B.12)
$$

\n
$$
\gtrsim y^{1-\alpha} \left(\frac{1}{1+|x-y|} \right)^{d+\gamma},
$$

as desired.

This concludes the proof. \Box

B.3. On a variant of Proposition [4.6](#page-30-1) involving powers of Λ_{κ} . We now discuss the possibility to prove a variant of the second version of the reversed Hardy inequality in Propo-sition [4.6,](#page-30-1) where the terms involving $\Lambda_0^{\frac{1-\gamma}{\alpha}}f$ are replaced with $\Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}}$. The idea to obtain an estimate involving $\Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}}f$ is to insert $1=\Lambda_{\kappa}^{-\frac{1-\gamma}{\alpha}}\Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}}$ in Duhamel's formula. To that end, we use Duhamel's formula in the form

$$
p_t(x, y) - \tilde{p}_t(x, y) = \kappa \int_0^t \int_{\mathbb{R}^d} \tilde{p}_{t-s}(x, z) |z|^{-\alpha} z \cdot \nabla_z p_s(z, y) dz ds
$$

=
$$
-\kappa \int_0^t \int_{\mathbb{R}^d} \nabla_z \left[\tilde{p}_{t-s}(x, z) |z|^{-\alpha} z \right] p_s(z, y) dz ds.
$$

Here, we integrated by parts to shift the z-derivative from $p_s(z, y)$ to $\tilde{p}_{t-s}(x, z)$ as derivative bounds for $e^{-t\Lambda_{\kappa}}$ are unavailable at the time of this writing and likely difficult to obtain. In

.

this situation, the heat kernel $e^{-t\Lambda_{\kappa}}$ can now act on a function f and we can write $e^{-t\Lambda_{\kappa}}f =$ $\Lambda_{\kappa}^{-\frac{1-\gamma}{\alpha}}e^{-t\Lambda_{\kappa}}\Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}}f$. By the product rule,

$$
\tilde{p}_t(x,y) - p_t(x,y) = \kappa \int_0^t ds \int_{\mathbb{R}^d} dz \left[|z|^{-\alpha} z \cdot \nabla_z \tilde{p}_{t-s}(x,z) + \frac{d-\alpha}{|z|^{\alpha}} \tilde{p}_{t-s}(x,z) \right] p_s(z,y).
$$

The second term has a $|z|^{-\alpha}$ -decay and can be treated as in [\[FMS21\]](#page-43-3), thereby giving rise to terms involving $M_t^{\gamma,0}$ and L_t^{γ} \tilde{t} . To handle the first term, we proceed similarly as in the proof of Proposition [4.6.](#page-30-1) The only difference is that in the analog of $Q_{1,1}^1$ (see [\(4.16\)](#page-31-0)), we use $e^{-\Lambda_{\kappa}} = \Lambda_{\kappa}^{-\frac{1-\gamma}{\alpha}} e^{-\Lambda_{\kappa}} \Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}}$ to get the estimate in terms of $|\Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}}f|$. We obtain, with $S_{x,1} = \{z \in$ \mathbb{R}^d : $|x|/16 < |z| < 4|x|$,

$$
\tilde{Q}_{1,1}^{1}f(x) := \kappa \int_{\mathbb{R}^{d}} \int_{S_{x,1}} |z|^{-\alpha} z \cdot (\nabla_{z} \tilde{p}_{1/2}(x, z)) \Lambda_{\kappa}^{-\frac{1-\gamma}{\alpha}} e^{-\frac{1}{2}\Lambda_{\kappa}}(z, y) \Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}} f(y) dz dy \n+ \kappa \int_{\mathbb{R}^{d}} \int_{0}^{1/2} \int_{S_{x,1}} |z|^{-\alpha} z \cdot (\nabla_{z} \tilde{p}_{1-s,1}(x, z)) \Lambda_{\kappa}^{-\frac{1-\gamma}{\alpha}} e^{-s\Lambda_{\kappa}}(z, y) \Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}} f(y) dz ds dy \n+ \kappa \int_{\mathbb{R}^{d}} \int_{1/2}^{1} \int_{S_{x,1}} |z|^{-\alpha} z \cdot (\nabla_{z} \tilde{p}_{1-s}(x, z)) \Lambda_{\kappa}^{-\frac{1-\gamma}{\alpha}+1} e^{-s\Lambda_{\kappa}}(z, y) \Lambda_{\kappa}^{\frac{1-\gamma}{\alpha}} f(y) dz ds dy.
$$
\n(B.13)

We now argue that the right-hand side is not expected to give rise to L_1^{γ} \int_{1}^{γ} or $M_{1}^{\gamma,\gamma}$. Let us consider, e.g., the second summand on the right-hand side of [\(B.13\)](#page-41-0), whose integral kernel is, in view of Lemma [2.10](#page-17-1) and [\(2.22\)](#page-17-3), bounded by a multiple of $|x|^{1-\alpha}$ times

$$
t^{-\frac{d+1}{\alpha}} s^{-\frac{d-\gamma\alpha}{\alpha}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha}+|x-z|}\right)^{d+1+\alpha} \cdot \left(\frac{s^{1/\alpha}}{s^{1/\alpha}+|z-y|}\right)^{d-1+\gamma} + t^{-\frac{d+1}{\alpha}} s^{-\frac{d-\gamma\alpha}{\alpha}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha}+|x-z|}\right)^{d+1+\alpha} \cdot \left(\frac{s^{1/\alpha}}{s^{1/\alpha}+|z-y|}\right)^{\beta-1+\gamma} \left(\frac{|y|}{s^{1/\alpha}}\right)^{\beta-d}.
$$
\n(B.14)

However, since γ < 1 and β < d, we can now not argue as in the proof of Proposition [4.6](#page-30-1) anymore, where we used Lemma [4.1.](#page-24-2) In particular, the z-integration is not expected to yield a function $F(x, y)$ satisfying $F(x, \cdot) \in L^1(\mathbb{R}^d)$ and $F(\cdot, y) \in L^1(\mathbb{R}^d)$ for all $x, y \in \mathbb{R}^d$, thereby leaving us behind with an integral kernel for which we cannot apply Schur tests.

On the other hand, if we assumed that [\(B.3\)](#page-38-1) holds, then $\nabla_x \Lambda_{\kappa}^{-\frac{1-\gamma}{\alpha}} e^{-t\Lambda_{\kappa}}(x, y)$ obeys a bound similar to those in Lemma [2.12,](#page-17-6) namely

$$
|\nabla_x \Lambda_\kappa^{-\frac{1-\gamma}{\alpha}} e^{-\Lambda_\kappa}(x, y)| \lesssim \left(\frac{1}{1+|x-y|}\right)^{d+\gamma} + \left(\frac{1}{1+|x-y|}\right)^{\beta+\gamma} |y|^{\beta-d}.\tag{B.15}
$$

In turn, this bound would allow us to proceed as in the proof of Proposition [4.6](#page-30-1) and show, for all $\gamma \in (0, 1), \beta \in (d - \gamma, d), t > 0$, and $x \in \mathbb{R}^d$,

$$
|(Q_t f)(x)| \lesssim \int_{\mathbb{R}^d} \left[L_t^{\gamma,1}(x, y) + M_t^{\gamma,0}(x, y) \right] |f(y)| dy
$$

+
$$
\int_{\mathbb{R}^d} L_t^{\gamma, \gamma}(x, y) |(\Lambda_\kappa^{\frac{1-\gamma}{\alpha}} f)(y)| dy
$$

+
$$
\int_{\mathbb{R}^d} M_t^{\gamma, \gamma}(x, y) |y|^{1-\gamma} (\Lambda_\kappa^{\frac{1-\gamma}{\alpha}} f)(y)| dy.
$$
 (B.16)

Consequently, by the proof of Theorem [1.4,](#page-5-1) we would obtain, for all $\gamma \in (0,1)$ such that $1 - \gamma \le \alpha s < \alpha - \gamma$, and $\beta \in (d - \gamma, d)$,

$$
\left\| \left(\int_0^\infty t^{-2s} \left| \left(t\Lambda_\kappa e^{-t\Lambda_\kappa} - t\Lambda_0 e^{-t\Lambda_0} \right) f \right|^2 \frac{dt}{t} \right)^{1/2} \right\|_{L^p(\mathbb{R}^d)} \lesssim \left\| \frac{\left| \Lambda_\kappa^{\frac{1-\gamma}{\alpha}} f(x) \right|}{|x|^{\alpha s + \gamma - 1}} \right\|_{L^p(\mathbb{R}^d)} + \left\| \frac{f}{|x|^{\alpha s}} \right\|_{L^p(\mathbb{R}^d)}.
$$
\n(B.17)

Appendix C. Conditional generalized Hardy inequality for gradient **PERTURBATIONS**

Our current generalized Hardy inequality in Theorem [1.5](#page-6-0) gives an upper bound of the scalar Hardy potential in terms of Λ_{κ} . Here, we prove a generalized Hardy inequality for the gradient perturbation in terms of Λ_{κ} under the assumption that suitable bounds for $\nabla e^{-t\Lambda_{\kappa}}$ are available.

Proposition C.1. Let $d \in \{3, 4, ...\}$, $\alpha \in (1, 2)$, $\beta \in ((d+\alpha)/2, d+\alpha)$, and $\kappa = \Psi(\beta)$ be defined by [\(1.7\)](#page-2-2). If [\(B.3\)](#page-38-1) holds true for some $\gamma > (2\alpha - d) \vee 0$, then, for any $p \in (1 \vee d/\beta, d/(\alpha - 1))$,

> $|||x|^{-\alpha}x \cdot \nabla f||_{L^p(\mathbb{R}^d)} \lesssim ||\Lambda_{\kappa}f||_{L^p(\mathbb{R}^d)}$ $(C.1)$

Note that the range of allowed p in Proposition [C.1](#page-42-4) would be slightly larger than that in Theorem [1.5.](#page-6-0)

Proof. We argue as in the proof of Theorem [1.5](#page-6-0) using Schur tests and bounds for the integral kernel of

$$
||x|^{-\alpha}x \cdot \nabla_x \Lambda_\kappa^{-1}(x, y)| = \left||x|^{-\alpha}x \int_0^\infty \nabla_x e^{-t\Lambda_\kappa}(x, y) dt\right|
$$

$$
\lesssim |x|^{1-\alpha} \int_0^\infty dt \, t^{-\frac{d+1}{\alpha}} \left(\frac{t^{1/\alpha}}{t^{1/\alpha} + |x-y|}\right)^{d+\gamma} \left(1 + \frac{|y|}{t^{1/\alpha}}\right)^{\beta-d}.
$$

Arguing as in the proof of Lemma [6.1,](#page-34-1) we get, using $\gamma > 2\alpha - d$,

$$
\left| |x|^{-\alpha} x \cdot \nabla_x \Lambda_\kappa^{-1}(x, y) \right| \lesssim |x|^{1-\alpha} |x-y|^{\alpha-1-d} \left(1 \wedge \frac{|y|}{|x-y|} \right)^{\beta-d}.
$$

Thus, it remains to perform Schur tests similar to those in Theorem [1.5.](#page-6-0) The conclusion follows by noting that the only difference between the relevant kernel here and in that proof is that α and s in the proof of Theorem [1.5](#page-6-0) have to replaced with $\alpha - 1$ and 1, respectively. \square

Acknowledgements. We are deeply grateful to Damir Kinzebulatov and Karol Szczypkowski for helpful discussions. T. A. Bui and X. T. Duong were supported by the research grant ARC DP220100285 from the Australian Research Council.

REFERENCES

- [Aus07] Pascal Auscher. On necessary and sufficient conditions for L^p -estimates of Riesz transforms associated to elliptic operators on \mathbb{R}^n and related estimates. Mem. Amer. Math. Soc., 186(871):xviii+75, 2007.
- [BBC03] Krzysztof Bogdan, Krzysztof Burdzy, and Zhen-Qing Chen. Censored stable processes. Probab. Theory Related Fields, 127(1):89–152, 2003.
- [BD23] The Anh Bui and Piero D'Ancona. Generalized Hardy operators. Nonlinearity, 36(1):171–198, 2023.
- [BG60] R. M. Blumenthal and R. K. Getoor. Some theorems on stable processes. Trans. Amer. Math. Soc., 95:263–273, 1960.
- [BJ07] Krzysztof Bogdan and Tomasz Jakubowski. Estimates of heat kernel of fractional Laplacian perturbed by gradient operators. Comm. Math. Phys., 271(1):179–198, 2007.
- [BJLPP22] Krzysztof Bogdan, Tomasz Jakubowski, Julia Lenczewska, and Katarzyna Pietruska-Pałuba. Optimal Hardy inequality for the fractional Laplacian on L^p . J. Funct. Anal., 282(8): Paper No. 109395, 31, 2022.
- [BM23] The Anh Bui and Konstantin Merz. Equivalence of Sobolev norms in Lebesgue spaces for Hardy operators in a half-space. arXiv e-prints, page [arXiv:2309.02928,](http://arxiv.org/abs/2309.02928) September 2023.
- [CT04] Rama Cont and Peter Tankov. Financial Modelling with Jump Processes. Chapman & Hall/CRC Financial Mathematics Series. Chapman & Hall/CRC, Boca Raton, FL, 2004.
- [Dav90] E. B. Davies. Heat Kernels and Spectral Theory, volume 92 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1990.
- [DLMF23] NIST Digital Library of Mathematical Functions. <https://dlmf.nist.gov/>, Release 1.1.9 of 2023- 03-15, 2023. F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. McClain, eds.
- [DR96] Xuan T. Duong and Derek W. Robinson. Semigroup kernels, Poisson bounds, and holomorphic functional calculus. J. Funct. Anal., 142(1):89–128, 1996.
- [FL19] Patrick J. Fitzsimmons and Liping Li. On the Dirichlet form of three-dimensional Brownian motion conditioned to hit the origin. Sci. China Math., 62(8):1477–1492, 2019.
- [FLS08] Rupert L. Frank, Elliott H. Lieb, and Robert Seiringer. Hardy-Lieb-Thirring inequalities for fractional Schrödinger operators. J. Amer. Math. Soc., $21(4):925-950$, 2008 .
- [FM23] Rupert L. Frank and Konstantin Merz. On Sobolev norms involving Hardy operators in a halfspace. J. Funct. Anal., 285(10):Paper No. 110104, 2023.
- [FMS21] Rupert L. Frank, Konstantin Merz, and Heinz Siedentop. Equivalence of Sobolev norms involving generalized Hardy operators. International Mathematics Research Notices, 2021(3):2284–2303, February 2021.
- [FMS23a] Rupert L. Frank, Konstantin Merz, and Heinz Siedentop. The Scott conjecture for large Coulomb systems: a review. Lett. Math. Phys., 113(1):Paper No. 11, 79, 2023.
- [FMS23b] Rupert L. Frank, Konstantin Merz, and Heinz Siedentop. Relativistic strong Scott conjecture: a short proof. In Density functionals for many-particle systems—mathematical theory and physical applications of effective equations, volume 41 of Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap., pages 69–79. World Sci. Publ., Hackensack, NJ, March [2023] ©2023.
- [FMSS20] Rupert L. Frank, Konstantin Merz, Heinz Siedentop, and Barry Simon. Proof of the strong Scott conjecture for Chandrasekhar atoms. Pure Appl. Funct. Anal., 5(6):1319–1356, December 2020.
- [FS08] Rupert L. Frank and Robert Seiringer. Non-linear ground state representations and sharp Hardy inequalities. J. Funct. Anal., 255(12):3407–3430, 2008.
- [GL90] Nicola Garofalo and Ermanno Lanconelli. Level sets of the fundamental solution and Harnack inequality for degenerate equations of Kolmogorov type. Trans. Amer. Math. Soc., 321(2):775– 792, 1990.
- [Har19] G. H. Hardy. Notes on some points in the integral calculus LI: On Hilbert's double-series theorem, and some connected theorems concerning the convergence of infinite series and integrals. Messenger of Mathematics, 48:107–112, 1919.
- [Har20] G. H. Hardy. Note on a theorem of Hilbert. Mathematische Zeitschrift, 6(3–4):314–317, 1920.
- [Her77] Ira W. Herbst. Spectral theory of the operator $(p^2 + m^2)^{1/2} - Ze^2/r$. Comm. Math. Phys., 53:285– 294, 1977.
- [Kat66] Tosio Kato. Perturbation Theory for Linear Operators, volume 132 of Grundlehren der mathematischen Wissenschaften. Springer-Verlag, Berlin, 1 edition, 1966.
- [KMS24] Damir Kinzebulatov, Kodjo Raphaël Madou, and Yuliy A. Semënov. On the supercritical fractional diffusion equation with Hardy-type drift. J. Anal. Math., 152(2):401–420, 2024.
- [KMV+17] Rowan Killip, Changxing Miao, Monica Visan, Junyong Zhang, and Jiqiang Zheng. The energycritical NLS with inverse-square potential. Discrete Contin. Dyn. Syst., 37(7):3831–3866, 2017.
- [KMV⁺18] R. Killip, C. Miao, M. Visan, J. Zhang, and J. Zheng. Sobolev spaces adapted to the Schrödinger operator with inverse-square potential. Math. Z., 288(3-4):1273–1298, 2018.
- [KMVZ17] Rowan Killip, Jason Murphy, Monica Visan, and Jiqiang Zheng. The focusing cubic NLS with inverse-square potential in three space dimensions. Differential Integral Equations, 30(3-4):161– 206, 2017.
- [KPS81] V. F. Kovalenko, M. A. Perelmuter, and Ya. A. Semenov. Schrödinger operators with $L_w^{l/2}(\mathbb{R}^l)$ potentials. J. Math. Phys., 22:1033–1044, 1981.
- [KRS08] R. Klages, G. Radons, and I.M. Sokolov. Anomalous Transport: Foundations and Applications. Wiley, 2008.
- [KS06] Panki Kim and Renming Song. Two-sided estimates on the density of Brownian motion with singular drift. Illinois J. Math., 50(1-4):635–688, 2006.
- [KS20] Damir Kinzebulatov and Yuliy A. Semënov. On the theory of the Kolmogorov operator in the spaces L^p and C_{∞} . Ann. Sc. Norm. Super. Pisa Cl. Sci. (5), 21:1573-1647, 2020.
- [KS23] Damir Kinzebulatov and Yuliy A. Semënov. Fractional Kolmogorov operator and desingularizing weights. Publ. Res. Inst. Math. Sci., 59(2):339–391, 2023.
- [KSS21] D. Kinzebulatov, Yu. A. Semënov, and K. Szczypkowski. Heat kernel of fractional Laplacian with Hardy drift via desingularizing weights. J. Lond. Math. Soc. (2), 104(4):1861–1900, 2021.
- [Lie81] Elliott H. Lieb. Thomas-fermi and related theories of atoms and molecules. Rev. Mod. Phys., 53(4):603–641, October 1981.
- [McI86] Alan McIntosh. Operators which have an H_{∞} functional calculus. In *Miniconference on Operator* Theory and Partial Differential Equations (North Ryde, 1986), volume 14 of Proc. Centre Math. Anal. Austral. Nat. Univ., pages 210–231. Austral. Nat. Univ., Canberra, 1986.
- [Mer21] Konstantin Merz. On scales of Sobolev spaces associated to generalized Hardy operators. Math. Z., 299(1):101–121, 2021.
- [Mer22] Konstantin Merz. On complex-time heat kernels of fractional Schrödinger operators via Phragmén-Lindelöf principle. J. Evol. Equ., $22(3)$: Paper No. 62, 30, 2022.
- [MK00] Ralf Metzler and Joseph Klafter. The random walk's guide to anomalous diffusion: a fractional dynamics approach. Physics Reports, 339(1):1–77, 2000.
- [MNS18] G. Metafune, L. Negro, and C. Spina. Sharp kernel estimates for elliptic operators with secondorder discontinuous coefficients. J. Evol. Equ., 18(2):467–514, 2018.
- [MS22] Konstantin Merz and Heinz Siedentop. Proof of the strong Scott conjecture for heavy atoms: the Furry picture. Annales Henri Lebesgue, 5:611–642, 2022.
- [MSS17] G. Metafune, M. Sobajima, and C. Spina. Kernel estimates for elliptic operators with second-order discontinuous coefficients. J. Evol. Equ., 17(1):485–522, 2017.
- [Sim84] B. Simon. Fifteen problems in mathematical physics. In Perspectives in Mathematics. Birkhäuser, 1984.
- [Sti19] Pablo Ra´ul Stinga. User's guide to the fractional Laplacian and the method of semigroups. In Handbook of Fractional Calculus with Applications. Vol. 2, pages 235–265. De Gruyter, Berlin, 2019.
- [Yaf99] D. Yafaev. Sharp constants in the Hardy-Rellich inequalities. Journ. Functional Analysis, 168(1):121–144, October 1999.
- [Zha95] Qi Zhang. A Harnack inequality for Kolmogorov equations. J. Math. Anal. Appl., 190(2):402–418, 1995.
- [Zha97] Qi S. Zhang. Gaussian bounds for the fundamental solutions of $\nabla(A\nabla u) + B\nabla u - u_t = 0$. Manuscripta Math., 93(3):381–390, 1997.

(The Anh Bui) School of Mathematical and Physical Sciences, Macquarie University, NSW 2109, Australia

Email address: the.bui@mq.edu.au

(Xuan Thinh Duong) School of Mathematical and Physical Sciences, Macquarie University, NSW 2109, Australia

Email address: xuan.duong@mq.edu.au

(Konstantin Merz) INSTITUT FÜR ANALYSIS UND ALGEBRA, TECHNISCHE UNIVERSITÄT BRAUNSCHWEIG, UNIVERSITÄTSPLATZ 2, 38106 BRAUNSCHWEIG, GERMANY, AND INSTITUT FÜR PARTIELLE DIFFEREN-TIALGLEICHUNGEN, TECHNISCHE UNIVERSITÄT BRAUNSCHWEIG, UNIVERSITÄTSPLATZ 2, 38106 BRAUNschweig, Germany

Email address: k.merz@tu-bs.de