

Spectra and transport of probability and energy densities in a \mathcal{PT} -symmetric square well with a delta-function potential

Francisco Ricardo Torres Arvizu,¹ Adrian Ortega,² and Hernán Larralde ^{*3}

¹*Instituto de Ciencias Físicas, UNAM, Av. Universidad s/n,
CP 62210 Cuernavaca Morelos, México*

²*Departamento de Física, Universidad de Guadalajara,
Blvd. Gral. Marcelino García Barragán 1421,
C.P. 44430, Guadalajara, Jalisco, México*

³*Instituto de Ciencias Físicas, UNAM, Av. Universidad s/n,
CP 62210 Cuernavaca Morelos, México*

We study the spectrum, eigenstates and transport properties of a simple \mathcal{PT} -symmetric model consisting in a finite, complex, square well potential with a delta potential at the origin. We show that as the strength of the delta potential increases, the system exhibits exceptional points accompanied by an accumulation of density associated with the break in the \mathcal{PT} -symmetry. We also obtain the density and energy density fluxes and analyze their transport properties. We find that in the \mathcal{PT} -symmetric phase transport is efficient, in the sense that all the density that flows into the system at the source, flows out at the sink, which is sufficient to derive a generalized unitary relation for the transmission and reflection coefficients.

I. INTRODUCTION

Since its introduction by Bender and Boettcher [1], \mathcal{PT} -symmetric quantum mechanics has attracted much interest due to the phenomena and applications that it can describe, see, for example [2–5] and references therein. One of the most studied aspects of non-Hermitian \mathcal{PT} -symmetric Hamiltonians is the appearance of exceptional points as a parameter of the system is varied. At such points, the \mathcal{PT} symmetry of the solutions breaks, the eigenvalues become complex [6], and two or more eigenvectors coalesce. As a result,

* Email: hernan@icf.unam.mx

the spectral decomposition of operators fails, and the Hamiltonian becomes defective [5, 7, 8].

To gain insight on the phenomenology that can arise in this kind of systems, it is desirable to have analytical models in which a system with a \mathcal{PT} -symmetric potential can be solved analytically (see for instance [9] and references therein for a general reference on non-Hermitian systems). For instance [10] addresses central potentials while references below address one-dimensional square well and step potentials. However, only a few \mathcal{PT} -symmetric potentials are solvable [5], perhaps the most explored have been one-dimensional square wells in their distinct versions. The first system of this type was a complex extension of the finite potential well where its bound states were calculated [11]. Due to the simplicity of this kind of systems, they have been studied in frameworks such as pseudo-Hermitian quantum mechanics [12] as well as from the perspective of supersymmetry [13], while other studies have focused on aspects such as the breaking of the \mathcal{PT} -symmetry in a step potential [14]. Later, modifications with periodic boundary conditions [15], discrete systems [16], two coupled \mathcal{PT} -symmetric square wells [17] and spatially antisymmetric [18] versions also have been investigated. Surprisingly, \mathcal{PT} -symmetric finite wells in an infinite interval, whose Hermitian form is traditionally addressed in almost any standard mechanics course [19], have barely been studied [20]. This model is interesting because it presents scattering states as well as an infinite set of bound state solutions [20]. In this work we consider a generalization of this model that includes a delta-potential at the origin, which gives rise to the appearance of exceptional points as well.

Traditionally, the physical meaning of the imaginary part of the potential is that it acts as source or drain of probability; thus, even in stationary states, fluxes are present in the system. Some studies on this subject have found that the transport of probability density in the \mathcal{PT} -symmetric phase in tight binding models is efficient, this is to say that in the \mathcal{PT} -symmetric phase the gain and loss of probability density in the system are globally balanced [21], whereas breaking the \mathcal{PT} symmetry causes the system to exhibit accumulation or depletion of density within the system [22, 23].

Regarding the scattering aspects of the non-Hermitian \mathcal{PT} -symmetric potential, it turns out that the transmission and reflection coefficients do not obey the usual unitary

rule. Instead, other “conservation rules” have been established [24–28]. Also, it has been reported that these coefficients present anomalous behaviour, i.e. they can take values greater than one [29], or even be infinite at certain values in the continuous spectrum [30, 31].

In this work, we study an extension of the (\mathcal{PT})-symmetric square potential well presented by Znojil [32] and Lévai [20]. Yet, unlike those systems, ours exhibits a broken \mathcal{PT} -symmetry phase. Our system consists of a finite \mathcal{PT} -symmetric complex potential well and a delta-potential at the origin modulated by a (real) coupling constant λ . This enables the system to exhibit a complexification of its eigenvalues as a function of this parameter, thereby displaying the characteristics specific to this phenomenon, such as the appearance of exceptional points. In addition to its spectral properties, we take also analyze its transport properties. For this, we compute the associated density currents for bound and scattering states, as well as the associated energy densities.

The outline of the paper is as follows: in Sec. II we present the solutions of the respective Schrödinger equations for bound and scattering states. For the bound states we discuss the spectra as a function of the parameters of the system. For scattering states, we find the reflection and transmission coefficients and we analyze their behavior. In Sec. III, first we discuss the continuity equations for the different quantum densities that are studied. Subsequently, the respective flows are calculated for this model for both the bound and scattering states. These fluxes are evaluated at the edges of the well to determine the efficiency of transport through the system. Finally, our results are summarized in Sec. IV.

II. THE MODEL AND ITS SOLUTIONS

We consider a finite, both in length and depth, \mathcal{PT} -symmetric square potential well [20, 32] in an infinite system, and add an extra term proportional to a Dirac delta function. This delta-potential acts as a barrier that, as the parameter associated with it increases, there comes a point where the transport from the source to the drain is no longer efficient, leading to a breakdown of the \mathcal{PT} -symmetry and the appearance of exceptional points. Specifically, we consider the following time independent Schrödinger equation

$$\left(-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} + V(x) + \lambda \delta(x) \right) \psi(x) = E \psi(x), \quad (1)$$

where λ is a real constant (the strength of the delta-potential) and $V(x)$ is the piecewise potential,

$$V(x) = \begin{cases} 0, & |x| < b, \\ V_0 + iV_I, & x \leq -b, \\ V_0 - iV_I, & x \geq b. \end{cases} \quad (2)$$

In this system, the wave function is subject to the boundary (matching) conditions

$$\begin{aligned} \psi(-b^-) &= \psi(-b^+), & \psi'(-b^-) &= \psi'(-b^+), \\ \psi(b^-) &= \psi(b^+), & \psi'(b^-) &= \psi'(b^+), \\ \psi(0^-) &= \psi(0^+), & \psi'(0^+) - \psi'(0^-) &= \Lambda\psi(0), \end{aligned} \quad (3)$$

where we have defined $\Lambda = \frac{2m\lambda}{\hbar^2}$ and the signs \pm denote the right or left limit, respectively, of the wavefunction at a given x . The matching conditions at the origin allow us to identify two types of basis functions: even (e) and odd (o) [33]

$$\psi_e(x) \propto \cos(kx) + \frac{\Lambda}{2k} \text{sgn}(x) \sin(kx), \quad (4)$$

$$\psi_o(x) \propto \sin(kx), \quad (5)$$

where $k = \sqrt{\frac{2mE}{\hbar^2}}$. Thus, inside the well, the \mathcal{PT} -symmetric wave function of the system can be written as a linear combination of the even and odd basis functions as [32]

$$\psi(x) = C_1\psi_e(x) + iC_2\psi_o(x), \quad (6)$$

with C_1, C_2 real constants. The complete solution of the Schödinger equations is

$$\psi(x) = \begin{cases} C_1 (\cos(kx) + \frac{\Lambda}{2k} \text{sgn}(x) \sin(kx)) + iC_2 \sin(kx), & -b < x < b, \\ A_1 e^{\alpha x} + A_2 e^{-\alpha x}, & x \leq -b, \\ B_1 e^{\tilde{\alpha} x} + B_2 e^{-\tilde{\alpha} x}, & x \geq b, \end{cases} \quad (7)$$

where $\alpha = \sqrt{v_0 + iv_I - k^2}$, $\tilde{\alpha} = \sqrt{v_0 - iv_I - k^2}$, $v_I = \frac{2mV_I}{\hbar^2}$ and $v_0 = \frac{2mV_0}{\hbar^2}$. While in principle k might be complex, in the \mathcal{PT} -symmetric phase k is real and $\tilde{\alpha} = \alpha^*$. Alternatively, we can write the real (α_R) and imaginary (α_I) part of α as

$$\alpha_R = \sqrt{\frac{\sqrt{(v_0 - k^2)^2 + v_I^2} + (v_0 - k^2)}{2}}, \quad (8)$$

$$\alpha_I = \sqrt{\frac{\sqrt{(v_0 - k^2)^2 + v_I^2} - (v_0 - k^2)}{2}}. \quad (9)$$

A. Discrete spectrum and bound states

The bound states are the normalizable solutions of the Schrödinger equation, therefore the wave function must vanish as $|x| \rightarrow \infty$. Taking the real parts of $\alpha, \tilde{\alpha}$ positive and making the coefficients $A_2, B_1 = 0$ in Eq. (7), we have an asymptotically vanishing wave function

$$\psi(x) = \begin{cases} C_1(\cos(kx) + \frac{\Lambda}{2k} \text{sgn}(x) \sin(kx)) + iC_2 \sin(kx), & -b < x < b, \\ A_1 e^{\alpha x}, & x \leq -b, \\ B_2 e^{-\tilde{\alpha} x}, & x \geq b. \end{cases} \quad (10)$$

The pseudo momentum k can be obtained by solving its associated transcendental equation which is obtained via the computation of the determinant arising from the boundary conditions Eq. (3). The equation is

$$(k^2 + \alpha\tilde{\alpha})\Lambda + (2k^2(\alpha + \tilde{\alpha}) + (k^2 - \alpha\tilde{\alpha})\Lambda) \cos(2kb) + k(2(\alpha\tilde{\alpha} - k^2) + (\alpha + \tilde{\alpha})\Lambda) \sin(2kb) = 0. \quad (11)$$

In the \mathcal{PT} -symmetric phase this equation can be written as

$$\underbrace{(2\Lambda(k \cos(kb) + \alpha_R \sin(kb)) + 4k(\alpha_R \cos(kb) - k \sin(kb)))}_{\text{Hermitian}} \underbrace{(k \cos(kb) + \alpha_R \sin(kb))}_{\text{non-Hermitian}} + 2\alpha_I^2 \sin(kb) (2k \cos(kb) + \Lambda \sin(kb)) = 0. \quad (12)$$

The first term in this expression, which we denote as the "Hermitian" part, factors into two independent equations corresponding to the two types of states expected in that case. These states can be identified as Λ -dependent (even) or Λ -independent (odd) states. In the \mathcal{PT} -symmetric system, the imaginary part of the potential couples the two kind of states in a new equation that cannot be factorized. This coupling, along with the strength of the parameter Λ associated with the Dirac delta, allows for the existence of exceptional points (Fig. 1). As the parameter Λ increases, the states coalesce, breaking the \mathcal{PT} -symmetry. In this figure, for reference, we have included the solutions for k when the delta-potential is absent ($\Lambda = 0$) in gray lines. In this case, there is no \mathcal{PT} -symmetry breaking, as noted in [20].

The exceptional points appear at different values of Λ as k grows. We can approximate their behavior when $k \gg v_0$ as follows. Expanding α_R and α_I for large values of k we have

$$\alpha_R \sim \frac{v_I}{2k}, \quad (13)$$

$$\alpha_I \sim k \left(1 - \frac{v_0}{2k^2}\right). \quad (14)$$

Substituting in Equation (12) and expanding to leading order in k we have

$$k\Lambda + (v_I \cos(2kb) - v_0 \sin(2kb)) = 0. \quad (15)$$

This equation has solution if we take $\Lambda \sim \chi/k$, where χ is constant

$$\chi + (v_I \cos(2kb) - v_0 \sin(2kb)) = 0, \quad (16)$$

and we find

$$\sin(2kb) = \frac{v_0\chi \pm \sqrt{v_I^2(v_0^2 + v_I^2 - \chi^2)}}{v_0^2 + v_I^2}. \quad (17)$$

From the square root, we note that real solutions only occur when

$$\chi < |\sqrt{v_0^2 + v_I^2}|. \quad (18)$$

Hence the curve that describes the location of the exceptional points for $k \gg 0$ is approximately $\kappa(\Lambda) = \sqrt{v_0^2 + v_I^2}/\Lambda$. We show this curve in figure (1), dashed purple line.

Another important aspect that can be highlighted about this system, shown in Fig. 1, is the presence of a discrete set of bound states with energy greater than V_0 . This fact has been reported previously in similar systems [20, 34]. Such states do not exist in the Hermitian case, which begs the question of what happens to them as the imaginary part of the potential vanishes. To answer this question, first we write the wave function in the \mathcal{PT} -symmetric phase obtaining the coefficients explicitly

$$\psi(x) = |C_1| \begin{cases} (\cos(kx) + \frac{\Lambda}{2k} \operatorname{sgn}(x) \sin(kx))(\alpha_R \sin(kb) + k \cos(kb)) + i\alpha_I (\cos(kb) + \frac{\Lambda}{2k} \sin(kb)) \sin(kx), & -b < x < b, \\ e^{\alpha(x+b)}(k \cos(kb) + \alpha^* \sin(kb)) \left(\cos(kb) + \frac{\Lambda}{2k} \sin(kb) \right), & x \leq -b, \\ e^{-\alpha^*(x-b)}(k \cos(kb) + \alpha \sin(kb)) \left(\cos(kb) + \frac{\Lambda}{2k} \sin(kb) \right), & x \geq b, \end{cases} \quad (19)$$

where the remaining constant C_1 accounts for the wavefunction normalization. With this expression we compute the probability density $\rho_D(x) = |\psi(x)|^2$, and using Eq. (11) we obtain

$$\rho_D(x) = |C_1|^2 \begin{cases} (\cos(kx) + \frac{\Lambda}{2k} \operatorname{sgn}(x) \sin(kx))^2 (\alpha_R \sin(kb) + k \cos(kb))^2 + \alpha_I^2 (\cos(kb) + \frac{\Lambda}{2k} \sin(kb))^2 \sin^2(kx), & -b < x < b, \\ e^{2\alpha(x+b)} k \left(k \cos(kb) + \alpha_R \sin(kb) \right) \left(\cos(kb) + \frac{\Lambda}{2k} \sin(kb) \right), & x \leq -b, \\ e^{-2\alpha(x-b)} k \left(k \cos(kb) + \alpha_R \sin(kb) \right) \left(\cos(kb) + \frac{\Lambda}{2k} \sin(kb) \right), & x \geq b. \end{cases} \quad (20)$$




FIG. 1: The blue curves represent the pseudo-momentum k for the bound states as a function of λ . These curves end at exceptional points at values approximated by $\kappa(\Lambda)$ in

Eq.(18) (purple line), for states with energies greater than $\sqrt{v_0}$ (red dashed line).

Additionally, we can see the solutions for $\Lambda = 0$ (gray lines). In the insets one can observe the complexification of the spectrum of the first six states as a function of the parameter

Λ , here k_R and k_I denote the real and imaginary parts of k .

Using the normalization condition [49]

$$\int_{-\infty}^{\infty} \rho_d(x) dx = 1, \quad (21)$$

we obtain

$$|C_1|^2 = \frac{4\alpha_R k^3}{D(k)}. \quad (22)$$

where $D(k)$ is given by

$$\begin{aligned} D(k) = & \alpha_R \left[2 \sin(kb) \left((4k^2 - \Lambda^2) \cos(kb) + 4k\Lambda \sin(kb) \right) (k \cos(kb) + \alpha_R \sin(kb))^2 \right. \\ & - 4k^2 \alpha_I^2 \sin(2kb) \left(\cos(kb) + \frac{\Lambda}{2k} \sin(kb) \right)^2 \\ & \left. + 2kb \left((4k^2 + \Lambda^2) (k \cos(kb) + \alpha_R \sin(kb))^2 + 4k^2 \alpha_I^2 \left(\cos(kb) + \frac{\Lambda}{2k} \sin(kb) \right)^2 \right) \right] \\ & + 8k^4 \left[\left(k \cos(kb) + \alpha_R \sin(kb) \right) \left(\cos(kb) + \frac{\Lambda}{2k} \sin(kb) \right) \right]. \end{aligned} \quad (23)$$

In order to understand how these states disappear while those with energies below V_0 remain, we focus on the behavior of the normalization constant Eq. (22) as $v_I \rightarrow 0$. Taking the limit of small v_I in Eq. (8) we find that

$$\alpha_R \sim \begin{cases} |v_0 - k^2|^{1/2}, & k^2 \leq v_0, \\ \frac{v_I}{2|v_0 - k^2|^{1/2}}, & k^2 > v_0. \end{cases} \quad (24)$$

Thus, as the imaginary potential decreases, α_R , and hence $|C_1|^2$, vanish as v_I decreases for $k^2 > v_0$, but not for $k^2 < v_0$. This reflects the fact that as the potential becomes real, the bound eigenstates with energies above V_0 become increasingly extended and their amplitude tends to zero to maintain normalization. This behaviour can be seen in Fig. 2. In this figure, one can see in a) the probability densities for bound states with $k^2 < v_0$ and for three different values of the imaginary part of the potential $v_I = 0.5, 5, 10$; even if the potential increases, the wavefunction remains more or less invariant. On the other hand, b) shows the case where $k^2 > v_0$, i.e. bound states above the maximum of the real part of the potential, again using the same values for v_I as in a); in this case, as v_I decreases, the wavefunctions become flatter.




FIG. 2: Density of the bound states for cases when $k^2 < V_0$ a) and $k^2 > V_0$ b) for some values of v_I . As the magnitude of the imaginary potential decreases, the bound states with $k^2 > v_0$ become more extended and their magnitude decreases accordingly.

B. Scattering States

In what follows we describe the properties of scattering states that occur in this system [26, 35]. The specific scattering processes can be seen schematically in Fig. 3. The solution




FIG. 3: Scattering process. a) Left-to-right incidence and b) right-to-left incidence. The figure also illustrates the meaning of the coefficients r_+ , r_- , t_+ and t_- .

of the Schrödinger equation for the left-to-right scattering case can be obtained making the

coefficient $B_1 = 0$ in Eq. (7), thus

$$\psi_+(x) = \begin{cases} C_1(\cos(kx) + \frac{\Lambda}{2k}\text{sgn}(x)\sin(kx)) + iC_2\sin(kx), & |x| < b, \\ A_1e^{\alpha x} + A_2e^{-\alpha x}, & x \leq -b, \\ B_2e^{-\alpha^*x}, & x \geq b. \end{cases} \quad (25)$$

We define the left-to-right reflection and transmission coefficients as

$$\mathbf{r}_+ = \frac{A_2}{A_1}, \quad (26)$$

$$\mathbf{t}_- = \frac{B_2}{A_1}. \quad (27)$$

The scattering solutions are not \mathcal{PT} -symmetric, and due to the presence of a source and a sink, the relation $|\mathbf{r}_+|^2 + |\mathbf{t}_-|^2 = 1$ is not satisfied. Instead we must consider the inverse process, i.e. the right-to-left scattering with wave function given by (c.f. Eq. (7))

$$\psi_-(x) = \begin{cases} C_1(\cos(kx) + \frac{\Lambda}{2k}\text{sgn}(x)\sin(kx)) + iC_2\sin(kx), & |x| < b, \\ A_2e^{-\alpha x}, & x \leq -b, \\ B_1e^{\alpha^*x} + B_2e^{-\alpha^*x}, & x \geq b. \end{cases} \quad (28)$$

The right-to-left reflection and transmission coefficients are

$$\mathbf{r}_- = \frac{B_2}{B_1}, \quad (29)$$

$$\mathbf{t}_+ = \frac{A_2}{B_1}. \quad (30)$$

The complete set of reflection and transmission coefficients can be explicitly computed through the transfer matrix method (see Appendix A), and yield

$$\begin{aligned} \mathbf{t}_+ &= \frac{4\alpha^*k^2}{-2k\sin(2kb)(|\alpha|^2+k^2+i\Lambda\alpha_I)+\cos(2kb)(|\alpha|^2\Lambda+k^2(-4i\alpha_I+\Lambda))+\Lambda(k^2-|\alpha|^2)}e^{2ib\alpha_I}, \\ \mathbf{r}_- &= \frac{(2k\sin(2kb)(\Lambda\alpha_R-(|\alpha|^2-k^2))+\cos(2kb)(|\alpha|^2\Lambda+k^2(4\alpha_R-\Lambda))-\Lambda(k^2+|\alpha|^2))}{-2k\sin(2kb)(|\alpha|^2+k^2+i\Lambda\alpha_I)+\cos(2kb)(|\alpha|^2\Lambda+k^2(-4i\alpha_I+\Lambda))+\Lambda(k^2-|\alpha|^2)}e^{2b\alpha^*}, \\ \mathbf{r}_+ &= -\frac{(2k\sin(2kb)(\Lambda\alpha_R+(|\alpha|^2-k^2))+\cos(2kb)(k^2(4\alpha_R+\Lambda)-|\alpha|^2\Lambda)+\Lambda(k^2+|\alpha|^2))}{-2k\sin(2kb)(|\alpha|^2+k^2+i\Lambda\alpha_I)+\cos(2kb)(|\alpha|^2\Lambda+k^2(-4i\alpha_I+\Lambda))+\Lambda(k^2-|\alpha|^2)}e^{-2b\alpha}, \\ \mathbf{t}_- &= -\frac{4\alpha k^2}{-2k\sin(2kb)(|\alpha|^2+k^2+i\Lambda\alpha_I)+\cos(2kb)(|\alpha|^2\Lambda+k^2(-4i\alpha_I+\Lambda))+\Lambda(k^2-|\alpha|^2)}e^{2ib\alpha_I}. \end{aligned} \quad (31)$$

From these equations we can compute the pseudo-transmissions $T_{\pm} = |\mathbf{t}_{\pm}|^2$ and pseudo-reflections $R_{\pm} = |\mathbf{r}_{\pm}|^2$, which are shown in Fig. 4. We note that the the left and right transmission are equal, $|\mathbf{t}_+(k)|^2 = |\mathbf{t}_-(k)|^2$. Actually, the pseudo-transmissions and




FIG. 4: a) Pseudo-transmission $T_{+/-}$, b) Left pseudo-reflection R_+ , c) Right pseudo-reflection R_- d) Module of the product of left and right pseudo-reflections as a function of k for the parameter $v_I = 10$, $v_I = 10$ and $b = 1$.

pseudo-reflection can show anomalous behavior, taking values greater than one as can be seen in Fig. 4. On the other hand, the left reflectivity (R_+) vanishes at the bound states as expected. This can be observed in Fig. 4 where we recover the spectrum plot in the (Λ, k) -plane. The figure also shows that there is a set of discrete values of k at which R_- vanishes, which correspond to non integrable reflection-less states.

Finally, it can also be verified that the reflection and transmission coefficients satisfy the

following identities reported in general for \mathcal{PT} -symmetric systems [28, 36]

$$\mathbf{r}_{\pm}(-k) = \mathbf{r}_{\pm}(k), \quad (32)$$

$$\mathbf{t}_{\pm}(-k) = \mathbf{t}_{\pm}(k). \quad (33)$$

Indeed, the system satisfies the "generalized unitary relation" [24, 28]

$$|\mathbf{t}_{\pm}(k)|^2 \pm |\mathbf{r}_{-}(k)\mathbf{r}_{+}(k)| = 1. \quad (34)$$

The sign is chosen according to whether the quantity, $1 - T = 1 - |\mathbf{t}_{+}(k)|^2$ [37] is positive or negative: in the regions where the pseudo-transmission presents an anomalous behavior $T > 1$, the negative sign will be chosen, otherwise, where $T < 1$ one chooses the positive one.

III. TRANSPORT PROPERTIES

In this section we focus on the analysis of the transport properties of the system. We discuss briefly how different definitions for the energy density yield different continuity equations, then we analyze both definitions in the context of our specific problem.

A. Probability density, energy density and their continuity equations

In general, we consider a particle in an arbitrary complex potential

$$V(x) = V_R(x) + iV_I(x). \quad (35)$$

The particle is described by the wavefunction $\psi(x, t)$ for which the probability density is $\rho_d(x, t) = |\psi(x, t)|^2$. This probability density satisfies the (standard) continuity equation

$$\frac{\partial \rho_d(x, t)}{\partial t} + \frac{\partial J_d(x, t)}{\partial x} = Q_d(x, t), \quad (36)$$

where the probability current is

$$J_d(x, t) = \frac{\hbar}{2mi} \left[\psi^*(x, t) \frac{\partial}{\partial x} \psi(x, t) - \psi(x, t) \frac{\partial}{\partial x} \psi^*(x, t) \right], \quad (37)$$

and

$$Q_d(x, t) = 2 \frac{V_I(x) \rho_d(x, t)}{\hbar}, \quad (38)$$

accounts for the source/drain terms. Under this perspective, the imaginary part of the potential is interpreted as giving rise to a source or sink of probability [38]. However, it is important to mention that a self-consistent conservation law has been proposed for the norm defined by the \mathcal{PT} inner product [39, 40]. However, this norm usually presents certain drawbacks, so for the purposes of this work, we will not adopt it. Instead, we will approach the transport properties from the perspective of the continuity equation (36).

Another quantity that satisfies a continuity equation in quantum mechanics, and is less explored, is the energy density [41–43]. Unlike the definition for the probability density, the definition for the energy density in quantum mechanics is not unique, different definitions yield different results and therefore different interpretations of the “flux” of energy in a quantum system. An initial attempt to define the energy density could be simply as

$$\rho^E(x, t) = \psi^*(x, t)H\psi(x, t). \quad (39)$$

However, this expression is not necessarily real, which makes its interpretation as a physical density troublesome. However, for \mathcal{PT} –symmetric states

$$\int V_I(x)\rho_d(x, t)dx = 0. \quad (40)$$

so V_I does not contribute to the expectation value of the energy. Instead, we note that the integrand of the previous equation is proportional to $Q_d(x, t)$ [27, 38, 44], thus, equation (40) can be interpreted as saying that in \mathcal{PT} –symmetric systems gain and loss of density is globally balanced [21, 22].

The condition (40) allows us to obtain a first acceptable definition of the energy density (ρ_1^E) for the \mathcal{PT} –symmetric phase as

$$\rho_1^E(x, t) = \frac{1}{2}(\psi^*(x, t)H\psi(x, t) + \psi(x, t)H^\dagger\psi^*(x, t)). \quad (41)$$

This definition yields real density, but has the drawback of not being necessarily positive even for non-negative potentials. Alternatively, another definition of energy density is

$$\rho_2^E(x, t) = \frac{\hbar^2}{2m} \left| \frac{\partial}{\partial x} \psi(x, t) \right|^2 + V_R(x)|\psi(x, t)|^2, \quad (42)$$

as can be checked by integrating by parts and using that the wave function vanishes at infinity.

Following the procedure shown in [43], it is possible to derive continuity equations for the two definitions of energy density,

$$\frac{\partial \rho_1^E(x, t)}{\partial t} + \frac{\partial}{\partial x} J_1^E(x, t) = Q^E(x, t), \quad (43)$$

$$\frac{\partial \rho_2^E(x, t)}{\partial t} + \frac{\partial}{\partial x} J_2^E(x, t) = Q^E(x, t), \quad (44)$$

where $Q^E(x, t) = 2\frac{V_I(x, t)\rho_2^E(x, t)}{\hbar}$ is the same in both cases, and the respective energy density fluxes J_1^E and J_2^E are given by

$$J_1^E(x, t) = \frac{\hbar}{4mi} \left[\psi^*(x, t) \frac{\partial}{\partial x} [H\psi(x, t)] - \psi(x, t) \frac{\partial}{\partial x} [H\psi(x, t)]^* \right. \\ \left. - \left(\frac{\partial}{\partial x} \psi^*(x, t) \right) H\psi(x, t) + \left(\frac{\partial}{\partial x} \psi(x, t) \right) [H\psi(x, t)]^* \right], \quad (45)$$

$$J_2^E(x, t) = \frac{\hbar}{2mi} \left[\left(\frac{\partial}{\partial x} \psi(x, t) \right) [H\psi(x, t)]^* - \left(\frac{\partial}{\partial x} \psi(x, t)^* \right) H\psi(x, t) \right], \quad (46)$$

which are similar to the energy density fluxes that appear in the Hermitian case [41–43].

Now we analyze first Sec. III A 1 the probability and energy transport properties of the bound states. Subsequently in Sec. III A 2, we apply this analysis for the scattering states.

1. Bound states

First we compute the probability density flux in the \mathcal{PT} –symmetric phase, i.e.

$$J_d(x) = |C_1|^2 \frac{\alpha_I \hbar}{m} k \left(k \cos(kb) + \alpha_R \sin(kb) \right) \left(\cos(kb) + \frac{\Lambda}{2k} \sin(kb) \right) \times \begin{cases} 1, & -b < x < b, \\ e^{2\alpha_R(x+b)}, & x \leq -b, \\ e^{-2\alpha_R(x-b)}, & x \geq b. \end{cases} \quad (47)$$

where $|C_1|^2$ is given in Eq.(22). We can see in this expression that the flux on the side of the source ($x \leq b$) increases as it approaches the edge of the well. Once there, it remains constant across the well, and it decreases on the other side of the well.

As a consequence of the stationary Schrödinger equation, the energy density fluxes in Eqs. (45) and (46) are

$$J_1^E(x) = J_2^E(x) = \frac{\hbar^2 k^2}{2m} J_d(x). \quad (48)$$

The source/drain terms of probability density and energy density are given by

$$Q_d(x) = \frac{2V_I|C_1|^2}{\hbar} \begin{cases} 0, & |x| < b, \\ e^{2\alpha_R(x+b)}k \left(k \cos(kb) + \alpha_R \sin(kb) \right) \left(\cos(kb) + \frac{\Lambda}{2k} \sin(kb) \right), & x \leq -b, \\ -e^{-2\alpha_R(x-b)}k \left(k \cos(kb) + \alpha_R \sin(kb) \right) \left(\cos(kb) + \frac{\Lambda}{2k} \sin(kb) \right), & x \geq b, \end{cases} \quad (49)$$

and

$$Q_E(x) = \frac{\hbar^2 k^2}{2m} Q_d(x). \quad (50)$$

The integral of this quantity over the entire system vanishes, which means there is no net gain nor loss of density or energy, as expected in the \mathcal{PT} -symmetric phase.

On the other hand, the energy density ρ_2^E is

$$\rho_2^E(x) = |C_1|^2 \frac{\hbar^2}{2m} \begin{cases} k^2 \left[\left(\Lambda \delta(x) + (-\sin(kx) + \frac{\Lambda}{2k} \operatorname{sgn}(x) \cos(kx)) \right)^2 (\alpha_R \sin(kb) + k \cos(kb))^2 \right. \\ \left. + \alpha_I^2 (\cos(kb) + \frac{\Lambda}{2k} \sin(kb))^2 \cos^2(kx) \right], & -b < x < b, \\ e^{2\alpha_R(x+b)} (2\alpha_R^2 + k^2) k \left(k \cos(kb) + \alpha_R \sin(kb) \right) \left(\cos(kb) + \frac{\Lambda}{2k} \sin(kb) \right), & x \leq -b, \\ e^{-2\alpha_R(x-b)} (2\alpha_R^2 + k^2) k \left(k \cos(kb) + \alpha_R \sin(kb) \right) \left(\cos(kb) + \frac{\Lambda}{2k} \sin(kb) \right), & x \geq b. \end{cases} \quad (51)$$

It is interesting to see how the potential induces a discontinuity in the energy density ρ_2^E , while the behavior of ρ_1^E is the same as that of the probability density.

2. Scattering states

In the scattering case, starting with the left to right incidence we have that the flux in the source side can be written as

$$J_{d+}(x) = \frac{|A_1|^2 \hbar}{m} \left[\alpha_I (e^{2\alpha_R x} - |\mathbf{r}_+(k)|^2 e^{-2\alpha_R x}) + i\alpha_R (\mathbf{r}_+(k) e^{-2i\alpha_I x} - \mathbf{r}_+^*(k) e^{2i\alpha_I x}) \right] \quad (52)$$

and the flux in the right side is

$$J_{d-}(x) = \frac{|A_1|^2 |\mathbf{t}_-(k)|^2 \hbar \alpha_I e^{-2\alpha_R x}}{m}. \quad (53)$$

Similarly, in the case of right to left incidence, the density flux on the right side of the well is

$$J_{d-}(x) = \frac{|B_1|^2 \hbar}{m} \left[\alpha_I (|\mathbf{r}_-(k)|^2 e^{-2\alpha_R x} - e^{2\alpha_R x}) + i\alpha_R (\mathbf{r}_-(k) e^{2i\alpha_I x} - \mathbf{r}_-^*(k) e^{-2i\alpha_I x}) \right], \quad (54)$$

and the flux on the left side, i.e. the flux for the transmitted wave is

$$J_{d_+}(x) = \frac{|B_1|^2 |\mathbf{t}_+(k)|^2 \hbar \alpha_I e^{-2\alpha_R x}}{m}. \quad (55)$$

Evaluating the fluxes at the edges of the well, we have the following conservation law

$$J_{d_+}(-b) = J_{d_-}(b). \quad (56)$$

As in the case for bound states, we have efficient transport also in this case. Relation (56) is also interesting because from it one can derive a kind of generalized unitary condition. In order to do so, we start with condition (56) and we evaluate Eq. (52) in $x = -b$, this yields

$$J_{d_+}(-b) = \alpha_I \frac{|A_1|^2 \hbar}{m} e^{-2\alpha_R b} \left[1 - (2k \sin(2kb) (\Lambda \alpha_R - (|\alpha|^2 - k^2)) + \cos(2kb) (|\alpha|^2 \Lambda + k^2 (4\alpha_R - \Lambda)) - \Lambda (k^2 + |\alpha|^2)) \times \right. \\ \left. \frac{(2k \sin(2kb) (\Lambda \alpha_R + (|\alpha|^2 - k^2)) + \cos(2kb) (k^2 (4\alpha_R + \Lambda) - |\alpha|^2 \Lambda) + \Lambda (k^2 + |\alpha|^2))}{((k^2 - |\alpha|^2) \Lambda - (k^2 + |\alpha|^2) (-\Lambda \cos(2bk) + 2k \sin(2bk)))^2 + (2k \alpha_I (2k \cos(2bk) + \Lambda \sin(2bk)))^2} \right]. \quad (57)$$

From Eq. (31) we can write

$$J_{d_+}(-b) = \alpha_I \frac{|A_1|^2 \hbar}{m} e^{-2\alpha_R b} \left[1 - |\mathbf{r}_+(k) \mathbf{r}_-(k)| \right]. \quad (58)$$

Finally, equating the last equation with Eq. (53) we obtain again the unitary relation Eq. (34), i.e.

$$|\mathbf{t}_\pm(k)|^2 \pm |\mathbf{r}_-(k) \mathbf{r}_+(k)| = 1. \quad (59)$$

In the case of Hermitian systems, this relation reduces to the standard probability conservation $|\mathbf{t}_\pm(k)|^2 + |\mathbf{r}_\pm(k)|^2 = 1$, in our case probability is not conserved in general.

IV. SUMMARY AND CONCLUSIONS

In this article, we investigate the spectral and transport properties of both energy and density in a \mathcal{PT} -symmetric system. For this purpose, we analyzed a relatively simple system which possesses both bound and scattering states. Specifically, the system we considered is a \mathcal{PT} -symmetric finite potential well with a delta-potential at the origin of strength λ . It is worth noting that without the delta potential, the eigenvalues of the system remain real [20] for any value of V_I , the imaginary part of the potential. However, the combination of λ and V_I allows the complexification of the eigenvalues and the

breaking of the \mathcal{PT} symmetry. This can be explained through the equation for the pseudo momentum k , there the imaginary part of the potential couples the values corresponding to eigenfunctions that would be odd (and hence independent of the delta potential) and those that would be even in the absence of V_I . This coupling becomes stronger as we increase the value of the parameters, finally producing exceptional points. We were also able to describe the approximate value at which the exceptional points appear in the limit $k \gg v_0$.

Another interesting aspect of the spectrum of these \mathcal{PT} -symmetric finite potential well systems is the presence of bound states with energies $E > V_0$. We also saw how these states disappear when the imaginary part of the potential goes to zero, which has to occur as we approach the Hermitian case. We found that the bound states with energy $E > V_0$ become infinitely extended causing their amplitude to vanish when the imaginary potential tends to zero, whereas those with energies $V_0 > E$ remain localized.

Regarding the transport properties of the system, the transmission and reflection coefficients were calculated. As expected, the zeros of the left reflection coefficient correspond to the eigenvalues of the bound states, because the incident wave and the transmitted wave both vanish asymptotically, so at the zeros of the transmission, the scattered and bound wave functions are the same. In the inverse process, the scattering from right to left, the states that correspond to the zeros of the reflection coefficient are a discrete set of reflectionless states that diverge to infinity.

The transport efficiency of the system can be quantified by evaluating the density fluxes at the edges of the well, that is, in the positions of the source and drain. In these regions, we can observe that, everything that comes from the source reaches the sink, the gain and loss balance, such that there is no accumulation or depletion of probability or energy in the system. This characteristic has also been reported for tight-binding Hamiltonians with gain and loss sites [22, 23]. However, as the strength of the delta increases it acts as a kind of barrier, which modulates the flow of probability and energy, interfering with the transport efficiency of the system, and leading to a break in the \mathcal{PT} -symmetry.

Since the system studied here can be solved exactly, it would be of interest to use our findings in extending the analysis to some other effects which might also be of potential

interest. These comprise, but are not limited to, the case of analyzing time-dependent wavepackets (e.g. Gaussian wavepackets) and its evolution throughout the \mathcal{PT} scattering system, the so-called topological energy transfer with exceptional points [45], the study of the jamming anomaly [46], etc.

Acknowledgements

Francisco Ricardo Torres Arvizu acknowledges support from CONAHCYT scholarship number 834573. Hernán Larralde and Francisco Ricardo Torres Arvizu acknowledges the National Autonomous University of México (UNAM) through the SUPPORT PROGRAM FOR RESEARCH AND TECHNOLOGICAL INNOVATION PROJECTS (PAPIIT) key IN103724. A. O. acknowledges support from the program “Apoyos para la Incorporación de Investigadoras e Investigadores Vinculada a la Consolidación Institucional de Grupos de Investigación 2023” from CONAHCYT, México.

Appendix A: Transfer matrix

The matching conditions can be summarize using the transfer matrices $\mathcal{M}^+(k)$, $\mathcal{M}^-(k)$ which are defined by the relations

$$\mathcal{M}^+(k) \begin{bmatrix} A_1 \\ A_2 \end{bmatrix} = \begin{bmatrix} B_2 \\ B_1 \end{bmatrix}, \quad \mathcal{M}^-(k) \begin{bmatrix} B_2 \\ B_1 \end{bmatrix} = \begin{bmatrix} A_1 \\ A_2 \end{bmatrix}, \quad (\text{A1})$$

it can easily see that $\mathcal{M}^+(k)^{-1} = \mathcal{M}^-(k)$. Thus the components of transfer matrix are

$$\begin{aligned} \mathcal{M}_{1,1}^+ &= \frac{e^{-2ib\alpha_I} (2k \sin(2kb) (2(|\alpha|+k^2) - i\Lambda\alpha_I) - \cos(2kb) (|\alpha|^2\Lambda + k^2(4i\alpha_I + \Lambda)) - \Lambda(k^2 - |\alpha|^2))}{4\alpha^*k^2}, \\ \mathcal{M}_{1,2}^+ &= \frac{e^{2b\alpha_R} (2k \sin(2kb) (\Lambda\alpha_R - (|\alpha|^2 - k^2)) + \cos(2kb) (|\alpha|^2\Lambda + k^2(4\alpha_R - \Lambda)) - \Lambda(k^2 + |\alpha|^2))}{4\alpha^*k^2}, \\ \mathcal{M}_{2,1}^+ &= \frac{e^{-2b\alpha_R} (-2k \sin(2kb) (\Lambda\alpha_R + (|\alpha|^2 - k^2)) + \cos(2kb) (k^2(4\alpha_R + \Lambda) - |\alpha|^2\Lambda) + \Lambda(k^2 + |\alpha|^2))}{4\alpha^*k^2}, \\ \mathcal{M}_{2,2}^+ &= \frac{e^{2ib\alpha_I} (-2k \sin(2kb) (|\alpha|^2 + k^2) + i\Lambda\alpha_I) + \cos(2kb) (|\alpha|^2\Lambda + k^2(-4i\alpha_I + \Lambda)) + \Lambda(k^2 - |\alpha|^2)}{4\alpha^*k^2}, \end{aligned} \quad (\text{A2})$$

Is well know that the transfer matrix (and its inverse) can be written in terms of the transmission and reflection coefficients (denoted \mathbf{r} , \mathbf{t} respectively), and considering the above

we have [35]

$$\mathcal{M}^+(k) = \begin{bmatrix} \mathbf{t}_-(k) - \mathbf{r}_+(k)\mathbf{r}_-(k)/\mathbf{t}_+(k) & \mathbf{r}_-(k)/\mathbf{t}_+(k) \\ -\mathbf{r}_+(k)/\mathbf{t}_+(k) & 1/\mathbf{t}_+(k) \end{bmatrix}, \quad (\text{A3})$$

$$\mathcal{M}^-(k) = \begin{bmatrix} 1/\mathbf{t}_-(k) & -\mathbf{r}_-(k)/\mathbf{t}_-(k) \\ \mathbf{r}_+(k)/\mathbf{t}_-(k) & \mathbf{t}_+(k) - \mathbf{r}_+(k)\mathbf{r}_-(k)/\mathbf{t}_-(k) \end{bmatrix}. \quad (\text{A4})$$

Also, the determinant of this matrix in this case is

$$\det(\mathcal{M}^+) = -\frac{\alpha}{\alpha^*} = \frac{\mathbf{t}_-(k)}{\mathbf{t}_+(k)}; \quad (\text{A5})$$

which means no reciprocity in transmission. Hence, we write the wave function of scattering process as

$$\psi_+(x) = A_1 \begin{cases} \frac{\left(k \cos(k(b-x)) + \alpha^* \sin(k(b-x))\right) \mathbf{t}_-(k) e^{-\alpha^* b}}{k}, & |x| < b, \\ e^{\alpha x} + \mathbf{r}_+(k) e^{-\alpha x}, & x \leq -b, \\ \mathbf{t}_-(k) e^{-\alpha^* x}, & x \geq b, \end{cases} \quad (\text{A6})$$

$$\psi_-(x) = B_1 \begin{cases} \frac{\left(k \cos(k(x+b)) - \alpha \sin(k(x+b))\right) \mathbf{t}_+(k) e^{\alpha b}}{k}, & |x| < b, \\ \mathbf{t}_+(k) e^{-\alpha x}, & x \leq -b, \\ e^{\alpha^* x} + \mathbf{r}_-(k) e^{-\alpha^* x}, & x \geq b. \end{cases} \quad (\text{A7})$$

- [1] C. M. Bender and S. Boettcher, Phys. Rev. Lett. **80**, 5243 (1998), URL <https://link.aps.org/doi/10.1103/PhysRevLett.80.5243>.
- [2] F. Bagarello, R. Passante, and C. Trapani, eds., *Non-Hermitian Hamiltonians in Quantum Physics. Springer Proceedings in Physics, Vol 184* (Springer, Cham, 2016).
- [3] R. El-Ganainy, K. G. Makris, M. Khajavikhan, Z. H. Musslimani, S. Rotter, and D. N. Christodoulides, Nature Physics **14**, 11 (2018), URL <https://doi.org/10.1038/nphys4323>.
- [4] D. Christodoulides, J. Yang, et al., *Parity-time symmetry and its applications*, vol. 280 (Springer, 2018).
- [5] C. M. Bender, *PT symmetry: In quantum and classical physics* (World Scientific, 2019).
- [6] A. A. Zyblovsky, A. P. Vinogradov, A. A. Pukhov, A. V. Dorofeenko, and A. A. Lisyansky, Physics-Uspekhi **57**, 1063 (2014).

- [7] T. Kato, *Perturbation theory for linear operators*, vol. 132 (Springer Science & Business Media, 2013).
- [8] W. D. Heiss, *Journal of Physics A: Mathematical and Theoretical* **45**, 444016 (2012), URL <https://doi.org/10.1088/1751-8113/45/44/444016>.
- [9] N. Moiseyev, *Non-Hermitian Quantum Mechanics* (Cambridge University Press, 2011).
- [10] G. Lévai, *J. of Phys.: Conf. Ser.* **4128**, 012045 (2008).
- [11] M. Znojil, *Physics Letters A* **285**, 7 (2001), ISSN 0375-9601, URL <https://www.sciencedirect.com/science/article/pii/S0375960101003012>.
- [12] A. Mostafazadeh and A. Batal, *Journal of Physics A: Mathematical and General* **37**, 11645 (2004), URL <https://dx.doi.org/10.1088/0305-4470/37/48/009>.
- [13] B. BAGCHI, S. MALLIK, H. BÍLA, V. JAKUBSKÝ, M. ZNOJIL, and C. QUESNE, *International Journal of Modern Physics A* **21**, 2173 (2006), <https://doi.org/10.1142/S0217751X0602951X>, URL <https://doi.org/10.1142/S0217751X0602951X>.
- [14] M. ZNOJIL and G. LÉVAI, *Modern Physics Letters A* **16**, 2273 (2001), <https://doi.org/10.1142/S0217732301005722>, URL <https://doi.org/10.1142/S0217732301005722>.
- [15] V. Jakubský and M. Znojil, *Czechoslovak Journal of Physics* **54**, 1101 (2004), URL <https://doi.org/10.1023/B:CJOP.0000044010.18569.bd>.
- [16] M. Znojil, *Journal of Physics A: Mathematical and General* **39**, 10247 (2006), URL <https://dx.doi.org/10.1088/0305-4470/39/32/S23>.
- [17] M. Znojil, *Journal of Physics A: Mathematical and General* **39**, 4047 (2006), URL <https://dx.doi.org/10.1088/0305-4470/39/15/011>.
- [18] C. Quesne, B. Bagchi, S. Mallik, H. Bila, V. Jakubsky, and M. Znojil, *Czechoslovak Journal of Physics* **55**, 1161 (2005), URL <https://doi.org/10.1007/s10582-005-0122-y>.
- [19] D. W. L. Sprung, H. Wu, and J. Martorell, *American Journal of Physics* **64**, 136 (1996), ISSN 0002-9505, https://pubs.aip.org/aapt/ajp/article-pdf/64/2/136/11786964/136_1_online.pdf, URL <https://doi.org/10.1119/1.18131>.
- [20] G. Lévai and J. Kovács, *Journal of Physics A: Mathematical and Theoretical* **52**, 025302 (2018).
- [21] S. Weigert, *Czechoslovak Journal of Physics* **54**, 1139 (2004), URL <https://doi.org/10.1023/B:CJOP.0000044010.18569.bd>.

- 1023/B:CJOP.0000044016.95629.a7.
- [22] A. Ortega, T. Stegmann, L. Benet, and H. Larralde, *Journal of Physics A: Mathematical and Theoretical* **53**, 445308 (2020).
- [23] A. Ortega, L. Benet, and H. Larralde, *Journal of Physics A: Mathematical and Theoretical* **55**, 015304 (2021).
- [24] L. Ge, Y. D. Chong, and A. D. Stone, *Phys. Rev. A* **85**, 023802 (2012), URL <https://link.aps.org/doi/10.1103/PhysRevA.85.023802>.
- [25] Z. Lin, H. Ramezani, T. Eichelkraut, T. Kottos, H. Cao, and D. N. Christodoulides, *Phys. Rev. Lett.* **106**, 213901 (2011), URL <https://link.aps.org/doi/10.1103/PhysRevLett.106.213901>.
- [26] F. Cannata, J.-P. Dedonder, and A. Ventura, *Annals of Physics* **322**, 397 (2007), ISSN 0003-4916, URL <https://www.sciencedirect.com/science/article/pii/S0003491606001229>.
- [27] Z. Ahmed, *Physics Letters A* **282**, 343 (2001), ISSN 0375-9601, URL <https://www.sciencedirect.com/science/article/pii/S0375960101002183>.
- [28] A. Mostafazadeh, *Journal of Physics A: Mathematical and Theoretical* **47**, 505303 (2014), URL <https://dx.doi.org/10.1088/1751-8113/47/50/505303>.
- [29] Z. Ahmed, *Physics Letters A* **324**, 152 (2004), ISSN 0375-9601, URL <https://www.sciencedirect.com/science/article/pii/S0375960104003214>.
- [30] A. Mostafazadeh, *Phys. Rev. Lett.* **102**, 220402 (2009), URL <https://link.aps.org/doi/10.1103/PhysRevLett.102.220402>.
- [31] L. Chaos-Cador and G. García-Calderón, *Phys. Rev. A* **87**, 042114 (2013), URL <https://link.aps.org/doi/10.1103/PhysRevA.87.042114>.
- [32] M. Znojil, *Journal of Mathematical Physics* **46**, 062109 (2005), ISSN 0022-2488, https://pubs.aip.org/aip/jmp/article-pdf/doi/10.1063/1.1925249/14868935/062109_1_online.pdf, URL <https://doi.org/10.1063/1.1925249>.
- [33] A. Mostafazadeh, *Journal of Physics A: Mathematical and General* **39**, 13495 (2006), URL <https://dx.doi.org/10.1088/0305-4470/39/43/008>.
- [34] G. Lévai and E. Magyari, *Journal of Physics A: Mathematical and Theoretical* **42**, 195302 (2009), URL <https://dx.doi.org/10.1088/1751-8113/42/19/195302>.
- [35] A. Mostafazadeh, in *Parity-time Symmetry and Its Applications* (Springer, 2018), pp. 75–121.

- [36] Z. Ahmed, Journal of Physics A: Mathematical and Theoretical **45**, 032004 (2011), URL <https://dx.doi.org/10.1088/1751-8113/45/3/032004>.
- [37] S. Garmon, M. Gianfreda, and N. Hatano, Physical Review A **92**, 022125 (2015).
- [38] P. Molinàs-Mata and P. Molinàs-Mata, Phys. Rev. A **54**, 2060 (1996), URL <https://link.aps.org/doi/10.1103/PhysRevA.54.2060>.
- [39] B. BAGCHI, C. QUESNE, and M. ZNOJIL, Modern Physics Letters A **16**, 2047 (2001), <https://doi.org/10.1142/S0217732301005333>, URL <https://doi.org/10.1142/S0217732301005333>.
- [40] G. S. Japaridze, Journal of Physics A: Mathematical and General **35**, 1709 (2002), URL <https://dx.doi.org/10.1088/0305-4470/35/7/315>.
- [41] J. Mathews, W. N., American Journal of Physics **42**, 214 (1974), ISSN 0002-9505, https://pubs.aip.org/aapt/ajp/article-pdf/42/3/214/11911083/214_1_online.pdf, URL <https://doi.org/10.1119/1.1987650>.
- [42] M. F. Ludovico, J. S. Lim, M. Moskalets, L. Arrachea, and D. Sánchez, Phys. Rev. B **89**, 161306 (2014), URL <https://link.aps.org/doi/10.1103/PhysRevB.89.161306>.
- [43] F. R. T. Arvizu, A. Ortega, and H. Larralde, Physica Scripta **98**, 125015 (2023), URL <https://dx.doi.org/10.1088/1402-4896/ad0c90>.
- [44] B. Wahlstrand, I. I. Yakimenko, and K.-F. Berggren, Phys. Rev. E **89**, 062910 (2014), URL <https://link.aps.org/doi/10.1103/PhysRevE.89.062910>.
- [45] H. Xu, D. Mason, L. Jiang, and J. G. E. Harris, Nature **553**, 80 (2016).
- [46] I. V. Barashenkov, D. A. Zezyulin, and V. V. Konotop, New J. Phys. **18**, 075015 (2016).
- [47] C. M. Bender, Contemporary Physics **46**, 277 (2005), <https://doi.org/10.1080/00107500072632>, URL <https://doi.org/10.1080/00107500072632>.
- [48] C. M. Bender, D. C. Brody, and H. F. Jones, Phys. Rev. Lett. **89**, 270401 (2002), URL <https://link.aps.org/doi/10.1103/PhysRevLett.89.270401>.
- [49] There are other norms used in these systems [20], which has been established by Bagchi [39] and is related to the \mathcal{PT} -symmetric inner product proposed by Bender [47]. Nevertheless, this norm is not necessarily positive, which is troublesome [5]. For other systems analogous to ours, the normalization constant has been calculated using the \mathcal{CPT} inner product [48], which is a positive norm, however, this normalization constant has only been possible to calculate it exactly for discrete systems [37] and only perturbatively for continuous systems [12].