

A UNIFIED STABILITY THEORY FOR CLASSICAL AND MONOTONE MARKOV CHAINS

TAKASHI KAMIHIGASHI AND JOHN STACHURSKI

ABSTRACT. This paper integrates two strands of the literature on stability of general state Markov chains: conventional, total variation based results and more recent order-theoretic results. First we introduce a complete metric over Borel probability measures based on “partial” stochastic dominance. We then show that many conventional results framed in the setting of total variation distance have natural generalizations to the partially ordered setting when this metric is adopted.

Keywords: Total variation, Markov chains, stochastic domination, coupling

MSC classifications Primary: 60J05, 60J99; secondary: 54E50, 06A06.

1. INTRODUCTION

Following the work of Wolfgang Doeblin [6, 7, 8], many classical results from Markov chain theory have built on fundamental connections between total variation distance, Markov chains and couplings. For some models, however, total variation convergence is too strong. In response, researchers have developed an alternative methodology based on monotonicity [10, 35, 1]. In this line of research, transition probabilities are assumed to have a form of monotonicity not required in the classical theory. At the same time, mixing conditions are generally weaker, as is the notion of convergence to the stationary distribution.¹

To give one example, consider a Markov chain $\{X_t\}$ defined by

$$(1) \quad X_{t+1} = \frac{X_t + W_{t+1}}{2}$$

where $\{W_t\}_{t \geq 1}$ is an IID Bernoulli(1/2) random sequence, taking values 0 and 1 with equal probability. For the state space take $S = [0, 1]$. Let $P^t(x, \cdot)$ be the distribution of X_t given $X_0 = x \in S$. Clearly, if X_t is a rational number in S , then so is X_{t+1} .

Date: October 2, 2024.

This paper was written in part while the second author was visiting RIEB at Kobe University. We have benefited from financial support from the Japan Society for the Promotion of Science (KAKENHI 15H05729) and Australian Research Council Discovery Grant DP120100321.

¹Further contributions to this approach can be found in [29, 16, 2, 19]. For some recent extensions and applications in economics see [20].

Similarly, if X_t is irrational, then so is X_{t+1} . Thus, if x and y are rational and irrational respectively, the distributions $P^t(x, \cdot)$ and $P^t(y, \cdot)$ are concentrated on disjoint sets, and hence, when $\|\cdot\|$ is the total variation norm,

$$(2) \quad \|P^t(x, \cdot) - P^t(y, \cdot)\| = 2$$

for all $t \in \mathbb{N}$. Total variation convergence fails for this class of models.

At the same time, the right hand side of (1) is increasing in the current state for each fixed value of the shock W_{t+1} . Moreover, trajectories mix in a monotone sense: A trajectory starting at $X_0 = 0$ can approach 1 with a suitable string of shocks and a trajectory starting at 1 can approach 0. Using these facts one can show using the results in [1], say, that a unique stationary distribution exists and the distribution of X_t converges to it in a complete metric defined over the Borel probability measures that is weaker than total variation convergence.

This is one example where monotone methods can be used to establish some form of stability, despite the fact that the classical conditions based around total variation convergence fail. Conversely, there are many models that the monotone methods developed in [1] and related papers cannot accommodate, while the classical theory based around total variation convergence handles them easily. One example is the simple “inventory” model

$$(3) \quad X_{t+1} = \begin{cases} (X_t - W_{t+1})_+ & \text{if } X_t > 0 \\ (K - W_{t+1})_+ & \text{if } X_t = 0, \end{cases}$$

where $x_+ := \max\{x, 0\}$. Again $\{W_t\}$ is IID. Assume that $\ln W_t$ is standard normal. The state space we take to be $S = [0, K]$. Figure 1 shows a typical trajectory when $K = 100$ and $X_0 = 50$.

On one hand, the monotone methods in [1] cannot be applied here because of a failure of monotonicity with respect to the standard ordering of \mathbb{R} . On the other hand, the classical theory based around total variation convergence is straightforward to apply. For example, one can use Doeblin’s condition (see, e.g., [24], theorem 16.2.3) to show the existence of a unique stationary distribution to which the distribution of X_t converges in total variation, regardless of the distribution of X_0 . In the terminology of [24], the process is uniformly ergodic.

The purpose of this paper is to show that both of these stability results (i.e, the two sets of results concerning the two models (1) and (3)), which were based on two hitherto separate approaches, can be derived from the same theoretical framework. More generally, we construct stability results that encompasses all uniformly ergodic models in the sense of [24] and all monotone models shown to be stable in [1], as well




FIGURE 1. A time series from the inventory model

as extending to other monotone or partially monotone models on state spaces other than \mathbb{R}^n .

We begin our analysis by introducing what is shown to be a complete metric γ on the set of Borel probability measures on a partially ordered Polish space that includes total variation distance, the Kolmogorov uniform distance and the Bhattacharya distance [1, 3] as special cases. We show that many fundamental concepts from conventional Markov chain theory using total variation distance and coupling have direct generalizations to the partially ordered setting when this new metric is adopted. Then, by varying the choice of partial order, we recover key aspects of both classical total variation based stability theory and monotone methods as special cases.²

After preliminaries, we begin with a discussion of “ordered” affinity, which generalizes the usual notion of affinity for measures. The concept of ordered affinity is then used

²There is one additional line of research that deals with Markov models for which the classical conditions of irreducibility and total variation convergence fail. In this line of analysis, irreducibility is replaced by an assumption that the law of motion for the state is itself contracting “on average,” and this contractivity is then passed on to an underlying metric over distributions that conforms in some way to the topology of the state space. See, for example, [4] or [34]. Such results can be used to show stability of our first example, which contracts on average with respect to the usual metric on \mathbb{R} . On the other hand, it cannot be applied to our second (i.e., inventory) example using the same metric, since the law of motion contains a jump. In [1] and [16] one can find many other applications where monotone methods can be used—including the results developed here—while the “contraction on average” conditions of [4] and [34] do not hold. In general, our results should be understood as complements rather than substitutes when compared to average contractions.

to define the total ordered variation metric. Throughout the paper, longer proofs are deferred to the appendix. The conclusion contains suggestions for future work.

2. PRELIMINARIES

Let S be a Polish (i.e., separable and completely metrizable) space, let \mathcal{O} be the open sets, let \mathcal{C} be the closed sets and let \mathcal{B} be the Borel sets. Let \mathcal{M}_s denote the set of all finite signed measures on (S, \mathcal{B}) . In other words, \mathcal{M}_s is all countably additive set functions from \mathcal{B} to \mathbb{R} . Let \mathcal{M} and \mathcal{P} be the finite measures and probability measures in \mathcal{M}_s , respectively. If κ and λ are in \mathcal{M}_s , then $\kappa \leq \lambda$ means that $\kappa(B) \leq \lambda(B)$ for all $B \in \mathcal{B}$. The symbol δ_x denotes the probability measure concentrated on $x \in S$.

Let bS be the set of all bounded \mathcal{B} -measurable functions from S into \mathbb{R} . If $h \in bS$ and $\lambda \in \mathcal{M}_s$, then $\lambda(h) := \int h d\lambda$. For f and g in bS , the statement $f \leq g$ means that $f(x) \leq g(x)$ for all $x \in S$. Let

$$H := \{h \in bS : -1 \leq h \leq 1\} \quad \text{and} \quad H_0 := \{h \in bS : 0 \leq h \leq 1\}.$$

The *total variation norm* of $\lambda \in \mathcal{M}_s$ is $\|\lambda\| := \sup_{h \in H} |\lambda(h)|$. Given μ and ν in \mathcal{P} , a random element (X, Y) taking values in $S \times S$ and defined on a common probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is called a *coupling* of (μ, ν) if $\mu = \mathbb{P} \circ X^{-1}$ and $\nu = \mathbb{P} \circ Y^{-1}$ (i.e., if the distribution of (X, Y) has marginals μ and ν respectively—see, e.g., [21] or [33]). The set of all couplings of (μ, ν) is denoted below by $\mathcal{C}(\mu, \nu)$. A sequence $\{\mu_n\} \subset \mathcal{P}$ converges to $\mu \in \mathcal{P}$ *weakly* if $\mu_n(h) \rightarrow \mu(h)$ as $n \rightarrow \infty$ for all continuous $h \in bS$. In this case we write $\mu_n \xrightarrow{w} \mu$.

Given μ and $\nu \in \mathcal{M}$, their measure theoretic *infimum* $\mu \wedge \nu$ is the largest element of \mathcal{M} dominated by both μ and ν . It can be defined by taking f and g to be densities of μ and ν respectively under the dominating measure $\lambda := \mu + \nu$ and defining $\mu \wedge \nu$ by $(\mu \wedge \nu)(B) := \int_B \min\{f(x), g(x)\} \lambda(dx)$ for all $B \in \mathcal{B}$. The total variation distance between μ and ν is related to $\mu \wedge \nu$ via $\|\mu - \nu\| = \|\mu\| + \|\nu\| - 2\|\mu \wedge \nu\|$. See, for example, [28]. For probability measures we also have

$$(4) \quad \sup_{B \in \mathcal{B}} \{\mu(B) - \nu(B)\} = \sup_{B \in \mathcal{B}} |\mu(B) - \nu(B)| = \|\mu - \nu\|/2.$$

The *affinity* between two measures μ, ν in \mathcal{M} is the value $\alpha(\mu, \nu) := (\mu \wedge \nu)(S)$. The following properties are elementary:

Lemma 2.1. *For all $(\mu, \nu) \in \mathcal{M} \times \mathcal{M}$ we have*

- (a) $0 \leq \alpha(\mu, \nu) \leq \min\{\mu(S), \nu(S)\}$
- (b) $\alpha(\mu, \nu) = \mu(S) = \nu(S)$ if and only if $\mu = \nu$.
- (c) $\alpha(c\mu, c\nu) = c\alpha(\mu, \nu)$ for all $c \geq 0$.

There are several other common representations of affinity. For example, when μ and ν are both probability measures, we have

$$(5) \quad \alpha(\mu, \nu) = 1 - \sup_{B \in \mathcal{B}} |\mu(B) - \nu(B)| = \max_{(X, Y) \in \mathcal{C}(\mu, \nu)} \mathbb{P}\{X = Y\}.$$

(See, e.g., [28, 21].) The second equality in (5) states that, if $(X, Y) \in \mathcal{C}(\mu, \nu)$, then $\mathbb{P}\{X = Y\} \leq \alpha(\mu, \nu)$, and, moreover, there exists a $(X, Y) \in \mathcal{C}(\mu, \nu)$ such that equality is attained. Any such coupling is called a *maximal* or *gamma* coupling. See theorem 5.2 of [21]. From (4) and (5) we obtain

$$(6) \quad \|\mu - \nu\| = 2(1 - \alpha(\mu, \nu)).$$

3. ORDERED AFFINITY

We next introduce a generalization of affinity when S has a partial order. We investigate its properties in detail, since both our metric and the stability theory presented below rely on this concept.

3.1. Preliminaries. As before, let S be a Polish space. A closed partial order \preceq on S is a partial order \preceq such that its graph

$$\mathbb{G} := \{(x, y) \in S \times S : x \preceq y\}$$

is closed in the product topology. In the sequel, a *partially ordered Polish space* is any such pair (S, \preceq) , where S is nonempty and Polish, and \preceq is a closed partial order on S . When no confusion arises, we denote it simply by S .³

For such a space S , we call $I \subset S$ *increasing* if $x \in I$ and $x \preceq y$ implies $y \in I$. We call $h: S \rightarrow \mathbb{R}$ *increasing* if $x \preceq y$ implies $h(x) \leq h(y)$. We let $i\mathcal{B}$, $i\mathcal{O}$ and $i\mathcal{C}$ denote the increasing Borel, open and closed sets, respectively, while ibS is the increasing functions in bS . In addition,

- $iH := H \cap ibS = \{h \in ibS : -1 \leq h \leq 1\}$ and
- $iH_0 := H_0 \cap ibS = \{h \in ibS : 0 \leq h \leq 1\}$.

If $B \in \mathcal{B}$, then $i(B)$ is all $y \in S$ such that $x \preceq y$ for some $x \in B$, while $d(B)$ is all $y \in S$ such that $y \preceq x$ for some $x \in B$. Given μ and ν in \mathcal{M} , we say that μ is *stochastically dominated* by ν and write $\mu \preceq_{sd} \nu$ if $\mu(S) = \nu(S)$ and $\mu(I) \leq \nu(I)$ for all $I \in i\mathcal{B}$. Equivalently, $\mu(S) = \nu(S)$ and $\mu(h) \leq \nu(h)$ for all h in iH or iH_0 . See [17].

³The partial order is assumed to be closed in the theory developed below because we build a metric over Borel probability measures that depends on this partial order and closedness of the order is required to show that the metric is complete.

One important partial order on S is the *identity order*, where $x \preceq y$ if and only if $x = y$. Then $i\mathcal{B} = \mathcal{B}$, $ibS = bS$, $iH = H$, $iH_0 = H_0$ and $\mu \preceq_{sd} \nu$ if and only if $\mu = \nu$.

Remark 3.1. Since S is a partially ordered Polish space, for any μ, ν in \mathcal{P} we have $\mu = \nu$ whenever $\mu(C) = \nu(C)$ for all $C \in i\mathcal{C}$, or, equivalently, $\mu(h) = \nu(h)$ for all continuous $h \in ibS$. See [17, lemma 1]. Hence $\mu \preceq_{sd} \nu$ and $\nu \preceq_{sd} \mu$ imply $\mu = \nu$.

Lemma 3.1. *If $\lambda \in \mathcal{M}_s$, then $\sup_{I \in i\mathcal{B}} \lambda(I) = \sup_{h \in iH_0} \lambda(h)$ and*

$$(7) \quad \sup_{h \in iH} |\lambda(h)| = \max \left\{ \sup_{h \in iH} \lambda(h), \sup_{h \in iH} (-\lambda)(h) \right\}.$$

The proof is in the appendix. One can easily check that

$$(8) \quad \lambda \in \mathcal{M}_s \text{ and } \lambda(S) = 0 \quad \implies \quad \sup_{h \in iH} \lambda(h) = 2 \sup_{h \in iH_0} \lambda(h).$$

3.2. Definition of Ordered Affinity. For each pair $(\mu, \nu) \in \mathcal{M} \times \mathcal{M}$, let

$$\Phi(\mu, \nu) := \{(\mu', \nu') \in \mathcal{M} \times \mathcal{M} : \mu' \leq \mu, \nu' \leq \nu, \mu' \preceq_{sd} \nu'\}.$$

We call $\Phi(\mu, \nu)$ the set of *ordered component pairs* for (μ, ν) . Here “ordered” means ordered by stochastic dominance. The set of ordered component pairs is always nonempty, since $(\mu \wedge \nu, \mu \wedge \nu)$ is an element of $\Phi(\mu, \nu)$.

Example 3.1. If μ and ν are two measures satisfying $\mu \preceq_{sd} \nu$, then $(\mu, \nu) \in \Phi(\mu, \nu)$.

Example 3.2. Given Bernoulli distributions $\mu = (\delta_1 + \delta_2)/2$ and $\nu = (\delta_0 + \delta_1)/2$, we have $(\mu', \nu') \in \Phi(\mu, \nu)$ when $\mu' = \nu' = \delta_1/2$.

We call an ordered component pair $(\mu', \nu') \in \Phi(\mu, \nu)$ a *maximal ordered component pair* if it has greater mass than all others; that is, if

$$\mu''(S) \leq \mu'(S) \quad \text{for all } (\mu'', \nu'') \in \Phi(\mu, \nu).$$

(We can restate this by replacing $\mu'(S)$ and $\mu''(S)$ with $\nu'(S)$ and $\nu''(S)$ respectively, since the mass of ordered component pairs is equal by the definition of stochastic dominance.) We let $\Phi^*(\mu, \nu)$ denote the set of maximal ordered component pairs for (μ, ν) . Thus, if

$$(9) \quad \alpha_O(\mu, \nu) := \sup\{\mu'(S) : (\mu', \nu') \in \Phi(\mu, \nu)\}.$$

then

$$\Phi^*(\mu, \nu) = \{(\mu', \nu') \in \Phi(\mu, \nu) : \mu'(S) = \alpha_O(\mu, \nu)\}.$$

Using the Polish space assumption, one can show that maximal ordered component pairs always exist:

Proposition 3.1. *The set $\Phi^*(\mu, \nu)$ is nonempty for all $(\mu, \nu) \in \mathcal{M} \times \mathcal{M}$.*

Proof. Fix $(\mu, \nu) \in \mathcal{M} \times \mathcal{M}$ and let $s := \alpha_O(\mu, \nu)$. From the definition, we can take sequences $\{\mu'_n\}$ and $\{\nu'_n\}$ in \mathcal{M} such that $(\mu'_n, \nu'_n) \in \Phi(\mu, \nu)$ for all $n \in \mathbb{N}$ and $\mu'_n(S) \uparrow s$. Since $\mu'_n \leq \mu$ and $\nu'_n \leq \nu$ for all $n \in \mathbb{N}$, Prohorov's theorem [11, theorem 11.5.4] implies that these sequences have convergent subsequences with $\mu'_{n_k} \xrightarrow{w} \mu'$ and $\nu'_{n_k} \xrightarrow{w} \nu'$ for some $\mu', \nu' \in \mathcal{M}$. We claim that (μ', ν') is a maximal ordered component pair.

Since $\mu'_{n_k} \xrightarrow{w} \mu'$ and $\mu'_n \leq \mu$ for all $n \in \mathbb{N}$, theorem 1.5.5 of [15] implies that for any Borel set B , we have $\mu'_{n_k}(B) \rightarrow \mu'(B)$ in \mathbb{R} . Hence $\mu'(B) \leq \mu(B)$ and, in particular, $\mu' \leq \mu$. An analogous argument gives $\nu' \leq \nu$. Moreover, the definition of $\Phi(\mu, \nu)$ and stochastic dominance imply that $\mu'_n(S) = \nu'_n(S)$ for all $n \in \mathbb{N}$, and therefore $\mu'(S) = \nu'(S)$. Also, for any $I \in i\mathcal{B}$, the fact that $\mu'_n(I) \leq \nu'_n(I)$ for all $n \in \mathbb{N}$ gives us $\mu'(I) \leq \nu'(I)$. Thus, $\mu' \preceq_{sd} \nu'$. Finally, $\mu'(S) = s$, since $\mu'_n(S) \uparrow s$. Hence (μ', ν') lies in $\Phi^*(\mu, \nu)$. \square

The value $\alpha_O(\mu, \nu)$ defined in (9) gives the mass of the maximal ordered component pair. We call it the *ordered affinity* from μ to ν . On an intuitive level, we can think of $\alpha_O(\mu, \nu)$ as the “degree” to which μ is dominated by ν in the sense of stochastic dominance. Since $(\mu \wedge \nu, \mu \wedge \nu)$ is an ordered component pair for (μ, ν) , we have

$$(10) \quad 0 \leq \alpha(\mu, \nu) \leq \alpha_O(\mu, \nu),$$

where $\alpha(\mu, \nu)$ is the standard affinity defined in section 2. In fact $\alpha_O(\mu, \nu)$ generalizes the standard the notion of affinity by extending it to arbitrary partial orders, as shown in the next lemma.

Lemma 3.2. *If \preceq is the identity order, then $\alpha_O = \alpha$ on $\mathcal{M} \times \mathcal{M}$.*

Proof. Fix $(\mu, \nu) \in \mathcal{M} \times \mathcal{M}$ and let \preceq be the identity order ($x \preceq y$ iff $x = y$). Then \preceq_{sd} also corresponds to equality, from which it follows that the supremum in (9) is attained by $\mu \wedge \nu$. Hence $\alpha_O(\mu, \nu) = \alpha(\mu, \nu)$. \square

3.3. Properties of Ordered Affinity. Let's list some elementary properties of α_O . The following list should be compared with lemma 2.1. It shows that analogous results hold for α_O as hold for α . (Lemma 2.1 is in fact a special case of lemma 3.3 with the partial order set to the identity order.)

Lemma 3.3. *For all $(\mu, \nu) \in \mathcal{M} \times \mathcal{M}$, we have*

- (a) $0 \leq \alpha_O(\mu, \nu) \leq \min\{\mu(S), \nu(S)\}$,
- (b) $\alpha_O(\mu, \nu) = \mu(S) = \nu(S)$ if and only if $\mu \preceq_{sd} \nu$, and
- (c) $c\alpha_O(\mu, \nu) = \alpha_O(c\mu, c\nu)$ whenever $c \geq 0$.

Proof. Fix $(\mu, \nu) \in \mathcal{M} \times \mathcal{M}$. Claim (a) follows directly from the definitions. Regarding claim (b), suppose first that $\mu \preceq_{sd} \nu$. Then $(\mu, \nu) \in \Phi(\mu, \nu)$ and hence $\alpha_O(\mu, \nu) = \mu(S)$. Conversely, if $\alpha_O(\mu, \nu) = \mu(S)$, then, since the only component $\mu' \leq \mu$ with $\mu'(S) = \mu(S)$ is μ itself, we must have $(\mu, \nu') \in \Phi(\mu, \nu')$ for some $\nu' \leq \nu$ with $\mu \preceq_{sd} \nu'$. But then $\mu(I) \leq \nu'(I) \leq \nu(I)$ for any $I \in i\mathcal{B}$. Hence $\mu \preceq_{sd} \nu$.

Claim (c) is trivial if $c = 0$, so suppose instead that $c > 0$. Fix $(\mu', \nu') \in \Phi(\mu, \nu)$ such that $\alpha_O(\mu, \nu) = \mu'(S)$. It is clear that $(c\mu', c\nu') \in \Phi(c\mu, c\nu)$, implying that

$$(11) \quad c\alpha_O(\mu, \nu) = c\mu'(S) \leq \alpha_O(c\mu, c\nu).$$

For reverse inequality, we can apply (11) again to get

$$\alpha_O(c\mu, c\nu) = c(1/c)\alpha_O(c\mu, c\nu) \leq c\alpha_O(\mu, \nu). \quad \square$$

3.4. Equivalent Representations. In (5) we noted that the affinity between two measures has several alternative representations. In our setting these results generalize as follows:

Theorem 3.1. *For all $(\mu, \nu) \in \mathcal{P} \times \mathcal{P}$, we have*

$$(12) \quad \alpha_O(\mu, \nu) = 1 - \sup_{I \in i\mathcal{B}} \{\mu(I) - \nu(I)\} = \max_{(X, Y) \in \mathcal{C}(\mu, \nu)} \mathbb{P}\{X \preceq Y\}.$$

Evidently (5) is a special case of (12) because (12) reduces to (5) when \preceq is set to equality. For example, when \preceq is equality,

$$\sup_{I \in i\mathcal{B}} \{\mu(I) - \nu(I)\} = \sup_{B \in \mathcal{B}} \{\mu(B) - \nu(B)\} = \sup_{B \in \mathcal{B}} |\mu(B) - \nu(B)|.$$

where the last step is from (4). Note also that, as shown in the proof of theorem 3.1, the supremum can also be written in terms of the open increasing sets $i\mathcal{O}$ or the closed decreasing sets $d\mathcal{C}$. In particular,

$$\sup_{I \in i\mathcal{B}} \{\mu(I) - \nu(I)\} = \sup_{I \in i\mathcal{O}} \{\mu(I) - \nu(I)\} = \sup_{D \in d\mathcal{C}} \{\nu(D) - \mu(D)\}.$$

One of the assertions of theorem 3.1 is the existence of a coupling $(X, Y) \in \mathcal{C}(\mu, \nu)$ attaining $\mathbb{P}\{X \preceq Y\} = \alpha_O(\mu, \nu)$. Let us refer to any such coupling as an *order maximal* coupling for (μ, ν) .

Example 3.3. For $(x, y) \in S \times S$, we have

$$\alpha_O(\delta_x, \delta_y) = \mathbb{1}\{x \preceq y\} = \mathbb{1}_{\mathbb{G}}(x, y),$$

as can easily be verified from the definition or either of the alternative representations in (12). The map $(x, y) \mapsto \mathbb{1}_{\mathbb{G}}(x, y)$ is measurable due to the Polish assumption. As a result, for any $(X, Y) \in \mathcal{C}(\mu, \nu)$ we have

$$\mathbb{E} \alpha_O(\delta_X, \delta_Y) = \mathbb{P}\{X \preceq Y\} \leq \alpha_O(\mu, \nu),$$

with equality when (X, Y) is an order maximal coupling.

3.5. Comments on Theorem 3.1. The existence of an order maximal coupling shown in theorem 3.1 implies two well-known results that are usually treated separately. One is the Nachbin–Strassen theorem (see, e.g., thm. 1 of [18] or ch. IV of [21]), which states the existence of a coupling $(X, Y) \in \mathcal{C}(\mu, \nu)$ attaining $\mathbb{P}\{X \preceq Y\} = 1$ whenever $\mu \preceq_{sd} \nu$. The existence of an order maximal coupling for each (μ, ν) in $\mathcal{P} \times \mathcal{P}$ implies this statement, since, under the hypothesis that $\mu \preceq_{sd} \nu$, we also have $\alpha_O(\mu, \nu) = 1$. Hence any order maximal coupling satisfies $\mathbb{P}\{X \preceq Y\} = 1$.

The other familiar result implied by the existence of an order maximal coupling is the existence of a maximal coupling in the standard sense (see the discussion of maximal couplings after (5) and the result on p. 19 of [21]). Indeed, if we take \preceq to be the identity order, then (12) reduces to (5), as already discussed.

4. TOTAL ORDERED VARIATION

Let S be a partially ordered Polish space. Consider the function on $\mathcal{P} \times \mathcal{P}$ given by

$$(13) \quad \gamma(\mu, \nu) := 2 - \alpha_O(\mu, \nu) - \alpha_O(\nu, \mu).$$

We call $\gamma(\mu, \nu)$ the *total ordered variation distance* between μ and ν . The natural comparison is with (6), which renders the same value if α_O is replaced by α . In particular, when \preceq is equality, ordered affinity reduces to affinity, and total ordered variation distance reduces to total variation distance. Since ordered affinities dominate affinities (see (10)), we have $\gamma(\mu, \nu) \leq \|\mu - \nu\|$ for all $(\mu, \nu) \in \mathcal{P} \times \mathcal{P}$.

Other, equivalent, representations of γ are available. For example, in view of (12), for any $(\mu, \nu) \in \mathcal{P} \times \mathcal{P}$ we have

$$(14) \quad \gamma(\mu, \nu) = \sup_{I \in i\mathcal{B}} (\mu - \nu)(I) + \sup_{I \in i\mathcal{B}} (\nu - \mu)(I),$$

By combining lemma 3.1 and (8), we also have

$$(15) \quad 2\gamma(\mu, \nu) = \sup_{h \in iH} (\mu - \nu)(h) + \sup_{h \in iH} (\nu - \mu)(h).$$

It is straightforward to show that

$$(16) \quad \sup_{I \in i\mathcal{B}} |\mu(I) - \nu(I)| \leq \gamma(\mu, \nu) \quad \text{and} \quad \sup_{D \in d\mathcal{B}} |\mu(D) - \nu(D)| \leq \gamma(\mu, \nu).$$

Lemma 4.1. *The function γ is a metric on \mathcal{P} .*

Proof. The claim that γ is a metric follows in a straightforward way from the definition or the alternative representation (14). For example, the triangle inequality is easy to verify using (14). Also, $\gamma(\mu, \nu) = 0$ implies $\mu = \nu$ by (14) and remark 3.1. \square

4.1. Connection to Other Modes of Convergence. As well as total variation, the metric γ is closely related to the so-called *Bhattacharya metric*, which is given by

$$(17) \quad \beta(\mu, \nu) := \sup_{h \in iH} |\mu(h) - \nu(h)|.$$

See [1, 3]. (In [3] the metric is defined by taking the supremum over iH_0 rather than iH , but the two definitions differ only by a positive scalar.) The Bhattacharya metric can be thought of as an alternative way to generalize total variation distance, in the sense that, like γ , the metric β reduces to total variation distance when \preceq is the identity order (since iH equals H under this order). From (7) we have

$$(18) \quad \frac{1}{2} \left[\sup_{h \in iH} \lambda(h) + \sup_{h \in iH} (-\lambda)(h) \right] \leq \sup_{h \in iH} |\lambda(h)| \leq \sup_{h \in iH} \lambda(h) + \sup_{h \in iH} (-\lambda)(h),$$

and from this and (15) we have

$$(19) \quad \gamma(\mu, \nu) \leq \beta(\mu, \nu) \leq 2\gamma(\mu, \nu).$$

Hence β and γ are equivalent metrics.

The metric γ is also connected to the Wasserstein metric [12, 13]. If ρ metrizes the topology on S , then the Wasserstein distance between probability measures μ and ν is

$$w(\mu, \nu) := \inf_{(X,Y) \in \mathcal{C}(\mu,\nu)} \mathbb{E} \rho(X, Y).$$

The total ordered variation metric can be compared as follows. Consider the "directed semimetric" $\hat{\rho}(x, y) := \mathbb{1}\{x \not\leq y\}$. In view of (12) we have

$$\gamma(\mu, \nu) = \inf_{(X,Y) \in \mathcal{C}(\mu,\nu)} \mathbb{E} \hat{\rho}(X, Y) + \inf_{(X,Y) \in \mathcal{C}(\mu,\nu)} \mathbb{E} \hat{\rho}(Y, X).$$

Thus, $\gamma(\mu, \nu)$ is found by summing two partial, "directed Wasserstein deviations." Summing the two directed differences from opposite directions yields a metric.

Proposition 4.1. *If $\{\mu_n\}_{n \geq 0} \subset \mathcal{P}$ is tight and $\gamma(\mu_n, \mu_0) \rightarrow 0$, then $\mu_n \xrightarrow{w} \mu_0$.*

Proof. Let $\{\mu_n\}$ and $\mu := \mu_0$ satisfy the conditions of the proposition. Take any subsequence of $\{\mu_n\}$ and observe that by Prohorov's theorem, this subsequence has a subsubsequence converging weakly to some $\nu \in \mathcal{P}$. Along this subsubsequence, for any continuous $h \in ibS$ we have both $\mu_n(h) \rightarrow \mu(h)$ and $\mu_n(h) \rightarrow \nu(h)$. This is sufficient for $\nu = \mu$ by remark 3.1. Thus, every subsequence of $\{\mu_n\}$ has a subsubsequence converging weakly to μ , and hence so does the entire sequence. \square

4.2. Completeness. To obtain completeness of (\mathcal{P}, γ) , we adopt the following additional assumption, which is satisfied if, say, compact sets are order bounded (i.e., lie in order intervals) and order intervals are compact. (For example, \mathbb{R}^n with the usual pointwise partial order has this property.)

Assumption 4.1. If $K \subset S$ is compact, then $i(K) \cap d(K)$ is also compact.

Theorem 4.1. *If assumption 4.1 holds, then (\mathcal{P}, γ) is complete.*

Remark 4.1. In [3] it was shown that β is a complete metric when $S = \mathbb{R}^n$. Due to equivalence of the metrics, theorem 4.1 extends this result to partially ordered Polish spaces where assumption 4.1 is satisfied.

5. APPLICATIONS

In this section we show that many results in classical and monotone Markov chain theory, hitherto treated separately, can be derived from the same set of results based around total ordered variation and ordered affinity.

Regarding notation, if $\{S_i\}$ are partially ordered Polish spaces over $i = 0, 1, 2, \dots$, we often use a common symbol \preceq for the partial order on any of these spaces. On products of these spaces we use the product topology and pointwise partial order. Once again, the symbol \preceq is used for the partial order. For example, if (x_0, x_1) and (y_0, y_1) are points in $S_0 \times S_1$, then $(x_0, x_1) \preceq (y_0, y_1)$ means that $x_0 \preceq y_0$ and $x_1 \preceq y_1$.

A function $P: (S_0, \mathcal{B}_1) \rightarrow [0, 1]$ is called a *Markov kernel from S_0 to S_1* if $x \mapsto P(x, B)$ is \mathcal{B}_0 -measurable for each $B \in \mathcal{B}_1$ and $B \mapsto P(x, B)$ is in \mathcal{P}_1 for all $x \in S_0$. If $S_0 = S_1 = S$, we will call P a *Markov kernel on S* , or just a Markov kernel. Following standard conventions (see, e.g., [24]), for any Markov kernel P from S_0 to S_1 , any $h \in bS_1$ and $\mu \in \mathcal{P}_0$, we define $\mu P \in \mathcal{P}_1$ and $Ph \in bS_0$ via

$$(\mu P)(B) = \int P(x, B)\mu(dx) \quad \text{and} \quad (Ph)(x) = \int h(y)P(x, dy).$$

Also, $\mu \otimes P$ denotes the joint distribution on $S_0 \times S_1$ defined by

$$(\mu \otimes P)(A \times B) = \int_A P(x, B)\mu(dx).$$

To simplify notation, we use P_x to represent the measure $\delta_x P = P(x, \cdot)$. P^m is the m -th composition of P with itself.

5.1. Order Affinity and Monotone Markov Kernels. Let S be a Polish space partially ordered by \preceq . A Markov kernel P is called *monotone* if $Ph \in ibS_0$ whenever $h \in ibS_1$. An equivalent condition is that $\mu P \preceq_{sd} \nu P$ whenever $\mu \preceq_{sd} \nu$; or just $P(x, \cdot) \preceq_{sd} P(y, \cdot)$ whenever $x \preceq y$. It is well-known (see, e.g., proposition 1 of [18]) that if $\mu \preceq_{sd} \nu$ and P is monotone, then $\mu \otimes P \preceq_{sd} \nu \otimes P$. Note that, when \preceq is the identity order, every Markov kernel is monotone.

Lemma 5.1. *If P is a monotone Markov kernel from S_0 to S_1 and μ, μ', ν and ν' are probabilities in \mathcal{P}_0 , then*

$$\mu' \preceq_{sd} \mu \text{ and } \nu \preceq_{sd} \nu' \implies \alpha_O(\mu P, \nu P) \leq \alpha_O(\mu' P, \nu' P).$$

Proof. Let P, μ, μ', ν and ν' have the stated properties. In view of the equivalently representation in (12), the claim will be established if

$$\sup_{I \in i\mathcal{B}} \{(\mu P)(I) - (\nu P)(I)\} \geq \sup_{I \in i\mathcal{B}} \{(\mu' P)(I) - (\nu' P)(I)\}.$$

This holds by the monotonicity of P and the order of μ, μ', ν and ν' . \square

Lemma 5.2. *If P is a monotone Markov kernel from S_0 to S_1 , then*

$$\alpha_O(\mu P, \nu P) \geq \alpha_O(\mu, \nu) \text{ for any } \mu, \nu \text{ in } \mathcal{P}_0.$$

Proof. Fix μ, ν in \mathcal{P}_0 and let $(\hat{\mu}, \hat{\nu})$ be a maximal ordered component pair for (μ, ν) . From monotonicity of P and the fact the Markov kernels preserve the mass of measures, it is clear that $(\hat{\mu}P, \hat{\nu}P)$ is an ordered component pair for $(\mu P, \nu P)$. Hence

$$\alpha_O(\mu P, \nu P) \geq (\hat{\mu}P)(S) = \hat{\mu}(S) = \alpha_O(\mu, \nu). \quad \square$$

On the other hand, for the joint distribution, the ordered affinity of the initial pair is preserved.

Lemma 5.3. *If P is a monotone Markov kernel from S_0 to S_1 , then*

$$\alpha_O(\mu \otimes P, \nu \otimes P) = \alpha_O(\mu, \nu). \text{ for any } \mu, \nu \text{ in } \mathcal{P}_0.$$

Proof. Fix μ, ν in \mathcal{P}_0 and let (X_0, X_1) and (Y_0, Y_1) be random pairs with distributions $\mu \otimes P$ and $\nu \otimes P$ respectively. We have

$$\mathbb{P}\{(X_0, X_1) \preceq (Y_0, Y_1)\} \leq \mathbb{P}\{X_0 \preceq Y_0\} \leq \alpha_O(\mu, \nu).$$

Taking the supremum over all couplings in $\mathcal{C}(\mu \otimes P, \nu \otimes P)$ shows that $\alpha_O(\mu \otimes P, \nu \otimes P)$ is dominated by $\alpha_O(\mu, \nu)$.

To see the reverse inequality, let $(\hat{\mu}, \hat{\nu})$ be a maximal ordered component pair for (μ, ν) . Monotonicity of P now gives $\hat{\mu} \otimes P \preceq_{sd} \hat{\nu} \otimes P$. Using this and the fact the

Markov kernels preserve the mass of measures, we see that $(\hat{\mu} \otimes P, \hat{\nu} \otimes P)$ is an ordered component pair for $(\mu \otimes P, \nu \otimes P)$. Hence

$$\alpha_O(\mu P, \nu P) \geq (\hat{\mu} \otimes P)(S_0 \times S_1) = \hat{\mu}(S_0) = \alpha_O(\mu, \nu). \quad \square$$

5.2. Monotone Markov Chains. Given $\mu \in \mathcal{P}$ and Markov kernel P on S , a stochastic process $\{X_t\}_{t \geq 0}$ taking values in $S^\infty := \times_{t=0}^\infty S$ will be called a Markov chain with initial distribution μ and kernel P if the distribution of $\{X_t\}$ on S^∞ is

$$\mathbf{Q}_\mu := \mu \otimes P \otimes P \otimes P \otimes \dots$$

(The meaning of the right hand side is clarified in, e.g., §III.8 of [21], p. 903 of [18], §3.4 of [24].) If P is a monotone Markov kernel, then $(x, B) \mapsto \mathbf{Q}_x(B) := \mathbf{Q}_{\delta_x}(B)$ is a monotone Markov kernel from S to S^∞ . See propositions 1 and 2 of [18].

There are various useful results about representations of Markov chains that are ordered almost surely. One is that, if the initial conditions satisfy $\mu \preceq_{sd} \nu$ and P is a monotone Markov kernel, then we can find Markov chains $\{X_t\}$ and $\{Y_t\}$ with initial distributions μ and ν and kernel P such that $X_t \preceq Y_t$ for all t almost surely. (See, e.g., theorem 2 of [18].) This result can be generalized beyond the case where μ and ν are stochastically ordered, using the results presented above. For example, let μ and ν be arbitrary initial distributions and let P be monotone, so that \mathbf{Q}_x is likewise monotone. By lemma 5.3 we have

$$\alpha_O(\mathbf{Q}_\mu, \mathbf{Q}_\nu) = \alpha_O(\mu \otimes \mathbf{Q}_x, \nu \otimes \mathbf{Q}_x) = \alpha_O(\mu, \nu).$$

In other words, the ordered affinity of the entire processes is given by the ordered affinity of the initial distributions. It now follows from theorem 3.1 that there exist Markov chains $\{X_t\}$ and $\{Y_t\}$ with initial distributions μ and ν and Markov kernel P such that

$$\mathbb{P}\{X_t \preceq Y_t, \forall t \geq 0\} = \alpha_O(\mu, \nu).$$

The standard result is a special case, since $\mu \preceq_{sd} \nu$ implies $\alpha_O(\mu, \nu) = 1$, and hence the sequences are ordered almost surely.

5.3. Nonexpansiveness. It is well-known that every Markov kernel is nonexpansive with respect to the total variation norm, so that

$$(20) \quad \|\mu P - \nu P\| \leq \|\mu - \nu\| \quad \text{for all } (\mu, \nu) \in \mathcal{P} \times \mathcal{P}.$$

An analogous result is true for γ when P is monotone. That is,

$$(21) \quad \gamma(\mu P, \nu P) \leq \gamma(\mu, \nu) \quad \text{for all } (\mu, \nu) \in \mathcal{P} \times \mathcal{P}.$$

The bound (21) follows directly from lemma 5.2. Evidently (20) be recovered from (21) by setting \preceq to equality.

Nonexpansiveness is interesting partly in its own right (we apply it in proofs below) and partly because it suggests that, with some additional assumptions, we can strengthen it to contractiveness. We expand on this idea below.

5.4. An Order Coupling Bound for Markov Chains. Doeblin [6] established and exploited the coupling inequality

$$(22) \quad \|\mu_t - \nu_t\| \leq 2\mathbb{P}\{X_j \neq Y_j \text{ for any } j \leq t\}$$

where μ_t and ν_t are the time t distributions of Markov chains $\{X_t\}$ and $\{Y_t\}$ generated by common Markov kernel P . Even when the state space is uncountable, the right hand side of (22) can often be shown to converge to zero by manipulating the joint distribution of (X_j, Y_j) to increase the chance of a meeting [9, 14, 26, 27, 33, 21, 25, 30, 24].

Consider the following generalization: given a monotone Markov kernel P on S and arbitrary $\mu, \nu \in \mathcal{P}$, we can construct Markov chains $\{X_t\}$ and $\{Y_t\}$ with common kernel P and respective initial conditions μ and ν such that

$$(23) \quad \gamma(\mu_t, \nu_t) \leq \mathbb{P}\{X_j \not\leq Y_j \text{ for any } j \leq t\} + \mathbb{P}\{Y_j \not\leq X_j \text{ for any } j \leq t\}.$$

This generalizes (22) because both the right and left hand side of (23) reduces to (22) if we take \leq to be equality.

To prove (23), we need only show that

$$(24) \quad 1 - \alpha_O(\mu P^t, \nu P^t) \leq \mathbb{P}\{X_j \not\leq Y_j \text{ for any } j \leq t\},$$

since, with (24) is established, we can reverse the roles of $\{X_t\}$ and $\{Y_t\}$ in (24) to obtain $1 - \alpha_O(\nu P^t, \mu P^t) \leq \mathbb{P}\{Y_j \not\leq X_j \text{ for any } j \leq t\}$ and then add this inequality to (24) to produce (23).

If $\{X_t\}$ and $\{Y_t\}$ are Markov chains with kernel P and initial conditions μ and ν , then (12) yields $\alpha_O(\mu P^t, \nu P^t) \geq \mathbb{P}\{X_t \leq Y_t\}$. Therefore, we need only construct such chains with the additional property that

$$(25) \quad \mathbb{P}\{X_j \not\leq Y_j \text{ for any } j \leq t\} = \mathbb{P}\{X_t \not\leq Y_t\}.$$

Intuitively we can do so by using a “conditional” version of the Nachbin–Strassen theorem, producing chains that, once ordered, remain ordered almost surely. This can be formalized as follows: By [22, theorem 2.3], there exists a Markov kernel M on $S \times S$ such that \mathbb{G} is absorbing for M (i.e., $M((x, y), \mathbb{G}) = 1$ for all (x, y) in \mathbb{G}),

$$P(x, A) = M((x, y), A \times S) \quad \text{and} \quad P(y, B) = M((x, y), S \times B)$$

for all $(x, y) \in S \times S$ and all $A, B \in \mathcal{B}$. Given M , let η be a distribution on $S \times S$ with marginals μ and ν , let $\mathbf{Q}_\eta := \eta \otimes M \otimes M \otimes \dots$ be the induced joint distribution, and let $\{(X_t, Y_t)\}$ have distribution \mathbf{Q}_η on $(S \times S)^\infty$. By construction, X_t has distribution

μP^t and Y_t has distribution νP^t . Moreover, (25) is valid because \mathbb{G} is absorbing for M , and hence $\mathbb{P}\{(X_j, Y_j) \notin \mathbb{G} \text{ for any } j \leq t\} = \mathbb{P}\{(X_t, Y_t) \notin \mathbb{G}\}$.

5.5. Uniform Ergodicity. Let S be a partially ordered Polish space satisfying assumption 4.1, and let P be a monotone Markov kernel on S . A distribution π is called *stationary* for P if $\pi P = \pi$. Consider the value

$$(26) \quad \sigma(P) := \inf_{(x,y) \in S \times S} \alpha_O(P_x, P_y),$$

which can be understood as an order-theoretic extension of the Markov–Dobrushin coefficient of ergodicity [5, 31]. It reduces to the usual notion when \preceq is equality.

Theorem 5.1. *If P is monotone, then*

$$(27) \quad \gamma(\mu P, \nu P) \leq (1 - \sigma(P)) \gamma(\mu, \nu) \quad \text{for all } (\mu, \nu) \in \mathcal{P} \times \mathcal{P}.$$

Thus, strict positivity of $\sigma(P)$ implies that $\mu \mapsto \mu P$ is a contraction map on (\mathcal{P}, γ) . Moreover, in many settings, the bound in (27) cannot be improved upon. For example,

Lemma 5.4. *If P is monotone, S is not a singleton and any x, y in S have a lower bound in S , then*

$$(28) \quad \forall \xi > \sigma(P), \exists \mu, \nu \in \mathcal{P} \text{ s.t. } \gamma(\mu P, \nu P) > (1 - \xi) \gamma(\mu, \nu).$$

The significance of theorem 5.1 is summarized in the next corollary.

Corollary 5.1. *Let P be monotone and let S satisfy assumption 4.1. If there exists an $m \in \mathbb{N}$ such that $\sigma(P^m) > 0$, then P has a unique stationary distribution π in \mathcal{P} , and*

$$(29) \quad \gamma(\mu P^t, \pi) \leq (1 - \sigma(P^m))^{\lfloor t/m \rfloor} \gamma(\mu, \pi) \quad \text{for all } \mu \in \mathcal{P}, t \geq 0.$$

Here $\lfloor x \rfloor$ is the largest $n \in \mathbb{N}$ with $n \leq x$.

Proof. Let P and μ be as in the statement of the theorem. The existence of a fixed point of $\mu \mapsto \mu P$, and hence a stationary distribution $\pi \in \mathcal{P}$, follows from theorem 5.1 applied to P^m , Banach’s contraction mapping theorem, and the completeness of (\mathcal{P}, γ) shown in theorem 4.1. The bound in (29) follows from (27) applied to P^m and the nonexpansiveness of P in the metric γ (see (21)). \square

5.6. Applications from the Introduction. Let us see how corollary 5.1 can be used to show stability of the two models discussed in the introduction, beginning with the monotone model in (1). The state space is $S = [0, 1]$ with its standard order and all assumptions are as per the discussion immediately following (1). We let P be the corresponding Markov kernel and consider the following coupling of (P_1, P_0) : Let W be a draw from the Bernoulli(1/2) distribution and let $V = 1 - W$. Then P_1 and P_0 are, respectively, the distributions of

$$X := \frac{1+W}{2} \quad \text{and} \quad Y := \frac{0+V}{2} = \frac{1-W}{2}.$$

Since $X \leq Y$ if and only if $W = 0$, we have, by lemma 5.1 and (12),

$$\alpha_O(P_x, P_y) \geq \alpha_O(P_1, P_0) \geq \mathbb{P}\{X \leq Y\} = \frac{1}{2}$$

for all $x, y \in S$. From the definition in (26) we then have $\sigma(P) \geq 1/2$, and globally stability in the γ metric follows from corollary 5.1.

Next we turn to the inventory model in (3), with state space $S = [0, K]$ and $\{W_t\}$ an IID process satisfying $\mathbb{P}\{W_t \geq w\} > 0$ for any real w . Let $\{X_t\}$ and $\{Y_t\}$ be generated by (3), with common shock sequence $\{W_t\}$, and respective initial conditions x, y in S . In view of (5), we have

$$\alpha(P_x, P_y) \geq \mathbb{P}\{X_1 = Y_1\} \geq \mathbb{P}\{W_1 \geq K\} =: \kappa > 0.$$

Letting \preceq be equality on S , we have $\alpha_O(P_x, P_y) = \alpha(P_x, P_y) \geq \kappa$, and hence $\sigma(P) \geq \kappa$. Globally stability in the γ metric follows from corollary 5.1.

5.7. General Applications. Next we show that theorem 5.1 and in particular corollary 5.1 cover as special cases two well known results on Markov chain stability from the classical literature on one hand and the more recent monotone Markov chain literature on the other.

Consider first the standard notion of uniform ergodicity: A Markov kernel P on S is called *uniformly ergodic* if it has a stationary distribution π and $\sup_{x \in S} \|P_x^t - \pi\| \rightarrow 0$ as $t \rightarrow \infty$. Uniform ergodicity was studied by Markov [23] in a countable state space and by Doeblin [7], Yoshida and Kakutani [36], Doob [9] and many subsequent authors in a general state space. It is defined and reviewed in chapter 16 of [24]. One of the most familiar equivalent conditions for uniform ergodicity [24, thm. 16.0.2] is the existence of an $m \in \mathbb{N}$ and a nontrivial $\phi \in \mathcal{M}$ such that $P_x^m \geq \phi$ for all x in S .

One can recover this result using corollary 5.1. Take \preceq to be equality, in which case every Markov operator is monotone, γ is total variation distance and assumption 4.1

is always satisfied. Moreover, $\sigma(P^m)$ reduces to the ordinary ergodicity coefficient of P^m , evaluated using the standard notion of affinity, and hence

$$\sigma(P^m) = \inf_{(x,y) \in S \times S} \alpha(P_x^m, P_y^m) = \inf_{(x,y) \in S \times S} (P_x^m \wedge P_y^m)(S) \geq \phi(S) > 0.$$

Thus, all the conditions of corollary 5.1 are satisfied, and

$$\sup_{x \in S} \|P_x^t - \pi\| = \sup_{x \in S} \gamma(P_x^t, \pi) \leq 2(1 - \sigma(P^m))^{\lfloor t/m \rfloor} \rightarrow 0 \quad (t \rightarrow \infty).$$

Now consider the setting of Bhattacharya and Lee [1], where $S = \mathbb{R}^n$, \preceq is the usual pointwise partial order \leq for vectors, and $\{g_t\}$ is a sequence of IID random maps from S to itself, generating $\{X_t\}$ via $X_t = g_t(X_{t-1}) = g_t \circ \dots \circ g_1(X_0)$. The corresponding Markov kernel is $P(x, B) = \mathbb{P}\{g_1(x) \in B\}$. The random maps are assumed to be order preserving on S , so that P is monotone. Bhattacharya and Lee use a “splitting condition,” which assumes existence of a $\bar{x} \in S$ and $m \in \mathbb{N}$ such that

- (a) $s_1 := \mathbb{P}\{g_m \circ \dots \circ g_1(y) \leq \bar{x}, \forall y \in S\} > 0$ and
- (b) $s_2 := \mathbb{P}\{g_m \circ \dots \circ g_1(y) \geq \bar{x}, \forall y \in S\} > 0.$

Under these assumptions, they show that $\sup_{x \in S} \beta(P_x^t, \pi)$ converges to zero exponentially fast in t , where β is the Bhattacharya metric introduced in (17). This finding extends earlier results by Dubins and Freedman [10] and Yahav [35] to multiple dimensions.

This result can be obtained as a special case of corollary 5.1. Certainly S is a partially ordered Polish space and assumption 4.1 is satisfied. Moreover, the ordered ergodicity coefficient $\sigma(P^m)$ is strictly positive. To see this, suppose that the splitting condition is satisfied at $m \in \mathbb{N}$. Pick any $x, y \in S$ and let $\{X_t\}$ and $\{Y_t\}$ be independent copies of the Markov chain, starting at x and y respectively. We have

$$\sigma(P^m) \geq \mathbb{P}\{X_m \leq Y_m\} \geq \mathbb{P}\{X_m \leq \bar{x} \leq Y_m\} = \mathbb{P}\{X_m \leq \bar{x}\} \mathbb{P}\{\bar{x} \leq Y_m\} \geq s_1 s_2.$$

The last term is strictly positive by assumption. Hence all the conditions of corollary 5.1 are satisfied, a unique stationary distribution π exists, and $\sup_{x \in S} \gamma(P_x^t, \pi)$ converges to zero exponentially fast in t . We showed in (19) that $\beta \leq 2\gamma$, so the same convergence holds for the Bhattacharya metric.

We can also recover a related convergence result due to Hopenhayn and Prescott [16, theorem 2] that is routinely applied to stochastic stability problems in economics. They assume that S is a compact metric space with a closed partial order and a least element a and greatest element b . They suppose that P is monotone, and that there exists an \bar{x} in S and an $m \in \mathbb{N}$ such that

$$(30) \quad P^m(a, [\bar{x}, b]) > 0 \quad \text{and} \quad P^m(b, [a, \bar{x}]) > 0.$$

In this setting, they show that P has a unique stationary distribution π and $\mu P^t \xrightarrow{w} \pi$ for any $\mu \in \mathcal{P}$ as $t \rightarrow \infty$. This result can be obtained from corollary 5.1. Under the stated assumptions, S is Polish and assumption 4.1 is satisfied. The coefficient $\sigma(P^m)$ is strictly positive because, if we let $\{X_t\}$ and $\{Y_t\}$ be independent copies of the Markov chain starting at b and a respectively, then, since $(X_m, Y_m) \in \mathcal{C}(P_b^m, P_a^m)$, we have

$$\alpha_O(P_b^m, P_a^m) \geq \mathbb{P}\{X_m \leq Y_m\} \geq \mathbb{P}\{X_m \leq \bar{x} \leq Y_m\} = \mathbb{P}\{X_m \leq \bar{x}\} \mathbb{P}\{\bar{x} \leq Y_m\}.$$

The last term is strictly positive by (30). Positivity of $\sigma(P^m)$ now follows from lemma 5.1, since $a \preceq x, y \preceq b$ for all $x, y \in S$. Hence, by corollary 5.1, there exists a unique stationary distribution π and $\gamma(\mu P^t, \pi) \rightarrow 0$ as $t \rightarrow \infty$ for any $\mu \in \mathcal{P}$. This convergence implies weak convergence by proposition 4.1 and compactness of S .

6. APPENDIX

The appendix collects remaining proofs. Throughout, in addition to notation defined above, cbS_0 denotes all continuous functions $h: S \rightarrow [0, 1]$, while

$$g(\mu, \nu) := \|\mu\| - \alpha_O(\mu, \nu) \quad \text{for each } \mu, \nu \in \mathcal{M}.$$

6.1. Proofs of Section 3 Results.

Proof of lemma 3.1. For the first equality, fix $\lambda \in \mathcal{M}_s$ and let

$$s(\lambda) := \sup_{I \in i\mathcal{B}} \lambda(I) \quad \text{and} \quad b(\lambda) := \sup_{h \in iH_0} \lambda(h).$$

Since $\mathbb{1}_I \in ibS$ for all $I \in i\mathcal{B}$, we have $b(\lambda) \geq s(\lambda)$. To see the reverse inequality, let $h \in iH_0$. Fix $n \in \mathbb{N}$. Let $r_j := j/n$ for $j = 0, \dots, n$. Define $h_n \in iH_0$ by

$$h_n(x) = \max\{r \in \{r_0, \dots, r_n\} : r \leq h(x)\}.$$

Since $h \leq h_n + 1/n$, we have

$$(31) \quad \lambda(h) \leq \lambda(h_n) + \frac{\|\lambda\|}{n}.$$

For $j = 0, \dots, n$, let $I_j := \{x \in S : h_n(x) \geq r_j\} \in i\mathcal{B}$. Note that

$$(32) \quad I_n = \{x \in S : h_n(x) = 1\} \subset I_{n-1} \subset \dots \subset I_0 = S.$$

We have $\lambda(h_n) = \lambda(I_n) + \sum_{j=1}^n r_{n-j} \lambda(I_{n-j} \setminus I_{n-j+1})$. We define $f_0, \dots, f_{n-1} \in iH_0$ and $A_0, \dots, A_{n-1} \in i\mathcal{B}$ as follows. Let $f_0 = h_n$ and $A_0 = I_n$. Evidently

$$\lambda(f_0) \geq \lambda(h_n), \quad \forall x \in A_0, f_0(x) = 1, \quad \forall x \in I_{n-1} \setminus A_0, f_0(x) = r_{n-1}.$$

Now suppose that for some $j \in \{0, 1, \dots, n-2\}$, we have

$$(33) \quad \lambda(f_j) \geq \lambda(h_n), \quad \forall x \in A_j, f_j(x) = 1, \quad \forall x \in I_{n-j-1} \setminus A_j, f_j(x) = r_{n-j-1}.$$

If $\lambda(I_{n-j-1} \setminus A_j) > 0$, then define

$$f_{j+1}(x) = \begin{cases} 1 & \text{if } x \in I_{n-j-1} \setminus A_j, \\ f_j(x) & \text{otherwise,} \end{cases} \quad \text{and} \quad A_{j+1} = I_{n-j-1}.$$

Note that in this case

$$\begin{aligned} \lambda(f_{j+1}) - \lambda(f_j) &= (1 - r_{n-j-1})\lambda(I_{n-j-1} \setminus A_j) > 0, \\ \forall x \in I_{n-j-2} \setminus A_{j+1}, \quad f_{j+1}(x) &= r_{n-j-2}. \end{aligned}$$

If $\lambda(I_{n-j-1} \setminus A_j) \leq 0$, then define

$$f_{j+1}(x) = \begin{cases} r_{n-j-2} & \text{if } x \in I_{n-j-1} \setminus A_j, \\ f_j(x) & \text{otherwise,} \end{cases} \quad \text{and} \quad A_{j+1} = A_j.$$

In this case $\lambda(f_{j+1}) - \lambda(f_j) = (r_{n-j-2} - r_{n-j-1})\lambda(I_{n-j-1} \setminus A_j) \geq 0$. We also have (6.1) by construction. Thus in both cases, we have (33) with j replaced by $j + 1$. Continuing this way, we see that (33) holds for all $j = 0, \dots, n - 1$.

Let $j = n - 1$ in (33). From the definition of r_j and (32) we have $r_0 = 0$ and $I_0 = S$. Thus

$$\lambda(f_{n-1}) \geq \lambda(h_n), \quad \forall x \in A_{n-1}, f_{n-1}(x) = 1, \quad \forall x \in S \setminus A_{n-1}, f_{n-1}(x) = 0.$$

Since $f_{n-1} = \mathbb{1}_{A_{n-1}}$ and $A_{n-1} = I_j$ for some $j \in \{0, \dots, n - 1\}$, recalling (31) we have

$$\lambda(h) - \frac{\|\lambda\|}{n} \leq \lambda(h_n) \leq \lambda(A_{n-1}) \leq s(\lambda).$$

Applying $\sup_{h \in iH_0}$ to the leftmost side, we see that $b(\lambda) - 1/n \leq s(\lambda)$. Since this is true for any $n \in \mathbb{N}$, we obtain $b(\lambda) \leq s(\lambda)$.

The claim (7) follows from $|a| = \max\{a, -a\}$ and interchange of max and sup. \square

Proof of theorem 3.1. Let (X, Y) be a coupling of (μ, ν) , and define

$$\mu'(B) := \mathbb{P}\{X \in B, X \preceq Y\} \quad \text{and} \quad \nu'(B) := \mathbb{P}\{Y \in B, X \preceq Y\}.$$

Clearly $\mu' \leq \mu$, $\nu' \leq \nu$ and $\mu'(S) = \mathbb{P}\{X \preceq Y\} = \nu'(S)$. Moreover, for any increasing set $I \in \mathcal{B}$ we clearly have $\mu'(I) = \nu'(I)$. Hence $(\mu', \nu') \in \Phi(\mu, \nu)$ and $\mathbb{P}\{X \preceq Y\} = \mu'(S) \leq \alpha_O(\mu, \nu)$. We now exhibit a coupling such that equality is attained. In doing so, we can assume that $a := \alpha_O(\mu, \nu) > 0$.⁴

To begin, observe that, by proposition 3.1, there exists a pair $(\mu', \nu') \in \Phi(\mu, \nu)$ with $\mu'(S) = \nu'(S) = a$. Let $\mu^r := \frac{\mu - \mu'}{1-a}$ and $\nu^r := \frac{\nu - \nu'}{1-a}$. By construction, $\mu^r, \nu^r, \mu'/a$ and ν'/a are probability measures satisfying

$$\mu = (1-a)\mu^r + a(\mu'/a) \quad \text{and} \quad \nu = (1-a)\nu^r + a(\nu'/a).$$

⁴If not, then for any $(X, Y) \in \mathcal{C}(\mu, \nu)$ we have $0 \leq \mathbb{P}\{X \preceq Y\} \leq \alpha_O(\mu, \nu) = 0$.

We construct a coupling (X, Y) as follows. Let U, X', Y', X^r and Y^r be random variables on a common probability space such that

- (a) $X' \stackrel{\mathcal{D}}{=} \mu'/a, Y' \stackrel{\mathcal{D}}{=} \nu'/a, X^r \stackrel{\mathcal{D}}{=} \mu^r$ and $Y^r \stackrel{\mathcal{D}}{=} \nu^r$
- (b) U is uniform on $[0, 1]$ and independent of (X', Y', X^r, Y^r) and
- (c) $\mathbb{P}\{X' \preceq Y'\} = 1$.

The pair in (c) can be constructed via the Nachbin–Strassen theorem [18, thm. 1], since $\mu'/a \preceq_{sd} \nu'/a$. Now let

$$X := \mathbb{1}\{U \leq a\}X' + \mathbb{1}\{U > a\}X^r \quad \text{and} \quad Y := \mathbb{1}\{U \leq a\}Y' + \mathbb{1}\{U > a\}Y^r.$$

Evidently $(X, Y) \in \mathcal{C}(\mu, \nu)$. Moreover, for this pair, we have

$$\mathbb{P}\{X \preceq Y\} \geq \mathbb{P}\{X \preceq Y, U \leq a\} = \mathbb{P}\{X' \preceq Y', U \leq a\}.$$

By independence the right hand side is equal to $\mathbb{P}\{X' \preceq Y'\}\mathbb{P}\{U \leq a\} = a$, so $\mathbb{P}\{X \preceq Y\} \geq a := \alpha_O(\mu, \nu)$. We conclude that

$$(34) \quad \alpha_O(\mu, \nu) = \max_{(X, Y) \in \mathcal{C}(\mu, \nu)} \mathbb{P}\{X \preceq Y\}.$$

Next, observe that, for any $(X, Y) \in \mathcal{C}(\mu, \nu)$ and $h \in ibS$, we have

$$\begin{aligned} \mu(h) - \nu(h) &= \mathbb{E} h(X) - \mathbb{E} h(Y) \\ &= \mathbb{E} [h(X) - h(Y)]\mathbb{1}\{X \preceq Y\} + \mathbb{E} [h(X) - h(Y)]\mathbb{1}\{X \not\preceq Y\} \\ &\leq \mathbb{E} [h(X) - h(Y)]\mathbb{1}\{X \not\preceq Y\}. \end{aligned}$$

Specializing to $h = \mathbb{1}_I$ for some $I \in i\mathcal{B}$, we have $\mu(I) - \nu(I) \leq \mathbb{P}\{X \not\preceq Y\} = 1 - \mathbb{P}\{X \preceq Y\}$. From this bound and (34), the proof of (12) will be complete if we can show that

$$(35) \quad \sup_{(X, Y) \in \mathcal{C}(\mu, \nu)} \mathbb{P}\{X \preceq Y\} \geq 1 - \sup_{I \in i\mathcal{B}} \{\mu(I) - \nu(I)\}.$$

To prove (35), let $\mathcal{B} \otimes \mathcal{B}$ be the product σ -algebra on $S \times S$ and let π_i be the i -th coordinate projection, so that $\pi_1(x, y) = x$ and $\pi_2(x, y) = y$ for any $(x, y) \in S \times S$. As usual, given $Q \subset S \times S$, we let $\pi_1(Q)$ be all $x \in S$ such that $(x, y) \in Q$, and similarly for π_2 . Recall that \mathcal{C} is the closed sets in S and $d\mathcal{C}$ is the decreasing sets in \mathcal{C} . Strassen's theorem [32] implies that, for any $\epsilon \geq 0$ and any closed set $K \subset S \times S$, there exists a probability measure ξ on $(S \times S, \mathcal{B} \otimes \mathcal{B})$ with marginals μ and ν such that $\xi(K) \geq 1 - \epsilon$ whenever

$$\nu(F) \leq \mu(\pi_1(K \cap (S \times F))) + \epsilon, \quad \forall F \in \mathcal{C}.$$

Note that if $F \in \mathcal{C}$, then, since \preceq is a closed partial order, so is the smallest decreasing set $d(F)$ that contains F . Let $\epsilon := \sup_{D \in d\mathcal{C}} \{\nu(D) - \mu(D)\}$, so that

$$\epsilon \geq \sup_{F \in \mathcal{C}} \{\nu(d(F)) - \mu(d(F))\} \geq \sup_{F \in \mathcal{C}} \{\nu(F) - \mu(d(F))\}.$$

Noting that $d(F)$ can be expressed as $\pi_1(\mathbb{G} \cap (S \times F))$, it follows that, for any $F \in \mathcal{C}$,

$$\nu(F) \leq \mu(\pi_1(\mathbb{G} \cap (S \times F))) + \epsilon.$$

Since \preceq is closed, \mathbb{G} is closed, and Strassen's theorem applies. From this theorem we obtain a probability measure ξ on the product space $S \times S$ such that $\xi(\mathbb{G}) \geq 1 - \epsilon$ and ξ has marginals μ and ν .

Because complements of increasing sets are decreasing and vice versa, we have

$$(36) \quad \sup_{I \in i\mathcal{B}} \{\mu(I) - \nu(I)\} \geq \sup_{D \in d\mathcal{C}} \{\nu(D) - \mu(D)\} = \epsilon \geq 1 - \xi(\mathbb{G}).$$

Now consider the probability space $(\Omega, \mathcal{F}, \mathbb{P}) = (S \times S, \mathcal{B} \otimes \mathcal{B}, \xi)$, and let $X = \pi_1$ and $Y = \pi_2$. We then have $\xi(\mathbb{G}) = \xi\{(x, y) \in S \times S : x \preceq y\} = \mathbb{P}\{X \preceq Y\}$. Combining this equality with (36) implies (35). \square

6.2. Proofs of Section 4 Results. We begin with an elementary lemma:

Lemma 6.1. *For any $\mu, \nu \in \mathcal{M}$, we have $\mu \leq \nu$ whenever $\mu(h) \leq \nu(h)$ for all $h \in cbS_0$.*

Proof. Suppose that $\mu(h) \leq \nu(h)$ for all $h \in cbS_0$. We claim that

$$(37) \quad \mu(F) \leq \nu(F) \text{ for any closed set } F \subset S.$$

To see this, let ρ be a metric compatible with the topology of S and let F be any closed subset of S . Let $f_\epsilon(x) := \max\{1 - \rho(x, F)/\epsilon, 0\}$ for $\epsilon > 0$, $x \in S$, where $\rho(x, F) = \inf_{y \in F} \rho(x, y)$. Since $\rho(\cdot, F)$ is continuous and $0 \leq f_\epsilon \leq 1$, we have $f_\epsilon \in cbS_0$. Let $F_\epsilon = \{x \in S : \rho(x, F) < \epsilon\}$ for $\epsilon > 0$. Note that $f_\epsilon(x) = 1$ for all $x \in F$, and that $f_\epsilon(x) = 0$ for all $x \notin F_\epsilon$. Thus,

$$(38) \quad \mu(F) \leq \mu(f_\epsilon) \leq \nu(f_\epsilon) \leq \nu(F_\epsilon).$$

Since $F = \bigcap_{\epsilon > 0} F_\epsilon$, we have $\lim_{\epsilon \downarrow 0} \nu(F_\epsilon) = \nu(F)$, so letting $\epsilon \downarrow 0$ in (38) yields $\mu(F) \leq \nu(F)$. Hence (37) holds.

Let $B \in \mathcal{B}$ and fix $\epsilon > 0$. Since all probability measures on a Polish space are regular, there exists a closed set $F \subset B$ such that $\mu(B) \leq \mu(F) + \epsilon$. Thus by (37), we have $\mu(B) \leq \mu(F) + \epsilon \leq \nu(F) + \epsilon \leq \nu(B) + \epsilon$. Since $\epsilon > 0$ is arbitrary, this yields $\mu(B) \leq \nu(B)$. Hence $\mu \leq \nu$. \square

Proof of theorem 4.1. Let $\{\mu_n\}$ be a Cauchy sequence in (\mathcal{P}, γ) . Our first claim is that $\{\mu_n\}$ is tight. To show this, fix $\epsilon > 0$. Let $\mu := \mu_N$ be such that

$$(39) \quad n \geq N \implies \gamma(\mu, \mu_n) < \epsilon.$$

Let K be a compact subset of S such that $\mu(K) \geq 1 - \epsilon$ and let $\bar{K} := i(K) \cap d(K)$. We have

$$\mu_n(\bar{K}^c) = \mu_n(i(K)^c \cup d(K)^c) \leq \mu_n(i(K)^c) + \mu_n(d(K)^c).$$

For $n \geq N$, this bound, (16), (39) and the definition of K yield

$$\mu_n(\bar{K}^c) < \mu(i(K)^c) + \mu(d(K)^c) + 2\epsilon \leq \mu(K^c) + \mu(K^c) + 2\epsilon \leq 4\epsilon.$$

Hence $\{\mu_n\}_{n \geq N}$ is tight. It follows that $\{\mu_n\}_{n \geq 1}$ is likewise tight. As a result, by Prohorov's theorem, it has a subsequence that converges weakly to some $\mu^* \in \mathcal{P}$. We aim to show that $\gamma(\mu_n, \mu^*) \rightarrow 0$.

To this end, fix $\epsilon > 0$ and let n_ϵ be such that $\gamma(\mu_m, \mu_{n_\epsilon}) < \epsilon$ whenever $m \geq n_\epsilon$. Fix $m \geq n_\epsilon$ and let $\nu := \mu_m$. For all $n \geq n_\epsilon$, we have $\gamma(\nu, \mu_n) < \epsilon$. Fixing any such $n \geq n_\epsilon$, we observe that since $g(\mu_n, \nu) < \epsilon$, there exists $(\tilde{\mu}_n, \tilde{\nu}_n) \in \Phi(\mu_n, \nu)$ with $\|\tilde{\mu}_n\| = \|\tilde{\nu}_n\| > 1 - \epsilon$. Multiplying $\tilde{\mu}_n$ and $\tilde{\nu}_n$ by $(1 - \epsilon)/\|\tilde{\mu}_n\| < 1$, denoting the resulting measures by $\tilde{\mu}_n$ and $\tilde{\nu}_n$ again, we have

$$(40) \quad \tilde{\mu}_n \leq \mu_n, \quad \tilde{\nu}_n \leq \nu, \quad \|\tilde{\mu}_n\| = \|\tilde{\nu}_n\| = 1 - \epsilon, \quad \tilde{\mu}_n \preceq_{sd} \tilde{\nu}_n.$$

Note that $\{\tilde{\nu}_n\}$ is tight. Since $\{\mu_n\}$ is tight, so is $\{\tilde{\mu}_n\}$. Thus there exist subsequences $\{\mu_{n_i}\}_{i \in \mathbb{N}}$, $\{\tilde{\mu}_{n_i}\}_{i \in \mathbb{N}}$, and $\{\tilde{\nu}_{n_i}\}_{i \in \mathbb{N}}$ of $\{\mu_n\}$, $\{\tilde{\mu}_n\}$, and $\{\tilde{\nu}_n\}$ respectively such that, for some $\tilde{\mu}^*, \tilde{\nu}^* \in \mathcal{M}$ with $\|\tilde{\mu}^*\| = \|\tilde{\nu}^*\| = 1 - \epsilon$, we have

$$\mu_{n_i} \xrightarrow{w} \mu^*, \quad \tilde{\mu}_{n_i} \xrightarrow{w} \tilde{\mu}^*, \quad \tilde{\nu}_{n_i} \xrightarrow{w} \tilde{\nu}^*, \quad \forall i \in \mathbb{N}, \quad \tilde{\mu}_{n_i} \preceq_{sd} \tilde{\nu}_{n_i}.$$

Given $h \in cbS_0$, since $\tilde{\mu}_{n_i}(h) \leq \mu_{n_i}(h)$ and $\tilde{\nu}_{n_i}(h) \leq \nu(h)$ for all $i \in \mathbb{N}$ by (40), we have $\tilde{\mu}^*(h) \leq \mu^*(h)$ and $\tilde{\nu}^*(h) \leq \nu(h)$ by weak convergence. Thus $\tilde{\mu}^* \leq \mu^*$ and $\tilde{\nu}^* \leq \nu$ by lemma 6.1. We have $\tilde{\mu}^* \preceq_{sd} \tilde{\nu}^*$ by [18, proposition 3]. It follows that $(\tilde{\mu}^*, \tilde{\nu}^*) \in \Phi(\mu^*, \nu)$. We have $g(\mu^*, \nu) \leq 1 - \|\tilde{\mu}^*\| = \epsilon$.

By a symmetric argument, we also have $g(\nu, \mu^*) \leq \epsilon$. Hence $\gamma(\nu, \mu^*) \leq 2\epsilon$. Recalling the definition of ν , we have now shown that, $\forall m \geq n_\epsilon$, $\gamma(\mu_m, \mu^*) \leq 2\epsilon$. Since ϵ was arbitrary this concludes the proof. \square

6.3. Proofs of Section 5 Results. We begin with some lemmata.

Lemma 6.2. *If P is monotone, then $\sigma(P) = \inf_{(\mu, \nu) \in \mathcal{P} \times \mathcal{P}} \alpha_O(\mu P, \nu P)$.*

Proof. Let P be a monotone Markov kernel. It suffices to show that the inequality $\sigma(P) \leq \inf_{(\mu, \nu) \in \mathcal{P} \times \mathcal{P}} \alpha_O(\mu P, \nu P)$ holds, since the reverse inequality is obvious. By

the definition of $\sigma(P)$ and the identities in (12), the claim will be established if we can show that

$$(41) \quad \sup_{x,y} \sup_{I \in i\mathcal{B}} \{P(x, I) - P(y, I)\} \geq \sup_{\mu, \nu} \sup_{I \in i\mathcal{B}} \{\mu P(I) - \nu P(I)\}.$$

where x and y are chosen from S and μ and ν are chosen from \mathcal{P} . Let s be the value of the right hand side of (41) and let $\epsilon > 0$. Fix $\mu, \nu \in \mathcal{P}$ and $I \in i\mathcal{B}$ such that $\mu P(I) - \nu P(I) > s - \epsilon$, or, equivalently,

$$\int \{P(x, I) - P(y, I)\} (\mu \times \nu)(dx, dy) > s - \epsilon.$$

From this expression we see that there are $\bar{x}, \bar{y} \in S$ such that $P(\bar{x}, I) - P(\bar{y}, I) > s - \epsilon$. Hence $\sup_{x,y} \sup_{I \in i\mathcal{B}} \{P(x, I) - P(y, I)\} \geq s$, as was to be shown. \square

Lemma 6.3. *If $\mu, \nu \in \mathcal{M}$ and $(\tilde{\mu}, \tilde{\nu})$ is an ordered component pair of (μ, ν) , then*

$$g(\mu P, \nu P) \leq g((\mu - \tilde{\mu})P, (\nu - \tilde{\nu})P).$$

Proof. Fix μ, ν in \mathcal{M} and $(\tilde{\mu}, \tilde{\nu}) \in \Phi(\mu, \nu)$. Consider the residual measures $\hat{\mu} := \mu - \tilde{\mu}$ and $\hat{\nu} := \nu - \tilde{\nu}$. Let (μ', ν') be a maximal ordered component pair of $(\hat{\mu}P, \hat{\nu}P)$, and define

$$\mu^* := \mu' + \tilde{\mu}P \quad \text{and} \quad \nu^* := \nu' + \tilde{\nu}P.$$

We claim that (μ^*, ν^*) is an ordered component pair for $(\mu P, \nu P)$. To see this, note that

$$\mu^* = \mu' + \tilde{\mu}P \leq \hat{\mu}P + \tilde{\mu}P = (\hat{\mu} + \tilde{\mu})P = \mu P,$$

and, similarly, $\nu^* \leq \nu P$. The measures μ^* and ν^* also have the same mass, since

$$\|\mu^*\| = \|\mu'\| + \|\tilde{\mu}P\| = \|\mu'\| + \|\tilde{\mu}\| = \|\nu'\| + \|\tilde{\nu}\| = \|\nu'\| + \|\tilde{\nu}P\| = \|\nu^*\|.$$

Moreover, since $\tilde{\mu} \preceq_{sd} \tilde{\nu}$ and P is monotone, we have $\tilde{\mu}P \preceq_{sd} \tilde{\nu}P$. Hence $\mu^* \preceq_{sd} \nu^*$, completing the claim that (μ^*, ν^*) is an ordered component pair for $(\mu P, \nu P)$. As a result,

$$\begin{aligned} g(\mu P, \nu P) &\leq \|\mu P\| - \|\mu^*\| = \|\mu P\| - \|\mu'\| - \|\tilde{\mu}P\| \\ &= \|\mu\| - \|\mu'\| - \|\tilde{\mu}\| = \|\hat{\mu}\| - \|\mu'\| = \|\hat{\mu}P\| - \|\mu'\| = g(\hat{\mu}P, \hat{\nu}P). \quad \square \end{aligned}$$

Proof of theorem 5.1. Let $\mu, \nu \in \mathcal{P}$. Let $(\tilde{\mu}, \tilde{\nu})$ be a maximal ordered component pair for (μ, ν) . Let $\hat{\mu} = \mu - \tilde{\mu}$ and $\hat{\nu} = \nu - \tilde{\nu}$ be the residuals. Since $\|\tilde{\mu}\| = \|\tilde{\nu}\|$, we have $\|\hat{\mu}\| = \|\hat{\nu}\|$. Suppose first that $\|\hat{\mu}\| > 0$. Then $\hat{\mu}P/\|\hat{\mu}\|$ and $\hat{\nu}P/\|\hat{\mu}\|$ are both in \mathcal{P} . Thus, by lemma 6.2,

$$1 - \alpha_O(\hat{\mu}P/\|\hat{\mu}\|, \hat{\nu}P/\|\hat{\mu}\|) \leq 1 - \sigma(P).$$

Applying the positive homogeneity property in lemma 3.3 yields

$$\|\hat{\mu}\| - \alpha_O(\hat{\mu}P, \hat{\nu}P) \leq (1 - \sigma(P)) \|\hat{\mu}\|,$$

Note that this inequality trivially holds if $\|\hat{\mu}\| = 0$. From the definition of g , we can write the same inequality as $g(\hat{\mu}P, \hat{\nu}P) \leq (1 - \sigma(P))g(\mu, \nu)$. If we apply lemma 6.3 to the latter we obtain

$$g(\mu P, \nu P) \leq (1 - \sigma(P))g(\mu, \nu).$$

Reversing the roles of μ and ν , we also have $g(\nu P, \mu P) \leq (1 - \sigma(P))g(\nu, \mu)$. Thus

$$\begin{aligned} \gamma(\mu P, \nu P) &= g(\mu P, \nu P) + g(\nu P, \mu P) \\ &\leq (1 - \sigma(P))[g(\mu, \nu) + g(\nu, \mu)] = (1 - \sigma(P))\gamma(\mu, \nu), \end{aligned}$$

verifying the claim in (27). \square

Proof of lemma 5.4. To see that (28) holds, fix $\xi > \sigma(P)$ and suppose first that $\sigma(P) = 1$. Then (28) holds because the right hand side of (28) can be made strictly negative by choosing $\mu, \nu \in \mathcal{P}$ to be distinct. Now suppose that $\sigma(P) < 1$ holds. It suffices to show that

$$(42) \quad \forall \epsilon > 0, \exists x, y \in S \text{ such that } y \preceq x, x \not\preceq y \text{ and } \alpha_0(P_x, P_y) < \sigma(P) + \epsilon.$$

Indeed, if we take (42) as valid, set $\epsilon := \xi - \sigma(P)$ and choose x and y to satisfy the conditions in (42), then we have $\gamma(P_x, P_y) = 2 - \alpha_0(P_x, P_y) - \alpha_0(P_y, P_x) > 2 - \xi - 1 = 1 - \xi = (1 - \xi)\gamma(\delta_x, \delta_y)$. Therefore (28) holds.

To show that (42) holds, fix $\epsilon > 0$. We can use $\sigma(P) < 1$ and the definition of $\sigma(P)$ as an infimum to choose an $\delta \in (0, \epsilon)$ and points $\bar{x}, \bar{y} \in S$ such that $\alpha_0(P_{\bar{x}}, P_{\bar{y}}) < \sigma(P) + \delta < 1$. Note that $\bar{x} \preceq \bar{y}$ cannot hold here, because then $\alpha_0(P_{\bar{x}}, P_{\bar{y}}) = 1$, a contradiction. So suppose instead that $\bar{x} \not\preceq \bar{y}$. Let y be a lower bound of \bar{x} and \bar{y} and let $x := \bar{x}$. We claim that (42) holds for the pair (x, y) .

To see this, observe that, by the monotonicity result in lemma 5.1 and $y \preceq \bar{y}$ we have $\alpha_0(P_x, P_y) = \alpha_0(P_{\bar{x}}, P_y) \leq \alpha_0(P_{\bar{x}}, P_{\bar{y}}) < \sigma(P) + \delta < \sigma(P) + \epsilon$. Moreover, $y \preceq x$ because $x = \bar{x}$ and y is by definition a lower bound of \bar{x} . Finally, $x \not\preceq y$ because if not then $\bar{x} = x \preceq y \preceq \bar{y}$, contradicting our assumption that $\bar{x} \not\preceq \bar{y}$. \square

REFERENCES

- [1] R.N. Bhattacharya and O. Lee. Asymptotics of a class of Markov processes which are not in general irreducible. *The Annals of Probability*, pages 1333–1347, 1988.
- [2] R.N. Bhattacharya, M. Majumdar, and N. Hashimzade. Limit theorems for monotone Markov processes. *Sankhya A*, 72(1):170–190, 2010.
- [3] Santanu Chakraborty and BV Rao. Completeness of the Bhattacharya metric on the space of probabilities. *Statistics & Probability Letters*, 36(4):321–326, 1998.
- [4] Persi Diaconis and David Freedman. Iterated random functions. *SIAM review*, 41(1):45–76, 1999.

- [5] Roland L Dobrushin. Central limit theorem for nonstationary Markov chains. *Theory of Probability and Its Applications*, 1(1):65–80, 1956.
- [6] Wolfgang Doeblin. Exposé de la théorie des chaînes simples constantes de Markov à un nombre fini d'états. *Mathématique de l'Union Interbalkanique*, 2(77-105):78–80, 1938.
- [7] Wolfgang Doeblin. Sur les propriétés asymptotiques de mouvements régis par certains types de chaînes simples. *Bull. Math. Soc. Roum. Sci.*, 39:(1) 57–115; (2), 3–61, 1937.
- [8] Wolfgang Doeblin. Éléments d'une théorie générale des chaînes simples constantes de Markoff. In *Annales Scientifiques de l'École Normale Supérieure*, volume 57, pages 61–111, 1940.
- [9] Joseph L Doob. *Stochastic processes*. Wiley New York, 1953.
- [10] Lester E Dubins and David A Freedman. Invariant probabilities for certain Markov processes. *The Annals of Mathematical Statistics*, 37(4):837–848, 1966.
- [11] R.M. Dudley. *Real Analysis and Probability*, volume 74. Cambridge University Press, 2002.
- [12] Alison L Gibbs and Francis Edward Su. On choosing and bounding probability metrics. *International statistical review*, 70(3):419–435, 2002.
- [13] Clark R Givens, Rae Michael Shortt, et al. A class of Wasserstein metrics for probability distributions. *The Michigan Mathematical Journal*, 31(2):231–240, 1984.
- [14] Theodore E Harris. The existence of stationary measures for certain Markov processes. In *Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability*, volume 2, pages 113–124, 1956.
- [15] O. Hernández-Lerma and J.B. Lasserre. *Markov Chains and Invariant Probabilities*. Springer, 2003.
- [16] Hugo A Hopenhayn and Edward C Prescott. Stochastic monotonicity and stationary distributions for dynamic economies. *Econometrica*, pages 1387–1406, 1992.
- [17] T. Kamae and U. Krengel. Stochastic partial ordering. *The Annals of Probability*, 6(6):1044–1049, 1978.
- [18] T. Kamae, U. Krengel, and G.L. O'Brien. Stochastic inequalities on partially ordered spaces. *The Annals of Probability*, pages 899–912, 1977.
- [19] T. Kamihigashi and J. Stachurski. An order-theoretic mixing condition for monotone Markov chains. *Statistics & Probability Letters*, 82(2):262–267, 2012.
- [20] Takashi Kamihigashi and John Stachurski. Stochastic stability in monotone economies. *Theoretical Economics*, 9(2):383–407, 2014.
- [21] T. Lindvall. *Lectures on the coupling method*. Dover Publications, 2002.
- [22] Motoya Machida and Alexander Shibakov. Monotone bivariate Markov kernels with specified marginals. *Proceedings of the American Mathematical Society*,

- 138(6):2187–2194, 2010.
- [23] Andrey Andreyevich Markov. Extension of the law of large numbers to dependent quantities. *Izv. Fiz.-Matem. Obsch. Kazan Univ.(2nd Ser)*, 15:135–156, 1906.
 - [24] Sean P Meyn and Richard L Tweedie. *Markov chains and stochastic stability*. Springer Science & Business Media, 2012.
 - [25] Esa Nummelin. *General irreducible Markov chains and non-negative operators*. Cambridge University Press, 2004.
 - [26] Steven Orey. Recurrent Markov chains. *Pacific J. Math*, 9(3):805–827, 1959.
 - [27] James W Pitman. Uniform rates of convergence for Markov chain transition probabilities. *Probability Theory and Related Fields*, 29(3):193–227, 1974.
 - [28] David Pollard. *A user’s guide to measure theoretic probability*, volume 8. Cambridge University Press, 2002.
 - [29] Assaf Razin and Joseph A Yahav. On stochastic models of economic growth. *International Economic Review*, pages 599–604, 1979.
 - [30] Daniel Revuz. *Markov chains*, volume 11. Elsevier, 2008.
 - [31] Eugene Seneta. Coefficients of ergodicity: structure and applications. *Advances in applied probability*, 11(03):576–590, 1979.
 - [32] V. Strassen. The existence of probability measures with given marginals. *The Annals of Mathematical Statistics*, pages 423–439, 1965.
 - [33] H. Thorisson. *Coupling, stationarity, and regeneration*. Springer New York, 2000.
 - [34] Wei Biao Wu and Xiaofeng Shao. Limit theorems for iterated random functions. *Journal of Applied Probability*, 41(2):425–436, 2004.
 - [35] Joseph A Yahav. On a fixed point theorem and its stochastic equivalent. *Journal of Applied Probability*, pages 605–611, 1975.
 - [36] K Yoshida and S Kakutani. Operator theoretical treatment of Markov processes and mean ergodic theorem. *Ann. Math*, 42, 1941.

RIEB, KOBE UNIVERSITY

Email address: tkamihig@rieb.kobe-u.ac.jp

RESEARCH SCHOOL OF ECONOMICS, AUSTRALIAN NATIONAL UNIVERSITY

Email address: john.stachurski@anu.edu.au