Distribution of a Unified (k_1, k_2, \ldots, k_m) -run

Savita Chaturvedi¹ and Amit N. Kumar²

^{1,2}Department of Mathematical Sciences, Indian Institute of Technology (BHU), Varanasi (Uttar Pradesh) - 221 005, India.
¹Email: savitachaturvedi.rs.mat23@itbhu.ac.in
²Email: amit.mat@itbhu.ac.in

Abstract

We explore a unified (k_1, k_2, \ldots, k_m) -run in multi-state trials, examining its distributional properties and waiting time distribution. Our study reveals that this particular run serves as a generalization encompassing various patterns. Additionally, we discuss various results pertaining to existing patterns as special cases. To illustrate our findings, we provide an application related to DNA frequent patterns.

Keywords: Multi-state trials; generating functions; exact distribution; waiting time distribution. **MSC 2020 Subject Classifications:** Primary: 62E99, 62E15; Secondary: 60E05, 60E10.

1 Introduction and Preliminaries

The concept of runs has extensive applications across various fields such as climatology, quality control, biology, computer science, and DNA sequence analysis, among many others. In a sequence of Bernoulli trials with possible outcomes of failure and success, the distribution of the number of occurrences of k-runs (defined as at least k consecutive successes) holds significant importance and has been extensively discussed in the literature. Several authors have studied the distributional properties of k-runs and its waiting time distribution along with various applications, see for more details, Aki *et al.* [1], Antzoulakos *et al.* [2], Antzoulakos and Chadjiconstantinidis [3], Balakrishnan and Koutras [4], Fu and Koutras [9], Makri *et al.* [20], Philippou *et al.* [21], and references therein. Additionally, the research on approximations related to runs has also been extensively explored in the literature. For more details, see Čekanavičius and Vellaisamy [24, 25, 26], Fu and Johnson [8], Kumar and Kumar [14], Kumar and Upadhye [15, 17], Kumar *et al.* [18], Liaudanskaitė and Čekanavičius [19], Upadhye and Kumar [22], Upadhye *et al.* [23], Vellaisamy [27], Wang and Xia [28], and references therein. Huang and Tsai [10] extended the concept of k-runs by introducing the notion of (k_1, k_2) -runs, which involve the pattern at least k_1 consecutive failures followed by at least k_2 consecutive successes. Subsequently, Dafnis *et al.* [6] explored three variations of (k_1, k_2) -runs, considering scenarios involving at least, exactly, and at most occurrences of runs. Further, Kumar and Upadhye [16] generalized these patterns to encompass three general types of runs. While previous studies focused on two-state trials in studying k-runs and (k_1, k_2) -runs, the demand across various applications necessitates the exploration of (k_1, k_2, \ldots, k_m) -runs in multi-state trials. For instance, in the DNA sequence analysis, it is crucial to represent sequences using four letters: A, G, C, and T, which correspond to a sequence of four-state trials. Kong [13] initiated this study by focusing the run at least k_1 consecutive 1's followed by k_2 consecutive 2's ... followed by k_m consecutive m's in a sequence of multi-state trials with labels $\{1, 2, \ldots, m\}$. Mathematically, let $\xi_1, \xi_2, \ldots, \xi_n$ be a sequence of independent and identically distributed (*iid*) multinomial random variables, each taking values from the set $\{1, 2, \ldots, m\}$, with probability $\mathbb{P}(\xi_i = j) = p_j$, for $j = 1, 2, \ldots, m$, and $\sum_{j=1}^m p_j = 1$. Define

$$\mathbb{I}_{i} := \left(\prod_{j_{1}=i}^{i+k_{1}-1} (2-\xi_{j_{1}})\right) \left(\prod_{j_{2}=i+k_{1}}^{i+k_{1}+k_{2}-1} (3-\xi_{j_{2}})\right) \dots \left(\prod_{j_{m}=i+k_{1}+\dots+k_{m-1}}^{i+k_{1}+\dots+k_{m}-1} (m+1-\xi_{j_{m}})\right).$$

Then,

$$X_{m;k_1,k_2,\dots,k_m}^{(n)} := \sum_{i=1}^{n-k_1-\dots-k_m+1} \mathbb{I}_i$$

represents the number of occurrences of the considered $(k_1, k_2, ..., k_m)$ -run. Kong [13] developed a generating function approach to derive the exact distribution of $X_{m;k_1,k_2,...,k_m}^{(n)}$. Additionally, Chadjiconstantinidis and Eryilmaz [5] have studied the waiting time distribution of $X_{m;k_1,k_2,...,k_m}^{(n)}$.

In this paper, we explore a unified $(k_1, k_2, ..., k_m)$ -run in a sequence of multi-state trials with labels $\{1, 2, ..., m\}$ defined as follows:

(**R**) at least ℓ_1 and at most k_1 consecutive 1's followed by at least ℓ_2 and at most k_2 consecutive 2's \ldots followed at least ℓ_m and at most k_m consecutive m's.

Here, we assume $k_i \ge \ell_i \ge 1$. Mathematically, let

$$\mathbb{I}_{i;r_1,r_2,\dots,r_m} := \left(\prod_{j_1=i}^{i+r_1-1} (2-\xi_{j_1})\right) \left(\prod_{j_2=i+r_1}^{i+r_1+r_2-1} (3-\xi_{j_2})\right) \dots \left(\prod_{j_m=i+r_1+\dots+r_{m-1}}^{i+r_1+\dots+r_m-1} (m+1-\xi_{j_m})\right),$$

where
$$r_i = s_i + \ell_i$$
, for $i = 1, 2, ..., m$, and $\mathbb{I}_i^* := \max_{\substack{r_i \le \ell_i + k_i \\ i = 1, 2, ..., m}} \mathbb{I}_{i;r_1, r_2, ..., r_m}$. Then
$$X_{m; \ell_1:k_1, \ell_2: k_2, ..., \ell_m: k_m}^{(n)} := \sum_{\substack{i=1 \\ i=1}}^{n-\ell_1 - \dots - \ell_m + 1} \mathbb{I}_i^*$$

represents the number of occurrences of the run (\mathbf{R}). To simplify the presentation of paper, we introduce the following notations:

- E_{k_i} denotes a run of at least k_i consecutive *i*'s.
- F_{k_i} denotes a run of exactly k_i consecutive *i*'s.
- G_{k_i} denotes a run of at most k_i consecutive *i*'s.
- E_{ℓ_i,k_i} denotes a run of at least ℓ_i and at most k_i consecutive *i*'s.

Moreover, we define the notation $E_i \prec E_j$ to represent the event E_i appears immediately after the event E_j . Therefore, the run (**R**) is equivalent to $E_{\ell_1,k_1} \prec E_{\ell_2,k_2} \prec \cdots \prec E_{\ell_m,k_m}$. Note that if $\ell_i = k_i$ and $\ell_i = 1$, for $i = 1, 2, \ldots, m$, then $E_{\ell_i,k_i} = F_{k_i}$ and $E_{\ell_i,k_i} = G_{k_i}$, respectively. In such cases, the run (**R**) simplifies to $F_{k_1} \prec F_{k_2} \prec \cdots \prec F_{k_m}$ and $G_{k_1} \prec G_{k_2} \prec \cdots \prec G_{k_m}$, respectively. It is worth noting that $E_{\ell_i,k_i} \to E_{\ell_i}$ when $k_i \to \infty$. Therefore, if $k_i \to \infty$, for $i = 1, 2, \ldots, m$, then the run (**R**) reduces to $E_{\ell_1} \prec E_{\ell_2} \prec \cdots \prec E_{\ell_m}$, which is studied by by Kong [13]. Thus, the run (**R**) can be transformed into a (k_1, k_2, \ldots, k_m) -run with any k_i , for $i = 1, 2, \ldots, m$, defined as at least, exactly, and at most k_i consecutive *i*'s. Therefore, the study of the run (**R**) suffices to study any type of (k_1, k_2, \ldots, k_m) -run with all at least, exactly, and at most cases.

We employ a modified methodology introduced by Kong [11, 13] to derive our results. The advantage of this method lies in its utilization of the generating functions of individual patterns to derive the generating function for complicated patterns of interest. The technique can be formulated as follows: Suppose a linear sequence comprising multiple objects with values drawn from the set $\{1, 2, ..., m\}$, where objects may occur repeatedly. Subsequently, we partition the sequence into distinct patterns of our interest, each forming specific blocks. These blocks are organized based on the pattern we wish to explore, ensuring relevance and coherence throughout the analysis. The relationship between blocks is characterized by the interaction w_{ij} , representing the influence of the *i*th type of block on the left and the *j*th type of block on the right. Furthermore, we denote the generating function as g_i associated with block *i*. The generating function for the entire system can be expressed as

$$G = 1 + \boldsymbol{e}\boldsymbol{M}^{-1}\boldsymbol{g}.$$
 (1.1)

Here, $e = (1 \ 1 \ \cdots \ 1)$ represents a row vector with all entries equal to 1, $g = (g_1 \ g_2 \ \cdots \ g_r)^T$ represents a column vector, where each entry corresponds to a generating function g_i , i = 1, 2, ..., r, and

$$\boldsymbol{M} := \begin{pmatrix} 1 & -g_1 w_{12} & -g_1 w_{13} & \cdots & -g_1 w_{1r} \\ -g_2 w_{21} & 1 & -g_2 w_{23} & \cdots & -g_2 w_{2r} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -g_r w_{r1} & -g_r w_{r2} & -g_r w_{r3} & \cdots & 1 \end{pmatrix}$$

When blocks are forbidden, the interaction is set to zero. To compute the distribution of $(k_1, k_2, ..., k_m)$ -runs, one approach is to initially consider two consecutive blocks together. Subsequently, by iteratively adding more blocks, we can obtain the generating function for $(k_1, k_2, ..., k_m)$ -runs. For further details, we refer the reader to Kong [11, 12, 13].

This paper is organized as follows: In Section 2, we derive the distributional properties of the random variable $X_{m;\ell_1:k_1,\ell_2:k_2,...,\ell_m:k_m}^{(n)}$. Our study reveals that this random variable serves as a generalization encompassing various patterns. Additionally, we discuss various results pertaining to existing patterns as special cases. In Section 3, we further explore the distributional properties of the waiting time distribution of $X_{m;\ell_1:k_1,\ell_2:k_2,...,\ell_m:k_m}^{(n)}$. Finally, in Section 4, we present an application to DNA frequent pattern, illustrating the practical implications of our results.

2 Distribution of $X_{m;\ell_1:k_1,\ell_2:k_2,...,\ell_m:k_m}^{(n)}$

In this section, we explore the run (\mathbf{R}) introduced in Section 1 and derive its distributional properties using the generating function approach proposed by Kong [13]. Our proofs are established through induction. Initially, we focus on the case of two-state trials, presenting several findings related to previous studies. Subsequently, we extend our analysis to multi-state trials, where we demonstrate the applicability of our results and how they encompass existing findings as special cases.

We begin to introduce some notations that will be useful for further finding. Throughout this paper, let $m \ge 2$, $\prod_{i}^{j} = 1$ and $\sum_{i}^{j} = 0$, for j < i. Recall that $\xi_{1}, \xi_{2}, \ldots, \xi_{n}$ is a sequence of *iid* multinomial random variables, each taking values from the set $\{1, 2, \ldots, m\}$, with probability $\mathbb{P}(\xi_{i} = j) = p_{j}$, for $i = 1, 2, \ldots, n, j = 1, 2, \ldots, m$, and $\sum_{j=1}^{m} p_{j} = 1$. The random variable $X_{m;\ell_{1}:k_{1},\ell_{2}:k_{2},\ldots,\ell_{m}:k_{m}}^{(n)}$ denotes the number of occurrences of the run $E_{\ell_{1},k_{1}} \prec E_{\ell_{2},k_{2}} \prec \cdots \prec E_{\ell_{m},k_{m}}$. Further, let us denote $p_{m,n}(r) = \mathbb{P}(X_{m;\ell_{1}:k_{1},\ell_{2}:k_{2},\ldots,\ell_{m}:k_{m}}^{(n)} = r)$. The generating function for the sequence of length n is represented as

$$\phi_{m,n}(w) = \sum_{j=0} p_{m,n}(j)w^j$$

and the double generating function is expressed as

$$\Phi_m(w,z) = \sum_{n=0}^{\infty} \phi_{m,n}(w) z^n = \sum_{n=0}^{\infty} \sum_{j=0}^{\infty} p_{m,n}(j) w^j z^n.$$
(2.1)

Next, we introduce some generating functions as follows:

$$g_i^{(1)} = \sum_{j=1}^{\ell_i - 1} (p_i z)^j = \frac{p_i z - (p_i z)^{\ell_i}}{1 - p_i z}, \quad i = 1, 2, \dots, m,$$

$$g_i^{(2)} = \sum_{j=k_1+1}^{\infty} (p_i z)^j = \frac{(p_i z)^{k_i + 1}}{1 - p_i z}, \quad i = 1, 2, \dots, m,$$

$$g_i^{(3)} = \sum_{j=\ell_i}^{k_i} (p_i z)^j = \frac{(p_i z)^{\ell_i} - (p_i z)^{k_i + 1}}{1 - p_i z}, \quad i = 1, 2, \dots, m.$$

Here, we assume $p_i z < 1$ for all i = 1, 2, ..., m. The following result provides the double generating function for the probabilities of the run $E_{\ell_1,k_1} \prec E_{\ell_2,k_2}$.

Theorem 2.1. The double generating function for the probabilities of the run (**R**) with m = 2 is given by

$$\Phi_2(w,z) = \frac{1}{1 - z - (w-1)((p_1 z)^{\ell_1} - (p_1 z)^{k_1 + 1})((p_2 z)^{\ell_2} - (p_2 z)^{k_2 + 1})}.$$
(2.2)

Proof. Let us divide the sequence into three distinct blocks as follows: the first block consists of at most $\ell_i - 1$ consecutive *i*'s, the second block consists of at least $k_i + 1$ consecutive *i*'s, and the third block consists of at least ℓ_i to at most k_i consecutive *i*'s, for i = 1, 2. Then the generating functions for these blocks are $g_i^{(1)}$, $g_i^{(2)}$, and $g_i^{(3)}$, respectively. Thus, $e = [1 \ 1 \ 1 \ 1 \ 1 \ 1]$,

$$\boldsymbol{M} = \begin{pmatrix} 1 & 0 & 0 & -g_1^{(1)} & -g_1^{(1)} & -g_1^{(1)} \\ 0 & 1 & 0 & -g_1^{(2)} & -g_1^{(2)} & -g_1^{(2)} \\ 0 & 0 & 1 & -g_1^{(3)} & -g_1^{(3)} & -g_1^{(3)}w \\ -g_2^{(1)} & -g_2^{(1)} & -g_2^{(1)} & 1 & 0 & 0 \\ -g_2^{(2)} & -g_2^{(2)} & -g_2^{(2)} & 0 & 1 & 0 \\ -g_2^{(3)} & -g_2^{(3)} & -g_2^{(3)} & 0 & 0 & 1 \end{pmatrix},$$

and $\boldsymbol{g} = [g_1^{(1)} g_1^{(2)} g_1^{(3)} g_2^{(1)} g_2^{(2)} g_2^{(3)}]^T$. Observe that the interaction is defined based on the adjacency of the blocks, where $w_{12} = 0$ indicates that these two blocks are forbidden to be adjacent to each other. The same argument applies to cases where the interaction is zero. Additionally, $w_{36} = w$ is used to track the number patterns of the run (**R**) with m = 2. Hence, substituting \boldsymbol{e} , \boldsymbol{M} and \boldsymbol{g} in (1.1), the

result follows.

Remark 2.1. Note that the double generating function presented in (2.2) differs from the one derived in (7) of Kumar and Upadhye [16].. The disparity arises from their consideration of a failure following the pattern $E_{\ell_1,k_1} \prec E_{\ell_2,k_2}$. A similar argument holds to the pattern $E_{\ell_1} \prec E_{\ell_2,k_2}$.

Next, the following corollary can be straightforwardly derived by letting $k_i \to \infty$, $\ell_i = k_i$, and $\ell_i = 1$ for i = 1, 2, in (2.2).

Corollary 2.1. (i) The double generating function for the probabilities of the run $E_{\ell_1} \prec E_{\ell_2}$ is given by

$$\Phi_2(w,z) = \frac{1}{1 - z - (w-1)(p_1 z)^{\ell_1} (p_2 z)^{\ell_2}}.$$
(2.3)

(ii) The double generating function for the probabilities of the run $F_{k_1} \prec F_{k_2}$ is given by

$$\Phi_2(w,z) = \frac{1}{1 - z - (w-1)(1 - p_1 z)(1 - p_2 z)(p_1 z)^{k_1} (p_2 z)^{k_2}}.$$
(2.4)

(iii) The double generating function for the probabilities of the run $G_{k_1} \prec G_{k_2}$ is given by

$$\Phi_2(w,z) = \frac{1}{1 - z - (w-1)(p_1 z)(p_2 z)(1 - (p_1 z)^{k_1})(1 - (p_2 z)^{k_2})}.$$
(2.5)

Remark 2.2. It is worth noting that the double generating functions provided in (2.3), (2.4), and (2.5) coincide with those derived by Dafnis et al. [6] in (3.1), (3.2), and (3.3), respectively.

Further, the following corollary can be obtained by taking $k_2 \rightarrow \infty$ in (2.2).

Corollary 2.2. The double generating function for the probabilities of the run $E_{\ell_1,k_1} \prec E_{\ell_2}$ is given by

$$\Phi_2(w,z) = \frac{1}{1 - z - (w-1)(p_1 z)^{\ell_1} (p_2 z)^{\ell_2} (1 - (p_1 z)^{k_1 - \ell_1 + 1})}.$$
(2.6)

Remark 2.3. Note that the double generating functions presented in (2.6) coincide with one derived by Kumar and Upadhye [16] in (3). Moreover, the obtained result in Theorem 2.1 extends to yield the double generating function for any type of (k_1, k_2) -runs with all at least, exactly, and at most cases.

It is important to highlight the advantage of this technique in connecting one pattern with another. If we aim to add one more pattern to previously obtained patterns, then information about the generating function of the pattern appearing at the right end of the sequence becomes crucial. This allows us to establish connections between adjacent blocks, facilitating the derivation of generating functions for more generalized patterns. For instance, in determining the generating function of the pattern $E_{\ell_1,k_1} \prec E_{\ell_2,k_2} \prec E_{\ell_3,k_3}$, we require the generating functions of the pattern $E_{\ell_1,k_1} \prec E_{\ell_2,k_2}$ appearing at the end of the sequence. Furthermore, we can derive the generating function of the run $E_{\ell_1,k_1} \prec E_{\ell_2,k_2} \prec E_{\ell_3,k_3}$ by adding E_{ℓ_3,k_3} to the right end sequence of $E_{\ell_1,k_1} \prec E_{\ell_2,k_2}$. This iterative process continues until we obtain the generating function of the pattern $E_{\ell_1,k_1} \prec E_{\ell_2,k_2} \prec \ldots \prec E_{\ell_m,k_m}$. Hence, our initial focus is on computing the generating functions of the pattern (**R**) at the right end of the sequence. The following theorem will provide the generating function of the pattern (**R**) that appears to the right end of the sequence.

Theorem 2.2. The generating function for the probabilities of the pattern $E_{\ell_1,k_1} \prec E_{\ell_2,k_2} \prec \ldots \prec E_{\ell_m,k_m}$ appearing at the right end of the sequence is given by

$$Y_m(z) = \frac{\prod_{i=1}^m ((p_i z)^{\ell_i} - (p_i z)^{k_i + 1})}{(1 - z \sum_{i=1}^m p_i) \prod_{i=2}^m (1 - p_i z)}.$$
(2.7)

The generating function for the complement of patterns that do not contain such a sequence is given by

$$N_m(z) = \frac{z \left(\sum_{i=1}^m p_i\right) \prod_{i=2}^m (1 - p_i z) - \prod_{i=1}^m ((p_i z)^{\ell_i} - (p_i z)^{k_i + 1})}{(1 - z \sum_{i=1}^m p_i) \prod_{i=2}^m (1 - p_i z)}.$$

Proof. We prove the result using induction of m. For m = 2, we have $e = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 \end{bmatrix}$,

$$\boldsymbol{M} := \begin{pmatrix} 1 & 0 & 0 & -g_1^{(1)} & -g_1^{(1)} & -g_1^{(1)} & -g_1^{(1)} \\ 0 & 1 & 0 & -g_1^{(2)} & -g_1^{(2)} & -g_1^{(2)} \\ 0 & 0 & 1 & -g_1^{(3)} & -g_1^{(3)} & -g_1^{(2)} \\ 0 & 0 & 1 & -g_1^{(3)} & -g_1^{(3)} & -g_1^{(3)} \\ -g_2^{(1)} & -g_2^{(1)} & -g_2^{(1)} & 1 & 0 & 0 \\ -g_2^{(2)} & -g_2^{(2)} & -g_2^{(2)} & 0 & 1 & 0 & 0 \\ -g_2^{(3)} & -g_2^{(3)} & -g_2^{(3)} & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

and $\boldsymbol{g} = [g_1^{(1)} g_1^{(2)} g_1^{(3)} g_2^{(1)} g_2^{(2)} g_2^{(3)} g_2^{(3)}]^T$. The matrix \boldsymbol{M} is generated using a similar concept applied in Theorem 2.1. Here, we introduce a special block E_{ℓ_3,k_3} that appears to the right of the sequence. It is important to note that this block has zero interaction with any other blocks. The variable u is used to track the pattern E_{ℓ_1,k_1} from the left and E_{ℓ_2,k_2} from the right, with the additional requirement of the block at the right of the sequence. Computing $\boldsymbol{e}\boldsymbol{M}^{-1}\boldsymbol{g}$ will provide the generating function in terms of a polynomial in u, where the coefficient of u yields to $Y_2(m)$. For more details, see Lemma 1 of Kong [11]. With some routine calculations, the coefficient of u in $\boldsymbol{e}\boldsymbol{M}^{-1}\boldsymbol{g}$ can be expressed as

$$Y_2(z) = \frac{\prod_{i=1}^2 ((p_i z)^{\ell_i} - (p_i z)^{k_i + 1})}{(1 - z) (1 - p_2 z)}.$$

The complement generating function $N_2(z)$ can be computed by subtracting $Y_2(z)$ from the full system. It is expressed as

$$N_2(z) = \frac{(p_1 + p_2)z}{1 - (p_1 + p_2)z} - Y_2(z) = \frac{z(p_1 + p_2)(1 - p_2z) - \prod_{i=1}^m ((p_i z)^{\ell_i} - (p_i z)^{k_i + 1})}{(1 - z(p_1 + p_2))(1 - p_2z)}$$

Hence, the result is true for m = 2. Assuming the results hold for m = k - 1, let us consider $e = [1 \ 1 \ 1 \ 1 \ 1 \ 1]$,

$$\boldsymbol{M} := \begin{pmatrix} 1 & 0 & -N_{k-1}(z) & -N_{k-1}(z) & -N_{k-1}(z) & -N_{k-1}(z) \\ 0 & 1 & -Y_{k-1}(z) & -Y_{k-1}(z) & -Y_{k-1}(z)u \\ -g_k^{(1)} & -g_k^{(1)} & 1 & 0 & 0 & 0 \\ -g_k^{(2)} & -g_k^{(2)} & 0 & 1 & 0 & 0 \\ -g_k^{(3)} & -g_k^{(3)} & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix},$$

and $\boldsymbol{g} = \begin{bmatrix} N_{k-1}(z) & Y_{k-1}(z) & g_k^{(1)} & g_k^{(2)} & g_k^{(3)} & g_k^{(3)} \end{bmatrix}^T$. Following the steps similar to the standard case m = 2, the coefficient of u is given by

$$Y_k(z) = \frac{g_k^{(3)} Y_{k-1} \left(1 + g_k^{(1)} + g_k^{(2)} + g_k^{(3)}\right)}{1 - \left(g_k^{(1)} + g_k^{(2)} + g_k^{(3)}\right) \left(Y_{k-1}(z) + N_{k-1}(z)\right)}.$$
(2.8)

The result can be verified for m = k by substituting the following identities in (2.8)

$$g_k^{(1)} + g_k^{(2)} + g_k^{(3)} = \frac{p_k z}{1 - p_k z}$$
 and $Y_{k-1}(z) + N_{k-1}(z) = \frac{(p_1 + p_2 + \dots + p_{k-1})z}{1 - (p_1 + p_2 + \dots + p_{k-1})z}.$

Hence, the results hold for any $m \ge 2$.

The following corollary can be easily verified by taking $k_i \to \infty$, for i = 1, 2, ..., m.

Corollary 2.3. The generating function for the probabilities of the pattern $E_{\ell_1} \prec E_{\ell_2} \prec \ldots \prec E_{\ell_m}$ appearing at the right end of the sequence is given by

$$Y_m(z) = \frac{\prod_{i=1}^m (p_i z)^{\ell_i}}{(1 - z \sum_{i=1}^m p_i) \prod_{i=2}^m (1 - p_i z)}$$

The generating function for the complement of patterns that do not contain such a sequence is given by

$$N_m(z) = \frac{z \left(\sum_{i=1}^m p_i\right) \prod_{i=2}^m (1-p_i z) - \prod_{i=1}^m (p_i z)^{\ell_i}}{(1-z \sum_{i=1}^m p_i) \prod_{i=2}^m (1-p_i z)}$$

Remark 2.4. Note that the generating functions provided in (2.7) coincide with Lemma 1 of Kong [13] for the pattern $E_{\ell_1} \prec E_{\ell_2} \prec \ldots \prec E_{\ell_m}$.

Finally, we are at a stage where we can derive our main result by using Theorems 2.1 and 2.2. The following theorem provide the generating function for the pattern $E_{\ell_1,k_1} \prec E_{\ell_2,k_2} \prec \ldots \prec E_{\ell_m,k_m}$.

Theorem 2.3. The double generating function for the probabilities of the run (**R**) is given by

$$\Phi_m(w,z) = \frac{\prod_{i=2}^{m-1} (1-p_i z)}{(1-z)\prod_{i=2}^{m-1} (1-p_i z) - (w-1)\prod_{i=1}^m ((p_i z)^{\ell_i} - (p_i z)^{k_i+1})}$$

Proof. We prove the result using induction of m. We have already proved the result for m = 2 in Theorem 2.1. For m = 3, let us consider $e = [1 \ 1 \ 1 \ 1 \ 1]$,

$$\boldsymbol{M} := \begin{pmatrix} 1 & 0 & -N_2(z) & -N_2(z) & -N_2(z) \\ 0 & 1 & -Y_2(z) & -Y_2(z) & -Y_2(z)w \\ -g_3^{(1)} & -g_3^{(1)} & 1 & 0 & 0 \\ -g_3^{(2)} & -g_3^{(2)} & 0 & 1 & 0 \\ -g_3^{(3)} & -g_3^{(3)} & 0 & 0 & 1 \end{pmatrix}$$

and $\boldsymbol{g} = \begin{bmatrix} N_2(z) & Y_2(z) & g_3^{(1)} & g_3^{(2)} & g_3^{(3)} \end{bmatrix}^T$. Following the steps similar to Theorem 2.1, we get

$$\Phi_3(w,z) = \frac{(1-p_2z)}{(1-z)(1-p_2z) - (w-1)\prod_{i=1}^3 ((p_iz)^{\ell_i} - (p_iz)^{k_i+1})}$$

Thus, the result is true for m = 3. Assume the result holds for m = k - 1. Now, consider $e = [1 \ 1 \ 1 \ 1 \ 1]$,

$$\boldsymbol{M} := \begin{pmatrix} 1 & 0 & -N_{k-1}(z) & -N_{k-1}(z) & -N_{k-1}(z) \\ 0 & 1 & -Y_{k-1}(z) & -Y_{k-1}(z) & -Y_{k-1}(z)w \\ -g_k^{(1)} & -g_k^{(1)} & 1 & 0 & 0 \\ -g_k^{(2)} & -g_k^{(2)} & 0 & 1 & 0 \\ -g_k^{(3)} & -g_k^{(3)} & 0 & 0 & 1 \end{pmatrix}$$

and $\boldsymbol{g} = \begin{bmatrix} N_{k-1}(z) & Y_{k-1}(z) & g_k^{(1)} & g_k^{(2)} & g_k^{(3)} \end{bmatrix}^T$. Substituting $N_{k-1}(z)$ and $Y_{k-1}(z)$ from Theorem 2.2 and following the steps similar as above, the result follows for m = k. Hence, the result holds for any $m \ge 2$.

The following corollary can be easily verified by letting $k_i \to \infty$, for i = 1, 2, ..., m in Theorem 2.3.

Corollary 2.4. The double generating function for the probabilities of the run $E_{\ell_1} \prec E_{\ell_2} \prec \cdots \prec E_{\ell_m}$ is given by

$$\Phi_m(w,z) = \frac{\prod_{i=2}^{m-1} (1-p_i z)}{(1-z)\prod_{i=2}^{m-1} (1-p_i z) - (w-1)\prod_{i=1}^m (p_i z)^{\ell_i}}.$$
(2.9)

Remark 2.5. Observe that the generating functions provided in (2.9) coincide with Theorem 1 of Kong [13] for the pattern $E_{\ell_1} \prec E_{\ell_2} \prec \ldots \prec E_{\ell_m}$.

As special cases, the following corollary can be proved by taking $\ell_i = k_i$ and $\ell_i = 1$, for i = 1, 2, ..., m in Theorem 2.3.

Corollary 2.5. (i) The double generating function for the probabilities of the run $F_{k_1} \prec F_{k_2} \prec \cdots \prec F_{k_m}$ is given by

$$\Phi_m(w,z) = \frac{\prod_{i=2}^{m-1} (1-p_i z)}{(1-z)\prod_{i=2}^{m-1} (1-p_i z) - (w-1)\prod_{i=1}^m (p_i z)^{\ell_i} (1-p_i z)}.$$

(ii) The double generating function for the probabilities of the run $G_{k_1} \prec G_{k_2} \prec \cdots \prec G_{k_m}$ is given by

$$\Phi_m(w,z) = \frac{\prod_{i=2}^{m-1} (1-p_i z)}{(1-z)\prod_{i=2}^{m-1} (1-p_i z) - (w-1)\prod_{i=1}^m (p_i z)(1-(p_i z)^{k_i})}.$$

- **Remark 2.6.** (*i*) Corollary 2.4 and Corollary 2.5 extend the patterns previously considered by Dafnis et al. [6] (for two-state trials) to multi-state trials.
 - (ii) Using a similar argument as previously discussed $(k_i \to \infty, \ell_i = k_i, \text{ and } \ell_i = 1, \text{ for } i \subseteq \{1, 2, \ldots, m\}$), we can derive the generating function for various types of (k_1, k_2, \ldots, k_m) -runs with all cases of at least, exactly, and at most occurrences.

Next, let α_j denote the coefficient of z^j in $\prod_{i=2}^{m-1}(1-p_iz)$, which is easy to compute given the value of m. It is important to note that $\alpha_0 = 1$. Therefore, we have

$$\prod_{i=2}^{m-1} (1 - p_i z) = 1 + \sum_{j=1}^{m-2} \alpha_j z^j.$$
(2.10)

Next, using Theorem 2.3 and the definition of double generating function given in (2.1), the following corollary can be easily verified.

Corollary 2.6. The recursive relation in the probability generation function $\phi_{m,n}(\cdot)$ is given by

$$\phi_{m,n}(w) + \sum_{j=1}^{m-2} (\alpha_j - \alpha_{j-1}) \phi_{m,n-j}(w) - \alpha_{m-2} \phi_{m,n-m+1}(w)$$

= $(w-1) \left(\prod_{i=1}^m p_i^{\ell_i}\right) \phi_{m,n-\ell}(w) - (w-1) \left(\prod_{i=1}^m p_i^{k_i+1}\right) \phi_{m,n-k-m}(w), \text{ for } n \ge \ell$

with initial condition $\phi_{m,n}(w) = 1$, for $n < \ell$ where $\ell := \ell_1 + \ell_2 + \dots + \ell_m$, $k := k_1 + k_2 + \dots + k_m$.

Using Corollary 2.6, we can easily derive the following result.

Corollary 2.7. The recursive relation in the probability mas function $p_{m,n}(\cdot)$ is given by

$$p_{m,n}(s) + \sum_{j=1}^{m-2} (\alpha_j - \alpha_{j-1}) p_{m,n-j}(s) - \alpha_{m-2} p_{m,n-m+1}(s) = \left(\prod_{i=1}^m p_i^{\ell_i}\right) (p_{m,n-\ell}(s-1) - p_{m,n-\ell}(s)) - \left(\prod_{i=1}^m p_i^{k_i+1}\right) (p_{m,n-k-m}(s-1) - p_{m,n-k-m}(s)),$$

for $s \ge 0$ and $n \ge \ell$ with initial condition $p_{m,0}(0) = 1$ and $p_{m,n}(s) = 0$ for s > 0 and $n < \ell$.

Next, let $\mu_{n,m,r} = \mathbb{E}((X_{m;\ell_1:k_1,\ell_2:k_2,\ldots,\ell_m:k_m}^{(n)})^r)$, the *r*th non-central moment of $X_{m;\ell_1:k_1,\ell_2:k_2,\ldots,\ell_m:k_m}^{(n)}$. The following result provide the recursive relation in $\mu_{n,m,r}$.

Corollary 2.8. The moments $\mu_{n,m,r}$ satisfies the following recursive relation

$$\mu_{m,n,r} + \sum_{j=1}^{m-2} (\alpha_j - \alpha_{j-1}) \mu_{m,n-j,r} - \alpha_{m-2} \mu_{m,n-m+1,r}$$

$$= \left(\prod_{i=1}^m p_i^{\ell_i}\right) \sum_{u=0}^{r-1} \binom{r}{u} \mu_{m,n-\ell,u} - \left(\prod_{i=1}^m p_i^{k_i+1}\right) \sum_{u=0}^{r-1} \binom{r}{u} \mu_{m,n-k-m,u}, \quad \text{for } r \ge 1 \text{ and } n \ge \ell$$

with initial condition $\mu_{m,n,0} = 1$ and $\mu_{m,n,r} = 0$ for $r \ge 1$ and $n < \ell$.

Remark 2.7. Note that if we are interested in a specific pattern related to at least, exactly, and at most cases then we can get the pattern by applying $k_i \to \infty$, $\ell_i = k_i$, and $\ell_i = 1$ for $i \subseteq \{1, 2, ..., m\}$. Consequently, all the results mentioned above can be derived accordingly for that particular pattern. Here, observe that the notation $i \subseteq \{1, 2, ..., m\}$ indicates that it corresponds to the pattern under consideration. For instance, if we are dealing the pattern $E_{\ell_1} \prec F_{\ell_2} \prec E_{\ell_3} \prec E_{\ell_4,k_4} \prec G_{k_5}$, then we set $k_1, k_3 \to \infty$, $\ell_2 = k_2$, and $\ell_5 = 1$ to derive the relevant results.

3 Waiting time Distribution of $X_{m;\ell_1:k_1,\ell_2:k_2,...,\ell_m:k_m}^{(n)}$

In this section, our focus lies on determining the waiting time distribution for the pattern (**R**). We adopt a similar technique used by Chadjiconstantinidis and Eryilmaz [5] to derive the results. Recall that $\xi_1, \xi_2, \ldots, \xi_n$ is a sequence of *iid* multinomial random variables, each taking values from the set $\{1, 2, \ldots, m\}$, with probability $\mathbb{P}(\xi_i = j) = p_j$, for $i = 1, 2, \ldots, n, j = 1, 2, \ldots, m$, and $\sum_{j=1}^m p_j = 1$. The random variable $X_{m;\ell_1:k_1,\ell_2:k_2,\ldots,\ell_m:k_m}^{(n)}$ denotes the number of occurrences of the run (**R**) with $p_{m,n}(s) = \mathbb{P}(X_{m;\ell_1:k_1,\ell_2:k_2,\ldots,\ell_m:k_m}^{(n)} = s)$. Further, let T_r denote the waiting time for the *r*th occurrence of the run (**R**). Thus, we express T_r as follows:

$$T_r = \sum_{i=1}^r W_i,\tag{3.1}$$

where $W_1 = T_1$ is the waiting time for the first occurrence of the pattern (**R**) and $W_i = T_i - T_{i-1}$, i = 2, 3, ..., r, are the inter-arrival time. Clearly, $T_r \ge r(\ell_1 + \ell_2 + \cdots + \ell_m)$. It is important to observe that since the underlying sequence consists of *iid* multi-state random variables, W_i 's are also *iid* random variables. It is easy to verified that

$$\mathbb{P}(T_r > n) = \mathbb{P}(X_{m;\ell_1:k_1,\ell_2:k_2,...,\ell_m:k_m}^{(n)} < r).$$

However, computing waiting time probabilities using the above expression is practically challenging. Therefore, we employ double generating function to compute the probability generating function (pgf) of T_r . Let us define

$$\psi_r(z) := \sum_{s=0}^{\infty} h_r(s) z^s = \sum_{s=0}^{\infty} \mathbb{P}(T_r = s) z^s,$$

Following the steps similar to the proof of Lemma 1 of Chadjiconstantinidis and Eryilmaz [5], the following lemma can be easily verified.

Lemma 3.1. Let ψ_1 represent the pgf for the waiting time random variable T_1 . Then

$$\psi_1(z) = 1 - (1-z)\Phi_m(0,z) = \frac{\prod_{i=1}^m ((p_i z)^{\ell_i} - (p_i z)^{k_i+1})}{(1-z)\prod_{i=2}^{m-1} (1-p_i z) + \prod_{i=1}^m ((p_i z)^{\ell_i} - (p_i z)^{k_i+1})}.$$

It is known that W_i 's are *iid* random variables. Therefore, using (3.1), the following theorem can be readily established.

Theorem 3.1. For $r \ge 1$, the pgf of T_r is given by

$$\psi_r(z) = \left(\frac{\prod_{i=1}^m ((p_i z)^{\ell_i} - (p_i z)^{k_i + 1})}{(1 - z)\prod_{i=2}^{m-1} (1 - p_i z) + \prod_{i=1}^m ((p_i z)^{\ell_i} - (p_i z)^{k_i + 1})}\right)^r.$$
(3.2)

Remark 3.1. If we allow $k_i \to \infty$, for i = 1, 2, ..., m then the pgf of T_r for the pattern $E_{\ell_1} \prec E_{\ell_2} \prec \cdots \prec E_{\ell_m}$ can be expressed as

$$\psi_r(z) = \left(\frac{\prod_{i=1}^m (p_i z)^{\ell_i}}{(1-z)\prod_{i=2}^{m-1} (1-p_i z) + \prod_{i=1}^m (p_i z)^{\ell_i}}\right)^r,$$

which coincides with Theorem 1 of Chadjiconstantinidis and Eryilmaz [5], as expected.

Further, by substituting $\ell_i = k_i$ and $\ell_i = 1$ for i = 1, 2, ..., m in Theorem 3.1, we obtain the following corollary.

Corollary 3.1. (i) For $r \ge 1$, the pgf of T_r for the pattern $F_{k_1} \prec F_{k_2} \prec \cdots \prec F_{k_m}$ is given by

$$\psi_r(z) = \left(\frac{\prod_{i=1}^m (p_i z)^{k_i} (1 - p_i z)}{(1 - z) \prod_{i=2}^{m-1} (1 - p_i z) + \prod_{i=1}^m (p_i z)^{k_i} (1 - p_i z)}\right)^r.$$

(i) For $r \ge 1$, the pgf of T_r for the pattern $G_{k_1} \prec G_{k_2} \prec \cdots \prec G_{k_m}$ is given by

$$\psi_r(z) = \left(\frac{\prod_{i=1}^m (p_i z)(1 - (p_i z)^{k_i})}{(1 - z)\prod_{i=2}^{m-1} (1 - p_i z) + \prod_{i=1}^m (p_i z)(1 - (p_i z)^{k_i})}\right)^r.$$

Using (2.10) in (3.2), the following corollary can be easily obtained.

Corollary 3.2. (i) The probability mass function of T_r satisfies the following recursive relation

$$h_r(s) + \sum_{j=1}^{m-1} (\alpha_j - \alpha_{j-1}) h_r(s-j) - \alpha_{m-2} h_r(s-m+1) + \prod_{i=1}^m p_i^{\ell_i} (h_r(s-\ell) - h_{r-1}(s-\ell))$$
$$= \prod_{i=1}^m p_i^{k_i+1} (h_r(s-k-m) - h_{r-1}(s-k-m)) - \prod_{i=1}^m p_i^{\ell_i} (h_r(s-\ell) - h_{r-1}(s-\ell)),$$

for $s \ge r\ell$, with initial conditions $h_0(s) = \delta_{s,0}$ and $h_r(s) = 0$, for $s < r\ell$.

(ii) Let $\mu_{r,n} = \mathbb{E}((T_r)^n)$. Then $\mu_{r,n}$ satisfies the following recursive relation

$$\mu_{r,n} - \sum_{u=0}^{n} \binom{n}{u} \mu_{r,u} + \sum_{j=1}^{m-2} \frac{n!}{(n-j)!} \alpha_{j} \mu_{r,n-j} - \sum_{j=1}^{m-2} \frac{n!}{(n-j)!} \alpha_{j} \sum_{u=0}^{n} \binom{n}{u} \mu_{r,n-j}$$

$$=\prod_{i=1}^{m} p_{i}^{k_{i}+1} \sum_{u=0}^{n} \binom{n}{u} (m+k)^{n-u} (\mu_{r,u} - \mu_{r-1,u}) - \prod_{i=1}^{m} p_{i}^{\ell_{i}} \sum_{u=0}^{n} \binom{n}{u} \ell^{n-u} (\mu_{r,u} - \mu_{r-1,u})$$

with initial condition $\mu_{0,n} = \delta_{n,0}$. Here, $\ell = \ell_1 + \ell_2 + \cdots + \ell_m$, $k = k_1 + k_2 + \cdots + k_m$, and $\delta_{s,0}$ is the Kronecker delta function.

Remark 3.2. It is worth to note that if our interest lies in specific patterns pertaining to at least, exactly, and at most cases, we can obtain these patterns by setting $k_i \to \infty$, $\ell_i = k_i$, and $\ell_i = 1$ for $i \subseteq \{1, 2, ..., m\}$. Consequently, all the results mentioned above can be derived accordingly for that particular pattern.

4 An Application to DNA Frequent Patterns

DNA sequencing is an important tool used to determine the nucleic acid sequence, that is, the order of nucleotides in DNA. The sequence consists of four letters: A, C, G, and T, which represent the nucleotides adenine, cytosine, guanine, and thymine, respectively. DNA frequent patterns refer to specific sub-patterns in DNA that occur more frequently than a manually set minimum support degree. For example, CG is a frequent pattern in the DNA sequence {C, G, A, T, C, G} when the minimum support degree is set to 2. If we increase the minimum support degree to more than 4 and aim to find the distribution of occurrences of DNA frequent patterns according to the studied patterns in this paper, then the distribution of (k_1, k_2, k_3, k_4) -runs becomes useful for studying the distribution of such DNA frequent patterns. For more details, see Deng *et al.* [7], Yildiz and Selale [29], and references therein. To illustrate this, let us consider the occurrences of A, C, G, and T in a DNA sequence with probabilities p_1, p_2, p_3 , and p_4 , respectively. Suppose n = 50 and the specific DNA frequent pattern is "ACCGT", that is, a (1, 2, 1, 1)-run with exact occurrences. From Corollary 2, the double generating function of the considered runs is given by:

$$\Phi_4(w,z) = \frac{\prod_{i=2}^3 (1-p_i z)}{(1-z)\prod_{i=2}^3 (1-p_i z) - (w-1)\prod_{i=1}^4 (p_i z)^{k_i} (1-p_i z)},$$
(4.1)

where $k_1 = 1, k_2 = 2, k_3 = k_4 = 1$. Note here that the minimum support degree is set to 5. Hence, the probability that the specific DNA frequent pattern occurs more than five times can be calculated using this expression. For instance, if $p_1 = 0.1, p_2 = 0.3, p_3 = 0.2$, and $p_4 = 0.4$, then the probabilities for the occurrences of the specific DNA frequent pattern are as follows:

| m | $\mathbb{P}(X^{50}_{4;1,2,1,1} = m)$ |
|----|--------------------------------------|
| 5 | 7.17960×10^{-9} |
| 6 | 2.68519×10^{-11} |
| 7 | 4.73516×10^{-14} |
| 8 | 3.03176×10^{-17} |
| 9 | 3.78619×10^{-21} |
| 10 | 6.34034×10^{-27} |

As expected, the probability is very small as we are considering the occurrences of specific patterns more than 4 times. Similarly, different types of patterns used in various applications can be analysed using this approach.

Data Availability

No data was used for the research described in the article.

Acknowledgements

The first author is financially supported by TA fellowship from IIT (BHU) Varanasi, India. The second author is partially supported by SERB Start-up Research Grant (File No. SRG/2023/000236) and MATRICS Research Grant (File No. MTR/2023/000200), India.

References

- Aki, S., Kuboki, H. and Hirano, K. (1984). On discrete distributions of order k. Ann. Inst. Statist. Math. 36, 431–440.
- [2] Antzoulakos, D. L., Bersimis, S. and Koutras, M. V. (2003). Waiting times associated with the sum of success run lengths. Mathematical and Statistical Methods in Reliability, in: Lindqvist, B., Doksum, K. (Eds.), 141–157. World Scientific, Singapore.
- [3] Antzoulakos, D. L. and Chadjiconstantinidis, S. (2001). Distributions of numbers of success runs of fixed length in Markov dependent trials. *Ann. Inst. Statist. Math.* **53**, 599–619.
- [4] Balakrishnan, N. and Koutras, M. V. (2002). *Runs and Scans with Applications*. John Wiley, New York.

- [5] Chadjiconstantinidis, S. and Eryilmaz, S. (2023). Computing waiting time probabilities related to (k_1, k_2, \ldots, k_l) pattern. *Statist. Papers* **64**, 1373–1390.
- [6] Dafnis, S. D., Antzoulakos, D. L. and Philippou, A. N. (2010). Distribution related to (k_1, k_2) events. J. Stat. Plan. Inference **140**, 1691–1700.
- [7] Deng, N., Chen, X., Li, D. and Xiong, C. (2019). Frequent patterns mining in dna sequence. *IEEE* Access 7, 108400–108410.
- [8] Fu, J. C. and Johnson, B. C. (2009). Approximate probabilities for runs and patterns in I.I.D. and Markov-dependent multistate trials. *Adv. in Appl. Probab.* 41, 292–308.
- [9] Fu, J. C. and Koutras, M. V. (1994). Distribution theory of runs: a Markov chain approach. J. *Amer. Statist. Assoc.* **89**, 1050–1058.
- [10] Huang, W. T. and Tsai, C. S. (1991). On a modified binomial distribution of order k. Statist. Probab. Lett. 11, 125–131.
- [11] Kong, Y. (2001). A simple method for evaluating partition functions of linear polymers. J. Phys. Chem. B 105, 10111–10114.
- [12] Kong, Y. (2006). Distribution of runs and longest runs: a new generating function approach. J. *Amer. Statist. Assoc.* **101**, 1253–1263.
- [13] Kong, Y. (2022). Multiple consecutive runs of multi-state trials: distributions of $(k_1, k_2, \ldots, k_\ell)$ patterns. J. Comput. Appl. Math. 403, Paper No. 113846, 10.
- [14] Kumar, A. N. and Kumar, P. (2024). A negative binomial approximation to the distribution of the sum of maxima of indicator random variables. *Statist. Probab. Lett.* 208, Paper No. 110040, 6.
- [15] Kumar, A. N. and Upadhye, N. S. (2017). On perturbations of stein operator. Comm. Statist. Theory Methods 46, 9284–9302.
- [16] Kumar, A. N. and Upadhye, N. S. (2019). Generalizations of distributions related to (k_1, k_2) -runs. *Metrika* **82**, 249–268.
- [17] Kumar, A. N. and Upadhye, N. S. (2022). On discrete Gibbs measure approximation to runs. *Comm. Statist. Theory Methods* 51, 1488–1513.
- [18] Kumar, A. N., Upadhye, N. S. and Vellaisamy, P. (2022). Approximations related to the sums of *m*-dependent random variables. *Braz. J. Probab. Stat.* 36, 349–368.

- [19] Liaudanskaitė, G. and Čekanavičius, V. (2022). Compound Poisson approximations to sums of extrema of Bernoulli variables. *Lith. Math. J.* 62, 481–499.
- [20] Makri, F. S., Philippou, A. N. and Psillakis, Z. M. (2007). Shortest and longest length of success runs in binary sequences. J. Statist. Plann. Inference 137, 2226–2239.
- [21] Philippou, A. N., Georghiou, C. and Philippou, G. N. (1985). A generalized geometric distribution and some of its properties. In: *Proceedings of the seventh conference on probability theory* (*Braşov, 1982*), VNU Sci. Press, Utrecht, 559–563.
- [22] Upadhye, N. S. and Kumar, A. N. (2018). Pseudo-binomial approximation to (k_1, k_2) -runs. *Statist. Probab. Lett.* **141**, 19–30.
- [23] Upadhye, N. S., Čekanavičius, V. and Vellaisamy, P. (2017). On Stein operators for discrete approximations. *Bernoulli* 23, 2828–2859.
- [24] Čekanavičius, V. and Vellaisamy, P. (2015). Discrete approximations for sums of *m*-dependent random variables. *ALEA Lat. Am. J. Probab. Math. Stat.* **12**, 765–792.
- [25] Čekanavičius, V. and Vellaisamy, P. (2019). On large deviations for sums of discrete *m*-dependent random variables. *Stochastics* **91**, 1092–1108.
- [26] Čekanavičius, V. and Vellaisamy, P. (2020). Lower bounds for discrete approximations to sums of *m*-dependent random variables. *Probab. Math. Statist.* 40, 23–35.
- [27] Vellaisamy, P. (2004). Poisson approximation for (k_1, k_2) -events via the Stein–Chen method. *J. Appl. Probab.* **41**, 1081–1092.
- [28] Wang, X. and Xia, A. (2008). On negative approximation to k-runs. J. Appl. Probab. 45, 456–471.
- [29] Yildiz, B. and Selale, H. (2011). Mining frequent patterns from microarray data. In: *Proceedings* of the 6th International Symposium on Health Informatics and Bioinformatics, 116–119.