Scaling Limits of Disorder Relevant Non-Binary Spin Systems

Liuyan Li $^{\rm 1}$

Vlad Margarint² Rongfeng Sun³

October 2, 2024

Abstract

In [7], Caravenna, Sun and Zygouras gave general criteria for the partition functions of binary valued spin systems with a relevant random field perturbation to have non-trivial continuum and weak disorder limits. In this work, we show how these criteria can be extended to non-binary valued spin systems.

MSC 2010. Primary: 82B44, Secondary: 82D60, 60K35. *Keywords.* Continuum Limit, Disorder Relevance, Polynomial Chaos, Wiener Chaos.

1 Introduction

We consider here equilibrium statistical mechanics models defined on a lattice, which interact with a random environment (disorder) in the form of a random external field. If we consider the random field as a perturbation of the underlying model without disorder, then such disorder perturbation is called a *relevant perturbation* if the presence of disorder, regardless of its strength, changes the large scale qualitative behaviour of the model. The disorder perturbation is called *irrelevant* if there is qualitative change of large scale behaviour only when the disorder is strong enough. Disorder relevance vs irrelevance often depends on the dimension of the underlying model. At the critical dimension, whether disorder is relevant or irrelevant is much more subtle and is often referred to as *marginal relevance* or *irrelevance*. We refer the reader to [9] for more background.

In [7], it was observed that for binary-valued spin systems experiencing a (non-marginal) relevant random field perturbation, it should be possible to tune the disorder strength to zero at a suitable rate as the lattice spacing tends to zero, such that one obtains a continuum model with non-trivial dependence on disorder. In [7], the authors formulated general convergence criteria for the partition functions of the disordered model to have non-trivial continuum limits, which opened the door to defining the continuum disordered model. Examples that fall into this framework include the directed polymer model (DPM) in dimension 1 + 1 [2, 1] which provided inspiration for [7], the disordered pinning model [7, 6], the long-range DPM in dimension 1 + 1 [7], and the random field perturbation of the critical 2-dim Ising model [7, 3].

One limitation of [7] is that the results were formulated for spin systems with binary valued spins, which played essential role in the proof. Our purpose here is to extend [7] and formulate convergence criteria for random field perturbations of non-binary valued spin systems, such

¹School of Statistics and Mathematics, Guangdong University of Finance and Economics, Guangzhou, 510320, P. R. China. E-mail: mathlily@gdufe.edu.cn

 $^{^2 {\}rm Office}$ 350E, Fretwell Building, University of North Carolina Charlotte, USA. Email: vmar-gari@charlotte.edu

³Department of Mathematics, National University of Singapore, 10 Lower Kent Ridge Road, 119076 Singapore. Email: matsr@nus.edu.sg

that their partition functions admit non-trivial continuum limits. One model with non-binary spins is the Blume-Capel model (see [4, 10] and the rereferences therein), although verifying our convergence criteria would require fine control of the model at criticality that is currently out of reach.

We also note here the parallel between disordered systems and the theory of singular stochastic partial differential equations (SPDEs) [11], where the driving noise plays the role of disorder perturbation of the underlying PDE, and the notions of sub-criticality, criticality, and super-criticality for singular SPDEs correspond respectively to the notions of disorder relevance, marginal relevance/irrelevance, and disorder irrelevance for disordered systems. The directed polymer model lies at the intersection of both worlds, and the interplay between the two different perspectives was instrumental in recent progress in defining the solution of the stochastic heat equation in the critical/marginal dimension d = 2 [8].

1.1 Setup

We will follow the same setup as in [7]. First we introduce the underlying spin system. For $d \geq 1$, we consider an open and simply connected domain $\Omega \subseteq \mathbb{R}^d$, and we define its lattice approximation by $\Omega_{\delta} := \Omega \cap (\delta \mathbb{Z})^d$ for $\delta > 0$. A spin $\sigma_x \in \mathbb{R}$ is assigned to each $x \in \Omega_{\delta}$, and let $P_{\Omega_{\delta}}^{\text{ref}}$ (with expectation $E_{\Omega_{\delta}}^{\text{ref}}$) be a probability measure on the spin configuration $\sigma = (\sigma_x)_{x \in \Omega_{\delta}}$. Typically, $P_{\Omega_{\delta}}^{\text{ref}}$ will be chosen to be the law of an equilibrium spin system at the critical point of a continuous phase transition, such that the spin field σ admits a non-trivial continuum limit as $\delta \downarrow 0$. Sometimes we will drop the sub and superscripts in $P_{\Omega_{\delta}}^{\text{ref}}$ to simplify the notation. For $x \in \Omega$, we will denote by x^{δ} the point in $\Omega_{\delta} = \Omega \cap (\delta \mathbb{Z})^d$ that is closest to x (fix any convention to break the tie if such x^{δ} is not unique).

Next, we introduce the random field (disorder), which is given by a family of i.i.d. random variables $\omega := (\omega_x)_{x \in \Omega_{\delta}}$ with $\mathbb{E}[\omega_x] = 0$ and $\mathbb{E}[\omega_x^2] = 1$. Probability and expectation for ω will be denoted by \mathbb{P} and \mathbb{E} . We assume that ω has finite log moment generating function

$$\phi(\lambda) := \log \mathbb{E}[e^{\lambda \omega_x}] \qquad \text{for all } |\lambda| < \lambda_0 \text{ for some } \lambda_0 > 0.$$
(1.1)

Note that under suitable scaling, the i.i.d. field $(\omega_x)_{x \in (\delta \mathbb{Z})^d}$ converges in the continuum limit to a white noise W on \mathbb{R}^d , which is a Gaussian process $W = (W(f))_{f \in L^2(\mathbb{R}^d)}$ with $\mathbb{E}[W(f)] = 0$ and $\operatorname{Cov}(W(f), W(g)) = \int_{\mathbb{R}^d} f(x)g(x) dx$.

Given the random field (disorder) ω and disorder strength $\lambda > 0$, we can then define the random field perturbation of $P_{\Omega_{\delta}}^{\text{ref}}$ through the following ω -dependent Gibbs measure

$$P^{\omega}_{\Omega_{\delta};\lambda}(\mathrm{d}\sigma) := \frac{e^{\sum_{x\in\Omega_{\delta}}\lambda\omega_{x}\sigma_{x}}}{Z^{\omega}_{\Omega_{\delta};\lambda}} P^{\mathrm{ref}}_{\Omega_{\delta}}(\mathrm{d}\sigma), \tag{1.2}$$

where the normalizing constant, called the partition function, is defined by

$$Z_{\Omega_{\delta};\lambda}^{\omega} := \mathbf{E}_{\Omega_{\delta}}^{\mathrm{ref}} \left[e^{\sum_{x \in \Omega_{\delta}} \lambda \omega_x \sigma_x} \right].$$
(1.3)

The question we want to address is whether there is a suitable choice of $\lambda = \lambda_{\delta} \downarrow 0$, sometimes called intermediate disorder scaling, such that under suitable centering and scaling, $Z_{\Omega_{\delta};\lambda_{\delta}}^{\omega}$ admits non-trivial distributional limits. This is usually the first step in showing that the random Gibbs measure $P_{\Omega_{\delta};\lambda}^{\omega}$ also has a non-trivial disordered continuum limit, where the disorder in the continuum limit is given by the white noise W. For binary-valued spin systems, this question was addressed in [7] where general convergence criteria were formulated. The goal of this paper is to consider the case where the spins σ_x are not binary-valued, although we still assume boundedness to avoid additional technical complications.

(A0) There exists K > 0 such that $P_{\Omega_{\delta}}^{\text{ref}}(\sigma_x \in [-K, K]) = 1$ for all $x \in \Omega_{\delta}$ and $\delta > 0$.

To motivate our main result, we first recall how the convergence of $Z_{\Omega_{\delta};\lambda_{\delta}}^{\omega}$ in law was established in [7, Section 8] for spin systems with $\sigma_x \in \{\pm 1\}$. The starting point is the expansion

$$Z_{\Omega_{\delta};\lambda_{\delta}}^{\omega} = \mathrm{E}_{\Omega_{\delta}}^{\mathrm{ref}} \Big[e^{\sum_{x \in \Omega_{\delta}} \lambda_{\delta} \omega_{x} \sigma_{x}} \Big] = \mathrm{E}_{\Omega_{\delta}}^{\mathrm{ref}} \Big[\prod_{x \in \Omega_{\delta}} e^{\lambda_{\delta} \omega_{x} \sigma_{x}} \Big] = \mathrm{E}_{\Omega_{\delta}}^{\mathrm{ref}} \Big[\prod_{x \in \Omega_{\delta}} (\cosh \lambda_{\delta} \omega_{x} + \sigma_{x} \sinh \lambda_{\delta} \omega_{x}) \Big] = \exp \Big\{ \sum_{x \in \Omega_{\delta}} \log \cosh \lambda_{\delta} \omega_{x} \Big\} \mathrm{E}_{\Omega_{\delta}}^{\mathrm{ref}} \Big[\prod_{x \in \Omega_{\delta}} (1 + \sigma_{x} \tanh \lambda_{\delta} \omega_{x}) \Big],$$
(1.4)

where we linearised $e^{\lambda_{\delta}\omega_x\sigma_x}$ in (1.4) since σ_x is binary valued. Note that $\sum_{x\in\Omega_{\delta}}\log\cosh\lambda_{\delta}\omega_x$ is a sum of i.i.d. random variables with

$$\mathbb{E}\Big[\sum_{x\in\Omega_{\delta}}\log\cosh\lambda_{\delta}\omega_x\Big] = |\Omega_{\delta}|\Big(\frac{\lambda_{\delta}^2}{2} + O(\lambda_{\delta}^4)\Big) \quad \text{and} \quad \mathbb{V}\mathrm{ar}\Big(\sum_{x\in\Omega_{\delta}}\log\cosh\lambda_{\delta}\omega_x\Big) \le C|\Omega_{\delta}|\lambda_{\delta}^4.$$

As long as $|\Omega_{\delta}|\lambda_{\delta}^4 = O(\delta^{-d}\lambda_{\delta}^4) \to 0$ as $\delta \downarrow 0$, i.e., $\lambda_{\delta} = o(\delta^{d/4})$, then

$$e^{-\frac{\lambda_{\delta}^{2}}{2}|\Omega_{\delta}|} \exp\left\{\sum_{x\in\Omega_{\delta}}\log\cosh\lambda_{\delta}\omega_{x}\right\} \to 1 \quad \text{in probability}, \tag{1.5}$$

and hence we can normalize $Z^\omega_{\Omega_\delta;\lambda_\delta}$ and then focus on the distributional limit of

$$\widetilde{Z}_{\Omega_{\delta};\lambda_{\delta}}^{\omega} := \mathrm{E}_{\Omega_{\delta}}^{\mathrm{ref}} \Big[\prod_{x \in \Omega_{\delta}} (1 + \sigma_x \tanh \lambda_{\delta} \omega_x) \Big] = 1 + \sum_{k=1}^{\infty} \sum_{I \subset \Omega_{\delta}, |I|=k} \mathrm{E}_{\Omega_{\delta}}^{\mathrm{ref}}[\sigma_I] \prod_{x \in I} \xi_{\delta}(x), \tag{1.6}$$

where $\sigma_I := \prod_{x \in I} \sigma_x$, and $\xi_{\delta}(x) := \tanh \lambda_{\delta} \omega_x$ is a family of i.i.d. random variables with

$$\mathbb{E}[\xi_{\delta}(x)] = O(\lambda_{\delta}^3) \quad \text{and} \quad \mathbb{V}\mathrm{ar}(\xi_{\delta}(x)) = \lambda_{\delta}^2 + O(\lambda_{\delta}^4) \qquad \text{by Taylor expansion.}$$
(1.7)

The expansion in (1.6) is called a polynomial chaos expansion in the family of random variables ξ_{δ} . Pretending that the mean of $\xi_{\delta}(x)$ is zero (because it is negligible as $\lambda_{\delta} \to 0$), it is an L^2 -orthogonal expansion and can be regarded as the discrete analogue of the Wiener-Itô chaos expansion w.r.t. a white noise W. In particular, if we match mean and variance (up to leading order) and make the approximation

$$\xi_{\delta}(x) \approx \lambda_{\delta} \delta^{-d/2} \int_{\Lambda_{\delta}(x)} W(\mathrm{d}y)$$

where $\Lambda_{\delta}(x)$ denotes the cube of side length δ centered at $x \in \Omega_{\delta}$. Then we have

$$\widetilde{Z}_{\Omega_{\delta};\lambda_{\delta}}^{\omega} \approx 1 + \sum_{k=1}^{\infty} \frac{1}{k!} \int \cdots \int_{\Omega^{k}} \lambda_{\delta}^{k} \delta^{-dk/2} \mathbb{E}_{\Omega_{\delta}}^{\mathrm{ref}}[\sigma_{x_{1}^{\delta}} \cdots \sigma_{x_{k}^{\delta}}] W(\mathrm{d}x_{1}) \cdots W(\mathrm{d}x_{k}), \qquad (1.8)$$

where x_i^{δ} the point in Ω_{δ} closest to $x_i \in \Omega$, and the factor 1/k! arises because Ω^k consists of k! sectors that give identical contributions by exchangeability. Suppose the spin field σ with law $P_{\Omega_{\delta}}^{\text{ref}}$ satisfies the assumption

(A1) There exists $\gamma > 0$ such that for any $k \in \mathbb{N}$, the rescaled k-spin correlation function

$$\psi_{\delta}(x_1,\ldots,x_k) := \mathbf{1}_{\{x_i^{\delta} \neq x_j^{\delta} \forall i \neq j\}} \,\delta^{-k\gamma} \operatorname{E}_{\Omega_{\delta}}^{\operatorname{ref}}[\sigma_{x_1^{\delta}}\ldots\sigma_{x_k^{\delta}}], \quad (x_1,\ldots,x_k) \in \Omega^k, \tag{1.9}$$

converges in $L^2(\Omega^k)$ to some limit $\psi_0: \Omega^k \to \mathbb{R}$. More precisely,

$$\lim_{\delta \downarrow 0} \|\psi_{\delta} - \psi_{0}\|_{L^{2}(\Omega^{k})}^{2} = 0.$$
(1.10)

Note that this assumption ensures that $P_{\Omega_{\delta}}^{\text{ref}}$ is the law of a spin system at the critical point of a continuous phase transition, which admits a non-trivial continuum limit as $\delta \downarrow 0$.

If (A1) is satisfied, then by [7, Theorem 2.3], we can choose $\lambda_{\delta} := \hat{\lambda} \delta^{\frac{d}{2} - \gamma}$ in (1.8) to obtain that

$$\widetilde{Z}^{\omega}_{\Omega_{\delta};\lambda_{\delta}} \xrightarrow{\text{dist}} 1 + \sum_{k=1}^{\infty} \frac{\widehat{\lambda}^{k}}{k!} \int \cdots \int_{\Omega^{k}} \psi_{0}(x_{1},\dots,x_{k}) \,\mathrm{d}W(x_{1})\cdots \,\mathrm{d}W(x_{k}), \tag{1.11}$$

provided this series is convergent in L^2 , and there is uniformity (in δ) in the convergence of the series \sum_k in (1.8) so that the series can be truncated at a large but fixed value of k = M as $\delta \downarrow 0$. This leads to the second assumption

(A2) For any $\lambda > 0$,

$$\lim_{M \to \infty} \limsup_{\delta \downarrow 0} \sum_{k=M+1}^{\infty} \frac{\hat{\lambda}^{2k}}{k!} \|\psi_{\delta}\|_{L^2(\Omega^k)}^2 = 0.$$
(1.12)

Remark 1.1 With $\lambda_{\delta} := \hat{\lambda} \delta^{\frac{d}{2}-\gamma}$, the condition $\lambda_{\delta} = o(\delta^{d/4})$ that ensures (1.5) becomes $\gamma < d/4$. When $P_{\Omega_{\delta}}^{\text{ref}}$ is the law of the critical 2-dim Ising model, assumptions (A1) and (A2) were verified in [7] with $\gamma = 1/8 < d/4 = 1/2$.

1.2 Main result

When the spins σ_x are not binary valued, the linearization step (1.4) can no longer be applied. Furthermore, it is not clear whether it is even possible to normalise the partition function as in (1.5) that could lead to a polynomial chaos expansion as in (1.6). Therefore we will consider instead the modified partition function

$$\widehat{Z}^{\omega}_{\Omega_{\delta};\lambda} := \mathbf{E}^{\mathrm{ref}}_{\Omega_{\delta}} \left[e^{\sum_{x \in \Omega_{\delta}} \left(\lambda \omega_x \sigma_x - \phi(\lambda \sigma_x) \right)} \right], \tag{1.13}$$

which automatically satisfies $\mathbb{E}[\widehat{Z}_{\Omega_{\delta};\lambda}] = 1$. When ω_x are i.i.d. standard Gaussian, such a normalization corresponds to replacing $e^{\lambda\omega_x\sigma_x}$ by the Wick exponential : $e^{\lambda\omega_x\sigma_x} := e^{\lambda\omega_x\sigma_x-\lambda^2\sigma_x^2/2}$. Although this changes the underlying Gibbs measure $P_{\Omega_{\delta},\lambda}^{\omega}$ defined in (1.2) because $\sum_x \phi(\lambda\sigma_x)$ depends on the spin configuration σ , there is intrinsic interest in such a Gibbs measure as seen in the case of the two-dimensional Parabolic Anderson Model (PAM) [15]. We will discuss this in more detail in Remark 1.4.

To show that $\widehat{Z}_{\Omega_{\delta};\lambda}^{\omega}$ defined in (1.13) has a non-trivial continuum limit, our basic strategy is to perform Taylor expansion. In contrast to (1.6), we will no longer have an L^2 -orthogonal expansion due to the presence of higher powers of $\lambda \omega_x \sigma_x$. But the leading order terms are expected to be comparable to the expansion in (1.6). Therefore most of the work goes into controlling the higher order terms in the expansion and show they are negligible as $\delta \downarrow 0$. This requires one more assumption to control the k-spin correlations when some of the k spins coincide:

(A3) For $k \ge 1$, let $x_1^{\delta}, \ldots, x_k^{\delta}$ be distinct points in Ω_{δ} . Let $r_1, \ldots, r_k \in \mathbb{N}$ and denote $(r_i)_2 := r_i \pmod{2} \in \{0, 1\}$. Then there exists a universal constant $C \ge 1$ such that

$$\mathbb{E}_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{i=1}^{k} \sigma_{x_{i}^{\delta}}^{r_{i}} \right] \le C^{\sum_{i=1}^{k} (r_{i} - (r_{i})_{2})} \left| \mathbb{E}_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{i=1}^{k} \sigma_{x_{i}^{\delta}}^{(r_{i})_{2}} \right] \right|.$$
(1.14)

In light of the scaling property in assumption (A1), we believe assumption (A3) can be weakened further by allowing a diverging constant $C = C(\delta)$, possibly as a negative power of δ . But we will not pursue it here.

Remark 1.2 For the Ising model with $\sigma_x \in \{\pm 1\}$, (1.14) holds trivially with C = 1. In general, when spins fuse, we expect there to be a pairing effect that allows one to replace σ_z^r by $\sigma_z^{(r)_2}$ for $r \in \mathbb{N}$. For a centred Gaussian field $(\sigma_{x^\delta})_{x^\delta \in \Omega_\delta}$ with a covariance $\mathbb{C}ov(\sigma_{x^\delta}, \sigma_{y^\delta})$ that decays polynomially in $|x^\delta - y^\delta|/\delta$, this can be verified using Wick's theorem. A similar result has been established for the critical site percolation on the planar triangular lattice, see [5, Theorem 1.1], even though Wick's theorem does not apply in this setting.

We are now ready to state our main result.

Theorem 1.3 Assume that the reference spin measure $P_{\Omega_{\delta}}^{\text{ref}}$ satisfies assumptions (A0)-(A3) for some $K, \gamma > 0$ and ψ_0 , with $\gamma < d/4$. Then with $\lambda = \lambda_{\delta} := \hat{\lambda} \delta^{\frac{d}{2} - \gamma}$, the normalized partition function $\widehat{Z}_{\Omega_{\delta};\lambda_{\delta}}^{\omega}$ defined in (1.13) converges in distribution to a non-trivial limit

$$\mathcal{Z}_{\Omega,\hat{\lambda}}^{W} = 1 + \sum_{k=1}^{\infty} \hat{\lambda}^{k} \int \cdots \int_{\Omega^{k}} \psi_{0}(x_{1}, \dots, x_{k}) \, \mathrm{d}W(x_{1}) \dots \, \mathrm{d}W(x_{k}), \tag{1.15}$$

where W is a white noise on \mathbb{R}^d and the series converges in L^2 .

Remark 1.4 (PAM) The Wick ordering of the exponential in (1.13) has appeared in the study of SPDEs such as the Parabolic Anderson Model (PAM)

$$\frac{\partial}{\partial t}u(t,x) = \frac{1}{2}\Delta u(t,x) + (u \cdot W)(t,x), \qquad t \ge 0, x \in \mathbb{R}^d, \tag{1.16}$$

where W is a white noise on \mathbb{R}^d . The equation is singular in $d \ge 2$ due to the product $u \cdot W$. Before the theory of regularity structures [11], this difficulty is often bypassed by replacing the singular product $u \cdot W$ by the Wick product $u \diamond W$ (see [14, 12, 13, 15]),

$$\frac{\partial}{\partial t}\hat{u}(t,x) = \frac{1}{2}\Delta\hat{u}(t,x) + (\hat{u}\diamond W)(t,x), \qquad t \ge 0, x \in \mathbb{R}^d.$$
(1.17)

By discretising time and space, we obtain approximations of u and \hat{u} which admit the Feynman-Kac representations

$$u^{\delta}(t,x) := \mathbf{E} \Big[e^{\sum_{z \in \delta \mathbb{Z}^d} \lambda_{\delta} \omega_z L(t^{\delta},z)} \Big],$$
$$\hat{u}^{\delta}(t,x) := \mathbf{E} \Big[e^{\sum_{z \in \delta \mathbb{Z}^d} \left(\lambda_{\delta} \omega_z L(t^{\delta},z) - \phi(\lambda_{\delta} L(t^{\delta},z)) \right)} \Big].$$

where $E[\cdot]$ denotes expectation for a random walk on $\delta \mathbb{Z}^d$ with local time $L(t^{\delta}, \cdot)$ at time $t^{\delta} := t\delta^{-2}$, and $(\omega_z)_{z \in \delta \mathbb{Z}^d}$ are i.i.d. standard normals that discretise the white noise W. We note that u^{δ} and \hat{u}^{δ} are instances of $Z^{\omega}_{\Omega_{\delta};\lambda}$ in (1.3) and $\hat{Z}^{\omega}_{\Omega_{\delta};\lambda}$ in (1.13), with spin values $\sigma_z = L(t^{\delta}, z)$ for $z \in \delta \mathbb{Z}^d$. Although \hat{u}^{δ} and its associated Gibbs measure (polymer measure) $\hat{P}^{\omega}_{\delta;\lambda_{\delta}}$ on the underlying random walk is less physical than u^{δ} and its associated polymer measure $P^{\omega}_{\delta;\lambda_{\delta}}$, it was recently pointed out in [15, Example 59] that in the planar case d = 2, one can recover the continuum limit of $P^{\omega}_{\delta;\lambda_{\delta}}$ from that of $\hat{P}^{\omega}_{\delta;\lambda_{\delta}}$ via a change of measure because under the measure $\hat{P}^{\omega}_{\delta;\lambda_{\delta}}$, the weight factor $e^{\sum_{z \in \delta \mathbb{Z}^d} \phi(\lambda_{\delta} L(t^{\delta}, z))}$ has a well-defined limit. This suggests that Theorem 1.3 could also be the first step towards identifying the continuum limit of $Z^{\omega}_{\Omega_{\delta;\lambda}}$.

Remark 1.5 (Unbounded Spins) We can formulate alternative convergence criteria in Theorem 1.3 without the assumption (A0) that the spins σ_x are uniformly bounded. The assumption $|\sigma_x| \leq K$ is only used in the proof of Theorem 1.3 to ensure that $\lambda_{\delta} K$ is sufficiently small as $\delta \downarrow 0$. Since $\lambda_{\delta} = \hat{\lambda} \delta^{\frac{d}{2} - \gamma}$, it is enough that $|\sigma_x| \leq K_{\delta}$ with $K_{\delta} \ll \delta^{\gamma - \frac{d}{2}}$. Therefore we can approximate $\hat{Z}^{\omega}_{\Omega_{\delta};\lambda}$ in (1.13) by

$$\widetilde{Z}_{\Omega_{\delta};\lambda}^{\omega} := \mathbb{E}_{\Omega_{\delta}}^{\mathrm{ref}} \left[e^{\sum_{x \in \Omega_{\delta}} \left(\lambda \omega_x \sigma_x - \phi(\lambda \sigma_x) \right)} \prod_{x \in \Omega_{\delta}} \mathbb{1}_{\{ |\sigma_x| \le K_{\delta} \}} \right].$$
(1.18)

The L^1 -norm of the error of this approximation can be bounded by

$$\mathbb{E}[\widehat{Z}^{\omega}_{\Omega_{\delta};\lambda} - \widetilde{Z}^{\omega}_{\Omega_{\delta};\lambda}] = \mathbb{P}^{\mathrm{ref}}_{\Omega_{\delta}}(|\sigma_{x}| > K_{\delta} \text{ for some } x \in \Omega_{\delta}).$$
(1.19)

Therefore the assumption (A0) can be replaced by the assumption (A0'): For some choice of $K_{\delta} \ll \delta^{\gamma - \frac{d}{2}}$, the r.h.s. of (1.19) tends to 0 as $\delta \downarrow 0$. For the same choice of K_{δ} , we can replace $P_{\Omega_{\delta}}^{\text{ref}}$ by the conditional law

$$\widetilde{\mathrm{P}}_{\Omega_{\delta}}^{\mathrm{ref}}(\cdot) = \mathrm{P}_{\Omega_{\delta}}^{\mathrm{ref}}(\cdot \,|\, \forall \, x \in \Omega_{\delta}, |\sigma_{x}| \le K_{\delta})$$

and rewrite

$$\widetilde{Z}_{\Omega_{\delta};\lambda}^{\omega} = \mathbb{P}_{\Omega_{\delta}}^{\mathrm{ref}} \left(|\sigma_{x}| \leq K_{\delta} \,\forall \, x \in \Omega_{\delta} \right) \cdot \widetilde{\mathrm{E}}_{\Omega_{\delta}}^{\mathrm{ref}} \left[e^{\sum_{x \in \Omega_{\delta}} \left(\lambda \omega_{x} \sigma_{x} - \phi(\lambda \sigma_{x}) \right)} \right].$$

Assumptions (A1)-(A3) should then also be modified accordingly with $E_{\Omega_{\delta}}^{ref}[\cdot]$ therein replaced by $\widetilde{E}_{\Omega_{\delta}}^{ref}[\cdot]$.

2 Proof of Theorem 1.3

In this section, we prove the main result of the paper. Recall from (1.13) the normalised partition function

$$\widehat{Z}^{\omega}_{\Omega_{\delta};\lambda} := \mathbf{E}^{\mathrm{ref}}_{\Omega_{\delta}} \left[e^{\sum_{x \in \Omega_{\delta}} \left(\lambda \omega_{x} \sigma_{x} - \phi(\lambda \sigma_{x}) \right)} \right], \quad \text{where} \quad \lambda = \lambda_{\delta} = \widehat{\lambda} \delta^{\frac{d}{2} - \gamma}.$$

For $x \in \Omega_{\delta}$, let $\eta_x = \eta_x(\sigma, \omega) := e^{\lambda \omega_x \sigma_x - \phi(\lambda \sigma_x)} - 1 - \lambda \omega_x \sigma_x$. The starting point of our analysis is the expansion

$$\widehat{Z}_{\Omega_{\delta};\lambda}^{\omega} = \mathbf{E}_{\Omega_{\delta}}^{\mathrm{ref}} \bigg[\prod_{x \in \Omega_{\delta}} (1 + \lambda \omega_{x} \sigma_{x} + \eta_{x}) \bigg]$$

$$= 1 + \sum_{k=1}^{|\Omega_{\delta}|} \sum_{\substack{\{x_{1}, \dots, x_{k}\} \subset \Omega_{\delta} \\ x_{i} \neq x_{j} \text{ for } i \neq j}} \mathbf{E}_{\Omega_{\delta}}^{\mathrm{ref}} \bigg[\prod_{i=1}^{k} (\lambda \omega_{x_{i}} \sigma_{x_{i}} + \eta_{x_{i}}) \bigg]$$

$$= 1 + \sum_{k=1}^{M} \sum_{\substack{\{x_{1}, \dots, x_{k}\} \subset \Omega_{\delta} \\ x_{i} \neq x_{j} \text{ for } i \neq j}} \mathbf{E}_{\Omega_{\delta}}^{\mathrm{ref}} \bigg[\prod_{i=1}^{k} (\lambda \omega_{x_{i}} \sigma_{x_{i}} + \eta_{x_{i}}) \bigg] + R_{M,\delta}, \qquad (2.1)$$

where we have truncated the sum at index $k = M \in \mathbb{N}$, and $R_{M,\delta}$ denotes the remainder. We will show in Section 2.1 that this truncated sum converges to the series in (1.15) truncated at k = M, i.e.,

$$\sum_{k=1}^{M} \sum_{\substack{\{x_1,\dots,x_k\} \subset \Omega_{\delta} \\ x_i \neq x_j \text{ for } i \neq j}} \operatorname{E}_{\Omega_{\delta}}^{\operatorname{ref}} \left[\prod_{i=1}^{k} (\lambda \omega_{x_i} \sigma_{x_i} + \eta_{x_i}) \right] \xrightarrow{\longrightarrow} \sum_{\delta \downarrow 0}^{M} \hat{\lambda}^k \int \cdots \int_{\Omega^k} \psi_0(x_1,\dots,x_k) \, \mathrm{d}W(x_1) \dots \, \mathrm{d}W(x_k),$$

$$(2.2)$$

and we will show in Section 2.2 that

$$\lim_{M \to \infty} \limsup_{\delta \downarrow 0} \mathbb{E}[R_{M,\delta}^2] = 0.$$
(2.3)

Since assumptions (A1)-(A2) imply that the series in (1.15) is convergent in L^2 , the conclusion of Theorem 1.3 follow immediately from (2.2) and (2.3).

2.1 Convergence of the truncated sum

In this subsection, we verify (2.2). In the truncated sum in (2.1), for each $1 \le k \le M$, we can further decompose the k-th term into

$$\sum_{\substack{\{x_1,\dots,x_k\}\subset\Omega_{\delta}\\x_i\neq x_j \text{ for } i\neq j}} \operatorname{E}_{\Omega_{\delta}}^{\operatorname{ref}} \Big[\prod_{i=1}^{k} (\lambda\omega_{x_i}\sigma_{x_i} + \eta_{x_i}) \Big] = \sum_{\substack{\{x_1,\dots,x_k\}\subset\Omega_{\delta}\\x_i\neq x_j \text{ for } i\neq j}} \lambda^k \operatorname{E}_{\Omega_{\delta}}^{\operatorname{ref}} [\sigma_{x_1}\dots\sigma_{x_k}] \prod_{i=1}^{k} \omega_{x_i} + \sum_{\substack{\{x_1,\dots,x_k\}\subset\Omega_{\delta}\\x_i\neq x_j \text{ for } i\neq j}} \sum_{\substack{(I,J)\vdash\{x_1,\dots,x_k\}\\I\neq\emptyset}} \operatorname{E}_{\Omega_{\delta}}^{\operatorname{ref}} \Big[\prod_{u\in I} \eta_u \prod_{v\in J} \lambda\omega_v \sigma_v \Big], \quad (2.4)$$

where $(I, J) \vdash \{x_1, \ldots, x_k\}$ denotes a partition with $I \cup J = \{x_1, \ldots, x_k\}$ and $I \cap J = \emptyset$.

The first term in (2.4) is in fact the dominant term. Indeed, if we sum it over $1 \le k \le M$, then by the choice of $\lambda = \hat{\lambda} \delta^{\frac{d}{2} - \gamma}$ and the definition of ψ_{δ} in (1.9), we have

$$\sum_{k=1}^{M} \sum_{\substack{\{x_1,\dots,x_k\} \subset \Omega_\delta \\ x_i \neq x_j \text{ for } i \neq j}} \lambda^k \mathrm{E}_{\Omega_\delta}^{\mathrm{ref}}[\sigma_{x_1}\dots\sigma_{x_k}] \prod_{i=1}^k \omega_{x_i} = \sum_{k=1}^{M} \hat{\lambda}^k \sum_{\substack{\{x_1,\dots,x_k\} \subset \Omega_\delta \\ x_i \neq x_j \text{ for } i \neq j}} \psi_\delta(x_1,\dots,x_k) \prod_{i=1}^k (\delta^{\frac{d}{2}} \omega_{x_i})$$
$$\implies \sum_{\delta \downarrow 0}^M \hat{\lambda}^k \int \cdots \int_{\Omega^k} \psi_0(x_1,\dots,x_k) \, \mathrm{d}W(x_1) \dots \, \mathrm{d}W(x_k),$$

where the convergence follows from assumption (A1) and [7, Theorem 2.3].

Therefore to prove (2.2), it only remains to show that for each $k \in \mathbb{N}$, the second term in (2.4) satisfies

$$\mathcal{E}_{k,\delta} := \sum_{\substack{\{x_1,\dots,x_k\} \subset \Omega_{\delta} \\ x_i \neq x_j \text{ for } i \neq j}} \sum_{\substack{(I,J) \vdash \{x_1,\dots,x_k\} \\ I \neq \emptyset}} \mathbb{E}_{\Omega_{\delta}}^{\text{ref}} \Big[\prod_{u \in I} \eta_u \prod_{v \in J} \lambda \sigma_v \omega_v \Big] \xrightarrow{\delta \downarrow 0} 0 \quad \text{in probability.}$$
(2.5)

To prove this, we first observe that for two different sets $\{x_1, \ldots, x_k\} \neq \{\tilde{x}_1, \ldots, \tilde{x}_k\}$, the corresponding summands in (2.5) are L^2 -orthogonal to each other.

Lemma 2.1 Let $I, J, \tilde{I}, \tilde{J} \subset \Omega_{\delta}$ satisfy $I \cap J = \emptyset$, $\tilde{I} \cap \tilde{J} = \emptyset$, and $I \cup J \neq \tilde{I} \cup \tilde{J}$. Then,

$$\mathbb{E}\left[\mathbb{E}_{\Omega_{\delta}}^{\mathrm{ref}}\left[\prod_{u\in I}\eta_{u}\prod_{v\in J}\lambda\sigma_{v}\omega_{v}\right]\mathbb{E}_{\Omega_{\delta}}^{\mathrm{ref}}\left[\prod_{u\in\tilde{I}}\eta_{u}\prod_{v\in\tilde{J}}\lambda\sigma_{v}\omega_{v}\right]\right]=0.$$

Proof. We can rewrite the expectation as

$$\mathbb{E}_{\Omega_{\delta}}^{\mathrm{ref},\otimes 2} \left[\mathbb{E} \left[\prod_{u \in I} \eta_{u}(\sigma,\omega) \prod_{v \in J} \lambda \sigma_{v} \omega_{v} \prod_{\tilde{u} \in \tilde{I}} \eta_{\tilde{u}}(\sigma',\omega) \prod_{\tilde{v} \in \tilde{J}} \lambda \sigma'_{\tilde{v}} \omega_{\tilde{v}} \right] \right],$$
(2.6)

where $E_{\Omega_{\delta}}^{\mathrm{ref},\otimes 2}$ denotes the expectation with respect to σ and σ' , two independent spin configurations with law $P_{\Omega_{\delta}}^{\mathrm{ref}}$. Recall that conditional on σ and σ' ,

$$\eta_x(\sigma,\omega) = e^{\lambda\omega_x\sigma_x - \phi(\lambda\sigma_x)} - 1 - \lambda\omega_x\sigma_x$$

and $\eta_x(\sigma',\omega)$ depend only on ω_x and hence are independent of $(\omega_y,\eta_y(\sigma,\omega),\eta_y(\sigma',\omega))$ for $y \neq x$. The assumption $I \cup J \neq \tilde{I} \cup \tilde{J}$ implies that there is some site $x \in I \cup J \cup \tilde{I} \cup \tilde{J}$ which appears exactly once in the product in (2.6). By the independence of $(\omega_y,\eta_y(\sigma,\omega),\eta_y(\sigma',\omega))$ for different $y \in \Omega_\delta$ and the fact that $\mathbb{E}[\eta_y(\sigma,\omega)] = \mathbb{E}[\eta_y(\sigma',\omega)] = \mathbb{E}[\omega_y] = 0$, it follows easily that the expectation in (2.6) equals 0.

Applying Lemma 2.1, to prove (2.5), it then suffices to show that

$$\mathbb{E}[\mathcal{E}_{k,\delta}^2] = \sum_{\substack{\{x_1,\dots,x_k\} \subset \Omega_\delta \\ x_i \neq x_j \text{ for } i \neq j}} \mathbb{E}\left[\left(\sum_{\substack{(I,J) \vdash \{x_1,\dots,x_k\} \\ I \neq \emptyset}} \operatorname{E}_{\Omega_\delta}^{\operatorname{ref}} \left[\prod_{u \in I} \eta_u \prod_{v \in J} \lambda \sigma_v \omega_v \right] \right)^2 \right] \xrightarrow{\delta \downarrow 0} 0.$$
(2.7)

Note that the sum over $(I, J) \vdash \{x_1, \ldots, x_k\}, I \neq \emptyset$, contains $2^k - 1$ terms. Using $(\sum_{i=1}^n a_i)^2 \le n \sum_{i=1}^n a_i^2$, we can bound

$$\mathbb{E}[\mathcal{E}_{k,\delta}^{2}] \leq 2^{k} \sum_{\substack{\{x_{1},\dots,x_{k}\}\subset\Omega_{\delta}\\x_{i}\neq x_{j} \text{ for } i\neq j}} \sum_{\substack{(I,J)\vdash\{x_{1},\dots,x_{k}\}\\I\neq\emptyset}} \mathbb{E}\left[\mathbb{E}_{\Omega_{\delta}}^{\text{ref}}\left[\prod_{u\in I}\eta_{u}\prod_{v\in J}\lambda\sigma_{v}\omega_{v}\right]^{2}\right] \\
= 2^{k} \sum_{\iota=1}^{k} \sum_{\substack{I,J\subset\Omega_{\delta},I\cap J=\emptyset\\|I|=\iota,|J|=k-\iota}} \mathbb{E}\left[\mathbb{E}_{\Omega_{\delta}}^{\text{ref}}\left[\prod_{u\in I}\eta_{u}\prod_{v\in J}\lambda\sigma_{v}\omega_{v}\right]^{2}\right] \\
= 2^{k} \sum_{\iota=1}^{k} \sum_{\substack{I,J\subset\Omega_{\delta},I\cap J=\emptyset\\|I|=\iota,|J|=k-\iota}} \mathbb{E}_{\Omega_{\delta}}^{\text{ref},\otimes 2}\left[\prod_{u\in I}\mathbb{E}[\eta_{u}(\sigma,\omega)\eta_{u}(\sigma',\omega)]\prod_{v\in J}\lambda^{2}\sigma_{v}\sigma'_{v}\right], \quad (2.8)$$

where σ' is an independent copy of σ with law $P_{\Omega_{\delta}}^{\text{ref}}$. Therefore to prove (2.7), it suffices to show that for each $1 \leq \iota \leq k$,

$$\sum_{\substack{I,J \subseteq \Omega_{\delta}, I \cap J = \emptyset\\|I| = \iota, |J| = k - \iota}} \mathbb{E}_{\Omega_{\delta}}^{\mathrm{ref}, \otimes 2} \left[\prod_{u \in I} \mathbb{E}[\eta_{u}(\sigma, \omega)\eta_{u}(\sigma', \omega)] \prod_{v \in J} \lambda^{2} \sigma_{v} \sigma_{v}' \right] \xrightarrow{\delta \downarrow 0} 0.$$
(2.9)

To see heuristically why (2.9) holds, recall from the discussion leading to (1.11) that $\lambda = \lambda_{\delta} = \hat{\lambda} \delta^{\frac{d}{2} - \gamma}$ is chosen such that if each spin σ_u is matched with a factor of λ and u is summed over Ω_{δ} , then the spin correlations $\mathrm{E}_{\Omega_{\delta}}^{\mathrm{ref}}[\sigma_{x_1^{\delta}} \dots \sigma_{x_k^{\delta}}]$ would be properly normalised and we will have convergence as in (1.11). However, if some of the spins among $\sigma_{x_i^{\delta}}$ coincide, then by assumption (A3), assigning one factor of λ to each spin gives more powers of λ than needed to normalize $\mathrm{E}_{\Omega_{\delta}}^{\mathrm{ref}}[\sigma_{x_1^{\delta}} \dots \sigma_{x_k^{\delta}}]$. This is exactly what happens when we expand $\mathbb{E}[\eta_u(\sigma, \omega)\eta_u(\sigma', \omega)]$ in powers of λ , σ_u and σ'_u . We will perform this expansion next, which is a bit involved.

Recall that
$$\eta_x(\sigma,\omega) = e^{\lambda\omega_x\sigma_x - \phi(\lambda\sigma_x)} - 1 - \lambda\omega_x\sigma_x$$
 and $\phi(a) = \log \mathbb{E}[e^{a\omega_x}]$. Therefore

$$\mathbb{E}[\eta_u(\sigma,\omega)\eta_u(\sigma',\omega)] = \mathbb{E}[(e^{\lambda\omega_u\sigma_u - \phi(\lambda\sigma_u)} - 1 - \lambda\omega_u\sigma_u)(e^{\lambda\omega_u\sigma'_u - \phi(\lambda\sigma'_u)} - 1 - \lambda\omega_u\sigma'_u)]$$

$$= e^{\phi(\lambda(\sigma_u + \sigma'_u)) - \phi(\lambda\sigma_u) - \phi(\lambda\sigma'_u)} - 1 + \lambda^2\sigma_u\sigma'_u$$

$$- \lambda\sigma_u\mathbb{E}[\omega_u e^{\lambda\omega_u\sigma'_u - \phi(\lambda\sigma'_u)}] - \lambda\sigma'_u\mathbb{E}[\omega_u e^{\lambda\omega_u\sigma_u - \phi(\lambda\sigma_u)}]. \quad (2.10)$$

Since $\phi(a)$ is assumed to be finite for all $|a| < a_0$ for some $a_0 > 0$ and $\phi(0) = 0$, ϕ must be analytic on the ball $|z| < z_0$ for some $z_0 > 0$ and has power series expansion

$$\phi(z) = \sum_{m=1}^{\infty} \frac{\kappa_m}{m!} z^m, \qquad (2.11)$$

where $\kappa_m = \frac{d^m \phi(z)}{dz^m} \Big|_{z=0}$ is the *m*-th cumulant of ω_x with $\kappa_1 = \mathbb{E}[\omega_x] = 0$ and $\kappa_2 = \mathbb{Var}(\omega_x^2) = 1$. Since $|\sigma_u| \leq K$ by assumption **(A0)**, for $\lambda = \lambda_\delta$ sufficiently small, we can rewrite the last term in (2.10) (similarly for the second last term) as

$$\lambda \sigma'_{u} \mathbb{E}[\omega_{u} e^{\lambda \omega_{u} \sigma_{u} - \phi(\lambda \sigma_{u})}] = \lambda \sigma'_{u} \frac{\mathrm{d}\phi(z)}{\mathrm{d}z} \Big|_{z = \lambda \sigma_{u}} = \lambda^{2} \sigma_{u} \sigma'_{u} + \sum_{m=3}^{\infty} \frac{\kappa_{m} \lambda^{m}}{(m-1)!} (\sigma_{u})^{m-1} \sigma'_{u}.$$
(2.12)

Substituting into (2.10) then yields

$$\mathbb{E}[\eta_u(\sigma,\omega)\eta_u(\sigma',\omega)] = e^{\phi(\lambda(\sigma_u+\sigma'_u))-\phi(\lambda\sigma_u)-\phi(\lambda\sigma'_u)} - 1 - \lambda^2 \sigma_u \sigma'_u -\sum_{m=3}^{\infty} \frac{\kappa_m \lambda^m}{(m-1)!} (\sigma'_u)^{m-1} \sigma_u - \sum_{m=3}^{\infty} \frac{\kappa_m \lambda^m}{(m-1)!} (\sigma_u)^{m-1} \sigma'_u.$$
(2.13)

Using (2.11) and the binomial expansion yields

$$\phi(\lambda(\sigma_u + \sigma'_u)) - \phi(\lambda\sigma_u) - \phi(\lambda\sigma'_u) = \sum_{m=2}^{\infty} \frac{\kappa_m}{m!} \lambda^m \sum_{l=1}^{m-1} \binom{m}{l} (\sigma_u)^{m-l} (\sigma'_u)^l.$$
(2.14)

Again by Taylor expansion, for $\lambda = \lambda_{\delta}$ sufficiently small, we have

$$e^{\phi(\lambda(\sigma_{u}+\sigma'_{u}))-\phi(\lambda\sigma_{u})-\phi(\lambda\sigma'_{u})} - 1$$

$$= \sum_{m=2}^{\infty} \frac{\kappa_{m}}{m!} \lambda^{m} \sum_{l=1}^{m-1} \binom{m}{l} (\sigma_{u})^{m-l} (\sigma'_{u})^{l} + \sum_{j=2}^{\infty} \frac{1}{j!} \left[\sum_{m=2}^{\infty} \frac{\kappa_{m}}{m!} \lambda^{m} \sum_{l=1}^{m-1} \binom{m}{l} (\sigma_{u})^{m-l} (\sigma'_{u})^{l} \right]^{j}.$$
(2.15)

Since $\kappa_2 = 1$, we can decompose the first term in (2.15) by separating the contributions from m = 2, m = 3 and $l \in \{1, m - 1\}$, versus $m \ge 4$ and $l \in \{2, \ldots, m - 2\}$ to rewrite it as

$$\lambda^{2}\sigma_{u}\sigma_{u}' + \sum_{m=3}^{\infty} \frac{\kappa_{m}\lambda^{m}}{(m-1)!} \Big((\sigma_{u})^{m-1}\sigma_{u}' + \sigma_{u}(\sigma_{u}')^{m-1} \Big) + \sum_{m=4}^{\infty} \frac{\kappa_{m}\lambda^{m}}{m!} \sum_{l=2}^{m-2} \binom{m}{l} (\sigma_{u})^{m-l} (\sigma_{u}')^{l}.$$

Substituting into (2.15) and then (2.13) gives

$$\mathbb{E}[\eta_u(\sigma,\omega)\eta_u(\sigma',\omega)] = \sum_{m=4}^{\infty} \frac{\kappa_m \lambda^m}{m!} \sum_{l=2}^{m-2} \binom{m}{l} (\sigma_u)^{m-l} (\sigma'_u)^l + \sum_{j=2}^{\infty} \frac{1}{j!} \left[\sum_{m=2}^{\infty} \frac{\kappa_m \lambda^m}{m!} \sum_{l=1}^{m-1} \binom{m}{l} (\sigma_u)^{m-l} (\sigma'_u)^l \right]^j.$$
(2.16)

Next we will expand the second term in the r.h.s. of (2.16) and perform resummation to write

$$\mathbb{E}[\eta_u(\sigma,\omega)\eta_u(\sigma',\omega)] = \sum_{m=4}^{\infty} \sum_{l=2}^{m-2} \lambda^m a_{m,l}(\sigma_u)^{m-l}(\sigma'_u)^l, \qquad (2.17)$$

where $a_{m,l}$ are constants to be determined later. To justify the resummation, we need to show absolute summability. For this, we first need to bound $\frac{|\kappa_m|}{m!}$. Since the power series for ϕ in (2.11) has a positive radius of convergence, there exists a finite constant C > 0 such that

$$\frac{|\kappa_m|}{m!} \le \mathcal{C}^m \quad \text{for any } m \ge 2.$$
(2.18)

Combined with the assumption (A0) that $|\sigma_u| \leq K$ for some finite K, we can bound

$$\sum_{m=2}^{\infty} \sum_{l=1}^{m-1} \left| \frac{\kappa_m \lambda^m}{m!} \binom{m}{l} (\sigma_u)^{m-l} (\sigma'_u)^l \right| \le \sum_{m=2}^{\infty} \mathcal{C}^m \lambda^m K^m \sum_{l=1}^{m-1} \binom{m}{l} \le \sum_{m=2}^{\infty} (2\lambda K\mathcal{C})^m,$$

which is convergent when $\lambda = \lambda_{\delta} = \hat{\lambda} \delta^{\frac{d}{2}-\gamma}$ is small enough. This implies that the r.h.s. of (2.16) is absolutely convergent, and we can rearrange terms into a power series in λ as in (2.17). To identify the constants $a_{m,l}$, for $j \geq 2$, we expand

$$\sum_{j=2}^{\infty} \frac{1}{j!} \left[\sum_{m=2}^{\infty} \frac{\kappa_m \lambda^m}{m!} \sum_{l=1}^{m-1} \binom{m}{l} (\sigma_u)^{m-l} (\sigma'_u)^l \right]^j$$

$$= \sum_{j=2}^{\infty} \frac{1}{j!} \sum_{m_1, \cdots, m_j \ge 2} \sum_{\substack{1 \le l_i \le m_i - 1, i = 1 \\ i = 1, \cdots, j}} \prod_{i=1}^j \left(\frac{\kappa_{m_i} \lambda^{m_i}}{m_i!} \binom{m_i}{l_i} (\sigma_u)^{(m_i - l_i)} (\sigma'_u)^{l_i} \right)$$

$$=: \sum_{j=2}^{\infty} \frac{1}{j!} \sum_{m=2j}^{\infty} \sum_{l=j}^{m-j} c_{j,m,l} \lambda^m (\sigma_u)^{m-l} (\sigma'_u)^l$$

$$=: \sum_{m=4}^{\infty} \lambda^m \sum_{l=2}^{m-2} d_{m,l} (\sigma_u)^{m-l} (\sigma'_u)^l, \qquad (2.19)$$

where for $j \ge 2$, $m \ge 2j$, and $j \le l \le m - j$,

$$c_{j,m,l} := \sum_{\substack{m_1, \cdots, m_j \ge 2\\m_1 + \cdots + m_j = m}} \sum_{\substack{1 \le l_i \le m_i - 1, i = 1\\i = 1, \cdots, j\\l_1 + \cdots + l_j = l}} \prod_{i=1}^{j} \frac{\kappa_{m_i}}{m_i!} \binom{m_i}{l_i}, \qquad (2.20)$$

and

$$d_{m,l} = \begin{cases} \sum_{j=2}^{l} \frac{1}{j!} c_{j,m,l}, & \text{if } 2 \le l \le \lfloor \frac{m}{2} \rfloor, \\ d_{m,m-l}, & \text{if } \lfloor \frac{m}{2} \rfloor < l \le m-2. \end{cases}$$

Combined with (2.16), it follows that (2.17) holds with

$$a_{m,l} := \frac{\kappa_m}{m!} \binom{m}{l} + d_{m,l} = a_{m,m-l}.$$
(2.21)

This concludes the expansion of $\mathbb{E}[\eta_u(\sigma, \omega)\eta_u(\sigma', \omega)]$ stated in (2.17).

Substituting (2.17) into (2.9), our goal is to show

$$\sum_{\substack{I,J \subset \Omega_{\delta}, I \cap J = \emptyset\\|I| = \iota, |J| = k - \iota}} \mathcal{E}_{\Omega_{\delta}}^{\mathrm{ref}, \otimes 2} \left[\prod_{u \in I} \left(\sum_{m=4}^{\infty} \sum_{l=2}^{m-2} \lambda^m a_{m,l} (\sigma_u)^{m-l} (\sigma'_u)^l \right) \prod_{v \in J} \lambda^2 \sigma_v \sigma'_v \right] \xrightarrow{\delta \downarrow 0} 0.$$
(2.22)

To control the convergence of the series, first recall from (2.18) that $\frac{|\kappa_m|}{m!} \leq C^m$. Therefore by (2.21), for $2 \leq l \leq \lfloor \frac{m}{2} \rfloor$, we have

$$|a_{m,l}| = |a_{m,m-l}| \le \mathcal{C}^m \binom{m}{l} + \sum_{j=2}^l \frac{1}{j!} |c_{j,m,l}|.$$
(2.23)

Recalling the definition of $c_{j,m,l}$ from (2.20), we have

$$|c_{j,m,l}| \le \mathcal{C}^m \sum_{\substack{m_1, \cdots, m_j \ge 2\\m_1 + \cdots + m_j = m}} \sum_{\substack{1 \le l_i \le m_i - 1, i = 1\\i = 1, \cdots, j}} \prod_{i=1}^j \binom{m_i}{l_i} \le \mathcal{C}^m \sum_{\substack{m_1, \cdots, m_j \ge 2\\m_1 + \cdots + m_j = m}} \prod_{i=1}^j 2^{m_i} = (2\mathcal{C})^m \binom{m-j-1}{j-1},$$

where the last identity holds by elementary combinatorial considerations. Substituting this into (2.23) then yields that, for $2 \le l \le \lfloor \frac{i}{2} \rfloor$,

$$|a_{m,l}| = |a_{m,m-l}| \le \mathcal{C}^m \binom{m}{l} + (2\mathcal{C})^m \sum_{j=2}^l \frac{1}{j!} \binom{m-j-1}{j-1} \le (2\mathcal{C})^m + (2\mathcal{C})^m \sum_{j=2}^l \frac{1}{j!} 2^{m-j-1} \le 2 \cdot (4\mathcal{C})^m + (2\mathcal{C})^m \sum_{j=2}^l \frac{1}{j!} 2^{m-j-1} \le 2 \cdot (4\mathcal{C})^m + (2\mathcal{C})^m + (2\mathcal{C}$$

Therefore

$$\sum_{l=2}^{m-2} |a_{m,l}| \le 2m \cdot (4\mathcal{C})^m \le (8\mathcal{C})^m.$$
(2.24)

Applying this bound and the assumption (A0) that $|\sigma_u| \leq K$ for any $u \in \Omega_{\delta}$, the expectation in (2.22) can be bounded from above by

$$\mathbb{E}_{\Omega_{\delta}}^{\mathrm{ref},\otimes 2} \left[\prod_{u \in I} \left(\sum_{m=4}^{\infty} \sum_{l=2}^{m-2} \left| \lambda^{m} a_{m,l}(\sigma_{u})^{m-l}(\sigma_{u}')^{l} \right| \right) \prod_{v \in J} \left| \lambda^{2} \sigma_{v} \sigma_{v}' \right| \right] \leq \left(\sum_{m=4}^{\infty} (8\mathcal{C}\lambda K)^{m} \right)^{|I|} (\lambda K)^{2|J|},$$

$$(2.25)$$

which is finite for $\lambda = \lambda_{\delta} := \hat{\lambda} \delta^{\frac{d}{2} - \gamma}$ sufficiently small. It follows that the expectation in (2.22) is absolutely convergent and can be expanded as

$$\mathbf{E}_{\Omega_{\delta}}^{\mathrm{ref},\otimes 2} \left[\prod_{u \in I} \left(\sum_{m=4}^{\infty} \sum_{l=2}^{m-2} \lambda^{m} a_{m,l}(\sigma_{u})^{m-l}(\sigma_{u}')^{l} \right) \prod_{v \in J} \lambda^{2} \sigma_{v} \sigma_{v}' \right] \\
= \mathbf{E}_{\Omega_{\delta}}^{\mathrm{ref},\otimes 2} \left[\sum_{m_{1},\cdots,m_{|I|} \geq 4} \sum_{\substack{2 \leq l_{i} \leq m_{i}-2, \\ i=1,\cdots,|I|}} \prod_{i=1}^{|I|} \left(\lambda^{m_{i}} a_{m_{i},l_{i}}(\sigma_{u_{i}})^{m_{i}-l_{i}}(\sigma_{u_{i}}')^{l_{i}} \right) \prod_{v \in J} \lambda^{2} \sigma_{v} \sigma_{v}' \right] \\
= \sum_{m_{1},\cdots,m_{|I|} \geq 4} \sum_{\substack{2 \leq l_{i} \leq m_{i}-2, \\ i=1,\cdots,|I|}} \mathbf{E}_{\Omega_{\delta}}^{\mathrm{ref},\otimes 2} \left[\prod_{i=1}^{|I|} \left(\lambda^{m_{i}} a_{m_{i},l_{i}}(\sigma_{u_{i}})^{m_{i}-l_{i}}(\sigma_{u_{i}}')^{l_{i}} \right) \prod_{v \in J} \lambda^{2} \sigma_{v} \sigma_{v}' \right], \quad (2.26)$$

where we have assumed that $I = \{u_1, \dots, u_{|I|}\}$. To show (2.22), it suffices to show

$$\sum_{\substack{I,J\subset\Omega_{\delta},I\cap J=\emptyset\\|I|=\iota,|J|=k-\iota\\ 2\leq l_i\leq m_i-2,\\i=1,\cdots,|I|}} \sum_{\substack{m_1,\cdots,m_{|I|}\geq 4\\ 2\leq l_i\leq m_i-2,\\i=1,\cdots,|I|}} \left| \mathrm{E}_{\Omega_{\delta}}^{\mathrm{ref},\otimes 2} \left[\prod_{i=1}^{|I|} \left(\lambda^{m_i} a_{m_i,l_i}(\sigma_{u_i})^{m_i-l_i}(\sigma_{u_i}')^{l_i} \right) \prod_{v\in J} \lambda^2 \sigma_v \sigma_v' \right] \right| \xrightarrow{\delta\downarrow 0} 0. \quad (2.27)$$

The absolute value in the left hand side of (2.27) can be bounded by

$$\lambda^{2|J|} \left(\prod_{i=1}^{|I|} \lambda^{m_i} |a_{m_i, l_i}| \right) \left| \mathbf{E}_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{i=1}^{|I|} (\sigma_{u_i})^{m_i - l_i} \prod_{v \in J} \sigma_v \right] \right| \left| \mathbf{E}_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{i=1}^{|I|} (\sigma'_{u_i})^{l_i} \prod_{v \in J} \sigma'_v \right] \right|.$$
(2.28)

Using assumption (A3), where $(r)_2 := r \pmod{2}$, there is a universal constant $C \ge 1$ such that

$$\left| \mathbf{E}_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{i=1}^{|I|} (\sigma_{u_i})^{m_i - l_i} \prod_{v \in J} \sigma_v \right] \right| \le C_{i=1}^{|I|} [m_i - l_i - (m_i - l_i)_2] \left| \mathbf{E}_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{i=1}^{|I|} \sigma_{u_i}^{(m_i - l_i)_2} \prod_{v \in J} \sigma_v \right] \right|$$

and

$$\left| \mathbf{E}_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{i=1}^{|I|} (\sigma_{u_i}')^{l_i} \prod_{v \in J} \sigma_v' \right] \right| \leq C_{i=1}^{|I|} \left| \mathbf{E}_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{i=1}^{|I|} (\sigma_{u_i}')^{(l_i)_2} \prod_{v \in J} \sigma_v' \right] \right|.$$

Let

$$I_{1} = \{ u_{i} \in I : (m_{i} - l_{i})_{2} = 1, 1 \leq i \leq |I| \},$$

$$I'_{1} = \{ u_{i} \in I : (l_{i})_{2} = 1, 1 \leq i \leq |I| \}.$$
(2.29)

Then we can write

$$\left| E_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{i=1}^{|I|} (\sigma_{u_{i}})^{m_{i}-l_{i}} \prod_{v \in J} \sigma_{v} \right] \right| \left| E_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{i=1}^{|I|} (\sigma'_{u_{i}})^{l_{i}} \prod_{v \in J} \sigma'_{v} \right] \right| \\
\leq c_{|I|} C^{\sum_{i=1}^{|I|} m_{i}} \left| E_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{u \in I_{1}} \sigma_{u} \prod_{v \in J} \sigma_{v} \right] \right| \left| E_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{u \in I_{1}'} \sigma'_{u} \prod_{v \in J} \sigma'_{v} \right] \right| \\
\leq c_{|I|} C^{\sum_{i=1}^{|I|} m_{i}} \left(E_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{u \in I_{1}} \sigma_{u} \prod_{v \in J} \sigma_{v} \right]^{2} + E_{\Omega_{\delta}}^{\mathrm{ref}} \left[\prod_{u \in I_{1}'} \sigma'_{u} \prod_{v \in J} \sigma'_{v} \right]^{2} \right),$$
(2.30)

where $c_{|I|} = C^{-\sum_{i=1}^{|I|} [(m_i - l_i)_2 + (l_i)_2]} \leq 1$. Inserting this bound into (2.28), the absolute value in the left hand side of (2.27) is bounded from above by

$$c_{|I|}\lambda^{2|J|} \left(\prod_{i=1}^{|I|} (\lambda C)^{m_i} |a_{m_i,l_i}|\right) \left(\operatorname{E}_{\Omega_{\delta}}^{\operatorname{ref}} \left[\prod_{u \in I_1} \sigma_u \prod_{v \in J} \sigma_v\right]^2 + \operatorname{E}_{\Omega_{\delta}}^{\operatorname{ref}} \left[\prod_{u \in I_1'} \sigma_u' \prod_{v \in J} \sigma_v'\right]^2\right).$$

Substitute this bound into (2.27). By the symmetry between $(\sigma, m_i - l_i, I_1)$ and (σ', l_i, I'_1) , the two terms in the sum above give equal contribution, and hence it suffices to show that

$$\sum_{\substack{I,J\subset\Omega_{\delta},I\cap J=\emptyset\\|I|=\iota,|J|=k-\iota\\I_1\subset I}}\sum_{\substack{2\leq l_i\leq m_i-2,\\i=1,\cdots,|I|}}\lambda^{2|J|} \left(\prod_{i=1}^{|I|}(\lambda C)^{m_i}|a_{m_i,l_i}|\right) \operatorname{E}_{\Omega_{\delta}}^{\operatorname{ref}}\left[\prod_{u\in I_1}\sigma_u\prod_{v\in J}\sigma_v\right]^2 \xrightarrow{\delta\downarrow 0} 0.$$
(2.31)

We can further decompose the sum above according to $\iota_1 := |I_1|$. Denote $I_2 := I \setminus I_1$, $\iota_2 := |I_2|$ and $\iota_3 = |J|$. Then it suffices to show that for any $\iota_1, \iota_2, \iota_3 \ge 0$ with $\iota := \iota_1 + \iota_2 \ge 1$ and

$$\sum_{\substack{I_1,I_2,J\subset\Omega_{\delta}\\I_1\cap I_2=\emptyset,I\cap J=\emptyset\\|I_1|=\iota_1,|I_2|=\iota_2,|J|=\iota_3}} \lambda^{2|J|} \left(\sum_{\substack{m_1,\cdots,m_{|I|}\geq 4\\2\leq l_i\leq m_i-2,\\i=1,\cdots,|I|}} \left(\prod_{i=1}^{|I|} (\lambda C)^{m_i} |a_{m_i,l_i}| \right) \right) \operatorname{E}_{\Omega_{\delta}}^{\operatorname{ref}} \left[\prod_{u\in I_1} \sigma_u \prod_{v\in J} \sigma_v \right]^2 \xrightarrow{\delta\downarrow 0} 0. \quad (2.32)$$

Using (2.24), the above sum over m_i and l_i can be bounded by

$$\sum_{\substack{m_1, \cdots, m_{|I|} \ge 4\\ 2 \le l_i \le m_i - 2,\\ i = 1, \cdots, |I|}} \left(\prod_{i=1}^{|I|} (\lambda C)^{m_i} |a_{m_i, l_i}| \right) = \prod_{i=1}^{|I|} \left(\sum_{m_i=4}^{\infty} (\lambda C)^{m_i} \sum_{l_i=2}^{m_i-2} |a_{m_i, l_i}| \right)$$

$$\leq \prod_{i=1}^{|I|} \left(\sum_{m_i=4}^{\infty} (8\lambda C\mathcal{C})^{m_i} \right) \le (\hat{C}\lambda)^{4|I|},$$
(2.33)

where $\hat{C} = 16CC$ and the bound holds for $\lambda = \lambda_{\delta} = \hat{\lambda}\delta^{\frac{d}{2}-\gamma} \leq \hat{C}^{-1}$. The l.h.s. of (2.32) can then be bounded by

$$\hat{C}^{4|I|}(\lambda)^{4|I|+2|J|} \sum_{\substack{I_1, I_2, J \subseteq \Omega_{\delta} \\ I_1 \cap I_2 = \emptyset, I \cap J = \emptyset \\ |I_1| = \iota_1, |I_2| = \iota_2, |J| = \iota_3}} \operatorname{E}_{\alpha_{\delta}}^{\operatorname{ref}} \left[\prod_{u \in I_1} \sigma_u \prod_{v \in J} \sigma_v \right]^2 \\
\leq \hat{C}^{4|I|}(\hat{\lambda}\delta^{\frac{d}{2}-\gamma})^{4|I|+2|J|}(|\Omega|\delta^{-d})^{|I_2|} \binom{k - |I_2|}{|I_1|} \delta^{(2\gamma-d)(k-\iota_2)} \sum_{G \subseteq \Omega_{\delta}, |G| = k-\iota_2} \delta^{d|G|} \left(\delta^{-\gamma|G|} \operatorname{E}_{\Omega_{\delta}}^{\operatorname{ref}} \left[\prod_{u \in G} \sigma_u \right] \right)^2 \\
\leq \tilde{C}^{\iota_1+\iota_2+\iota_3} \delta^{d(\iota_1+\iota_2)-2\gamma(\iota_1+2\iota_2)} \sum_{G \subseteq \Omega_{\delta}, |G| = \iota_1+\iota_3} \delta^{d|G|} \left(\delta^{-\gamma|G|} \operatorname{E}_{\Omega_{\delta}}^{\operatorname{ref}} \left[\prod_{u \in G} \sigma_u \right] \right)^2 \\
= \tilde{C}^{\iota_1+\iota_2+\iota_3} \delta^{d(\iota_1+\iota_2)-2\gamma(\iota_1+2\iota_2)} \frac{\|\psi_{\delta}\|_{L^2(\Omega^{\iota_1+\iota_3})}^2}{(\iota_1+\iota_3)!}, \qquad (2.34)$$

where \tilde{C} does not depend on δ , ι_1 , ι_2 or ι_3 , and $\|\psi_{\delta}\|^2_{L^2(\Omega^{\iota_1+\iota_3})} \to \|\psi_0\|^2_{L^2(\Omega^{\iota_1+\iota_3})}$ as $\delta \downarrow 0$ by assumption **(A1)**. Therefore (2.32) holds if

$$d(\iota_1+\iota_2)-2\gamma(\iota_1+2\iota_2)>0 \quad \Longleftrightarrow \quad \gamma<\frac{d}{2}\cdot\frac{\iota_1+\iota_2}{\iota_1+2\iota_2},$$

which holds for all $\iota_1, \iota_2 \ge 0$ with $\iota_1 + \iota_2 \ge 1$ if and only if $\gamma < d/4$. This is part of the assumption in Theorem 1.3 and hence the proof of (2.7) and (2.5) is complete.

2.2 Control of the remainder

In this section, we conclude the proof of Theorem 1.3 by proving (2.3), that is, the remainder

$$R_{M,\delta} = \sum_{k=M+1}^{\infty} \sum_{\substack{\{x_1,\dots,x_k\} \subset \Omega_\delta \\ x_i \neq x_j \text{ for } i \neq j}} \operatorname{E}_{\Omega_\delta}^{\operatorname{ref}} \left[\prod_{i=1}^k (\lambda \omega_{x_i} \sigma_{x_i} + \eta_{x_i}) \right]$$
$$= \sum_{k=M+1}^{\infty} \sum_{\substack{\{x_1,\dots,x_k\} \subset \Omega_\delta \\ x_i \neq x_j \text{ for } i \neq j}} \sum_{\substack{I \subset \{x_1,\dots,x_k\} \setminus I \\ J = \{x_1,\dots,x_k\} \setminus I}} \operatorname{E}_{\Omega_\delta}^{\operatorname{ref}} \left[\prod_{u \in I} \eta_u \prod_{v \in J} \lambda \sigma_v \omega_v \right]$$

satisfies $\lim_{M\to\infty} \limsup_{\delta\downarrow 0} \mathbb{E}[R_{M,\delta}^2] = 0$. By the same calculations in (2.7) and (2.8), we can apply Lemma 2.1 and the inequality $(\sum_{i=1}^n a_i)^2 \leq n \sum_{i=1}^n a_i^2$ to obtain

$$\mathbb{E}[R_{M,\delta}^2] \leq \sum_{k=M+1}^{\infty} \sum_{\substack{\{x_1,\dots,x_k\} \subset \Omega_{\delta} \\ x_i \neq x_j \text{ for } i \neq j}} 2^k \sum_{\substack{I \subset \{x_1,\dots,x_k\} \\ J = \{x_1,\dots,x_k\} \setminus I}} \mathbb{E}_{\Omega_{\delta}}^{\operatorname{ref},\otimes 2} \left[\prod_{u \in I} \mathbb{E}[\eta_u(\sigma,\omega)\eta_u(\sigma',\omega)] \prod_{v \in J} \lambda^2 \sigma_v \sigma'_v \right], \quad (2.35)$$

where σ' is the independent copy of σ . We decompose the r.h.s of (2.35) into two parts corresponding respectively to $I = \emptyset$, which gives the dominant contribution, and $I \neq \emptyset$:

$$S_{M,\delta}^{(0)} := \sum_{k=M+1}^{\infty} 2^k \lambda_{\substack{\{x_1,\dots,x_k\} \subset \Omega_\delta \\ x_i \neq x_j \text{ for } i \neq j}}^{2k} \operatorname{E}_{\Omega_\delta}^{\operatorname{ref}}[\sigma_{x_1}\dots\sigma_{x_k}]^2,$$

$$S_{M,\delta}^{(1)} := \sum_{k=M+1}^{\infty} 2^k \sum_{\substack{\{x_1,\dots,x_k\} \subset \Omega_\delta \\ x_i \neq x_j \text{ for } i \neq j}} \sum_{\substack{I \subset \{x_1,\dots,x_k\} \\ |I| \ge 1}} \operatorname{E}_{\Omega_\delta}^{\operatorname{ref},\otimes 2} \left[\prod_{u \in I} \mathbb{E}[\eta_u(\sigma,\omega)\eta_u(\sigma',\omega)] \prod_{v \in J} \lambda^2 \sigma_v \sigma'_v \right]. \quad (2.36)$$

It follows directly from assumption (A2) that $\lim_{M\to\infty} \limsup_{\delta\downarrow 0} S_{M,\delta}^{(0)} = 0.$

To prove (2.3), it only remains to show that

$$\lim_{M \to \infty} \limsup_{\delta \downarrow 0} S_{M,\delta}^{(1)} = 0.$$
(2.37)

In fact, we will show that $\limsup_{\delta \downarrow 0} S_{M,\delta}^{(1)} = 0$. Our strategy is similar to the proof of (2.9), except now we need to control the sum over $k \ge M + 1$.

Recall from (2.17) the expansion

$$\mathbb{E}[\eta_u(\sigma,\omega)\eta_u(\sigma',\omega)] = \sum_{m=4}^{\infty} \sum_{l=2}^{m-2} \lambda^m a_{m,l}(\sigma_u)^{m-l} (\sigma'_u)^l.$$

Substituting this into (2.36) gives

$$S_{M,\delta}^{(1)} = \sum_{k=M+1}^{\infty} 2^k \sum_{\substack{\{x_1,\dots,x_k\} \subset \Omega_\delta \\ x_i \neq x_j \text{ for } i \neq j}} \sum_{\substack{I \subset \{x_1,\dots,x_k\} \\ |I| \ge 1}} \mathbb{E}_{\Omega_\delta}^{\text{ref},\otimes 2} \left[\prod_{u \in I} \left(\sum_{m=4}^{\infty} \sum_{l=2}^{m-2} \lambda^m a_{m,l}(\sigma_u)^{m-l}(\sigma_u')^l \right) \prod_{v \in J} \lambda^2 \sigma_v \sigma_v' \right].$$

Let us first consider the contributions from a fixed $k \ge M + 1$. Denote $I = \{u_1, \dots, u_{|I|}\} \ne \emptyset$ and $J = \{x_1, \ldots, x_k\} \setminus I$. By the same calculations as those leading to (2.34), we have the bound

$$\sum_{\substack{\{x_{1},\dots,x_{k}\}\subset\Omega_{\delta}\\x_{i}\neq x_{j} \text{ for } i\neq j}} \sum_{I\subset\{x_{1},\dots,x_{k}\}} \left| E_{\Omega_{\delta}}^{\operatorname{ref},\otimes 2} \left[\prod_{u\in I} \left(\sum_{m=4}^{\infty} \sum_{l=2}^{m-2} \lambda^{m} a_{m,l}(\sigma_{u})^{m-l}(\sigma_{u}')^{l} \right) \prod_{v\in J} \lambda^{2} \sigma_{v} \sigma_{v}' \right] \right| \\
\leq \sum_{\substack{\iota_{1},\iota_{2},\iota_{3}\geq 0\\\iota_{1}+\iota_{2}\geq 1\\\iota_{1}+\iota_{2}+\iota_{3}=k}} \sum_{\substack{I_{1},I_{2},J\subset\Omega_{\delta}\\I_{1}\cap I_{2}=\emptyset,I\cap J=\emptyset\\I_{1}\cap I_{2}=\emptyset,I\cap J=\emptyset}} \lambda^{2|J|} \left(\prod_{i=1}^{|I|} (\lambda C)^{m_{i}} |a_{m_{i},l_{i}}| \right) E_{\Omega_{\delta}}^{\operatorname{ref}} \left[\prod_{u\in I_{1}} \sigma_{u} \prod_{v\in J} \sigma_{v} \right]^{2} \\
\leq \sum_{\substack{\iota_{1},\iota_{2},\iota_{3}\geq 0\\\iota_{1}+\iota_{2}\geq 1\\\iota_{1}+\iota_{2}+\iota_{3}=k}} \sum_{\substack{I_{1},I_{2},J\subset\Omega_{\delta}\\I_{1}\cap I_{2}=\emptyset,I\cap J=\emptyset\\\iota_{1}+\iota_{2}=\iota_{3}=k}} \lambda^{2|J|} (\hat{C}\lambda)^{4|I|} E_{\Omega_{\delta}}^{\operatorname{ref}} \left[\prod_{u\in I_{1}} \sigma_{u} \prod_{v\in J} \sigma_{v} \right]^{2} \\
\leq \sum_{\substack{\iota_{1},\iota_{2},\iota_{3}\geq 0\\\iota_{1}+\iota_{2}\geq 1\\\iota_{1}+\iota_{2}+\iota_{3}=k}} \tilde{C}^{\iota_{1}+\iota_{2}+\iota_{3}}} \delta^{d(\iota_{1}+\iota_{2})-2\gamma(\iota_{1}+2\iota_{2})} \frac{\|\psi_{\delta}\|_{L^{2}(\Omega^{\iota_{1}+\iota_{3}})}^{2}}{(\iota_{1}+\iota_{3})!}.$$
(2.38)

Summing this bound over $k \ge M + 1$ then gives

$$\begin{split} |S_{M,\delta}^{(1)}| &\leq \sum_{k=M+1}^{\infty} 2^k \sum_{\substack{\iota_1, \iota_2, \iota_3 \geq 0\\ \iota_1 + \iota_2 \geq 1\\ \iota_1 + \iota_2 + \iota_3 = k}} \tilde{C}^{\iota_1 + \iota_2 + \iota_3} \delta^{d(\iota_1 + \iota_2) - 2\gamma(\iota_1 + 2\iota_2)} \frac{\|\psi_\delta\|_{L^2(\Omega^{\iota_1 + \iota_3})}^2}{(\iota_1 + \iota_3)!} \\ &\leq \sum_{\iota_1, \iota_3 \geq 0} \frac{\tilde{C}^{\iota_1 + \iota_3}}{(\iota_1 + \iota_3)!} \|\psi_\delta\|_{L^2(\Omega^{\iota_1 + \iota_3})}^2 \sum_{\substack{\iota_2 \geq M + 1 - \iota_1 - \iota_3\\ \iota_1 + \iota_2 \geq 1}} (\tilde{C} \delta^{d - 4\gamma})^{\iota_2} \delta^{(d - 2\gamma)\iota_1} \\ &\leq 2\tilde{C} \delta^{d - 4\gamma} \sum_{m=0}^{\infty} \frac{(2\tilde{C})^m}{m!} \|\psi_\delta\|_{L^2(\Omega^m)}^2, \end{split}$$

where the sum is uniformly bounded as $\delta \downarrow 0$ by assumptions (A1) and (A2). Since we assume $\gamma < d/4$, (2.37) follows immediately. This concludes the proof of (2.3).

Acknowledgements. L. Li acknowledges the support of the China Scholarship Council program (Project ID: 202206370088). V. Margarint would like to thank the National University of Singapore for the hospitality. R. Sun is supported by NUS grant A-8001448-00-00. We thank Federico Camia for helpful discussions on the assumption (A3).

References

- T. Alberts, K. Khanin, and J. Quastel. The continuum directed random polymer. J. Stat. Phys., 154(1-2):305–326, 2014.
- [2] T. Alberts, K. Khanin, and J. Quastel. The intermediate disorder regime for directed polymers in dimension 1 + 1. Ann. Probab., 42(3):1212–1256, 2014.
- [3] A. Bowditch and R. Sun. The two-dimensional continuum random field Ising model. Ann. Probab., 50(2):419–454, 2022.
- [4] P. Butera and M. Pernici. The Blume-Capel model for spins S = 1 and 3/2 in dimensions d = 2 and 3. *Phys. A*, 507:22–66, 2018.

- [5] F. Camia and Y. Feng. Conformally covariant probabilities, operator product expansions, and logarithmic correlations in two-dimensional critical percolation, arXiv:2407.04246, 2024.
- [6] F. Caravenna, R. Sun, and N. Zygouras. The continuum disordered pinning model. Probab. Theory Related Fields, 164(1-2):17–59, 2016.
- [7] F. Caravenna, R. Sun, and N. Zygouras. Polynomial chaos and scaling limits of disordered systems. J. Eur. Math. Soc. (JEMS), 19(1):1–65, 2017.
- [8] F. Caravenna, R. Sun, and N. Zygouras. The critical 2d stochastic heat flow. Invent. Math., 233(1):325–460, 2023.
- [9] G. Giacomin. Disorder and critical phenomena through basic probability models, volume 2025 of Lecture Notes in Mathematics. Springer, Heidelberg, 2011. Lecture notes from the 40th Probability Summer School held in Saint-Flour, 2010, École d'Été de Probabilités de Saint-Flour. [Saint-Flour Probability Summer School].
- [10] T. S. Gunaratnam, D. Krachun, and C. Panagiotis. Existence of a tricritical point for the Blume-Capel model on Z^d. Probab. Math. Phys., 5(3):785–845, 2024.
- [11] M. Hairer. A theory of regularity structures. Invent. Math., 198(2):269–504, 2014.
- [12] Y. Hu. Chaos expansion of heat equations with white noise potentials. Potential Anal., 16(1):45–66, 2002.
- [13] Y.-Z. Hu and J.-A. Yan. Wick calculus for nonlinear Gaussian functionals. Acta Math. Appl. Sin. Engl. Ser., 25(3):399–414, 2009.
- [14] D. Nualart and M. Zakai. Generalized Brownian functionals and the solution to a stochastic partial differential equation. J. Funct. Anal., 84(2):279–296, 1989.
- [15] J. Quastel, A. Ramirez, and B. Virag. KPZ fluctuations in the planar stochastic heat equation, arXiv:2210.13607, 2022.