

Thermodynamics of dyonic black holes in minimal supergravity

Rong-Gen Cai^{1,2,3,*}, Li Li^{1,3,4,†} and Jun-Kun Zhao^{1‡}

¹*CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China*

²*School of Physical Science and Technology, Ningbo University, Ningbo, 315211, China*

³*School of Fundamental Physics and Mathematical Sciences,*

Hangzhou Institute for Advanced Study, UCAS, Hangzhou 310024, China and

⁴*School of Physical Sciences, University of Chinese Academy of Sciences, No.19A Yuquan Road, Beijing 100049, China*

Thermodynamics of black holes offers a promising avenue for exploring the quantum nature of black holes and quantum gravity. In this Letter, we investigate the thermodynamic properties of dyonic black holes in the five-dimensional Einstein-Maxwell-Chern-Simons theory, obtained from IIB supergravity. We demonstrate that the standard form of the first law of thermodynamics is inconsistent with the quantum statistical relation widely adopted in black hole physics. By employing the on-shell variation of the Euclidean action and the Iyer-Wald formalism, we resolve this discrepancy and derive both the standard form of the first law and Smarr formula for the dyonic black holes. Furthermore, our findings are corroborated by numerical tests and are consistent with general hydrodynamic expectations.

Introduction.—Black hole might be one of the simplest yet most complex object known in physics. Often regarded as the “atom” of the 21st century physics, it offers a promising window into the nature of gravity. While many classical properties of black holes have been uncovered over the last century, understanding their quantum features remains a significant challenge. The discovery of Hawking radiation [1] revealed that black holes are thermodynamic systems with finite temperature. Specifically, the entropy of a black hole is equal to a quarter of its event horizon area in Planck units. This famous area law suggests the holographic nature of quantum gravity, *i.e.*, the description of a volume of space can be thought as encoded on a lower dimensional boundary to that region. Nevertheless, understanding the microstates of a quantum black hole and resolving the black hole information paradox remain big problems.

Black hole thermodynamics lies at the intersection of gravitation, statistics physics and quantum mechanics. The fundamental thermodynamic quantity, the free energy W of a black hole, can be calculated using the path integral approach and is expressed as the product of the temperature T and its Euclidean on-shell action I_E in a semi-classical approximation [2]:

$$W = T I_E, \quad (1)$$

This relationship is known as the quantum statistical relation [3] which serves as a cornerstone of the black hole thermodynamics and has become a standard framework for discussing the thermodynamics of black holes in textbooks. For example, it has significant applications in studying black hole thermodynamics in Anti-de Sitter (AdS) spacetime, such as the Hawking-Page transition [4], liquid–gas phase transition [5, 6]. Furthermore,

the thermodynamic properties of black holes play a crucial role in applying holography to strongly coupled quantum many-body systems [7–10].

Even though our understanding of black hole thermodynamics is primarily theoretical, the fact that the identification in equation (1) yields thermodynamics potential consistent with the laws of black hole thermodynamics is rather remarkable and by no means obvious, particularly for gravitational systems involving matter fields. In this Letter, we demonstrate that, despite the extensive research, our understanding of black hole thermodynamics remains incomplete. We will focus on a black hole within the framework of Einstein-Maxwell-Chern-Simons theory. This is a “top-down” model, meaning that the action can be derived from a consistent truncation of supergravity or M-theory. Black hole thermodynamics in this theory has attracted substantial interests, including black hole in Gödel universe [11, 12], rotating black holes [13, 14], helical black holes [15]. Of particular interest is the dyonic black holes in AdS spacetime [16–18], which allows intriguing phenomenology, such as magnetic-field-driven quantum phase transition and the chiral transports [19–21]. Although the dyonic solutions were reported decades ago, their thermodynamics properties have not been thoroughly investigated.

In the presence of an external magnetic field, the standard laws of thermodynamics, as presented in textbooks, are expressed as [22]

$$\begin{aligned} \delta w &= -s\delta T - \rho\delta\mu - M_B\delta B, \\ \delta\epsilon &= T\delta s + \mu\delta\rho - M_B\delta B, \end{aligned} \quad (2)$$

where $w = W/V$ is the free energy density in a fixed spatial volume V , and the energy density ϵ is given by the thermodynamic relation $\epsilon = w + Ts + \mu\rho$ [23]. Here T, μ and B denote temperature, chemical potential and magnetic field, respectively, while their conjugate quantities are the entropy density s , charge density ρ , and

* cairg@itp.ac.cn

† liliphy@itp.ac.cn

‡ junkunzhao@itp.ac.cn

magnetization M_B . In this study, we shall investigate the thermodynamics of the dyonic black holes within the top-down Einstein-Maxwell theory that includes a Chern-Simons term. We will demonstrate that the conventional treatment of black hole thermodynamics, particularly the quantum statistical relation (1), does not yield the standard laws of thermodynamics as articulated in (2). The aim of this Letter is then to address this inconsistent and derive both the generalized Smarr formula and the first law using the method of on-shell variation for Euclidean action as well as the Iyer-Wald formalism. Our analytical results are further supported by numerics.

Model and violation of thermodynamics first law.—The action of the five-dimensional Einstein-Maxwell-Chern-Simons theory reads

$$S = \frac{1}{16\pi G} \int d^5x \sqrt{-g} \left(R + \frac{12}{L^2} - \frac{1}{4} F_{ab} F^{ab} + \frac{k}{24} \epsilon^{abcde} A_a F_{bc} F_{de} \right), \quad (3)$$

where $F = dA$ is the field strength of the $U(1)$ gauge field A_a , G is the Newton's constant, L is the AdS radius, and k is the Chern-Simons coupling. The action is invariant under the transformation $(k, A_a) \leftrightarrow (-k, -A_a)$, allowing us to restrict our focus to $k \geq 0$. When $k = k_{\text{susy}} = \frac{2}{\sqrt{3}}$, this action corresponds to the bosonic part of minimal supergravity and can be derived from a consistent truncation of Type IIB supergravity or M-theory [24, 25]. More generally, k can be treated as a free parameter from the bottom-up perspective, and our discussion will be independent of the value of k . Henceforth, we shall set $16\pi G = \hbar = c = k_B = L = 1$.

One obtains the equations of motion:

$$G_{ab} - \frac{6}{L^2} g_{ab} - \frac{1}{2} (F_{ac} F_b{}^c - \frac{1}{4} g_{ab} F_{cd} F^{cd}) = 0, \quad (4)$$

$$\nabla_b F^{ba} + \frac{k}{8} \epsilon^{abcde} F_{bc} F_{de} = 0, \quad (5)$$

for which the dyonic black holes are expressed as

$$ds^2 = \frac{1}{r^2} \left[- (f e^{-\chi} - h^2 p^2) dt^2 + 2ph^2 dt dz + dx^2 + dy^2 + h^2 dz^2 + \frac{dr^2}{f} \right], \quad (6)$$

$$A = A_t dt + \frac{B}{2} (x dy - y dx) - A_z dz,$$

where f, χ, h, p, A_t and A_z are functions of the radial coordinate r . The constant B is nothing but the constant background magnetic field perpendicular to the x - y plane. The asymptotically AdS boundary is located at $r = 0$, while the black hole horizon is assumed at $r = r_h$.

The system admits an analytical black hole solution when $B = 0$:

$$f = 1 - \left(1 + \frac{\mu^2 r_h^2}{3} \right) \frac{r^4}{r_h^4} + \frac{\mu^2}{3r_h^4} r^6, \quad A_t = \mu \left(1 - \frac{r^2}{r_h^2} \right), \quad (7)$$

with $h = 1, \chi = p = A_z = 0$. This is nothing but the standard electrically charged AdS Reissner-Nordström black hole with chemical potential μ . However, for a non-zero magnetic field $B \neq 0$, one must solve the system numerically. Interestingly, the interplay between the electric charge and magnetic field leads to a quantum phase transition when the magnetic field crosses a critical value B_c [16–18].

For the dyonic black holes that asymptotically approach to AdS_5 , one has the the following asymptotic expansion as $r \rightarrow 0$,

$$\begin{aligned} f(r) &= 1 + \dots + f_4 r^4 + \dots, \\ \chi(r) &= \chi_0 + \dots, \\ h(r) &= 1 + \dots + h_4 r^4 + \dots, \\ p(r) &= p_4 r^4 + \dots, \\ A_t(r) &= e^{-\chi_0/2} \left(\mu - \frac{\rho}{2} r^2 + \dots \right), \\ A_z(r) &= A_{z2} r^2 + \dots. \end{aligned} \quad (8)$$

The boundary data are specified by seven parameters $f_4, \chi_0, h_4, p_4, \mu, \rho$ and A_{z2} . The physical interpretation of these parameters will be clear later. Near the event horizon, we impose regularity conditions with $f(r_h) = A_t(r_h) = p(r_h) = 0$. The temperature and entropy density are given by

$$T = - \frac{e^{\chi_0/2}}{4\pi} f' e^{-\chi/2} \Big|_{r=r_h}, \quad s = \frac{4\pi h}{r^3} \Big|_{r=r_h}. \quad (9)$$

The latter is from the famous Bekenstein-Hawking area law. We will finally set $\chi_0 = 0$ to ensure that the Hawking temperature of the black hole corresponds to be the temperature of the boundary field theory.

We now calculate the on-shell Euclidean action to analyse the thermodynamics of the black hole solutions. We consider the total Euclidean action I_E defined as

$$I_E = I + I_{bdy}, \quad (10)$$

where $I = -iS$ and I_{bdy} represents the Euclidean boundary action, including counter terms. Other physical quantities of the system can be obtained from the expectation values of the boundary stress-energy tensor $\langle T_{\mu\nu} \rangle$ and current $\langle J^\mu \rangle$. For more details, please refer to supplementary material [26]. The non-zero components are

$$\begin{aligned} \epsilon = \langle T_{tt} \rangle &= -3f_4 + 8h_4 + \frac{B^2}{4}, \\ \mathcal{P}_\perp = \langle T_{xx} \rangle = \langle T_{yy} \rangle &= -f_4 - \frac{B^2}{6}, \\ \mathcal{P}_\parallel = \langle T_{zz} \rangle &= -f_4 + 8h_4 + \frac{B^2}{12}, \\ \langle T_{tz} \rangle = \langle T_{zt} \rangle &= 4p_4 = -\frac{k}{2} B \mu^2, \end{aligned} \quad (11)$$

as well as

$$\langle J^t \rangle = \rho, \quad \langle J^z \rangle = -2A_{z2} = kB\mu, \quad (12)$$

where we have chosen the covariant current by omitting the Chern-Simons contribution. The momentum density $\langle T_{tz} \rangle$ and the current density $\langle J^z \rangle$ arise from the chiral anomaly. The transverse pressure \mathcal{P}_\perp differs from the longitudinal pressure \mathcal{P}_\parallel due to the magnetic field.

The free energy can be obtained using the quantum statistical relation (1). A detailed calculation reveals that w can be expressed in three equivalent forms

$$w = \epsilon - Ts - \mu\rho - \frac{kB}{3} \int_0^{r_h} A_t A'_z dr, \quad (13)$$

$$= -\mathcal{P}_\perp + B \int_0^{r_h} \left[\frac{B}{r} \left(e^{-\frac{\chi}{2} h} - 1 \right) + \frac{2k}{3} A_t A'_z \right] dr + B^2 \ln r_h, \quad (14)$$

$$= -\mathcal{P}_\parallel + \frac{kB}{3} \int_0^{r_h} A'_t A_z dr, \quad (15)$$

These expressions lead to the following Smarr-type relations:

$$\epsilon + \mathcal{P}_\parallel = Ts + \mu\rho, \quad (16)$$

$$\mathcal{P}_\parallel = \mathcal{P}_\perp - \left(\int_0^{r_h} \left[\frac{B}{r} \left(e^{-\frac{\chi}{2} h} - 1 \right) + k A_t A'_z \right] dr + B \ln r_h \right) B. \quad (17)$$

We highlight that the expression for the free energy in equation (13) includes a nontrivial bulk integration term that is proportional to the Chern-Simons coupling k , signaling a violation of the standard law of thermodynamics (2). Furthermore, a direct calculation of the entropy and magnetic susceptibility $\chi_B \equiv M_B/B$ from the free energy reveals that

$$-\left(\frac{\partial w}{\partial T} \right)_{B,\mu} \neq s, \quad -\frac{1}{B} \left(\frac{\partial w}{\partial B} \right)_{T,\mu} \neq \chi_B^{hydro}, \quad (18)$$

where s is the Bekenstein-Hawking entropy and χ_B^{hydro} is the magnetic susceptibility derived from field theory and hydrodynamics [27–29] (see also (36) and supplementary material). This discrepancy is clearly illustrated in Fig. 1. Such significant deviations indicate that the standard first law of black hole thermodynamics, as typically presented in textbooks, does not apply to the Einstein-Maxwell-Chern-Simons theory, suggesting an inconsistency within the system.

The cure of first law.—To address the issues above, we will conduct a thorough analysis and establish a consistent formulation of the first law of thermodynamics. Our approach will employ two independent methods: the variation of Euclidean on-shell action and the Iyer-Wald formalism. The computations are quite involved, please refer to the supplemental material for more details.

In Euclidean signature, the black hole horizon is smooth when the Euclidean time $\tau = it$ is periodic, with a period given by $\Delta\tau = 4\pi/(f'e^{-\chi/2})|_{r=r_h}$. Consequently, the only remaining boundary is the AdS boundary. The variation of the total on-shell Euclidean action




FIG. 1. Numerical tests of the first law of thermodynamics for dyonic black holes obtained by the quantum statistic relation (1). **Top:** Comparison of the Bekenstein-Hawking entropy s and $-(\partial w / \partial T)|_{\mu, B}$ at $B = 0.33$. **Bottom:** Comparison of magnetic susceptibility from hydrodynamics χ_B^{hydro} and $-(\partial w / \partial B)|_{\mu, T}$ at $T = 0.005$. One can find significant deviation. The plots are for minimal supergravity with $k = k_{susy} = \frac{2}{\sqrt{3}}$ and $\mu = 1$.

is given by

$$\delta I_E = \Delta\tau V \left\{ e^{-\frac{\chi_0}{2}} \left[-\rho\delta\mu - (e^{\chi_0}\epsilon - \mu\rho) \frac{\delta\chi_0}{2} \right] - k\delta Q_{cs} - e^{-\chi_0/2} M_B \delta B \right\}, \quad (19)$$

where $V = \int dx dy dz$. The quantity Q_{cs} , arising from the Chern-Simons term of (3), is given by

$$Q_{cs} = \frac{B}{6} \int_0^{r_h} (A'_z A_t - A_z A'_t) dr, \quad (20)$$

and M_B is defined as

$$M_B = - \left(\int_0^{r_h} \left[\frac{B}{r} \left(e^{-\chi/2} h - 1 \right) + \frac{k}{2} (A'_z A_t - A_z A'_t) \right] dr + B \ln r_h \right). \quad (21)$$

As we will show the latter is nothing other than the magnetization.

Note that the temperature is given by $T = \frac{e^{\chi_0/2}}{\Delta\tau}$. Therefore, the variation of χ_0 yields $\delta\chi_0 = 2\frac{\delta T}{T}$ with $\Delta\tau$ held fixed [30]. Using the quantum statistical relation (1), we obtain

$$\delta w = - \left(s + e^{\frac{\chi_0}{2}} \frac{kQ_{cs}}{T} \right) \delta T - \rho \delta \mu - e^{\frac{\chi_0}{2}} k \delta Q_{cs} - M_B \delta B. \quad (22)$$

It is manifest that the deviation from the standard form of the first law is attributed to the bulk integration Q_{cs} . Note also that (22) is independent of the specific definition of black hole energy. Nevertheless, after performing the Legendre transformation $\tilde{w} = w + e^{\chi_0/2} k Q_{cs}$, we recover the standard thermodynamic relation:

$$\tilde{w} = \epsilon - Ts - \mu\rho, \quad (23)$$

$$\delta\tilde{w} = -s\delta T - \rho\delta\mu - M_B\delta B, \quad (24)$$

where M_B is the magnetization of the system. Moreover, this identification also yields the expected relation

$$\mathcal{P}_{\parallel} = \mathcal{P}_{\perp} + M_B B = -\tilde{w}, \quad (25)$$

by using (17).

We have demonstrated that in order to uphold the standard laws of thermodynamics, as expressed in (23) and (24), the quantum statistical relation (1) must be modified. This conclusion can be definitively proven using the Iyer-Wald formalism which is a powerful framework for studying black hole thermodynamics [31, 32]. For example, it provides an elegant derivation of the extended black hole thermodynamics [33] and thermodynamics of black holes with scalar hair [34]. We will show that (23) and (24) can be derived exactly from the Iyer-Wald formalism.

Following the Iyer-Wald procedure [32], the variation of the Lagrangian 5-form $\mathbf{L} = \mathcal{L}\epsilon$ of (3) under a general variation of the dynamical fields $\phi = (g_{ab}, A_a)$ is expressed as

$$\delta\mathbf{L} = \mathbf{E}\delta\phi + d\Theta, \quad (26)$$

where \mathbf{E} represents the equations of motion (4) and (5), and Θ is the symplectic potential form. For a variation induced by an infinitesimal diffeomorphism $\delta_\xi x^a = \xi^a(x)$, Noether's theorem guarantees the existence of a Noether current 4-form defined by

$$\mathbf{J} \equiv \Theta(\phi, \mathcal{L}_\xi\phi) - \xi \cdot \mathbf{L}, \quad (27)$$

where $\xi \cdot$ denotes the contraction of ξ^a with the first index of \mathbf{L} . A standard calculation [35] shows that $d\mathbf{J} = -\mathbf{E}\mathcal{L}_\xi\phi$, which implies that \mathbf{J} is closed on-shell for any ξ^a . Thus, there exists a Noether charge 3-form \mathbf{Q} , locally constructed from ϕ and ξ^a , such that $\mathbf{J} = d\mathbf{Q}$ when the fields ϕ satisfy the equations of motion. Moreover, if $\delta\phi$ satisfies the linearized equations and ξ^a is a

symmetry of the solution, *i.e.* $\mathcal{L}_\xi\phi = 0$, we obtain the fundamental identity

$$\omega = \delta\mathbf{J} - d(\xi \cdot \Theta) = d(\delta\mathbf{Q} - \xi \cdot \Theta) = 0, \quad (28)$$

where $\omega = \omega(\phi, \delta\phi, \mathcal{L}_\xi\phi)$ denotes the symplectic current.

We choose the time-like Killing vector $\xi^a = (\partial_t)^a = \delta_t^a$, which vanishes at the event horizon $r = r_h$. Let Σ be a $t = \text{const.}$ space-like hypersurface with the horizon $r = r_h$ as its interior boundary. Thus, the boundary of this hypersurface, $\partial\Sigma$, includes the contributions from the non-compact x, y, z directions, namely

$$\begin{aligned} \partial\Sigma = & S_{r=r_h} \cup S_{r=0} \cup S_{x=L_x/2} \cup S_{x=-L_x/2} \cup \\ & S_{y=L_y/2} \cup S_{y=-L_y/2} \cup S_{z=L_z/2} \cup S_{z=-L_z/2}. \end{aligned}$$

Without loss of generality, we regulate the boundaries by considering finite limits (L_x, L_y, L_z) along the three spatial directions, giving a spatial volume $V = L_x L_y L_z$. By integrating (28) over the hypersurface Σ , we derive the first law of black hole thermodynamics:

$$\delta\epsilon = T\delta s + \mu\delta\rho - M_B\delta B, \quad (29)$$

where M_B is defined as in (21).

In addition to the fundamental identity (28), we have

$$d\mathbf{Q} = -\xi \cdot \mathbf{L}, \quad (30)$$

when the dynamical fields ϕ satisfy the equations of motion and ξ is a Killing vector (*i.e.* $\mathcal{L}_\xi\phi = 0$). By integrating (30) over the hypersurface Σ and applying Stokes's theorem, we obtain

$$\int_{\partial\Sigma} \mathbf{Q} = - \int_{\Sigma} \xi \cdot \mathbf{L}. \quad (31)$$

From this, we obtain explicitly that

$$\tilde{w} = \epsilon - Ts - \mu\rho = w + kQ_{cs}, \quad (32)$$

which is the expected thermodynamic relation, confirming (23). Combining (29) and (32), we obtain

$$\delta\tilde{w} = -s\delta T - \rho\delta\mu - M_B\delta B, \quad (33)$$

which matches exactly the results obtained from the on-shell variation of the Euclidean action, as given in (23) and (24). Considering the scaling symmetry together with the first law, one obtains the (generalized) Smarr relation

$$4\epsilon - 3(Ts + \mu\rho) + 2M_B B = \frac{B^2}{2}. \quad (34)$$

The left-hand side of this equation is non-zero in the presence of magnetic field B , indicating a derivation from the standard Smarr relation due to the chiral anomaly. This situation is analogous to the Kerr-AdS case, where the breaking of scale invariance leads to a generalized Smarr formula [3].

Furthermore, using the expressions for the bulk on-shell action (see supplementary material), we obtain from (31) that $\mathcal{P}_{\parallel} = \mathcal{P}_{\perp} + M_B B$ which matches exactly with (25). By using (11), we derive an explicit expression for the magnetization:

$$M_B = \frac{\mathcal{P}_{\parallel} - \mathcal{P}_{\perp}}{B} = \frac{B}{4} + \frac{8h_4}{B}, \quad (35)$$

in addition to $-(\partial\tilde{w}/\partial B)|_{\mu,T}$ from (33). As a consistency check, we find that the magnetic susceptibility obtained from field theory and hydrodynamics [27–29]:

$$\chi_B^{hydro} \equiv \frac{M_B}{B} = \frac{1}{4} + \frac{8h_4}{B^2}, \quad (36)$$

is in exact agreement with the one from our result (35).




FIG. 2. Numerical verification of the thermodynamics of dyonic black holes in minimal supergravity: $\delta\tilde{w} = -s\delta T - \rho\delta\mu - M_B\delta B$. **Top:** Comparison of the Bekenstein-Hawking entropy s of (9) and $-(\partial\tilde{w}/\partial T)|_{\mu,B}$ at $B = 0.33$. **Bottom:** Comparison of magnetization M_B of (35) and $-(\partial\tilde{w}/\partial B)|_{\mu,T}$ at $T = 0.005$. The agreement is perfect within numerical errors. The plots are generated for $k = k_{\text{susy}} = \frac{2}{\sqrt{3}}$ and $\mu = 1$.

We now verify our results using the full numerical computations presented in Fig. 2. The top panel of Fig. 2 illustrates the temperature dependence of entropy density s both from the Bekenstein-Hawking entropy (9) and the one derived from the first law (33). These two results are in perfect agreement, in sharp contrast to the result from the quantum statistical relation (1), as shown in Fig. 1. The bottom panel of Fig. 2 depicts the magnetization as a function of magnetic field, where the analytical expression (21) quantitatively matches the numerical results obtained from the first law (33).

Conclusion.—We have investigated the thermodynamics of dyonic black hole in five-dimensional Einstein-Maxwell gravity with a Chern-Simons term, obtained from IIB supergravity. Surprisingly, we have found that the textbook results for the first law of thermodynamics break down when applying the quantum statistical relation (1), as illustrated in Fig. 1. Using two independent methods, the on-shell variation of the total Euclidean action and the Iyer-Wald formalism, we have resolved this issue and established the standard first law of thermodynamics, which agrees with field theory and hydrodynamics, and has been validated through numerical tests (see Fig. 2). We have explicitly demonstrated that the free energy of the system should be expressed as $\tilde{w} = w + kQ_{cs}$ of (32), rather than w derived from the on-shell action (1), which has been widely adopted in the literature. It is important to note that Q_{cs} of (20) is a non-local bulk integration that cannot be eliminated by adding any local boundary term when computing the on-shell Euclidean action via (1).

Our work highlights the necessity for a deeper understanding of black hole thermodynamics, particularly in the presence of magnetic fields. Specifically, a correct formulation of the first law of black hole thermodynamics is crucial for studying magnetic phenomena, such as magnetization, the magnetocaloric effect, and quantum phase transitions induced by magnetic fields in holography. The methods employed here are applicable to any geometric theory of gravity. We anticipate that similar results may arise in other systems, such as [36]. Furthermore, it is of significant interest to provide a microscopic interpretation for our findings from both the supergravity perspective and the dual boundary quantum field theory.

Acknowledgments—We thank Mirjam Cvetič, Hong Lu, Peng-Zhang He, Yong Xiao, Amos Yarom and Hua Zhan for useful discussions. This work was partly supported by the National Natural Science Foundation of China Grants No. 12075298, No. 12122513, No. 11991052, No. 12047503, and No.12247156.

[1] S. W. Hawking, Particle Creation by Black Holes, *Commun. Math. Phys.* **43**, 199 (1975), [Erratum: *Commun.*

mun.Math.Phys. **46**, 206 (1976)].

- [2] G. W. Gibbons and S. W. Hawking, Action Integrals and Partition Functions in Quantum Gravity, *Phys. Rev. D* **15**, 2752 (1977).
- [3] G. W. Gibbons, M. J. Perry, and C. N. Pope, The First law of thermodynamics for Kerr-anti-de Sitter black holes, *Class. Quant. Grav.* **22**, 1503 (2005), [arXiv:hep-th/0408217](#).
- [4] S. W. Hawking and D. N. Page, Thermodynamics of Black Holes in anti-De Sitter Space, *Commun. Math. Phys.* **87**, 577 (1983).
- [5] A. Chamblin, R. Emparan, C. V. Johnson, and R. C. Myers, Charged AdS black holes and catastrophic holography, *Phys. Rev. D* **60**, 064018 (1999), [arXiv:hep-th/9902170](#).
- [6] D. Kubiznak and R. B. Mann, P-V criticality of charged AdS black holes, *JHEP* **07**, 033, [arXiv:1205.0559 \[hep-th\]](#).
- [7] J. Casalderrey-Solana, H. Liu, D. Mateos, K. Rajagopal, and U. A. Wiedemann, *Gauge/String Duality, Hot QCD and Heavy Ion Collisions* (Cambridge University Press, 2014) [arXiv:1101.0618 \[hep-th\]](#).
- [8] M. Ammon and J. Erdmenger, *Gauge/gravity duality: Foundations and applications* (Cambridge University Press, Cambridge, 2015).
- [9] J. Zaanen, Y.-W. Sun, Y. Liu, and K. Schalm, *Holographic Duality in Condensed Matter Physics* (Cambridge Univ. Press, 2015).
- [10] S. A. Hartnoll, A. Lucas, and S. Sachdev, *Holographic Quantum Matter* (MIT Press, 2018).
- [11] G. Barnich and G. Compere, Conserved charges and thermodynamics of the spinning Godel black hole, *Phys. Rev. Lett.* **95**, 031302 (2005), [arXiv:hep-th/0501102](#).
- [12] E. G. Gimon and A. Hashimoto, Black holes in Godel universes and pp waves, *Phys. Rev. Lett.* **91**, 021601 (2003), [arXiv:hep-th/0304181](#).
- [13] J. B. Gutowski and H. S. Reall, Supersymmetric AdS(5) black holes, *JHEP* **02**, 006, [arXiv:hep-th/0401042](#).
- [14] Z. W. Chong, M. Cvetič, H. Lu, and C. N. Pope, General non-extremal rotating black holes in minimal five-dimensional gauged supergravity, *Phys. Rev. Lett.* **95**, 161301 (2005), [arXiv:hep-th/0506029](#).
- [15] A. Donos and J. P. Gauntlett, Black holes dual to helical current phases, *Phys. Rev. D* **86**, 064010 (2012), [arXiv:1204.1734 \[hep-th\]](#).
- [16] E. D'Hoker and P. Kraus, Magnetic Field Induced Quantum Criticality via new Asymptotically AdS₅ Solutions, *Class. Quant. Grav.* **27**, 215022 (2010), [arXiv:1006.2573 \[hep-th\]](#).
- [17] E. D'Hoker and P. Kraus, Holographic Metamagnetism, Quantum Criticality, and Crossover Behavior, *JHEP* **05**, 083, [arXiv:1003.1302 \[hep-th\]](#).
- [18] E. D'Hoker and P. Kraus, Quantum Criticality via Magnetic Branes, *Lect. Notes Phys.* **871**, 469 (2013), [arXiv:1208.1925 \[hep-th\]](#).
- [19] J. Erdmenger, M. Haack, M. Kaminski, and A. Yarom, Fluid dynamics of R-charged black holes, *JHEP* **01**, 055, [arXiv:0809.2488 \[hep-th\]](#).
- [20] N. Banerjee, J. Bhattacharya, S. Bhattacharyya, S. Dutta, R. Loganayagam, and P. Surowka, Hydrodynamics from charged black branes, *JHEP* **01**, 094, [arXiv:0809.2596 \[hep-th\]](#).
- [21] D. T. Son and P. Surowka, Hydrodynamics with Triangle Anomalies, *Phys. Rev. Lett.* **103**, 191601 (2009), [arXiv:0906.5044 \[hep-th\]](#).
- [22] L. D. Landau, E. Lifshitz, and L. P. Pitaevskii, *Electrodynamics of Continuous Media*, (Butterworth-Heinemann, 1995).
- [23] Alternatively, we can define a total energy $\epsilon^{tot} = \epsilon + M_B B$. Then, the first law of thermodynamics is given by $\delta\epsilon^{tot} = T\delta s + \mu\delta\rho + B\delta M_B$.
- [24] A. Buchel and J. T. Liu, Gauged supergravity from type IIB string theory on Y**p,q manifolds, *Nucl. Phys. B* **771**, 93 (2007), [arXiv:hep-th/0608002](#).
- [25] J. P. Gauntlett, E. O. Colgain, and O. Varela, Properties of some conformal field theories with M-theory duals, *JHEP* **02**, 049, [arXiv:hep-th/0611219](#).
- [26] The supplementary material provides more details for the definition of thermodynamic quantities, the variation of the Euclidean action, the computation of Iyer-Wald formalism and an overview of magnetic susceptibility from hydrodynamics, supporting the results made in the main text, which also includes relevant references [37–41].
- [27] P. Kovtun, Thermodynamics of polarized relativistic matter, *JHEP* **07**, 028, [arXiv:1606.01226 \[hep-th\]](#).
- [28] K. Jensen, R. Loganayagam, and A. Yarom, Anomaly inflow and thermal equilibrium, *JHEP* **05**, 134, [arXiv:1310.7024 \[hep-th\]](#).
- [29] M. Ammon, M. Kaminski, R. Koirala, J. Leiber, and J. Wu, Quasinormal modes of charged magnetic black branes & chiral magnetic transport, *JHEP* **04**, 067, [arXiv:1701.05565 \[hep-th\]](#).
- [30] A. Donos and J. P. Gauntlett, On the thermodynamics of periodic AdS black branes, *JHEP* **10**, 038, [arXiv:1306.4937 \[hep-th\]](#).
- [31] R. M. Wald, Black hole entropy is the Noether charge, *Phys. Rev. D* **48**, R3427 (1993), [arXiv:gr-qc/9307038](#).
- [32] V. Iyer and R. M. Wald, Some properties of Noether charge and a proposal for dynamical black hole entropy, *Phys. Rev. D* **50**, 846 (1994), [arXiv:gr-qc/9403028](#).
- [33] Y. Xiao, Y. Tian, and Y.-X. Liu, Extended Black Hole Thermodynamics from Extended Iyer-Wald Formalism, *Phys. Rev. Lett.* **132**, 021401 (2024), [arXiv:2308.12630 \[gr-qc\]](#).
- [34] L. Li, On Thermodynamics of AdS Black Holes with Scalar Hair, *Phys. Lett. B* **815**, 136123 (2021), [arXiv:2008.05597 \[gr-qc\]](#).
- [35] J. Lee and R. M. Wald, Local symmetries and constraints, *J. Math. Phys.* **31**, 725 (1990).
- [36] E. D'Hoker and P. Kraus, Charge Expulsion from Black Brane Horizons, and Holographic Quantum Criticality in the Plane, *JHEP* **09**, 105, [arXiv:1202.2085 \[hep-th\]](#).
- [37] J. P. Gauntlett, J. Sonner, and T. Wiseman, Quantum Criticality and Holographic Superconductors in M-theory, *JHEP* **02**, 060, [arXiv:0912.0512 \[hep-th\]](#).
- [38] R. M. Wald, On identically closed forms locally constructed from a field, *J. Math. Phys.* **31**, 2378 (1990).
- [39] S. Hollands and R. M. Wald, Stability of Black Holes and Black Branes, *Commun. Math. Phys.* **321**, 629 (2013), [arXiv:1201.0463 \[gr-qc\]](#).
- [40] M. Ammon, J. Leiber, and R. P. Macedo, Phase diagram of 4D field theories with chiral anomaly from holography, *JHEP* **03**, 164, [arXiv:1601.02125 \[hep-th\]](#).
- [41] I. Amado, K. Landsteiner, and F. Pena-Benitez, Anomalous transport coefficients from Kubo formulas in Holography, *JHEP* **05**, 081, [arXiv:1102.4577 \[hep-th\]](#).

SUPPLEMENTARY MATERIAL

This supplementary material provides a detailed analysis referenced in the main text. We present the equations of motion for the dyonic black holes and compute all thermodynamic quantities. We then outline the on-shell variation of the Euclidean action and the Iyer-Wald formalism in detail. We also derive the Smarr relation and offer a brief overview of the hydrodynamic description of the chiral charged fluid in the presence of a strong external magnetic field.

A: Equations of motion and thermodynamics

The explicit form of the equations of motion for the dyonic black holes is given by

$$\begin{aligned}
& \left[\frac{e^{\chi/2} h}{r} (A'_t + p A'_z) \right]' + k B A'_z = 0, \\
& \left[\frac{e^{\chi/2}}{r h} (p h^2 A'_t - (f e^{-\chi} - h^2 p^2) A'_z) \right]' - k B A'_t = 0, \\
& \frac{f''}{f} - \chi'' + \left(\frac{\chi'}{2} - \frac{3f'}{2f} \right) \chi' - \frac{3e^\chi (h^2 p'^2 + r^2 (A'_t + p A'_z)^2)}{2f} + \frac{6h'}{r h} + \frac{r^2 A_z'^2}{2h^2} - \frac{3B^2 r^2}{2f} + \frac{12 - 12f}{r^2 f} = 0, \\
& \frac{f''}{f} - \chi'' - \left(\frac{h'}{h} + \frac{3\chi'}{2} + \frac{3}{r} \right) \frac{f'}{f} + \left(\frac{h'}{h} + \frac{\chi'}{2} \right) \chi' - \frac{e^\chi (2h^2 p'^2 + r^2 (A'_t + p A'_z)^2)}{f} + \frac{6h'}{r h} = 0, \\
& h'' + \left(\frac{f'}{f} - \frac{\chi'}{2} - \frac{3}{r} \right) h' - \frac{B^2 r^2 h}{2f} + \frac{e^\chi h^3 p'^2}{2f} + \frac{r^2 A_z'^2}{2h} = 0, \\
& p'' + \left(\frac{3h'}{h} + \frac{\chi'}{2} - \frac{3}{r} \right) p' - \frac{r^2 A'_z (A'_t + p A'_z)}{h^2} = 0,
\end{aligned} \tag{A1}$$

where the prime denotes the derivative with respect to r . The above equations of motion has the following scaling symmetry:

$$(r, t, x, y, z) \rightarrow \lambda(r, t, x, y, z), \quad (A_t, A_z) \rightarrow \lambda^{-1}(A_t, A_z), \quad B \rightarrow \lambda^{-2} B, \quad (f, \chi, h, p) \rightarrow (f, \chi, h, p), \tag{A2}$$

with λ a positive constant.

To solve the above equations, we first need to determine the asymptotic expansions of the bulk fields. Near the AdS boundary $r = 0$, we obtain the following expansions:

$$\begin{aligned}
f(r) &= 1 + \frac{B^2}{2} r^4 \ln r + f_4 r^4 + \dots, \\
\chi(r) &= \chi_0 + \frac{B^2}{4} r^4 \ln r + \left(\frac{B^2}{12} + 2h_4 \right) r^4 + \dots, \\
h(r) &= 1 + \frac{B^2}{8} r^4 \ln r + h_4 r^4 + \dots, \\
p(r) &= p_4 r^4 + \dots, \\
A_t(r) &= e^{-\chi_0/2} \left(\mu - \frac{\rho}{2} r^2 + \dots \right), \\
A_z(r) &= A_{z2} r^2 + \dots.
\end{aligned} \tag{A3}$$

Note that χ_0 should be set to zero to ensure that the Hawking temperature of the black hole matches the temperature of the boundary field theory. However, to account for temperature variations in the Euclidean action, χ_0 will be retained as a free parameter and will be set to zero only in the final expressions presented in the main text.

Near the event horizon $r = r_h$, we impose the conditions $f(r_h) = A_t(r_h) = 0$ to ensure a smooth horizon. Additionally, by applying the α symmetry transformation: $z \rightarrow z - \alpha t, p \rightarrow p - \alpha, A_t \rightarrow A_t + \alpha A_z$ with α being a

constant, we can set $p(r_h) = 0$. Near the black hole horizon, the field expansions are given by:

$$\begin{aligned}
f(r) &= f_1(r_h - r) + \dots, \\
\chi(r) &= \chi_1 + \dots, \\
h(r) &= h_1 + \dots, \\
p(r) &= p_1(r_h - r) + \dots, \\
A_t(r) &= A_{t1}(r_h - r) + \dots, \\
A_z(r) &= A_{z1} + \dots.
\end{aligned} \tag{A4}$$

Note that the scaling symmetry (A2) can be used to fix the location of the black hole horizon to $r_h = 1$ that is convenient in numerics. The interplay between a finite charge density and a background magnetic field leads to a quantum phase transition at an intermediate magnetic field strength [16, 17].

With the background solutions determined, we can study the thermodynamic properties of the dyonic black holes. The on-shell action suffers from divergence and we need to regulate it by adding appropriate counter terms at the AdS boundary. The renormalized on-shell action is expressed as:

$$S_{ren} = S + S_{bdy} = S + \frac{1}{16\pi G} \int d^4x \sqrt{-\gamma} \left[2K - 6 - \frac{\hat{R}}{2} + \frac{\ln r}{4} (\hat{R}_{\mu\nu} \hat{R}^{\mu\nu} - \frac{\hat{R}^2}{3} - F^2) \right], \tag{A5}$$

where $\gamma_{\mu\nu}$ is the induced metric at the conformal boundary $r \rightarrow 0$, K is the trace of the extrinsic curvature, and $\hat{R}_{\mu\nu}$ denotes the Ricci tensor associated with $\gamma_{\mu\nu}$.

To obtain analytical expressions for the thermodynamic variables, we can utilize the properties of the equations of motion and the Ricci tensor to rewrite the on shell action in terms of surface terms. The Einstein equation implies that its (i, i) component satisfies

$$2R^i_i = R + \frac{12}{L^2} - \frac{1}{4}F^2 + F^i_c F_i^c, \quad (i = t, x, y, z), \tag{A6}$$

where there is no summation over i in this expression. Consequently, the Lagrangian density \mathcal{L} can be rewritten as

$$\mathcal{L} = R + \frac{12}{L^2} - \frac{1}{4}F^2 + \frac{k}{24} \epsilon^{abcde} A_a F_{bc} F_{de} = 2R^i_i - F^i_c F_i^c + \frac{k}{24} \epsilon^{abcde} A_a F_{bc} F_{de}, \tag{A7}$$

where i represents one of the coordinates (t, x, y, z) . Using the identity $R^a_b \xi^b = \nabla_b \nabla^a \xi^b$ for any killing vector ξ^a , we can express R^i_i as a total derivative. Specifically, we have

$$\begin{aligned}
R^a_b \xi^b &= \nabla_b \nabla^a \xi^b = -\nabla_b \nabla^b \xi^a, \\
&= -\frac{1}{\sqrt{-g}} \partial_b (\sqrt{-g} g^{bc} \nabla_c \xi^a), \\
&= -\frac{1}{\sqrt{-g}} \partial_r (\sqrt{-g} g^{rc} \nabla_c \xi^a),
\end{aligned} \tag{A8}$$

where we have used the fact that the background fields depend only on the radial coordinate r . For each killing vector ∂_t, ∂_x and ∂_z , we obtain

$$\begin{aligned}
2\sqrt{-g}R^t_t &= \left[\frac{e^{-\chi/2} h}{r^4} (-rf' + re^\chi h^2 pp' + f(2 + r\chi')) \right]', \\
2\sqrt{-g}R^x_x &= 2\sqrt{-g}R^y_y = \left[\frac{2e^{-\chi/2} fh}{r^4} \right]', \\
2\sqrt{-g}R^z_z &= \left[\frac{e^{-\chi/2}}{r^4} (2f(h - rh') - re^\chi h^3 pp') \right]'.
\end{aligned} \tag{A9}$$

Using the expression for the Lagrangian density (A7), we can convert the bulk action into a total derivative. After some calculations, the on-shell action is given by

$$S = \int d^4x \int dr \left[2\sqrt{-g}R^t_t + (A_t N_t + \frac{k}{3} B A_t A_z)' + \frac{k}{3} B A_t A_z' \right], \tag{A10}$$

$$= \int d^4x \int dr \left[2\sqrt{-g}R^x_x - \frac{B^2 e^{-\chi/2} h}{r} + \frac{k}{3} B (A'_t A_z - A_t A'_z) \right], \tag{A11}$$

$$= \int d^4x \int dr \left[2\sqrt{-g}R^z_z + (A_z N_z - \frac{k}{3} B A_t A_z)' - \frac{k}{3} B A'_t A_z \right], \tag{A12}$$

where $N_t \equiv \frac{e^{\chi/2} h}{r} (A'_t + p A'_z)$ and $N_z \equiv \frac{e^{\chi/2}}{r h} (p h^2 A'_t - (f e^{-\chi} - h^2 p^2) A'_z)$. Additionally, we have $N'_t = -k B A'_z$ and $N'_z = k B A'_t$ as derived from the Maxwell equation.

By substituting (A10-A12) and the UV and IR expansions into (A5), the renormalized on-shell action reads

$$\begin{aligned} \frac{S_{ren}}{\Delta T V} &= 3f_4 - 8h_4 - \frac{B^2}{4} + \mu\rho + e^{-\frac{\chi}{2}} f_1 h_1 + \frac{kB}{3} \int_0^{r_h} A_t A'_z dr, \\ &= -f_4 - \frac{B^2}{6} - B^2 \ln r_h - B \int_0^{r_h} \left[\frac{B}{r} \left(e^{-\frac{\chi}{2}} h - 1 \right) - \frac{k}{3} (A'_t A_z - A_t A'_z) \right] dr, \\ &= -f_4 + 8h_4 + \frac{B^2}{12} - \frac{kB}{3} \int_0^{r_h} A'_t A_z dr, \end{aligned} \quad (A13)$$

where $\Delta T = \int dt$ and $V = \int dx dy dz$.

The boundary stress tensor and current can be obtained by varying the on-shell action with respect to the boundary metric $\gamma_{\mu\nu}$ and the boundary value of gauge field. The stress tensor is given by

$$\langle T_{\mu\nu} \rangle = \lim_{r \rightarrow 0} \frac{1}{r^2} \left[-2K_{\mu\nu} + 2(K-3)\gamma_{\mu\nu} + \ln r \left(F_{\mu\rho} F_{\nu}{}^\rho - \frac{\gamma_{\mu\nu}}{4} F^2 \right) \right], \quad (A14)$$

for which the non-zero components of $\langle T_{\mu\nu} \rangle$ are:

$$\begin{aligned} \epsilon = \langle T_{tt} \rangle &= -3f_4 + 8h_4 + \frac{B^2}{4}, & \mathcal{P}_\perp = \langle T_{xx} \rangle = \langle T_{yy} \rangle &= -f_4 - \frac{B^2}{6}, \\ \mathcal{P}_\parallel = \langle T_{zz} \rangle &= -f_4 + 8h_4 + \frac{B^2}{12}, & \langle T_{tz} \rangle = \langle T_{zt} \rangle &= 4p_4. \end{aligned} \quad (A15)$$

Similarly, the current dual to the gauge field is given by

$$\langle J^\mu \rangle = \lim_{r \rightarrow 0} \sqrt{-\gamma} \left[n_r \left(F^{\mu r} + \frac{k}{6} \epsilon^{r\mu\alpha\beta\gamma} A_\alpha F_{\beta\gamma} \right) + \nabla_\alpha F^{\alpha\mu} \ln r \right]. \quad (A16)$$

The non-zero components of the current are:

$$\langle J^t \rangle = \rho, \quad \langle J^z \rangle = -2A_{z2}, \quad (A17)$$

where we have chosen the covariant current by simply dropping the Chern-Simons contribution in $\langle J^z \rangle$. Note that, combining the equations of motion for gauge field and the asymptotic expansions at UV and IR, we can obtain $A_{z2} = -kB\mu/2$ and thus $\langle J^z \rangle = kB\mu$. Further, the chiral anomaly leads to $p_4 = -kB\mu^2/8$ *i.e.* the momentum density $\langle T_{tz} \rangle = -kB\mu^2/2$ [40, 41]. As numerically checked, $\langle J^z \rangle$ and $\langle T_{tz} \rangle$ are indeed constant for fixed magnetic field and are independent of the temperature.

To obtain the on-shell Euclidean action I_E , we perform an analytic continuation by Wick-rotating t to the Euclidean time coordinate $\tau = it$. To ensure that both the metric and gauge field are real in the Euclidean signature, we also set $A_{t(E)} = -iA_t$, $p_{(E)} = -ip$. Consequently, the Euclidean metric and gauge field are given by

$$\begin{aligned} ds_{(E)}^2 &= \frac{1}{r^2} \left[(f e^{-\chi} + h^2 p_{(E)}^2) d\tau^2 + 2p_{(E)} h^2 d\tau dz + dx^2 + dy^2 + h^2 dz^2 + \frac{dr^2}{f} \right], \\ A_{(E)} &= A_{t(E)} d\tau - \frac{B}{2} y dx + \frac{B}{2} x dy - A_z dz. \end{aligned} \quad (A18)$$

To keep the Chern-Simons term real in the Euclidean signature, we also choose $k_{(E)} = ik$. Thus, the Chern-Simons term becomes

$$\sqrt{g_{(E)}} \frac{k_{(E)}}{24} \epsilon^{abcde} A_{a(E)} F_{bc(E)} F_{de(E)} = \frac{k_{(E)} B}{3} \left(A'_{t(E)} A_z - A_{t(E)} A'_z \right). \quad (A19)$$

Then, the Euclidean action is defined as $I_E = -iS_{ren}$ and the free energy density can be expressed as

$$\begin{aligned} w &= -\frac{T \ln Z}{V} = \frac{TI_E}{V}, \\ &= \epsilon - Ts - \mu\rho - \frac{kB}{3} \int_0^{r_h} A_t A'_z dr, \\ &= -\langle T_{xx} \rangle + B \int_0^{r_h} \left[\frac{B}{r} \left(e^{-\frac{\chi}{2}} h - 1 \right) + \frac{2k}{3} A_t A'_z \right] dr + B^2 \ln r_h, \\ &= -\langle T_{zz} \rangle + \frac{kB}{3} \int_0^{r_h} A'_t A_z dr, \end{aligned} \quad (A20)$$

where we have used the relation $\int_0^{r_h} A_t A'_z dr = -\int_0^{r_h} A'_t A_z dr$, which follows from the boundary condition $A_t(r = r_h) = A_z(r = 0) = 0$.

B: Variation of Euclidean action

Substituting the background ansatz into the original action S , the effective action of the system is given by

$$S_{eff} = \int d^5x \frac{e^{-\chi/2} h}{r^3} \left\{ -f'' - \frac{2fh''}{h} + f\chi'' + f' \left(\frac{8}{r} - \frac{2h'}{h} + \frac{3\chi'}{2} \right) + \left(\frac{h'}{rh} - \frac{\chi'}{2r} \right) (8f + rf\chi') \right. \\ \left. + \frac{1}{2} e^\chi [h^2 p'^2 + r^2 (A'_t + pA'_z)^2] + \frac{12 - 20f}{r^2} - \frac{r^2}{2} \left(B^2 + \frac{fA_z'^2}{h^2} \right) \right\} + \int d^5x \frac{kB}{3} (A'_t A_z - A_t A'_z). \quad (\text{B1})$$

Next, we consider analytical continuation to Euclidean time τ by performing a Wick rotation $t = -i\tau$. The Euclidean action is given by

$$I_E = I + I_{bdy}, \quad (\text{B2})$$

where $I = -iS = -iS_{eff}$ and I_{bdy} is the Euclidean boundary action obtained from (A5). Consequently, the variation of the total on-shell Euclidean action can be separated into two parts: $\delta I_E = \delta I + \delta I_{bdy}$. The on-shell variation of I yields

$$\frac{\delta I}{\Delta\tau V} = - \int dr \partial_r \left\{ \frac{e^{-\chi/2} h}{r^3} \left[\left(\frac{5}{r} - \frac{h'}{h} + \chi' \right) \delta f - \delta f' \right] + \frac{2e^{-\chi/2} f}{r^3} \left(\frac{\delta h}{r} - \delta h' \right) \right. \\ \left. + \frac{e^{-\chi/2} fh}{r^3} \left[\left(\frac{f'}{2f} - \frac{\chi'}{2} - \frac{1}{u} \right) \delta\chi + \delta\chi' \right] + \frac{e^{\chi/2} h^3 p'}{r^3} \delta p + \frac{kB}{3} A_z \delta A_t \right. \\ \left. + \frac{e^{\chi/2} h (A'_t + pA'_z)}{r} (\delta A_t + p\delta A_z) - \left(\frac{kB}{3} A_t + \frac{e^{-\chi/2} f A'_z}{rh} \right) \delta A_z \right\} \\ - \int dr \left[\frac{kB}{3} (A'_z \delta A_t - \delta A_z A'_t) - \left(\frac{B e^{-\chi/2} h}{r} + \frac{k}{3} (A_t A'_z - A'_t A_z) \right) \delta B \right], \quad (\text{B3})$$

while the variation from boundary terms I_{bdy} is given by

$$\frac{\delta I_{bdy}}{\Delta\tau V} = - \left\{ \frac{e^{-\chi/2} h}{r^3} \left[\left(\frac{8}{r} - \frac{12 + B^2 r^4 \ln r}{4r\sqrt{f}} - \frac{2h'}{h} + \chi' \right) \delta f - \delta f' \right] \right. \\ \left. + \frac{e^{-\chi/2} f}{r^3} \left[\left(\frac{8}{r} - \frac{12 + B^2 r^4 \ln r}{2r\sqrt{f}} - \frac{f'}{f} + \chi' \right) \delta h - 2\delta h' \right] \right. \\ \left. + \frac{e^{-\chi/2} fh}{2r^3} \left[\left(-\frac{8}{r} + \frac{12 + B^2 r^4 \ln r}{2r\sqrt{f}} + \frac{f'}{f} + \frac{2h'}{h} - \chi' \right) \delta\chi + 2\delta\chi' \right] \right. \\ \left. - \left(B e^{-\chi/2} \sqrt{f} h \ln r \right) \delta B \right\} \Big|_{r \rightarrow 0}. \quad (\text{B4})$$

Here, $\Delta\tau \equiv \int d\tau$ and $V \equiv \int dx dy dz$.

Note that in the variation of I , there is a bulk integral term given by $k \int \frac{B}{3} (A'_z \delta A_t - \delta A_z A'_t) dr$. This suggests a limitation in the method used for obtaining the background equations directly from the variation of the effective action (B1), as discussed, for example, in [37]. Nevertheless, it is possible to convert this bulk integral into a total derivative term by observing that

$$A'_z \delta A_t - \delta A_z A'_t = \frac{1}{2} \delta (A'_z A_t - A_z A'_t) - \frac{1}{2} [\delta A_z A_t - A_z \delta A_t]'. \quad (\text{B5})$$

Thus, we obtain

$$\int dr \frac{kB}{3} (A'_z \delta A_t - \delta A_z A'_t) \\ = k\delta \left[\frac{B}{6} \int dr (A'_z A_t - A_z A'_t) \right] - \left[\frac{k}{6} \int dr (A'_z A_t - A_z A'_t) \right] \delta B - \frac{kB}{6} [\delta A_z A_t - A_z \delta A_t]_0^{r_h}, \\ = k\delta \left[\frac{B}{6} \int dr (A'_z A_t - A_z A'_t) \right] - \left[\frac{k}{6} \int dr (A'_z A_t - A_z A'_t) \right] \delta B, \quad (\text{B6})$$

where we have used the boundary conditions $A_t(r = r_h) = A_z(r = 0) = 0$.

In the Euclidean metric, the spacetime near the black brane horizon resembles a cigar geometry, and hence the only remaining boundary is the AdS boundary. Using the asymptotic expansions, the variation of the total on-shell Euclidean action is given by

$$\delta I_E = \delta I + \delta I_{bdy} = \Delta\tau V \left\{ e^{-\frac{\chi_0}{2}} \left[-\rho\delta\mu - (e^{\chi_0}\epsilon - \mu\rho) \frac{\delta\chi_0}{2} \right] - k\delta Q_{cs} - e^{-\chi_0/2} M_B \delta B \right\}, \quad (B7)$$

where Q_{cs} is defined as

$$Q_{cs} = \frac{B}{6} \int_0^{r_h} (A'_z A_t - A_z A'_t) dr, \quad (B8)$$

and M_B is given by

$$M_B = - \left(\int_0^{r_h} \left[\frac{B}{r} \left(e^{-\chi/2} h - 1 \right) + \frac{k}{2} (A'_z A_t - A_z A'_t) \right] dr + B \ln r_h \right). \quad (B9)$$

C: Iyer-Wald formalism

The variation of Lagrangian 5-form $\mathbf{L} = \mathcal{L}\epsilon$ is given by

$$\delta\mathbf{L} = \mathbf{E}\delta\phi + d\Theta, \quad (C1)$$

where $\mathbf{E}\delta\phi = (\mathbf{E}_g)^{ab}\delta g_{ab} + (\mathbf{E}_A)^a\delta A_a$ represents the equations of motion for the dynamical fields $\phi = (g_{ab}, A_a)$ and Θ is the symplectic potential form defined as

$$\Theta = \epsilon_{aa_1 a_2 a_3 a_4} \left[(g^{ac} g^{bd} - g^{ab} g^{cd}) \nabla_b \delta g_{cd} - \left(F^{ae} + \frac{k}{6} \epsilon^{abcde} A_b F_{cd} \right) \delta A_e \right]. \quad (C2)$$

The symplectic current $(n-1)$ form can be defined from Θ as

$$\omega(\phi, \delta_1\phi, \delta_2\phi) = \delta_2\Theta(\phi, \delta_1\phi) - \delta_1\Theta(\phi, \delta_2\phi), \quad (C3)$$

where $\delta_1\phi$ and $\delta_2\phi$ are variations of the dynamical fields.

Considering an infinitesimal diffeomorphism $\delta_\xi x^a = \xi^a(x)$, one can associate with it a Noether current 4-form, defined by

$$\mathbf{J} \equiv \Theta(\phi, \mathcal{L}_\xi\phi) - \xi \cdot \mathbf{L}, \quad (C4)$$

where $\xi \cdot$ denotes the contraction of ξ^a with the first index of \mathbf{L} . Under the infinitesimal diffeomorphism transformations $\delta_\xi x^a = \xi^a$, the variation of the dynamical fields equals their Lie derivatives: $\delta_\xi g_{ab} = \mathcal{L}_\xi g_{ab}$, $\delta_\xi A_e = \mathcal{L}_\xi A_e$. Consequently, the Noether current is found to be

$$\mathbf{J} = \epsilon_{aefgh} \left[2E_R^{abcd} \nabla_d (\nabla_b \xi_c + \nabla_c \xi_b) - E_A^{ai} \left(\xi^j \nabla_j A_i + (\nabla_i \xi^j) A_j \right) - \xi^a \mathcal{L} \right], \quad (C5)$$

where

$$E_R^{abcd} = \partial R / \partial R_{abcd} = \frac{1}{2} (g^{ac} g^{bd} - g^{ad} g^{bc}), \quad E_A^{ai} = \left(F^{ai} + \frac{k}{6} \epsilon^{abcdi} A_b F_{cd} \right). \quad (C6)$$

Moreover, a standard calculation [35] shows that

$$d\mathbf{J} = -\mathbf{E}\mathcal{L}_\xi\phi, \quad (C7)$$

which implies that \mathbf{J} is closed for all ξ^a once the equations of motion are satisfied. Thus, there exists a Noether charge 3-form \mathbf{Q} , locally constructed from ϕ and ξ^a , such that $\mathbf{J} = d\mathbf{Q}$ provided ϕ solve the equations of motion. Therefore, using the algorithm of [38] for obtaining \mathbf{Q} from \mathbf{J} , we find the Noether charge

$$\mathbf{Q} = -\epsilon_{abfgh} \left[\nabla^a \xi^b + \frac{1}{2} \left(F^{ab} + \frac{k}{6} \epsilon^{aijkb} A_i F_{jk} \right) A_c \xi^c \right]. \quad (C8)$$

With all the necessary ingredients in place, we can now obtain the fundamental identity following [32, 39]. To begin, considering the variation of (C4), and using (C1) and (C3), we have

$$\begin{aligned}
\delta\mathbf{J} &= \delta\Theta(\phi, \mathcal{L}_\xi\phi) - \xi \cdot \delta\mathbf{L}, \\
&= \delta\Theta(\phi, \mathcal{L}_\xi\phi) - \xi \cdot (\mathbf{E}\delta\phi + d\Theta), \\
&= \delta\Theta(\phi, \mathcal{L}_\xi\phi) - \mathcal{L}_\xi\Theta(\phi, \delta\phi) + d(\xi \cdot \Theta) - \xi \cdot \mathbf{E}\delta\phi, \\
&= -\omega(\phi, \delta\phi, \mathcal{L}_\xi\phi) + d(\xi \cdot \Theta) - \xi \cdot \mathbf{E}\delta\phi,
\end{aligned} \tag{C9}$$

where we have used the Cartan's magic formula $\mathcal{L}_\xi\mu = \xi \cdot d\mu + d(\xi \cdot \mu)$ for any differential form μ and vector field ξ^a in the third line. When ξ^a corresponds to a symmetry of the dynamical fields (*i.e.* $\mathcal{L}_\xi\phi = 0$), it follows that $\omega(\phi, \delta\phi, \mathcal{L}_\xi\phi) = 0$. If ϕ satisfies the equations of motion, then $\mathbf{E} = 0$ and $\mathbf{J} = d\mathbf{Q}$. Moreover, if $\delta\phi$ satisfies the linearized equations of motion, we find that $\delta\mathbf{J} = d\delta\mathbf{Q}$. Consequently, we obtain the fundamental identity

$$\omega(\phi, \delta\phi, \mathcal{L}_\xi\phi) = \delta\mathbf{J} - d(\xi \cdot \Theta) = d(\delta\mathbf{Q} - \xi \cdot \Theta) = 0. \tag{C10}$$

Let us choose ξ^a to be the time-like Killing vector $\xi^a = (\partial_t)^a = \delta_t^a$, which vanishes at the event horizon $r = r_h$. Let Σ be a $t = \text{const.}$ space-like hypersurface with the horizon $r = r_h$ serving as its interior boundary. Then, the boundary of this hypersurface, $\partial\Sigma$, including the boundaries of the non-compact x, y, z directions, is expressed as

$$\partial\Sigma = S_{r=r_h} \cup S_{r=0} \cup S_{x=L_x/2} \cup S_{x=-L_x/2} \cup S_{y=L_y/2} \cup S_{y=-L_y/2} \cup S_{z=L_z/2} \cup S_{z=-L_z/2}, \tag{C11}$$

where we have considered the regulation (L_x, L_y, L_z) along three spatial directions. Here $S_{r=r_h}$ denotes the boundary at $r = r_h$ and $S_{r=0}$ represents the boundary at $r = 0$. The remaining terms account for the boundaries at the edges of the non-compact directions in the x, y and z coordinates.

Assume that $n_{(t)a} \propto \partial_a t$ denotes the time-like unit normal vector of the hypersurface Σ and $n_{(i)a} \propto \partial_a x_i$ with $(x_i = x, y, z, r)$ represents the space-like unit normal vectors of S_i . The induced volume element and the projection tensor associated with each S_i can be constructed as follows:

$$\hat{\epsilon}_{(i)abc} = n_{(i)}^d n_{(t)}^e \epsilon_{edabc}, \quad P_{ab} = g_{ab} + n_{(t)a} n_{(t)b} - n_{(i)a} n_{(i)b}, \tag{C12}$$

with $\hat{\epsilon}_{(i)}^{abc} \hat{\epsilon}_{(i)abc} = 3!$ and $P_{ab} n_{(t)}^a = P_{ab} n_{(i)}^a = 0$. Note that the hypersurface S_i ($i = x, y, z, r$) is parameterized by the equations $x^a = x^a(y^i)$ with one of the spatial coordinates $y^i = (x, y, z, r)$ held fixed. Thus, the vectors $e_i^a = (\partial x^a)/(\partial y^i)$, which are tangent to curves in S_i , satisfy $e_i^a n_{(i)a} = 0$. We can project any tensor field onto the hypersurface S_i using e_i^a . For instance, the induced metric of the hypersurface is given by $h_{(i)ij} = g_{ab} e_i^a e_j^b = P_{ab} e_i^a e_j^b$.

Note that, the restriction of a bulk 3-form to a three dimensional hypersurface is proportional to the (induced) volume element of that hypersurface. For instance, we have $\mathbf{Q}|_i = Q_i \hat{\epsilon}_{(i)}$, where the scalar field Q_i is given by $Q_i = \mathbf{Q}_{abc} \hat{\epsilon}_{(i)}^{abc}/6$. Therefore, we can compute the restriction of $(\delta\mathbf{Q} - \xi \cdot \Theta)$ to each hypersurface S_i . Specifically, we have $(\delta\mathbf{Q} - \xi \cdot \Theta)|_i = (\delta Q - \xi \cdot \Theta)_i \epsilon_{(i)}$. Along each non-compact boundary, we find

$$\begin{aligned}
(\delta Q - \xi \cdot \Theta)_x &= x \left[\left(\frac{B e^{-\chi/2} h}{2r} + \frac{k}{6} (2A_t A'_z - A'_t A_z) \right) \delta B + \frac{kB}{6} (A_t \delta A_z)' \right], \\
(\delta Q - \xi \cdot \Theta)_y &= \frac{y}{x} (\delta Q - \xi \cdot \Theta)_x, \\
(\delta Q - \xi \cdot \Theta)_z &= 0.
\end{aligned} \tag{C13}$$

By integrating the fundamental identity (C10) over the hypersurface Σ , we obtain

$$\begin{aligned}
\delta H = 0 &= \int_\Sigma \omega = \int_\Sigma d(\delta\mathbf{Q} - \xi \cdot \Theta) = \int_{\partial\Sigma} (\delta\mathbf{Q} - \xi \cdot \Theta) \\
&= \int_{S_{r=r_h}} (\delta Q - \xi \cdot \Theta)_r \sqrt{h_{(r)}} dx dy dz - \int_{S_{r=\epsilon}} (\delta Q - \xi \cdot \Theta)_r \sqrt{h_{(r)}} dx dy dz \\
&\quad + \int_{S_{x=\frac{L_x}{2}}} (\delta Q - \xi \cdot \Theta)_x \sqrt{h_{(x)}} dy dz dr - \int_{S_{x=-\frac{L_x}{2}}} (\delta Q - \xi \cdot \Theta)_x \sqrt{h_{(x)}} dy dz dr \\
&\quad + \int_{S_{y=\frac{L_y}{2}}} (\delta Q - \xi \cdot \Theta)_y \sqrt{h_{(y)}} dx dz dr - \int_{S_{y=-\frac{L_y}{2}}} (\delta Q - \xi \cdot \Theta)_y \sqrt{h_{(y)}} dx dz dr \\
&= 3\delta f_4 - 8\delta h_4 - \frac{B}{2} \delta B + e^{-\frac{\chi_1}{2}} f_1 \left(\frac{\delta h_1}{r_h^3} - \frac{3h_1 \delta r_h}{r_h^4} \right) + \mu \delta \rho + B(\ln \epsilon) \delta B \\
&\quad + \int_\epsilon^{r_h} dr \left(\frac{B e^{-\chi/2} h}{r} + \frac{k}{2} (A'_z A_t - A_z A'_t) \right) \delta B,
\end{aligned} \tag{C14}$$

where we have used $\int_0^{r_h} (A_t \delta A_z)' dr = (A_t \delta A_z)|_0^{r_h} = 0$. Note that, the variation of the black hole entropy density is

$$\delta s = \delta \left(\frac{4\pi h_1}{r_h^3} \right) = 4\pi \left(\frac{\delta h_1}{r_h^3} - \frac{3h_1 \delta r_h}{r_h^4} \right). \quad (\text{C15})$$

and the variation of energy density ϵ in (A15) reads

$$\delta \epsilon = -3\delta f_4 + 8\delta h_4 + \frac{B}{2}\delta B. \quad (\text{C16})$$

By substituting the above equations into (C14), we find

$$0 = -\delta \epsilon + T\delta s + \mu\delta \rho + \left(\int_0^{r_h} dr \left[\frac{B}{r} \left(e^{-\chi/2} h - 1 \right) + \frac{k}{2} (A'_z A_t - A_z A'_t) \right] + B \ln r_h \right) \delta B. \quad (\text{C17})$$

Thus, we arrive at the first law of black hole thermodynamics

$$\delta \epsilon = T\delta s + \mu\delta \rho - M_B \delta B, \quad (\text{C18})$$

as shown in the main text.

On the other hand, using the Noether current \mathbf{J} , we can also determine the expected free energy \tilde{w} of the system. The conditions that ϕ solve the background equations of motion and ξ is a Killing vector (*i.e.* $\mathcal{L}_\xi \phi = 0$) imply

$$d\mathbf{Q} = \mathbf{J} = \Theta(\phi, \mathcal{L}_\xi \phi) - \xi \cdot \mathbf{L} = -\xi \cdot \mathbf{L}. \quad (\text{C19})$$

Integrating the identity $d\mathbf{Q} = -\xi \cdot \mathbf{L}$ over the hypersurface Σ and applying the Stokes's theorem, we find

$$\int_{\partial\Sigma} \mathbf{Q} = - \int_{\Sigma} \xi \cdot \mathbf{L}. \quad (\text{C20})$$

Consequently, we have

$$\left[Q_r(r_h) - Q_r(\epsilon) \right] \sqrt{h(r)} V = V \int_{\epsilon}^{r_h} \sqrt{-g} \mathcal{L} dr - V \int_{\epsilon}^{r_h} \frac{kB}{3} A_t A'_z dr, \quad (\text{C21})$$

where $Q_r = \mathbf{Q}_{abc} \hat{\epsilon}_{(r)}^{abc} / 6$ and $V = \int dx dy dz = L_x L_y L_z$. Adding the boundary term S_{bdy} of (A5) to both sides of the above equation and taking $\epsilon \rightarrow 0$, we obtain

$$\epsilon - Ts - \mu\rho = \tilde{w}. \quad (\text{C22})$$

Finally, using the expressions (A11) for the bulk on-shell action, we obtain from (C20) and (A15) that

$$\mathcal{P}_{\parallel} = \mathcal{P}_{\perp} + M_B B. \quad (\text{C23})$$

D: Smarr relation

From the scaling symmetry (A2), one can find that, under the scale transformation $r \rightarrow \lambda \hat{r}$ with λ a positive constant, the relevant physical quantities scale as

$$s = \lambda^3 \hat{s}, \quad \rho = \lambda^3 \hat{\rho}, \quad B = \lambda^2 \hat{B}, \quad (\text{D1})$$

while the energy density ϵ acquires an anomalous scaling transformation

$$\epsilon = \lambda^4 \hat{\epsilon} - \frac{\hat{B}^2}{2} \lambda^4 \ln \lambda. \quad (\text{D2})$$

Thus, when we express ϵ as a function of s, ρ and B , we have

$$\epsilon(\lambda^3 \hat{s}, \lambda^3 \hat{\rho}, \lambda^2 \hat{B}) = \lambda^4 \hat{\epsilon}(\hat{s}, \hat{\rho}, \hat{B}) - \frac{\hat{B}^2}{2} \lambda^4 \ln \lambda. \quad (\text{D3})$$

Taking derivative of both side of this equation with respect to λ and then setting λ to 1, we arrive at

$$\left(\frac{\partial\epsilon}{\partial s}\right)_{\rho,B} (3s) + \left(\frac{\partial\epsilon}{\partial\rho}\right)_{s,B} (3\rho) + \left(\frac{\partial\epsilon}{\partial B}\right)_{s,\rho} (2B) = 4\epsilon - \frac{B^2}{2}. \quad (\text{D4})$$

Taking advantage of the first law of thermodynamics (C18), we can derive the (generalized) Smarr relation

$$-4\epsilon + 3(Ts + \mu\rho) - 2M_B B = -\frac{B^2}{2}. \quad (\text{D5})$$

The right hand side, which is non-zero when $B \neq 0$, indicates a derivation from the standard Smarr formula due to the chiral anomaly. According to (C22), this Smarr relation effectively expresses the trace anomaly of the boundary stress-energy tensor *i.e.* $T^\mu_\mu = -B^2/2$.

E: Magnetic susceptibility from hydrodynamics

In this section, we provide a brief overview of the hydrodynamic description of the 3+1 dimensional chiral charged fluid in the presence of a strong external magnetic field, following the works of [27–29]. Relativistic hydrodynamics is crucial for describing finite-temperature interacting relativistic field theories at large distances and long time scales. This framework has found applications across various fields, including nuclear physics, condensed matter physics, astrophysics, and cosmology. The hydrodynamic variables include the local temperature $T(x)$, the local chemical potential $\mu(x)$ and the local fluid velocity $u^\mu(x)$ (normalized such that $u^\mu u_\mu = -1$). The stress-energy tensor $T^{\mu\nu}$ and current J^μ are expressed through constitutive relations in terms of hydrodynamic variables and their derivatives.

In the presence of external background field and anomalous effects, the hydrodynamic equations governing the stress-energy tensor $T^{\mu\nu}$ and the axial current J^μ are given by

$$\nabla_\mu T^{\mu\nu} = F^{\nu\lambda} J_\lambda, \quad (\text{E1})$$

$$\nabla_\mu J^\mu = \frac{k}{8} E^\mu B_\mu, \quad (\text{E2})$$

where $E^\mu = F^{\mu\nu} u_\nu$ and $B^\mu = \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} u_\nu F_{\alpha\beta}$ represent the electric and magnetic fields in (3+1) dimensions, respectively [27]. To leading order in the derivative expansion, the constitutive relations are expressed as

$$\langle T_{EFT}^{\mu\nu} \rangle = \epsilon_0 u^\mu u^\nu + P_0 \Delta^{\mu\nu} + q^\mu u^\nu + q^\nu u^\mu + M^{\mu\alpha} g_{\alpha\beta} F^{\beta\nu} + u^\mu u^\alpha (M_{\alpha\beta} F^{\beta\nu} - F_{\alpha\beta} M^{\beta\nu}) + \mathcal{O}(\partial), \quad (\text{E3})$$

$$\langle J_{EFT}^\mu \rangle = n_0 u^\mu + \xi_B B^\mu + \mathcal{O}(\partial), \quad (\text{E4})$$

where ϵ_0 is the energy density, P_0 is the pressure, and $\Delta^{\mu\nu} = g^{\mu\nu} + u^\mu u^\nu$. The terms $\xi_B B^\mu$ and $q^\mu = \xi_V B^\mu$ represent chiral transport coefficients arising from the chiral anomaly, while the polarization tensor $M^{\mu\nu} = \chi_{BB} \epsilon^{\mu\nu\alpha\beta} B_\alpha u_\beta$ is defined as the variation of the generating functional with respect to the field strength [27].

In thermal equilibrium, where we choose $u^\mu = (u^t, u^x, u^y, u^z) = (1, 0, 0, 0)$ and assume $B^\mu \propto \vec{z}$, the energy-momentum tensor and the gauge current are given by

$$\langle T_{EFT}^{\mu\nu} \rangle = \begin{pmatrix} \epsilon_0 & 0 & 0 & \xi_V^{(0)} B \\ 0 & P_0 - \chi_B B^2 & 0 & 0 \\ 0 & 0 & P_0 - \chi_B B^2 & 0 \\ \xi_V^{(0)} B & 0 & 0 & P_0 \end{pmatrix} + \mathcal{O}(\partial), \quad (\text{E5})$$

$$\langle J_{EFT}^\mu \rangle = (n_0, 0, 0, \xi_B^{(0)} B) + \mathcal{O}(\partial), \quad (\text{E6})$$

where the subscript “0” and superscript “(0)” denote that these quantities are evaluated in thermal equilibrium. Note that the thermodynamic relation $\epsilon_0 = -P_0 + Ts + \mu\rho$ has been employed to derive (E5), which holds under the identification $P_0 = -\tilde{w}$.

Finally, by comparing the results for the stress-energy tensor (E5) with the gravitational computations (A15), we obtain the expression for the magnetic susceptibility χ_B .

$$\chi_B B^2 = \langle T^{zz} \rangle - \langle T^{xx} \rangle = \frac{B^2}{4} + 8h_4. \quad (\text{E7})$$

Thus, the magnetic susceptibility from the hydrodynamic description reads

$$\chi_B = \frac{1}{4} + \frac{8h_4}{B^2}, \quad (\text{E8})$$

which is what we have shown in the main text.