EXCESS DECAY FOR QUASILINEAR EQUATIONS IN THE HEISENBERG GROUP AND CONSEQUENCES

ARKA MALLICK AND SWARNENDU SIL

ABSTRACT. We study regularity results for the solutions of quasilinear subelliptic p-Laplace type equation in Heisenberg groups. We prove somewhat surprising excess decay estimates for the constant coefficient homogeneous equation. Excess decay estimates, while well known in the Euclidean case, due to the celebrated works of Uraltseva and Uhlenbeck, was not known in the setting of Heisenberg groups until now and this lack of excess decay estimate is often attributed to the noncommutativity of the horizontal vector fields. Our results show that, in spite of the this noncommutative feature, excess decay estimates analogous to the Euclidean case hold.

To illustrate the potency of our excess decay estimates, we prove two results. First is a Hölder continuity result that extends the presently known results to the full range $1 < p < \infty$. The second is a sharp borderline continuity result for the horizontal gradient of the solution. More precisely, we show that if $u \in HW^{1,p}_{loc}$ satisfies

div_{III} $(a(x)|\mathfrak{X}u|^{p-2}\mathfrak{X}u) \in L^{(Q,1)}_{\text{loc}}$

in a domain of \mathbb{H}_n , where \mathbb{H}_n is the Heisenberg group with homogeneous dimension $Q = 2n + 2$, for $1 \lt p \lt \infty$, with uniformly positive, bounded, Dini continuous scalar function a , then $\mathfrak{X}u$ is continuous. This generalizes the classical result by Stein and the work of Kuusi-Mingione for linear and quasilinear equations, respectively, in the Euclidean case and Folland-Stein for the linear case in the Heisenberg group setting. This result is new even when either a is constant or when the equation is homogeneous.

1. Introduction and main results

Let $n \geq 1$ be an integer and let \mathbb{H}_n denote the *n*-th Heisenberg group over the reals. Let $Q := 2n + 2$ stands for the homogeneous dimension of \mathbb{H}_n . Heisenberg groups \mathbb{H}_n are important objects in mathematics and mathematical physics alike. From the point of view of mathematics, they appear in diverse areas such as representation theory, harmonic analysis, complex geometry, subRiemannian geometry and metric geometry. They are the simplest nonAbelian examples of stratified nilpotent Lie groups, which are homogeneous groups and also metric groups. They are also the simplest examples of Carnot groups and Carnot-Caratheodory spaces, apart from Euclidean spaces. They are CR manifolds, which also serve as models for boundaries of domains in \mathbb{C}^{n+1} . The study of subLaplacians on Heisenberg

Date: October 2, 2024.

²⁰²⁰ Mathematics Subject Classification. Primary 35B65, 35H20, 35J62; Secondary 35H10, 35J70, 35J75.

Key words and phrases. Heisenberg group, p-Laplacian, quasilinear subelliptic equations, excess decay estimate, nonlinear Stein theorem, Lorentz space, continuity of horizontal gradient.

S. Sil's research is partially supported by SERB MATRICS MTR/2023/000885.

2 A. MALLICK AND S. SIL

groups are also interesting from the point of view PDEs, as these are differential operators defined by the so-called Hörmander vector fields and are hypoelliptic, but not elliptic. Nonetheless, the study of these subLaplacians and the related singular and fractional integrals is well developed thanks to several seminal contributions, e.g. [\[27\]](#page-63-0), [\[23\]](#page-63-1), [\[28\]](#page-63-2), [\[16\]](#page-63-3), [\[17\]](#page-63-4), [\[14\]](#page-63-5), [\[18\]](#page-63-6), [\[41\]](#page-64-0) and many others (see the book [\[1\]](#page-62-0) and the references therein). Subelliptic p -Laplace type equations has also been studied intensively for quite some time, see the book [\[40\]](#page-64-1), also [\[4\]](#page-62-1) and the recent survey [\[3\]](#page-62-2). Our present contribution seeks to add significant momentum to the understanding of p-subLaplacians on Heisenberg groups.

The main concern of the present article is to remove perhaps the single most significant roadblock to the progress of regularity theory for p -Laplace type equations in Heisenberg groups - namely the absence of an excess decay estimate.

Excess decay in the Euclidean case: If $u \in W_{loc}^{1,p}(\Omega)$ is a local weak solution to the homogeneous p-Laplace equation

$$
\operatorname{div}\left(\left|\nabla u\right|^{p-2} \nabla u \right) = 0 \qquad \text{in } \Omega,
$$

where $\Omega \subset \mathbb{R}^n$ is a domain in the *n*-dimensional Euclidean space, we have the celebrated excess decay estimate (see [\[21\]](#page-63-7), [\[22\]](#page-63-8), also [\[47\]](#page-64-2), [\[46\]](#page-64-3) and [\[32\]](#page-64-4))

$$
(1) \qquad \int_{B_{\rho}} \left| V\left(\nabla u\right) - \left(V\left(\nabla u\right)\right)_{\rho} \right|^2 \leq C \left(\frac{\rho}{R}\right)^{2\beta} \int_{B_R} \left| V\left(\nabla u\right) - \left(V\left(\nabla u\right)\right)_R \right|^2,
$$

for any balls $B_\rho \subset B_R \subset \Omega$, for some $\beta \in (0,1)$, where $V(\xi) := |\xi|^{\frac{p-2}{2}} \xi$ and $C \equiv C(n, p) > 0$ is a constant. Later, DiBenedetto-Manfredi [\[8\]](#page-63-9) obtained a different type of excess decay estimate, in terms of ∇u itself. They showed the excess decay estimate

(2)
$$
\int_{B_{\rho}} \left| \nabla u - (\nabla u)_{\rho} \right|^2 \leq C \left(\frac{\rho}{R} \right)^{2\alpha} \int_{B_R} \left| \nabla u - (\nabla u)_R \right|^2,
$$

again for any balls $B_\rho \subset B_R \subset \Omega$, for some $\alpha \in (0,1)$ and for some constant $C \equiv C(n, p) > 0$. The Hölder continuity of ∇u follows immediately from either of these excess decay estimates. The key to proving (1) is a type of reverse Hölder inequality (see [\[21\]](#page-63-7), [\[22\]](#page-63-8)), which follows from the relevant Caccioppoli inequality in these case. However, the proof of [\(2\)](#page-1-1) goes via an iteration argument, which is adapted in $[8]$ from its parabolic counterparts in $[6]$, $[7]$. A Caccioppoli type inequality is also instrumental in this case as well. These excess decay estimates have also proved pivotal for the subsequent development of the regularity theory for quasilinear equations and systems with coefficients and non-zero right hand sides (see e.g. [\[8\]](#page-63-9), [\[10\]](#page-63-12) [\[11\]](#page-63-13), [\[12\]](#page-63-14), [\[30\]](#page-63-15), [\[31\]](#page-63-16) etc).

Setting of Heisenberg groups : Coming to the setting of Heisenberg groups, if one considers the analogous equation

$$
\operatorname{div}_{\mathbb{H}}\left(|\mathfrak{X}u|^{p-2}\mathfrak{X}u\right) = 0 \qquad \qquad \text{in } \Omega,
$$

where div_H denotes the horizontal divergence, $\mathfrak{X}u$ denotes the horizontal gradient and now $\Omega \subset \mathbb{H}_n$ is a domain in the Heisenberg group \mathbb{H}_n , the main difference from the Euclidean case is that although $\mathbb{H}_n \simeq \mathbb{R}^{2n+1}$ as sets, only the horizontal derivatives $\mathfrak{X}_1, \ldots, \mathfrak{X}_{2n}$ appears in the equation, not the full Euclidean gradient $(X_1u, \ldots, X_{2n}u, Tu)$. Perhaps the even more significant difference is that unlike the Euclidean derivatives, the horizontal derivatives do not commute and the vertical derivative Tu is the only nontrivial commutator

(3)
$$
Tu = X_i X_{n+i} u - X_{n+i} X_i u \quad \text{for all } 1 \le i \le n.
$$

Practically all the difficulties of the Heisenberg group setting, compared to the Euclidean case, can be traced back to the difficulty of estimating the vertical derivative Tu , which appears as soon as one tries to write down the equations satisfied by the horizontal first derivatives X_iu . These equations (see [\(52\)](#page-17-0) and [\(53\)](#page-17-1)) are still subelliptic, but they are inhomogeneous, whereas one obtains only homogeneous equations in the Euclidean case. Reverse Hölder type inequalities, as mentioned above, follows from Caccippoli inequalities for homogeneous elliptic equations, but for inhomogeneous equations, they are considerably harder to obtain and do not, in general, hold. In this case, this task is even more problematic due to the fact that the right hand side contains terms involving the vertical derivative Tu and as the expression for Tu in [\(3\)](#page-2-0) shows, morally, Tu is more like a second order derivative than a first order derivative.

Due to these difficulties, the progress in the Heisenberg group setting was much slower. First result was due to Capogna [\[2\]](#page-62-3), who proved a regularity result which was enough to conclude C^{∞} regularity in the nondegenerate case. After this, next significant breakthrough was achieved in Domokos [\[9\]](#page-63-17), who proved a higher integrability estimate for Tu when $1 < p < 4$. For a long time, all available regularity results were restricted to a small range of p , since these relied crucially on the result by Domokos ([\[36\]](#page-64-5), [\[35\]](#page-64-6), [\[37\]](#page-64-7) etc).

Zhong's mixed Caccioppoli and oscillation control by energy : The first result to break this barrier is due to Zhong [\[50\]](#page-64-8). In this remarkable work, Zhong proved a number of mixed Caccioppoli inequalities and used them to prove the the Hölder continuity of $\mathfrak{X}u$ for $p \geq 2$. Later, this program was carried to completion in Mukherjee-Zhong [\[39\]](#page-64-9), where Hölder continuity of $\mathfrak{X}u$ was proved also for the range $1 < p < 2$, using the ideas of Tolksdorf [\[45\]](#page-64-10).

However, both these results established Hölder continuity by deriving oscillation estimates by 'energy'. More precisely, the estimate is

(4)
$$
|\mathfrak{X}u(x)-\mathfrak{X}u(y)| \leq c\left(\frac{\rho}{R}\right)^{\alpha_1}\left(\int_{B_R}|\mathfrak{X}u|^p\ dx\right)^{\frac{1}{p}}, \text{ for all } x, y \in B_\rho,
$$

for any metric balls $B_\rho \subset B_{R/2} \subset B_R \subset\subset \Omega$. This is considerably weaker than estimates of the form [\(1\)](#page-1-0) or [\(2\)](#page-1-1), as the energy in B_R can be much larger than the average osicillation in B_R . For proving Hölder continuity of \mathfrak{X}_u , [\(4\)](#page-2-1) is sufficient, but this is not quite strong enough so that one can derive other results using the same perturbation techniques as in the Euclidean case. Indeed, the only perturbation type results that is known so far is the following: If $2 - 1/Q < p < \infty$ and u is a weak solution to the equation

$$
\operatorname{div}_{\mathbb{H}}\left(a(x)\left|\mathfrak{X}u\right|^{p-2}\mathfrak{X}u\right) = f \qquad \qquad \text{in } \Omega,
$$

where a is uniformly positive, bounded, Hölder continuous and f is Hölder continuous, then $\mathfrak{X}u$ is Hölder continuous. This was proved in Mukherjee-Sire[\[38\]](#page-64-11) for the range $p \geq 2$ and in Yu [\[49\]](#page-64-12) for the range $2 - 1/Q < p < 2$, using Wolff potential bounds. In [\[38\]](#page-64-11), the authors noted (see Proposition 3.1 in [\[38\]](#page-64-11)) that the mixed Caccioppoli inequalities of Zhong only yields the significantly weaker estimate

(5)
$$
\int_{B_{\rho}} \left| \nabla u - (\nabla u)_{\rho} \right| dx \leq C \left(\frac{\rho}{R} \right)^{\alpha} \left[\int_{B_{R}} \left| \nabla u - (\nabla u)_{R} \right| dx + \chi R^{\alpha} \right],
$$

where

$$
\chi = \frac{1}{R_0^{\alpha}} \sup_{B_{R_0}} |\mathfrak{X} u|,
$$

for any metric balls $B_\rho \subset B_R \subset\subset B_{R_0} \subset \Omega$. The authors in [\[38\]](#page-64-11) also asserted, both in the Introduction and in Remark 3.2, that the appearance of the extra term with $\chi \neq 0$ is unavoidable and seemed to suggest this as a manifestation of the appearance of the terms involving Tu in the equations for X_iu , which are evidently absent in the Euclidean case.

An excess decay estimate in Heisenberg groups : Obviously, if there were a genuine lack of Euclidean-like excess decay estimates, it is not at all unreasonable to ascribe that lack to the noncommutative nature of the vector fields. Our main contribution in the present work, somewhat surprisingly in this sense, asserts that Euclidean type excess decay estimates indeed hold in the Heisenberg group setting as well, the noncommutativity of the vector fields notwithstanding.

Theorem 1. Suppose u solves the constant coefficient homogeneous p-Laplace equation in a domain $\Omega \subset \mathbb{H}_n$, with $1 < p < \infty$,

$$
\operatorname{div}_{\mathbb{H}}\left(|\mathfrak{X}u|^{p-2}\mathfrak{X}u\right) = 0 \qquad \qquad \text{in } \Omega.
$$

Let $1 \le q \le 2$ be a real number. Then there exists constants $\alpha \in (0,1)$, and $C_0 \ge 1$, both depending only on n and p, such that for any $x_0 \in \Omega$ and any $0 < \rho < R$ such that $B_R(x_0) \subset \subset \Omega$, we have

$$
(6) \qquad \left(\frac{f}{f_{B_{\rho}(x_0)}}\left|\mathfrak{X} u-(\mathfrak{X} u)_{x_0,\rho}\right|^q\right)^{\frac{1}{q}} \leq C_0 \left(\frac{\rho}{R}\right)^{\alpha} \left(\frac{f}{f_{B_R(x_0)}}\left|\mathfrak{X} u-(\mathfrak{X} u)_{x_0,R}\right|^q\right)^{\frac{1}{q}}.
$$

We also have a excess decay estimate for $V(\mathfrak{X}u)$.

Theorem 2. Suppose u solves the constant coefficient homogeneous p-Laplace equation in a domain $\Omega \subset \mathbb{H}_n$, with $1 < p < \infty$,

$$
\operatorname{div}_{\mathbb{H}}\left(\left|\mathfrak{X} u\right|^{p-2} \mathfrak{X} u\right) = 0 \qquad \qquad in \ \Omega.
$$

Let $1 \le q \le 2$ be a real number. There exists constants $\beta \in (0,1)$ and $C_{0,V} \ge 1$, both depending only on n and p, such that for any $x_0 \in \Omega$ and any $0 < \rho < R$ such that $B_R(x_0) \subset \subset \Omega$, we have

$$
(7) \quad \left(\int_{B_{\rho}(x_0)} \left| V(\mathfrak{X} u) - (V(\mathfrak{X} u))_{x_0,\rho} \right|^q dx \right)^{\frac{1}{q}} \leq C_{0,V} \left(\frac{\rho}{R}\right)^{\beta} \left(\int_{B_R(x_0)} \left| V(\mathfrak{X} u) - (V(\mathfrak{X} u))_{x_0,R} \right|^q dx \right)^{\frac{1}{q}}.
$$

The balls above are understood to be metric balls (see Section [2.1](#page-8-0) and [2.2](#page-8-1)) and for any \mathbb{R}^{2n} -valued function g, $(g)_{x_0,R}$ denotes the integral average of g over the metric ball $B_R(x_0)$ (see Notation [8\)](#page-9-0). The two theorems above are exact analogues of their Euclidean counterpart. We would also like to point out that while both results are known in the Euclidean setting, the passage from the exponent 2 in [\(2\)](#page-1-1) to any exponent $1 \le q \le 2$ was achieved later, by an argument due to Lieberman [\[33\]](#page-64-13). For (1) , analogous passage can be achived by reverse Hölder inequalities established in [\[21\]](#page-63-7) and [\[22\]](#page-63-8). We also note that the proof of [\(1\)](#page-1-0) for $1 < p < 2$ follows via a duality argument in [\[22\]](#page-63-8). Our argument yields the full result for the full range of p in one go, without needing additional arguments and thus yields a new proof for the Euclidean case as well, as long as an oscillation to energy bound like [\(4\)](#page-2-1) is available (for eaxmple, Tolksdorf [\[45\]](#page-64-10) proved such estimates independently of excess decay estimates). Note also that the inequality in Theorem [1](#page-3-0) for $q = 1$ is exactly [\(5\)](#page-3-1) with $\chi = 0$.

As we mentioned above, in the Euclidean case, Hölder continuity of gradient is proved using the excess decay estimates. The main idea behind our excess decay estimates is the converse question: Suppose we already know that the solutions are Hölder continuous and we have an oscillation estimate by energy. When can we derive the excess decay estimate?

The main insight of our estimate is that this is possible as soon as we have a linear-like Caccioppoli inequality in the nondegenerate regime. The form of the Caccioppoli inequalities in the general case does not matter at all. While the Caccioppoli inequalities derived by Zhong in the general case is of mixed type, in the nondegenerate regime, we indeed have a Caccioppoli inequality which looks exactly like the one for a linear homogeneous equation (see [\(51\)](#page-17-2)). This itself is somewhat surprising, as the equations [\(52\)](#page-17-0) and [\(53\)](#page-17-1) are not homogeneous. These Caccioppoli inequalities imply reverse Hölder inequalities in the standard manner (see [\(65\)](#page-21-0)), but of course, again for the nondegenerate regime only.

Our iteration lemma (Lemma [40](#page-28-0)) is similar in spirit to Lemma 4.4 and Lemma 4.5 in [\[6\]](#page-63-10) (see also [\[7\]](#page-63-11)). However, unlike in their situations, we do not have a suitable Caccioppoli inequality for the equation in general. Thus our lemma is weaker as we always need the nondegeneracy condition $\lambda/4B \leq |\mathfrak{X}u| \leq A\lambda$. Due to this extra hypotheses, our proofs of Theorems [1](#page-3-0) and [2](#page-3-2) diverge significantly from the proof of [\(2\)](#page-1-1) given in [\[8\]](#page-63-9). Our result is sufficient to derive excess decay if oscillation estimate by energy is known beforehand. But for the same reasons, it cannot furnish an independent proof of the Hölder continuity result, while their's can. We would also like to point out that the idea of treating the nondegenerate regime and the degenerate regime separately is also there in DiBenedetto-Manfredi [\[8\]](#page-63-9) (also earlier in Evans [\[13\]](#page-63-18)). However, in the degenerate regime, one usually obtains a sup decay of the form

(8)
$$
\sup_{B_{\sigma R}} |\nabla u| \leq \eta \sup_{B_R} |\nabla u|
$$

for some constants $\eta, \sigma \in (0, 1)$. By iterating this, one can of course conclude the existence of an integer $i \geq 1$ such that one has

$$
\sup_{B_{\sigma^i R}} |\nabla u| \le \eta^i \sup_{B_R} |\nabla u| < \int_{B_R} |\nabla u - (\nabla u)_R| \, dx.
$$

However, apriori, this integer *i* depends on $f_{B_R} |\nabla u - (\nabla u)_R| dx$. The novelty in our proof is that one can actually reduce to the case where it is possible to determine an integer i_a , apriori in terms of n and p alone such that this happens (compare Claim [45](#page-35-0)).

Consequences of excess decay estimates : Our excess decay estimates make the problem in the Heisenberg setting amenable to Euclidean perturbation techniques. As an illustration, we have the following result.

Theorem 3 (Hölder estimates). Let $1 < p < \infty$ and $0 < \mu_1, \mu_2 < 1$ be real numbers. Let $\Omega \subset \mathbb{H}_n$ be open. Suppose

- (i) $a: \Omega \to [\nu, L]$ is $C_{loc}^{0,\mu_1}(\Omega)$, where $0 < \nu < L < \infty$ and
- (ii) $F \in C_{loc}^{0,\mu_2} (\Omega; \mathbb{R}^{2n})$.

Let $u \in HW_{loc}^{1,p}(\Omega)$ be a local weak solution to

(9)
$$
\operatorname{div}_{\mathbb{H}}\left(a(x)|\mathfrak{X}u|^{p-2}\mathfrak{X}u\right) = \operatorname{div}_{\mathbb{H}} F \qquad \text{in } \Omega.
$$

Suppose

$$
\mu := \min \left\{ \mu_1, \mu_2 \right\} < \min \left\{ \beta, \alpha \right\},\
$$

where $\alpha, \beta \in (0,1)$ are the exponents given by Theorems [1](#page-3-0) and [\(2\)](#page-3-2) respectively. Then we have

$$
\mathfrak{X}u \in C_{loc}^{0,\min\left\{\mu,\frac{\mu}{p-1}\right\}}\left(\Omega;\mathbb{R}^{2n}\right).
$$

As an immediate consequence, we have the following, which generalizes the re-sults in [\[38\]](#page-64-11) and [\[49\]](#page-64-12) to the full range $1 < p < \infty$.

Corollary 4. Let $1 < p < \infty$ and let $\Omega \subset \mathbb{H}_n$ be open. Suppose

(i)
$$
a: \Omega \to [\nu, L]
$$
 is $C_{loc}^{0,\mu_1}(\Omega)$, where $0 < \nu < L < \infty$, $0 < \mu_1 < 1$ and (ii) $f \in L_{loc}^q(\Omega)$ with $q > Q$.

Let $u \in HW_{loc}^{1,p}(\Omega)$ be a local weak solution to

(10)
$$
\operatorname{div}_{\mathbb{H}}\left(a(x)|\mathfrak{X}u|^{p-2}\mathfrak{X}u\right) = f \qquad \qquad in \ \Omega.
$$

Suppose

$$
\mu:=\min\left\{\mu_1,1-\frac{Q}{q}\right\}<\min\left\{\beta,\alpha\right\},
$$

where $\alpha, \beta \in (0,1)$ are the exponents given by Theorems [1](#page-3-0) and [\(2\)](#page-3-2) respectively. Then we have

$$
\mathfrak{X}u \in C_{loc}^{0,\min\left\{\mu,\frac{\mu}{p-1}\right\}}\left(\Omega;\mathbb{R}^{2n}\right).
$$

The Hölder spaces above are understood to be Folland-Stein classes (see Section [2.2](#page-8-1)). We emphasize that our proof is via the classical Camapanto method and does neither require any perturbation argument nor any potential estimates.

Subelliptic nonlinear Stein theorem : The efficacy of our excess decay estimate goes beyond easy Campanato style perturbation arguments. In [\[43\]](#page-64-14), Stein proved the proved the borderline Sobolev embedding result which states that for $n \geq 2, u \in L^1(\mathbb{R}^n)$ and $\nabla u \in L^{(n,1)}(\mathbb{R}^n;\mathbb{R}^n)$ implies u is continuous. Coupled with standard Calderon-Zygmund estimates, which extend to Lorentz spaces, this implies $u \in C^1(\mathbb{R}^n)$ if $\Delta u \in L^{(n,1)}(\mathbb{R}^n)$. The search for a nonlinear generalization of this result culminated in Kuusi-Mingione [\[31\]](#page-63-16), where the authors proved the following general quasilinear vectorial version

(11)
$$
\operatorname{div} (a(x) |\nabla u|^{p-2} \nabla u) \in L^{(n,1)}(\mathbb{R}^n; \mathbb{R}^N) \qquad \Longrightarrow \qquad u \in C^1(\mathbb{R}^n; \mathbb{R}^N),
$$

where a is a uniformly positive Dini continuous function and $1 < p < \infty$. For the case of Heisenberg groups, even the linear result that

 $\text{div}_{\mathbb{H}}(\mathfrak{X}u)) \in L^{(Q,1)} \implies \mathfrak{X}u \text{ is continuous,}$

is difficult to find in the literature. However, this can be shown combining the ideas from [\[43\]](#page-64-14) with the ones from Folland-Stein [\[18\]](#page-63-6), Folland-Stein [\[17\]](#page-63-4) and Folland [\[14\]](#page-63-5), see Section [2.5](#page-13-0) for details. We prove the following nonlinear Stein type theorem for Hesienberg groups.

Theorem 5 (Subelliptic Nonlinear Stein theorem). Let $1 < p < \infty$ and let $\Omega \subset \mathbb{H}_n$ be open. Suppose that

- (i) $f \in L_{loc}^{(Q,1)}(\Omega)$, where $Q = 2n + 2$ is the homogeneous dimension of \mathbb{H}_n ,
- (ii) $a : \Omega \to [\gamma, L]$, where $0 < \gamma < L < \infty$, is Dini continuous.
- Let $u \in HW_{loc}^{1,p}(\Omega)$ be a local weak solution to the equation

(12)
$$
\operatorname{div}_{\mathbb{H}}(a(x)|\mathfrak{X}u|^{p-2}\mathfrak{X}u)) = f \qquad \qquad in \ \Omega.
$$

Then $\mathfrak{X}u$ is continuous in Ω .

The conditions on the coefficient is sharp already in the Euclidean case, as shown by Jin et al [\[26\]](#page-63-19). Cianchi [\[5\]](#page-62-4) showed the Lorentz space $L^{(n,1)}$ is also sharp in the Euclidean case. The homogeneous dimension Q is the natural replacement of n in the setting of Heisenberg groups. As far as we are aware, this result is new even when $f \equiv 0$ or when a is a constant function. This result is nontrivial already in the Euclidean case, where it is the end product of a series of works, starting with Duzaar-Mingione [\[11\]](#page-63-13) and culminating in Kuusi-Mingione [\[29\]](#page-63-20), Kuusi-Mingione [\[30\]](#page-63-15), Kuusi-Mingione [\[31\]](#page-63-16). In the beautiful work [\[31\]](#page-63-16), the authors derived suitable form of the comparison estimates in the nondegenerate regime. This, coupled with an exit time argument and the Euclidean excess decay, this is sufficient for the result.

The delicate comparison estimates in [\[31\]](#page-63-16), however, extends immediately to the Heisenberg group setting, as they are ultimately a consequence of the structure of the p-Laplace equation. The missing pieces of the puzzle that remained are the follwing:

- The excess decay estimate which were not available, until now that is, in this setting.
- An intermediate continuity results for the homogeneous p-Laplace equation with Dini continuous coefficients (see Theorem 2 in $|31|$). This is a crucial intermediate step in their proof. In the Euclidean case, this result was obtained in the parabolic case in [\[29\]](#page-63-20).

We show, that the excess decay estimate is really the only missing key ingredient. With our excess decay estimates at hand, we prove the following.

Theorem 6. Let $1 < p < \infty$ and let $\Omega \subset \mathbb{H}_n$ be open. Let $w \in HW^{1,p}_{loc}(\Omega)$ be a local weak solution of

(13)
$$
\operatorname{div}_{\mathbb{H}}(a(x)|\mathfrak{X}w|^{p-2}\mathfrak{X}w)) = 0 \qquad \text{in } \Omega,
$$

where $a : \Omega \to [\gamma, L]$, where $0 < \gamma < L < \infty$, is Dini continuous. Then $\mathfrak{X}w$ is continuous in Ω . Moreover, if $\omega(\cdot)$ denotes the modulus of continuity associated with the function a, then

(i) there exists a constant $C_2 \equiv C_2(n, p, \gamma, L, \omega(\cdot)) \geq 1$ and a positive radius $R_1 = R_1(n, p, \gamma, L, \omega(\cdot)) > 0$ such that if $R \leq R_1$, then the estimate

(14)
$$
\sup_{B(x,R/2)} |\mathfrak{X}w| \leq C_2 \int_{B(x,R)} |\mathfrak{X}w|,
$$

holds whenever $B(x, R) \subset \Omega$. If $a(\cdot)$ is a constant function, the estimate holds without any restriction on R.

(ii) Assume that the inequality

(15)
$$
\sup_{B(x,R/2)} |\mathfrak{X}w| \leq A\lambda
$$

hold for some $A \geq 1$ and $\lambda > 0$. Then for any $\delta \in (0,1)$ there exists a positive constant $\sigma_1 \equiv \sigma_1(n, p, \gamma, L, \omega(\cdot), A, \delta) \in (0, \frac{1}{4})$ such that for every $\sigma \leq \sigma_1$, we have,

(16)
$$
\sup_{x,y\in B(x,\sigma R)} |\mathfrak{X}w(x)-\mathfrak{X}w(y)| \leq \delta\lambda.
$$

This now takes care of all the missing pieces and the arguments of Kuusi-Mingione [\[31\]](#page-63-16) can be adapted, essentially verbatim with the obvious notational change, to our setting to establish Theorem [5.](#page-6-0)

Future directions : We fully expect that with the present contribution at hand, a full-blown 'nonlinear potential theory' or a 'nonlinear Calderon-Zygmund theory', as it is usually called in the Euclidean case, for the Heisenberg group case, is just around the corner now. In most cases, the lack of an excess decay estimate was the only real missing ingredient till date. Consequently, in our opinion, our estimates would enable most, if not all, of the gradient regularity results for the p-Laplacian in the Euclidean case to be extended to cover the horizontal gradient in the Heisenberg group setting. This would potentially alter the entire landscape of the field and allow the Heisenberg group case to catch up to the Euclidean case, which right now has a significant lead in development.

Our excess decay estimate is also instrumental for establishing estimates in mean oscillation spaces, namely BMO and VMO. This will be taken up in an upcoming work.

Finally we remark that there is nothing that is inherently scalar in our arguments. As soon as we have an oscillation estimate by energy for the p-Laplacian system in Heisenberg groups, our argument would work the same and produce an excess decay estimate and consequently, both Theorem [3,](#page-5-0) Corollary [4](#page-5-1) and Theorem [5](#page-6-0) for p-Laplacian systems as well. In view of our results, we conjecture that these results hold for systems too.

In the present contribution, we only focus on the p -Laplace equation with or without coefficents in the Heisenberg groups. Although we have not paid any particular attention to this at present, but we believe most, if not all, of our techniques extend to the case of a general nonlinearity

$\text{div}_{\mathbb{H}} a(x, \mathfrak{X} u) = f,$

where the nonlinearity $a: \Omega \times \mathbb{R}^{2n} \to \mathbb{R}^{2n}$ statisfies a p-Laplace type structural assumptions or perhaps even more general nonlinearities. Whether our estimates also extend to cases beyond the Heisenberg group, for example, to Carnot groups of Step 2, remains to be seen and is an interesting question for the future.

Organization : The rest of the article is organized as follows. In Section [2,](#page-8-2) we describe our notations and record the preliminary materials. Section [3](#page-15-0) proves our main estimates for the constant coefficient homogeneous equation, namely the excess decay estimate. Section [4](#page-47-0) proves Theorem [3](#page-5-0) and Corollary [4.](#page-5-1) Section [5](#page-53-0) proves Theorem [6.](#page-6-1) Section [6](#page-59-0) provides a sketch of how to adapt the proof of [\[31\]](#page-63-16) in our setting to prove Theorem [5.](#page-6-0) Our presentation in this section is deliberately extremely brief. The argument in [\[31\]](#page-63-16) is itself somewhat long. Armed with our excess decay estimates, the proof in our case is an exact replica of their argument with the obvious changes in notations. Hence, we refrain from the verbatim repetitions that would add little else apart from additional pages. Thus, we strongly suggest the readers to read Section [6](#page-59-0) alongside the article [\[31\]](#page-63-16) for better comprehension.

2. Preliminaries

2.1. Heisenberg groups.

Definition 7. For $n \geq 1$, the Heisenberg Group denoted by \mathbb{H}_n , is identified with the Euclidean space \mathbb{R}^{2n+1} with the group operation

(17)
$$
x \cdot y := \left(x_1 + y_1, \ \ldots, \ x_{2n} + y_{2n}, \ t + s + \frac{1}{2} \sum_{i=1}^n (x_i y_{n+i} - x_{n+i} y_i)\right)
$$

for every $x = (x_1, \ldots, x_{2n}, t), y = (y_1, \ldots, y_{2n}, s) \in \mathbb{H}_n$.

 \mathbb{H}_n with the group operation [\(17\)](#page-8-3) along with the smooth structure of \mathbb{R}^{2n+1} forms a non Abelian simply connected stratified nilpotent Lie group, whose left invariant vector fields corresponding to the canonical basis of the Lie algebra, are

$$
X_i = \partial_{x_i} - \frac{x_{n+i}}{2} \partial_t, \quad X_{n+i} = \partial_{x_{n+i}} + \frac{x_i}{2} \partial_t,
$$

for every $1 \leq i \leq n$ and the only non zero commutator $T = \partial_t$. We have

(18)
$$
[X_i, X_{n+i}] = T \text{ and } [X_i, X_j] = 0 \ \forall j \neq n+i.
$$

We call X_1, \ldots, X_{2n} as horizontal vector fields and T as the vertical vector field. For a scalar function $f : \mathbb{H}_n \to \mathbb{R}$, we denote $\mathfrak{X} f = (X_1 f, \dots, X_{2n} f)$ and $\mathfrak{X} \mathfrak{X} f =$ $(X_i(X_i f))_{i,j}$ as the Horizontal gradient and Horizontal Hessian, respectively. From [\(18\)](#page-8-4), we have the following trivial but nevertheless, an important inequality $|Tf| \leq$ $2|\mathfrak{X}\mathfrak{X}f|$. For a vector valued function $F = (f_1, \ldots, f_{2n}) : \mathbb{H}_n \to \mathbb{R}^{2n}$, the *Horizontal* divergence is defined as

$$
\operatorname{div}_{\mathbb{H}}(F) = \sum_{i=1}^{2n} X_i f_i.
$$

The Euclidean gradient of a function $g: \mathbb{R}^k \to \mathbb{R}$, shall be denoted by $\nabla g =$ (D_1g, \ldots, D_kg) and the Hessian matrix by $\nabla^2 g$.

2.2. Function spaces on \mathbb{H}_n . The *Carnot-Carathieodory metric* (henceforth CCmetric), which is a left-invariant metric on \mathbb{H}_n , denoted by $d_{\rm CC}$, is defined as the length of the shortest horizontal curves, connecting two points. This is equivalent to the Korànyi metric, denoted as

$$
d_{\mathbb{H}_n}(x, y) = ||y^{-1} \cdot x||_{\mathbb{H}_n},
$$

.

where the Korànyi norm for $x = (x_1, \ldots, x_{2n}, t) \in \mathbb{H}_n$ is

(19)
$$
||x||_{\mathbb{H}_n} := \left(\sum_{i=1}^{2n} x_i^2 + |t|\right)^{\frac{1}{2}}
$$

Throughout this article we use CC-metric balls, which are defined to be

$$
B_r(x) := \{ y \in \mathbb{H}_n : d_{\rm CC}(x, y) < r \} \qquad \text{for } r > 0 \text{ and } x \in \mathbb{H}_n.
$$

However, by virtue of the equivalence of the metrics, all assertions for CC-balls can be restated to Korànyi balls.

The bi-invariant Haar measure of \mathbb{H}_n is just the Lebesgue measure of \mathbb{R}^{2n+1} . We would consistently use the following notations.

Notation 8. For a measurable set $A \subset \mathbb{H}_n$, we denote the $(2n + 1)$ - dimensional Lebesgue measure as |A|. Let $A \subset \mathbb{H}_n$ be any measurable subset with positive measure and let $g: A \to \mathbb{R}^N$ be a measurable function, taking values in any Euclidean space \mathbb{R}^N , $N \geq 1$. We denote its integral average by the notation

$$
(g)_A := \int_A g(x) dx := \frac{1}{|A|} \int_A g(x) dx.
$$

If $A = B(x_0, R)$ for $x_0 \in \mathbb{H}_n$ and $R > 0$, then $(g)_{x_0, R}$ and $(g)_{A}$ denotes the same quantity. If x_0 is fixed in the context, then $(g)_{x_0,R}$ and $(g)_R$ denotes the same quantity.

We record an important property of the integral averages that we would use throughout the rest.

Proposition 9. Let $A \subset \mathbb{H}_n$ be any measurable subset with positive and finite measure and let $g: A \to \mathbb{R}^N$ be a measurable function, taking values in any Euclidean space \mathbb{R}^N , $N \geq 1$. Then for any $\xi \in \mathbb{R}^N$ and any $1 \leq q < \infty$, we have

(20)
$$
\left(\int_A |g - (g)_A|^q \ dx\right)^{\frac{1}{q}} \le 2 \left(\int_A |g - \xi|^q \ dx\right)^{\frac{1}{q}}.
$$

The non-isotropic dilations are the group homorphisms given by

(21)
$$
\delta_R(x_1,\ldots,x_{2n},t) := (Rx_1,\ldots,Rx_{2n},R^2t),
$$

where $R > 0$ and we have the following relation

$$
(22) \t\t B(0,R) = \delta_R B(0,1).
$$

The equivalence between the Korànyi metric and the CC-metric along with the natural scaling [\(21\)](#page-9-1), gives a natural candidate for the homogeneous dimension of the group \mathbb{H}^n , which is $2n + 2$. Throughout this article we denote $Q := 2n + 2$. The Hausdorff dimension with respect to the metric d_{CC} is also Q. Since d_{CC} is left invariant, so we have the following crucial property of metric balls.

Proposition 10. For any CC-metric ball $B_r \subset \mathbb{H}_n$ with $r > 0$, there exists a constant $c = c(n) > 0$ such that

$$
|B_r| = c(n)r^Q.
$$

Let $\Omega \subset \mathbb{H}_n$ be open. For $1 \leq p \leq \infty$, the usual L^p spaces $L^p(\Omega)$ is defined in the usual manner and extended componentwise to vector-valued functions.

Definition 11 (Horizontal Sobolev spaces). For $1 \leq p < \infty$, the Horizontal Sobolev space $HW^{1,p}(\Omega)$ consists of functions $u \in L^p(\Omega)$ such that the distributional horizontal gradient $\mathfrak{X}u$ is in $L^p(\Omega,\mathbb{R}^{2n})$. $HW^{1,p}(\Omega)$ is a Banach space with respect to the norm

(24)
$$
||u||_{HW^{1,p}(\Omega)} = ||u||_{L^p(\Omega)} + ||\mathfrak{X}u||_{L^p(\Omega, \mathbb{R}^{2n})}.
$$

We define $HW^{1,p}_{loc}(\Omega)$ as its local variant and $HW^{1,p}_0(\Omega)$ as the closure of $C_0^{\infty}(\Omega)$ in $HW^{1,p}(\Omega)$ with respect to the norm in [\(24\)](#page-9-2). The second order Horizontal Sobolev space $HW^{2,p}(\Omega)$ is the Banach space

$$
HW^{2,p}(\Omega) := \left\{ u \in HW^{1,p}(\Omega) : \mathfrak{X}u \in HW^{1,p}(\Omega, \mathbb{R}^{2n}) \right\},
$$

equipped with the norm

$$
||u||_{HW^{2,p}(\Omega)} = ||u||_{HW^{1,p}(\Omega)} + ||\mathfrak{XX}u||_{L^p(\Omega,\mathbb{R}^{2n}\times\mathbb{R}^{2n})}.
$$

The definition is extended componentwise to vector valued functions as well.

Definition 12 (Folland-Stein classes). For $0 < \alpha \leq 1$, we define

(25)
$$
C^{0,\alpha}(\Omega) = \left\{ u \in L^{\infty}(\Omega) : \sup_{x,y \in \Omega, x \neq y} \frac{|u(x) - u(y)|}{d_{CC}(x,y)^{\alpha}} < +\infty \right\}
$$

for $0 < \alpha \leq 1$, which are Banach spaces with the norm

(26)
$$
||u||_{C^{0,\alpha}(\Omega)} = ||u||_{L^{\infty}(\Omega)} + [u]_{C^{0,\alpha}(\Omega)},
$$

where the Folland-Stein seminorm is defined as

(27)
$$
[u]_{C^{0,\alpha}(\Omega)} := \sup_{x,y \in \Omega, x \neq y} \frac{|u(x) - u(y)|}{d_{CC}(x,y)^{\alpha}}.
$$

These have standard extensions to classes $C^{k,\alpha}(\Omega)$ for $k \in \mathbb{N}$, which consists of functions having horizontal derivatives up to order k in $C^{0,\alpha}(\Omega)$. The local counterparts are denoted as $C^{k,\alpha}_{\text{loc}}(\Omega)$. Hölder spaces with respect to homogeneous metrics are sometimes called Folland-Stein classes, as they first appeared in Folland-Stein [\[18\]](#page-63-6) and denoted by Γ^{α} or $\Gamma^{0,\alpha}$ in some literature. However, by a harmless abuse of notation, we would continue to use the classical notation. We also recall the definition of modulus of continuity of a uniformly continuous function in our metric setting.

Definition 13. Let $\Omega \subset \mathbb{H}_n$ be an open subset and let $a : \Omega \to \mathbb{R}$ be a uniformly continuous function. The modulus of continuity of a, denoted $\omega_{a,\Omega}$ is a concave increasing function $\omega_{a,\Omega} : [0,\infty) \to [0,\infty)$, defined by

$$
\omega_{a,\Omega}(r) := \sup_{\substack{x,y \in \Omega, \\ d_{CC}(x,y) \le r}} |a(x) - a(y)| \quad \text{for all } r > 0,
$$

and we set $\omega_{a,\Omega}(0) = 0$ by convention.

We would often denote the modulus of continuity by ω_a when the domain is clear and also just ω when the function is also clear from the context. It is easy to see that by definition, $a \in C^{0,\alpha}(\Omega)$ if and only if $\omega_{a,\Omega}(r) \leq Cr^{\alpha}$ for some constant $C > 0$ and the smallest such constant is equal to the seminorm $[a]_{C^{0,\alpha}(\Omega)}$. We now recall the definition of Dini continuity.

Definition 14 (Dini continuity). Let $\Omega \subset \mathbb{H}_n$ be an open subset. A function $a: \Omega \to \mathbb{R}$ is called Dini continuous if a is uniformly continuous in Ω and its modulus of continuity satisfies

(28)
$$
\int_0^\infty \omega_{a,\Omega}(\rho) \frac{\mathrm{d}\rho}{\rho} < \infty.
$$

Now, assume $\Omega \subset \mathbb{H}_n$ is any open subset such that there exists a constant $A = A(\Omega) > 0$ such that for any $x_0 \in \Omega$ and any $r < \text{diam }\Omega$, we have

$$
|B_r(x_0) \cap \Omega| \ge Ar^Q.
$$

This property is clearly satisfied if Ω itself is a metric ball.

Definition 15 (Morrey and Campanato spaces). For $1 \le p < \infty$ and $\lambda \ge 0$, $L^{p,\lambda}(\Omega)$ stands for the Morrey space of all $u \in L^p(\Omega)$ such that

$$
||u||_{\mathcal{L}^{p,\lambda}(\Omega)}^p := \sup_{\substack{x_0 \in \Omega, \\ \rho > 0}} \rho^{-\lambda} \int_{B_{\rho}(x_0) \cap \Omega} |u|^p < \infty,
$$

endowed with the norm $||u||_{L^{p,\lambda}}$ and $\mathcal{L}^{p,\lambda}(\Omega)$ denotes the Campanato space of all $u \in L^p(\Omega)$ such that

$$
[u]_{\mathcal{L}^{p,\lambda}(\Omega)}^p := \sup_{\substack{x_0 \in \Omega, \\ \rho > 0}} \rho^{-\lambda} \int_{B_{\rho}(x_0) \cap \Omega} |u - (u)_{\rho,x_0,\Omega}|^p < \infty,
$$

endowed with the norm $||u||_{\mathcal{L}^{p,\lambda}} := ||u||_{L^p} + [u]_{\mathcal{L}^{p,\lambda}}$. Here

$$
(u)_{\rho,x_0,\Omega} = \frac{1}{|(B_{\rho}(x_0) \cap \Omega)|} \int_{B_{\rho}(x_0) \cap \Omega} u = \int_{B_{\rho}(x_0) \cap \Omega} u.
$$

These definitions are extended componentwise for vector-valued functions. We record the following facts, which are proved exactly in the same manner as their Euclidean counterparts in view of [\(23\)](#page-9-3).

Proposition 16. For $0 \leq \lambda < Q$, we have

$$
L^{p,\lambda}(\Omega) \simeq \mathcal{L}^{p,\lambda}(\Omega) \quad \text{with equivalent norms.}
$$

For $Q < \lambda \leq Q + p$, we have

$$
C^{0,\frac{\lambda-Q}{p}}(\Omega) \simeq \mathcal{L}^{p,\lambda}(\Omega)
$$
 with equivalent seminorms.

When $p > Q$, we also have the following analogue of Morrey's theorem.

Proposition 17. Let $\Omega \subset \mathbb{H}_n$ be open and let $u \in HW^{1,p}_{loc}(\Omega)$ with $p > Q$. Then $u \in C^{0, \frac{p-Q}{p}}(\Omega)$ and for any metric ball $B_{2R} \subset \Omega$, we have the estimate

$$
||u||_{C^{0,\frac{p-Q}{p}}(\overline{B_{R/2}})} \leq C(R,Q,s) ||u||_{HW^{1,p}(\Omega)}.
$$

Proof. By the Poincaré inequality (21) , we have

$$
\int_{B_{\rho}} |u - (u)_{\rho}|^p \leq c\rho^p \int_{B_{\rho}} |\mathfrak{X} u|^p \leq c\rho^p \int_{B_R} |\mathfrak{X} u|^p,
$$

for any $0 < \rho < R$, whenever the ball $B_{2R} \subset \Omega$. This estimate implies, in the standard way, that $u \in \mathcal{L}^{p,p}_{loc}(\Omega)$. Since $p > Q$, the second conclusion of Proposition [16](#page-11-0) implies the result.

For details on classical Morrey and Campanato spaces, we refer to [\[20\]](#page-63-21) and for the sub-elliptic setting we refer to [\[4\]](#page-62-1).

2.3. Lorentz spaces and a series.

Definition 18 (Lorentz spaces). For $1 \leq p < \infty$, and $0 < \theta \leq \infty$, A measurable function $u : \Omega \to \mathbb{R}$ is said to belong to the Lorentz space $L^{(p,\theta)}(\Omega)$ if

$$
||u||_{L^{(p,\theta)}(\Omega)}^{\theta} := \int_0^{\infty} (t^p | \{x \in \Omega : |u| > t\}|)^{\frac{\theta}{p}} \frac{dt}{t} < \infty
$$

for $0 < \theta < \infty$ and if

$$
||u||_{L^{(p,\infty)}(\Omega)}^p := \sup_{t>0} (t^p | \{x \in \Omega : |u| > t\}|) < \infty.
$$

The quantities $||u||_{L(p,\theta)}(\Omega)$ and $||u||_{L(p,\infty)}(\Omega)$ are not norms. They are quasinorms which makes the Lorentz space a quasi-Banach space. For different properties of Lorentz spaces, see [\[24\]](#page-63-22), [\[44\]](#page-64-15). Now we need the following crucial result, which is proved analogously to Lemma 1 in [\[31\]](#page-63-16).

Lemma 19. Let $f \in L^{(Q,1)}(\mathbb{H}_n)$ and for any $1 < q < Q$, define

$$
S_q(x_0, r, \sigma) := \sum_{j=1}^{\infty} r_j \left(\int_{B_j} |f|^q \right)^{\frac{1}{q}},
$$

where $x_0 \in \mathbb{H}_n$, $\sigma \in (0, 1/4)$, $r > 0$ and

$$
B_j := B(x_0, r_j) \qquad and \qquad r_j := \sigma^j r.
$$

Then for any $1 < q < Q$, there exists a constant $c \equiv c(Q, q) > 0$ such that for any $r > 0$ and any $\sigma \in (0, 1/4)$, we have

(29)
$$
S_q(x_0, r, \sigma) \leq c\sigma^{1-\frac{Q}{q}} \|f\|_{L^{(Q,1)}(\mathbb{H}_n)}.
$$

2.4. Sobolev inequalities. We would need the Sobolev inequality for Heisenberg groups, which we write in a scale-invariant form.

Proposition 20 (Scale-invariant Sobolev inequality). Let $B_R \subset \mathbb{H}_n$ be any ball of radius $R > 0$. Let $1 \leq s < Q$. Then there exists a constant $c > 0$, depending only on n and s such that

$$
(30) \qquad \left(\int_{B_R} |u|^{\frac{Q_s}{Q-s}}\right)^{\frac{Q-s}{Q_s}} \le c \left(\int_{B_R} |\mathfrak{X}u|^s + \frac{1}{R^s} \int_{B_R} |u|^s\right)^{\frac{1}{s}},
$$

for any $u \in HW^{1,s}(B_R)$.

We would need the following Poincaré inequality.

Proposition 21 (Poincaré inequality with means). Let $B_R \subset \mathbb{H}_n$ be any ball of radius $R > 0$. Let $1 \leq s < \infty$. Then there exists a constant $c > 0$, depending only on n and s such that for any $u \in HW^{1,s}(B_R)$, we have

(31)
$$
\left(\int_{B_R} |u - (u)_{B_R}|^s \right)^{\frac{1}{s}} \le cR \left(\int_{B_R} |\mathfrak{X} u|^s \right)^{\frac{1}{s}}.
$$

We would also use the following Poincaré-Sobolev type inequalities.

Proposition 22 (Poincaré-Sobolev inequality). Let $B_R \subset \mathbb{H}_n$ be any ball of radius $R > 0$. Let $1 \leq s < Q$. Then there exists a constant $c > 0$, depending only on n and s such that for any $u \in HW^{1,s}(B_R)$, we have

(32)
$$
\left(\int_{B_R} |u|^{\frac{Q_s}{Q-s}}\right)^{\frac{Q-s}{Q_s}} \le cR\left(\int_{B_R} |\mathfrak{X} u|^s\right)^{\frac{1}{s}}.
$$

Proposition 23 (Poincaré-Sobolev inequality with means). Let $B_R \subset \mathbb{H}_n$ be any ball of radius $R > 0$. Let $1 \leq s < Q$. Then there exists a constant $c > 0$, depending only on n and s such that for any $u \in HW^{1,s}(B_R)$, we have

.

(33)
$$
\left(\oint_{B_R} |u - (u)_{B_R}|^{\frac{Q_s}{Q-s}}\right)^{\frac{Q-s}{Q_s}} \le cR\left(\oint_{B_R} |\mathfrak{X} u|^s\right)^{\frac{1}{s}}
$$

The scale invariant forms of (30) , (31) , (32) , (33) can be derived by using their corresponding versions when $R = 1$, along with [\(21\)](#page-9-1) and [\(22\)](#page-9-4). For the case $R = 1$, we can use the Poincaré inequality established in [\[25\]](#page-63-23) and the global density result in [\[19,](#page-63-24) Theorem 1.7] to derive [\(31\)](#page-12-2). Additionally, in this case, by using [\[34,](#page-64-16) Theorem 2.1] and [\[19,](#page-63-24) Theorem 1.7] we can obtain [\(30\)](#page-12-1), [\(32\)](#page-13-1) and [\(33\)](#page-13-2).

2.5. Linear subelliptic Stein theorem. A borderline case of the Sobolev inequality is the following result.

Theorem 24. Let $u \in L^1(\Omega)$ be such that $\mathfrak{X}u \in L_{loc}^{(Q,1)}(\Omega;\mathbb{R}^{2n})$, where $\Omega \subset \mathbb{H}_n$ is open. Then u is continuous in Ω .

A PDE version of the above result is the following, which can be termed as the linear Stein theorem for Heisenberg groups.

Theorem 25. Let $\Omega \subset \mathbb{H}_n$ be open. Let $u \in HW^{1,2}_{loc}(\Omega)$ be such that

$$
\operatorname{div}_{\mathbb{H}} \mathfrak{X} u \in L_{loc}^{(Q,1)}(\Omega) .
$$

Then $\mathfrak{X}u$ is continuous in Ω .

Proof. We prove only the case $\Omega = \mathbb{H}_n$ under the additional assumption that $u \in$ $HW^{1,2}(\mathbb{H}_n)$, div_{$\mathbb{H} \mathfrak{X}u \in L^{(Q,1)}(\mathbb{H}_n)$. Since our statements are local, the general} case follows from this by usual localization arguments and bootstrapping the L^p estimates ([\[15,](#page-63-25) page 917]). Now the main point is that the subLaplacian has a fundamental solution (see [\[14\]](#page-63-5)), we call it $\mathcal N$ in analogy with the Newtonian potential. Thus, we can write

$$
u = \mathcal{N} * (\operatorname{div}_{\mathbb{H}} \mathfrak{X} u) \qquad \text{in } \mathbb{H}_n.
$$

This implies,

$$
\mathfrak{X} u = (\mathfrak{X} \mathcal{N}) * (\operatorname{div}_{\mathbb{H}} \mathfrak{X} u) \qquad \text{in } \mathbb{H}_n.
$$

Note that in both the equations above, the convolution is the group convolution in \mathbb{H}_n . Now, the kernel $K = \mathfrak{X} \mathcal{N}$ is homogeneous of degree $-(Q-1)$ and is essentially the analogue of the kernels of the Riesz potentials I_1 in the Euclidean case. Thus, one can easily prove (see Folland [\[14\]](#page-63-5)) $K \in L^{\left(\frac{Q}{Q-1}, \infty\right)}(\mathbb{H}_n)$. Since $\text{div}_{\mathbb{H}} \mathfrak{X} u \in$ $L^{(Q,1)}(\mathbb{H}_n)$ by assumption and $L^{(Q,1)}$ is the dual space of $L^{(\frac{Q}{Q-1},\infty)}$, for every fixed $y \in \mathbb{H}_n$, we have

$$
\int_{\mathbb{H}_n} K\left(x^{-1} \cdot y\right) \left[\operatorname{div}_{\mathbb{H}} \mathfrak{X} u\right](y) \, \mathrm{d}y \leq c \left\| K \right\|_{L^{\left(\frac{Q}{Q-1},\infty\right)}\left(\mathbb{H}_n\right)} \left\| \operatorname{div}_{\mathbb{H}} \mathfrak{X} u\right\|_{L^{\left(Q,1\right)}\left(\mathbb{H}_n\right)}.
$$

Thus, the map

$$
x \mapsto K * (\operatorname{div}_{\mathbb{H}} \mathfrak{X} u)(x)
$$

is bounded and consequently, continuous. This completes the proof. \Box

Proof of Theorem [24.](#page-13-3) The proof is the same as above. We skip the details. \square

2.6. Minimization and weak formulation.

Proposition 26. Let $\Omega \subset \mathbb{H}_n$ be open and let $a : \Omega \to [\gamma, L]$, where $0 < \gamma < L < \infty$, be a measurable map. Let $1 < p < \infty$. Then for any ball $B_R \subset\subset \Omega$ and any given function $u_0 \in HW^{1,p}(B_R)$, the boundary value problem

(34)
$$
\begin{cases} \operatorname{div}_{\mathbb{H}} \left(a(x) \left| \mathfrak{X} u \right|^{p-2} \mathfrak{X} u \right) = 0 & \text{in } B_R, \\ u = u_0 & \text{on } \partial B_R, \end{cases}
$$

admits a unique weak solution $u \in u_0 + HW_0^{1,p}(B_R)$. Moreover, the solution u is the unique minimizer to the minimization problem

$$
m = \inf \left\{ \frac{1}{p} \int_{B_R} a(x) \left| \mathfrak{X} u \right|^p : u \in u_0 + HW_0^{1,p}(B_R) \right\}.
$$

Proof. In view of the Poincaré-Sobolev inequality (32) , the proof in the Euclidean case works muta
dis mutandis. $\hfill \square$

2.7. The auxiliary mapping V . As in the Euclidean case, we need an auxialiary mapping V to deal with the structure of the p -Laplace equation. We define the mapping $V : \mathbb{R}^{2n} \to \mathbb{R}^{2n}$ by

(35)
$$
V(z) := |z|^{\frac{p-2}{2}} z,
$$

which is a locally Lipschitz bijection from \mathbb{R}^{2n} into itself. We summarize the relevant properties of the map in the following.

Lemma 27. For any $p > 1$, there exists a constant $c_V \equiv c_V(n, p) > 0$ such that

(36)
$$
\frac{|z_1 - z_2|}{c_V} \le \frac{|V(z_1) - V(z_2)|}{(|z_1| + |z_2|)^{\frac{p-2}{2}}} \le c_V |z_1 - z_2|,
$$

for any $z_1, z_2 \in \mathbb{R}^{2n}$, not both zero. This implies the classical monotonicity estimate

$$
(37) \qquad (|z_1|+|z_2|)^{p-2}|z_1-z_2|^2 \le c_M \left\langle |z_1|^{p-2} z_1 - |z_2|^{p-2} z_2, z_1-z_2 \right\rangle,
$$

with a constant $c_M = c_M(n, p) > 0$ for all $p > 1$ and all $z_1, z_2 \in \mathbb{R}^{2n}$. Moreover, if $1 < p \leq 2$, there exists a constant $c = c(n, p) > 0$ such that for any $z_1, z_2 \in \mathbb{R}^{2n}$,

(38)
$$
|z_1 - z_2| \le c |V(z_1) - V(z_2)|^{\frac{2}{p}} + c |V(z_1) - V(z_2)| |z_2|^{\frac{2-p}{2}}.
$$

The estimates (36) and (37) are classical (cf. Lemma 2.1, $[22]$). The estimate [\(38\)](#page-14-2) follows from this (cf. Lemma 2, [\[31\]](#page-63-16)).

Notation 28. We use the symbol c to denote a generic positive constant $c > 0$ in our estimates, unless specifically indicated otherwise, whose value can change from a line to the next.

16 A. MALLICK AND S. SIL

3. Excess decay for subelliptic p-harmonic functions

In this section, we are concerned with regularity results for homogeneous p subLaplace equation in Heisenberg groups. Namely, we shall consider

(39)
$$
\operatorname{div}_{\mathbb{H}}\left(|\mathfrak{X}u|^{p-2}\mathfrak{X}u\right)=0 \qquad \qquad \text{in } \Omega.
$$

We note that the equation can also be written as

(40)
$$
\operatorname{div}_{\mathbb{H}}\left(Df(\mathfrak{X}u)\right)=\sum_{i=1}^{2n}X_i\big(D_if(\mathfrak{X}u)\big)=0\qquad\text{in }\Omega,
$$

where $Df = (D_1f, D_2f, \ldots, D_{2n}f)$ is the Euclidean gradient of f and f is given by

$$
f(z) := \frac{1}{p} |z|^p, \quad \forall z \in \mathbb{R}^{2n}
$$

.

We compute

(41)
$$
D_j D_i f(z) = |z|^{p-2} \delta_{ij} + (p-2)|z|^{p-4} z_i z_j
$$
, for $i, j = 1, ..., 2n$ and $z \in \mathbb{R}^{2n}$,
where δ_{ij} denotes the Kronecker delta function. Hence, we deduce

(42)
$$
\sum_{i,j=1}^{2n} D_j D_i f(z) \eta_j \eta_i \ge c |z|^{p-2} |\eta|^2 \quad \text{for all } z, \eta \in \mathbb{R}^{2n}.
$$

3.1. Hölder continuity results. We begin by recording the known results that we would use. The following theorem is due to Zhong [\[50\]](#page-64-8).

Theorem 29. Let $1 < p < \infty$ and $u \in HW^{1,p}(\Omega)$ be a weak solution of equation [\(39\)](#page-15-1). Then $\mathfrak{X}u \in L^{\infty}_{loc}(\Omega;\mathbb{R}^{2n})$. Moreover, for any ball $B_{2r}(x_0) \subset \Omega$, we have that

(43)
$$
\sup_{B_r(x_0)} |\mathfrak{X} u| \leq c_p \Big(\int_{B_{2r}(x_0)} |\mathfrak{X} u|^p \ dx \Big)^{\frac{1}{p}},
$$

where $c_p > 0$ depends only on n, p.

By a standard interpolation technique, this last result easily implies the following.

Theorem 30. Let $1 < p < \infty$ and $u \in HW^{1,p}(\Omega)$ be a weak solution of equation [\(39\)](#page-15-1). Then $\mathfrak{X}u \in L^{\infty}_{loc}(\Omega;\mathbb{R}^{2n})$. Moreover, for any $0 < q < \infty$ and any ball $B_{2r}(x_0) \subset \Omega$, we have the estimate

(44)
$$
\sup_{B_r(x_0)} |\mathfrak{X} u| \leq c_q \Big(\int_{B_{2r}(x_0)} |\mathfrak{X} u|^q \ dx \Big)^{\frac{1}{q}},
$$

where $c_q > 0$ depends only on n, p and q.

The following is due to Zhong [\[50\]](#page-64-8) (for $p \geq 2$) and Mukherjee-Zhong [\[39\]](#page-64-9) (for $1 < p < 2$).

Theorem 31. Let $1 < p < \infty$ and $u \in HW^{1,p}(\Omega)$ be a weak solution of equation [\(39\)](#page-15-1). Then the horizontal gradient $\mathfrak{X}u$ is locally Hölder continuous. Moreover, there is a positive exponent $\alpha_1 = \alpha_1(n, p) < 1$ such that for any ball $B_{r_0} \subset\subset \Omega$ and any $0 < r \leq r_0$, we have

(45)
$$
|\mathfrak{X}u(x)-\mathfrak{X}u(y)| \leq c_h \left(\frac{r}{r_0}\right)^{\alpha_1} ||\mathfrak{X}u||_{\infty,r_0}, \quad \text{for all } x, y \in B_r,
$$

where $c_h > 0$ depends only on n, p, where $\|\mathfrak{X}u\|_{\infty,r_0} := \sup_{B_{r_0}} |\mathfrak{X}u|$.

Theorem [31](#page-15-2) and Theorem [29](#page-15-3) easily implies the following result.

Theorem 32. Let $1 < p < \infty$ and $u \in HW^{1,p}(\Omega)$ be a weak solution of equation [\(39\)](#page-15-1) Then for any ball $B_{r_0} \subset \subset \Omega$ and any $0 < r \leq r_0$, we have

(46)
$$
\sup_{x,y\in B_r} |V(\mathfrak{X}u)(x)-V(\mathfrak{X}u)(y)| \leq c_{h,V} \left(\frac{r}{r_0}\right)^{\beta_1} \|\mathfrak{X}u\|_{\infty,r_0}^{\frac{p}{2}},
$$

where the constants $c_{h,V} > 0$ and $\beta_1 \in (0,1)$, both depends only on n and p.

Proof. We have

$$
\sup_{x,y\in B_r} |V(\mathfrak{X}u)(x) - V(\mathfrak{X}u)(y)|^2
$$
\n
$$
\leq c_V \sup_{x,y\in B_r} \left[(|\mathfrak{X}u(x)| + |\mathfrak{X}u(y)|)^{p-2} |\mathfrak{X}u(x) - \mathfrak{X}u(y)|^2 \right]
$$
\n
$$
\leq c_M c_V \sup_{x,y\in B_r} \left[\left\langle |\mathfrak{X}u(x)|^{p-2} \mathfrak{X}u(x) - |\mathfrak{X}u(y)|^{p-2} \mathfrak{X}u(y), \mathfrak{X}u(x) - \mathfrak{X}u(y) \right\rangle \right]
$$
\n
$$
\leq c_M c_V \sup_{x,y\in B_r} \left[\left| |\mathfrak{X}u(x)|^{p-2} \mathfrak{X}u(x) - |\mathfrak{X}u(y)|^{p-2} \mathfrak{X}u(y) \right| |\mathfrak{X}u(x) - \mathfrak{X}u(y)| \right]
$$
\n
$$
\leq c_M c_V \sup_{x,y\in B_r} \left[\left(|\mathfrak{X}u(x)|^{p-1} + |\mathfrak{X}u(y)|^{p-1} \right) |\mathfrak{X}u(x) - \mathfrak{X}u(y)| \right]
$$
\n
$$
\leq 2^{p-1} c_M c_V \left| |\mathfrak{X}u| \right|_{\infty, r_0}^{p-1} \left(\sup_{x,y\in B_r} |\mathfrak{X}u(x) - \mathfrak{X}u(y)| \right)
$$
\n
$$
\stackrel{(45)}{\leq} 2^{p-1} c_M c_V c_h \left(\frac{r}{r_0} \right)^{\alpha_1} \left| |\mathfrak{X}u| \right|_{\infty, r_0}^p.
$$

Thus, setting $\beta_1 := \alpha_1/2$, we have

$$
\sup_{x,y\in B_r} |V(\mathfrak{X} u)(x) - V(\mathfrak{X} u)(y)| \leq c \left(\frac{r}{r_0}\right)^{\beta_1} {\|\mathfrak{X} u\|}_{\infty,r_0}^{\frac{p}{2}}.
$$

This completes the proof.

3.2. Caccioppoli inequalities in the nondegenerate regime. We begin by deriving some Caccioppoli type inequalities under the assumption that $|\mathfrak{X}u|$ is both bounded above and below in a fixed scale $\lambda > 0$.

Lemma 33. Let $1 < p < \infty$ and let $\Omega \subset \mathbb{H}_n$ be open. Let $u \in HW^{1,p}_{loc}(\Omega)$ be a weak solution to

div^H |Xu| ^p−² Xu = 0 in Ω.

Let $\tilde{\Omega} \subset\subset \Omega$ be any open subset. Suppose we have

(47)
$$
\frac{\lambda}{4B} \leq |\mathfrak{X}u| \leq A\lambda \qquad \text{in } \tilde{\Omega},
$$

for some constants $A, B \ge 1$ and $\lambda > 0$. Let $B_\rho \subset B_R \subset \tilde{\Omega}$ be concentric balls of radius $0 < \rho < R < 1$. Let $\eta \in C_c^{\infty}(B_R)$ satisfy $0 \leq \eta \leq 1$ in B_R with $\eta \equiv 1$ in B_{ρ} and $|\mathfrak{X}\eta| \leq c/(R-\rho)$ for some absolute constant $c > 1$. Then there exists a

$$
\sqcup
$$

constant $C_0 \equiv C_0 (A, B, n, p) \geq 1$, but independent of λ , R, ρ or u, such that for any $\zeta \in \mathbb{R}$ and any $\xi \in \mathbb{R}^{2n}$, we have the following inequalitites.

(48)
$$
\int_{B_R} \eta^4 |\mathfrak{X} T u|^2 dx \leq \frac{C}{(R - \rho)^2} \int_{B_R} \eta^2 |T u - \zeta|^2,
$$

(49)
$$
\int_{B_R} \eta^4 |\mathfrak{X} T u|^2 dx \leq \frac{C}{(R-\rho)^4} \int_{B_R} |\mathfrak{X} u - \xi|^2 dx,
$$

(50)
$$
\int_{B_R} \eta^2 |Tu|^2 dx \leq \frac{C}{(R-\rho)^2} \int_{B_R} |\mathfrak{X}u - \xi|^2 dx,
$$

(51)
$$
\int_{B_R} \eta^2 |\mathfrak{XX} u|^2 \, dx \leq \frac{C}{(R-\rho)^2} \int_{B_R} |\mathfrak{X} u - \xi|^2 \, dx.
$$

Proof. Note that by [\(47\)](#page-16-0) and Capogna's regularity result in [\[2\]](#page-62-3), we have

$$
\mathfrak{X} u \in HW_{\text{loc}}^{1,2}\left(B_R; \mathbb{R}^{2n}\right) \text{ and } Tu \in HW_{\text{loc}}^{1,2}\left(B_R\right) \cap L_{\text{loc}}^{\infty}\left(B_R\right).
$$

One can easily check that these regularity assertions are enough to justify our calculations in this lemma. Observe also that it is easy to verify using integration by parts and the definition of weak derivative that

$$
Tu \in HW_{\text{loc}}^{1,2}(B_R) \Rightarrow \mathfrak{X}Tu \in L_{\text{loc}}^2(B_R; \mathbb{R}^{2n})
$$

$$
\Rightarrow T\mathfrak{X}u \in L_{\text{loc}}^2(B_R; \mathbb{R}^{2n}) \text{ as well and } \mathfrak{X}Tu = T\mathfrak{X}u.
$$

We would not comment on this further.

Claim 34. For every $l = 1, 2, ..., n$, $\psi_l := X_l u$ is a weak solution of $(52) \sum_{1}^{2n}$ $X_i\big(D_jD_if(\mathfrak{X}u)X_j\psi_l\big) + \sum^{2n}$ $X_i(D_{n+l}D_i f(\mathfrak{X}u)Tu) + T(D_{n+l} f(\mathfrak{X}u)) = 0,$

 $i=1$

and $\psi_{n+l} = X_{n+l}u$ is a weak solution of

 $i,j=1$

(53)
$$
\sum_{i,j=1}^{2n} X_i(D_j D_i f(\mathfrak{X} u) X_j \psi_{n+l}) - \sum_{i=1}^{2n} X_i(D_l D_i f(\mathfrak{X} u) T u) - T(D_l f(\mathfrak{X} u)) = 0.
$$

Furthermore, Tu is a weak solution of

 \sim

(54)
$$
\sum_{i,j=1}^{2n} X_i(D_j D_i f(\mathfrak{X} u) X_j(T u)) = 0.
$$

This can be verified by a direct calculation, see Lemma 3.1 and Lemma 3.2 in [\[50\]](#page-64-8). Now we plugg $\varphi = \eta^4 (Tu - \zeta)$ with $\zeta \in \mathbb{R}$ as a test function in the weak formulation of [\(54\)](#page-17-3) to deduce

$$
0 = \int_{\Omega} \eta^4 \sum_{i,j=1}^{2n} D_j D_i f(\mathfrak{X}u) X_j(Tu) X_i(Tu) dx
$$

+
$$
4 \int_{\Omega} \eta^3 \sum_{i,j=1}^{2n} D_j D_i f(\mathfrak{X}u) X_j(Tu) X_i \eta(Tu - \zeta) dx.
$$

Hence, using [\(42\)](#page-15-5) and Young's inequality with $\varepsilon > 0$, we have

$$
\int_{\Omega} \eta^4 |\mathfrak{X} u|^{p-2} |\mathfrak{X} u|^2 dx
$$
\n
$$
\leq c \int_{\Omega} \eta^3 |\mathfrak{X} u|^{p-2} |\mathfrak{X} \eta| |\mathfrak{X} u| |Tu - \zeta| dx
$$
\n
$$
\leq c \varepsilon \int_{\Omega} \eta^4 |\mathfrak{X} u|^{p-2} |\mathfrak{X} u|^2 dx + C_{\varepsilon} \int_{\Omega} \eta^2 |\mathfrak{X} u|^{p-2} |\mathfrak{X} \eta|^2 |Tu - \zeta|^2 dx.
$$

Choosing $\varepsilon > 0$ small enough, we have

$$
\int_{\Omega} \eta^4 |\mathfrak{X} u|^{p-2} |\mathfrak{X} Tu|^2 \ dx \leq C \int_{\Omega} \eta^2 |\mathfrak{X} \eta|^2 |\mathfrak{X} u|^{p-2} |Tu - \zeta|^2 \ dx.
$$

In view of [\(47\)](#page-16-0), this immediately implies [\(48\)](#page-17-4). Now to derive [\(49\)](#page-17-5), we begin by deducing a crucial estimate for Tu. For any $\xi \in \mathbb{R}^{2n}$, integrating by parts and using Young's inequality with $\varepsilon > 0$, we have

$$
\int_{\Omega} \eta^2 |Tu|^2 dx
$$

\n
$$
= \int_{\Omega} \eta^2 Tu (X_l X_{n+l} u - X_{n+l} X_l u) dx
$$

\n
$$
= \int_{\Omega} \eta^2 Tu [X_l (X_{n+l} u - \xi_{n+l}) - X_{n+l} (X_l u - \xi_l)] dx
$$

\n
$$
= - \int_{\Omega} X_l (\eta^2 Tu) (X_{n+l} u - \xi_{n+l}) dx + \int_{\Omega} X_{n+l} (\eta^2 Tu) (X_l u - \xi_l) dx
$$

\n
$$
= -2 \int_{\Omega} \eta X_l \eta (Tu) (X_{n+l} u - \xi_{n+l}) dx - \int_{\Omega} \eta^2 X_l Tu (X_{n+l} u - \xi_{n+l}) dx
$$

\n
$$
+ 2 \int_{\Omega} \eta X_{n+l} \eta (Tu) (X_l u - \xi_l) dx + \int_{\Omega} \eta^2 X_{n+l} Tu (X_l u - \xi_l) dx
$$

\n
$$
\leq 4 \int_{\Omega} \eta |\mathfrak{X} \eta| |Tu| |\mathfrak{X} u - \xi| + 2 \int_{\Omega} \eta^2 |\mathfrak{X} Tu| |\mathfrak{X} u - \xi|
$$

\n
$$
\leq \varepsilon \int_{\Omega} \eta^2 |Tu|^2 dx + C_{\varepsilon} \int_{\Omega} |\mathfrak{X} \eta|^2 |\mathfrak{X} u - \xi|^2 dx + 2 \int_{\Omega} \eta^2 |\mathfrak{X} Tu| |\mathfrak{X} u - \xi|.
$$

Choosing $\varepsilon = 1/2$, this implies

(55)
$$
\int_{\Omega} \eta^2 |Tu|^2 dx \leq C \left(\int_{\Omega} |\mathfrak{X}\eta|^2 |\mathfrak{X}u - \xi|^2 dx + \int_{\Omega} \eta^2 |\mathfrak{X}Tu||\mathfrak{X}u - \xi| \right).
$$

Combining this with [\(48\)](#page-17-4) with the choice $\zeta = 0$ and using Young's inequality with $\delta > 0$, we arrive at

$$
\int_{B_R} \eta^4 |\mathfrak{X} T u|^2 dx
$$
\n
$$
\leq \frac{C}{(R-\rho)^2} \left(\int_{B_R} |\mathfrak{X} \eta|^2 |\mathfrak{X} u - \xi|^2 dx + \int_{B_R} \eta^2 |\mathfrak{X} T u| |\mathfrak{X} u - \xi| dx \right)
$$
\n
$$
\leq \frac{C}{(R-\rho)^2} \int_{B_R} |\mathfrak{X} \eta|^2 |\mathfrak{X} u - \xi|^2 dx + C\delta^2 \int_{B_R} \eta^4 |\mathfrak{X} T u|^2 dx
$$
\n
$$
+ \frac{C}{(R-\rho)^4 \delta^2} \int_{B_R} |\mathfrak{X} u - \xi|^2 dx.
$$

Choose $\delta^2 = \frac{1}{2C}$ we have ˆ B_R $\eta^4 |\mathfrak{X} T u|^2 dx \leq \frac{C}{\sqrt{2}}$ $(R-\rho)^2$ ˆ B_R $|\mathfrak{X}\eta|^2 |\mathfrak{X}u-\xi|^2 dx + \frac{C}{\sqrt{2}}$ $(R-\rho)^4$ ˆ B_R $|\mathfrak{X} u - \xi|^2 dx$.

Thus we have proved [\(49\)](#page-17-5). On the other hand, using this back in [\(55\)](#page-18-0), we deduce

$$
\int_{B_R} \eta^2 |Tu|^2 dx
$$
\n
$$
\leq C \left(\int_{B_R} |\mathfrak{X}\eta|^2 |\mathfrak{X}u - \xi|^2 dx + \int_{B_R} \eta^2 |\mathfrak{X}Tu||\mathfrak{X}u - \xi| \right)
$$
\n
$$
\leq \frac{C}{(R - \rho)^2} \int_{B_R} |\mathfrak{X}u - \xi|^2 dx + \left(\int_{B_R} \eta^4 |\mathfrak{X}Tu|^2 dx \right)^{\frac{1}{2}} \left(\int_{B_R} |\mathfrak{X}u - \xi|^2 dx \right)^{\frac{1}{2}}
$$
\n
$$
\stackrel{(49)}{\leq} \frac{C}{(R - \rho)^2} \int_{B_R} |\mathfrak{X}u - \xi|^2 dx + \frac{C}{(R - \rho)^2} \int_{B_R} |\mathfrak{X}u - \xi|^2 dx.
$$

This proves [\(50\)](#page-17-6).

Now we finally turn our attention to [\(51\)](#page-17-2). Fix $l \in \{1, 2, ..., n\}$ and use $\varphi =$ $\eta^2(X_l u - \xi_l)$ as a test function in [\(52\)](#page-17-0) to deduce

$$
\int_{B_R} \sum_{i,j=1}^{2n} \eta^2 D_j D_i f(\mathfrak{X}u) X_j X_l u X_i (X_l u - \xi_l) dx
$$

= $-2 \int_{B_R} \sum_{i,j=1}^{2n} \eta D_j D_i f(\mathfrak{X}u) X_j X_l u X_i \eta (X_l u - \xi_l) dx$
 $- \int_{B_R} \sum_{i=1}^{2n} D_{n+l} D_i f(\mathfrak{X}u) T u X_i (\eta^2 (X_l u - \xi_l)) dx$
(56) $+ \int_{B_R} T(D_{n+l} f(\mathfrak{X}u)) (\eta^2 (X_l u - \xi_l)) dx := I_1^l + I_2^l + I_3^l.$

Similarly, we can deduce from [\(53\)](#page-17-1) that for all $l \in \{n+1, \ldots, 2n\}$,

$$
\int_{B_R} \sum_{i,j=1}^{2n} \eta^2 D_j D_i f(\mathfrak{X}u) X_j X_l u X_i (X_l u - \xi_l) dx
$$

= $- 2 \int_{B_R} \sum_{i,j=1}^{2n} \eta D_j D_i f(\mathfrak{X}u) X_j X_l u X_i \eta (X_l u - \xi_l) dx$
+ $\int_{B_R} \sum_{i=1}^{2n} D_{l-n} D_i f(\mathfrak{X}u) T u X_i (\eta^2 (X_l u - \xi_l)) dx$
(57) $- \int_{B_R} T(D_{l-n} f(\mathfrak{X}u)) (\eta^2 (X_l u - \xi_l)) dx := I_1^l + I_2^l + I_3^l.$

Summing the above equations for all l from 1 to $2n$, we arrive at

(58)
$$
\int_{B_R} \sum_{i,j,l} \eta^2 D_j D_i f(\mathfrak{X} u) X_j X_l u X_i (X_l u - \xi_l) dx = \sum_l (I_1^l + I_2^l + I_3^l),
$$

where we sum for i, j, l from 1 to 2n. By [\(42\)](#page-15-5), we deduce

(59)
$$
\int_{B_R} \eta^2 |\mathfrak{X} u|^{p-2} |\mathfrak{X} \mathfrak{X} u|^2 dx \leq C \sum_l \left(I_1^l + I_2^l + I_3^l \right)
$$

Now, we have

$$
\begin{aligned} \left|I_1^l\right| &\leq C \int_{B_R} \eta |\mathfrak{X} u|^{p-2} |\mathfrak{XX} u| |\mathfrak{X} \eta| |\mathfrak{X} u - \xi| \, dx \\ (60) \qquad &\leq \varepsilon \int_{B_R} \eta^2 |\mathfrak{X} u|^{p-2} |\mathfrak{XX} u|^2 \, dx + C_{\varepsilon} \int_{B_R} |\mathfrak{X} \eta|^2 |\mathfrak{X} u|^{p-2} |\mathfrak{X} u - \xi|^2 \, dx. \end{aligned}
$$

Now for I_2^l , using the fact that $|Tu| \leq C |\mathfrak{XX}u|$, we have

$$
\begin{aligned}\n|I_2^l| &\leq C \int_{B_R} \eta |\mathfrak{X}u|^{p-2} |Tu| |\mathfrak{X}\eta| |\mathfrak{X}u - \xi| \, dx + C \int_{B_R} \eta^2 |\mathfrak{X}u|^{p-2} |Tu| |\mathfrak{X}\mathfrak{X}u| \, dx \\
&\leq C \int_{B_R} \eta |\mathfrak{X}u|^{p-2} |\mathfrak{X}\mathfrak{X}u| |\mathfrak{X}\eta| |\mathfrak{X}u - \xi| \, dx + \varepsilon \int_{B_R} \eta^2 |\mathfrak{X}u|^{p-2} |\mathfrak{X}\mathfrak{X}u|^2 \, dx \\
&\quad + C_{\varepsilon} \int_{B_R} \eta^2 |\mathfrak{X}u|^{p-2} |Tu|^2 \, dx\n\end{aligned}
$$
\n(i)

\n
$$
\leq \varepsilon \int \eta^2 |\mathfrak{X}u|^{p-2} |\mathfrak{X}\mathfrak{X}u|^2 \, dx + C_{\varepsilon} \int \eta^2 |\mathfrak{X}u|^{p-2} |Tu|^2 \, dx
$$

(61)
$$
\leq \varepsilon \int_{\Omega} \eta^2 |\mathfrak{X} u|^{p-2} |\mathfrak{XX} u|^2 dx + C_{\varepsilon} \int_{\Omega} \eta^2 |\mathfrak{X} u|^{p-2} |Tu|^2 dx
$$

$$
+ C_{\varepsilon} \int_{\Omega} |\mathfrak{X} \eta|^2 |\mathfrak{X} u|^{p-2} |\mathfrak{X} u - \xi|^2 dx.
$$

Now, for I_3^l , when $1 \leq l \leq n$, we compute

$$
T(D_{n+l}f(\mathfrak{X}u)) = T\left(|\mathfrak{X}u|^{p-2}\right)X_{n+l}u + |\mathfrak{X}u|^{p-2}TX_{n+l}u
$$

=
$$
(p-2)|\mathfrak{X}u|^{p-4}\langle \mathfrak{X}u,T\mathfrak{X}u\rangle X_{n+l}u + |\mathfrak{X}u|^{p-2}TX_{n+l}u.
$$

Similarly, when $n + 1 \leq l \leq 2n$, we have

$$
T(D_{l-n}f(\mathfrak{X}u)) = T\left(|\mathfrak{X}u|^{p-2}\right)X_{l-n}u + |\mathfrak{X}u|^{p-2}TX_{l-n}u
$$

=
$$
(p-2)|\mathfrak{X}u|^{p-4}\langle \mathfrak{X}u, T\mathfrak{X}u\rangle X_{l-n}u + |\mathfrak{X}u|^{p-2}TX_{l-n}u.
$$

Hence, in either case we have,

(62)
$$
\left|I_3^l\right| \leq C \int_{\Omega} \eta^2 |\mathfrak{X} u|^{p-2} |\mathfrak{X} Tu| |\mathfrak{X} u - \xi| dx.
$$

Combining [\(59\)](#page-20-0), [\(60\)](#page-20-1), [\(61\)](#page-20-2) and [\(62\)](#page-20-3) and choosing $\varepsilon > 0$ sufficiently small, we deduce

$$
\int_{\Omega} \eta^2 |\mathfrak{X}u|^{p-2} |\mathfrak{XX}u|^2 dx
$$

\n
$$
\leq C \int_{\Omega} \eta^2 |\mathfrak{X}u|^{p-2} |Tu|^2 dx + C \int_{\Omega} |\mathfrak{X}\eta|^2 |\mathfrak{X}u|^{p-2} |\mathfrak{X}u - \xi|^2 dx
$$

\n(63)
\n
$$
+ C \int_{\Omega} \eta^2 |\mathfrak{X}u|^{p-2} |\mathfrak{X}Tu| |\mathfrak{X}u - \xi| dx.
$$

Now, combining [\(63\)](#page-20-4), [\(49\)](#page-17-5), [\(50\)](#page-17-6) and [\(47\)](#page-16-0), we deduce

$$
\int_{B_R} \eta^2 |\mathfrak{XX} u|^2 dx
$$
\n
$$
\leq \frac{C}{(R-\rho)^2} \int_{B_R} |\mathfrak{X} u - \xi|^2 dx + C \int_{B_R} \eta^2 |\mathfrak{X} Tu| |\mathfrak{X} u - \xi| dx
$$
\n
$$
\leq \frac{C}{(R-\rho)^2} \int_{B_R} |\mathfrak{X} u - \xi|^2 dx + C \left(\int_{B_R} \eta^4 |\mathfrak{X} Tu|^2 dx \right)^{\frac{1}{2}} \left(\int_{B_R} |\mathfrak{X} u - \xi|^2 dx \right)^{\frac{1}{2}}
$$
\n
$$
\leq \frac{C}{(R-\rho)^2} \int_{B_R} |\mathfrak{X} u - \xi|^2 dx.
$$

This proves (51) and completes the proof of the Lemma.

3.3. Reverse Hölder inequalities. Our Caccioppoli inequalities in Lemma [33](#page-16-1) leads us to a few reverse Hölder inequalities in the nondegenerate regime.

Lemma 35. Let $1 < p < \infty$ and let $\Omega \subset \mathbb{H}_n$ be open. Let $u \in HW^{1,p}_{loc}(\Omega)$ be a weak solution to

$$
\operatorname{div}_{\mathbb{H}}\left(\left|\mathfrak{X}u\right|^{p-2}\mathfrak{X}u\right)=0\qquad \qquad \text{in }\Omega.
$$

Let $\tilde{\Omega} \subset\subset \Omega$ be any open subset. Suppose we have

(64)
$$
\frac{\lambda}{4B} \leq |\mathfrak{X}u| \leq A\lambda \qquad \text{in } \tilde{\Omega},
$$

for some constants $A, B \ge 1$ and $\lambda > 0$. Then there exists a constant $C_{RH} \equiv$ $C_{RH}(A, B, n, p) \geq 1$, such that for any $\xi \in \mathbb{R}^{2n}$, and for any ball $B_R \subset \tilde{\Omega}$ with $R > 0$, we have the following reverse Hölder inequality

(65)
$$
\left(\oint_{B_{R/2}} |\mathfrak{X} u - \xi|^{\frac{2Q}{Q-2}} dx\right)^{\frac{Q-2}{2Q}} \leq C_{RH} \int_{B_R} |\mathfrak{X} u - \xi| dx.
$$

In particular, for any exponent $1 \leq q \leq 2Q/(Q-2)$, we have the following estimate

(66)
$$
\left(\int_{B_{R/2}} |\mathfrak{X} u - \xi|^q \, dx\right)^{\frac{1}{q}} \leq C_{RH} \left(\int_{B_R} |\mathfrak{X} u - \xi|^q \, dx\right)^{\frac{1}{q}}.
$$

Proof. In view of [\(64\)](#page-21-1), the Caccippoli inequality [\(51\)](#page-17-2) holds for any ball $B \subset \Omega$. Now fix concentric balls $B_\rho \subset B_r \subset \tilde{\Omega}$. Let $\eta \in C_c^\infty(B_R)$, $0 \leq \eta \leq 1$ in B_R with $\eta \equiv 1$ in B_ρ and $|\mathfrak{X}\eta| \leq c/(R-\rho)$ for some constant $c > 1$. Then $\eta(\mathfrak{X}u - \xi) \in$ $HW_0^{1,2} (B_R; \mathbb{R}^{2n})$ for any $\xi \in \mathbb{R}^{2n}$. Thus we have

$$
\int_{B_{\rho}} |\mathfrak{X}u - \xi|^{\frac{2Q}{Q-2}} \leq \int_{B_R} |\eta(\mathfrak{X}u - \xi)|^{\frac{2Q}{Q-2}}
$$
\n
$$
\leq C \left(\int_{B_R} |\mathfrak{X}[\eta(\mathfrak{X}u - \xi)]|^2 \right)^{\frac{Q}{Q-2}}
$$
\n
$$
\leq C \left(\int_{B_R} \eta^2 |\mathfrak{XX}u|^2 + \int_{B_R} |\langle \mathfrak{X}\eta, \mathfrak{X}u - \xi \rangle|^2 \right)^{\frac{Q}{Q-2}}
$$
\n(67)\n
$$
\leq C (R - \rho)^{-\frac{2Q}{Q-2}} \left(\int_{B_R} |\mathfrak{X}u - \xi|^2 \right)^{\frac{Q}{Q-2}}.
$$

This implies, using the Hölder inequality and the Young's inequality,

$$
\begin{split} \left(\int_{B_{\rho}}|\mathfrak{X} u-\xi|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2Q}} &\leq \frac{C}{(R-\rho)}\left(\int_{B_{R}}|\mathfrak{X} u-\xi|^{2}\right)^{\frac{1}{2}}\\ &\leq \frac{C}{(R-\rho)}\left(\int_{B_{R}}|\mathfrak{X} u-\xi|^{\frac{2Q}{Q+2}}|\mathfrak{X} u-\xi|^{\frac{4}{Q+2}}\right)^{\frac{1}{2}}\\ &\leq \frac{C}{(R-\rho)}\left(\int_{B_{R}}|\mathfrak{X} u-\xi|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2(Q+2)}}\left(\int_{B_{R}}|\mathfrak{X} u-\xi|\right)^{\frac{2}{Q+2}}\\ &\leq \frac{1}{2}\left(\int_{B_{R}}|\mathfrak{X} u-\xi|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2Q}}+\frac{C}{(R-\rho)^{\frac{Q+2}{2}}}\int_{B_{R}}|\mathfrak{X} u-\xi|\,. \end{split}
$$

Thus, setting

$$
\phi\left(r\right):=\left(\int_{B_{r}}\left|\mathfrak{X} u-\xi\right|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2Q}}
$$

and appying Lemma 8.18 in [\[20\]](#page-63-21), we deduce

$$
\left(\int_{B_{\rho}}|\mathfrak{X} u-\xi|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2Q}}\leq C\left(R-\rho\right)^{-\frac{Q+2}{2}}\int_{B_{R}}|\mathfrak{X} u-\xi|
$$

for every $0 < \rho < R$. Setting $\rho = R/2$, we arrive at

$$
\left(\oint_{B_{R/2}}|\mathfrak{X} u-\xi|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2Q}}\leq C\int_{B_R}|\mathfrak{X} u-\xi|.
$$

This proves [\(65\)](#page-21-0) and [\(66\)](#page-21-2) follows from this using Hölder inequality on both sides. This completes the proof.

As a consequence, we have the following result for $V(\mathfrak{X}u)$, which is not quite as sharp when $1 < p < 2$, but this would suffice for our purposes.

Lemma 36. Let $1 < p < \infty$ and let $\Omega \subset \mathbb{H}_n$ be open. Let $u \in HW^{1,p}_{loc}(\Omega)$ be a weak solution to

$$
\operatorname{div}_{\mathbb{H}}\left(|\mathfrak{X}u|^{p-2}\mathfrak{X}u\right)=0 \qquad \qquad in \ \Omega.
$$

Let $\tilde{\Omega} \subset\subset \Omega$ be any open subset. Suppose we have

(68)
$$
\frac{\lambda}{4B} \leq |\mathfrak{X}u| \leq A\lambda \qquad \text{in } \tilde{\Omega},
$$

for some constants $A, B \geq 1$ and $\lambda > 0$. Let $\xi \in \mathbb{R}^{2n}$ be a constant vector if $2 \leq p < \infty$ and

(69)
$$
|\xi| \le A\lambda \quad \text{if } 1 < p < 2.
$$

Then there exists a constant $C_{RH} = C_{RH,V}(A, B, n, p) \geq 1$, such that for any ball $B_R \subset \tilde{\Omega}$ with $R > 0$, we have

(70)
$$
\left(\int_{B_{R/2}} |V(\mathfrak{X}u) - V(\xi)|^{\frac{2Q}{Q-2}} dx\right)^{\frac{Q-2}{2Q}} \leq C_{RH,V} \int_{B_R} |V(\mathfrak{X}u) - V(\xi)| dx.
$$

In particular, for any exponent $1 \leq q \leq 2Q/(Q-2)$, we have the following estimate

$$
(71)\quad \left(\oint_{B_{R/2}} |V(\mathfrak{X}u) - V(\xi)|^q \mathrm{d}x\right)^{\frac{1}{q}} \leq C_{RH,V}\left(\oint_{B_R} |V(\mathfrak{X}u) - V(\xi)|^q \mathrm{d}x\right)^{\frac{1}{q}},
$$

for any $\xi \in \mathbb{R}^n$ which satisfies [\(69\)](#page-22-0).

Proof. We first tackle the case $p \geq 2$, where no bound on ξ is needed. To derive the inequalities for $V(\mathfrak{X}u)$, observe that $V(\cdot)$ is smooth away from the origin and in view of the pointwise bounds [\(68\)](#page-22-1), we can justify the computation of derivatives and deduce the inequality

$$
\left|\mathfrak{X}\left[V\left(\mathfrak{X} u\right)\right]\right| \leq c\left|\mathfrak{X} u\right|^{\frac{p-2}{2}}\left|\mathfrak{X} \mathfrak{X} u\right|
$$

for some constant $c > 0$, which depends only on n and p. This, combined with the Caccippoli inequality [\(51\)](#page-17-2) implies, again in view of the pointwise bounds, that $V(\mathfrak{X}u) \in HW_{\text{loc}}^{1,2}(\tilde{\Omega})$. Thus, for B_{ρ}, B_{R} and η as defined earlier, we have $\eta(V(\mathfrak{X}u) - V(\xi)) \in HW_0^{1,2} (B_R; \mathbb{R}^{2n})$ for any $\xi \in \mathbb{R}^{2n}$. Thus we have

$$
\int_{B_{\rho}} |V(\mathfrak{X}u) - V(\xi)|^{\frac{2Q}{Q-2}} \n\leq \int_{B_{R}} \eta^{\frac{2Q}{Q-2}} |V(\mathfrak{X}u) - V(\xi)|^{\frac{2Q}{Q-2}} \n\leq C \left(\int_{B_{R}} |\mathfrak{X}[\eta(V(\mathfrak{X}u) - V(\xi))]|^{2} \right)^{\frac{Q}{Q-2}} \n= C \left(\int_{B_{R}} \eta^{2} |\mathfrak{X}[V(\mathfrak{X}u)]|^{2} + \int_{B_{R}} |\langle \mathfrak{X}\eta, V(\mathfrak{X}u) - V(\xi) \rangle|^{2} \right)^{\frac{Q}{Q-2}} \n\leq C \left(\int_{B_{R}} \eta^{2} |\mathfrak{X}u|^{p-2} |\mathfrak{X}\mathfrak{X}u|^{2} + \int_{B_{R}} |\langle \mathfrak{X}\eta, V(\mathfrak{X}u) - V(\xi) \rangle|^{2} \right)^{\frac{Q}{Q-2}} \n\leq C \left(\lambda^{p-2} \int_{B_{R}} \eta^{2} |\mathfrak{X}\mathfrak{X}u|^{2} + \frac{1}{(R-\rho)^{2}} \int_{B_{R}} |V(\mathfrak{X}u) - V(\xi)|^{2} \right)^{\frac{Q}{Q-2}} \n\leq \frac{C}{(R-\rho)^{\frac{2Q}{Q-2}}} \left(\int_{B_{R}} \lambda^{p-2} |\mathfrak{X}u - \xi|^{2} + \int_{B_{R}} |V(\mathfrak{X}u) - V(\xi)|^{2} \right)^{\frac{Q}{Q-2}}.
$$

Now in view of [\(68\)](#page-22-1) and [\(69\)](#page-22-0), we deduce

$$
\lambda^{p-2} |\mathfrak{X}u - \xi|^2 \le \max \left\{ (4B)^{p-2}, (2A)^{2-p} \right\} (|\mathfrak{X}u| + |\xi|)^{p-2} |\mathfrak{X}u - \xi|^2
$$

$$
\stackrel{(36)}{\le} c_V^2 \max \left\{ (4B)^{p-2}, (2A)^{2-p} \right\} |V(\mathfrak{X}u) - V(\xi)|^2.
$$

Using this in the last estimate, we arrive at

$$
\int_{B_{\rho}}\left|V\left(\mathfrak{X} u\right)-V\left(\xi\right)\right|^{\frac{2Q}{Q-2}}\leq C\left(R-\rho\right)^{-\frac{2Q}{Q-2}}\left(\int_{B_{R}}\left|V\left(\mathfrak{X} u\right)-V\left(\xi\right)\right|^2\right)^{\frac{Q}{Q-2}}.
$$

The desired estimates now follow from this in the same way as in Lemma [35.](#page-21-3) \Box

3.4. Estimates in the nondegenerate regime. Some of the Lemmas we would prove in this subsection are similar in spirit to some Lemmas in [\[6\]](#page-63-10) (compare Lemma [37,](#page-24-0) [39](#page-26-0) and [40](#page-28-0) with Lemma 4.3, 4.4 and 4.5 in [\[6\]](#page-63-10) respectively). However, our results are valid only the nondegenerate regime.

Lemma 37. Let $1 < p < \infty$. Suppose $u \in HW^{1,p}_{loc}(B_{2R})$ be a weak solution of

$$
\operatorname{div}_{\mathbb{H}}\left(\left|\mathfrak{X} u\right|^{p-2} \mathfrak{X} u\right) = 0 \qquad \qquad \text{in } B_{2R}.
$$

Let $A, B \geq 1$ and $\lambda > 0$ be constants and suppose

(72)
$$
\frac{\lambda}{4B} \leq |\mathfrak{X}u| \leq A\lambda \quad in \ B_R.
$$

Let $\xi \in \mathbb{R}^{2n}$ be any vector satisfying

$$
\frac{\lambda}{4B} \le |\xi| \le A\lambda.
$$

Let $v \in HW^{1,p}(B_{R/2})$ be the unique weak solution of the Dirichlet boundary value problem for the linear equation

$$
\begin{cases} \operatorname{div}_{\mathbb{H}} \left[\left| \xi \right|^{p-2} \mathfrak{X} v + (p-2) \left| \xi \right|^{p-4} \langle \xi, \mathfrak{X} v \rangle \xi \right] = 0 & \text{in } B_{R/2}, \\ v = u & \text{on } \partial B_{R/2}. \end{cases}
$$

Then there exists a constant $C \equiv C(Q, p, A, B) \ge 1$ such that we have the estimate

(73)
$$
\int_{B_{R/2}} |\mathfrak{X} u - \mathfrak{X} v|^2 \leq C \left(\frac{1}{A^2 \lambda^2} \int_{B_R} |\mathfrak{X} u - \xi|^2 \right)^{\frac{2}{Q-2}} \int_{B_R} |\mathfrak{X} u - \xi|^2.
$$

Proof. From the equation satisfied by u , we can write

$$
\begin{split}\n\text{div}_{\mathbb{H}} \left[|\xi|^{p-2} \mathfrak{X} u + (p-2) |\xi|^{p-4} \langle \xi, \mathfrak{X} u \rangle \xi \right] \\
&= \text{div}_{\mathbb{H}} \left[|\xi|^{p-2} \mathfrak{X} u + (p-2) |\xi|^{p-4} \langle \xi, \mathfrak{X} u \rangle \xi - |\mathfrak{X} u|^{p-2} \mathfrak{X} u \right] \\
&= \text{div}_{\mathbb{H}} \left[|\xi|^{p-2} \xi - |\mathfrak{X} u|^{p-2} \mathfrak{X} u + |\xi|^{p-2} \left(\mathfrak{X} u - \xi \right) + (p-2) |\xi|^{p-4} \langle \xi, \mathfrak{X} u - \xi \rangle \xi \right] \\
&\quad + (p-2) \text{div}_{\mathbb{H}} \left(|\xi|^{p-2} \xi \right) \\
&= \text{div}_{\mathbb{H}} \left[|\xi|^{p-2} \xi - |\mathfrak{X} u|^{p-2} \mathfrak{X} u + |\xi|^{p-2} \left(\mathfrak{X} u - \xi \right) + (p-2) |\xi|^{p-4} \langle \xi, \mathfrak{X} u - \xi \rangle \xi \right] \\
&:= \text{div}_{\mathbb{H}} F.\n\end{split}
$$

Setting $w := u - v \in HW_0^{1,p}(B_{R/2})$, we see that this implies

$$
\operatorname{div}_{\mathbb{H}}\left[\left|\xi\right|^{p-2} \mathfrak{X} w + (p-2)\left|\xi\right|^{p-4} \left\langle \xi, \mathfrak{X} w \right\rangle \xi\right] = \operatorname{div}_{\mathbb{H}} F \qquad \text{in } B_{R/2}.
$$

Plugging w as a test function, we deduce

$$
|\xi|^{p-2} \int_{B_{R/2}} |\mathfrak{X} w|^2 + (p-2) |\xi|^{p-4} \int_{B_{R/2}} |\langle \xi, \mathfrak{X} w \rangle|^2 = \int_{B_{R/2}} \langle F, \mathfrak{X} w \rangle.
$$

Now, if $p \geq 2$, clearly we have

$$
|\xi|^{p-2} \int_{B_{R/2}} |\mathfrak{X}w|^2 \leq |\xi|^{p-2} \int_{B_{R/2}} |\mathfrak{X}w|^2 + (p-2) |\xi|^{p-4} \int_{B_{R/2}} |\langle \xi, \mathfrak{X}w \rangle|^2.
$$

On the other hand, if $1 < p < 2$, then we have

$$
(p-2) |\xi|^{p-4} \int_{B_{R/2}} |\langle \xi, \mathfrak{X} w \rangle|^2 \ge (p-2) |\xi|^{p-2} \int_{B_{R/2}} |\mathfrak{X} w|^2
$$

.

Hence in this case, we have

$$
(p-1)|\xi|^{p-2}\int_{B_{R/2}}|\mathfrak{X}w|^2\leq|\xi|^{p-2}\int_{B_{R/2}}|\mathfrak{X}w|^2+(p-2)|\xi|^{p-4}\int_{B_{R/2}}|\langle \xi,\mathfrak{X}w\rangle|^2.
$$

Thus, in either case, we have,

$$
\int_{B_{R/2}} |\mathfrak{X}w|^2 \leq \frac{C}{\lambda^{p-2}} \min\left\{1, p-1\right\} |\xi|^{p-2} \int_{B_{R/2}} |\mathfrak{X}w|^2
$$
\n
$$
\leq \frac{C}{\lambda^{p-2}} \left[|\xi|^{p-2} \int_{B_{R/2}} |\mathfrak{X}w|^2 + (p-2) |\xi|^{p-4} \int_{B_{R/2}} |\langle \xi, \mathfrak{X}w \rangle|^2 \right]
$$
\n
$$
\leq \frac{C}{\lambda^{p-2}} \int_{B_{R/2}} \langle F, \mathfrak{X}w \rangle
$$
\n
$$
\leq \frac{C}{\lambda^{p-2}} \int_{B_{R/2}} |F| |\mathfrak{X}w| \leq \delta^2 \frac{C}{\lambda^{p-2}} \int_{B_{R/2}} |\mathfrak{X}w|^2 + \frac{1}{4\delta^2} \frac{C}{\lambda^{p-2}} \int_{B_{R/2}} |F|^2,
$$

where C depends on A, B, p. Choosing $\delta > 0$ such that $C \delta^2 \lambda^{2-p} = 1/2$, we deduce

(74)
$$
\int_{B_{R/2}} |\mathfrak{X}w|^2 \leq \frac{C}{\lambda^{2p-4}} \int_{B_{R/2}} |F|^2.
$$

Now, from the definition of F , we have (see Lemma 4.3. in [\[6\]](#page-63-10)),

(75)
$$
|F| \leq \frac{c(A,B)}{\lambda} (|\mathfrak{X}u| + |\xi|)^{p-2} |\mathfrak{X}u - \xi|^2.
$$

Thus, we have, with constants $c \equiv c(A, B) > 0$,

(76)
$$
\int_{B_{R/2}} |F|^2 \le c(A, B) \lambda^{2p-6} \int_{B_{R/2}} |\mathfrak{X} u - \xi|^4
$$

$$
\le c\lambda^{2p-2-\frac{2Q}{Q-2}} \int_{B_{R/2}} |\mathfrak{X} u - \xi|^{\frac{2Q}{Q-2}}.
$$

Now, applying [\(65\)](#page-21-0), we have

$$
\int_{B_{R/2}} |\mathfrak{X} u - \xi|^{2Q \over Q-2} \leq C_{RH}^{2Q \over Q-2} \left(\int_{B_R} |\mathfrak{X} u - \xi|^2 \right)^{Q \over Q-2}.
$$

Using this in [\(76\)](#page-25-0), we conclude that there exists a constant $C \equiv C(A, B, n, p) > 0$ such that

$$
\int_{B_{R/2}} |F|^2 \le C\lambda^{2p-2-\frac{2Q}{Q-2}} \left(\int_{B_R} |\mathfrak{X} u - \xi|^2\right)^{\frac{2}{Q-2}} \int_{B_R} |\mathfrak{X} u - \xi|^2.
$$

Combining this with [\(74\)](#page-25-1), we have

$$
\int_{B_{R/2}} |\mathfrak{X}w|^2 \le C\lambda \frac{[(2p-2-\frac{2Q}{Q-2})+(4-2p)]}{\left(\int_{B_R} |\mathfrak{X}u-\xi|^2\right)^{\frac{2}{Q-2}}} \int_{B_R} |\mathfrak{X}u-\xi|^2
$$

$$
\le C\lambda^{-\frac{4}{Q-2}} \left(\int_{B_R} |\mathfrak{X}u-\xi|^2\right)^{\frac{2}{Q-2}} \int_{B_R} |\mathfrak{X}u-\xi|^2.
$$

This implies [\(73\)](#page-24-1) and completes the proof. \square

Lemma 38. Let $1 < p < \infty$, $A, B \ge 1$, and $\lambda > 0$ be constants and let $\xi \in \mathbb{R}^{2n}$ be any vector satisfying

$$
\frac{\lambda}{4B} \le |\xi| \le A\lambda.
$$

Let $v \in HW^{1,p}(B_{R/2})$ be a weak solution of the linear equation

$$
\operatorname{div}_{\mathbb{H}}\left[\left|\xi\right|^{p-2} \mathfrak{X} v + (p-2)\left|\xi\right|^{p-4} \left\langle \xi, \mathfrak{X} v \right\rangle \xi\right] = 0 \qquad \text{in } B_{R/2}.
$$

Then there exists a constant $C \equiv C(Q, p, A, B) > 0$ such that for any $0 < \rho \le R/2$, we have the estimate

(77)
$$
\int_{B_{\rho}} \left| xv - (xv)_{\rho} \right|^2 \leq C \left(\frac{\rho}{R} \right)^{Q+2} \int_{B_{R/2}} \left| xv - (xv)_{R/2} \right|^2.
$$

Proof. The proof can be found in Theorem 3.2 in Xu-Zuily [\[48\]](#page-64-17). As is well known, the constant in the estimate, depends only upon the ellipticity ratio, which in this case is A/B , i.e. independent of λ .

Lemma 39. Let $1 < p < \infty$. Suppose $u \in HW^{1,p}_{loc}(B_{2R})$ be a weak solution of

$$
\operatorname{div}_{\mathbb{H}}\left(\left|\mathfrak{X} u\right|^{p-2} \mathfrak{X} u\right) = 0 \qquad \qquad \text{in } B_{2R}.
$$

Let $A, B \geq 1$ be constants and suppose

$$
\frac{\lambda}{4B} \le |\mathfrak{X}u| \le A\lambda \qquad \text{in } B_R.
$$

Then for any $\delta_0 > 0$ with $64A^2B^2\delta_0 < 1$, there exist $\varepsilon, \tau \in (0,1)$, depending only on Q , p , A , B , and δ_0 , satisfying

(78)
$$
0 < \frac{\log \delta_0}{2 \log \tau} < 1,
$$

such that if $\xi^0 \in \mathbb{R}^{2n}$ is a vector satisfying

(i) $\frac{\lambda}{4R} \leq |\xi^0| \leq A\lambda$, (ii) 1 B_R $|\mathfrak{X}u-\xi^0|$ $2^2 \leq \varepsilon A^2 \lambda^2$,

then there exists a vector $\xi^1 \in \mathbb{R}^{2n}$ satisfying

(79)
$$
\frac{\lambda}{8B} \le |\xi^1| \le 2A\lambda,
$$

(80)
$$
\int_{B_{\tau R}} |\mathfrak{X} u - \xi^1|^2 \leq \varepsilon A^2 \lambda^2,
$$

and

(81)
$$
\int_{B_{\tau R}} |\mathfrak{X} u - \xi^1|^2 \leq \delta_0 \tau^Q \int_{B_R} |\mathfrak{X} u - \xi^0|^2.
$$

Proof. We apply Lemma [37](#page-24-0) with the choice $\xi = \xi^0$ to deduce

(82)
$$
\int_{B_{R/2}} |\mathfrak{X} u - \mathfrak{X} v|^2 \leq C_1 \varepsilon^{\frac{2}{Q-2}} \int_{B_R} |\mathfrak{X} u - \xi_0|^2
$$

for some constant $C_1 \equiv C_1(Q, p, A, B) > 1$. Now we estimate

(83)
$$
\int_{B_{R/2}} |\mathfrak{X}v - \xi^{0}|^{2} \le 2 \int_{B_{R/2}} |\mathfrak{X}u - \xi^{0}|^{2} + 2 \int_{B_{R/2}} |\mathfrak{X}u - \mathfrak{X}v|^{2} \le C_{2} \left(1 + \varepsilon^{\frac{2}{Q-2}}\right) \int_{B_{R}} |\mathfrak{X}u - \xi^{0}|^{2},
$$

for some constant $C_2 \equiv C_2(Q, p, A, B) > 1$. Now, by Lemma [38,](#page-26-2) for any $0 < \rho <$ $R/2$, we have

(84)
$$
\int_{B_{\rho}} \left| \mathfrak{X}v - (\mathfrak{X}v)_{\rho} \right|^{2} \leq C_{3} \left(\frac{\rho}{R} \right)^{Q+2} \int_{B_{R/2}} \left| \mathfrak{X}v - (\mathfrak{X}v)_{R/2} \right|^{2} \leq C_{3} \left(\frac{\rho}{R} \right)^{Q+2} \int_{B_{R/2}} \left| \mathfrak{X}v - \xi^{0} \right|^{2},
$$

for some constant $C_3 \equiv C_3(Q, p, A, B) > 1$. Hence, setting $\rho = \tau R/2$ and

$$
\xi^1 := (\mathfrak{X}v)_{\rho} = (\mathfrak{X}v)_{\tau R/2},
$$

we have

$$
\int_{B_{\tau R/2}} |\mathfrak{X}u - \xi^1|^2 \le 2 \int_{B_{\tau R/2}} |\mathfrak{X}v - \xi^1|^2 + 2 \int_{B_{\tau R/2}} |\mathfrak{X}u - \mathfrak{X}v|^2
$$
\n
$$
\stackrel{(84)}{\le} 2C_3 \tau^{Q+2} \int_{B_{R/2}} |\mathfrak{X}v - \xi^0|^2 + 2 \int_{B_{R/2}} |\mathfrak{X}u - \mathfrak{X}v|^2
$$
\n
$$
\stackrel{(82),(83)}{\le} C_4 \left[\tau^{Q+2} \left(1 + \varepsilon^{\frac{2}{Q-2}} \right) + \varepsilon^{\frac{2}{Q-2}} \right] \int_{B_R} |\mathfrak{X}u - \xi^0|^2
$$
\n
$$
\le C_4 \left[\tau^{Q+2} + \varepsilon^{\frac{2}{Q-2}} \right] \int_{B_R} |\mathfrak{X}u - \xi^0|^2.
$$

Here, again $C_4 \equiv C_4(Q, p, A, B) > 1$ is a constant. Now we choose $\varepsilon > 0$ sufficiently small (depending on τ) such that we have

$$
(85) \qquad \qquad \varepsilon^{\frac{2}{Q-2}} = \tau^{Q+2}.
$$

This choice implies that

(86)
$$
\int_{B_{\tau R/2}} |\mathfrak{X} u - \xi^1|^2 \leq C_M \tau^{Q+2} \int_{B_R} |\mathfrak{X} u - \xi^0|^2,
$$

where

(87)
$$
C_M := \max\{C_1, C_2, C_3, 2C_4\}.
$$

Now we choose $\tau > 0$ small enough to have

$$
(88) \t\t 2^{Q+2}C_M\tau^2 \le \delta_0.
$$

Note that this implies (78) . This proves (81) . Assumption (ii) and (81) implies [\(80\)](#page-26-5). Now observe that since $Q \geq 4$, we have

$$
\frac{(Q-2)(Q+2)}{2} - Q = \frac{Q^2 - 2Q - 4}{2} = \frac{(Q-1)^2 - 5}{2} \ge 2.
$$

Thus, we deduce

(89)
$$
2^{Q+2}\tau^{\frac{(Q-2)(Q+2)}{2}-Q} \leq 2^{Q+2}\tau^2 \leq 2^{Q+2}C_M\tau^2 \leq \delta_0.
$$

Using this, we deduce

$$
|\xi^{1} - \xi^{0}|^{2} \leq \int_{B_{\tau R/2}} |\mathfrak{X}v - \xi^{0}|^{2}
$$

\n
$$
\leq 2 \int_{B_{\tau R/2}} |\mathfrak{X}u - \xi^{0}|^{2} + 2 \int_{B_{\tau R/2}} |\mathfrak{X}u - \mathfrak{X}v|^{2}
$$

\n
$$
\leq 2^{Q+1}\tau^{-Q} \int_{B_{R}} |\mathfrak{X}u - \xi^{0}|^{2} + 2\tau^{-Q} \int_{B_{R/2}} |\mathfrak{X}u - \mathfrak{X}v|^{2}
$$

\n(90)
\n
$$
\leq (82), (85), (88)
$$

\n
$$
\leq (2^{Q+1}\tau^{-Q} + \delta_{0}) \int_{B_{R}} |\mathfrak{X}u - \xi^{0}|^{2}
$$

\n
$$
\leq 2^{Q+2}\tau^{-Q}\varepsilon A^{2} \lambda^{2} \leq 2^{Q+2}\tau^{\frac{(Q-2)(Q+2)}{2} - Q} A^{2} \lambda^{2} \leq \delta_{0} A^{2} \lambda^{2}
$$

Hence, we have

$$
\left|\xi^1 - \xi^0\right|^2 \le \delta_0 A^2 \lambda^2
$$

Then we have

$$
\left(\frac{1}{4B} - \sqrt{\delta_0}A\right)\lambda \le |\xi^0| - |\xi^1 - \xi^0| \le |\xi^1| \le |\xi^0| + |\xi^1 - \xi^0| \le \left(1 + \sqrt{\delta_0}\right)A\lambda.
$$

Since $64A^2B^2\delta_0 < 1$, we have

$$
\left(\frac{1}{4B} - \sqrt{\delta_0}A\right)\lambda > \frac{1}{B}\left(\frac{1}{4} - \frac{1}{8}\right)\lambda > \frac{\lambda}{8B} \quad \text{and}
$$

$$
\left(1 + \sqrt{\delta_0}\right)A\lambda \le \left(1 + \frac{1}{8AB}\right)A\lambda \le \left(1 + \frac{1}{8}\right)A\lambda \le 2A\lambda.
$$

This proves (79) and completes the proof.

Lemma 40. Let $1 < p < \infty$. Suppose $u \in HW^{1,p}_{loc}(B_{2R})$ be a weak solution of

$$
\operatorname{div}_{\mathbb{H}}\left(|\mathfrak{X}u|^{p-2}\mathfrak{X}u\right)=0\qquad \qquad \text{in }B_{2R}.
$$

Let $A, B \geq 1$ be constants and suppose

$$
\frac{\lambda}{4B} \le |\mathfrak{X}u| \le A\lambda \qquad \text{in } B_R.
$$

Then there exist constants $\delta_0, \varepsilon, \tau \in (0,1)$, depending only on Q, p, A, B, satisfying

(91)
$$
0 < \frac{\log \delta_0}{2 \log \tau} < 1,
$$

such that if $\xi^0 \in \mathbb{R}^{2n}$ is a vector satisfying

(i)
$$
\frac{\lambda}{4B} \le |\xi^0| \le A\lambda
$$
,
(ii) $\oint_{B_R} |\mathfrak{X}u - \xi^0|^2 \le \varepsilon A^2 \lambda^2$,

then there exists a sequence of vectors $\{\xi^s\}_{s\in\mathbb{N}}$ in \mathbb{R}^{2n} such that for all integers $s \geq 0$, we have

(92)
$$
\frac{\lambda}{8B} \le |\xi^s| \le 2A\lambda,
$$

.

30 A. MALLICK AND S. SIL

(93)
$$
\int_{B_{\tau^s R}} |\mathfrak{X} u - \xi^s|^2 \leq \varepsilon A^2 \lambda^2,
$$

and

(94)
$$
\int_{B_{\tau^{s+1}R}} |\mathfrak{X} u - \xi^{s+1}|^2 \leq \delta_0 \tau^Q \int_{B_{\tau^{s}R}} |\mathfrak{X} u - \xi^s|^2.
$$

Proof. We plan to construct the sequence inductively using Lemma [39.](#page-26-0) To do this, we only need to show that it is possible to choose $\delta_0 > 0$ such that given ξ^s , the vector ξ_{s+1} constructed by Lemma [39](#page-26-0) also satisfies the weaker bounds

$$
\frac{\lambda}{8B} \le |\xi^{s+1}| \le 2A\lambda,
$$

instead of the significantly worse bounds

$$
\frac{\lambda}{8^s B} \le \left| \xi^{s+1} \right| \le 2^s A \lambda.
$$

This is easy to achieve. Indeed, recalling the estimates in Lemma [39,](#page-26-0) particularly [\(90\)](#page-28-1), we deduce

$$
\begin{split} \left|\xi^{s+1} - \xi^s\right|^2 &\le 2^{Q+2}\tau^{-Q} \oint_{B_{\tau^s R}} |\mathfrak{X} u - \xi^s|^2 \\ &\le 2^{Q+2}\tau^{-Q} \delta_0^s \oint_{B_R} \left|\mathfrak{X} u - \xi^0\right|^2 \\ &\le 2^{Q+2}\tau^{-Q} \delta_0^s \varepsilon A^2 \lambda^2 \stackrel{(85)}{\le} 2^{Q+2}\tau \frac{(Q+2)(Q-2)}{2} - Q \delta_0^s A^2 \lambda^2 \stackrel{(89)}{\le} \delta_0^{s+1} A^2 \lambda^2. \end{split}
$$

Taking square root and summing over s, we deduce

$$
\left|\xi^{s+1} - \xi^0\right| \le \left(\sum_{i=1}^\infty \delta_0^{\frac{i}{2}}\right) A\lambda.
$$

 \Box

Now, [\(92\)](#page-28-2) is satisfied for all $s \ge 0$ with the choice $\delta_0 = 1/(1 + 8AB)^2$

Now we are ready to prove our main lemma.

Lemma 41. Let $1 < p < \infty$. Suppose $u \in HW^{1,p}_{loc}(B_{2R})$ be a weak solution of $\operatorname{div}_{\mathbb{H}}\left(|\mathfrak{X} u|^{p-2}\mathfrak{X} u\right)$ in B_{2R} .

Let $A, B \geq 1$ be constants and suppose

(95)
$$
\frac{\lambda}{4B} \leq |\mathfrak{X}u| \leq A\lambda \quad in B_R.
$$

Then there exist constants $\varepsilon_0, \alpha_0 \in (0,1)$ and $C_0 \geq 1$, all depending only on Q , p, A and B such that if

(96)
$$
\frac{\lambda}{8B} \le |(\mathfrak{X}u)_{\sigma R}| \le A\lambda
$$

and

(97)
$$
\int_{B_{\sigma R}} |\mathfrak{X}u - (\mathfrak{X}u)_{\sigma R}|^2 \leq \varepsilon_0 A^2 \lambda^2
$$

hold for some constant $0 < \sigma \leq 1$, then for any $0 < \rho < \sigma R$, we have

$$
(98) \qquad \int_{B_{\rho}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\rho} \right|^{2} \leq C_{0} \sigma^{-Q-2\alpha_{0}} \left(\frac{\rho}{R} \right)^{2\alpha_{0}} \int_{B_{R}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{R} \right|^{2}.
$$

Proof. Set

$$
\xi^0 := (\mathfrak{X} u)_{\sigma R} \, .
$$

Now note that [\(95\)](#page-29-0) clearly implies

(99)
$$
\frac{\lambda}{8B} \le \frac{\lambda}{4B} \le |\mathfrak{X}u| \le A\lambda \quad \text{in } B_{\sigma R}.
$$

By [\(96\)](#page-29-1) we also have

(100)
$$
\frac{\lambda}{8B} \le |(\mathfrak{X}u)_{\sigma R}| \le A\lambda.
$$

Hence, [\(99\)](#page-30-0), [\(100\)](#page-30-1) and [\(97\)](#page-29-2) implies that the hypotheses of Lemma [40](#page-28-0) are satisfied for our choice of ξ^0 in $B_{\sigma R}$ with the same A and B being replaced by 2B, if $\varepsilon_0 = \varepsilon$, where $\varepsilon \in (0, 1)$ is the constant given by Lemma [40.](#page-28-0) Thus, there exists $\delta_0, \tau \in (0, 1)$, both depending only on Q , p , A and B such that there exists a sequence of vectors ${\{\xi^s\}}_{s\in\mathbb{N}}$ such that for all integers $s\geq 0$, we have $\frac{\lambda}{16B}\leq |\xi^s|\leq 4A\lambda$ and

$$
\int_{B_{\tau^{s+1}\sigma R}} \left| \mathfrak{X} u - \xi^{s+1} \right|^2 \leq (\delta_0)^{s+1} \int_{B_{\sigma R}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\sigma R} \right|^2.
$$

Now, let $0 < \rho < \sigma R$. Then there exists an integer $s_0 \geq 0$ such that

$$
\tau^{s_0+1}\sigma R < \rho \le \tau^{s_0}\sigma R.
$$

Suppose $s_0 \geq 1$. Then we have

$$
\int_{B_{\rho}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\rho} \right|^{2} \leq \int_{B_{\rho}} |\mathfrak{X} u - \xi^{s_{0}}|^{2}
$$
\n
$$
\leq \left(\frac{\tau^{s_{0}} \sigma R}{\rho} \right)^{Q} \int_{B_{\tau^{s_{0}} \sigma R}} |\mathfrak{X} u - \xi^{s_{0}}|^{2}
$$
\n
$$
\leq \left(\frac{\tau^{s_{0}} \sigma R}{\rho} \right)^{Q} (\delta_{0})^{s_{0}} \int_{B_{\sigma R}} |\mathfrak{X} u - (\mathfrak{X} u)_{\sigma R}|^{2}
$$
\n
$$
\leq \tau^{-Q} (\delta_{0})^{s_{0}} \int_{B_{\sigma R}} |\mathfrak{X} u - (\mathfrak{X} u)_{\sigma R}|^{2}.
$$

Now set

$$
\alpha_0 := \frac{\log \delta_0}{2 \log \tau} \in (0, 1), \text{ in view of (91)}.
$$

Now, we deduce

$$
\int_{B_{\rho}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\rho} \right|^{2} \leq \tau^{-Q} \left(\delta_{0} \right)^{s_{0}} \left(\frac{\sigma R}{\rho} \right)^{2\alpha_{0}} \left(\frac{\rho}{\sigma R} \right)^{2\alpha_{0}} \int_{B_{\sigma R}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\sigma R} \right|^{2}
$$
\n
$$
\leq \tau^{-Q} \left(\delta_{0} \right)^{s_{0}} \left(\delta_{0} \right)^{-s_{0}-1} \left(\frac{\rho}{\sigma R} \right)^{2\alpha_{0}} \int_{B_{\sigma R}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\sigma R} \right|^{2}
$$
\n
$$
\leq \tau^{-Q} \delta_{0}^{-1} \left(\frac{\rho}{\sigma R} \right)^{2\alpha_{0}} \int_{B_{\sigma R}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{R} \right|^{2}
$$
\n
$$
\leq \tau^{-Q} \delta_{0}^{-1} \sigma^{-Q-2\alpha_{0}} \left(\frac{\rho}{R} \right)^{2\alpha_{0}} \int_{B_{R}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{R} \right|^{2}.
$$

Thus, we conclude by setting $C = C(Q, p, A, B) := \tau^{-Q} \delta_0^{-1}$. If $s_0 = 0$, then for the same choice of α_0 as above, we clearly have,

$$
\int_{B_{\rho}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\rho} \right|^{2} \leq \int_{B_{\rho}} |\mathfrak{X} u - (\mathfrak{X} u)_{\sigma R}|^{2}
$$
\n
$$
\leq \left(\frac{\sigma R}{\rho} \right)^{Q-2\alpha_{0}} \left(\frac{\rho}{\sigma R} \right)^{2\alpha_{0}} \int_{B_{\sigma R}} |\mathfrak{X} u - (\mathfrak{X} u)_{\sigma R}|^{2}
$$
\n
$$
\leq \tau^{2\alpha_{0} - Q} \sigma^{-Q-2\alpha_{0}} \left(\frac{\rho}{R} \right)^{2\alpha_{0}} \int_{B_{R}} |\mathfrak{X} u - (\mathfrak{X} u)_{R}|^{2}.
$$

Thus, we have the estimate in this case as well for the same

$$
C = C(Q, p, A, B) := \tau^{-Q} \delta_0^{-1} = \tau^{2\alpha_0 - Q}.
$$

This completes the proof.

We also have the following result.

Lemma 42. Let
$$
1 < p < \infty
$$
. Suppose $u \in HW_{loc}^{1,p}(B_{2R})$ be a weak solution of

$$
\operatorname{div}_{\mathbb{H}}\left(\left|\mathfrak{X}u\right|^{p-2}\mathfrak{X}u\right)=0\qquad \qquad \text{in }B_{2R}.
$$

Let $A, B \geq 1$ be constants and suppose

(101)
$$
\frac{\lambda}{4B} \le |V(\mathfrak{X}u)|^{\frac{2}{p}} \le A\lambda \quad in B_R.
$$

Then there exist constants $\theta_0, \alpha_0 \in (0, 1)$ and $C_1 \geq 1$, all depending only on Q, p, A and B, such that if

(102)
$$
\frac{\lambda}{8B} \le |(V(\mathfrak{X}u))_{\sigma R}|^{\frac{2}{p}} \le A\lambda
$$

and

(103)
$$
\oint_{B_{\sigma R}} |V(\mathfrak{X}u) - (V(\mathfrak{X}u))_{\sigma R}|^2 \leq \theta_0 A^p \lambda^p,
$$

hold for some constant $0 < \sigma \leq 1$, then for any $0 < \rho < \sigma R$, we have (104)

$$
\int_{B_{\rho}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_\rho\right|^2\leq C_1\sigma^{-Q-2\alpha_0}\left(\frac{\rho}{R}\right)^{2\alpha_0}\int_{B_R}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_R\right|^2.
$$

Proof. Set

(105)
$$
\xi^{0} := V^{-1} ((V (\mathfrak{X} u))_{\sigma R}).
$$

Observe that this implies

$$
\left|\xi^0\right|^{\frac{p}{2}} = \left|\left(V\left(\mathfrak{X}u\right)\right)_{\sigma R}\right|.
$$

Thus, clearly, we have the bounds

$$
\frac{\lambda}{8B} \le |\xi^0| \le A\lambda.
$$

As $|V(\mathfrak{X}u)|^{\frac{2}{p}} = |\mathfrak{X}u|$, we also have the obvious bounds

$$
\frac{\lambda}{8B} \le \frac{\lambda}{4B} \le |\mathfrak{X}u| \le A\lambda \quad \text{in } B_{\sigma R}.
$$

$$
\Box
$$

Now, using [\(36\)](#page-14-0), we have

$$
\int_{B_{\sigma R}} \left| \mathfrak{X} u - \xi^0 \right|^2 \le c_0 \int_{B_{\sigma R}} \left| V \left(\mathfrak{X} u \right) - V \left(\xi^0 \right) \right|^2 \le c_0 \theta_0 A^2 \lambda^2,
$$

for a constant $c_0 = c_0 (Q, p, A, B) > 0$. We set

$$
\theta_0 = \frac{\varepsilon}{c_0},
$$

where ε is the constant given by Lemma [40](#page-28-0) for our choice of ξ^0 in $B_{\sigma R}$ with the same A and B being replaced by 2B. Thus, there exists $\delta_0, \tau \in (0,1)$, both depending only on Q, p, A and B such that there exists a sequence of vectors $\{\xi^s\}_{s\in\mathbb{N}}$ such that for all integers $s \geq 0$, we have

(106)
$$
\frac{\lambda}{16B} \le |\xi^s| \le 2A\lambda
$$

and

$$
\oint_{B_{\tau^{s+1}\sigma R}} |\mathfrak{X} u - \xi^{s+1}|^2 \leq (\delta_0)^{s+1} \int_{B_{\sigma R}} |\mathfrak{X} u - \xi^0|^2.
$$

This implies, again by [\(36\)](#page-14-0), for some constant $C \equiv C(Q, p, A, B) > 0$ we have

$$
\int_{B_{\tau^{s+1}\sigma R}} |V(\mathfrak{X}u) - V(\xi^{s+1})|^2 \le C(\lambda)^{2-p} \int_{B_{\tau^{s+1}\sigma R}} |\mathfrak{X}u - \xi^{s+1}|^2
$$

\n
$$
\le C(\delta_0)^{s+1} \int_{B_{\sigma R}} \lambda^{2-p} |\mathfrak{X}u - \xi^0|^2
$$

\n
$$
\le C(\delta_0)^{s+1} \int_{B_{\sigma R}} |V(\mathfrak{X}u) - V(\xi^0)|^2.
$$

Since $V(\xi^0) = (V(\mathfrak{X}u))_{\sigma R}$, we arrive at

$$
\int_{B_{\tau^{s+1}\sigma R}} \left| V\left(\mathfrak{X} u\right) - V\left(\xi^{s+1}\right) \right|^2 \leq C \left(\delta_0\right)^{s+1} \int_{B_{\sigma R}} \left| V\left(\mathfrak{X} u\right) - \left(V\left(\mathfrak{X} u\right)\right)_{\sigma R} \right|^2.
$$

From this, via an interpolation argument similar to the one used above in Lemma [41](#page-29-3) , we have

$$
\int_{B_{\rho}} \left| V(\mathfrak{X} u) - V(\xi^{s+1}) \right|^2 \leq C\sigma^{-Q-2\alpha_0} \left(\frac{\rho}{R} \right)^{2\alpha_0} \int_{B_R} \left| V(\mathfrak{X} u) - (V(\mathfrak{X} u))_R \right|^2.
$$
\nThis completes the proof.

3.5. Proofs of Theorem [1](#page-3-0) and Theorem [2.](#page-3-2) We now prove our crucial excess decay estimates.

Proof. (of Theorem [1](#page-3-0)) We can and henceforth assume $x_0 = 0$. Note that, it is enough to prove [\(6\)](#page-3-3), for $0 < \rho < \kappa R$, for some constant $\kappa \in (0,1)$ depending only n and p, as the estimate is trivial otherwise. First, we fix $C_{RH} > 0$ which is given by Lemma [35](#page-21-3) with the choice $A = B = 1$. So, clearly C_{RH} depends only on n and p. Next, let $B = B(n, p) \ge 1$ be defined by $B := 2c_1$, where c_1 is the constant given by Theorem [30](#page-15-7) with the choice $q = 1$. Let $\varepsilon_0, \alpha_0 \in (0, 1)$ be the constants given by Lemma [41](#page-29-3) for the choices of $A = 2C_{RH}$ and $B = 2c_1$. Let $\alpha_1 \in (0,1)$ be the exponent in Theorem [31.](#page-15-2) Set

$$
\alpha := \min \left\{ \alpha_0, \alpha_1 \right\}.
$$

Now we claim the following.

Claim 43. Without loss of generality, we can assume that the we have

(107)
$$
\frac{1}{B} \|\mathfrak{X}u\|_{\infty, R/2} \leq |(\mathfrak{X}u)_R| \leq 4 \|\mathfrak{X}u\|_{\infty, R/2}
$$

and

(108)
$$
\left(\int_{B_{R/2}}|\mathfrak{X} u - (\mathfrak{X} u)_R|^q\right)^{\frac{1}{q}} \leq \varepsilon_0 \left\|\mathfrak{X} u\right\|_{\infty, R/2}.
$$

Proof of Claim [43:](#page-33-0) Indeed, if (108) is violated, then by Theorem [31,](#page-15-2) for any $0 < \rho < R/2$, we have

$$
\left(\oint_{B_{\rho}}\left|\mathfrak{X} u-(\mathfrak{X} u)_{\rho}\right|^{q} dx\right)^{\frac{1}{q}} \leq \sup_{x,y\in B_{\rho}}\left|\mathfrak{X} u(x)-\mathfrak{X} u(y)\right|
$$

$$
\leq c_{h}\left(\frac{\rho}{R}\right)^{\alpha_{1}}\left\|\mathfrak{X} u\right\|_{\infty,R/2}
$$

$$
< \frac{c_{h}}{\varepsilon_{0}}\left(\frac{\rho}{R}\right)^{\alpha}\left(\oint_{B_{R/2}}\left|\mathfrak{X} u-(\mathfrak{X} u)_{R}\right|^{q}\right)^{\frac{1}{q}}
$$

$$
\leq \frac{2^{Q}c_{h}}{\varepsilon_{0}}\left(\frac{\rho}{R}\right)^{\alpha}\left(\oint_{B_{R}}\left|\mathfrak{X} u-(\mathfrak{X} u)_{R}\right|^{q}\right)^{\frac{1}{q}}
$$

which implies our desired estimate. So we can assume [\(108\)](#page-33-1) holds. Now note that

,

$$
\left| \left(\mathfrak{X} u \right)_{R} \right|^{2} = \left\langle \left(\mathfrak{X} u \right)_{R}, \left(\mathfrak{X} u \right)_{R} \right\rangle = \left\langle \left(\mathfrak{X} u \right)_{R}, \left(\mathfrak{X} u \right)_{R} - \mathfrak{X} u \left(x \right) \right\rangle + \left\langle \left(\mathfrak{X} u \right)_{R}, \mathfrak{X} u \left(x \right) \right\rangle,
$$

for a.e. $x \in B_{R/2}$. Using Cauchy-Schwarz, integrating and finally using Young's inequality, we deduce

$$
\begin{aligned} \left| \left(\mathfrak{X} u \right)_R \right|^2 &\leq \left| \left(\mathfrak{X} u \right)_R \right| \left(\int_{B_{R/2}} \left| \mathfrak{X} u - \left(\mathfrak{X} u \right)_R \right| + \int_{B_{R/2}} \left| \mathfrak{X} u \right| \right) \\ &\overset{(108)}{\leq} \left(\varepsilon_0 + 1 \right) \left| \left(\mathfrak{X} u \right)_R \right| \left\| \mathfrak{X} u \right\|_{\infty, R/2} \\ &\leq \frac{1}{2} \left| \left(\mathfrak{X} u \right)_R \right|^2 + 2 \left(\varepsilon_0 + 1 \right)^2 \left\| \mathfrak{X} u \right\|_{\infty, R/2}^2. \end{aligned}
$$

Hence, we deduce

$$
|(\mathfrak{X} u)_R| \leq 2 \left(\varepsilon_0+1\right) \|\mathfrak{X} u\|_{\infty, R/2} \leq 4 \|\mathfrak{X} u\|_{\infty, R/2}.
$$

Thus, the only way [\(107\)](#page-33-2) can fail is if we have

$$
|(\mathfrak{X}u)_R|<\frac{1}{B}\|\mathfrak{X}u\|_{\infty,R/2}.
$$

But then by Theorem [30](#page-15-7) (with ' $q = 1$ '), Hölder's inequality and the triangle inequality for the L^q norm, we find,

$$
\begin{aligned} \|\mathfrak{X}u\|_{\infty,R/2} &\leq c_1 \left(\int_{B_R} |\mathfrak{X}u|^q\right)^{\frac{1}{q}} \leq c_1 \left(\int_{B_R} |\mathfrak{X}u - (\mathfrak{X}u)_R|^q\right)^{\frac{1}{q}} + c_1 \left|(\mathfrak{X}u)_R\right| \\ &< c_1 \left(\int_{B_R} |\mathfrak{X}u - (\mathfrak{X}u)_R|^q\right)^{\frac{1}{q}} + \frac{c_1}{B} \|\mathfrak{X}u\|_{\infty,R/2} \,. \end{aligned}
$$

Thus, recalling $B = 2c_1$, we have

$$
\|\mathfrak{X}u\|_{\infty,R/2}\leq B\left(\int_{B_R}|\mathfrak{X}u-(\mathfrak{X}u)_R|^q\right)^{\frac{1}{q}}.
$$

Hence, as before, for any $0<\rho< R/2,$ we have

$$
\left(\oint_{B_{\rho}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\rho} \right|^{q} dx \right)^{\frac{1}{q}} \leq c \left(\frac{\rho}{R}\right)^{\alpha_{1}} \left\| \mathfrak{X} u \right\|_{\infty, R/2}
$$

$$
\leq c B \left(\frac{\rho}{R}\right)^{\alpha} \left(\oint_{B_{R}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{R} \right|^{q} \right)^{\frac{1}{q}},
$$

which again yields the desired estimate. This completes the proof of Claim [43.](#page-33-0)

From now on, we shall assume that [\(107\)](#page-33-2) and [\(108\)](#page-33-1) holds. Now we claim

Claim 44. There exists a constant $\tau_1 \in (0,1)$, depending only on n and p such that at least one of the following alternatives occur.

(i) *The nondegenerate alternative:* We have

(109)
$$
|\mathfrak{X} u| \geq \frac{1}{4} ||\mathfrak{X} u||_{\infty, R/2} \quad in \ B_{\tau_1 R/2}.
$$

(ii) *The degenerate alternative:* We have

(110)
$$
\|\mathfrak{X} u\|_{\infty, \tau_1 R/2} \leq \frac{1}{2} \|\mathfrak{X} u\|_{\infty, R/2}.
$$

Proof of Claim [44:](#page-34-0) By Theorem [31,](#page-15-2) for any $0 < \tau < 1$, we have

$$
\sup_{x,y\in B_{\tau R/2}}|\mathfrak{X} u(x)-\mathfrak{X} u(y)|\leq c_h\tau^{\alpha_1}||\mathfrak{X} u||_{\infty,R/2}.
$$

Now, choose $\tau_1 \in (0,1)$ sufficiently small such that

$$
c_h \tau_1^{\alpha_1} = \frac{1}{4}.
$$

Now, if [\(109\)](#page-34-1) is violated, then there exists a point $\bar{x} \in B_{\tau_1 R/2}$ such that

$$
\left|\mathfrak{X}u\left(\bar{x}\right)\right|<\frac{1}{4}\left\Vert \mathfrak{X}u\right\Vert _{\infty,R/2}.
$$

But then, for any $x \in B_{\tau_1 R/2}$, by Theorem [31,](#page-15-2) we deduce

$$
\begin{aligned} |\mathfrak{X}u(x)| &\leq |\mathfrak{X}u(\bar{x})| + |\mathfrak{X}u(x) - \mathfrak{X}u(\bar{x})| \\ &\leq \frac{1}{4} ||\mathfrak{X}u||_{\infty, R/2} + c_h \tau_1^{\alpha_1} ||\mathfrak{X}u||_{\infty, R/2} = \frac{1}{2} ||\mathfrak{X}u||_{\infty, R/2} \,. \end{aligned}
$$

This completes the proof of Claim [44.](#page-34-0)

Now we define

$$
R_i := \tau_1^i \frac{R}{2} \qquad \text{for all integers } i \ge 0.
$$

Thus, by Claim [44,](#page-34-0) at each level i , at least one of the following alternatives must occur.

(a) The nondegenerate alternative: We have

(111)
$$
|\mathfrak{X}u| \geq \frac{1}{4} ||\mathfrak{X}u||_{\infty,R_i} \quad \text{in } B_{R_{i+1}}.
$$

(b) The degenerate alternative: We have

(112)
$$
\|\mathfrak{X} u\|_{\infty, R_{i+1}} \leq \frac{1}{2} \|\mathfrak{X} u\|_{\infty, R_i}.
$$

Next, we choose $i_a \geq 1$ to be first integer for which we have

$$
(113) \t\t\t 2ia-1 \ge B.
$$

Clearly, i_a depends only on B and thus ultimately, only on n and p.

Claim 45. Without loss of generality, we can assume that [\(111\)](#page-34-2) occurs for the first time for some $0 \leq i < i_a$.

Proof of Claim [45:](#page-35-0) If Claim [45](#page-35-0) is false, then (112) must have occured for all $0 \leq i \leq i_a - 1$. But this implies, we have,

$$
\|\mathfrak{X}u\|_{\infty,R_{i_a}} \leq \left(\frac{1}{2}\right)^{i_a} \|\mathfrak{X}u\|_{\infty,R/2}.
$$

This now implies

(114)
$$
\left(\int_{B_{R_{i_a}}} |\mathfrak{X} u - (\mathfrak{X} u)_R|^q\right)^{\frac{1}{q}} \ge \left(\int_{B_{R_{i_a}}} |\mathfrak{X} u|^q\right)^{\frac{1}{q}}.
$$

Indeed, if [\(114\)](#page-35-2) is false, then recalling [\(107\)](#page-33-2), we have

$$
\frac{1}{B} \|\mathfrak{X}u\|_{\infty,R/2} \leq |(\mathfrak{X}u)_R|
$$
\n
$$
= \int_{B_{R_{i_a}}} |(\mathfrak{X}u)_R| dx
$$
\n
$$
\overset{\text{Hölder}}{\leq} \left(\int_{B_{R_{i_a}}} |\mathfrak{X}u - (\mathfrak{X}u)_R|^q dx\right)^{\frac{1}{q}} + \left(\int_{B_{R_{i_a}}} |\mathfrak{X}u|^q\right)^{\frac{1}{q}}
$$
\n
$$
< 2 \left(\int_{B_{R_{i_a}}} |\mathfrak{X}u|^q\right)^{\frac{1}{q}} \leq 2 \|\mathfrak{X}u\|_{\infty,R_{i_a}} \leq 2 \left(\frac{1}{2}\right)^{i_a} \|\mathfrak{X}u\|_{\infty,R/2},
$$

which contradicts [\(113\)](#page-35-3). Hence [\(114\)](#page-35-2) holds. Now using Theorem [30](#page-15-7) (with $q = 1$) together with [\(114\)](#page-35-2) and the fact $1 \le q \le 2$, we find

$$
\begin{split} \left\| \mathfrak{X} u \right\|_{\infty, R_{i_a}/2} &\leq c_1 \int_{B_{R_{i_a}}} \left| \mathfrak{X} u \right| \leq c_1 \left(\int_{B_{R_{i_a}}} \left| \mathfrak{X} u \right|^q \right)^{\frac{1}{q}} \\ &\leq c_1 \left(\int_{B_{R_{i_a}}} \left| \mathfrak{X} u - (\mathfrak{X} u)_R \right|^q \right)^{\frac{1}{q}} \\ &\leq c_1 \left(\frac{R}{R_{i_a}} \right)^{\frac{Q}{q}} \left(\int_{B_R} \left| \mathfrak{X} u - (\mathfrak{X} u)_R \right|^q \right)^{\frac{1}{q}} \\ &\leq c_1 2^Q \tau_1^{-i_a} Q \left(\int_{B_R} \left| \mathfrak{X} u - (\mathfrak{X} u)_R \right|^q \right)^{\frac{1}{q}} . \end{split}
$$

Combining this with Theorem [31,](#page-15-2) for any $0 < \rho < R_{i_a}/2$, we have

$$
\left(\oint_{B_{\rho}}\left|\mathfrak{X} u-(\mathfrak{X} u)_{\rho}\right|^{q} dx\right)^{\frac{1}{q}} \leq \sup_{x,y\in B_{\rho}}\left|\mathfrak{X} u(x)-\mathfrak{X} u(y)\right|
$$

$$
\leq c_{h} \left(\frac{\rho}{R_{i_{a}}}\right)^{\alpha_{1}}\left\|\mathfrak{X} u\right\|_{\infty,R_{i_{a}}/2}
$$

$$
\leq c_{h} 2^{\alpha_{1}} \tau_{1}^{-\alpha_{1} i_{a}} \left(\frac{\rho}{R}\right)^{\alpha_{1}}\left\|\mathfrak{X} u\right\|_{\infty,R_{i_{a}}/2}
$$

$$
\leq c_{h} c_{1} 2^{2Q} \tau_{1}^{-i_{a}(Q+\alpha_{1})} \left(\frac{\rho}{R}\right)^{\alpha_{1}} \left(\oint_{B_{R}}\left|\mathfrak{X} u-(\mathfrak{X} u)_{R}\right|^{q}\right)^{\frac{1}{q}}.
$$

This together with the fact that if $\frac{R_{i_a}}{2} \leq \rho \leq R$, then $\frac{\tau_1^{i_a}}{2} \leq \frac{\rho}{R} \leq 1$, yields our desired estimate. This proves Claim [45.](#page-35-0)

Thus, we can assume that there exists a smallest integer $0 \le i_0 < i_a$ such that [\(111\)](#page-34-2) occurs for $i = i_0$. Thus, we have

(115)
$$
\frac{1}{4} ||\mathfrak{X}u||_{\infty, R_{i_0}} \leq |\mathfrak{X}u| \leq ||\mathfrak{X}u||_{\infty, R_{i_0}} \quad \text{in } B_{R_{i_0+1}}.
$$

Now we claim

Claim 46. There exists a constant $\sigma \in (0,1)$, depending only on n, p and ε_0 , and thus ultimately only on n and p, such that we have

(116)
$$
\left(\int_{B_{\sigma R_{i_0+1}}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\sigma R_{i_0+1}} \right|^q \right)^{\frac{1}{q}} \leq \frac{\varepsilon_0}{16} \left\| \mathfrak{X} u \right\|_{\infty, R_{i_0}}.
$$

Proof of Claim [46:](#page-36-0) Let $c_h > 0$ be the constant given by Theorem [31.](#page-15-2) Choose $\sigma \in (0, 1)$ sufficiently small such that

$$
c_h \sigma^{\alpha_1} = \frac{\varepsilon_0}{16}.
$$

Then, by Theorem [31,](#page-15-2) we deduce

$$
\left(\oint_{B_{\sigma R_{i_0+1}}}\left|\mathfrak{X} u - (\mathfrak{X} u)_{\sigma R_{i_0+1}}\right|^q\right)^{\frac{1}{q}} \leq \sup_{x,y \in B_{\sigma R_{i_0+1}}} \left|\mathfrak{X} u(x) - \mathfrak{X} u(y)\right|
$$

$$
\leq c_h \sigma^{\alpha_1} \left\|\mathfrak{X} u\right\|_{\infty, R_{i_0+1}}
$$

$$
\leq \frac{\varepsilon_0}{16} \left\|\mathfrak{X} u\right\|_{\infty, R_{i_0+1}} \leq \frac{\varepsilon_0}{16} \left\|\mathfrak{X} u\right\|_{\infty, R_{i_0}}
$$

This proves Claim [46.](#page-36-0)

Now we are in a position to complete the proof of the Theorem. Observe that exactly one of the following can occur. Either we have

(117)
$$
\left| \left(\mathfrak{X} u \right)_{\frac{1}{2} \sigma R_{i_0+1}} \right| \geq \frac{1}{8B} \left\| \mathfrak{X} u \right\|_{\infty, R_{i_0}},
$$

or we must have

(118)
$$
\left| (\mathfrak{X} u)_{\frac{1}{2}\sigma R_{i_0+1}} \right| < \frac{1}{8B} ||\mathfrak{X} u||_{\infty, R_{i_0}}.
$$

.

If [\(118\)](#page-36-1) holds, then by Theorem [30](#page-15-7) (with ' $q = 1$ '), we have

$$
\|\tilde{x}u\|_{\infty, \frac{1}{4}\sigma R_{i_0+1}} \leq c_1 \int_{B_{\frac{1}{2}\sigma R_{i_0+1}}} |\tilde{x}u|
$$

\n
$$
\leq c_1 \left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}} |\tilde{x}u|^q \right)^{\frac{1}{q}}
$$

\n
$$
\leq c_1 \left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}} \left| \tilde{x}u - (\tilde{x}u)_{\frac{1}{2}\sigma R_{i_0+1}} \right|^q \right)^{\frac{1}{q}} + c_1 \left| (\tilde{x}u)_{\frac{1}{2}\sigma R_{i_0+1}} \right|
$$

\n
$$
\leq c_1 \left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}} \left| \tilde{x}u - (\tilde{x}u)_{\frac{1}{2}\sigma R_{i_0+1}} \right|^q \right)^{\frac{1}{q}} + \frac{c_1}{8B} \|\tilde{x}u\|_{\infty, R_{i_0}}
$$

\n(119)
\n
$$
\leq c_1 \left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}} \left| \tilde{x}u - (\tilde{x}u)_{\frac{1}{2}\sigma R_{i_0+1}} \right|^q \right)^{\frac{1}{q}} + \frac{c_1}{2B} \|\tilde{x}u\|_{\infty, \frac{1}{4}\sigma R_{i_0+1}},
$$

where in the last line, we have used the bound

(120)
$$
\frac{1}{4} \|\mathfrak{X}u\|_{\infty, R_{i_0}} \le \|\mathfrak{X}u\|_{\infty, \frac{1}{4}\sigma R_{i_0+1}},
$$

which is implied by [\(115\)](#page-36-2). Now by our choice of $B = 2c_1$, [\(119\)](#page-37-0) implies

$$
\|\mathfrak{X} u\|_{\infty,\frac{1}{4}\sigma R_{i_0+1}/2} \leq \frac{4c_1}{3}\left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}}\left|\mathfrak{X} u-(\mathfrak{X} u)_{\frac{1}{2}\sigma R_{i_0+1}}\right|^{q}\right)^{\frac{1}{q}}.
$$

 $\mathbf{1}$

Once again, combined with Theorem [31,](#page-15-2) this implies, for any $0 < \rho \leq \sigma R_{i_a}/4 \leq$ $\sigma R_{i_0+1}/4$, we have

$$
\left(\oint_{B_{\rho}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\rho} \right|^{q} dx \right)^{\frac{1}{q}}\n\leq c_{h} \left(\frac{\rho}{\sigma R_{i_{0}+1}}\right)^{\alpha_{1}} \left\| \mathfrak{X} u \right\|_{\infty, \frac{1}{4}\sigma R_{i_{0}+1}}\n\leq \frac{4c_{h}c_{1}}{3} \left(\frac{\rho}{\sigma R_{i_{0}+1}}\right)^{\alpha_{1}} \left(\oint_{B_{\frac{1}{2}\sigma R_{i_{0}+1}}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\frac{1}{2}\sigma R_{i_{0}+1}} \right|^{q} \right)^{\frac{1}{q}}\n\leq \frac{(20)}{3} \frac{8c_{h}c_{1}}{3} \left(\frac{\rho}{\sigma R_{i_{0}+1}}\right)^{\alpha_{1}} \left(\oint_{B_{\frac{1}{2}\sigma R_{i_{0}+1}}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{R} \right|^{q} \right)^{\frac{1}{q}}\n\leq c \left(\frac{R}{\sigma R_{i_{0}+1}}\right)^{\frac{\alpha}{q}+\alpha_{1}} \left(\frac{\rho}{R}\right)^{\alpha_{1}} \left(\oint_{B_{R}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{R} \right|^{q} \right)^{\frac{1}{q}}.
$$

Now, since σ , τ_1 , α_1 are fixed constants and i_a is a fixed integer, all depending only on *n* and *p*, with $\tau_1 \in (0,1)$ and $i_0 < i_a$, we note that

$$
\left(\frac{R}{\sigma R_{i_0+1}}\right)^{\frac{Q}{q}+\alpha_1}=\left(\frac{1}{\sigma\tau_1^{i_0+1}}\right)^{\frac{Q}{q}+\alpha_1}\leq \left(\frac{1}{\sigma\tau_1^{i_a}}\right)^{\frac{Q}{q}+\alpha_1}\leq \left(\frac{1}{\sigma\tau_1^{i_a}}\right)^{Q+\alpha_1},
$$

where the last term on the right is again a fixed constant, depending only on n and p. Hence, [\(121\)](#page-37-1) now easily yields our desired estimate. Hence it only remains to settle the case when [\(117\)](#page-36-3) holds. Observe that [\(116\)](#page-36-4) implies

$$
\left(\int_{B_{\sigma R_{i_0+1}}}\left|\mathfrak{X} u-(\mathfrak{X} u)_{\sigma R_{i_0+1}}\right|^q\right)^{\frac{1}{q}}\leq \frac{\varepsilon_0}{16}\left\|\mathfrak{X} u\right\|_{\infty,R_{i_0}}\leq \varepsilon_0\left\|\mathfrak{X} u\right\|_{\infty,R_{i_0+1}},
$$

where once again we have used the bound (120) . Combining this with the Hölder and reverse Hölder inequality, we deduce

$$
\left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}}\left|\mathfrak{X} u - (\mathfrak{X} u)_{\frac{1}{2}\sigma R_{i_0+1}}\right|^2\right)^{\frac{1}{2}}\n\leq 2\left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}}\left|\mathfrak{X} u - (\mathfrak{X} u)_{\sigma R_{i_0+1}}\right|^2\right)^{\frac{1}{2}}\n\leq 2\left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}}\left|\mathfrak{X} u - (\mathfrak{X} u)_{\sigma R_{i_0+1}}\right|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2Q}}\n\leq 2\left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}}\left|\mathfrak{X} u - (\mathfrak{X} u)_{\sigma R_{i_0+1}}\right|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2Q}}\n\leq 2C_{RH}\int_{B_{\sigma R_{i_0+1}}}\left|\mathfrak{X} u - (\mathfrak{X} u)_{\sigma R_{i_0+1}}\right|^q\right)^{\frac{1}{q}}\leq 2C_{RH}\varepsilon_0\left\|\mathfrak{X} u\right\|_{\infty,R_{i_0+1}}.
$$

Thus, recalling $\varepsilon_0 \in (0,1)$, we have

$$
\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\frac{1}{2}\sigma R_{i_0+1}} \right|^2 \leq \varepsilon_0^2 (2C_{RH})^2 \left\| \mathfrak{X} u \right\|_{\infty, R_{i_0+1}}^2
$$
\n
$$
\leq \varepsilon_0 (2C_{RH})^2 \left\| \mathfrak{X} u \right\|_{\infty, R_{i_0+1}}^2.
$$

By virtue of [\(115\)](#page-36-2), we also trivially have (123)

$$
\frac{1}{4B} \|\mathfrak{X}u\|_{\infty, R_{i_0+1}} \leq \frac{1}{4} \|\mathfrak{X}u\|_{\infty, R_{i_0}} \leq |\mathfrak{X}u| \leq 2C_{RH} \|\mathfrak{X}u\|_{\infty, R_{i_0+1}} \quad \text{in } B_{R_{i_0+1}}.
$$

Now note that [\(117\)](#page-36-3) immediately implies

$$
(124) \frac{1}{8B} \|\mathfrak{X}u\|_{\infty, R_{i_0+1}} \le \frac{1}{8B} \|\mathfrak{X}u\|_{\infty, R_{i_0}} \le \left| (\mathfrak{X}u)_{\frac{1}{2}\sigma R_{i_0+1}} \right| \le 2C_{RH} \|\mathfrak{X}u\|_{\infty, R_{i_0+1}}.
$$

Now (123) (124) and (122) implies that the hypotheses of Lamm-41 are satisfied.

Now, [\(123\)](#page-38-0), [\(124\)](#page-38-1) and [\(122\)](#page-38-2) implies that the hypotheses of Lemma [41](#page-29-3) are satisfied with R replaced by $R_{i_0+1}/2$, $A = 2C_{RH}$, $B = 2c_1$ and $\lambda = {\|\mathfrak{X}u\|}_{\infty, R_{i_0+1}}$, which we

can always assume to be positive, as otherwise there is nothing to prove. Hence, Lemma [41](#page-29-3) implies that for any $0 < \rho \leq \sigma R_{i_a}/2 \leq \sigma R_{i_0+1}/2$, we have (125)

$$
\int_{B_\rho}\left|\mathfrak X u-(\mathfrak X u)_\rho\right|^2\leq C_0\sigma^{-Q-2\alpha_0}\left(\frac{\rho}{R_{i_0+1}}\right)^{2\alpha_0}\int_{B_{\frac{1}{2}R_{i_0+1}}}\left|\mathfrak X u-(\mathfrak X u)_{\frac{1}{2}R_{i_0+1}}\right|^2.
$$

From here, to derive our desired estimate, we use Hölder inequality on one side and reverse Hölder inequality on the other. More precisely, by [\(125\)](#page-39-0), for any $0 < \rho \leq$ $\sigma R_{i_a}/2 \leq \sigma R_{i_0+1}/2$, we have

$$
\left(\int_{B_{\rho}}\left|\mathfrak{X} u-(\mathfrak{X} u)_{\rho}\right|^{q}\right)^{\frac{1}{q}}
$$
\n
$$
\stackrel{\text{Hölder}}{\leq} \left(\int_{B_{\rho}}\left|\mathfrak{X} u-(\mathfrak{X} u)_{\rho}\right|^{2}\right)^{\frac{1}{2}}
$$
\n
$$
\stackrel{\text{Hölder}}{\leq} \left(\int_{G}\sigma^{-\frac{Q}{2}-\alpha_{0}}\left(\frac{\rho}{R_{i_{0}+1}}\right)^{\alpha_{0}}\left(\int_{B_{\frac{1}{2}R_{i_{0}+1}}}\left|\mathfrak{X} u-(\mathfrak{X} u)_{\frac{1}{2}R_{i_{0}+1}}\right|^{2}\right)^{\frac{1}{2}}
$$
\n
$$
\stackrel{\text{Hölder}}{\leq} C_{0}\sigma^{-\frac{Q}{2}-\alpha_{0}}\left(\frac{\rho}{R_{i_{0}+1}}\right)^{\alpha_{0}}\left(\int_{\frac{1}{2}B_{R_{i_{0}+1}}}\left|\mathfrak{X} u-(\mathfrak{X} u)_{\frac{1}{2}R_{i_{0}+1}}\right|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2Q}}
$$
\n
$$
\stackrel{\text{(20)}}{\leq} 2C_{0}\sigma^{-\frac{Q}{2}-\alpha_{0}}\left(\frac{\rho}{R_{i_{0}+1}}\right)^{\alpha_{0}}\left(\int_{B_{\frac{1}{2}R_{i_{0}+1}}} \left|\mathfrak{X} u-(\mathfrak{X} u)_{R_{i_{0}+1}}\right|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2Q}}
$$
\n
$$
\stackrel{\text{(115),(66)}}{\leq} 2C_{RH}C_{0}\sigma^{-\frac{Q}{2}-\alpha_{0}}\left(\frac{\rho}{R_{i_{0}+1}}\right)^{\alpha_{0}}\int_{B_{R_{i_{0}+1}}} \left|\mathfrak{X} u-(\mathfrak{X} u)_{R_{i_{0}+1}}\right|^{2}\right)^{\frac{1}{q}}
$$
\n
$$
\stackrel{\text{(20)}}{\leq} 4C_{RH}C_{0}\sigma^{-\frac{Q}{2}-\alpha_{0}}\left(\frac{\rho}{R_{i_{0}+1}}\right)^{\alpha_{0}}\left(\int_{
$$

Since C_{RH} , C_0 , σ , α_0 , τ_1 , i_a are all fixed, which ultimately depend only on n and p, we derive our desired estimate. This completes the proof. \Box

Now we prove the corresponding result for $V(\mathfrak{X}u)$. The proof is completely analogous to the proof of Theorem [1.](#page-3-0) One only has to use Theorem [32](#page-16-2) instead of Theorem [31](#page-15-2) and Lemma [42](#page-31-0) instead of Lemma [41.](#page-29-3) For the convenience of readers we provide complete details.

Proof. (of Theorem [2](#page-3-2)) We can and henceforth assume $x_0 = 0$. As remarked in the proof of Theorem [1,](#page-3-0) it is enough to prove [\(7\)](#page-3-4), for $0 < \rho < \kappa R$, for some constant $\kappa \in (0,1)$ depending only n and p, as the estimate is trivial otherwise. First, we fix $C_{RH,V} > 0$ which is given by Lemma [36](#page-22-2) with the choice $A = B = 1$. So, clearly $C_{RH,V}$ depends only on n and p. Next, let $B = B(n,p) \ge 1$ be defined by $B := 2^{1+\frac{1}{p}} c_{p/2}$, where $c_{p/2}$ is the constant given by Theorem [30](#page-15-7) with the choice $q = p/2$. Let $\theta_0, \alpha_0 \in (0, 1)$ be the constants given by Lemma [42](#page-31-0) for the choices of $A = 2C_{RH,V}$ and $B = 2^{1+\frac{1}{p}}c_{p/2}$. Let $\beta_1 \in (0,1)$ be the exponent in Theorem [32.](#page-16-2) Set

$$
\beta := \min \left\{ \alpha_0, \beta_1 \right\}.
$$

Claim 47. Without loss of generality, we can assume that the we have

(126)
$$
\frac{1}{B} \|\mathfrak{X}u\|_{\infty, R/2} \le |(V(\mathfrak{X}u))_{R}|^{\frac{2}{p}} \le 16 \|\mathfrak{X}u\|_{\infty, R/2}
$$

and

(127)
$$
\left(\int_{B_{R/2}} |V(\mathfrak{X} u) - (V(\mathfrak{X} u))_R|^q\right)^{\frac{1}{q}} \leq \theta_0 \left\|\mathfrak{X} u\right\|_{\infty, R/2}^{\frac{p}{2}}.
$$

Proof of Claim [47:](#page-40-0) Indeed, if (127) is violated, then by Theorem [32,](#page-16-2) for any $0 < \rho < R/2$, we have

$$
\left(\oint_{B_{\rho}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\rho}\right|^{q} dx\right)^{\frac{1}{q}}
$$
\n
$$
\leq \sup_{x,y\in B_{\rho}}\left|V\left(\mathfrak{X} u\right)\left(x\right)-V\left(\mathfrak{X} u\right)\left(y\right)\right|
$$
\n
$$
\leq c_{h,V}\left(\frac{\rho}{R}\right)^{\beta_{1}}\left\|\mathfrak{X} u\right\|_{\infty,R/2}^{\frac{p}{2}}
$$
\n
$$
\leq \frac{c_{h,V}}{\theta_{0}}\left(\frac{\rho}{R}\right)^{\beta}\left(\oint_{B_{R/2}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{R}\right|^{q}\right)^{\frac{1}{q}}
$$
\n
$$
\leq \frac{2^{Q}c_{h,V}}{\theta_{0}}\left(\frac{\rho}{R}\right)^{\beta}\left(\oint_{B_{R}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{R}\right|^{q}\right)^{\frac{1}{q}},
$$

which implies our desired estimate. So we can assume [\(127\)](#page-40-1) holds. Now note that

$$
\begin{aligned} \left| \left(V \left(\mathfrak{X} u \right) \right)_R \right|^2 &= \left\langle \left(V \left(\mathfrak{X} u \right) \right)_R, \left(V \left(\mathfrak{X} u \right) \right)_R \right\rangle \\ &= \left\langle \left(V \left(\mathfrak{X} u \right) \right)_R, \left(V \left(\mathfrak{X} u \right) \right)_R - V \left(\mathfrak{X} u \right) \left(x \right) \right\rangle + \left\langle \left(V \left(\mathfrak{X} u \right) \right)_R, V \left(\mathfrak{X} u \right) \left(x \right) \right\rangle, \end{aligned}
$$

for a.e. $x \in B_{R/2}$. Using Cauchy-Schwarz, integrating and Hölder, we deduce

$$
\begin{aligned} \left| \left(V \left(\mathfrak{X} u \right) \right)_R \right|^2 &\leq \left| \left(V \left(\mathfrak{X} u \right) \right)_R \right| \left(\int_{B_{R/2}} \left| V \left(\mathfrak{X} u \right) - \left(V \left(\mathfrak{X} u \right) \right)_R \right| + \int_{B_{R/2}} \left| V \left(\mathfrak{X} u \right) \right| \right) \\ &\leq \left(\theta_0 + 1 \right) \left| \left(V \left(\mathfrak{X} u \right) \right)_R \right| \left\| \mathfrak{X} u \right\|_{\infty, R/2}^{\frac{p}{2}} . \end{aligned}
$$

Since, $0 < \theta_0 < 1$ so, using Young's inequality in the last estimate we derive

$$
\begin{split} \left| \left(V \left(\mathfrak{X} u \right) \right)_R \right|^{\frac{2}{p}} &\leq \left(\theta_0 + 1 \right)^{\frac{1}{p}} \left| \left(V \left(\mathfrak{X} u \right) \right)_R \right|^{\frac{1}{p}} \left\| \mathfrak{X} u \right\|_{\infty, R/2}^{\frac{1}{2}} \\ &\leq 2^{\frac{1}{p}} \left| \left(V \left(\mathfrak{X} u \right) \right)_R \right|^{\frac{1}{p}} \left\| \mathfrak{X} u \right\|_{\infty, R/2}^{\frac{1}{2}} \\ &\leq \frac{1}{2} \left| \left(V \left(\mathfrak{X} u \right) \right)_R \right|^{\frac{2}{p}} + 2^{\frac{2}{p} + 1} \left\| \mathfrak{X} u \right\|_{\infty, R/2} . \end{split}
$$

Hence, we deduce

$$
|(V(\mathfrak{X} u))_R|^{\frac{2}{p}} \leq 2^{\frac{2}{p}+2} \|\mathfrak{X} u\|_{\infty, R/2} \leq 16 \|\mathfrak{X} u\|_{\infty, R/2}.
$$

Thus, the only way [\(126\)](#page-40-2) can fail is if we have

$$
\left| \left(V \left(\mathfrak{X} u \right) \right)_R \right|^{\frac{2}{p}} < \frac{1}{B} \left\| \mathfrak{X} u \right\|_{\infty, R/2}
$$

.

But then by Theorem [30](#page-15-7) (with ' $q = p/2$ '), Hölder's inequality and the Minkowski inequality, we find,

$$
\begin{split} \|\mathfrak{X}u\|_{\infty,R/2} &\leq c_{p/2} \left(\int_{B_R} |\mathfrak{X}u|^{\frac{p}{2}}\right)^{\frac{2}{p}} \\ &\leq c_{p/2} \left(\int_{B_R} |V(\mathfrak{X}u)|^q\right)^{\frac{2}{pq}} \\ &\leq c_{p/2} \left[\left(\int_{B_R} |V(\mathfrak{X}u) - (V(\mathfrak{X}u))_R|^q\right)^{\frac{1}{q}} + |(V(\mathfrak{X}u))_R|\right]^{\frac{2}{p}} \\ &\leq 2^{\frac{1}{p}} c_{p/2} \left[\left(\int_{B_R} |V(\mathfrak{X}u) - (V(\mathfrak{X}u))_R|^q\right)^{\frac{2}{q}} + |(V(\mathfrak{X}u))_R|^2\right]^{\frac{1}{p}} \\ &\leq 2^{\frac{1}{p}} c_{p/2} \left(\int_{B_R} |V(\mathfrak{X}u) - (V(\mathfrak{X}u))_R|^q\right)^{\frac{2}{p}} + 2^{\frac{1}{p}} c_{p/2} \left|(V(\mathfrak{X}u))_R\right|^{\frac{2}{p}} \\ &< 2^{\frac{1}{p}} c_{p/2} \left(\int_{B_R} |V(\mathfrak{X}u) - (V(\mathfrak{X}u))_R|^q\right)^{\frac{2}{p}} + \frac{2^{\frac{1}{p}} c_{p/2}}{B} \|\mathfrak{X}u\|_{\infty, R/2} \,. \end{split}
$$

Thus, using the choice of $B = 2^{1 + \frac{1}{p}} c_{p/2}$, we have

$$
\|\mathfrak{X}u\|_{\infty,R/2} \leq B\left(\int_{B_R} |V(\mathfrak{X}u) - (V(\mathfrak{X}u))_R|^q\right)^{\frac{2}{pq}}.
$$

Hence, as before, for any $0 < \rho < R/2$, we have

$$
\left(\oint_{B_{\rho}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\rho}\right|^{q} dx\right)^{\frac{1}{q}}
$$
\n
$$
\leq c_{h,V}\left(\frac{\rho}{R}\right)^{\beta_{1}}\left\|\mathfrak{X} u\right\|_{\infty,R/2}^{\frac{p}{2}} \leq c_{h,V}B^{\frac{p}{2}}\left(\frac{\rho}{R}\right)^{\beta_{1}}\left(\oint_{B_{R}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{R}\right|^{q}\right)^{\frac{1}{q}},
$$

which again yields the desired estimate. This completes the proof of Claim [47.](#page-40-0)

From now on, we shall assume that [\(126\)](#page-40-2) and [\(127\)](#page-40-1) holds. Now we by Claim [44](#page-34-0) we assert the existence of a constant $\tau_1 \in (0,1)$, depending only on n and p such that at least one of the following alternatives occur.

(i) The nondegenerate alternative: We have

(128)
$$
|\mathfrak{X}u| \geq \frac{1}{4} ||\mathfrak{X}u||_{\infty, R/2} \quad \text{in } B_{\tau_1 R/2}.
$$

(ii) The degenerate alternative: We have

(129)
$$
\|\mathfrak{X} u\|_{\infty, \tau_1 R/2} \leq \frac{1}{2} \|\mathfrak{X} u\|_{\infty, R/2}.
$$

Now we define

$$
R_i := \tau_1^i \frac{R}{2} \qquad \text{for all integers } i \ge 0.
$$

Thus, by Claim 44 , at each level i, at least one of the following alternatives must occur.

(a) The nondegenerate alternative: We have

(130)
$$
|\mathfrak{X}u| \geq \frac{1}{4} ||\mathfrak{X}u||_{\infty, R_i} \quad \text{in } B_{R_{i+1}}.
$$

(b) The degenerate alternative: We have

(131)
$$
\|\mathfrak{X}u\|_{\infty,R_{i+1}} \leq \frac{1}{2} \|\mathfrak{X}u\|_{\infty,R_i}.
$$

Next, we choose $i_a \geq 1$ to be first integer for which we have

(132)
$$
2^{i_a - 1} \ge 2^{\frac{2}{p}} B.
$$

Clearly, i_a depends only on B and thus ultimately, only on n and p. Now we claim the following.

Claim 48. Without loss of generality, we can assume that [\(130\)](#page-42-0) occurs for the first time for some $0 \leq i < i_a$.

Proof of Claim [48:](#page-42-1) If Claim [48](#page-42-1) is false, then [\(131\)](#page-42-2) must have occured for all $0 \leq i \leq i_a - 1$. But this implies, we have,

$$
\|\mathfrak{X}u\|_{\infty,R_{i_a}} \leq \left(\frac{1}{2}\right)^{i_a} \|\mathfrak{X}u\|_{\infty,R/2}.
$$

This now implies

(133)
$$
\left(\oint_{B_{R_{i_a}}} |V(\mathfrak{X}u) - (V(\mathfrak{X}u))_R|^q\right)^{\frac{1}{q}} \ge \left(\oint_{B_{R_{i_a}}} |V(\mathfrak{X}u)|^q\right)^{\frac{1}{q}}.
$$

Indeed, if [\(133\)](#page-42-3) is false, then recalling [\(126\)](#page-40-2), we have

$$
\frac{1}{B^{\frac{p}{2}}} ||\mathfrak{X}u||_{\infty,R/2}^{\frac{p}{2}} \leq |(V (\mathfrak{X} u))_{R}|
$$
\n
$$
= \int_{B_{R_{i_a}}} |(V (\mathfrak{X} u))_{R}| \, dx
$$
\n
$$
\overset{\text{Hölder}}{\leq} \left(\int_{B_{R_{i_a}}} |V (\mathfrak{X} u) - (V (\mathfrak{X} u))_{R}|^{q} \, dx \right)^{\frac{1}{q}} + \left(\int_{B_{R_{i_a}}} |V (\mathfrak{X} u)|^{q} \right)^{\frac{1}{q}}
$$
\n
$$
< 2 \left(\int_{B_{R_{i_a}}} |V (\mathfrak{X} u)|^{q} \right)^{\frac{1}{q}} \leq 2 ||\mathfrak{X}u||_{\infty,R_{i_a}}^{\frac{p}{2}} \leq 2 \left(\frac{1}{2} \right)^{\frac{p}{2} i_a} ||\mathfrak{X}u||_{\infty,R/2}^{\frac{p}{2}},
$$

which is a contradiction, by (132) . Hence (133) holds. Now using Theorem [30](#page-15-7) (with 'q=p/2') together with [\(133\)](#page-42-3) and the fact $1 \le q \le 2$, we find

$$
\begin{split} \|\mathfrak{X}u\|_{\infty,R_{i_{a}}/2}&\leq c_{p/2}\left(\int_{B_{R_{i_{a}}}}|\mathfrak{X}u|^\frac{p}{2}\right)^\frac{2}{p}\\ &\leq c_{p/2}\left(\int_{B_{R_{i_{a}}}}|V\left(\mathfrak{X}u\right)|^{q}\right)^\frac{2}{pq}\\ &\leq c_{p/2}\left(\int_{B_{R_{i_{a}}}}|V\left(\mathfrak{X}u\right)-\left(V\left(\mathfrak{X}u\right)\right)_{R}|^{q}\right)^\frac{2}{pq}\\ &\leq c_{p/2}\left(\frac{R}{R_{i_{a}}}\right)^\frac{2Q}{pq}\left(\int_{B_{R}}|V\left(\mathfrak{X}u\right)-\left(V\left(\mathfrak{X}u\right)\right)_{R}|^{q}\right)^\frac{2}{pq}\\ &\leq c_{p/2}2^\frac{2Q}{p}\tau_{1}^{-\frac{2i_{a}Q}{p}}\left(\int_{B_{R}}|V\left(\mathfrak{X}u\right)-\left(V\left(\mathfrak{X}u\right)\right)_{R}|^{q}\right)^\frac{2}{pq}. \end{split}
$$

Combining this with Theorem [32,](#page-16-2) for any $0 < \rho < R_{i_a}/2$, we have

$$
\left(\oint_{B_{\rho}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\rho}\right|^{q}dx\right)^{\frac{1}{q}}
$$
\n
$$
\leq \sup_{x,y\in B_{\rho}}\left|V\left(\mathfrak{X} u\right)\left(x\right)-V\left(\mathfrak{X} u\right)\left(y\right)\right|
$$
\n
$$
\leq c\left(\frac{\rho}{R_{i_{a}}}\right)^{\beta_{1}}\left\|\mathfrak{X} u\right\|_{\infty,R_{i_{a}}/2}^{\frac{p}{2}}
$$
\n
$$
\leq c\tau_{1}^{-\beta_{1}i_{a}}\left(\frac{\rho}{R}\right)^{\beta_{1}}\left\|\mathfrak{X} u\right\|_{\infty,R_{i_{a}}/2}^{\frac{p}{2}}
$$
\n
$$
\leq c c_{\rho/2}^{\frac{p}{2}}\tau_{1}^{-i_{a}\left(Q+\beta_{1}\right)}\left(\frac{\rho}{R}\right)^{\beta_{1}}\left(\oint_{B_{R}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{R}\right|^{q}\right)^{\frac{1}{q}}.
$$

This together with the fact that if $\frac{R_{i_a}}{2} \leq \rho \leq R$, then $\frac{\tau_1^{i_a}}{2} \leq \frac{\rho}{R} \leq 1$, yields our desired estimate. This proves Claim [48.](#page-42-1)

Thus, we can assume that there exists a smallest integer $0 \leq i_0 < i_a$ such that [\(130\)](#page-42-0) occurs for $i = i_0$. Thus, we have

(134)
$$
\frac{1}{4} ||\mathfrak{X}u||_{\infty, R_{i_0}} \leq |\mathfrak{X}u| \leq ||\mathfrak{X}u||_{\infty, R_{i_0}} \quad \text{in } B_{R_{i_0+1}}.
$$

Now we claim

Claim 49. There exists a constant $\sigma \in (0,1)$, depending only on n, p and θ_0 , and thus ultimately only on n and p, such that we have

(135)
$$
\left(\int_{B_{\sigma R_{i_0+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\sigma R_{i_0+1}}\right|^q\right)^{\frac{1}{q}} \leq \frac{\theta_0}{16}\left\|\mathfrak{X} u\right\|_{\infty,R_{i_0}}^{\frac{p}{2}}.
$$

Proof of Claim [49:](#page-43-0) Let $c_{h,V} > 0$ be the constant given by Theorem [32.](#page-16-2) Choose $\sigma \in (0, 1)$ sufficiently small such that

$$
c_{h,V}\sigma^{\beta_1}=\frac{\theta_0}{16}.
$$

Clearly, σ ultimately depends only on n and p. Then, by Theorem [32,](#page-16-2) we deduce

$$
\left(\int_{B_{\sigma R_{i_{0}+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\sigma R_{i_{0}+1}}\right|^{q}\right)^{\frac{1}{q}}\leq \sup_{x,y\in B_{\sigma R_{i_{0}+1}}}\left|V\left(\mathfrak{X} u\right)\left(x\right)-V\left(\mathfrak{X} u\right)\left(y\right)\right|
$$
\n
$$
\leq c_{h,V}\sigma^{\beta_{1}}\left\|\mathfrak{X} u\right\|_{\infty,R_{i_{0}+1}}^{\frac{p}{2}}\leq \frac{\theta_{0}}{16}\left\|\mathfrak{X} u\right\|_{\infty,R_{i_{0}+1}}^{\frac{p}{2}}\leq \frac{\theta_{0}}{16}\left\|\mathfrak{X} u\right\|_{\infty,R_{i_{0}}}^{\frac{p}{2}}.
$$

This proves Claim [49.](#page-43-0) Now we are in a position to complete the proof of the Theorem. Note that, we trivially have for any $0 < \kappa \leq 1$

(136)
$$
\left| V^{-1} \left[(V(\mathfrak{X} u))_{\kappa R_{i_0+1}} \right] \right| = \left| (V(\mathfrak{X} u))_{\kappa R_{i_0+1}} \right|^{\frac{2}{p}} \leq \|\mathfrak{X} u\|_{\infty, R_{i_0}}.
$$

Now, observe that exactly one of the following can occur. Either we have

(137)
$$
\left| (V(\mathfrak{X}u))_{\frac{1}{2}\sigma R_{i_0+1}} \right|^{\frac{2}{p}} \geq \frac{1}{8B} \left\| \mathfrak{X}u \right\|_{\infty, R_{i_0}},
$$

or we must have

(138)
$$
\left| (V(\mathfrak{X}u))_{\frac{1}{2}\sigma R_{i_0+1}} \right|^{\frac{2}{p}} < \frac{1}{8B} \left\| \mathfrak{X}u \right\|_{\infty, R_{i_0}}.
$$

If [\(138\)](#page-44-0) holds, then by Theorem [30,](#page-15-7) we have

$$
\|\mathfrak{X}u\|_{\infty,\frac{1}{4}\sigma R_{i_0+1}} \leq c_{p/2} \left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}} |\mathfrak{X}u|^{\frac{p}{2}} \right)^{\frac{2}{p}} \leq c_{p/2} \left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}} |V(\mathfrak{X}u)|^q \right)^{\frac{2}{pq}}.
$$

Now, denoting $\Upsilon := (V(\mathfrak{X}u))_{\frac{1}{2}\sigma R_{i_0+1}}$ as a temporary shorthand and using the bound

(139)
$$
\frac{1}{4} \|\mathfrak{X} u\|_{\infty, R_{i_0}} \le \|\mathfrak{X} u\|_{\infty, \frac{1}{4}\sigma R_{i_0+1}},
$$

which is implied by [\(134\)](#page-43-1), in the last line of the following, we deduce

$$
\|\mathfrak{X}u\|_{\infty,\frac{1}{4}\sigma R_{i_0+1}} \leq c_{p/2} \left[\left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}} |V(\mathfrak{X}u) - \mathfrak{T}|^q \right)^{\frac{1}{q}} + |\mathfrak{T}| \right]^{\frac{2}{p}} \right]
$$

$$
\leq \frac{(138)}{\leq} 2^{\frac{1}{p}} c_{p/2} \left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}} |V(\mathfrak{X}u) - \mathfrak{T}|^q \right)^{\frac{2}{pq}} + \frac{2^{\frac{1}{p}} c_{p/2}}{8B} \|\mathfrak{X}u\|_{\infty, R_{i_0}}
$$

$$
\leq 2^{\frac{1}{p}} c_p \left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}} |V(\mathfrak{X}u) - \mathfrak{T}|^q \right)^{\frac{2}{pq}} + \frac{2^{\frac{1}{p}} c_{p/2}}{2B} \|\mathfrak{X}u\|_{\infty, \frac{1}{4}\sigma R_{i_0+1}}.
$$

Now by our choice of $B = 2^{1+\frac{1}{p}}c_{p/2}$ together with [\(140\)](#page-44-1) implies

$$
\|\mathfrak{X}u\|_{\infty,\frac{1}{4}\sigma R_{i_0+1}/2} \leq \frac{2^{2+\frac{1}{p}}c_{p/2}}{3}\left(\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}}\left|V\left(\mathfrak{X}u\right)-\left(V\left(\mathfrak{X}u\right)\right)_{\frac{1}{2}\sigma R_{i_0+1}}\right|^{q}\right)^{\frac{2}{pq}}.
$$

Once again, combined with Theorem [32,](#page-16-2) this implies, for any $0 < \rho \leq \sigma R_{i_a}/4 \leq$ $\sigma R_{i_0+1}/4$, we have

$$
\left(\oint_{B_{\rho}}\left|V\left(\mathfrak{X}u\right)-\left(V\left(\mathfrak{X}u\right)\right)_{\rho}\right|^{q}dx\right)^{\frac{1}{q}}\n\leq c\left(\frac{\rho}{\sigma R_{i_{0}+1}}\right)^{\beta_{1}}\left\|\mathfrak{X}u\right\|_{\infty,\frac{1}{4}\sigma R_{i_{0}+1}}^{\frac{p}{2}}\n\leq c\left(\frac{\rho}{\sigma R_{i_{0}+1}}\right)^{\beta_{1}}\left\|\mathfrak{X}u\right\|_{\infty,\frac{1}{4}\sigma R_{i_{0}+1}}^{\frac{p}{2}}\n\leq \frac{2^{p+2}c_{p/2}^{E}c}{3^{\frac{p}{2}}}\left(\frac{\rho}{\sigma R_{i_{0}+1}}\right)^{\beta_{1}}\left(\oint_{B_{\frac{1}{2}\sigma R_{i_{0}+1}}} \left|V\left(\mathfrak{X}u\right)-\left(V\left(\mathfrak{X}u\right)\right)_{\frac{1}{2}\sigma R_{i_{0}+1}}\right|^{q}\right)^{\frac{1}{q}}\n\leq \frac{(20)}{3^{\frac{p}{2}}}\frac{2^{p+3}c_{p/2}^{E}c}{3^{\frac{p}{2}}}\left(\frac{\rho}{\sigma R_{i_{0}+1}}\right)^{\beta_{1}}\left(\oint_{B_{\frac{1}{2}\sigma R_{i_{0}+1}}} \left|V\left(\mathfrak{X}u\right)-\left(V\left(\mathfrak{X}u\right)\right)_{R}\right|^{q}\right)^{\frac{1}{q}}.
$$
\n(141)
$$
\leq c\left(\frac{R}{\sigma R_{i_{0}+1}}\right)^{\frac{q}{q}+\beta_{1}}\left(\frac{\rho}{R}\right)^{\beta_{1}}\left(\oint_{B_{R}} \left|V\left(\mathfrak{X}u\right)-\left(V\left(\mathfrak{X}u\right)\right)_{R}\right|^{q}\right)^{\frac{1}{q}}.
$$

Now, since σ , τ_1 , β_1 are fixed constants and i_a is a fixed integer, all depending only on *n* and *p*, with $\sigma, \tau_1 \in (0, 1)$ and $i_0 < i_a$, we note that

$$
\left(\frac{R}{\sigma R_{i_0+1}}\right)^{\frac{Q}{q}+\alpha_1}=\left(\frac{1}{\sigma\tau_1^{i_0+1}}\right)^{\frac{Q}{q}+\alpha_1}\leq \left(\frac{1}{\sigma\tau_1^{i_a}}\right)^{\frac{Q}{q}+\alpha_1}\leq \left(\frac{1}{\sigma\tau_1^{i_a}}\right)^{Q+\alpha_1}
$$

,

where the last term on the right is again a fixed constant, depending only on n and p. Hence, [\(141\)](#page-45-0) now easily yields our desired estimate. Hence it only remains to settle the case when [\(137\)](#page-44-2) holds. Observe that [\(135\)](#page-43-2) implies

$$
\left(\int_{B_{\sigma R_{i_0+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\sigma R_{i_0+1}}\right|^q\right)^{\frac{1}{q}}\leq \frac{\theta_0}{16}\left\|\mathfrak{X} u\right\|_{\infty,R_{i_0}}^{\frac{p}{2}}\leq \theta_0\left\|\mathfrak{X} u\right\|_{\infty,R_{i_0+1}}^{\frac{p}{2}},
$$

where once again we have used the bound (139) . Combining this with the Hölder and reverse Hölder inequality, we deduce

$$
\left(\int_{B_{\frac{1}{2}\sigma R_{i_{0}+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\frac{1}{2}\sigma R_{i_{0}+1}}\right|^{2}\right)^{\frac{1}{2}}\n\leq 2\left(\int_{B_{\frac{1}{2}\sigma R_{i_{0}+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\sigma R_{i_{0}+1}}\right|^{2}\right)^{\frac{1}{2}}\n\leq 2\left(\int_{B_{\frac{1}{2}\sigma R_{i_{0}+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\sigma R_{i_{0}+1}}\right|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2Q}}\n\leq 2\left(\int_{B_{\frac{1}{2}\sigma R_{i_{0}+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\sigma R_{i_{0}+1}}\right|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2Q}}\n\leq 2C_{RH,V}\int_{B_{\sigma R_{i_{0}+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\sigma R_{i_{0}+1}}\right|^{q}\right)^{\frac{1}{q}}\leq 2C_{RH,V}\theta_{0}\left\|\mathfrak{X} u\right\|_{\infty,R_{i_{0}+1}}^{\frac{p}{2}}.
$$

Thus, recalling $\theta_0 \in (0,1)$, we have

$$
\int_{B_{\frac{1}{2}\sigma R_{i_0+1}}} \left| V(\mathfrak{X} u) - (V(\mathfrak{X} u))_{\frac{1}{2}\sigma R_{i_0+1}} \right|^2 \leq \theta_0^2 (2C_{RH,V})^2 \left\| \mathfrak{X} u \right\|_{\infty, R_{i_0+1}}^p
$$
\n(142)\n
$$
\leq \theta_0 (2C_{RH,V})^p \left\| \mathfrak{X} u \right\|_{\infty, R_{i_0+1}}^p.
$$

By virtue of [\(134\)](#page-43-1), we also trivially have

$$
\frac{1}{4B} \|\mathfrak{X}u\|_{\infty, R_{i_0+1}} \leq \frac{1}{4} \|\mathfrak{X}u\|_{\infty, R_{i_0}} \leq |\mathfrak{X}u| = |V(\mathfrak{X}u)|^{\frac{2}{p}} \leq 2C_{RH,V} \|\mathfrak{X}u\|_{\infty, R_{i_0+1}} \text{ in } B_{R_{i_0+1}}.
$$

Now note that [\(137\)](#page-44-2) immediately implies

$$
\frac{1}{8B} \|\mathfrak{X}u\|_{\infty, R_{i_0+1}} \leq \frac{1}{8B} \|\mathfrak{X}u\|_{\infty, R_{i_0}} \leq \left| (V(\mathfrak{X}u))_{\frac{1}{2}\sigma R_{i_0+1}} \right|^{\frac{2}{p}} \leq 2C_{RH,V} \|\mathfrak{X}u\|_{\infty, R_{i_0+1}}.
$$
\n(144)

Now, [\(143\)](#page-46-0), [\(144\)](#page-46-1) and [\(142\)](#page-46-2) implies that the hypotheses of Lemma [42](#page-31-0) are satisfied with $A = 2C_{RH,V}$ and $B = 2^{1+\frac{1}{p}}c_{p/2}$, R replaced by $R_{i_0+1}/2$, and $\lambda =$ $\|\mathfrak{X}u\|_{\infty,R_{i_0+1}}$, which we can always assume to be positive, as otherwise there is noth-ing to prove. Hence, Lemma [42](#page-31-0) implies that for any $0 < \rho \leq \sigma R_{i_a}/2 \leq \sigma R_{i_0+1}/2$, we have

$$
\int_{B_{\rho}} \left| V(\mathfrak{X}u) - (V(\mathfrak{X}u))_{\rho} \right|^{2}
$$
\n
$$
\leq C\sigma^{-Q-2\alpha_{0}} \left(\frac{\rho}{R_{i_{0}+1}} \right)^{2\alpha_{0}} \int_{B_{\frac{1}{2}R_{i_{0}+1}}} \left| V(\mathfrak{X}u) - (V(\mathfrak{X}u))_{\frac{1}{2}R_{i_{0}+1}} \right|^{2}.
$$

From here, to derive our desired estimate, we use Hölder inequality on one side and reverse Hölder inequality on the other. More precisely, by [\(145\)](#page-46-3), for any $0 < \rho \leq$ $\sigma R_{i_a}/2 \leq \sigma R_{i_0+1}/2$, we have

$$
\left(\int_{B_{\rho}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\rho}\right|^{q}\right)^{\frac{1}{q}}
$$
\nHölder\n
$$
\leq \left(\int_{B_{\rho}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\rho}\right|^{2}\right)^{\frac{1}{2}}
$$
\n
$$
\leq C\sigma^{-\frac{Q}{2}-\alpha_{0}}\left(\frac{\rho}{R_{i_{0}+1}}\right)^{\alpha_{0}}\left(\int_{B_{\frac{1}{2}R_{i_{0}+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\frac{1}{2}R_{i_{0}+1}}\right|^{2}\right)^{\frac{1}{2}}
$$
\nHölder\n
$$
\leq C\sigma^{-\frac{Q}{2}-\alpha_{0}}\left(\frac{\rho}{R_{i_{0}+1}}\right)^{\alpha_{0}}\left(\int_{\frac{1}{2}B_{R_{i_{0}+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\frac{1}{2}R_{i_{0}+1}}\right|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2Q}}
$$
\n
$$
\leq 2C\sigma^{-\frac{Q}{2}-\alpha_{0}}\left(\frac{\rho}{R_{i_{0}+1}}\right)^{\alpha_{0}}\left(\int_{B_{\frac{1}{2}R_{i_{0}+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{R_{i_{0}+1}}\right|^{\frac{2Q}{Q-2}}\right)^{\frac{Q-2}{2Q}}.
$$

Now by using [\(134\)](#page-43-1), [\(136\)](#page-44-4) and [\(70\)](#page-22-3), we continue the estimation

$$
\left(\int_{B_{\rho}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{\rho}\right|^{q}\right)^{\frac{1}{q}}
$$
\n
$$
\leq 2C_{RH,V}C\sigma^{-\frac{Q}{2}-\alpha_{0}}\left(\frac{\rho}{R_{i_{0}+1}}\right)^{\alpha_{0}}\int_{B_{R_{i_{0}+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{R_{i_{0}+1}}\right|
$$
\nHölder\n
$$
\leq 2C_{RH,V}C\sigma^{-\frac{Q}{2}-\alpha_{0}}\left(\frac{\rho}{R_{i_{0}+1}}\right)^{\alpha_{0}}\left(\int_{B_{R_{i_{0}+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{R_{i_{0}+1}}\right|^{q}\right)^{\frac{1}{q}}
$$
\n
$$
\leq 4C_{RH,V}C\sigma^{-\frac{Q}{2}-\alpha_{0}}\left(\frac{\rho}{R_{i_{0}+1}}\right)^{\alpha_{0}}\left(\int_{B_{R_{i_{0}+1}}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{R}\right|^{q}\right)^{\frac{1}{q}}
$$
\n
$$
\leq 4C_{RH,V}C\left(\frac{R}{R_{i_{0}+1}}\right)^{\frac{Q}{q}}\sigma^{-\frac{Q}{2}-\alpha_{0}}\left(\frac{\rho}{R_{i_{0}+1}}\right)^{\alpha_{0}}\left(\int_{B_{R}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{R}\right|^{q}\right)^{\frac{1}{q}}
$$
\n
$$
\leq 4C_{RH,V}C\left(\frac{R}{R_{i_{0}+1}}\right)^{\frac{Q}{q}+\alpha_{0}}\sigma^{-\frac{Q}{2}-\alpha_{0}}\left(\frac{\rho}{R}\right)^{\alpha_{0}}\left(\int_{B_{R}}\left|V\left(\mathfrak{X} u\right)-\left(V\left(\mathfrak{X} u\right)\right)_{R}\right|^{q}\right)^{\frac{1}{q}}
$$
\n
$$
\leq 4C_{RH,V}C\left(\frac{2
$$

Since $C_{RH,V}, \sigma, \alpha_0, \tau_1, i_a$ are all fixed, which ultimately depend only on n and p, we derive our desired estimate. This completes the proof. \Box

4. HÖLDER CONTINUITY

4.1. **Preliminary estimates.** Let $x_0 \in \Omega$ and $0 < R < 1$ be such that $B(x_0, 2R) \subset$ Ω. Clearly, if $u ∈ HW^{1,p}_{loc}(\Omega)$ is a weak solution to [\(9\)](#page-5-2), then $u ∈ HW^{1,p}(B_R)$ is a weak solution to

(146)
$$
\operatorname{div}_{\mathbb{H}}(a(x)|\mathfrak{X}u|^{p-2}\mathfrak{X}u) = \operatorname{div}_{\mathbb{H}} F \qquad \text{in } B_R.
$$

Now we define $w \in u + HW_0^{1,p}(B_R)$ to be the unique solution of

(147)
$$
\begin{cases} \operatorname{div}_{\mathbb{H}} \left(a(x) | \mathfrak{X}w \right|^{p-2} \mathfrak{X}w = 0 & \text{in } B_R, \\ w = u & \text{on } \partial B_R, \end{cases}
$$

and $v \in w + HW_0^{1,p}(B_R)$ to be the unique solution of

(148)
$$
\begin{cases} \operatorname{div}_{\mathbb{H}} \left(a(x_0) | \mathfrak{X} v |^{p-2} \mathfrak{X} v \right) = 0 & \text{in } B_R, \\ v = w & \text{on } \partial B_R. \end{cases}
$$

Now we record an easy comparison estimate, whose proof follows from the weak formulation of [\(146\)](#page-47-1) and [\(147\)](#page-47-2), the estimates [\(36\)](#page-14-0), [\(37\)](#page-14-1) and Young's inequality.

Lemma 50. Let $1 < p < \infty$, u be as in [\(146\)](#page-47-1) and w be as in [\(147\)](#page-47-2). Then for any $\xi \in \mathbb{R}^{2n}$, we have the following inequality

(149)
$$
\int_{B_R} |V(\mathfrak{X} u) - V(\mathfrak{X} w)|^2 \leq c \int_{B_R} |F - \xi| |\mathfrak{X} u - \mathfrak{X} w|,
$$

where $c > 0$ is a constant depending on n, p, ν and L.

The following is proved analogously to Lemma 3 in [\[42\]](#page-64-18).

Lemma 51. Let $1 < p < \infty$, w be as in [\(147\)](#page-47-2) and v be as in [\(148\)](#page-47-3). Then there exists a constant $c \equiv c(n, p, \gamma, L)$ such that we have the inequality

(150)
$$
\int_{B_R} |V(\mathfrak{X}v) - V(\mathfrak{X}w)|^2 \leq c \left[\omega(R)\right]^2 \int_B |\mathfrak{X}w|^p.
$$

4.2. Proofs of Theorem [3](#page-5-0) and Corollary [4.](#page-5-1) Now we are in a position to prove the Hölder continuity results.

Proof. (of Theorem [3](#page-5-0)) Since all our estimates are local, we can assume $a \in$ $C^{0,\mu_1}(\Omega)$, $F \in C^{0,\mu_2}(\Omega;\mathbb{R}^{2n})$ and $u \in HW^{1,p}(\Omega)$. We first focus on the case $1 < p \leq 2$, the other case being much easier. We divide the proof in two steps.

Step 1: First, we are going to show $\mathfrak{X}u \in L^{p,Q-\delta}_{loc}(\Omega,\mathbb{R}^{2n})$ for every $\delta > 0$, under the assumptions of the theorem. We first choose $x_0 \in \Omega$ and radii $\rho, R > 0$ such that $0 < 4\rho < R$ and $B_R(x_0) \subset\subset \Omega$. Now we define the functions w and v the same way as before in B_R by using [\(147\)](#page-47-2) and [\(148\)](#page-47-3) respectively. Now using [\(43\)](#page-15-8), by standard calculations, for any $1 < p < \infty$, we have

(151)
$$
\int_{B_{\rho}} |\mathfrak{X}u|^p \leq c \left(\frac{\rho}{R}\right)^Q \int_{B_R} |\mathfrak{X}u|^p + c \int_{B_R} |\mathfrak{X}u - \mathfrak{X}v|^p.
$$

Now, since $1 < p \leq 2$, using [\(38\)](#page-14-2), we deduce

$$
\int_{B_R} |\mathfrak{X}w - \mathfrak{X}v|^p \le c \int_{B_R} |V(\mathfrak{X}w) - V(\mathfrak{X}v)|^2 + c \int_{B_R} |V(\mathfrak{X}w) - V(\mathfrak{X}v)|^p |\mathfrak{X}w|^{\frac{p(2-p)}{2}}.
$$

Using Young's inequality with $\varepsilon > 0$, together with [\(150\)](#page-48-0), for $1 < p < 2$, we deduce

$$
\int_{B_R} |\mathfrak{X}w - \mathfrak{X}v|^p \leq \varepsilon \int_{B_R} |\mathfrak{X}w|^p + c \left(1 + \varepsilon^{\frac{p-2}{p}}\right) \int_{B_R} |V(\mathfrak{X}w) - V(\mathfrak{X}v)|^2
$$

$$
\leq \varepsilon \int_{B_R} |\mathfrak{X}w|^p + c \left(1 + \varepsilon^{\frac{p-2}{p}}\right) \left[\omega(R)\right]^2 \int_{B_R} |\mathfrak{X}w|^p.
$$

The inequality above trivially holds for $p = 2$. Note that w minimizes the functional

$$
w \mapsto \frac{1}{p} \int_{B_R} a(x) |\mathfrak{X}w|^p
$$
 in $u + HW_0^{1,p}(B_R)$.

Thus, by minimality and the bounds on a , we have

(152)
$$
\frac{\gamma}{p} \int_{B_R} |\mathfrak{X}w|^p \leq \frac{1}{p} \int_{B_R} a(x) |\mathfrak{X}w|^p \leq \frac{1}{p} \int_{B_R} a(x) |\mathfrak{X}u|^p \leq \frac{L}{p} \int_{B_R} |\mathfrak{X}u|^p.
$$

Using this in the last estimate, we arrive at

(153)
$$
\int_{B_R} |\mathfrak{X}w - \mathfrak{X}v|^p \leq \frac{cL}{\gamma} \left(\varepsilon + \varepsilon^{\frac{p-2}{p}} \left[\omega(R) \right]^2 \right) \int_{B_R} |\mathfrak{X}u|^p.
$$

For the other term, we have, again using [\(38\)](#page-14-2)

$$
\int_{B_R} |\mathfrak{X} u - \mathfrak{X} w|^p \le c \int_{B_R} |V(\mathfrak{X} u) - V(\mathfrak{X} w)|^2 + \int_{B_R} |V(\mathfrak{X} u) - V(\mathfrak{X} w)|^p |\mathfrak{X} u|^{\frac{p(2-p)}{2}} \n\le \varepsilon \int_{B_R} |\mathfrak{X} u|^p + c \left(1 + \varepsilon^{\frac{p-2}{p}}\right) \int_{B_R} |V(\mathfrak{X} u) - V(\mathfrak{X} w)|^2 \n\overset{(149)}{\le} \varepsilon \int_{B_R} |\mathfrak{X} u|^p + c \varepsilon^{\frac{p-2}{p}} \int_{B_R} |F - \xi| |\mathfrak{X} u - \mathfrak{X} w| \n\le \frac{1}{2} \int_{B_R} |\mathfrak{X} u - \mathfrak{X} w|^p + \varepsilon \int_{B_R} |\mathfrak{X} u|^p + c \varepsilon^{\frac{p-2}{p-1}} \int_{B_R} |F - \xi|^{\frac{p}{p-1}}.
$$

Thus, we deduce

(154)
$$
\int_{B_R} |\mathfrak{X} u - \mathfrak{X} w|^p \leq 2\varepsilon \int_{B_R} |\mathfrak{X} u|^p + c\varepsilon^{\frac{p-2}{p-1}} \int_{B_R} |F - \xi|^{\frac{p}{p-1}}.
$$

Combining [\(154\)](#page-49-0) and [\(153\)](#page-48-2), with the shorthand

(155)
$$
C_{\varepsilon,R} := 2^{p-1} \left[\left(2 + \frac{cL}{\gamma} \right) \varepsilon + \frac{cL}{\gamma \varepsilon^{\frac{2-p}{p}}} \left[\omega(R) \right]^2 \right],
$$

for some $c \equiv c(n, p, \gamma, L) > 0$, coming from [\(154\)](#page-49-0) and [\(153\)](#page-48-2), we have,

(156)
$$
\int_{B_R} |\mathfrak{X} u - \mathfrak{X} v|^p \leq 2^{p-1} \left(\int_{B_R} |\mathfrak{X} u - \mathfrak{X} w|^p + \int_{B_R} |\mathfrak{X} w - \mathfrak{X} v|^p \right) \leq C_{\varepsilon, R} \int_{B_R} |\mathfrak{X} u|^p + c \varepsilon^{\frac{p-2}{p-1}} \int_{B_R} |F - \xi|^{\frac{p}{p-1}}.
$$

Now, plugging this with the choice $\xi = 0$, in [\(151\)](#page-48-3) we have

$$
\int_{B_{\rho}} |\mathfrak{X}u|^{p} \leq c \left(\frac{\rho}{R}\right)^{Q} \int_{B_{R}} |\mathfrak{X}u|^{p} + C_{\varepsilon,R} \int_{B_{R}} |\mathfrak{X}u|^{p} + c \varepsilon^{\frac{p-2}{p-1}} \int_{B_{R}} |F|^{\frac{p}{p-1}} \leq c \left(\frac{\rho}{R}\right)^{Q} \int_{B_{R}} |\mathfrak{X}u|^{p} + C_{\varepsilon,R} \int_{B_{R}} |\mathfrak{X}u|^{p} + c \varepsilon^{\frac{p-2}{p-1}} R^{Q} \|F\|_{L^{\infty}(\Omega; \mathbb{R}^{2n})}^{\frac{p}{p-1}}
$$

.

Now we use the standard iteration lemma (see Lemma 5.13 in [\[20\]](#page-63-21)). We fix $\delta > 0$ and use Lemma 5.13 in [\[20\]](#page-63-21) with $\alpha = Q, \beta = Q - \delta$ and

$$
\phi(r) := \int_{B_r} |\mathfrak{X} u|^p.
$$

Now, in view of [\(155\)](#page-49-1), we first choose $\varepsilon > 0$ small enough and then choose R small enough (depending on ε) such that $C_{\varepsilon,R}$ above is less then the threshold ε_0 , given by the interation Lemma. This yields,

$$
\frac{1}{\rho^{Q-\delta}}\int_{B_{\rho}}|\mathfrak{X} u|^p\leq c\left(\|\mathfrak{X} u\|_{L^p(\Omega;\mathbb{R}^{2n})}^p+\|F\|_{L^{\infty}(\Omega;\mathbb{R}^{2n})}^{\frac{p}{p-1}}\right).
$$

Note that, here $c > 0$ depends on n, p, ν, L and δ . By the standard covering argument, this implies

$$
\|\mathfrak{X}u\|_{\mathrm{L}^{p,Q-\delta}(\Omega_2;\mathbb{R}^{2n})}\leq c\left(\|\mathfrak{X}u\|_{L^p(\Omega_1;\mathbb{R}^{2n})}+\|F\|_{L^{\infty}(\Omega_1;\mathbb{R}^{2n})}^{\frac{1}{p-1}}\right),
$$

for any $\Omega_2 \subset\subset \Omega_1 \subset\subset \Omega$, where the constant c now depends also on Ω_2 and Ω_1 .

Step 2: Now we finish the proof of Hölder continuity of $\mathfrak{X}u$. With the Morrey bound at our disposal, we return to the comparison estimates and estimate them differently this time. We have the following estimate for some constant $c \equiv c(n, p, \nu, L) > 0$,

$$
\int_{B_R} |\mathfrak{X}w - \mathfrak{X}v|^p
$$
\n
$$
\leq c \int_{B_R} |V(\mathfrak{X}w) - V(\mathfrak{X}v)|^2 + c \int_{B_R} |V(\mathfrak{X}w) - V(\mathfrak{X}v)|^p |\mathfrak{X}w|^{\frac{p(2-p)}{2}}
$$
\n
$$
\leq c \int_{B_R} |V(\mathfrak{X}w) - V(\mathfrak{X}v)|^2 + c \left(\int_{B_R} |V(\mathfrak{X}w) - V(\mathfrak{X}v)|^2 \right)^{\frac{p}{2}} \left(\int_{B_R} |\mathfrak{X}w|^p \right)^{\frac{2-p}{2}}
$$
\n
$$
\leq c \left[\omega(R) \right]^2 \int_{B_R} |\mathfrak{X}w|^p + \left([\omega(R)]^2 \int_{B_R} |\mathfrak{X}w|^p \right)^{\frac{p}{2}} \left(\int_{B_R} |\mathfrak{X}w|^p \right)^{\frac{2-p}{2}}
$$
\n
$$
\leq c \left([\omega(R)]^2 + [\omega(R)]^p \right) \int_{B_R} |\mathfrak{X}w|^p.
$$

In view of [\(152\)](#page-48-4), this implies

(157)
$$
\int_{B_R} |\mathfrak{X}w - \mathfrak{X}v|^p \le c \left(\left[\omega(R) \right]^2 + \left[\omega(R) \right]^p \right) \int_{B_R} |\mathfrak{X}u|^p.
$$

As a is $C^{0,\mu}$ and $p \leq 2$, this implies

(158)
$$
\int_{B_R} |\mathfrak{X}w - \mathfrak{X}v|^p \le c(n, p, \nu, L, \delta) R^{p\mu + Q - \delta} ||\mathfrak{X}u||_{L^{p,Q-\delta}}^p.
$$

For the other comparison estimate, for some constant $c \equiv c(n, p, \nu, L) > 0$, we have

$$
\int_{B_R} |\mathfrak{X} u - \mathfrak{X} w|^p
$$
\n
$$
\leq c \int_{B_R} |V(\mathfrak{X} u) - V(\mathfrak{X} w)|^2 + c \int_{B_R} |V(\mathfrak{X} u) - V(\mathfrak{X} w)|^p |\mathfrak{X} u|^{\frac{p(2-p)}{2}}
$$
\n
$$
\leq c \int_{B_R} |V(\mathfrak{X} u) - V(\mathfrak{X} w)|^2 + c \left(\int_{B_R} |V(\mathfrak{X} u) - V(\mathfrak{X} w)|^2 \right)^{\frac{p}{2}} \left(\int_{B_R} |\mathfrak{X} u|^p \right)^{\frac{2-p}{2}}.
$$

Now using [\(149\)](#page-48-1) on the right, we deduce

$$
\int_{B_R} |\mathfrak{X} u - \mathfrak{X} w|^p
$$
\n
$$
\leq c \int_{B_R} |F - \xi| |\mathfrak{X} u - \mathfrak{X} w| + c \left(\int_{B_R} |F - \xi| |\mathfrak{X} u - \mathfrak{X} w| \right)^{\frac{p}{2}} \left(\int_{B_R} |\mathfrak{X} u|^p \right)^{\frac{2-p}{2}}
$$
\n
$$
\leq c \int_{B_R} |F - \xi| |\mathfrak{X} u - \mathfrak{X} w|
$$
\n
$$
+ c \left(\int_{B_R} |F - \xi| \frac{p}{p-1} \right)^{\frac{p-1}{2}} \left(\int_{B_R} |\mathfrak{X} u - \mathfrak{X} w|^p \right)^{\frac{1}{2}} \left(\int_{B_R} |\mathfrak{X} u|^p \right)^{\frac{2-p}{2}}
$$
\n
$$
\leq \frac{1}{2} \int_{B_R} |\mathfrak{X} u - \mathfrak{X} w|^p + c \int_{B_R} |F - \xi|^\frac{p}{p-1}
$$
\n
$$
+ c \left(\int_{B_R} |F - \xi|^\frac{p}{p-1} \right)^{p-1} \left(\int_{B_R} |\mathfrak{X} u|^p \right)^{2-p}.
$$

Thus, we arrive at

(159)
$$
\int_{B_R} |\mathfrak{X} u - \mathfrak{X} w|^p \le c \int_{B_R} |F - \xi|^{\frac{p}{p-1}} + c \left(\int_{B_R} |F - \xi|^{\frac{p}{p-1}} \right)^{p-1} \left(\int_{B_R} |\mathfrak{X} u|^p \right)^{2-p}
$$

for some constant $c \equiv c(n, p, \nu, L) > 0$. Now choosing $\xi = (F)_{B_R}$ and recalling the fact that $F \in C^{0,\mu}$ and the Campanato characterization of Hölder continuity, we have

$$
\int_{B_R} |\mathfrak{X} u - \mathfrak{X} w|^p
$$
\n
$$
\leq R^{Q + \frac{p\mu}{p-1}} [F]_{\mathcal{L}^{\frac{p}{p-1}, Q + \frac{p\mu}{p-1}}}^{\frac{p}{p-1}} + cR^{[(p-1)Q + p\mu]} [F]_{\mathcal{L}^{\frac{p}{p-1}, Q + \frac{p\mu}{p-1}}}^p \cdot R^{(Q-\delta)(2-p)} \|\mathfrak{X} u\|_{\mathcal{L}^{p,Q-\delta}}^{p(2-p)}
$$
\n
$$
\leq R^{Q + \frac{p\mu}{p-1}} [F]_{\mathcal{L}^{\frac{p}{p-1}, Q + \frac{p\mu}{p-1}}}^{\frac{p}{p-1}} + R^{Q + p\mu - \delta(2-p)} [F]_{\mathcal{L}^{\frac{p}{p-1}, Q + \frac{p\mu}{p-1}}}^p \|\mathfrak{X} u\|_{\mathcal{L}^{p,Q-\delta}}^{p(2-p)}.
$$

Now if $1 < p < 2$, we apply Young's inequality on the last term on the right with the the exponents $(1/(p-1), 1/(2-p))$, to arrive at

$$
(160) \qquad \int_{B_R} |\mathfrak{X} u - \mathfrak{X} w|^p \le c R^{Q + p\mu - \delta(2-p)} \left([F]_{\mathcal{L}^{\frac{p}{p-1}, Q + \frac{p\mu}{p-1}}}^{\frac{p}{p-1}} + ||\mathfrak{X} u||_{\mathcal{L}^{p, Q - \delta}}^p \right),
$$

where we used the fact that since $p < 2$, we have $Q + \frac{p\mu}{p-1} > Q + p\mu - \delta(2-p)$. Note that, [\(160\)](#page-51-0) holds trivially if $p = 2$. Now, since $1 < p \leq 2$, using [\(6\)](#page-3-3) of Theorem [1](#page-3-0) with $q = p$ we have, in the standard way,

$$
\int_{B_{\rho}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\rho} \right|^{p} \le c \left(\frac{\rho}{R} \right)^{Q + p\alpha} \int_{B_{R}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{R} \right|^{p} + \int_{B_{R}} \left| \mathfrak{X} u - \mathfrak{X} v \right|^{p}.
$$

Utilizing [\(160\)](#page-51-0) and [\(158\)](#page-50-0) in this, we deduce

$$
\int_{B_{\rho}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{\rho} \right|^{p} \leq c \left(\frac{\rho}{R} \right)^{Q+p\alpha} \int_{B_{R}} \left| \mathfrak{X} u - (\mathfrak{X} u)_{R} \right|^{p} + cR^{Q+p\mu-\delta} \left\| \mathfrak{X} u \right\|_{\mathbb{L}^{p,Q-\delta}}^{p} \n+ cR^{Q+p\mu-\delta(2-p)} \left(\left[F \right]_{\mathcal{L}}^{\frac{p}{p-1},Q+\frac{p\mu}{p-1}} + \left\| \mathfrak{X} u \right\|_{\mathbb{L}^{p,Q-\delta}}^{p} \right).
$$

Note that for $1 < p \leq 2$, the smallest exponent of R on the right is $Q + p\mu - \delta$. Thus, choosing $0 < \delta < p\mu$, using Lemma 5.13 in [\[20\]](#page-63-21) and standard covering arguments, together with Camapanto's characterization of Hölder continuity, the estimate above implies $\mathfrak{X}u \in C_{\text{loc}}^{0,\mu-\frac{\delta}{p}}$ for any $0 < \delta < p\mu$. But then we also have $\mathfrak{X}u \in L^{\infty}_{\text{loc}}$ and consequently, $\mathfrak{X}u \in L^{p,Q}_{\text{loc}}$. Now we go back to [\(157\)](#page-50-1) and [\(159\)](#page-51-1) and derive [\(158\)](#page-50-0) and [\(160\)](#page-51-0) with $\delta = 0$ with constants depending only on n, p, ν and L. Consequently, we have proved $\mathfrak{X}u \in C^{0,\mu}_{loc}$. Tracing back our estimates, we have the estimate

$$
\left\|\mathfrak{X}u\right\|_{C^{0,\mu}\left(\overline{\Omega_2};\mathbb{R}^{2n}\right)} \leq C\left(\left\|\mathfrak{X}u\right\|_{L^p\left(\Omega_1;\mathbb{R}^{2n}\right)}+\left\|F\right\|_{C^{0,\mu}\left(\overline{\Omega_1};\mathbb{R}^{2n}\right)}\right),
$$

for any $\Omega_2 \subset\subset \Omega_1 \subset\subset \Omega$, where the constant C now depends also on Ω_2 and Ω_1 .

Now we prove the case $p > 2$. This is quite easy. We use Lemma [50](#page-48-5) to derive

$$
\int_{B_R} |V(\mathfrak{X}u) - V(\mathfrak{X}w)|^2 \le c \int_{B_R} |F - \xi| |\mathfrak{X}u - \mathfrak{X}w|
$$
\n
$$
\le \varepsilon \int_{B_R} |\mathfrak{X}u - \mathfrak{X}w|^p + c \int_{B_R} |F - \xi|^{\frac{p}{p-1}}
$$
\n
$$
\stackrel{(36)}{\le} c_V \varepsilon \int_{B_R} |V(\mathfrak{X}u) - V(\mathfrak{X}w)|^2 + c \int_{B_R} |F - \xi|^{\frac{p}{p-1}}
$$

Choosing $\varepsilon > 0$ small enough, we have

(161)
$$
\int_{B_R} |V(\mathfrak{X}u) - V(\mathfrak{X}w)|^2 \leq c \int_{B_R} |F - \xi|^{\frac{p}{p-1}}.
$$

Similarly, using [\(152\)](#page-48-4) in [\(150\)](#page-48-0), we obtain

(162)
$$
\int_{B_R} |V(\mathfrak{X}w) - V(\mathfrak{X}v)|^2 \le c \left[\omega(R)\right]^2 \int_{B_R} |\mathfrak{X}u|^p = c \left[\omega(R)\right]^2 \int_{B_R} |V(\mathfrak{X}u)|^2.
$$
 Now, using (151) and choosing $\xi = 0$ we have

$$
\int_{B_{\rho}} |V(\mathfrak{X}u)|^2 = \int_{B_{\rho}} |\mathfrak{X}u|^p \le c \left(\frac{\rho}{R}\right)^Q \int_{B_R} |\mathfrak{X}u|^p + c \int_{B_R} |\mathfrak{X}u - \mathfrak{X}v|^p
$$

$$
\stackrel{(36)}{\le} c \left(\frac{\rho}{R}\right)^Q \int_{B_R} |V(\mathfrak{X}u)|^2 + c \int_{B_R} |V(\mathfrak{X}u) - V(\mathfrak{X}v)|^2
$$

$$
\stackrel{(161),(162)}{\le} c \left[\left(\frac{\rho}{R}\right)^Q + [\omega(R)]^2\right] \int_{B_R} |V(\mathfrak{X}u)|^2 + c \int_{B_R} |F|^{\frac{p}{p-1}}.
$$

As before, choosing R small enough and using the iteration Lemma 5.13 in [\[20\]](#page-63-21), this implies $V(\mathfrak{X}u) \in L^{2,Q-\delta}_{loc}$ for every $\delta > 0$. Now, we have, using [\(7\)](#page-3-4) of Theorem [2](#page-3-2) with $q=2$,

$$
\int_{B_{\rho}} \left| V(\mathfrak{X}u) - (V(\mathfrak{X}u))_{\rho} \right|^{2}
$$
\n
$$
\leq c \left(\frac{\rho}{R} \right)^{Q+2\beta} \int_{B_{R}} |V(\mathfrak{X}u) - (V(\mathfrak{X}u))_{R}|^{2} + cR^{Q+2\mu-\delta} ||V(\mathfrak{X}u)||_{\mathbb{L}^{2,Q-\delta}}^{2}
$$
\n
$$
+ cR^{Q+\frac{p\mu}{p-1}} [F]_{\mathcal{L}^{\frac{p}{p-1},Q+\frac{p\mu}{p-1}}}^{\frac{p}{p-1}}.
$$

Recall that $\mu < \beta$. Also, since $p > 2$ implies $p/(p-1) < 2$, we can choose $\delta > 0$ small enough such that the minimum power of R on the right is $Q + \frac{p\mu}{p-1}$. This, by standard iteration lemma and covering arguments imply $V(\mathfrak{X}u) \in \mathcal{L}_{loc}^{2,Q+\frac{p\mu}{p-1}}$. Hence, using [\(36\)](#page-14-0) we conclude that $\mathfrak{X}u \in \mathcal{L}_{loc}^{p,Q+\frac{p\mu}{p-1}}$. By Campanato's characterization, this implies $\mathfrak{X}u \in C_{\text{loc}}^{0, \frac{\mu}{\mu-1}}$. This completes the proof.

Proof. (of Corollary [4\)](#page-5-1) This follows immediately from Theorem [3](#page-5-0) as given any $f \in L^q_{\text{loc}}(\Omega)$, with $q > Q$, we can always write it as a divergence of a locally Hölder continuous vector field. More precisely, we can find $F \in C^{0, \frac{q-Q}{q}}_{\text{loc}}(\Omega; \mathbb{R}^{2n})$ such that div $_{\mathbb{H}} F = f$ in Ω . Indeed, we just find a solution of

$$
\operatorname{div}_{\mathbb{H}} (\mathfrak{X} \psi) = f \text{ in } \Omega,
$$

and set $F = \mathfrak{X}\psi$. By standard L^p estimates for the subLaplacian, (see [\[15,](#page-63-25) page 917]) we deduce $F \in HW^{1,q}_{loc}(\Omega;\mathbb{R}^{2n})$. Since $q > Q$, by Proposition [17,](#page-11-1) we have $F \in C_{\text{loc}}^{0, \frac{q-Q}{q}}(\Omega; \mathbb{R}^{2n})$. This completes the proof.

5. Homogeneous equation with Dini coefficients

5.1. General setting and comparison estimates. In this section, our goal is to prove Theorem [6.](#page-6-1) Let $B(x_0, 2R) \subset \Omega$ be a fixed ball and for $i \geq 0$, we set

(163)
$$
B_i \equiv B(x_0, R_i), \qquad R_i := \sigma^i R, \qquad \sigma \in (0, \frac{1}{2}).
$$

Let w be a weak solution of [\(13\)](#page-6-2). We define the maps $v_i \in w + HW_0^{1,p}(B_i)$ to be the unique solution of

(164)
$$
\begin{cases} \operatorname{div}_{\mathbb{H}}(a(x_0)|\mathfrak{X}v_i|^{p-2}\mathfrak{X}v_i)) = 0 & \text{in } B_i, \\ v_i = w & \text{on } \partial B_i. \end{cases}
$$

Also, we define the quantities, for $i \geq 0$ and $r \geq 1$,

$$
m_i(G) := |(G)_{B_i}|
$$
 and $E_r(G, B_i) := \left(\oint_{B_i} |G - (G)_{B_i}|^r\right)^{\frac{1}{r}}.$

Now we record the following comparison estimate, which is proved exactly as Lemma 3 in [\[42\]](#page-64-18).

Lemma 52. Assume that w, v_i satisfy [\(13\)](#page-6-2) and [\(164\)](#page-53-1) respectively and $i \geq 0$. Then there exists a constant $c_4 \equiv c_4(n, p, \gamma, L)$ such that we have the inequality

(165)
$$
\int_{B_i} |V(\mathfrak{X} v_i) - V(\mathfrak{X} w)|^2 \le c_4 \left[\omega(R_i) \right]^2 \int_{B_i} |\mathfrak{X} w|^p.
$$

5.2. Proof of Theorem [6.](#page-6-1) Now we prove the following pointwise estimate, which immediately implies the sup estimate.

Theorem 53. Let $w \in HW_{loc}^{1,p}(\Omega)$ be be a local weak solution of [\(13\)](#page-6-2) with $1 <$ $p < \infty$. Then there exists a constant $c = c(n, p, \gamma, L, \omega(\cdot)) \geq 1$ and a positive radius $R_1 = R_1(n, p, \gamma, L, \omega(\cdot)) > 0$ such that the pointwise estimate

(166)
$$
|\mathfrak{X}w(x)| \leq c \left(\int_{B(x,R)} |\mathfrak{X}w|^p \right)^{\frac{1}{p}},
$$

holds whenever $B(x, 2R) \subset \Omega$, $2R \le R_1$ and x is a Lebesgue point of $\mathfrak{X}w$. If $a(\cdot)$ is a constant function, the estimate holds without any restriction on R.

Proof. Step 1: Choice of constants. We pick an arbitrary point $x_0 \in \Omega$ and a arbitrary positive radius $R_1 > 0$ such that $B(x_0, R_1) \subset \Omega$. We pick $0 < R < R_1/2$ and for now set $B(x_0, R)$ as our starting ball and consider the chain of shrinking balls as explained in [\(163\)](#page-53-2) for some parameter $\sigma \in (0, \frac{1}{4})$. We shall soon make specific choices of both the parameters R_1 and σ . We define the constant λ as

(167)
$$
\lambda^{\frac{p}{2}} := H_1 \left(\int_{B(x_0,R)} |V(\mathfrak{X}w)|^2 \right)^{\frac{1}{2}},
$$

where H_1 will be chosen soon. Clearly, we can assume $\lambda > 0$. In view of Theorem [2,](#page-3-2) we choose $\sigma \in (0, \frac{1}{4})$ small enough such that

(168)
$$
C_{0,V} \sigma^{\beta} \leq 4^{-(Q+4)}.
$$

Now that we have chosen σ , we set

(169)
$$
H_1 := 10^{5Q} \sigma^{-2Q}.
$$

Note that H_1 depends only on n and p. Now, we chose the radius $R_1 > 0$ small enough such that we have

(170)
$$
\omega(R_1) + \int_0^{2R_1} \omega(\varrho) \frac{d\varrho}{\varrho} \le \frac{\sigma^{2Q}}{6^Q 10^6 c_4}.
$$

Note that R_1 depends on n, p, γ, L and $\omega(\cdot)$. Also, if $a(\cdot)$ is a constant function, the dependence on $\omega(\cdot)$ is redundant. With this, we have chosen all the relevant parameters.

Step 2: Excess decay. We now want to prove

Claim 54 (excess decay estimate). If

(171)
$$
\left(\int_{B_i} |\mathfrak{X}w|^p\right)^{\frac{1}{p}} \leq \lambda,
$$

then

(172)
$$
E_2(V(\mathfrak{X}w), B_{i+1}) \leq \frac{1}{4} E_2(V(\mathfrak{X}w), B_i) + \frac{2c_4\lambda^{\frac{p}{2}}}{\sigma^Q} \omega(R_i),
$$

where c_4 is given by Lemma [52.](#page-53-3)

By the excess decay estimate [\(7\)](#page-3-4) of Theorem [2](#page-3-2) with $q = 2$ and the choice of σ in [\(168\)](#page-54-0), we have,

(173)
$$
E_2(V(\mathfrak{X}v_i), B_{i+1}) \leq \frac{1}{4^{Q+4}} E_2(V(\mathfrak{X}v_i), B_i).
$$

Now using the property of the mean, triangle inequality and elementary estimates along with Lemma [52,](#page-53-3) we have,

$$
E_2(V(\mathfrak{X}w), B_{i+1}) \le E_2(V(\mathfrak{X}v_i), B_{i+1}) + \sigma^{-Q/2} \left(\int_{B_i} |V(\mathfrak{X}v_i) - V(\mathfrak{X}w)|^2 \right)^{\frac{1}{2}}
$$

(174)

$$
\le \sum_{i=1}^{(165),(171)} E_2(V(\mathfrak{X}v_i), B_{i+1}) + \sigma^{-Q} c_4 \omega(R_i) \lambda^{\frac{p}{2}}.
$$

Also similarly,

(175)
$$
E_2(V(\mathfrak{X}v_i), B_i) \stackrel{(165),(171)}{\leq} E_2(V(\mathfrak{X}w), B_i) + c_4 \omega(R_i) \lambda^{\frac{p}{2}}.
$$

Combining [\(173\)](#page-54-2), [\(174\)](#page-54-3) and [\(175\)](#page-54-4), we have [\(172\)](#page-54-5).

Step 3: Control on composite quantities. Now we want to show, by induction that

(176)
$$
m_i(V(\mathfrak{X}w)) + E_2(V(\mathfrak{X}w), B_i) \le \lambda^{\frac{p}{2}} \quad \text{for all } i \ge 1.
$$

This is true for $i = 1$ by our choice of H_1 in [\(169\)](#page-54-6) and elementary estimates. Thus we assume this is true for all $j \in \{1, \ldots, i\}$ and prove it for $i+1.$ But we have ,

$$
\int_{B_j} |\mathfrak{X}w|^p = \int_{B_j} |V(\mathfrak{X}w)|^2 \leq [m_j(V(\mathfrak{X}w))]^2 + [E_2(V(\mathfrak{X}w), B_j)]^2
$$

$$
\leq [m_j(V(\mathfrak{X}w)) + E_2(V(\mathfrak{X}w), B_j)]^2.
$$

Thus the induction hypotheses implies the bound

$$
\left(\oint_{B_j} |\mathfrak{X}w|^p\right)^{\frac{1}{p}} \leq \lambda \text{ for all } j \in \{1, \dots, i\}.
$$

Thus, by the estimates (172) for each j and summing, we obtain,

$$
\sum_{j=2}^{i+1} E_2(V(\mathfrak{X} w), B_j) \leq \frac{1}{2} \sum_{j=1}^i E_2(V(\mathfrak{X} w), B_j) + \frac{2c_4 \lambda^{\frac{p}{2}}}{\sigma^Q} \sum_{j=1}^i \omega(R_j).
$$

Note that

$$
\int_0^{2R} \omega(\varrho) \frac{d\varrho}{\varrho} = \sum_{i=0}^\infty \int_{R_{i+1}}^{R_i} \omega(\varrho) \frac{d\varrho}{\varrho} + \int_R^{2R} \omega(\varrho) \frac{d\varrho}{\varrho}
$$

\n
$$
\geq \sum_{i=0}^\infty \omega(R_{i+1}) \int_{R_{i+1}}^{R_i} \frac{d\varrho}{\varrho} + \omega(R) \int_R^{2R} \frac{d\varrho}{\varrho}
$$

\n(177)
\n
$$
= \log(\frac{1}{\sigma}) \sum_{i=0}^\infty \omega(R_{i+1}) + \omega(R) \log 2 \geq \log 2 \sum_{i=0}^\infty \omega(R_i).
$$

Thus, we deduce, using [\(167\)](#page-53-5), [\(169\)](#page-54-6), [\(177\)](#page-55-0) and [\(170\)](#page-54-7),

(178)
$$
\sum_{j=1}^{i+1} E_2(V(\mathfrak{X}w), B_j) \leq 2E_2(V(\mathfrak{X}w), B_1) + \frac{4c_4\lambda^{\frac{p}{2}}}{\sigma^Q} \sum_{j=1}^i \omega(R_j) \leq \frac{\sigma^Q\lambda^{\frac{p}{2}}}{100}.
$$

Now, using this and writing the difference of averages as a telescoping sum, we deduce,

$$
m_{i+1}(V(\mathfrak{X}w)) - m_1(V(\mathfrak{X}w)) = \sum_{j=1}^i (m_{j+1}(V(\mathfrak{X}w)) - m_j(V(\mathfrak{X}w)))
$$

$$
\leq \sum_{j=1}^i \int_{B_{j+1}} |V(\mathfrak{X}w) - (V(\mathfrak{X}w))_{B_j}|
$$

$$
\leq \sigma^{-Q} \sum_{j=1}^i E_2(V(\mathfrak{X}w), B_j) \leq \frac{\lambda^{\frac{p}{2}}}{100}.
$$

Now [\(178\)](#page-55-1) and [\(179\)](#page-55-2) together yields [\(176\)](#page-54-8).

Step 4: Finally, by [\(176\)](#page-54-8), we have, for $i \geq 1$,

$$
\int_{B_i} |\mathfrak{X}w|^p = \int_{B_i} |V(\mathfrak{X}w)|^2 \leq [m_i(V(\mathfrak{X}w))]^2 + [E_2(V(\mathfrak{X}w), B_i)]^2
$$

$$
\leq [m_i(V(\mathfrak{X}w)) + E_2(V(\mathfrak{X}w), B_i)]^2 \leq \lambda^p.
$$

But this implies, if x_0 is a Lebesgue point of $|\mathfrak{X}w|^p$, then we have,

(180)
$$
|\mathfrak{X}w(x_0)|^p = \lim_{i \to \infty} \int_{B_i} |\mathfrak{X}w|^p \leq \lambda^p.
$$

This immediately implies [\(166\)](#page-53-6) and completes the proof. \Box

Proof of Theorem [6.](#page-6-1) Pick $\Omega_1 \subset\subset \Omega$ and set $\lambda := \|\mathfrak{X}w\|_{L^\infty(\Omega_1)} + 1$. Clearly, $\lambda < \infty$ by Theorem [53.](#page-53-7) We first prove that $V(\mathfrak{X}w)$ is continuous. The strategy of the proof is to show that $V(\mathfrak{X}w)$ is the locally uniform limit of a net of continuous maps, defined by the averages

$$
x \mapsto (V(\mathfrak{X}w))_{B(x,\rho)}.
$$

To do this, we pick any $\Omega' \subset\subset \Omega_1$. We show that for every $\varepsilon > 0$, there exists a radius

$$
0 < r_{\varepsilon} \le \text{dist}(\Omega', \partial \Omega_1) / 1000 = R^*,
$$

depending only on $n, p, \gamma, L, \omega(\cdot), \varepsilon$ such that for every $x \in \Omega'$, the estimate

(181)
$$
\left| (V(\mathfrak{X}w))_{B(x,\rho)} - (V(\mathfrak{X}w))_{B(x,\rho)} \right| \leq \lambda^{\frac{p}{2}} \varepsilon \quad \text{holds for every } \rho, \rho \in (0,r_{\varepsilon}].
$$

This would imply that the sequence of maps $x \mapsto (V(\mathfrak{X}w))_{B(x,\rho)}$ are uniformly Cauchy and would conclude the continuity of $V(\mathfrak{X}w)$.

Step 1: Choice of constants. We fix $\varepsilon > 0$. Now we choose the constants as in the proof of boundedness, but in the the scale ε . More precisely, we choose now, we choose $\sigma \in (0, \frac{1}{4})$ small enough such that

(182)
$$
C_{0,V} \sigma^{\beta} \leq \frac{\varepsilon}{4^{(Q+4)}},
$$

where $C_{0,V}$ and β are constants given by Theorem [2.](#page-3-2) Clearly, σ depend only on ε, n and $p.$ Now, we fix a radius $R_1>0$ small enough such that we have

(183)
$$
\omega(R_1) + \int_0^{2R_1} \omega(\varrho) \frac{d\varrho}{\varrho} \le \frac{\varepsilon \sigma^{2Q}}{6^Q 10^6 c_4}.
$$

Note that R_1 depends on $n, p, \gamma, L, \omega(\cdot)$, and this time, also on ε . Also, if $a(\cdot)$ is a constant function, the dependence on $\omega(\cdot)$ is redundant. With this, we have chosen all the relevant parameters.

Step 2: Smallness of the excess. Fix $R < \min\{R^*, R_1\}$ and consider the chain of shrinking balls as described in [\(163\)](#page-53-2). Proceeding exactly as in Step 2 of the proof of Theorem [53,](#page-53-7) we obtain the following.

For any $i \geq 1$, we have

(184)
$$
E_2(V(\mathfrak{X} w), B_{i+1}) \leq \frac{\varepsilon}{4} E_2(V(\mathfrak{X} w), B_i) + \frac{2c_4\lambda^{\frac{p}{2}}}{\sigma^Q} \omega(R_i).
$$

Now in view of [\(177\)](#page-55-0) and [\(183\)](#page-56-0) we have from [\(184\)](#page-56-1)

$$
E_2(V(\mathfrak{X} w), B_{i+1}) \leq \frac{\varepsilon}{4} E_2(V(\mathfrak{X} w), B_i) + \frac{\varepsilon \sigma^Q \lambda^{\frac{p}{2}}}{3^Q 10^6}.
$$

Now iterating this we obtain,

$$
E_2(V(\mathfrak{X} w), B_{i+1}) \leq \left(\frac{\varepsilon}{4}\right)^i E_2(V(\mathfrak{X} w), B_1) + \frac{\varepsilon \sigma^Q \lambda^{\frac{p}{2}}}{3^Q 10^6} \sum_{j=0}^\infty \left(\frac{\varepsilon}{4}\right)^i.
$$

Now we choose $i_0 \equiv i_0(n, p, \gamma, L, \omega(\cdot), \varepsilon) \geq 2$, such that $\varepsilon^{i_0-1} \leq \sigma^Q$. Then this implies for any $i \geq i_0 + 1$ we have

$$
E_2(V(\mathfrak{X}w), B_i) \leq \varepsilon \sigma^Q \lambda^{\frac{p}{2}}.
$$

This now easily implies that, given $\varepsilon \in (0,1)$, there exists a positive radius $r_{\varepsilon} =$ $r_{\varepsilon}(n, p, \gamma, L, \omega(\cdot), \varepsilon)$ such that we have

(185)
$$
E_2(V(\mathfrak{X}w), B_\rho) \leq \lambda^{\frac{p}{2}} \varepsilon,
$$

whenever $0 < \rho \leq r_{\varepsilon}$ and $B_{\rho} \subset\subset \Omega$.

Step 3: Cauchy estimate.

Now we finish the proof. First we fix a radius one last time. Given $\varepsilon \in (0,1)$, using [\(185\)](#page-57-0) we choose a radius $R_3 \equiv R_3(n, p, \gamma, L, \omega(\cdot), \varepsilon)$ such that

(186)
$$
\sup_{0<\rho\leq R_3}\sup_{x\in\Omega_0}E_2(V(\mathfrak{X}w),B_\rho)\leq \frac{\sigma^{4Q}\lambda^{\frac{p}{2}}\varepsilon}{10^{10}}.
$$

We set

$$
R_0 := \frac{\min\left\{R^*, R_3, R_1\right\}}{16}
$$

and again consider the chain of shrinking balls as described in [\(163\)](#page-53-2) with the starting radius R_0 . Now we want to show that given two integers $2 \leq i_1 < i_2$, we have the estimate p

(187)
$$
\left| (V(\mathfrak{X}w))_{B_{i_1}} - (V(\mathfrak{X}w))_{B_{i_2}} \right| \leq \frac{\lambda^{\frac{\kappa}{2}} \varepsilon}{10}.
$$

Note that this will complete the proof, since for any $0 < \rho < \varrho \leq \sigma^2 R_0$, there exist integers such that

$$
\sigma^{i_1+1}R_0 < \varrho \le \sigma^{i_1}R_0 \quad \text{and } \sigma^{i_2+1}R_0 < \rho \le \sigma^{i_2}R_0.
$$

Also, we have the easy estimates

$$
\begin{aligned} \left| (V(\mathfrak{X}w))_{B_{\varrho}} - (V(\mathfrak{X}w))_{B_{i_1+1}} \right| &\leq \int_{B_{i_1+1}} \left| V(\mathfrak{X}w) - (V(\mathfrak{X}w))_{B_{\varrho}} \right| \\ &\leq \frac{|B_{\varrho}|}{|B_{i_1+1}|} \int_{B_{\varrho}} \left| V(\mathfrak{X}w) - (V(\mathfrak{X}w))_{B_{\varrho}} \right| \\ &\leq \sigma^{-Q} E_2(V(\mathfrak{X}w), B_{\varrho}) \stackrel{(186)}{\leq} \frac{\lambda^{\frac{p}{2}} \varepsilon}{1000} \end{aligned}
$$

and similarly

$$
\left| (V(\mathfrak{X} w))_{B_{\rho}} - (V(\mathfrak{X} w))_{B_{i_2+1}} \right| \leq \frac{\lambda^{\frac{p}{2}} \varepsilon}{1000}.
$$

These two estimates combined with [\(187\)](#page-57-2) will establish [\(181\)](#page-56-2). Thus it only remains to establish [\(187\)](#page-57-2). Summing up [\(184\)](#page-56-1) from $i = i_1 - 1$ to $i_2 - 2$, and using [\(186\)](#page-57-1), [\(183\)](#page-56-0) and [\(177\)](#page-55-0) we obtain

$$
\sum_{i=i_1}^{i_2-1} E_2(V(\mathfrak{X} w), B_i) \leq 2E_2(V(\mathfrak{X} w), B_{i_1-1}) + \frac{4c_4\lambda^{\frac{p}{2}}}{\sigma^Q} \sum_{i=1}^{\infty} \omega(R_i) \leq \frac{\sigma^{2Q} \lambda^{\frac{p}{2}} \varepsilon}{50}.
$$

This yields [\(187\)](#page-57-2) via the elementary estimate

$$
\left| (V(\mathfrak{X} w))_{B_{i_1}} - (V(\mathfrak{X} w))_{B_{i_2}} \right| \leq \sum_{i=i_1}^{i_2-1} \int_{B_{i+1}} |V(dw) - (V(\mathfrak{X} w))_{B_i}|
$$

$$
\leq \sigma^{-Q} \sum_{i=i_1}^{i_2-1} E_2(V(\mathfrak{X} w), B_i).
$$

This establishes [\(181\)](#page-56-2).

Step 4: Final conclusions. Now we prove Theorem [6.](#page-6-1) Note that [\(181\)](#page-56-2) implies $V(\mathfrak{X}w)$ is continuous. [\(14\)](#page-6-3) follows from Theorem [53.](#page-53-7) Next, fix $x \in \Omega$, A, R and λ satisfying [\(15\)](#page-7-0). Now there exists $0 < r_{\varepsilon} < R/16$ which depends only on $n, p, \gamma, L, \omega(\cdot), \varepsilon$, such that for every $\tilde{x} \in B(x, R)$

(188)
$$
\left| (V(\mathfrak{X}w))_{B(\tilde{x},\rho)} - (V(\mathfrak{X}w))_{B(\tilde{x},\rho)} \right| \leq (A\lambda)^{\frac{p}{2}} \varepsilon \quad \text{and}
$$

(189)
$$
E_2(V(\mathfrak{X}w), B_\rho) \le (A\lambda)^{\frac{p}{2}} \varepsilon,
$$

whenever $0 < \rho, \varrho \leq r_{\varepsilon}$. The proof is similar to [\(181\)](#page-56-2) and [\(185\)](#page-57-0). Since $V(\mathfrak{X}w)$ is continuous, by letting $\rho \to 0$ in [\(188\)](#page-58-0), we obtain

$$
\left| V(\mathfrak{X} w) \left(\tilde{x} \right) - \left(V(\mathfrak{X} w) \right)_{B(\tilde{x},\varrho)} \right| \leq \left(A \lambda \right)^{\frac{p}{2}} \varepsilon \quad \text{ holds for every } \varrho \in (0,r_{\varepsilon}].
$$

Choose $\sigma_1 > 0$ sufficiently small such that $\sigma_1 R \leq r_{\varepsilon}/64$. Now we have

$$
B(\tilde{y}, r_{\varepsilon}/8) \subset B(\tilde{x}, r_{\varepsilon}) \subset B(x, R/4), \quad \text{ for any } \tilde{y}, \tilde{x} \in B(x, \sigma_1 R).
$$

We estimate

$$
\left| (V(\mathfrak{X}w))_{B(\tilde{y},r_{\varepsilon}/8)} - (V(\mathfrak{X}w))_{B(\tilde{x},r_{\varepsilon}/8)} \right|
$$

\n
$$
\leq \int_{B(\tilde{y},r_{\varepsilon}/8)} \left| V(\mathfrak{X}w) - (V(\mathfrak{X}w))_{B(\tilde{x},r_{\varepsilon}/8)} \right|
$$

\n
$$
\leq 8^Q \int_{B(\tilde{x},r_{\varepsilon})} \left| V(\mathfrak{X}w) - (V(\mathfrak{X}w))_{B(\tilde{x},r_{\varepsilon}/8)} \right|
$$

\n
$$
\leq 8^Q \int_{B(\tilde{x},r_{\varepsilon})} \left| V(\mathfrak{X}w) - (V(\mathfrak{X}w))_{B(\tilde{x},r_{\varepsilon})} \right|
$$

\n
$$
+ 8^Q \left| (V(\mathfrak{X}w))_{B(\tilde{x},r_{\varepsilon})} - (V(\mathfrak{X}w))_{B(\tilde{x},r_{\varepsilon}/8)} \right|
$$

\n
$$
\leq 2 \cdot 8^{2Q} \int_{B(\tilde{x},r_{\varepsilon})} \left| V(\mathfrak{X}w) - (V(\mathfrak{X}w))_{B(\tilde{x},r_{\varepsilon})} \right|
$$

\n
$$
\leq 16^{2Q} \left(\int_{B(\tilde{x},r_{\varepsilon})} \left| V(\mathfrak{X}w) - (V(\mathfrak{X}w))_{B(\tilde{x},r_{\varepsilon})} \right|^2 \right)^{\frac{1}{2}} \stackrel{(189)}{\leq} 16^{2Q} (A\lambda)^{\frac{p}{2}} \varepsilon.
$$

Thus, by triangle inequality, we have

$$
\begin{aligned} \left| V(\mathfrak{X}w) \left(\tilde{x} \right) - V(\mathfrak{X}w) \left(\tilde{y} \right) \right| \\ &\leq \left| V(\mathfrak{X}w) \left(\tilde{x} \right) - \left(V(\mathfrak{X}w) \right)_{B(\tilde{x}, r_{\varepsilon}/8)} \right| + \left| V(\mathfrak{X}w) \left(\tilde{y} \right) - \left(V(\mathfrak{X}w) \right)_{B(\tilde{y}, r_{\varepsilon}/8)} \right| \\ &\quad + \left| \left(V(\mathfrak{X}w) \right)_{B(\tilde{y}, r_{\varepsilon}/8)} - \left(V(\mathfrak{X}w) \right)_{B(\tilde{x}, r_{\varepsilon}/8)} \right|, \end{aligned}
$$

for any $\tilde{x}, \tilde{y} \in B(x, \sigma_1 R)$. Thus, we deduce

(190)
$$
|V(\mathfrak{X}w)(\tilde{x}) - V(\mathfrak{X}w)(\tilde{y})| \leq 48^{2Q} (A\lambda)^{\frac{p}{2}} \varepsilon.
$$

Now, we first prove [\(16\)](#page-7-1) when $p \geq 2$. For any given $\delta > 0$, clearly we can assume

$$
\max\left\{ \left|\mathfrak{X}w\left(\tilde{x}\right)\right|,\left|\mathfrak{X}w\left(\tilde{y}\right)\right|\right\} \geq \frac{\delta\lambda}{2},
$$

as the estimate follows trivially otherwise. Using the fact that $p \geq 2$, we deduce

$$
\begin{aligned} |\mathfrak{X}w\left(\tilde{x}\right) - \mathfrak{X}w\left(\tilde{y}\right)| &\leq c_V \frac{|V(\mathfrak{X}w)\left(\tilde{x}\right) - V(\mathfrak{X}w)\left(\tilde{y}\right)|}{\left(|\mathfrak{X}w\left(\tilde{x}\right)| + |\mathfrak{X}w\left(\tilde{y}\right)|\right)^{\frac{p-2}{2}}} \\ &\leq c_V 2^{\frac{p}{2}-1} \left(\delta \lambda\right)^{1-\frac{p}{2}} |V(\mathfrak{X}w)\left(\tilde{x}\right) - V(\mathfrak{X}w)\left(\tilde{y}\right)| \\ &\leq c_V 2^{\frac{p}{2}-1} 48^{2Q} A^{\frac{p}{2}} \delta^{1-\frac{p}{2}} \lambda \varepsilon. \end{aligned}
$$

On the other hand, for $1 < p < 2$, we have

$$
\begin{aligned} |\mathfrak{X}w\left(\tilde{x}\right)-\mathfrak{X}w\left(\tilde{y}\right)| &\leq c_V \frac{|V(\mathfrak{X}w)\left(\tilde{x}\right)-V(\mathfrak{X}w)\left(\tilde{y}\right)|}{\left(|\mathfrak{X}w\left(\tilde{x}\right)|+|\mathfrak{X}w\left(\tilde{y}\right)|\right)^{\frac{p-2}{2}}} \\ &\leq c_V \left(2A\lambda\right)^{\frac{2-p}{2}} \left|V(\mathfrak{X}w)\left(\tilde{x}\right)-V(\mathfrak{X}w)\left(\tilde{y}\right)\right| \stackrel{(190)}{\leq} c_V 2^{\frac{p-2}{2}} 48^{2Q} A\lambda \varepsilon. \end{aligned}
$$

Thus, we choose

$$
\varepsilon := \begin{cases} \frac{\delta^{\frac{p}{2}}}{c_V 2^{\frac{p}{2} - 1} 48^{2Q} A^{\frac{p}{2}}} & \text{if } p \ge 2, \\ \frac{\delta}{c_V 2^{\frac{p-2}{2}} 48^{2Q} A} & \text{if } 1 < p < 2. \end{cases}
$$

Then we can determine the radius r_{ε} and consequently σ_1 depending on this choice of ε which depends only on $n, p, \gamma, L, \omega(\cdot), \delta$ as desired. This proves [\(16\)](#page-7-1), which implies the continuity of $\mathfrak{X}w$. This completes the proof.

6. Nonlinear Stein theorem

6.1. **General setting.** Let $u \in HW^{1,p}_{loc}(\Omega)$ be a weak solution to [\(12\)](#page-6-4). Pick $x_0 \in \Omega$ and $0 < R < 1$ such that $B(x_0, 2R) \subset\subset \Omega$. Then clearly, $u \in HW^{1,p}(B_{2R})$ and satisfies

(191)
$$
\operatorname{div}_{\mathbb{H}}(a(x)|\mathfrak{X}u|^{p-2}\mathfrak{X}u) = f \qquad \text{in } B_{2R}.
$$

For $j \geq 0$, we set

(192)
$$
B_j := B(x_0, r_j), \qquad r_j := \sigma^j R, \quad \sigma \in (0, 1/4).
$$

With a slight abuse of notation we denote

$$
cB_j := B(x_0, cr_j), \text{ for any } c > 0.
$$

For $j \geq 0$, we define $w_j \in u + HW_0^{1,p}(B_j)$ to be the unique solution of

(193)
$$
\begin{cases} \operatorname{div}_{\mathbb{H}} \left(a(x) | \mathfrak{X} w_j |^{p-2} \mathfrak{X} w_j \right) = 0 & \text{in } B_j, \\ w_j = u & \text{on } \partial B_j, \end{cases}
$$

and $v_j \in w_j + HW_0^{1,p}(B(x_0, R_j/2))$ to be the unique solution of

(194)
$$
\begin{cases} \operatorname{div}_{\mathbb{H}} \left(a(x_0) |\mathfrak{X} v_j|^{p-2} \mathfrak{X} v_j \right) = 0 & \text{in } B(x_0, R_j/2), \\ v_j = w_j & \text{on } \partial \left(B(x_0, R_j/2) \right). \end{cases}
$$

For $1 < p < \infty$, p' denotes the number $\frac{p}{p-1}$ and let $p_0 > p$ be any real number. Set

(195)
$$
p^* := \begin{cases} \frac{Qp}{Q-p}, & \text{if } p < Q \\ p_0, & \text{if } p \ge Q, \end{cases} \quad \text{and} \quad q := \begin{cases} \frac{Qp}{Q+p} & \text{if } p \ge 2 \\ \frac{Qp}{Qp-Q+p} & \text{if } 1 < p < 2. \end{cases}
$$

6.2. New ingredients. Now we record the new ingredients supplied by our analysis in the present work. The first one is a consequence of Theorem [6.](#page-6-1)

Lemma 55. Let $w_j \in u + HW_0^{1,p}(B_j)$ to be the unique weak solution of [\(193\)](#page-59-2) for $j \geq 0$. Then there exists a constant $C_2 \equiv C_2(n, p, \gamma, L, \omega(\cdot)) \geq 1$ and a positive radius $R_1 = R_1(n, p, \gamma, L, \omega(\cdot)) > 0$ such that if $R \le R_1$, then the following holds:

(196)
$$
\sup_{\frac{1}{2}B_j} |\mathfrak{X} w_j| \leq C_2 \int_{B_j} |\mathfrak{X} w_j|.
$$

Furthermore, for any $A \geq 1$ and any $\delta \in (0,1)$, there exists a positive constant $\sigma_1 \equiv \sigma_1(n, p, \gamma, L, \omega(\cdot), A, \delta) \in (0, \frac{1}{4})$ such that for any $\lambda > 0$,

(197)
$$
\sup_{\frac{1}{2}B_j} |\mathfrak{X} w_j| \leq A\lambda \implies \sup_{x,y \in \sigma_1 B_j} |\mathfrak{X} w_j(x) - \mathfrak{X} w_j(y)| \leq \delta \lambda.
$$

The following follows from Theorems [30](#page-15-7) and [31](#page-15-2) and does not need our results.

Lemma 56. Let $v_j \in w_j + HW_0^{1,p} (B(x_0, R_j/2))$ to be the unique weak solution of [\(194\)](#page-60-0) for $j \geq 0$. Then there exists a constant $C_3 \equiv C_3(n,p) \geq 1$ such that

(198)
$$
\sup_{B(x_0,R_j/4)} |\mathfrak{X} v_j| \leq C_3 \int_{B(x_0,R_j/2)} |\mathfrak{X} v_j|.
$$

For every $A \geq 1$ there exists a constants $C_4 \equiv C_4(n, p, A) \equiv \tilde{C}_4(n, p)A$ and $\alpha \equiv$ $\alpha(n, p) \in (0, 1)$, such that for any $\lambda > 0$ and for any $\tau \in (0, 1/4)$, we have

(199)
$$
\sup_{B(x_0,R_j/4)} |\mathfrak{X} v_j| \leq A\lambda \implies \sup_{x,y \in B(x_0,\tau R_j)} |\mathfrak{X} v_j(x) - \mathfrak{X} v_j(y)| \leq C_4 \tau^{\alpha} \lambda.
$$

The second new ingredient is the following, which is just Theorem [1.](#page-3-0)

Lemma 57. Let v_j be as in [\(194\)](#page-60-0) for $j \ge 0$. Given $\bar{\varepsilon} > 0$, there exists $\sigma_2 \equiv$ $\sigma_2(n, p, \gamma, L, \bar{\varepsilon}) \in (0, 1/4)$, such that if $\sigma \in (0, \sigma_2]$ then for all $t \in [0, 1]$, we have

$$
(200) \qquad \left(\int_{B_{\sigma R_j}} \left| \mathfrak{X} v_j - (\mathfrak{X} v_j)_{\sigma R_j} \right|^t \right)^{\frac{1}{t}} \leq \bar{\varepsilon} \left(\int_{B_{\frac{1}{2}R_j}} \left| \mathfrak{X} v_j - (\mathfrak{X} v_j)_{\frac{1}{2}R_j} \right|^t \right)^{\frac{1}{t}}.
$$

6.3. Comparison estimates. We begin by recording a number of comparison estimates. The following two can be proved analogously to Lemma 5 in [\[42\]](#page-64-18) and Lemma 3 in [\[42\]](#page-64-18) and is valid for $1 < p < \infty$.

Lemma 58. Let u be as in [\(191\)](#page-59-3) and w_j be as in [\(193\)](#page-59-2) and $j \ge 0$. There exists a constant $c_5 \equiv c_5 (n, p, \gamma, L)$ such that for any $p > 1$, the following inequality holds.

(201)
$$
\int_{B_j} (|\mathfrak{X} u| + |\mathfrak{X} w_j|)^{p-2} |\mathfrak{X} u - \mathfrak{X} w_j|^2 \leq c_5 \int_{B_j} |f| |u - w_j|.
$$

Lemma 59. Let w_j, v_j be as before and $j \geq 0$. Then there exists a constant $c \equiv$ $c(n, p, \gamma, L)$ such that we have the inequality

(202)
$$
\int_{\frac{1}{2}B_j}|V(\mathfrak{X}v_j)-V(\mathfrak{X}w_j)|^2 \leq c\left[\omega(r_j)\right]^2 \int_{\frac{1}{2}B_j}|\mathfrak{X}w_j|^p.
$$

The following three lemmas are proved analogously to Lemma 7, Lemma 8 and Lemma 9 of [\[42\]](#page-64-18), respectively, using Lemma [55,](#page-60-1) Lemma [56](#page-60-2) and Lemma [57](#page-60-3) in place of Theorem 9, Theorem 8 and Lemma 2, respectively, of [\[42\]](#page-64-18). These provide the comparison estimates for the case $p > 2$.

Lemma 60. Let w_j be as in [\(193\)](#page-59-2), v_j be as in [\(194\)](#page-60-0) with $p > 2$ and $j \ge 0$. Then there exists a constant $C_7 \equiv C_7(n, p, \gamma, L)$ such that we have the inequality

(203)
$$
\left(\int_{\frac{1}{2}B_j}|\mathfrak{X}v_j-\mathfrak{X}w_j|^p\right)^{\frac{1}{p}}\leq C_7\left[\omega(r_j)\right]^{\frac{2}{p}}\left(\int_{\frac{1}{2}B_j}|\mathfrak{X}w_j|^p\right)^{\frac{1}{p}}.
$$

Lemma 61. Let u be as in [\(191\)](#page-59-3) with $p > 2$ and w_j be as in [\(193\)](#page-59-2) and $j \ge 0$. There exists a constant $C_5 \equiv C_5(n, p, \tilde{q}, \gamma, L)$ such that the following inequality

,

(204)
$$
\left(\int_{B_j} |\mathfrak{X} u - \mathfrak{X} w_j|^p \right)^{\frac{1}{p}} \leq C_5 \left(r_j^q \int_{B_j} |f|^q \right)^{\frac{1}{q(p-1)}}
$$

holds for every $\tilde{q} \geq (p^*)'$ when $p < Q$ and for every $\tilde{q} > 1$ when $p \geq Q$. Moreover, when $j \geq 1$, there exists another constant $C_6 \equiv C_6(n, p, \tilde{q}, \gamma, L, \sigma) > 0$ such that the following inequality holds

(205)
$$
\left(\oint_{B_j} |\mathfrak{X} w_{j-1} - \mathfrak{X} w_j|^p\right)^{\frac{1}{p}} \leq C_6 \left(r_{j-1}^q \oint_{B_{j-1}} |f|^{\tilde{q}}\right)^{\frac{1}{q(p-1)}}.
$$

Lemma 62. Let u be as in [\(191\)](#page-59-3) with $p > 2$ and v_j, w_j be as before in [\(193\)](#page-59-2), [\(194\)](#page-60-0), respectively, for $j \ge 1$ and let q be as in [\(195\)](#page-60-4). Suppose for some $\lambda > 0$ we have

(206)
$$
r_{j-1} \left(\int_{B_{j-1}} |f|^q \right)^{\frac{1}{q}} \leq \lambda^{p-1}
$$

and for some constants $A, B \geq 1$, the estimates

(207)
$$
\sup_{\frac{1}{2}B_j} |\mathfrak{X} w_j| \le A\lambda \quad and \quad \frac{\lambda}{B} \le |\mathfrak{X} w_{j-1}| \le A\lambda \quad in \ B_j
$$

hold. Then there exists a constant $C_8 \equiv C_8(n, p, \gamma, L, A, B, \sigma) > 0$ such that

(208)
$$
\left(\int_{\frac{1}{2}B_j} |\mathfrak{X} u - \mathfrak{X} v_j|^{p'}\right)^{\frac{1}{p'}} \leq C_8 \omega(r_j) \lambda + C_8 \lambda^{2-p} r_{j-1} \left(\int_{B_{j-1}} |f|^q\right)^{\frac{1}{q}}.
$$

The following Lemma is proved analogously to Lemma 10 in [\[42\]](#page-64-18), using Lemma [56](#page-60-2) and Lemma [57](#page-60-3) in place of Theorem 8 and Lemma 2, respectively, of [\[42\]](#page-64-18). This provides the comparison estimate for the case $1 < p \leq 2$.

Lemma 63. Let u be as in [\(191\)](#page-59-3) with $1 < p \le 2$ and v_i, w_j be as before in [\(193\)](#page-59-2), [\(194\)](#page-60-0), respectively, for $j \ge 1$ and let q be as in [\(195\)](#page-60-4). Suppose for some $\lambda > 0$, we have

$$
r_j\left(\mathop{\hskip0pt{1}}\nolimits_{B_j}|f|^q\right)^{\frac{1}{q}} \leq \lambda^{p-1} \qquad \text{and} \qquad \left(\mathop{\hskip0pt{1}}\nolimits_{B_j}|\mathfrak{X} u|^p\right)^{\frac{1}{p}} \leq \lambda.
$$

Then there exists a constant $C_8 \equiv C_8(n, p, \gamma, L) > 0$ such that

$$
(209) \qquad \left(\oint_{\frac{1}{2}B_j} |\mathfrak{X}u - \mathfrak{X}v_j|^p\right)^{\frac{1}{p}} \leq C_8 \omega(r_j) \lambda + C_8 \lambda^{2-p} r_j \left(\oint_{B_j} |f|^q\right)^{\frac{1}{q}}.
$$

Moreover, there exists another constant $C_5 = C_5(n, p, \nu, L) > 0$ such that

(210)
$$
\left(\int_{B_j} |\mathfrak{X} u - \mathfrak{X} w_j|^p \right)^{\frac{1}{p}} \leq C_5 \lambda^{2-p} r_j \left(\int_{B_j} |f|^q \right)^{\frac{1}{q}}.
$$

6.4. Proof of Theorem [5.](#page-6-0)

Theorem 64. Let u be as in Theorem [5,](#page-6-0) solves the equation (12) . Then $\mathfrak{X}u$ is locally bounded in Ω . Moreover, there exists a constant $c \geq 1$ and a positive radius R_0 , both depending only on Q, p, γ, L and $\omega(\cdot)$ such that the pointwise estimate

(211)
$$
|\mathfrak{X}u(x_0)| \leq c \left(\int_{B(x_0,r)} |\mathfrak{X}u|^s \right)^{\frac{1}{s}} + c||f||_{L^{(Q,1)}(B(x_0,r))}^{\frac{1}{p-1}}
$$

holds whenever $B(x_0, 2r) \subset \Omega$, x_0 is a Lebesgue point of $\mathfrak{X}u$ and $2r \leq R_0$. Estimate [\(211\)](#page-62-5) holds with no restriction on r when the coefficient function $a(\cdot)$ is constant.

Proof. The proof of the theorem now proceeds analogously to the proof of Theorem 4 in [\[31\]](#page-63-16), where D is replaced by \mathfrak{X}, n is replaced by Q and the constants are replaced by analogous constants. We use Lemma [55,](#page-60-1) Lemma [56,](#page-60-2) Lemma [57,](#page-60-3) Lemma [61,](#page-61-0) Lemma [60,](#page-61-1) Lemma [62,](#page-61-2) Lemma [63](#page-62-6) and Lemma [19](#page-12-3) as substitutes for Theorem 2, Theorem 3, Lemma 3, Lemma 4, Lemma 5, Lemma 6, Lemma 7 and Lemma 1, respectively, of [\[31\]](#page-63-16). We remark that our excess decay estimate is crucial for proving these replacements.

Proof of Theorem [5.](#page-6-0) Now the proof of Theorem [5](#page-6-0) follows exactly analogously to the proof of Theorem 1 in [\[31\]](#page-63-16), with the same replacements as above. \Box

REFERENCES

- [1] BONFIGLIOLI, A., LANCONELLI, E., AND UGUZZONI, F. Stratified Lie groups and potential theory for their sub-Laplacians. Springer Monographs in Mathematics. Springer, Berlin, 2007.
- [2] Capogna, L. Regularity of quasi-linear equations in the Heisenberg group. Comm. Pure Appl. Math. 50, 9 (1997), 867–889.
- [3] CAPOGNA, L., CITTI, G., AND ZHONG, X. Regularity theory of quasilinear elliptic and parabolic equations in the Heisenberg group. Vietnam. J. Math 52, 4 (2024), 807–827.
- [4] CAPOGNA, L., DANIELLI, D., PAULS, S. D., AND TYSON, J. T. An introduction to the Heisenberg group and the sub-Riemannian isoperimetric problem, vol. 259 of Progress in Mathematics. Birkhäuser Verlag, Basel, 2007.
- [5] CIANCHI, A. Maximizing the L^{∞} norm of the gradient of solutions to the Poisson equation. J. Geom. Anal. 2, 6 (1992), 499–515.

64 A. MALLICK AND S. SIL

- [6] DIBENEDETTO, E., AND FRIEDMAN, A. Regularity of solutions of nonlinear degenerate parabolic systems. J. Reine Angew. Math. 349 (1984), 83–128.
- [7] DIBENEDETTO, E., AND FRIEDMAN, A. Hölder estimates for nonlinear degenerate parabolic systems. J. Reine Angew. Math. 357 (1985), 1–22.
- DIBENEDETTO, E., AND MANFREDI, J. On the higher integrability of the gradient of weak solutions of certain degenerate elliptic systems. Amer. J. Math. 115, 5 (1993), 1107–1134.
- [9] DOMOKOS, A. Differentiability of solutions for the non-degenerate p-Laplacian in the Heisenberg group. J. Differential Equations 204, 2 (2004), 439–470.
- [10] Duzaar, F., and Mingione, G. Gradient estimates via linear and nonlinear potentials. J. Funct. Anal 259, 11 (2010), 2961–2998.
- [11] Duzaar, F., and Mingione, G. Gradient continuity estimates. Calc. Var. Partial Differential Equations 39, 3-4 (2010), 379–418.
- [12] Duzaar, F., and Mingione, G. Gradient estimates via non-linear potentials. Amer. J. Math 133, 4 (2011), 1093–1149.
- [13] EVANS, L. C. A new proof of local $C^{1,\alpha}$ regularity for solutions of certain degenerate elliptic p.d.e. J. Differential Equations 45, 3 (1982), 356–373.
- [14] FOLLAND, G. B. Subelliptic estimates and function spaces on nilpotent Lie groups. Ark. Mat. 13, 2 (1975), 161–207.
- [15] Folland, G. B. Applications of analysis on nilpotent groups to partial differential equations. Bull. Amer. Math. Soc. 83, 5 (1977), 912–930.
- [16] FOLLAND, G. B., AND KOHN, J. J. The Neumann problem for the Cauchy-Riemann complex, vol. No. 75 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1972.
- [17] FOLLAND, G. B., AND STEIN, E. M. Estimates for the $\bar{\partial}_b$ complex and analysis on the Heisenberg group. Comm. Pure Appl. Math. 27 (1974), 429–522.
- [18] FOLLAND, G. B., AND STEIN, E. M. Hardy spaces on homogeneous groups, vol. 28 of Mathematical Notes. Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1982.
- [19] Garofalo, N., and Nhieu, Duy-M. Lipschitz continuity, global smooth approximations and extension theorems for Sobolev functions in Carnot-Carathéodory spaces. Journal d'Analyse Mathématique 74, (1998), 67-97.
- [20] Giaquinta, M., and Martinazzi, L. An introduction to the regularity theory for elliptic systems, harmonic maps and minimal graphs, second ed., vol. 11 of Appunti. Scuola Normale Superiore di Pisa (Nuova Serie) [Lecture Notes. Scuola Normale Superiore di Pisa (New Series)]. Edizioni della Normale, Pisa, 2012.
- [21] Giaquinta, M., and Modica, G. Remarks on the regularity of the minimizers of certain degenerate functionals. Manuscripta Math. 57, 1 (1986), 55–99.
- [22] HAMBURGER, C. Regularity of differential forms minimizing degenerate elliptic functionals. J. Reine Angew. Math. 431 (1992), 7–64.
- [23] HÖRMANDER, L. Hypoelliptic second order differential equations. Acta Math. 119 (1967), 147–171.
- [24] HUNT, R. A. On $L(p, q)$ spaces. *Enseign. Math.* (2) 12 (1966), 249–276.
- [25] JERISON, D. The Poincaré inequality for vector fields satisfying Hörmander's condition. Duke Math. J. 53, 2 (1986), 503–523.
- [26] Jin, T., Maz'ya, V., and Van Schaftingen, J. Pathological solutions to elliptic problems in divergence form with continuous coefficients. C. R. Math. Acad. Sci. Paris 347, 13-14 (2009), 773–778.
- [27] KOHN, J. J., AND NIRENBERG, L. Non-coercive boundary value problems. Comm. Pure Appl. Math. 18 (1965), 443–492.
- [28] KORÁNYI, A., AND VÁGI, S. Singular integrals on homogeneous spaces and some problems of classical analysis. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3) 25 (1971), 575–648 (1972).
- [29] Kuusi, T., and Mingione, G. New perturbation methods for nonlinear parabolic problems. J. Math. Pures Appl. (9) 98, 4 (2012), 390–427.
- [30] Kuusi, T., and Mingione, G. Linear potentials in nonlinear potential theory. Arch. Ration. Mech. Anal. 207, 1 (2013), 215–246.
- [31] Kuusi, T., and Mingione, G. A nonlinear Stein theorem. Calc. Var. Partial Differential Equations 51, 1-2 (2014), 45–86.
- [32] LADYZHENSKAYA, O. A., AND URAL'TSEVA, N. N. Linear and quasilinear elliptic equations. Academic Press, New York-London, 1968. Translated from the Russian by Scripta Technica, Inc, Translation editor: Leon Ehrenpreis.
- [33] LIEBERMAN, G. M. The natural generalization of the natural conditions of Ladyzhenskaya and Uraltseva for elliptic equations. Comm. Partial Differential Equations 16, 2-3 (1991), 311–361.
- [34] Lu, G. The sharp Poincaré inequality for free vector fields: an endpoint result. Revista Matemática Iberoamericana 10, 2 (1994), 453-466.
- [35] MANFREDI, J. J., AND MINGIONE, G. Regularity results for quasilinear elliptic equations in the Heisenberg group. Math. Ann. 339, 3 (2007), 485–544.
- [36] MARCHI, S. $C^{1,\alpha}$ local regularity for the solutions of the p-Laplacian on the Heisenberg group for $2 \le p \le 1 + \sqrt{5}$. Z. Anal. Anwendungen 20, 3 (2001), 617–636. Erratum: Z. Anal. Anwendungen 22, (2003) 471–472.
- [37] MINGIONE, G., ZATORSKA-GOLDSTEIN, A., AND ZHONG, X. Gradient regularity for elliptic equations in the Heisenberg group. Adv. Math. 222, 1 (2009), 62–129.
- [38] MUKHERJEE, S., AND SIRE, Y. Regularity of inhomogeneous quasi-linear equations on the Heisenberg group. Anal. Theory Appl. 37, 4 (2021), 520–540.
- [39] MUKHERJEE, S., AND ZHONG, X. $C^{1,\alpha}$ -regularity for variational problems in the Heisenberg group. Anal. PDE 14, 2 (2021), 567–594.
- [40] RICCIOTTI, D. p-Laplace equation in the Heisenberg group. SpringerBriefs in Mathematics. Springer, [Cham]; BCAM Basque Center for Applied Mathematics, Bilbao, 2015. Regularity of solutions, BCAM SpringerBriefs.
- [41] Roncal, L., and Thangavelu, S. Hardy's inequality for fractional powers of the sublaplacian on the Heisenberg group. Adv. Math. 302, (2016), 106–158.
- [42] SiL, S. Nonlinear Stein theorem for differential forms. Calc. Var. Partial Differential Equations 58, 4 (2019), Paper No. 154.
- [43] STEIN, E. M. Editor's note: the differentiability of functions in \mathbb{R}^n . Ann. of Math. (2) 113, 2 (1981), 383–385.
- [44] STEIN, E. M., AND WEISS, G. Introduction to Fourier analysis on Euclidean spaces. Princeton University Press, Princeton, N.J., 1971. Princeton Mathematical Series, No. 32.
- [45] TOLKSDORF, P. Regularity for a more general class of quasilinear elliptic equations. J. Differential Equations 51, 1 (1984), 126–150.
- [46] Uhlenbeck, K. Regularity for a class of non-linear elliptic systems. Acta Math. 138, 3-4 (1977), 219–240.
- [47] URAL'CEVA, N. N. Degenerate quasilinear elliptic systems. Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 7 (1968), 184–222.
- [48] Xu, C.-J., AND ZUILY, C. Higher interior regularity for quasilinear subelliptic systems. Calc. Var. Partial Differential Equations 5, 4 (1997), 323–343.
- [49] Yu, C. Regularity for quasi-linear p-laplacian type non-homogeneous equations in the heisenberg group. Mathematics 10, 21 (2022).
- [50] Zhong, X. Regularity for variational problems in the Heisenberg group. arXiv e-prints (2018), [arXiv:1711.03284.](http://arxiv.org/abs/1711.03284)

Department of Mathematics, Indian Institute of Science, Bangalore, India. Email address: arkamallick@iisc.ac.in

Department of Mathematics, Indian Institute of Science, Bangalore, India. Email address: swarnendusil@iisc.ac.in