

Absorbing State Phase Transitions and Stability of Long-Range Coherence in Dissipative Quantum State Preparation

Matthew Wampler and Nigel R. Cooper

T.C.M. Group, Cavendish Laboratory, University of Cambridge, J.J. Thomson Avenue, Cambridge CB3 0HE, United Kingdom

We investigate the absorbing state phase transition in a purely dissipative quantum reaction-diffusion model where the absorbing state itself has long range phase coherence. The model may also be viewed as a dissipative quantum state preparation procedure for the (generalized) W-state with errors. The ‘error’ Lindblad jump operators preserve the W-state as a dark state, but nonetheless act to decohere the system and induce the phase transition. We find cases where the preparation protocol is either fragile or robust against weak error quantum jump rates, and show that remnants of the coherence persist even after the phase transition to the decohering phase.

Introduction—The dynamics of non-equilibrium quantum many-body systems exhibit striking phenomena which would be rare, if not impossible, in classical or equilibrium settings. Interest in this area has been especially spurred by recent experimental advances – across a variety of platforms – in the control of quantum systems at the atom-by-atom (or qubit-by-qubit) level [1–5].

Often, evolution in these systems has been viewed as a competition between the coherent Hamiltonian dynamics which may contribute to the quantum nature of a system, and the decohering effects of coupling to the environment or measurements which destroy entanglement and other quantum signatures. However, even in systems with no Hamiltonian dynamics, it is possible to generate entanglement solely through dissipation [6, 7]. Additionally, the irreversible nature of dissipation allows for the one-way cooling of quantum systems to a desired steady state as well as provides the possibility for interactive control of a system through feedback. These properties – which allow dissipative dynamics to be especially robust to perturbation – make it a key tool in quantum information [8], showing up in quantum state preparation and stabilization [9–17], error correction [18–22], and as distinct models of universal quantum computation [6, 23].

In condensed matter, the same experimental advancements in engineered coupling to the environment and quantum control have led to the discovery of new phases of matter and dynamical phase transitions which are unique to the non-equilibrium setting [24–29]. Non-equilibrium phase transitions are present in a wide variety of contexts [30–32]. An important class of systems which exhibit such transitions are reaction-diffusion models, where particles (or reactants) move through a system by diffusion and react when they meet. Here, the system may exhibit phase transitions to absorbing states (i.e. configurations which may be approached by the dynamics, but not left) [33]. There has been a flurry of recent interest in quantum extensions to reaction-diffusion models [34–40] where it has been found that quantum effects may significantly alter the properties of the phase transition and system dynamics.

In this paper, we provide a prototypical example of a quantum reaction-diffusion model where the absorbing state itself has long range phase coherence. We find, furthermore, this coherence may be induced entirely through dissipative means, with no Hamiltonian evolution. The model may simultaneously be viewed as a dissipative quantum state preparation procedure – preparing the long range entangled absorbing state – with additional ‘error’ environmental couplings that ruin the preparation protocol. The absorbing state phase transition, from this perspective, may be viewed as a probe into the stability of purely dissipative quantum state preparation. Interestingly, we also find that even when the error terms are strong enough to induce a phase transition, some quantum coherence still persists in the steady state.

The model consists of local Lindbladian dynamics that includes the generalized W-state as a dark state. These are a class of states given by

$$|W_{\text{Gen}}\rangle = \frac{1}{\sqrt{N}} \sum_{a=0}^{N-1} e^{i\phi_a} |\downarrow_0 \dots \downarrow_{a-1} \uparrow_a \downarrow_{a+1} \dots \downarrow_{N-1}\rangle \quad (1)$$

where a labels the sites of the lattice, and various choices of phase ϕ_a give, for example, the W-state (all $\phi_a = 0$) or vortex lattices (phase winding around closed loops). While the (generalized) W-state is a dark state of the Lindbladian (to be defined below), some terms are “attractive” in the sense that they drive the system towards the W-state (the W-state preparation protocol) while other terms are “repulsive” and drive states in the neighborhood of the W-state towards a different state (in this case, the maximally mixed state). Competition between these terms induces a phase transition away from the absorbing W-state.

The Model—The model we consider in this work is a lattice of two level systems where pairs of lattice sites are coupled (via a set of environments) to undergo dissipative evolution. We restrict to the case well-described by Lindbladian dynamics, given by the master equation for the density operator ρ

$$\dot{\rho} = \sum_{\ell} \gamma_{\ell} \left(L_{\ell} \rho L_{\ell}^{\dagger} - \frac{1}{2} \{ L_{\ell}^{\dagger} L_{\ell}, \rho \} \right). \quad (2)$$




FIG. 1. Our model and dissipative W-state preparation. a) Quantum Jump operators for the model written in terms of total spin eigenstates $|s, m_s\rangle$ for 2-site pairs. The γ_1, γ_2 terms prepare the W-state, while γ_3 terms induce a phase transition to a mixed state. b) Points represent states and arrows are quantum jumps in the Lindblad dynamics. c) Preparation of W-state using ladder between symmetry sectors.

The model is purely dissipative (with real $\gamma_\ell > 0$), which is why no coherent, Hamiltonian evolution appears in (2).

In particular, we consider 2-site jump operators grouped into the following four sets:

$$L_1 = \{|1, 0\rangle_{a,b}\langle 0, 0|_{a,b}\}, \quad (3a)$$

$$L_2 = \{|1, 0\rangle_{a,b}\langle 1, 1|_{a,b}\}, \quad (3b)$$

$$L_3 = \{|0, 0\rangle_{a,b}\langle 1, 1|_{a,b}\}, \quad (3c)$$

$$L_{3'} = \{|1, 1\rangle_{a,b}\langle 0, 0|_{a,b}\}. \quad (3d)$$

We will consider two cases: all-to-all jump operators— with any pair of sites a, b — as well as nearest-neighbor (NN) where a, b are on adjacent lattice sites (with coordination number Z). We have written 2-site states in terms of total spin quantum numbers $|s, m_s\rangle$.

In the Lindbladian (2), terms L_1, L_2, L_3 , and $L_{3'}$ have coefficients $\gamma_1, \gamma_2, \gamma_3$ and $\gamma_{3'}$ respectively. Throughout the work, we will take $\gamma_3 = \gamma_{3'}$. The model is summarized in Fig. 1a.

It may be seen that this Lindbladian represents a quantum reaction-diffusion model by viewing the spin-up excitations as hard-core bosonic particles. From this viewpoint, L_1 conserves particle number (total S_z) and— since the dissipative evolution favors states where a particle is in a coherent superposition on neighboring sites— will tend to spread the particle throughout the system, i.e. it is the quantum analog of the ‘diffusion’ term. On the other hand, the rest of the jump operators correspond to prototypical reaction processes. Namely, ‘coagulation’ terms L_2 and L_3 correspond to processes where two particles collide (i.e. are on adjacent sites) and decay down to a single particle state, while ‘branching’ terms $L_{3'}$ correspond to Lindblad jumps from a single

particle state to a two particle state. Two-site dissipative dynamics similar to that generated by the quantum jumps above have been realized across a variety of experimental platforms, for example [41–43].

From the quantum state preparation perspective, the model exhibits competition between quantum jumps where the W-state is either an ‘attractive’ (L_1, L_2) or ‘repulsive’ ($L_3, L_{3'}$) dark state. For attractive dark states of a jump L , we have $L^\dagger|W\rangle \neq 0$. This implies there is some attractive basin of states which flow to the W-state under L . For repulsive dark states, $L|W\rangle = L^\dagger|W\rangle = 0$. Here, states in the neighborhood of the W-state will never jump to the W-state, but will instead flow to some other steady state.

Dissipative W-State Preparation—We now show that Lindbladian dynamics with L_1 and L_2 prepares the (generalized) W-state. In full generality, this is not possible as the generalized W-state falls outside the class of states which may be prepared by local, purely dissipative Lindbladians [11]. However, several strategies have been explored to extend the set of preparable states [12, 44–47] allowing for experimental W-state realizations [48–50].

The strategy we use in this work is to restrict to a subspace of allowed initial states. Jump operators which commute with a symmetry of the target state may be used for dissipative preparation within the subspace of states that share that symmetry (see Fig. 1). Further jump operators may be added which take other states into the target symmetry sector, thereby increasing the space of initial states which prepare the target state.

We first describe how the W-state may be prepared using this strategy, and then will comment on how the protocol is altered for the generalized W-state. The W-state is a superposition of states with a single spin-up excitation, therefore let us focus on the subspace of states with conserved total S_z . Within the 1-spin-up ($S_z = -\frac{N}{2} + 1$) sector, the W-state is the only state with maximal total spin. Therefore, if local jump operators which preserve S_z but increase total spin are applied throughout the system, then the 1-spin-up subspace will approach the W-state. The jump operators L_1 (3a) achieve precisely this. These jump operators each locally lock the phase between the two sites such that the collective effect of all the jump operators is to obtain global phase coherence. Protocols of this type were previously considered in [10, 51] where a similar set of jump operators was used in the context of lattice ultracold bosonic atoms.

The operators L_2 may now be added to connect other symmetry sectors to the target $S_z = -\frac{N}{2} + 1$ sector by lowering S_z . The full protocol is then summarized in Fig. 1c. The S_z conserving terms L_1 push each S_z -sector towards the corresponding globally phase coherent state. The L_2 terms then lower S_z in the system until there are fewer than two spin-ups. In this way, any initial state will converge to the W-state except for one— the vacuum (i.e. completely spin-down state). Like the W-state, the

vacuum $|\Downarrow \dots \Downarrow\rangle$ is a dark state of L_1 and L_2 . Given [11], this is as close to a full dissipative preparation of the W-state as one can get. Namely, there is only one state which the dissipative attractor dynamics does not take to the W-state, which means a random initial state will converge to the W-state with probability 1 in the thermodynamic limit.

To prepare the generalized W-state, note that the generalized W-state is simply a local gauge transformation of the W-state, i.e.

$$|W_{\text{Gen}}\rangle = U_{\text{gauge}}|W\rangle = \prod_a e^{i\phi_a \sigma_a^+ \sigma_a^-} |W\rangle \quad (4)$$

with σ_a^\pm the spin raising/lowering operators and ϕ_a desired phase to imprint on site a . A preparation strategy may then be constructed by applying this same gauge transformation to the jump operators, $L_\ell \rightarrow U_{\text{gauge}} L_\ell U_{\text{gauge}}^\dagger$, such that the dynamics under (2) is preserved except to the new absorbing state $|W_{\text{Gen}}\rangle$.

Absorbing State Phase Transition—We now investigate what happens when $\gamma_3 \neq 0$. Initially, we will focus on the case where the jump operators not only act on nearest neighbors (NN), but also act between every 2-site pair in the system. This choice is motivated by two main reasons.

First, the W-state is all-to-all symmetric and so leveraging an all-to-all preparation procedure might be expected to improve upon preparation time. We indeed find (see below) that this is the case: the all-to-all Lindbladian for W-state preparation is gapped (implying constant preparation time) while the NN case has a gap which closes polynomially in system size. We will show, however, that this improvement for all-to-all preparation comes at a cost; the all-to-all procedure is fragile to $O(\frac{1}{N})$ error quantum jump rates (whereas the transition occurs for error jump rates $O(1)$ in the NN case).

Second, any all-to-all Lindbladian (or general open quantum dynamics) may be represented by a set of generalized Dicke Operators, which act as closed operations within the space of generalized Dicke states (see [52] for details). Specifically, any permutation symmetric density matrix may be represented as a linear combination of generalized Dicke states

$$\rho = \sum_{q, q_z, \sigma_z} \alpha_{q, q_z, \sigma_z} \mathcal{D}_{q, q_z, \sigma_z} \quad (5)$$

where the generalized Dicke states $\mathcal{D}_{q, q_z, \sigma_z}$ are defined in terms of three quantum numbers q , q_z , and σ_z . The number of $\mathcal{D}_{q, q_z, \sigma_z}$ is only $O(N^3)$, and therefore scales much more favorably than the exponential size of the full density matrix. Generalized Dicke states extend to open systems the more commonly used Dicke states, i.e. the set of permutation symmetric pure states whose utility was first introduced in the context of quantum optics [53]. We will focus here on the main results from the

generalized Dicke analysis; calculation details may be found in the Appendix.

Using generalized Dicke operators, it is possible to see explicitly that the only two dark states of the system with $\gamma_3 = 0$ are the vacuum and the W-state. Furthermore, any initial state that has zero overlap with the vacuum will converge to the W-state. The operators L_3 , $L_{3'}$ on the other hand have 3 important steady states: the vacuum, the W-state, and the maximally mixed state. Importantly, as remarked in our introduction of the model, both the vacuum and W-state are repulsive dark states. Initial states with zero overlap in the vacuum or W-state will flow to the maximally mixed state.

To numerically study the phase transition induced by these competing jump operators, we examine the gap Δ between the first 2 eigenvalues (corresponding to the vacuum and W-state) and the next eigenvalue closest to 0. The gap closes as γ_3 is increased (and $\gamma_1 = \gamma_2 = 1$). We plot the gap in Fig. 2a as a function of $N\gamma_3$. The gap closing for such small values of γ_3 - scaling with $\frac{1}{N}$ in the system size - implies that the W-state preparation strategy is quite fragile to the addition of L_3 , $L_{3'}$.

A priori, this fragility may be surprising as, for example, L_1 and $L_{3'}$ jump from the same state (the singlet state) and act on the same number of site-pairs. Similarly for L_2 , L_3 , and the $|1, 1\rangle$ state. There is, therefore, no $O(N)$ difference in the rate at which cohering vs. decohering quantum jumps occur. This would seem to suggest a transition should occur when $\frac{\gamma_3}{\gamma_1} = O(1)$ not $O(\frac{1}{N})$.

To see why this is not the case, it is helpful to look at the NN model and consider evolution of the expectation value $\langle P^{1,0}, |1, -1\rangle \rangle$ where $P^{1,0}, |1, -1\rangle = \frac{1}{N-1} \sum_{\langle a, b \rangle} (|1, 0\rangle_{a, b} \langle 1, 0|_{a, b} + |1, -1\rangle_{a, b} \langle 1, -1|_{a, b})$. Physically, $\langle P^{1,0}, |1, -1\rangle \rangle$ represents the expectation value that neighboring sites will be in either the $|1, 0\rangle$ or $|1, -1\rangle$ state. This is maximal for the W-state and serves as an order parameter for the transition. As an example, consider evolution of this expectation value for any ρ in the $S_z = -\frac{N}{2} + 1$ sector (see Appendix F for derivation with any ρ) given by

$$\frac{d}{dt} \langle P^{1,0}, |1, -1\rangle \rangle = \left(\gamma_1 - \gamma_3 \frac{Z-1}{2} \right) \langle P^{0,0} \rangle \quad (6)$$

where $\langle P^{0,0} \rangle$ is the expectation value that neighboring sites are in the $|0, 0\rangle$ state. Intuitively, the extra factor of Z occurs for the decohering terms because—for any pair of sites a, b —the jump operators can decohere a, b by acting on a and any neighbor $c \neq b$ (or similarly, sites b and a neighbor $c \neq a$). However, to make a, b coherent, L_1 must act directly on sites a, b . In the all-to-all limit, $Z \rightarrow N$ and the decohering terms dominate by a factor of N , implying any non-zero γ_3 will ruin the W-state preparation procedure.

We now investigate the properties of the steady state in the regime where the gap closes. Here, the




FIG. 2. Absorbing state phase transition in the all-to-all system. a) Gap to first non-zero eigenvalue. Here, $\gamma_1 = \gamma_2 = 1$ and numerics are done within the $\sigma_z = 0$ subspace to access larger system sizes. b) Proportion of Steady state (with totally spin-up initial state) in each S_z symmetry sector. Specifically, the coefficients $\alpha_{\frac{N}{2}, q_z, 0}$ for each generalized Dicke state in the steady state. Numerics were done with $N = 80$ and run for $t = 1000$. c) Long range correlations in the steady state. Transition point is consistent with closing of the gap in a).

steady state will be a linear combination of the W-state and the eigendensity-matrix corresponding to the third 0 eigenvalue. The relative weights of the linear combination may in general depend on the initial state and the values of $\gamma_1, \gamma_2, \gamma_3$. In Fig. 2b, we compare the steady states in this regime, taking as our initial state the totally spin polarized system $|\uparrow\uparrow \dots \uparrow\rangle$. There, we plot the distribution of coefficients $\alpha_{q, q_z, \sigma_z}$ (5) of the steady state for fixed $q = \frac{N}{2}, \sigma_z = 0$. These generalized Dicke state coefficients correspond to the sum of all diagonal terms of the full density matrix within the sector with spin magnetic quantum number $S_z = q_z$ (generalized Dicke states with $\sigma_z = 0$ have $\frac{N}{2} + q_z$ spin-ups and the $q = \frac{N}{2}, \sigma_z = 0$ coefficients make up the entire diagonal of the density matrix $\sum_{q_z} \alpha_{\frac{N}{2}, q_z, 0} = 1$). These coefficients provide a probe into the weight of each of the total S_z sectors in the final mixed density matrix.

When $\gamma_1, \gamma_2 \ll \gamma_3$, the steady state (labeled as ‘mixed’ in Fig. 2b) is the maximally mixed state (minus the all spin-down state). From counting the number of states in each total S_z sector, the distribution of coefficients for this ‘mixed’ state is given by $\alpha_{q, q_z, \sigma_z}^{\text{Mixed}} = \delta_{q, \frac{N}{2}} \delta_{\sigma_z, 0} \frac{\binom{N}{\frac{N}{2} + q_z}}{2^N - 1}$, where δ is the Kronecker delta and the -1 comes from the removal of the all spin-down state. As γ_3 decreases, the distribution shifts from the maximally mixed state to a distribution centered around smaller and smaller q_z . Additionally, the weight of the W-state increases as γ_3 becomes small (with $\alpha_{\frac{N}{2}, -\frac{N}{2} + 1, 0}$ increasing towards 1). In the regime where the gap opens, the steady state becomes the W-state.

As a measure of coherence in the system, we consider the two-point correlator $\langle X_a X_b \rangle$. Using generalized Dicke operators, it is possible to show that

$$\langle X_a X_b \rangle = \frac{1}{\binom{N}{2}} \sum_{q_z} \sum_{\ell=-1}^1 \alpha_{\frac{N}{2}-1, q_z+\ell, \ell} \quad \forall a \neq b \quad (7)$$

For the W-state, we have that $\langle X_a X_b \rangle_W = \frac{2}{N}$. In Fig. 2c, we compare $\langle X_a X_b \rangle$ for the steady state of our model to that of the W-state. The ratio $\frac{\langle X_a X_b \rangle}{\langle X_a X_b \rangle_W}$ departs from 1 at the same value of γ_3 where the gap closes (compare to Fig. 2a). Coherence still persists, however, after the transition. The steady state, therefore, does not just converge to the steady state of L_3 (the maximally mixed state) after the transition. Instead, the steady state is a mixture of states with coherence less than, but comparable to, that of the W-state. It is only in the limit $\gamma_3 \gg \gamma_1, \gamma_2$ that this coherence approaches 0 and the steady state becomes maximally mixed.

We now contrast the above all-to-all results with the NN version of the model. As previously discussed, the transition in the NN model occurs when $\frac{\gamma_3}{\gamma_1} = O(1)$. Furthermore, unlike the all-to-all transition which goes from gapped to gapless, the NN transition is from a polynomial gap to an exponential gap in system size. The replacement of the fixed gap (for all-to-all) with a polynomial gap is because the local jump operators in the NN model need time at least on order of the system size to generate coherence across the entire system. In the mixed phase, the NN model also exhibits coherence, but in this case the coherence decays exponentially in m for $\langle X_a X_{a+m} \rangle$ again due to locality. Numerics for small system sizes supporting these results are provided in Appendix F.

Conclusions—We have investigated a purely dissipative quantum reaction-diffusion model which exhibits competing terms that act to either cohere or decohere the system. The cohering terms may also be viewed as a dissipative quantum state preparation protocol for the generalized W-state, while the decohering terms act as errors to the preparation procedure. The transition occurs in the all-to-all version of the model for any non-zero rate of the decohering jump operators, implying the state preparation procedure is fragile to errors. For the NN version of the model,

however, the phase transition occurs when the error rate is inversely proportional to the coordination number instead of inversely proportional to the system size, implying the state preparation procedure is robust to weak error rates. We also find in both cases that some quantum coherence still persists even in the phase where the decohering terms dominate, only converging to the maximally mixed state when the error rate becomes infinite.

This model serves as a prototypical example of a quantum reaction-diffusion model where the absorbing state is long range coherent. It also serves as an example of how error quantum jumps may act to ruin dissipative quantum state preparation procedures even when the

target state is preserved as a dark state. This is most extreme in the all-to-all case where the preparation procedure fails for any non-zero error rate.

ACKNOWLEDGMENTS

This work was supported by a Simons Investigator Award [Grant No. 511029] and the Engineering and Physical Sciences Research Council [grant number EP/V062654/1]. For the purpose of open access, the authors have applied a creative commons attribution (CC BY) licence to any author accepted manuscript version arising.

-
- [1] J. T. Barreiro, M. Müller, P. Schindler, D. Nigg, T. Monz, M. Chwalla, M. Hennrich, C. F. Roos, P. Zoller, and R. Blatt, An open-system quantum simulator with trapped ions, *Nature* **470**, 486 (2011).
- [2] M. Müller, S. Diehl, G. Pupillo, and P. Zoller, Engineered open systems and quantum simulations with atoms and ions, in *Advances in Atomic, Molecular, and Optical Physics*, *Advances In Atomic, Molecular, and Optical Physics*, Vol. 61, edited by P. Berman, E. Arimondo, and C. Lin (Academic Press, 2012) pp. 1–80.
- [3] T. Langen, R. Geiger, and J. Schmiedmayer, Ultracold atoms out of equilibrium, *Annual Review of Condensed Matter Physics* **6**, 201 (2015).
- [4] M. Kjaergaard, M. E. Schwartz, J. Braumüller, P. Krantz, J. I.-J. Wang, S. Gustavsson, and W. D. Oliver, Superconducting qubits: Current state of play, *Annual Review of Condensed Matter Physics* **11**, 369 (2020).
- [5] T. D. Farokh Mivehvar, Francesco Piazza and H. Ritsch, Cavity QED with quantum gases: new paradigms in many-body physics, *Advances in Physics* **70**, 1 (2021), <https://doi.org/10.1080/00018732.2021.1969727>.
- [6] F. Verstraete, M. M. Wolf, and J. Ignacio Cirac, Quantum computation and quantum-state engineering driven by dissipation, *Nature Physics* **5**, 633 (2009).
- [7] M. Ippoliti, M. J. Gullans, S. Gopalakrishnan, D. A. Huse, and V. Khemani, Entanglement phase transitions in measurement-only dynamics, *Phys. Rev. X* **11**, 011030 (2021).
- [8] P. M. Harrington, E. J. Mueller, and K. W. Murch, Engineered dissipation for quantum information science, *Nature Reviews Physics* **4**, 660 (2022).
- [9] S. O. Valenzuela, W. D. Oliver, D. M. Berns, K. K. Berggren, L. S. Levitov, and T. P. Orlando, Microwave-induced cooling of a superconducting qubit, *Science* **314**, 1589 (2006), <https://www.science.org/doi/pdf/10.1126/science.1134008>.
- [10] B. Kraus, H. P. Büchler, S. Diehl, A. Kantian, A. Micheli, and P. Zoller, Preparation of entangled states by quantum Markov processes, *Phys. Rev. A* **78**, 042307 (2008).
- [11] F. Ticozzi and L. Viola, Stabilizing entangled states with quasi-local quantum dynamical semigroups, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **370**, 5259 (2012), <https://royalsocietypublishing.org/doi/pdf/10.1098/rsta.2011.0485>.
- [12] F. Ticozzi and L. Viola, Steady-state entanglement by engineered quasi-local Markovian dissipation, *Quantum Info. Comput.* **14**, 0265 (2014).
- [13] N. Didier, F. Qassemi, and A. Blais, Perfect squeezing by damping modulation in circuit quantum electrodynamics, *Phys. Rev. A* **89**, 013820 (2014).
- [14] Z. Leghtas, S. Touzard, I. M. Pop, A. Kou, B. Vlastakis, A. Petrenko, K. M. Sliwa, A. Narla, S. Shankar, M. J. Hatridge, M. Reagor, L. Frunzio, R. J. Schoelkopf, M. Mirrahimi, and M. H. Devoret, Confining the state of light to a quantum manifold by engineered two-photon loss, *Science* **347**, 853 (2015), <https://www.science.org/doi/pdf/10.1126/science.aaa2085>.
- [15] Y. Liu, S. Shankar, N. Ofek, M. Hatridge, A. Narla, K. M. Sliwa, L. Frunzio, R. J. Schoelkopf, and M. H. Devoret, Comparing and combining measurement-based and driven-dissipative entanglement stabilization, *Phys. Rev. X* **6**, 011022 (2016).
- [16] S. Boutin, C. K. Andersen, J. Venkatraman, A. J. Ferris, and A. Blais, Resonator reset in circuit QED by optimal control for large open quantum systems, *Phys. Rev. A* **96**, 042315 (2017).
- [17] F. Nathan, L. O'Brien, K. Noh, M. H. Matheny, A. L. Grimsmo, L. Jiang, and G. Refael, Self-correcting GKP qubit and gates in a driven-dissipative circuit (2024), arXiv:2405.05671 [cond-mat.mes-hall].
- [18] E. Kapit, The upside of noise: engineered dissipation as a resource in superconducting circuits, *Quantum Science and Technology* **2**, 033002 (2017).
- [19] F. Reiter, A. S. Sørensen, P. Zoller, and C. A. Muschik, Dissipative quantum error correction and application to quantum sensing with trapped ions, *Nature Communications* **8**, 1822 (2017).
- [20] J. M. Gertler, B. Baker, J. Li, S. Shirol, J. Koch, and C. Wang, Protecting a bosonic qubit with autonomous quantum error correction, *Nature* **590**, 243 (2021).
- [21] B. de Neeve, T.-L. Nguyen, T. Behrle, and J. P. Home, Error correction of a logical grid state qubit by dissipative pumping, *Nature Physics* **18**, 296 (2022).
- [22] V. V. Sivak, A. Eickbusch, B. Royer, S. Singh,

- I. Tsioutsios, S. Ganjam, A. Miano, B. L. Brock, A. Z. Ding, L. Frunzio, S. M. Girvin, R. J. Schoelkopf, and M. H. Devoret, Real-time quantum error correction beyond break-even, *Nature* **616**, 50 (2023).
- [23] H. J. Briegel, D. E. Browne, W. Dür, R. Raussendorf, and M. Van den Nest, Measurement-based quantum computation, *Nature Physics* **5**, 19 (2009).
- [24] D. F. Walls, P. D. Drummond, S. S. Hassan, and H. J. Carmichael, Non-Equilibrium Phase Transitions in Cooperative Atomic Systems, *Progress of Theoretical Physics Supplement* **64**, 307 (1978), <https://academic.oup.com/ptps/article-pdf/doi/10.1143/PTPS.64.307/5292058/64-307.pdf>.
- [25] E. G. Dalla Torre, E. Demler, T. Giamarchi, and E. Altman, Quantum critical states and phase transitions in the presence of non-equilibrium noise, *Nature Physics* **6**, 806 (2010).
- [26] E. G. D. Torre, S. Diehl, M. D. Lukin, S. Sachdev, and P. Strack, Keldysh approach for nonequilibrium phase transitions in quantum optics: Beyond the Dicke model in optical cavities, *Phys. Rev. A* **87**, 023831 (2013).
- [27] J. Klinder, H. Keßler, M. Wolke, L. Mathey, and A. Hemmerich, Dynamical phase transition in the open Dicke model, *Proceedings of the National Academy of Sciences* **112**, 3290 (2015), <https://www.pnas.org/doi/pdf/10.1073/pnas.1417132112>.
- [28] M. Fitzpatrick, N. M. Sundaresan, A. C. Y. Li, J. Koch, and A. A. Houck, Observation of a dissipative phase transition in a one-dimensional circuit QED lattice, *Phys. Rev. X* **7**, 011016 (2017).
- [29] B. Skinner, J. Ruhman, and A. Nahum, Measurement-induced phase transitions in the dynamics of entanglement, *Phys. Rev. X* **9**, 031009 (2019).
- [30] A.-L. Barabási and H. E. Stanley, *Fractal Concepts in Surface Growth* (Cambridge University Press, 1995).
- [31] S. Maerivoet and B. De Moor, Cellular automata models of road traffic, *Physics Reports* **419**, 1 (2005).
- [32] T. E. Harris, Contact Interactions on a Lattice, *The Annals of Probability* **2**, 969 (1974).
- [33] H. Hinrichsen, Non-equilibrium critical phenomena and phase transitions into absorbing states, *Advances in Physics* **49**, 815 (2000), <https://doi.org/10.1080/00018730050198152>.
- [34] M. van Horsen and J. P. Garrahan, Open quantum reaction-diffusion dynamics: Absorbing states and relaxation, *Phys. Rev. E* **91**, 032132 (2015).
- [35] M. Marcuzzi, M. Buchhold, S. Diehl, and I. Lesanovsky, Absorbing state phase transition with competing quantum and classical fluctuations, *Phys. Rev. Lett.* **116**, 245701 (2016).
- [36] R. Gutiérrez, C. Simonelli, M. Archimi, F. Castellucci, E. Arimondo, D. Ciampini, M. Marcuzzi, I. Lesanovsky, and O. Morsch, Experimental signatures of an absorbing-state phase transition in an open driven many-body quantum system, *Phys. Rev. A* **96**, 041602 (2017).
- [37] F. Carollo, E. Gillman, H. Weimer, and I. Lesanovsky, Critical behavior of the quantum contact process in one dimension, *Phys. Rev. Lett.* **123**, 100604 (2019).
- [38] C.-J. Lin and L. Zou, Reaction-diffusion dynamics in a fibonacci chain: Interplay between classical and quantum behavior, *Phys. Rev. B* **103**, 174305 (2021).
- [39] G. Peretto, F. Carollo, J. P. Garrahan, and I. Lesanovsky, Reaction-limited quantum reaction-diffusion dynamics, *Phys. Rev. Lett.* **130**, 210402 (2023).
- [40] G. Peretto, F. Carollo, J. P. Garrahan, and I. Lesanovsky, Quantum reaction-limited reaction-diffusion dynamics of annihilation processes, *Phys. Rev. E* **108**, 064104 (2023).
- [41] J. T. Barreiro, M. Müller, P. Schindler, D. Nigg, T. Monz, M. Chwalla, M. Hennrich, C. F. Roos, P. Zoller, and R. Blatt, An open-system quantum simulator with trapped ions, *Nature* **470**, 486 (2011).
- [42] S. Shankar, M. Hatridge, Z. Leghtas, K. M. Sliwa, A. Narla, U. Vool, S. M. Girvin, L. Frunzio, M. Mirrahimi, and M. H. Devoret, Autonomously stabilized entanglement between two superconducting quantum bits, *Nature* **504**, 419 (2013).
- [43] Z. Leghtas, S. Touzard, I. M. Pop, A. Kou, B. Vlastakis, A. Petrenko, K. M. Sliwa, A. Narla, S. Shankar, M. J. Hatridge, M. Reagor, L. Frunzio, R. J. Schoelkopf, M. Mirrahimi, and M. H. Devoret, Confining the state of light to a quantum manifold by engineered two-photon loss, *Science* **347**, 853 (2015), <https://www.science.org/doi/pdf/10.1126/science.aaa2085>.
- [44] A. Pechen and H. Rabitz, Teaching the environment to control quantum systems, *Phys. Rev. A* **73**, 062102 (2006).
- [45] P. Scaramuzza and F. Ticozzi, Switching quantum dynamics for fast stabilization, *Phys. Rev. A* **91**, 062314 (2015).
- [46] T. Grigoletto and F. Ticozzi, Stabilization via feedback switching for quantum stochastic dynamics, *IEEE Control Systems Letters* **6**, 235 (2022).
- [47] S. Morales, Y. Gefen, I. Gornyi, A. Zazunov, and R. Egger, Engineering unsteerable quantum states with active feedback, *Phys. Rev. Res.* **6**, 013244 (2024).
- [48] C. Aron, M. Kulkarni, and H. E. Türeci, Photon-mediated interactions: A scalable tool to create and sustain entangled states of n atoms, *Phys. Rev. X* **6**, 011032 (2016).
- [49] C.-M. Halati, A. Sheikhan, and C. Kollath, Cavity-induced artificial gauge field in a Bose-Hubbard ladder, *Phys. Rev. A* **96**, 063621 (2017).
- [50] D. C. Cole, J. J. Wu, S. D. Erickson, P.-Y. Hou, A. C. Wilson, D. Leibfried, and F. Reiter, Dissipative preparation of W states in trapped ion systems, *New Journal of Physics* **23**, 073001 (2021).
- [51] S. Diehl, A. Micheli, A. Kantian, B. Kraus, H. P. Büchler, and P. Zoller, Quantum states and phases in driven open quantum systems with cold atoms, *Nature Physics* **4**, 878 (2008).
- [52] S. Hartmann, Generalized Dicke states, *Quantum Info. Comput.* **16**, 1333–1348 (2016).
- [53] R. H. Dicke, Coherence in spontaneous radiation processes, *Phys. Rev.* **93**, 99 (1954).
- [54] B. A. Chase and J. M. Geremia, Collective processes of an ensemble of spin-1/2 particles, *Phys. Rev. A* **78**, 052101 (2008).
- [55] H. Georgi, *Lie Algebras In Particle Physics* (Taylor and Francis, 2000).

Appendix A: Generalized Dicke States and Operators

We briefly review here generalized Dicke states and operators. See [52] for further details and relations. Consider dynamics under Lindblad operators (or, more generally, Kraus operators) which act symmetrically on a set of N sites. As first noted (independently) in [52, 54], both analytical and numerical investigations into such scenarios may be dramatically simplified by leveraging the fact that the set of Lindblad operators of this type is closed under $SU(4)$ transformations. As we will see below, the number of basis states for the space of symmetric density matrices scales only polynomially in N , providing an exponential improvement over the $2^N \times 2^N$ scaling of the full density matrix. In analogy with Dicke states – the set of fully permutation symmetric pure states (closed under $SU(2)$ operations instead of $SU(4)$) that form the appropriate basis for Hamiltonians which act symmetrically on all sites – the basis of fully symmetric density matrices are referred to as generalized Dicke states and the generators of $SU(4)$ which act on them are referred to as generalized Dicke operators.

Specifically, the set of all permutation symmetric Lindblad operators may be generated by:

$$\begin{aligned}
\mathcal{Q}_\pm[\rho] &:= \sum_{j=1}^N \sigma_j^\pm \rho \sigma_j^\mp; & \mathcal{Q}_z[\rho] &:= \frac{1}{4} \sum_{j=1}^N (\sigma_j^z \rho + \rho \sigma_j^z) \\
\mathcal{S}_\pm[\rho] &:= \sum_{j=1}^N \sigma_j^\pm \rho \sigma_j^\pm; & \mathcal{S}_z[\rho] &:= \frac{1}{4} \sum_{j=1}^N (\sigma_j^z \rho - \rho \sigma_j^z) \\
\mathcal{M}_\pm[\rho] &:= \sum_{j=1}^N \sigma_j^\pm \rho \frac{1 + \sigma_j^z}{2}; & \mathcal{M}_z[\rho] &:= \frac{1}{2} \sum_{j=1}^N \sigma_j^z \rho \frac{1 + \sigma_j^z}{2} \\
\mathcal{N}_\pm[\rho] &:= \sum_{j=1}^N \sigma_j^\pm \rho \frac{1 - \sigma_j^z}{2}; & \mathcal{N}_z[\rho] &:= \frac{1}{2} \sum_{j=1}^N \sigma_j^z \rho \frac{1 - \sigma_j^z}{2} \\
\mathcal{U}_\pm[\rho] &:= \sum_{j=1}^N \frac{1 + \sigma_j^z}{2} \rho \sigma_j^\mp; & \mathcal{U}_z[\rho] &:= \frac{1}{2} \sum_{j=1}^N \frac{1 + \sigma_j^z}{2} \rho \sigma_j^z \\
\mathcal{V}_\pm[\rho] &:= \sum_{j=1}^N \frac{1 - \sigma_j^z}{2} \rho \sigma_j^\mp; & \mathcal{V}_z[\rho] &:= \frac{1}{2} \sum_{j=1}^N \frac{1 - \sigma_j^z}{2} \rho \sigma_j^z
\end{aligned} \tag{A1}$$

Of the 18 generalized Dicke operators listed above, 15 are linearly independent and correspond to the generators of $SU(4)$.

The generalized Dicke states are fundamentally represented in terms of the single site density operators

$$\begin{aligned}
u &= |1\rangle\langle 1|, & d &= |0\rangle\langle 0| \\
s &= |1\rangle\langle 0|, & c &= |0\rangle\langle 1|
\end{aligned} \tag{A2}$$

where u, d, s, c , i.e. up, down, strange, and charm, are a reference to the $SU(4)$ four-flavor quark model [55]. Specifically, the generalized Dicke states are given by

$$\mathcal{D}_{q, q_z, \sigma_z} = \mathcal{S} (u^\alpha d^\beta s^\gamma c^\delta) \tag{A3}$$

where \mathcal{S} is the symmetrizer and $\alpha + \beta + \gamma + \delta = N$. The quantum numbers q, q_z, σ_z are given in terms of $\alpha, \beta, \gamma, \delta$ by

$$q = \frac{\alpha + \beta}{2}, \tag{A4}$$

$$q_z = \frac{\alpha - \beta}{2}, \tag{A5}$$

$$\sigma_z = \frac{\gamma - \delta}{2}. \tag{A6}$$

The range of possible q, q_z, σ_z are $q = 0, \frac{1}{2}, \dots, \frac{N}{2}$; $q_z = -q, -q + 1, \dots, q$; and $\sigma_z = -\frac{N}{2} + q, -\frac{N}{2} + q + 1, \dots, \frac{N}{2} - q$. By simple counting arguments, this implies there are a total of $\frac{1}{6}(N+1)(N+2)(N+3) = O(N^3)$ generalized Dicke states.

It is possible to denote the action of the generalized Dicke operators (A1) onto the states (A3). Namely,

Operator	(q, q_z, σ_z)	$+, -$	Operator	Coeff.
\mathcal{Q}_\pm	$(0, \pm 1, 0)$	β, α	\mathcal{Q}_z	$\frac{\beta - \alpha}{2} = -q_z$
Σ_\pm	$(0, 0, \pm 1)$	δ, γ	Σ_z	$\frac{\delta - \gamma}{2} = -\sigma_z$
\mathcal{M}_\pm	$(\mp \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2})$	β, γ	\mathcal{M}_z	$\frac{\beta - \gamma}{2}$
\mathcal{N}_\pm	$(\pm \frac{1}{2}, \pm \frac{1}{2}, \pm \frac{1}{2})$	δ, α	\mathcal{N}_z	$\frac{\delta - \alpha}{2}$
\mathcal{U}_\pm	$(\mp \frac{1}{2}, \pm \frac{1}{2}, \mp \frac{1}{2})$	β, δ	\mathcal{U}_z	$\frac{\beta - \delta}{2}$
\mathcal{V}_\pm	$(\pm \frac{1}{2}, \pm \frac{1}{2}, \mp \frac{1}{2})$	γ, α	\mathcal{V}_z	$\frac{\gamma - \alpha}{2}$

(A7)

where the table above denotes how each operator alters the quantum numbers and what coefficient it adds to the generalized Dicke state. For example,

$$\mathcal{Q}_+ \mathcal{D}_{q, q_z, \sigma_z} = \beta \mathcal{D}_{q, q_z + 1, \sigma_z}, \quad (\text{A8})$$

$$\mathcal{Q}_z \mathcal{D}_{q, q_z, \sigma_z} = -q_z \mathcal{D}_{q, q_z, \sigma_z}. \quad (\text{A9})$$

Due to the algebra above, it will often be useful to represent permutation symmetric Lindbladians as transformations on a 3 dimensional lattice with each dimension corresponding to a quantum number of the generalized Dicke states. Each site is labeled by a generalized Dicke state

$$\mathcal{D}_{q, q_z, \sigma_z} \equiv |q, q_z, \sigma_z\rangle, \quad (\text{A10})$$

and a general symmetric Lindbladian \mathcal{L} will have matrix elements of the form

$$\mathcal{L} = \sum_{\substack{q, q_z, \sigma_z \\ q', q'_z, \sigma'_z}} \mathcal{L}(q', q'_z, \sigma'_z; q, q_z, \sigma_z) |q', q'_z, \sigma'_z\rangle \langle q, q_z, \sigma_z|. \quad (\text{A11})$$

In general, a density matrix will take the form of a linear combination of generalized Dicke states

$$\rho = \sum_{q, q_z, \sigma_z} \alpha_{q, q_z, \sigma_z} |q, q_z, \sigma_z\rangle. \quad (\text{A12})$$

We will often omit quantum numbers on the coefficients for density matrices confined to a subspace with a fixed quantum number. For example, if the evolution of a density matrix is confined within the $\sigma_z = 0$ sector, we will write its coefficients as α_{q, q_z} . We note that the $\alpha_{q, q_z, \sigma_z}$ are constrained by the condition $\text{Tr } \rho = 1$, in particular

$$\sum_{q_z} \alpha_{\frac{N}{2}, q_z, 0} = 1. \quad (\text{A13})$$

Appendix B: All-to-All Model with Generalized Dicke Operators

We here write the Lindbladian for the all-to-all version of our model in terms of generalized Dicke operators. We subdivide the Lindbladian into terms

$$\mathcal{L} = \mathcal{L}_1 + \mathcal{L}_2 + \mathcal{L}_3 + \mathcal{L}_{3'} \quad (\text{B1})$$

with each \mathcal{L}_α denoting the part of the Lindblad superoperator which corresponds to the set of jump operators (3a), (3b), (3c), or (3d) in the main text (with all pairs i, j instead of nearest neighbors $\langle i, j \rangle$). In other words, we define

$$\mathcal{L}_\alpha [\rho] = \sum_{L \in \mathcal{L}_\alpha} \gamma_\alpha \left(L \rho L^\dagger - \frac{1}{2} \{L^\dagger L, \rho\} \right). \quad (\text{B2})$$

where each $L \in L_\alpha$ corresponds to the jump operator of form L_α acting on a given site pair i, j .

In terms of generalized Dicke operators, the operators \mathcal{L}_α may be written

$$\mathcal{L}_1 = \gamma_1 \{ \mathcal{M}_+ \mathcal{N}_- + \mathcal{U}_+ \mathcal{V}_- - 2\mathcal{S}\mathcal{C} \} \quad (\text{B3})$$

$$\mathcal{L}_2 = \gamma_2 \left\{ \mathcal{N}_- \mathcal{V}_- + \mathcal{Q}_- \left[\frac{N}{4} - \mathcal{Q}_z + \frac{1}{2}(\mathcal{M}_z - \mathcal{N}_z) - 1 \right] - \left(\frac{N}{2} - \mathcal{Q}_z \right) \left(\frac{N}{2} - \mathcal{Q}_z - 1 \right) - \Sigma_z^2 \right\} \quad (\text{B4})$$

$$\begin{aligned} \mathcal{L}_3 + \mathcal{L}_{3'} = \gamma_3 \left\{ (\mathcal{Q}_- + \mathcal{Q}_+) \left[\frac{N}{4} - \mathcal{Q}_z + \frac{1}{2}(\mathcal{M}_z - \mathcal{N}_z) \right] - \mathcal{Q}_- - N \left(\frac{N}{2} - \mathcal{Q}_z \right) \right. \\ \left. - \mathcal{N}_+ \mathcal{V}_+ - \mathcal{N}_- \mathcal{V}_- + \frac{1}{2} [(\mathcal{M}_+ + \mathcal{N}_+) (\mathcal{M}_- + \mathcal{N}_-) + (\mathcal{U}_+ + \mathcal{V}_+) (\mathcal{U}_- + \mathcal{V}_-)] \right\}. \end{aligned} \quad (\text{B5})$$

where \mathcal{S} and \mathcal{C} are given by

$$\mathcal{S} = \frac{N}{4} + \mathcal{N}_z - \frac{1}{2}\mathcal{Q}_z - \frac{3}{2}\Sigma_z \quad (\text{B6})$$

$$\mathcal{C} = \frac{N}{4} + \mathcal{N}_z - \frac{1}{2}\mathcal{Q}_z + \frac{1}{2}\Sigma_z \quad (\text{B7})$$

and correspond to the the number operators for the number of s and c states in the generalized Dicke state, i.e.

$$\mathcal{S}\mathcal{D}_{q,q_z,\sigma_z} = \left(\frac{N}{2} - q + \sigma_z \right) \mathcal{D}_{q,q_z,\sigma_z} = \gamma \mathcal{D}_{q,q_z,\sigma_z} \quad (\text{B8})$$

$$\mathcal{C}\mathcal{D}_{q,q_z,\sigma_z} = \left(\frac{N}{2} - q - \sigma_z \right) \mathcal{D}_{q,q_z,\sigma_z} = \delta \mathcal{D}_{q,q_z,\sigma_z}. \quad (\text{B9})$$

Equations (B3), (B4), and (B5) were found by writing the jump operators in terms of Pauli operators

$$|1,0\rangle_{i,j}\langle 0,0|_{i,j} = \frac{\sigma_i^z - \sigma_j^z - i\sigma_i^y\sigma_j^x + i\sigma_i^x\sigma_j^y}{4}, \quad (\text{B10})$$

$$|1,0\rangle_{i,j}\langle 1,1|_{i,j} = \frac{1}{\sqrt{2}} \left[\sigma_i^- \left(\frac{I - \sigma_j^z}{2} \right) + \left(\frac{I - \sigma_i^z}{2} \right) \sigma_j^- \right], \quad (\text{B11})$$

$$|0,0\rangle_{i,j}\langle 1,1|_{i,j} = \frac{1}{\sqrt{2}} \left[\sigma_i^- \left(\frac{I - \sigma_j^z}{2} \right) - \left(\frac{I - \sigma_i^z}{2} \right) \sigma_j^- \right], \quad (\text{B12})$$

$$|1,1\rangle_{i,j}\langle 0,0|_{i,j} = \frac{1}{\sqrt{2}} \left[\sigma_i^+ \left(\frac{I - \sigma_j^z}{2} \right) - \left(\frac{I - \sigma_i^z}{2} \right) \sigma_j^+ \right]. \quad (\text{B13})$$

in Eq. (B2) and using $\sum_{i,j} \text{s.t. } i \neq j = \sum_{i,j} - \sum_{i} \text{s.t. } i=j$. Above, we have written 2-site states in terms of total spin eigenstates $|s, m_s\rangle$ given by

$$\begin{aligned} |1,1\rangle_{ij} &= |\uparrow\uparrow\rangle_{ij}, \quad |1,0\rangle_{ij} = \frac{1}{\sqrt{2}}(|\downarrow\uparrow\rangle_{ij} + |\uparrow\downarrow\rangle_{ij}), \\ |1,-1\rangle_{ij} &= |\downarrow\downarrow\rangle_{ij}, \quad |0,0\rangle_{ij} = \frac{1}{\sqrt{2}}(|\downarrow\uparrow\rangle_{ij} - |\uparrow\downarrow\rangle_{ij}). \end{aligned} \quad (\text{B14})$$

Appendix C: W-State Preparation

In this section, we show that $\mathcal{L}_1 + \mathcal{L}_2$ together prepare the W-state. Let us begin by solving for the steady state of \mathcal{L}_1 . By acting with \mathcal{L}_1 on the state $\mathcal{D}_{q,q_z,\sigma_z}$, the matrix elements of \mathcal{L}_1 in the representation (A11) may be found to be

$$\begin{aligned}\mathcal{L}_1(q-1; q) &= 2(q+q_z)(q-q_z) \\ \mathcal{L}_1(q; q) &= -2\left(\frac{N}{2}-q+\sigma_z\right)\left(\frac{N}{2}-q-\sigma_z\right)\end{aligned}\tag{C1}$$

where we have denoted terms which leave q_z, σ_z invariant as $\mathcal{L}_1(q-1; q) \equiv \mathcal{L}_1(q-1, q_z, \sigma_z; q, q_z, \sigma_z)$. We will refer to the first term above, which takes $q \rightarrow q-1$, as the ‘‘hopping’’ term and the second term above, which leaves q, q_z, σ_z invariant, as the ‘‘on-site’’ term. Note, as expected, (C1) does not alter q_z and thus there will be a different, independent solution for each symmetry sector q_z . Similarly, σ_z is also invariant under the action of the Lindbladian. We therefore focus on the evolution of q to find the steady state of the dynamics. The steady state, for any given sector of q_z, σ_z , will be some linear combination of q generalized Dicke states

$$\mathcal{D}_{\text{steady}} = \sum_q \alpha_q \mathcal{D}_{q, q_z, \sigma_z}.\tag{C2}$$

The steady state condition $\mathcal{L}_1 \mathcal{D}_{\text{steady}} = 0$ then implies that the on-site term times α_q must be equal and opposite to the hopping term times α_{q+1} for all q .

Instead of solving directly for α_q , we solve for a related quantity, $\alpha_q^{(N)}$ defined below, which will help make the interpretation of the result more transparent. We remark that each $\alpha_{q, q_z, \sigma_z}$ corresponds to the sum of all the elements within the q, q_z, σ_z sector of the full density matrix [see (A12)]. Comparing different α_q is thus obscured by the fact that the different q, q_z, σ_z sectors vary in size. For a more direct comparison we define $\alpha_q^{(N)}$ as the weight of a single matrix element within the q, q_z, σ_z sector of the density matrix. In particular, let $\mathcal{N}(q, q_z, \sigma_z)$ be the number of density matrix elements within the q, q_z, σ_z sector, i.e.

$$\begin{aligned}\mathcal{N}(q, q_z, \sigma_z) &= \binom{N}{\alpha} \binom{N-\alpha}{\beta} \binom{N-\alpha-\beta}{\gamma} \\ &= \binom{N}{q+q_z} \binom{N-q-q_z}{q-q_z} \binom{N-2q}{\frac{N}{2}-q+\sigma_z}.\end{aligned}\tag{C3}$$

where the above is calculated using (A3) and basic counting arguments.

Therefore, we may now define

$$\alpha_q = \mathcal{N}(q, q_z, \sigma_z) \alpha_q^{(N)},\tag{C4}$$

It is possible to work directly in a basis which yields $\alpha_q^{(N)}$ instead of α_q by renormalizing the generalized Dicke states

$$|q, q_z, \sigma_z\rangle^{(N)} = \mathcal{N}(q, q_z, \sigma_z) |q, q_z, \sigma_z\rangle,\tag{C5}$$

$$\langle q, q_z, \sigma_z |^{(N)} = \frac{\langle q, q_z, \sigma_z |}{\mathcal{N}(q, q_z, \sigma_z)}.\tag{C6}$$

Writing any \mathcal{L} in this basis yields

$$\mathcal{L} = \sum_{\substack{q', q'_z, \sigma'_z \\ q', q'_z, \sigma'_z}} \mathcal{L}^{(N)}(q', q'_z, \sigma'_z; q, q_z, \sigma_z) |q', q'_z, \sigma'_z\rangle^{(N)} \langle q, q_z, \sigma_z |^{(N)}\tag{C7}$$

$$\mathcal{L}^{(N)}(q', q'_z, \sigma'_z; q, q_z, \sigma_z) = \mathcal{L}(q', q'_z, \sigma'_z; q, q_z, \sigma_z) \frac{\mathcal{N}(q, q_z, \sigma_z)}{\mathcal{N}(q', q'_z, \sigma'_z)}.\tag{C8}$$

Using that $\frac{\mathcal{N}(q, q_z, \sigma_z)}{\mathcal{N}(q-1, q_z, \sigma_z)} = \frac{(\frac{N}{2}-q+\sigma_z+1)(\frac{N}{2}-q-\sigma_z+1)}{(q+q_z)(q-q_z)}$, rewriting (C1) in the renormalized basis gives

$$\begin{aligned}\mathcal{L}_1^{(N)}(q; q+1) &= 2\left(\frac{N}{2}-q+\sigma_z\right)\left(\frac{N}{2}-q-\sigma_z\right), \\ \mathcal{L}_1^{(N)}(q; q) &= -2\left(\frac{N}{2}-q+\sigma_z\right)\left(\frac{N}{2}-q-\sigma_z\right).\end{aligned}\tag{C9}$$

In the renormalized basis, the hopping term is equal and opposite to the onsite term. Letting $\mathcal{D}_{\text{steady}}^{(N)} = \sum_q \alpha_q^{(N)} \mathcal{D}_{q,q_z,\sigma_z}^{(N)}$, the steady state condition implies

$$\begin{aligned} \mathcal{L}_1 \mathcal{D}_{\text{steady}}^{(N)} &= 0 \\ \implies \alpha_q^{(N)} \mathcal{L}_1^{(N)}(q; q) + \alpha_{q+1}^{(N)} \mathcal{L}_1^{(N)}(q; q+1) &= 0 \quad \forall q \\ \implies \alpha_{q+1}^{(N)} &= \alpha_q^{(N)} \end{aligned} \quad (\text{C10})$$

Note that, since the generalized Dicke states which correspond to the diagonal of the full density matrix all have $q = \frac{N}{2}$ and since $\text{Tr} \rho = 1$, we have that $\alpha_{\frac{N}{2}}^{(N)} = \frac{1}{\mathcal{N}(\frac{N}{2}, q_z, 0)} = \frac{1}{\binom{N}{\frac{N}{2} + q_z}}$. Therefore,

$$\alpha_q^{(N)} = \frac{1}{\binom{N}{\frac{N}{2} + q_z}}. \quad (\text{C11})$$

For the case $q_z = -\frac{N}{2} + 1$ and $\sigma_z = 0$, this is precisely given by the W state

$$\rho_W = |W\rangle\langle W|. \quad (\text{C12})$$

More generally, the steady states corresponding to any $\sigma_z = 0$ sector are given by the pure Dicke state corresponding to the given q_z . The $\sigma_z \neq 0$ terms correspond to superpositions between the different particle number sectors. We note that all such $\sigma_z \neq 0$ generalized Dicke states have zero trace, and so must be accompanied (in linear combination) by $\sigma_z = 0$ generalized Dicke states to be a steady state. Furthermore, the hermiticity of the density matrix requires that the weight of $\mathcal{D}_{q,q_z,\sigma_z}$ and $\mathcal{D}_{q,q_z,-\sigma_z}$ must be related by a complex conjugate in the steady state.

As a technical note, the renormalized basis breaks down at the boundaries of the 3D lattice. Before renormalizing, any physical matrix element $\mathcal{L}(q', q'_z, \sigma'_z; q, q_z, \sigma_z)$ will be 0 if it maps a site on the lattice to a q', q'_z, σ'_z which falls outside the range of possible values for the quantum numbers. However, $\mathcal{N}(q', q'_z, \sigma'_z)$ is undefined for any choice of q', q'_z, σ'_z which is not within the allowed ranges. This leaves $\mathcal{L}^{(N)}(q', q'_z, \sigma'_z; q, q_z, \sigma_z)$ ill-defined. For example, at lattice site $|\frac{N}{2}, -\frac{N}{2}, 0\rangle$, the non-renormalized matrix element $\mathcal{L}_1(\frac{N}{2} - 1, \frac{N}{2}) = 0$ as this would make q_z lie outside its allowed range of $-q$ to q . However, the corresponding renormalized matrix element $\mathcal{L}_1^{(N)}(\frac{N}{2} - 1, \frac{N}{2}) = 2 \neq 0$. It is therefore necessary to reset all such boundary terms to 0 after renormalizing.

Let us now add \mathcal{L}_2 to the dynamics. First consider the $\sigma_z = 0$ sector. Here, the steady state is expanded in terms of generalized Dicke states as

$$\mathcal{D}_{\text{steady}}^{(N)} = \sum_q \alpha_{q,q_z}^{(N)} |q, q_z, \sigma_z\rangle^{(N)} \quad (\text{C13})$$

with the coefficients given by the system of equations $(\mathcal{L}_1 + \mathcal{L}_2) \mathcal{D}_{\text{steady}}^{(N)} = 0$. Specifically, we have

$$\begin{aligned} &\alpha_{q,q_z}^{(N)} \left(\mathcal{L}_1^{(N)}(q, q_z; q, q_z) + \mathcal{L}_2^{(N)}(q, q_z; q, q_z) \right) + \alpha_{q+1,q_z}^{(N)} \mathcal{L}_1^{(N)}(q, q_z; q+1, q_z) \\ &+ \alpha_{q,q_z+1}^{(N)} \mathcal{L}_2^{(N)}(q, q_z; q, q_z+1) + \alpha_{q+1,q_z+1}^{(N)} \mathcal{L}_2^{(N)}(q, q_z; q+1, q_z+1) = 0 \quad \forall q, q_z \end{aligned} \quad (\text{C14})$$

In the system of equations above, the term $\alpha_{\frac{N}{2}, \frac{N}{2}}^{(N)}$ only appears in the equation $\alpha_{\frac{N}{2}, \frac{N}{2}}^{(N)} \left(\mathcal{L}_1^{(N)}(\frac{N}{2}, \frac{N}{2}; \frac{N}{2}, \frac{N}{2}) + \mathcal{L}_2^{(N)}(\frac{N}{2}, \frac{N}{2}; \frac{N}{2}, \frac{N}{2}) \right) = -\gamma_2 N(N-1) \alpha_{\frac{N}{2}, \frac{N}{2}}^{(N)} = 0$. This implies $\alpha_{\frac{N}{2}, \frac{N}{2}}^{(N)} = 0$. By induction, all terms $\alpha_{q,q_z}^{(N)} = 0$ when $q_z > -\frac{N}{2} + 1$. This may be seen by using (C14) and taking the inductive step $q \rightarrow q-1$ if $q > q_z$ and $q \rightarrow \frac{N}{2}$, $q_z \rightarrow q_z - 1$ otherwise. In each inductive step, we are left with $\alpha_{q,q_z}^{(N)} \left(\mathcal{L}_1^{(N)}(q, q_z; q, q_z) + \mathcal{L}_2^{(N)}(q, q_z; q, q_z) \right) = 0$ as all $\alpha_{q,q_z}^{(N)}$ for larger q, q_z are 0 by the inductive hypothesis. When $q_z > -\frac{N}{2} + 1$, $\left(\mathcal{L}_1^{(N)}(q, q_z; q, q_z) + \mathcal{L}_2^{(N)}(q, q_z; q, q_z) \right) \neq 0$ which implies all $\alpha_{q,q_z}^{(N)} = 0$ for $q_z > -\frac{N}{2} + 1$. When $q_z \leq -\frac{N}{2} + 1$, all $\mathcal{L}_2^{(N)}$ matrix elements are 0, thus the longtime dynamics is completely determined by \mathcal{L}_1 in these sectors. As we saw previously in this section, \mathcal{L}_1 prepares a W -state for $q_z = -\frac{N}{2} + 1$ and the vacuum for $q_z = -\frac{N}{2}$.

Since there is no hopping term which connects the $q_z = -\frac{N}{2}$ sector to the rest of the lattice, then any initial state starting outside this sector (and with $\sigma_z = 0$) will converge to the W-state. Outside of the $\sigma_z = 0$ sector, we can apply a similar inductive argument, except this time there are no states where the on-site term is 0 (except for $\sigma_z = \pm\frac{1}{2}$ and $q_z = -\frac{N}{2} + \frac{1}{2}$ which correspond to states in a superposition between the vacuum and the 1 particle sector). This means, for any initial state which does not include the vacuum, all of the coefficients with $\sigma_z \neq 0$ will be 0 and the system will converge to the W-state.

Appendix D: Absorbing State Phase Transition

The matrix elements for \mathcal{L}_1 , \mathcal{L}_2 , and $\mathcal{L}_3 + \mathcal{L}_{3'}$, after renormalizing by $\mathcal{N}(q, q_z, \sigma_z)$, are given by

$$\begin{aligned}\mathcal{L}_1^{(N)}(q-1, q_z; q, q_z) &= 2\gamma_1 \left[\left(\frac{N}{2} - q + 1 \right)^2 - \sigma_z^2 \right] \\ \mathcal{L}_1^{(N)}(q, q_z; q, q_z) &= -2\gamma_1 \left[\left(\frac{N}{2} - q \right)^2 - \sigma_z^2 \right]\end{aligned}\tag{D1}$$

$$\begin{aligned}\mathcal{L}_2^{(N)}(q, q_z - 1; q, q_z) &= \gamma_2 (q + q_z - 1)(q - q_z + 1) \\ \mathcal{L}_2^{(N)}(q-1, q_z - 1; q, q_z) &= \gamma_2 \left[\left(\frac{N}{2} - q + 1 \right)^2 - \sigma_z^2 \right] \\ \mathcal{L}_2^{(N)}(q, q_z; q, q_z) &= -\gamma_2 \left(\frac{N}{2} + q_z \right) \left(\frac{N}{2} + q_z - 1 \right) - \gamma_2 \sigma_z^2\end{aligned}\tag{D2}$$

$$\begin{aligned}\mathcal{L}_3^{(N)}(q, q_z + 1; q, q_z) &= \gamma_3 (q + q_z)(q + q_z + 1) \\ \mathcal{L}_3^{(N)}(q, q_z - 1; q, q_z) &= \gamma_3 (q + q_z - 1)(q - q_z + 1) \\ \mathcal{L}_3^{(N)}(q+1, q_z + 1; q, q_z) &= -\gamma_3 (q + q_z + 2)(q + q_z + 1) \\ \mathcal{L}_3^{(N)}(q-1, q_z - 1; q, q_z) &= -\gamma_3 \left[\left(\frac{N}{2} - q + 1 \right)^2 - \sigma_z^2 \right] \\ \mathcal{L}_3^{(N)}(q+1, q_z; q, q_z) &= \gamma_3 (q + q_z + 1)(q - q_z + 1) \\ \mathcal{L}_3^{(N)}(q-1, q_z; q, q_z) &= \gamma_3 \left[\left(\frac{N}{2} - q + 1 \right)^2 - \sigma_z^2 \right] \\ \mathcal{L}_3^{(N)}(q, q_z; q, q_z) &= -\gamma_3 \left[(N-1) \left(\frac{N}{2} + q_z \right) - 2q \left(\frac{N}{2} - q \right) \right]\end{aligned}\tag{D3}$$

It is straightforward to check from these matrix elements that the W-state, written in terms of renormalized generalized Dicke states as

$$\mathcal{D}_W^{(N)} = \frac{1}{N} \left| \frac{N}{2} - 1, -\frac{N}{2} + 1, 0 \right\rangle^{(N)} + \frac{1}{N} \left| \frac{N}{2}, -\frac{N}{2} + 1, 0 \right\rangle^{(N)},\tag{D4}$$

is individually a steady state of \mathcal{L}_1 , \mathcal{L}_2 , and \mathcal{L}_3 . Furthermore, the vacuum $|\frac{N}{2}, -\frac{N}{2}, 0\rangle$ is also a steady state. The Lindbladian \mathcal{L}_3 , on the other hand, has another steady state - the maximally mixed state (within the subspace of all states excluding the vacuum). Writing in the renormalized generalized Dicke basis, the maximally mixed (sans vacuum) state is given by

$$\mathcal{D}_{\text{mixed}}^{(N)} = \sum_{q_z \neq -\frac{N}{2}} \frac{1}{2^N - 1} \left| \frac{N}{2}, q_z, 0 \right\rangle^{(N)}.\tag{D5}$$

Using (D3), we have $\mathcal{L}_3^{(N)} \mathcal{D}_{\text{mixed}}^{(N)} = 0$, i.e. it is indeed a steady state.

We remark that a phase transition will still occur even if $\gamma_2 = 0$. This may be understood in the following way. If $\gamma_1 \gg \gamma_3$, then between each quantum jump from γ_3 terms, the contribution from γ_1 will push each q_z sector to the corresponding pure Dicke state. If the pure Dicke state is reached for a given q_z sector, then \mathcal{L}_3 can no longer raise q_z (as Dicke states contain no singlets) and \mathcal{L}_3 lower q_z out of that sector. Thus, now γ_3 acts to lower q_z until the system is within the 1 particle subspace. The addition of the \mathcal{L}_2 term then only serves to increase the critical value of γ_3 where the transition takes place.

Appendix E: Coherence in the Mixed Steady State

We here derive Eq. (7) from the manuscript. First, we use permutation symmetry to rewrite $\langle X_a X_b \rangle$ in terms of all-to-all operators. Namely,

$$\langle X_a X_b \rangle = \frac{1}{N(N-1)} \sum_{a \neq b} \langle X_a X_b \rangle = \frac{1}{N(N-1)} \left[\left\langle \left(\sum_a X_a \right) \left(\sum_b X_b \right) \right\rangle - N \right]. \quad (\text{E1})$$

Let us now rewrite $\langle (\sum_a X_a) (\sum_b X_b) \rangle$ in terms of generalized Dicke states and operators. Note that

$$\left(\sum_a X_a \right) \rho = (\mathcal{M}_+ + \mathcal{N}_+ + \mathcal{M}_- + \mathcal{N}_-) [\rho] \quad (\text{E2})$$

and that $\text{Tr} = \delta_{q,0} \delta_{\sigma_z,0} \sum_{q_z} \langle q, q_z, \sigma_z |$. We therefore have

$$\left\langle \left(\sum_a X_a \right) \left(\sum_b X_b \right) \right\rangle = \delta_{q,0} \delta_{\sigma_z,0} \sum_{q_z} \alpha_{q,q_z,\sigma_z} \langle q, q_z, \sigma_z | (\mathcal{M}_+ + \mathcal{N}_+ + \mathcal{M}_- + \mathcal{N}_-)^2 | q, q_z, \sigma_z \rangle \quad (\text{E3})$$

$$= N + 2 \sum_{q_z} \sum_{\ell=-1}^1 \alpha_{\frac{N}{2}-1, q_z+\ell, \ell}. \quad (\text{E4})$$

Therefore, combining equations (E1) and (E4) we have

$$\langle X_a X_b \rangle = \frac{1}{\binom{N}{2}} \sum_{q_z} \sum_{\ell=-1}^1 \alpha_{\frac{N}{2}-1, q_z+\ell, \ell}. \quad (\text{E5})$$

which is (7) in the main text.

Appendix F: Nearest-Neighbor Results

Below are small system size numerics supporting the NN results discussed in the main manuscript. Namely, Fig. 3 shows the transition from exponential gap to power law gap in system size. Fig. 4 shows correlations in the mixed phase. Similar to the all-to-all case, the NN model exhibits correlations in the mixed phase which become stronger as γ_3 approaches the critical value. However, in this case the correlations are short range, decaying exponentially with distance. Numerics were performed on a system of 10 sites with open boundary conditions. Correlations are for the eigendensity matrix corresponding to the mixed steady state. As a technical point, the true steady state is a linear combination of the eigendensity matrices corresponding to the three 0 eigenvalues. Since the eigenvalues are degenerate, one must take a general linear combination ρ of the three steady eigendensity matrices and orthonormalize such that $\text{Tr} \rho = 1$, $\text{Tr} \rho |W\rangle \langle W| = 0$, $\text{Tr} \rho |0\rangle \langle 0| = 0$, and such that ρ is positive semi-definite to find the mixed steady state.

We here elaborate on why the transition occurs when $\frac{\gamma_3}{\gamma_1} = O(\frac{1}{Z})$ for a lattice with uniform coordination number Z . As discussed in the main letter, it is helpful to consider the expectation value $\langle P^{(1,0),|1,-1)} \rangle$, where




FIG. 3. Decay of gap for increasing system size. Here, $\gamma_3 = 1$ and $\gamma_1 = \gamma_2 = \gamma$. In comparing the log-linear plot (left) and log-log plot (right) a transition from exponential to power law decay appears near $\gamma = 0.1$.




FIG. 4. Decay of correlations for eigen-density matrix corresponding to the first non-zero eigenvalue.

$$P^{(|1,0\rangle,|1,-1\rangle)} = \frac{1}{N-1} \sum_{\langle a,b \rangle} P_{a,b}^{(|1,0\rangle,|1,-1\rangle)} \quad (\text{F1})$$

$$\frac{1}{N-1} \sum_{\langle a,b \rangle} (|1,0\rangle_{a,b} \langle 1,0|_{a,b} + |1,-1\rangle_{a,b} \langle 1,-1|_{a,b}) \quad (\text{F2})$$

$$= \frac{1}{N-1} \sum_{\langle a,b \rangle} \frac{1}{2} (P_a^0 + P_b^0 + \sigma_a^+ \sigma_b^- + \sigma_a^- \sigma_b^+) \quad (\text{F3})$$

where $P_a^0 = (\frac{1+Z_a}{2})$ is the projector of site a onto the state \downarrow . We similarly define $P^{(|S\rangle)} = \frac{1}{N-1} \sum_{\langle a,b \rangle} P_{a,b}^{(|S\rangle)}$ with $P_{a,b}^{(|S\rangle)} = |S\rangle_{a,b} \langle S|_{a,b}$ for any two-site state $|S\rangle \in \{|0,0\rangle, |1,-1\rangle, |1,0\rangle, |1,1\rangle\}$.

The evolution of $\langle P^{(|1,0\rangle,|1,-1\rangle)} \rangle$ is given by

$$\frac{d}{dt}\langle P^{1,0},|1,-1\rangle\rangle = \text{Tr} \left\{ P^{1,0},|1,-1\rangle \frac{d\rho}{dt} \right\} \quad (\text{F4})$$

$$= \frac{1}{N-1} \sum_{\langle a,b \rangle} \sum_{\langle c,d \rangle} \gamma_j \text{Tr} \left\{ \rho \left[L_{j;a,b}^\dagger P_{c,d}^{1,0},|1,-1\rangle L_{j;a,b} - \frac{1}{2} \left(L_{j;a,b}^\dagger L_{j;a,b} P_{c,d}^{1,0},|1,-1\rangle + P_{c,d}^{1,0},|1,-1\rangle L_{j;a,b}^\dagger L_{j;a,b} \right) \right] \right\} \quad (\text{F5})$$

Let us first discuss the case where $a = c$ and $b = d$. For cohering terms like L_1 , we have that

$$L_{1;a,b}^\dagger P_{a,b}^{1,0},|1,-1\rangle L_{1;a,b} - \frac{1}{2} \left(L_{1;a,b}^\dagger L_{1;a,b} P_{a,b}^{1,0},|1,-1\rangle + P_{a,b}^{1,0},|1,-1\rangle L_{1;a,b}^\dagger L_{1;a,b} \right) \quad (\text{F6})$$

$$= |0,0\rangle\langle 1,0| P^{1,0},|1,-1\rangle |1,0\rangle\langle 0,0| - \frac{1}{2} \left(|0,0\rangle\langle 0,0| P^{1,0},|1,-1\rangle + P^{1,0},|1,-1\rangle |0,0\rangle\langle 0,0| \right) \quad (\text{F7})$$

$$= |0,0\rangle\langle 0,0| \quad (\text{F8})$$

where in the second line we dropped the a, b subscripts for brevity. Thus, the contribution from the L_1 on a, b and $P_{a,b}^{1,0},|1,-1\rangle$ term in the sum (F5) is

$$\gamma_1 \text{Tr} \{ \rho |0,0\rangle\langle 0,0| \} \quad (\text{F9})$$

$$= \gamma_1 \langle P_{a,b}^{0,0} \rangle. \quad (\text{F10})$$

Similarly, the contribution from L_2 on a, b is $\gamma_2 \langle P_{a,b}^{1,1} \rangle$. For the decohering terms (for example, taking first $L_{3'}$ and again dropping a, b subscripts for brevity), however, we have that

$$L_{3'}^\dagger P^{1,0},|1,-1\rangle L_{3'} - \frac{1}{2} \left(L_{3'}^\dagger L_{3'} P^{1,0},|1,-1\rangle + P^{1,0},|1,-1\rangle L_{3'}^\dagger L_{3'} \right) \quad (\text{F11})$$

$$= |0,0\rangle\langle 1,1| P^{1,0},|1,-1\rangle |1,1\rangle\langle 0,0| - \frac{1}{2} \left(|0,0\rangle\langle 0,0| P^{1,0},|1,-1\rangle + P^{1,0},|1,-1\rangle |0,0\rangle\langle 0,0| \right) \quad (\text{F12})$$

$$= 0. \quad (\text{F13})$$

The contribution from $L_{3'}$ is similarly 0. We therefore have that the contribution to (F5) from the $a = b, c = d$ terms (after summing over a, b), denoted by $\frac{d}{dt}\langle P^{1,0},|1,-1\rangle\rangle|_{a=b,c=d}$, is given by

$$\frac{d}{dt}\langle P^{1,0},|1,-1\rangle\rangle|_{a=b,c=d} = \gamma_1 \langle P^{0,0} \rangle + \gamma_2 \langle P^{1,1} \rangle \quad (\text{F14})$$

We now consider when it is not the case that $a = c$ and $b = d$. If $a \neq c, d$ and $b \neq c, d$, then $[L_{\alpha;a,b}, P_{c,d}^{1,0},|1,-1\rangle] = 0$, implying these terms in the sum (F5) are 0. Now consider terms where either $a = c$ or $b = d$. Specifically, for a jump operator acting on the sites a, b , we must consider the projectors $\sum_{s.t. \langle a,d \rangle} P_{a,d}^{1,0},|1,-1\rangle + \sum_{s.t. \langle b,c \rangle} P_{b,c}^{1,0},|1,-1\rangle$. From now on we will write \sum_c or \sum_d for brevity. Before we calculate the contribution to the sum (F5) from these terms, it is helpful to establish a few identities. Namely,

$$\begin{aligned} & \left\{ |0,0\rangle_{a,b} \langle 0,0|_{a,b}, \left(\sum_d P_{a,d}^{1,0},|1,-1\rangle + \sum_c P_{b,c}^{1,0},|1,-1\rangle \right) \right\} \\ &= \left(Z - 1 + \sum_c P_c^0 + \sum_d P_d^0 \right) |0,0\rangle_{a,b} \langle 0,0|_{a,b} \\ &+ \frac{1}{2\sqrt{2}} \left[|1,0\rangle_{a,b} \langle 0,0|_{a,b} \left(\sum_c \sigma_c^- - \sum_d \sigma_d^- \right) - |1,0\rangle_{a,b} \langle 1,1|_{a,b} \left(\sum_c \sigma_c^+ - \sum_d \sigma_d^+ \right) + h.c. \right] \end{aligned} \quad (\text{F15a})$$

$$\begin{aligned} & \left\{ |1, 1\rangle_{a,b} \langle 1, 1|_{a,b} \left(\sum_d P_{a,d}^{[1,0],[1,-1]} + \sum_c P_{b,c}^{[1,0],[1,-1]} \right) \right\} \\ &= \left(\sum_c P_c^0 + \sum_d P_d^0 \right) |1, 1\rangle_{a,b} \langle 1, 1|_{a,b} + \frac{1}{2} \left[|1, 1\rangle_{a,b} \left(\langle 10|_{a,b} \sum_c \sigma_c^- + \langle 01|_{a,b} \sum_d \sigma_d^- \right) + h.c. \right] \end{aligned} \quad (\text{F15b})$$

$$\langle 1, 0|_{a,b} \left(\sum_d P_{a,d}^{[1,0],[1,-1]} + \sum_c P_{b,c}^{[1,0],[1,-1]} \right) |1, 0\rangle_{a,b} = \frac{1}{2} \left(Z - 1 + \sum_c P_c^0 + \sum_d P_d^0 \right), \quad (\text{F15c})$$

$$\langle 0, 0|_{a,b} \left(\sum_d P_{a,d}^{[1,0],[1,-1]} + \sum_c P_{b,c}^{[1,0],[1,-1]} \right) |0, 0\rangle_{a,b} = \frac{1}{2} \left(Z - 1 + \sum_c P_c^0 + \sum_d P_d^0 \right), \quad (\text{F15d})$$

$$\langle 1, 1|_{a,b} \left(\sum_d P_{a,d}^{[1,0],[1,-1]} + \sum_c P_{b,c}^{[1,0],[1,-1]} \right) |1, 1\rangle_{a,b} = \frac{1}{2} \left(\sum_c P_c^0 + \sum_d P_d^0 \right), \quad (\text{F15e})$$

where in the above relations we have used that $P_{a,b}^{[1,0],[1,-1]} = \frac{1}{2} (P_a^0 + P_b^0 + \sigma_a^+ \sigma_b^- + \sigma_a^- \sigma_b^+)$.

Using (F15), we find the following for L_1 , L_2 , L_3 , and $L_{3'}$:

L	$L_{a,b}^\dagger \left(\sum_d P_{a,d}^{[1,0],[1,-1]} + \sum_c P_{b,c}^{[1,0],[1,-1]} \right) L_{a,b} - \frac{1}{2} \left\{ L_{a,b}^\dagger L_{a,b} \sum_d P_{a,d}^{[1,0],[1,-1]} + \sum_c P_{b,c}^{[1,0],[1,-1]} \right\}$
L_1	$-\frac{1}{4\sqrt{2}} \left[1, 0\rangle_{a,b} \langle 0, 0 _{a,b} (\sum_c \sigma_c^- - \sum_d \sigma_d^-) - 1, 0\rangle_{a,b} \langle 1, 1 _{a,b} (\sum_c \sigma_c^+ - \sum_d \sigma_d^+) + h.c. \right]$
L_2	$\frac{Z-1}{2} 1, 1\rangle_{a,b} \langle 1, 1 _{a,b} - \frac{1}{4} \left[1, 1\rangle_{a,b} \left(\langle 10 _{a,b} \sum_c \sigma_c^- + \langle 01 _{a,b} \sum_d \sigma_d^- \right) + h.c. \right]$
L_3	$\frac{Z-1}{2} 1, 1\rangle_{a,b} \langle 1, 1 _{a,b} - \frac{1}{4} \left[1, 1\rangle_{a,b} \left(\langle 10 _{a,b} \sum_c \sigma_c^- + \langle 01 _{a,b} \sum_d \sigma_d^- \right) + h.c. \right]$
$L_{3'}$	$-\frac{Z-1}{2} 0, 0\rangle_{a,b} \langle 0, 0 _{a,b} - \frac{1}{4\sqrt{2}} \left[1, 0\rangle_{a,b} \langle 0, 0 _{a,b} (\sum_c \sigma_c^- - \sum_d \sigma_d^-) - 1, 0\rangle_{a,b} \langle 1, 1 _{a,b} (\sum_c \sigma_c^+ - \sum_d \sigma_d^+) + h.c. \right]$

In the above table, the term $\sum_c \sigma_c^- - \sum_d \sigma_d^-$ and its hermitian conjugate appear several times. In the all-to-all limit, we have that $\sum_c \sigma_c^- - \sum_d \sigma_d^- = 0$ as the neighbors of site a and site b are the same (i.e. all sites). We now explain why these terms are also negligible for the NN model in the thermodynamic limit. Let us first look at the term $|1, 0\rangle_{a,b} \langle 0, 0|_{a,b} (\sum_c \sigma_c^- - \sum_d \sigma_d^-)$. This term's contribution in equation (F5) is $\frac{1}{N-1} \sum_{\langle a,b \rangle} \text{Tr} \left\{ \langle 0, 0|_{a,b} \rho |1, 0\rangle_{a,b} (\sum_c \sigma_c^- - \sum_d \sigma_d^-) \right\}$. For a general, random state, this is 0 as the state will on average have no asymmetry between the neighbors d of a and the neighbors c of b . Specifically, suppose $\text{Tr} \left\{ \langle 0, 0|_{a,b} \rho |1, 0\rangle_{a,b} \sum_c \sigma_c^- \right\} = \xi_{abc}$ where ξ_{abc} is some normal distribution with mean μ_{abc} and variance $\text{Var}(\xi_{abc})$ determined by the random state ρ (and similarly ξ_{abd} for the σ_d^- expectation value). We therefore have $\text{Tr} \left\{ \langle 0, 0|_{a,b} \rho |1, 0\rangle_{a,b} (\sum_c \sigma_c^- - \sum_d \sigma_d^-) \right\} = \xi_{abc} - \xi_{abd} \equiv \xi_{ab}$ where ξ_{ab} is a normal distribution with $\mu_{ab} = \mu_{abc} - \mu_{abd}$ and $\text{Var}(\xi_{ab}) = \text{Var}(\xi_{abc}) + \text{Var}(\xi_{abd})$. If ξ_{abc} and ξ_{abd} are i.i.d. we have that $\mu_{ab} = 0$ and $\text{Var}(\xi_{ab}) = 2\text{Var}(\xi_{abc})$. Summing over a, b we get $\frac{1}{N-1} \sum_{\langle a,b \rangle} \text{Tr} \left\{ \langle 0, 0|_{a,b} \rho |1, 0\rangle_{a,b} (\sum_c \sigma_c^- - \sum_d \sigma_d^-) \right\} = \frac{1}{N-1} \sum_{\langle a,b \rangle} \xi_{ab} \equiv \xi$ where (again assuming i.i.d. of ξ_{ab}) ξ has mean $\mu = 0$ and variance $\text{Var}(\xi) = \frac{2}{N-1} \text{Var}(\xi_{abc})$. Thus, ξ is precisely 0 in the thermodynamic limit. Therefore, we will treat the terms $\sum_c \sigma_c^- - \sum_d \sigma_d^-$ as negligible.

Now equipped with the solution to $L_{a,b}^\dagger P_{c,d}^{[1,0],[1,-1]} L_{a,b} - \frac{1}{2} \left(L_{a,b}^\dagger L_{a,b} P_{c,d}^{[1,0],[1,-1]} + P_{c,d}^{[1,0],[1,-1]} L_{a,b}^\dagger L_{a,b} \right)$ for all possible values of a, b, c , and d , we find that (F5) becomes

$$\frac{d}{dt} \langle P^{[1,0],[1,-1]} \rangle = \gamma_1 \langle P^{[0,0]} \rangle + \gamma_2 \langle P^{[1,1]} \rangle - \gamma_{3'} \frac{Z-1}{2} \langle P^{[0,0]} \rangle + (\gamma_2 + \gamma_3) \frac{Z-1}{2} \langle P^{[1,1]} \rangle - \frac{\gamma_2 + \gamma_3}{2} \langle T \rangle \quad (\text{F16})$$

where $T = \frac{1}{N-1} \sum_{\langle a,b \rangle} \frac{1}{4} \left[|1, 1\rangle_{a,b} \left(\langle 10|_{a,b} \sum_c \sigma_c^- + \langle 01|_{a,b} \sum_d \sigma_d^- \right) + h.c. \right]$.

The terms $\gamma_1 \langle P^{[0,0]} \rangle + \gamma_2 \langle P^{[1,1]} \rangle$ come from the cohering quantum jumps acting in cases $a = c, b = d$ and correspond to an increase in the coherence of the system. The term $(\gamma_2 + \gamma_3) \frac{Z-1}{2} \langle P^{[1,1]} \rangle$ corresponds to the γ_2 and γ_3 terms lowering total S_z and thereby increasing the number of pairs in the $|1, -1\rangle$ state. The rest of the terms correspond to cases where a coherent state on any two sites a, b is destroyed by jump operators which act between either a or b and a neighboring site.

For an initial density matrix in the single spin-up sector, equation (F16) becomes (6) in the main text.