## <span id="page-0-3"></span>LOW-REGULARITY GLOBAL SOLUTION OF THE INHOMOGENEOUS NONLINEAR SCHRÖDINGER EQUATIONS IN MODULATION SPACES

## DIVYANG G. BHIMANI, DIKSHA DHINGRA<sup>[1](#page-0-0)</sup>, VIJAY KUMAR SOHANI

Abstract. The study of low regularity Cauchy data for nonlinear dispersive PDEs has successfully been achieved using modulation spaces  $M^{p,q}$  in recent years. In this paper, we study the inhomogeneous nonlinear Schrödinger equation (INLS)

$$
iu_t + \Delta u \pm |x|^{-b} |u|^\alpha u = 0
$$
 (b,  $\alpha > 0$ )

on whole space  $\mathbb{R}^n$  in modulation spaces. In the subcritical regime  $(0 < \alpha < \frac{4-2b}{n})$ , we establish local well-posedness in  $L^2 + M^{\alpha+2, \frac{\alpha+2}{\alpha+1}} (\supset L^2 + H^s$  for  $s > \frac{n\alpha}{2(\alpha+2)}$ . By adapting Bourgain's high-low decomposition method, we establish global well-posedness in  $M^{p, \frac{p}{p-1}}$  with  $2 < p$  and p sufficiently close to 2. This is the first global well-posedness result for INLS on modulation spaces, which contains certain Sobolev  $H^s$  ( $0 < s < 1$ ) and  $L_s^p$ –Sobolev spaces.

#### 1. INTRODUCTION

1.1. **Background.** We study the inhomogeneous nonlinear Schrödinger equation (INLS for short) of the following form

<span id="page-0-1"></span>
$$
\begin{cases} iu_t(x,t) + \Delta u(x,t) + \mu |x|^{-b} (|u|^\alpha u)(x,t) = 0 \\ u(x,0) = u_0(x) \end{cases} (x,t) \in \mathbb{R}^n \times \mathbb{R}, \quad (1.1)
$$

where  $u(x, t) \in \mathbb{C}, \mu = \pm 1, \alpha, b > 0$  and  $\Delta$  is a Laplacian operator. In this paper, we assume that  $0 < b < \min\{2, n\}$  unless it is explicitly specified. The parameters  $\mu = 1$ (resp.  $\mu = -1$ ) corresponds to the focusing (resp. defocusing) case.

The classical nonlinear Schrödinger equation (NLS for short,  $b = 0$  case) has been extensively studied over the past three decades. See [\[17](#page-19-0), [41,](#page-20-0) [48](#page-20-1), [52\]](#page-20-2). This appeared in nonlinear optics [\[29,](#page-19-1) [42](#page-20-3)] and Bose-Einstein condensate (BEC) [\[20](#page-19-2)]. The NLS governing beam propagation does not support stable high power propagation in a homogeneous bulk medium. In [\[29](#page-19-1), [34,](#page-20-4) [42](#page-20-3)], it was proposed that stable high power propagation could be achieved in plasma by sending a preliminary laser beam to create a channel with reduced electron density. This channel mitigates the nonlinear effects within it, thereby allowing for stable propagation of the high-power beam. Considering these conditions, the beam propagation can be modeled by the inhomogeneous NLS of the form

<span id="page-0-2"></span>
$$
iu_t + \Delta u + K(x)|u|^\alpha u = 0.
$$
\n(1.2)

Here u is the electric field in optics,  $\alpha > 0$  is the power of nonlinear interaction, and the potential  $K(x)$  is proportional to the electron density. INLS  $(1.1)$  plays an important role as a limiting equation in the analysis of  $(1.2)$  with  $K(x) \sim |x|^{-b}$  as  $|x| \to \infty$ ; see [\[28](#page-19-3)]. INLS [\(1.1\)](#page-0-1) has also sparked significant theoretical interest on nonlinear phenomena in BEC with spatially inhomogeneous interactions; see [\[5\]](#page-18-0). Due to its vast applications,

<sup>1</sup>Corresponding author: Diksha Dhingra.

<sup>1</sup>Mathematics Subject Classification (2020): Primary 35Q55, 35Q60, 42B37; Secondary 35A01.

<span id="page-0-0"></span>Keywords: inhomogeneous nonlinear Schrödinger equation, global well-posedness, Bourgain's high-low decomposition method, modulation spaces.

<span id="page-1-1"></span>in recent years, the theory of well-posedness, scattering, blow-up analysis, etc. for INLS [\(1.1\)](#page-0-1) has been studied extensively. See e.g. [\[1,](#page-18-1) [2,](#page-18-2) [4,](#page-18-3) [14](#page-19-4)[–16](#page-19-5), [21](#page-19-6)[–23,](#page-19-7) [26](#page-19-8)[–28,](#page-19-3) [31,](#page-19-9) [32,](#page-19-10) [43,](#page-20-5) [44](#page-20-6)].

Formally, the solution to INLS [\(1.1\)](#page-0-1) conserves both mass and energy:

$$
M[u(t)] = \int_{\mathbb{R}^n} |u(x,t)|^2 dx = M[u_0],
$$
\n
$$
E[u(t)] = \int_{\mathbb{R}^n} |\nabla u(x,t)|^2 dx - \frac{\mu}{\alpha + 2} |||x|^{-b} |u|^{\alpha + 2}||_{L^1_x} = E[u_0].
$$
\n(1.3)

If  $u = u(x, t)$  is a solution of INLS [\(1.1\)](#page-0-1), then so is  $u_\lambda(x, t) = \lambda^{\frac{2-b}{\alpha}} u(\lambda x, \lambda^2 t)$  (scal-ing) also a solution of INLS [\(1.1\)](#page-0-1) with initial data  $u_{\lambda}(x,0) = \lambda^{\frac{2-b}{\alpha}} u(\lambda x,0)$ . Computing homogeneous Sobolev norm we have

<span id="page-1-0"></span>
$$
||u_{\lambda}(\cdot,0)||_{\dot{H}^{s}} = \lambda^{s-\frac{n}{2}+\frac{2-b}{\alpha}}||u_{0}||_{\dot{H}^{s}}.
$$

The  $\dot{H}^s$ -norm is invariant under the above scaling for  $s = s_b = \frac{n}{2} - \frac{2-b}{\alpha}$  (called critical Sobolev index). We say INLS  $(1.1)$  is

$$
L^{2}(\dot{H}^{1}) - \begin{cases} \text{subcritical} & \text{if } s_{b} < 0 \ (s_{b} = 1) \\ \text{critical} & \text{if } s_{b} = 0 \ (s_{b} = 1) \\ \text{supercritical} & \text{if } s_{b} > 0 \ (s_{b} > 1). \end{cases}
$$

We have mass critical or  $L^2$ -critical case if  $s_b = 0$  ( $\alpha = \frac{4-2b}{n}$ ). If  $s_b = 1$  ( $\alpha = \frac{4-2b}{n-2}$  $\frac{4-2b}{n-2},$ we have energy critical or  $\dot{H}^1$ – critical case. Finally, the problem is known to be masssupercritical and energy-subcritical or intercritical if  $0 < s_b < 1$ . For  $b = 0$ , we write  $s_0 = s_c$ . To state some known results, we set notations:

$$
\alpha_s = \begin{cases} \frac{4-2b}{n-2s} & \text{if } 0 \le s < \frac{n}{2} \\ +\infty & \text{if } s \ge \frac{n}{2} \end{cases}, \quad \alpha_s^* = \begin{cases} \frac{4-2b}{n-2s} & \text{if } s < \frac{n}{2} \\ +\infty & \text{if } s = \frac{n}{2} \end{cases},
$$
  

$$
\tilde{2} = \begin{cases} \frac{n}{3} & \text{if } n = 1, 2, 3 \\ 2 & \text{if } n \ge 4 \end{cases} \text{ and } \quad \hat{2} = \begin{cases} \min\{2, 1 + \frac{n-2s}{2}\} & \text{if } n \ge 3 \\ n-s & \text{if } n = 1, 2. \end{cases}
$$

Now, we briefly summarize some known results of INLS [\(1.1\)](#page-0-1):

- Genoud and Stuart [\[28,](#page-19-3) Theorem 1.1] proved:
	- locally well-posed (LWP for short) in  $H^1$  for  $0 < \alpha < \alpha_1, \mu = -1$ .
	- globally well-posed (GWP for short) for small data in  $H^1$  when  $\alpha_0 \leq \alpha$  $\alpha_1, \mu = -1.$
	- GWP for any data in  $H^1$  for  $0 < \alpha < \alpha_0, \mu = -1$ .
	- GWP in  $H^1$  for  $\mu = 1$  (see [\[17](#page-19-0)]).
- $-$  Guzmán [\[31,](#page-19-9) Theorems 1.4, 1.8 and 1.9] proved:
	- LWP in  $H^s$  for  $\max\{0, s_b\} < s \le \min\left\{1, \frac{n}{2}\right\}$ ,  $0 < b < \tilde{2}$ , and  $0 < \alpha < \alpha_s^*$ .

 $\frac{2}{3}, \frac{2}{3}$ – GWP for small data in  $H^s$  for  $s_b < s \leq \min\left\{1, \frac{n}{2}\right\}$  $\left\{\frac{a}{2}\right\},\,0 < b < \tilde{2},\,\frac{4-2b}{n} < \alpha < \alpha_s^*$ .

- GWP in  $L^2$  for  $0 < \alpha < \frac{4-2b}{n}$ . (For  $b = 0$ , see [\[49](#page-20-7), Theorem 1.1.])
- JinMyoung An et al. [\[2](#page-18-2), Theorems 1.4 and 1.6], [\[1,](#page-18-1) Theorem 1.3] expanded the range for s and  $0 < b < 2$ . Specifically, INLS [\(1.1\)](#page-0-1) is:
	- LWP in  $H^s$   $(0 \le s < \min\{n, \frac{n}{2} + 1\})$  for  $0 < \alpha < \alpha_s$ .
	- GWP for small data in  $H^s$   $(0 < s < \min\{n, \frac{n}{2} + 1\})$  for  $\alpha_0 < \alpha < \alpha_s$ .
- Genoud [\[27](#page-19-11), Theorem 1] proved focusing mass-critical INLS  $(1.1)$  is GWP in  $H<sup>1</sup>$ having data below ground state. This result was extended to the intercritical case in [\[24,](#page-19-12) Theorem 1.5]. See [\[21](#page-19-6)] and [\[22\]](#page-19-13). We also refer the reader to [\[14\]](#page-19-4) for

<span id="page-2-4"></span>the global behaviour of solutions to 3D focusing INLS [\(1.1\)](#page-0-1) at the mass-energy threshold.

- Murphy [\[44\]](#page-20-6) studied scattering below the ground state for the intercritical nonradial case in  $H^1$ . See [\[16,](#page-19-5) [23](#page-19-7), [32\]](#page-19-10) for more results on scattering. On the other hand, for blow-up analysis we refer (among others) to papers by Merle [\[43\]](#page-20-5) and Banica-Carles- Duyckaerts [\[4](#page-18-3)] and Cardoso-Farah [\[15\]](#page-19-14).

Finally, taking these known results into account, in the next remark we would like to highlight the following points:

- <span id="page-2-0"></span>**Remark 1.1.** (1) The problem of GWP for large data in  $H^s$  with  $0 < s < 1$  remains unsolved.
	- (2) Most authors studied INLS  $(1.1)$  in the L<sup>2</sup>-based Sobolev spaces. It is natural to investigate similar theory in  $L_s^p$ –Sobolev spaces.

We shall see soon that modulation spaces will provide us an appropriate framework for the concern raised in Remark [1.1.](#page-2-0)

1.2. **Modulation spaces.** In the past two decades,  $M_s^{p,q}$ -spaces have been extensively explored as spaces for low regularity Cauchy data in the study of dispersive PDEs. See [\[3](#page-18-4), [6](#page-18-5)[–12,](#page-19-15) [18,](#page-19-16) [19](#page-19-17), [47](#page-20-8), [51](#page-20-9), [52\]](#page-20-2).

Let  $\rho : \mathbb{R}^n \to [0,1]$  be a smooth function satisfying  $\rho(\xi) = 1$  if  $|\xi|_{\infty}^2 \leq \frac{1}{2}$  $|\xi|_{\infty}^2 \leq \frac{1}{2}$  $|\xi|_{\infty}^2 \leq \frac{1}{2}$  $rac{1}{2}$  and  $\rho(\xi) = 0$  if  $|\xi|_{\infty} \geq 1$ . Let  $\rho_k$  be a translation of  $\rho$ , that is  $\rho_k(\xi) = \rho(\xi - k)$   $(k \in \mathbb{Z}^n)$ . Denote

$$
\sigma_k(\xi) = \frac{\rho_k(\xi)}{\sum_{l \in \mathbb{Z}^n} \rho_l(\xi)} \quad (k \in \mathbb{Z}^n).
$$

The frequency-uniform decomposition operators can be defined by

$$
\Box_k = \mathcal{F}^{-1} \sigma_k \mathcal{F} \quad (k \in \mathbb{Z}^n)
$$

where  $\mathcal F$  and  $\mathcal F^{-1}$  denote the Fourier and inverse Fourier transform respectively. The weighted modulation spaces  $M_s^{p,q}$   $(1 \leq p, q \leq \infty, s \in \mathbb{R})$  is defined as follows:

$$
M_s^{p,q} = M_s^{p,q}(\mathbb{R}^n) = \left\{ f \in \mathcal{S}'(\mathbb{R}^n) : ||f||_{M_s^{p,q}} = ||\|\Box_k f\|_{L_x^p} (1+|k|)^s\|_{\ell_k^q} < \infty \right\}.
$$

For  $s = 0$ , we write  $M_0^{p,q} = M^{p,q}$ . See Remark [1.8.](#page-5-0) For  $p = q = 2$ , modulation spaces coincide with Sobolev spaces, i.e.  $M_s^{2,2} = H^s$  ( $s \in \mathbb{R}$ ). For  $p \in [1,\infty]$ , we denote  $p'$  the Hölder conjugate, i.e.  $\frac{1}{p} + \frac{1}{p'}$  $\frac{1}{p'} = 1$ . By Lemma [2.1](#page-6-0) [\(1\)](#page-6-1) and [\(2\)](#page-6-2), we have

<span id="page-2-2"></span>
$$
H^s \subset M^{\alpha+2, (\alpha+2)'}
$$
 for  $s > \frac{n\alpha}{2(\alpha+2)}$ . (1.4)

Recall  $L_s^p$ -Sobolev norm:  $||f||_{L_s^p} = ||\mathcal{F}^{-1}\langle \cdot \rangle^s \mathcal{F}f||_{L^p}$ . By Proposition [2.2,](#page-6-3) for  $s > n \left( \frac{1}{p'}\right)$  $\frac{1}{p'}-\frac{1}{p}$ p  $\setminus$ and  $p \in [2, \infty]$ , we have

<span id="page-2-3"></span>
$$
L_s^p \hookrightarrow M^{p,p'} \hookrightarrow L^p. \tag{1.5}
$$

In fact, modulation spaces accommodate rougher Cauchy data compared to any fractional Bessel potential space. See e.g  $[7,$  Chapter 6,  $[52,$  Chapter 6. The free Schrödinger propagator  $e^{-it\Delta}$ :  $M^{p,q} \to M^{p,q}$  is a bounded operator for all p, q, see Proposition [2.3,](#page-6-4) while it is unbounded in  $L_s^p$  Sobolev spaces for  $p \neq 2$ . Another great advantage is that  $e^{-it\Delta}$  enjoys a truncated decay in modulation spaces, but its decay from  $L^{p'}$  to  $L^p$ contains singularity at  $t = 0$ . See Proposition [2.3.](#page-6-4) Using these facts, Wang and Hudzik in their seminal work in [\[51\]](#page-20-9) proved NLS  $(b = 0, \alpha \in 2\mathbb{N})$  is GWP for small data in  $M^{2,1}$ . This enables us to consider the lower-regularity initial data in modulation spaces for a

<span id="page-2-1"></span> ${}^{2}$ Define  $|\xi|_{\infty} = \max\{|\xi_{i}| : \xi = (\xi_{1}, ..., \xi_{n})\}.$ 

<span id="page-3-2"></span>class of data out of the critical Sobolev spaces  $H^{s_c}$ . See Proposition [2.2.](#page-6-3) Later taking algebra property (Lemma [2.1](#page-6-0) [\(4\)](#page-6-5)) into account, in  $[6, 12]$  $[6, 12]$ , it is proved that NLS is LWP in  $M_s^{p,1}$   $(1 \le p \le \infty, s \ge 0)$  and in  $M_s^{p,q}$   $(1 \le p, q \le \infty, s > \frac{n}{q'})$  via fixed point argument. Guo [\[30](#page-19-18)] proved 1D cubic NLS is LWP in  $M^{2,p}$  ( $2 \le p < \infty$ ) and later Oh-Wang [\[45](#page-20-10)] established global existence for this result. Chaichenets et al. in [\[18](#page-19-16), [19\]](#page-19-17) established global well-posesness for cubic NLS in  $M^{p,p'}$  for p sufficiently close to 2.

On the other hand, Bhimani et al. in [\[9](#page-18-7), [10](#page-18-8)] established GWP in  $M^{p,q} \cap L^2$  for Hartree equation (and also for Hartree-Fock eq.), i.e. NLS with nonlinearity  $H_b(u) = (|\cdot|^{-b} * |u|^2)u$ . The key step in their work is to get the trilinear estimate for  $H_b(u)$  and this was possible due to the regularizing effect of convolution. However, it seems their method does not apply to treat nonlinearity  $|\cdot|^{-b}|u|^\alpha u$  (due to just multiplication by singular potential). We also mention that, for  $M_s^{p,q}$  with some negative regularity  $(s < 0)$ , strong ill-posedness theory (norm inflation with infinite loss of regularity) is developed for NLS and Hartree equations in [\[8](#page-18-9), [11](#page-18-10)].

In spite of these progresses and ongoing interest for adapting modulation spaces in dispersive PDEs, there are no results for INLS [\(1.1\)](#page-0-1) so far. In view of this and the aforementioned discussion of the previous section (see Remark [1.1\)](#page-2-0), we are inspire to study INLS [\(1.1\)](#page-0-1) in modulation spaces.

1.3. Main results. We are now ready to state LWP result in the following theorem. To this end, we denote

<span id="page-3-0"></span>
$$
\tilde{b} = \begin{cases}\n\frac{3-\sqrt{7}}{2} & \text{if } n = 1 \\
2-\sqrt{2} & \text{if } n = 2 \\
\frac{n+6-\sqrt{(n+6)^2-32}}{4} & \text{if } n \ge 3.\n\end{cases}
$$
\n(1.6)

<span id="page-3-1"></span>**Theorem 1.2** (Local well-posedness). Let  $0 < \alpha < \frac{4-2b}{n}$ ,  $\tilde{b}$  be as in [\(1.6\)](#page-3-0) and

$$
\begin{cases} 0 < b \leq \tilde{b} & \text{if } n \neq 2 \\ 0 < b < \tilde{b} & \text{if } n = 2. \end{cases}
$$

Assume that  $u_0 \in L^2 + M^{\alpha+2, (\alpha+2)^t}$ . Then there exists  $T^* = T^*(||u_0||_{L^2 + M^{\alpha+2, (\alpha+2)^t}}, n, \alpha) > 0$ and a unique maximal solution u of INLS [\(1.1\)](#page-0-1) such that

$$
u \in C([0,T^*), L^2) \cap L^{\frac{4(\alpha+2)}{n\alpha}}([0,T^*), L^{\alpha+2}) + C([0,T^*), M^{\alpha+2,(\alpha+2)'}).
$$

Moreover,

- (1) (Blow-up alternative) If  $T^* < \infty$ , then  $||u(\cdot, t)||_{L^2+M^{\alpha+2}, (\alpha+2)'} \to \infty$  as  $t \to T^*$ .
- (2) (Lipschitz continuity) The mapping  $u_0 \mapsto u(t)$  is locally Lipschitz from  $L^2$  +  $M^{\alpha+2,(\alpha+2)' }$  to  $C([0,T'],L^2) \cap L^{\frac{4(\alpha+2)}{n\alpha}}([0,T'],L^{\alpha+2})+C([0,T'],M^{\alpha+2,(\alpha+2)'})$  for  $T' <$  $T^*$ .

Up to now we cannot solve INLS [\(1.1\)](#page-0-1) in  $L^p$ -Sobolev spaces or in  $L^p$  but in  $M^{p,p'}$ for  $p \neq 2$ . See Remark [1.1,](#page-2-0) [\(1.4\)](#page-2-2) and [\(1.5\)](#page-2-3). Thus, Theorem [1.2](#page-3-1) complements the points mentioned in Remark [1.1.](#page-2-0)

Remark 1.3. We briefly mention key ideas to prove Theorem [1.2.](#page-3-1)

- By utilizing the boundedness of the operator  $e^{it\Delta}$  in modulation spaces (Proposition [2.3\)](#page-6-4), we treat the linear part.
- The main difficulty is to handle the spatially decaying factor  $|\cdot|^{-b}$  in the nonlinearity as  $|\cdot|^{-b}$  does not belong to any  $L^p$  spaces. We handle this by decomposing

<span id="page-4-2"></span> $\mathbb{R}^n = B \cup B^c$  into two parts. Specifically,

$$
\begin{cases} |\cdot|^{-b}\in L^\gamma(B) & if \ \frac{n}{\gamma}-b>0\\ |\cdot|^{-b}\in L^\gamma(B^c) & if \ \frac{n}{\gamma}-b<0. \end{cases}
$$

- This introduces additional complexity as we need to select different admissible pairs for the regions inside the ball i.e. in  $B$  and outside the ball i.e. in  $B<sup>c</sup>$ . The restriction on  $\alpha$  and b comes due to Lemma [3.1,](#page-8-0) see Remark [3.2.](#page-10-0)
- We invoke Strichartz estimates (Theorem [2.6\)](#page-7-0) to run the fixed point argument via Banach contraction principle. To this end, we used embedding  $M^{p,p'} \subset L^p$ , which also justifies the selection of the exponent  $p = \alpha + 2$ .
- We shall see that the life span of the solution  $T$  depends on the size of the initial data  $u_0$  in the  $L^2 + M^{\alpha+2, (\alpha+2)'}$  norm since the problem is "subcritical".

The local solution established in Theorem [1.2](#page-3-1) can be extended to a global one under certain restriction on exponent p. To this end, we denote

$$
p_{\max} := \begin{cases} \frac{4\alpha + 8 - n\alpha}{2\alpha + 2 + b + \frac{n(4 - 2b - n\alpha)}{2(\alpha + 2)(n + 2 - b)}} & \text{if } \alpha - \frac{n\alpha^2}{4(\alpha + 2)} - \frac{4 - 2b - n\alpha}{4} + \frac{n(4 - 2b - n\alpha)}{4(\alpha + 2)(n + 2 - b)} > 0\\ \alpha + 2 & \text{otherwise.} \end{cases}
$$

<span id="page-4-0"></span>We are now ready to state GWP result in the next theorem.

**Theorem 1.4** (Global well-posedness). Let  $0 < \alpha < \frac{4-2b}{n}$ ,  $\tilde{b}$  be as in [\(1.6\)](#page-3-0) and

$$
\begin{cases} 0 < b \leq \tilde{b} & if \ n \neq 2 \\ 0 < b < \tilde{b} & if \ n = 2. \end{cases}
$$

Assume that  $u_0 \in M^{p,p'}$  for  $p \in (2, p_{max})$ . Then INLS [\(1.1\)](#page-0-1) has a unique solution u satisfying

$$
u \in C(\mathbb{R}, L^2) \cap L_{loc}^{\frac{4(\alpha+2)}{n\alpha}}(\mathbb{R}, L^{\alpha+2}) + C(\mathbb{R}, M^{\alpha+2, (\alpha+2)'}).
$$

Theorem [1.4](#page-4-0) is the first GWP result for INLS [\(1.1\)](#page-0-1) in modulation spaces. We note that existing results does not address GWP in  $H^s$  for  $0 < s < 1$  in the mass-subcritical case without any restriction on the initial data being small. See Remark [1.1.](#page-2-0) Since  $H^s \subset M^{p,p'}$ for some  $s \in (0,1)$  (see [\(1.4\)](#page-2-2)), Theorem [1.4](#page-4-0) partially addresses this gap and complement known results for INLS  $(1.1)$ .

In [\[13](#page-19-19)], Bourgain introduced a general scheme, so called Bourgain's high-low decomposition method to establish GWP for NLS in  $H^s$  for  $s > \frac{3}{5}$ . The idea is to split initial data  $u_0$  between two suitable function spaces and solve in each of them a different NLS. And then combine the solutions to get a function that solves original problem. See Remark [1.5,](#page-4-1) [\[37](#page-20-11), Section 3.2] and [\[48](#page-20-1), Section 3.9] for more details. Later, Vargas and Vega [\[50](#page-20-12)] adapted this method to establish global existence for 1D cubic NLS even if  $||u_0||_{L^2} = \infty$ . See [\[33,](#page-20-13) [38\]](#page-20-14) for more related works. Recently, Chaichenets et al. in [\[19,](#page-19-17) Theorem 3] and [\[18](#page-19-16)] successfully adapted this method to establish GWP for NLS (i.e. INLS [\(1.1\)](#page-0-1) with  $b = 0$ ) in modulation spaces. Our method of the proof is inspired by these results.

<span id="page-4-1"></span>Remark 1.5. We briefly mention key ideas to prove Theorem [1.4.](#page-4-0)

- We split the initial data into two parts using Lemma [2.5:](#page-7-1)

$$
u_0 = \phi_0 + \psi_0 \in L^2 + M^{(\alpha+2), (\alpha+2)'}
$$

We say part in  $L^2$  having high frequency in the sense that  $\|\phi_0\|_{L^2} \lesssim_p N^{\beta}$  for some  $\beta > 0$  and part in  $M^{(\alpha+2),(\alpha+2)'}$  having low frequency in the sense that  $\|\psi_0\|_{M^{(\alpha+2), (\alpha+2)'} \lesssim_p N^{-1}$ . See [\(4.1\)](#page-12-0) and [\(4.2\)](#page-12-1).

.

- <span id="page-5-2"></span>- By Theorem [1.2,](#page-3-1) we get local existence in  $L^2 + M^{(\alpha+2),(\alpha+2)'}$ . Guzmán [\[31,](#page-19-9) Theorem 1.8] proved INLS  $(1.1)$  is GWP in  $L^2$ . This proof depends on the fact that the solution enjoys the conservation of mass (stated in  $(1.3)$ ) and  $(4.5)$ ). However, there is no known useful conservation law for the solution in terms of  $M^{p,q}$ -spaces norm that help us to extend the solution globally in time.
- In order to handle this situation, we construct a solution  $u$  of INLS  $(1.1)$  in the form

$$
u = (v_0 + w_0) + e^{it\Delta} \psi_0.
$$

Here  $v_0$  is the  $L^2$ - global solution of [\(1.1\)](#page-0-1) with data  $\phi_0$ , see [\(4.5\)](#page-12-2). While  $e^{it\Delta}\psi_0 \in$  $M^{(\alpha+2),(\alpha+2)'}$  for  $t \in \mathbb{R}$  is the linear evolution of  $\psi_0$ , see Proposition [2.3](#page-6-4) and  $w_0$  is the nonlinear interaction term; see  $(4.11)$ . See  $(4.10)$  for more details.

- Since  $\|\psi_0\|_{M^{(\alpha+2),(\alpha+2)'}}\lesssim_p N^{-1}$  can be made small, we get  $v_0 + w_0$  close to  $v_0$  in  $L^2$ . Consequently,  $M(v_0 + w_0)$ , although no longer conserved, grows slowly enough to yield a global solution. See Remark [4.4](#page-13-0) for more details on the proof of Theorem [1.4.](#page-4-0)

**Remark 1.6.** The restriction on p essentially comes due to our choice of suitable norm size of the high frequency part of initial data. We refer to Remark [4.5](#page-16-0) for details.

**Remark 1.7.** Theorem [1.4](#page-4-0) may be interpreted as follows: for data  $u_0 \in M^{p,p'}$ , the solution  $u(\cdot, t)$  lies in a larger modulation spaces

$$
(L^2 \cap L^{\alpha+2}) + M^{\alpha+2,(\alpha+2)'} \supset M^{p,p'}.
$$

We have this loss of regularity as we rely on decomposition of initial data and we do not know whether Schrödinger propogator  $e^{it\Delta}$  gain any regularity on  $M^{p,q}$  (although it is bounded). See [\(4.10\)](#page-12-4) and Proposition [2.3.](#page-6-4)

<span id="page-5-0"></span>Remark 1.8. In the early 1980s Feichtinger [\[25](#page-19-20)] introduced a class of Banach spaces, the so called modulation spaces, which allow a measurement of space variable and Fourier transform variable of a function or distribution on  $\mathbb{R}^n$  simultaneously using the shorttime Fourier transform (STFT). The STFT of a tempered distribution  $f \in \mathcal{S}'(\mathbb{R}^n)$  with respect to a window function  $0 \neq g \in \mathcal{S}(\mathbb{R}^n)$  (Schwartz space) is defined by

$$
V_g f(x,\omega) = \int_{\mathbb{R}^n} f(t) \overline{g(t-x)} e^{-2\pi i \omega \cdot t} dt,
$$

whenever the integral exists. It is known that

$$
||f||_{M_s^{p,q}} \asymp ||||V_g f(x,\omega)||_{L_x^p} (1+|\omega|^2)^{s/2}||_{L_\omega^q},
$$

see  $[25]$ ,  $[51$ , Proposition 2.1].

This paper is organised as follows. Preliminaries and notations are introduced in Section [2,](#page-5-1) which will be helpful in the sequel. In Section [3,](#page-7-2) we obtain the estimates of the nonlinear term and the proof of the Theorem [1.2](#page-3-1) is also discussed. In Section [4,](#page-12-5) the proof for Theorem [1.4](#page-4-0) is presented.

### 2. Notations and Preliminaries

<span id="page-5-1"></span>Notations. The symbol  $X \leq Y$  means  $X \leq CY$  for some constant  $C > 0$ . While  $X \approx Y$ means  $C^{-1}X \le Y \le CX$  for some constant  $C > 0$ . The symbol  $A \hookrightarrow B$  denotes the <span id="page-6-9"></span>continuous embedding of the topological linear space  $A$  into  $B$ . The norm of the spacetime Lebesgue spaces  $L^q([0,T], L^r(\mathbb{R}^n))$  is defined as

$$
||u||_{L_T^qL^r} := ||u||_{L^q([0,T],L^r(\mathbb{R}^n))} = \left(\int_0^T ||u(\cdot,t)||_{L^r(\mathbb{R}^n)}^q dt\right)^{\frac{1}{q}}.
$$

We simply write  $||u||_{L^qL^r}$  in place of  $||u||_{L^q(\mathbb{R},L^r(\mathbb{R}^n))}$ . The unit ball centred at 0 is denoted by  $B = B(0, 1)$  and  $B^c = \mathbb{R}^n \setminus B$ .

2.1. Basic Properties of  $M^{p,q}$ -spaces. Let us briefly recall some relevant facts concerning the modulation spaces.

<span id="page-6-2"></span><span id="page-6-1"></span><span id="page-6-0"></span>**Lemma 2.1** ([\[7](#page-18-6), [52\]](#page-20-2)). Let  $p, q, p_i, q_i \in [1, \infty], (i = 0, 1, 2)$  and  $s_1, s_2 \in \mathbb{R}$ . (1)  $M_{s_1}^{p_1,q_1} \hookrightarrow M_{s_2}^{p_2,q_2}$  whenever  $p_1 \leq p_2, q_1 \leq q_2, s_2 \leq s_1$ .

<span id="page-6-6"></span>(2)  $M_{s_1}^{p,q_1}$  →  $M_{s_2}^{p,q_2}$  whenever  $q_2 < q_1, s_1 - s_2 > \frac{n}{q_2}$  $\frac{n}{q_2} - \frac{n}{q_1}$  $\frac{n}{q_1}$ .

<span id="page-6-5"></span> $(3)$   $M^{p,q_1} \hookrightarrow L^p \hookrightarrow M^{p,q_2}$  for  $q_1 \leq \min\{p, p'\}$  and  $q_2 \geq \max\{p, p'\}.$ 

(4) If  $\frac{1}{p_1} + \frac{1}{p_2}$  $\frac{1}{p_2}=\frac{1}{p_0}$  $\frac{1}{p_0}$  and  $\frac{1}{q_1} + \frac{1}{q_2}$  $\frac{1}{q_2} = 1 + \frac{1}{q_0}, \text{ then } ||fg||_{M^{p_0,q_0}} \lesssim ||f||_{M^{p_1,q_1}} ||g||_{M^{p_2,q_2}}.$ 

<span id="page-6-3"></span>**Proposition 2.2** (examples). Let  $1 \leq p, q \leq \infty, s_1, s_2 \in \mathbb{R}$ .

- (1) The  $M_s^{p,q}$ -spaces are invariant under Fourier transform only when  $p = q$ . See for e.g. [\[46,](#page-20-15) p. 2084]. It follows that  $M^{p,p'} \neq H^s = M_s^{\tilde{2},2}$  for  $p \neq 2, s \in \mathbb{R}$ .
- <span id="page-6-7"></span>(2) For  $p > 2$  and  $s_b < 0$ ,  $H^{s_b} \not\subset M^{p,p'}$ .
- (3)  $M_{s_c}^{2,1}$  ⊂  $H^{s_c}$  and  $M_{s}^{2,1}$  ⊄  $H^{s_c}$  for  $s < s_c = s_0$ .
- (4) (See [\[39\]](#page-20-16)).Denote

$$
\tau(p,q) = \max\left\{0, n\left(\frac{1}{q} - \frac{1}{p}\right), n\left(\frac{1}{q} + \frac{1}{p} - 1\right)\right\}.
$$

Then  $L_{s_1}^p \subset M_{s_2}^{p,q}$  if and only if one of the following conditions is satisfied:

$$
(i) \ q \ge p > 1, s_1 \ge s_2 + \tau(p, q); \quad (ii) \ p > q, s_1 > s_2 + \tau(p, q);
$$

(iii)  $p = 1, q = \infty, s_1 > s_2 + \tau(1, \infty);$  (iv)  $p = 1, q \neq \infty, s_1 > s_2 + \tau(1, q).$ 

(5) For completely sharp embedding between  $M^{p,q}_{s_1}$  and  $H^s$ , see [\[40](#page-20-17)], [\[35,](#page-20-18) Theorems 1.3 and 1.4].

*Proof.* In fact, if  $H^{s_b} \subset M^{p,p'}$ , then by Lemma [2.1](#page-6-0)[\(3\)](#page-6-6), we have  $L^2 \subset L^p$ , which is a contradiction. For part  $(3)$ , see e.g. [\[47,](#page-20-8) Section 1], [\[3](#page-18-4), Section 3], [\[40,](#page-20-17) [52\]](#page-20-2).

<span id="page-6-4"></span>**Proposition 2.3** (See Proposition 4.1 in [\[51\]](#page-20-9), [\[7\]](#page-18-6)). Let  $p, q \in [1, \infty]$  and  $s \in \mathbb{R}$ . Denote  $the Schrödinger propagator by$ 

$$
e^{it\Delta} f(x) := \int_{\mathbb{R}^n} e^{i\pi t |\xi|^2} \hat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi \quad (f \in \mathcal{S}, t \in \mathbb{R}).
$$
  
(1) 
$$
\|e^{it\Delta} f\|_{M_s^{p,q}} \lesssim_{n,s} (1+t^2)^{-\frac{n}{2}(\frac{1}{2}-\frac{1}{p})} \|f\|_{M_s^{p',q}} \quad (2 \le p \le \infty).
$$
  
(2) 
$$
\|e^{it\Delta} f\|_{M_s^{p,q}} \lesssim_{n,s} (1+t^2)^{\frac{n}{2}|\frac{1}{2}-\frac{1}{p}|} \|f\|_{M_s^{p,q}}.
$$

<span id="page-6-8"></span>**Lemma 2.4** (see e.g. Lemma 3.9 in  $[18]$ ). Denote

$$
G(u, v, w) = |u + v|^{\alpha} (u + v) - |u + w|^{\alpha} (u + w)
$$

for  $\alpha > 0$  and  $u, v, w \in \mathbb{C}$ . Then

$$
|G(u,v,w)| \lesssim_{\alpha} (|u|^{\alpha} + |v|^{\alpha} + |w|^{\alpha})|v-w|.
$$

We simply write  $G(u, v, 0) = G(u, v)$ .

<span id="page-7-5"></span><span id="page-7-1"></span>**Lemma 2.5** (Interpolation; see Theorem 6.1 (D) in [\[25](#page-19-20)] and Proposition 5.1 in [\[18](#page-19-16)]). Let  $u \in M^{p,p'}, p \in (2,r)$  and  $N > 0$ . Then there exists  $v \in L^2$  and  $w \in M^{r,r'}$  such that

 $u = v + w$ 

with the following property:

$$
\begin{cases} ||v||_{L^2} \leq C ||u||_{M^{p,p'}} N^{\beta} \\ ||w||_{M^{r,r'}} \leq C ||u||_{M^{p,p'}} \frac{1}{N} \end{cases}, \text{ where } \beta = \frac{\frac{1}{2} - \frac{1}{p}}{\frac{1}{p} - \frac{1}{r}}.
$$

<span id="page-7-3"></span>**Definition 2.1.** We say that a pair  $(q, r)$  is *admissible* if

$$
\frac{1}{q} = \frac{n}{2} \left( \frac{1}{2} - \frac{1}{r} \right) \quad \text{where } \begin{cases} 2 \le r \le \frac{2n}{n-2} & \text{if } n \ge 3, \\ 2 \le r < \infty \\ 2 \le r \le \infty & \text{if } n = 1. \end{cases}
$$

The set of all admissible pairs is denoted by

 $\mathcal{A} = \{(q, r) : (q, r)$  is admissible pair}.

<span id="page-7-0"></span>**Theorem 2.6** (Strichartz estimates, [\[17,](#page-19-0) [36](#page-20-19), [41\]](#page-20-0)). Let  $f \in L^2$ ,  $(\tilde{q}, \tilde{r})$ ,  $(q, r) \in \mathcal{A}$  and  $g \in L^{\tilde{q}'}(\mathbb{R}, L^{\tilde{r}'})$ . Then

$$
||e^{-it\Delta}f||_{L^qL^r} \lesssim_{n,r} ||f||_{L^2}
$$

$$
\left\| \int_0^t e^{-i(t-s)\Delta}g(x,s)ds \right\|_{L^qL^r} \lesssim_{n,r,\tilde{r}} ||g||_{L^{\tilde{q}'}L^{\tilde{r}'}}.
$$

3. LOCAL WELL-POSEDNESS IN  $L^2 + M^{\alpha+2,(\alpha+2)'}$ 

<span id="page-7-2"></span>In this section, we shall prove Theorem [1.2.](#page-3-1) To this end, we shall briefly introduce some notations and prove Lemma [3.1.](#page-8-0)

Consider the Banach space  $X(T)$  expressed as

$$
X(T) := X_1(T) + X_2(T)
$$
\n(3.1)

where

$$
X_1(T) := C([0, T], L^2) \cap L^{\frac{4(\alpha+2)}{n\alpha}}([0, T], L^{\alpha+2})
$$

equipped with the norm

$$
||v||_{X_1(T)} = \max \left\{ ||v||_{L_T^{\infty}L^2}, ||v||_{L_T^{\frac{4(\alpha+2)}{n\alpha}}L^{\alpha+2}} \right\}
$$

and

$$
X_2(T) := C([0,T], M^{\alpha+2,(\alpha+2)'}).
$$

The norm on  $X(T)$  is given as

$$
||u||_{X(T)} = \inf_{\substack{u=v+w\\v\in X_1(T)\\w\in X_2(T)}} (||v||_{X_1(T)} + ||w||_{X_2(T)}).
$$

Denote

<span id="page-7-4"></span>
$$
Y(T) := L^{\frac{4(\alpha+2)}{n\alpha}}([0,T], L^{\alpha+2}).
$$
\n(3.2)

The following lemma provides an estimate for the nonlinearity  $|x|^{-b}|u|^\alpha u$  which will play a crucial role in our analysis.

<span id="page-8-0"></span>**Lemma 3.1.** Let  $0 < \alpha < \frac{4-2b}{n}$ ,  $\tilde{b}$  be as in [\(1.6\)](#page-3-0) and

$$
\begin{cases} 0 < b \leq \tilde{b} & \text{if } n \neq 2 \\ 0 < b < \tilde{b} & \text{if } n = 2. \end{cases}
$$

Then

$$
\inf_{(\gamma,\rho)\in\mathcal{A}}\|\ |x|^{-b}|u|^\alpha v\|_{L_T^{\gamma'}L^{\rho'}}\lesssim (T^{1-\frac{n\alpha}{4}}+T^{\frac{4-2b-n\alpha}{4}-\frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}})\|u\|_{Y(T)}^\alpha\|v\|_{Y(T)}.
$$

*Proof.* Recall  $B = B(0, 1)$ . Note that

$$
\inf_{(\gamma,\rho)\in\mathcal{A}} \| |x|^{-b} |u|^\alpha v \|_{L_T^{\gamma'} L^{\rho'}} \le \inf_{(\gamma,\rho)\in\mathcal{A}} \| |x|^{-b} |u|^\alpha v \|_{L_T^{\gamma'} L^{\rho'}(B^c)} + \inf_{(\gamma,\rho)\in\mathcal{A}} \| |x|^{-b} |u|^\alpha v \|_{L_T^{\gamma'} L^{\rho'}(B)}
$$
  
=  $A_1 + A_2$ .

We need to find an admissible pair  $(\gamma_1, \rho_1)$  to estimate  $A_1$  by  $Y(T)$  norm. Using Hölder's inequality twice, we obtain

$$
A_1 \lesssim \| |u|^{\alpha} v \|_{L_T^{\gamma'_1} L^{\rho'_1}(B^c)}
$$
  
\n
$$
\leq T^{1 - \frac{n\alpha}{4}} \| u \|_{Y(T)}^{\alpha} \| v \|_{Y(T)}
$$
\n(3.3)

with  $\rho'_1$  satisfying Hölder conditions

<span id="page-8-4"></span>
$$
\frac{1}{\rho_1'} = \frac{\alpha}{\alpha + 2} + \frac{1}{\alpha + 2}.\tag{3.4}
$$

This gives value of  $\rho'_1 = \frac{\alpha+2}{\alpha+1}$  (and  $\rho_1 = \alpha+2$ ). Since  $(\gamma_1, \rho_1)$  is an admissible pair <sup>[3](#page-8-1)</sup>, we get

$$
\gamma_1' = \frac{4(\alpha + 2)}{4(\alpha + 2) - n\alpha}
$$

While  $\gamma'_1$  satisfies Hölder conditions <sup>[4](#page-8-2)</sup>

<span id="page-8-5"></span>
$$
\frac{1}{\gamma_1'} = \frac{1}{\frac{4(\alpha+2)}{4(\alpha+2)-n\alpha}} = \frac{1}{\nu} + \frac{\alpha}{\frac{4(\alpha+2)}{n\alpha}} + \frac{1}{\frac{4(\alpha+2)}{n\alpha}}.
$$
\n(3.5)

<span id="page-8-8"></span>.

Here  $\frac{1}{\nu}$  represents the exponent of T. Solving for  $\frac{1}{\nu}$ [5](#page-8-3) , we get

<span id="page-8-7"></span><span id="page-8-6"></span>
$$
\frac{1}{\nu} = 1 - \frac{n\alpha}{4}.
$$

Similarly, we need to find an admissible pair  $(\gamma_2, \rho_2)$  to estimate  $A_2$  by  $Y(T)$  norm. Applying Hölder's inequality twice, we obtain

$$
A_2 \le || |x|^{-b} |u|^\alpha v ||_{L_T^{\gamma_2'} L^{\rho_2'}(B)}
$$
  
\n
$$
\le || ||x|^{-b} ||_{L^{\gamma_3}(B)} ||u||_{L^{\alpha+2}}^{\alpha} ||v||_{L^{\alpha+2}} ||_{L_T^{\gamma_2'}} \tag{3.6}
$$

$$
\leq T^{\frac{1}{q_1}} \| |x|^{-b} \|_{L^{\gamma_3}(B)} \| u \|_{Y(T)}^{\alpha} \| v \|_{Y(T)}.
$$
\n(3.7)

<sup>&</sup>lt;sup>3</sup>It is easy to check that  $\rho_1 = \alpha + 2$  satisfies all conditions in Definition [2.1](#page-7-3) for the bound on  $\alpha$ .

<span id="page-8-1"></span><sup>&</sup>lt;sup>4</sup>Each Hölder exponent in [\(3.4\)](#page-8-4) and [\(3.5\)](#page-8-5) lies in the interval [1,  $\infty$ ] by the hypothesis on  $\alpha$ .

<span id="page-8-3"></span><span id="page-8-2"></span><sup>&</sup>lt;sup>5</sup>Note that  $\frac{1}{\nu}$  is positive by the hypothesis on  $\alpha$ .

Here  $\rho'_2, \gamma'_2, q_1$  and  $\gamma_3$  satisfies following conditions:

<span id="page-9-0"></span>
$$
\frac{1}{\rho_2'} = \frac{1}{\gamma_3} + \frac{\alpha}{\alpha + 2} + \frac{1}{\alpha + 2}
$$
\n(3.8)

$$
\frac{1}{\gamma_2'} = \frac{1}{q_1} + \frac{n\alpha^2}{4(\alpha+2)} + \frac{n\alpha}{4(\alpha+2)}
$$
(3.9)

$$
\frac{1}{\gamma_2'} = 1 - \frac{1}{2} \left( \frac{n}{\rho_2'} - \frac{n}{2} \right)
$$
\n(3.10)

$$
\frac{1}{q_1} > 0 \tag{3.11}
$$

$$
\frac{n}{\gamma_3} > b. \tag{3.12}
$$

Note that  $(3.8)$  and  $(3.9)$  is due to Hölder's inequality for space and time variable applied to get [\(3.6\)](#page-8-6) and [\(3.7\)](#page-8-7) respectively <sup>[6](#page-9-1)</sup>. Since  $(\gamma_2, \rho_2)$  is an admissible pair, we have [\(3.10\)](#page-9-0). The exponent of  $T$  needs to be positive in  $(3.7)$ , hence we have  $(3.11)$ . Condition  $(3.12)$ is required as  $||x|^{-b}||_{L^{\gamma_3}(B)} < \infty$  if and only if  $\frac{n}{\gamma_3} > b$ . From [\(3.8\)](#page-9-0) and [\(3.12\)](#page-9-0),

$$
\frac{n}{\rho_2'} - \frac{n(\alpha + 1)}{\alpha + 2} > b. \tag{3.13}
$$

Solving for  $\alpha^7$  $\alpha^7$ ,

$$
\alpha < \frac{2n - 2b\rho_2' - n\rho_2'}{n\rho_2' - n + b\rho_2'}.\tag{3.14}
$$

By our hypothesis,  $0 < \alpha < \frac{4-2b}{n}$ . Solving for  $\rho'_2$ 

$$
\frac{2n - 2b\rho_2' - n\rho_2'}{n\rho_2' - n + b\rho_2'} = \frac{4 - 2b}{n}
$$

yields

$$
\rho_2' = \frac{2n(n+2-b)}{n(n+4) + 2b(2-b)}.\tag{3.15}
$$

Inserting the value of  $\rho'_2$  in [\(3.10\)](#page-9-0), we get

$$
\gamma_2' = \frac{4(n+2-b)}{(2-b)(n+4-2b)}.\tag{3.16}
$$

Taking Remark [3.2](#page-10-0) into account, we have

$$
(\gamma_2, \rho_2) = \left(\frac{4(n+2-b)}{2n+4b+bn-2b^2}, \frac{2n(n+2-b)}{n^2-2nb-4b+2b^2}\right) \in \mathcal{A}.
$$

From [\(3.9\)](#page-9-0), we obtain

$$
\frac{4}{q_1} - (4 - 2b) = \frac{(2 - b)(n + 4 - 2b)}{n + 2 - b} - \frac{n\alpha(\alpha + 1)}{\alpha + 2} - (4 - 2b)
$$

$$
= \frac{-n(4 - 2b - n\alpha)}{(n + 2 - b)(\alpha + 2)} - n\alpha.
$$

This gives

<span id="page-9-3"></span>
$$
\frac{1}{q_1} = \frac{4 - 2b - n\alpha}{4} - \frac{n(4 - 2b - n\alpha)}{4(n + 2 - b)(\alpha + 2)}.
$$
\n(3.17)

<span id="page-9-2"></span><span id="page-9-1"></span><sup>&</sup>lt;sup>6</sup>The Hölder exponents in [\(3.8\)](#page-9-0) and [\(3.9\)](#page-9-0) lies in the interval  $[1, \infty]$  due to the bound on  $\alpha$  and b.

<span id="page-10-6"></span>Since  $0 < \alpha < \frac{4-2b}{n}$  and  $0 < b < \min\{2, n\}$ , note that  $\frac{1}{q_1}$  in  $(3.17)$  is a positive quantity. Substituting the value of  $\frac{1}{q_1}$  in [\(3.7\)](#page-8-7), we have

$$
A_2 \lesssim T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}} \|u\|_{Y(T)}^{\alpha} \|v\|_{Y(T)}.
$$
\n(3.18)

<span id="page-10-0"></span>Combining [\(3.3\)](#page-8-8) and [\(3.18\)](#page-10-1), we have the claim.

**Remark 3.2.** In order to estimate  $A_2$  by  $Y(T)$  norm in Lemma [3.1,](#page-8-0) we used below admissible pair  $(\gamma_2, \rho_2)$ , which imposes the restriction on b.

(1) Under the hypothesis of Lemma [3.1,](#page-8-0) we have

$$
(\gamma_2, \rho_2) = \left(\frac{4(n+2-b)}{2n+4b+bn-2b^2}, \frac{2n(n+2-b)}{n^2-2nb-4b+2b^2}\right) \in \mathcal{A}.
$$

(2) For  $n \geq 3$ , we have

$$
\left(0 < b \le \frac{n + 6 - \sqrt{(n + 6)^2 - 32}}{4}\right) \implies 2 \le \rho_2 \le \frac{2n}{n - 2}.
$$

(3) When  $n = 1, 2$ , we have  $2 \le \rho_2 \le \infty$  and  $2 \le \rho_2 < \infty$  respectively for b satisfying

$$
\begin{cases} 0 < b \leq \frac{3-\sqrt{7}}{2} & \text{if } n = 1 \\ 0 < b < 2-\sqrt{2} & \text{if } n = 2. \end{cases}
$$

<span id="page-10-5"></span>**Remark 3.3.** For  $T \leq 1$  and  $\alpha > 0$ , by Lemma [2.1\(](#page-6-0)[3\)](#page-6-6), we have

$$
X(T) = C_T L^2 \cap Y(T) + C_T M^{(\alpha+2), (\alpha+2)'} \hookrightarrow Y(T)^9.
$$

Thus, we have

$$
\inf_{(\gamma,\rho)\in\mathcal{A}} \| |x|^{-b} |u|^\alpha v \|_{L_T^{\gamma'}L^{\rho'}} \lesssim_n (T^{1-\frac{n\alpha}{4}} + T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}}) \|u\|_{X(T)}^{\alpha} \|v\|_{X(T)}
$$

where b and  $\alpha$  be as in Lemma [3.1.](#page-8-0)

**Remark 3.4.** For  $b = 0$ , we only need to evaluate  $A_1$  in Lemma [3.1](#page-8-0) as we do not have any singularity in this case. Cf. [\[18](#page-19-16), [19](#page-19-17)].

**Remark 3.5.** In contrast to  $L^2$ -GWP result of Guzmán (see  $(4.5)$ ), we need to narrow the range of b, from  $0 < b < \min\{2, n\}^{10}$  $0 < b < \min\{2, n\}^{10}$  $0 < b < \min\{2, n\}^{10}$  to  $0 < b \leq \tilde{b}$ . Revisit Remark [3.2](#page-10-0) for details.

**Proof of Theorem [1.2](#page-3-1).** By Duhamel's principle, INLS  $(1.1)$  is equivalent to the integral equation

$$
u(t) = e^{it\Delta}u_0 + i\mu \int_0^t e^{i(t-s)\Delta} |x|^{-b} (|u|^\alpha u)(s) ds := \Lambda(u)(t).
$$

Let a and  $T$  be a positive real numbers (to be chosen later). Define

$$
B(a,T) := \{ u \in X(T) : ||u||_{X(T)} \le a \}.
$$

<span id="page-10-1"></span>

<sup>&</sup>lt;sup>8</sup>It is easy to compute that  $2 \leq \rho_2$  for  $0 < b < 2$ .

<span id="page-10-2"></span><sup>&</sup>lt;sup>9</sup>Note that  $\|\cdot\|_{Y(T)} \lesssim_n \|\cdot\|_{X(T)}$ .

<span id="page-10-4"></span><span id="page-10-3"></span><sup>&</sup>lt;sup>10</sup>In [\[31](#page-19-9), Lemma 3.1],  $\rho_2 = \frac{4-2b+2n}{n-b}$  with  $\rho_2 \le \frac{2n}{n-2}$  for  $n \ge 3$  and  $\rho_2 > \frac{2n}{n-b}$  which would be fulfilled only if  $b \leq \min\{2, n\}.$ 

We will show that  $\Lambda$  is a contraction map on  $B(a,T)$ . Firstly, we consider the linear evolution of  $u_0$  where  $u_0 = v_0 + w_0 \in L^2 + M^{\alpha+2, (\alpha+2)'}$ ,  $v_0 \in L^2$  and  $w_0 \in M^{\alpha+2, (\alpha+2)'}$ . Assume that  $T \leq 1$ . Using Theorem [2.6](#page-7-0) and Proposition [2.3,](#page-6-4) we have

<span id="page-11-0"></span>
$$
||e^{it\Delta}u_0||_{X(T)} \le ||e^{it\Delta}v_0||_{X_1(T)} + ||e^{it\Delta}w_0||_{X_2(T)}
$$
  

$$
\lesssim_{n,\alpha} ||v_0||_{L^2} + (1+T^2)^{\frac{n}{2}(\frac{1}{2}-\frac{1}{\alpha+2})} ||w_0||_{M^{\alpha+2,(\alpha+2)'}}
$$
  

$$
\lesssim_n ||u_0||_{L^2+M^{\alpha+2,(\alpha+2)'}}.
$$
 (3.19)

This suggests the choice of  $a = C(n, \alpha) ||u_0||_{L^2 + M^{\alpha+2, (\alpha+2)'}}$ . Using  $X_1(T) \hookrightarrow X(T)$ , Theo-rem [2.6](#page-7-0) (with  $r \in {\alpha + 2, 2}$ ) and Remark [3.3,](#page-10-5) we have

$$
\left| \left| \int_{0}^{t} e^{i(t-\tau)\Delta} |x|^{-b} (|u|^{\alpha} u)(\tau) d\tau \right| \right|_{X(T)} \lesssim \left| \left| \int_{0}^{t} e^{i(t-\tau)\Delta} |x|^{-b} (|u|^{\alpha} u)(\tau) d\tau \right| \right|_{X_{1}(T)} \n\lesssim_{n,\alpha,b} |||u|^{\alpha} u||_{L_{T}^{\gamma'_{1}} L^{\rho'_{1}}(B^{c})} + |||x|^{-b} |u|^{\alpha} u||_{L_{T}^{\gamma'_{2}} L^{\rho'_{2}}(B)} \n\lesssim_{n} (T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}}) ||u||_{X(T)}^{\alpha+1}.
$$
\n(3.20)

Taking

<span id="page-11-1"></span>
$$
T := \min\left\{1, \ C(n,\alpha,b) \|u_0\|_{L^2+M^{\alpha+2,(\alpha+2)'}\atop{L^2+M^{\alpha+2,(\alpha+2)'}}} \right\},\tag{3.21}
$$

we combine [\(3.19\)](#page-11-0) and [\(3.20\)](#page-11-1) to conclude  $\Lambda(u) \in B(a,T)$ . Similarly, one can show that  $\Lambda(u)$  is a contraction mapping. In fact, using Lemma [2.4](#page-6-8) for  $G(0, u, v)$  and the previous argument employed to bound the integral part, for  $u, v \in B(a,T)$ , we have

$$
\|\Lambda(u) - \Lambda(v)\|_{X(T)} \lesssim \left\| \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} G(0, u, v)(\tau) d\tau \right\|_{X_1(T)}
$$
  

$$
\lesssim_{n, \alpha, b} \| (|u|^\alpha + |v|^\alpha) |u - v| \|_{L_T^{\gamma'_1} L^{\rho'_1}(B^c)} + \| |x|^{-b} (|u|^\alpha + |v|^\alpha) |u - v| \|_{L_T^{\gamma'_2} L^{\rho'_2}(B)}
$$
  

$$
\lesssim_n (T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}}) (||u||_{X(T)}^{\alpha} + ||v||_{X(T)}^{\alpha}) ||u - v||_{X(T)}.
$$
 (3.22)

By the choice of a and  $T^{-11}$  $T^{-11}$  $T^{-11}$ , we have

$$
\|\Lambda(u) - \Lambda(v)\|_{X(T)} \le \frac{1}{2} \|u - v\|_{X(T)}.
$$

Thus, by the contraction mapping theorem, we obtain a unique fixed point for  $\Lambda$ , which is a solution to INLS  $(1.1)$ . By the standard argument, one can establish the blow-up alternative.

For Lipschitz continuity, let  $v_0, w_0 \in V$ , where V is a neighborhood of  $u_0$ . Denote by v and w the unique maximal solutions of INLS  $(1.1)$  over the interval  $[0, T^*)$  with initial values  $v_0$  and  $w_0$ , respectively. Thus, v and  $w \in X(T')$  for  $T' < T^*$ . Hence

$$
\|v - w\|_{X(T')} = \|e^{it\Delta}v_0 - e^{it\Delta}w_0 + \Lambda(v) - \Lambda(w)\|_{X(T')}
$$
  

$$
\lesssim_{n,\alpha} \|v_0 - w_0\|_{L^2 + M^{\alpha+2,(\alpha+2)'}} + \frac{1}{2} \|v - w\|_{X(T')}.
$$
Thus,  $||v - w||_{X(T')}$  
$$
\lesssim_{n,\alpha} \|v_0 - w_0\|_{L^2 + M^{\alpha+2,(\alpha+2)'}}
$$

for any  $v_0, w_0 \in V \subset L^2 + M^{\alpha+2, (\alpha+2)'}$ . This concludes local Lipschitz continuity.  $\Box$ 

<span id="page-11-2"></span><sup>&</sup>lt;sup>11</sup>In the expression of a and  $T, C(n, \alpha, b)$  is choosen sufficiently small to ensure that  $\Lambda(u)$  is a contraction map on  $B(a,T)$ .

# 4. GLOBAL WELL-POSEDNESS IN  $M^{p,p'}$

<span id="page-12-10"></span><span id="page-12-5"></span>In this section, we shall prove Theorem [1.4.](#page-4-0) We start by decomposing (using Lemma [2.5\)](#page-7-1) initial data  $u_0 \in M^{p,p'} \subset L^2 + M^{(\alpha+2),(\alpha+2)'}$  into two parts such that the size of  $M^{(\alpha+2),(\alpha+2)'}$  –data can be controlled by arbitrary small quantity. Specifically, for any  $N > 1$  and given  $u_0 \in M^{p,p'}$ , there exists  $\phi_0 \in L^2$ ,  $\psi_0 \in M^{(\alpha+2),(\alpha+2)'}$  (depending on N) such that

<span id="page-12-0"></span>
$$
u_0 = \phi_0 + \psi_0 \tag{4.1}
$$

with

<span id="page-12-1"></span>
$$
\|\phi_0\|_{L^2} \lesssim N^{\beta}, \quad \|\psi_0\|_{M^{(\alpha+2), (\alpha+2)'}} \lesssim \frac{1}{N}
$$
 (4.2)

where

<span id="page-12-9"></span>
$$
\beta = \frac{\frac{1}{2} - \frac{1}{p}}{\frac{1}{p} - \frac{1}{\alpha + 2}}.\tag{4.3}
$$

Firstly, we consider INLS [\(1.1\)](#page-0-1) with initial data  $\phi_0$ :

<span id="page-12-6"></span>
$$
\begin{cases}\ni\partial_t v_0 + \Delta v_0 + \mu |x|^{-b} |v_0|^\alpha v_0 = 0\\v_0(\cdot, 0) = \phi_0 \in L^2.\end{cases}
$$
\n(4.4)

We recall that, in  $[31,$  Theorem 1.8, Guzmán proved that  $(4.4)$  has a unique global solution

<span id="page-12-2"></span>
$$
v_0 \in C(\mathbb{R}, L^2) \cap L^q_{loc}(\mathbb{R}, L^r)
$$
\n
$$
(4.5)
$$

<span id="page-12-8"></span>
$$
\sup_{(q,r)\in\mathcal{A}} \|v_0\|_{L^q_{loc}L^r} \lesssim_{n,r} \|\phi_0\|_{L^2}.
$$
\n(4.6)

Now, consider the modified INLS [\(1.1\)](#page-0-1) associated with the evolution of  $\psi_0$ :

<span id="page-12-7"></span>
$$
\begin{cases}\ni\partial_t w + \Delta w + \mu |x|^{-b} (|w + v_0|^{\alpha} (w + v_0) - |v_0|^{\alpha} v_0) = 0 \\
w(\cdot, 0) = \psi_0 \in M^{\alpha + 2, (\alpha + 2)'}.\n\end{cases} \tag{4.7}
$$

The solution of above I.V.P [\(4.7\)](#page-12-7) is given as

$$
w = e^{it\Delta}\psi_0 + w_0. \tag{4.8}
$$

The nonlinear interaction  $w_0$  corresponding to  $\psi_0$  can be expressed as

$$
w_0 = i\mu \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} (|w + v_0|^\alpha (w + v_0) - |v_0|^\alpha v_0) (\tau) d\tau.
$$
 (4.9)

Formally, we may rewrite the solution to INLS  $(1.1)$  corresponding to data  $u_0$  in  $(4.1)$  as follows

<span id="page-12-4"></span><span id="page-12-3"></span>
$$
u = v_0 + w = v_0 + e^{it\Delta}\psi_0 + w_0.
$$
\n(4.10)

In view of [\(4.5\)](#page-12-2) and Proposition [2.3,](#page-6-4) we notice that  $v_0$  and  $e^{it\Delta}\psi_0$  are globally defined in appropriate spaces. In order to establish desire global existence, we first understand the time interval of existence for  $w_0$ . To this end, we may rewrite

$$
w_0 = i\mu \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} G(v_0 + e^{i\tau\Delta} \psi_0, w_0)(\tau) d\tau + i\mu \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} G(v_0, e^{i\tau\Delta} \psi_0)(\tau) d\tau,
$$
\n(4.11)

with G as defined in Lemma [2.4.](#page-6-8) And more generally, we have the following local wellposedness result for perturb integral equation [\(4.11\)](#page-12-3).

<span id="page-13-4"></span>**Proposition 4.1.** Let  $\phi \in L^2$ ,  $\psi \in M^{(\alpha+2), (\alpha+2)'}$  and  $Y(T)$  be as in [\(3.2\)](#page-7-4). Assume that b and  $\alpha$  be as in Theorem [1.4.](#page-4-0) Denote by v the  $L^2$ -global solution (as in [\(4.5\)](#page-12-2)) for initial value  $\phi$ . Then there exists a constant  $C = C(n, \alpha, b) > 0$  such that integral equation

$$
w = i\mu \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} G(v + e^{i\tau\Delta} \psi, w)(\tau) d\tau + i\mu \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} G(v, e^{i\tau\Delta} \psi)(\tau) d\tau
$$

has a unique solution  $w \in Y(T)$  provided T satisfying

<span id="page-13-6"></span>
$$
T \le 1\tag{4.12}
$$

$$
T \leq C \left( \|\phi\|_{L^2} + \|\psi\|_{M^{\alpha+2,(\alpha+2)'}} \right)^{-\frac{\alpha}{4 - 2b - n\alpha} - \frac{n(4 - 2b - n\alpha)}{4(\alpha + 2)(n + 2 - b)}} \tag{4.13}
$$

$$
T \leq C \left( \|\psi\|_{M^{\alpha+2, (\alpha+2)'}} \right)^{-\frac{\alpha}{4 - 2b - n\alpha} - \frac{n(4 - 2b - n\alpha)}{4} + \frac{n\alpha^2}{4(\alpha + 2)(n + 2 - b)} + \frac{n\alpha^2}{4(\alpha + 2)}}.
$$
\n
$$
(4.14)
$$

<span id="page-13-7"></span>**Remark 4.2.** Since  $0 < \alpha < \frac{4-2b}{n}$  and  $0 < b < \tilde{b}^{12}$  $0 < b < \tilde{b}^{12}$  $0 < b < \tilde{b}^{12}$ , it follows that  $\frac{n}{(\alpha+2)(n+2-b)} < 1$ . Thus, the exponents on right hand side of conditions  $(4.13)$  and  $(4.14)$  involving  $L^2$  and  $M^{\alpha+2,(\alpha+2)'}$ - norms are both negative.

<span id="page-13-5"></span>**Corollary 4.3** (growth of perturb solution). Under the hypothesis of Proposition [4.1,](#page-13-4) there exists a constant  $C(n, \alpha, b)$  satisfying

<span id="page-13-3"></span><span id="page-13-2"></span>
$$
||w||_{L^{\infty}_TL^2} \lesssim_{n,\alpha,b} T^{\frac{n\alpha}{4(\alpha+2)}} ||\psi||_{M^{\alpha+2,(\alpha+2)'}}.
$$

We postpone the proof of Proposition [4.1](#page-13-4) and Corollary [4.3](#page-13-5) for the moment. And once we have these tools, we are ready to prove Theorem [1.4.](#page-4-0) We start with the following remark.

<span id="page-13-0"></span>Remark 4.4 (proof strategy). The following key points are in order to establish a global existence.

- Revisit Remark [1.5.](#page-4-1) In view of the discussion in the para below [\(4.10\)](#page-12-4), we are left to handle nonlinear interaction  $w_0$ . This we shall do using Proposition [4.1](#page-13-4) and Corollary [4.3.](#page-13-5)
- To this end, taking  $\phi = \phi_0$  and  $\psi = \psi_0$ , note that the solution of  $w_0$  exists on time-interval [0,  $T(N)$ ] with  $T = T(N)$  satisfying conditions [\(4.12\)](#page-13-6) to [\(4.14\)](#page-13-3).
- We extend this time-interval by the iterative scheme, which involve the following initial value decomposition:
	- At each iteration  $k \geq 1$ , we update the initial value  $\phi_k \in L^2$  using the new smoother term  $w_{k-1}(kT)$  and  $v_{k-1}(kT)$  obtained from the previous iteration at the initial time  $t = kT$  and merge  $\phi_k$  with  $\psi_k = e^{ikT\Delta} \psi_0$  to form the new approximation. See [\(4.16\)](#page-14-0).
	- Solving INLS [\(1.1\)](#page-0-1) with  $\phi_k \in L^2$  yields a solution  $v_k$  as described in [\(4.5\)](#page-12-2)-[\(4.6\)](#page-12-8). Further, solving modified INLS [\(4.7\)](#page-12-7) with initial value  $\psi_k$  and merging  $v_k$ , we get solution to INLS [\(1.1\)](#page-0-1) in the interval  $[kT,(k+1)T]$ . See [\(4.17\)](#page-14-1).
	- We repeat the iteration by redefining  $\phi_{k+1}$  and  $\psi_{k+1}$ , and maintaining control of the nonlinear interaction term  $w_k((k+1)T)$  using Corollary [4.3](#page-13-5) so that it can be absorbed into  $v_k((k+1)T)$  without loosing mass conservation of  $\phi_{k+1}$ .
- Finally, we shall observe that  $kT(N)$  can be made large enough under certain restriction on p. See Remark [4.5.](#page-16-0) This together with the blow-up alternative (Theorem [1.2\)](#page-3-1), yields a global solution.

<span id="page-13-1"></span><sup>&</sup>lt;sup>12</sup>Note that  $\tilde{b} < \min\{2, n\}$  for all *n*.

**Proof of Theorem [1.4](#page-4-0).** If possible, we assume that, for  $u_0 \in M^{p,p'}$ , the solution es-tablished in Theorem [1.2](#page-3-1) is not global in time. So we have the maximal time  $T^* < \infty$ . In this case, we shall produce a solution u of INLS  $(1.1)$  (to be defined in  $(4.17)$  below), which will exist on a larger interval  $[0, T_1]$  for  $T_1 > T^*$ . This will lead to a contradiction to the maximal time interval  $[0, T^*$ ).

Denote the constant from Proposition [4.1](#page-13-4) by  $C = C(n, \alpha, b)$  and put

<span id="page-14-3"></span>
$$
T = T(N) = (3CN^{\beta})^{-\frac{\alpha}{4 - 2b - n\alpha} - \frac{n(4 - 2b - n\alpha)}{4(\alpha + 2)(n + 2 - b)}}.
$$
\n(4.15)

By splitting the initial data (see [\(4.1\)](#page-12-0)) and using Proposition [4.1](#page-13-4) for  $\phi = \phi_0$  and  $\psi = \psi_0$ , the solution of INLS  $(1.1)$  exists in the interval  $[0, T(N)]$  and it is of the form

$$
u = v_0 + e^{it\Delta}\psi_0 + w_0,
$$

see [\(4.10\)](#page-12-4). We wish to extend our solution to the interval  $[T(N), 2T(N)]$  by similar procedure but with the new initial data as the sum of the following two functions:

$$
\phi_1 = v_0(T) + w_0(T)
$$
 and  $\psi_1 = e^{iT\Delta} \psi_0$ .

More generally, we wish to extend our solution further by considering the following iterative procedure :

– We define  $\phi_k$  and  $\psi_k$  for  $k \geq 1$  (for  $k = 0$ ,  $\phi_0$  and  $\psi_0$  are defined in [\(4.1\)](#page-12-0)) as follows:

<span id="page-14-0"></span>
$$
\phi_k = v_{k-1}(kT) + w_{k-1}(kT)
$$
 and  $\psi_k = e^{ikT\Delta}\psi_0,$ \n(4.16)

where

$$
w_{k-1}(kT) = i\mu \int_0^{kT} e^{i(kT-\tau)\Delta} |x|^{-b} G(v_{k-1} + e^{ik\tau\Delta} \psi_0, w_{k-1})(\tau) d\tau + i\mu \int_0^{kT} e^{i(kT-\tau)\Delta} |x|^{-b} G(v_{k-1}, e^{ik\tau\Delta} \psi_0)(\tau) d\tau.
$$

– Assume for  $kT \leq T^{*13}, \phi = \phi_k$  $kT \leq T^{*13}, \phi = \phi_k$  $kT \leq T^{*13}, \phi = \phi_k$  and  $\psi = e^{ikT\Delta}\psi_0$ , T satisfy all three conditions [\(4.12\)](#page-13-6), [\(4.13\)](#page-13-2) and [\(4.14\)](#page-13-3) of Proposition [4.1,](#page-13-4) where  $k \in \{0, 1, \dots, K - 1\}$ .

– Let  $v_k$  be INLS evolution of  $\phi_k$ , and by construction

<span id="page-14-1"></span>
$$
u(\cdot, t) = v_k(\cdot, t - kT) + w_k(\cdot, t - kT) + e^{it\Delta}\psi_0, \quad \text{if } t \in [kT, (k+1)T]
$$
(4.17)

defines a solution of INLS  $(1.1)$  for  $k \in \{0, 1, \dots, K-1\}.$ 

We shall show that the iterative process ends with  $KT > T^*$ . Since  $v_K$  and  $e^{it\Delta}\psi_0$  are globally defined in appropriate spaces, we are left to handle nonlinear interaction term  $w_K$  to extend the solution at K<sup>th</sup> iteration. To do this, we shall use Proposition [4.1](#page-13-4) with  $\phi = \phi_K$  and  $\psi = e^{iKT\Delta}\psi_0$ .

In view of Remark [4.2,](#page-13-7)  $T = T(N) \rightarrow 0$  as  $N \rightarrow \infty$  (see [\(4.15\)](#page-14-3)), and so the smallness condition  $(4.12)$  is satisfied independently of k for large N. Using Proposition [2.3](#page-6-4) and [\(4.2\)](#page-12-1), we have

<span id="page-14-4"></span>
$$
\|e^{it\Delta}\psi_0\|_{L^{\infty}([0,T^*+1],M^{\alpha+2,(\alpha+2)'})} \lesssim_{n,T^*} \|\psi_0\|_{M^{\alpha+2,(\alpha+2)'}} \lesssim_n \frac{1}{N} \xrightarrow{N \to \infty} 0.
$$
 (4.18)

Inserting  $e^{ikT\Delta}\psi_0$  in the right hand side of [\(4.14\)](#page-13-3), we have

$$
\left(\|e^{ikT\Delta}\psi_0\|_{M^{\alpha+2,(\alpha+2)'}}\right)^{-\frac{\alpha}{4-2b-n\alpha}-\frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}+\frac{n\alpha^2}{4(\alpha+2)}}\gtrsim_n N^{\frac{\alpha}{4-2b-n\alpha}-\frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}+\frac{n\alpha^2}{4(\alpha+2)}}\xrightarrow{N\to\infty}\infty.
$$

<span id="page-14-2"></span><sup>13</sup>Note that  $(K-1)T \leq T^*$ . Since  $T \leq 1$ , we will have  $KT \leq T^* + 1$ .

Since the lower bound is independent of k and  $T \xrightarrow{N \to \infty} 0$ , condition [\(4.14\)](#page-13-3) holds for sufficiently large  $N$ .

Thus, we either have  $KT > T^*$  or condition [\(4.13\)](#page-13-2) fails in the last iterative step  $k = K$ , i.e.

<span id="page-15-0"></span>
$$
3CN^{\beta} < \|\phi_K\|_{L^2} + \|e^{iKT\Delta}\psi_0\|_{M^{\alpha+2,(\alpha+2)'}}.\tag{4.19}
$$

Considering [\(4.18\)](#page-14-4), [\(4.19\)](#page-15-0) can be written as

<span id="page-15-1"></span>
$$
3CN^{\beta} < \|\phi_K\|_{L^2} + CN^{\beta}.\tag{4.20}
$$

We claim that even under condition  $(4.20)$ , we will have  $KT > T^*$ . This clearly lead to a contradiction to the definition of  $T^*$ .

In view of the construction of  $\phi_k$  and Corollary [4.3,](#page-13-5) we note that  $\phi_k \in L^2$  for  $k \in L^2$  $\{0, 1, \dots, K-1\}$ . Now exploiting the conservation  $(1.3)$  and Corollary [4.3](#page-13-5) (for  $w = w_k$ ) and  $\psi = e^{ikT\Delta}\psi_0, 0 \le k \le K - 1$ , we have

$$
\|\phi_K\|_{L^2} \le \|v_{K-1}\|_{L^{\infty}_{[(K-1)T,KT]}L^2} + \|w_{K-1}\|_{L^{\infty}_{[(K-1)T,KT]}L^2}
$$
\n
$$
= \|\phi_{K-1}\|_{L^2} + \|w_{K-1}\|_{L^{\infty}_{[(K-1)T,KT]}L^2}
$$
\n
$$
\le \|v_{K-2}\|_{L^{\infty}_{[(K-2)T,(K-1)T]}L^2} + \|w_{K-2}\|_{L^{\infty}_{[(K-2)T,(K-1)T]}L^2} + \|w_{K-1}\|_{L^{\infty}_{[(K-1)T,KT]}L^2}
$$
\n
$$
= \|\phi_{K-2}\|_{L^2} + \|w_{K-2}\|_{L^{\infty}_{[(K-2)T,(K-1)T]}L^2} + \|w_{K-1}\|_{L^{\infty}_{[(K-1)T,KT]}L^2}
$$
\n
$$
\le \cdots \le \|\phi_0\|_{L^2} + \sum_{k=0}^{K-1} \|w_k\|_{L^{\infty}_{[kT,(k+1)T]}L^2}
$$
\n
$$
\le_{n,\alpha,b} CN^{\beta} + T^{\frac{n\alpha}{4(\alpha+2)}} \sum_{k=0}^{K-1} \|e^{ikT\Delta}\psi_0\|_{M^{\alpha+2,(\alpha+2)'}}
$$
\n
$$
\le_{n,T^*} CN^{\beta} + T^{\frac{n\alpha}{4(\alpha+2)}} K \frac{C}{N}.
$$
\n(4.21)

In the last two inequalities, we have used  $(4.2)$  and  $(4.18)$ . Thus, using  $(4.15)$ ,  $(4.20)$  can be expressed as

$$
KT \gtrsim_{n,\alpha,b,T^*} N^{1+\beta} T^{1-\frac{n\alpha}{4(\alpha+2)}} \approx N^{1+\beta \left(1-\frac{\alpha \left(1-\frac{n\alpha}{4(\alpha+2)}\right)}{4-\frac{n(4-2b-n\alpha)}{4-\frac
$$

Note that N can be taken arbitrarily large. For any  $\beta$  satisfying

$$
0 < \beta < \begin{cases} \eta & \text{if } \alpha - \frac{n\alpha^2}{4(\alpha+2)} - \frac{4-2b-n\alpha}{4} + \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)} > 0 \\ \infty & \text{otherwise} \end{cases} \tag{4.23}
$$

where

<span id="page-15-3"></span><span id="page-15-2"></span>
$$
\eta = \frac{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}}{\alpha - \frac{n\alpha^2}{4(\alpha+2)} - \frac{4-2b-n\alpha}{4} + \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}},
$$

the exponent of N is positive in  $(4.22)$ , we get  $KT > T^*$ . This concludes the proof of Theorem [1.4.](#page-4-0) <span id="page-16-0"></span>Remark 4.5. (1) Recall β defined in terms of p in [\(4.3\)](#page-12-9). The range of β in [\(4.23\)](#page-15-3) in turn decides the range of p. Thus, we have  $p \in (2, p_{max})$  and

$$
p_{\max} = \begin{cases} \frac{4\alpha + 8 - n\alpha}{2\alpha + 2 + b + \frac{n(4 - 2b - n\alpha)}{2(\alpha + 2)(n + 2 - b)}} & \text{if } \beta = \eta\\ \alpha + 2 & \text{if } \beta = \infty. \end{cases}
$$

This justifies the choice of  $p$  in Theorem [1.4.](#page-4-0)

(2) It is easy to verify that  $2 < p_{max} = \frac{2}{1}$  $\frac{2}{1-\frac{4\zeta}{4\alpha+8-n\alpha}}$  when  $\beta = \eta$  and  $\zeta = \frac{4-2b-n\alpha}{4}$  $n(4-2b-n\alpha)$  $\frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}$ , since  $0<\frac{4\zeta}{4\alpha+8-n\alpha}<1$  by our assumption on b and  $\alpha$ .

We shall now prove Proposition [4.1](#page-13-4) and then Corollary [4.3.](#page-13-5)

**Proof of Proposition [4.1](#page-13-4).** Recall  $Y(T)$  defined in [\(3.2\)](#page-7-4). Define

$$
B(A, T) = \{ u \in Y(T) : ||u||_{Y(T)} \le A \}
$$

such that  $w \in B(A, T)$  with  $A > 0$  (to be choosen later), T be the minimum of the righthand sides of the conditions  $(4.12)$ ,  $(4.13)$  and  $(4.14)(w.l.o.g.$  we may assume). Further, define

$$
\Gamma(w) := i\mu \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} G(v + e^{it\Delta} \psi, w) d\tau + i\mu \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} G(v, e^{it\Delta} \psi) d\tau.
$$

Firstly, we need to show that  $\Gamma(w) \in B(A, T)$ .

Using Theorem [2.6,](#page-7-0) Lemma [2.4](#page-6-8) and Lemma [3.1](#page-8-0) under the assumption [\(4.12\)](#page-13-6), for any  $v_1, v_2 \in \mathbb{C}$ , we have

$$
\left\| \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} G(v_1, v_2)(\tau) d\tau \right\|_{Y(T)} \lesssim_{n, \alpha, b} |||v_1|^\alpha v_2||_{L_T^{\gamma'_1} L^{\rho'_1}(B^c)} + |||x|^{-b}|v_1|^\alpha v_2||_{L_T^{\gamma'_2} L^{\rho'_2}(B)}
$$
  
+  $|||v_2|^{\alpha+1}||_{L_T^{\gamma'_1} L^{\rho'_1}(B^c)} + |||x|^{-b}|v_2|^{\alpha+1}||_{L_T^{\gamma'_2} L^{\rho'_2}(B)}$   
(4.24)

<span id="page-16-2"></span><span id="page-16-1"></span>
$$
\lesssim \left( T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}} \right) \left( \|v_1\|_{Y(T)}^\alpha \|v_2\|_{Y(T)} + \|v_2\|_{Y(T)}^{\alpha+1} \right) \tag{4.25}
$$
\n
$$
\leq \left( T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)} + \frac{n\alpha}{4(\alpha+2)}} \right) \|v_1\|_\alpha \|v_1\|_\alpha
$$

$$
\lesssim_{\alpha} \left( T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)} + \frac{n\alpha}{4(\alpha+2)}} \right) \|v_1\|_{Y(T)}^{\alpha} \|v_2\|_{L_T^{\infty} L^{\alpha+2}} + \left( T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)} + \frac{n\alpha(\alpha+1)}{4(\alpha+2)}} \right) \|v_2\|_{L_T^{\infty} L^{\alpha+2}}^{\alpha+1}.
$$
\n(4.26)

In the last inequality, we have used the embedding  $L_T^{\infty} \hookrightarrow L_T^{\frac{4(\alpha+2)}{n\alpha}}$  i.e.  $\|\cdot\|_{L_T^{\frac{4(\alpha+2)}{n\alpha}}} \leq$  $T^{\frac{n\alpha}{4(\alpha+2)}}$   $\|\cdot\|_{L^{\infty}_T}.$ 

Using the estimate [\(4.26\)](#page-16-1) for  $v_1 = v, v_2 = e^{i\tau \Delta} \psi$  along with [\(4.6\)](#page-12-8), Lemma [2.1](#page-6-0) [\(3\)](#page-6-6) and Proposition [2.3](#page-6-4) under the assumption [\(4.12\)](#page-13-6), we obtain

$$
\left\| \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} G(v, e^{i\tau \Delta} \psi)(\tau) d\tau \right\|_{Y(T)}
$$

$$
\leq_{\alpha,n,b} \left( T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)} + \frac{n\alpha}{4(\alpha+2)}} \right) ||v||_{Y(T)}^{\alpha} ||e^{i\tau \Delta} \psi||_{L_T^{\infty} L^{\alpha+2}} \n+ \left( T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)} + \frac{n\alpha(\alpha+1)}{4(\alpha+2)}} \right) ||e^{i\tau \Delta} \psi||_{L_T^{\infty}, L^{\alpha+2}}^{\alpha+1} \n\leq_{\alpha,n} \left( T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)} + \frac{n\alpha}{4(\alpha+2)}} \right) ||\phi||_{L^2}^{\alpha} ||\psi||_{M^{\alpha+2, (\alpha+2)'}} \n+ \left( T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)} + \frac{n\alpha(\alpha+1)}{4(\alpha+2)}} \right) ||\psi||_{M^{\alpha+2, (\alpha+2)'}}^{\alpha+1} \n= T^{\frac{n\alpha}{4(\alpha+2)}} ||\psi||_{M^{\alpha+2, (\alpha+2)'}} \left( T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}} ||\phi||_{L^2}^{\alpha} \n+ T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)} + \frac{n\alpha^2}{4(\alpha+2)}} ||\psi||_{M^{\alpha+2, (\alpha+2)'}}^{\alpha} \right)
$$
\n
$$
\leq_{\alpha,n,b} T^{\frac{n\alpha}{4(\alpha+2)}} ||\psi||_{M^{\alpha+2, (\alpha+2)'}}.
$$

The last inequality follows due to our assumptions  $(4.13)$  and  $(4.14)$ . This suggests the choice of

<span id="page-17-0"></span>
$$
A = \frac{3}{C(n, \alpha, b)} T^{\frac{n\alpha}{4(\alpha+2)}} \|\psi\|_{M^{\alpha+2, (\alpha+2)'}}.
$$
 (4.27)

where  $C = C(n, \alpha, b)$  is the same constant as in [\(4.13\)](#page-13-2) and [\(4.14\)](#page-13-3), choosen such that

<span id="page-17-1"></span>
$$
\left| \left| \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} G(v, e^{i\tau \Delta} \psi)(\tau) d\tau \right| \right|_{Y(T)} \le \frac{A}{3}
$$
 (4.28)

holds. Using the estimate [\(4.25\)](#page-16-2) for  $v_1 = v + e^{i\tau \Delta} \psi$  and  $v_2 = w$  along with [\(4.6\)](#page-12-8), Lemma [2.1](#page-6-0) [\(3\)](#page-6-6) and Proposition [2.3](#page-6-4) under the assumption [\(4.12\)](#page-13-6), we have

$$
\left\| \int_{0}^{t} e^{i(t-\tau)\Delta} |x|^{-b} G(v + e^{i\tau \Delta} \psi, w)(\tau) d\tau \right\|_{Y(T)}
$$
  

$$
\leq_{\alpha,n,b} T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}} \left( \|v + e^{i\tau \Delta} \psi\|_{Y(T)}^{\alpha} \|w\|_{Y(T)} + \|w\|_{Y(T)}^{\alpha+1} \right)
$$
  

$$
\leq_{\alpha,n} \|w\|_{Y(T)} \left\{ T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)}} \left( (\|\phi\|_{L^{2}} + \|\psi\|_{M^{\alpha+2,(\alpha+2)'}})^{\alpha} + \|w\|_{Y(T)}^{\alpha} \right) \right\}
$$
  

$$
\leq_{\alpha,n,b} \|w\|_{Y(T)} \left\{ \frac{1}{3} + T^{\frac{4-2b-n\alpha}{4} - \frac{n(4-2b-n\alpha)}{4(\alpha+2)(n+2-b)} + \frac{n\alpha^{2}}{4(\alpha+2)}} \|\psi\|_{M^{\alpha+2,(\alpha+2)'} }^{\alpha} \right\}
$$

In the last inequality, we have used  $(4.13)$  in the first summand (choosing C in  $(4.13)$ ) small enough) and substitute the norm of w in  $Y(T)$  to the power of  $\alpha$  by  $A^{\alpha}$ ,  $(A$  given in [\(4.27\)](#page-17-0)) in the second summand. Consider second summand of last inequality under the assumption  $(4.14)$  (choosing C small enough in  $(4.14)$ ) to get

<span id="page-17-2"></span>
$$
\left\| \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} G(v + e^{i\tau \Delta} \psi, w)(\tau) d\tau \right\|_{Y(T)} \le \frac{2A}{3}.
$$
 (4.29)

Combining [\(4.28\)](#page-17-1) and [\(4.29\)](#page-17-2), we can say that  $\Gamma(w)$  belongs to  $B(A, T)$ . Contractivity of Γ follows similarly. Thus, by the Banach fixed-point theorem, we get a unique fixed point w to the integral equation [\(4.11\)](#page-12-3) on the time-interval  $[0, T]$ .

**Proof of Corollary [4.3](#page-13-5).** The proof follows from the Strichartz estimates (Theorem [2.6\)](#page-7-0) by replacing the norm defined with respect to  $Y(T)$  by  $L_T^{\infty}L^2$ . Using an admissible pair ( $\infty$ , 2) on the left hand side and the same pairs on the right hand side of [\(4.25\)](#page-16-2) and [\(4.26\)](#page-16-1) in the proof of Proposition [4.1,](#page-13-4) we obtain

$$
||w||_{L_T^{\infty}L^2} \leq \left| \left| \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} G(v + e^{it\Delta} \psi, w)(\tau) d\tau \right| \right|_{L_T^{\infty}L^2}
$$
  
+ 
$$
\left| \left| \int_0^t e^{i(t-\tau)\Delta} |x|^{-b} G(v, e^{i\tau\Delta} \psi)(\tau) d\tau \right| \right|_{L_T^{\infty}L^2}
$$
  

$$
\lesssim_{\alpha,n,b} T^{\frac{n\alpha}{4(\alpha+2)}} ||\psi||_{M^{\alpha+2,(\alpha+2)'}}.
$$

This completes the proof of Corollary [4.3.](#page-13-5)

Acknowledgments: The second author acknowledges the financial support from the University Grants Commission (UGC), India (file number 201610135365) for pursuing Ph.D. program. The second and third authors would like to express their gratitude to the Bhaskaracharya Mathematics Laboratory and the Brahmagupta Mathematics Library within the Department of Mathematics at IIT Indore, which are supported by the DST FIST Project (file number  $SR/FST/MS-I/2018/26$ ). The third author would like to thankfully acknowledge the financial support from the Matrics Project of DST (file number 2018/001166).

#### **REFERENCES**

- <span id="page-18-1"></span>[1] J. M. An and J. M. Kim. Small data global well-posedness and scattering for the inhomogeneous nonlinear Schrödinger equation in  $H^s(\mathbb{R}^n)$ . Z. Anal. Anwend., 40(4):453-475, 2021. [2](#page-1-1)
- <span id="page-18-2"></span>[2] J.M. An and J.M. Kim. Local well-posedness for the inhomogeneous nonlinear Schrödinger equation in  $H^s(\mathbb{R}^n)$ . Nonlinear Anal. Real World Appl., 59: Paper No. 103268, 21, 2021. [2](#page-1-1)
- <span id="page-18-4"></span>[3] L. Zhao B. X. Wang and B. Guo. Isometric decomposition operators, function spaces  $E_{p,q}^{\lambda}$  and applications to nonlinear evolution equations. J. Funct. Anal., 233(1):1–39, 2006. [3,](#page-2-4) [7](#page-6-9)
- <span id="page-18-3"></span>[4] Valeria Banica, R´emi Carles, and Thomas Duyckaerts. Minimal blow-up solutions to the mass-critical inhomogeneous NLS equation. Comm. Partial Differential Equations, 36(3):487–531, 2011. [2,](#page-1-1) [3](#page-2-4)
- <span id="page-18-0"></span>[5] J. Belmonte-Beitia, Víctor M Pérez-García, Vadym Vekslerchik, and Pedro J Torres. Lie symmetries and solitons in nonlinear systems with spatially inhomogeneous nonlinearities. Physical review letters, 98(6):064102, 2007. [1](#page-0-3)
- <span id="page-18-5"></span> $[6]$  A. Bényi and K. A. Okoudjou. Local well-posedness of nonlinear dispersive equations on modulation spaces. Bull. Lond. Math. Soc., 41(3):549–558, 2009. [3,](#page-2-4) [4](#page-3-2)
- <span id="page-18-6"></span>[7]  $\AA$ . Bényi and K. A. Okoudjou. *Modulation spaces—with applications to pseudo*differential operators and nonlinear Schrödinger equations. Applied and Numerical Harmonic Analysis. Birkhäuser/Springer, New York, [2020] ©2020. [3,](#page-2-4) [7](#page-6-9)
- <span id="page-18-9"></span>[8] D. G. Bhimani and R. Carles. Norm inflation for nonlinear Schrödinger equations in Fourier-Lebesgue and modulation spaces of negative regularity. J. Fourier Anal. Appl., 26(6):Paper No.78, 34, 2020. [4](#page-3-2)
- <span id="page-18-7"></span>[9] D. G. Bhimani, M. Grillakis, and K. A. Okoudjou. The Hartree-Fock equations in modulation spaces. Comm. Partial Differential Equations, 45(9):1088–1117, 2020. [4](#page-3-2)
- <span id="page-18-8"></span>[10] D. G. Bhimani, H. Hajaiej, and S. Haque. The mixed fractional Hartree equations in Fourier amalgam and modulation spaces. arXiv:2302.10683, 2023. [4](#page-3-2)
- <span id="page-18-10"></span>[11] D. G. Bhimani and S. Haque. Strong ill-posedness for fractional Hartree and cubic NLS equations. J. Funct. Anal., 285(11):Paper No. 110157, 47, 2023. [4](#page-3-2)

- <span id="page-19-15"></span>[12] D. G. Bhimani and P. K. Ratnakumar. Functions operating on modulation spaces and nonlinear dispersive equations. J. Funct. Anal., 270(2):621–648, 2016. [3,](#page-2-4) [4](#page-3-2)
- <span id="page-19-19"></span>[13] J. Bourgain. Global solutions of nonlinear Schrödinger equations, volume 46 of American Mathematical Society Colloquium Publications. American Mathematical Society, Providence, RI, 1999. [5](#page-4-2)
- <span id="page-19-4"></span>[14] L. Campos and J. Murphy. Threshold solutions for the intercritical inhomogeneous NLS. SIAM J. Math. Anal., 55(4):3807–3843, 2023. [2](#page-1-1)
- <span id="page-19-14"></span>[15] M. Cardoso and L. G. Farah. Blow-up solutions of the intercritical inhomogeneous NLS equation: the non-radial case. Math. Z., 303(3):Paper No. 63, 18, 2023. [3](#page-2-4)
- <span id="page-19-5"></span>[16] M. Cardoso, L. G. Farah, C. M. Guzmán, and Jason Murphy. Scattering below the ground state for the intercritical non-radial inhomogeneous NLS. Nonlinear Anal. Real World Appl., 68:Paper No. 103687, 19, 2022. [2,](#page-1-1) [3](#page-2-4)
- <span id="page-19-0"></span>[17] T. Cazenave. Semilinear Schrödinger equations, volume 10 of Courant Lecture Notes in Mathematics. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2003. [1,](#page-0-3) [2,](#page-1-1) [8](#page-7-5)
- <span id="page-19-16"></span>[18] L. Chaichenets. *Modulation spaces and nonlinear Schrödinger equations*. PhD thesis, KIT-Bibliothek, 2018. [3,](#page-2-4) [4,](#page-3-2) [5,](#page-4-2) [7,](#page-6-9) [8,](#page-7-5) [11](#page-10-6)
- <span id="page-19-17"></span>[19] L. Chaichenets, D. Hundertmark, P. Kunstmann, and N. Pattakos. On the existence of global solutions of the one-dimensional cubic NLS for initial data in the modulation space  $M_{n,q}(\mathbb{R})$ . J. Differential Equations, 263(8):4429-4441, 2017. [3,](#page-2-4) [4,](#page-3-2) [5,](#page-4-2) [11](#page-10-6)
- <span id="page-19-2"></span>[20] F. Dalfovo, S. Giorgini, L. P. Pitaevskii, and S. Stringari. Theory of bose-einstein condensation in trapped gases. Reviews of modern physics, 71(3):463, 1999. [1](#page-0-3)
- <span id="page-19-6"></span>[21] V. D. Dinh. Blowup of  $H^1$  solutions for a class of the focusing inhomogeneous nonlinear Schrödinger equation. Nonlinear Anal., 174:169–188, [2](#page-1-1)018. 2
- <span id="page-19-13"></span>[22] V. D. Dinh and S. Keraani. Long time dynamics of nonradial solutions to inhomogeneous nonlinear schrödinger equations. SIAM Journal on Mathematical Analysis, 53(4):4765–4811, 2021. [2](#page-1-1)
- <span id="page-19-7"></span>[23] L. G. Farah and C. M. Guzmán. Scattering for the radial 3D cubic focusing inhomogeneous nonlinear Schrödinger equation. *J. Differential Equations*,  $262(8):4175-4231$ , 2017. [2,](#page-1-1) [3](#page-2-4)
- <span id="page-19-12"></span>[24] L.G. Farah. Global well-posedness and blow-up on the energy space for the inhomogeneous nonlinear Schrödinger equation. J. Evol. Equ.,  $16(1):193-208$  $16(1):193-208$  $16(1):193-208$ ,  $2016.2$
- <span id="page-19-20"></span>[25] H. G. Feichtinger. Modulation spaces on locally compact abelian groups. Technical report, University of Vienna, 1983, Proceedings of the International Conference on Wavelet and Applications, 2002, New Delhi Allied Publishers, India, 2003. [6,](#page-5-2) [8](#page-7-5)
- <span id="page-19-8"></span>[26] F. Genoud. Bifurcation and stability of travelling waves in self-focusing planar waveguides. Adv. Nonlinear Stud., 10(2):357–400, 2010. [2](#page-1-1)
- <span id="page-19-11"></span>[27] F. Genoud. An inhomogeneous,  $L^2$ -critical, nonlinear Schrödinger equation. Z. Anal. Anwend., 31(3):283–290, 2012. [2](#page-1-1)
- <span id="page-19-3"></span>[28] F. Genoud and C. A. Stuart. Schrödinger equations with a spatially decaying nonlinearity: existence and stability of standing waves. *Discrete Contin. Dyn. Syst.*,  $21(1):137-186$  $21(1):137-186$ , 2008. [1,](#page-0-3) 2
- <span id="page-19-1"></span>[29] T. S. Gill. Optical guiding of laser beam in nonuniform plasma. Pramana, 55:835– 842, 2000. [1](#page-0-3)
- <span id="page-19-18"></span>[30] S. Guo. On the 1D cubic nonlinear Schrödinger equation in an almost critical space. J. Fourier Anal. Appl., 23(1):91–124, 2017. [4](#page-3-2)
- <span id="page-19-9"></span>[31] C. M. Guzmán. On well posedness for the inhomogeneous nonlinear Schrödinger equation. Nonlinear Anal. Real World Appl., 37:249–286, 2017. [2,](#page-1-1) [6,](#page-5-2) [11,](#page-10-6) [13](#page-12-10)
- <span id="page-19-10"></span>[32] C. M. Guzmán and J. Murphy. Scattering for the non-radial energy-critical inhomogeneous NLS. J. Differential Equations, 295:187–210, 2021. [2,](#page-1-1) [3](#page-2-4)
- <span id="page-20-13"></span>[33] R. Hyakuna and M. Tsutsumi. On existence of global solutions of Schrödinger equations with subcritical nonlinearity for  $\tilde{L}^p$ -initial data. *Proc. Amer. Math. Soc.*, 140(11):3905–3920, 2012. [5](#page-4-2)
- <span id="page-20-4"></span>[34] Y. V. Kartashov, B. A. Malomed, V. A. Vysloukh, M. R. Belić, and L. Torner. Rotating vortex clusters in media with inhomogeneous defocusing nonlinearity. Optics *letters*,  $42(3):446-449, 2017.$  $42(3):446-449, 2017.$  $42(3):446-449, 2017.$  1
- <span id="page-20-18"></span>[35] T. Kato. The inclusion relations between  $\alpha$ -modulation spaces and  $L^p$ -Sobolev spaces or local Hardy spaces. J. Funct. Anal., 272(4):1340–1405, 2017. [7](#page-6-9)
- <span id="page-20-19"></span>[36] M. Keel and T. Tao. Endpoint Strichartz estimates. Amer. J. Math., 120(5):955–980, 1998. [8](#page-7-5)
- <span id="page-20-11"></span>[37] C. E. Kenig. On the work of Jean Bourgain in nonlinear dispersive equations. Bull. Amer. Math. Soc. (N.S.), 58(2):173–189, 2021. [5](#page-4-2)
- <span id="page-20-14"></span>[38] C. E. Kenig, G. Ponce, and L. Vega. Global well-posedness for semi-linear wave equations. Comm. Partial Differential Equations, 25(9-10):1741–1752, 2000. [5](#page-4-2)
- <span id="page-20-16"></span>[39] M. Kobayashi and M. Sugimoto. The inclusion relation between Sobolev and modulation spaces. J. Funct. Anal., 260(11):3189–3208, 2011. [7](#page-6-9)
- <span id="page-20-17"></span>[40] M. Kobayashi and M. Sugimoto. The inclusion relation between Sobolev and modulation spaces. J. Funct. Anal., 260(11):3189–3208, 2011. [7](#page-6-9)
- <span id="page-20-0"></span>[41] F. Linares and G. Ponce. Introduction to Nonlinear Dispersive Equations. Springer, New York, 2020. [1,](#page-0-3) [8](#page-7-5)
- <span id="page-20-3"></span>[42] C. S. Liu and V. K. Tripathi. Laser guiding in an axially nonuniform plasma channel. Physics of plasmas, 1(9):3100–3103, 1994. [1](#page-0-3)
- <span id="page-20-5"></span>[43] Franck Merle. Nonexistence of minimal blow-up solutions of equations  $iu_t = -\Delta u$  $k(x)|u|^{4/N}u$  in  $\mathbf{R}^N$ . Ann. Inst. H. Poincaré Phys. Théor., 64(1):33–85, 1996. [2,](#page-1-1) [3](#page-2-4)
- <span id="page-20-6"></span>[44] J. Murphy. A simple proof of scattering for the intercritical inhomogeneous NLS. Proc. Amer. Math. Soc., 150(3):1177–1186, 2022. [2,](#page-1-1) [3](#page-2-4)
- <span id="page-20-10"></span>[45] T. Oh and Y. Wang. Global well-posedness of the one-dimensional cubic nonlinear Schrödinger equation in almost critical spaces. J. Differential Equations,  $269(1):612-$ 640, 2020. [4](#page-3-2)
- <span id="page-20-15"></span>[46] M. Ruzhansky, M. Sugimoto, J. Toft, and N. Tomita. Changes of variables in modulation and Wiener amalgam spaces. *Math. Nachr.*,  $284(16):2078-2092$  $284(16):2078-2092$  $284(16):2078-2092$ ,  $2011$ . 7
- <span id="page-20-8"></span>[47] M. Ruzhansky, B. X. Wang, and H. Zhang. Global well-posedness and scattering for the fourth order nonlinear Schrödinger equations with small data in modulation and Sobolev spaces. J. Math. Pures Appl. (9), 105(1):31–65, 2016. [3,](#page-2-4) [7](#page-6-9)
- <span id="page-20-1"></span>[48] T. Tao. Nonlinear dispersive equations, volume 106 of CBMS Regional Conference Series in Mathematics. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2006. Local and global analysis. [1,](#page-0-3) [5](#page-4-2)
- <span id="page-20-7"></span>[49] Yoshio Tsutsumi.  $L^2$ -solutions for nonlinear Schrödinger equations and nonlinear groups. Funkcial. Ekvac., 30(1):115–125, 1987. [2](#page-1-1)
- <span id="page-20-12"></span>[50] A. Vargas and L. Vega. Global wellposedness for 1D non-linear Schrödinger equation for data with an infinite  $L^2$  norm. J. Math. Pures Appl.  $(9)$ , 80 $(10)$ :1029-1044, 2001. [5](#page-4-2)
- <span id="page-20-9"></span>[51] B. X. Wang and H. Hudzik. The global Cauchy problem for the NLS and NLKG with small rough data. J. Differential Equations,  $232(1):36-73$  $232(1):36-73$  $232(1):36-73$ ,  $2007$ . [3,](#page-2-4) [6,](#page-5-2) 7
- <span id="page-20-2"></span>[52] B. X. Wang, Z. Huo, C. Hao, and Z. Guo. Harmonic analysis method for nonlinear evolution equations. I. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2011. [1,](#page-0-3) [3,](#page-2-4) [7](#page-6-9)

Divyang G. Bhimani, Department of Mathematics, Indian Institute of Science Education and Research, Pune 411008, India

Email address: divyang.bhimani@iiserpune.ac.in

Diksha Dhingra, Department of Mathematics, Indian Institute of Technology, Indore, 452020, India

Email address: dikshadd1996@gmail.com

Vijay Kumar Sohani, Department of Mathematics, Indian Institute of Technology, Indore 452020, India

Email address: vsohani@iiti.ac.in