# SPECTRAL ESTIMATES ON HYPERBOLIC SURFACES AND A NECESSARY CONDITION FOR OBSERVABILITY OF THE HEAT SEMIGROUP ON MANIFOLDS

#### ALIX DELEPORTE, MARC ROUVEYROL

ABSTRACT. This article is a continuation of  $\lbrack \text{Rou24} \rbrack$ . We study the concentration properties of spectral projectors on manifolds, in connection with the uncertainty principle. In [\[Rou24\]](#page-30-0), the second author proved an optimal uncertainty principle for the spectral projector of the Laplacian on the hyperbolic half-plane. The aim of the present work is to generalize this condition to surfaces with hyperbolic ends. In particular, we tackle the case of cusps, in which the volume of balls of fixed radius is not bounded from below. We establish that spectral estimates hold from sets satisfying a thickness condition, with a proof based on propagation of smallness estimates of Carleman and Logunov–Malinnikova type. We also prove the converse, namely the necessary character of the thickness condition, on any smooth manifold with Ricci curvature bounded from below.

#### CONTENTS



Key words and phrases. spectral estimates, propagation of smallness, hyperbolic surfaces, thick sets, heat equation.

#### <span id="page-1-0"></span>2 ALIX DELEPORTE, MARC ROUVEYROL

#### 1. Introduction and main result

#### <span id="page-1-1"></span>1.1. Main result.

1.1.1. Notation and main results. On a non-compact Riemannian manifold  $(M, g)$  without boundary, endowed with a smooth metric g, consider the Laplace–Beltrami operator  $-\Delta_q$ with domain  $C_c^{\infty}(M)$ . The corresponding quadratic form is  $q: D(q) \to \mathbb{R}$  given by

$$
D(q) = H^{1}(M) = \{ u \in L^{2}(M), \nabla_{g} u \in L^{2}(M) \}, \quad q(u) = \int_{M} |\nabla_{g} u|_{g}^{2} d \operatorname{vol}_{g}.
$$

We denote by  $-\Delta_g$  (with the convention  $\Delta_g \leq 0$ ) the Friedrichs extension of  $(-\Delta_g, C_c^{\infty}(M))$ , with domain  $D(-\Delta_g) = \{u \in H^1(M), \Delta_g u \in L^2(M)\}\.$  We refer to [\[RS80\]](#page-30-1) and [\[Lew22\]](#page-29-0) for spectral-theoretic considerations.

 $\sqrt{-\Delta_g}$  is well-defined and admits a spectral measure dm<sub>Λ</sub>. Any function  $u \in L^2(M)$  can be The Laplace–Beltrami operator is non-negative and self-adjoint, so that its square root decomposed against this spectral measure and  $\sqrt{-\Delta_g}$  admits a functional calculus:

(1.1) 
$$
u = \int_0^{+\infty} dm_{\lambda} u
$$
,  $\phi(\sqrt{-\Delta_g})u = \int_0^{+\infty} \phi(\lambda) dm_{\lambda} u$   

$$
\langle \phi(\sqrt{-\Delta_g})u, \varphi(\sqrt{-\Delta_g})u \rangle_{L^2(M)} = \int_0^{+\infty} \phi(\lambda) \varphi(\lambda) \langle dm_{\lambda} u, u \rangle
$$

for any  $u \in L^2(M)$ ,  $\phi, \varphi \in \mathcal{L}^{\infty}(\mathbb{R}^+)$ . We then define the spectral projector on the frequency window [0,  $\Lambda$ ],  $\Lambda > 0$ , by

(1.2) 
$$
\Pi_{\Lambda} u = \mathbb{1}_{\sqrt{-\Delta_g} \leq \Lambda} u = \int_0^{\Lambda} dm_{\lambda} u,
$$

so that

<span id="page-1-3"></span>(1.3) 
$$
\|\phi(\sqrt{-\Delta_g})\Pi_{\Lambda}u\|_{L^2(M)} \leq \sup_{\lambda\in[0,\Lambda]}|\phi(\lambda)| \|\Pi_{\Lambda}u\|_{L^2(M)}.
$$

The present article aims at studying uncertainty principles for spectrally localized functions in Ran( $\Pi_{\Lambda}$ ), in the specific case where M is a hyperbolic surface, or a surface with hyperbolic ends. It is a natural sequel of the article [\[Rou24\]](#page-30-0) by the second author, which dealt with the case of the hyperbolic half-plane  $M = \left(\mathbb{H}, \frac{dx^2 + dy^2}{dt^2}\right)$  $\frac{y^2+dy^2}{y^2}$ , for hyperbolic surfaces are quotients of  $\mathbb H$ by suitable isometry groups. To avoid diluting the main theorems in geometric discussions, we next state the results of the article, and postpone definition of the precise geometric framework to the next subsection.

The uncertainty principle we prove takes the form of an inequality in the spirit of Logvinenko and Sereda's spectral projector estimates [\[LS74,](#page-30-2) [Kov00,](#page-29-1) [Kov01\]](#page-29-2). Precisely, we say that a Borel subset  $\omega \subset M$  satisfies a *spectral estimate* if there exists some constant  $C(\omega)$ , independent of u and the frequency threshold  $\Lambda$ , such that

<span id="page-1-2"></span>(Spec) 
$$
\forall u \in L^2(M), \forall \Lambda > 0, \quad \|\Pi_{\Lambda}u\|_{L^2(M)} \leq Ce^{C\Lambda}\|\Pi_{\Lambda}u\|_{L^2(\omega)}.
$$

In [\[Rou24\]](#page-30-0), we proved that in the case  $M = \mathbb{H}$ , the spectral estimate is equivalent to the so-called thickness condition:

<span id="page-2-3"></span>**Definition 1.1.** – Let  $R > 0$ . We say that  $\omega \subset M$  is thick at scale R (or R-thick) if there exists some volume ratio  $\delta(R) > 0$  such that

<span id="page-2-0"></span>(Thick) 
$$
\forall z \in M, \frac{\text{vol}(B_z(R) \cap \omega)}{\text{vol}(B_z(R))} \ge \delta.
$$

We shall say that  $\omega$  is thick if there exists some  $R > 0$  such that  $\omega$  is R-thick.

<span id="page-2-1"></span>The first of our results states that thickness of  $\omega$  is sufficient for a spectral estimate to hold:

**Theorem 1.** – Assume M is a surface with hyperbolic ends. Take  $\omega$  a Borel subset of M. If  $\omega$ satisfies [\(Thick\)](#page-2-0) for some parameters  $(R, \delta)$ , then  $\omega$  satisfies [\(Spec\)](#page-1-2) for some constant  $C(\omega)$ .

Section [2](#page-11-1) is dedicated to the proof of Theorem [1.](#page-2-1) The next theorem states that the converse of Theorem [1](#page-2-1) also holds, namely that the thickness assumption is also necessary for a spectral estimate to hold. Only a Ricci curvature lower bound is required, making the assumptions on M much weaker.

<span id="page-2-2"></span>**Theorem 2.** – Let  $(M, g)$  be a complete, boundaryless, smooth Riemannian manifold, not necessarily compact, of arbitrary dimension, with Ricci curvature bounded from below:

(1.4) 
$$
\text{Ric} \ge -Kg, \quad K > 0.
$$

in the sense of quadratic forms. If  $\omega$  satisfies [\(Spec\)](#page-1-2), then  $\omega$  is thick at scale R for some sufficiently large  $R > 0$ .

Theorem [2](#page-2-2) is proved in Section [3.](#page-22-0) A complete bibliographical discussion of previous similar results and of elements of proof will come in the next subsections of this introduction, but we already make some comments about our main results. They are the object of the following remarks.

### Remarks

- (1) As explained in the next subsection, surfaces with hyperbolic ends do satisfy the Ricci curvature lower bound Ric  $\geq -Kq$ ,  $K > 0$ . Consequently, Theorem [2](#page-2-2) implies that the converse of Theorem [1](#page-2-1) holds.
- (2) In both Theorem [1](#page-2-1) and Theorem [2,](#page-2-2) the metric g is assumed to be smooth. It is likely that this assumption could be weakened in Theorem [2,](#page-2-2) as the natural framework in which the assumption

$$
\operatorname{Ric}(g) + Kg \ge 0
$$

is well-defined is that of Lipschitz metrics (with a distributional definition of the Ricci tensor). Ric $(g)$  + Kg would then be a positive measure, implying no gain of regularity on g even in harmonic charts. However, the proof of Theorem [2](#page-2-2) relies on heat kernel estimates which have only been proved for smooth metrics, to the best of the authors' knowledge.

(3) One interesting feature of surfaces with hyperbolic ends is that the denominator  $vol(B_z(R))$  which appears in the denominator of [\(Thick\)](#page-2-0) can tend to zero, in the asymptotic in which  $z$  is sent to the end of a cusp. Thus, unlike in preceding works [\[BM21,](#page-28-0) [Rou24\]](#page-30-0), there is no uniform lower bound for the volume of  $\omega$  inside a given ball of radius R.

- (4) In a recent article [\[LBM24\]](#page-29-3), Kévin Le Balc'h and Jérémy Martin prove an equivalent of Theorem [1](#page-2-1) for operators of the form  $-\text{div}(A(x)\nabla) + V(x), V \in L^{\infty}$ , on the full flat space. Their proof, like ours, is based on Logunov–Malinnikova estimates for the gradients of solutions to elliptic equations [\[LM18b\]](#page-29-4), in the spirit of Burq and Moyano [\[BM21,](#page-28-0) [BM23\]](#page-28-1). It would be interesting to combine the two techniques to treat operators of the form  $-\Delta_q + V$ , with  $V \in L^{\infty}(M)$ .
- (5) The following volume doubling estimate holds on any Riemannian manifold with Ricci curvature bounded from below and is a consequence of the Bishop–Gromov inequality [\[CGT82,](#page-28-2) [GHL04\]](#page-28-3): there exists a constant  $C_D > 0$  such that for any  $z \in M, R > 0$ ,

$$
\text{vol}(B_z(2R)) \le C_D e^{C_D R} \text{vol}(B_z(R)).
$$

The doubling constant  $C_D$  depends only on the dimension of the manifold and the curvature lower bound. The  $e^{C_D R}$  factor is obtained by comparison theorems between the growth of balls on manifolds with Ricci curvature bounded from below and on the hyperbolic space with constant curvature equal to the lower bound, on which it is the correct asymptotic for large R. Since the critical mass ratio  $\delta$  of Assumption [1.1](#page-2-3) is allowed to depend on R,  $R_0$ -thickness of  $\omega$  for some  $R_0 > 0$  implies R-thickness for any  $R > R_0$ .

- (6) In a manifold with infinite diameter, a thick set cannot be compact.
- (7) By the volume doubling estimate above, if  $\omega$  is R-thick and K is a compact subset of  $\omega$ , then  $\omega \setminus K$  remains thick (typically at scale diam(K) + 2R).

<span id="page-3-0"></span>1.1.2. Hyperbolic surfaces and hyperbolic ends. We now give all the relevant definitions related to the geometry of hyperbolic surfaces and surfaces with hyperbolic ends. The key elements are the classification of hyperbolic ends given in Theorem [3](#page-4-0) and the realizations of funnels and cusps as quotients of appropriate subsets of the hyperbolic half-plane (depicted in Figure [1\)](#page-4-1). The presentation follows Chapter 2 of David Borthwick's excellent book on the topic [\[Bor16\]](#page-28-4), keeping only information relevant to the article. For complementary material on spectral theory of hyperbolic surfaces, we refer to [\[Bor16,](#page-28-4) [Ber11\]](#page-28-5) and the references therein.

The hyperbolic upper half-plane will be denoted  $\mathbb{H} = \{(x, y) \in \mathbb{R}^2, y > 0\}$ . The hyperbolic metric on  $\mathbb H$  is  $g_{\mathbb H} = \frac{dx^2 + dy^2}{u^2}$  $\frac{d+dy^2}{y^2}$ . The corresponding Riemannian measure is  $d \text{ vol}_{\mathbb{H}} = \frac{dxdy}{y^2}$  $\frac{xay}{y^2}$  and the Laplace–Beltrami operator on H is given by

$$
\Delta_{\mathbb{H}} = y^2 \left( \partial_x^2 + \partial_y^2 \right).
$$

Geodesics for the metric  $g_{\parallel}$  are vertical straight lines and arcs of circles centered on the x-axis. Geodesic balls are Euclidean balls with center shifted upwards with respect to the hyperbolic center. Precisely, given a point  $z = (x, y) \in \mathbb{H}$  and a radius  $R > 0$ , we have

(1.5) 
$$
B_z^{\mathbb{H}}(R) = B_{(x,y\cosh(R))}^{euc}(y\sinh(R)).
$$

where  $B^{\mathbb{H}}$  denotes a geodesic ball and  $B^{euc}$  denotes a Euclidean ball.

The group of isometries of  $(\mathbb{H}, g_{\mathbb{H}})$ , known as Möbius transformations, is isomorphic to the group of  $2 \times 2$  real matrices with determinant 1, quotiented by  $\{\pm I_2\}$ . The common notation for this group is  $PSL_2(\mathbb{R})$ . Given a matrix

$$
T = \begin{pmatrix} a & b \\ c & d \end{pmatrix}
$$



<span id="page-4-1"></span>FIGURE 1. Left: fundamental domain for a funnel (delimited by three geodesic arcs, the top one of length  $l$ ) in the hyperbolic half-plane  $\mathbb H$  and its images by the corresponding cyclic Fuchsian group. Right: fundamental domain  $[0, 1] \times [a, +\infty]$  for a cusp in the hyperbolic half-plane H and its images by the Fuchsian group  $\langle z \mapsto z + 1 \rangle$ .

in  $PSL_2(\mathbb{R})$ , the corresponding isometry is the complex function

$$
z = x + iy \mapsto \frac{az + b}{cz + d}.
$$

Discrete subgroups of  $PSL_2(\mathbb{R})$  are called Fuchsian. They are exactly the groups of isometries acting properly discontinuously on H.

We call a *hyperbolic surface* any complete, 2-dimensional Riemannian manifold with Gaussian curvature constant equal to −1. As a consequence of Hopf's theorem on the classification of manifolds with constant sectional curvature [\[Lee97,](#page-29-5) Theorem 11.12 and Corollary 11.13], any hyperbolic surface X is isometric to a quotient of  $\mathbb H$  by some Fuchsian group  $\Gamma$ .

In [\[Rou24\]](#page-30-0), the second author proved the equivalence between [\(Thick\)](#page-2-0) and [\(Spec\)](#page-1-2) for  $\Delta_{\mathbb{H}}$ . A main contribution of the present work is to generalize this to geometrically finite hyperbolic surfaces, the definition of which we now give. A *fundamental domain*  $\mathcal{F} \subset \mathbb{H}$  for a Fuchsian group  $\Gamma$  is a closed subset satisfying

$$
\bigcup_{T\in\Gamma}T\mathcal{F}=\mathbb{H},\quad\forall T\in\Gamma\setminus\{I\},\text{Int}(\mathcal{F})\cup\text{Int}(T\mathcal{F})=\varnothing.
$$

We say that a Fuchsian group is geometrically finite if it admits a fundamental domain which is a geodesically convex polygon with a a finite number of sides. This is equivalent to the hyperbolic surface  $\Gamma \setminus \mathbb{H}$  having finite genus (number of holes) and a finite number of ends.

The structure of geometrically finite hyperbolic surfaces is well-understood. H and its quotients by cyclic Fuchsian groups are called elementary surfaces. The classification result for non-elementary geometrically finite hyperbolic surfaces is the following [\[Bor16,](#page-28-4) Theorem 2.23]:

<span id="page-4-0"></span>**Theorem 3** (Classification of hyperbolic ends).  $-Any$  non-elementary, geometrically finite hyperbolic surface M admits a compact region K such that  $M \setminus K$  is a finite, disjoint union of cusps and funnels.

Namely, (non-elementary, geometrically finite) hyperbolic surfaces can be decomposed as a disjoint union of elements of the three following type:

- a compact core, known as the truncated Nielsen region;
- funnels;

• cusps.

Here, given a hyperbolic cylinder  $\langle z \mapsto e^l z \rangle \setminus \mathbb{H}$ , where  $l > 0$  is the length of its delimiting geodesic  $\{z \in \mathbb{H}, \text{Im}(z) = 0\}$ , a funnel is a half-cylinder, namely a connected component of the cylinder minus the geodesic.

Given a parabolic cylinder  $\langle z \mapsto z + 1 \rangle \backslash H$ , a cusp is "the small end" of the cylinder. More precisely, Denoting  $\mathbb{S}_{\theta}^1 = \mathbb{Z} \setminus \mathbb{R}$  the circle of length 1, cusps are isometric to  $\mathbb{S}_{\theta}^1 \times (a, +\infty)_y$  for some  $a > 0$  depending on the cusp, endowed with the metric  $\frac{d\theta^2 + dy^2}{y^2}$ .

We depict fundamental domains for funnels and cusps in Figure [1.](#page-4-1) Note that the local injectivity radius at a point (the largest radius of a geodesic ball centered on that point and onto which the exponential map is injective) goes to zero asymptotically at the end of a cusp but grows to infinity in a funnel. In particular, the volume of balls of fixed radius centered at a given point tends to 0 towards the end of a cusp but is asymptotically constant towards the end of a funnel.

In the article, funnels will be denoted F or  $F_j$  and cusps will be denoted C or  $C_j$ .

From now on, we will always imply that the geometric finiteness property holds when mentioning hyperbolic surfaces.

The nature of the compact core does not intervene in the proof of our results. Thus, though we are motivated by hyperbolic surfaces, our main result is stated for *surfaces with hyperbolic* ends. The precise framework for Theorem [1](#page-2-1) is thus the following.

**Definition 1.2.** – We say that a two-dimensional Riemannian manifold M endowed with a smooth metric q is a surface with hyperbolic ends if there exists some compact region  $K \subset M$ , still called the compact core, such that  $M \setminus K$  is a finite union of funnels and cusps.

We insist on the fact that in this definition, no geometric assumption is made in the compact  $\operatorname{core} K$ .

We end this overview by recalling the behavior of the spectrum of the Laplace–Beltrami  $\Delta_g$  on a hyperbolic surface. Note that cusps have finite volume while the volume of a funnel is infinite. For compact hyperbolic surfaces, like on any compact manifold, the spectrum is made of discrete eigenvalues and there exists an orthonormal basis of eigenvectors in  $L^2(M)$ . We shall not discuss this case further as it was covered by Burq and Moyano in [\[BM23\]](#page-28-1), in the case of Dirichlet or Neumann boundary conditions,  $W^{2,\infty}$  manifolds and Lipschitz metrics. For non-compact, finite-volume hyperbolic surfaces (meaning they have no funnels and a finite number of cusps), the Laplace–Beltrami operator has absolutely continuous spectrum  $\frac{1}{4}$  $\frac{1}{4}$ ,  $\infty$ ), a discrete spectrum consisting of finitely many eigenvalues in  $[0, \frac{1}{4}]$  $\frac{1}{4}$ , and may have up to infinitely many embedded eigenvalues in  $\left[\frac{1}{4}\right]$  $(\frac{1}{4}, \infty)$  [\[Bor16,](#page-28-4) Theorem 3.12]. These embedded eigenvalues are very unstable, in the sense that they are destroyed by certain deformations of  $\Gamma$ [\[PS85\]](#page-30-3). There are no such embedded eigenvalues in the infinite-area case [\[Bor16,](#page-28-4) Proposition 7.5].

Lastly, we emphasize that Theorem [1](#page-2-1) for hyperbolic surfaces cannot be deduced immediately from the main result of [\[Rou24\]](#page-30-0) on the full half-plane H, as the Γ-periodic extension of a function in  $L^2(\Gamma \setminus \mathbb{H}) \setminus \{0\}$  does not belong to  $L^2(\mathbb{H})$ . Should there exist a strategy based on spectral inequalities on the full half-plane and periodic extensions from  $\Gamma \setminus \mathbb{H}$  to  $\mathbb{H}$ , we believe that our local-to-gobal, chart-based method for proving Theorem [1](#page-2-1) is more robust in the perspective of generalizations to a larger class of manifolds.

<span id="page-5-0"></span>1.2. Context. This bibliographical overview is divided into two parts. First, we present a hopefully not too partial history of spectral estimates and their applications, with an attempt to cover the most recent advances in the field. Next, we cover some bibliography on the propagation of smallness inequality toolbox that is used throughout the article and its connection to spectral estimates.

1.2.1. Spectral estimates on manifolds, motivations and applications. The study of spectral estimates of the form [\(Spec\)](#page-1-2) from thick sets goes back at least to Logvinenko–Sereda and Kacnel'son [\[LS74,](#page-30-2) [Kac73\]](#page-29-6), in connection with uncertainty principles for the Fourier transform. Their methods are complex-analytic and were improved by Nazarov and Kovrizhkin, among others [\[Naz93,](#page-30-4) [Kov00,](#page-29-1) [Kov01,](#page-29-2) [Kov03\]](#page-29-7). In particular, Kovrizhkin made explicit the dependence of the constant  $Ce^{C\Lambda}$  appearing in [\(Spec\)](#page-1-2), in terms of the thickness parameters  $\delta$  and R.

In their seminal article [\[LR95\]](#page-29-8), Lebeau and Robbiano established that spectral estimates allowed to prove controllability results for the heat equation on compact manifolds, both with and without boundary. In the boundary-free case, they proved exact null-controllabilty of the heat equation for  $L^2$  initial data and arbitrarily short time when the sensor set  $\omega$  is open. The equivalence between controllability and observability and the link with spectral estimates were developed in an abstract setting by Miller [\[Mil10\]](#page-30-5), with explicit dependence of the controllability and observability costs. Unlike in [\[LR95\]](#page-29-8), where exact controllability is shown directly and observability is given as a corollary, most recent controllability results for the heat equation in the spirit of Lebeau and Robbiano now prove observability directly. The telescoping series method of Phung and Wang [\[PW13\]](#page-30-6) provides a robust proof that spectral estimates imply observability from positive-measure time sets via appropriate interpolation estimates. Subsequent developments of this method for Borel sensor sets  $\omega$  on the flat space include [\[AE13,](#page-27-1) [AEWZ14\]](#page-27-2). The sharp character of the thickness condition for observability of the heat equation on the flat space  $\mathbb{R}^d$  was proved simultaneously and independently in [\[EV18\]](#page-28-6) and [\[WWZZ19\]](#page-30-7). For additional results on spectral estimates and controllability of parabolic equation from thick sets in non-compact subdomains of the flat space we refer to [\[ES21\]](#page-28-7), the review article  $[**ENS**<sup>+</sup>20]$  and the references therein.

The Lebeau–Robbiano strategy for controllability generalizes to other parabolic equations beyond heat, including linear thermoelasticity systems [\[LZ98,](#page-30-8) [BN02\]](#page-28-9), the Kolmogorov equation [\[LRM16\]](#page-29-9), and parabolic equations involving quadratic (possibly non-self-adjoint) operators [\[BJPS21,](#page-28-10) [MPS23\]](#page-30-9). The explicit dependence in  $\delta$  of the constant appearing in [\(Spec\)](#page-1-2) provided by Kovrizhkin [\[Kov01\]](#page-29-2) has also allowed to prove controllability results for weakly dissipative parabolic equations, typically involving fractional Laplacians with exponents smaller than  $\frac{1}{2}$ [\[AK24\]](#page-28-11). The thickness condition on  $\omega$  required for controllability of weakly dissipative equations involves thickness at asymptotically small scales (see [\[AK24,](#page-28-11) Definition 3]). In a recent article [\[BT24\]](#page-28-12), Kovrizhkin's spectral estimate with explicit dependence of the constant was generalized to Banach-space valued functions, opening up possible applications to parabolic systems and parabolic equations with a parameter.

Propagation of smallness (typically Carleman inequality-based) and spectral inequalities have also proved key to controlling the [0,Λ] frequency range of Schrödinger [\[Mil04b,](#page-30-10) [SSY25\]](#page-30-11) and wave [\[BJ16\]](#page-28-13) equations for arbitrarily large thresholds  $\Lambda$ . Controllability in the highfrequency asymptotic regime is then provided by other methods, typically of semiclassical nature.

Since the original article by Lebeau and Robbiano [\[LR95\]](#page-29-8), the theory of spectral estimates on manifolds has encountered numerous developments, with applications to both controllability results and spectral geometry. In [\[JL99\]](#page-29-10), the authors provide a spectral estimate from open sensor sets on compact manifolds, both with and without boundary, and use it to obtain

bounds on the  $(d-1)$ -Hausdorff measure of nodal sets of sums of eigenfunctions of the Laplacian in the spirit of [\[DF88,](#page-28-14) [DF90\]](#page-28-15). Like in [\[LR95\]](#page-29-8), the proof of the spectral inequality is based on interior and boundary Carleman estimates (the latter when the sensor set  $\omega$  is a subset of the boundary). In recent years, the spectral estimates of [\[LR95,](#page-29-8) [JL99\]](#page-29-10) have been generalized to the case where  $\omega$  is merely a Borel subset of the interior of the compact manifold, on the sphere [\[DV24\]](#page-28-16) and on general compact manifolds with or without boundary [\[BM21\]](#page-28-0). In the latter, Burq and Moyano use Logunov and Malinnikova's propagation of smallness estimates for gradients of harmonic functions [\[LM18b,](#page-29-4) Theorem 5.1] as a replacement for previously Carleman-based propagation of smallness to derive controllability properties from Borel and zero-measure sets on compact Riemannian manifolds with  $W^{2,\infty}$  metrics. The passage from the boundary-free case to the case of Dirichlet or Neumann boundary conditions is performed using a double-manifold reflection argument.

The present article is part of a program aiming at generalizing spectral estimate results from thick sets to non-compact geometric settings by adapting the strategy first introduced by Burq and Moyano. Miller [\[Mil05a,](#page-30-12) [Mil05b\]](#page-30-13) was the first to tackle that problem using the Lebeau– Robbiano method and to provide a necessary condition for spectral estimates on non-compact manifolds with Ricci curvature bounded below. The next steps in that direction focused on the full space  $\mathbb{R}^d$  endowed with analytic [\[LM19\]](#page-29-11) and bounded [\[BM23\]](#page-28-1) perturbations of the flat metric. The first instance of a result on non-compact, non-flat manifolds was provided by Rose and Tautenhahn [\[RT24\]](#page-30-14), using different, Brownian-motion and heat-kernel based methods, in a general geometric framework (Ricci curvature bounded from below, namely the assumption of Theorem [2\)](#page-2-2). Their result holds for frequency ranges  $[0, \Lambda]$  below a frequency threshold constrained by the curvature lower bound, thus not allowing to control the heat equation. The first high-frequency result for spectral estimates from thick (Borel) sensor sets on a non-compact, non-flat Riemannian flat was provided by the second author in the specific case of the hyperbolic half-plane  $\mathbb{H}$  [\[Rou24\]](#page-30-0). In that article, the spectral estimate is deduced from [\(Thick\)](#page-2-0) by adapting the approach of Burq and Moyano [\[BM23,](#page-28-1) [BM21\]](#page-28-0) on a suitably defined covering of the manifold, while the necessary character of the thickness condition is shown using heat kernels in the spirit of [\[WWZZ19\]](#page-30-7). The present article can be seen as a continuation of [\[Rou24\]](#page-30-0). The main contributions made here and absent from [\[Rou24\]](#page-30-0) are the following:

- (1) in Sections [2.2](#page-14-0) and [2.3,](#page-17-0) we tackle the fact that the denominator  $vol(B_z(R))$  (with R fixed) can go to zero on a manifold with null global injectivity radius (in our case, when z tends to the end of a cusp). In  $\mathbb{H}$ , that denominator was constant with respect to z and included in the  $\delta(R)$  parameter. We hope that the periodic lifting argument provided may be generalized to a larger class of manifolds, relaxing the constantcurvature assumption.
- (2) in Section [2.4,](#page-20-0) we prove that controllability of the heat equation on a manifold with ends only requires a sensor set  $\omega$  active in the ends. The idea of that section is to use Carleman-based propagation of smallness estimates to propagate smallness from any end to the compact core.
- (3) in Section [3,](#page-22-0) we generalize the idea of combining observability estimates and heat kernel bounds on manifolds developed in [\[Rou24,](#page-30-0) Section 2.3] to manifolds with smooth metric and Ricci curvature bounded from below. Again, attention must be paid to recovering the small denominator  $vol(B_z(R))$  which could be ignored on H.

We now mention two questions which which we deem interesting but chose not to address in the article. The first one concerns the dependence of the constant  $Ce^{C\Lambda}$  in [\(Spec\)](#page-1-2) on the geometry and the thickness parameters  $(R, \delta)$ , which has applications of its own as discussed earlier. On compact manifolds, this (more specifically, the dependence of the observability cost for the heat equation) was studied by Miller [\[Mil04a\]](#page-30-15). The results were refined by Lau-rent and Léautaud [\[LL21\]](#page-29-12) in the case where the sensor set  $\omega$  is a ball of decreasing radius. Concerning the behavior of observability and control costs for the heat equation as a function of the constants appearing in the spectral estimate, we refer to [\[WWZZ19,](#page-30-7) [NTTV20\]](#page-30-16) and the references therein. Lastly, Theorem 5.1 of [\[RT24\]](#page-30-14) gives an explicit dependence of the spectral estimate constant in terms of the Ricci curvature lower bound and the thickness parameters for frequencies smaller than a specific (and again, explicit) threshold. Note that the thickness parameters used in [\[RT24\]](#page-30-14) are not exactly those of the present article: the  $(R, \rho)$ -thickness condition used in  $[RT24]$ , under which the sensor set  $\omega$  must contain a ball of radius  $\rho$  inside every ball of radius  $R$ , is similar to [\(Thick\)](#page-2-0) in spirit but strictly implies it.

The other main question that we chose not to tackle in the present article is that of the sensitivity of the thickness condition to potentials. Understanding spectral estimates for the operator  $-\Delta + V$  opens up applications to controllability of the heat equation with a potential (see for example [\[BJPS21,](#page-28-10) [MPS23\]](#page-30-9) in the harmonic case) and random Schrödinger operators [\[RMV13,](#page-30-17) [Kle13\]](#page-29-13). On the full space  $\mathbb{R}^d$ , it has been proved [\[LBM24\]](#page-29-3) that spectral estimates from sets satisfying [\(Thick\)](#page-2-0) still hold for  $-\text{div}(A(x)\nabla u) + V$  with V an  $L^{\infty}$  potential and A uniformly elliptic with Lipschitz entries. Like that of Theorem [1,](#page-2-1) their proof is based on the Logunov–Malinnikova propagation of smallness estimate for gradients. Previous results in the case of bounded potentials used an analytic framework [\[LM19\]](#page-29-11) or a decay assumption at infinity [\[Zhu24,](#page-30-18) Theorem 2]. In the case of unbounded, polynomially growing potentials of the full space, it has been shown that spectral estimates hold from sets of decaying density [\[AS23,](#page-28-17) [DSV24,](#page-28-18) [Wan24,](#page-30-19) [Zhu24\]](#page-30-18). The idea in that case is to combine propagation of smallness arguments with frequency-quantitative decay (Agmon) estimates for eigenfunctions [\[Agm82,](#page-27-3) [Alp24\]](#page-28-19). We hope to study Agmon estimates on non-compact manifolds and decaying thickness conditions in presence of a potential in forthcoming work.

<span id="page-8-0"></span>1.2.2. About propagation of smallness inequalities and their application on manifolds. We conclude this bibliographical overview by covering the propagation of smallness tools used to prove Theorem [1.](#page-2-1) Precise statements of the corresponding theorems can be found in Appendix [A.](#page-26-0) Because the proof of Theorem [2](#page-2-2) is of a different, geometric-analytic nature, discussion of the tools used to prove that result is postponed to Section [3.](#page-22-0)

The main propagation of smallness tool used in the article is the result by Logunov and Malinnikova [\[LM18b,](#page-29-4) Theorem 5.1] for gradients of harmonic functions. The proof is a combination of estimates for the critical sets of solutions of elliptic equations [\[CNV15\]](#page-28-20), properties of the doubling index of these solutions and combinatorial lemmas, which were used by the same authors to obtain results on nodal set bounds [\[LM18a,](#page-29-14) [Log18a,](#page-29-15) [Log18b\]](#page-29-16). The estimate holds on subdomains of the Euclidean space and the constant involved depends on the Lipschitz and ellipticity constants of the metric as well as the Hausdorff measure  $\delta$  of the set from which smallness is propagated. For these reasons, we need to apply the Logunov–Malinnikova estimate in charts satisfying at least the following properties:

(1) the charts must contain a ball of radius  $R$  so that the thickness assumption [\(Thick\)](#page-2-0) may allow to recover a uniform mass of the sensor set  $\omega$  inside the chart;

#### 10 ALIX DELEPORTE, MARC ROUVEYROL

- (2) the atlas is locally uniformly finite, meaning each point of the manifold lies at the intersection of a uniformly bounded number of charts;
- (3) the domain of the charts must be uniform, and the metric must be controlled in terms of ellipticity and Lipchitz constants in the corresponding coordinate system.

In the way it is used in the present article, the Logunov–Malinnikova estimate can be thought of as a replacement for Carleman-estimate based propagation of smallness (see for example [\[JL99,](#page-29-10) Lemmas 14.4, 14.5]). One major improvement in the former is that it requires only very mild assumptions on the set from which smallness propagates (positivity of some  $(d-1-\epsilon)$  Hausdorff content), while the Carleman method requires open sets – or open sets of the boundary, allowing for propagation of smallness from a  $(d-1)$ -dimensional set. The usual proof of Carleman estimates relies on pseudodifferential calculus with a large parameter and a commutator estimate based on the Gårding inequality. This gives a weighted unique continuation estimate (the Carleman inequality), and a good choice of the weight function allows to deduce propagation of smallness results (see [\[LR95,](#page-29-8) Section 3] or [\[JL99,](#page-29-10) Sections 14.3, 14.4]). A comprehensive overview of Carleman estimates and their applications to control problems is the book of Le Rousseau, Lebeau and Robbiano [\[LRLR22a,](#page-29-17) [LRLR22b\]](#page-29-18). For a shorter introduction (in French), we refer the reader to [\[Leb15\]](#page-29-19).

1.3. Strategy of proof. The proof of the sufficient condition consists in applying propagation of smallness inequalities in charts, to a harmonic function which contains all relevant information on  $\Pi_{\Lambda} u$ . To that end, we introduce the harmonic extension

(1.6) 
$$
v_{\Lambda}(t,\cdot) = \int_{\lambda=0}^{\Lambda} \frac{\sinh(\lambda t)}{\lambda} dm_{\lambda} u,
$$

which satisfies

$$
(1.7) \t v_{\Lambda}(t=0,\cdot)=0, \quad \partial_t v_{\Lambda}(t=0,\cdot)=\Pi_{\Lambda} u, \quad \nabla_g v_{\Lambda}(t=0,\cdot)=0, \quad \partial_t^2 + \Delta_g v_{\Lambda}=0.
$$



<span id="page-9-0"></span>Figure 2. Sketch of a hyperbolic surface of genus 1 with two cusps. In purple dashed lines, the decomposition into its compact core and cusps  $M = K \sqcup \mathcal{C}_1 \sqcup \mathcal{C}_2$ . In blue full lines, the regions  $\tilde{C}_1$ ,  $\tilde{C}_2$  where we prove propagation of smallness estimates, and the compact region  $\tilde{K}$  overlapping  $\tilde{C}_1$  and  $\tilde{C}_2$ .

The proof of Theorem [1](#page-2-1) adapts Burq and Moyano's strategy for Lipschitz metrics on the full space  $\mathbb{R}^d$  [\[BM21\]](#page-28-0) (itself inspired by the pioneering work of Jerison and Lebeau [\[JL99\]](#page-29-10)). The general idea is to apply propagation of smallness results for the harmonic extension  $v_{\Lambda}$ in local charts and to derive [\(Spec\)](#page-1-2) from summing these estimates together. We now describe the steps of the proof of Theorem [1](#page-2-1) and how they differ from [\[BM23,](#page-28-1) [BM21\]](#page-28-0) and the second author's previous work [\[Rou24\]](#page-30-0).

The first step is to apply Logunov and Malinnikova's propagation of smallness inequality [\[LM18b\]](#page-29-4) for gradients in a suitably chosen rectangle of the hyperbolic half-plane H. We refer to section [1.2.2](#page-8-0) for a quick bibliographical overview of this result. This is essentially the only step where the explicit expression of the metric is used, as it allows to construct charts appropriate for applying the Logunov–Malinnikova estimate with uniform constants. The inequality obtained (Lemma [2.1\)](#page-11-3) and its proof are close to those used in [\[Rou24\]](#page-30-0).

We then decompose the surface with hyperbolic ends into its cusps  $\mathcal{C}_j$ , funnels  $F_j$ , and a compact core K. The propagation of smallness inequality for rectangles in  $\mathbb H$  is extended to funnels and cusps using periodic lifting arguments based on their respective structure (see Figure [1\)](#page-4-1). Specifically, the surface is decomposed as depicted in Figure [2:](#page-9-0) we prove propagation of smallness from the intersection of  $\omega$  with slightly smaller subsets  $\tilde{F}_j$ ,  $\tilde{C}_j$  of the funnels and cusps, respectively.

We then denote K an open, relatively compact enlargement of K that intersects every  $\tilde{F}$ ,  $\tilde{C}$ . In that relatively compact set, we use Carleman-based propagation of smallness results from [\[LRLR22b\]](#page-29-18) and [\[JL99\]](#page-29-10) to reduce the question of the compact core to propagation of smallness inequalities in the ends. We finish by patching together estimates in every tilded subset to obtain the spectral estimate.

The proof of Theorem [2](#page-2-2) is inspired by a similar result from [\[WWZZ19\]](#page-30-7) on the flat space. Indeed, the spectral estimate implies an observability inequality for solutions of the free heat equation as stated in Lemma [3.4.](#page-24-0) The thickness inequality can then be recovered by testing this observability inequality for short times with a well-chosen heat kernel. Key elements of the proof are heat kernel bounds and volume doubling estimates that hold on manifolds with smooth metrics and Ricci curvature bounded from below. As remarked earlier, we hope that the regularity assumption on the metric  $q$  can be weakened, but we believe that a lower bound on Ricci curvature is a crucial condition in Theorem [2.](#page-2-2) Indeed, there are examples of ill-behaved heat kernels when that assumption is dropped.

The plan of the article is the following. In Section [2,](#page-11-1) we prove Theorem [1.](#page-2-1) The proof is divided into four subsections. In the first, we prove a propagation of smallness result for rectangles in the hyperbolic half-plane. In subsections [2.2](#page-14-0) and [2.3,](#page-17-0) we show how this result extends to funnels and cusps using periodic lifting arguments. We finish Section [2](#page-11-1) by showing propagation of smallness from an end to the compact core and patching estimates on funnels, cusps and the compact core together to deduce the spectral estimate. Section [3](#page-22-0) is dedicated to the proof of Theorem [2.](#page-2-2) The Appendix contains statements and comments on the propagation of smallness inequalities used throughout Section [2.](#page-11-1)

<span id="page-10-0"></span>Notation. Throughout the rest of the article, we denote by z a generic element in the manifold M,  $d_g(z, z')$  the geodesic distance between two points  $z, z' \in M$  and  $d \text{vol}_g$  the Riemannian measure corresponding to the metric g.  $B_z(R)$  denotes a geodesic ball of center z and radius R.

The symbol  $\lesssim$  means that the left-hand-side is bounded by the right-hand-side up to an implicit multiplicative constant which is independent of relevant quantities. The dependence of implicit constants is specified as an index when deemed necessary.  $A \simeq B$  means that both  $A \lesssim B$  and  $B \lesssim A$  hold.

<span id="page-11-0"></span>Acknowledgements. M.R. would like to thank Ruoyu P.T. Wang for pointing out the interest of studying spectral estimates on surfaces with cusps, and Nicolas Burq and Luc Robbiano for interesting discussions about the topic of this article. M.R. acknowledges funding from a CDSN PhD grant of ENS Paris-Saclay, via the Hadamard Doctoral School of Mathematics. A.D. acknowledges funding by the ERC-2022-ADG grant GEOEDP (project 101097171).

#### 2. THICKNESS OF  $\omega$  IMPLIES THE SPECTRAL ESTIMATE

<span id="page-11-2"></span><span id="page-11-1"></span>2.1. Propagation of smallness on a rectangle of the hyperbolic half-plane. As explained in the introduction, we aim to obtain propagation of smallness estimates on both funnels and cusps. These estimates will in turn derive from an inequality localized on rectangles of the hyperbolic half-plane, the proof of which is borrowed from [\[Rou24\]](#page-30-0). These estimates will be applied with w and u equal to appropriate periodic liftings to  $\mathbb H$  of  $v_\Lambda$  and  $\Pi_{\Lambda}u$ , respectively. The statement is the following:

<span id="page-11-3"></span>**Lemma 2.1.** – Fix some  $R' > 0$  and denote, for any positive  $\kappa$ ,

(2.1)  
\n
$$
\mathcal{R}_{\kappa} = (-\kappa, \kappa)_{x} \times \left(\kappa 2^{-R'}, \kappa \left(\frac{3}{2}\right)^{R'}\right)_{y},
$$
\n
$$
\mathcal{R}_{\kappa}^{\eta} = (- (1 + \eta)\kappa, (1 + \eta)\kappa)_{x} \times \left((1 - \eta)\kappa 2^{-R'}, (1 + \eta)\kappa \left(\frac{3}{2}\right)^{R'}\right)_{y},
$$

rectangles of the hyperbolic half-plane  $\left(\mathbb{H}, \frac{dx^2+dy^2}{dt^2}\right)$  $\frac{+dy^2}{y^2}$ . Let  $\mathfrak{u} : \mathcal{R}^{\kappa}_{\eta} \to \mathbb{R}$  and  $w : \mathbb{R}_t \times \mathcal{R}^{\kappa}_{\eta} \to \mathbb{R}$  satisfy

$$
(\partial_t^2 + \Delta_{\mathbb{H}})w = 0, \quad \partial_t w(t = 0) = \mathfrak{u}, \quad \nabla_{\mathbb{H}} w(t = 0) = 0.
$$

Consider a subset  $\hat{\omega} \subset \mathcal{R}_{\kappa}$  satisfying  $\text{vol}_{\mathbb{H}}(\hat{\omega}) \geq \delta$ . Then for any  $\eta > 0$ , there exist a constant  $C(R', \eta, \delta) > 0$ , an exponent  $\alpha(R') \in (0, 1)$ , both independent of  $(\Lambda, \kappa)$ , and a fixed integer l such that

$$
(2.2) \quad ||u||_{L^{2}(\mathcal{R}_{\kappa})}^{2} \leq C(R', \eta) ||u||_{L^{2}(\tilde{\omega})}^{2\alpha}
$$
  
\$\times \left( \sum\_{p,q=0}^{l} ||\partial\_{t}^{p}(-\Delta\_{\mathbb{H}})^{q}w||\_{L^{2}((-T,T)\times\mathcal{R}\_{\kappa}^{n})}^{2} + ||\nabla\_{\mathbb{H}}\partial\_{t}^{p}(-\Delta\_{\mathbb{H}})^{q}w||\_{L^{2}((-T,T)\times\mathcal{R}\_{\kappa}^{n})}^{2} \right)^{1-\alpha}.

Besides, the same bound holds with  $\|\partial_t w\|_{L^2((-\frac{T}{2},\frac{T}{2})\times \mathcal{R}_\kappa)}$  in the left-hand-side.

*Proof.* In this proof only,  $L^2$  indices indicate norms integrated against the Lebesgue measure, while  $L^2_{\mathbb{H}}$  indices denote norms taken against the hyperbolic measure  $d \text{vol}_{\mathbb{H}} = \frac{dx dy}{n^2}$  $\frac{xay}{y^2}$  or the measure  $\frac{dtdxdy}{y^2}$  $\frac{dxdy}{y^2}$  when also integrating over the auxiliary variable t.

We perform the change of variables

$$
(X,Y) = \phi_{\kappa}(x,y) = \kappa^{-1}(x,y)
$$

so that  $(X, Y)$  spans the rectangle  $\mathcal{R}_1 = (-1, 1)_X \times \left( \left( \frac{1}{2} \right)$  $\left(\frac{1}{2}\right)^{R'}$  ,  $\left(\frac{3}{2}\right)$  $\left(\frac{3}{2}\right)^{R'}$  $_Y$  when  $(x, y)$  spans  $\mathcal{R}_{\kappa}$ . This change of coordinates preserves the hyperbolic Laplacian:

$$
\Delta_{\mathbb{H}} = y^2(\partial_x^2 + \partial_y^2) = \kappa^2 Y^2(\kappa^{-2}\partial_x^2 + 2\kappa^{-2}\partial_y^2) = Y^2(\partial_x^2 + \partial_y^2).
$$

We then define  $W^{\kappa}(t, X, Y) = w(t, \phi_{\kappa}^{-1}(X, Y)),$  which satisfies

$$
(\partial_t^2 + Y^2 \Delta_{X,Y}) W^{\kappa}(t, X, Y) = (\partial_t^2 + \Delta_{\mathbb{H}}) w(t, x, y) = 0, \quad t \in \mathbb{R}, (X, Y) \in \mathcal{R}_1.
$$

Multiplying by  $\frac{1}{Y^2}$ , we get

$$
(\partial_t \frac{1}{Y^2} \partial_t + \Delta_{X,Y}) W^{\kappa} = 0 \text{ over } \mathcal{R}_1,
$$

which in turn can be written  $\text{div}(A\nabla_{t,X,Y}W^{\kappa})=0$  for

$$
A(t, X, Y) = \begin{pmatrix} \frac{1}{Y^2} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.
$$

It is clear that for any  $T > 0$ , A is uniformly elliptic with Lipschitz coefficients over the box  $(-T, T)_t \times \mathcal{R}_1 = (-T, T)_t \times (-1, 1)_X \times ((\frac{1}{2})$  $\frac{1}{2}\big)^{\tilde{R}'}$  ,  $\big(\frac{3}{2}\big)$  $\left(\frac{3}{2}\right)^{R'}$  $_Y$ , and its ellipticity and Lipschitz constants are independent of  $T, \kappa$ .

Fix  $0 < T_1 < T_2$ . We denote

(2.3)  $\mathcal{K} = (-T_1, T_1) \times \mathcal{R}_1, \Omega = (-T_2, T_2) \times \mathcal{R}_1^{\eta}, F_{\kappa} = \phi_{\kappa}(\widehat{\omega}), E_{\kappa} = \{0\} \times F_{\kappa}.$ Then by the assumption on  $\widehat{\omega}$ ,

$$
\int_{\widehat{\omega}} \frac{dxdy}{y^2} = \int_{\phi_{\kappa}(\widehat{\omega})} \frac{dXdY}{Y^2} \ge \delta.
$$

Thus,

(2.4) 
$$
\text{vol}_{euc}(\phi_{\kappa}(\widehat{\omega})) \ge \frac{\delta}{4^{R'}},
$$

the right-hand-side of which we still denote δ. By Logunov and Malinnikova's theorem [\[LM18b,](#page-29-4) Theorem 5.1, the statement of which is recalled in Appendix  $A$ , we get

<span id="page-12-0"></span>(2.5) 
$$
\sup_{\mathcal{K}} |\nabla_{t,X,Y} W^{\kappa}| \leq C \sup_{E_{\kappa}} |\nabla_{t,X,Y} W^{\kappa}|^{\alpha} \sup_{\Omega} |\nabla_{t,X,Y} W^{\kappa}|^{1-\alpha}.
$$

The constants  $C > 0, 0 < \alpha < 1$  depend on  $\delta$  but do not depend on u,  $\Lambda$ , or  $\kappa$ .

We change the  $L^{\infty}$  norms to  $L^2$  norms using a trick taken from [\[BM21,](#page-28-0) Section 2]. Assume that  $\nabla W^{\kappa}|\mathcal{K}$  is not identically zero and let

$$
a = \left(\epsilon \frac{\sup_{\mathcal{K}} |\nabla_{t,X,Y} W^{\kappa}|}{\sup_{\Omega} |\nabla_{t,X,Y} W^{\kappa}|^{1-\alpha}}\right)^{\frac{1}{\alpha}}, \quad F_{\kappa}' = \{(X,Y) \in F_{\kappa}, |\nabla_{t,X,Y} W^{\kappa}(t=0,X,Y)| \leq a\}.
$$

Let  $E'_{\kappa} = \{0\} \times F'_{\kappa}$ . One necessarily has  $\mathrm{vol}_{euc}(F'_{\kappa}) \leq \frac{\delta}{2}$  $\frac{0}{2}$ . Indeed, otherwise, the Logunov– Malinnikova estimate [\(2.5\)](#page-12-0) holds with identical constants but replacing  $E_{\kappa}$  with  $E'_{\kappa}$ . This gives

(2.6) 
$$
\sup_{\mathcal{K}} |\nabla_{t,X,Y} W^{\kappa}| \leq C a^{\alpha} \sup_{\Omega} |\nabla_{t,X,Y} W^{\kappa}|^{1-\alpha} \leq C \epsilon \sup_{\mathcal{K}} |\nabla_{t,X,Y} W^{\kappa}|.
$$

For  $C\epsilon$  < 1, we obtain that  $\nabla_{t,X,Y}W^{\kappa}|\kappa$  is identically zero, hence a contradiction.

As a consequence,

$$
(2.7) \int_{F_{\kappa}} |\nabla_{t,X,Y} W^{\kappa}(t=0,X,Y)|^{2} dX dY \geq \int_{F_{\kappa}\setminus F'_{\kappa}} |\nabla W^{\kappa}(t=0,X,Y)|^{2} dX dY
$$
  
\n
$$
\geq a^{2} \frac{\delta}{2} \geq \frac{\delta}{2} \left( \epsilon \frac{\sup_{K} |\nabla_{t,X,Y} W^{\kappa}|}{\sup_{K} |\nabla_{t,X,Y} W^{\kappa}|^{1-\alpha}} \right)^{\frac{2}{\alpha}}.
$$
  
\nSince  $\nabla_{t,X,Y} W^{\kappa}(t=0) = \begin{pmatrix} u \circ \phi_{\kappa}^{-1} \\ 0 \\ \vdots \\ 0 \end{pmatrix}$ , denoting  $U^{\kappa} = u \circ \phi_{\kappa}^{-1}$ , we get  
\n
$$
\int_{\mathcal{R}_{1}} |U^{\kappa}(X,Y)|^{2} dX dY \leq \sup_{t \in (-T_{1},T_{1})} \int_{\mathcal{R}_{1}} |\nabla_{t,X,Y} W^{\kappa}|^{2} dX dY
$$
  
\n
$$
\lesssim \sup_{K} |\nabla_{t,X,Y} W^{\kappa}|^{2}
$$
  
\n
$$
(2.8)
$$
  
\n
$$
\lesssim \left( \int_{F_{\kappa}} |\nabla_{t,X,Y} W^{\kappa}(t=0,X,Y)|^{2} dX dY \right)^{\alpha} \sup_{\Omega} |\nabla_{t,X,Y} W^{\kappa}|^{2-2\alpha}.
$$

<span id="page-13-1"></span>The implicit multiplicative constant in the last inequality depends only on  $\delta$ ,  $\epsilon$ ,  $T_1$ ,  $T_2$ ,  $R'$ ,  $\eta$ and numerical constants, and does not depend on  $\Lambda, u, \kappa$ .

Recall that since y is roughly of size  $\kappa$  over  $\mathcal{R}_{\kappa}$ , we have for any function  $f \in L^2(\mathcal{R}_1)$ ,

<span id="page-13-0"></span>
$$
\int_{\mathcal{R}_1} |f(X,Y)|^2 dX dY = \int_{\mathcal{R}_{\kappa}} \kappa^{-2} |f \circ \phi_{\kappa}(x,y)|^2 dxdy \simeq \int_{\mathcal{R}_{\kappa}} |f \circ \phi_{\kappa}(x,y)|^2 \frac{dxdy}{y^2}.
$$

Applying this to  $f = U^{\kappa}_{\Lambda}$  and  $f = U^{\kappa}_{\Lambda} \mathbb{1}_{\phi_{\kappa}(\widehat{\omega})}$ , we get

(2.9) 
$$
||U^{\kappa}_{\Lambda}||_{L^{2}(\mathcal{R})} \simeq ||\mathfrak{u}||_{L^{2}_{\mathbb{H}}(\mathcal{R}_{\kappa})}, \quad ||U^{\kappa}_{\Lambda}||_{L^{2}(F_{\kappa})} \simeq ||\mathfrak{u}||_{L^{2}_{\mathbb{H}}(\widehat{\omega})}.
$$

Hence,

(2.10) 
$$
\| \mathfrak{u} \|_{L^2_{\mathbb{H}}(\mathcal{R}_\kappa)}^2 \lesssim \| \mathfrak{u} \|_{L^2_{\mathbb{H}}(\widehat{\omega})}^{2\alpha} \sup_{\Omega} |\nabla_{t,X,Y} W^{\kappa}|^{2-2\alpha}.
$$

We now adapt the Sobolev injection process of [\[Rou24,](#page-30-0) Section 2.2], itself inspired by [\[BM23\]](#page-28-1), to bound  $\sup_{\Omega} |\nabla_{t,X,Y} W^{\kappa}|^2$ . Recall that

$$
\Omega = (-T_2, T_2) \times \mathcal{R}_1^{\eta}.
$$

By the Sobolev injection in dimension 3, we have

$$
\sup_{\Omega} |\nabla_{t,X,Y} W^{\kappa}|^2 \leq \|\chi \nabla_{t,X,Y} W^{\kappa}\|_{H^2(\Omega^{(1)})}^2
$$

with  $\chi$  a smooth cutoff function supported in

$$
\Omega^{(1)} = (-T_3, T_3) \times \mathcal{R}_1^{2\eta}
$$

and equal to 1 over  $\Omega$ , for some  $T_3 > T_2$ . We can then apply elliptic regularity (see [\[Eva22,](#page-28-21) Section 6.3]) to the operator  $-(\partial_t^2 + Y^2 \Delta_{X,Y})$ , which is uniformly elliptic over  $\Omega^{(1)}$ , to obtain

$$
(2.11) \quad \|\chi \nabla_{t,X,Y} W^{\kappa}\|_{H^{2}(\Omega^{(1)})}^{2} \lesssim \|(1 - \partial_{t}^{2} - Y^{2} \Delta_{X,Y}) \chi \nabla_{t,X,Y} W^{\kappa}\|_{L^{2}(\Omega^{(1)})}^{2}.
$$

$$
\lesssim \|(1 - \partial_{t}^{2} - Y^{2} \Delta_{X,Y}) \chi \partial_{t} W^{\kappa}\|_{L^{2}(\Omega^{(1)})}^{2} + \|(1 - \partial_{t}^{2} - Y^{2} \Delta_{X,Y}) \chi \nabla_{X,Y} W^{\kappa}\|_{L^{2}(\Omega^{(1)})}^{2}.
$$

The first term is bounded up to some multiplicative constant by

$$
(2.12) \quad \|\partial_t W^{\kappa}\|_{L^2(\Omega^{(1)})}^2 + \|\partial_t^2 W^{\kappa}\|_{L^2(\Omega^{(1)})}^2 + \|\partial_t^3 W^{\kappa}\|_{L^2(\Omega^{(1)})}^2 + \|\nabla_{X,Y}\partial_t W^{\kappa}\|_{L^2(\Omega^{(1)})}^2 + \|(Y^2 \Delta_{X,Y})\partial_t W^{\kappa}\|_{L^2(\Omega^{(1)})}^2
$$

and the second term by

(2.13)

$$
\|\nabla_{X,Y}W^{\kappa}\|_{L^{2}(\Omega^{(1)})}^{2} + \|\nabla_{X,Y}\partial_{t}W^{\kappa}\|_{L^{2}(\Omega^{(1)})}^{2} + \|\nabla_{X,Y}\partial_{t}^{2}W^{\kappa}\|_{L^{2}(\Omega^{(1)})}^{2} + \|Y^{2}\nabla_{X,Y}W^{\kappa}\|_{L^{2}(\Omega^{(1)})}^{2} + \|Y^{2}\nabla_{X,Y}^{2}W^{\kappa}\|_{L^{2}(\Omega^{(1)})}^{2} + \|Y\Delta_{X,Y}W^{\kappa}\|_{L^{2}(\Omega^{(1)})}^{2} + \|\nabla_{X,Y}(Y^{2}\Delta_{X,Y}W^{\kappa})\|_{L^{2}(\Omega^{(1)})}^{2}
$$

where for the last four terms we used the identity

$$
Y^2 \Delta(\chi \nabla f) = Y^2 (\Delta \chi) \nabla f + Y^2 \nabla^2 f(\nabla \chi) - \chi \Delta f \begin{pmatrix} 0 \\ 2Y \end{pmatrix} + \chi \nabla (Y^2 \Delta f).
$$

Here  $\nabla^2 f$  denotes the Hessian matrix of a function f. We again swell  $\Omega^{(1)}$  by  $\eta$  in the  $(X, Y)$ variables to bound the Hessian of  $W^{\kappa}$  (in the  $(X,Y)$  variables) by its Laplacian and lower order terms:

$$
\|\nabla_{X,Y}^2 W^{\kappa}\|_{L^2(\Omega^{(1)})}^2 \lesssim \|W^{\kappa}\|_{L^2(\Omega^{(2)})}^2 + \|\nabla_{X,Y} W^{\kappa}\|_{L^2(\Omega^{(2)})}^2 + \|\Delta_{X,Y} W^{\kappa}\|_{L^2(\Omega^{(2)})}^2
$$

with  $\Omega^{(2)} = (-T_3, T_3) \times \mathcal{R}_1^{3\eta}$  and an implicit multiplicative constant independent of  $(\kappa, \Lambda)$ .

Switching back to coordinates  $(x, y)$  and using  $(2.9)$ , we deduce the following estimates, which allow to conclude the proof:

$$
||Y^{l}\partial_{t}^{m}W^{\kappa}||_{L^{2}(\Omega^{(2)})}^{2} = ||\partial_{t}^{m}w||_{L_{\mathbb{H}}^{2}((-T_{3},T_{3})\times\mathcal{R}_{\kappa}^{3\eta})}^{2}, \qquad (l,m) \in \{0,1,2\}^{2},
$$
  
\n
$$
||Y^{l}\Delta_{X,Y}W^{\kappa}||_{L^{2}(\Omega^{(2)})}^{2} \simeq ||\Delta_{\mathbb{H}}w||_{L_{\mathbb{H}}^{2}((-T_{3},T_{3})\times\mathcal{R}_{\kappa}^{3\eta})}^{2}, \qquad l \in \{1,2\},
$$
  
\n
$$
||\nabla_{X,Y}(Y^{2}\Delta_{X,Y}W^{\kappa})||_{L^{2}(\Omega^{(2)})}^{2} \simeq ||y\nabla_{x,y}(y^{2}\Delta_{x,y}w)||_{L_{\mathbb{H}}^{2}((-T_{3},T_{3})\times\mathcal{R}_{\kappa}^{3\eta})}^{2}
$$
  
\n
$$
\simeq ||\nabla_{\mathbb{H}}\Delta_{\mathbb{H}}w||_{L_{\mathbb{H}}^{2}((-T_{3},T_{3})\times\mathcal{R}_{\kappa}^{3\eta})}^{2}.
$$

Note that in inequality [\(2.8\)](#page-13-1),  $\|\partial_t W^{\kappa}\|_{L^2(\mathcal{K})}^2$  is bounded by the right-hand-side of the second line, so that the inequalities we have proved for  $U^{\kappa}$  (resp.  $\mu$ ) in the rest of the proof also hold for  $\partial_t W^{\kappa}$  (resp.  $\partial_t w$ ). This implies the second part of the result.

<span id="page-14-0"></span>2.2. Propagation of smallness in a funnel. This subsection and the next are dedicated to proving that the propagation of smallness result of Lemma [2.1](#page-11-3) extends to funnels and cusps as long as  $\omega$  satisfies the thickness condition [\(1.1\)](#page-2-3). In the simpler case of a funnel, the proof consists in covering the funnel (up to a compact remainder) with rectangles  $\mathcal{R}_{\kappa}$ . The rectangles may spill on a finite number of images of the funnel by elements of its corresponding Fuchsian group, as depicted in Figure [3.](#page-16-0)

The following lemma is easily checked using that  $B_{(x,y)}(R) \subset \mathbb{H}$  is the Euclidean ball with center  $(x, y \cosh(R))$  and radius  $y \sinh(R)$ :

<span id="page-15-2"></span>**Lemma 2.2.** – Consider R' defined by the one-to-one relation

<span id="page-15-3"></span>(2.14) 
$$
\tanh(R) = \min\left(1 - 2^{-R'}, \left(\frac{3}{2}\right)^{R'} - 1\right),
$$

Let  $j, k \in \mathbb{Z}$  be two integers and denote (2.15)

$$
\mathcal{R}_{j,k} = \mathcal{R}_{j,k}(R') := (2^{R'j}k - 2^{R'j}, 2^{R'j}k + 2^{R'j})_x \times (2^{R'(j-1)}, 3^{R'} \times 2^{R'(j-1)})_y = \mathcal{R}_{2^{R'j}} + 2^{R'j} \begin{pmatrix} k \\ 0 \end{pmatrix},
$$

a horizontal translation of  $\mathcal{R}_{2R'j}$ . Then

$$
(2.16) \t\t Bz(R) \subset \mathcal{R}_{j,k}
$$

where  $z = \left(2^{R'j}k, \frac{2^{R'j}}{\cosh(j)}\right)$  $\cosh(R)$  $\big)$ . In particular, by [\(Thick\)](#page-2-0),  $vol(\omega \cap \mathcal{R}_{i,k}) \geq vol(\omega \cap B_z(R)) \geq \delta vol(B_z(R)).$ 

Since the volume of any ball in a funnel is bounded from below independently of  $z$ , we can make the assumption that the intersection of  $\omega$  with each rectangle  $\mathcal{R}_{j,k}$  has measure bounded from below by some constant which we still denote  $\delta$ . Since there is a finite number of funnel ends in the surface M, this  $\delta$  can still be chosen independent of the funnel.

Consider a funnel F and the domain isometric to F that is symmetric with respect to the y-axis, delimited by three geodesic arcs and the x-axis (see the colored domain in Figure [3\)](#page-16-0). In the proof of Theorem [4,](#page-15-0) we will identify  $F$  and this fundamental domain. The images of the funnel by its corresponding hyperbolic element of  $PSL_2(\mathbb{R})$  (which has fixed points A and  $B \in \{y=0\}$  and preserves the geodesic arc  $\widehat{AB}$ ) are depicted as the white domains of Figure [3.](#page-16-0) We now prove the following result

<span id="page-15-0"></span>**Theorem 4** (Propagation of smallness in a funnel). – Let  $D > 0$ . There exists  $\tilde{F}$  some complement in  $F$  of a relatively compact set of  $M$  such that the following inequality holds for some constant  $C > 0$  and any  $D > 0$ :

<span id="page-15-1"></span>
$$
(2.17) \quad ||\partial_t v_\Lambda||^2_{L^2((-\frac{T}{2},\frac{T}{2})\times\tilde{F})} \le C(R',\delta) \Bigg(e^{\frac{2-2\alpha}{2\alpha}D\Lambda} ||\Pi_\Lambda u||^2_{L^2(F\cap\omega)}\n+ e^{-D\Lambda} \sum_{p,q=0}^l \Big( ||\partial_t^p(-\Delta_g)^qv_\Lambda||^2_{L^2((-T,T)\times F)} + ||\nabla_g\partial_t^p(-\Delta_g)^qv_\Lambda||^2_{L^2((-T,T)\times F)}\Big)\Bigg)
$$

Note that all the compact remainder sets that are not dealt with in the left-hand-side of  $(2.17)$  will be tackled in subsection [2.4](#page-20-0) when studying propagation of smallness from the ends to the compact core.

*Proof.* Denote by  $\gamma \in PSL_2(\mathbb{R})$  the isometry of  $\mathbb H$  for which F (or rather, its projection if  $\mathbb H$ as depicted in Figure [3\)](#page-16-0) is a fundamental domain.

We fix some  $j_0 \in \mathbb{Z}$  such that for every  $j \leq j_0$ , the projection of the rectangle  $\mathcal{R}_{2R'j}$  onto F does not self-overlap. Such a  $j_0$  exists as in a funnel, the local injectivity radius at some point  $(x, y)$  tends to infinity as y tends to zero. We fix

$$
\tilde{F} = \{ (x, y) \in F, y < 3^{R'} \times 2^{R'(j-1)} \}.
$$



<span id="page-16-0"></span>Figure 3. The fundamental domain of a funnel (in orange), see Figure [1.](#page-4-1) Apart from a compact neighborhood of its geodesic boundary, we cover the funnel with rectangle rectangle horizontal translations of some  $\mathcal{R}_{\kappa}$  for various  $\kappa$  (in purple), adapted to the thickness condition at scale R and the hyperbolic scaling. For better legibility, we have drawn the rectangles as not overlapping each other, though each of them overlaps a finite, bounded uniformly number of others.

In Figure [3,](#page-16-0)  $\tilde{F}$  is the intersection of F (identified with the colored domain) with the surface spanned by the purple rectangles.

We then choose a family of rectangles  $(\mathcal{R}_{j,k})_{j\leq j_0,|k|\leq k_j}$ , with the  $k_j$  large enough to ensure that

$$
\tilde{F} \subset \bigcup_{j \leq j_0} \bigcup_{|k| \leq k_j} \mathcal{R}_{j,k},
$$

and small enough to guarantee that  $\bigcup_{j\leq j_0}\bigcup_{|k|\leq k_j}\mathcal{R}_{j,k}$  is contained in  $F\cup \gamma F\cup \gamma^{-1}F$ . This is the situation depicted in Figure  $3$  (the union of all rectangles is a subset of  $F$  and the two fundamental domains for  $\gamma$  that neighbor it). Such a family of rectangles exists as long as  $j_0$ is chosen small enough.

We lift  $\Pi_{\Lambda}u|_F$  to the domain of  $\mathbb H$  delimited by the x-axis and the geodesic arc  $\widehat{AB}$ , setting  $\Pi_{\Lambda}u(\gamma z) = \Pi_{\Lambda}u(z)$ . We still denote this extension by  $\Pi_{\Lambda}u$  in order not to multiply notation. We apply the same procedure to its harmonic extension  $v_\Lambda$ . Since the lifted  $\Pi_\Lambda u$  is  $\gamma$ -invariant, the lifted  $v_{\Lambda}$  remains harmonic with respect to  $\partial_t^2 + \Delta_{\mathbb{H}}$  in each of the rectangles  $\mathcal{R}_{j,k}$ .

One can also check that each set  $\mathcal{R}_{j,k}$  intersected with  $\omega \cup \gamma \omega \cup \gamma^{-1} \omega$  has measure at least  $\delta$ . Since every  $\mathcal{R}_{j,k}$  is a horizontal translation of  $\mathcal{R}_{2R'j}$  and the hyperbolic metric in invariant by horizontal translation, Lemma [2.1](#page-11-3) applies to  $(w, \mathfrak{u}) = (v_{\Lambda}, \Pi_{\Lambda}u)$  in each rectangle and Young's inequality  $a^{2\alpha}b^{2-2\alpha} \leq C(a^2+b^2)$  yields

$$
\begin{split} \|\partial_t v\|^2_{L^2((-\frac{T}{2},\frac{T}{2})\times \tilde{F})}&\leq \sum_{j\leq j_0}\sum_{|k|\leq k_j}\|\partial_t v\|^2_{L^2((-\frac{T}{2},\frac{T}{2})\times \mathcal{R}_{j,k})}\\ &\lesssim \sum_{j,k}\|\Pi_{\Lambda} u\|_{L^2(\mathcal{R}_{j,k}\cap (\omega\cup \gamma \omega\cup \gamma^{-1}\omega))}\times \Bigg(\sum_{p,q=0}^l \Big(\|\partial_t^p (-\Delta_g)^qv_\Lambda\|^2_{L^2((-T,T)\times \mathcal{R}_{j,k}^\eta)}\Big)\\ &\quad+\|\nabla_g\partial_t^p (-\Delta_g)^qv_\Lambda\|^2_{L^2((-T,T)\times \mathcal{R}_{j,k}^\eta)}\Big)\Bigg)^{1-\alpha}\\ &\lesssim \sum_{j,k}\Big(e^{\frac{2-2\alpha}{2\alpha}D\Lambda}\|\Pi_{\Lambda} u\|^2_{L^2(\mathcal{R}_{j,k}\cap (\omega\cup \gamma \omega\cup \gamma^{-1}\omega))}\\ &\quad+e^{-D\Lambda}\|\nabla_g\partial_t^p (-\Delta_g)^qv_\Lambda\|^2_{L^2((-T,T)\times \mathcal{R}_{j,k}^\eta)}\Bigg)+e^{-D\Lambda}\|\nabla_g\partial_t^p (-\Delta_g)^qq v_\Lambda\|^2_{L^2((-T,T)\times \mathcal{R}_{j,k}^\eta)}\Bigg)\,. \end{split}
$$

Here,  $\mathcal{R}_{j,k}^{\eta}$  denotes the rectangle

$$
\mathcal{R}_{2^{R'j}}^{\eta}+2^{R'j}\begin{pmatrix}k\\0\end{pmatrix}.
$$

For  $j_0$  and  $\eta$  small enough, we have

$$
\bigcup_{j\leq j_0,|k|\leq k_j} \mathcal{R}_{j,k}^{\eta}\subset (F\cup \gamma F\cup \gamma^{-1}F)\ .
$$

The result then derives from this property, the  $\gamma$ -invariance of the  $L^2$  norm, and the fact that each point is covered by a finite, uniformly bounded number of rectangles  $\mathcal{R}_{j,k}^{\eta}$ .

### <span id="page-17-0"></span>2.3. Propagation of smallness in a cusp. Recall that a given cusp  $\mathcal C$  is isometric to

$$
\left(\mathbb{S}_{\theta}^{1}\times (a,+\infty)_{y},\frac{d\theta^{2}+dy^{2}}{y^{2}}\right)
$$

for some  $a > 0$  and  $\mathcal{S}_{\theta}^1 = \mathbb{R}/\mathbb{Z}$ . The strategy to obtain a propagation of smallness result on this cusp is similar to that developed on funnels, with the additional difficulty that the volume of balls is not bounded from below. Specifically, the volume of some ball  $B_{(\theta,y)}(R)$  is of order 1  $\frac{1}{y}$  (times an R-dependent multiplicative constant) when y goes to infinity. Thus, the thickness condition in a cusp becomes

$$
\text{vol}(B_{(\theta,y)}(R) \cap \omega) \ge \frac{\delta}{y}
$$

for some positive  $\delta$ . To recover a uniform lower bound on the measure of the 2-dimensional set and uniform constants in the Logunov–Malinnikova estimate, we thus need to periodically lift the cusp  $\mathcal{C}$ , and to consider rectangles covering a number of periods that depends on their y coordinate.



<span id="page-18-0"></span>FIGURE 4. A rectangle  $\tilde{\mathcal{R}}_j$  in a cusp, its periodic extension  $\mathcal{R}_{[2^{R'j}]}$ , and their enlargements by  $\eta$ . See Figure [1](#page-4-1) and Section [1.1.2](#page-3-0) for explanations on the representation of a cusp in  $\mathbb{H}^2$ .

<span id="page-18-2"></span>**Theorem 5.** – There exists  $\tilde{C}$  some complement in C of a relatively compact set such that the following inequality holds for some constant  $C > 0$  and any  $D > 0$ :

<span id="page-18-1"></span>
$$
(2.18)\quad \|\partial_t v_\Lambda\|_{L^2((-\frac{T}{2},\frac{T}{2})\times\tilde{C})}^2 \le C(R',\delta) \left(e^{\frac{2-2\alpha}{2\alpha}D\Lambda} \|\Pi_\Lambda u\|_{L^2(\mathcal{C}\cap\omega)}^2 + e^{-D\Lambda} \sum_{p,q=0}^l \left(\|\partial_t^p(-\Delta_g)^qv_\Lambda\|_{L^2((-T,T)\times\mathcal{C})}^2 + \|\nabla_g\partial_t^p(-\Delta_g)^qv_\Lambda\|_{L^2((-T,T)\times\mathcal{C})}^2\right)\right)
$$

Proof. The main difference with the proof for funnels is that the projection of a rectangle  $\mathcal{R}_{j,k}$  onto the cusp self-overlaps for large j, specifically for any j such that  $2^{R'j+1} > 1$ . Fix  $j_0 \geq \lceil \log_2\left(-\frac{1}{R'}\right) \rceil$  and denote for  $j \geq j_0$ 

(2.19) 
$$
\tilde{\mathcal{R}}_j = \pi_{\mathcal{C}}\left(\mathcal{R}_{\lceil 2^{jR'} \rceil}\right) = \mathbb{S}_{\theta} \times \left(\lceil 2^{jR'} \rceil 2^{-R'}, \lceil 2^{jR'} \rceil (3/2)^{R'}\right)_y,
$$

$$
\pi_{\mathcal{C}} : (x, y) \in \mathbb{H} \mapsto (x \mod 1, y) \in \mathcal{C}.
$$

 $j_0$  is defined so that the  $\tilde{\mathcal{R}}_j$  are horizontal bands inside the cusp  $\mathcal{C}$ . Like before, we also denote for  $\eta > 0$ ,

$$
\tilde{\mathcal{R}}_j^{\eta} = \mathbb{S}_{\theta} \times \left( (1 - \eta) \lceil 2^{jR'} \rceil 2^{-R'}, (1 + \eta) \lceil 2^{jR'} \rceil (3/2)^{R'} \right)_y.
$$

Besides, it is possible to increase  $j_0$  so that the  $(\tilde{\mathcal{R}}_j^{\eta}$  $\overline{ }$ overlap vertically a finite, bounded  $j \geq j_0$  $(\eta$ -dependent) number of times. From now on, we assume  $j_0$  satisfies this property and denote

<span id="page-19-0"></span>
$$
\tilde{\mathcal{C}} = \mathcal{S}_{\theta}^1 \times (2^{R'j_0-1}, +\infty) = \bigcup_{j \ge j_0} \tilde{\mathcal{R}}_j,
$$

the complement in  $\mathcal C$  of which is a relatively compact set. There hold the norm inequalities

$$
(2.20) \t\t\t || \cdot ||_{L^2(\tilde{C})}^2 \simeq \sum_{j \ge j_0} || \cdot ||_{L^2(\tilde{\mathcal{R}}_j)}^2 \lesssim \sum_{j \ge j_0} || \cdot ||_{L^2(\tilde{\mathcal{R}}_j^{\eta})}^2 \lesssim_{\eta} ||.||_{L^2(C)}^2.
$$

We now lift  $\mathcal{C} \simeq \mathbb{S}_{\theta}^1 \times (a, +\infty)_y$  periodically to a subset  $\mathbb{R}_x \times (a, +\infty)_y$  of the hyperbolic halfplane H. We extend periodically in  $\theta$  the metric g, the function  $\Pi_{\Lambda} u$ , its harmonic extension  $v_\Lambda$  and the thick set  $\omega$ , and will denote the extensions  $g = g_{\parallel\parallel} = \frac{dx^2 + dy^2}{u^2}$  $\frac{1+ay^2}{y^2}, \underline{\Pi_\Lambda u}, \underline{v_\Lambda}, \underline{\omega}.$ 

By the thickness property, the measure of  $\omega$  inside each band  $\tilde{\mathcal{R}}_j$  of the cusp is bounded from below with decaying bound:

$$
\mathrm{vol}_g(\omega \cap \tilde{\mathcal{R}}_j) \gtrsim_{\delta} \frac{1}{\lceil 2^{R'j} \rceil}.
$$

Indeed, one can check similarly to Lemma [2.2](#page-15-2) that

$$
B_{\left(0,\frac{\lceil 2^{R^\prime}j\rceil}{\cosh(R)}\right)}(R)\subset\tilde{\mathcal{R}}_j
$$

where  $R'$  is defined by  $(2.14)$ . Thus,

$$
\mathcal{R}_{\lceil 2^{R'j} \rceil} = \lceil 2^{R'j} \rceil \left[ (-1,1) \times \left( \left( \frac{1}{2} \right)^{R'}, \left( \frac{3}{2} \right)^{R'} \right) \right]
$$

contains exactly  $\lceil 2^{R'j} \rceil$  copies of  $\omega \cap \tilde{\mathcal{R}}_j$ , and  $\text{vol}_{g_{\mathbb{H}}} \left( \underline{\omega} \cap \mathcal{R}_{\lceil 2^{R'j} \rceil} \right)$  is bounded from below uniformly in  $j$ . This procedure is illustrated in Figure  $\hat{4}$ .

We now apply Lemma [2.1](#page-11-3) for rectangles  $\mathcal{R}_{\lceil 2^{R'j} \rceil}$  and the set  $\underline{\omega}$ , yielding

$$
(2.21) \quad \|\partial_t \underline{v_\Lambda}\|_{L^2\left((-\frac{T}{2},\frac{T}{2})\times \mathcal{R}_{\lceil 2^{R^\prime} j\rceil}\right)}^2 \lesssim \|\underline{\Pi_\Lambda u}\|_{L^2\left(\underline{\omega} \cap \mathcal{R}_{\lceil 2^{R^\prime} j\rceil}\right)}^{2\alpha} \times \left(\sum_{p,q=0}^l \|\partial_t^p (-\Delta_{\mathbb{H}})^q \underline{v_\Lambda}\|_{L^2\left((-T,T)\times \mathcal{R}_{\lceil 2^{R^\prime} j\rceil}^{\eta}\right)}^2 + \|\nabla_{\mathbb{H}} \partial_t^p (-\Delta_{\mathbb{H}})^q \underline{v_\Lambda}\|_{L^2\left((-T,T)\times \mathcal{R}_{\lceil 2^{R^\prime} j\rceil}^{\eta}\right)}^2\right)^{1-\alpha}.
$$

By periodicity and fixing  $\eta \ll 1$ , it is possible to bound the left-hand-side from below and right-hand-side from above to get

$$
(2.22) \quad 2 \times \lceil 2^{R'j} \rceil ||\partial_t v_{\Lambda}||_{L^2\left((- \frac{T}{2}, \frac{T}{2}) \times \tilde{\mathcal{R}}_j\right)}^2 \lesssim (2\lceil 2^{R'j} \rceil)^\alpha \lVert \Pi_{\Lambda} u \rVert_{L^2(\omega \cap \tilde{\mathcal{R}}_j)}^{2\alpha} \times (4\lceil 2^{R'j} \rceil)^{1-\alpha} \times \left( \sum_{p,q=0}^l ||\partial_t^p (-\Delta_g)^q v_{\Lambda}||_{L^2((- T,T) \times \tilde{\mathcal{R}}_j^n)}^2 + ||\nabla_g \partial_t^p (-\Delta_g)^q v_{\Lambda}||_{L^2((- T,T) \times \tilde{\mathcal{R}}_j^n)}^2 \right)^{1-\alpha}.
$$

The  $[2^{R'j}]$  constants simplify, hence we get a bound for  $\|\partial_t v_\Lambda\|_{L^2((-T,T)\times \tilde{\mathcal{R}}_j)}^2$  with jindependent constant. The rest of the proof consists in applying Young's inequality, summing over j, and using bound  $(2.20)$  to derive  $(2.18)$ , similarly to the case of funnels.

<span id="page-20-0"></span>2.4. General surface with hyperbolic ends. Carleman on the compact core  $+$  good decomposition  $+$  sections 2.2, 2.3.

The last preliminary propagation of smallness result needed to conclude the proof consists in bounding the harmonic extension  $\partial_v v_\Lambda$  on the compact core of the manifold.

As noted in the introduction, the thickness assumption on  $\omega$  is unchanged when removing a relatively compact subset from  $\omega$ , up to increasing the thickness scale R. Thus, the intersection of the compact core of the manifold (in which we do not assume hyperbolicity properties) with the sensor set is allowed to be empty under our assumptions. What we show is propagation of smallness from an end (in which the sensor set is active) to the compact core. A physical interpretation of the following result is that to control the heat equation on a non-compact manifold, one only need to control what happens at infinity.

<span id="page-20-1"></span>**Theorem 6** (Propagation of smallness in the ends is sufficient).  $-$  Let U, V be open subsets of a smooth manifold M with  $U \subset\subset V$ . Let X be an open subset of U, then for any  $T > 0$ , there exist some constant  $C(U, V, X, T) > 0$  and some  $\theta \in (0, 1)$  such that

(2.23) k∂tvΛkL2((<sup>−</sup> <sup>T</sup> 2 , T 2 )×U) ≤ Ck∂tvΛk θ H<sup>1</sup> t,z((−T,T)×V ) kΠΛuk 1−θ L2(X) .

This result is inspired by Theorem 9.1 of [\[LRLR22b\]](#page-29-18), but the authors obtain a bound by the  $L^2$  norm of the full harmonic extension of  $\Pi_{\Lambda}u$  ( $\partial_t v_{\Lambda}$  in our case) rather than  $\Pi_{\Lambda}u$  itself. We combine their statement with an adaptation of a Jerison and Lebeau's propagation of smallness results from the boundary [\[JL99,](#page-29-10) Lemma 14.5] to our setting.

*Proof of Theorem [6.](#page-20-1)* The first step of the proof is to propagate smallness for the harmonic extension  $\partial_t v_\Lambda(t, z)$  from a small enough chart inside X to U (times appropriate intervals in the t variable). This is a standard procedure in Carleman inequality-based propagation of smallness estimates. A general statement, proofs and generalizations can be found in [\[LRLR22b,](#page-29-18) Chapter 9]. We state a version of their result suited to our setting in Theorem [8](#page-27-4) of Appendix [A.](#page-26-0) Set  $m = \frac{\text{dist}(U, V^c)}{3} > 0$ , which is positive by the relative compactness property of U. Applying Theorem [8](#page-27-4) with  $R_1 = 0, f = 0, U = \left(-\frac{7}{2}\right)$  $\frac{T}{2},\frac{T}{2}$  $\frac{T}{2}$   $\times$  U,  $\mathcal{V} = (-T, T) \times U_m$ , with  $T > 0$ ,  $U_m = \bigcup_{z \in U} B_z(m)$ , and the solution  $w = \partial_t v_{\Lambda}$  of

$$
(\partial_t^2 + \Delta_g)w = 0,
$$

we get

<span id="page-20-3"></span>
$$
(2.24) \t\t ||\partial_t v_\Lambda||_{L^2((-\frac{T}{2},\frac{T}{2})\times U)}^2 \leq C ||\partial_t v_\Lambda||_{H^1_{t,z}((-T,T)\times U_m)}^{2-2\beta} ||\partial_t v_\Lambda||_{L^2((-T,T)\times X)}^{2\beta}.
$$

The second step is to apply Jerison and Lebeau's propagation of smallness from the bound-ary (see Theorem [9](#page-27-5) of [A](#page-26-0)ppendix A for statement and comments) to bound  $\|\partial_t v_\Lambda\|_{L^2((-\frac{T}{2},\frac{T}{2})\times X)}$ by  $\|\Pi_{\Lambda}u\|_{L^2(X)}$ . We do so with  $F = v_{\Lambda}$ , on both sides of the  $\{t = 0\}$  surface, with  $B$  a ball such that  $B \cap \{t = 0\} = \{t = 0\} \times X$ . We get

<span id="page-20-2"></span>
$$
(2.25) \t\t ||\partial_t v_\Lambda||^2_{L^2((-\frac{T}{2},\frac{T}{2})\times X)} \leq C ||\partial_t v_\Lambda(t=0)||^{2\gamma}_{L^2(X_m)} ||v_\Lambda||^{2-2\gamma}_{H^1_{t,z}((-T,T)\times V)}
$$

The result then follows from injecting  $(2.25)$  in  $(2.24)$  and shrinking X, m appropriately. The exponent  $\theta$  of the theorem corresponds to  $\beta \gamma \in (0, 1)$ .

.

From there on, the proof consists in summing all the propagation of smallness estimates in patches (the compact core, funnels and cusps) and deriving the spectral estimate from Young and functional calculus inequalities, in the spirit of [\[BM21\]](#page-28-0) and [\[Rou24\]](#page-30-0).

We apply Theorem [6](#page-20-1) with  $U = \tilde{K}$  defined in the introduction,  $V = M$ , and X an open set that is contained in the intersection of  $\tilde{K}$  with one of the ends  $\tilde{F}$  or  $\tilde{C}$  defined in Sections [2.2,](#page-14-0) [2.3.](#page-17-0) That end will be denoted  $P_0$ . In what follows,  $P$  (for patch) will denote either a funnel F or a cusp C and  $\tilde{P}$  will denote the domains  $\tilde{F}$ ,  $\tilde{C}$  of Theorems [4](#page-15-0) and [5.](#page-18-2) This yields

$$
\|\partial_t v_\Lambda\|_{L^2((-\frac{T}{2},\frac{T}{2})\times\tilde{K})}^2 \le \|\Pi_\Lambda u\|_{L^2(P_0)}^{2\gamma} \|\partial_t v_\Lambda\|_{H^1_{t,z}((-T,T)\times M)}^{2-2\gamma}
$$
  

$$
\lesssim e^{\frac{2-2\gamma}{2\gamma}D\Lambda} \|\Pi_\Lambda u\|_{L^2(P_0)}^2 + e^{-D\Lambda} \|\partial_t v_\Lambda\|_{H^1_{t,z}((-T,T)\times M)}^2.
$$

<span id="page-21-0"></span>Recall that, by Theorems  $4$  and  $5$ , the following estimate holds on any funnel or cusp  $P$ :

<span id="page-21-1"></span>
$$
(2.27) \quad ||\partial_t v_\Lambda||_{L^2((-\frac{T}{2},\frac{T}{2})\times \tilde{P})}^2 \lesssim_{R,\delta} \left(e^{\frac{2-2\alpha}{2\alpha}D\Lambda} ||\Pi_\Lambda u||_{L^2(P\cap\omega)}^2 + e^{-D\lambda} \sum_{p,q=0}^l \left( ||\partial_t^p (-\Delta_g)^q v_\Lambda||_{L^2((-T,T)\times P)}^2 + ||\nabla_g \partial_t^p (-\Delta_g)^q v_\Lambda||_{L^2((-T,T)\times P)}^2 \right) \right).
$$

By the statement of Lemma  $2.1$  with  $\mu$  in the left-hand-side, this inequality also holds when replacing  $\|\partial_t v_\Lambda\|_L^2$  $L^2((-\frac{T}{2},\frac{T}{2})\times\tilde{P})$  by  $\|\Pi_\Lambda u\|_{L^2(\tilde{P})}^2$ .

In inequalities  $(2.26)$  and  $(2.27)$ , D is a fixed, large positive constant which will be specified at the very end of the proof. We now sum  $(2.26)$  and  $(2.27)$  over all funnels and cusps P to obtain [\(Spec\)](#page-1-2). First of all, by positivity of the projector-valued measure, we have for all  $t \in \mathbb{R}$ 

<span id="page-21-3"></span>
$$
\|\Pi_\Lambda u\|_{L^2(M)}^2 = \int_0^\Lambda \langle dm_\lambda u, u \rangle \le \int_0^\Lambda \cosh(\lambda t)^2 \langle dm_\lambda u, u \rangle = \|\partial_t v_\Lambda(t,.)\|_{L^2(M)}^2.
$$

Integrating over  $t \in \left(-\frac{7}{2}\right)$  $\frac{T}{2},\frac{T}{2}$  $(\frac{1}{2})$  gives

(2.28) 
$$
\|\Pi_{\Lambda} u\|_{L^2(M)} \lesssim_T \|\partial_t v_{\Lambda}\|_{L^2((-\frac{T}{2},\frac{T}{2})\times M)}
$$

We now decompose  $\|\partial_t v_\Lambda\|_{L^2(M)}^2$  as the sum over funnels, cusps and the compact core. Note that funnels and cusps are disjoint from one another, so that

.

$$
\|\partial_t v_\Lambda\|_{L^2\left((-\frac{T}{2},\frac{T}{2})\times M\right)}^2\leq \left(\sum_{\text{end }P}\|\partial_t v_\Lambda\|_{L^2\left((-\frac{T}{2},\frac{T}{2})\times \tilde{P}\right)}^2\right)+\|\partial_t v_\Lambda\|_{L^2\left((-\frac{T}{2},\frac{T}{2})\times \tilde{K}\right)}^2.
$$

Using  $(2.26)$ , we bound the norm over the compact core:

$$
\begin{aligned} \left\|\partial_t v_\Lambda\right\|_{L^2\left((-\frac{T}{2},\frac{T}{2})\times M\right)}^2 &\lesssim \left(\sum_{\text{end }P}\left\|\partial_t v_\Lambda\right\|_{L^2\left((-\frac{T}{2},\frac{T}{2})\times \tilde{P}\right)}^2\right) + e^{-D\Lambda} \|\partial_t v_\Lambda\|_{H_{t,z}^1((-T,T)\times M)}^2 \\ &+ e^{\frac{2-2\gamma}{2\gamma}D\Lambda} \|\Pi_\Lambda u\|_{L^2\left((-\frac{T}{2},\frac{T}{2})\times P_0\right)} . \end{aligned}
$$

We apply propagation of smallness in funnels and cusps as given by  $(2.27)$  to bound the sum by  $L^2$  norms on the entire manifold: we get for some constant  $K > 0$  independent of D

<span id="page-21-2"></span>
$$
(2.29) \quad ||\partial_t v_{\Lambda}||^2_{L^2((-\frac{T}{2},\frac{T}{2})\times M)} \lesssim e^{K D\Lambda} ||\Pi_{\Lambda} u||_{L^2(\omega)} + e^{-D\Lambda} \sum_{p,q=0}^l \left( ||\partial_t^p (-\Delta_g)^q v_{\Lambda}||^2_{L^2((-T,T)\times M)} + ||\nabla_g \partial_t^p (-\Delta_g)^q v_{\Lambda}||^2_{L^2((-T,T)\times M)} \right).
$$

The term  $\|\partial_t v_\Lambda\|_{H^1_{t,z}((-T,T)\times M)}^2$  is absorbed in the sum.

We now bound the gradient terms: for a given function  $h \in D(-\Delta_q)$ . Ignoring the time integral for now, one has for fixed  $t$ 

<span id="page-22-1"></span>
$$
\|\nabla_g h\|_{L^2(M)}^2 = \langle h, -\Delta_g h \rangle_{L^2(M)} \le \frac{1}{2} \left( \|h\|_{L^2(M)}^2 + \|\nabla_g h\|_{L^2(M)}^2 \right).
$$

Thus, the sum in [\(2.29\)](#page-21-2) can be replaced by

(2.30) 
$$
\sum_{p,q=0}^{l+1} \left( \|\partial_t^p (-\Delta_g)^q v_\Lambda\|_{L^2((-T,T)\times M)}^2 \right).
$$

This step simplifies the method developed in [\[Rou24\]](#page-30-0), in which gradient term bounds were performed locally by defining appropriate cutoffs in charts.

Recall that

(2.31) 
$$
\partial_t^p ((-\Delta_g)^q v_\Lambda) = \int_0^\Lambda \lambda^{p+2q-1} \sinh^{(p)}(\lambda t) dm_\lambda u, \quad p, q \in \mathbb{N}.
$$

Thus, using the functional calculus bound [\(1.3\)](#page-1-3) and the bound

$$
\Lambda^{3l+2}(\sinh(\Lambda T) + \cosh(\Lambda T)) \le e^{K_0 \Lambda}
$$

for some sufficiently large  $K_0$ , we get

$$
\sum_{p,q=0}^{l+1} \left( \|\partial_t^p (-\Delta_g)^q v_\Lambda\|_{L^2((-T,T)\times M)}^2 \right) \le e^{K_1 \Lambda} \|\Pi_\Lambda u\|_{L^2(M)}^2, \quad K_1 > 0.
$$

Using  $(2.28)$ ,  $(2.29)$  and  $(2.30)$ , we obtain

$$
(2.32) \t\t ||\Pi_{\Lambda}u||_{L^{2}(M)}^{2} \leq C(e^{K D\Lambda} \|\Pi_{\Lambda}u\|_{L^{2}(\omega)}^{2} + e^{(K_{1}-D)\Lambda} \|\Pi_{\Lambda}u\|_{L^{2}(M)}^{2}).
$$

Fixing  $D = 2K_1$ , we obtain the desired spectral inequality for  $\Lambda$  larger than some  $\Lambda_0$  such that  $Ce^{-K_1\Lambda_0} = \frac{1}{2}$  $\frac{1}{2}$ . For  $\Lambda \in [0, \Lambda_0]$ , we apply the result to  $\Pi_{\Lambda_0} \Pi_{\Lambda} u = \Pi_{\Lambda} u$ . This ends the proof of Theorem [1.](#page-2-1)

### 3. Spectral estimates imply thickness

<span id="page-22-0"></span>In this section, we prove that thickness is necessary to spectral estimates on manifolds with Ricci curvature bounded from below. Theorem [2](#page-2-2) generalizes the necessity of Logvinenko and Sereda's thickness condition [\[LS74\]](#page-30-2) (see for example [\[HJ95,](#page-29-20) Chapter 3.4]). The proof is a generalization of those of [\[WWZZ19,](#page-30-7) Lemma 2.5] and [\[Rou24,](#page-30-0) Section 2.3]. It is based on the fact that spectral estimates imply an observability inequality for solutions of the heat equation. The idea is then to test this observability inequality with heat kernels. The Ricci curvature lower bound implies Gaussian estimates from above and below for the heat kernel as well as estimates on the volume of balls, both of which are used in the proof.

We start with a few key inequalities on manifolds with Ricci curvature bounded from below. The first one is a volume doubling property which is a consequence of [\[CGT82\]](#page-28-2) (see also [\[Heb96,](#page-29-21) Theorem 1.5], [\[GHL04,](#page-28-3) Theorem 4.19]):

**Lemma 3.1** (Volume doubling). – There exists a positive constant  $C_D(n, K) > 0$  such that for any  $z \in M$  and every  $r > 0$ , one has

<span id="page-22-2"></span>(3.1) 
$$
\text{vol}(B_z(2r)) \leq C_D e^{C_D r} \text{vol}(B_z(r)).
$$

Iterating [\(3.1\)](#page-22-2) gives for any  $n \in \mathbb{N}$ :

(3.2) 
$$
\text{vol}(B_z(2^n r)) \le (C_D)^n \exp(C_D(2^n - 1)r) \text{ vol}(B_z(r)).
$$

<span id="page-23-1"></span><span id="page-23-0"></span>The following is a consequence of the volume doubling estimate:

**Lemma 3.2** (Quotient of Gaussian integrals). – Consider two positive numbers  $\alpha, \beta$ . Then

(3.3) 
$$
\inf_{z_0 \in M} \frac{\int_M e^{-\alpha d(z, z_0)^2} d\text{vol}(z)}{\int_M e^{-\beta d(z, z_0)^2} d\text{vol}(z)} > 0.
$$

Proof of Lemma [3.2.](#page-23-0) The idea of the proof is to compare the integral of a Gaussian centered around a point  $z_0$  with the volume of the ball  $B_{z_0}(1)$ . We start by bounding the numerator from below:

<span id="page-23-4"></span>
$$
\int_M e^{-\alpha d(z,z_0)^2} d\operatorname{vol}(z) \ge e^{-\alpha} \operatorname{vol}(B_{z_0}(1)).
$$

We now bound the denominator from above by decomposing the integral on a sum of integrals over annuli and using the volume dilation estimate  $(3.2)$ :

$$
\int_{M} e^{-\beta d(z,z_0)^2} d\operatorname{vol}(z)
$$
\n
$$
\leq \int_{B_{z_0}(1)} e^{-\beta d(z,z_0)^2} d\operatorname{vol}(z) + \sum_{n=0}^{+\infty} \int_{B_{z_0}(2^{n+1}) \setminus B_{z_0}(2^n)} e^{-\beta d(z,z_0)^2} d\operatorname{vol}(z)
$$
\n
$$
\leq \operatorname{vol}(B_{z_0}(1)) + \sum_{n=0}^{+\infty} e^{-\beta 4^n} \operatorname{vol}(B_{z_0}(2^{n+1}))
$$
\n
$$
\leq \operatorname{vol}(B_{z_0}(1)) + \left(\sum_{n=0}^{+\infty} \exp(2^{n+1}C_D - 4^n \beta) (C_D)^{n+1}\right) \operatorname{vol}(B_{z_0}(1)).
$$

The sum converges, so that we get the  $z_0$ -independent lower bound:

$$
\frac{\int_M e^{-\alpha d(z,z_0)^2} d\operatorname{vol}(z)}{\int_M e^{-\beta d(z,z_0)^2} d\operatorname{vol}(z)} \ge \frac{e^{-\alpha}}{1 + \left(\sum_{n=0}^{+\infty} \exp\left(2^{n+1}C_D - 4^n\beta\right) (C_D)^{n+1}\right)}.
$$

This proves Lemma [3.2.](#page-23-0) Note that the numerator only needs to be an integral of a nonnegative, radially decaying function, rather than of a Gaussian.

Lower bounds on Ricci curvature also ensure the following Gaussian (in terms of the geodesic distance) upper and lower bounds on the heat kernel [\[Bau22,](#page-28-22) Theorem 2.35]:

**Lemma 3.3** (Gaussian bounds for the heat kernel). – Denote  $p(t, z, z_0)$  the heat kernel of M at time t between two points z,  $z_0$  in M. There exist constants  $C_1, C_2$  depending only on  $(n, K)$  such that the following estimates hold:

<span id="page-23-3"></span>(3.4) 
$$
p(z, z_0, t) \leq \frac{C_1}{\text{vol}(B_{z_0}(\sqrt{t}))} \exp\left(-\frac{d(z, z_0)^2}{5t} + C_2(t + d(z, z_0)^2)\right),
$$

<span id="page-23-2"></span>(3.5) 
$$
p(z, z_0, t) \ge \frac{C_1^{-1}}{\text{vol}(B_{z_0}(\sqrt{t}))} \exp\left(-\frac{d(z, z_0)^2}{3t} - C_2(t + d(z, z_0)^2)\right).
$$

For proof of these geometric inequalities, we refer to  $\lceil \text{Bau22} \rceil$  and the references therein, in particular [\[LY86,](#page-30-20) [Gri09\]](#page-29-22).

<span id="page-24-1"></span>We now consider a Borel subset  $\omega \subset M$  satisfying the spectral estimate, which we recall:

(3.6) 
$$
\forall u \in L^2, \forall \Lambda > 0, \quad \|\Pi_{\Lambda}u\|_{L^2(M)} \leq Ce^{C\Lambda}\|\Pi_{\Lambda}u\|_{L^2(\omega)}.
$$

The spectral estimate  $(3.6)$  implies the following final-time observability estimate from  $(0,T)\times\omega$  (see [\[Mil10,](#page-30-5) [PW13,](#page-30-6) [WWZZ19,](#page-30-7) [BM23\]](#page-28-1) for proof and refined statements, in particular concerning the time set):

<span id="page-24-0"></span>**Lemma 3.4** (Observability). – Assume  $\omega$  satisfies the spectral estimate property [\(3.6\)](#page-24-1) and consider  $u(t, z)$  a solution of the heat equation

(3.7) 
$$
\partial_t u - \Delta_g u = 0 \text{ in } \mathbb{R}_t^+ \times M,
$$

$$
u|_{t=0} = u_0 \text{ in } M.
$$

for some  $u_0 \in L^2(M)$ . For any  $T > 0$ , there exists some constant  $C_{obs}(T) > 0$  such that

<span id="page-24-2"></span>(3.8) 
$$
\int_M |u(T,z)|^2 d\,\text{vol}(z) \leq C_{obs}(T) \int_0^T \int_{\omega} |u(s,z)|^2 d\,\text{vol}(z) ds.
$$

A proof of (a stronger version of) Lemma [3.4](#page-24-0) on the flat space can be found in [\[WWZZ19,](#page-30-7) Lemmas 2.3 and 2.4]. It adapts easily to the manifold case, as explained in [\[Rou24,](#page-30-0) Appendix  $B$ .

Fix  $t_K > 0$  a time such that

$$
-\beta_K = -\frac{2}{5t_K} + 2C_2 < -4C_D,
$$

for example  $t_K = \frac{1}{10(C_2+1)}$  $\frac{1}{10(C_2+2C_D)}$ . Note that  $t_K$  depends only on the dimension and curvature bound of the manifold M. We fix some  $z_0 \in M$  and apply the observability inequality [\(3.8\)](#page-24-2) at time  $T = \frac{t_K}{2}$  for the initial condition

(3.9) 
$$
u_0(z) = p\left(z, z_0, \frac{t_K}{2}\right), \quad z \in M.
$$

By the semi-group property for the heat equation, we get

(3.10) 
$$
u(s, z) = p\left(z, z_0, s + \frac{t_K}{2}\right), \quad s \ge 0, z \in M,
$$

thus

$$
(3.11) \qquad \int_M |p(z, z_0, t_K)|^2 d\operatorname{vol}(z) \le C_{obs} \int_0^{\frac{t_K}{2}} \int_\omega \left| p\left(z, z_0, \frac{t_K}{2} + s\right) \right|^2 d\operatorname{vol}(z) ds.
$$

We bound the left-hand-side from below using [\(3.5\)](#page-23-2):

<span id="page-24-3"></span>
$$
\int_{M} |p(z, z_0, t_K)|^2 d\operatorname{vol}(z) \ge \frac{C_1^{-2}}{\operatorname{vol}(B_{z_0}(\sqrt{t_K}))^2} \int_{M} \exp^{-2C_2 t_K - \alpha_K d(z, z_0)^2} d\mu(z)
$$

$$
\ge \frac{m_K}{\operatorname{vol}(B_{z_0}(\sqrt{t_K}))^2} \int_{M} e^{-\alpha_K d(z, z_0)^2} d\mu(z)
$$

with  $m_K$  and  $\alpha_K = 2\left(C_2 + \frac{1}{3t_K}\right)$  two constants depending on the dimension and the curvature lower bound. In the rest of the proof, constants indexed by a  $K$  are only dependent on those geometric features. They do not depend on  $z_0$  or the parameter R that will be introduced later on.

We now bound the right-hand-side from above using the Gaussian upper bound [\(3.4\)](#page-23-3). We then introduce some radius  $R > 0$ , which will be fixed independently of  $z_0$  at the end of the proof. We split the integral over  $\omega$  into its  $B_{z_0}(R)$  and  $B_{z_0}(R)^c$  parts:

$$
\int_{0}^{t_{K}/2} \int_{\omega} |p(z, z_{0}, t + s)|^{2} dvol(z) ds
$$
\n
$$
\leq \int_{0}^{t_{K}/2} \int_{\omega} \frac{C_{1}^{2}}{\text{vol}\left(B_{z_{0}}\left(\sqrt{\frac{t_{K}}{2} + s}\right)\right)^{2}} e^{C_{2}(t_{K}+2s)} \exp^{-\beta_{K}d(z, z_{0})^{2}} dvol(z)
$$
\n
$$
\leq \frac{C_{1}^{2} e^{2C_{2}t_{K}}}{\text{vol}\left(B_{z_{0}}\left(\sqrt{\frac{t_{K}}{2}}\right)\right)^{2}} \left(\int_{\omega \cap B_{z_{0}}(R)} e^{-\beta_{K}d(z, z_{0})^{2}} dvol(z) + \int_{\omega \cap B_{z_{0}}(R)^{c}} e^{-\beta_{K}d(z, z_{0})^{2}} dvol(z)\right)
$$
\n
$$
\leq \frac{M_{K}}{\text{vol}\left(B_{z_{0}}\left(\sqrt{\frac{t_{K}}{2}}\right)\right)^{2}} \left(\text{vol}(\omega \cap B_{z_{0}}(R)) + e^{-\frac{\beta_{K}}{2}R^{2}} \int_{\omega \cap B_{z_{0}}(R)^{c}} e^{-\frac{\beta_{K}}{2}d(z, z_{0})^{2}} dvol(z)\right)
$$
\n
$$
\leq \frac{M_{K}}{\text{vol}\left(B_{z_{0}}\left(\sqrt{\frac{t_{K}}{2}}\right)\right)^{2}} \left(\text{vol}(\omega \cap B_{z_{0}}(R)) + e^{-\frac{\beta_{K}}{2}R^{2}} \int_{M} e^{-\frac{\beta_{K}}{2}d(z, z_{0})^{2}} dvol(z)\right).
$$

Plugging the lower and upper bounds in [\(3.11\)](#page-24-3) yields

$$
(3.12) \quad \int_M e^{-\alpha_K d(z,z_0)^2} d\mu(z) \le M'_K \left( \text{vol}(\omega \cap B_{z_0}(R)) + e^{-\frac{\beta_K}{2}R^2} \int_M e^{-\frac{\beta_K}{2} d(z,z_0)^2} d\text{vol}(z) \right).
$$

where

$$
M'_{K} = \frac{M_{K}}{m_{k}} \sup_{z_{0} \in M} \frac{\text{vol}(B_{z_{0}}(\sqrt{t_{K}}))^{2}}{\text{vol}\left(B_{z_{0}}\left(\sqrt{\frac{t_{K}}{2}}\right)\right)^{2}}
$$

can be chosen independently of  $z_0$  thanks to the volume doubling estimate  $(3.1)$ .

We conclude the proof of Theorem [2](#page-2-2) using Lemma [3.2.](#page-23-0) By the lemma, for  $\alpha = \alpha_K$  and  $\beta = \frac{\beta_K}{2}$  $\frac{2K}{2}$ , it is possible to find some  $R = R_K > 0$  such that

$$
e^{-\frac{\beta_K}{2}R^2} < \frac{1}{2} \inf_{z_0 \in M} \frac{\int_M e^{-\alpha_K d(z, z_0)^2} d\operatorname{vol}(z)}{\int_M e^{-\frac{\beta_K}{2} d(z, z_0)^2} d\operatorname{vol}(z)}.
$$

For such an  $R$ ,  $(3.3)$  yields

(3.13) 
$$
\int_M e^{-\alpha_K d(z,z_0)^2} d\mu(z) \leq 2M'_K \left( \text{vol}(\omega \cap B_{z_0}(R)) \right),
$$

hence we recover the thickness condition

(3.14) 
$$
\delta \operatorname{vol}(B_{z_0}(R)) \le \operatorname{vol}(\omega \cap B_{z_0}(R))
$$

for

$$
\delta = \frac{1}{2M'_K}e^{-\alpha_K R^2}
$$

.



Figure 5. An example of setting where the Logunov–Malinnikova inequality allows to propagate the smallness of gradients from a 2 $-$ dimensional set E to a larger 3–dimensional box K. The auxiliary t introduced in the paper can be interpreted as the horizontal variable.

 $\delta$  depends only on R, the dimension n of the manifold, and the curvature lower bound.

We finish this section by noting that the proof that observability implies a thickness assumption may generalize to other semigroups  $e^{-tH}$ . The only assumptions needed are the volume doubling estimate, which depends on the metric rather than the operator considered, and an integral kernel of the propagator satisfying a Gaussian pointwise bound from above and a positive radially decaying bound from below.

## <span id="page-26-0"></span>Appendix A. Statement of propagation of smallness inequalities used in the paper

The goal of this appendix is to recall the different propagation of smallness inequalities that are used throughout the paper. We first present the Logunov–Malinnikova theorem on propagation of smallness for gradients of harmonic functions [\[LM18b,](#page-29-4) Theorem 5.1], which is used crucially in charts in Section [2.1.](#page-11-2)

**Theorem 7** ([\[LM18b\]](#page-29-4)). – Consider  $\Omega$  a bounded domain in  $\mathbb{R}^{d+1}$ . There exists a constant  $c \in (0,1)$  that depends only on the dimension such that the following holds: take any u solution of an elliptic equation in divergence form

(A.1) div(A∇v) = 0 over Ω

where  $A = (a_{i,j}(x))_{i,j}$  is a symmetric, uniformly elliptic matrix with Lipschitz entries:

$$
\Lambda_1^{-1}|\zeta|^2 \le \langle A\zeta, \zeta \rangle \le \Lambda_1|\zeta|^2, \quad |a_{i,j}(x) - a_{i,j}(y)| \le \Lambda_2|x - y|.
$$

Let m,  $\delta$ ,  $\rho$  be positive numbers, and assume sets  $E, K \subset \Omega$  satisfy

$$
C_{\mathcal{H}}^{d-c+\delta}(E) > m, \quad \text{dist}(E, \partial \Omega) > \rho, \text{dist}(\mathcal{K}, \partial \Omega) > \rho.
$$

Then there exist  $C, \gamma > 0$  depending only on  $m, \delta, \rho, \Lambda_1, \Lambda_2, \Omega$  such that

$$
\sup_{\mathcal{K}} |\nabla v| \leq C \sup_{E} |\nabla v|^{\gamma} \sup_{\Omega} |\nabla v|^{1-\gamma}.
$$

In particular, the result holds for  $\delta = c$  ie if E is a set of Hausdorff dimension d. An illustration of this result is presented if Figure [A](#page-26-0) in the case  $d = 2, \delta = c$ .

In Section [2.4,](#page-20-0) which is dedicated to bounds on the compact core of a manifolds, we use Carleman-inequality based propagation of smallness from open sets in the spirit of Lebeau and Robbiano [\[LR95\]](#page-29-8). The first of these results can be found in the textbook [\[LRLR22b,](#page-29-18) Theorem 9.1] (see also [\[LRLR22a,](#page-29-17) Lemma 5.3] for the underlying result applied in charts). The proof is based on interior Carleman estimates applied in ordered local charts along paths connecting a finite number of points in the compact core to a point in the observation set. We state it in the specific case  $l = 0$ :

<span id="page-27-4"></span>**Theorem 8** ([\[LRLR22b\]](#page-29-18)). – Let  $(M, g)$  be a d-dimensional, possibly non-compact Riemannian manifold. Consider the operator  $P = -\Delta_g + R_1$  with  $R_1$  a first-order differential operator on M. Let  $U, V$  be two open subsets of M such that U is connected,  $\overline{U} \cap \partial M = \emptyset$  and  $U \subset\subset V \subset\subset \mathcal{M}.$ 

Let X be an open subset of U. There exist  $C > 0$  and  $\beta \in (0,1)$  such that

(A.2) 
$$
||w||_{H^1(\mathcal{U})} \leq C||w||_{H^1(\mathcal{V})}^{1-\beta} (||f||_{L^2(\mathcal{V})} + ||w||_{L^2(\mathcal{X})})^{\beta},
$$

for any  $w, f \in H^2(\mathcal{V}) \times L^2(\mathcal{V})$  satisfying  $Pw = f$ .

We apply this result to the harmonic extension  $w = \partial_t v_{\Lambda}$ , which is a solution of an elliptic equation. However, to get a bound by  $\Pi_{\Lambda}u = \partial_t v_{\Lambda}(t=0)$  in the right-hand side, propagation of smallness from the boundary is also needed. This the object of the next theorem. We take the statement from [\[JL99,](#page-29-10) Lemma 14.5]. The idea of the proof is to apply a boundary Carleman estimate (itself coming from [\[LR95,](#page-29-8) [Rob95\]](#page-30-21)) to the harmonic extension multiplied by appropriate cutoffs localizing in concentric sectors of a half-ball  $B \cap \{t > 0\}$ . The statement is the following:

<span id="page-27-5"></span>**Theorem 9** ([\[JL99\]](#page-29-10)). – Denote  $X_T = [-T, T] \times M$  for  $T > 0$ . Let B be a ball in  $X_2$  centered centered on  $0 \times M$ . Let  $B^+ = \{(t, z) \in B, t > 0\}$ . Suppose that  $V \subset B^+$ , and F is a solution of  $(\partial_t^2 + \Delta_g)F = 0$  satisfying  $F(0, x) = 0$ . There exist  $\gamma \in (0, 1)$  and C depending only on V, B and M such that

(A.3) 
$$
||F||_{L^{2}(V)} \leq C||\partial_{t} F(t=0)||_{L_{z}^{2}(B \cap t=0)}^{\gamma} ||F||_{H^{1}(B^{+})}^{1-\gamma}.
$$

Note that though the work of Jerison–Lebeau deals with compact manifolds with boundary, this lemma and its proof are local and avoid the boundary. The proof given in [\[JL99\]](#page-29-10) deals with the flat Laplacian. As noted in [\[LR95,](#page-29-8) Section 3.A], the inequality for any operator of the form  $\partial_t^2 + \Delta_g$ .

#### <span id="page-27-0"></span>**REFERENCES**

- <span id="page-27-1"></span>[AE13] Jone Apraiz and Luis Escauriaza. Null-control and measurable sets. ESAIM Control Optim. Calc. Var., 19(1):239–254, 2013.
- <span id="page-27-2"></span>[AEWZ14] Jone Apraiz, Luis Escauriaza, Gengsheng Wang, and Can Zhang. Observability inequalities and measurable sets. J. Eur. Math. Soc. (JEMS), 16(11):2433–2475, 2014.
- <span id="page-27-3"></span>[Agm82] Shmuel Agmon. Lectures on exponential decay of solutions of second-order elliptic equations: bounds on eigenfunctions of N-body Schrödinger operators, volume 29 of Mathematical Notes. Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1982.
- <span id="page-28-11"></span>[AK24] Paul Alphonse and Armand Koenig. Null-controllability for weakly dissipative heat-like equations. Evol. Equ. Control Theory, 13(3):973–988, 2024.
- <span id="page-28-19"></span>[Alp24] Paul Alphonse. Null-controllability of evolution equations associated with fractional Shubin operators through quantitative Agmon estimates. Ann. Inst. Fourier (Grenoble), 74(4):1671–1720, 2024.
- <span id="page-28-17"></span>[AS23] Paul Alphonse and Albrecht Seelmann. Quantitative spectral inequalities for the anisotropic shubin operators and applications to null-controllability, 2023.
- <span id="page-28-22"></span>[Bau22] Fabrice Baudoin. Geometric inequalities on Riemannian and sub-Riemannian manifolds by heat semigroups techniques. In New trends on analysis and geometry in metric spaces, volume 2296 of Lecture Notes in Math., pages  $7-91$ . Springer, Cham, [2022]  $\odot$ 2022.
- <span id="page-28-5"></span>[Ber11] Nicolas Bergeron. Le spectre des surfaces hyperboliques. Savoirs Actuels (Les Ulis). [Current Scholarship (Les Ulis)]. EDP Sciences, Les Ulis; CNRS Éditions, Paris, 2011.
- <span id="page-28-13"></span>[BJ16] Nicolas Burq and Romain Joly. Exponential decay for the damped wave equation in unbounded domains. Commun. Contemp. Math., 18(6):1650012, 27, 2016.
- <span id="page-28-10"></span>[BJPS21] Karine Beauchard, Philippe Jaming, and Karel Pravda-Starov. Spectral estimates for finite combinations of Hermite functions and null-controllability of hypoelliptic quadratic equations. Studia  $Math., 260(1):1-43, 2021.$
- <span id="page-28-0"></span>[BM21] Nicolas Burq and Iván Moyano. Propagation of smallness and spectral estimates, 2021.
- <span id="page-28-1"></span>[BM23] Nicolas Burq and Iván Moyano. Propagation of smallness and control for heat equations. J. Eur. Math. Soc. (JEMS), 25(4):1349–1377, 2023.
- <span id="page-28-9"></span>[BN02] Assia Benabdallah and Maria Grazia Naso. Null controllability of a thermoelastic plate. Abstr. Appl. Anal., 7(11):585–599, 2002.
- <span id="page-28-4"></span>[Bor16] David Borthwick. Spectral theory of infinite-area hyperbolic surfaces, volume 318 of Progress in Mathematics. Birkhäuser/Springer, [Cham], second edition, 2016.
- <span id="page-28-12"></span>[BT24] Clemens Bombach and Martin Tautenhahn. A Logvinenko-Sereda theorem for vector-valued functions and application to control theory, 2024.
- <span id="page-28-2"></span>[CGT82] Jeff Cheeger, Mikhail Gromov, and Michael Taylor. Finite propagation speed, kernel estimates for functions of the Laplace operator, and the geometry of complete Riemannian manifolds. J. Differential Geometry, 17(1):15–53, 1982.
- <span id="page-28-20"></span>[CNV15] Jeff Cheeger, Aaron Naber, and Daniele Valtorta. Critical sets of elliptic equations. Comm. Pure Appl. Math., 68(2):173–209, 2015.
- <span id="page-28-14"></span>[DF88] Harold Donnelly and Charles Fefferman. Nodal sets of eigenfunctions on Riemannian manifolds. Invent. Math., 93(1):161–183, 1988.
- <span id="page-28-15"></span>[DF90] Harold Donnelly and Charles Fefferman. Nodal sets of eigenfunctions: Riemannian manifolds with boundary. In Analysis, et cetera, pages 251–262. Academic Press, Boston, MA, 1990.
- <span id="page-28-18"></span>[DSV24] Alexander Dicke, Albrecht Seelmann, and Ivan Veselić. Spectral inequality with sensor sets of decaying density for Schrödinger operators with power growth potentials. Partial Differ. Equ. Appl., 5(2):Paper No. 7, 2024.
- <span id="page-28-16"></span>[DV24] Alexander Dicke and Ivan Veselić. Spherical logvinenko–sereda–kovrijkine type inequality and nullcontrollability of the heat equation on the sphere. Arch. Math., Sep 2024.
- <span id="page-28-8"></span>[ENS<sup>+</sup>20] Michela Egidi, Ivica Nakić, Albrecht Seelmann, Matthias Täufer, Martin Tautenhahn, and Ivan Veselić. Null-controllability and control cost estimates for the heat equation on unbounded and large bounded domains. In Control theory of infinite-dimensional systems, volume 277 of Oper. Theory Adv. Appl., pages 117–157. Birkhäuser/Springer, Cham, [2020] ©2020.
- <span id="page-28-7"></span>[ES21] Michela Egidi and Albrecht Seelmann. An abstract Logvinenko-Sereda type theorem for spectral subspaces. J. Math. Anal. Appl., 500(1):Paper No. 125149, 32, 2021.
- <span id="page-28-6"></span>[EV18] Michela Egidi and Ivan Veselić. Sharp geometric condition for null-controllability of the heat equation on  $\mathbb{R}^d$  and consistent estimates on the control cost. Arch. Math. (Basel), 111(1):85–99, 2018.
- <span id="page-28-21"></span>[Eva22] Lawrence C. Evans. Partial differential equations, volume 19. American Mathematical Society, 2022.
- <span id="page-28-3"></span>[GHL04] Sylvestre Gallot, Dominique Hulin, and Jacques Lafontaine. Riemannian geometry. Universitext. Springer-Verlag, Berlin, third edition, 2004.

<span id="page-29-22"></span><span id="page-29-21"></span><span id="page-29-20"></span><span id="page-29-19"></span><span id="page-29-18"></span><span id="page-29-17"></span><span id="page-29-16"></span><span id="page-29-15"></span><span id="page-29-14"></span><span id="page-29-13"></span><span id="page-29-12"></span><span id="page-29-11"></span><span id="page-29-10"></span><span id="page-29-9"></span><span id="page-29-8"></span><span id="page-29-7"></span><span id="page-29-6"></span><span id="page-29-5"></span><span id="page-29-4"></span><span id="page-29-3"></span><span id="page-29-2"></span><span id="page-29-1"></span><span id="page-29-0"></span>

- <span id="page-30-2"></span>[LS74] Vladimir N. Logvinenko and Ju. F. Sereda. Equivalent norms in spaces of entire functions of exponential type. Teor. Funk. Anal. i Priložen., 20:102–111, 175, 1974.
- <span id="page-30-20"></span>[LY86] Peter Li and Shing-Tung Yau. On the parabolic kernel of the Schrödinger operator. Acta Math., 156(3-4):153–201, 1986.
- <span id="page-30-8"></span>[LZ98] Gilles Lebeau and Enrique Zuazua. Null-controllability of a system of linear thermoelasticity. Arch. Rational Mech. Anal., 141(4):297–329, 1998.
- <span id="page-30-15"></span>[Mil04a] Luc Miller. Geometric bounds on the growth rate of null-controllability cost for the heat equation in small time. J. Differential Equations, 204(1):202–226, 2004.
- <span id="page-30-10"></span>[Mil04b] Luc Miller. How violent are fast controls for Schrödinger and plate vibrations? Arch. Ration. Mech. Anal., 172(3):429–456, 2004.
- <span id="page-30-12"></span>[Mil05a] Luc Miller. On the null-controllability of the heat equation in unbounded domains. Bull. Sci. Math., 129(2):175–185, 2005.
- <span id="page-30-13"></span>[Mil05b] Luc Miller. Unique continuation estimates for the Laplacian and the heat equation on non-compact manifolds. Math. Res. Lett., 12(1):37–47, 2005.
- <span id="page-30-5"></span>[Mil10] Luc Miller. A direct Lebeau-Robbiano strategy for the observability of heat-like semigroups. Discrete Contin. Dyn. Syst. Ser. B, 14(4):1465–1485, 2010.
- <span id="page-30-9"></span>[MPS23] Jérémy Martin and Karel Pravda-Starov. Spectral inequalities for combinations of Hermite functions and null-controllability for evolution equations enjoying Gelfand-Shilov smoothing effects. J. Inst. Math. Jussieu, 22(6):2533–2582, 2023.
- <span id="page-30-4"></span>[Naz93] Fedor L. Nazarov. Local estimates for exponential polynomials and their applications to inequalities of the uncertainty principle type. Algebra i Analiz, 5(4):3–66, 1993.
- <span id="page-30-16"></span>[NTTV20] Ivica Nakić, Matthias Täufer, Martin Tautenhahn, and Ivan Veselić. Sharp estimates and homogenization of the control cost of the heat equation on large domains. ESAIM Control Optim. Calc. Var., 26:Paper No. 54, 26, 2020.
- <span id="page-30-3"></span>[PS85] Ralph S. Phillips and Peter Sarnak. On cusp forms for co-finite subgroups of  $PSL(2, \mathbf{R})$ . Invent. Math., 80(2):339–364, 1985.
- <span id="page-30-6"></span>[PW13] Kim Dang Phung and Gengsheng Wang. An observability estimate for parabolic equations from a measurable set in time and its applications. J. Eur. Math. Soc. (JEMS), 15(2):681–703, 2013.
- <span id="page-30-17"></span>[RMV13] Constanza Rojas-Molina and Ivan Veselić. Scale-free unique continuation estimates and applications to random Schrödinger operators. Comm. Math. Phys., 320(1):245–274, 2013.
- <span id="page-30-21"></span>[Rob95] Luc Robbiano. Fonction de coût et contrôle des solutions des équations hyperboliques. Asymptotic Anal.,  $10(2):95-115$ , 1995.
- <span id="page-30-0"></span>[Rou24] Marc Rouveyrol. Spectral estimate for the Laplace-Beltrami operator on the hyperbolic half-plane, 2024.
- <span id="page-30-1"></span>[RS80] Michael Reed and Barry Simon. Methods of modern mathematical physics. I. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, second edition, 1980. Functional analysis.
- <span id="page-30-14"></span>[RT24] C. Rose and M. Tautenhahn. Unique continuation estimates on manifolds with Ricci curvature bounded below. J. Math. Phys., 65(4):Paper No. 041503, 16, 2024.
- <span id="page-30-11"></span>[SSY25] Pei Su, Chenmin Sun, and Xu Yuan. Quantitative observability for one-dimensional Schrödinger equations with potentials. J. Funct. Anal., 288(2):Paper No. 110695, 2025.
- <span id="page-30-19"></span>[Wan24] Yunlei Wang. Quantitative 2d propagation of smallness and control for 1d heat equations with power growth potentials, 2024.
- <span id="page-30-7"></span>[WWZZ19] Gengsheng Wang, Ming Wang, Can Zhang, and Yubiao Zhang. Observable set, observability, interpolation inequality and spectral inequality for the heat equation in  $\mathbb{R}^n$ . J. Math. Pures Appl. (9), 126:144–194, 2019.
- <span id="page-30-18"></span>[Zhu24] Jiuyi Zhu. Spectral inequalities for Schrödinger equations with various potentials, 2024.

Laboratoire de Mathématiques d'Orsay, Université Paris-Saclay, Bâtiment 307, 91405 Orsay CEDEX

 $Email \ address:$  marc.rouveyrol@universite-paris-saclay.fr