Covariant interacting fractons

Erica Bertolini®* ebertolini@stp.dias.ie Hyungrok Kim (金炯錄)®[†] h.kim2@herts.ac.uk

3rd October 2024

Abstract

We show that there exists a natural analogue of the Yang–Mills equations using the Frölicher–Nijenhuis bracket between vector-valued differential forms. The gauge field is a rank-two tensor, and when one constrains it to be symmetric, then the system exhibits fractonic behaviours. In the linearised limit, the constrained equations of motion reduce to those of the covariant fracton model [1, 2, 3].

Contents

1	Introduction and summary	1
2	Mathematical background2.1Frölicher–Nijenhuis bracket2.2Twisting	3 3 4
3	 Motivation and idea 3.1 A review of Yang–Mills theory	5 5 5
4	Covariant interacting fractonic gauge theory4.1Non-symmetric theory4.2Symmetric theory	6 6 8

1 Introduction and summary

In recent years, a new kind of quasiparticle has emerged from the physics literature: *fractons* [4, 5, 6, 7]. Originating in lattice models in the context of spin glasses

^{*}School of Theoretical Physics, Dublin Institute for Advanced Studies, 10 Burlington Road, D04 C932, Dublin, Ireland.

Scoil na Fisice Teoiriciúla, Institiúid Ard-Léinn Bhaile Átha Cliath, 10 Bóthar Burlington, do4 c932, Baile Átha Cliath, Éire.

[†]Department of Mathematics and Theoretical Physics, University of Hertfordshire, Hatfield, Hertfordshire AL10 9AB, United Kingdom.

[8] and quantum information [9, 10], fractons have quickly attracted the interest of a wide variety of physicists from condensed matter to mathematical physics and have been influential in gauge theories and other quantum field theories [11, 12, 13, 14, 15, 16, 17, 18, 19, 20]. The main characteristic of fracton quasiparticles is immobility, which is also the reason for the name. Indeed a fracton is defined as a fraction of a mobile quasiparticle, and in isolation it cannot move at all. Only dipole-like excitations are free to displace, or, in general have fewer constraints on their motion [11, 12, 6, 7]. These additional constraints defines other fracton-related quasiparticles such as *lineons* and *planons*, which can move in a one- or two-dimensional subspace respectively. The restricted-motion feature, which unites all fracton theories, is shared by many physical systems and models, and it is one of the reasons for which fractons are so popular nowadays. For instance limited mobility, or complete immobility, can be harnessed for developing quantum memories [9, 10], or used as a mapping/duality to study topological defects in elastic media [21, 22, 23, 24, 25, 26, 27, 28]. It is also a characteristic found in Carrollian theories, which thus seem to display some fractonic behaviour [29, 30]. The other reason for attracting so much interest is found in the tensorial nature of fractonic theories. In gauge theory [11, 12] these models are indeed typically described in terms of a rank-2 symmetric tensor $A_{ij}(x)$ (with *i*, *j* spatial indices), which transforms under the gauge transformation

$$\delta_{\epsilon} A_{ij} = \partial_i \partial_j \epsilon \tag{1}$$

and share strong similarities with the electromagnetic Maxwell theory, of which they represent higher-rank generalisations. A generalised Gauss law is typically postulated as

$$\partial_i \partial_j E^{ij} = \rho, \tag{2}$$

where $E^{ij}(x)$ is a symmetric electric tensor field, implying dipole moment conservation through

$$D^{i} \coloneqq \int \mathrm{d}^{d}x \, x^{i} \rho = -\int \mathrm{d}^{d}x \, \partial_{j} E^{ij} = \oint \mathrm{d}^{d-1}x \, (\cdots), \tag{3}$$

that is, the dipole moment cannot change except through a nonzero flux at the boundary. This encodes the immobility of the fractonic charge $\rho(x)$ [6, 7] since, if a single charge were to move, it would change the total dipole moment of the system. The tensorial nature of the gauge field also hints towards connections with the theory of linearised gravity [31, 32, 33, 34, 35], which emerges naturally when the covariant fracton theory is taken into account [1, 2, 36, 37]. The covariant extension

$$\delta A_{\mu\nu} = \partial_{\mu}\partial_{\nu}\epsilon \tag{4}$$

of the fractonic transformation (1) is indeed a particular case of the infinitesimal diffeomorphisms that defines linearised gravity, sometimes called longitudinal diffeomorphisms [38]. Thus an action invariant under (4) would naturally carry a linearised gravity term. This covariant formulation [1] gives rise to the definition of an invariant rank-3 field strength $F_{\mu\nu\rho}(x)$, through which the Maxwell analogy and fracton phenomenology of [11, 12, 39] is reproduced and expanded from first principles. The covariant theory of fractons [1] is free (or at least quantum-electrodynamics-like) in the sense that the gauge field $A_{\mu\nu}(x)$ does

not interact directly with itself. However, the existence of a generalised invariant field strength $F_{\mu\nu\rho}(x)$ suggests a mathematically natural 'nonabelianisation' of the above theory through the use of the Frölicher–Nijenhuis bracket [40, 41] (reviewed in [42, §8]), which is the focus of this paper.

In an interacting theory of fractons where the gauge field interacts with itself, the gauge field modes themselves are fractonic, so that there may be constraints on asymptotic in- and out-states. This may be interesting from an amplitude-theoretic point of view and may possibly signal new loopholes to Weinberg–Witten-type no-go theorems [43].

The analysis presented in this paper is also of independent mathematical interest. The Frölicher–Nijenhuis bracket is a fundamental geometric structure, which has recently appeared in the context of integrable models such as self-dual Yang–Mills theory [44]. In fact, the Frölicher–Nijenhuis bracket generalises to arbitrary Lie algebroids [45], which appear in gauge theory in many contexts [46, 47, 48, 49, 50, 51, 52], and both the fractonic case at hand and ordinary Yang–Mills theory may be seen as special cases (for the tangent Lie algebroid and a Lie algebroid bundle, respectively) of a more general construction associated to general Lie algebroid. We thus see, again, that strong analogies manifest themselves between fractons and Maxwell/Yang–Mills theories.

Our interacting fractonic model is defined on flat space, as gauge invariance breaks on curved space: the field strength *F* only transforms tensorially if one assumes that the metric $g_{\mu\nu}$ is flat and also transforms under diffeomorphism. Thus, to write an action principle one cannot have $g_{\mu\nu}$ as a background (since it must transform), and the equations of motion for $g_{\mu\nu}$ must ensure that it remains flat (since otherwise gauge invariance fails). As a consequence, for the scope of the analysis presented here, we only postulate an equation of motion. For efforts at fractonic behaviours on curved spaces, see [53, 54, 55]. We also do not discuss issues regarding the classical or quantum stability of our model.

Acknowledgements Hyungrok Kim thanks Simon-Raphael Fischer[®] and Fridrich Valach III[®] for helpful discussion.

2 Mathematical background

In the following, we will need to make use of the Frölicher–Nijenhuis bracket on vector-valued differential forms and the language of twisting, which we review briefly.

2.1 Frölicher–Nijenhuis bracket

Let *M* be a smooth manifold. The graded vector space of vector-valued differential forms $\Omega^{\bullet}(M; TM) = \bigoplus_{i=0}^{d} \Omega^{i}(M)$ becomes a graded Lie algebra with respect to the *Frölicher–Nijenhuis bracket* [40, 41] (reviewed in [42, §8]):

$$[\phi \otimes X, \psi \otimes Y] = (\phi \land \psi) \otimes \mathcal{L}_X Y + (\phi \land \mathcal{L}_X \psi) \otimes Y - (\mathcal{L}_Y \phi \land \psi) \otimes X + (-1)^p (d\phi \land i_X \psi) \otimes Y + (-1)^p (i_Y \phi \land d\psi) \otimes X$$
(5)

for vector fields $X, Y \in \Gamma(TM)$ (where $\Gamma(-)$ denotes the space of sections of a vector bundle) and homogeneous differential forms $\phi \in \Omega^p(M), \psi \in \Omega^q(M)$,

where $\mathcal{L}_X(-)$ is the Lie derivative of a tensor field along a vector field, and $i_X(-)$ is the interior derivative of a differential form along a vector field.

In particular, between two (1,1)-tensors K_{ν}^{μ} and L_{ν}^{μ} , we have

$$[K, L]^{\rho}_{\mu\nu} = 2K^{\sigma}_{[\mu]}\partial_{\sigma}L^{\rho}_{[\nu]} + 2L^{\sigma}_{[\mu]}\partial_{\sigma}K^{\rho}_{[\nu]} - 2K^{\rho}_{\sigma}\partial_{[\mu}L^{\sigma}_{\nu]} - 2L^{\rho}_{\sigma}\partial_{[\mu}K^{\sigma}_{\nu]}.$$
 (6)

where antisymmetrisations are normalised.

Notice that, when one of the arguments is a (1,0)-tensor (i.e. a vector field), it reduces to the usual Lie derivative:

$$[X, -] = \mathcal{L}_X \qquad (X \in \Gamma(TM)). \tag{7}$$

Suppose that *M* is equipped with a Riemannian metric *g* whose Riemann curvature vanishes. Then, using the induced Levi-Civita connection ∇ , we may define the covariant exterior derivative

$$d^{\nabla} \colon \Omega^{\bullet}(M; TM) \to \Omega^{\bullet+1}(M; TM)$$
(8)

that squares to zero, and then $\Omega^{\bullet}(M; TM)$ forms a differential graded Lie algebra. Note that, when the curvature of *g* does not vanish, then d^{∇} need not square to zero.

2.2 Twisting

A curved¹ differential graded Lie algebra (g, r, d, [-, -]) is a \mathbb{Z} -graded Lie algebra (g, [-, -]) together with a linear map

$$d: \mathfrak{g} \to \mathfrak{g} \tag{9}$$

of degree one and an element $r \in g$ of degree two, called the *curvature*, such that d is a graded derivation with respect to the Lie bracket [-, -] and

$$d(d(x)) = [r, x] \tag{10}$$

for any $x \in \mathfrak{g}$. This is the special case of the notion of a curved L_{∞} -algebra [56, 57, 58] ($\mathfrak{g}, \mu_0, \mu_1, \mu_2, \ldots$), which is a graded vector space \mathfrak{g} equipped with totally graded-antisymmetric multilinear maps $\mu_i \colon \mathfrak{g}^{\otimes i} \to \mathfrak{g}$ that satisfy the Jacobi identity up to homotopy. Then a curved differential graded Lie algebra is the same as a curved L_{∞} -algebra in which $\mu_i = 0$ except for $i \in \{0, 1, 2\}$; then μ_0, μ_1 , and μ_2 correspond to r, d, and [-, -], respectively.

Given a differential graded Lie algebra $(g, d_g, [-, -]_g)$ and an element $Q \in g^1$ of degree one, then the *twist*² of g by Q is the curved differential graded Lie algebra $g_Q := (g, dQ + \frac{1}{2}[Q, Q]_g, d_g + [Q, -]_g, [-, -]_g)$. When $[Q, Q]_g = 0$, then this is a differential graded Lie algebra.

¹The nomenclature comes from an analogy with the fact that, when one considers e.g. differential forms valued in a vector bundle with connection, the (covariant) exterior derivative operator does not in general square to zero anymore but rather to a curvature-dependent term.

²This is a special case of the twisting of a curved L_{∞} -algebra [56, 57, 58].

3 Motivation and idea

3.1 A review of Yang–Mills theory

Linearised Yang–Mills theory (that is, a direct sum of copies of Maxwell theory) admits a natural nonabelianisation in the form of Yang–Mills theory. Let us recall how this works. The field strength in linearised Yang–Mills theory is

$$F^a_{\mu\nu} = 2\partial_{[\mu}A^a_{\nu]}.\tag{11}$$

In the non-Abelian theory, this is modified to

$$F^{a}_{\mu\nu} = 2\partial_{[\mu}A^{a}_{\nu]} + f^{a}{}_{bc}A^{b}_{\mu}A^{c}_{\nu}.$$
 (12)

It is convenient to use the notation of Lie-algebra-valued differential forms, in terms of which we have

$$F = dA + \frac{1}{2}[A, A].$$
 (13)

That is, the field strength is fixed by the structure of a differential graded Lie algebra on the space of g-valued differential forms $\Omega^{\bullet}(M) \otimes \mathfrak{g} = \bigoplus_{i=0}^{\dim M} \Omega^{i}(M) \otimes \mathfrak{g}$, where \mathfrak{g} is the colour Lie algebra, M is spacetime, and \bullet is a placeholder for the form degree. Furthermore, this fixes the structure of gauge transformations and Bianchi identities:

$$\delta_{\alpha}A = d_{A}\alpha \qquad d_{A}F = 0 \qquad d_{A} := d + \frac{1}{2}[A, -] \qquad \delta_{\alpha}F = [\alpha, F].$$
(14)

The procedure of replacing d with d_A goes by the name of *twisting* [56, 57, 58] as discussed in section 2.2. After this, we no longer have a differential graded Lie algebra in the usual sense since

$$d_A^2 = [F, -] \neq 0,$$
(15)

but we speak of a *curved* differential graded Lie algebra. Given this, the equation of motion for the theory is fixed to be

$$\mathbf{d}_A \star F = \mathbf{0}.\tag{16}$$

3.2 Nonabelianising the covariant fracton model: the idea

The covariant fracton model [1] is a free theory whose fundamental field is a symmetric tensor $A_{\mu\nu}(x)$. An invariant field strength with one derivative can be defined, which is of the form [3, eq. 7.2.16]

$$F_{\mu\nu\rho} = a_1 \partial_\mu A_{\nu\rho} + a_2 \partial_\rho A_{\mu\nu} - (a_1 + a_2) \partial_\nu A_{\mu\rho} \tag{17}$$

for some suitable parameters $a_1, a_2 \in \mathbb{R}$. For any value of a_1, a_2 , the field strength $F_{\mu\nu\rho}(x)$ satisfies a Bianchi identity [3, p. 83]

$$0 = \partial_{\mu}(F_{\beta\nu\rho} - F_{\beta\rho\nu}) + \partial_{\nu}(F_{\beta\rho\mu} - F_{\beta\mu\rho}) + \partial_{\rho}(F_{\beta\mu\nu} - F_{\beta\nu\mu}) = 6\partial_{[\mu|}F_{\beta|\nu\rho]}.$$
 (18)

This has the form of an exterior derivative, except that the index β does not participate in the antisymmetrisation. Thus, it is natural to regard *F* as a T**M*-valued two-form, similar to how the Yang–Mills field strength is a Lie-algebra-valued two-form. This then means that $A_{\mu\nu}(x)$ should also be regarded as a T**M*-valued one-form.

There are however three problems that arise in this case, which are related.

- 1. There is no obvious Lie bracket for T**M*-valued differential forms (unlike Lie-algebra-valued differential forms).
- 2. A T**M*-valued one-form will *not* generally be antisymmetric between its two indices.
- 3. The gauge parameter should naturally be a T**M*-valued zero-form, i.e. an ordinary one-form, which is bigger than the scalar field gauge parameter of the covariant fracton model.

We resolve these interrelated problems as follows.

- 1. Unlike T**M*-valued forms, there *does* exist a natural Lie bracket on T*M*-valued forms: the Frölicher–Nijenhuis bracket [40, 41] (reviewed in [42, §8]). Thus, we work with T*M*-valued forms, and initially ignore the symmetry property of the T*M*-valued one-form gauge field $A^{\mu}{}_{\nu}(x)$. Therefore, $\Omega^{\bullet}(M; TM)$ is a graded Lie algebra. For this to be a *differential* graded Lie algebra, we fix a flat connection on T*M*.
- 2. Having formulated this theory, then we will impose the symmetry requirement with respect to a pseudo-Riemannian metric:

$$g_{\mu\nu}A^{\nu}{}_{\rho} = g_{\rho\nu}A^{\nu}{}_{\mu}.$$
 (19)

3. The constraint (19) will then naturally reduce the gauge symmetry from $\Omega^0(M; TM)$ to $\Omega^0(M)$, i.e. it will require the gauge parameter to be a scalar as for the covariant fracton theory.

4 Covariant interacting fractonic gauge theory

As mentioned in section 3, we first construct a nonlinear equation of motion for a (1,1)-tensor field $A^{\mu}{}_{\nu}$ in section 4.1. Then we constrain it to be symmetric in section 4.2.

4.1 Non-symmetric theory

Let *M* be a smooth manifold equipped with a flat Riemannian metric $g_{\mu\nu}$ (such as Minkowski space). Then $(\Omega^{\bullet}(M; TM), d^{\nabla}, [-, -])$ is a differential graded Lie algebra, where [-, -] is the Frölicher–Nijenhuis bracket (5).

Consider a (1,1)-tensor

$$A^{\mu}{}_{\nu} \in \Omega^1(M; \mathrm{T}M). \tag{20}$$

Then we may twist, as discussed in section 2.2, to obtain the curved differential graded Lie algebra

$$(\Omega^{\bullet}(M; \mathrm{T}M), F_A, \mathrm{d}_A^{\vee}, [-\wedge -])$$
(21)

where

$$\mathbf{d}_{A}^{\nabla} \coloneqq \mathbf{d}^{\nabla} + [A, -] \tag{22}$$

and

$$F_A = \mathrm{d}^{\nabla} A + \frac{1}{2} [A, A] \tag{23}$$

is the curvature.³ In particular, we have

$$(d_A^{\nabla})^2 = [F_A, -].$$
 (24)

We postulate the infinitesimal gauge symmetry

$$\delta_{\epsilon} A = \mathrm{d}_A \epsilon \tag{25}$$

for a vector gauge parameter $\epsilon \in \Omega^0(M; TM) = \Gamma(TM)$ which, in explicit component notation, is

$$(\delta_{\epsilon}A)^{\mu}{}_{\nu} = \nabla_{\nu}\epsilon^{\mu} - \mathcal{L}_{\epsilon}A^{\nu} = \partial_{\nu}\epsilon^{\mu} - \epsilon^{\rho}\partial_{\rho}A^{\mu}{}_{\nu} + \partial_{\rho}\epsilon^{\mu}A^{\rho}{}_{\nu} - \partial_{\nu}\epsilon^{\rho}A^{\mu}{}_{\rho}, \qquad (26)$$

and define the field strength

$$F_A := \mathrm{d}A + \frac{1}{2}[A, A] \in \Omega^2(X; E).$$
(27)

Explicitly,

$$F^{\rho}{}_{\mu\nu} = \nabla_{\mu}A^{\rho}{}_{\nu} - \nabla_{\nu}A^{\rho}{}_{\mu} + O(A^2),$$
(28)

where ∇_{μ} is the (Riemannian) covariant derivative of a tensor field. This is, to linear order, similar to the field strength in [3, (7.2.16)] with $(a_1, a_2) = (1, 0)$, which however depends on a symmetric rank-2 tensor, while here $A^{\mu}{}_{\rho}$ is an arbitrary rank (1,1) tensor. We shall discuss the symmetric case in section 4.2.

Under a gauge transformation, the field strength *F* then transforms covariantly:

$$\delta_{\epsilon}F = -\mathcal{L}_{\epsilon}F. \tag{29}$$

The fact that it is not invariant reminds us of the field strength in Yang–Mills theory (14). If we interpret the gauge parameter ϵ^{μ} as an infinitesimal diffeomorphism, then this implies that $F^{\mu}{}_{\nu\rho}$ transforms tensorially.

Now, we may postulate the equation of motion

$$\nabla^{\nu}F^{\mu}{}_{\nu\rho} = 0. \tag{30}$$

This is a diffeomorphism-invariant equation as long as we also transform $g_{\mu\nu}$ under diffeomorphisms, i.e.

$$\delta_{\epsilon}g_{\mu\nu} \coloneqq -\mathcal{L}_{\epsilon}g_{\mu\nu} = -\nabla_{\mu}\epsilon_{\nu} - \nabla_{\nu}\epsilon_{\mu}. \tag{31}$$

³If one wishes, one can work with a rescaled version of the Frölicher–Nijenhuis bracket, where the right-hand side of (5) comes with an additional factor of *g*; then then the $g \rightarrow 0$ limit corresponds to the free theory.

4.2 Symmetric theory

To make contact with the covariant fracton model, we now constrain $A^{\mu}{}_{\nu}$ to be symmetric. That is, we impose the following constraint:

$$g_{\mu\nu}A^{\nu}{}_{\rho} = g_{\rho\nu}A^{\nu}{}_{\mu}.$$
 (32)

This constraint is not gauge-invariant for arbitrary ϵ^{μ} since the right-hand side of (26) need not be symmetric. However, it *is* gauge-invariant if we restrict to 'diffeomorphisms' of the form

$$\epsilon^{\mu} = \partial^{\mu} \lambda \tag{33}$$

(known as the longitudinal diffeomorphisms [38]) for some smooth function $\lambda \in C^{\infty}(M)$, so that the resulting gauge transformation is

$$\delta_{\epsilon} A_{\mu\nu} = \partial_{\mu} \partial_{\nu} \lambda + \cdots, \qquad (34)$$

which to linearised order agrees with the covariant fracton gauge transformation (4). Now, we have the symmetrised equation of motion

$$\nabla^{\mu}F_{(\rho|\mu|\nu)} = 0. \tag{35}$$

The linearised equation of motion (with $g_{\mu\nu} = \eta_{\mu\nu}$ the Minkowski metric) is

$$0 = \partial^{\mu}(\partial_{\mu}A_{\rho\nu} - \partial_{\nu}A_{\rho\mu}) + \partial^{\mu}(\partial_{\mu}A_{\nu\rho} - \partial_{\rho}A_{\nu\mu}) = 2\partial^{2}A_{\rho\nu} - \partial_{\nu}\partial^{\mu}A_{\rho\mu} - \partial^{\mu}\partial_{\rho}A_{\nu\mu},$$
(36)

which is the equation of motion found in the covariant fracton theory of fractons [1, 2, 3].

References

- [1] Erica Bertolini and Nicola Maggiore. Maxwell theory of fractons. *Physical Review D*, 106(12):125008, 2022. arXiv:2209.01485, doi:10.1103/PhysRevD.106.125008.
- [2] Alberto Blasi and Nicola Maggiore. The theory of symmetric tensor field: From fractons to gravitons and back. *Physics Letters B*, 833:137304, 2022. arXiv:2207.05956, doi:10.1016/j.physletb.2022.137304.
- [3] Erica Bertolini. *Notes from the Bulk*. PhD thesis, Università di Genova, March 2024. doi:10.15167/bertolini-erica_phd2024-03-25.
- [4] Sagar Vijay, Jeongwan Haah (하정완), and Liang Fu (傅亮). A new kind of topological quantum order: A dimensional hierarchy of quasiparticles built from stationary excitations. *Physical Review B*, 92(23):235136, December 2015. arXiv:1505.02576, doi:10.1103/PhysRevB.92.235136.
- [5] Sagar Vijay, Jeongwan Haah (하정완), and Liang Fu (傅亮). Fracton topological order, generalized lattice gauge theory and duality. *Physical Review B*, 94(23):235157, December 2016. arXiv:1603.04442, doi:10.1103/PhysRevB.94.235157.
- [6] Rahul Mahajan Nandkishore and Michael A. Hermele. Fractons. Annual Review of Condensed Matter Physics, 10:295–313, 2019. arXiv:1803.11196, doi:10.1146/annurev-conmatphys-031218-013604.

- [7] Michael Pretko, Xie Chen (陈谐), and Yizhi You. Fracton phases of matter. *International Journal of Modern Physics A*, 35(06):2030003, 2020. arXiv:2001.01722, doi:10.1142/S0217751X20300033.
- [8] Claudio Chamon. Quantum glassiness. Physical Review Letters, 94(4):040402, February 2005. arXiv:cond-mat/0404182, doi:10.1103/physrevlett.94.040402.
- [9] Jeongwan Haah (하정완). Local stabilizer codes in three dimensions without string logical operators. *Physical Review A*, 83(4):042330, April 2011. arXiv:1101.1962, doi:10.1103/physreva.83.042330.
- [10] Sergey Borisovich Bravyi (Сергей Борисович Бравый) and Jeongwan Haah (하정완). Quantum self-correction in the 3D cubic code model. *Physical Review Letters*, 111(20):200501, November 2013. doi:10.1103/PhysRevLett.111.200501.
- [11] Michael Pretko. Subdimensional particle structure of higher rank U(1) spin liquids. *Physical Review B*, 95(11):115139, 2017. arXiv:1604.05329, doi:10.1103/PhysRevB.95.115139.
- [12] Michael Pretko. Generalized electromagnetism of subdimensional particles: A spin liquid story. *Physical Review B*, 96(3):035119, July 2017. arXiv:1606.08857, doi:10.1103/PhysRevB.96.035119.
- [13] Han Yan (闫寒). Hyperbolic fracton model, subsystem symmetry, and holography. *Physical Review B*, 99(15):155126, April 2019. arXiv:1807.05942, doi:10.1103/PhysRevB.99.155126.
- [14] Yizhi You, Fiona J. Burnell, and Taylor L. Hughes. Multipolar topological field theories: Bridging higher order topological insulators and fractons. *Physical Review B*, 103(24):245128, June 2021. arXiv:1909.05868, doi:10.1103/PhysRevB.103.245128.
- [15] Nathan Seiberg and Shu-Heng Shao (邵書珩). Exotic symmetries, duality, and fractons in 2+1-dimensional quantum field theory. *SciPost Physics*, 10(2):027, February 2021. arXiv:2003.10466, doi:10.21468/SciPostPhys.10.2.027.
- [16] Nathan Seiberg and Shu-Heng Shao (邵書珩). Exotic U(1) symmetries, duality, and fractons in 3+1-dimensional quantum field theory. *SciPost Physics*, 9(4):046, October 2020. arXiv:2004.00015, doi:10.21468/SciPostPhys.9.4.046.
- [17] Kevin Slagle. Foliated quantum field theory of fracton order. *Physical Review Letters*, 126(10):101603, March 2021. arXiv:2008.03852, doi:10.1103/PhysRevLett.126.101603.
- [18] Alessio Caddeo, Carlos Hoyos Badajoz, and Daniele Musso. Emergent dipole gauge fields and fractons. *Physical Review D*, 106(11):L111903, December 2022. arXiv:2206.12877, doi:10.1103/PhysRevD.106.L111903.
- [19] Xiaoyang Huang. A Chern-Simons theory for dipole symmetry. SciPost Physics, 15(4):153, October 2023. arXiv:2305.02492, doi:10.21468/SciPostPhys.15.4.153.

- [20] Alfredo Héctor Pérez Donoso, Stefan Prohazka, and Ali Seraj. Fracton infrared triangle. *Physical Review Letters*, 133(2):021603, July 2024. arXiv:2310.16683, doi:10.1103/PhysRevLett.133.021603.
- [21] Michael Pretko and Leo Radzihovsky (Лев Радзиховский). Fractonelasticity duality. *Physical Review Letters*, 120(19):195301, May 2018. arXiv:1711.11044, doi:10.1103/PhysRevLett.120.195301.
- [22] Michael Pretko, Zhengzheng Zhai, and Leo Radzihovsky (Лев Радзиховский). Crystal-to-fracton tensor gauge theory dualities. *Physical Review B*, 100(13):134113, October 2019. arXiv:1907.12577, doi:10.1103/PhysRevB.100.134113.
- [23] Andrey Gromov (Андрей Громов), Andrew Lucas, and Rahul Mahajan Nandkishore. Fracton hydrodynamics. *Physical Review Research*, 2(3):033124, July 2020. arXiv:2003.09429, doi:10.1103/PhysRevResearch.2.033124.
- [24] Jian-Keng Yuan (袁键铿), Shuai A. Chen (陈帅), and Peng Ye (叶鹏). Fractonic superfluids. *Physical Review Research*, 2(2):023267, June 2020. arXiv:1911.02876, doi:10.1103/PhysRevResearch.2.023267.
- [25] Juven C. Wang (王浚帆) and Shing-Tung Yau (丘成桐). Non-abelian gauged fracton matter field theory: Sigma models, superfluids, and vortices. *Physical Review Research*, 2(4):043219, November 2020. arXiv:1912.13485, doi:10.1103/PhysRevResearch.2.043219.
- [26] Kevin John Torres Grosvenor, Carlos Hoyos Badajoz, Francisco Peña-Benítez, and Piotr Surówka. Hydrodynamics of ideal fracton fluids. *Physical Review Research*, 3(4):043186, December 2021. arXiv:2105.01084, doi:10.1103/PhysRevResearch.3.043186.
- [27] Aleksander Głódkowski, Francisco Peña-Benítez, and Piotr Surówka. Hydrodynamics of dipole-conserving fluids. *Physical Review E*, 107(3):034142, March 2023. arXiv:2212.06848, doi:10.1103/PhysRevE.107.034142.
- [28] Lazaros Tsaloukidis (Λάζαρος Τσαλουχίδης) and Piotr Surówka. Elastic Liénard-Wiechert potentials of dynamical dislocations from tensor gauge theory in 2+1 dimensions. *Physical Review B*, 109(10):104118, March 2024. arXiv:2302.14092, doi:10.1103/PhysRevB.109.104118.
- [29] José Figueroa-O'Farrill, Alfredo Héctor Pérez Donoso, and Stefan Prohazka. Carroll/fracton particles and their correspondence. *Journal* of High Energy Physics, 06:207, June 2023. arXiv:2305.06730, doi:10.1007/JHEP06(2023)207.
- [30] José Figueroa-O'Farrill, Alfredo Héctor Pérez Donoso, and Stefan Prohazka. Quantum Carroll/fracton particles. *Journal of High Energy Physics*, 10:041, October 2023. arXiv:2307.05674, doi:10.1007/JHEP10(2023)041.
- [31] Zheng-Cheng Gu (顾正澄) and Xiao-Gang Wen (文小刚). A lattice bosonic model as a quantum theory of gravity, June 2006. arXiv:gr-qc/0606100, doi:10.48550/arXiv.gr-qc/0606100.

- [32] Cenke Xu (许岑珂). Gapless bosonic excitation without symmetry breaking: An algebraic spin liquid with soft gravitons. *Physical Review B*, 74(22):224433, December 2006. doi:10.1103/PhysRevB.74.224433.
- [33] Zheng-Cheng Gu (顾正澄) and Xiao-Gang Wen (文小刚). Emergence of helicity ±2 modes (gravitons) from qubit models. Nuclear Physics B, 863(1):90–129, October 2012. arXiv:0907.1203, doi:10.1016/j.nuclphysb.2012.05.010.
- [34] Cenke Xu (许岑珂) and Petr Hořava. Emergent gravity at a Lifshitz point from a Bose liquid on the lattice. *Physical Review D*, 81:104033, May 2010. arXiv:1003.0009, doi:10.1103/PhysRevD.81.104033.
- [35] Michael Pretko. Emergent gravity of fractons: Mach's principle revisited. *Physical Review D*, 96(2):024051, May 2017. arXiv:1702.07613, doi:10.1103/PhysRevD.96.024051.
- [36] Erica Bertolini, Alberto Blasi, Andrea Damonte, and Nicola Maggiore. Gauging fractons and linearized gravity. *Symmetry*, 15(4):945, April 2023. arXiv:2304.10789, doi:10.3390/sym15040945.
- [37] Erica Bertolini, Nicola Maggiore, and Giandomenico Palumbo. Covariant fracton gauge theory with boundary. *Physical Review D*, 108(2):025009, July 2023. arXiv:2306.13883, doi:10.1103/PhysRevD.108.025009.
- [38] Denis Dalmazi and Rafael Robson Lino dos Santos. The dimensional reduction of linearized spin-2 theories invariant under transverse diffeomorphisms. *The European Physics Journal C*, 81(6):547, June 2021. arXiv:2010.12051, doi:10.1140/epjc/s10052-021-09297-0.
- [39] Abhinav Prem, Michael Pretko, and Rahul Mahajan Nandkishore. Emergent phases of fractonic matter. *Physical Review B*, 97(8):085116, February 2018. arXiv:1709.09673, doi:10.1103/PhysRevB.97.085116.
- [40] Alfred Frölicher and Albert Nijenhuis. Theory of vector-valued differential forms: Part I. Derivations in the graded ring of differential forms. *Indagationes Mathematicae (Proceedings)*, 59:338–350, 1956. doi:10.1016/S1385-7258(56)50046-7.
- [41] Alfred Frölicher and Albert Nijenhuis. Invariance of vector form operations under mappings. *Commentarii Mathematici Helvetici*, 34:227–248, 1960. doi:10.1007/BF02565938.
- [42] Ivan Kolář, Peter Wolfram Michor, and Jan Slovák. Natural Operations in Differential Geometry. Springer-Verlag, Berlin, Heidelberg, February 1993. doi:10.1007/978-3-662-02950-3.
- [43] Steven Weinberg and Edward Witten. Limits on massless particles. *Physics Letters B*, 96(1–2):59–62, October 1980. doi:10.1016/0370-2693(80)90212-9.
- [44] Folkert Müller-Hoissen. Frölicher-Nijenhuis geometry and integrable matrix PDE systems, September 2024. arXiv:2409.01328, doi:10.48550/arXiv.2409.01328.

- [45] Antonio De Nicola and Ivan Yudin. Covariant Lie derivatives and Frölicher–Nijenhuis bracket on Lie Algebroids. International Journal of Geometric Methods in Modern Physics, 12(09):1560018, October 2015. arXiv:1412.2533, doi:10.1142/S021988781560018X.
- [46] Martin Bojowald, Alexei Kotov (Олексій Котов), and Thomas Strobl. Lie algebroid morphisms, poisson sigma models, and off-shell closed gauge symmetries. *Journal of Geometry and Physics*, 54(4):400–426, August 2005. arXiv:math/0406445, doi:10.1016/j.geomphys.2004.11.002.
- [47] Christoph Mayer and Thomas Strobl. Lie algebroid Yang–Mills with matter fields. *Journal of Geometry and Physics*, 59(12):1613–1623, December 2009. arXiv:0908.3161, doi:10.1016/j.geomphys.2009.07.018.
- [48] Alexei Kotov (Олексій Котов) and Thomas Strobl. Curving Yang-Mills-Higgs gauge theories. *Physical Review D*, 92(8):085032, October 2015. arXiv:1510.07654, doi:10.1103/PhysRevD.92.085032.
- [49] Simon-Raphael Fischer. Curved Yang–Mills–Higgs gauge theories in the case of massless gauge bosons. *Journal of Geometry and Physics*, 162:104104, April 2021. arXiv:2008.01507, doi:10.1016/j.geomphys.2021.104104.
- [50] Simon-Raphael Fischer. Geometry of curved Yang-Mills-Higgs gauge theories. PhD thesis, Université de Genève and Université Claude Bernard Lyon 1, June 2021. arXiv:2104.02175, doi:10.13097/archive-ouverte/unige:152555.
- [51] Simon-Raphael Fischer, Mehran Jalali Farahani, Hyungrok Kim, and Christian Saemann. Adjusted connections I: Differential cocycles for principal groupoid bundles with connection, June 2024. arXiv:2406.16755, doi:10.48550/arXiv.2406.16755.
- [52] Simon-Raphael Fischer, Mehran Jalali Farahani, Hyungrok Kim, and Christian Saemann. Topological classification of symmetry breaking and vacuum degeneracy, June 2024. arXiv:2406.17634, doi:10.48550/arXiv.2406.17634.
- [53] Kevin Slagle, Abhinav Prem, and Michael Pretko. Symmetric tensor gauge theories on curved spaces. *Annals of Physics*, 410:167910, November 2019. arXiv:1807.00827, doi:10.1016/j.aop.2019.167910.
- [54] Evangelos Afxonidis (Ευάγγελος Αυξωνίδης), Alessio Caddeo, Carlos Hoyos Badajoz, and Daniele Musso. Fracton gravity from spacetime dipole symmetry. *Physical Review D*, 109(6):065013, March 2024. arXiv:2311.01818, doi:10.1103/PhysRevD.109.065013.
- [55] Davide Rovere. Anomalies in covariant fracton theories, June 2024. arXiv:2406.06686, doi:10.48550/arXiv.2406.06686.
- [56] Andreas Kraft and Jonas Schnitzer. An introduction to L_{∞} -algebras and their homotopy theory for the working mathematician. *Reviews in Mathematical Physics*, 36(01):2330006, 2024. arXiv:2207.01861, doi:10.1142/S0129055X23300066.

- [57] Vladimir Viktorovich Dotsenko (Владимир Викторович Доценко), Sergey Viktorovich Shadrin (Сергей Викторович Шадрин), and Bruno Vallette. The twisting procedure, October 2018. arXiv:1810.02941, doi:10.48550/arXiv.1810.02941.
- [58] Vladimir Viktorovich Dotsenko (Владимир Викторович Доценко), Sergey Viktorovich Shadrin (Сергей Викторович Шадрин), and Bruno Vallette. *Maurer–Cartan Methods in Deformation Theory: the Twisting Procedure*, volume 488 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, UK, August 2023. arXiv:2212.11323, doi:10.1017/9781108963800.