STABILITY OF A CLASS OF SUPERCRITICAL VOLUME-FILLING CHEMOTAXIS-FLUID MODEL NEAR COUETTE FLOW

LILI WANG, WENDONG WANG, AND YI ZHANG

ABSTRACT. Consider a class of chemotaxis-fluid model incorporating a volumefilling effect in the sense of Painter and Hillen (Can. Appl. Math. Q. 2002; 10(4): 501-543), which is a supercritical parabolic-elliptic Keller-Segel system. As shown by Winkler et al., for any given mass, there exists a corresponding solution of the same mass that blows up in either finite or infinite time. In this paper, we investigate the stability properties of the two dimensional Patlak-Keller-Segel-type chemotaxis-fluid model near the Couette flow (Ay, 0) in $\mathbb{T} \times \mathbb{R}$, and show that the solutions are global in time as long as the initial cell mass $M < \frac{2\pi}{\sqrt{3}}$ and the shear flow is sufficiently strong (A is large enough).

Keywords: Keller-Segel; Chemotaxis-fluid model; Couette flow; Enhanced dissipation; Suppression; Blow-up

1. INTRODUCTION

Consider the following two-dimensional parabolic-elliptic chemotaxis-fluid model in $\mathbb{T} \times \mathbb{R}$ with $\mathbb{T} = [0, 2\pi)$:

$$\begin{cases} \partial_t n + u \cdot \nabla n = \nabla \cdot (\phi(n)\nabla n) - \nabla \cdot (\psi(n)\nabla c), \\ \Delta c + n - c = 0, \\ \partial_t u + u \cdot \nabla u + \nabla P = \Delta u + n\nabla \Phi_0, \quad \nabla \cdot u = 0, \\ (n, u)\big|_{t=0} = (n_{\rm in}, u_{\rm in}), \end{cases}$$
(1.1)

where the function n represents the cell density, c denotes the chemoattractant density, and u denotes the velocity of fluid. In addition, P is the pressure and Φ_0 is the potential function. The functions ϕ and ψ are assumed to belong to $C^2([0,\infty))$ and to satisfy $\phi > 0$ on $[0,\infty)$, $\psi(0) = 0$ and $\psi > 0$ on $(0,\infty)$. Based on a biased random walk analysis, Hillen and Painter [19] derived a functional link between the self-diffusivity $\phi(n)$ and the chemotactic sensitivity $\psi(n)$ by assuming that the cells' movement is inhibited near points where the cells are densely packed. That, in a non-dimensional version, takes the form

$$\phi(n) = Q(n) - uQ'(n), \quad \psi(n) = nQ(n),$$

Date: October 4, 2024.

where Q(n) denotes the density-dependent probability for a cell to find space somewhere in its current neighborhood. Since this probability is basically unknown, different choices for Q are conceivable, each of these providing a certain version of (1.1) that incorporates this so-called volume-filling effect (see [27]). Here we choose Q(n) = n + 1. Assume $\Phi_0 = y$ as Zeng-Zhang-Zi in [30]. Then (1.1) becomes

$$\begin{cases} \partial_t n + u \cdot \nabla n = \Delta n - \nabla \cdot ((n^2 + n) \nabla c), \\ \Delta c + n - c = 0, \\ \partial_t u + u \cdot \nabla u + \nabla P = \Delta u + ne_2, \quad \nabla \cdot u = 0, \\ (n, u)\big|_{t=0} = (n_{\rm in}, u_{\rm in}). \end{cases}$$
(1.2)

Let us briefly recall some developments on the system (1.1). If u = 0 and $\Phi_0 = 0$, the system (1.1) is reduced to the classical parabolic-elliptic Patlak-Keller-Segel system, which is proposed as a macroscopic model for chemotactic cell migration and was jointly developed by Patlak [20], Keller and Segel [12]. In details, the classical Patlak-Keller-Segel (PKS) system is stated as follows:

$$\begin{cases} \partial_t n = \nabla \cdot (\phi(n)\nabla n) - \nabla \cdot (\psi(n)\nabla c), \\ \tau \partial_t c = \triangle c + n - c, \end{cases}$$
(1.3)

where $\tau = 1, 0$ corresponds to the parabolic-parabolic case and the parabolic-elliptic case, respectively. For $\phi(n) = 1$ and $\psi(n) = n$, (1.3) is the most common formulation of the Patlak-Keller-Segel model, and the last term of $(1.3)_1$ has a critical Fujita exponent in 2D. As long as the dimension of space is higher than one, the solutions of the classical PKS system may blow up in finite time. The 2D PKS model of parabolic-parabolic has a critical mass of 8π , if the cell mass $M := ||n_{in}||_{L^1}$ is less than 8π , Calvez-Corrias [2] obtained the solutions of the system are global in time, while the cell mass is greater than 8π , the solutions will blow up in finite time by Schweyer in [21]. Moreover, the 2D parabolic-elliptic Patlak-Keller-Segel system is globally well-posed if and only if the total mass $M \leq 8\pi$ by Wei in [26]. When the spatial dimension is higher than two, the solutions of the PKS system will blow up for any initial mass, meaning that no mass threshold for aggregation exists in that case (for example, see [16], [17], [23], [28] etc.). Generally, for the supercritical case of $\psi(n) = n(n+1)^{\alpha}$ with $\alpha > 0$, blow-up phenomena may occur for any initial mass in 2D. For example, Winkler in [27] mentioned a blow-up criterion:

$$\frac{\psi(n)}{\phi(n)} \ge C_0 n \tag{1.4}$$

for some $C_0 > 0$ and sufficiently large n in the two-dimensional case, though he considered the parabolic-parabolic case at that time. When the spatial dimension is equal to two or greater than or equal to 3, Horstmann-Winkler [9] showed that there exist solutions of (1.3) that become unbounded for $\alpha > 0$. For more related results, we refer to [24, 25, 27, 29] and the references therein.

An interesting question: Can the stabilizing effect of the moving fluid suppress the finite-time blow-up?

As we all know, when considering $\phi(n) = 1$ and $\psi(n) = n$, some progresses have been made in proving the suppression of the chemotactic blow-up by the presence of fluid flow as follows:

2D case. For the parabolic-elliptic PKS system in 2D, Kiselev-Xu [13] suppressed the blow-up by stationary relaxation enhancing flows or time-dependent Yao-Zlatos near-optimal mixing flows in \mathbb{T}^d . He [8] investigated the suppression of blow-up by a large strictly monotone shear flow for the parabolic-parabolic PKS model in $\mathbb{T} \times \mathbb{R}$. For the coupled PKS-NS system, Zeng-Zhang-Zi firstly considered the 2D PKS-NS system near the Couette flow in $\mathbb{T} \times \mathbb{R}$ [30]. Li-Xiang-Xu [14] studied the suppression of blow-up in PKS-NS system via the Poiseuille flow in $\mathbb{T} \times \mathbb{R}$, and Cui-Wang considered Poiseuille flow with the Navier-slip boundary of PKS-NS system and obtained the solutions are global without any smallness condition [5].

3D case. For the PKS system of parabolic-elliptic case in 3D, there are few research results in this area. Bedrossian-He investigated the suppression of blow-up by shear flows in \mathbb{T}^3 and $\mathbb{T} \times \mathbb{R}^2$ by assuming the initial mass is less than 8π in [1]. Shi-Wang [22] considered the suppression effect of the flow $(y, y^2, 0)$ in $\mathbb{T}^2 \times \mathbb{R}$, and Deng-Shi-Wang [7] proved the Couette flow with a sufficiently large amplitude prevents the blow-up of solutions in the whole space. Besides, for the stability effect of buoyancy, Hu-Kiselev-Yao considered the blow-up suppression for the PKS system coupled with a fluid flow that obeys Darcy's law for incompressible porous media via buoyancy force [11]; see also [10]. For the 3D coupled PKS-NS system, Cui-Wang-Wang [6] considered the Couette flow in a finite channel, and obtained the solutions of the 3D linearized PKS-NS system are global as long as the initial cell mass is sufficiently small.

However, for $\phi(n) = 1$ and $\psi(n) = n(n+1)^{\alpha}$ with $\alpha > 0$, up to now it is still unknown whether the blow-up does not happen provided that the amplitude of some shear flow is sufficiently large. In this paper, we consider $\phi(n) = 1$, $\psi(n) = n(n+1)$ and the main goal is to investigate these types of issues.

Before expressing our main theorem, we first introduce a perturbation v around the two-dimensional Couette flow (Ay, 0), which v(t, x, y) = u(t, x, y) - (Ay, 0) satisfying $v|_{t=0} = v_{in} = (v_{1,in}, v_{2,in})$. Then we rewrite the system (1.2) into

$$\begin{cases} \partial_t n + Ay \partial_x n + v \cdot \nabla n - \Delta n = -\nabla \cdot (n^2 \nabla c) - \nabla (n \nabla c), \\ \Delta c + n - c = 0, \\ \partial_t v + Ay \partial_x v + \begin{pmatrix} Av_2 \\ 0 \end{pmatrix} + v \cdot \nabla v - \Delta v + \nabla P = \begin{pmatrix} 0 \\ n \end{pmatrix}, \\ \nabla \cdot v = 0. \end{cases}$$
(1.5)

To deal with the pressure P, we introduce the vorticity ω and the stream function Φ as follows

$$\omega := \partial_y v_1 - \partial_x v_2, \quad v = \nabla^{\perp} \Phi = (\partial_y \Phi, -\partial_x \Phi).$$

Then ω satisfies

$$\partial_t \omega + Ay \partial_x \omega - \Delta \omega + v \cdot \nabla \omega = -\partial_x n.$$

After the time re-scaling $t \mapsto \frac{t}{A}$, we get

$$\begin{cases} \partial_t n + y \partial_x n - \frac{1}{A} \bigtriangleup n = -\frac{1}{A} \nabla \cdot (vn) - \frac{1}{A} \nabla \cdot (n^2 \nabla c) - \frac{1}{A} \nabla \cdot (n \nabla c), \\ \bigtriangleup c + n - c = 0, \\ \partial_t \omega + y \partial_x \omega - \frac{1}{A} \bigtriangleup \omega = -\frac{1}{A} (\partial_x n + v \cdot \nabla \omega), \\ \nabla \cdot v = 0. \end{cases}$$
(1.6)

Our main result is stated as follows.

Theorem 1.1. Assume that the initial data $n_{\text{in}} \in L^1 \cap L^\infty(\mathbb{T} \times \mathbb{R})$, $v_{\text{in}} \in H^1(\mathbb{T} \times \mathbb{R})$ and the initial cell mass $M = ||n_{\text{in}}||_{L^1}$ satisfies $C^4_*M^2 < 3$, where C_* is the sharp Sobolev constant of

$$\|f\|_{L^{4}(\mathbb{R})} \leq C_{*} \|f\|_{L^{1}(\mathbb{R})}^{\frac{1}{2}} \|\partial_{y}f\|_{L^{2}(\mathbb{R})}^{\frac{1}{2}}.$$
(1.7)

Then there exists a positive constant A_1 depending on $||n_{in}||_{L^1 \cap L^{\infty}(\mathbb{T} \times \mathbb{R})}$ and $||v_{in}||_{H^1(\mathbb{T} \times \mathbb{R})}$, such that if $A \ge A_1$, the solution of (1.6) is global in time.

Remark 1.1. For $\phi(n) = 1$ and $\psi(n) = n(n+1)^{\alpha}$ with a critical Fujita exponent $\alpha > 0$ in the system (1.1), as shown by Winkler [27] or [4], for any given mass, there exists a corresponding solution of the same mass that blows up in either finite or infinite time. Theorem 1.1 shows that the Couette flow (sufficiently large) can suppress the blow-up of a positive initial mass for $\alpha = 1$ and the constant C_* comes from the estimate of $\|n_0\|_{L^{\infty}L^2}$ (see (3.11)), which seems to be exactly necessary. Hence, it is still unknown whether the Couette flow can suppress the blow-up for the case of $\alpha > 1$.

On the other hand, one can compute the detailed value of the above possible critical initial mass. For the following Gagliardo-Nirenberg interpolation inequality

$$\begin{cases} \|f\|_{L^{m+1}(\mathbb{R}^n)} \leq C_{q,m,p} \|f\|_{L^{q+1}(\mathbb{R}^n)}^{1-\theta} \|\nabla f\|_{L^p(\mathbb{R}^n)}^{\theta},\\ \theta = \frac{pn(m-q)}{(m+1)[n(p-q-1)+p(q+1)]}, \end{cases}$$

Nagy [18] (see also [3, 15]) obtained the best constant $C_{q,m,p}$ for the one-dimensional case in 1941. In particular, $C_{0,3,2} = \left(\frac{4\pi^2}{9}\right)^{-\frac{1}{4}}$ for p = 2, q = 0, m = 3 and n = 1, which corresponds to (1.7). Thus, we can give a specific upper bound for the initial cell mass M as follows.

Theorem 1.2. Assume that the initial data $n_{\text{in}} \in L^1 \cap L^\infty(\mathbb{T} \times \mathbb{R})$, $v_{\text{in}} \in H^1(\mathbb{T} \times \mathbb{R})$ and the initial cell mass $M = ||n_{\text{in}}||_{L^1} < \frac{2\pi}{\sqrt{3}}$. Then there exists a positive constant A_1 depending on $||n_{\text{in}}||_{L^1 \cap L^\infty(\mathbb{T} \times \mathbb{R})}$ and $||v_{\text{in}}||_{H^1(\mathbb{T} \times \mathbb{R})}$, such that if $A \ge A_1$, the solution of (1.6) is global in time.

Remark 1.2. Theorem 1.2 shows that the Couette flow (sufficiently large) can suppress the blow-up of a positive initial mass, whose upper bound is greater than $\frac{2\pi}{\sqrt{3}}$. It's interesting whether the initial mass is sharp.

Finally, we will list some notations used in this paper. Notations:

• Define the Fourier transform by

$$f(t, x, y) = \sum_{k \in \mathbb{Z}} \hat{f}(t, y) e^{ikx},$$

where $\hat{f}(t,y) = \frac{1}{|\mathbb{T}|} \int_{\mathbb{T}} f(t,x,y) e^{-ikx} dx.$

• For a given function f = f(t, x, y), write its x-part zero mode and x-part non-zero mode by

$$P_0 f = f_0 = \frac{1}{|\mathbb{T}|} \int_{\mathbb{T}} f(t, x, y) dx$$
, and $P_{\neq} f = f_{\neq} = f - f_0$.

Especially, we use $v_{k,0}$ and $v_{k,\neq}$ to represent the zero mode and non-zero mode of the velocity $v_k(k = 1, 2)$, respectively. Similarly, $\omega_{k,0}$ and $\omega_{k,\neq}$ represent the zero mode and non-zero mode of the vorticity $\omega_k(k = 1, 2)$.

• The norm of the L^p space and the time-space norm $||f||_{L^q L^p}$ are defined as

$$\|f\|_{L^{p}(\mathbb{T}\times\mathbb{R})} = \left(\int_{\mathbb{T}\times\mathbb{R}} |f|^{p} dx dy\right)^{\frac{1}{p}}, \quad \|f\|_{L^{q}L^{p}} = \left\|\|f\|_{L^{p}(\mathbb{T}\times\mathbb{R})}\right\|_{L^{q}(0,t)}.$$

• For $a \ge 0$, define the norm $||f||_{X_a}$ by

$$\begin{split} \|f\|_{X_a}^2 &= \|e^{aA^{-\frac{1}{3}t}}f\|_{L^{\infty}L^2}^2 + \|e^{aA^{-\frac{1}{3}t}}\nabla^{\perp}\triangle^{-1}\partial_x f\|_{L^2L^2}^2 \\ &+ \frac{1}{A^{\frac{1}{3}}}\|e^{aA^{-\frac{1}{3}t}}f\|_{L^2L^2}^2 + \frac{1}{A}\|e^{aA^{-\frac{1}{3}t}}\nabla f\|_{L^2L^2}^2. \end{split}$$

• Denote by M the total mass $||n(t)||_{L^1}$. Clearly, integration by parts and divergence theorem yield that

$$M := \|n(t)\|_{L^1} = \|n_{\rm in}\|_{L^1}.$$

• Throughout this paper, denote by C a positive constant independent of A, t and the initial data, and it may be different from line to line.

The rest part of this paper is organized as follows. In Section 2, the key idea and the proof of **Theorem 1.1** are presented. Section 3 is devoted to providing a collection of elementary lemmas including space-time estimates, which are essential for the proof of **Proposition 2.1-2.2**. In Section 4, we finish the proof of **Proposition 2.1**, where the smallness of the initial mass is necessary. The proof of **Proposition 2.2** is established in Section 5. In Section 6, we complete the proof of **Theorem 1.2**.

2. Key ideas and proof of Theorem 1.1

For given functions f and g, there hold

$$(fg)_0 = f_0g_0 + (f_{\neq}g_{\neq})_0, \quad (fg)_{\neq} = f_0g_{\neq} + f_{\neq}g_0 + (f_{\neq}g_{\neq})_{\neq}.$$

As the enhanced dissipation of fluid only affects the non-zero mode, and it is essential to separate the zero mode and the non-zero mode of system (1.6) as follows:

$$\begin{pmatrix}
\partial_t n_0 - \frac{1}{A} \Delta n_0 = -\frac{1}{A} \nabla \cdot (v_{\neq} n_{\neq})_0 - \frac{1}{A} \left(\partial_y (n_0^2 \partial_y c_0) + \partial_y ((n_{\neq}^2)_0 \partial_y c_0) \right) \\
- \frac{2}{A} \nabla \cdot (n_{\neq} n_0 \nabla c_{\neq})_0 - \frac{1}{A} \nabla \cdot ((n_{\neq}^2)_{\neq} \nabla c_{\neq})_0 \\
- \frac{1}{A} \partial_y (n_0 \partial_y c_0) - \frac{1}{A} \nabla \cdot (n_{\neq} \nabla c_{\neq})_0, \\
\partial_t \omega_0 - \frac{1}{A} \Delta \omega_0 = -\frac{1}{A} \nabla \cdot (v_{\neq} \omega_{\neq})_0, \\
v_0 = (\partial_y (\partial_{yy})^{-1} \omega_0, 0),
\end{cases}$$
(2.1)

and

$$\begin{cases}
\partial_t n_{\neq} + y \partial_x n_{\neq} - \frac{1}{A} \bigtriangleup n_{\neq} = -\frac{1}{A} \left(\nabla \cdot (v_0 n_{\neq}) + \nabla \cdot (v_{\neq} n_0) + \nabla \cdot (v_{\neq} n_{\neq})_{\neq} \right) \\
- \frac{2}{A} \partial_y (n_{\neq} n_0 \partial_y c_0) - \frac{1}{A} \left(\partial_y ((n_{\neq}^2)_{\neq} \partial_y c_0) + \nabla \cdot (n_0^2 \nabla c_{\neq}) \right) \\
- \frac{1}{A} \left(\nabla \cdot ((n_{\neq}^2)_0 \nabla c_{\neq}) + \nabla \cdot ((n_{\neq}^2)_{\neq} \nabla c_{\neq})_{\neq} \right) - \frac{2}{A} \nabla \cdot (n_{\neq} n_0 \nabla c_{\neq})_{\neq} \\
- \frac{1}{A} \left(\nabla \cdot (n_0 \nabla c_{\neq}) + \partial_y (n_{\neq} \partial_y c_0) + \nabla \cdot (n_{\neq} \nabla c_{\neq})_{\neq} \right), \\
- \bigtriangleup c_{\neq} + c_{\neq} = n_{\neq}, \\
\partial_t \omega_{\neq} + y \partial_x \omega_{\neq} - \frac{1}{A} \bigtriangleup \omega_{\neq} = -\frac{1}{A} \partial_x n_{\neq} - \frac{1}{A} \left(\nabla \cdot (v_0 \omega_{\neq}) + \nabla \cdot (v_{\neq} \omega_0) + \nabla \cdot (v_{\neq} \omega_{\neq})_{\neq} \right), \\
v_{\neq} = \nabla^{\perp} \bigtriangleup^{-1} \omega_{\neq}.
\end{cases}$$
(2.2)

Remark 2.1. Note that zero mode and non-zero mode can be controlled by their own functions. That is, for any $1 \le p \le \infty$, there hold

$$||f_0||_{L^p(\mathbb{T}\times\mathbb{R})} \le ||f||_{L^p(\mathbb{T}\times\mathbb{R})},$$

and

 $\|f_{\neq}\|_{L^p(\mathbb{T}\times\mathbb{R})} \le \|f\|_{L^p(\mathbb{T}\times\mathbb{R})} + \|f_0\|_{L^p(\mathbb{T}\times\mathbb{R})} \le 2\|f\|_{L^p(\mathbb{T}\times\mathbb{R})}.$

Remark 2.2. Due to div $v_0 = 0$, we have $\partial_y v_{2,0} = 0$ and $v_{2,0} = 0$. Then the zero mode of v_1 satisfies

$$\partial_t v_{1,0} - \frac{1}{A} \partial_{yy} v_{1,0} = -\frac{1}{A} \partial_y (v_{2,\neq} v_{1,\neq})_0.$$
(2.3)

We introduce an energy functional:

$$E(t) := \|\omega_{\neq}\|_{X_a} + \|n_{\neq}\|_{X_a},$$

with the initial norm

$$E_{\rm in} := \|\omega_{{\rm in},\neq}\|_{L^2} + \|n_{{\rm in},\neq}\|_{L^2}.$$

Let T be the terminal point of the largest range [0, T] such that the following hypothesis hold

$$E(t) \le 2K_{\neq},\tag{2.4}$$

$$||n||_{L^{\infty}L^{\infty}} \le 2K_{\infty},\tag{2.5}$$

for any $t \in [0, T]$, where K_{\neq} and K_{∞} will be calculated during the calculation.

The following propositions are key to obtaining the main results. Combining them with the local well-posedness of the system (1.6), we can deduce the global existence of the solution.

Proposition 2.1. Assume that the initial date $n_{in} \in L^1 \cap L^\infty(\mathbb{T} \times \mathbb{R})$ and $v_{in} \in H^1(\mathbb{T} \times \mathbb{R})$, under the conditions of (2.4) and (2.5), there exist a positive constant K_{\neq} depending on E_{in} , and a positive constant A_2 depending on K_{\neq}, K_{∞}, M and $\|v_{in}\|_{H^1}$, such that if $A \geq A_2$, there holds

$$E(t) \le K_{\neq},$$

for all $t \in [0, T]$.

Proposition 2.2. Assume that the initial date $n_{\text{in}} \in L^1 \cap L^\infty(\mathbb{T} \times \mathbb{R})$, $v_{\text{in}} \in H^1(\mathbb{T} \times \mathbb{R})$ and $C^4_*M^2 < 3$, under the conditions of (2.4) and (2.5), there exists a positive constant K_∞ depending on $\|n_{\text{in}}\|_{L^1 \cap L^\infty}$, such that

$$\|n\|_{L^{\infty}L^{\infty}} \le K_{\infty}$$

for all $t \in [0, T]$.

Proof of Theorem 1.1. Taking $A_1 = \max\{A_2, A_3\}$ and combining **Proposition 2.1** and **Proposition 2.2**, the proof is complete.

3. A Priori estimates

3.1. Space-time estimate. The following space-time estimate plays an important role to bound the non-zero modes of the solution to the system (1.6).

Lemma 3.1 (See Proposition 3.1, [30]). Let f satisfies

$$\partial_t f + y \partial_x f - \frac{1}{A} \Delta f = \partial_x f_1 + f_2 + \operatorname{div} f_3.$$
(3.1)

If $P_0f = P_0f_1 = P_0f_2 = P_0f_3 = 0$, then for $a \in [0, 4]$, we have

$$\|f\|_{X_a}^2 \le C\left(\|f(0)\|_{L^2}^2 + \|e^{aA^{-\frac{1}{3}t}}\nabla f_1\|_{L^2L^2}^2 + A^{\frac{1}{3}}\|e^{aA^{-\frac{1}{3}t}}f_2\|_{L^2L^2}^2 + A\|e^{aA^{-\frac{1}{3}t}}f_3\|_{L^2L^2}^2\right).$$

3.2. Elliptic estimates. We estimate c by elliptic energy method, which is similar as in [5] or [30].

Lemma 3.2. Let c_0 and n_0 be the zero mode of c and n, respectively, satisfying $-\partial_{uu}c_0 + c_0 = n_0.$

Then there hold

$$\begin{aligned} \|\partial_y^2 c_0(t)\|_{L^2} + \|\partial_y c_0(t)\|_{L^2} + \|c_0\|_{L^2} &\leq C \|n_0(t)\|_{L^2}, \\ \|\partial_y c_0(t)\|_{L^4} &\leq C \|n_0(t)\|_{L^2}, \end{aligned}$$

and

$$||c_0(t)||_{L^{\infty}} \le C ||n_0(t)||_{L^2}, \quad ||\partial_y c_0(t)||_{L^{\infty}} \le C ||n_0(t)||_{L^2},$$

for any $t \geq 0$.

Proof. The basic energy estimates yield

$$\|\partial_y^2 c_0(t)\|_{L^2}^2 + \|\partial_y c_0(t)\|_{L^2}^2 + \|c_0(t)\|_{L^2}^2 \le C \|n_0(t)\|_{L^2}^2.$$

Furthermore, using the Gagliardo-Nirenberg inequality, we have

$$\begin{aligned} \|\partial_y c_0(t)\|_{L^4} &\leq C \|\partial_y c_0(t)\|_{L^2}^{\frac{3}{4}} \|\partial_y^2 c_0(t)\|_{L^2}^{\frac{1}{4}} \leq C \|n_0(t)\|_{L^2}, \\ \|c_0(t)\|_{L^{\infty}} &\leq C \|c_0(t)\|_{L^2}^{\frac{1}{2}} \|\partial_y c_0(t)\|_{L^2}^{\frac{1}{2}} \leq C \|n_0(t)\|_{L^2}, \end{aligned}$$

and

$$\|\partial_y c_0(t)\|_{L^{\infty}} \le C \|\partial_y c_0(t)\|_{L^2}^{\frac{1}{2}} \|\partial_y^2 c_0(t)\|_{L^2}^{\frac{1}{2}} \le C \|n_0(t)\|_{L^2}.$$

Lemma 3.3. Let c_{\neq} and n_{\neq} be the non-zero mode of c and n, respectively, satisfying

$$-\triangle c_{\neq} + c_{\neq} = n_{\neq}.$$

Then there hold

 $\|\triangle c_{\neq}(t)\|_{L^{2}} + \|\nabla c_{\neq}(t)\|_{L^{2}} \le C \|n_{\neq}(t)\|_{L^{2}},$

and

 $\|\nabla c_{\neq}(t)\|_{L^4} \le C \|n_{\neq}(t)\|_{L^2},$

for any $t \geq 0$.

Proof. By integrating by parts, we have

$$\|\triangle c_{\neq}(t)\|_{L^{2}}^{2} + \|\nabla c_{\neq}(t)\|_{L^{2}}^{2} + \|c_{\neq}(t)\|_{L^{2}}^{2} \le C \|n_{\neq}(t)\|_{L^{2}}^{2}$$

Using the Gagliardo-Nirenberg inequality, we obtain

$$\|\nabla c_{\neq}(t)\|_{L^{4}} \leq C \|c_{\neq}(t)\|_{L^{2}}^{\frac{1}{4}} \|\triangle c_{\neq}(t)\|_{L^{2}}^{\frac{3}{4}} \leq C \|n_{\neq}(t)\|_{L^{2}}.$$

3.3. A Priori estimates for non-zero mode.

Lemma 3.4. Let f be a function such that $f_{\neq} \in H^1(\mathbb{T} \times \mathbb{R})$, there holds

$$||f_{\neq}||_{L^{2}(\mathbb{T}\times\mathbb{R})} \leq C ||\partial_{x}f_{\neq}||_{L^{2}(\mathbb{T}\times\mathbb{R})} \leq C ||\nabla f_{\neq}||_{L^{2}(\mathbb{T}\times\mathbb{R})}$$

Proof. It follows from Poincaré's inequality immediately and we omit it.

Lemma 3.5. Let v_{\neq} is determined by $(2.2)_4$. Then there holds

$$||e^{aA^{-\frac{1}{3}t}}v_{\neq}||_{L^{2}L^{\infty}} \le CA^{\frac{1}{4}}||\omega_{\neq}||_{X_{a}}.$$

Proof. Recall that $v = \nabla^{\perp} \Phi = (\partial_y \Phi, -\partial_x \Phi)$ and $\Delta \Phi = \omega$, and using the Fourier series, we get

$$v_{1,\neq} = \sum_{k\neq 0} \widehat{\partial}_y \widehat{\Phi}_{\neq}(k,y) e^{ikx},$$

$$v_{2,\neq} = \sum_{k\neq 0} \widehat{\partial}_x \widehat{\Phi}_{\neq}(k,y) e^{ikx} = -\sum_{k\neq 0} ik \widehat{\Phi}_{\neq}(k,y) e^{ikx},$$

and

$$\omega_{\neq} = \sum_{k \neq 0} (\partial_y^2 - k^2) \widehat{\Phi}_{\neq}(k, y) e^{ikx},$$

where $\hat{\Phi}_{\neq} = \frac{1}{2\pi} \int_{\mathbb{T}} \Phi_{\neq}(x, y) e^{-ikx} dx$. Then direct calculations indicate that

$$\|v_{1,\neq}\|_{L^2}^2 = 2\pi \int_{\mathbb{R}} \sum_{k\neq 0} \left(\widehat{\partial_y \Phi_{\neq}}(k,y) \overline{\widehat{\partial_y \Phi_{\neq}}}(k,y)\right) dy = 2\pi \sum_{k\neq 0} \|\widehat{\Phi}'_{\neq}(k,\cdot)\|_{L^2}^2, \qquad (3.2)$$

$$\|v_{2,\neq}\|_{L^2}^2 = 2\pi \int_{\mathbb{R}} \sum_{k\neq 0} \left(|k|^2 \widehat{\Phi}_{\neq}(k,y) \overline{\widehat{\Phi}}_{\neq}(k,y) \right) dy = 2\pi \sum_{k\neq 0} |k|^2 \|\widehat{\Phi}_{\neq}(k,\cdot)\|_{L^2}^2, \quad (3.3)$$

and

$$\begin{split} \|\omega_{\neq}\|_{L^{2}}^{2} &= 2\pi \int_{\mathbb{R}} \sum_{k \neq 0} \left((\partial_{y}^{2} - k^{2}) \widehat{\Phi}_{\neq}(k, y) (\partial_{y}^{2} - k^{2}) \overline{\widehat{\Phi}}_{\neq}(k, y) \right) dy \\ &= \sum_{k \neq 0} 2\pi \left(|k|^{4} \|\widehat{\Phi}_{\neq}(k, \cdot)\|_{L^{2}}^{2} + 2|k|^{2} \|\widehat{\Phi}_{\neq}'(k, \cdot)\|_{L^{2}}^{2} + \|\widehat{\Phi}_{\neq}''(k, \cdot)\|_{L^{2}}^{2} \right). \end{split}$$
(3.4)

It follows from (3.2)-(3.4) that

$$\|\nabla v_{\neq}\|_{L^{2}} \le C \|\omega_{\neq}\|_{L^{2}}.$$
(3.5)

Consider that $v_{\neq} = \sum_{k \neq 0} \hat{v}_{\neq}(k, y) e^{ikx}$, using the Gagliardo-Nirenberg inequality, we obtain

$$\|v_{\neq}\|_{L^{\infty}} \leq \sum_{k \neq 0} \|\widehat{v}_{\neq}(k,y)\|_{L^{\infty}_{y}} \leq C \sum_{k \neq 0} \|\widehat{v}_{\neq}(k,\cdot)\|_{L^{2}}^{\frac{1}{2}} \|\partial_{y}\widehat{v}_{\neq}(k,\cdot)\|_{L^{2}}^{\frac{1}{2}}.$$

Due to Hölder's inequality, there holds

$$\begin{aligned} \|v_{\neq}\|_{L^{\infty}} &\leq C \left(\sum_{k \neq 0} |k|^{1+2\varepsilon} \|\widehat{v}_{\neq}(k, \cdot)\|_{L^{2}} \|\partial_{y}\widehat{v}_{\neq}(k, \cdot)\|_{L^{2}} \right)^{\frac{1}{2}} \left(\sum_{k \neq 0} \frac{1}{|k|^{1+2\varepsilon}} \right)^{\frac{1}{2}} \\ &\leq C \left(\sum_{k \neq 0} |k|^{2\varepsilon} \|\widehat{\nabla v}_{\neq}(k, \cdot)\|_{L^{2}}^{2} \right)^{\frac{1}{2}}, \end{aligned}$$

where $\varepsilon \in (0, \frac{1}{2}]$. Using Hölder's inequality again and (3.5), we have

$$\begin{aligned} \|v_{\neq}\|_{L^{\infty}}^{2} &\leq C \sum_{k \neq 0} \|k \widehat{\nabla} v_{\neq}(k, \cdot)\|_{L^{2}}^{2\varepsilon} \|\widehat{\nabla} v_{\neq}(k, \cdot)\|_{L^{2}}^{2(1-\varepsilon)} \\ &\leq C \|\partial_{x} \nabla v_{\neq}\|_{L^{2}}^{2\varepsilon} \|\nabla v_{\neq}\|_{L^{2}}^{2(1-\varepsilon)} \leq C \|\partial_{x} \omega_{\neq}\|_{L^{2}}^{2\varepsilon} \|\omega_{\neq}\|_{L^{2}}^{2(1-\varepsilon)}, \end{aligned}$$

which follows that

$$\|e^{aA^{-\frac{1}{3}t}}v_{\neq}\|_{L^{2}L^{\infty}} \leq C \|e^{aA^{-\frac{1}{3}t}}\partial_{x}\omega_{\neq}\|_{L^{2}L^{2}}^{\varepsilon} \|e^{aA^{-\frac{1}{3}t}}\omega_{\neq}\|_{L^{2}L^{2}}^{1-\varepsilon} \leq CA^{\frac{1+2\varepsilon}{6}} \|\omega_{\neq}\|_{X_{a}},$$

for any $\varepsilon \in (0, \frac{1}{2}]$.

We complete the proof by selecting $\varepsilon = \frac{1}{4}$.

3.4. A Priori estimates for zero mode.

Lemma 3.6. Under the assumptions (2.4)-(2.5) and $C_*^4M^2 < 3$, there exists a positive constant A_3 depending on K_{\neq}, K_{∞} and M, such that if $A \ge A_3$, there hold

$$\|n_0\|_{L^{\infty}L^2} \le C\left(\|n_{\mathrm{in},0}\|_{L^2} + \frac{M^3}{(3 - C_*^4 M^2)^{\frac{3}{4}}} + 1\right) := H_1, \tag{3.6}$$

$$\|n_0\|_{L^{\infty}L^4} \le C\left(\|n_{\mathrm{in},0}\|_{L^4} + \|n_{\mathrm{in},0}\|_{L^2}^{\frac{3}{2}} + \frac{M^{\frac{9}{2}}}{(3 - C_*^4 M^2)^{\frac{9}{8}}} + 1\right) := H_2, \tag{3.7}$$

$$\|\omega_0\|_{L^{\infty}L^2} + A^{-\frac{1}{2}} \|\partial_y \omega_0\|_{L^2L^2} \le C \left(\|\omega_{\mathrm{in},0}\|_{L^2} + 1\right) := H_3, \tag{3.8}$$

$$\|v_{1,0}\|_{L^{\infty}L^{\infty}} \le C \left(\|v_{\mathrm{in},0}\|_{L^{2}} + \|\omega_{\mathrm{in},0}\|_{L^{2}} + 1\right) := H_{4}.$$
(3.9)

Proof. **I. Estimate of** $||n_0||_{L^{\infty}L^2}$. Multiplying both sides of $(2.1)_1$ by $2n_0$ and integrating it with y over \mathbb{R} , we obtain

$$\begin{aligned} \frac{d}{dt} \|n_0\|_{L^2}^2 &+ \frac{2}{A} \|\partial_y n_0\|_{L^2}^2 \\ &= \frac{2}{A} \int_{\mathbb{R}} (v_{\neq} n_{\neq})_0 \nabla n_0 dy + \frac{2}{A} \int_{\mathbb{R}} n_0^2 \partial_y c_0 \partial_y n_0 dy + \frac{2}{A} \int_{\mathbb{R}} (n_{\neq}^2)_0 \partial_y c_0 \partial_y n_0 dy \\ &+ \frac{4}{A} \int_{\mathbb{R}} (n_{\neq} n_0 \partial_y c_{\neq})_0 \partial_y n_0 dy + \frac{2}{A} \int_{\mathbb{R}} ((n_{\neq}^2)_{\neq} \partial_y c_{\neq})_0 \partial_y n_0 dy \\ &+ \frac{2}{A} \int_{\mathbb{R}} n_0 \partial_y c_0 \partial_y n_0 dy + \frac{2}{A} \int_{\mathbb{R}} (n_{\neq} \nabla c_{\neq})_0 \partial_y n_0 dy \\ &\leq \frac{5\delta}{A} \|\partial_y n_0\|_{L^2}^2 + \frac{C(\delta)}{A} \|(v_{\neq} n_{\neq})_0\|_{L^2}^2 + \frac{C(\delta)}{A} \|(n_{\neq}^2)_0 \partial_y c_0\|_{L^2}^2 \\ &+ \frac{C(\delta)}{A} \|(n_{\neq} n_0 \partial_y c_{\neq})_0\|_{L^2}^2 + \frac{C(\delta)}{A} \|((n_{\neq}^2)_{\neq} \partial_y c_{\neq})_0\|_{L^2}^2 + \frac{C(\delta)}{A} \|(n_{\neq} \nabla c_{\neq})_0\|_{L^2}^2 \\ &+ \frac{2}{A} \int_{\mathbb{R}} n_0^2 \partial_y c_0 \partial_y n_0 dy + \frac{2}{A} \int_{\mathbb{R}} n_0 \partial_y c_0 \partial_y n_0 dy, \end{aligned}$$
(3.10)

where δ is a positive constant. Due to integration by parts and $(2.1)_2$, there holds

$$\frac{2}{A} \int_{\mathbb{R}} n_0^2 \partial_y c_0 \partial_y n_0 dy + \frac{2}{A} \int_{\mathbb{R}} n_0 \partial_y c_0 \partial_y n_0 dy$$

$$= -\frac{2}{3A} \int_{\mathbb{R}} n_0^3 \partial_{yy} c_0 dy - \frac{1}{A} \int_{\mathbb{R}} n_0^2 \partial_{yy} c_0 dy$$

$$= \frac{2}{3A} \int_{\mathbb{R}} n_0^4 dy - \frac{2}{3A} \int_{\mathbb{R}} c_0 n_0^3 dy + \frac{1}{A} \int_{\mathbb{R}} n_0^3 dy - \frac{1}{A} \int_{\mathbb{R}} c_0 n_0^2 dy$$

$$:= I_1 + I_2 + I_3 + I_4.$$
(3.11)

Using the Gagliardo-Nirenberg inequality

$$\|n_0\|_{L^4} \le C_* \|n_0\|_{L^1}^{\frac{1}{2}} \|\partial_y n_0\|_{L^2}^{\frac{1}{2}}, \quad \|c_0\|_{L^{\infty}} \le C \|c_0\|_{L^2}^{\frac{1}{2}} \|\partial_y c_0\|_{L^2}^{\frac{1}{2}}, \qquad (3.12)$$
$$\|n_0\|_{L^3} \le C \|n_0\|_{L^2}^{\frac{5}{6}} \|\partial_y n_0\|_{L^2}^{\frac{1}{6}},$$

and Young's inequality, Lemma 3.2, we get

$$I_{1} \leq \frac{2}{3A} C_{*}^{4} \|n_{0}\|_{L^{1}}^{2} \|\partial_{y}n_{0}\|_{L^{2}}^{2} \leq \frac{2}{3A} C_{*}^{4} M^{2} \|\partial_{y}n_{0}\|_{L^{2}}^{2},$$

$$I_{2} \leq \frac{2}{3A} \|c_{0}\|_{L^{\infty}} \|n_{0}\|_{L^{3}}^{3} \leq \frac{C}{A} \|c_{0}\|_{L^{2}}^{\frac{1}{2}} \|\partial_{y}c_{0}\|_{L^{2}}^{\frac{1}{2}} \|n_{0}\|_{L^{2}}^{\frac{5}{2}} \|\partial_{y}n_{0}\|_{L^{2}}^{\frac{1}{2}},$$

$$\leq \frac{C}{A} \|n_{0}\|_{L^{2}}^{\frac{7}{2}} \|\partial_{y}n_{0}\|_{L^{2}}^{\frac{1}{2}} \leq \frac{\delta}{A} \|\partial_{y}n_{0}\|_{L^{2}}^{2} + \frac{C(\delta)}{A} \|n_{0}\|_{L^{2}}^{\frac{14}{3}},$$

$$I_{3} \leq \frac{C}{A} \|n_{0}\|_{L^{2}}^{\frac{5}{2}} \|\partial_{y}n_{0}\|_{L^{2}}^{\frac{1}{2}} \leq \frac{\delta}{A} \|\partial_{y}n_{0}\|_{L^{2}}^{2} + \frac{C(\delta)}{A} \|n_{0}\|_{L^{2}}^{\frac{10}{3}},$$

and

$$I_4 \le \frac{1}{A} \|c_0\|_{L^{\infty}} \|n_0\|_{L^2}^2 \le \frac{C}{A} \|n_0\|_{L^2}^3.$$

Combining with $I_1 - I_4$, (3.11) yields that

$$\frac{2}{A} \int_{\mathbb{R}} n_0^2 \partial_y c_0 \partial_y n_0 dy + \frac{2}{A} \int_{\mathbb{R}} n_0 \partial_y c_0 \partial_y n_0 dy$$

$$\leq \frac{C(\delta)}{A} \left(\|n_0\|_{L^2}^{\frac{14}{3}} + \|n_0\|_{L^2}^{\frac{10}{3}} + \|n_0\|_{L^2}^3 \right) + \left(2\delta + \frac{2}{3}C_*^4 M^2 \right) \frac{\|\partial_y n_0\|_{L^2}^2}{A},$$

substituting it into (3.10), we get

$$\frac{d}{dt} \|n_0\|_{L^2}^2 \leq -\left(2 - 7\delta - \frac{2}{3}C_*^4 M^2\right) \frac{\|\partial_y n_0\|_{L^2}^2}{A} + \frac{C(\delta)}{A} \|n_0\|_{L^2}^{\frac{14}{3}} + \frac{C(\delta)}{A} \|n_0\|_{L^2}^{\frac{10}{3}}
+ \frac{C}{A} \|n_0\|_{L^2}^3 + \frac{C(\delta)}{A} \left(\|v_{\neq} n_{\neq}\|_{L^2}^2 + \|(n_{\neq}^2)_{\neq} \partial_y c_{\neq}\|_{L^2}^2\right)
+ \frac{C(\delta)}{A} \left(\|n_{\neq} n_0 \partial_y c_{\neq}\|_{L^2}^2 + \|(n_{\neq}^2)_0 \partial_y c_0\|_{L^2}^2 + \|n_{\neq} \nabla c_{\neq}\|_{L^2}^2\right).$$
(3.13)

Notice that $C_*^4 M^2 < 3$, then letting

$$7\delta = 1 - \frac{1}{3}C_*^4M^2, \quad \tau = 2 - 7\delta - \frac{2}{3}C_*^4M^2 = 1 - \frac{1}{3}C_*^4M^2 > 0,$$

and using the Nash inequality

$$-\|\partial_y n_0\|_{L^2}^2 \le -C \frac{\|n_0\|_{L^2}^6}{\|n_0\|_{L^1}^4} \le -C \frac{\|n_0\|_{L^2}^6}{M^4},$$

it follows from (3.13) that

$$\frac{d}{dt} \|n_0\|_{L^2}^2 \leq -\frac{C\tau}{AM^4} \|n_0\|_{L^2}^3 \left(\|n_0\|_{L^2}^3 - \frac{M^4}{\tau} \|n_0\|_{L^2}^{\frac{5}{3}} - \frac{M^4}{\tau} \|n_0\|_{L^2}^{\frac{1}{3}} - \frac{M^4}{\tau} \right)
+ \frac{C}{A} \left(\|v_{\neq} n_{\neq}\|_{L^2}^2 + \|(n_{\neq}^2)_{\neq} \partial_y c_{\neq}\|_{L^2}^2 + \|n_{\neq} n_0 \partial_y c_{\neq}\|_{L^2}^2 \right)
+ \frac{C}{A} \left(\|(n_{\neq}^2)_0 \partial_y c_0\|_{L^2}^2 + \|n_{\neq} \nabla c_{\neq}\|_{L^2}^2 \right)
\leq -\frac{C\tau}{AM^4} \|n_0\|_{L^2}^3 \left(\|n_0\|_{L^2}^3 - \left(\frac{M^4}{\tau}\right)^{\frac{9}{4}} - 1 \right)
+ \frac{C}{A} \left(\|v_{\neq} n_{\neq}\|_{L^2}^2 + \|(n_{\neq}^2)_{\neq} \partial_y c_{\neq}\|_{L^2}^2 + \|n_{\neq} n_0 \partial_y c_{\neq}\|_{L^2}^2 \right)
+ \frac{C}{A} \left(\|(n_{\neq}^2)_0 \partial_y c_0\|_{L^2}^2 + \|n_{\neq} \nabla c_{\neq}\|_{L^2}^2 \right).$$
(3.14)

For all $t \ge 0$, we denote

$$\begin{aligned} G(t) &:= \frac{C}{A} \int_0^t \left(\|v_{\neq} n_{\neq}\|_{L^2}^2 + \|(n_{\neq}^2)_{\neq} \partial_y c_{\neq}\|_{L^2}^2 + \|n_{\neq} n_0 \partial_y c_{\neq}\|_{L^2}^2 \\ &+ \|(n_{\neq}^2)_0 \partial_y c_0\|_{L^2}^2 + \|n_{\neq} \nabla c_{\neq}\|_{L^2}^2 \right) ds. \end{aligned}$$

Due to Lemma 3.4 and (3.5), we find

$$\|v_{\neq}\|_{L^{2}} \le C \|\nabla v_{\neq}\|_{L^{2}} \le C \|\omega_{\neq}\|_{L^{2}}, \tag{3.15}$$

and notice that

$$\|n_0\|_{L^{\infty}L^2} \le \|n_0\|_{L^{\infty}L^1}^{\frac{1}{2}} \|n_0\|_{L^{\infty}L^{\infty}}^{\frac{1}{2}} \le CM^{\frac{1}{2}}K_{\infty}^{\frac{1}{2}},$$
(3.16)

then using assumptions (2.4)-(2.5), Lemma 3.2 and Lemma 3.3, direct calculations indicate that

$$\begin{split} G(t) &\leq \frac{C}{A} \left(\|v_{\neq}\|_{L^{2}L^{2}}^{2} \|n_{\neq}\|_{L^{\infty}L^{\infty}}^{2} + \|n_{\neq}^{2}\|_{L^{\infty}L^{\infty}}^{2} \|\partial_{y}c_{\neq}\|_{L^{2}L^{2}}^{2} \right) \\ &+ \frac{C}{A} \left(\|n_{\neq}n_{0}\|_{L^{\infty}L^{\infty}}^{2} \|\partial_{y}c_{\neq}\|_{L^{2}L^{2}}^{2} + \|(n_{\neq}^{2})_{0}\|_{L^{2}L^{2}}^{2} \|\partial_{y}c_{0}\|_{L^{\infty}L^{\infty}}^{2} \right) \\ &+ \frac{C}{A} \|n_{\neq}\|_{L^{\infty}L^{\infty}}^{2} \|\nabla c_{\neq}\|_{L^{2}L^{2}}^{2} \\ &\leq \frac{C}{A} \left(\|\omega_{\neq}\|_{L^{2}L^{2}}^{2} \|n_{\neq}\|_{L^{\infty}L^{\infty}}^{2} + \|n\|_{L^{\infty}L^{\infty}}^{4} \|n_{\neq}\|_{L^{2}L^{2}}^{2} \right) \\ &+ \frac{C}{A} \|n_{\neq}\|_{L^{\infty}L^{\infty}}^{2} \|n_{\neq}\|_{L^{2}L^{2}}^{2} \|n_{0}\|_{L^{\infty}L^{2}}^{2} + \frac{C}{A} \|n_{\neq}\|_{L^{\infty}L^{\infty}}^{2} \|n_{\neq}\|_{L^{2}L^{2}}^{2} \\ &\leq \frac{C}{A^{\frac{2}{3}}} \left(K_{\infty}^{2} \|\omega_{\neq}\|_{X_{a}}^{2} + K_{\infty}^{4} \|n_{\neq}\|_{X_{a}}^{2} + MK_{\infty}^{3} \|n_{\neq}\|_{X_{a}}^{2} + K_{\infty}^{2} \|n_{\neq}\|_{X_{a}}^{2} \right) \\ &\leq \frac{C}{A^{\frac{2}{3}}} \left(K_{\neq}^{2}K_{\infty}^{2} + K_{\neq}^{2}K_{\infty}^{4} + MK_{\neq}^{2}K_{\infty}^{3} \right) \leq C, \end{split}$$

provided that

$$A \ge \left(K_{\neq}^2 K_{\infty}^2 + K_{\neq}^2 K_{\infty}^4 + M K_{\neq}^2 K_{\infty}^3 \right)^{\frac{3}{2}}.$$

Then (3.14) can be written as follows

$$\frac{d}{dt}\left(\|n_0\|_{L^2}^2 - G(t)\right) \le -\frac{C\tau}{AM^4} \|n_0\|_{L^2}^3 \left(\|n_0\|_{L^2}^3 - \frac{M^9}{\tau^{\frac{9}{4}}} - 1\right),$$

which implies that

$$\|n_0\|_{L^{\infty}L^2} \le C\left(\|n_{\mathrm{in},0}\|_{L^2} + \frac{M^3}{\tau^{\frac{3}{4}}} + 1\right) = C\left(\|n_{\mathrm{in},0}\|_{L^2} + \frac{M^3}{(3 - C_*^4 M^2)^{\frac{3}{4}}} + 1\right).$$

Hence (3.6) holds.

II. Estimate of $||n_0||_{L^{\infty}L^4}$. Multiplying both sides of $(2.1)_1$ by $4n_0^3$ and integrating it with y over \mathbb{R} , we obtain

$$\begin{aligned} \frac{d}{dt} \|n_0\|_{L^4}^4 &+ \frac{3}{A} \|\partial_y n_0^2\|_{L^2}^2 \\ &= \frac{6}{A} \int_{\mathbb{R}} (v_{\neq} n_{\neq})_0 n_0 \nabla n_0^2 dy + \frac{4}{A} \int_{\mathbb{R}} n_0^2 \partial_y c_0 \partial_y n_0^3 dy + \frac{6}{A} \int_{\mathbb{R}} (n_{\neq}^2)_0 \partial_y c_0 n_0 \partial_y n_0^2 dy \\ &+ \frac{12}{A} \int_{\mathbb{R}} (n_{\neq} n_0 \partial_y c_{\neq})_0 n_0 \partial_y n_0^2 dy + \frac{6}{A} \int_{\mathbb{R}} ((n_{\neq}^2)_{\neq} \partial_y c_{\neq})_0 n_0 \partial_y n_0^2 dy \\ &+ \frac{6}{A} \int_{\mathbb{R}} (n_{\neq} \nabla c_{\neq})_0 n_0 \nabla n_0^2 dy + \frac{4}{A} \int_{\mathbb{R}} n_0 \partial_y c_0 \partial_y n_0^3 dy \\ &\leq \frac{1}{A} \|\partial_y n_0^2\|_{L^2}^2 + \frac{45}{A} \|(v_{\neq} n_{\neq})_0 n_0\|_{L^2}^2 + \frac{45}{A} \|(n_{\neq}^2)_0 \partial_y c_0 n_0\|_{L^2}^2 \\ &+ \frac{180}{A} \|(n_{\neq} n_0 \partial_y c_{\neq})_0 n_0\|_{L^2}^2 + \frac{45}{A} \|((n_{\neq}^2)_{\neq} \partial_y c_{\neq})_0 n_0\|_{L^2}^2 + \frac{45}{A} \|(n_{\neq} \nabla c_{\neq})_0 n_0\|_{L^2}^2 \\ &+ \frac{4}{A} \int_{\mathbb{R}} n_0^2 \partial_y c_0 \partial_y n_0^3 dy + \frac{4}{A} \int_{\mathbb{R}} n_0 \partial_y c_0 \partial_y n_0^3 dy. \end{aligned}$$
(3.17)

Similarly as (3.11), we find that

$$\frac{4}{A} \int_{\mathbb{R}} n_0^2 \partial_y c_0 \partial_y n_0^3 dy + \frac{4}{A} \int_{\mathbb{R}} n_0 \partial_y c_0 \partial_y n_0^3 dy$$

$$= -\frac{12}{5A} \int_{\mathbb{R}} n_0^5 \partial_{yy} c_0 dy - \frac{3}{A} \int_{\mathbb{R}} n_0^4 \partial_{yy} c_0 dy$$

$$= \frac{12}{5A} \int_{\mathbb{R}} n_0^6 dy - \frac{12}{5A} \int_{\mathbb{R}} c_0 n_0^5 dy + \frac{3}{A} \int_{\mathbb{R}} n_0^5 dy - \frac{3}{A} \int_{\mathbb{R}} c_0 n_0^4 dy$$

$$:= J_1 + J_2 + J_3 + J_4.$$
(3.18)

Applying the Gagliardo-Nirenberg inequality, Young's inequality and Lemma 3.2, there hold

$$J_{1} = \frac{12}{5A} \|n_{0}^{2}\|_{L^{3}}^{3} \leq \frac{C}{A} \|n_{0}^{2}\|_{L^{2}}^{\frac{5}{2}} \|\partial_{y}n_{0}^{2}\|_{L^{2}}^{\frac{1}{2}} \leq \frac{1}{4A} \|\partial_{y}n_{0}^{2}\|_{L^{2}}^{2} + \frac{C}{A} \|n_{0}^{2}\|_{L^{2}}^{\frac{10}{3}},$$

$$J_{2} \leq \frac{12}{5A} \|c_{0}\|_{L^{\infty}} \|n_{0}^{2}\|_{L^{\frac{5}{2}}}^{\frac{5}{2}} \leq \frac{C}{A} \|n_{0}\|_{L^{2}} \|n_{0}^{2}\|_{L^{2}}^{\frac{9}{4}} \|\partial_{y}n_{0}^{2}\|_{L^{2}}^{\frac{1}{4}},$$

$$\leq \frac{1}{4A} \|\partial_{y}n_{0}^{2}\|_{L^{2}}^{2} + \frac{C}{A} \|n_{0}\|_{L^{\infty}L^{2}}^{\frac{8}{7}} \|n_{0}^{2}\|_{L^{2}}^{\frac{18}{7}},$$

$$J_{3} \leq \frac{C}{A} \|n_{0}^{2}\|_{L^{2}}^{\frac{9}{4}} \|\partial_{y}n_{0}^{2}\|_{L^{2}}^{\frac{1}{4}} \leq \frac{1}{4A} \|\partial_{y}n_{0}^{2}\|_{L^{2}}^{2} + \frac{C}{A} \|n_{0}^{2}\|_{L^{2}}^{\frac{17}{7}},$$

and

$$J_4 \le \frac{3}{A} \|c_0\|_{L^{\infty}} \|n_0^2\|_{L^2}^2 \le \frac{C}{A} \|n_0\|_{L^{\infty}L^2} \|n_0^2\|_{L^2}^2.$$

Combing with $J_1 - J_4$, (3.18) follows that

$$\frac{4}{A} \int_{\mathbb{R}} n_0^2 \partial_y c_0 \partial_y n_0^3 dy + \frac{4}{A} \int_{\mathbb{R}} n_0 \partial_y c_0 \partial_y n_0^3 dy \\
\leq \frac{1}{A} \|\partial_y n_0^2\|_{L^2}^2 + \frac{C}{A} \|n_0^2\|_{L^2}^{\frac{10}{3}} + \frac{C}{A} (H_1^{\frac{8}{7}} + 1) \|n_0^2\|_{L^2}^{\frac{18}{7}} + \frac{C}{A} H_1 \|n_0^2\|_{L^2}^2.$$
(3.19)

Due to the Nash inequality

$$-\|\partial_y n_0^2\|_{L^2}^2 \le -C \frac{\|n_0^2\|_{L^2}^6}{\|n_0^2\|_{L^1}^4} \le -C \frac{\|n_0^2\|_{L^2}^6}{H_1^8},$$

and (3.19), we get from (3.17) that

$$\begin{aligned} \frac{d}{dt} \|n_{0}\|_{L^{4}}^{4} \\ &\leq -\frac{C}{AH_{1}^{8}} \|n_{0}^{2}\|_{L^{2}}^{6} + \frac{C}{A} \|n_{0}^{2}\|_{L^{2}}^{\frac{10}{3}} + \frac{C}{A} (H_{1}^{\frac{8}{7}} + 1) \|n_{0}^{2}\|_{L^{2}}^{\frac{18}{7}} + \frac{C}{A} H_{1} \|n_{0}^{2}\|_{L^{2}}^{2} \\ &+ \frac{C}{A} \|(v_{\neq}n_{\neq})_{0}n_{0}\|_{L^{2}}^{2} + \frac{C}{A} \|(n_{\neq}^{2})_{0}\partial_{y}c_{0}n_{0}\|_{L^{2}}^{2} + \frac{C}{A} \|(n_{\neq}n_{0}\partial_{y}c_{\neq})_{0}n_{0}\|_{L^{2}}^{2} \\ &+ \frac{C}{A} \|((n_{\neq}^{2})_{\neq}\partial_{y}c_{\neq})_{0}n_{0}\|_{L^{2}}^{2} + \frac{C}{A} \|(n_{\neq}\nabla c_{\neq})_{0}n_{0}\|_{L^{2}}^{2} \\ &\leq -\frac{C \|n_{0}^{2}\|_{L^{2}}^{2}}{AH_{1}^{8}} \left(\|n_{0}^{2}\|_{L^{2}}^{4} - H_{1}^{8}\|n_{0}^{2}\|_{L^{2}}^{\frac{4}{3}} - (H_{1}^{\frac{64}{7}} + H_{1}^{8})\|n_{0}^{2}\|_{L^{2}}^{\frac{4}{7}} - H_{1}^{9} \right) \\ &+ \frac{C}{A} \|(v_{\neq}n_{\neq})_{0}n_{0}\|_{L^{2}}^{2} + \frac{C}{A} \|(n_{\neq}^{2})_{0}\partial_{y}c_{0}n_{0}\|_{L^{2}}^{2} + \frac{C}{A} \|(n_{\neq}n_{0}\partial_{y}c_{\neq})_{0}n_{0}\|_{L^{2}}^{2} \\ &+ \frac{C}{A} \|((n_{\neq}^{2})_{\neq}\partial_{y}c_{\neq})_{0}n_{0}\|_{L^{2}}^{2} + \frac{C}{A} \|(n_{\neq}\nabla c_{\neq})_{0}n_{0}\|_{L^{2}}^{2} \\ &\leq -\frac{C \|n_{0}^{2}\|_{L^{2}}^{2}}{AH_{1}^{8}} \left(\|n_{0}^{2}\|_{L^{2}}^{4} - H_{1}^{12} - 1 \right) + \frac{C}{A} \|(v_{\neq}n_{\neq})_{0}n_{0}\|_{L^{2}}^{2} \\ &+ \frac{C}{A} \|((n_{\neq}^{2})_{0}\partial_{y}c_{0}n_{0}\|_{L^{2}}^{2} + \frac{C}{A} \|(n_{\neq}n_{0}\partial_{y}c_{\neq})_{0}n_{0}\|_{L^{2}}^{2} \\ &+ \frac{C}{A} \|((n_{\neq}^{2})_{\phi}\partial_{y}c_{\phi}n_{0}\|_{L^{2}}^{2} + \frac{C}{A} \|(n_{\neq}n_{0}\partial_{y}c_{\neq})_{0}n_{0}\|_{L^{2}}^{2} \end{aligned}$$

Denote

$$M(t) := \frac{C}{A} \int_0^t \left(\| (v_{\neq} n_{\neq})_0 n_0 \|_{L^2}^2 + \| (n_{\neq}^2)_0 \partial_y c_0 n_0 \|_{L^2}^2 + \| (n_{\neq} n_0 \partial_y c_{\neq})_0 n_0 \|_{L^2}^2 \right) \\ + \| ((n_{\neq}^2)_{\neq} \partial_y c_{\neq})_0 n_0 \|_{L^2}^2 + \| (n_{\neq} \nabla c_{\neq})_0 n_0 \|_{L^2}^2 \right) ds, \ \forall t \ge 0.$$

It follows from assumptions (2.4)-(2.5), (3.16), Lemma 3.2 and Lemma 3.3 that

$$\begin{split} M(t) &\leq \frac{C}{A} \left(\|v_{\neq}\|_{L^{2}L^{2}}^{2} \|n_{\neq}\|_{L^{\infty}L^{\infty}}^{2} \|n_{0}\|_{L^{\infty}L^{\infty}}^{2} + \|n_{\neq}\|_{L^{2}L^{2}}^{2} \|n_{\neq}\|_{L^{\infty}L^{\infty}}^{2} \|n_{0}\partial_{y}c_{0}\|_{L^{\infty}L^{\infty}}^{2} \right) \\ &+ \frac{C}{A} \left(\|\partial_{y}c_{\neq}\|_{L^{2}L^{2}}^{2} \|n_{\neq}n_{0}\|_{L^{\infty}L^{\infty}}^{2} \|n_{0}\|_{L^{\infty}L^{\infty}}^{2} + \|\partial_{y}c_{\neq}\|_{L^{2}L^{2}}^{2} \|n_{\neq}^{2}\|_{L^{\infty}L^{\infty}}^{2} \|n_{0}\|_{L^{\infty}L^{\infty}}^{2} \right) \\ &+ \frac{C}{A} \|n_{\neq}\|_{L^{\infty}L^{\infty}}^{2} \|\nabla c_{\neq}\|_{L^{2}L^{2}}^{2} \|n_{0}\|_{L^{\infty}L^{\infty}}^{2} \\ &\leq \frac{C}{A} \left(\|\omega_{\neq}\|_{L^{2}L^{2}}^{2} \|n\|_{L^{\infty}L^{\infty}}^{4} + \|n_{\neq}\|_{L^{2}L^{2}}^{2} \|n\|_{L^{\infty}L^{\infty}}^{4} \|n_{0}\|_{L^{\infty}L^{2}}^{2} \right) \\ &+ \frac{C}{A} \left(\|n_{\neq}\|_{L^{2}L^{2}}^{2} \|n\|_{L^{\infty}L^{\infty}}^{6} + \|n_{\neq}\|_{L^{2}L^{2}}^{2} \|n\|_{L^{\infty}L^{\infty}}^{4} \right) \\ &\leq \frac{C}{A^{\frac{2}{3}}} \left(\|\omega_{\neq}\|_{X_{a}}^{2} K_{\infty}^{4} + \|n_{\neq}\|_{X_{a}}^{2} M K_{\infty}^{5} + \|n_{\neq}\|_{X_{a}}^{2} K_{\infty}^{6} + \|n_{\neq}\|_{X_{a}}^{2} K_{\infty}^{4} \right) \\ &\leq \frac{C}{A^{\frac{2}{3}}} \left(K_{\neq}^{2} K_{\infty}^{4} + M K_{\neq}^{2} K_{\infty}^{5} + K_{\neq}^{2} K_{\infty}^{6} \right) \\ &\leq C, \end{split}$$

provided that

$$A \ge \left(K_{\neq}^{2} K_{\infty}^{4} + M K_{\neq}^{2} K_{\infty}^{5} + K_{\neq}^{2} K_{\infty}^{6}\right)^{\frac{3}{2}}.$$

Hence (3.20) can be rewritten as follows

$$\frac{d}{dt} \left(\|n_0\|_{L^4}^4 - M(t) \right) \le -\frac{C \|n_0^2\|_{L^2}^2}{AH_1^8} \left(\|n_0^2\|_{L^2}^4 - H_1^{12} - 1 \right),$$

which implies that

$$||n_0||_{L^{\infty}L^4} \le C\left(||n_{\mathrm{in},0}||_{L^4} + H_1^{\frac{3}{2}} + 1\right),$$

this gives (3.7).

III. Estimate of ω_0 . As ω_0 satisfies

$$\partial_t \omega_0 - \frac{1}{A} \partial_{yy} \omega_0 = -\frac{1}{A} \partial_y (v_{2,\neq} \omega_{\neq})_0,$$

multiplying it by $2\omega_0$ and integrating with y over \mathbb{R} , we obtain

$$\begin{aligned} \frac{d}{dt} \|\omega_0\|_{L^2}^2 + \frac{2}{A} \|\partial_y \omega_0\|_{L^2}^2 &= \frac{2}{A} \int_{\mathbb{R}} (v_{2,\neq} \omega_{\neq})_0 \partial_y \omega_0 dy \\ &\leq \frac{1}{A} \|\partial_y \omega_0\|_{L^2}^2 + \frac{C}{A} \|v_{2,\neq} \omega_{\neq}\|_{L^2}^2, \end{aligned}$$

which follows that

$$\|\omega_0\|_{L^{\infty}L^2}^2 + \frac{1}{A} \|\partial_y \omega_0\|_{L^2L^2}^2 \le \|\omega_{\mathrm{in},0}\|_{L^2}^2 + \frac{C}{A} \|v_{2,\neq}\|_{L^2L^{\infty}}^2 \|\omega_{\neq}\|_{L^{\infty}L^2}^2.$$
(3.21)

Using **Lemma 3.5**, if $A \ge K_{\neq}^8$, (3.21) implies that

$$\|\omega_0\|_{L^{\infty}L^2}^2 + \frac{1}{A} \|\partial_y \omega_0\|_{L^2L^2}^2 \le C \left(\|\omega_{\mathrm{in},0}\|_{L^2}^2 + \frac{\|\omega_{\neq}\|_{X_a}^4}{A^{\frac{1}{2}}} \right) \le C \left(\|\omega_{\mathrm{in},0}\|_{L^2}^2 + 1 \right).$$

This gives (3.8).

IV. Estimate of $v_{1,0}$. From (2.3), we find $v_{1,0}$ satisfies

$$\partial_t v_{1,0} - \frac{1}{A} \partial_{yy} v_{1,0} = -\frac{1}{A} \partial_y (v_{2,\neq} v_{1,\neq})_0,$$

and the basic energy estimates indicate that

$$\begin{aligned} \frac{d}{dt} \|v_{1,0}\|_{L^2}^2 &+ \frac{2}{A} \|\partial_y v_{1,0}\|_{L^2}^2 = \frac{2}{A} \int_{\mathbb{R}} (v_{2,\neq} v_{1,\neq})_0 \partial_y v_{1,0} dy \\ &\leq \frac{1}{A} \|\partial_y v_{1,0}\|_{L^2}^2 + \frac{1}{A} \|(v_{2,\neq} v_{1,\neq})_0\|_{L^2}^2 \end{aligned}$$

Integrating the above inequality with t, we get

$$\|v_{1,0}\|_{L^{\infty}L^{2}}^{2} + \frac{1}{A} \|\partial_{y}v_{1,0}\|_{L^{2}L^{2}}^{2} \leq \|v_{1,0}(0)\|_{L^{2}}^{2} + \frac{C}{A} \|v_{1,\neq}\|_{L^{2}L^{\infty}}^{2} \|v_{2,\neq}\|_{L^{\infty}L^{2}}^{2}.$$
(3.22)

Using Lemma 3.4 and Lemma 3.5, we have

$$\|v_{2,\neq}\|_{L^{\infty}L^{2}} \le C \|\partial_{x}v_{2,\neq}\|_{L^{\infty}L^{2}} \le C \|\omega_{\neq}\|_{L^{\infty}L^{2}} \le C \|\omega_{\neq}\|_{X_{a}},$$

and $||v_{1,\neq}||_{L^2L^{\infty}} \leq CA^{\frac{1}{4}} ||\omega_{\neq}||_{X_a}$. If $A \geq K_{\neq}^8$, then (3.22) yields that

$$\|v_{1,0}\|_{L^{\infty}L^{2}}^{2} + \frac{1}{A} \|\partial_{y}v_{1,0}\|_{L^{2}L^{2}}^{2} \leq \|v_{1,0}(0)\|_{L^{2}}^{2} + \frac{C}{A^{\frac{1}{2}}} \|\omega_{\neq}\|_{X_{a}}^{4} \leq C\left(\|v_{\mathrm{in},0}\|_{L^{2}}^{2} + 1\right). \quad (3.23)$$

Due to $\omega_0 = \partial_y v_{1,0}$, using Gagliardo-Nirenberg inequality, there holds

$$\|v_{1,0}\|_{L^{\infty}L^{\infty}} \le C \|v_{1,0}\|_{L^{\infty}L^{2}}^{\frac{1}{2}} \|\partial_{y}v_{1,0}\|_{L^{\infty}L^{2}}^{\frac{1}{2}} = C \|v_{1,0}\|_{L^{\infty}L^{2}}^{\frac{1}{2}} \|\omega_{0}\|_{L^{\infty}L^{2}}^{\frac{1}{2}}.$$

Combining this with (3.8) and (3.23), we obtain

$$\|v_{1,0}\|_{L^{\infty}L^{\infty}} \leq C \left(\|v_{\mathrm{in},0}\|_{L^{2}} + \|\omega_{\mathrm{in},0}\|_{L^{2}} + 1\right).$$

This proves (3.9).

4. The estimate of E(t) and proof of Proposition 2.1

Proof of Proposition 2.1. I. Estimate $\|\omega_{\neq}\|_{X_a}$. Applying Lemma 3.1 to $(2.2)_3$, and using (2.4), Lemma 3.5 and Lemma 3.6, we get

$$\begin{split} \|\omega_{\neq}\|_{X_{a}} \leq C \|\omega_{\mathrm{in},\neq}\|_{L^{2}} + \frac{C}{A} \|e^{aA^{-\frac{1}{3}t}} \nabla n_{\neq}\|_{L^{2}L^{2}} + \frac{C}{A^{\frac{1}{2}}} \|e^{aA^{-\frac{1}{3}t}} v_{1,0}\omega_{\neq}\|_{L^{2}L^{2}} \\ &+ \frac{C}{A^{\frac{1}{2}}} \left(\|e^{aA^{-\frac{1}{3}t}} v_{\neq}\omega_{0}\|_{L^{2}L^{2}} + \|e^{aA^{-\frac{1}{3}t}} (v_{\neq}\omega_{\neq})_{\neq}\|_{L^{2}L^{2}} \right) \\ \leq C \|\omega_{\mathrm{in},\neq}\|_{L^{2}} + \frac{C}{A} \|e^{aA^{-\frac{1}{3}t}} \nabla n_{\neq}\|_{L^{2}L^{2}} + \frac{C}{A^{\frac{1}{2}}} \|e^{aA^{-\frac{1}{3}t}} \omega_{\neq}\|_{L^{2}L^{2}} \|v_{1,0}\|_{L^{\infty}L^{\infty}} \\ &+ \frac{C}{A^{\frac{1}{2}}} \left(\|e^{aA^{-\frac{1}{3}t}} v_{\neq}\|_{L^{2}L^{\infty}} \|\omega_{0}\|_{L^{\infty}L^{2}} + \|e^{aA^{-\frac{1}{3}t}} v_{\neq}\|_{L^{2}L^{\infty}} \|\omega_{\neq}\|_{L^{\infty}L^{2}} \right) \\ \leq C \left(\|\omega_{\mathrm{in},\neq}\|_{L^{2}} + \frac{1}{A^{\frac{1}{2}}} \|n_{\neq}\|_{X_{a}} + \frac{H_{4}}{A^{\frac{1}{3}}} \|\omega_{\neq}\|_{X_{a}} + \frac{H_{3} + \|\omega_{\neq}\|_{X_{a}}}{A^{\frac{1}{4}}} \|\omega_{\neq}\|_{X_{a}} \right) \\ \leq C \left(\|\omega_{\mathrm{in},\neq}\|_{L^{2}} + \frac{H_{3}^{2} + H_{4}^{2} + K_{\neq}^{2} + 1}{A^{\frac{1}{4}}} \right). \end{split}$$

II. Estimate $||n_{\neq}||_{X_a}$. Applying **Lemma 3.1** to $(2.2)_1$, one deduces

$$\begin{split} \|n_{\neq}\|_{X_{a}} \leq C \|n_{\mathrm{in},\neq}\|_{L^{2}} + \frac{C}{A^{\frac{1}{2}}} \left(\|e^{aA^{-\frac{1}{3}t}}n_{\neq}n_{0}\partial_{y}c_{0}\|_{L^{2}L^{2}} + \|e^{aA^{-\frac{1}{3}t}}(n_{\neq}^{2})_{\neq}\partial_{y}c_{0}\|_{L^{2}L^{2}} \right) \\ &+ \frac{C}{A^{\frac{1}{2}}} \left(\|e^{aA^{-\frac{1}{3}t}}n_{0}^{2}\nabla c_{\neq}\|_{L^{2}L^{2}} + \|e^{aA^{-\frac{1}{3}t}}(n_{\neq}^{2})_{0}\nabla c_{\neq}\|_{L^{2}L^{2}} \right) \\ &+ \frac{C}{A^{\frac{1}{2}}} \left(\|e^{aA^{-\frac{1}{3}t}}(n_{\neq}^{2})_{\neq}\nabla c_{\neq}\|_{L^{2}L^{2}} + \|e^{aA^{-\frac{1}{3}t}}n_{\phi}\nabla c_{\neq}\|_{L^{2}L^{2}} \right) \\ &+ \frac{C}{A^{\frac{1}{2}}} \left(\|e^{aA^{-\frac{1}{3}t}}v_{1,0}n_{\neq}\|_{L^{2}L^{2}} + \|e^{aA^{-\frac{1}{3}t}}v_{\phi}n_{0}\|_{L^{2}L^{2}} \right) \\ &+ \frac{C}{A^{\frac{1}{2}}} \left(\|e^{aA^{-\frac{1}{3}t}}(v_{\neq}n_{\neq})_{\neq}\|_{L^{2}L^{2}} + \|e^{aA^{-\frac{1}{3}t}}n_{0}\nabla c_{\neq}\|_{L^{2}L^{2}} \right) \\ &+ \frac{C}{A^{\frac{1}{2}}} \left(\|e^{aA^{-\frac{1}{3}t}}n_{\phi}\partial_{y}c_{0}\|_{L^{2}L^{2}} + \|e^{aA^{-\frac{1}{3}t}}(n_{\phi}\nabla c_{\phi})_{\phi}\|_{L^{2}L^{2}} \right) \\ &+ \frac{C}{A^{\frac{1}{2}}} \left(\|e^{aA^{-\frac{1}{3}t}}n_{\phi}\partial_{y}c_{0}\|_{L^{2}L^{2}} + \|e^{aA^{-\frac{1}{3}t}}(n_{\phi}\nabla c_{\phi})_{\phi}\|_{L^{2}L^{2}} \right) . \end{split}$$

$$(4.2)$$

Using (3.16), assumptions (2.4)-(2.5), Lemma 3.2, Lemma 3.3, Lemma 3.5 and Lemma 3.6, it follows from (4.2) that

$$\begin{split} \|n_{\neq}\|_{X_{a}} \leq & C \|n_{\mathrm{in},\neq}\|_{L^{2}} + \frac{C}{A^{\frac{1}{2}}} \|e^{aA^{-\frac{1}{3}}t} n_{\neq}\|_{L^{2}L^{2}} \left(\|n_{0}\|_{L^{\infty}L^{\infty}} + \|n_{\neq}\|_{L^{\infty}L^{\infty}}\right) \|\partial_{y}c_{0}\|_{L^{\infty}L^{\infty}} \\ & + \frac{C}{A^{\frac{1}{2}}} \|e^{aA^{-\frac{1}{3}}t} \nabla c_{\neq}\|_{L^{2}L^{2}} \left(\|n_{\neq}\|_{L^{\infty}L^{\infty}}^{2} + \|n_{0}\|_{L^{\infty}L^{\infty}}^{2}\right) \\ & + \frac{C}{A^{\frac{1}{2}}} \|v_{1,0}\|_{L^{\infty}L^{\infty}} \|e^{aA^{-\frac{1}{3}}t} n_{\neq}\|_{L^{2}L^{2}} + \frac{C}{A^{\frac{1}{2}}} \left(\|n_{0}\|_{L^{\infty}L^{2}} + \|n_{\neq}\|_{L^{\infty}L^{2}}\right) \|e^{aA^{-\frac{1}{3}}t} v_{\neq}\|_{L^{2}L^{\infty}} \\ & + \frac{C}{A^{\frac{1}{2}}} \left(\|n_{0}\|_{L^{\infty}L^{\infty}} + \|n_{\neq}\|_{L^{\infty}L^{\infty}}\right) \|e^{aA^{-\frac{1}{3}}t} \nabla c_{\neq}\|_{L^{2}L^{2}} + \frac{C}{A^{\frac{1}{2}}} \|e^{aA^{-\frac{1}{3}}t} n_{\neq}\|_{L^{2}L^{2}} \|\partial_{y}c_{0}\|_{L^{\infty}L^{\infty}} \\ & \leq & C \|n_{\mathrm{in},\neq}\|_{L^{2}} + \frac{C}{A^{\frac{1}{3}}} \left(M^{\frac{1}{2}}K^{\frac{3}{2}}_{\infty} + K^{2}_{\infty} + H_{4} + K_{\infty} + M^{\frac{1}{2}}K^{\frac{1}{2}}_{\infty}\right) \|n_{\neq}\|_{X_{a}} \\ & + \frac{C}{A^{\frac{1}{4}}} \left(M^{\frac{1}{2}}K^{\frac{1}{2}}_{\infty} + K_{\neq}\right) \|\omega_{\neq}\|_{X_{a}} \\ & \leq & C \left(\|n_{\mathrm{in},\neq}\|_{L^{2}} + \frac{K^{4}_{\infty} + K^{2}_{\neq} + M^{4} + H^{2}_{4} + 1}{A^{\frac{1}{4}}}\right). \end{split}$$

To sum up, we conclude that

$$E(t) \le C\left(E_{\rm in} + \frac{K_{\infty}^4 + K_{\neq}^2 + M^4 + H_3^2 + H_4^2 + 1}{A^{\frac{1}{4}}}\right),\tag{4.3}$$

where

$$E_{\rm in} = \|\omega_{{\rm in},\neq}\|_{L^2} + \|n_{{\rm in},\neq}\|_{L^2}.$$

Let us denote $A_2 := (K_{\infty}^4 + K_{\neq}^2 + M^4 + H_3^2 + H_4^2 + 1)^4$. Thus if $A \ge A_2$, (4.3) implies that

$$E(t) \le C \left(E_{\rm in} + 1 \right) := K_{\neq}.$$

We complete the proof.

5. The L^{∞} estimate of the density and proof of Proposition 2.2

Proof of Propsition 2.2. We use mathematical induction to obtain the L^{∞} estimate of the density. Firstly, from (2.4) and (3.6), it is easy to know that

$$||n||_{L^{\infty}L^{2}} \le ||n_{0}||_{L^{\infty}L^{2}} + ||n_{\neq}||_{L^{\infty}L^{2}} \le H_{1} + 2K_{\neq} := D_{2}.$$
(5.1)

Step I. Estimate $||n||_{L^{\infty}L^4}$. It follows from Lemma 3.6 that

$$\|n_0\|_{L^{\infty}L^4} \le H_2,\tag{5.2}$$

then we only need to estimate $||n_{\neq}||_{L^{\infty}L^4}$. Multiplying $(2.2)_1$ by $4n_{\neq}^3$ and integrating the resulting equation over \mathbb{R} , noting that $\nabla \cdot v_0 = 0$, there holds

$$\begin{aligned} \frac{d}{dt} \|n_{\neq}^{2}\|_{L^{2}}^{2} + \frac{3}{A} \|\nabla n_{\neq}^{2}\|_{L^{2}}^{2} \\ &= \frac{6}{A} \int_{\mathbb{R}} n_{\neq} v_{\neq} n_{0} \nabla n_{\neq}^{2} dy + \frac{6}{A} \int_{\mathbb{R}} n_{\neq} (v_{\neq} n_{\neq})_{\neq} \nabla n_{\neq}^{2} dy + \frac{12}{A} \int_{\mathbb{R}} n_{\neq}^{2} n_{0} \nabla c_{0} \cdot \nabla n_{\neq}^{2} dy \\ &+ \frac{6}{A} \int_{\mathbb{R}} (n_{\neq}^{2})_{\neq} n_{\neq} \nabla c_{0} \cdot \nabla n_{\neq}^{2} dy + \frac{6}{A} \int_{\mathbb{R}} n_{\neq} n_{0}^{2} \nabla c_{\neq} \cdot \nabla n_{\neq}^{2} dy + \frac{6}{A} \int_{\mathbb{R}} n_{\neq} (n_{\neq}^{2})_{0} \nabla c_{\neq} \cdot \nabla n_{\neq}^{2} dy \\ &+ \frac{6}{A} \int_{\mathbb{R}} n_{\neq} ((n_{\neq}^{2})_{\neq} \nabla c_{\neq})_{\neq} \nabla n_{\neq}^{2} dy + \frac{12}{A} \int_{\mathbb{R}} n_{\neq} (n_{\neq} n_{0} \nabla c_{\neq})_{\neq} \nabla n_{\neq}^{2} dy \\ &+ \frac{6}{A} \int_{\mathbb{R}} n_{\neq} n_{0} \nabla c_{\neq} \cdot \nabla n_{\neq}^{2} dy + \frac{6}{A} \int_{\mathbb{R}} n_{\neq}^{2} \nabla c_{0} \cdot \nabla n_{\neq}^{2} dy + \frac{6}{A} \int_{\mathbb{R}} n_{\neq} (n_{\neq} \nabla c_{\neq})_{\neq} \nabla n_{\neq}^{2} dy \\ &\leq \frac{2}{A} \|\nabla n_{\neq}^{2}\|_{L^{2}}^{2} + \frac{C}{A} \left(\|n_{\neq} v_{\neq} n_{0}\|_{L^{2}}^{2} + \|n_{\neq} (v_{\neq} n_{\neq})_{\neq}\|_{L^{2}}^{2} + \|n_{\varphi}^{2} n_{0} \partial_{y} c_{0}\|_{L^{2}}^{2} \right) \\ &+ \frac{C}{A} \left(\|n_{\varphi}^{2}|_{\neq} n_{\varphi} \partial_{y} c_{0}\|_{L^{2}}^{2} + \|n_{\varphi} (n_{\varphi} \nabla c_{\varphi})_{\neq}\|_{L^{2}}^{2} + \|n_{\varphi} n_{0} \nabla c_{\varphi}\|_{L^{2}}^{2} \right) \\ &+ \frac{C}{A} \left(\|n_{\varphi}^{2}|_{y} c_{0}\|_{L^{2}}^{2} + \|n_{\varphi} (n_{\varphi} \nabla c_{\varphi})_{\neq}\|_{L^{2}}^{2} \right) \\ &+ \frac{C}{A} \left(\|n_{\varphi}^{2}\partial_{y} c_{0}\|_{L^{2}}^{2} + \|n_{\varphi} (n_{\varphi} \nabla c_{\varphi})_{\neq}\|_{L^{2}}^{2} \right). \end{aligned}$$

$$(5.3)$$

For any $t \ge 0$, denote

$$\begin{split} N(t) &:= \frac{C}{A} \int_0^t \left(\|n_{\neq} v_{\neq} n_0\|_{L^2}^2 + \|n_{\neq} (v_{\neq} n_{\neq})_{\neq}\|_{L^2}^2 + \|n_{\neq}^2 n_0 \partial_y c_0\|_{L^2}^2 \\ &+ \|(n_{\neq}^2)_{\neq} n_{\neq} \partial_y c_0\|_{L^2}^2 + \|n_{\neq} n_0^2 \nabla c_{\neq}\|_{L^2}^2 + \|n_{\neq} (n_{\neq}^2)_0 \nabla c_{\neq}\|_{L^2}^2 \\ &+ \|n_{\neq} ((n_{\neq}^2)_{\neq} \nabla c_{\neq})_{\neq}\|_{L^2} + \|n_{\neq} (n_{\neq} n_0 \nabla c_{\neq})_{\neq}\|_{L^2}^2 + \|n_{\neq} n_0 \nabla c_{\neq}\|_{L^2}^2 \\ &+ \|n_{\neq}^2 \partial_y c_0\|_{L^2}^2 + \|n_{\neq} (n_{\neq} \nabla c_{\neq})_{\neq}\|_{L^2}^2 \right) ds, \end{split}$$

then using the Hölder's inequality, assumptions (2.4)-(2.5), (3.16), Lemma 3.2 and Lemma 3.3, we get

$$\begin{split} N(t) &\leq \frac{C}{A} \|v_{\neq}\|_{L^{2}L^{2}}^{2} \|n\|_{L^{\infty}L^{\infty}}^{4} + \frac{C}{A} \|n_{\neq}\|_{L^{2}L^{2}}^{2} \|n\|_{L^{\infty}L^{\infty}}^{4} \|\partial_{y}c_{0}\|_{L^{\infty}L^{\infty}}^{2} \\ &+ \frac{C}{A} \|\nabla c_{\neq}\|_{L^{2}L^{2}}^{2} \left(\|n\|_{L^{\infty}L^{\infty}}^{6} + \|n\|_{L^{\infty}L^{\infty}}^{4}\right) + \frac{C}{A} \|n_{\neq}\|_{L^{2}L^{2}}^{2} \|n\|_{L^{\infty}L^{\infty}}^{2} \|\partial_{y}c_{0}\|_{L^{\infty}L^{\infty}}^{2} \\ &\leq \frac{C}{A^{\frac{2}{3}}} \|\omega_{\neq}\|_{X_{a}}^{2} K_{\infty}^{4} + \frac{C}{A^{\frac{2}{3}}} \|n_{\neq}\|_{X_{a}}^{2} \left(MK_{\infty}^{5} + K_{\infty}^{6} + K_{\infty}^{4} + MK_{\infty}^{3}\right) \\ &\leq \frac{C}{A^{\frac{2}{3}}} \left(K_{\neq}^{2}K_{\infty}^{4} + MK_{\neq}^{2}K_{\infty}^{3} + MK_{\neq}^{2}K_{\infty}^{5} + K_{\neq}^{2}K_{\infty}^{6}\right) \leq C, \end{split}$$

provided that

$$A \ge \left(K_{\neq}^2 K_{\infty}^4 + M K_{\neq}^2 K_{\infty}^3 + M K_{\neq}^2 K_{\infty}^5 + K_{\neq}^2 K_{\infty}^6 \right)^{\frac{3}{2}}.$$

Hence (5.3) implies that

$$\frac{d}{dt} \left(\|n_{\neq}^2\|_{L^2}^2 - N(t) \right) \le -\frac{1}{A} \|\nabla n_{\neq}^2\|_{L^2}^2 \le 0.$$
(5.4)

By integrating both sides of (5.4) with t, we obtain

$$\|n_{\neq}\|_{L^{\infty}L^{4}} \le C\left(\|n_{\text{in}}\|_{L^{4}}+1\right) := H_{5}.$$
(5.5)

Combining the estimates of (3.7) and (5.5), one deduces

$$||n||_{L^{\infty}L^4} \le ||n_0||_{L^{\infty}L^4} + ||n_{\neq}||_{L^{\infty}L^4} \le H_2 + H_5 := D_4.$$
(5.6)

Step II. Estimate $||n||_{L^{\infty}L^p}$. For $p = 2^j$ with j > 2, multiplying $(1.6)_1$ by $2pn^{2p-1}$, and integrating by parts the resulting equation over $\mathbb{T} \times \mathbb{R}$, using $\nabla \cdot v = 0$, $(1.6)_2$ and $||c||_{L^2} \leq ||n||_{L^2}$, one deduces

$$\frac{d}{dt} \|n^{p}\|_{L^{2}}^{2} + \frac{2(2p-1)}{pA} \|\nabla n^{p}\|_{L^{2}}^{2}
= \frac{2p(2p-1)}{A(2p+1)} \int_{\mathbb{T}\times\mathbb{R}} \nabla c \cdot \nabla n^{2p+1} dx dy + \frac{2(2p-1)}{A} \int_{\mathbb{T}\times\mathbb{R}} n^{p} \nabla c \cdot \nabla n^{p} dx dy \qquad (5.7)
:= S_{1} + S_{2}.$$

For S_1 , using $(1.6)_2$, the Gagliardo-Nirenberg inequality and the Young's inequality, there holds

$$S_{1} = -\frac{2p(2p-1)}{A(2p+1)} \int_{\mathbb{T}\times\mathbb{R}} n^{2p+1} \triangle c dx dy$$

$$= \frac{2p(2p-1)}{A(2p+1)} \int_{\mathbb{T}\times\mathbb{R}} n^{2p+2} dx dy - \frac{2p(2p-1)}{A(2p+1)} \int_{\mathbb{T}\times\mathbb{R}} n^{2p+1} c dx dy$$

$$\leq C \frac{2p(2p-1)}{A(2p+1)} \left(\|n^{p}\|_{L^{2}}^{2} \|\nabla n^{p}\|_{L^{2}}^{\frac{2}{p}} + \|n\|_{L^{2}} \|n^{p}\|_{L^{2}} \|\nabla n^{p}\|_{L^{2}}^{\frac{p+1}{p}} \right)$$

$$\leq \frac{2p-1}{2pA} \|\nabla n^{p}\|_{L^{2}}^{2} + \frac{C(2p-1)(p-1)}{Ap} \left(\frac{p}{2p+1} \right)^{\frac{p}{p-1}} \|n^{p}\|_{L^{2}}^{\frac{2p}{2p-1}}$$

$$+ \frac{C(p-1)(2p-1)}{Ap(p+1)} \left(\frac{p(p+1)}{2p+1} \right)^{\frac{2p}{p-1}} \|n^{p}\|_{L^{2}}^{\frac{2p}{p-1}}.$$

(5.8)

For S_2 , by Hölder's inequality, Young's inequality and Gagliardo-Nirenberg inequality, one obtain

$$S_{2} \leq \frac{2(2p-1)}{A} \|\nabla n^{p}\|_{L^{2}} \|n^{p}\nabla c\|_{L^{2}}$$

$$\leq \frac{2p-1}{4pA} \|\nabla n^{p}\|_{L^{2}}^{2} + \frac{2p(2p-1)}{A} \|n^{p}\nabla c\|_{L^{2}}^{2}$$

$$\leq \frac{2p-1}{4pA} \|\nabla n^{p}\|_{L^{2}}^{2} + \frac{Cp(2p-1)}{A} \|n^{p}\|_{L^{2}} \|\nabla n^{p}\|_{L^{2}} \|\nabla c\|_{L^{4}}^{2}$$

$$\leq \frac{2p-1}{2pA} \|\nabla n^{p}\|_{L^{2}}^{2} + \frac{Cp^{3}(2p-1)}{A} \|n^{p}\|_{L^{2}}^{2} \|\nabla c\|_{L^{4}}^{4}.$$
(5.9)

Combining (5.8) with (5.9), we get by (5.7) that

$$\frac{d}{dt} \|n^{p}\|_{L^{2}}^{2} \leq -\frac{1}{A} \|\nabla n^{p}\|_{L^{2}}^{2} + \frac{Cp}{A} \|n^{p}\|_{L^{2}}^{\frac{2p}{p-1}} + \frac{Cp^{3}}{A} \|n\|_{L^{2}}^{\frac{2p}{p-1}} \|n^{p}\|_{L^{2}}^{\frac{2p}{p-1}} + \frac{Cp^{4}}{A} \|n^{p}\|_{L^{2}}^{2} \|\nabla c\|_{L^{4}}^{4}.$$
(5.10)

Thanks to Nash inequality

$$-\|\nabla n^p\|_{L^2}^2 \le -C\frac{\|n^p\|_{L^2}^4}{\|n^p\|_{L^1}^2},$$

(5.10) yields that

$$\begin{split} \frac{d}{dt} \|n^{p}\|_{L^{2}}^{2} &\leq -\frac{C}{A} \frac{\|n^{p}\|_{L^{2}}^{4}}{\|n^{p}\|_{L^{1}}^{2}} + \frac{Cp}{A} \|n^{p}\|_{L^{2}}^{\frac{2p}{p-1}} + \frac{Cp^{3}}{A} \|n\|_{L^{2}}^{\frac{2p}{p-1}} \|n^{p}\|_{L^{2}}^{\frac{2p}{p-1}} \\ &\quad + \frac{Cp^{4}}{A} \|n^{p}\|_{L^{2}}^{2} \|\nabla c\|_{L^{4}}^{4} \\ &\leq -\frac{C\|n^{p}\|_{L^{2}}^{2}}{A\|n^{p}\|_{L^{1}}^{2}} \left(\|n^{p}\|_{L^{2}}^{2} - p\|n^{p}\|_{L^{1}}^{2} \|n^{p}\|_{L^{2}}^{\frac{2p}{p-1}} - p^{3}\|n\|_{L^{2}}^{\frac{2p}{p-1}} \|n^{p}\|_{L^{1}}^{2} \|n^{p}\|_{L^{2}}^{\frac{2}{p-1}} \\ &\quad - p^{4}\|n^{p}\|_{L^{1}}^{2} \left(\|n^{p}\|_{L^{2}}^{2} - p\frac{p-1}{p-2}\|n^{p}\|_{L^{1}}^{2(p-1)} - p^{3}\|n\|_{L^{2}}^{\frac{2p}{p-1}} \|n^{p}\|_{L^{2}}^{\frac{2p}{p-1}} \\ &\quad - p^{4}\|n^{p}\|_{L^{1}}^{2} \left(\|n^{p}\|_{L^{2}}^{2} - p\frac{p-1}{p-2}\|n^{p}\|_{L^{1}}^{2(p-1)} - p^{3(p-1)}\|n\|_{L^{2}}^{\frac{2p}{p-1}} \|n^{p}\|_{L^{1}}^{\frac{2(p-1)}{p-2}} \\ &\quad - p^{4}\|n^{p}\|_{L^{1}}^{2} \|\nabla c\|_{L^{4}}^{4} \right), \end{split}$$

which follows that

$$\begin{aligned} \|n^{p}\|_{L^{\infty}L^{2}}^{2} \leq C \Biggl(\|n^{p}(0)\|_{L^{2}}^{2} + p^{\frac{p-1}{p-2}} \|n^{p}\|_{L^{\infty}L^{1}}^{\frac{2(p-1)}{p-2}} + p^{\frac{3(p-1)}{p-2}} \|n\|_{L^{\infty}L^{2}}^{\frac{2p}{p-2}} \|n^{p}\|_{L^{\infty}L^{1}}^{\frac{2(p-1)}{p-2}} \\ + p^{4} \|n^{p}\|_{L^{\infty}L^{1}}^{2} \|\nabla c\|_{L^{\infty}L^{4}}^{4} \Biggr). \end{aligned}$$

$$(5.11)$$

Using (5.1) and elliptic estimation, we rewrite (5.11) into

$$\begin{split} \|n\|_{L^{\infty}L^{2p}}^{2p} &\leq C\left(\|n_{\mathrm{in}}\|_{L^{2p}}^{2p} + p^{\frac{p-1}{p-2}}\|n\|_{L^{\infty}L^{p}}^{\frac{2p(p-1)}{p-2}} + p^{\frac{3(p-1)}{p-2}}D_{2}^{\frac{2p}{p-2}}\|n\|_{L^{\infty}L^{p}}^{\frac{2p(p-1)}{p-2}} + p^{4}D_{4}^{4}\|n\|_{L^{\infty}L^{p}}^{2p}\right) \\ &\leq \tilde{C}\left(1 + p^{4}\|n\|_{L^{\infty}L^{p}}^{\frac{2p(p-1)}{p-2}} + \frac{p-2}{p-1}p^{\frac{4(p-1)}{p-2}}\|n\|_{L^{\infty}L^{p}}^{\frac{2p(p-1)}{p-2}}\right) \\ &\leq \tilde{C}\max\left(1, p^{8}\|n\|_{L^{\infty}L^{p}}^{\frac{2p(p-1)}{p-2}}\right), \end{split}$$

$$(5.12)$$

where \tilde{C} is a positive constant depending on $||n_{\rm in}||_{L^{\infty}}$, D_2 and D_4 . **Step III. Estimate** $||n||_{L^{\infty}L^{\infty}}$. Recall $p = 2^j$ for $j \ge 2$, then using iterative estimation, it follows from (5.12) that

$$\begin{split} \sup_{t\geq 0} \int_{\mathbb{T}\times\mathbb{R}} |n|^{2^{j+1}} dx dy \\ \leq \tilde{C} \max\left[1, 2^{8j} \left(\sup_{t\geq 0} \int_{\mathbb{T}\times\mathbb{R}} |n|^{2^{j}} dx dy \right)^{2 \left(\frac{2^{j}-1}{2^{j}-2}\right)} \right] \\ \leq \tilde{C} \max\left\{ 1, 2^{8j} \left[\tilde{C} \max\left(1, 2^{8(j-1)} \left(\sup_{t\geq 0} \int_{\mathbb{T}\times\mathbb{R}} |n|^{2^{j-1}} dx dy \right)^{2 \left(\frac{2^{j-1}-1}{2^{j-1}-2}\right)} \right) \right]^{2 \left(\frac{2^{j}-1}{2^{j-2}}\right)} \right\} \\ \leq \tilde{C} + \tilde{C}^{1+\frac{2^{j}-1}{2^{j-1}-1}} 2^{8j} \\ + \dots + \tilde{C}^{1+\sum_{k=1}^{j-2}k} \prod_{l=1}^{k} \frac{2^{j-l+1}-1}{2^{j-l+1}-2} 2^{8j+8\sum_{k=1}^{j-3}2^{k}(j-k)} \prod_{l=1}^{k} \frac{2^{j-l+1}-1}{2^{j-l+1}-2}} p_{4}^{4\sum_{k=1}^{j-1}2^{k}} \prod_{l=1}^{k} \frac{2^{j-l+1}-1}{2^{j-l+1}-2} \\ + \tilde{C}^{1+\sum_{k=1}^{j-2}2^{k}} \prod_{l=1}^{k} \frac{2^{j-l+1}-1}{2^{j-l+1}-2} 2^{8j+8\sum_{k=1}^{j-2}2^{k}(j-k)} \prod_{l=1}^{k} \frac{2^{j-l+1}-1}{2^{j-l+1}-2}} p_{4}^{4\sum_{k=1}^{j-1}2^{k}} \prod_{l=1}^{k} \frac{2^{j-l+1}-1}{2^{j-l+1}-2} \\ \leq \tilde{C}^{1+\sum_{k=1}^{j-2}2^{k}} \prod_{l=1}^{k} \frac{2^{j-l+1}-1}{2^{j-l+1}-2}} 2^{8j+8\sum_{k=1}^{j-2}2^{k}(j-k)} \prod_{l=1}^{k} \frac{2^{j-l+1}-1}{2^{j-l+1}-2}} p_{4}^{4\sum_{k=1}^{j-1}2^{k}} \prod_{l=1}^{k} \frac{2^{j-l+1}-1}{2^{j-l+1}-2}} \\ + (j-1)\tilde{C}^{1+\sum_{k=1}^{j-2}2^{k}} \prod_{l=1}^{k} \frac{2^{j-l+1}-1}{2^{j-l+1}-2}} 2^{8j+8\sum_{k=1}^{j-3}2^{k}(j-k)} \prod_{l=1}^{k} \frac{2^{j-l+1}-1}{2^{j-l+1}-2}}. \end{split}$$
(5.13)

Due to

$$\sum_{k=1}^{j-1} 2^k \prod_{l=1}^k \frac{2^{j-l+1}-1}{2^{j-l+1}-2}$$

= $\sum_{k=1}^{j-1} 2^k \left(\frac{2^j-1}{2^j-2}\right) \times \left(\frac{2^{j-1}-1}{2^{j-1}-2}\right) \times \dots \times \left(\frac{2^{j-k+2}-1}{2^{j-k+2}-2}\right) \times \left(\frac{2^{j-k+1}-1}{2^{j-k+1}-2}\right)$
= $\sum_{k=1}^{j-1} 2^k \frac{2^j-1}{2^k (2^{j-k}-1)} = \sum_{k=1}^{j-1} \frac{2^j-1}{2^{j-k}-1},$

and

$$8j + 8\sum_{k=1}^{j-2} 2^k (j-k) \prod_{l=1}^k \frac{2^{j-l+1}-1}{2^{j-l+1}-2} = 8j + 8\sum_{k=1}^{j-2} (j-k) \frac{2^{j+1}-2}{2^{j+1-k}-2},$$

there hold

$$0 \le \lim_{j \to \infty} \frac{1}{2^{j+1}} \left(\sum_{k=1}^{j-1} 2^k \prod_{l=1}^k \frac{2^{j-l+1} - 1}{2^{j-l+1} - 2} \right) = \lim_{j \to \infty} \left[\frac{1}{2} \left(1 - \frac{1}{2^j} \right) \sum_{k=1}^{j-1} \frac{1}{2^{j-k} - 1} \right]$$
$$\le \frac{1}{2} \sum_{n=0}^{\infty} \frac{1}{2^n} = 1,$$

and

$$0 \leq \lim_{j \to \infty} \frac{1}{2^{j+1}} \left[8j + 8 \sum_{k=1}^{j-2} 2^k (j-k) \prod_{l=1}^k \frac{2^{j-l+1} - 1}{2^{j-l+1} - 2} \right]$$
$$= 8 \lim_{j \to \infty} \frac{j}{2^{j+1}} + 8 \lim_{j \to \infty} \left(\frac{2^{j+1} - 2}{2^{j+1}} \sum_{k=1}^{j-2} \frac{j-k}{2^{j+1-k} - 2} \right) \leq 4 \sum_{n=0}^{\infty} \frac{n+1}{2^n} = 12,$$

which along with (5.13) implies that

$$\|n\|_{L^{\infty}L^{\infty}} \le \hat{C} := K_{\infty},$$

where \hat{C} is a positive constant depending on $||n_{\text{in}}||_{L^1 \cap L^\infty}$.

To sum up, we complete the proof.

6. Proof of Theorem 1.2

Proof of Theorem 1.2. Recall (3.12) and (3.13) in the proof of **Lemma 3.6**, where we use the interpolation inequality

$$\|n_0\|_{L^4} \le C_* \|n_0\|_{L^1}^{\frac{1}{2}} \|\partial_y n_0\|_{L^2}^{\frac{1}{2}},$$

and $C_*^4 M^2 < 3$. From **Remark 1.1**, notice that $C_* = \left(\frac{4\pi^2}{9}\right)^{-\frac{1}{4}}$, which follows that $M < \frac{2\pi}{\sqrt{3}}$. Then combining it with **Theorem 1.1**, the proof is complete.

Acknowledgement

The authors would like to thank Professor Wei Wang and Dr. Shikun Cui for some helpful communications. W. Wang was supported by National Key R&D Program of China (No. 2023YFA1009200) and NSFC under grant 12071054 and 12471219.

DECLARATION OF COMPETING INTEREST

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

DATA AVAILABILITY

No data was used in this paper.

References

- [1] Bedrossian J. and He S. (2017). Suppression of blow-up in Patlak–Keller–Segel via shear flows. SIAM Journal on Mathematical Analysis, 49(6), 4722-4766.
- [2] Calvez V. and Corrias L. (2008). The parabolic-parabolic Keller-Segel model in R². Communications in Mathematical Sciences, 6(2), 417-447.
- [3] Carlen A. and Loss M. (1993). Sharp constant in Nash's inequality. International Mathematics Research Notices, 7, 213-215.
- [4] Cieślak, Tomasz; Stinner, Christian; New critical exponents in a fully parabolic quasilinear Keller-Segel system and applications to volume filling models. (English summary) J. Differential Equations 258 (2015), no. 6, 2080-2113.
- [5] Cui S. and Wang W. (2023). Suppression of blow-up in multi-species Patlak-Keller-Segel-Navier-Stokes system via the Poiseuille flow in a finite channel. arXiv:2311.18519.
- [6] Cui S., Wang L. and Wang W. (2024). Suppression of blow-up in Patlak-Keller-Segel system coupled with linearized Navier-Stokes equations via the 3D Couette flow. arXiv:2401.15982.
- [7] Deng S., Shi B. and Wang W. (2024). Suppression of blow-up in 3-D Keller-Segel model via Couette flow in whole space. arXiv:2311.18590.
- [8] He S. (2018). Suppression of blow-up in parabolic-parabolic Patlak-Keller-Segel via strictly monotone shear flows. Nonlinearity, 31(8), 3651.
- [9] Horstmanna D., Winkler M. (2005). Boundedness vs. blow-up in a chemotaxis system. J. Differential Equations, 215, 52-107.
- [10] Hu Z. (2023). Suppression of Chemotactic Singularity via Viscous Flow with Large Buoyancy. arXiv:2311.10003.

- [11] Hu Z., Kiselev A. and Yao Y. (2023) Suppression of chemotactic singularity by buoyancy. arXiv:2305.01036.
- [12] Keller E.F. and Segel L.A. (1970). Initiation of slime mold aggregation viewed as an instability. Journal of Theoretical Biology, 26(3), 399.
- [13] Kiselev A. and Xu X. (2016). Suppression of chemotactic explosion by mixing. Archive for Rational Mechanics and Analysis, 222, 1077-1112.
- [14] Li H., Xiang Z. and Xu X. (2023). Suppression of blow-up in Patlak-Keller-Segel-Navier-Stokes system via the Poiseuille flow, arXiv:2312.01069.
- [15] Liu J. and Wang J. (2017). One the best constant for Gagliardo-Nirenberg interpolation inequalities, arXiv:1712.10208v1.
- [16] Nagai T. (1995). Blow-up of radially symmetric solutions to a chemotaxis system. Adv. Math. Sci. Appl. 5, no. 2, 581–601.
- [17] Nagai T. (2000). Behavior of solutions to a parabolic-elliptic system modelling chemotaxis. J. Korean Math. Soc. 37, 721-732.
- [18] Nagy B. V. Sz. (1941). Ucber integralungleichungen zwischen einer funktion aun ihrer ableitung. Acta Univ. Szeged. Sect. Sci. Math. 10, 64-74.
- [19] Painter K. and Hillen T. (2002). Volume-filling and quorum-sensing in models for chemosensitive movement. Canadian Applied Mathematics Quarterly. 10 (4), 501-543.
- [20] Patlak C.S. (1953). Random walk with persistence and external bias. The bulletin of mathematical biophysics, 15, 311-338.
- [21] Schweyer R. (2014). Stable blow-up dynamic for the parabolic-parabolic Patlak-Keller-Segel model. arXiv:1403.4975.
- [22] Shi B. and Wang W. (2024). Enhanced dissipation and blow-up suppression for the three dimensional Keller-Segel equation with the plane Couette-Poiseuille flow. J. Differential Equations, 403, 368-405.
- [23] Souplet P. and Winkler M. (2019). Blow-up profiles for the parabolic-elliptic Keller-Segel system in dimensions $n \geq 3$. Commun. Math. Phys. 367, 665–681.
- [24] Stinner C. and Winkler M. (2024). A critical exponent in a quasilinear Keller–Segel system with arbitrarily fast decaying diffusivities accounting for volume-filling effects. J. Evol. Equ. 24-26.
- [25] Wang W. and Liu Z. (2024). Globally bounded solutions in a 2D chemotaxis-Navier-Stokes system with general sensitivity and nonlinear production. Z. Angew. Math. Phy. 75:74.
- [26] Wei D. (2018). Global well-posedness and blow-up for the 2-D Patlak-Keller-Segel equation. Journal of Functional Analysis, 274(2), 388-401.
- [27] Winkler M. (2009). Does a 'volume-filling effect' always prevent chemotactic collapse? Math. Methods Appl. Sci. 33, 12-24.
- [28] Winkler M. (2013). Finite-time blow-up in the higher-dimensional parabolicparabolic Keller-Segel system. Journal de Mathematiques Pures et Appliquees, 100(5), 748-767.

- [29] Winkler M. and Djie K C. (2010). Boundedness and finite-time collapse in a chemotaxis system with volume-filling effect. Nonlinear Anal. 72, 1044-1064.
- [30] Zeng L., Zhang Z. and Zi R. (2021). Suppression of blow-up in Patlak-Keller-Segel-Navier-Stokes system via the couette flow. Journal of Functional Analysis, 280(10), 108967.

(Lili Wang) School of Mathematical Sciences, Dalian University of Technology, Dalian, 116024, China

Email address: wanglili_@mail.dlut.edu.cn

(Wendong Wang) SCHOOL OF MATHEMATICAL SCIENCES, DALIAN UNIVERSITY OF TECHNOL-OGY, DALIAN, 116024, CHINA

 $Email \ address: \verb|wendong@dlut.edu.cn||$

(Yi Zhang) School of Mathematical Sciences, Dalian University of Technology, Dalian, 116024, China

Email address: zysx@mail.dlut.edu.cn