# Towards new relativistic doubly $\kappa$ -deformed D=4 quantum phase spaces

Jerzy Lukierski<sup>a</sup>, Stjepan Meljanac<sup>b</sup>, Salvatore Mignemi<sup>c</sup>, Anna Pachoł<sup>d</sup> and Mariusz Woronowicz<sup>a</sup> <sup>a</sup> Institute of Theoretical Physics, Wrocław University,

pl. Maxa Borna 9, 50-205 Wrocław, Poland

<sup>b</sup> Division of Theoretical Physics, Rudjer Bošković Institute,

Bijenička c.54, HR-10002 Zagreb, Croatia

<sup>c</sup> Dipartimento di Matematica, Università di Cagliari via Ospedale 72,

09124 Cagliari, Italy and INFN, Sezione di Cagliari, 09042 Monserrato, Italy

<sup>d</sup> Department of Microsystems, University of South-Eastern Norway, Campus Vestfold, Norway

We propose new noncommutative (NC) models of quantum phase spaces, containing a pair of  $\kappa$ -deformed Poincaré algebras, with two independent  $\kappa$  and  $\tilde{\kappa}$ -deformations in space-time and fourmomenta sectors. The first quantum phase space can be obtained by contractions  $M, R \to \infty$  of recently introduced doubly  $\kappa$ -deformed  $(\kappa, \tilde{\kappa})$ -Yang models, with the parameters M, R describing inverse space-time and fourmomenta curvatures and constant four-vectors  $a_{\mu}, b_{\mu}$  determining one of nine types of  $(\kappa, \tilde{\kappa})$  deformations. The second model is provided by the non-linear doubly  $\kappa$ -deformed TSR algebra spanned by 14 coset  $\hat{o}(1,5)/\hat{o}(2)$  generators. The basic algebraic difference between the two models is the following: the first one, described by  $\hat{o}(1,5)$  Lie algebra can be supplemented by the Hopf algebra structure, while the second model contains the quantum phase space commutators  $[\hat{x}_{\mu}, \hat{q}_{\nu}]$  describing the quantum-deformed Heisenberg algebra relations with the standard numerical  $i\hbar\eta_{\mu\nu}$  term.

#### INTRODUCTION I.

The canonical D = 4 relativistic phase space algebra is generated by commuting space-time  $x_{\mu}$ and fourmomenta  $q_{\mu}$  coordinates and can be extended by the Lorentz symmetry generators  $\hat{M}_{\mu\nu}$ . We denote such algebra by  $H^{(1,3)} = (x_{\mu}, q_{\mu}, \hat{M}_{\mu\nu})$ , with space-time coordinates and fourmomenta satisfying the Heisenberg canonical commutation relations:

$$[x_{\mu}, q_{\nu}] = i\hbar\eta_{\mu\nu} \tag{1}$$

where  $\mu, \nu = 0, 1, 2, 3, \eta_{\mu\nu} = \text{diag}(-1, 1, 1, 1)$  and Lorentz algebra

$$[\hat{M}_{\mu\nu}, \hat{M}_{\rho\tau}] = i\hbar(\eta_{\mu\rho}\hat{M}_{\nu\tau} - \eta_{\mu\tau}\hat{M}_{\nu\rho} + \eta_{\nu\tau}\hat{M}_{\mu\rho} - \eta_{\nu\rho}\hat{M}_{\mu\tau})$$
(2)

defining the relativistic covariance relations:

$$[\hat{M}_{\mu\nu}, x_{\rho}] = i\hbar(\eta_{\mu\rho}x_{\nu} - \eta_{\nu\rho}x_{\mu}), \qquad (3)$$

$$[\hat{M}_{\mu\nu}, q_{\rho}] = i\hbar(\eta_{\mu\rho}q_{\nu} - \eta_{\nu\rho}q_{\mu}).$$
(4)

From the set of the generators of algebra  $H^{(1,3)}$  one can construct the pair of overlapping classical Poincaré algebras: one generated by  $(x_{\mu}, \hat{M}_{\mu\nu})$  and the second one by  $(q_{\mu}, \hat{M}_{\mu\nu})$ . The relations (1)-(4) are covariant under the following Born map B [1–3]

$$B: \quad x_{\mu} \to q_{\mu}, \quad q_{\mu} \to -x_{\mu}, \quad \hat{M}_{\mu\nu} \leftrightarrow \hat{M}_{\mu\nu}. \tag{5}$$

and it follows from relations (1)-(4) that the algebra  $H^{(1,3)}$  is Born-selfdual.

Our first aim will be to describe the deformations of the algebra  $H^{(1,3)}$  which lead to the noncanonical algebra  $\hat{H}^{(1,3)}$  with noncommutative (NC) quantum phase space coordinates  $(\hat{x}_{\mu}, \hat{q}_{\mu})$ . Due to the presence of quantum gravity (QG) effects (see e.g. [4]) we replace  $x_{\mu} \to \hat{x}_{\mu}$ ,  $q_{\mu} \to \hat{q}_{\mu}$ , and after the quantization we get from  $H^{(1,3)}$  the quantum relativistic algebra  $H^{(1,3)} \to \hat{H}^{(1,3)} =$  $(\hat{x}_{\mu}, \hat{q}_{\mu}, \hat{M}_{\mu\nu})$ . Then in general  $[\hat{x}_{\mu}, \hat{x}_{\nu}] \neq 0$  and  $[\hat{q}_{\mu}, \hat{q}_{\mu}] \neq 0$ , while the commutator  $[\hat{x}_{\mu}, \hat{q}_{\nu}]$  becomes significantly different from the relation (1). We consider two well-known NC models which provide the particular choices of the algebras  $\hat{H}^{(1,3)}$ . The first example of such algebra has been proposed by Yang [5] as an extension of the Snyder model [6]. The defining relations of D = 4 Yang model are as follows:

$$[\hat{x}_{\mu}, \hat{x}_{\nu}] = \frac{i\hbar}{M^2} \hat{M}_{\mu\nu}, \tag{6}$$

$$[\hat{q}_{\mu}, \hat{q}_{\nu}] = \frac{i\hbar}{R^2} \hat{M}_{\mu\nu},\tag{7}$$

$$[\hat{x}_{\mu}, \hat{q}_{\nu}] = i\hbar\eta_{\mu\nu}\hat{r} \tag{8}$$

where the (real) parameters M and R describe the curvatures of NC space-time and NC fourmomenta space<sup>1</sup>. The additional generator  $\hat{r}$  satisfies the relations:

$$[\hat{r}, \hat{x}_{\mu}] = \frac{i\hbar}{M^2} \hat{q}_{\mu}, \qquad [\hat{r}, \hat{q}_{\mu}] = -\frac{i\hbar}{R^2} \hat{x}_{\mu}$$

$$\tag{9}$$

and it follows that  $\hat{r}$  describes the generator of particular  $\hat{o}(2)$  internal symmetries <sup>2</sup>.

If we supplement the Born map (5) with the following additional mappings:

$$B: \qquad M \leftrightarrow R, \qquad \hat{r} \leftrightarrow \hat{r} \tag{10}$$

it is easy to see that the Yang model (6)-(9) is Born self-dual. After the following assignment of the generators:

$$\hat{M}_{\mu4} = M\hat{x}_{\mu}, \ \hat{M}_{\mu5} = R\hat{q}_{\mu}, \ \hat{M}_{45} = MR\hat{r}$$
(11)

the relations (6)-(9) and (2) permit to interpret the D = 4 Yang algebra as algebraically equivalent to  $\hat{o}(1,5)$  Lie algebra:

$$[\hat{M}_{AB}, \hat{M}_{CD}] = i\hbar(\eta_{AC}\hat{M}_{BD} - \eta_{AD}\hat{M}_{BC} + \eta_{BD}\hat{M}_{AC} - \eta_{BC}\hat{M}_{AD})$$
(12)

where  $\eta_{AB} = diag(-1, 1, ..., 1)$  and A, B = 0, 1, ..., 5.

Another example of NC models which will be considered here is provided by the NC geometry of  $\kappa$ -deformed Minkowski space-time accompanied by Born-dual  $\tilde{\kappa}$ -deformed fourmomenta space.

<sup>&</sup>lt;sup>1</sup> For the notions and the use of NC Riemannian geometry see e.g. [7–10] in quantum gravity models and for example string theory [11]

<sup>&</sup>lt;sup>2</sup> The standard  $\hat{o}(2)$  relations with  $\hat{o}(2)$  generator  $\hat{I}$  are  $[I, \hat{X}_{\mu}] = i\hbar\hat{Q}_{\mu}$ ,  $[I, \hat{Q}_{\mu}] = -i\hbar\hat{X}_{\mu}$  and can be obtained from (9) by the redefinition:  $\hat{I} = MR\hat{r}$ ,  $\hat{X}_{\mu} = M\hat{x}_{\mu}$ ,  $\hat{Q}_{\mu} = R\hat{q}_{\mu}$ .

In order to describe all three possible types of  $\kappa$ -deformations of NC Minkowski space-time (timelike, standard or light-like) one introduces the constant fourvector  $a_{\mu}$  which selects the quantized direction in space-time [12]-[14]. The  $\kappa$ -Minkowski commutation relations are:

$$[\hat{x}_{\mu}, \hat{x}_{\nu}] = \frac{i\hbar}{\kappa} (\hat{a}_{\mu}\hat{x}_{\nu} - a_{\nu}\hat{x}_{\mu}) \tag{13}$$

where  $a_{\mu}$  can be chosen in threefold way as  $a^2 = a_{\mu}a^{\mu} = (1, 0, -1)^3$ . By introducing the constant fourvector  $b_{\mu}$  describing quantization direction in quantum fourmomentum space  $\hat{q}_{\mu}$  one can introduce the following analogous  $\tilde{\kappa}$ -deformation of NC fourmomenta<sup>4</sup>:

$$[\hat{q}_{\mu}, \hat{q}_{\nu}] = i\hbar\tilde{\kappa}(b_{\mu}\hat{q}_{\nu} - b_{\nu}\hat{q}_{\mu}).$$
(14)

It should be pointed out that recently [15] (see also [16], [17]) the double  $\kappa$ -deformation of the Yang model<sup>5</sup> has been introduced, containing in its subalgebras the  $\kappa$ -deformation (13) of NC coordinates and the  $\tilde{\kappa}$ -deformation (14) of NC fourmomentum space. It was shown [15] how to obtain doubly  $\kappa$ -deformed Yang models from  $\kappa$ -deformed Snyder models [18]-[26] by applying the generalized Born map. In this paper we propose new models of relativistic doubly  $\kappa$ -deformed D=4 quantum phase spaces which will be obtained in two different ways. First way starts from the doubly  $\kappa$ -deformed Yang models ([15], [16], [17]), which will be recalled in Sec. 2. We consider the following two alternative derivations of new models:

i) Firstly, we perform the contraction procedure  $(M, R) \to \infty$  in the doubly  $\kappa$ -deformed Yang model and obtain new relativistic doubly  $\kappa$ -deformed D=4 quantum phase space, depending on two parameters  $\kappa$  and  $\tilde{\kappa}$ .

ii) Secondly, we use the Lie algebraic description of doubly  $\kappa$ -deformed Yang model as represented by  $\hat{o}(1,5;g)$  Lie algebra which can be obtained from  $\hat{o}(1,5) \equiv \hat{o}(1,5,\eta)$  by replacing diagonal pseudoorthogonal metric  $\eta_{AB}$  by suitably chosen constant parameter-dependent symmetric metric  $g \equiv g_{AB}^{6}$ . We properly fix the parameter-dependent metric g to obtain the new model of relativistic doubly  $\kappa$ -deformed D=4 quantum phase space, depending on three independent parameters  $\kappa, \tilde{\kappa}$ and  $\rho$ .

In both cases i) and ii) the obtained doubly  $\kappa$ -deformed quantum phase spaces are algebraically equivalent to  $\hat{o}(1,5)$  Lie algebra. i.e. to the original Yang model.

Another possible algebra with NC coordinates and NC momenta is provided by the Triply Special Relativity (TSR) model [27]. It is spanned by 14 generators of the coset  $\hat{o}(1,5)/\hat{o}(2)$  and depends on three independent fundamental parameters. Often, TSR model is also called Snyder-de Sitter (SdS) model since it is obtained as a generalization of the Snyder model (flat space-time with curved momentum space) to the model with curved both NC space-time and NC fourmomenta. Additionally, in TSR model, one considers appropriate deformations of the Heisenberg phase space commutator  $[\hat{x}_{\mu}, \hat{p}_{\nu}]$ , which in the quantum-mechanical limit, without QG corrections, reduces to the canonical Heisenberg commutation relations.

The main novelty of the present paper is in Sec. 3, where we propose the modification of TSR model by introducing doubly  $\kappa$ -deformation in the quantum phase space. In such a case the  $(\kappa, \tilde{\kappa})$ -deformed quantum phase space reduces to the canonical Heisenberg relations in the contraction limit  $\kappa \to \infty$  and  $\tilde{\kappa} \to 0$ .

<sup>&</sup>lt;sup>3</sup> The relations (13),(14) introduce two mass-like parameters  $\kappa$ ,  $\tilde{\kappa}$  and two constant fourvectors  $a_{\mu}$ ,  $b_{\mu}$  limited by threefold constraints  $a^2 = (1, 0, -1)$  and  $b^2 = (1, 0, -1)$  which specify  $3 \times 3 = 9$  types of  $(\kappa, \tilde{\kappa})$ -deformed pairs of Poincaré algebras. In general cases, as we will show in Sec. 2, still there should appear a third parameter  $\rho$ .

 $<sup>^4</sup>$  In general case the mass-like parameters  $\kappa$  and  $\tilde{\kappa}$  are independent.

<sup>&</sup>lt;sup>5</sup> In shortcut notation it is also called  $(\kappa, \tilde{\kappa})$ -Yang model, see [15].

<sup>&</sup>lt;sup>6</sup> In fact we choose the algebras  $\hat{o}(1,5;g)$  which are isomorphic to the classical Lie algebra  $\hat{o}(1,5)$ 

# II. DOUBLY $\kappa$ -DEFORMED D=4 QUANTUM PHASE SPACES FROM YANG MODELS

Many efforts to generalize Snyder and Yang models appeared in the literature, for example the  $\kappa$ -Minkowski extension of Snyder model was firstly proposed in [18] (see also [19]-[26]) while  $\kappa$ -Minkowski type extensions of Yang model in both coordinates and momenta sectors were proposed recently in [15] (see also [16], [17]). In such extensions of the Yang model we need to introduce a pair of different  $\kappa$ -Minkowski terms, respectively in NC space-time and quantum fourmomenta, i.e. we need two independent mass-like parameters  $\kappa$  and  $\tilde{\kappa}$  resulting in  $(\kappa, \tilde{\kappa})$ -Yang model, which can be defined by the following set of commutation relations (see [15]):

$$[\hat{x}_{\mu}, \hat{x}_{\nu}] = i\hbar \left[ \frac{1}{M^2} \hat{M}_{\mu\nu} + \frac{1}{\kappa} (a_{\mu} \hat{x}_{\nu} - a_{\nu} \hat{x}_{\mu}) \right],$$
(15)

$$[\hat{q}_{\mu}, \hat{q}_{\nu}] = i\hbar \left[ \frac{\hat{M}_{\mu\nu}}{R^2} + \tilde{\kappa} (b_{\mu}\hat{q}_{\nu} - b_{\nu}\hat{q}_{\mu}) \right]$$
(16)

and

$$[\hat{M}_{\mu\nu}, \hat{x}_{\rho}] = i\hbar \left[ \eta_{\mu\rho} \hat{x}_{\nu} - \eta_{\nu\rho} \hat{x}_{\mu} + \frac{1}{\kappa} (a_{\mu} \hat{M}_{\rho\nu} - a_{\nu} \hat{M}_{\rho\mu}) \right],$$
(17)

$$[\hat{M}_{\mu\nu}, \hat{q}_{\rho}] = i\hbar \left[ \eta_{\mu\rho} \hat{q}_{\nu} - \eta_{\nu\rho} \hat{q}_{\mu} + \tilde{\kappa} (b_{\mu} \hat{M}_{\rho\nu} - b_{\nu} \hat{M}_{\rho\mu}) \right].$$
(18)

We see that in the relations (15), (16) the noncommutativity of quantum space-time and fourmomenta is described by the sum of the noncommutativities from the Yang model and from  $\kappa$ deformation. Further, it can be shown that the relations (17), (18) can be justified by the calculation of Jacobi identities. The remaining quantum phase space relations are [15], [16]<sup>7</sup>:

$$[\hat{x}_{\mu}, \hat{q}_{\nu}] = i\hbar \left( \eta_{\mu\nu} \hat{r} + \tilde{\kappa} b_{\mu} \hat{x}_{\nu} - \frac{a_{\nu}}{\kappa} \hat{q}_{\mu} + \frac{\rho}{MR} \hat{M}_{\mu\nu} \right), \tag{19}$$

$$[\hat{r}, \hat{x}_{\mu}] = i\hbar \left(\frac{1}{M^2}\hat{q}_{\mu} - \frac{1}{MR}\rho\hat{x}_{\mu} - \frac{a_{\mu}}{\kappa}\hat{r}\right),\tag{20}$$

$$[\hat{r}, \hat{q}_{\mu}] = i\hbar \left( -\frac{1}{R^2} \hat{x}_{\mu} + \frac{1}{MR} \rho \hat{q}_{\mu} - \tilde{\kappa} b_{\mu} \hat{r} \right).$$
(21)

$$\mu,\nu \rightarrow i,j; \quad \hbar=1; \quad \frac{1}{R}=\alpha \quad \frac{1}{M}=\beta;$$

and the generators

$$\hat{x}_{\nu} \to \tilde{X}_i; \quad \hat{q}_{\mu} \to \tilde{P}_i; \quad \hat{r} \to \hat{h}; \quad a_{\mu} \to \kappa \beta a_{\mu}; \quad b_{\mu} \to (\alpha/\tilde{\kappa})b_{\mu}.$$

<sup>&</sup>lt;sup>7</sup> To link the notations used in [15] and [16] we need the following convention matching, where the left hand side contains the notation used in [15] and the right hand side corresponds to the one from [16]:

Additionally, we have

$$[\hat{r}, \hat{M}_{\mu\nu}] = -i\hbar \left[ \frac{1}{\kappa} (a_{\mu}\hat{q}_{\nu} - a_{\nu}\hat{q}_{\mu}) - \tilde{\kappa}(b_{\mu}\hat{x}_{\nu} - b_{\nu}\hat{x}_{\mu}) \right].$$
(22)

The relation (22) shows that the internal and Lorentzian generators do not commute with each other. The  $\kappa, \tilde{\kappa}$ -dependence of the Yang model can be explained by the generalized Born map  $\tilde{B}$ , obtained if we supplement (5) and (10) by the following relations<sup>8</sup>

$$\tilde{B}: a_{\mu} \to b_{\mu}, \quad b_{\mu} \to -a_{\mu}, \quad \kappa \leftrightarrow \frac{1}{\tilde{\kappa}}.$$
(23)

Both maps (5), (23) describe pseudo-involutions, satisfying the relations  $B^2 = \tilde{B}^4 = 1$ . It can be checked that the relations (19)-(22) are self dual under the generalized Born map (5), extended by (23).

# A. Doubly $\kappa$ -deformed D=4 quantum phase spaces by contraction

To obtain the new model of quantum phase space, including both NC coordinates and NC momenta, and which is dependent on two additional parameters  $\kappa$  and  $\tilde{\kappa}$ , we perform the contraction when  $M \to \infty$  and  $R \to \infty$  in doubly  $(\kappa, \tilde{\kappa})$ -Yang model, described by (15)-(22). We call this new model doubly  $\kappa$ -Poincaré algebra. It is given by relations<sup>9</sup>

$$[\hat{x}_{\mu}, \hat{x}_{\nu}] = \frac{i\hbar}{\kappa} (\hat{a}_{\mu}\hat{x}_{\nu} - a_{\nu}\hat{x}_{\mu}), \qquad [\hat{q}_{\mu}, \hat{q}_{\nu}] = i\hbar\tilde{\kappa} (b_{\mu}\hat{q}_{\nu} - b_{\nu}\hat{q}_{\mu}), \tag{24}$$

$$[\hat{M}_{\mu\nu}, \hat{x}_{\rho}] = i\hbar \left[ \eta_{\mu\rho} \hat{x}_{\nu} - \eta_{\nu\rho} \hat{x}_{\mu} + \frac{1}{\kappa} (a_{\mu} \hat{M}_{\rho\nu} - a_{\nu} \hat{M}_{\rho\mu}) \right],$$
(25)

$$[\hat{M}_{\mu\nu}, \hat{q}_{\rho}] = i\hbar \left[ \eta_{\mu\rho} \hat{q}_{\nu} - \eta_{\nu\rho} \hat{q}_{\mu} + \tilde{\kappa} (b_{\mu} \hat{M}_{\rho\nu} - b_{\nu} \hat{M}_{\rho\mu}) \right], \qquad (26)$$

$$[\hat{x}_{\mu}, \hat{q}_{\nu}] = i\hbar \left( \eta_{\mu\nu} \hat{r} + \tilde{\kappa} b_{\mu} \hat{x}_{\nu} - \frac{a_{\nu}}{\kappa} \hat{q}_{\mu} \right), \qquad (27)$$

$$[\hat{r}, \hat{x}_{\mu}] = -i\hbar \frac{a_{\mu}}{\kappa} \hat{r}, \qquad [\hat{r}, \hat{q}_{\mu}] = -i\hbar \tilde{\kappa} b_{\mu} \hat{r}, \qquad (28)$$

$$[\hat{r}, \hat{M}_{\mu\nu}] = -i\hbar \left[ \frac{1}{\kappa} (a_{\mu}\hat{q}_{\nu} - a_{\nu}\hat{q}_{\mu}) - \tilde{\kappa} (b_{\mu}\hat{x}_{\nu} - b_{\nu}\hat{x}_{\mu}) \right].$$
(29)

<sup>&</sup>lt;sup>8</sup> In standard Born duality (e.g. for Yang algebra (6)-(9)) the algebra is self-dual due to the exchange of generators and constants responsible for agreement of dimensions of the generators (see (5) and (10)). The double  $\kappa$  deformed Yang algebra (15)-(22) is self-dual in the above sense only in the case when the fourvectors a = b = 0. For other values of a and b the double  $\kappa$  deformed Yang algebra is self-dual only if we additionally require the change of the structure constants of algebra (corresponding to  $a \to b, b \to -a$ ) and we name such an extension of Born duality as generalized Born duality.

<sup>&</sup>lt;sup>9</sup> The relations (13), (14) are repeated here for convenience and clarity of presentation of the new model.

#### **B.** Doubly $\kappa$ -deformed D = 4 quantum phase spaces by fixing the metric $g_{AB}$ in $\hat{o}(1, 5, g_{AB})$

It is already known since 1947 (see [5]) that the algebra describing D = 4 Yang model, i.e. (6)-(9) is spanned by the  $\hat{o}(1,5)$  algebra. The generalizations of this description, by replacing  $\eta_{AB}$  by more general metric  $g_{AB}$  ( $\eta_{AB} \rightarrow g_{AB}$ ) have been proposed, e.g. in [29]. Doubly ( $\kappa, \tilde{\kappa}$ )-Yang models, which we recalled in Sec. 2 and which are described by eqs (15)-(22), have the Lie algebra structure of  $\hat{o}(1,5;g_{AB}^{(Y)})$  (see [15]):

$$[\hat{M}_{AB}, \hat{M}_{CD}] = i\hbar (g_{AC}^{(Y)} \hat{M}_{BD} - g_{AD}^{(Y)} \hat{M}_{BC} + g_{BD}^{(Y)} \hat{M}_{AC} - g_{BC}^{(Y)} \hat{M}_{AD})$$
(30)

where the symmetric metric components  $g_{AB}^{(Y)}$  with the signature (-1, 1, ..., 1) depend on five parameters <sup>10</sup>  $(M, R, \kappa, \tilde{\kappa}, \rho)$  where  $(M = \lambda^{-1}, \tilde{M} = R^{-1})$  is the pair of mass parameters (or equivalently the pair of length parameters  $\lambda = M^{-1}, R$ ), the mass-like parameters  $(\kappa, \tilde{\kappa})$  and the dimensionless parameter  $\rho$  and a pair of constant dimensionless four-vectors  $a_{\mu}, b_{\mu}, (\mu = 0, 1, 2, 3)$ which respectively determine the type of  $\kappa$ -dependence in D = 4 quantum space-time and D = 4 quantum fourmomenta sectors of Yang algebra. The metric  $g_{AB}^{(Y)}$  is determined by the following assignments of the generators:

$$\hat{M}_{AB} = \left(\hat{M}_{\mu\nu}, \ \hat{M}_{\mu4} = M\hat{x}_{\mu}, \ \hat{M}_{\mu5} = R\hat{q}_{\mu}, \ \hat{M}_{45} = MR\hat{r}\right)$$
(31)

where  $[M_{AB}] = L^0$  (dimensionless), in consistency with relation (30), with  $\hat{M}_{\mu\nu}$  describing D = 4Lorentz algebra and the scalar  $\hat{r}$  providing the generator of the  $\hat{o}(2)$  internal symmetries. Relations (30) describe the  $(\kappa, \tilde{\kappa})$ -Yang model if we insert the following components of the D = 6 metric tensor:

$$g_{AB}^{(Y)} = \begin{pmatrix} \eta_{\mu\nu} & g_{\mu4}^{(Y)} & g_{\mu5}^{(Y)} \\ g_{4\nu}^{(Y)} & g_{44}^{(Y)} & g_{45}^{(Y)} \\ g_{5\nu}^{(Y)} & g_{54}^{(Y)} & g_{55}^{(Y)} \end{pmatrix}$$
(32)

where<sup>11</sup>

$$g_{\mu4}^{(Y)} = g_{4\mu}^{(Y)} = \frac{M}{\kappa} a_{\mu}, \quad g_{\mu5}^{(Y)} = g_{5\mu}^{(Y)} = R\tilde{\kappa}b_{\mu}, \quad g_{45}^{(Y)} = g_{54}^{(Y)} = \rho, \quad g_{44}^{(Y)} = g_{55}^{(Y)} = 1.$$
(33)

Note  $g_{AB}^{(Y)}$  are dimensionless  $([g_{AB}^{(Y)}] = L^0)$  in consistency with relations (30). To obtain the new model of doubly  $\kappa$ -deformed quantum phase space (doubly  $\kappa$ -Poincaré algebra), depending on three parameters  $\kappa$ ,  $\tilde{\kappa}$  and  $\rho$  we modify the metric  $g_{AB}^{(Y)}$  by setting  $g_{44} = 0 = g_{55} = 0$ . We note that this new quantum phase space will preserve Lorentz covariance and we consider the same assignment of generators (31) but only change the metric  $g_{AB}^{(Y)}$  to the following

$$\tilde{g}_{AB} = \begin{pmatrix} \eta_{\mu\nu} & g_{\mu4} = \frac{M}{\kappa} a_{\mu} & g_{\mu5} = R\tilde{\kappa} b_{\mu} \\ g_{4\nu} = \frac{M}{\kappa} a_{\nu}^T & 0 & \rho \\ g_{5\nu} = R\tilde{\kappa} b_{\nu}^T & \rho & 0 \end{pmatrix}$$
(34)

<sup>&</sup>lt;sup>10</sup> Note that in Sec. 2B the parameters M, R are kept finite.

<sup>&</sup>lt;sup>11</sup> We note that the metric  $g_{AB}$  with fixed  $\eta_{\mu\nu}$  can in the most general case be built up from 11 free parameters with arbitrary values of  $g_{44}$  and  $g_{55}$  (see also [16]). However, following our choice in [15] we are using the version with only 9 free parameters, fixing 2 parameters by the relations  $g_{44} = 1$  and  $g_{55} = 1$ .

$$\det \tilde{g}_{AB} = \rho^2 - 2\rho M R \frac{\tilde{\kappa}}{\kappa} \left( a_\mu b^\mu \right) + \left( M R \frac{\tilde{\kappa}}{\kappa} \right)^2 \left[ \left( a_\mu b^\mu \right)^2 - a^2 b^2 \right] \neq 0.$$
(35)

Now the relations of  $\hat{o}(1, 5, \tilde{g}_{AB})$ :

$$[\hat{M}_{AB}, \hat{M}_{CD}] = i\hbar(\tilde{g}_{AC}\hat{M}_{BD} - \tilde{g}_{AD}\hat{M}_{BC} + \tilde{g}_{BD}\hat{M}_{AC} - \tilde{g}_{BC}\hat{M}_{AD})$$
(36)

describe doubly  $\kappa$ -deformed quantum phase space algebra given by (24), (25), (26) and (2) with the exception of the commutation relation between NC coordinates and NC momenta which now becomes:

$$[\hat{x}_{\mu}, \hat{q}_{\nu}] = i\hbar(\eta_{\mu\nu}\hat{r} + \tilde{\kappa}b_{\mu}\hat{x}_{\mu} - \frac{a_{\nu}}{\kappa}\hat{q}_{\mu} + \frac{\rho}{MR}\hat{M}_{\mu\nu})$$
(37)

Note that (37) provides a more general version of the doubly  $\kappa$ -deformed phase space than the one obtained in the previous section, cf. (27). Because of the last term, proportional to the parameter  $\rho$ , the two models of relativistic D = 4 doubly  $\kappa$ -deformed quantum phase spaces, called doubly  $\kappa$ -Poincaré algebras (obtained in Sec. 2A and 2B) are different.

We can obtain from the relation (37) the version from Sec. 2A by setting  $\rho = 0$  in (34), while keeping det  $\tilde{g}_{AB} \neq 0$ . These conditions put certain restrictions on the possible choices of the constant fourvectors  $a_{\mu}$  and  $b_{\mu}$ , namely  $(a_{\mu}b^{\mu})^2 \neq a^2b^2$ . One of the allowed cases would be  $a_{\mu}b^{\mu} = 0$ ,  $a^2 \neq 0$ ,  $b^2 \neq 0$  (see (35)). In particular these restrictions imply that  $b^{\mu}$  cannot be proportional to  $a^{\mu}$ .

#### C. Doubly $\kappa$ -deformation directly from Yang model

In Sections 2.A and 2.B we performed the reductions of doubly  $(\kappa, \tilde{\kappa})$ -Yang models [15] (15)-(22) to obtain double- $\kappa$  deformation. In this Section we present a more straightforward possibility, by starting from the Yang algebra (2), (6)-(9) and introducing the change of basis resulting in the double- $\kappa$  deformation. To obtain the full set of Yang algebra commutators we add to (2), (6)-(9) the relation  $[\hat{r}, \hat{M}_{\mu\nu}] = 0$  what implies that  $\hat{r}$  describes an Abelian internal  $\hat{o}(2)$  symmetry generator. Then, we can change the basis of (2), (6)-(9) by the following transformation (see e.g.[30])

$$\tilde{x}_{\mu} = \hat{x}_{\mu} + \frac{1}{\kappa} a^{\rho} M_{\mu\rho}, \qquad \tilde{q}_{\mu} = \hat{q}_{\mu} + \tilde{\kappa} b^{\rho} M_{\mu\rho}, \qquad \tilde{M}_{\mu\nu} = \hat{M}_{\mu\nu}$$
(38)

where  $\kappa$ ,  $\tilde{\kappa}$  are real parameters and  $a_{\mu}$ ,  $b_{\mu}$  real fourvectors as before. At this stage we do not specify the dependence of the  $\hat{r}$  generator on the remaining Yang algebra generators  $\tilde{x}_{\mu}, \tilde{q}_{\mu}, \tilde{M}_{\mu\nu}$  generators. The algebra of the transformed generators (38) looks as follows:

$$[\tilde{x}_{\mu}, \tilde{x}_{\nu}] = i\hbar(\frac{1}{M^2} + \frac{a^2}{\kappa^2})\tilde{M}_{\mu\nu} + \frac{i\hbar}{\kappa}(a_{\mu}\tilde{x}_{\nu} - a_{\nu}\tilde{x}_{\mu}),$$
(39)

and

$$[\tilde{p}_{\mu}, \tilde{p}_{\nu}] = i\hbar(\frac{1}{R^2} + \tilde{\kappa}^2 b^2)\tilde{M}_{\mu\nu} + i\hbar\tilde{\kappa}(b_{\mu}\tilde{p}_{\nu} - b_{\nu}\tilde{p}_{\mu}), \qquad (40)$$

where  $a^2 = a^{\mu}a_{\mu}$  and  $b^2 = b^{\mu}b_{\mu}$  and we additionally have:

$$[\tilde{x}_{\mu}, \tilde{q}_{\nu}] = i\hbar\eta_{\mu\nu}(\hat{r} + \frac{1}{\kappa}a^{\rho}\tilde{q}_{\rho} - \tilde{\kappa}b^{\rho}\tilde{x}_{\rho} - \frac{\tilde{\kappa}}{\kappa}a^{\rho}b^{\sigma}\tilde{M}_{\rho\sigma}) + i\hbar\left[\tilde{\kappa}b_{\mu}\tilde{x}_{\nu} - \frac{1}{\kappa}a_{\nu}\tilde{q}_{\mu} + \frac{\tilde{\kappa}}{\kappa}a^{\rho}b_{\rho}\tilde{M}_{\mu\nu}\right], \quad (41)$$

$$[\tilde{M}_{\mu\nu}, \tilde{x}_{\rho}] = i\hbar(\eta_{\mu\rho}\tilde{x}_{\nu} - \eta_{\nu\rho}\tilde{x}_{\mu}) + \frac{i\hbar}{\kappa}(a_{\nu}\tilde{M}_{\mu\rho} - a_{\mu}\tilde{M}_{\nu\rho}), \qquad (42)$$

$$[\tilde{M}_{\mu\nu}, \tilde{p}_{\rho}] = i\hbar(\eta_{\mu\rho}\tilde{p}_{\nu} - \eta_{\nu\rho}\tilde{p}_{\mu}) + i\hbar\tilde{\kappa}(b_{\nu}\tilde{M}_{\mu\rho} - b_{\mu}\tilde{M}_{\nu\rho}).$$
(43)

To write the remaining commutation relations we should specify the transformation of the generator  $\hat{r}$ . Let us consider two cases:

1. If we put simply  $\tilde{r} = \hat{r}$ , this leads to the following modified commutators of Yang algebra

$$[\tilde{r}, \tilde{x}_{\mu}] = \frac{i\hbar}{M^2} (\tilde{q}_{\mu} - \tilde{\kappa} b^{\rho} \tilde{M}_{\mu\rho}), \qquad [\tilde{r}, \tilde{q}_{\mu}] = -\frac{i\hbar}{R^2} (\tilde{x}_{\mu} - \frac{1}{\kappa} a^{\rho} \tilde{M}_{\mu\rho}), \qquad [\tilde{r}, \tilde{M}_{\mu\nu}] = 0.$$
(44)

In such a case the internal symmetry and Lorentzian generators are still commuting with each other.

2. We can introduce the following formula for the modified generator  $\hat{r}$  (compare with the first term on the RHS of (41))

$$\tilde{r} = \hat{r} + \frac{1}{\kappa} a^{\rho} \tilde{q}_{\rho} - \tilde{\kappa} b^{\rho} \tilde{x}_{\rho} - \frac{\tilde{\kappa}}{\kappa} a^{\rho} b^{\sigma} \tilde{M}_{\rho\sigma}.$$
(45)

In this case the commutator (41) can be rewritten as follows:

$$[\tilde{x}_{\mu}, \tilde{q}_{\nu}] = i\hbar \left[ \eta_{\mu\nu} \tilde{r} + \tilde{\kappa} b_{\mu} \tilde{x}_{\nu} - \frac{1}{\kappa} a_{\nu} \tilde{q}_{\mu} + \frac{\tilde{\kappa}}{\kappa} a^{\rho} b_{\rho} \tilde{M}_{\mu\nu} \right].$$
(46)

We can also calculate all the remaining commutators containing the  $\tilde{r}$  generator

$$[\tilde{r}, \tilde{x}_{\mu}] = i\hbar \left[ -\frac{1}{\kappa} a_{\mu} \tilde{r} - \frac{\tilde{\kappa}}{\kappa} a^{\rho} b_{\rho} \tilde{x}_{\mu} + (\frac{1}{M^2} + \frac{a^2}{\kappa^2}) \tilde{q}_{\mu} \right],$$
(47)

$$[\tilde{r}, \tilde{q}_{\mu}] = i\hbar \left[ \tilde{\kappa} b_{\mu} \tilde{r} + \frac{\tilde{\kappa}}{\kappa} a^{\rho} b_{\rho} \tilde{q}_{\mu} - \left(\frac{1}{R^2} + \tilde{\kappa}^2 b^2\right) \tilde{x}_{\mu} \right], \tag{48}$$

$$[\tilde{M}_{\mu\nu}, \tilde{r}] = i\hbar \left[ \frac{1}{\kappa} (a_{\mu}\tilde{q}_{\nu} - a_{\nu}\tilde{q}_{\mu}) - \tilde{\kappa} (b_{\mu}\tilde{x}_{\nu} - b_{\nu}\tilde{x}_{\mu}) \right]$$
(49)

and we see from (49) that the internal symmetry generator  $\hat{r}$  and Lorentzian generators do not commute with each other in the new basis.

Further, to obtain the double  $\kappa$ -deformed Poincaré algebra we require the following conditions:

$$\frac{1}{M^2} + \frac{a^2}{\kappa^2} = 0, \qquad \frac{1}{R^2} + \tilde{\kappa}^2 b^2 = 0$$
(50)

to remove the terms proportional to  $\tilde{M}_{\mu\nu}$  (i.e. the curvature terms) on the RHS of eqs. (39) and (40).

Subsequently, we will consider only two cases<sup>12</sup> (see e.g. [31]):

# 1. Doubly time-like $\kappa$ deformations

If we require  $a^2 = -\frac{\kappa^2}{M^2}$  then the curvature terms on the RHS of (39) vanish and we get pure  $\kappa$ -deformation type of commutation for coordinates. Analogously, the condition  $b^2 = -\frac{1}{R^2 \tilde{\kappa}^2}$  leads to the new  $\tilde{\kappa}$  type commutator for momenta. The time-like  $\kappa$ -deformations for coordinates and momenta are imposed by the conditions  $a^2 = -1$  and  $b^2 = -1$ , which due to (50) imply the following relations between parameters:  $\kappa = M$  and  $\tilde{\kappa} = R^{-1}$ . We obtain:

$$[\tilde{x}_{\mu}, \tilde{x}_{\nu}] = \frac{i\hbar}{M} (a_{\mu}\tilde{x}_{\nu} - a_{\nu}\tilde{x}_{\mu}), \qquad (51)$$

$$[\tilde{p}_{\mu}, \tilde{p}_{\nu}] = \frac{i\hbar}{R} (b_{\mu}\tilde{p}_{\nu} - b_{\nu}\tilde{p}_{\mu}), \qquad (52)$$

$$[\tilde{M}_{\mu\nu}, \tilde{x}_{\rho}] = i\hbar(\eta_{\mu\rho}\tilde{x}_{\nu} - \eta_{\nu\rho}\tilde{x}_{\mu}) + \frac{i\hbar}{M}(a_{\nu}\tilde{M}_{\mu\rho} - a_{\mu}\tilde{M}_{\nu\rho}),$$
(53)

$$[\tilde{M}_{\mu\nu}, \tilde{p}_{\rho}] = i\hbar(\eta_{\mu\rho}\tilde{p}_{\nu} - \eta_{\nu\rho}\tilde{p}_{\mu}) + \frac{i\hbar}{R}(b_{\nu}\tilde{M}_{\mu\rho} - b_{\mu}\tilde{M}_{\nu\rho}).$$
(54)

In the case  $\tilde{r} = \hat{r}$ , the commutators (51)-(54) are supplemented by the relations

$$[\tilde{x}_{\mu}, \tilde{q}_{\nu}] = i\eta_{\mu\nu}(\tilde{r} + \frac{1}{M}a^{\rho}\tilde{q}_{\rho} - \frac{1}{R}b^{\rho}\tilde{x}_{\rho} - \frac{1}{MR}a^{\rho}b^{\sigma}\tilde{M}_{\rho\sigma}) + \frac{i}{R}b_{\mu}\tilde{x}_{\nu} - \frac{i}{M}a_{\nu}\tilde{q}_{\mu} + \frac{i}{MR}a^{\rho}b_{\rho}\tilde{M}_{\mu\nu},$$
(55)

$$[\tilde{r}, \tilde{x}_{\mu}] = \frac{i\hbar}{M^2} (\tilde{q}_{\mu} - \frac{1}{R} b^{\rho} \tilde{M}_{\mu\rho}), \qquad [\tilde{r}, \tilde{q}_{\mu}] = -\frac{i\hbar}{R^2} (\tilde{x}_{\mu} - \frac{1}{M} a^{\rho} \tilde{M}_{\mu\rho}), \qquad [\tilde{r}, \tilde{M}_{\mu\nu}] = 0.$$
(56)

If we consider  $\tilde{r}$  generator defined by the formula (45), the commutators (51)-(54) should be supplemented by the following ones:

$$[\tilde{x}_{\mu}, \tilde{q}_{\nu}] = i\hbar \left[ \eta_{\mu\nu} \tilde{r} + \frac{1}{R} b_{\mu} \tilde{x}_{\nu} - \frac{1}{M} a_{\nu} \tilde{q}_{\mu} + \frac{1}{MR} a^{\rho} b_{\rho} \tilde{M}_{\mu\nu} \right],$$
(57)

$$[\tilde{r}, \tilde{x}_{\mu}] = i\hbar \left[ -\frac{1}{M} a_{\mu} \tilde{r} - \frac{1}{MR} a^{\rho} b_{\rho} \tilde{x}_{\mu} \right],$$
(58)

<sup>&</sup>lt;sup>12</sup> There are 9 choices of double  $\kappa$ -deformations, which could be considered from mathematical point of view, which correspond to  $3 \times 3 = 9$  choices of fourvectors  $a_{\mu}$ ,  $b_{\mu}$  with their covariant lengths  $a^2$ ,  $b^2$  satisfying the three-fold choice (-1, 0, 1).

10

$$[\tilde{r}, \tilde{q}_{\mu}] = i\hbar \left[ \frac{1}{R} b_{\mu} \tilde{r} + \frac{1}{MR} a^{\rho} b_{\rho} \tilde{q}_{\mu} \right],$$
(59)

$$[\tilde{M}_{\mu\nu}, \tilde{r}] = i\hbar \left[ \frac{1}{M} (a_{\mu}\tilde{q}_{\nu} - a_{\nu}\tilde{q}_{\mu}) - \frac{1}{R} (b_{\mu}\tilde{x}_{\nu} - b_{\nu}\tilde{x}_{\mu}) \right].$$
(60)

# 2. Doubly light-cone $\kappa$ deformations

One can obtain light-cone  $\kappa$ -deformation for space-time coordinates by imposing  $a^2 = 0$ . In such a case we have to take the additional limit  $M \to \infty$  to remove the curvature terms from (39). Analogously (see (40)), for the fourmomenta we should impose the conditions  $b^2 = 0$  and  $R \to \infty$ . We obtain the pair of relations (cf. (13), (14)):

$$[\tilde{x}_{\mu}, \tilde{x}_{\nu}] = \frac{i\hbar}{\kappa} (a_{\mu}\tilde{x}_{\nu} - a_{\nu}\tilde{x}_{\mu}), \qquad (61)$$

$$[\tilde{p}_{\mu}, \tilde{p}_{\nu}] = i\hbar\tilde{\kappa}(b_{\mu}\tilde{p}_{\nu} - b_{\nu}\tilde{p}_{\mu}).$$
(62)

When  $\tilde{r} = \hat{r}$ , the formulas (44) take the form

$$[\tilde{r}, \tilde{x}_{\mu}] = [\tilde{r}, \tilde{q}_{\mu}] = [\tilde{r}, \tilde{M}_{\mu\nu}] = 0$$
(63)

and because the commutators (41)-(43) are not M, R -dependent they remain unchanged. Moreover, from (63) it follows that  $\tilde{r} = \hat{r} \sim I$  (i.e., it is proportional to the identity operator) and the algebra presented here will correspond to the algebra presented in the next section (see Sec. 3, doubly light-cone  $\kappa$  deformations).

If we consider generator  $\tilde{r}$  defined by the formula (45), the commutators (61)-(62) are supplemented by the following ones:

$$[\tilde{r}, \tilde{x}_{\mu}] = i\hbar \left[ -\frac{1}{\kappa} a_{\mu} \tilde{r} - \frac{\tilde{\kappa}}{\kappa} a^{\rho} b_{\rho} \tilde{x}_{\mu} \right],$$
(64)

$$[\tilde{r}, \tilde{q}_{\mu}] = i\hbar \left[ \tilde{\kappa} b_{\mu} \tilde{r} + \frac{\tilde{\kappa}}{\kappa} a^{\rho} b_{\rho} \tilde{q}_{\mu} \right],$$
(65)

and we get the formulae (42)-(43), (46), (49).

# III. FROM TSR TO NEW DOUBLY $\kappa$ DEFORMED PHASE SPACE

Another way to obtain the doubly  $\kappa$  deformed phase spaces is to use a similar approach as the one presented in the previous Sec. 2.C but now in the case of Triply Special Relativity (TSR) algebra, which was firstly formulated in 2004 [27] by the following set of commutation relations:

$$[\hat{x}_{\mu}, \hat{x}_{\nu}] = \frac{i\hbar}{M^2} \hat{M}_{\mu\nu}, \qquad [\hat{p}_{\mu}, \hat{p}_{\nu}] = \frac{i\hbar}{R^2} \hat{M}_{\mu\nu}, \qquad M, R \in \mathbb{R}$$
(66)

$$[\hat{M}_{\mu\nu}, \hat{x}_{\rho}] = i\hbar(\eta_{\mu\rho}\hat{x}_{\nu} - \eta_{\nu\rho}\hat{x}_{\mu}), \qquad (67)$$

$$[\hat{M}_{\mu\nu}, \hat{p}_{\rho}] = i\hbar(\eta_{\mu\rho}\hat{p}_{\nu} - \eta_{\nu\rho}\hat{p}_{\mu}), \qquad (68)$$

$$[\hat{M}_{\mu\nu}, \hat{M}_{\rho\tau}] = i\hbar(\eta_{\mu\rho}\hat{M}_{\nu\tau} - \eta_{\mu\tau}\hat{M}_{\nu\rho} + \eta_{\nu\tau}\hat{M}_{\mu\rho} - \eta_{\nu\rho}\hat{M}_{\mu\tau}),$$
(69)

$$[\hat{x}_{\mu}, \hat{p}_{\nu}] = i\hbar \hat{g}_{\mu\nu}, \qquad \hat{g}_{\mu\nu} = \eta_{\mu\nu} + \frac{1}{R^2} \hat{x}_{\mu} \hat{x}_{\nu} + \frac{1}{M^2} \hat{p}_{\mu} \hat{p}_{\nu} + \frac{1}{MR} (\hat{x}_{\mu} \hat{p}_{\nu} + \hat{p}_{\mu} \hat{x}_{\nu} - \hat{M}_{\mu\nu}). \tag{70}$$

Formulas (66)-(70) satisfy Jacobi identities. We point out that (70) is not unique, there are infinitely many relations which would satisfy the required Jacobi identities. Note that the subalgebras spanned by the pairs  $(\hat{M}, \hat{x})$  and  $(\hat{M}, \hat{p})$  are just the standard de Sitter algebras.

The generators  $\hat{g} = \{\hat{x}_{\mu}, \hat{p}_{\mu}, \hat{M}_{\mu\nu}\}$  are Hermitian i.e.  $\hat{g} = \hat{g}^{\dagger}$ . This leads to the following condition

$$\hat{g}_{\mu\nu} = \hat{g}^{\dagger}_{\mu\nu}.\tag{71}$$

After some straightforward manipulations we obtain that:

$$g_{\mu\nu} - g^{\dagger}_{\mu\nu} = \frac{i\hbar}{MR} (\frac{2}{MR} \hat{M}_{\mu\nu} + \hat{g}_{\mu\nu} - \hat{g}_{\nu\mu}).$$
(72)

We can compute the expression  $\hat{g}_{\mu\nu} - \hat{g}_{\nu\mu}$  appearing above, by using (70) and we get [39]:

$$(\hat{g}_{\mu\nu} - \hat{g}_{\nu\mu}) = -\frac{2}{MR}\hat{M}_{\mu\nu}.$$
(73)

It means that Hermitian conditions for  $\hat{g}$  and  $\hat{g}_{\mu\nu}$  are consistent with TSR algebra without any additional assumptions.

Now, we can change the basis of the TSR algebra (66)-(70) using linear transformations (38), what leads to the following modified commutators for coordinates and momenta

$$[\tilde{x}_{\mu}, \tilde{x}_{\nu}] = i\hbar(\frac{1}{M^2} + \frac{a^2}{\kappa^2})\tilde{M}_{\mu\nu} + \frac{i\hbar}{\kappa}(a_{\mu}\tilde{x}_{\nu} - a_{\nu}\tilde{x}_{\mu}),$$
(74)

and

$$[\tilde{p}_{\mu}, \tilde{p}_{\nu}] = i\hbar (\frac{1}{R^2} + \tilde{\kappa}^2 b^2) \tilde{M}_{\mu\nu} + i\hbar \tilde{\kappa} (b_{\mu} \tilde{p}_{\nu} - b_{\nu} \tilde{p}_{\mu}),$$
(75)

where  $a^2 = a^{\mu}a_{\mu}$  and  $b^2 = b^{\mu}b_{\mu}$ , as before. The remaining commutators of TSR algebra in the new basis described by the formulae (38) have the following form:

$$[\tilde{M}_{\mu\nu}, \tilde{x}_{\rho}] = i\hbar(\eta_{\mu\rho}\tilde{x}_{\nu} - \eta_{\nu\rho}\tilde{x}_{\mu}) + \frac{i\hbar}{\kappa}(a_{\nu}\tilde{M}_{\mu\rho} - a_{\mu}\tilde{M}_{\nu\rho}),$$
(76)

$$[\tilde{M}_{\mu\nu}, \tilde{p}_{\rho}] = i\hbar(\eta_{\mu\rho}\tilde{p}_{\nu} - \eta_{\nu\rho}\tilde{p}_{\mu}) + i\hbar\tilde{\kappa}(b_{\nu}\tilde{M}_{\mu\rho} - b_{\mu}\tilde{M}_{\nu\rho}),$$
(77)

$$[\tilde{x}_{\mu}, \tilde{p}_{\nu}] = i\hbar \tilde{g}_{\mu\nu},\tag{78}$$

$$\tilde{g}_{\mu\nu} = \hat{g}_{\mu\nu}(\tilde{g}) + \eta_{\mu\nu}(\frac{1}{\kappa}a^{\rho}\tilde{p}_{\rho} - \tilde{\kappa}b^{\rho}\tilde{x}_{\rho} + \frac{\tilde{\kappa}}{\kappa}a^{\rho}b^{\sigma}\tilde{M}_{\sigma\rho}) + \tilde{\kappa}b_{\mu}\tilde{x}_{\nu} - \frac{1}{\kappa}a_{\nu}\tilde{p}_{\mu} + \frac{\tilde{\kappa}}{\kappa}ab\tilde{M}_{\mu\nu},$$
(79)

11

where we used the shortcut notation for:

$$\hat{g}_{\mu\nu}(\tilde{g}) = \tilde{h}_{\mu\nu} + \left(\frac{1}{\kappa R}a^{\rho} + \frac{\tilde{\kappa}}{M}b^{\rho}\right)\left(\frac{1}{\kappa R}a^{\sigma} + \frac{\tilde{\kappa}}{M}b^{\sigma}\right)\tilde{M}_{\mu\rho}\tilde{M}_{\nu\sigma}$$

$$- \left(\frac{1}{R}\tilde{x}_{\mu} + \frac{1}{M}\tilde{p}_{\mu}\right)\left(\frac{1}{\kappa R}a^{\rho} + \frac{\tilde{\kappa}}{M}b^{\rho}\right)\tilde{M}_{\nu\rho} - \left(\frac{1}{\kappa R}a^{\rho} + \frac{\tilde{\kappa}}{M}b^{\rho}\right)\tilde{M}_{\mu\rho}\left(\frac{1}{R}\tilde{x}_{\nu} + \frac{1}{M}\tilde{p}_{\nu}\right),$$

$$(80)$$

and

$$\tilde{h}_{\mu\nu} = \eta_{\mu\nu} + \frac{1}{R^2} \tilde{x}_{\mu} \tilde{x}_{\nu} + \frac{1}{M^2} \tilde{p}_{\mu} \tilde{p}_{\nu} + \frac{1}{MR} (\tilde{x}_{\mu} \tilde{p}_{\nu} + \tilde{p}_{\mu} \tilde{x}_{\nu} - \tilde{M}_{\mu\nu}).$$
(81)

One can obtain double  $\kappa$ -deformed phase space by removing the terms proportional to  $\tilde{M}_{\mu\nu}$  on the RHS of (74) and (75).

Following our discussion in Sec. 2C we will focus only on the two types of double  $\kappa$ -deformations:

#### 1. Doubly time-like $\kappa$ deformations

Similarly as in Sec. 2C (point 1) to obtain the  $\kappa$ -deformed commutator (13) for coordinates we must require  $a^2 = -\frac{\kappa^2}{M^2}$  on the RHS of (74). Analogously, the  $\kappa$ -type commutator (14) for momenta requires setting  $b^2 = -\frac{1}{R^2 \tilde{\kappa}^2}$ . While  $a^2 = -1$  and  $b^2 = -1$  give the timelike (or standard)  $\kappa$ -deformations for coordinates and momenta. This leads to the following relationships between parameters:  $\kappa = M$  and  $\tilde{\kappa} = R^{-1}$ . This way we obtain the known formulae (see (13), (14)):

$$[\tilde{x}_{\mu}, \tilde{x}_{\nu}] = \frac{i\hbar}{M} (a_{\mu}\tilde{x}_{\nu} - a_{\nu}\tilde{x}_{\mu}), \qquad (82)$$

and

$$[\tilde{p}_{\mu}, \tilde{p}_{\nu}] = \frac{i\hbar}{R} (b_{\mu}\tilde{p}_{\nu} - b_{\nu}\tilde{p}_{\mu}).$$
(83)

By imposing additional conditions on fourvectors  $a_{\mu}$  and  $b_{\mu}$ , we are able to simplify the formulas (79)-(81). The simplifications obtained by removing Lorentz generator  $\tilde{M}_{\mu\nu}$  from the right hand side of the commutator (78) can be obtained by putting  $a_{\mu} = -b_{\mu}$ . In this case we can supplement the commutators (82) and (83) by

$$[\tilde{M}_{\mu\nu}, \tilde{x}_{\rho}] = i\hbar(\eta_{\mu\rho}\tilde{x}_{\nu} - \eta_{\nu\rho}\tilde{x}_{\mu}) + \frac{i\hbar}{M}(a_{\nu}\tilde{M}_{\mu\rho} - a_{\mu}\tilde{M}_{\nu\rho}), \tag{84}$$

$$[\tilde{M}_{\mu\nu}, \tilde{p}_{\rho}] = i\hbar(\eta_{\mu\rho}\tilde{p}_{\nu} - \eta_{\nu\rho}\tilde{p}_{\mu}) + \frac{i\hbar}{R}(b_{\nu}\tilde{M}_{\mu\rho} - b_{\mu}\tilde{M}_{\nu\rho}), \tag{85}$$

$$[\tilde{x}_{\mu}, \tilde{p}_{\nu}] = i\hbar \tilde{g}_{\mu\nu},\tag{86}$$

where

$$\tilde{g}_{\mu\nu} = \eta_{\mu\nu} (1 + \frac{1}{M} a^{\rho} \tilde{p}_{\rho} + \frac{1}{R} a^{\rho} \tilde{x}_{\rho}) - \frac{1}{R} a_{\mu} \tilde{x}_{\nu} - \frac{1}{M} a_{\nu} \tilde{p}_{\mu} 
+ \frac{1}{R^{2}} \tilde{x}_{\mu} \tilde{x}_{\nu} + \frac{1}{M^{2}} \tilde{p}_{\mu} \tilde{p}_{\nu} + \frac{1}{MR} (\tilde{x}_{\mu} \tilde{p}_{\nu} + \tilde{p}_{\mu} \tilde{x}_{\nu}).$$
(87)

#### 2. Doubly light-cone $\kappa$ deformations

Similarly, as in Sec. 2C (point 2) to obtain the light-cone  $\kappa$ -deformation for space-time coordinates we impose  $a^2 = 0$ . In such a case one should take the additional limit  $M \to \infty$  in order to remove the curvature terms proportional to  $M_{\mu\nu}$  from (74). Analogously for momenta (see (75)) we have the conditions  $b^2 = 0$  and  $R \to \infty$ . In this last case the relations (13), (14) are supplemented by formulae (76)-(79), with the condition  $\hat{g}_{\mu\nu}(\tilde{g}) = \eta_{\mu\nu}$ . If we additionally put  $b_{\mu} = \lambda a_{\mu}, \lambda \in \mathbb{R}$ , one can remove Lorentz generator  $\tilde{M}_{\mu\nu}$  from the right hand side of the commutator (78).

# IV. FINAL REMARKS

In this paper we propose new models of doubly  $\kappa$ -deformed Poincaré algebras with two independent  $\kappa$ -deformations for NC quantum space-time (13) and NC quantum fourmomenta coordinates (14), where we use two independent deformation parameters  $\kappa$  and  $\tilde{\kappa}$ . In standard approach, one obtains the  $\kappa$ -deformations of space-time coordinates sector by employing the Hopf duality of the  $\kappa$ -deformed Poincaré group and  $\kappa$ -deformed Poincaré algebra. This leads to the description of quantum phase space by the Heisenberg double construction [32]-[37]. Here, the new doubly  $\kappa$ -deformed quantum phase spaces are resulting from various approaches. We have discussed the following:

i) the contraction procedure of double  $\kappa$ -deformations of Yang model with built in  $\hat{o}(2)$  internal symmetry generator  $\hat{r}$ , what permits to describe algebraically the models with Lie algebraic  $\hat{o}(1,5)$ structure. Then by exploring the Lie algebra structure  $\hat{o}(1,5;g)$  with the constant parameterdependent symmetric metric, we have obtained the doubly  $\kappa$ -deformed quantum phase space by properly fixing some of the parameters of the metric (namely  $g_{44} = g_{55} = 0$ ). Additionally, we have also proposed a change of basis directly from the Yang model to obtain the doubly  $\kappa$ -deformed quantum phase space as doubly  $\kappa$ -deformed Poincaré algebras (13), (14).

ii) As another approach we have proposed a change of basis in the doubly  $\kappa$ -deformed non-linear TSR model [27] which can be obtained from Yang model by expressing bi-linearly the fifteenth generator  $\hat{r}$  in terms of the remaining fourteen physical generators  $\hat{x}_{\mu}, \hat{p}_{\mu}, \hat{M}_{\mu\nu}$ .

As special cases we discuss doubly time-like and doubly light-cone  $\kappa$ -deformed quantum phase spaces.

It should be mentioned that the original TSR model [27] has been recently generalized [23], [38]-[40] and at present these generalizations are under our consideration. Moreover, the quantum phase spaces with the presence of noncommutativity in both space-time coordinates and fourmomenta sectors have been recently considered in the context of Extended Generalized Uncertainty Principle (EGUP) and the Liouville theorem [41]. The canonical commutation relations of quantum phase space which involve both NC space-time coordinates and NC fourmomenta, in general, will lead to various types of EGUPs (see e.g. [42]) and one can expect that the modifications to the Heisenberg uncertainty relation may result in new thermodynamical properties of statistical systems [41], [43], [44].

#### ACKNOWLEDGEMENTS

J. Lukierski, A. Pachoł and M. Woronowicz acknowledge the support of the Polish NCN grant 2022/45/B/ST2/01067. S. Mignemi recognizes the support of Gruppo Nazionale di Fisica Matem-

atica. A. Pachoł acknowledges COST Action CaLISTA CA21109.

- [1] M. Born, Proceedings of the Royal Society (London) A165, 291 (1938)
- [2] M. Born, Rev. Mod. Phys. 21, 463 (1949)
- [3] L. Freidel, R. G. Leigh and D. Minic, Phys. Lett. B 730 (2014), 302 [arXiv:1307.7080 [hep-th]].
- [4] S. Doplicher, K. Fredenhagen and J. E. Roberts, Commun. Math. Phys. 172 (1995), 187 [arXiv:0303037 [hep-th]].
- [5] C.N. Yang, Phys. Rev. D 47 (1947), 874
- [6] H. S. Snyder, Phys. Rev. 71, 38 (1947)
- [7] A. Kehagias, J. Madore, J. Mourad and G. Zoupanos, J. Math. Phys. 36 (1995), 5855 [arXiv:hep-th/9502017 [hep-th]].
- [8] P. Aschieri, M. Dimitrijevic, F. Meyer and J. Wess, Class. Quant. Grav. 23 (2006), 1883 [arXiv:hep-th/0510059 [hep-th]].
- [9] E.J. Beggs and S. Majid, Quantum Riemannian Geometry, Grundlehren der mathematischen Wissenschaften, vol. 355, Springer (2020) 809pp.
- [10] S. Majid, "Quantum Gravity: are we there yet?," [arXiv:2402.18536 [gr-qc]].
- [11] N. Seiberg and E. Witten, JHEP 09 (1999), 032 [arXiv:hep-th/9908142 [hep-th]].
- [12] P. Kosinski, P. Maslanka, 'From Field Theory to Quantum Groups' proceedings, ed. B. Jancewicz, J. Sobczyk, World Scientific, Singapore 1996 [arXiv:q-alg/9512018 [q-alg]]
- J. Lukierski, V. Lyakhovsky and M. Mozrzymas, Phys. Lett. B 538 (2002), 375 [arXiv:hep-th/0203182 [hep-th]].
- [14] P. Kosinski, P. Maslanka, J. Lukierski and A. Sitarz, Contribution to: Conference on Topics in Mathematical Physics, General Relativity, and Cosmology on the Occasion of the 75th Birthday of Jerzy F. Plebanski, 255 [arXiv:hep-th/0307038 [hep-th]].
- [15] J. Lukierski, S. Meljanac, S. Mignemi, A. Pachoł and M. Woronowicz, Phys. Lett. B 854 (2024), 138729 [arXiv:2311.16994 [hep-th]].
- [16] T. Martinić -Bilać, S. Meljanac, S.Mignemi, Eur. Phys. J. C (2024) 84, 846 [arXiv: 2404.01792 [hep-th]].
- [17] J. Lukierski and A. Pachoł, PoS CORFU2023 (2024), 247 [arXiv:2405.16497 [hep-th]].
- [18] S. Meljanac, D. Meljanac, A. Samsarov and M. Stojic, [arXiv:0909.1706 [math-ph]].
- [19] S. Meljanac, D. Meljanac, A. Samsarov and M. Stojic, Mod. Phys. Lett. A 25 (2010), 579 [arXiv:0912.5087 [hep-th]].
- [20] S. Meljanac, D. Meljanac, A. Samsarov and M. Stojic, Phys. Rev. D 83 (2011), 065009 [arXiv:1102.1655].
- [21] S. Meljanac and S. Mignemi, Phys. Lett. B 814, 136117 (2021) [arXiv:2101.05275 [hep-th]].
- [22] S. Meljanac and S. Mignemi, Phys. Rev. D 104, no.8, 086006 (2021) [arXiv:2106.08131 [physics.gen-ph]].
- [23] S. Meljanac and S. Mignemi, Phys. Lett. B 833 (2022), 137289 [arXiv:2206.04772 [hep-th]].
- [24] S. Meljanac and S. Mignemi, Phys. Rev. D 104, no.8, 086006 (2021) [arXiv:2106.08131 [physics.gen-ph]].
- [25] S. Meljanac and S. Mignemi, Symmetry 15 (2023) no.7, 1373 [arXiv:2208.10242 [quant-ph]].
- [26] J. Lukierski, S. Meljanac, S. Mignemi and A. Pachol, Phys. Lett. B 838 (2023), 137709 [arXiv:2208.06712 [hep-th]].
- [27] J. Kowalski-Glikman and L. Smolin, Phys. Rev. D 70 (2004), 065020 [arXiv:hep-th/0406276 [hep-th]].
- [28] S. Zakrzewski, J. Phys. A27, 2075 (1994);
- S. Majid, H. Ruegg, Phys. Lett. B 334 (1994), 348, [arXiv:hep-th/9405107].
- [29] A. Borowiec and A. Pachol, Eur. Phys. J. C 74 (2014) no.3, 2812 [arXiv:1311.4499 [math-ph]].
- [30] J. Kowalski-Glikman and S. Nowak, Int. J. Mod. Phys. D 12 (2003), 299 [arXiv:hep-th/0204245 [hep-th]].
- [31] A. Blaut, M. Daszkiewicz, J. Kowalski-Glikman and S. Nowak, Phys. Lett. B 582 (2004), 82 [arXiv:hep-th/0312045 [hep-th]].
- [32] P. Kosinski and P. Maslanka, (1994) [arXiv:hep-th/9411033 [hep-th]];

- [33] S. Giller, C. Gonera, P. Kosinski, and P. Maslanka, Acta Phys. Pol. B, 27, 2171 (1996) [arXiv:q-alg/9602006 [q-alg]]
- [34] J. Lukierski and A. Nowicki, Proc. 21st Intl. Colloquium on Group Theoretical Methods in Physics (Group 24: Physical and mathematical aspects of symmetries, V. K. Dobrev and H. D. Doebner, eds.), Heron Press, Sofia , 186 (1997) [arXiv:q-alg/9702003 [q-alg]];
- [35] G. Amelino-Camelia, J. Lukierski, and A. Nowicki, Phys. Atomic Nuclei, 61, 1811 (1998) [arXiv:hep-th/9706031 [hep-th]];
- [36] A. Borowiec, A. Pachol, Theor. Math. Phys. 169 (2), 1620 (2011);
- [37] S. Meljanac and A. Pachoł, Symmetry 13 (2021) no.6, 1055 [arXiv:2101.02512 [hep-th]].
- [38] S. Meljanac and R. Štrajn, SIGMA 18 (2022), 022 [arXiv:2112.12038 [math-ph]].
- [39] S. Meljanac, S. Mignemi, J.Math.Phys. 64 (2023) 2, 023505, [arXiv:2211.11755 [gr-qc]]
- [40] T. M. Bilać, S. Meljanac and S. Mignemi, [arXiv:2407.10616 [hep-th]].
- [41] A. Pachoł [arXiv:2409.05110 [hep-th]].
- [42] F. Wagner, G. Varao, I.P. Lobo, V.B. Bezerra, Phys. Rev. D 108(6), 066008 (2023).
- [43] A. Pachoł, A. Wojnar, Eur. Phys. J. C 83, 1097 (2023), [arXiv:2307.03520 [gr-qc]].
- [44] A. Pachoł, A. Wojnar, Class. Quant. Grav. 40, 195021 (2023), [arXiv:2304.08215 [gr-qc]].