# **Scattering of mesons and emergence of tetraquarks in two-dimensional QCD**

*Hagop* Sazdjian<sup>1,[∗](#page-0-0)</sup>

<sup>1</sup>Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France

**Abstract.** Scattering of two mesons is considered in the framework of twodimensional QCD in the large-*N<sup>c</sup>* limit with four different quark flavors. The scattering takes place through two coupled channels, corresponding to direct and quark-exchange processes, which are of order  $O(N_c^{-2})$  and  $O(N_c^{-1})$ , respectively. Finiteness of the scattering amplitudes to order  $N_c^{-2}$  is pointed out. The theory reduces, at low energies, to an effective theory of mesons, interacting by a quark-exchange process, by means of a contact term. The unitarization of the scattering amplitudes leads to the emergence of a tetraquark bound state, located very close to the lowest two-meson elastic threshold.

### **1 Introduction**

Two-dimensional QCD, in the large- $N_c$  limit, first introduced and analyzed by 't Hooft [\[1](#page-6-0)– [4\]](#page-6-1), is an efficient tool for probing many properties of hadrons related to the confinement of quarks. In two spacetime dimensions, confinement is a natural property of QCD, due to the strong infrared behavior of the gluon propagator. In the large-*N<sup>c</sup>* limit, inelasticity and pair creation effects are damped, in which case the confinement properties of the theory become more salient. In noncovariant gauges (light-cone, axial), gluon self-interactions and ghosts disappear. Finally, in the many-body case, the interactions between hadronic clusters are free of long-range van der Waals forces.

We investigate, in that framework, the properties of the theory in the four-body sector, made of two quarks and two antiquarks, each with different flavor, and search for the possible existence of tetraquark bound states. The investigation is concentrated on the properties of two-meson scattering amplitudes. Reviews on the large-*N<sup>c</sup>* limit can be found in [\[5](#page-6-2), [6\]](#page-6-3) and more specifically in relation with the tetraquark problem in [\[7\]](#page-6-4).

#### **2 Quark-antiquark sector**

The quark fields are considered in the fundamental representation of the color-gauge group *SU(N<sub>c</sub>)*. The light-cone coordinates and momenta are defined as  $x^{\pm} = \frac{1}{\sqrt{2}}(x^0 \pm x^1)$ ,  $p_{\pm} =$  $\frac{1}{\sqrt{2}}(p_0 \pm p_1)$ . In the light-cone gauge, the gluon field satisfies *A*− = 0. Because of the particular structure of the quark-gluon vertex in the light-cone gauge, where only the Dirac matrix γ<sup>−</sup> appears, the Dirac matrices can be factorized from the *N<sup>c</sup>* -leading expressions of all dynamical quantities.

<span id="page-0-0"></span><sup>∗</sup> e-mail: hagop.sazdjian@ijclab.in2p3.fr

In two dimensions, the coupling constant  $g$  of the theory has the dimension of mass. One may introduce the string tension  $\sigma$  according to the definition

$$
\sigma \equiv \frac{g^2 N_c}{4}.\tag{1}
$$

(g has been defined with the one-index notation of the gluon fields.) In the large- $N_c$  limit, the string tension is assumed to remain finite, which means that the coupling constant squared should decrease like 1/*Nc*.

The gluon propagator contributes with its ++ component:

<span id="page-1-2"></span>
$$
D(q) \equiv D_{++}(q) = \frac{i}{q_-^2}.
$$
 (2)

In two dimensions, this propagator is infrared singular. To avoid infrared divergences in intermediate calculations, we introduce a small mass-like parameter  $\lambda$  (behaving like the − component of a momentum) in the propagator, playing the role of an infrared cutoff; to simplify notations,  $\lambda$  will be most often replaced by an inverse quantity,  $\Lambda$ :

<span id="page-1-3"></span>
$$
\frac{i}{q_{-}^{2}} \longrightarrow \frac{i}{q_{-}^{2} + \lambda^{2}}, \qquad \Lambda \equiv \frac{\sigma}{\lambda}.
$$
 (3)

One expects that physical quantities will remain finite when the limits  $\lambda \to 0$  or  $\Lambda \to \infty$  are taken.

The quark propagator, after taking into account self-energy radiative corrections coming from the large-*N<sup>c</sup>* leading part, represented by the one-gluon contribution in its integral equation, is

<span id="page-1-1"></span>
$$
S(p) = \frac{ip_{-}}{2p_{+}p_{-} - |p_{-}|\Lambda - m'^{2} + i\epsilon}, \qquad m'^{2} = m^{2} - \frac{2\sigma}{\pi}.
$$
 (4)

One notices that in the limit  $\Lambda \to \infty$ , the renormalized mass of the quark goes to infinity, indicating that the quark ceases to be an observable free particle with a finite mass.

For quark-antiquark systems in the color-singlet sector, one can write down the corresponding Bethe–Salpeter equation, where the kernel is given, at leading order of *N<sup>c</sup>* , by the one-gluon exchange diagram. Designating by *r* the total momentum and by *p* the quark momentum, one introduces the bound state wave function  $\phi(r, p)$ . Noticing that the interaction in the light-cone gauge is lightlike instantaneous, one further introduces the instantaneous wave function  $\varphi$  defined as

<span id="page-1-0"></span>
$$
\varphi(r, p_{-}) = \int \frac{dp_{+}}{2\pi} \phi(r, p). \tag{5}
$$

ϕ satisfies spectral conditions that imply that the momentum component *p*<sup>−</sup> should be positive and bounded by *r*<sup>−</sup> , which itself is positive. Defining then the reduced variable *x* as

<span id="page-1-4"></span>
$$
x = \frac{p_-}{r_-}, \qquad 0 \le x \le 1,\tag{6}
$$

the equation satisfied by  $\varphi$  becomes

$$
\left[r^2 - \frac{m_1^2}{x} - \frac{m_2^2}{(1-x)}\right]\varphi(x) = -\left(\frac{2\sigma}{\pi}\right)\int_0^1 dy \frac{(\varphi(y) - \varphi(x))}{(y-x)^2},\tag{7}
$$

where  $m_1$  and  $m_2$  are the quark and antiquark masses, respectively. This is the 't Hooft equation [\[2](#page-6-5)]. One notices that the infrared cutoff has disappeared, related to the fact that the eigenvalues  $r^2$ , which are to be determined, correspond to the physical meson masses.

The spectrum of the eigenfunctions and eigenvalues corresponds to an infinite tower of meson states, characterized by a discrete quantum number *n*, with a linear increase with *n* of the higher masses squared, a feature that is typical of a confinement regime.

The quark-antiquark scattering amplitude  $\mathcal T$  has been calculated in [\[3\]](#page-6-6). Its expression is

<span id="page-2-1"></span>
$$
\mathcal{T}(r, x, x') = -\frac{2\sigma}{N_c} \frac{1}{r_-^2 (x - x')^2} + \frac{1}{N_c} \sum_n \frac{\phi_n(r, x) \phi_n^*(r, x')}{(r^2 - r_n^2)},
$$
\n(8)

where  $\phi_n$  is obtained from  $\phi_n$  after factorizing in the latter the free quark-antiquark Green's function. One notices that apart from the one-gluon exchange term (the first term), which cannot be absorbed into products of wave functions, the scattering amplitude is entirely given by the meson wave function contributions. The scattering amplitude  $\mathcal T$  is cutoff dependent through the complete wave functions  $\phi$  or  $\phi$ , which, contrary to the instantaneous wave function  $\varphi$  [Eq. [\(5\]](#page-1-0), are cutoff dependent. Therefore,  $\tau$  is not a directly observable quantity. Furthermore, it explicitly contains the long-range one-gluon exchange interaction. In this connection, one of the basic requirements from the theory is that in observable processes, like meson-meson scattering, the cutoff dependences, as well as the long-range contributions, disappear from the final expressions.

## **3 Two-quark–two-antiquark systems**

We now consider systems made of two quarks and two antiquarks. The quark flavors are assumed different from each other; this case avoids mixing problems with ordinary meson states. The quarks are denoted with labels 1 and 3, with masses  $m_1$  and  $m_3$ , and the antiquarks with labels  $\bar{2}$  and  $\bar{4}$ , with masses  $m_2$  and  $m_4$ . The Green's function, corresponding to the four ingoing and outgoing particles, is schematically represented in Fig. [1.](#page-2-0)



<span id="page-2-0"></span>Figure 1. Four-body Green's function, with two quarks, 1 and 3, and two antiquarks,  $\bar{2}$  and  $\bar{4}$ .

We are interested in the color-singlet sectors of the incoming and outgoing particles in the Green's function, describing scattering processes of mesons. We have two sets of mesonic clusters:  $(1\overline{2})(3\overline{4})$  and  $(1\overline{4})(3\overline{2})$ . One can then distinguish four different channels in the scattering process, two of which are called "direct", the ingoing and outgoing clusters being the same, and the two others "recombination", the outgoing clusters having undergone a quark exchange. The four different processes are:

<span id="page-2-2"></span>
$$
(1\overline{2}) + (3\overline{4}) \longrightarrow (1\overline{2}) + (3\overline{4}), \quad \text{direct channel 1} \quad (D1), \tag{9}
$$

 $(1\bar{2}) + (3\bar{4}) \rightarrow (1\bar{4}) + (3\bar{2}),$  recombination channel 1 (*R*1), (10)

 $(1\bar{4}) + (3\bar{2}) \longrightarrow (1\bar{4}) + (3\bar{2}),$  direct channel 2 (*D*2), (11)

$$
(1\overline{4}) + (3\overline{2}) \longrightarrow (1\overline{2}) + (3\overline{4}), \quad \text{recombination channel } 2 \quad (R2). \tag{12}
$$

Integral equations for the Green's functions in the color-singlet sectors at leading orders of *N<sup>c</sup>* can be obtained following the method used in the quark-antiquark case. Their detailed

derivation is left to a separate work. We will be content by indicating here some of the qualitative aspects that characterize them. Similar derivations and related technical problems can be found in  $[8-11]$ .

The integral equations relative to the four channels decouple into two subsets of equations, the first concerning channel 1 and the second, channel 2. Inside each subset, one has two coupled integral equations relative to the direct and recombination processes. Actually, one passes from one subset to the other by appropriate permutations of indices. It is therefore sufficient to concentrate on one subset, channel 1, say.

Specifically, one finds the following large-*N<sup>c</sup>* behaviors for the scattering amplitudes:

$$
\mathcal{T}_{R1} = O(N_c^{-1}) + O(N_c^{-3}), \tag{13}
$$

$$
\mathcal{T}_{D1} = O(N_c^{-2}) + O(N_c^{-4}). \tag{14}
$$

The on-mass shell scattering amplitudes are obtained by projecting the off-mass shell scattering amplitudes on the ingoing and outgoing wave functions.

# **4** Finiteness of the scattering amplitudes to order  $1/N_c^2$

An important test for the theory is the verification that the meson-meson scattering amplitudes are finite quantities in the limit  $\Lambda \to \infty$ . The infrared cutoff is present in the quark propagators, the gluon propagator and the external wave functions. Another test is the verification that the meson-meson scattering amplitudes are free of long-range van der Waals type forces.

The calculations take into account the following features. Each quark propagator carries a damping factor  $\Lambda^{-1}$ . The gluon propagator, upon integration, produces a diverging factor  $\Lambda$ . Each external meson wave function carries a diverging power Λ. Each quark loop integration with respect to the + component of the loop momentum removes one power from the damping factors. Previous calculations of this type have been presented in [\[3](#page-6-6)].

Generally, individual diagrams are linearly divergent in Λ. One therefore has to consider combinations of several diagrams having similar structures.

The recombination channel process receives, at order  $N_c^{-1}$ , contributions from three diagrams, one of them being free of gluon propagators and the two others containing one gluon propagator. It turns out that the sum of the three diagrams is finite in the limit  $\Lambda \to \infty$ . The finite part has the structure of an effective four-meson contact term with smooth momentum dependence, excluding any presence of long-range forces. The effective coupling constant of the contact term is provided by an overlapping integral of the four external meson wave functions.

Since the next-to-leading term in the recombination channel, generated by the leading terms in the corresponding Green's function, is of order  $N_c^{-3}$ , one should check whether nonleading terms of the  $1/N_c$  expansion might produce divergences at order  $1/N_c^2$ . These arise from quark-loop corrections to the gluon propagator, from vertex corrections and from crossed diagrams.

The quark-loop correction can be calculated using the propagator [\(4\)](#page-1-1). The result is a modification of the gluon propagator  $(2)-(3)$  $(2)-(3)$  to the form [\[3\]](#page-6-6)

$$
D(q) = \frac{i}{q_{-}^{2} + \lambda^{2} + \lambda |q_{-}| N_{f}/(\pi N_{c})},
$$
\n(15)

where  $N_f$  is the number of quark flavors. The net effect of this change is a renormalization of the cutoff parameter  $\Lambda$  in the following way:

$$
\Lambda \longrightarrow \Lambda \left(1 - \frac{N_f}{\pi^2 N_c}\right). \tag{16}
$$

The same change of  $\Lambda$  also occurs in the quark propagator [\(4\)](#page-1-1),  $m^2$  remaining unchanged. Therefore, the cancellation mechanism between gluon and quark propagator divergences remains unaffected.

Vertex corrections and gluon crossed diagrams produce generally leading corrections of order  $1/N_c$ . However, in the present case, the latter vanish due to the spectral conditions that the meson wave functions should satisfy [Eqs.  $(5)$  and  $(6)$ ].

Therefore, the finiteness of the recombination channel scattering amplitude remains true also at order  $N_c^{-2}$ .

The direct channel process receives, at order  $N_c^{-2}$ , contributions from 65 diagrams, containing up to four gluon propagators. These diagrams can be be grouped, according to their structure, into five categories, which can be studied independently from each other. The net result is that the divergent parts cancel out in each category and one remains with a finite part which is nothing but the generalized two-meson unitarity diagram generated by the finite part of the recombination process. (The two-meson loops are actually infinite in number, involving all mesons of the quark-antiquark spectra, [Eq. [\(8\)](#page-2-1)].)

The above results establish the finiteness of the two-meson scattering amplitudes to order  $N_c^{-2}$  and provide the equivalent low-energy effective meson theory, which reduces to the recombination process contact term and the generation from it of unitarity diagrams.

#### **5 Unitarized scattering amplitudes**

The extension of the above study to higher orders than  $N_c^{-2}$  becomes out of reach, because of the increasing number of diagrams and also because of the appearance of new types of diagrams. For the last category, we might observe that even if they produce new types of effective interactions, they could not compete, at large  $N_c$ , with those found up to order  $N_c^{-2}$ . It is therefore not unreasonable to ignore, for a first study, such possible new effects.

Concerning the higher-order contributions generated by the leading-order diagrams through the Green's functions, we observe that the results obtained above indicated the tendancy of the theory to unitarize the effective contact interaction generated by the recombination process. Assuming that the resulting effective theory satisfies the unitarity property, one is entitled to complete the results obtained up to order  $N_c^{-2}$  by unitarizing the corresponding scattering amplitudes.

We undertake that operation using two approximations. First, we keep in the series of intermediate meson states that contribute to the unitarity loop diagrams only the ground state mesons, neglecting their radial excitations. The latter contribute with dampted mass factors and with smaller four-meson couplings, due to the smallness of the overlapping integrals between ground state and radial wave functions. Second, we neglect the momentum dependence of the effective four-meson coupling of the ground-state mesons, by treating it as a constant. With these approximations, the relevant equations are reduced to coupled four-channel equations [Eqs.  $(9)-(12)$  $(9)-(12)$  $(9)-(12)$ ], that could be dealt with in matrix form.

Designating by  $K_R$  the real coupling constant of the recombination channel (unitarity requiring the equality  $K_{R1} = K_{R2}$ , the scattering amplitude and the kernel of its integral equation take the following matrix forms:

<span id="page-4-0"></span>
$$
\mathcal{T} = \begin{pmatrix} \mathcal{T}_{D1} & \mathcal{T}_{R2} \\ \mathcal{T}_{R1} & \mathcal{T}_{D2} \end{pmatrix}, \quad K = \begin{pmatrix} 0 & K_R \\ K_R & 0 \end{pmatrix}, \quad J = \begin{pmatrix} J_{1\bar{2},3\bar{4}} & 0 \\ 0 & J_{1\bar{4},3\bar{2}} \end{pmatrix}, \tag{17}
$$

where we have also introduced the two-meson loop functions  $J_{1\bar{2},3\bar{4}}$  and  $J_{1\bar{4},3\bar{2}}$ :

<span id="page-5-0"></span>
$$
J_{a\bar{b},c\bar{d}}(s, M_{a\bar{b}}, M_{c\bar{d}}) = -\frac{i}{2\sqrt{-\lambda_{a\bar{b},c\bar{d}}}} \Big[ 1 - \frac{1}{\pi} \arctan \Big( \frac{\sqrt{-\lambda_{a\bar{b},c\bar{d}}}}{s - (M_{a\bar{b}}^2 + M_{c\bar{d}}^2)} \Big) \Big],
$$
  

$$
\lambda_{a\bar{b},c\bar{d}} = \Big( s - (M_{a\bar{b}} + M_{c\bar{d}})^2 \Big) \Big( s - (M_{a\bar{b}} - M_{c\bar{d}})^2 \Big).
$$
 (18)

When the interaction is represented by a contact term, as is the case here, the unitarization operation reduces to the iteration of the interaction by means of the meson loops. One has

<span id="page-5-1"></span>
$$
\mathcal{T} = (1 - iKJ)^{-1}K
$$
  
= 
$$
\frac{1}{\left(1 + K_R J_{1\bar{4},3\bar{2}} K_R J_{1\bar{2},3\bar{4}}\right)} \left( \begin{array}{cc} iK_R J_{1\bar{4},3\bar{2}} K_R & K_R \\ K_R & iK_R J_{1\bar{2},3\bar{4}} K_R \end{array} \right),
$$
 (19)

from which one may identify, with [\(17\)](#page-4-0), the different components of  $\mathcal{T}$ .

### **6 Tetraquarks**

Tetraquark bound states may occur at the zeros of the denominator of  $\mathcal T$ . This leads to the equation

$$
1 + K_R J_{1\bar{4},3\bar{2}} K_R J_{1\bar{2},3\bar{4}} = 0. \tag{20}
$$

This equation has generally one bound state solution, which lies very close to the lowest elastic two-meson threshold, with binding energies of the order of 1 MeV or much less, according to the mass configurations of the mesons. The smallness of the binding energy is traced back to the smallness of the coupling constant  $K_R$ , which, within a rough estimate, is of the order of  $10(2\sigma/\pi)$ , with respect to some combinations of the meson masses. In the case of two heavy quarks, the latter ratio effectively contributes with a power of 4.

For the particular example of the configuration  $(c\bar{s}b\bar{u})$ , which leads, in the pseudoscalar sector, to the two two-meson clusters  $(D^+B^-)$  and  $(D^0\bar{B}^0_s)$ , the binding energy comes out of the order of 0.02 MeV with respect to the  $(D^0 \bar{B}^0_s)$  elastic threshold.

Existence of tetraquark bound states with four different quark flavors, containing *b* and *c* quarks, has been observed in Lattice QCD calculations [\[12](#page-6-9)[–14\]](#page-6-10).

Of particular interest is the comparison of the coupling constants of the tetraquark state to the two meson clusters. Assuming for definiteness that  $(M_{1\bar{2}} + M_{3\bar{4}}) > (M_{1\bar{4}} + M_{3\bar{2}})$ , and denoting the tetraquark binding energy by  $E_{TB}$  and  $\Delta M = (M_{1\bar{2}} + M_{3\bar{4}}) - (M_{1\bar{4}} + M_{3\bar{2}})$ , one observes from Eqs. [\(18\)](#page-5-0) and [\(19\)](#page-5-1), for nonrelativistic systems, that the coupling constant squared of the tetraquark to the clusters (12)(34) is proportional to  $E_{TB}^{-1/2}$ , while that to the clusters (14)(32) is proportional to  $(\Delta M)^{-1/2}$ . The ratio of these quantities for the example considered above is of the order of 30, which means that the tetraquark couples more strongly to the heavier two-meson clusters than to its neighboring ones.

The latter property would suggest at first sight that the tetraquark is essentially a molecular system made of the heavier cluster ingredients, here  $(1\bar{2})(3\bar{4})$ . However, such an interpretation, neglecting the second set of clusters, would not explain in a natural way the particular position of the tetraquark pole in the vicinity of the lighter clusters. The two two-meson clusters seem to play an equal role. This situation is due to the quark-exchange mechanism, which is the basic interaction ingredient of the two-meson clusters, and which needs the presence of the two distinct two-meson clusters to operate.

# **7 Conclusion**

Two-dimensional QCD at large *N<sup>c</sup>* allows a consistent treatment of the confining interactions, with possible control of the infrared divergences. In the sector of four different flavors, the theory reduces, at low energies, to an effective theory of mesons, where the basic interaction is provided by a quark-exchange four-meson contact term. The unitarization of the scattering amplitude in all channels of interaction leads to the prediction of the existence of a tetraquark bound state lying very close to the lightest two-meson elastic threshold.

The author thanks W. Lucha and D. Melikhov for stimulating discussions on the subject. This research has received financial support from the EU research and innovation programme Horizon 2020, under Grant agreement No. 824093.

## **References**

- <span id="page-6-0"></span>[1] G. 't Hooft, A planar diagram theory for strong interactions, Nucl. Phys. B 72, 461 (1974). 10.1016/[0550-3213\(74\)90154-0](https://doi.org/10.1016/0550-3213(74)90154-0)
- <span id="page-6-5"></span>[2] G. 't Hooft, A two-dimensional model for mesons, Nucl. Phys. B 75, 461 (1974). 10.1016/[0550-3213\(74\)90088-1](https://doi.org/10.1016/0550-3213(74)90088-1)
- <span id="page-6-6"></span>[3] C.G. Callan, Jr., N. Coote, D.J. Gross, Two-dimensional Yang-Mills theory: a model of quark confinement, Phys. Rev. D 13, 1649 (1976). 10.1103/[PhysRevD.13.1649](https://doi.org/10.1103/PhysRevD.13.1649)
- <span id="page-6-1"></span>[4] M.B. Einhorn, Form-factors and deep inelastic scattering in two-dimensional Quantum Chromodynamics, Phys. Rev. D 14, 3451 (1976). 10.1103/[PhysRevD.14.3451](https://doi.org/10.1103/PhysRevD.14.3451)
- <span id="page-6-2"></span>[5] E. Witten, Baryons in the 1/*N* expansion, Nucl. Phys. B 160, 57 (1979). 10.1016/[0550-3213\(79\)90232-3](https://doi.org/10.1016/0550-3213(79)90232-3)
- <span id="page-6-3"></span>[6] S. Coleman, Aspects of Symmetry (Cambridge University Press, Cambridge, U.K., 1985), ISBN 9780521318273, 0521318270
- <span id="page-6-4"></span>[7] W. Lucha, D. Melikhov, H. Sazdjian, Tetraquarks in large-*N<sup>c</sup>* QCD, Prog. Part. Nucl. Phys. 120, 103867 (2021), 2102.02542. 10.1016/[j.ppnp.2021.103867](https://doi.org/10.1016/j.ppnp.2021.103867)
- <span id="page-6-7"></span>[8] W. Heupel, G. Eichmann, C.S. Fischer, Tetraquark bound states in a Bethe-Salpeter approach, Phys. Lett. B 718, 545 (2012), 1206.5129. 10.1016/[j.physletb.2012.11.009](https://doi.org/10.1016/j.physletb.2012.11.009)
- [9] G. Eichmann, C.S. Fischer, W. Heupel, The light scalar mesons as tetraquarks, Phys. Lett. B 753, 282 (2016), 1508.07178. 10.1016/[j.physletb.2015.12.036](https://doi.org/10.1016/j.physletb.2015.12.036)
- [10] G. Eichmann, C.S. Fischer, W. Heupel, N. Santowsky, P.C. Wallbott, Four-quark states from functional methods, Few Body Syst. 61, 38 (2020), 2008.10240. 10.1007/[s00601-020-01571-3](https://doi.org/10.1007/s00601-020-01571-3)
- <span id="page-6-8"></span>[11] A.N. Kvinikhidze, B. Blankleider, Covariant tetraquark equations in quantum field theory, Phys. Rev. D 106, 054024 (2022), 2102.09558. 10.1103/[PhysRevD.106.054024](https://doi.org/10.1103/PhysRevD.106.054024)
- <span id="page-6-9"></span>[12] C. Alexandrou, J. Finkenrath, T. Leontiou, S. Meinel, M. Pflaumer, M. Wagner, Shallow bound states and hints for broad resonances with quark content  $\bar{b}\bar{c}ud$  in  $B - \bar{D}$  and *B*<sup>\*</sup> –  $\bar{D}$  scattering from Lattice QCD, Phys. Rev. Lett. 132, 151902 (2024), 2312.02925. 10.1103/[PhysRevLett.132.151902](https://doi.org/10.1103/PhysRevLett.132.151902)
- [13] M. Padmanath, A. Radhakrishnan, N. Mathur, Bound isoscalar axial-vector  $bc\bar{u}\bar{d}$ tetraquark *Tbc* from Lattice QCD using two-meson and diquark-antidiquark variational basis, Phys. Rev. Lett. 132, 201902 (2024), 2307.14128. 10.1103/Phys-RevLett.132.201902
- <span id="page-6-10"></span>[14] N. Mathur, M. Padmanath,  $\bar{b}\bar{c}q_1q_2$  four-quark states from Lattice QCD, PoS LAT-TICE2021, 443 (2022), 2111.01147. 10.22323/[1.396.0443](https://doi.org/10.22323/1.396.0443)