

INVARIANT MEASURES AND THEIR LIMITING BEHAVIOR OF THE LANDAU-LIFSHITZ-BLOCH EQUATION IN UNBOUNDED DOMAIN

DAIWEN HUANG, ZHAOYANG QIU, AND BIXIANG WANG

ABSTRACT. This paper deals with the existence and limiting behavior of invariant measures of the stochastic Landau-Lifshitz-Bloch equation driven by linear multiplicative noise and additive noise defined in the entire space \mathbb{R}^d for $d = 1, 2$, which describes the phase spins in ferromagnetic materials around the Curie temperature. We first establish the existence and uniqueness of solutions by a domain expansion method. We then prove the existence of invariant measures by the weakly Feller argument. In the case $d = 1$, we show the uniform tightness of the set of all invariant measures of the stochastic equation, and prove any limit of a sequence of invariant measures of the perturbed equation must be an invariant measure of the limiting system. The cut-off arguments, stopping time techniques and uniform tail-ends estimates of solutions are developed to overcome the difficulty caused by the high-order nonlinearity and the non-compactness of Sobolev embeddings in unbounded domain.

1. INTRODUCTION

The Landau-Lifshitz-Bloch (LLB) equation is a model for simulating the heat-assisted magnetic recording in ferromagnetic materials around the Curie temperature, which is derived from the Landau-Lifshitz-Gilbert equation at low temperatures and the Ginzburg-Landau theory of phase transitions. And the LLB equation explains the light-induced demagnetization with powerful femtosecond lasers, see, e.g., [1, 18] for the physical backgrounds. Let \mathcal{D} be a domain in \mathbb{R}^d with $d \leq 2$. Then a specific form of the LLB equation in \mathcal{D} is given by

$$\frac{\partial \mathbf{u}}{\partial t} = \gamma \mathbf{u} \times \mathbf{H}_{eff} + \mathbf{H}_{eff},$$

where the effective field \mathbf{H}_{eff} has the form

$$\mathbf{H}_{eff} = \Delta \mathbf{u} - (1 + \kappa |\mathbf{u}|^2) \mathbf{u},$$

and $\mathbf{u} = (u_1, u_2, u_3)$ is the average spin polarization, $\gamma > 0$ represents the gyromagnetic ratio and $\kappa > 0$ is a constant related to the temperature. Without loss of generality, we will assume $\kappa = \gamma = 1$. The notation \times denotes the vector cross product, and $|\cdot|$ is the Euclidean norm of \mathbb{R}^3 . Then the deterministic LLB equation can be written more precisely as

$$d\mathbf{u} = \Delta \mathbf{u} dt + \mathbf{u} \times \Delta \mathbf{u} dt - (1 + |\mathbf{u}|^2) \mathbf{u} dt.$$

Date: October 4, 2024.

2020 Mathematics Subject Classification. 35Q35, 76D05, 35R60, 60F10.

Key words and phrases. stochastic Landau-Lifshitz-Bloch equation, unbounded domain, well-posedness, invariant measure, limiting behavior, tail-ends estimates.

For the deterministic LLB equation, Le [32] first proved the existence of weak solutions in a smooth bounded domain using the Galerkin approximation scheme and the method of compactness. Then Guo et al. [25] established the well-posedness of strong solutions of the equation on a finite-dimensional closed Riemannian manifold. In this paper, we will consider the stochastic LLB equation driven by a linear multiplicative noise and additive noise.

1.1. Motivations. In the ferromagnetic theory, a significant issue is to describe the transition between different equilibrium states induced by thermal fluctuation, which was proposed initially by Néel [35]. The random fluctuation is often introduced into the equation by perturbing the effective field H_{eff} using the Gaussian noise, see [19]; that is, H_{eff} is replaced by $H_{eff} + \frac{\circ dW}{dt}$ where $W(t) = \sum_{k=1}^{\infty} \mathbf{f}_k W_k(t)$ is a Wiener noise defined in a complete filtered probability space $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$, $\{\mathbf{f}_k\}_{k=1}^{\infty}$ is a sequence of functions satisfying

$$\sum_{k=1}^{\infty} \|\mathbf{f}_k\|_{W^{1,\infty}(\mathbb{R}^d, \mathbb{R}^3) \cap H^1(\mathbb{R}^d, \mathbb{R}^3)} < \infty, \quad (1.1)$$

and $\{W_k\}_{k \geq 1}$ is a sequence of independent real-valued Wiener processes.

Consider the case where the ferromagnet fills in the entire space \mathbb{R}^d , then the stochastic LLB equation in \mathbb{R}^d reads as

$$d\mathbf{u} = \Delta \mathbf{u} dt + \mathbf{u} \times \Delta \mathbf{u} dt - (1 + |\mathbf{u}|^2) \mathbf{u} dt + \sum_{k=1}^{\infty} (\mathbf{u} \times \mathbf{f}_k + \mathbf{f}_k) \circ dW_k, \quad (1.2)$$

which is driven by a linear multiplicative noise and additive noise in the sense of the Stratonovich differential. As explained in [16], such noise can deduce a Boltzmann distribution valid for the full range of temperatures.

Note that the Stratonovich stochastic equation (1.2) is equivalent to the Itô equation in \mathbb{R}^d :

$$d\mathbf{u} = \Delta \mathbf{u} dt + \mathbf{u} \times \Delta \mathbf{u} dt - (1 + |\mathbf{u}|^2) \mathbf{u} dt + \frac{1}{2} \sum_{k=1}^{\infty} (\mathbf{u} \times \mathbf{f}_k) \times \mathbf{f}_k dt + \sum_{k=1}^{\infty} (\mathbf{u} \times \mathbf{f}_k + \mathbf{f}_k) dW_k, \quad (1.3)$$

which is supplemented with the initial data:

$$\mathbf{u}(0, x) = \mathbf{u}_0(x), \quad x \in \mathbb{R}^d.$$

Due to its physical importance and mathematical challenges, the stochastic LLB model has been studied recently by several authors. When the equation is defined in a bounded domain \mathcal{D} , the existence of martingale solutions as well as the existence of a unique strong solutions of (1.3) with initial data in $H^1(\mathcal{D})$ were proved by Jiang et al. [26] and Brzeźniak et al. [4] respectively. In addition, the existence of invariant measures of (1.3) was proved in [4] by the weakly Feller method. When the equation is driven by a transport noise, the existence and the regularity of solutions were established by Qiu et al. [36]. Brzeźniak et al. [3] considered the large deviations and transitions between equilibria for stochastic Landau-Lifshitz-Gilbert equation. Note that in all these articles, the stochastic equation is defined in a bounded domain in \mathbb{R}^d or the torus. As far as the authors

are aware, it seems that there is no result available in the literature on the well-posedness and invariant measures of (1.3) when the domain is unbounded. The main goal of the present paper is to deal with this problem and study the existence and the limiting behavior of invariant measures of (1.3) defined in the entire space \mathbb{R}^d . The reader is referred to [12, 13, 17, 20, 21, 23, 24, 27] and [2, 5, 6, 9, 10, 15, 28, 29, 34, 37] for the well-posedness and invariant measures of stochastic partial differential equations defined in bounded domains and unbounded domains, respectively. It is worth mentioning that the main difficulty in the case of unbounded domains lies in the fact that Sobolev embeddings are no longer compact which introduces an essential obstacle for establishing the tightness of distributions of a family of solutions.

1.2. Main results. As mentioned above, the main purpose of this paper is to investigate the existence and limiting behavior of invariant measures of (1.3). To that end, we first establish the existence and uniqueness of solutions of (1.3) in \mathbb{R}^d for $d = 1, 2$, which is stated below.

Theorem 1.1. *Let $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P}, W)$ be a given stochastic basis. If (1.1) holds and $\mathbf{u}_0 \in H^1(\mathbb{R}^d, \mathbb{R}^3)$ with $d = 1, 2$, then the stochastic LLB equation (1.3) admits a unique global solution \mathbf{u} in the following sense:*

(i). *The process \mathbf{u} is weakly continuous, $H^1(\mathbb{R}^d, \mathbb{R}^3)$ -valued, \mathcal{F}_t -progressively measurable such that*

$$\mathbf{u} \in L^p(\Omega; L^\infty(0, T; H^1(\mathbb{R}^d, \mathbb{R}^3)) \cap L^2(0, T; H^2(\mathbb{R}^d, \mathbb{R}^3))), \quad \forall p \geq 1, \quad (1.4)$$

and

$$\mathbf{u} \times \Delta \mathbf{u} \in L^{pr}(\Omega; L^r(0, T; L^2(\mathbb{R}^d, \mathbb{R}^3))), \quad \forall p \geq 1, r \in [1, 2). \quad (1.5)$$

(ii). *For all $t \in [0, T]$, the following equality holds in $L^2(\mathbb{R}^d, \mathbb{R}^3)$, \mathbb{P} -almost surely,*

$$\begin{aligned} \mathbf{u}(t) = & \mathbf{u}_0 + \int_0^t \Delta \mathbf{u}(s) ds + \int_0^t \mathbf{u}(s) \times \Delta \mathbf{u}(s) ds \\ & - \int_0^t (1 + |\mathbf{u}(s)|^2) \mathbf{u}(s) ds + \frac{1}{2} \sum_{k=1}^{\infty} \int_0^t (\mathbf{u}(s) \times \mathbf{f}_k) \times \mathbf{f}_k ds + \sum_{k=1}^{\infty} \int_0^t (\mathbf{u}(s) \times \mathbf{f}_k + \mathbf{f}_k) dW_k. \end{aligned} \quad (1.6)$$

In the case of bounded domains [4, 26], the existence of solutions of the stochastic LLB equation was proved by the finite-dimensional approximation scheme and the compactness argument, where the compactness of Sobolev embeddings was used to show the tightness of approximate solutions. Since we consider the stochastic equation (1.3) in the unbounded domain \mathbb{R}^d and the Sobolev embeddings are no longer compact in unbounded domains, the arguments of [4, 26] do not apply to our case. In this paper, we will adopt a domain expansion method to show the existence of solutions of (1.3). More precisely, we first consider an approximate solution \mathbf{u}^n defined in the bounded domain $\mathcal{D}_n = \{x \in \mathbb{R}^d : |x| < n\}$, then derive the uniform estimates of approximate solutions, and finally take the limit of the sequence $\{\mathbf{u}^n\}_{n=1}^{\infty}$ to obtain a solution of (1.3) defined in \mathbb{R}^d . However, in order to pass to the limit of approximate solutions, we need to establish the tightness of distributions of

the sequence, for which we must overcome the difficulty caused by the non-compactness of Sobolev embeddings in \mathbb{R}^d . We will apply the idea of uniform tail-ends estimates to solve the problem by showing that the approximate solutions are uniformly small on the complement of \mathcal{D}_n when n is sufficiently large as in [9, 37] for the reaction-diffusion equation. By the uniform smallness of the tails of solutions in $L^2(\mathbb{R}^d) \setminus \mathcal{D}_n$ (see Lemma 2.3 (iv)) and the compactness of embeddings in bounded domains, we then prove the tightness of distributions of the approximate sequence (see Lemma 2.4). Finally, we show the limit of the approximate solutions is a martingale solution of the stochastic LLB equation in \mathbb{R}^d , which along with the pathwise uniqueness of solutions and the Yamada-Watanabe theorem implies the existence of a unique strong solution.

With the well-posedness in hand, we then investigate the existence and uniform tightness of invariant measures of (1.3) with respect to noise intensity. More precisely, we consider the following equation with a parameter $\varepsilon \in [0, 1]$:

$$d\mathbf{u}^\varepsilon = \Delta \mathbf{u}^\varepsilon dt + \mathbf{u}^\varepsilon \times \Delta \mathbf{u}^\varepsilon dt - (1 + |\mathbf{u}^\varepsilon|^2) \mathbf{u}^\varepsilon dt + \varepsilon \sum_{k=1}^{\infty} (\mathbf{u}^\varepsilon \times \mathbf{f}_k + \mathbf{f}_k) \circ dW_k, \quad (1.7)$$

in \mathbb{R}^d , where ε represents the intensity of noise.

The main result of the paper on the invariant measures of the stochastic LLB equation (1.7) is presented below.

Theorem 1.2. *If (1.1) holds, then:*

- (i). *For $d = 1, 2$, the stochastic equation (1.3) has at least one invariant measure in $H^1(\mathbb{R}^d, \mathbb{R}^3)$.*
- (ii). *For $d = 1$, the set $\bigcup_{\varepsilon \in [0, 1]} \mathcal{I}^\varepsilon$ is tight in $H^1(\mathbb{R}, \mathbb{R}^3)$ where \mathcal{I}^ε is the collection of all invariant measures of (1.7) corresponding to ε .*

Furthermore, for any $\varepsilon_n \rightarrow \varepsilon_0 \in [0, 1]$ and $\mu^{\varepsilon_n} \in \mathcal{I}^{\varepsilon_n}$, there exists a subsequence $\{\mu^{\varepsilon_{n_k}}\}_{k=1}^{\infty}$ of $\{\mu^{\varepsilon_n}\}_{n=1}^{\infty}$ and $\mu^{\varepsilon_0} \in \mathcal{I}^{\varepsilon_0}$ such that $\mu^{\varepsilon_{n_k}} \rightarrow \mu^{\varepsilon_0}$ weakly.

Note that Theorem 1.2 (ii) implies that any limit of a sequence of invariant measures of (1.7) must be an invariant measure of the limiting equation.

Based on the uniform tail-ends estimates of solutions in $L^2(\Omega, C([0, T], L^2(\mathbb{R}^d)))$, we will prove Theorem 1.2 (i) for the existence of invariant measures of (1.3) by the weakly Feller method as in [4, 31].

The most difficult part of the paper is to prove Theorem 1.2 (ii), and currently we only obtain that result for $d = 1$. For $d = 2$, Theorem 1.2 (ii) remains open.

Next, we explain why the proof of Theorem 1.2 (ii) is so challenging. In order to establish the uniform tightness of the set $\bigcup_{\varepsilon \in [0, 1]} \mathcal{I}^\varepsilon$ in $H^1(\mathbb{R}^d, \mathbb{R}^3)$, we must derive the uniform tail-ends estimates of solutions in $H^1(\mathbb{R}^d, \mathbb{R}^3)$ rather than in $L^2(\mathbb{R}^d, \mathbb{R}^3)$, because the uniform tail-ends estimates in $L^2(\mathbb{R}^d, \mathbb{R}^3)$ are not sufficient in this case. To obtain the uniform tail-ends estimates of solutions

in $H^1(\mathbb{R}^d, \mathbb{R}^3)$, we need to use the energy equation of solutions in $H^1(\mathbb{R}^d, \mathbb{R}^3)$. But, the energy equation of solutions in $H^1(\mathbb{R}^d, \mathbb{R}^3)$ is not available for $d = 2$. To understand this issue, let's rewrite (1.6) as

$$\mathbf{u}(t) = \mathbf{u}_0 + \int_0^t \mathbf{F}(\mathbf{u}(s)) ds + \sum_{k=1}^{\infty} \int_0^t (\mathbf{u}(s) \times \mathbf{f}_k + \mathbf{f}_k) dW_k, \quad (1.8)$$

where

$$\mathbf{F}(\mathbf{u}) = \Delta \mathbf{u} + \mathbf{u} \times \Delta \mathbf{u} - (1 + |\mathbf{u}|^2) \mathbf{u} + \frac{1}{2} \sum_{k=1}^{\infty} (\mathbf{u} \times \mathbf{f}_k) \times \mathbf{f}_k.$$

The high-order nonlinear term $\mathbf{u} \times \Delta \mathbf{u}$ has lowest regularity among all terms in $\mathbf{F}(\mathbf{u})$, and hence it dominates the regularity of $\mathbf{F}(\mathbf{u})$. If we assume

$$\mathbf{u} \times \Delta \mathbf{u} \in L^2(0, T; L^2(\mathbb{R}^d, \mathbb{R}^3)), \quad \text{P-almost surely,} \quad (1.9)$$

then by (1.4) we find that $\mathbf{F}(\mathbf{u}) \in L^2(0, T; L^2(\mathbb{R}^d, \mathbb{R}^3))$ P-almost surely, and hence by (1.4) and [30, Theorem 3.2] we infer that $\mathbf{u} \in C([0, T], H^1(\mathbb{R}^d, \mathbb{R}^3))$ P-almost surely, and \mathbf{u} satisfies the energy equation in $H^1(\mathbb{R}^d, \mathbb{R}^3)$:

$$\begin{aligned} \|\mathbf{u}(t)\|_{H^1(\mathbb{R}^d, \mathbb{R}^3)}^2 &= \|\mathbf{u}_0\|_{H^1(\mathbb{R}^d, \mathbb{R}^3)}^2 + 2 \int_0^t ((I - \Delta) \mathbf{u}(s), \mathbf{F}(\mathbf{u}(s)))_{L^2(\mathbb{R}^d, \mathbb{R}^3)} ds \\ &+ \sum_{k=1}^{\infty} \int_0^t \|\mathbf{u}(s) \times \mathbf{f}_k + \mathbf{f}_k\|_{H^1(\mathbb{R}^d, \mathbb{R}^3)}^2 ds + 2 \sum_{k=1}^{\infty} \int_0^t ((I - \Delta) \mathbf{u}(s), \mathbf{u}(s) \times \mathbf{f}_k + \mathbf{f}_k)_{L^2(\mathbb{R}^d, \mathbb{R}^3)} dW_k, \end{aligned} \quad (1.10)$$

for all $t \in [0, T]$, P-almost surely.

Unfortunately, by (1.5) we find that for $d = 2$, $\mathbf{u} \times \Delta \mathbf{u} \in L^r(0, T; L^2(\mathbb{R}^d, \mathbb{R}^3))$ only for $r < 2$ but not for $r = 2$, P-almost surely, and hence the condition (1.9) is not satisfied for $d = 2$. Consequently, we cannot obtain the energy equation (1.10) for \mathbf{u} in $H^1(\mathbb{R}^2, \mathbb{R}^3)$, and thus we are unable to establish the uniform tail-ends estimates of solutions in $H^1(\mathbb{R}^2, \mathbb{R}^3)$. We may formally derive the uniform tail-ends estimates in $H^1(\mathbb{R}^2, \mathbb{R}^3)$, but those calculations cannot be justified due to the lower regularity of $\mathbf{F}(\mathbf{u})$.

Nevertheless, if $d = 1$, then we have the embedding $H^1(\mathbb{R}, \mathbb{R}^3) \hookrightarrow L^\infty(\mathbb{R}, \mathbb{R}^3)$, and hence by (1.4) we infer that $\mathbf{u} \times \Delta \mathbf{u} \in L^2(0, T; L^2(\mathbb{R}, \mathbb{R}^3))$, P-almost surely; that is, $\mathbf{u} \times \Delta \mathbf{u}$ satisfies (1.9) for $d = 1$. As a result, we find that $\mathbf{u} \in C([0, T], H^1(\mathbb{R}, \mathbb{R}^3))$ P-almost surely, and \mathbf{u} satisfies the energy equation in $H^1(\mathbb{R}, \mathbb{R}^3)$. This is why we are able to obtain the uniform tail-ends estimates of solutions in $H^1(\mathbb{R}, \mathbb{R}^3)$, but not in $H^1(\mathbb{R}^2, \mathbb{R}^3)$ (see Lemma 4.5).

On the other hand, in order to prove a limit of a sequence of invariant measures of (1.7) is an invariant measure of the limiting equation, we must establish the convergence of solutions in probability in $H^1(\mathbb{R}^d, \mathbb{R}^3)$. Again, since the solutions have no sufficient regularity for $d = 2$, we are unable to obtain the convergence of solutions in probability in $H^1(\mathbb{R}^2, \mathbb{R}^3)$. However, if $d = 1$, by the embedding $H^1(\mathbb{R}, \mathbb{R}^3) \hookrightarrow L^\infty(\mathbb{R}, \mathbb{R}^3)$, the convergence of solutions in probability in $H^1(\mathbb{R}, \mathbb{R}^3)$ can be established (see Lemma 4.11).

We remark that by the idea of uniform tail-ends estimates, we can also prove the existence of invariant measures of (1.2) for $d = 1$ by the Feller property of the Markov semigroup with respect to the strong topology instead of the weak topology (see Lemma 4.7), which provides an alternative approach to show the existence of invariant measures for $d = 1$.

The rest of this paper is organized as follows. In section 2, we establish the existence and uniqueness of global solutions by the domain expansion method. In section 3, we prove the existence of invariant measures by the weakly Feller method. In the last section, we prove Theorem 1.2 and the existence of invariant measures for $d = 1$ by the Feller property of solutions with respect to the strong topology.

For later sections, we recall the following Skorokhod-Jakubowski representation theorem from [2, 22].

Proposition 1.1. *Suppose X is a topological space, and there exists a sequence of continuous functions $g_n : X \rightarrow \mathbb{R}$ that separates points of X . If $\{\mu_n\}_{n=1}^\infty$ is a tight sequence of probability measures on $(X, \mathcal{B}(X))$, then there exists a subsequence $\{\mu_{n_k}\}_{k=1}^\infty$ of $\{\mu_n\}_{n=1}^\infty$, a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$, X -valued random variables v_k and v such that the law of v_k is μ_{n_k} for all $k \in \mathbb{N}$ and $v_k \rightarrow v$ $\tilde{\mathbb{P}}$ -almost surely as $k \rightarrow \infty$.*

2. EXISTENCE AND UNIQUENESS OF SOLUTIONS

In this section, we prove the global well-posedness of strong solutions of (1.3) defined in \mathbb{R}^d with $d = 1, 2$. Due to the non-compactness of the standard Sobolev embeddings in unbounded domains, we adopt a domain expansion argument to show the existence of solutions of (1.3). Specifically, we first consider the approximate solutions of equation (1.3) defined in a bounded domain $\mathcal{D}_n = \{x \in \mathbb{R}^d : |x| < n\}$ for every $n \in \mathbb{N}$, then take the limit of these approximate solutions as $n \rightarrow \infty$ to obtain a solution of (1.3) defined in the entire space \mathbb{R}^d .

2.1. Construction of approximate solutions. In this subsection, we will define a sequence of approximate solutions for (1.3). We first introduce the Sobolev spaces that will be used in the sequel. Let \mathcal{D} be an open set in \mathbb{R}^d with $d = 1, 2$. Given $p \geq 1$, let

$$L^p(\mathcal{D}, \mathbb{R}^3) = \left\{ \mathbf{u} : \mathcal{D} \rightarrow \mathbb{R}^3 : \int_{\mathcal{D}} |\mathbf{u}(x)|^p dx < \infty \right\}.$$

The norm of $L^p(\mathcal{D}, \mathbb{R}^3)$ is denoted by $\|\cdot\|_{L^p(\mathcal{D}, \mathbb{R}^3)}$. In particular, for $p = 2$, the inner product and the norm of $L^2(\mathcal{D}, \mathbb{R}^3)$ are written as $(\cdot, \cdot)_{L^2(\mathcal{D}, \mathbb{R}^3)}$ and $\|\cdot\|_{L^2(\mathcal{D}, \mathbb{R}^3)}$, respectively. For every $k \in \mathbb{N}$ and $p \geq 1$, let

$$W^{k,p}(\mathcal{D}, \mathbb{R}^3) = \left\{ \mathbf{u} \in L^p(\mathcal{D}, \mathbb{R}^3) : \partial^\alpha \mathbf{u} \in L^p(\mathcal{D}, \mathbb{R}^3), |\alpha| \leq k \right\}.$$

The norm of $W^{k,p}(\mathcal{D}, \mathbb{R}^3)$ is denoted by $\|\cdot\|_{W^{k,p}(\mathcal{D}, \mathbb{R}^3)}$. Let $W_0^{k,p}(\mathcal{D}, \mathbb{R}^3)$ be the closure of $C_0^\infty(\mathcal{D}, \mathbb{R}^3)$ with respect to the topology of $W^{k,p}(\mathcal{D}, \mathbb{R}^3)$. For $p = 2$, we write $H^k(\mathcal{D}, \mathbb{R}^3) = W^{k,2}(\mathcal{D}, \mathbb{R}^3)$ which

is a Hilbert space with inner product $(\cdot, \cdot)_{H^k(\mathcal{D}, \mathbb{R}^3)} = \sum_{|\alpha| \leq k} (\partial^\alpha \cdot, \partial^\alpha \cdot)_{L^2(\mathcal{D}, \mathbb{R}^3)}$. Similarly, we also denote $W_0^{k,2}(\mathcal{D}, \mathbb{R}^3)$ by $H_0^k(\mathcal{D}, \mathbb{R}^3)$. If no confusion occurs, we will use $L^p(\mathcal{D})$ for $L^p(\mathcal{D}, \mathbb{R}^3)$, and use similar notations for other spaces.

Let Y be a separable Hilbert space with inner product $(\cdot, \cdot)_Y$. Denote by Y_w the space Y endowed with weak topology. Given $T > 0$ let

$$C([0, T], Y_w) = \{\mathbf{u} : \mathbf{u} \text{ is a continuous function from } [0, T] \text{ to } Y_w\}.$$

which is equipped with the following topology: $\mathbf{u}_n \rightarrow \mathbf{u}$ in $C([0, T], Y_w)$ if for every $y \in Y$, $\lim_{n \rightarrow \infty} \sup_{t \in [0, T]} |(\mathbf{u}_n(t) - \mathbf{u}(t), y)_Y| = 0$. Given $R > 0$, let $C([0, T], Y_w^R)$ be a subset of $C([0, T], Y_w)$ given by

$$C([0, T], Y_w^R) = \left\{ \mathbf{u} \in C([0, T], Y_w) : \sup_{t \in [0, T]} \|\mathbf{u}(t)\|_Y \leq R \right\}.$$

Since the weak topology of Y on the closed ball of radius R centered at the origin is metrizable, we see that $C([0, T], Y_w^R)$ is a complete metric space.

To construct approximate solutions for (1.3), we choose a smooth function $\theta : \mathbb{R}^d \rightarrow \mathbb{R}$ such that $0 \leq \theta(x) \leq 1$ for all $x \in \mathbb{R}^d$ and

$$\theta(x) = \begin{cases} 0, & \text{if } |x| \geq \frac{3}{4}, \\ 1, & \text{if } |x| \leq \frac{1}{2}. \end{cases} \quad (2.1)$$

Given $n \in \mathbb{N}$, let $\theta_n(x) = \theta\left(\frac{x}{n}\right)$ for all $x \in \mathbb{R}^d$.

For every $n \in \mathbb{N}$, consider the solution \mathbf{u}^n of the equation defined in $\mathcal{D}_n = \{x \in \mathbb{R}^d : |x| < n\}$:

$$\begin{cases} d\mathbf{u}^n = \Delta \mathbf{u}^n dt + \mathbf{u}^n \times \Delta \mathbf{u}^n dt - (1 + |\mathbf{u}^n|^2) \mathbf{u}^n dt + \sum_{k=1}^{\infty} (\mathbf{u}^n \times \mathbf{f}_k + \mathbf{f}_k) \circ dW_k, \\ \mathbf{u}^n|_{\partial \mathcal{D}_n} = 0, \\ \mathbf{u}^n(0, x) = \theta_n(x) \mathbf{u}_0(x), \quad x \in \mathbb{R}^d, \end{cases} \quad (2.2)$$

where $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$.

By [4], we know that if (1.1) holds, then for every $n \in \mathbb{N}$ and $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$, problem (2.2) has a unique solution \mathbf{u}^n which is a weakly continuous $H^1(\mathcal{D}_n)$ -valued progressively measurable stochastic process such that for every $T > 0$ and $p \geq 2$,

$$\mathbf{u}^n \in L^p(\Omega; L^\infty(0, T; H_0^1(\mathcal{D}_n)) \cap L^2(0, T; H^2(\mathcal{D}_n))).$$

The uniform estimates of the sequence $\{\mathbf{u}^n\}_{n=1}^\infty$ are collected in the following lemma which can be found in [4].

Lemma 2.1. *Let (1.1) hold and $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$. Then the sequence $\{\mathbf{u}^n\}_{n=1}^\infty$ of solutions of (2.2) has the properties:*

(i). For every $p \geq 1$, $R > 0$ and $T > 0$, there exists a positive number $C_1 = C_1(p, R, T)$ independent of $n \in \mathbb{N}$ such that for all $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$ with $\|\mathbf{u}_0\|_{H^1(\mathbb{R}^d)} \leq R$,

$$\mathbb{E} \left(\sup_{s \in [0, T]} \|\mathbf{u}^n(s)\|_{H_0^1(\mathcal{D}_n)}^{2p} \right) + \mathbb{E} \left(\int_0^T \|\mathbf{u}^n(s)\|_{H^2(\mathcal{D}_n)}^2 ds \right)^p \leq C_1, \quad (2.3)$$

and

$$\mathbb{E} \left(\int_0^T \|\mathbf{u}^n(s) \times \Delta \mathbf{u}^n(s)\|_{L^2(\mathcal{D}_n)}^{\frac{4}{3}} ds \right)^p \leq C_1. \quad (2.4)$$

(ii). For every $R > 0$ and $T > 0$, there exists a positive number $C_2 = C_2(R, T)$ independent of $n \in \mathbb{N}$ such that for all $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$ with $\|\mathbf{u}_0\|_{H^1(\mathbb{R}^d)} \leq R$,

$$\mathbb{E} \left(\|\mathbf{u}^n\|_{W^{\frac{3}{16}, 16}(0, T; L^2(\mathcal{D}_n))}^{16} \right) \leq C_2. \quad (2.5)$$

(iii). For every $R > 0$, there exists a positive number $C_3 = C_3(R)$ independent of $n \in \mathbb{N}$ such that for all $t \geq 0$ and $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$ with $\|\mathbf{u}_0\|_{H^1(\mathbb{R}^d)} \leq R$,

$$\mathbb{E} \left(\|\mathbf{u}^n(t)\|_{H_0^1(\mathcal{D}_n)}^2 \right) + \int_0^t \mathbb{E} \left(\|\mathbf{u}^n(s)\|_{H^2(\mathcal{D}_n)}^2 \right) ds \leq C_3 + C_3 t. \quad (2.6)$$

Next, we derive the uniform estimates on the tails of the sequence $\{\mathbf{u}^n\}_{n=1}^\infty$, which will be used to overcome the non-compactness of Sobolev embeddings on unbounded domains.

Lemma 2.2. *If (1.1) holds, then for every $\varepsilon > 0$, $T > 0$ and $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$, there exists $M = M(\varepsilon, T, \mathbf{u}_0) \geq 1$ such that for all $m \geq M$ and $n \in \mathbb{N}$,*

$$\mathbb{E} \left(\sup_{t \in [0, T]} \int_{m < |x| < n} |\mathbf{u}^n(t, \mathbf{u}_0)(x)|^2 dx \right) < \varepsilon.$$

Proof. Let $\phi(x) = 1 - \theta(x)$ for all $x \in \mathbb{R}^d$, where θ is the smooth function given by (2.1). Given $m \in \mathbb{N}$, denote by $\phi_m(x) = \phi(\frac{x}{m})$. For all $n, m \in \mathbb{N}$, by Itô's formula, we get for all $r, t \in [0, T]$ with $r \leq t$, P-almost surely,

$$\begin{aligned} & \|\phi_m \mathbf{u}^n(r)\|_{L^2(\mathcal{D}_n)}^2 - \|\phi_m \mathbf{u}^n(0)\|_{L^2(\mathcal{D}_n)}^2 \\ &= 2 \int_0^r (\Delta \mathbf{u}^n(s), \phi_m^2 \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)} ds + 2 \int_0^r (\mathbf{u}^n(s) \times \Delta \mathbf{u}^n(s), \phi_m^2 \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)} ds \\ & \quad - 2 \int_0^r ((1 + |\mathbf{u}^n(s)|^2) \mathbf{u}^n(s), \phi_m^2 \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)} ds + 2 \sum_{k \geq 1} \int_0^r (\phi_m \mathbf{f}_k, \phi_m \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)} dW_k \\ & \quad + \sum_{k \geq 1} \int_0^r \|\phi_m (\mathbf{u}^n(s) \times \mathbf{f}_k + \mathbf{f}_k)\|_{L^2(\mathcal{D}_n)}^2 ds + \sum_{k \geq 1} \int_0^r ((\mathbf{u}^n(s) \times \mathbf{f}_k) \times \mathbf{f}_k, \phi_m^2 \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)} ds, \end{aligned} \quad (2.7)$$

where we have used the identity $(a \times b, a) = 0$ to cancel the multiplicative noise.

For the first term on the right-hand side of (2.7) we have for all $r \leq t$,

$$\begin{aligned}
& 2 \int_0^r (\Delta \mathbf{u}^n(s), \phi_m^2 \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)} ds = -2 \int_0^r (\nabla \mathbf{u}^n(s), \nabla(\phi_m^2 \mathbf{u}^n(s)))_{L^2(\mathcal{D}_n)} ds \\
& = -2 \int_0^r (\nabla \mathbf{u}^n(s), \phi_m^2 \nabla \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)} ds - \frac{4}{m} \int_0^r (\nabla \mathbf{u}^n(s), \phi_m \nabla \phi(x/m) \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)} ds \\
& \leq \frac{c_1}{m} \int_0^t \|\mathbf{u}^n(s)\|_{H_0^1(\mathcal{D}_n)}^2 ds,
\end{aligned} \tag{2.8}$$

where $c_1 > 0$ is a constant independent of both n and m . For the second and third terms on the right-hand side of (2.7) we have

$$2(\mathbf{u}^n(s) \times \Delta \mathbf{u}^n(s), \phi_m^2 \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)} = 2(\mathbf{u}^n(s) \times (\phi_m^2 \Delta \mathbf{u}^n(s)), \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)} = 0, \tag{2.9}$$

and

$$-2((1 + |\mathbf{u}^n(s)|^2)\mathbf{u}^n(s), \phi_m^2 \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)} = -2\|\phi_m \mathbf{u}^n(s)\|_{L^2(\mathcal{D}_n)}^2 - 2(|\mathbf{u}^n(s)|^2, \phi_m^2 |\mathbf{u}^n(s)|^2)_{L^2(\mathcal{D}_n)} \leq 0. \tag{2.10}$$

For the stochastic term on the right-hand side of (2.7), by the BDG inequality we get

$$\begin{aligned}
2\mathbb{E} \left(\sup_{0 \leq r \leq t} \left| \int_0^r \sum_{k \geq 1} (\phi_m \mathbf{f}_k, \phi_m \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)} dW_k \right| \right) & \leq 2c_2 \mathbb{E} \left(\int_0^t \sum_{k \geq 1} |(\phi_m \mathbf{f}_k, \phi_m \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)}|^2 ds \right)^{\frac{1}{2}} \\
& \leq c_2 \sum_{k \geq 1} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R}^d)}^2 + c_2 \int_0^t \mathbb{E} \left(\|\phi_m \mathbf{u}^n(s)\|_{L^2(\mathcal{D}_n)}^2 \right) ds.
\end{aligned} \tag{2.11}$$

For the last two terms on the right-hand side of (2.7) we have for all $0 \leq r \leq t$,

$$\begin{aligned}
& \sum_{k \geq 1} \int_0^r \|\phi_m(\mathbf{u}^n(s) \times \mathbf{f}_k + \mathbf{f}_k)\|_{L^2(\mathcal{D}_n)}^2 ds + \sum_{k \geq 1} \int_0^r ((\mathbf{u}^n(s) \times \mathbf{f}_k) \times \mathbf{f}_k, \phi_m^2 \mathbf{u}^n(s))_{L^2(\mathcal{D}_n)} ds \\
& \leq 2 \sum_{k \geq 1} \int_0^t \|(\phi_m \mathbf{u}^n(s)) \times \mathbf{f}_k\|_{L^2(\mathcal{D}_n)}^2 ds + 2t \sum_{k \geq 1} \|\phi_m \mathbf{f}_k\|_{L^2(\mathcal{D}_n)}^2 \\
& \quad + \sum_{k \geq 1} \int_0^t \|((\phi_m \mathbf{u}^n(s)) \times \mathbf{f}_k) \times \mathbf{f}_k\|_{L^2(\mathcal{D}_n)} \|\phi_m \mathbf{u}^n(s)\|_{L^2(\mathcal{D}_n)} ds \\
& \leq 3 \sum_{k \geq 1} \|\mathbf{f}_k\|_{L^\infty(\mathbb{R}^d)}^2 \int_0^t \|\phi_m \mathbf{u}^n(s)\|_{L^2(\mathcal{D}_n)}^2 ds + 2t \sum_{k \geq 1} \|\phi_m \mathbf{f}_k\|_{L^2(\mathcal{D}_n)}^2.
\end{aligned} \tag{2.12}$$

It follows from (2.7)-(2.12) that for all $t \in [0, T]$ and $n, m \in \mathbb{N}$,

$$\begin{aligned}
& \mathbb{E} \left(\sup_{0 \leq r \leq t} \|\phi_m \mathbf{u}^n(r)\|_{L^2(\mathcal{D}_n)}^2 \right) \leq \mathbb{E} \left(\|\phi_m \mathbf{u}^n(0)\|_{L^2(\mathcal{D}_n)}^2 \right) \\
& \quad + \left(c_2 + 3 \sum_{k \geq 1} \|\mathbf{f}_k\|_{L^\infty(\mathbb{R}^d)}^2 \right) \int_0^t \mathbb{E} \left(\|\phi_m \mathbf{u}^n(s)\|_{L^2(\mathcal{D}_n)}^2 \right) ds
\end{aligned}$$

$$+(2T + c_2) \sum_{k \geq 1} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R}^d)}^2 + \frac{c_1}{m} \int_0^t \mathbb{E} \left(\|\mathbf{u}^n(s)\|_{H_0^1(\mathcal{D}_n)}^2 \right) ds. \quad (2.13)$$

By (2.3) and (2.13) we find that there exists $c_2 = c_2(T, \mathbf{u}_0) > 0$ such that for all $t \in [0, T]$ and $n, m \in \mathbb{N}$,

$$\begin{aligned} & \mathbb{E} \left(\sup_{0 \leq r \leq t} \|\phi_m \mathbf{u}^n(r)\|_{L^2(\mathcal{D}_n)}^2 \right) \leq \mathbb{E} \left(\|\phi_m \mathbf{u}^n(0)\|_{L^2(\mathcal{D}_n)}^2 \right) \\ & + \left(c_2 + 3 \sum_{k \geq 1} \|\mathbf{f}_k\|_{L^\infty(\mathbb{R}^d)}^2 \right) \int_0^t \mathbb{E} \left(\sup_{0 \leq r \leq s} \|\phi_m \mathbf{u}^n(r)\|_{L^2(\mathcal{D}_n)}^2 \right) ds \\ & + (2T + c_2) \sum_{k \geq 1} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R}^d)}^2 + \frac{c_1 c_2}{m}. \end{aligned} \quad (2.14)$$

By (2.14) and Gronwall's lemma we obtain that for all $t \in [0, T]$ and $n, m \in \mathbb{N}$,

$$\mathbb{E} \left(\sup_{0 \leq r \leq t} \|\phi_m \mathbf{u}^n(r)\|_{L^2(\mathcal{D}_n)}^2 \right) \leq \left(\|\phi_m \mathbf{u}_0\|_{L^2(\mathbb{R}^d)}^2 + (2T + c_2) \sum_{k \geq 1} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R}^d)}^2 + \frac{c_1 c_2}{m} \right) e^{c_3 T}, \quad (2.15)$$

where $c_3 = c_2 + 3 \sum_{k \geq 1} \|\mathbf{f}_k\|_{L^\infty(\mathbb{R}^d)}^2$. Note that as $m \rightarrow \infty$,

$$\begin{aligned} & \|\phi_m \mathbf{u}_0\|_{L^2(\mathbb{R}^d)}^2 + (2T + c_2) \sum_{k \geq 1} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R}^d)}^2 \\ & \leq \int_{|x| > \frac{1}{2}m} |\mathbf{u}_0(x)|^2 dx + (2T + c_2) \int_{|x| > \frac{1}{2}m} \sum_{k \geq 1} |\mathbf{f}_k(x)|^2 dx \rightarrow 0, \end{aligned}$$

which along with (2.15) shows that for every $\varepsilon > 0$, there exists $M = M(\varepsilon, T, \mathbf{u}_0) \geq 1$, such that for all $m \geq M$, $n \in \mathbb{N}$ and $t \in [0, T]$,

$$\mathbb{E} \left(\sup_{0 \leq r \leq t} \int_{\frac{3}{4}m < |x| < n} |\mathbf{u}^n(r, \mathbf{u}_0)(x)|^2 dx \right) \leq \mathbb{E} \left(\sup_{0 \leq r \leq t} \|\phi_m \mathbf{u}^n(r, \mathbf{u}_0)\|_{L^2(\mathcal{D}_n)}^2 \right) < \varepsilon,$$

as desired. \square

Let $\mathbf{v}^n(t, x) = \theta_n(x) \mathbf{u}^n(t, x)$ for all $n \in \mathbb{N}$, $t \geq 0$ and $x \in \mathcal{D}_n$. We then extend $\mathbf{v}^n(t, x)$ for $x \in \mathbb{R}^d \setminus \mathcal{D}_n$ by zero extension, and thus consider $\mathbf{v}^n(t, x)$ as a function defined for all $t \geq 0$ and $x \in \mathbb{R}^d$. By (2.1) we find that the sequence $\{\mathbf{v}^n\}_{n=1}^\infty$ enjoys the uniform estimates as established by Lemmas 2.1 and 2.2 which are presented below.

Lemma 2.3. *Let (1.1) hold and $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$. Then the sequence $\{\mathbf{v}^n\}_{n=1}^\infty$ has the properties:*

(i). *For every $p \geq 1$, $R > 0$ and $T > 0$, there exists a positive number $C_1 = C_1(p, R, T)$ independent of $n \in \mathbb{N}$ such that for all $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$ with $\|\mathbf{u}_0\|_{H^1(\mathbb{R}^d)} \leq R$,*

$$\mathbb{E} \left(\sup_{s \in [0, T]} \|\mathbf{v}^n(s, \mathbf{u}_0)\|_{H^1(\mathbb{R}^d)}^{2p} \right) + \mathbb{E} \left(\int_0^T \|\mathbf{v}^n(s, \mathbf{u}_0)\|_{H^2(\mathbb{R}^d)}^2 ds \right)^p \leq C_1, \quad (2.16)$$

and

$$\mathbb{E} \left(\int_0^T \|\mathbf{v}^n(s, \mathbf{u}_0) \times \Delta \mathbf{v}^n(s, \mathbf{u}_0)\|_{L^2(\mathbb{R}^d)}^{\frac{4}{3}} ds \right)^p \leq C_1. \quad (2.17)$$

(ii). For every $R > 0$ and $T > 0$, there exists a positive number $C_2 = C_2(R, T)$ independent of $n \in \mathbb{N}$ such that for all $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$ with $\|\mathbf{u}_0\|_{H^1(\mathbb{R}^d)} \leq R$,

$$\mathbb{E} \left(\|\mathbf{v}^n(\cdot, \mathbf{u}_0)\|_{W^{\frac{3}{16}, 16}(0, T; L^2(\mathbb{R}^d))}^{16} \right) \leq C_2. \quad (2.18)$$

(iii). For every $R > 0$, there exists a positive number $C_3 = C_3(R)$ independent of $n \in \mathbb{N}$ such that for all $t \geq 0$ and $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$ with $\|\mathbf{u}_0\|_{H^1(\mathbb{R}^d)} \leq R$,

$$\mathbb{E} \left(\|\mathbf{v}^n(t, \mathbf{u}_0)\|_{H^1(\mathbb{R}^d)}^2 \right) + \int_0^t \mathbb{E} \left(\|\mathbf{v}^n(s, \mathbf{u}_0)\|_{H^2(\mathbb{R}^d)}^2 \right) ds \leq C_3 + C_3 t. \quad (2.19)$$

(iv). For every $\varepsilon > 0$, $T > 0$ and $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$, there exists $M = M(\varepsilon, T, \mathbf{u}_0) \geq 1$ such that for all $m \geq M$ and $n \in \mathbb{N}$,

$$\mathbb{E} \left(\sup_{t \in [0, T]} \int_{|x| > m} |\mathbf{v}^n(t, \mathbf{u}_0)(x)|^2 dx \right) < \varepsilon. \quad (2.20)$$

2.2. Tightness of laws of approximate solutions. In this subsection, we establish the tightness of the sequence of distributions of $\{\mathbf{v}^n(\cdot, \mathbf{u}_0)\}_{n=1}^\infty$, which is given below.

Lemma 2.4. *If (1.1) holds, then the sequence of distributions of $\{\mathbf{v}^n(\cdot, \mathbf{u}_0)\}_{n=1}^\infty$ is tight in the space $C([0, T]; L^2(\mathbb{R}^d)) \cap C([0, T]; H^1_w(\mathbb{R}^d)) \cap L^2_w(0, T; H^2(\mathbb{R}^d))$, where $L^2_w(0, T; H^2(\mathbb{R}^d))$ is the space $L^2(0, T; H^2(\mathbb{R}^d))$ endowed with the weak topology.*

Proof. By (2.16), (2.18) and Chebyshev's inequality, we deduce that for every $T > 0$ and $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$,

$$\begin{aligned} & \mathbb{P} \left(\sup_{t \in [0, T]} \|\mathbf{v}^n(t, \mathbf{u}_0)\|_{H^1(\mathbb{R}^d)}^2 + \int_0^T \|\mathbf{v}^n(t, \mathbf{u}_0)\|_{H^2(\mathbb{R}^d)}^2 dt + \|\mathbf{v}^n(t, \mathbf{u}_0)\|_{W^{\frac{3}{16}, 16}(0, T; L^2(\mathbb{R}^d))}^2 > R \right) \\ & \leq \frac{1}{R} \mathbb{E} \left(\sup_{t \in [0, T]} \|\mathbf{v}^n(t, \mathbf{u}_0)\|_{H^1(\mathbb{R}^d)}^2 + \int_0^T \|\mathbf{v}^n(t, \mathbf{u}_0)\|_{H^2(\mathbb{R}^d)}^2 dt + \|\mathbf{v}^n(t, \mathbf{u}_0)\|_{W^{\frac{3}{16}, 16}(0, T; L^2(\mathbb{R}^d))}^2 \right) \\ & \leq \frac{c_1}{R}, \end{aligned}$$

where $c_1 = c_1(T, \mathbf{u}_0) > 0$. Therefore, for every $\varepsilon > 0$, there exists $R(\varepsilon) > 0$ such that

$$\mathbb{P} \left(\sup_{t \in [0, T]} \|\mathbf{v}^n(t, \mathbf{u}_0)\|_{H^1(\mathbb{R}^d)}^2 + \int_0^T \|\mathbf{v}^n(t, \mathbf{u}_0)\|_{H^2(\mathbb{R}^d)}^2 dt + \|\mathbf{v}^n(t, \mathbf{u}_0)\|_{W^{\frac{3}{16}, 16}(0, T; L^2(\mathbb{R}^d))}^2 > R(\varepsilon) \right) < \frac{1}{2} \varepsilon. \quad (2.21)$$

On the other hand, by (2.20), for every $\varepsilon > 0$, $T > 0$ and $m \in \mathbb{N}$, we find that there exists $N_m \in \mathbb{N}$ depending only on m , T and ε such that for all $n \in \mathbb{N}$,

$$\mathbb{E} \left(\sup_{t \in [0, T]} \int_{|x| > \frac{1}{2} N_m} |\mathbf{v}^n(t, \mathbf{u}_0)(x)|^2 dx \right) \leq \frac{\varepsilon}{2^{2m+1}}. \quad (2.22)$$

By (2.22) and Chebyshev's inequality we obtain that for all $n \in \mathbb{N}$,

$$\mathbb{P} \left(\sup_{t \in [0, T]} \int_{|x| > \frac{1}{2} N_m} |\mathbf{v}^n(t, \mathbf{u}_0)(x)|^2 dx > \frac{1}{2^m} \right) \leq \frac{\varepsilon}{2^{m+1}},$$

and hence for all $n \in \mathbb{N}$,

$$\mathbb{P} \left(\bigcup_{m=1}^{\infty} \left\{ \sup_{t \in [0, T]} \int_{|x| > \frac{1}{2} N_m} |\mathbf{v}^n(t, \mathbf{u}_0)(x)|^2 dx > \frac{1}{2^m} \right\} \right) \leq \sum_{m=1}^{\infty} \frac{\varepsilon}{2^{m+1}} \leq \frac{\varepsilon}{2}. \quad (2.23)$$

Recall that $\phi(x) = 1 - \theta(x)$ and $\phi_m(x) = \phi\left(\frac{x}{m}\right)$ for all $x \in \mathbb{R}^d$. By (2.23) we get that for all $n \in \mathbb{N}$,

$$\mathbb{P} \left(\left\{ \sup_{t \in [0, T]} \|\phi_{N_m} \mathbf{v}^n(t, \mathbf{u}_0)\|_{L^2(\mathbb{R}^d)}^2 \leq \frac{1}{2^m}, \text{ for all } m \in \mathbb{N} \right\} \right) \geq 1 - \frac{\varepsilon}{2}. \quad (2.24)$$

Define the sets

$$\mathcal{S}_1^\varepsilon = \left\{ \mathbf{v} : \sup_{t \in [0, T]} \|\mathbf{v}(t)\|_{H^1(\mathbb{R}^d)}^2 + \int_0^T \|\mathbf{v}(t)\|_{H^2(\mathbb{R}^d)}^2 dt + \|\mathbf{v}\|_{W^{\frac{3}{16}, 16}(0, T; L^2(\mathbb{R}^d))}^2 \leq R(\varepsilon) \right\}, \quad (2.25)$$

$$\mathcal{S}_2^\varepsilon = \left\{ \mathbf{v} : \sup_{t \in [0, T]} \|\phi_{N_m} \mathbf{v}(t)\|_{L^2(\mathbb{R}^d)}^2 \leq \frac{1}{2^m}, \text{ for all } m \in \mathbb{N} \right\}, \quad (2.26)$$

and

$$\mathcal{S}^\varepsilon = \mathcal{S}_1^\varepsilon \cap \mathcal{S}_2^\varepsilon.$$

It follows from (2.21), (2.24)-(2.26) that for all $n \in \mathbb{N}$,

$$\mathbb{P}(\mathbf{v}^n(t, \mathbf{u}_0) \in \mathcal{S}^\varepsilon) > 1 - \varepsilon.$$

It remains to verify that the set \mathcal{S}^ε is compact in

$$C([0, T]; L^2(\mathbb{R}^d)) \cap C([0, T]; H_w^1(\mathbb{R}^d)) \cap L_w^2(0, T; H^2(\mathbb{R}^d)).$$

Note that every topology of $C([0, T]; L^2(\mathbb{R}^d))$, $C([0, T]; H_w^1(\mathbb{R}^d))$ and $L_w^2(0, T; H^2(\mathbb{R}^d))$ is metrizable on \mathcal{S}^ε . The precompactness of \mathcal{S}^ε in $L_w^2(0, T; H^2(\mathbb{R}^d))$ follows directly from its boundedness in $L^2(0, T; H^2(\mathbb{R}^d))$.

We now show that the set \mathcal{S}^ε is precompact in $C([0, T]; L^2(\mathbb{R}^d))$ by the Arzela-Ascoli theorem for which we need to check that \mathcal{S}^ε is equicontinuous in $C([0, T]; L^2(\mathbb{R}^d))$ and for every $t \in [0, T]$, the set $\{\mathbf{v}(t) : \mathbf{v} \in \mathcal{S}^\varepsilon\}$ is precompact in $L^2(\mathbb{R}^d)$. The equicontinuity of \mathcal{S}^ε in $C([0, T]; L^2(\mathbb{R}^d))$ follows from the embedding $W^{\frac{3}{16}, 16}(0, T; L^2(\mathbb{R}^d)) \hookrightarrow C([0, T]; L^2(\mathbb{R}^d))$.

On the other hand, by (2.26) we see that for every $\eta > 0$, there exists $m_0 \in \mathbb{N}$ such that for all $\mathbf{v} \in \mathcal{S}^\varepsilon$,

$$\sup_{t \in [0, T]} \|\phi_{N_{m_0}} \mathbf{v}(t)\|_{L^2(\mathbb{R}^d)}^2 < \frac{\eta^2}{16}. \quad (2.27)$$

Note that for every $t \in [0, T]$, the set $\{\mathbf{v}(t) : \mathbf{v} \in \mathcal{S}^\varepsilon\}$ is bounded in $H^1(\mathbb{R}^d)$. By the compactness of the embedding $H^1(\mathcal{D}_{m_0}) \hookrightarrow L^2(\mathcal{D}_{m_0})$, we see that the set $\{(1 - \phi_{N_{m_0}})\mathbf{v}(t) : \mathbf{v} \in \mathcal{S}^\varepsilon\}$ is precompact in $L^2(\mathcal{D}_{m_0})$, and hence it has a finite open cover of radius $\frac{1}{4}\eta$ in $L^2(\mathcal{D}_{m_0})$, which along with (2.27) implies that the set $\{\mathbf{v}(t) : \mathbf{v} \in \mathcal{S}^\varepsilon\}$ has a finite open cover of radius η in $L^2(\mathbb{R}^d)$. Due to the arbitrariness of $\eta > 0$, we infer that the set $\{\mathbf{v}(t) : \mathbf{v} \in \mathcal{S}^\varepsilon\}$ is precompact in $L^2(\mathbb{R}^d)$, and hence the set \mathcal{S}^ε is precompact in $C([0, T]; L^2(\mathbb{R}^d))$ by the Arzela-Ascoli theorem.

We finally prove the set \mathcal{S}^ε is precompact in $C([0, T]; H_w^1(\mathbb{R}^d))$. Let $\{\mathbf{z}_n\}_{n=1}^\infty$ be a sequence in \mathcal{S}^ε . We will prove $\{\mathbf{z}_n\}_{n=1}^\infty$ has a convergent subsequence in $C([0, T]; H_w^1(\mathbb{R}^d))$. Since \mathcal{S}^ε is precompact in $C([0, T]; L^2(\mathbb{R}^d))$, we know that there exists a subsequence $\{\mathbf{z}_{n_k}\}_{k=1}^\infty$ of $\{\mathbf{z}_n\}_{n=1}^\infty$ and $\mathbf{z} \in C([0, T]; L^2(\mathbb{R}^d))$ such that $\mathbf{z}_{n_k} \rightarrow \mathbf{z}$ in $C([0, T]; L^2(\mathbb{R}^d))$. By the construction of \mathcal{S}^ε , we infer that $\mathbf{z} \in \mathcal{S}^\varepsilon$, and for every $\psi \in H^2(\mathbb{R}^d)$,

$$\begin{aligned} \sup_{t \in [0, T]} \left| (\mathbf{z}_{n_k}(t) - \mathbf{z}(t), \psi)_{H^1(\mathbb{R}^d)} \right| &= \sup_{t \in [0, T]} \left| (\mathbf{z}_{n_k}(t) - \mathbf{z}(t), (I - \Delta)\psi)_{L^2(\mathbb{R}^d)} \right| \\ &\leq \sup_{t \in [0, T]} \|\mathbf{z}_{n_k}(t) - \mathbf{z}(t)\|_{L^2(\mathbb{R}^d)} \|(I - \Delta)\psi\|_{L^2(\mathbb{R}^d)} \rightarrow 0. \end{aligned}$$

By the density of $H^2(\mathbb{R}^d)$ in $H^1(\mathbb{R}^d)$ and the uniform boundedness of \mathcal{S}^ε in $H^1(\mathbb{R}^d)$, we find that for every $\psi \in H^1(\mathbb{R}^d)$,

$$\sup_{t \in [0, T]} \left| (\mathbf{z}_{n_k}(t) - \mathbf{z}(t), \psi)_{H^1(\mathbb{R}^d)} \right| \rightarrow 0,$$

and thus $\mathbf{z}_{n_k} \rightarrow \mathbf{z}$ in $C([0, T]; H_w^1(\mathbb{R}^d))$, as desired. \square

By Lemma 2.4 and the Skorokhod-Jakubowski representation theorem, we obtain the following result.

Lemma 2.5. *If (1.1) holds, then there exist a stochastic basis $(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}_{t \geq 0}, \tilde{\mathbb{P}})$, random variables $(\tilde{\mathbf{v}}, \tilde{W})$ and $(\tilde{\mathbf{v}}^n, \tilde{W}^n)$, $n \in \mathbb{N}$, defined on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}_{t \geq 0}, \tilde{\mathbb{P}})$ such that in the space $(C([0, T]; L^2(\mathbb{R}^d)) \cap C([0, T]; H_w^1(\mathbb{R}^d)) \cap L_w^2(0, T; H^2(\mathbb{R}^d))) \times C([0, T]; \mathbb{R}^\infty)$:*

(i). *The law of $(\tilde{\mathbf{v}}^n, \tilde{W}^n)$ is the same as (\mathbf{v}^n, W) for every $n \in \mathbb{N}$.*

(ii). *$(\tilde{\mathbf{v}}^n, \tilde{W}^n) \rightarrow (\tilde{\mathbf{v}}, \tilde{W})$, $\tilde{\mathbb{P}}$ -almost surely.*

(iii). *The random variables $\tilde{\mathbf{v}}$ and $\tilde{\mathbf{v}}^n$ share the same uniform estimates with \mathbf{v}^n as given by Lemma 2.3. In particular, for every $n \in \mathbb{N}$, $T > 0$ and $p \geq 1$,*

$$\tilde{\mathbb{E}} \left(\sup_{t \in [0, T]} \|\tilde{\mathbf{v}}^n(t)\|_{H^1(\mathbb{R}^d)}^{2p} \right) + \tilde{\mathbb{E}} \left(\int_0^T \|\tilde{\mathbf{v}}^n(t)\|_{H^2(\mathbb{R}^d)}^2 dt \right)^p \leq C_1, \quad (2.28)$$

and

$$\tilde{\mathbb{E}} \left(\sup_{t \in [0, T]} \|\tilde{\mathbf{v}}(t)\|_{H^1(\mathbb{R}^d)}^{2p} \right) + \tilde{\mathbb{E}} \left(\int_0^T \|\tilde{\mathbf{v}}(t)\|_{H^2(\mathbb{R}^d)}^2 dt \right)^p \leq C_1, \quad (2.29)$$

where $C_1 > 0$ is the same number as in (2.16).

(iv). $\tilde{\mathbf{v}}^n \rightarrow \tilde{\mathbf{v}}$ in $L^2(\tilde{\Omega}; L^2(0, T; H^1(\mathbb{R}^d)))$.

Proof. We will verify that all conditions of Proposition 1.1 are fulfilled. We first show that there exists a sequence of continuous real-valued functions defined in each of the space appearing in

$$(C([0, T]; L^2(\mathbb{R}^d)) \cap C([0, T]; H_w^1(\mathbb{R}^d)) \cap L_w^2(0, T; H^2(\mathbb{R}^d))) \times C([0, T]; \mathbb{R}^\infty),$$

which separate points. Such continuous functions indeed exist in $C([0, T]; L^2(\mathbb{R}^d))$ and $C([0, T]; \mathbb{R}^\infty)$ since they are both separable Banach spaces.

Since $L^2(0, T; H^2(\mathbb{R}^d))$ is a separable Hilbert space, any countable dense subset of $(L^2(0, T; H^2(\mathbb{R}^d)))^*$ forms a sequence of continuous functions on $L_w^2(0, T; H^2(\mathbb{R}^d))$ that separate points.

Let $\{\psi_n\}_{n=1}^\infty$ be a dense subset of $H^1(\mathbb{R}^d)$ and $\{r_j\}_{j=1}^\infty$ be the set of all rational numbers in $[0, T]$. Given $n, j \in \mathbb{N}$, define $g_{n,j} : C([0, T]; H_w^1(\mathbb{R}^d)) \rightarrow \mathbb{R}$ by

$$g_{n,j}(\xi) = (\xi(r_j), \psi_n)_{H^1(\mathbb{R}^d)}, \quad \forall \xi \in C([0, T]; H_w^1(\mathbb{R}^d)).$$

Then $g_{n,j}$ is continuous on $C([0, T]; H_w^1(\mathbb{R}^d))$ and the sequence $\{g_{n,j}\}_{n,j=1}^\infty$ separates the points of $C([0, T]; H_w^1(\mathbb{R}^d))$.

On the other hand, by Lemma 2.4 we see that the distributions of the sequence $\{(\mathbf{v}^n, W)\}_{n=1}^\infty$ are tight in $(C([0, T]; L^2(\mathbb{R}^d)) \cap C([0, T]; H_w^1(\mathbb{R}^d)) \cap L_w^2(0, T; H^2(\mathbb{R}^d))) \times C([0, T]; \mathbb{R}^\infty)$, and hence by Proposition 1.1, there exist a subsequence of $\{(\mathbf{v}^n, W)\}_{n=1}^\infty$ (which is not relabeled), a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$, random variables $(\tilde{\mathbf{v}}, \tilde{W})$ and $(\tilde{\mathbf{v}}^n, \tilde{W}^n)$, $n \in \mathbb{N}$, defined on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ such that (i) and (ii) are fulfilled. Then the uniform estimates of (2.28) and (2.29) follows from (i) and Lemma 2.3 immediately.

Finally, we show (iv). By (ii), we know that $\tilde{\mathbf{v}}^n \rightarrow \tilde{\mathbf{v}}$ in $C([0, T]; L^2(\mathbb{R}^d))$, $\tilde{\mathbb{P}}$ -almost surely. Then, by (2.28)-(2.29) and Vitali's theorem, we find that $\tilde{\mathbf{v}}^n \rightarrow \tilde{\mathbf{v}}$ in $L^2(\tilde{\Omega}; C([0, T]; L^2(\mathbb{R}^d)))$, and hence

$$\begin{aligned} & \tilde{\mathbb{E}} \int_0^T \|\tilde{\mathbf{v}}^n(t) - \tilde{\mathbf{v}}(t)\|_{H^1(\mathbb{R}^d)}^2 dt = \tilde{\mathbb{E}} \int_0^T (\tilde{\mathbf{v}}^n(t) - \tilde{\mathbf{v}}(t), (I - \Delta)(\tilde{\mathbf{v}}^n(t) - \tilde{\mathbf{v}}(t)))_{L^2(\mathbb{R}^d)} dt \\ & \leq \tilde{\mathbb{E}} \int_0^T \|\tilde{\mathbf{v}}^n(t) - \tilde{\mathbf{v}}(t)\|_{H^2(\mathbb{R}^d)} \|\tilde{\mathbf{v}}^n(t) - \tilde{\mathbf{v}}(t)\|_{L^2(\mathbb{R}^d)} dt \\ & \leq \tilde{\mathbb{E}} \left(\sup_{t \in [0, T]} \|\tilde{\mathbf{v}}^n(t) - \tilde{\mathbf{v}}(t)\|_{L^2(\mathbb{R}^d)} \int_0^T \|\tilde{\mathbf{v}}^n(t) - \tilde{\mathbf{v}}(t)\|_{H^2(\mathbb{R}^d)} dt \right) \\ & \leq T^{\frac{1}{2}} \left[\tilde{\mathbb{E}} \left(\sup_{t \in [0, T]} \|\tilde{\mathbf{v}}^n(t) - \tilde{\mathbf{v}}(t)\|_{L^2(\mathbb{R}^d)}^2 \right) \right]^{\frac{1}{2}} \left[\tilde{\mathbb{E}} \int_0^T \|\tilde{\mathbf{v}}^n(t) - \tilde{\mathbf{v}}(t)\|_{H^2(\mathbb{R}^d)}^2 dt \right]^{\frac{1}{2}} \end{aligned}$$

$$\leq T^{\frac{1}{2}} \left[\tilde{\mathbb{E}} \left(\sup_{t \in [0, T]} \|\tilde{\mathbf{v}}^n(t) - \tilde{\mathbf{v}}(t)\|_{L^2(\mathbb{R}^d)}^2 \right) \right]^{\frac{1}{2}} \left[2\tilde{\mathbb{E}} \int_0^T (\|\tilde{\mathbf{v}}^n(t)\|_{H^2(\mathbb{R}^d)}^2 + \|\tilde{\mathbf{v}}(t)\|_{H^2(\mathbb{R}^d)}^2) dt \right]^{\frac{1}{2}} \rightarrow 0,$$

that is, $\tilde{\mathbf{v}}^n \rightarrow \tilde{\mathbf{v}}$ in $L^2(\tilde{\Omega}; L^2(0, T; H^1(\mathbb{R}^d)))$, which completes the proof. \square

2.3. Passage to the limit of approximate solutions. We are now in a position to complete the proof of Theorem 1.1 by showing the process $\tilde{\mathbf{v}}$ of the limit of the sequence $\{\tilde{\mathbf{v}}\}_{n=1}^\infty$ is a solution of (1.1).

Proof of Theorem 1.1. If $\varphi \in C_0^\infty(\mathbb{R}^d)$, then by (2.2) we have, for all $t \in [0, T]$ and $n \in \mathbb{N}$,

$$\begin{aligned} (\mathbf{u}^n(t), \varphi)_{L^2(\mathcal{D}_n)} &= (\mathbf{u}^n(0), \varphi)_{L^2(\mathcal{D}_n)} + \int_0^t (\Delta \mathbf{u}^n(s), \varphi)_{L^2(\mathcal{D}_n)} ds \\ &+ \int_0^t (\mathbf{u}^n(s) \times \Delta \mathbf{u}^n(s), \varphi)_{L^2(\mathcal{D}_n)} ds - \int_0^t ((1 + |\mathbf{u}^n(s)|^2) \mathbf{u}^n(s), \varphi)_{L^2(\mathcal{D}_n)} ds \\ &+ \frac{1}{2} \sum_{k \geq 1} \int_0^t ((\mathbf{u}^n(s) \times \mathbf{f}_k) \times \mathbf{f}_k, \varphi)_{L^2(\mathcal{D}_n)} ds + \sum_{k \geq 1} \int_0^t (\mathbf{u}^n(s) \times \mathbf{f}_k + \mathbf{f}_k, \varphi)_{L^2(\mathcal{D}_n)} dW_k. \end{aligned} \quad (2.30)$$

Since φ has a compact support in \mathbb{R}^d , there exists $n_0 \in \mathbb{N}$ such that $\varphi(x) = 0$ for all $x \in \mathbb{R}^d \setminus \mathcal{D}_{n_0}$. Therefore, we have for all $n \geq n_0$,

$$(\mathbf{u}^n(t), \varphi)_{L^2(\mathcal{D}_n)} = \int_{\mathcal{D}_{n_0}} \mathbf{u}^n(t, x) \varphi(x) dx. \quad (2.31)$$

Recall that $\mathbf{v}^n(t, x) = \theta_n(x) \mathbf{u}^n(t, x)$ and $\theta_n(x) = 1$ for $|x| \leq \frac{1}{2}n$, we know that $\mathbf{v}^n(t, x) = \mathbf{u}^n(t, x)$ for all $t \geq 0$ and $|x| \leq \frac{1}{2}n$, and hence for all $n \geq 2n_0$,

$$\int_{\mathcal{D}_{n_0}} \mathbf{u}^n(t, x) \varphi(x) dx = \int_{\mathcal{D}_{n_0}} \mathbf{v}^n(t, x) \varphi(x) dx = \int_{\mathbb{R}^d} \mathbf{v}^n(t, x) \varphi(x) dx. \quad (2.32)$$

Notice that $\mathbf{v}^n(t, x)$ is defined for all $x \in \mathbb{R}^d$, while $\mathbf{u}^n(t, x)$ is defined only for $x \in \mathcal{D}_n$. By (2.31)-(2.32) we get, for all $n \geq 2n_0$,

$$(\mathbf{u}^n(t), \varphi)_{L^2(\mathcal{D}_n)} = (\mathbf{v}^n(t), \varphi)_{L^2(\mathbb{R}^d)}. \quad (2.33)$$

By the same reason, for all $n \geq 2n_0$, the inner product in $L^2(\mathcal{D}_n)$ of every term in (2.30) can be replaced by the inner product in $L^2(\mathbb{R}^d)$, and the variable \mathbf{u}^n can be replaced by \mathbf{v}^n . Consequently, for all $t \in [0, T]$ and $n \geq 2n_0$, we have,

$$\begin{aligned} (\mathbf{v}^n(t), \varphi)_{L^2(\mathbb{R}^d)} &= (\theta_n \mathbf{u}_0, \varphi)_{L^2(\mathbb{R}^d)} + \int_0^t (\Delta \mathbf{v}^n(s), \varphi)_{L^2(\mathbb{R}^d)} ds \\ &+ \int_0^t (\mathbf{v}^n(s) \times \Delta \mathbf{v}^n(s), \varphi)_{L^2(\mathbb{R}^d)} ds - \int_0^t ((1 + |\mathbf{v}^n(s)|^2) \mathbf{v}^n(s), \varphi)_{L^2(\mathbb{R}^d)} ds \\ &+ \frac{1}{2} \sum_{k \geq 1} \int_0^t ((\mathbf{v}^n(s) \times \mathbf{f}_k) \times \mathbf{f}_k, \varphi)_{L^2(\mathbb{R}^d)} ds + \sum_{k \geq 1} \int_0^t (\mathbf{v}^n(s) \times \mathbf{f}_k + \mathbf{f}_k, \varphi)_{L^2(\mathbb{R}^d)} dW_k. \end{aligned} \quad (2.34)$$

By Lemma 2.5 (i), we see that $(\tilde{\mathbf{v}}^n, \tilde{W}^n)$ has the same law as (\mathbf{v}^n, W) for every $n \in \mathbb{N}$. Then by the argument of [8, 38], we find that $(\tilde{\mathbf{v}}^n, \tilde{W}^n)$ also satisfies system (2.34). From now on, we will deal with the variables $(\tilde{\mathbf{v}}^n, \tilde{W}^n)$ instead of (\mathbf{v}^n, W) , and for the sake of simplicity in notation, we will use \mathbf{v}^n for both the variables \mathbf{v}^n and $\tilde{\mathbf{v}}^n$, and use W^n for \tilde{W}^n .

By Lemma 2.5 (iv) we infer that for every $\psi \in L^\infty(\Omega \times [0, T])$,

$$\begin{aligned}
& \mathbb{E} \left(\int_0^T \psi(t) \left(\int_0^t ((\Delta \mathbf{v}^n(s), \varphi)_{L^2(\mathbb{R}^d)} - (\Delta \mathbf{v}(s), \varphi)_{L^2(\mathbb{R}^d)}) ds \right) dt \right) \\
& \leq \mathbb{E} \left(\int_0^T \psi(t) \left(\int_0^t \|\nabla \mathbf{v}^n(s) - \nabla \mathbf{v}(s)\|_{L^2(\mathbb{R}^d)} \|\nabla \varphi\|_{L^2(\mathbb{R}^d)} ds \right) dt \right) \\
& \leq T \|\psi\|_{L^\infty(\Omega \times [0, T])} \|\nabla \varphi\|_{L^2(\mathbb{R}^d)} \mathbb{E} \left(\int_0^T \|\nabla \mathbf{v}^n(s) - \nabla \mathbf{v}(s)\|_{L^2(\mathbb{R}^d)} ds \right) \\
& \leq T^{\frac{3}{2}} \|\psi\|_{L^\infty(\Omega \times [0, T])} \|\nabla \varphi\|_{L^2(\mathbb{R}^d)} \|\mathbf{v}^n - \mathbf{v}\|_{L^2(\Omega, L^2(0, T; H^1(\mathbb{R}^d)))} \rightarrow 0, \tag{2.35}
\end{aligned}$$

as $n \rightarrow \infty$.

For the third term on the right-hand side of (2.34), we have for all $t \in [0, T]$,

$$\begin{aligned}
& \left| \int_0^t (\mathbf{v}^n(s) \times \Delta \mathbf{v}^n(s) - \mathbf{v}(s) \times \Delta \mathbf{v}(s), \varphi)_{L^2(\mathbb{R}^d)} ds \right| \\
& = \left| \int_0^t ((\mathbf{v}^n(s) - \mathbf{v}(s)) \times \Delta \mathbf{v}^n(s) + \mathbf{v}(s) \times \Delta(\mathbf{v}^n(s) - \mathbf{v}(s)), \varphi)_{L^2(\mathbb{R}^d)} ds \right| \\
& \leq \left| \int_0^t ((\mathbf{v}^n(s) - \mathbf{v}(s)) \times \Delta \mathbf{v}^n(s), \varphi)_{L^2(\mathbb{R}^d)} ds \right| \\
& \quad + \left| \int_0^t (\nabla \mathbf{v}(s) \times \nabla(\mathbf{v}^n(s) - \mathbf{v}(s)), \varphi)_{L^2(\mathbb{R}^d)} + (\mathbf{v}(s) \times \nabla(\mathbf{v}^n(s) - \mathbf{v}(s)), \nabla \varphi)_{L^2(\mathbb{R}^d)} ds \right| \\
& \leq \|\varphi\|_{L^\infty(\mathbb{R}^d)} \int_0^t \|\mathbf{v}^n(s) - \mathbf{v}(s)\|_{L^2(\mathbb{R}^d)} \|\Delta \mathbf{v}^n(s)\|_{L^2(\mathbb{R}^d)} ds \\
& \quad + \|\varphi\|_{L^\infty(\mathbb{R}^d)} \int_0^t \|\nabla \mathbf{v}(s)\|_{L^2(\mathbb{R}^d)} \|\nabla \mathbf{v}^n(s) - \nabla \mathbf{v}(s)\|_{L^2(\mathbb{R}^d)} ds \\
& \quad + \|\nabla \varphi\|_{L^\infty(\mathbb{R}^d)} \int_0^t \|\mathbf{v}(s)\|_{L^2(\mathbb{R}^d)} \|\nabla \mathbf{v}^n(s) - \nabla \mathbf{v}(s)\|_{L^2(\mathbb{R}^d)} ds \\
& \leq c_1 \left(\int_0^T \|\mathbf{v}^n(t) - \mathbf{v}(t)\|_{H^1(\mathbb{R}^d)}^2 dt \right)^{\frac{1}{2}} \left(\int_0^T (\|\mathbf{v}^n(t)\|_{H^2(\mathbb{R}^d)}^2 + \|\mathbf{v}(t)\|_{H^1(\mathbb{R}^d)}^2) dt \right)^{\frac{1}{2}},
\end{aligned}$$

where $c_1 > 0$ depends only on φ . Then by Lemma 2.5 (iii)-(iv) we obtain that for any $\psi \in L^\infty(\Omega \times [0, T])$,

$$\begin{aligned}
& \mathbb{E} \left(\int_0^T \psi(t) \left(\int_0^t (\mathbf{v}^n(s) \times \Delta \mathbf{v}^n(s) - \mathbf{v}(s) \times \Delta \mathbf{v}(s), \varphi)_{L^2(\mathbb{R}^d)} ds \right) dt \right) \\
& \leq \|\psi\|_{L^\infty(\Omega \times [0, T])} \mathbb{E} \left(\int_0^T \left| \int_0^t (\mathbf{v}^n(s) \times \Delta \mathbf{v}^n(s) - \mathbf{v}(s) \times \Delta \mathbf{v}(s), \varphi)_{L^2(\mathbb{R}^d)} ds \right| dt \right)
\end{aligned}$$

$$\begin{aligned}
&\leq c_1 T \|\psi\|_{L^\infty(\Omega \times [0, T])} \mathbb{E} \left(\left(\int_0^T \|\mathbf{v}^n(t) - \mathbf{v}(t)\|_{H^1(\mathbb{R}^d)}^2 dt \right)^{\frac{1}{2}} \left(\int_0^T \left(\|\mathbf{v}^n(t)\|_{H^2(\mathbb{R}^d)}^2 + \|\mathbf{v}(t)\|_{H^1(\mathbb{R}^d)}^2 \right) dt \right)^{\frac{1}{2}} \right) \\
&\leq c_2 \|\mathbf{v}^n - \mathbf{v}\|_{L^2(\Omega, L^2(0, T; H^1(\mathbb{R}^d)))} \left(\|\mathbf{v}^n\|_{L^2(\Omega, L^2(0, T; H^2(\mathbb{R}^d)))} + \|\mathbf{v}\|_{L^2(\Omega, L^2(0, T; H^1(\mathbb{R}^d)))} \right) \rightarrow 0, \quad (2.36)
\end{aligned}$$

as $n \rightarrow \infty$, where $c_2 = c_1 T \|\psi\|_{L^\infty(\Omega \times [0, T])}$.

For the fourth term on the right-hand side of (2.34), by the embedding $H^1(\mathbb{R}^d) \hookrightarrow L^4(\mathbb{R}^d)$, we have

$$\begin{aligned}
&\left| \int_0^t (|\mathbf{v}^n(s)|^2 \mathbf{v}^n(s) - |\mathbf{v}(s)|^2 \mathbf{v}(s), \varphi)_{L^2(\mathbb{R}^d)} ds \right| \\
&\leq \frac{3}{2} \|\varphi\|_{L^\infty(\mathbb{R}^d)} \int_0^t \left(\|\mathbf{v}^n(s)\|_{L^4(\mathbb{R}^d)}^2 + \|\mathbf{v}(s)\|_{L^4(\mathbb{R}^d)}^2 \right) \|\mathbf{v}^n(s) - \mathbf{v}(s)\|_{L^2(\mathbb{R}^d)} ds \\
&\leq \frac{3}{2} T^{\frac{1}{2}} \|\varphi\|_{L^\infty(\mathbb{R}^d)} \left(\sup_{s \in [0, T]} \|\mathbf{v}^n(s)\|_{H^1(\mathbb{R}^d)}^2 + \sup_{s \in [0, T]} \|\mathbf{v}(s)\|_{H^1(\mathbb{R}^d)}^2 \right) \|\mathbf{v}^n - \mathbf{v}\|_{L^2(0, T; L^2(\mathbb{R}^d))}. \quad (2.37)
\end{aligned}$$

By (2.37) and Lemma 2.5 (iii)-(iv) we obtain that for all $\psi \in L^\infty(\Omega \times [0, T])$,

$$\begin{aligned}
&\mathbb{E} \left(\int_0^T \psi(t) \left(\int_0^t (|\mathbf{v}^n(s)|^2 \mathbf{v}^n(s) - |\mathbf{v}(s)|^2 \mathbf{v}(s), \varphi)_{L^2(\mathbb{R}^d)} ds \right) dt \right) \\
&\leq \|\psi\|_{L^\infty(\Omega \times [0, T])} \mathbb{E} \left(\int_0^T \left| \int_0^t (|\mathbf{v}^n(s)|^2 \mathbf{v}^n(s) - |\mathbf{v}(s)|^2 \mathbf{v}(s), \varphi)_{L^2(\mathbb{R}^d)} ds \right| dt \right) \\
&\leq c_3 \mathbb{E} \left(\left(\sup_{s \in [0, T]} \|\mathbf{v}^n(s)\|_{H^1(\mathbb{R}^d)}^2 + \sup_{s \in [0, T]} \|\mathbf{v}(s)\|_{H^1(\mathbb{R}^d)}^2 \right) \|\mathbf{v}^n - \mathbf{v}\|_{L^2(0, T; L^2(\mathbb{R}^d))} \right) \\
&\leq c_3 \left(\|\mathbf{v}^n\|_{L^4(\Omega, L^\infty(0, T; H^1(\mathbb{R}^d)))}^2 + \|\mathbf{v}\|_{L^4(\Omega, L^\infty(0, T; H^1(\mathbb{R}^d)))}^2 \right) \|\mathbf{v}^n - \mathbf{v}\|_{L^2(\Omega, L^2(0, T; L^2(\mathbb{R}^d)))} \rightarrow 0, \quad (2.38)
\end{aligned}$$

as $n \rightarrow \infty$, where $c_3 = \frac{3}{2} T^{\frac{3}{2}} \|\varphi\|_{L^\infty(\mathbb{R}^d)} \|\psi\|_{L^\infty(\Omega \times [0, T])}$.

Similarly, for the fifth term on the right-hand side of (2.34) we have for all $\psi \in L^\infty(\Omega \times [0, T])$,

$$\mathbb{E} \left(\int_0^T \psi(t) \left(\sum_{k \geq 1} \int_0^t ((\mathbf{v}^n(s) \times \mathbf{f}_k) \times \mathbf{f}_k - (\mathbf{v}(s) \times \mathbf{f}_k) \times \mathbf{f}_k, \varphi)_{L^2(\mathbb{R}^d)} ds \right) dt \right) \rightarrow 0, \quad (2.39)$$

as $n \rightarrow \infty$.

We next consider the convergence of the last term on the right-hand side of (2.34). By (1.1) and Lemma 2.5 (iv), we see that for almost all $\omega \in \Omega$,

$$\sum_{k \geq 1} \int_0^t (\mathbf{v}^n(s) \times \mathbf{f}_k - \mathbf{v}(s) \times \mathbf{f}_k, \varphi)_{L^2(\mathbb{R}^d)} ds \rightarrow 0, \quad \text{as } n \rightarrow \infty,$$

in $L^2(0, T; \mathbb{R})$, which along with [14, Lemma 2.1] shows that

$$\sum_{k \geq 1} \int_0^t ((\mathbf{v}^n(s) \times \mathbf{f}_k + \mathbf{f}_k), \varphi)_{L^2(\mathbb{R}^d)} dW_k^n \rightarrow \sum_{k \geq 1} \int_0^t ((\mathbf{v}(s) \times \mathbf{f}_k + \mathbf{f}_k), \varphi)_{L^2(\mathbb{R}^d)} dW_k, \quad (2.40)$$

in $L^2(0, T)$ in probability as $n \rightarrow \infty$. Moreover, by (2.28) and (1.1) we have

$$\sup_{n \geq 1} \mathbb{E} \left(\int_0^T ((\mathbf{v}^n(t) \times \mathbf{f}_k + \mathbf{f}_k), \varphi)^4 dt \right) < \infty,$$

which together with (2.40) yields

$$\sum_{k \geq 1} \int_0^t ((\mathbf{v}^n(s) \times \mathbf{f}_k + \mathbf{f}_k), \varphi)_{L^2(\mathbb{R}^d)} dW_k^n \rightarrow \sum_{k \geq 1} \int_0^t ((\mathbf{v}(s) \times \mathbf{f}_k + \mathbf{f}_k), \varphi)_{L^2(\mathbb{R}^d)} dW_k, \quad (2.41)$$

in $L^2(\Omega \times [0, T])$. Therefore, for any $\psi \in L^\infty(\Omega \times [0, T])$ we get from (2.41) that

$$\begin{aligned} & \mathbb{E} \left(\int_0^T \psi(t) \left(\sum_{k \geq 1} \int_0^t ((\mathbf{v}^n(s) \times \mathbf{f}_k + \mathbf{f}_k), \varphi)_{L^2(\mathbb{R}^d)} dW_k^n \right) dt \right) \\ & \rightarrow \mathbb{E} \left(\int_0^T \psi(t) \left(\sum_{k \geq 1} \int_0^t ((\mathbf{v}(s) \times \mathbf{f}_k + \mathbf{f}_k), \varphi)_{L^2(\mathbb{R}^d)} dW_k \right) dt \right). \end{aligned} \quad (2.42)$$

Taking the limit of (2.34) as $n \rightarrow \infty$, by (2.35)-(2.36), (2.38)-(2.39), (2.42) and Lemma 2.5 (iv), we obtain that for any $\psi \in L^\infty(\Omega \times [0, T])$ and $\varphi \in C_0^\infty(\mathbb{R}^d)$,

$$\begin{aligned} & \mathbb{E} \left(\int_0^T \psi(t) (\mathbf{v}(t), \varphi)_{L^2(\mathbb{R}^d)} dt \right) = \mathbb{E} \left(\int_0^T \psi(t) (\mathbf{u}_0, \varphi)_{L^2(\mathbb{R}^d)} dt \right) \\ & + \mathbb{E} \left(\int_0^T \psi(t) \left(\int_0^t (\Delta \mathbf{v}(s) + \mathbf{v}(s) \times \Delta \mathbf{v}(s) - (1 + |\mathbf{v}(s)|^2) \mathbf{v}(s), \varphi)_{L^2(\mathbb{R}^d)} ds \right) dt \right) \\ & + \frac{1}{2} \mathbb{E} \left(\int_0^T \psi(t) \left(\sum_{k \geq 1} \int_0^t ((\mathbf{v}(s) \times \mathbf{f}_k) \times \mathbf{f}_k, \varphi)_{L^2(\mathbb{R}^d)} ds \right) dt \right) \\ & + \mathbb{E} \left(\int_0^T \psi(t) \left(\sum_{k \geq 1} \int_0^t ((\mathbf{v}(s) \times \mathbf{f}_k + \mathbf{f}_k), \varphi)_{L^2(\mathbb{R}^d)} dW_k \right) dt \right). \end{aligned} \quad (2.43)$$

By the arbitrariness of $\psi \in L^\infty(\Omega \times [0, T])$, $\varphi \in C_0^\infty(\mathbb{R}^d)$ and the density of $C_0^\infty(\mathbb{R}^d)$ in $L^2(\mathbb{R}^d)$, we infer from (2.43) that \mathbf{v} satisfies (1.6), and thus $(\Omega, \mathcal{F}, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P}, W, \mathbf{v})$ is a global martingale solution of the stochastic equation (1.3).

Note that (1.4) follows from the regularity of $\tilde{\mathbf{v}}$ given by (2.29). In addition, by (2.17) we have

$$\mathbb{E} \left(\int_0^T \|\mathbf{v}(s, \mathbf{u}_0) \times \Delta \mathbf{v}(s, \mathbf{u}_0)\|_{L^2(\mathbb{R}^d)}^{\frac{4}{3}} ds \right)^p < \infty, \quad \forall p \geq 1,$$

which shows that (1.5) is true for $r \in [1, \frac{4}{3}]$. Next, we prove (1.5) is also true for $r \in [\frac{4}{3}, 2)$. We just consider the case $d = 2$ since $d = 1$ is simpler.

Given $r \in [\frac{4}{3}, 2)$ and $d = 2$, let $a = \frac{2}{r} - 1$ and $q = \frac{2}{a} - 2$. Since $r \in [\frac{4}{3}, 2)$, we have $a \in (0, \frac{1}{2}]$ and $q \geq 2$. Recall the following Gagliardo-Nirenberg inequality for $d = 2$:

$$\|\mathbf{v}\|_{L^\infty(\mathbb{R}^2)} \leq c_1 \|\Delta \mathbf{v}\|_{L^2(\mathbb{R}^2)}^a \|\mathbf{v}\|_{L^q(\mathbb{R}^2)}^{1-a}, \quad (2.44)$$

where $c_1 > 0$ is a constant. Since $H^1(\mathbb{R}^2) \hookrightarrow L^q(\mathbb{R}^2)$ for any $q \geq 2$, by (2.44) we get

$$\|\mathbf{v}\|_{L^\infty(\mathbb{R}^2)} \leq c_2 \|\Delta \mathbf{v}\|_{L^2(\mathbb{R}^2)}^a \|\mathbf{v}\|_{H^1(\mathbb{R}^2)}^{1-a}, \quad (2.45)$$

where $c_2 > 0$ is a constant. Then by (2.45) we have

$$\begin{aligned} & \int_0^T \|\mathbf{v}(s, \mathbf{u}_0) \times \Delta \mathbf{v}(s, \mathbf{u}_0)\|_{L^2(\mathbb{R}^2)}^r ds \leq \int_0^T \|\mathbf{v}(s, \mathbf{u}_0)\|_{L^\infty(\mathbb{R}^2)}^r \|\Delta \mathbf{v}(s, \mathbf{u}_0)\|_{L^2(\mathbb{R}^2)}^r ds \\ & \leq c_2^r \int_0^T \|\mathbf{v}(s, \mathbf{u}_0)\|_{H^1(\mathbb{R}^2)}^{(1-a)r} \|\Delta \mathbf{v}(s, \mathbf{u}_0)\|_{L^2(\mathbb{R}^2)}^{(a+1)r} ds \leq c_2^r \sup_{s \in [0, T]} \|\mathbf{v}(s, \mathbf{u}_0)\|_{H^1(\mathbb{R}^2)}^{(1-a)r} \int_0^T \|\Delta \mathbf{v}(s, \mathbf{u}_0)\|_{L^2(\mathbb{R}^2)}^2 ds, \end{aligned}$$

which along with (1.4) shows that for any $p \geq 1$,

$$\begin{aligned} & \mathbb{E} \left(\int_0^T \|\mathbf{v}(s, \mathbf{u}_0) \times \Delta \mathbf{v}(s, \mathbf{u}_0)\|_{L^2(\mathbb{R}^2)}^r ds \right)^p \\ & \leq c_2^{pr} \mathbb{E} \left(\sup_{s \in [0, T]} \|\mathbf{v}(s, \mathbf{u}_0)\|_{H^1(\mathbb{R}^2)}^{(1-a)pr} \left(\int_0^T \|\Delta \mathbf{v}(s, \mathbf{u}_0)\|_{L^2(\mathbb{R}^2)}^2 ds \right)^p \right) \\ & \leq c_2^{pr} \left(\mathbb{E} \left(\sup_{s \in [0, T]} \|\mathbf{v}(s, \mathbf{u}_0)\|_{H^1(\mathbb{R}^2)}^{2(1-a)pr} \right) \right)^{\frac{1}{2}} \left(\mathbb{E} \left(\left(\int_0^T \|\Delta \mathbf{v}(s, \mathbf{u}_0)\|_{L^2(\mathbb{R}^2)}^2 ds \right)^{2p} \right) \right)^{\frac{1}{2}} < \infty, \end{aligned}$$

and hence (1.5) is valid for all $p \geq 1$ and $r \in [1, 2)$.

Finally the pathwise uniqueness of solutions of (1.3) follows from the continuity of solutions with respect to initial data in $L^2(\mathbb{R}^d)$ which is established in Lemma 2.6 below. This along with the Yamada-Watanabe theorem completes the proof of Theorem 1.1.

Next, we prove the continuity of solutions of (1.3) in initial data in $L^2(\mathbb{R}^d)$ which will be used to establish the Feller property of the Markov semigroup associated with (1.3).

Lemma 2.6. *Suppose (1.1) holds, $T > 0$ and $\mathbf{u}_0, \mathbf{u}_{0,n} \in H^1(\mathbb{R}^d)$ such that $\mathbf{u}_{0,n} \rightarrow \mathbf{u}_0$ in $H^1(\mathbb{R}^d)$. Then the solutions $\mathbf{u}(\cdot, \mathbf{u}_{0,n})$ and $\mathbf{u}(\cdot, \mathbf{u}_0)$ of (1.3) satisfy, for any $p \geq 1$,*

$$\lim_{n \rightarrow \infty} \mathbb{E} \left(\sup_{t \in [0, T]} \|\mathbf{u}(t, \mathbf{u}_{0,n}) - \mathbf{u}(t, \mathbf{u}_0)\|_{L^2(\mathbb{R}^d)}^p \right) = 0.$$

Proof. Let $\mathbf{u}_n(t) = \mathbf{u}(t, \mathbf{u}_{0,n})$, $\mathbf{u}(t) = \mathbf{u}(t, \mathbf{u}_0)$, and $\mathbf{v}_n(t) = \mathbf{u}(t, \mathbf{u}_{0,n}) - \mathbf{u}(t, \mathbf{u}_0)$ for $t \in [0, T]$. Then by (1.3) we get

$$\begin{aligned} d\mathbf{v}_n &= (\Delta \mathbf{v}_n + \mathbf{v}_n \times \Delta \mathbf{u}_n + \mathbf{u} \times \Delta \mathbf{v}_n - \mathbf{v}_n - (|\mathbf{u}_n|^2 \mathbf{u}_n - |\mathbf{u}|^2 \mathbf{u})) dt \\ & \quad + \frac{1}{2} \sum_{k \geq 1} (\mathbf{v}_n \times \mathbf{f}_k) \times \mathbf{f}_k dt + \sum_{k \geq 1} (\mathbf{v}_n \times \mathbf{f}_k) dW_k. \end{aligned} \quad (2.46)$$

By Itô's formula we obtain

$$\begin{aligned} & d\|\mathbf{v}_n\|_{L^2(\mathbb{R}^d)}^2 + 2 \left(\|\nabla \mathbf{v}_n\|_{L^2(\mathbb{R}^d)}^2 + \|\mathbf{v}_n\|_{L^2(\mathbb{R}^d)}^2 \right) dt \\ &= 2(\mathbf{v}_n, (\mathbf{u} \times \Delta \mathbf{v}_n))_{L^2(\mathbb{R}^d)} dt - 2(\mathbf{v}_n, (|\mathbf{u}_n|^2 \mathbf{u}_n - |\mathbf{u}|^2 \mathbf{u}))_{L^2(\mathbb{R}^d)} dt. \end{aligned} \quad (2.47)$$

Note that

$$-2 (\mathbf{v}_n, (|\mathbf{u}_n|^2 \mathbf{u}_n - |\mathbf{u}|^2 \mathbf{u}))_{L^2(\mathbb{R}^d)} = -2 \int_{\mathbb{R}^d} (\mathbf{u}_n - \mathbf{u}) \cdot (|\mathbf{u}_n|^2 \mathbf{u}_n - |\mathbf{u}|^2 \mathbf{u}) dx \leq 0.$$

Then by (2.3) we get for all $t \in [0, T]$, P-almost surely,

$$\begin{aligned} & \|\mathbf{v}_n(t)\|_{L^2(\mathbb{R}^d)}^2 + 2 \int_0^t \|\nabla \mathbf{v}_n(s)\|_{L^2(\mathbb{R}^d)}^2 ds + 2 \int_0^t \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R}^d)}^2 ds \\ & \leq \|\mathbf{v}_n(0)\|_{L^2(\mathbb{R}^d)}^2 + 2 \int_0^t (\mathbf{v}_n(s), (\mathbf{u}(s) \times \Delta \mathbf{v}_n(s)))_{L^2(\mathbb{R}^d)} ds. \end{aligned} \quad (2.48)$$

For the last term on the right-hand side of (2.48) we will use the inequality for $d = 2$:

$$\|\mathbf{v}\|_{L^4(\mathbb{R}^d)} \leq c_1 \|\nabla \mathbf{v}\|_{L^2(\mathbb{R}^d)}^{\frac{1}{2}} \|\mathbf{v}\|_{L^2(\mathbb{R}^d)}^{\frac{1}{2}}, \quad \forall \mathbf{v} \in H^1(\mathbb{R}^d), \quad (2.49)$$

where $c_1 > 0$ is a constant. Then by (2.49) we get

$$\begin{aligned} & 2 \int_0^t (\mathbf{v}_n(s), (\mathbf{u}(s) \times \Delta \mathbf{v}_n(s)))_{L^2(\mathbb{R}^d)} ds = -2 \int_0^t (\mathbf{v}_n(s), (\nabla \mathbf{u}(s) \times \nabla \mathbf{v}_n(s)))_{L^2(\mathbb{R}^d)} ds \\ & \leq 2 \int_0^t \|\mathbf{v}_n(s)\|_{L^4(\mathbb{R}^d)} \|\nabla \mathbf{u}(s)\|_{L^4(\mathbb{R}^d)} \|\nabla \mathbf{v}_n(s)\|_{L^2(\mathbb{R}^d)} ds \\ & \leq 2c_1^2 \int_0^t \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R}^d)}^{\frac{1}{2}} \|\mathbf{u}(s)\|_{H^1(\mathbb{R}^d)}^{\frac{1}{2}} \|\mathbf{u}(s)\|_{H^2(\mathbb{R}^d)}^{\frac{1}{2}} \|\nabla \mathbf{v}_n(s)\|_{L^2(\mathbb{R}^d)}^{\frac{3}{2}} ds \\ & \leq \int_0^t \|\nabla \mathbf{v}_n(s)\|_{L^2(\mathbb{R}^d)}^2 ds + c_2 \int_0^t \|\mathbf{u}(s)\|_{H^1(\mathbb{R}^d)}^2 \|\mathbf{u}(s)\|_{H^2(\mathbb{R}^d)}^2 \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R}^d)}^2 ds, \end{aligned} \quad (2.50)$$

where $c_2 > 0$ is an absolute constant. By (2.48) and (2.50) we get for all $t \in [0, T]$, P-almost surely,

$$\begin{aligned} & \|\mathbf{v}_n(t)\|_{L^2(\mathbb{R}^d)}^2 + \int_0^t \|\nabla \mathbf{v}_n(s)\|_{L^2(\mathbb{R}^d)}^2 ds + 2 \int_0^t \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R}^d)}^2 ds \\ & \leq \|\mathbf{v}_n(0)\|_{L^2(\mathbb{R}^d)}^2 + c_2 \int_0^t \|\mathbf{u}(s)\|_{H^1(\mathbb{R}^d)}^2 \|\mathbf{u}(s)\|_{H^2(\mathbb{R}^d)}^2 \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R}^d)}^2 ds. \end{aligned} \quad (2.51)$$

By (2.51) and Gronwall's inequality we get for all $t \in [0, T]$, P-almost surely,

$$\begin{aligned} & \|\mathbf{v}_n(t)\|_{L^2(\mathbb{R}^d)}^2 \leq \|\mathbf{v}_n(0)\|_{L^2(\mathbb{R}^d)}^2 e^{c_2 \int_0^t \|\mathbf{u}(s)\|_{H^1(\mathbb{R}^d)}^2 \|\mathbf{u}(s)\|_{H^2(\mathbb{R}^d)}^2 ds} \\ & \leq \|\mathbf{v}_n(0)\|_{L^2(\mathbb{R}^d)}^2 e^{c_2 \sup_{s \in [0, T]} \|\mathbf{u}(s)\|_{H^1(\mathbb{R}^d)}^2 \int_0^T \|\mathbf{u}(s)\|_{H^2(\mathbb{R}^d)}^2 ds} = c_3 \|\mathbf{v}_n(0)\|_{L^2(\mathbb{R}^d)}^2, \end{aligned} \quad (2.52)$$

where

$$c_3 = e^{c_2 \sup_{s \in [0, T]} \|\mathbf{u}(s)\|_{H^1(\mathbb{R}^d)}^2 \int_0^T \|\mathbf{u}(s)\|_{H^2(\mathbb{R}^d)}^2 ds}.$$

By (2.29) we know that $c_3 < \infty$, P-almost surely. Then by (2.52) we see that P-almost surely,

$$\lim_{n \rightarrow \infty} \|\mathbf{v}_n\|_{C([0, T], L^2(\mathbb{R}^d))} = 0.$$

On the other hand, by (2.29) we see that if $\mathbf{u}_{0, n} \rightarrow \mathbf{u}_0$ in $H^1(\mathbb{R}^d)$, then the sequence $\{\mathbf{v}_n\}_{n=1}^{\infty}$ is uniformly integrable in $L^p(\Omega, C([0, T], L^2(\mathbb{R}^d)))$ for any $p \geq 1$, and hence by Vitali's theorem, we get for all $p \geq 1$,

$$\lim_{n \rightarrow \infty} \mathbb{E} \left(\|\mathbf{v}_n\|_{C([0, T], L^2(\mathbb{R}^d))}^p \right) = 0,$$

which completes the proof. \square

3. EXISTENCE OF INVARIANT MEASURES

In this section, we prove the existence of invariant measures of the Markov semigroup associated with the solution operators of the stochastic equation (1.3). Given $t \geq 0$ and $\mathbf{u}_0 \in H^1(\mathbb{R}^d)$, denote by $\mathbf{P}_t(\mathbf{u}_0, \cdot)$ the transition probability given by

$$\mathbf{P}_t(\mathbf{u}_0, \mathcal{O}) = \mathbb{P}(\mathbf{u}(t, \mathbf{u}_0) \in \mathcal{O}), \quad \forall \mathcal{O} \in \mathcal{B}(H^1(\mathbb{R}^d)),$$

where $\mathbf{u}(t, \mathbf{u}_0)$ is the solution of (1.3) starting from the initial point \mathbf{u}_0 .

For any bounded Borel function $\varphi \in \mathcal{B}_b(H^1(\mathbb{R}^d))$, define the Markov transition semigroup

$$(\mathbf{P}_t \varphi)(\mathbf{u}_0) = \mathbb{E}[\varphi(\mathbf{u}(t, \mathbf{u}_0))], \quad \mathbf{u}_0 \in H^1(\mathbb{R}^d).$$

A probability measure μ on $\mathcal{B}(H^1(\mathbb{R}^d))$ is an invariant measure of $\{\mathbf{P}_t\}_{t \geq 0}$ if

$$\int_{H^1(\mathbb{R}^d)} \mathbf{P}_t \varphi d\mu = \int_{H^1(\mathbb{R}^d)} \varphi d\mu, \quad \text{for all } t \geq 0, \varphi \in \mathcal{B}_b(H^1(\mathbb{R}^d)).$$

We will apply the weakly Feller method [31] to establish the existence of invariant measures of $\{\mathbf{P}_t\}_{t \geq 0}$ in $H^1(\mathbb{R}^d)$, for which we need the following convergence result as given by Lemma 2.5.

Lemma 3.1. *Suppose (1.1) holds, $\mathbf{u}_0, \mathbf{u}_{0,n} \in H^1(\mathbb{R}^d)$ such that $\{\mathbf{u}_{0,n}\}_{n=1}^\infty$ weakly converges to \mathbf{u}_0 in $H^1(\mathbb{R}^d)$. Let $\mathbf{u}^n(\cdot, \mathbf{u}_{0,n})$ be the unique solution of (1.3) with respect to initial data $\mathbf{u}_{0,n}$. Then there exist a stochastic basis $(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}_{t \geq 0}, \tilde{\mathbb{P}})$, random variables $(\tilde{\mathbf{u}}, \tilde{W})$ and $(\tilde{\mathbf{u}}^n, \tilde{W}^n)$, $n \in \mathbb{N}$, defined on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}_{t \geq 0}, \tilde{\mathbb{P}})$ such that in the space $(C([0, T]; L^2(\mathbb{R}^d)) \cap C([0, T]; H_w^1(\mathbb{R}^d)) \cap L_w^2(0, T; H^2(\mathbb{R}^d))) \times C([0, T]; \mathbb{R}^\infty)$, up to a subsequence:*

- (i). *The law of $(\tilde{\mathbf{u}}^n, \tilde{W}^n)$ is the same as (\mathbf{u}^n, W) for every $n \in \mathbb{N}$.*
- (ii). *$(\tilde{\mathbf{u}}^n, \tilde{W}^n) \rightarrow (\tilde{\mathbf{u}}, \tilde{W})$, $\tilde{\mathbb{P}}$ -almost surely.*
- (iii). *$(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}_{t \geq 0}, \tilde{\mathbf{u}}, \tilde{W}, \tilde{\mathbb{P}})$ is a solution of (1.3).*

Proof. Since $\{\mathbf{u}_{0,n}\}_{n=1}^\infty$ weakly converges to \mathbf{u}_0 in $H^1(\mathbb{R}^d)$, we find that the sequence $\{\mathbf{u}_{0,n}\}_{n=1}^\infty$ is bounded in $H^1(\mathbb{R}^d)$, and thus all the uniform estimates established in the previous section are valid for the sequence $\{\mathbf{u}^n(\cdot, \mathbf{u}_{0,n})\}_{n=1}^\infty$. Then the desired result follows from the same argument as Lemma 2.5. The details are omitted here. \square

Next, we show the existence of invariant measures of (1.3).

Theorem 3.1. *If (1.1) holds, then the stochastic equation (1.3) has an invariant measure in $H^1(\mathbb{R}^d)$.*

Proof. By Lemma 3.1, one can verify that the Markov semigroup $\{\mathbf{P}_t\}_{t \geq 0}$ is weakly Feller in $H^1(\mathbb{R}^d)$ in the sense that if $\{\mathbf{u}_{0,n}\}_{n=1}^\infty$ is convergent to \mathbf{u}_0 in $H^1(\mathbb{R}^d)$ weakly, and $\varphi : H^1(\mathbb{R}^d) \rightarrow \mathbb{R}$ is bounded and weakly continuous, then for all $t \geq 0$,

$$(\mathbf{P}_t \varphi)(\mathbf{u}_{0,n}) \rightarrow (\mathbf{P}_t \varphi)(\mathbf{u}_0), \quad \text{as } n \rightarrow \infty.$$

Moreover, the tightness of time averaged measures on $(H^1(\mathbb{R}^d), bw)$ could be obtained by the estimates (2.19). Then, we obtain the existence of at least one invariant measure for (1.3) following the Maslowski-Seidler theorem, see, e.g., [4, 31] for more details. \square

In the next section, we study the limit of invariant measures of (1.3) as the noise intensity ε approaches $\varepsilon_0 \in [0, 1]$.

4. LIMITING BEHAVIOR OF INVARIANT MEASURES

In this section, we investigate the uniform tightness of the collection of all invariant measures of (1.7) for $\varepsilon \in [0, 1]$ in the one-dimensional case. More precisely, we will show the union $\bigcup_{\varepsilon \in [0, 1]} \mathcal{I}_\varepsilon$ is tight in $H^1(\mathbb{R})$ where \mathcal{I}_ε is the set of all invariant measures of (1.7) corresponding to ε . To that end, we must establish the uniform tail-ends estimates of solutions in $H^1(\mathbb{R})$ in order to overcome the difficulty introduced by the non-compactness of the Sobolev embeddings on unbounded domains. Furthermore, by the idea of uniform tail-ends estimates, we will provide an alternative method to prove the existence of invariant measures of (1.7) by the Feller property of the Markov semigroup with respect to the strong topology instead of the weak topology.

We first establish the following uniform estimates of solutions which will be used for the forthcoming tail-ends arguments.

Lemma 4.1. *If (1.1) holds, then for every $\mathbf{u}_0 \in H^1(\mathbb{R})$, the solution $\mathbf{u}(t, \mathbf{u}_0)$ of (1.3) satisfies, for all $t \geq 0$,*

$$\begin{aligned} \mathbf{E} \left(\|\mathbf{u}(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 \right) + \int_0^t e^{s-t} \mathbf{E} \left(\|\mathbf{u}(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 + \int_{\mathbb{R}} |\mathbf{u}(s, \mathbf{u}_0)|^2 |\nabla \mathbf{u}(s, \mathbf{u}_0)|^2 dx \right) ds \\ \leq C e^{-t} \|\mathbf{u}_0\|_{H^1(\mathbb{R})}^2 + C, \end{aligned} \quad (4.1)$$

and

$$\int_0^t \mathbf{E} \left(\|\mathbf{u}(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 + \int_{\mathbb{R}} |\mathbf{u}(s, \mathbf{u}_0)|^2 |\nabla \mathbf{u}(s, \mathbf{u}_0)|^2 dx \right) ds \leq C \|\mathbf{u}_0\|_{H^1(\mathbb{R})}^2 + Ct, \quad (4.2)$$

where $C > 0$ is a constant depending only on $\{\mathbf{f}_k\}_{k=1}^\infty$.

Proof. By Itô's formula, we get from (1.3) that for all $t > 0$, P-almost surely,

$$\begin{aligned} d\|\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + 2 \left(\|\nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + \|\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + \|\mathbf{u}(t)\|_{L^4(\mathbb{R})}^4 \right) dt \\ = \sum_{k=1}^{\infty} \|f_k\|_{L^2(\mathbb{R})}^2 dt + 2 \sum_{k=1}^{\infty} (\mathbf{u}(t), f_k)_{L^2(\mathbb{R})} dW_k, \end{aligned}$$

and hence

$$\frac{d}{dt}\mathbf{E}\left(\|\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2\right) + 2\mathbf{E}\left(\|\nabla\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + \|\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + \|\mathbf{u}(t)\|_{L^4(\mathbb{R})}^4\right) = \sum_{k=1}^{\infty}\|f_k\|_{L^2(\mathbb{R})}^2. \quad (4.3)$$

By Gronwall's inequality, we obtain for all $t \geq 0$, P-almost surely,

$$\begin{aligned} \mathbf{E}\left(\|\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2\right) + \int_0^t e^{s-t}\mathbf{E}\left(2\|\nabla\mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 + \|\mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 + 2\|\mathbf{u}(s)\|_{L^4(\mathbb{R})}^4\right) ds \\ \leq e^{-t}\|\mathbf{u}_0\|_{L^2(\mathbb{R})}^2 + \sum_{k=1}^{\infty}\|f_k\|_{L^2(\mathbb{R})}^2. \end{aligned} \quad (4.4)$$

Note that

$$\begin{aligned} ((\nabla\mathbf{u} \times \mathbf{f}_k) \times \mathbf{f}_k, \nabla\mathbf{u}) &= -(\nabla\mathbf{u} \times \mathbf{f}_k, \nabla\mathbf{u} \times \mathbf{f}_k), \\ ((\mathbf{u} \times \nabla\mathbf{f}_k) \times \mathbf{f}_k, \nabla\mathbf{u}) &= -(\nabla\mathbf{u} \times \mathbf{f}_k, \mathbf{u} \times \nabla\mathbf{f}_k), \end{aligned}$$

hence, by Itô's formula, we also have for all $t > 0$, P-almost surely,

$$\begin{aligned} d\|\nabla\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + 2\left(\|\Delta\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + \|\nabla\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + \int_{\mathbb{R}}|\mathbf{u}(t)|^2|\nabla\mathbf{u}(t)|^2 dx\right) dt \\ + 4\int_{\mathbb{R}}|\mathbf{u}(t)\nabla\mathbf{u}(t)|^2 dx dt = 2\sum_{k=1}^{\infty}(\nabla\mathbf{u}(t), \mathbf{u}(t) \times \nabla\mathbf{f}_k + \nabla\mathbf{f}_k) dW_k \\ + \sum_{k=1}^{\infty}(\nabla\mathbf{u}(t), (\mathbf{u}(t) \times \mathbf{f}_k) \times \nabla\mathbf{f}_k + \mathbf{f}_k \times \nabla\mathbf{f}_k)_{L^2(\mathbb{R})} dt \\ + \sum_{k=1}^{\infty}(\mathbf{u}(t) \times \nabla\mathbf{f}_k + \nabla\mathbf{f}_k, \nabla\mathbf{u}(t) \times \mathbf{f}_k + \mathbf{u}(t) \times \nabla\mathbf{f}_k + \nabla\mathbf{f}_k)_{L^2(\mathbb{R})} dt, \end{aligned}$$

and then

$$\begin{aligned} \frac{d}{dt}\mathbf{E}\left(\|\nabla\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2\right) + 2\mathbf{E}\left(\|\Delta\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + \|\nabla\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + \int_{\mathbb{R}}|\mathbf{u}(t)|^2|\nabla\mathbf{u}(t)|^2 dx\right) \\ \leq \sum_{k=1}^{\infty}\mathbf{E}(\nabla\mathbf{u}(t), (\mathbf{u}(t) \times \mathbf{f}_k) \times \nabla\mathbf{f}_k + \mathbf{f}_k \times \nabla\mathbf{f}_k)_{L^2(\mathbb{R})} \\ + \sum_{k=1}^{\infty}\mathbf{E}(\mathbf{u}(t) \times \nabla\mathbf{f}_k + \nabla\mathbf{f}_k, \nabla\mathbf{u}(t) \times \mathbf{f}_k + \mathbf{u}(t) \times \nabla\mathbf{f}_k + \nabla\mathbf{f}_k)_{L^2(\mathbb{R})}. \end{aligned} \quad (4.5)$$

For the first term on the right-hand side of (4.5) we have

$$\begin{aligned} \sum_{k=1}^{\infty}\mathbf{E}(\nabla\mathbf{u}(t), (\mathbf{u}(t) \times \mathbf{f}_k) \times \nabla\mathbf{f}_k + \mathbf{f}_k \times \nabla\mathbf{f}_k)_{L^2(\mathbb{R})} \\ \leq \sum_{k=1}^{\infty}\mathbf{E}\left(\|\mathbf{f}_k\|_{L^\infty(\mathbb{R})}\|\nabla\mathbf{f}_k\|_{L^\infty(\mathbb{R})}\|\nabla\mathbf{u}(t)\|_{L^2(\mathbb{R})}\|\mathbf{u}(t)\|_{L^2(\mathbb{R})}\right) \\ + \sum_{k=1}^{\infty}\mathbf{E}\left(\|\mathbf{f}_k\|_{L^\infty(\mathbb{R})}\|\nabla\mathbf{f}_k\|_{L^2(\mathbb{R})}\|\nabla\mathbf{u}(t)\|_{L^2(\mathbb{R})}\right) \end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{2} \mathbb{E} \left(\|\nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) + \left(\sum_{k=1}^{\infty} \|\mathbf{f}_k\|_{W^{1,\infty}(\mathbb{R})}^2 \right)^2 \mathbb{E} \left(\|\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) \\
&\quad + \left(\sum_{k=1}^{\infty} \|\mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\nabla \mathbf{f}_k\|_{L^2(\mathbb{R})} \right)^2. \tag{4.6}
\end{aligned}$$

For the last term on the right-hand side of (4.5) we have

$$\begin{aligned}
&\sum_{k=1}^{\infty} \mathbb{E} (\mathbf{u}(t) \times \nabla \mathbf{f}_k + \nabla \mathbf{f}_k, \nabla \mathbf{u}(t) \times \mathbf{f}_k + \mathbf{u}(t) \times \nabla \mathbf{f}_k + \nabla \mathbf{f}_k)_{L^2(\mathbb{R})} \\
&\leq \sum_{k=1}^{\infty} \mathbb{E} \left(\|\mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\nabla \mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\mathbf{u}\|_{L^2(\mathbb{R})} \|\nabla \mathbf{u}\|_{L^2(\mathbb{R})} \right) \\
&+ \sum_{k=1}^{\infty} \mathbb{E} \left(\|\mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\nabla \mathbf{f}_k\|_{L^2(\mathbb{R})} \|\nabla \mathbf{u}\|_{L^2(\mathbb{R})} + \|\nabla \mathbf{f}_k\|_{L^\infty(\mathbb{R})}^2 \|\mathbf{u}\|_{L^2(\mathbb{R})}^2 \right) \\
&\quad + 2 \sum_{k=1}^{\infty} \mathbb{E} \left(\|\nabla \mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\nabla \mathbf{f}_k\|_{L^2(\mathbb{R})} \|\mathbf{u}\|_{L^2(\mathbb{R})} + \|\nabla \mathbf{f}_k\|_{L^2(\mathbb{R})}^2 \right) \\
&\leq \frac{1}{2} \mathbb{E} \left(\|\nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) + 2 \left(\sum_{k=1}^{\infty} \|\mathbf{f}_k\|_{W^{1,\infty}(\mathbb{R})}^2 \right)^2 \mathbb{E} \left(\|\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) \\
&+ 2 \left(\sum_{k=1}^{\infty} \|\mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\nabla \mathbf{f}_k\|_{L^2(\mathbb{R})} \right)^2 + \mathbb{E} \left(\|\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) + 2 \sum_{k=1}^{\infty} \|\nabla \mathbf{f}_k\|_{L^2(\mathbb{R})}^2. \tag{4.7}
\end{aligned}$$

It follows from (4.5)-(4.7) that

$$\begin{aligned}
&\frac{d}{dt} \mathbb{E} \left(\|\nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) + \mathbb{E} \left(2\|\Delta \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + \|\nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + 2 \int_{\mathbb{R}} |\mathbf{u}(t)|^2 |\nabla \mathbf{u}(t)|^2 dx \right) \\
&\leq c_1 \mathbb{E} \left(\|\mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) + c_2, \tag{4.8}
\end{aligned}$$

where c_1 and c_2 are positive numbers depending only on $\{\mathbf{f}_k\}_{k=1}^{\infty}$. By Gronwall's inequality we get for all $t \geq 0$,

$$\begin{aligned}
&\mathbb{E} \left(\|\nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) + 2 \int_0^t e^{s-t} \mathbb{E} \left(\|\Delta \mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 + \int_{\mathbb{R}} |\mathbf{u}(s)|^2 |\nabla \mathbf{u}(s)|^2 dx \right) ds \\
&\leq e^{-t} \|\nabla \mathbf{u}_0\|_{L^2(\mathbb{R})}^2 + c_1 \int_0^t e^{s-t} \mathbb{E} \left(\|\mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 \right) ds + c_2. \tag{4.9}
\end{aligned}$$

By (4.4) and (4.9) we get for all $t \geq 0$,

$$\begin{aligned}
&\mathbb{E} \left(\|\nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) + 2 \int_0^t e^{s-t} \mathbb{E} \left(\|\Delta \mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 + \int_{\mathbb{R}} |\mathbf{u}(s)|^2 |\nabla \mathbf{u}(s)|^2 dx \right) ds \\
&\leq c_3 e^{-t} \|\mathbf{u}_0\|_{H^1(\mathbb{R})}^2 + c_3,
\end{aligned}$$

where $c_3 > 0$ depends only on $\{\mathbf{f}_k\}_{k=1}^{\infty}$, which along with (4.4) yields (4.1).

Next, we prove (4.2). By integrating (4.3) on $(0, t)$ we obtain

$$2 \int_0^t \mathbb{E} \left(\|\nabla \mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 + \|\mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 \right) ds \leq \mathbb{E} \left(\|\mathbf{u}_0\|_{L^2(\mathbb{R})}^2 \right) + t \sum_{k=1}^{\infty} \|f_k\|_{L^2(\mathbb{R})}^2. \quad (4.10)$$

On the other hand, by integrating (4.8) on $(0, t)$ we get

$$\begin{aligned} & 2 \int_0^t \mathbb{E} \left(\|\Delta \mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 + \int_{\mathbb{R}} |\mathbf{u}(s)|^2 |\nabla \mathbf{u}(s)|^2 dx \right) ds \\ & \leq \mathbb{E} \left(\|\nabla \mathbf{u}_0\|_{L^2(\mathbb{R})}^2 \right) + c_1 \int_0^t \mathbb{E} \left(\|\mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 \right) ds + c_2 t. \end{aligned} \quad (4.11)$$

It follows from (4.10)-(4.11) that for all $t \geq 0$,

$$\int_0^t \mathbb{E} \left(\|\mathbf{u}(s)\|_{H^2(\mathbb{R})}^2 + \int_{\mathbb{R}} |\mathbf{u}(s)|^2 |\nabla \mathbf{u}(s)|^2 dx \right) ds \leq c_4 \|\mathbf{u}_0\|_{H^1(\mathbb{R})}^2 + c_4 t,$$

where $c_4 > 0$ depends only on $\{\mathbf{f}_k\}_{k=1}^{\infty}$, which implies (4.2) and thus completes the proof. \square

As an immediate consequence of Lemma 4.1, we obtain the following uniform estimates which are useful when studying the tightness of the set of all invariant measures of (1.3).

Lemma 4.2. *If (1.1) holds, then for every $R > 0$, there exists $T = T(R) > 0$ such that for all $t \geq T$, the solution $\mathbf{u}(t, \mathbf{u}_0)$ of (1.3) with $\|\mathbf{u}_0\|_{H^1(\mathbb{R})} \leq R$ satisfies*

$$\mathbb{E} \left(\|\mathbf{u}(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 \right) + \int_0^t e^{s-t} \mathbb{E} \left(\|\mathbf{u}(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 + \int_{\mathbb{R}} |\mathbf{u}(s, \mathbf{u}_0)|^2 |\nabla \mathbf{u}(s, \mathbf{u}_0)|^2 dx \right) ds \leq C,$$

and

$$\frac{1}{t} \int_0^t \mathbb{E} \left(\|\mathbf{u}(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 + \int_{\mathbb{R}} |\mathbf{u}(s, \mathbf{u}_0)|^2 |\nabla \mathbf{u}(s, \mathbf{u}_0)|^2 dx \right) ds \leq C,$$

where $C > 0$ is a constant depending only on $\{\mathbf{f}_k\}_{k=1}^{\infty}$.

We now derive the uniform estimates on the tails of solutions in $L^2(\mathbb{R})$.

Lemma 4.3. *If (1.1) holds, then for every $\varepsilon > 0$ and $\mathbf{u}_0 \in H^1(\mathbb{R})$, there exists $M = M(\varepsilon, \mathbf{u}_0) \geq 1$ such that for all $m \geq M$ and $t \geq 0$, the solution $\mathbf{u}(t, \mathbf{u}_0)$ of (1.3) satisfies*

$$\mathbb{E} \left(\int_{|x|>m} |\mathbf{u}(t, \mathbf{u}_0)(x)|^2 dx \right) < \varepsilon.$$

Proof. As before, let θ be the cut-off function given by (2.1), $\phi(x) = 1 - \theta(x)$, and $\phi_m(x) = \phi(\frac{x}{m})$ for $x \in \mathbb{R}$ and $m \in \mathbb{N}$.

By (1.3) and Itô's formula, we get for all $t \geq 0$, P-almost surely,

$$\begin{aligned} & \|\phi_m \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 - \|\phi_m \mathbf{u}_0\|_{L^2(\mathbb{R})}^2 \\ & = 2 \int_0^t (\Delta \mathbf{u}(s), \phi_m^2 \mathbf{u}(s))_{L^2(\mathbb{R})} ds + 2 \int_0^t (\mathbf{u}(s) \times \Delta \mathbf{u}(s), \phi_m^2 \mathbf{u}(s))_{L^2(\mathbb{R})} ds \end{aligned}$$

$$\begin{aligned}
& -2 \int_0^t ((1 + |\mathbf{u}(s)|^2)\mathbf{u}(s), \phi_m^2 \mathbf{u}(s))_{L^2(\mathbb{R})} ds + 2 \sum_{k=1}^{\infty} \int_0^t (\phi_m \mathbf{f}_k, \phi_m \mathbf{u}(s))_{L^2(\mathbb{R})} dW_k \\
& + \sum_{k=1}^{\infty} \int_0^t \|\phi_m(\mathbf{u}(s) \times \mathbf{f}_k + \mathbf{f}_k)\|_{L^2(\mathbb{R})}^2 ds + \sum_{k=1}^{\infty} \int_0^t ((\mathbf{u}(s) \times \mathbf{f}_k) \times \mathbf{f}_k, \phi_m^2 \mathbf{u}(s))_{L^2(\mathbb{R})} ds. \quad (4.12)
\end{aligned}$$

By (4.12) we get for all $t > 0$, P-almost surely,

$$\begin{aligned}
& \frac{d}{dt} \mathbb{E} \left(\|\phi_m \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) - 2\mathbb{E} \left(\Delta \mathbf{u}(t), \phi_m^2 \mathbf{u}(t) \right)_{L^2(\mathbb{R})} + 2\mathbb{E} \left((1 + |\mathbf{u}(t)|^2)\mathbf{u}(t), \phi_m^2 \mathbf{u}(t) \right)_{L^2(\mathbb{R})} \\
& = \sum_{k=1}^{\infty} \mathbb{E} \left(\|\phi_m(\mathbf{u}(t) \times \mathbf{f}_k + \mathbf{f}_k)\|_{L^2(\mathbb{R})}^2 \right) + \sum_{k=1}^{\infty} \mathbb{E} \left((\mathbf{u}(t) \times \mathbf{f}_k) \times \mathbf{f}_k, \phi_m^2 \mathbf{u}(t) \right)_{L^2(\mathbb{R})}. \quad (4.13)
\end{aligned}$$

Similar to (2.8) we have

$$2\mathbb{E} \left(\Delta \mathbf{u}(t), \phi_m^2 \mathbf{u}(t) \right)_{L^2(\mathbb{R})} \leq \frac{c_1}{m} \mathbb{E} \left(\|\mathbf{u}(t)\|_{H^1(\mathbb{R})}^2 \right), \quad (4.14)$$

where $c_1 > 0$ is a constant independent of m . For the last two terms on the right-hand side of (4.13), we have

$$\sum_{k=1}^{\infty} \mathbb{E} \left(\|\phi_m(\mathbf{u}(t) \times \mathbf{f}_k + \mathbf{f}_k)\|_{L^2(\mathbb{R})}^2 \right) = \sum_{k=1}^{\infty} \mathbb{E} \left(\|\phi_m(\mathbf{u}(t) \times \mathbf{f}_k)\|_{L^2(\mathbb{R})}^2 + \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})}^2 \right),$$

and

$$\sum_{k=1}^{\infty} \mathbb{E} \left((\mathbf{u}(t) \times \mathbf{f}_k) \times \mathbf{f}_k, \phi_m^2 \mathbf{u}(t) \right)_{L^2(\mathbb{R})} = - \sum_{k=1}^{\infty} \mathbb{E} \left(\|\phi_m(\mathbf{u}(t) \times \mathbf{f}_k)\|_{L^2(\mathbb{R})}^2 \right).$$

Taking sum we have

$$\sum_{k=1}^{\infty} \mathbb{E} \left(\|\phi_m(\mathbf{u}(t) \times \mathbf{f}_k + \mathbf{f}_k)\|_{L^2(\mathbb{R})}^2 \right) + \sum_{k=1}^{\infty} \mathbb{E} \left((\mathbf{u}(t) \times \mathbf{f}_k) \times \mathbf{f}_k, \phi_m^2 \mathbf{u}(t) \right)_{L^2(\mathbb{R})} = \sum_{k=1}^{\infty} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})}^2. \quad (4.15)$$

By (4.13)-(4.15) we obtain for all $t > 0$ and $m \in \mathbb{N}$,

$$\frac{d}{dt} \mathbb{E} \left(\|\phi_m \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) + 2\mathbb{E} \left(\|\phi_m \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) \leq \frac{c_1}{m} \mathbb{E} \left(\|\mathbf{u}(t)\|_{H^1(\mathbb{R})}^2 \right) + \sum_{k=1}^{\infty} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})}^2. \quad (4.16)$$

Solving (4.16) we get for all $t \geq 0$ and $m \in \mathbb{N}$,

$$\begin{aligned}
& \mathbb{E} \left(\|\phi_m \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) + \int_0^t e^{s-t} \mathbb{E} \left(\|\phi_m \mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 \right) ds \\
& \leq e^{-t} \mathbb{E} \left(\|\phi_m \mathbf{u}_0\|_{L^2(\mathbb{R})}^2 \right) + \frac{c_1}{m} \int_0^t e^{s-t} \mathbb{E} \left(\|\mathbf{u}(s)\|_{H^1(\mathbb{R})}^2 \right) ds + \sum_{k=1}^{\infty} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})}^2. \quad (4.17)
\end{aligned}$$

By (4.17) and Lemma 4.1 we obtain for all $t \geq 0$ and $m \in \mathbb{N}$,

$$\begin{aligned}
& \mathbb{E} \left(\|\phi_m \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) + \int_0^t e^{s-t} \mathbb{E} \left(\|\phi_m \mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 \right) ds \\
& \leq \|\phi_m \mathbf{u}_0\|_{L^2(\mathbb{R})}^2 + \frac{c_2}{m} (1 + \|\mathbf{u}_0\|_{H^1(\mathbb{R})}^2) + \sum_{k=1}^{\infty} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})}^2,
\end{aligned}$$

where $c_2 > 0$ depends only on $\{\mathbf{f}_k\}_{k=1}^\infty$. Since $\sum_{k=1}^\infty \|\mathbf{f}_k\|_{L^2(\mathbb{R})}^2 < \infty$, we see that

$$\|\phi_m \mathbf{u}_0\|_{L^2(\mathbb{R})}^2 + \sum_{k=1}^\infty \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})}^2 \leq \int_{|x| > \frac{1}{2}m} |\mathbf{u}_0(x)|^2 dx + \int_{|x| > \frac{1}{2}m} \sum_{k=1}^\infty |\mathbf{f}_k(x)|^2 dx \rightarrow 0,$$

as $m \rightarrow \infty$, which together with (4.17) indicates that for every $\varepsilon > 0$, there exists $M = M(\varepsilon, \mathbf{u}_0) \geq 1$, such that for all $m \geq M$ and $t \geq 0$,

$$\begin{aligned} & \mathbb{E} \left(\int_{|x| > \frac{3}{4}m} |\mathbf{u}(t, \mathbf{u}_0)(x)|^2 dx \right) + \int_0^t e^{s-t} \mathbb{E} \left(\|\phi_m \mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 \right) ds \\ & \leq \mathbb{E} \left(\|\phi_m \mathbf{u}(t, \mathbf{u}_0)\|_{L^2(\mathbb{R})}^2 \right) + \int_0^t e^{s-t} \mathbb{E} \left(\|\phi_m \mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 \right) ds < \varepsilon, \end{aligned} \quad (4.18)$$

which completes the proof. \square

Following the proof of Lemma 4.3, by using Lemma 4.2 instead of Lemma 4.1, we obtain the following uniform tail-ends estimates.

Lemma 4.4. *If (1.1) holds, then for every $\varepsilon > 0$ and $R > 0$, there exist $T = T(\varepsilon, R) > 0$ and $M = M(\varepsilon) \geq 1$ such that for all $t \geq T$ and $m \geq M$, the solution $\mathbf{u}(t, \mathbf{u}_0)$ of (1.3) with $\|\mathbf{u}_0\|_{H^1(\mathbb{R})} \leq R$, satisfies*

$$\mathbb{E} \left(\int_{|x| > m} |\mathbf{u}(t, \mathbf{u}_0)(x)|^2 dx \right) < \varepsilon.$$

Next, we improve Lemma 4.3 and derive the uniform estimates on the tails of solutions in $H^1(\mathbb{R})$.

Lemma 4.5. *If (1.1) holds, then for every $\varepsilon > 0$, and $\mathbf{u}_0 \in H^1(\mathbb{R})$, there exists $M = M(\varepsilon, \mathbf{u}_0) \geq 1$ such that for all $m \geq M$, the solution $\mathbf{u}(t, \mathbf{u}_0)$ of (1.3) satisfies*

$$\sup_{t \geq 0} \mathbb{E} \left(\int_{|x| > m} (|\mathbf{u}(t, \mathbf{u}_0)(x)|^2 + |\nabla \mathbf{u}(t, \mathbf{u}_0)(x)|^2) dx \right) < \varepsilon.$$

Proof. Let ϕ_m be the same cut-off function as in the proof of Lemma 4.3. By (1.3) and Itô's formula, we get

$$\begin{aligned} d\|\phi_m \nabla \mathbf{u}\|_{L^2(\mathbb{R})}^2 &= -2 (\Delta \mathbf{u}, \operatorname{div}(\phi_m^2 \nabla \mathbf{u}))_{L^2(\mathbb{R})} dt - 2 (\mathbf{u} \times \Delta \mathbf{u}, \operatorname{div}(\phi_m^2 \nabla \mathbf{u}))_{L^2(\mathbb{R})} dt \\ &\quad - 2(\phi_m \nabla(1 + |\mathbf{u}|^2) \mathbf{u}, \phi_m \nabla \mathbf{u})_{L^2(\mathbb{R})} dt \\ &\quad + \sum_{k=1}^\infty (\phi_m \nabla((\mathbf{u} \times \mathbf{f}_k) \times \mathbf{f}_k), \phi_m \nabla \mathbf{u})_{L^2(\mathbb{R})} dt \\ &\quad + 2 \sum_{k=1}^\infty (\phi_m \nabla(\mathbf{u} \times \mathbf{f}_k + \mathbf{f}_k), \phi_m \nabla \mathbf{u})_{L^2(\mathbb{R})} dW_k \\ &\quad + \sum_{k=1}^\infty \|\phi_m \nabla(\mathbf{u} \times \mathbf{f}_k + \mathbf{f}_k)\|_{L^2(\mathbb{R})}^2 dt, \end{aligned}$$

and hence for all $t > 0$, P-almost surely,

$$\begin{aligned}
\frac{d}{dt} \mathbb{E} \left(\|\phi_m \nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) &= -2\mathbb{E} \left((\Delta \mathbf{u}(t), \operatorname{div} (\phi_m^2 \nabla \mathbf{u}(t)))_{L^2(\mathbb{R})} \right) \\
&\quad - 2\mathbb{E} \left((\mathbf{u}(t) \times \Delta \mathbf{u}(t), \operatorname{div} (\phi_m^2 \nabla \mathbf{u}(t)))_{L^2(\mathbb{R})} \right) \\
&\quad - 2\mathbb{E} \left((\phi_m \nabla(1 + |\mathbf{u}(t)|^2) \mathbf{u}(t), \phi_m \nabla \mathbf{u}(t))_{L^2(\mathbb{R})} \right) \\
&\quad + \sum_{k=1}^{\infty} \mathbb{E} \left((\phi_m \nabla((\mathbf{u}(t) \times \mathbf{f}_k) \times \mathbf{f}_k), \phi_m \nabla \mathbf{u}(t))_{L^2(\mathbb{R})} \right) \\
&\quad + \sum_{k=1}^{\infty} \mathbb{E} \left(\|\phi_m \nabla(\mathbf{u}(t) \times \mathbf{f}_k + \mathbf{f}_k)\|_{L^2(\mathbb{R})}^2 \right). \tag{4.19}
\end{aligned}$$

For the first term on the right-hand side of (4.19), by Hölder's inequality we have

$$-2\mathbb{E} \left((\Delta \mathbf{u}(t), \operatorname{div} (\phi_m^2 \nabla \mathbf{u}(t)))_{L^2(\mathbb{R})} \right) \leq -2\mathbb{E} \left(\|\phi_m \Delta \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) + \frac{c_1}{m} \|\mathbf{u}(t)\|_{H^2}^2, \tag{4.20}$$

where $c_1 > 0$ is independent of m .

For the second term on the right-hand side of (4.19) we have

$$\begin{aligned}
&-2\mathbb{E} \left((\mathbf{u}(t) \times \Delta \mathbf{u}(t), \operatorname{div} (\phi_m^2 \nabla \mathbf{u}(t)))_{L^2(\mathbb{R})} \right) \\
&= -2\mathbb{E} \left((\mathbf{u}(t) \times \Delta \mathbf{u}(t), \phi_m^2 \Delta \mathbf{u}(t))_{L^2(\mathbb{R})} \right) - 2\mathbb{E} \left((\mathbf{u}(t) \times \Delta \mathbf{u}(t), (\nabla \phi_m^2) \nabla \mathbf{u}(t))_{L^2(\mathbb{R})} \right) \\
&= -2\mathbb{E} \left((\mathbf{u}(t) \times \Delta \mathbf{u}(t), (\nabla \phi_m^2) \nabla \mathbf{u}(t))_{L^2(\mathbb{R})} \right) \\
&\leq \frac{c_2}{m} \mathbb{E} \left(\|\Delta \mathbf{u}(t)\|_{L^2(\mathbb{R})} \left(\int_{\mathbb{R}} |\mathbf{u}(t)|^2 |\nabla \mathbf{u}(t)|^2 dx \right)^{\frac{1}{2}} \right) \\
&\leq \frac{c_2}{2m} \mathbb{E} \left(\|\Delta \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) + \frac{c_2}{2m} \mathbb{E} \left(\int_{\mathbb{R}} |\mathbf{u}(t)|^2 |\nabla \mathbf{u}(t)|^2 dx \right), \tag{4.21}
\end{aligned}$$

where $c_2 > 0$ is independent of m .

For the third term on the right-hand side of (4.19) we have

$$\begin{aligned}
&-2\mathbb{E} \left((\phi_m \nabla(1 + |\mathbf{u}(t)|^2) \mathbf{u}(t), \phi_m \nabla \mathbf{u}(t))_{L^2(\mathbb{R})} \right) \\
&= -2\mathbb{E} \left(\|\phi_m \nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + \int_{\mathbb{R}} \phi_m^2 |\mathbf{u}(t)|^2 |\nabla \mathbf{u}(t)|^2 dx \right) - 4\mathbb{E} \left(\int_{\mathbb{R}} \phi_m^2 |\mathbf{u}(t) \nabla \mathbf{u}(t)|^2 dx \right) \\
&\leq -2\mathbb{E} \left(\|\phi_m \nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right). \tag{4.22}
\end{aligned}$$

For the last two terms on the right-hand side of (4.19) we have

$$\begin{aligned}
&\sum_{k=1}^{\infty} \mathbb{E} \left((\phi_m \nabla((\mathbf{u}(t) \times \mathbf{f}_k) \times \mathbf{f}_k), \phi_m \nabla \mathbf{u}(t))_{L^2(\mathbb{R})} \right) + \sum_{k=1}^{\infty} \mathbb{E} \left(\|\phi_m \nabla(\mathbf{u}(t) \times \mathbf{f}_k + \mathbf{f}_k)\|_{L^2(\mathbb{R})}^2 \right) \\
&= \sum_{k=1}^{\infty} \mathbb{E} \left((\phi_m^2 \nabla \mathbf{u}(t), (\mathbf{u}(t) \times \mathbf{f}_k) \times \nabla \mathbf{f}_k + \mathbf{f}_k \times \nabla \mathbf{f}_k)_{L^2(\mathbb{R})} \right)
\end{aligned}$$

$$+ \sum_{k=1}^{\infty} \mathbb{E} \left((\phi_m^2(\mathbf{u}(t) \times \nabla \mathbf{f}_k + \nabla \mathbf{f}_k), \nabla \mathbf{u}(t) \times \mathbf{f}_k + \mathbf{u}(t) \times \nabla \mathbf{f}_k + \nabla \mathbf{f}_k)_{L^2(\mathbb{R})} \right). \quad (4.23)$$

For the first term on the right-hand side of (4.23) we have

$$\begin{aligned} & \sum_{k=1}^{\infty} \mathbb{E} (\phi_m^2 \nabla \mathbf{u}(t), (\mathbf{u}(t) \times \mathbf{f}_k) \times \nabla \mathbf{f}_k + \mathbf{f}_k \times \nabla \mathbf{f}_k)_{L^2(\mathbb{R})} \\ & \leq \sum_{k=1}^{\infty} \mathbb{E} (\|\mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\nabla \mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\phi_m \nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})} \|\phi_m \mathbf{u}(t)\|_{L^2(\mathbb{R})}) \\ & \quad + \sum_{k=1}^{\infty} \mathbb{E} (\|\nabla \mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})} \|\phi_m \nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}) \\ & \leq \frac{1}{2} \mathbb{E} (\phi_m \|\nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2) + \left(\sum_{k=1}^{\infty} \|\mathbf{f}_k\|_{W^{1,\infty}(\mathbb{R})}^2 \right)^2 \mathbb{E} (\|\phi_m \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2) \\ & \quad + \left(\sum_{k=1}^{\infty} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})} \|\nabla \mathbf{f}_k\|_{L^\infty(\mathbb{R})} \right)^2. \end{aligned} \quad (4.24)$$

For the second term on the right-hand side of (4.23) we have

$$\begin{aligned} & \sum_{k=1}^{\infty} \mathbb{E} (\phi_m^2(\mathbf{u}(t) \times \nabla \mathbf{f}_k + \nabla \mathbf{f}_k), \nabla \mathbf{u}(t) \times \mathbf{f}_k + \mathbf{u}(t) \times \nabla \mathbf{f}_k + \nabla \mathbf{f}_k)_{L^2(\mathbb{R})} \\ & \leq \sum_{k=1}^{\infty} \mathbb{E} (\|\mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\nabla \mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\phi_m \mathbf{u}\|_{L^2(\mathbb{R})} \|\phi_m \nabla \mathbf{u}\|_{L^2(\mathbb{R})}) \\ & + \sum_{k=1}^{\infty} \mathbb{E} (\|\nabla \mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})} \|\phi_m \nabla \mathbf{u}\|_{L^2(\mathbb{R})} + \|\nabla \mathbf{f}_k\|_{L^\infty(\mathbb{R})}^2 \|\phi_m \mathbf{u}\|_{L^2(\mathbb{R})}^2) \\ & \quad + 2 \sum_{k=1}^{\infty} \mathbb{E} (\|\nabla \mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\phi_m \nabla \mathbf{f}_k\|_{L^2(\mathbb{R})} \|\phi_m \mathbf{u}\|_{L^2(\mathbb{R})} + \|\phi_m \nabla \mathbf{f}_k\|_{L^2(\mathbb{R})}^2) \\ & \leq \frac{1}{2} \mathbb{E} (\|\phi_m \nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2) + 2 \left(\sum_{k=1}^{\infty} \|\mathbf{f}_k\|_{W^{1,\infty}(\mathbb{R})}^2 \right)^2 \mathbb{E} (\|\phi_m \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2) \\ & + 2 \left(\sum_{k=1}^{\infty} \|\nabla \mathbf{f}_k\|_{L^\infty(\mathbb{R})} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})} \right)^2 + \mathbb{E} (\|\phi_m \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2) + 2 \sum_{k=1}^{\infty} \|\phi_m \nabla \mathbf{f}_k\|_{L^2(\mathbb{R})}^2. \end{aligned} \quad (4.25)$$

By (4.23)-(4.25) we get

$$\begin{aligned} & \sum_{k=1}^{\infty} \mathbb{E} ((\phi_m \nabla((\mathbf{u}(t) \times \mathbf{f}_k) \times \mathbf{f}_k), \phi_m \nabla \mathbf{u}(t))_{L^2(\mathbb{R})}) + \sum_{k=1}^{\infty} \mathbb{E} (\|\phi_m \nabla(\mathbf{u}(t) \times \mathbf{f}_k + \mathbf{f}_k)\|_{L^2(\mathbb{R})}^2) \\ & \leq \mathbb{E} (\|\phi_m \nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2) + c_3 \mathbb{E} (\|\phi_m \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2) + 2 \sum_{k=1}^{\infty} \|\phi_m \nabla \mathbf{f}_k\|_{L^2(\mathbb{R})}^2 \end{aligned}$$

$$+3 \left(\sum_{k=1}^{\infty} \|\nabla \mathbf{f}_k\|_{L^\infty(\mathbb{R})}^2 \right) \left(\sum_{k=1}^{\infty} \|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})}^2 \right), \quad (4.26)$$

where $c_3 > 0$ depends only on $\{\mathbf{f}_k\}_{k=1}^{\infty}$.

It follows from (4.19)-(4.22) and (4.26) that

$$\begin{aligned} & \frac{d}{dt} \mathbf{E} \left(\|\phi_m \nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) + \mathbf{E} \left(2\|\phi_m \Delta \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 + \|\phi_m \nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) \\ & \leq \frac{c_4}{m} \mathbf{E} \left(\|\mathbf{u}(t)\|_{H^2(\mathbb{R})}^2 + \int_{\mathbb{R}} |\mathbf{u}(t)|^2 |\nabla \mathbf{u}(t)|^2 dx \right) + c_4 \mathbf{E} \left(\|\phi_m \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) \\ & \quad + c_4 \sum_{k=1}^{\infty} \left(\|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})}^2 + \|\phi_m \nabla \mathbf{f}_k\|_{L^2(\mathbb{R})}^2 \right), \end{aligned}$$

where $c_4 > 0$ depends only on $\{\mathbf{f}_k\}_{k=1}^{\infty}$. Solving the inequality, we get for all $t \geq 0$,

$$\begin{aligned} & \mathbf{E} \left(\|\phi_m \nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) \leq e^{-t} \mathbf{E} \left(\|\phi_m \nabla \mathbf{u}_0\|_{L^2(\mathbb{R})}^2 \right) \\ & \quad + \frac{c_4}{m} \int_0^t e^{s-t} \mathbf{E} \left(\|\mathbf{u}(s)\|_{H^2(\mathbb{R})}^2 + \int_{\mathbb{R}} |\mathbf{u}(s)|^2 |\nabla \mathbf{u}(s)|^2 dx \right) ds \\ & \quad + c_4 \int_0^t e^{s-t} \mathbf{E} \left(\|\phi_m \mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 \right) ds + c_4 \sum_{k=1}^{\infty} \left(\|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})}^2 + \|\phi_m \nabla \mathbf{f}_k\|_{L^2(\mathbb{R})}^2 \right). \end{aligned} \quad (4.27)$$

By (4.27) and Lemma 4.1 we obtain for all $t \geq 0$ and $m \in \mathbb{N}$,

$$\begin{aligned} & \mathbf{E} \left(\|\phi_m \nabla \mathbf{u}(t)\|_{L^2(\mathbb{R})}^2 \right) \leq \|\phi_m \nabla \mathbf{u}_0\|_{L^2(\mathbb{R})}^2 + \frac{c_5}{m} (1 + \|\mathbf{u}_0\|_{H^1(\mathbb{R})}^2) \\ & \quad + c_4 \int_0^t e^{s-t} \mathbf{E} \left(\|\phi_m \mathbf{u}(s)\|_{L^2(\mathbb{R})}^2 \right) ds + c_4 \sum_{k=1}^{\infty} \left(\|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})}^2 + \|\phi_m \nabla \mathbf{f}_k\|_{L^2(\mathbb{R})}^2 \right), \end{aligned} \quad (4.28)$$

where $c_5 > 0$ depends only on $\{\mathbf{f}_k\}_{k=1}^{\infty}$. By (1.1) we find that

$$\begin{aligned} & \|\phi_m \mathbf{u}_0\|_{H^1(\mathbb{R})}^2 + c_4 \sum_{k=1}^{\infty} \left(\|\phi_m \mathbf{f}_k\|_{L^2(\mathbb{R})}^2 + \|\phi_m \nabla \mathbf{f}_k\|_{L^2(\mathbb{R})}^2 \right) \\ & \leq \int_{|x| > \frac{1}{2}m} (|\mathbf{u}_0(x)|^2 + |\nabla \mathbf{u}_0(x)|^2) dx + \int_{|x| > \frac{1}{2}m} \sum_{k=1}^{\infty} (|\mathbf{f}_k(x)|^2 + |\nabla \mathbf{f}_k(x)|^2) dx \rightarrow 0, \end{aligned}$$

as $m \rightarrow \infty$, which together with (4.18) and (4.28) implies that for every $\varepsilon > 0$, there exists $M = M(\varepsilon, \mathbf{u}_0) \geq 1$, such that for all $m \geq M$ and $t \geq 0$,

$$\mathbf{E} \left(\int_{|x| > \frac{3}{4}m} |\nabla \mathbf{u}(t, \mathbf{u}_0)(x)|^2 dx \right) \leq \mathbf{E} \left(\|\phi_m \nabla \mathbf{u}(t, \mathbf{u}_0)\|_{L^2(\mathbb{R})}^2 \right) < \varepsilon,$$

which together with Lemma 4.3 completes the proof. \square

By Lemma 4.4 and the argument of Lemma 4.5, we also have the following uniform estimates.

Lemma 4.6. *If (1.1) holds, then for every $\varepsilon > 0$ and $R > 0$, there exist $T = T(\varepsilon, R) > 0$ and $M = M(\varepsilon) \geq 1$ such that for all $t \geq T$ and $m \geq M$, the solution $\mathbf{u}(t, \mathbf{u}_0)$ of (1.3) with $\|\mathbf{u}_0\|_{H^1(\mathbb{R})} \leq R$, satisfies*

$$\mathbb{E} \left(\int_{|x|>m} (|\mathbf{u}(t, \mathbf{u}_0)(x)|^2 + |\nabla \mathbf{u}(t, \mathbf{u}_0)(x)|^2) dx \right) < \varepsilon.$$

4.1. Feller property. In this section, we establish the Feller property of the Markov transition semigroup $\{\mathbf{P}_t\}_{t \geq 0}$ in terms of the norm topology instead of the weak topology in $H^1(\mathbb{R})$.

Lemma 4.7. *If (1.1) holds, then the family $\{\mathbf{P}_t\}_{t \geq 0}$ is Feller in $H^1(\mathbb{R})$; that is, $\mathbf{P}_t \varphi \in C_b(H^1(\mathbb{R}))$ for every $\varphi \in C_b(H^1(\mathbb{R}))$ and $t \geq 0$.*

Proof. Let $t_0 > 0$, $\varphi \in C_b(H^1(\mathbb{R}))$ and $\mathbf{u}_0, \mathbf{u}_{0,n} \in H^1(\mathbb{R})$ such that $\mathbf{u}_{0,n} \rightarrow \mathbf{u}_0$ in $H^1(\mathbb{R})$. We only need to show $(\mathbf{P}_{t_0} \varphi)(\mathbf{u}_{0,n}) \rightarrow (\mathbf{P}_{t_0} \varphi)(\mathbf{u}_0)$; that is,

$$\lim_{n \rightarrow \infty} \mathbb{E}(\varphi(\mathbf{u}(t_0, \mathbf{u}_{0,n}))) = \mathbb{E}(\varphi(\mathbf{u}(t_0, \mathbf{u}_0))), \quad (4.29)$$

where $\mathbf{u}(\cdot, \mathbf{u}_{0,n})$ and $\mathbf{u}(\cdot, \mathbf{u}_0)$ are solutions of (1.3) with initial data $\mathbf{u}_{0,n}$ and \mathbf{u}_0 , respectively. Since $\mathbf{u}_{0,n} \rightarrow \mathbf{u}_0$ in $H^1(\mathbb{R})$, we see that the sequence $\{\mathbf{u}_{0,n}\}_{n=1}^\infty$ is bounded in $H^1(\mathbb{R})$, and thus by the uniform estimates of solutions we infer that

$$\mathbb{E} \left(\sup_{0 \leq t \leq t_0} \|\mathbf{u}(t, \mathbf{u}_{0,n})\|_{H^1(\mathbb{R})}^2 + \int_0^{t_0} \|\mathbf{u}(s, \mathbf{u}_{0,n})\|_{H^2(\mathbb{R})}^2 ds \right) \leq c, \quad (4.30)$$

and

$$\mathbb{E} \left(\sup_{0 \leq t \leq t_0} \|\mathbf{u}(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 + \int_0^{t_0} \|\mathbf{u}(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 ds \right) \leq c, \quad (4.31)$$

where $c > 0$ is independent of $n \in \mathbb{N}$. It follows from (4.30)-(4.31) that for every $R > 0$,

$$\mathbb{P} \left(\sup_{0 \leq t \leq t_0} \|\mathbf{u}(t, \mathbf{u}_{0,n})\|_{H^1(\mathbb{R})}^2 + \int_0^{t_0} \|\mathbf{u}(s, \mathbf{u}_{0,n})\|_{H^2(\mathbb{R})}^2 ds > R \right) \leq \frac{c}{R},$$

and

$$\mathbb{P} \left(\sup_{0 \leq t \leq t_0} \|\mathbf{u}(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 + \int_0^{t_0} \|\mathbf{u}(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 ds > R \right) \leq \frac{c}{R}.$$

Therefore, for every $\delta > 0$, there exists $R_\delta > 0$ such that for all $n \in \mathbb{N}$,

$$\mathbb{P} \left(\sup_{0 \leq t \leq t_0} \|\mathbf{u}(t, \mathbf{u}_{0,n})\|_{H^1(\mathbb{R})}^2 + \int_0^{t_0} \|\mathbf{u}(s, \mathbf{u}_{0,n})\|_{H^2(\mathbb{R})}^2 ds > R_\delta \right) \leq \frac{1}{2} \delta, \quad (4.32)$$

and

$$\mathbb{P} \left(\sup_{0 \leq t \leq t_0} \|\mathbf{u}(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 + \int_0^{t_0} \|\mathbf{u}(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 ds > R_\delta \right) \leq \frac{1}{2} \delta. \quad (4.33)$$

Given $n \in \mathbb{N}$, define the stopping times:

$$\tau_{1,n} = \inf \left\{ t \geq 0; \|\mathbf{u}(t, \mathbf{u}_{0,n})\|_{H^1(\mathbb{R})}^2 + \int_0^t \|\mathbf{u}(s, \mathbf{u}_{0,n})\|_{H^2(\mathbb{R})}^2 ds > R_\delta \right\},$$

and

$$\tau_{2,n} = \inf \left\{ t \geq 0; \|\mathbf{u}(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 + \int_0^t \|\mathbf{u}(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 ds > R_\delta \right\}.$$

As usual, $\inf \emptyset = +\infty$. Denote by $\tau_n = \tau_{1,n} \wedge \tau_{2,n}$.

Let $\mathbf{v}_n(t) = \mathbf{u}(t, \mathbf{u}_{0,n}) - \mathbf{u}(t, \mathbf{u}_0)$ for $t \geq 0$. Then by (2.46) we get for all $t \geq 0$, P-almost surely,

$$\begin{aligned} & \|\nabla \mathbf{v}_n(t)\|_{L^2(\mathbb{R})}^2 + 2 \int_0^t \|\Delta \mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 ds + 2 \int_0^t \|\nabla \mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 ds \\ &= \|\nabla \mathbf{v}_{0,n}\|_{L^2(\mathbb{R})}^2 - 2 \int_0^t (\Delta \mathbf{v}_n(s), \mathbf{v}_n(s) \times \Delta \mathbf{u}(s, \mathbf{u}_{0,n})) ds \\ & \quad - 2 \int_0^t (\nabla \mathbf{v}_n(s), \nabla (|\mathbf{u}(s, \mathbf{u}_{0,n})|^2 \mathbf{u}(s, \mathbf{u}_{0,n}) - |\mathbf{u}(s, \mathbf{u}_0)|^2 \mathbf{u}(s, \mathbf{u}_0))) ds \\ & \quad - \sum_{k=1}^{\infty} \int_0^t (\Delta \mathbf{v}_n(s), (\mathbf{v}_n(s) \times \mathbf{f}_k) \times \mathbf{f}_k) ds + \sum_{k=1}^{\infty} \int_0^t \|\nabla(\mathbf{v}_n(s) \times \mathbf{f}_k)\|_{L^2(\mathbb{R})}^2 ds \\ & \quad + 2 \sum_{k=1}^{\infty} \int_0^t (\nabla \mathbf{v}_n(s), \mathbf{v}_n(s) \times \nabla \mathbf{f}_k) dW_k. \end{aligned} \quad (4.34)$$

By (4.34) we have for all $t \geq 0$,

$$\begin{aligned} & \mathbb{E} \left(\|\nabla \mathbf{v}_n(t \wedge \tau_n)\|_{L^2(\mathbb{R})}^2 \right) + 2\mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\Delta \mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 ds \right) + 2\mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\nabla \mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 ds \right) \\ &= \mathbb{E} \left(\|\nabla \mathbf{v}_{0,n}\|_{L^2(\mathbb{R})}^2 \right) - 2\mathbb{E} \left(\int_0^{t \wedge \tau_n} (\Delta \mathbf{v}_n(s), \mathbf{v}_n(s) \times \Delta \mathbf{u}(s, \mathbf{u}_{0,n})) ds \right) \\ & \quad - 2\mathbb{E} \left(\int_0^{t \wedge \tau_n} (\nabla \mathbf{v}_n(s), \nabla (|\mathbf{u}(s, \mathbf{u}_{0,n})|^2 \mathbf{u}(s, \mathbf{u}_{0,n}) - |\mathbf{u}(s, \mathbf{u}_0)|^2 \mathbf{u}(s, \mathbf{u}_0))) ds \right) \\ & \quad - \mathbb{E} \left(\sum_{k=1}^{\infty} \int_0^{t \wedge \tau_n} (\Delta \mathbf{v}_n(s), (\mathbf{v}_n(s) \times \mathbf{f}_k) \times \mathbf{f}_k) ds \right) \\ & \quad + \mathbb{E} \left(\sum_{k=1}^{\infty} \int_0^{t \wedge \tau_n} \|\nabla(\mathbf{v}_n(s) \times \mathbf{f}_k)\|_{L^2(\mathbb{R})}^2 ds \right). \end{aligned} \quad (4.35)$$

To deal with the second term on the right-hand side of (4.35), we will use the following embedding inequality:

$$\|\mathbf{u}\|_{L^\infty(\mathbb{R})} \leq c_1 \|\mathbf{u}\|_{L^2(\mathbb{R})}^{\frac{1}{2}} \|\nabla \mathbf{u}\|_{L^2(\mathbb{R})}^{\frac{1}{2}}, \quad \forall \mathbf{u} \in H^1(\mathbb{R}), \quad (4.36)$$

where $c_1 > 0$ is an absolute constant. Note that for all $t \in [0, t_0]$,

$$\begin{aligned} & \left| -2\mathbb{E} \left(\int_0^{t \wedge \tau_n} (\Delta \mathbf{v}_n(s), \mathbf{v}_n(s) \times \Delta \mathbf{u}(s, \mathbf{u}_{0,n})) ds \right) \right| \\ & \leq 2\mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\Delta \mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \|\mathbf{v}_n(s)\|_{L^\infty(\mathbb{R})} \|\Delta \mathbf{u}(s, \mathbf{u}_{0,n})\|_{L^2(\mathbb{R})} ds \right) \\ & \leq \frac{1}{4}\mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\Delta \mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 ds \right) + 4\mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\mathbf{v}_n(s)\|_{L^\infty(\mathbb{R})}^2 \|\Delta \mathbf{u}(s, \mathbf{u}_{0,n})\|_{L^2(\mathbb{R})}^2 ds \right). \end{aligned} \quad (4.37)$$

For the second term on the right-hand side of (4.37), by (4.36) we have for all $t \in [0, t_0]$,

$$\begin{aligned}
& 4\mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\mathbf{v}_n(s)\|_{L^\infty(\mathbb{R})}^2 \|\Delta \mathbf{u}(s, \mathbf{u}_{0,n})\|_{L^2(\mathbb{R})}^2 ds \right) \\
& \leq 4c_1^2 \mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \|\nabla \mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \|\Delta \mathbf{u}(s, \mathbf{u}_{0,n})\|_{L^2(\mathbb{R})}^2 ds \right) \\
& \leq 4c_1^2 \mathbb{E} \left(\sup_{s \in [0, t_0]} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \int_0^{t \wedge \tau_n} \|\nabla \mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \|\Delta \mathbf{u}(s, \mathbf{u}_{0,n})\|_{L^2(\mathbb{R})}^2 ds \right) \\
& \leq 4c_1^2 \mathbb{E} \left(\sup_{s \in [0, t_0]} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \int_0^{t \wedge \tau_n} (\|\mathbf{u}(s, \mathbf{u}_{0,n})\|_{H^1(\mathbb{R})} + \|\mathbf{u}(s, \mathbf{u}_0)\|_{H^1(\mathbb{R})}) \|\Delta \mathbf{u}(s, \mathbf{u}_{0,n})\|_{L^2(\mathbb{R})}^2 ds \right) \\
& \leq 8c_1^2 R_\delta^{\frac{1}{2}} \mathbb{E} \left(\sup_{s \in [0, t_0]} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \int_0^{t \wedge \tau_n} \|\Delta \mathbf{u}(s, \mathbf{u}_{0,n})\|_{L^2(\mathbb{R})}^2 ds \right) \\
& \leq 8c_1^2 R_\delta^{\frac{3}{2}} \mathbb{E} \left(\sup_{s \in [0, t_0]} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \right). \tag{4.38}
\end{aligned}$$

By (4.37)-(4.38) we get for all $t \in [0, t_0]$,

$$\begin{aligned}
& \left| -2\mathbb{E} \left(\int_0^{t \wedge \tau_n} (\Delta \mathbf{v}_n(s), \mathbf{v}_n(s) \times \Delta \mathbf{u}(s, \mathbf{u}_{0,n})) ds \right) \right| \\
& \leq \frac{1}{4} \mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\Delta \mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 ds \right) + 8c_1^2 R_\delta^{\frac{3}{2}} \mathbb{E} \left(\sup_{s \in [0, t_0]} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \right). \tag{4.39}
\end{aligned}$$

For the third term on the right-hand side of (4.35) we have for all $t \in [0, t_0]$,

$$\begin{aligned}
& 2\mathbb{E} \left(\int_0^{t \wedge \tau_n} (\Delta \mathbf{v}_n(s), |\mathbf{u}(s, \mathbf{u}_{0,n})|^2 \mathbf{u}(s, \mathbf{u}_{0,n}) - |\mathbf{u}(s, \mathbf{u}_0)|^2 \mathbf{u}(s, \mathbf{u}_0)) ds \right) \\
& \leq 2\mathbb{E} \left(\int_0^{t \wedge \tau_n} \left(\int_{\mathbb{R}} |\Delta \mathbf{v}_n(s)| |\mathbf{v}_n(s)| (|\mathbf{u}(s, \mathbf{u}_{0,n})|^2 + |\mathbf{u}(s, \mathbf{u}_0)|^2 + |\mathbf{u}(s, \mathbf{u}_{0,n})| |\mathbf{u}(s, \mathbf{u}_0)|) dx \right) ds \right) \\
& \leq 3\mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\Delta \mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})} (\|\mathbf{u}(s, \mathbf{u}_{0,n})\|_{L^\infty(\mathbb{R})}^2 + \|\mathbf{u}(s, \mathbf{u}_0)\|_{L^\infty(\mathbb{R})}^2) ds \right) \\
& \leq 3c_1^2 \mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\Delta \mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})} (\|\mathbf{u}(s, \mathbf{u}_{0,n})\|_{H^1(\mathbb{R})}^2 + \|\mathbf{u}(s, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2) ds \right) \\
& \leq 3c_1^2 R_\delta \mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\Delta \mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})} ds \right) \\
& \leq \frac{1}{4} \mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\Delta \mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 ds \right) + 9c_1^4 R_\delta^2 \mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 ds \right). \tag{4.40}
\end{aligned}$$

For the last two terms on the right-hand side of (4.35) we have

$$-\mathbb{E} \left(\sum_{k=1}^{\infty} \int_0^{t \wedge \tau_n} (\Delta \mathbf{v}_n(s), (\mathbf{v}_n(s) \times \mathbf{f}_k) \times \mathbf{f}_k) ds \right)$$

$$\begin{aligned}
& + \mathbb{E} \left(\sum_{k=1}^{\infty} \int_0^{t \wedge \tau_n} \|\nabla(\mathbf{v}_n(s) \times \mathbf{f}_k)\|_{L^2(\mathbb{R})}^2 ds \right) \\
& \leq \sum_{k=1}^{\infty} \|\mathbf{f}_k\|_{L^\infty(\mathbb{R})}^2 \mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\Delta \mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})} ds \right) \\
& + 2 \sum_{k=1}^{\infty} \|\mathbf{f}_k\|_{W^{1,\infty}(\mathbb{R})}^2 \mathbb{E} \left(\int_0^{t \wedge \tau_n} (\|\nabla \mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 + \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2) ds \right) \\
& \leq \frac{1}{4} \mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\Delta \mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 ds \right) + c_2 \mathbb{E} \left(\int_0^{t \wedge \tau_n} (\|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 + \|\nabla \mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2) ds \right), \tag{4.41}
\end{aligned}$$

where $c_2 > 0$ depends only on $\{\mathbf{f}_k\}_{k=1}^\infty$.

It follows from (4.35) and (4.39)-(4.41) that for all $t \in [0, t_0]$,

$$\begin{aligned}
& \mathbb{E} \left(\|\nabla \mathbf{v}_n(t \wedge \tau_n)\|_{L^2(\mathbb{R})}^2 \right) \leq \|\nabla \mathbf{v}_n(0)\|_{L^2(\mathbb{R})}^2 \\
& + c_3 R_\delta^{\frac{3}{2}} \mathbb{E} \left(\sup_{s \in [0, t_0]} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \right) + c_3 R_\delta^2 \mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\mathbf{v}_n(s)\|_{H^1(\mathbb{R})}^2 ds \right), \tag{4.42}
\end{aligned}$$

where $c_3 > 0$ depends only on $\{\mathbf{f}_k\}_{k=1}^\infty$.

On the other hand, by (2.51) we have for all $t \in [0, t_0]$,

$$\begin{aligned}
& \mathbb{E} \left(\|\mathbf{v}_n(t \wedge \tau_n)\|_{L^2(\mathbb{R})}^2 \right) \leq \|\mathbf{v}_n(0)\|_{L^2(\mathbb{R})}^2 \\
& + c_4 \mathbb{E} \left(\int_0^{t \wedge \tau_n} \|\mathbf{u}(s, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 \|\mathbf{u}(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 ds \right) \\
& \leq \|\mathbf{v}_n(0)\|_{L^2(\mathbb{R})}^2 + c_4 R_\delta \mathbb{E} \left(\sup_{s \in [0, t_0]} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 \int_0^{t \wedge \tau_n} \|\mathbf{u}(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 ds \right) \\
& \leq \|\mathbf{v}_n(0)\|_{L^2(\mathbb{R})}^2 + c_4 R_\delta^2 \mathbb{E} \left(\sup_{s \in [0, t_0]} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 \right), \tag{4.43}
\end{aligned}$$

where $c_4 > 0$ is an absolute constant.

By (4.42)-(4.43) we obtain for all $t \in [0, t_0]$,

$$\begin{aligned}
& \mathbb{E} \left(\|\mathbf{v}_n(t \wedge \tau_n)\|_{H^1(\mathbb{R})}^2 \right) \leq \|\mathbf{v}_n(0)\|_{H^1(\mathbb{R})}^2 + c_3 R_\delta^2 \mathbb{E} \left(\int_0^t \|\mathbf{v}_n(s \wedge \tau_n)\|_{H^1(\mathbb{R})}^2 ds \right) \\
& + c_3 R_\delta^{\frac{3}{2}} \mathbb{E} \left(\sup_{s \in [0, t_0]} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \right) + c_4 R_\delta^2 \mathbb{E} \left(\sup_{s \in [0, t_0]} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 \right). \tag{4.44}
\end{aligned}$$

Apply Gronwall's inequality to (4.44) to obtain that for all $t \in [0, t_0]$,

$$\begin{aligned}
& \mathbb{E} \left(\|\mathbf{v}_n(t \wedge \tau_n)\|_{H^1(\mathbb{R})}^2 \right) \leq e^{c_3 R_\delta^2 t} \|\mathbf{v}_n(0)\|_{H^1(\mathbb{R})}^2 \\
& + e^{c_3 R_\delta^2 t} R_\delta^{\frac{3}{2}} \left(c_3 \mathbb{E} \left(\sup_{s \in [0, t_0]} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})} \right) + c_4 R_\delta^{\frac{1}{2}} \mathbb{E} \left(\sup_{s \in [0, t_0]} \|\mathbf{v}_n(s)\|_{L^2(\mathbb{R})}^2 \right) \right),
\end{aligned}$$

which along with Lemma 2.6 implies that

$$\lim_{n \rightarrow \infty} \mathbb{E} \left(\|\mathbf{v}_n(t_0 \wedge \tau_n)\|_{H^1(\mathbb{R})}^2 \right) = 0,$$

that is,

$$\lim_{n \rightarrow \infty} \mathbb{E} \left(\|\mathbf{u}(t_0 \wedge \tau_n, \mathbf{u}_{0,n}) - \mathbf{u}(t_0 \wedge \tau_n, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 \right) = 0. \quad (4.45)$$

Given $\eta > 0$, by (4.32)-(4.33) we get

$$\begin{aligned} & \mathbb{P}(\omega \in \Omega : \|\mathbf{u}(t_0, \mathbf{u}_{0,n}) - \mathbf{u}(t_0, \mathbf{u}_0)\|_{H^1(\mathbb{R})} > \eta) \\ &= \mathbb{P}(\omega \in \Omega : \tau_n < t_0, \|\mathbf{u}(t_0, \mathbf{u}_{0,n}) - \mathbf{u}(t_0, \mathbf{u}_0)\|_{H^1(\mathbb{R})} > \eta) \\ & \quad + \mathbb{P}(\omega \in \Omega : \tau_n \geq t_0, \|\mathbf{u}(t_0, \mathbf{u}_{0,n}) - \mathbf{u}(t_0, \mathbf{u}_0)\|_{H^1(\mathbb{R})} > \eta) \\ &\leq \delta + \mathbb{P}(\omega \in \Omega : \tau_n \geq t_0, \|\mathbf{u}(t_0, \mathbf{u}_{0,n}) - \mathbf{u}(t_0, \mathbf{u}_0)\|_{H^1(\mathbb{R})} > \eta) \\ &\leq \delta + \mathbb{P}(\omega \in \Omega : \|\mathbf{u}(t_0 \wedge \tau_n, \mathbf{u}_{0,n}) - \mathbf{u}(t_0 \wedge \tau_n, \mathbf{u}_0)\|_{H^1(\mathbb{R})} > \eta) \\ &\leq \delta + \eta^{-2} \mathbb{E} \left(\|\mathbf{u}(t_0 \wedge \tau_n, \mathbf{u}_{0,n}) - \mathbf{u}(t_0 \wedge \tau_n, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 \right). \end{aligned} \quad (4.46)$$

First letting $n \rightarrow \infty$ and then $\delta \rightarrow 0$ in (4.46), we get for every $\eta > 0$,

$$\lim_{n \rightarrow \infty} \mathbb{P}(\omega \in \Omega : \|\mathbf{u}(t_0, \mathbf{u}_{0,n}) - \mathbf{u}(t_0, \mathbf{u}_0)\|_{H^1(\mathbb{R})} > \eta) = 0, \quad (4.47)$$

that is, $\mathbf{u}(t_0, \mathbf{u}_{0,n}) \rightarrow \mathbf{u}(t_0, \mathbf{u}_0)$ in $H^1(\mathbb{R})$ in probability. Then (4.29) follows from (4.47), the fact $\varphi \in C_b(H^1(\mathbb{R}))$ and the dominated convergence theorem. \square

4.2. Tightness of time averaged measures. In this subsection, we prove the tightness of the time averaged measures μ_n , $n \in \mathbb{N}$, which is defined by

$$\mu_n(\Gamma) = \frac{1}{n} \int_0^n \mathbb{P}(\mathbf{u}(t, \mathbf{u}_0) \in \Gamma) dt, \quad (4.48)$$

where $\Gamma \in \mathcal{B}(H^1(\mathbb{R}))$, and $\mathbf{u}(\cdot, \mathbf{u}_0)$ is the solution of (1.3) with initial data $\mathbf{u}_0 \in H^1(\mathbb{R})$.

Lemma 4.8. *If (1.1) holds, then the sequence $\{\mu_n\}_{n=1}^\infty$ given by (4.48) is tight in $H^1(\mathbb{R})$.*

Proof. Note that, there exists a constant $c_1 > 0$ depending only on \mathbf{u}_0 such that for all $n \in \mathbb{N}$,

$$\frac{1}{n} \int_0^n \mathbb{E} \left(\|\mathbf{u}(t, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 \right) dt \leq c_1. \quad (4.49)$$

On the other hand, by Lemma 4.5 we find that for every $\varepsilon > 0$ and $m \in \mathbb{N}$, there exists a positive integer n_m depending on m, ε and \mathbf{u}_0 such that for all $t \geq 0$,

$$\mathbb{E} \left(\int_{|x| \geq \frac{1}{2} n_m} (|\mathbf{u}(t, \mathbf{u}_0)(x)|^2 + |\nabla \mathbf{u}(t, \mathbf{u}_0)(x)|^2) dx \right) < 2^{-4m} \varepsilon. \quad (4.50)$$

Let ϕ_m be the cut-off function used in the proof of Lemma 4.3, and

$$\mathbf{v}_m(t) = \phi_{n_m} \mathbf{u}(t, \mathbf{u}_0), \quad \mathbf{w}_m(t) = (1 - \phi_{n_m}) \mathbf{u}(t, \mathbf{u}_0).$$

Then by (4.50) we have for all $t \geq 0$,

$$\mathbb{E} \left(\|\mathbf{v}_m(t)\|_{H^1(\mathbb{R})}^2 \right) < 2^{-4m} \varepsilon. \quad (4.51)$$

By (4.49) we find that there exists a constant $c_2 > 0$ depending only on \mathbf{u}_0 such that for all $n \in \mathbb{N}$,

$$\frac{1}{n} \int_0^n \mathbb{E} \left(\|\mathbf{w}_m(t)\|_{H^2(\mathbb{R})}^2 \right) dt \leq c_2. \quad (4.52)$$

Define the sets:

$$\mathcal{W}_m^\varepsilon := \left\{ \mathbf{u} \in H^2(\mathbb{R}) : \mathbf{u} = 0 \text{ for } |x| \geq n_m \text{ and } \|\mathbf{u}\|_{H^2(\mathbb{R})} \leq \frac{2^m \sqrt{c_2}}{\sqrt{\varepsilon}} \right\},$$

and

$$\mathcal{Y}_m^\varepsilon := \left\{ \mathbf{u} \in H^1(\mathbb{R}) : \|\mathbf{u} - \mathbf{v}\|_{H^1(\mathbb{R})} \leq 2^{-m} \text{ for some } \mathbf{v} \in \mathcal{W}_1^\varepsilon \right\},$$

and

$$\mathcal{Y}^\varepsilon = \bigcap_{m=1}^{\infty} \mathcal{Y}_m^\varepsilon.$$

Then \mathcal{Y}^ε is compact in $H^1(\mathbb{R})$. In addition, by (4.51) and (4.52) one can verify that $\mu_n(\mathcal{Y}^\varepsilon) > 1 - \varepsilon$ for all $n \in \mathbb{N}$, and hence $\{\mu_n\}_{n=1}^{\infty}$ is tight in $H^1(\mathbb{R})$. \square

Based on Lemma 4.8, by the Krylov-Bogoliubov theory, we also obtain the existence of invariant measures of (1.3) by the Feller property (Lemma 4.7) with respect to the norm topology instead of the weak topology in $H^1(\mathbb{R})$.

4.3. Limits of invariant measures. In this subsection, we consider the limits of invariant measures of (1.7) with respect to ε . Given $\varepsilon \in [0, 1]$, denote by \mathcal{I}^ε the set of all invariant measures of equation (1.7) corresponding to ε . We will prove the union $\bigcup_{\varepsilon \in [0, 1]} \mathcal{I}^\varepsilon$ is tight in $H^1(\mathbb{R})$. To that end, we recall that all the uniform estimates established in the previous sections are valid for the solutions of (1.7) which are actually uniform for all $\varepsilon \in [0, 1]$. In particular, Lemmas 4.2 and 4.6 apply to the solutions of (1.7) which are presented below.

Lemma 4.9. *If (1.1) holds, then for every $R > 0$, there exists $T = T(R) > 0$ such that for all $t \geq T$ and $\varepsilon \in [0, 1]$, the solution $\mathbf{u}^\varepsilon(t, \mathbf{u}_0)$ of (1.7) with $\|\mathbf{u}_0\|_{H^1(\mathbb{R})} \leq R$ satisfies*

$$\mathbb{E} \left(\|\mathbf{u}^\varepsilon(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 \right) + \int_0^t e^{s-t} \mathbb{E} \left(\|\mathbf{u}^\varepsilon(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 + \int_{\mathbb{R}} |\mathbf{u}^\varepsilon(s, \mathbf{u}_0)|^2 |\nabla \mathbf{u}^\varepsilon(s, \mathbf{u}_0)|^2 dx \right) ds \leq C,$$

and

$$\frac{1}{t} \int_0^t \mathbb{E} \left(\|\mathbf{u}^\varepsilon(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 + \int_{\mathbb{R}} |\mathbf{u}^\varepsilon(s, \mathbf{u}_0)|^2 |\nabla \mathbf{u}^\varepsilon(s, \mathbf{u}_0)|^2 dx \right) ds \leq C,$$

where $C > 0$ is a constant depending on $\{\mathbf{f}_k\}_{k=1}^{\infty}$ but not on \mathbf{u}_0 or ε .

Lemma 4.10. *If (1.1) holds, then for every $\delta > 0$ and $R > 0$, there exist $T = T(\delta, R) > 0$ and $M = M(\delta) \geq 1$ such that for all $t \geq T$, $m \geq M$ and $\varepsilon \in [0, 1]$, the solution $\mathbf{u}^\varepsilon(t, \mathbf{u}_0)$ of (1.7) with $\|\mathbf{u}_0\|_{H^1(\mathbb{R})} \leq R$ satisfies*

$$\mathbb{E} \left(\int_{|x| > m} (|\mathbf{u}^\varepsilon(t, \mathbf{u}_0)(x)|^2 + |\nabla \mathbf{u}^\varepsilon(t, \mathbf{u}_0)(x)|^2) dx \right) < \delta.$$

Next, We prove the union $\bigcup_{\varepsilon \in [0,1]} \mathcal{I}^\varepsilon$ is tight in $H^1(\mathbb{R})$.

Proposition 4.1. *If (1.1) holds, then the set $\bigcup_{\varepsilon \in [0,1]} \mathcal{I}^\varepsilon$ is tight in $H^1(\mathbb{R})$.*

Proof. By Lemma 4.9 we find that for every $\mathbf{u}_0 \in H^1(\mathbb{R})$, there exists $T_1 = T_1(\mathbf{u}_0) > 0$ such that for all $t \geq T_1$ and $\varepsilon \in [0, 1]$, the solution $\mathbf{u}^\varepsilon(t, \mathbf{u}_0)$ of (1.7) satisfies

$$\frac{1}{t} \int_0^t \mathbb{E} \left(\|\mathbf{u}^\varepsilon(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 \right) ds \leq c_1, \quad (4.53)$$

where $c_1 > 0$ is a constant depending on $\{\mathbf{f}_k\}_{k=1}^\infty$ but not on \mathbf{u}_0 or ε . By Lemma 4.10 we infer that for every $\mathbf{u}_0 \in H^1(\mathbb{R})$, $\delta > 0$ and $m \in \mathbb{N}$, there exists $T_2 = T_2(\delta, m, \mathbf{u}_0) > T_1$ and $n_m = n_m(\delta, m) \in \mathbb{N}$ such that for all $t \geq T_2$ and $\varepsilon \in [0, 1]$,

$$\mathbb{E} \left(\int_{|x| \geq \frac{1}{2}n_m} (|\mathbf{u}^\varepsilon(t, \mathbf{u}_0)(x)|^2 + |\nabla \mathbf{u}^\varepsilon(t, \mathbf{u}_0)(x)|^2) dx \right) < 2^{-4m} \delta. \quad (4.54)$$

Let ϕ_m be the cut-off function as in the proof of Lemma 4.3, and

$$\mathbf{v}_m^\varepsilon(t) = \phi_{n_m} \mathbf{u}^\varepsilon(t, \mathbf{u}_0), \quad \mathbf{w}_m^\varepsilon(t) = (1 - \phi_{n_m}) \mathbf{u}^\varepsilon(t, \mathbf{u}_0).$$

By (4.54) we have for all $t \geq T_2$ and $\varepsilon \in [0, 1]$,

$$\mathbb{E} \left(\|\mathbf{v}_m^\varepsilon(t)\|_{H^1(\mathbb{R})}^2 \right) < 2^{-4m} \delta. \quad (4.55)$$

By (4.53) we infer that there exists a constant $c_2 > 0$ depending only on $\{\mathbf{f}_k\}_{k=1}^\infty$ but not on \mathbf{u}_0 or ε such that for all $t \geq T_2$, $m \in \mathbb{N}$ and $\varepsilon \in [0, 1]$,

$$\frac{1}{t} \int_0^t \mathbb{E} \left(\|\mathbf{w}_m^\varepsilon(s)\|_{H^2(\mathbb{R})}^2 \right) ds \leq c_2. \quad (4.56)$$

Denote by

$$\mathcal{W}_1^{m,\delta} = \left\{ \mathbf{u} \in H^2(\mathbb{R}) : \mathbf{u} = 0 \text{ for } |x| \geq n_m \text{ and } \|\mathbf{u}\|_{H^2(\mathbb{R})} \leq \frac{2^m \sqrt{c_2}}{\sqrt{\delta}} \right\},$$

and

$$\mathcal{W}_2^{m,\delta} = \left\{ \mathbf{u} \in H^1(\mathbb{R}) : \|\mathbf{u} - \mathbf{v}\|_{H^1(\mathbb{R})} \leq 2^{-m} \text{ for some } \mathbf{v} \in \mathcal{W}_1^{m,\delta} \right\},$$

and

$$\mathcal{W}^\delta = \bigcap_{m=1}^{\infty} \mathcal{W}_2^{m,\delta}.$$

Then \mathcal{W}^δ is compact in $H^1(\mathbb{R})$, and for every $\mu^\varepsilon \in \mathcal{I}^\varepsilon$,

$$\mu^\varepsilon(\mathcal{W}^\delta) = \lim_{n \rightarrow \infty} \mu^\varepsilon(\mathcal{W}_n^\delta), \quad (4.57)$$

where $\mathcal{W}_n^\delta = \bigcap_{m=1}^n \mathcal{W}_2^{m,\delta}$. For every $n \in \mathbb{N}$, by the invariance of μ^ε , Fubini's theorem and Fatou's lemma, we have

$$\begin{aligned} & \mu^\varepsilon(H^1(\mathbb{R}) \setminus \mathcal{W}_n^\delta) \\ &= \int_{H^1(\mathbb{R})} \mathbb{P}(\mathbf{u}^\varepsilon(s, \mathbf{u}_0) \notin \mathcal{W}_n^\delta) d\mu^\varepsilon(\mathbf{u}_0) \end{aligned}$$

$$\begin{aligned}
&= \limsup_{t \rightarrow \infty} \int_{H^1(\mathbb{R})} \left(\frac{1}{t} \int_0^t \mathbb{P}(\mathbf{u}^\varepsilon(s, \mathbf{u}_0) \notin \mathcal{W}_n^\delta) ds \right) d\mu^\varepsilon(\mathbf{u}_0) \\
&\leq \int_{H^1(\mathbb{R})} \limsup_{t \rightarrow \infty} \left(\frac{1}{t} \int_0^t \sum_{m=1}^n \mathbb{P}(\mathbf{u}^\varepsilon(s, \mathbf{u}_0) \notin \mathcal{W}_2^{m, \delta}) ds \right) d\mu^\varepsilon(\mathbf{u}_0) \\
&\leq \int_{H^1(\mathbb{R})} \limsup_{t \rightarrow \infty} \left(\frac{1}{t} \int_0^t \sum_{m=1}^n \mathbb{P}(\mathbf{w}_m^\varepsilon(s) \notin \mathcal{W}_1^{m, \delta}) ds \right) d\mu^\varepsilon(\mathbf{u}_0) \\
&\quad + \int_{H^1(\mathbb{R})} \limsup_{t \rightarrow \infty} \left(\frac{1}{t} \int_0^t \sum_{m=1}^n \mathbb{P} \left(\mathbf{w}_m^\varepsilon(s) \in \mathcal{W}_1^{m, \delta} \text{ and } \|\mathbf{v}_m^\varepsilon(s)\|_{H^1(\mathbb{R})} > \frac{1}{2^m} \right) ds \right) d\mu^\varepsilon(\mathbf{u}_0) \\
&\leq \sum_{m=1}^n \frac{\delta}{2^{2m} c_2} \int_{H^1(\mathbb{R})} \limsup_{t \rightarrow \infty} \left(\frac{1}{t} \int_0^t \mathbb{E} \|\mathbf{w}_m^\varepsilon(s)\|_{H^2(\mathbb{R})}^2 ds \right) d\mu^\varepsilon(\mathbf{u}_0) \\
&\quad + \sum_{m=1}^n 2^{2m} \int_{H^1(\mathbb{R})} \limsup_{t \rightarrow \infty} \left(\frac{1}{t} \int_0^t \mathbb{E} \|\mathbf{v}_m^\varepsilon(s)\|_{H^1(\mathbb{R})}^2 ds \right) d\mu^\varepsilon(\mathbf{u}_0). \tag{4.58}
\end{aligned}$$

For the first term on the right-hand side of (4.58), by (4.56) we have

$$\begin{aligned}
&\sum_{m=1}^n \frac{\delta}{2^{2m} c_2} \int_{H^1(\mathbb{R})} \limsup_{t \rightarrow \infty} \left(\frac{1}{t} \int_0^t \mathbb{E} \|\mathbf{w}_m^\varepsilon(s)\|_{H^2(\mathbb{R})}^2 ds \right) d\mu^\varepsilon(\mathbf{u}_0) \\
&\leq \sum_{m=1}^n \frac{\delta}{2^{2m}} < \frac{\delta}{3}. \tag{4.59}
\end{aligned}$$

For the second term on the right-hand side of (4.58), by (4.55) we have

$$\sum_{m=1}^n 2^{2m} \int_{H^1(\mathbb{R})} \limsup_{t \rightarrow \infty} \left(\frac{1}{t} \int_0^t \mathbb{E} \|\mathbf{v}_m^\varepsilon(s)\|_{H^1(\mathbb{R})}^2 ds \right) d\mu^\varepsilon(\mathbf{u}_0) \leq \sum_{m=1}^n \frac{\delta}{2^{2m}} < \frac{\delta}{3}. \tag{4.60}$$

Then, from (4.58)-(4.60), we obtain for every $n \in \mathbb{N}$,

$$\mu^\varepsilon(\mathcal{W}_n^\delta) > 1 - \delta. \tag{4.61}$$

By (4.57) and (4.61) we get for every $\mu^\varepsilon \in \mathcal{I}^\varepsilon$,

$$\mu^\varepsilon(\mathcal{W}^\delta) \geq 1 - \delta,$$

which completes the proof. \square

We now show the uniform convergence in probability of solutions of (1.7).

Lemma 4.11. *If (1.1) holds, then for every $L > 0$, $T > 0$, $\delta > 0$ and $\varepsilon_0 \in [0, 1]$, the solution $\mathbf{u}^\varepsilon(t, \mathbf{u}_0)$ of (1.7) satisfies*

$$\lim_{\varepsilon \rightarrow \varepsilon_0} \sup_{\|\mathbf{u}_0\|_{H^1(\mathbb{R})} \leq L} \mathbb{P} \left(\sup_{t \in [0, T]} \|\mathbf{u}^\varepsilon(t, \mathbf{u}_0) - \mathbf{u}^{\varepsilon_0}(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})} \geq \delta \right) = 0. \tag{4.62}$$

Proof. Note that there exists $c_1 = c_1(L, T) > 0$ such that for all $\varepsilon \in [0, 1]$ and $\mathbf{u}_0 \in H^1(\mathbb{R})$ with $\|\mathbf{u}_0\|_{H^1(\mathbb{R})} \leq L$, the solutions of (1.7) satisfy

$$\mathbb{E} \left(\sup_{0 \leq t \leq T} \|\mathbf{u}^\varepsilon(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 + \int_0^T \|\mathbf{u}^\varepsilon(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 ds \right) \leq c_1. \quad (4.63)$$

It follows from (4.63) that for every $\eta > 0$, there exists $R = R(\eta, L, T) > 0$ such that for all $\varepsilon \in [0, 1]$ and $\mathbf{u}_0 \in H^1(\mathbb{R})$ with $\|\mathbf{u}_0\|_{H^1(\mathbb{R})} \leq L$,

$$\mathbb{P} \left(\sup_{0 \leq t \leq T} \|\mathbf{u}^\varepsilon(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 + \int_0^T \|\mathbf{u}^\varepsilon(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 ds > R \right) \leq \frac{1}{2}\eta. \quad (4.64)$$

Define the stopping times by

$$\tau_{1,\varepsilon} = \inf \left\{ t \geq 0; \|\mathbf{u}^\varepsilon(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 + \int_0^t \|\mathbf{u}^\varepsilon(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 ds > R \right\},$$

and

$$\tau_{1,\varepsilon_0} = \inf \left\{ t \geq 0; \|\mathbf{u}^{\varepsilon_0}(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 + \int_0^t \|\mathbf{u}^{\varepsilon_0}(s, \mathbf{u}_0)\|_{H^2(\mathbb{R})}^2 ds > R \right\}.$$

As usual, $\inf \emptyset = +\infty$. Let $\tau_\varepsilon = \tau_{1,\varepsilon} \wedge \tau_{1,\varepsilon_0}$. Then by (4.64) we find that

$$\mathbb{P}(\tau_\varepsilon < T) \leq \eta. \quad (4.65)$$

By (1.7) we have for all $t \in [0, T]$, P-almost surely,

$$\begin{aligned} d(\mathbf{u}^\varepsilon(t) - \mathbf{u}^{\varepsilon_0}(t)) &= \Delta(\mathbf{u}^\varepsilon(t) - \mathbf{u}^{\varepsilon_0}(t))dt + (\mathbf{u}^\varepsilon(t) - \mathbf{u}^{\varepsilon_0}(t)) \times \Delta \mathbf{u}^\varepsilon(t)dt \\ &+ \mathbf{u}^{\varepsilon_0} \times \Delta(\mathbf{u}^\varepsilon(t) - \mathbf{u}^{\varepsilon_0}(t))dt - (\mathbf{u}^\varepsilon(t) - \mathbf{u}^{\varepsilon_0}(t))dt - (|\mathbf{u}^\varepsilon(t)|^2 \mathbf{u}^\varepsilon(t) - |\mathbf{u}^{\varepsilon_0}(t)|^2 \mathbf{u}^{\varepsilon_0}(t)) dt \\ &+ \frac{1}{2}(\varepsilon^2 - \varepsilon_0^2) \sum_{k=1}^{\infty} (\mathbf{u}^\varepsilon(t) \times \mathbf{f}_k + \mathbf{f}_k)dt + \frac{1}{2}\varepsilon_0^2 \sum_{k=1}^{\infty} ((\mathbf{u}^\varepsilon(t) - \mathbf{u}^{\varepsilon_0}(t)) \times \mathbf{f}_k) dt \\ &+ (\varepsilon - \varepsilon_0) \sum_{k=1}^{\infty} (\mathbf{u}^\varepsilon(t) \times \mathbf{f}_k + \mathbf{f}_k)dW_k + \varepsilon_0 \sum_{k=1}^{\infty} ((\mathbf{u}^\varepsilon(t) - \mathbf{u}^{\varepsilon_0}(t)) \times \mathbf{f}_k) dW_k. \end{aligned} \quad (4.66)$$

By (4.66) and Itô's formula we get for all $t \in [0, T]$, P-almost surely,

$$\begin{aligned} &\|\mathbf{u}^\varepsilon(t) - \mathbf{u}^{\varepsilon_0}(t)\|_{H^1(\mathbb{R})}^2 \\ &+ 2 \int_0^t \left(\|\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)\|_{H^1(\mathbb{R})}^2 + \|\nabla \mathbf{u}^\varepsilon(s) - \nabla \mathbf{u}^{\varepsilon_0}(s)\|_{L^2(\mathbb{R})}^2 + \|\Delta \mathbf{u}^\varepsilon(s) - \Delta \mathbf{u}^{\varepsilon_0}(s)\|_{L^2(\mathbb{R})}^2 \right) ds \\ &= -2 \int_0^t (\Delta(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), (\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)) \times \Delta \mathbf{u}^\varepsilon(s))_{L^2(\mathbb{R})} ds \\ &\quad + 2 \int_0^t (\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s), \mathbf{u}^{\varepsilon_0}(s) \times \Delta(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)))_{L^2(\mathbb{R})} ds \\ &- 2 \int_0^t ((I - \Delta)(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), |\mathbf{u}^\varepsilon(s)|^2 \mathbf{u}^\varepsilon(s) - |\mathbf{u}^{\varepsilon_0}(s)|^2 \mathbf{u}^{\varepsilon_0}(s))_{L^2(\mathbb{R})} ds \\ &\quad + (\varepsilon^2 - \varepsilon_0^2) \sum_{k=1}^{\infty} \int_0^t ((I - \Delta)(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), \mathbf{u}^\varepsilon(s) \times \mathbf{f}_k + \mathbf{f}_k)_{L^2(\mathbb{R})} ds \end{aligned}$$

$$\begin{aligned}
& +\varepsilon_0^2 \sum_{k=1}^{\infty} \int_0^t ((I - \Delta)(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), (\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)) \times \mathbf{f}_k)_{L^2(\mathbb{R})} ds \\
& + \sum_{k=1}^{\infty} \int_0^t \|(\varepsilon - \varepsilon_0)(\mathbf{u}^\varepsilon(s) \times \mathbf{f}_k + \mathbf{f}_k) + \varepsilon_0(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)) \times \mathbf{f}_k\|_{H^1(\mathbb{R})}^2 ds \\
& + 2(\varepsilon - \varepsilon_0) \sum_{k=1}^{\infty} \int_0^t ((I - \Delta)(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), \mathbf{u}^\varepsilon(s) \times \mathbf{f}_k + \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \\
& + 2\varepsilon_0 \sum_{k=1}^{\infty} \int_0^t ((I - \Delta)(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), (\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)) \times \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k. \tag{4.67}
\end{aligned}$$

For the first term on the right-hand side of (4.67), by Young's inequality we have

$$\begin{aligned}
& -2 \int_0^t (\Delta(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), (\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)) \times \Delta \mathbf{u}^\varepsilon(s))_{L^2(\mathbb{R})} ds \\
& \leq 2 \int_0^t \|\Delta(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s))\|_{L^2(\mathbb{R})} \|\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)\|_{L^\infty(\mathbb{R})} \|\Delta \mathbf{u}^\varepsilon(s)\|_{L^2(\mathbb{R})} ds \\
& \leq \int_0^t \|\Delta(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s))\|_{L^2(\mathbb{R})}^2 ds + \int_0^t \|\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)\|_{H^1(\mathbb{R})}^2 \|\Delta \mathbf{u}^\varepsilon(s)\|_{L^2(\mathbb{R})}^2 ds. \tag{4.68}
\end{aligned}$$

Similar to (4.40)-(4.41) and (4.68), using Young's inequality to estimate all the other terms on the right-hand side of (4.67), we obtain for all $\varepsilon, \varepsilon_0 \in [0, 1]$ and $0 \leq t \leq t_1 \leq T$, P-almost surely,

$$\begin{aligned}
& \|\mathbf{u}^\varepsilon(t) - \mathbf{u}^{\varepsilon_0}(t)\|_{H^1(\mathbb{R})}^2 \leq c_2(\varepsilon - \varepsilon_0)^2 \int_0^t \left(1 + \|\mathbf{u}^\varepsilon(s)\|_{H^1(\mathbb{R})}^2\right) ds \\
& + c_2 \int_0^t \|\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)\|_{H^1(\mathbb{R})}^2 \left(1 + \|\mathbf{u}^\varepsilon(s)\|_{H^2(\mathbb{R})}^2 + \|\mathbf{u}^\varepsilon(s)\|_{H^1(\mathbb{R})}^4 + \|\mathbf{u}^{\varepsilon_0}(s)\|_{H^1(\mathbb{R})}^4\right) ds \\
& + 2(\varepsilon - \varepsilon_0) \sum_{k=1}^{\infty} \int_0^t ((I - \Delta)(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), \mathbf{u}^\varepsilon(s) \times \mathbf{f}_k + \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \\
& + 2\varepsilon_0 \sum_{k=1}^{\infty} \int_0^t (\nabla(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), (\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)) \times \nabla \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k, \tag{4.69}
\end{aligned}$$

where $c_2 > 0$ depends only on $\{\mathbf{f}_k\}_{k=1}^{\infty}$. By (4.69), we obtain for all $\varepsilon, \varepsilon_0 \in [0, 1]$ and $0 \leq t \leq t_1 \leq T$, P-almost surely,

$$\begin{aligned}
& \sup_{r \in [0, t]} \|\mathbf{u}^\varepsilon(r \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(r \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})}^2 \leq c_2(\varepsilon - \varepsilon_0)^2 \int_0^{t \wedge \tau_\varepsilon} \left(1 + \|\mathbf{u}^\varepsilon(s)\|_{H^1(\mathbb{R})}^2\right) ds \\
& + c_2 \int_0^{t \wedge \tau_\varepsilon} \|\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)\|_{H^1(\mathbb{R})}^2 \left(1 + \|\mathbf{u}^\varepsilon(s)\|_{H^2(\mathbb{R})}^2 + \|\mathbf{u}^\varepsilon(s)\|_{H^1(\mathbb{R})}^4 + \|\mathbf{u}^{\varepsilon_0}(s)\|_{H^1(\mathbb{R})}^4\right) ds \\
& + 2|\varepsilon - \varepsilon_0| \left(\sup_{r \in [0, t]} \left| \sum_{k=1}^{\infty} \int_0^{r \wedge \tau_\varepsilon} ((I - \Delta)(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), \mathbf{u}^\varepsilon(s) \times \mathbf{f}_k + \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \right| \right) \\
& + 2\varepsilon_0 \left(\sup_{r \in [0, t]} \left| \sum_{k=1}^{\infty} \int_0^{r \wedge \tau_\varepsilon} (\nabla(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), (\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)) \times \nabla \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \right| \right)
\end{aligned}$$

$$\begin{aligned}
&\leq c_2(\varepsilon - \varepsilon_0)^2 \int_0^{t_1 \wedge \tau_\varepsilon} \left(1 + \|\mathbf{u}^\varepsilon(s)\|_{H^1(\mathbb{R})}^2\right) ds \\
+ c_2 \int_0^t \sup_{r \in [0, s]} &\|\mathbf{u}^\varepsilon(r \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(r \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})}^2 \mathbf{1}_{(0, \tau_\varepsilon)}(s) \left(1 + \|\mathbf{u}^\varepsilon(s)\|_{H^2(\mathbb{R})}^2 + \|\mathbf{u}^\varepsilon(s)\|_{H^1(\mathbb{R})}^4 + \|\mathbf{u}^{\varepsilon_0}(s)\|_{H^1(\mathbb{R})}^4\right) ds \\
&+ 2|\varepsilon - \varepsilon_0| \left(\sup_{r \in [0, t_1]} \left| \sum_{k=1}^{\infty} \int_0^{r \wedge \tau_\varepsilon} ((I - \Delta)(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), \mathbf{u}^\varepsilon(s) \times \mathbf{f}_k + \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \right| \right) \\
&+ 2\varepsilon_0 \left(\sup_{r \in [0, t_1]} \left| \sum_{k=1}^{\infty} \int_0^{r \wedge \tau_\varepsilon} (\nabla(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), (\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)) \times \nabla \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \right| \right). \quad (4.70)
\end{aligned}$$

Now for a fixed $t_1 \in [0, T]$, applying Gronwall's lemma to (4.70) for $t \in [0, t_1]$, we obtain for all $\varepsilon, \varepsilon_0 \in [0, 1]$ and $t \in [0, t_1]$, P-almost surely,

$$\begin{aligned}
&\sup_{r \in [0, t]} \|\mathbf{u}^\varepsilon(r \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(r \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})}^2 \\
&\leq c_2(\varepsilon - \varepsilon_0)^2 \varphi_\varepsilon(t) \int_0^{t_1 \wedge \tau_\varepsilon} \left(1 + \|\mathbf{u}^\varepsilon(s)\|_{H^1(\mathbb{R})}^2\right) ds \\
&+ 2|\varepsilon - \varepsilon_0| \varphi_\varepsilon(t) \left(\sup_{r \in [0, t_1]} \left| \sum_{k=1}^{\infty} \int_0^{r \wedge \tau_\varepsilon} ((I - \Delta)(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), \mathbf{u}^\varepsilon(s) \times \mathbf{f}_k + \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \right| \right) \\
&+ 2\varepsilon_0 \varphi_\varepsilon(t) \left(\sup_{r \in [0, t_1]} \left| \sum_{k=1}^{\infty} \int_0^{r \wedge \tau_\varepsilon} (\nabla(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), (\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)) \times \nabla \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \right| \right), \quad (4.71)
\end{aligned}$$

where for all $t \in [0, t_1]$,

$$\begin{aligned}
\varphi_\varepsilon(t) &= e^{\int_0^{t \wedge \tau_\varepsilon} c_2(1 + \|\mathbf{u}^\varepsilon(s)\|_{H^2(\mathbb{R})}^2 + \|\mathbf{u}^\varepsilon(s)\|_{H^1(\mathbb{R})}^4 + \|\mathbf{u}^{\varepsilon_0}(s)\|_{H^1(\mathbb{R})}^4) ds} \\
&\leq e^{\int_0^{t \wedge \tau_\varepsilon} c_2(1 + 2R^2 + \|\mathbf{u}^\varepsilon(s)\|_{H^2(\mathbb{R})}^2) ds} \leq e^{c_2(1 + 2R^2)T + c_2R}. \quad (4.72)
\end{aligned}$$

Let $c_3 = e^{c_2(1 + 2R^2)T + c_2R}$. Then by (4.71)-(4.72) we get for all $\varepsilon, \varepsilon_0 \in [0, 1]$ and $t_1 \in [0, T]$, P-almost surely,

$$\begin{aligned}
&\sup_{r \in [0, t_1]} \|\mathbf{u}^\varepsilon(r \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(r \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})}^2 \leq c_2 c_3 (\varepsilon - \varepsilon_0)^2 (1 + R) t_1 \\
&+ 2|\varepsilon - \varepsilon_0| c_3 \left(\sup_{r \in [0, t_1]} \left| \sum_{k=1}^{\infty} \int_0^{r \wedge \tau_\varepsilon} ((I - \Delta)(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), \mathbf{u}^\varepsilon(s) \times \mathbf{f}_k + \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \right| \right) \\
&+ 2\varepsilon_0 c_3 \left(\sup_{r \in [0, t_1]} \left| \sum_{k=1}^{\infty} \int_0^{r \wedge \tau_\varepsilon} (\nabla(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), (\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)) \times \nabla \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \right| \right). \quad (4.73)
\end{aligned}$$

Taking the expectation of (4.73) to obtain for all $\varepsilon, \varepsilon_0 \in [0, 1]$ and $t_1 \in [0, T]$,

$$\begin{aligned}
&\mathbb{E} \left(\sup_{r \in [0, t_1]} \|\mathbf{u}^\varepsilon(r \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(r \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})}^2 \right) \leq c_2 c_3 (\varepsilon - \varepsilon_0)^2 (1 + R) T \\
&+ 2|\varepsilon - \varepsilon_0| c_3 \mathbb{E} \left(\sup_{r \in [0, t_1]} \left| \sum_{k=1}^{\infty} \int_0^{r \wedge \tau_\varepsilon} ((I - \Delta)(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), \mathbf{u}^\varepsilon(s) \times \mathbf{f}_k + \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \right| \right)
\end{aligned}$$

$$+2\varepsilon_0 c_3 \mathbb{E} \left(\sup_{r \in [0, t_1]} \left| \sum_{k=1}^{\infty} \int_0^{r \wedge \tau_\varepsilon} (\nabla(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), (\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)) \times \nabla \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \right| \right). \quad (4.74)$$

Using the BDG inequality, Young's inequality and (1.1), we get

$$\begin{aligned} & 2|\varepsilon - \varepsilon_0| c_3 \mathbb{E} \left(\sup_{r \in [0, t_1]} \left| \sum_{k=1}^{\infty} \int_0^{r \wedge \tau_\varepsilon} ((I - \Delta)(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), \mathbf{u}^\varepsilon(s) \times \mathbf{f}_k + \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \right| \right) \\ & + 2\varepsilon_0 c_3 \mathbb{E} \left(\sup_{r \in [0, t_1]} \left| \sum_{k=1}^{\infty} \int_0^{r \wedge \tau_\varepsilon} (\nabla(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), (\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)) \times \nabla \mathbf{f}_k)_{L^2(\mathbb{R})} dW_k \right| \right) \\ & \leq 2|\varepsilon - \varepsilon_0| c_3 c_4 \mathbb{E} \left(\sum_{k=1}^{\infty} \int_0^{t_1 \wedge \tau_\varepsilon} |((I - \Delta)(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), \mathbf{u}^\varepsilon(s) \times \mathbf{f}_k + \mathbf{f}_k)|^2 ds \right)^{\frac{1}{2}} \\ & \quad + 2\varepsilon_0 c_3 c_4 \mathbb{E} \left(\sum_{k=1}^{\infty} \int_0^{t_1 \wedge \tau_\varepsilon} |(\nabla(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)), (\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)) \times \nabla \mathbf{f}_k)|^2 ds \right)^{\frac{1}{2}} \\ & \leq 2|\varepsilon - \varepsilon_0| c_3 c_4 \mathbb{E} \left(\sum_{k=1}^{\infty} \int_0^{t_1 \wedge \tau_\varepsilon} \|\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)\|_{H^1(\mathbb{R})}^2 \|\mathbf{u}^\varepsilon(s) \times \mathbf{f}_k + \mathbf{f}_k\|_{H^1(\mathbb{R})}^2 ds \right)^{\frac{1}{2}} \\ & + 2\varepsilon_0 c_3 c_4 \mathbb{E} \left(\int_0^{t_1 \wedge \tau_\varepsilon} \|\nabla(\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s))\|_{L^2(\mathbb{R})}^2 \|\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)\|_{L^2(\mathbb{R})}^2 \sum_{k=1}^{\infty} \|\nabla \mathbf{f}_k\|_{L^\infty(\mathbb{R})}^2 ds \right)^{\frac{1}{2}} \\ & \leq 2|\varepsilon - \varepsilon_0| c_5 \mathbb{E} \left(\int_0^{t_1 \wedge \tau_\varepsilon} \|\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)\|_{H^1(\mathbb{R})}^2 (1 + \|\mathbf{u}^\varepsilon(s)\|_{H^1(\mathbb{R})}^2) ds \right)^{\frac{1}{2}} \\ & \quad + 2\varepsilon_0 c_6 \mathbb{E} \left(\int_0^{t_1 \wedge \tau_\varepsilon} \|\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)\|_{H^1(\mathbb{R})}^2 \|\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)\|_{L^2(\mathbb{R})}^2 ds \right)^{\frac{1}{2}} \\ & \leq 2|\varepsilon - \varepsilon_0| c_5 (1 + R)^{\frac{1}{2}} \mathbb{E} \left(\int_0^{t_1 \wedge \tau_\varepsilon} \|\mathbf{u}^\varepsilon(s) - \mathbf{u}^{\varepsilon_0}(s)\|_{H^1(\mathbb{R})}^2 ds \right)^{\frac{1}{2}} \\ & + 2\varepsilon_0 c_6 \mathbb{E} \left(\sup_{r \in [0, t_1]} \|\mathbf{u}^\varepsilon(r \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(r \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})} \left(\int_0^{t_1} \|\mathbf{u}^\varepsilon(s \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(s \wedge \tau_\varepsilon)\|_{L^2(\mathbb{R})}^2 ds \right)^{\frac{1}{2}} \right) \\ & \leq |\varepsilon - \varepsilon_0|^2 c_5^2 (1 + R) + \mathbb{E} \left(\int_0^{t_1} \|\mathbf{u}^\varepsilon(s \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(s \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})}^2 ds \right) \\ & + \frac{1}{2} \mathbb{E} \left(\sup_{r \in [0, t_1]} \|\mathbf{u}^\varepsilon(r \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(r \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})}^2 \right) + 2\varepsilon_0^2 c_6^2 \mathbb{E} \left(\int_0^{t_1} \|\mathbf{u}^\varepsilon(s \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(s \wedge \tau_\varepsilon)\|_{L^2(\mathbb{R})}^2 ds \right) \\ & \leq |\varepsilon - \varepsilon_0|^2 c_5^2 (1 + R) + (1 + 2\varepsilon_0^2 c_6^2) \int_0^{t_1} \mathbb{E} \left(\sup_{r \in [0, s]} \|\mathbf{u}^\varepsilon(r \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(r \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})}^2 \right) ds \\ & \quad + \frac{1}{2} \mathbb{E} \left(\sup_{r \in [0, t_1]} \|\mathbf{u}^\varepsilon(r \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(r \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})}^2 \right), \quad (4.75) \end{aligned}$$

where all the positive constants c_4 , c_5 and c_6 depend only on $\{\mathbf{f}_k\}_{k=1}^{\infty}$.

It follows from (4.74)-(4.75) that for all $\varepsilon, \varepsilon_0 \in [0, 1]$ and $t_1 \in [0, T]$,

$$\begin{aligned} & \mathbb{E} \left(\sup_{r \in [0, t_1]} \|\mathbf{u}^\varepsilon(r \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(r \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})}^2 \right) \leq 2c_2c_3(\varepsilon - \varepsilon_0)^2(1 + R)T \\ & + 2|\varepsilon - \varepsilon_0|^2c_5^2(1 + R) + 2(1 + 2\varepsilon_0^2c_6^2) \int_0^{t_1} \mathbb{E} \left(\sup_{r \in [0, s]} \|\mathbf{u}^\varepsilon(r \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(r \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})}^2 \right) ds. \end{aligned} \quad (4.76)$$

Applying Gronwall's inequality to (4.76) for $t_1 \in [0, T]$, we get for all $\varepsilon, \varepsilon_0 \in [0, 1]$ and $t_1 \in [0, T]$,

$$\begin{aligned} & \mathbb{E} \left(\sup_{r \in [0, t_1]} \|\mathbf{u}^\varepsilon(r \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(r \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})}^2 \right) \\ & \leq (2c_2c_3(\varepsilon - \varepsilon_0)^2(1 + R)T + 2|\varepsilon - \varepsilon_0|^2c_5^2(1 + R)) e^{2(1+2\varepsilon_0^2c_6^2)t_1}, \end{aligned}$$

and hence for all $\varepsilon, \varepsilon_0 \in [0, 1]$ and $t_1 = T$, we obtain

$$\begin{aligned} & \mathbb{E} \left(\sup_{r \in [0, T]} \|\mathbf{u}^\varepsilon(r \wedge \tau_\varepsilon) - \mathbf{u}^{\varepsilon_0}(r \wedge \tau_\varepsilon)\|_{H^1(\mathbb{R})}^2 \right) \\ & \leq 2(\varepsilon - \varepsilon_0)^2(1 + R) (c_2c_3T + c_5^2) e^{2(1+2\varepsilon_0^2c_6^2)T}. \end{aligned} \quad (4.77)$$

By (4.65) and Chebyshev's inequality, we get for all $\varepsilon, \varepsilon_0 \in [0, 1]$ and $\mathbf{u}_0 \in H^1(\mathbb{R})$ with $\|\mathbf{u}_0\|_{H^1(\mathbb{R})} \leq L$,

$$\begin{aligned} & \mathbb{P} \left(\sup_{t \in [0, T]} \|\mathbf{u}^\varepsilon(t, \mathbf{u}_0) - \mathbf{u}^{\varepsilon_0}(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})} \geq \delta \right) \\ & \leq \mathbb{P} \left(\left\{ \sup_{t \in [0, T]} \|\mathbf{u}^\varepsilon(t, \mathbf{u}_0) - \mathbf{u}^{\varepsilon_0}(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})} \geq \delta \right\} \cap \{\tau_\varepsilon \geq T\} \right) \\ & \quad + \mathbb{P} \left(\left\{ \sup_{t \in [0, T]} \|\mathbf{u}^\varepsilon(t, \mathbf{u}_0) - \mathbf{u}^{\varepsilon_0}(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})} \geq \delta \right\} \cap \{\tau_\varepsilon < T\} \right) \\ & \leq \mathbb{P} \left(\sup_{t \in [0, T]} \|\mathbf{u}^\varepsilon(t \wedge \tau_\varepsilon, \mathbf{u}_0) - \mathbf{u}^{\varepsilon_0}(t \wedge \tau_\varepsilon, \mathbf{u}_0)\|_{H^1(\mathbb{R})} \geq \delta \right) + \mathbb{P}(\tau_\varepsilon < T) \\ & \leq \frac{1}{\delta^2} \mathbb{E} \left(\sup_{t \in [0, T]} \|\mathbf{u}^\varepsilon(t \wedge \tau_\varepsilon, \mathbf{u}_0) - \mathbf{u}^{\varepsilon_0}(t \wedge \tau_\varepsilon, \mathbf{u}_0)\|_{H^1(\mathbb{R})}^2 \right) + \eta. \end{aligned} \quad (4.78)$$

By (4.77)-(4.78) we find that for all $\varepsilon, \varepsilon_0 \in [0, 1]$,

$$\begin{aligned} & \sup_{\|\mathbf{u}_0\|_{H^1(\mathbb{R})} \leq L} \mathbb{P} \left(\sup_{t \in [0, T]} \|\mathbf{u}^\varepsilon(t, \mathbf{u}_0) - \mathbf{u}^{\varepsilon_0}(t, \mathbf{u}_0)\|_{H^1(\mathbb{R})} \geq \delta \right) \\ & \leq 2(\varepsilon - \varepsilon_0)^2(1 + R)\delta^{-2} (c_2c_3T + c_5^2) e^{2(1+2\varepsilon_0^2c_6^2)T} + \eta. \end{aligned}$$

First letting $\varepsilon \rightarrow \varepsilon_0$ and then $\eta \rightarrow 0$, we obtain (4.62), which completes the proof. \square

We are now in a position to prove Theorem 1.2.

Proof of Theorem 1.2. Note that Theorem 1.2 (i) follows from Theorem 3.1. It remains to show Theorem 1.2 (ii).

By Proposition 4.1 we know the set $\bigcup_{\varepsilon \in (0,1]} \mathcal{I}^\varepsilon$ is tight in $H^1(\mathbb{R})$. Then for any sequence $\{\mu^{\varepsilon_n}\}_{n=1}^\infty \subseteq \bigcup_{\varepsilon \in (0,1]} \mathcal{I}^\varepsilon$ with $\varepsilon_n \rightarrow \varepsilon_0 \in [0, 1]$, there exists a subsequence (still denoted by $\{\mu^{\varepsilon_n}\}_{n=1}^\infty$) and a probability measure μ^{ε_0} in $H^1(\mathbb{R})$ such that $\mu^{\varepsilon_n} \rightarrow \mu^{\varepsilon_0}$ weakly. By Lemma 4.11, we find from [33] (or [7] for $\varepsilon_0 = 0$) that μ^{ε_0} is an invariant measure of the limiting equation (1.7) with $\varepsilon = \varepsilon_0$, which concludes the proof.

ACKNOWLEDGMENTS

D. Huang is supported by the National Natural Science Foundation of China (Grant No. 12471228). Z. Qiu is supported by the National Natural Science Foundation of China (Grant No. 12401305), the National Science Foundation for Colleges and Universities in Jiangsu Province (Grant No. 24KJB110011) and the National Science Foundation of Jiangsu Province (Grant No. BK20240721).

REFERENCES

- [1] U. Atxitia, O. Chubykalo-Fesenko, N. Kazantseva, D. Hinzke, U. Nowak, R.W. Chantrell: Micromagnetic modeling of laser-induced magnetization dynamics using the Landau-Lifshitz-Bloch equation, *Appl. Phys. Lett.* 91(23):232507 (2007)
- [2] Z. Brzeźniak, L. Motyl: Existence of a martingale solution of the stochastic Navier-Stokes equations in unbounded 2D and 3D domains, *J. Differential Equations* 254:1627-1685 (2013)
- [3] Z. Brzeźniak, B. Goldys, T. Jegaraj: Large deviations and transitions between equilibria for stochastic Landau-Lifshitz-Gilbert equation, *Arch. Rational Mech. Anal.* 226:497-558 (2017)
- [4] Z. Brzeźniak, B. Goldys, K.N. Le: Existence of a unique solution and invariant measures for the stochastic Landau-Lifshitz-Bloch equation, *J. Differential Equations* 269(11):9471-9507 (2020)
- [5] Z. Brzeźniak, M. Ondrejat, J. Seidler: Invariant measures for stochastic nonlinear beam and wave equations, *J. Differential Equations* 260:4157-4179 (2016)
- [6] Z. Brzeźniak, E. Motyl, M. Ondrejat: Invariant measure for the stochastic Navier-Stokes equations in unbounded 2D domains, *Ann. Probab.* 45:3145-3201 (2017)
- [7] L. Chen, Z. Dong, J. Jiang, J. Zhai: On limiting behavior of stationary measures for stochastic evolution systems with small noise intensity, *Sci. China Math.* 63(8):1463-1504 (2020)
- [8] M. Chen, D. Wang, H. Wang: Martingale solution for the three-dimensional stochastic non-homogeneous incompressible Navier-Stokes equations driven by Lévy noise, *J. Funct. Anal.* 276:2007-2051 (2019)
- [9] Z. Chen, B. Wang: Invariant measures of fractional stochastic delay reaction-diffusion equations on unbounded domain, *Nonlinearity* 34(6):3969 (2021)
- [10] Z. Chen, B. Wang: Existence, exponential mixing and convergence of periodic measures of fractional stochastic delay reaction-diffusion equations on \mathbb{R}^n , *J. Differential Equations* 336:505-564 (2022)
- [11] Z. Chen, B. Wang: Limit measures of stochastic Schrödinger lattice systems, *Proc. Amer. Math. Soc.* 150(4):1669-1684 (2022)
- [12] P. Constantin, N. Glatt-Holtz, V. Vicol: Unique ergodicity for fractionally dissipated, stochastically forced 2D Euler equations, *Commun. Math. Phys.* 330:819-857 (2014)
- [13] G. Da Prato, A. Debussche: Ergodicity for the 3D stochastic Navier-Stokes equations, *J. Math. Pures Appl.* 82(8):877-947 (2003)
- [14] A. Debussche, N. Glatt-Holtz, R. Temam: Local martingale and pathwise solutions for an abstract fluids model, *Phys. D: Nonlinear Phenomena* 240(14-15):1123-1144 (2011)

- [15] J. Eckmann, M. Hairer: Invariant measures for stochastic partial differential equations in unbounded domains, *Nonlinearity* 14:133-151 (2001)
- [16] R.F.L. Evans, D. Hinzke, U. Atxitia, U. Nowak, R.W. Chantrell, O. Chubykalo-Fesenko: Stochastic form of the Landau-Lifshitz-Bloch equation, *Phys. Rev. B* 85:014433 (2012)
- [17] F. Flandoli: Dissipativity and invariant measures for stochastic Navier-Stokes equations, *NoDEA*, 1:403-423 (1994)
- [18] D.A. Garanin: Generalized equation of motion for a ferromagnet, *Phys. A* 172(3):470-491 (1991)
- [19] D.A. Garanin, O. Chubykalo-Fesenko: Thermal fluctuations and longitudinal relaxation of single-domain magnetic particles at elevated temperatures, *Phys. Rev. B* 70:212409 (2004)
- [20] B. Gess, J.M. Tölle: Ergodicity and local limits for stochastic local and nonlocal p -Laplace equations, *SIAM J. Math. Anal.* 48(6):4094-4125 (2016)
- [21] B. Gess, P.E. Souganidis: Long-time behavior, invariant measures, and regularizing effects for stochastic scalar conservation Laws, *Comm. Pure App. Math.* 70(8): 1562-1597 (2017)
- [22] A. Jakubowski: The almost sure Skorokhod representation for subsequences in nonmetric spaces, *Theory Probab. Appl.* 42(1):167-175 (1998)
- [23] H. Gao, H. Liu: Well-posedness and invariant measures for a class of stochastic 3D Navier-Stokes equations with damping driven by jump noise, *J. Differential Equations* 267:5938-5975 (2019)
- [24] M. Hairer: Exponential mixing properties of stochastic PDEs through asymptotic coupling, *Probab. Theory Relat. Field* 124(3):345-380 (2002)
- [25] Z. Jia, B. Guo: Landau-Lifshitz-Bloch equation on Riemannian manifold, *Frontiers Math. China* 14:45-76 (2019)
- [26] S. Jiang, Q. Ju, H. Wang: Martingale weak solutions of the stochastic Landau-Lifshitz-Bloch equation, *J. Differential Equations* 266(5):2542-2574 (2019)
- [27] J. Huang, Y. Zheng, T. Shen, C. Guo: Asymptotic properties of the 2D stochastic fractional Boussinesq equations driven by degenerate noise, *J. Differential Equations*, 310:362-403 (2022)
- [28] J. Kim: On the stochastic Burgers equation with polynomial nonlinearity in the real line, *Discrete Contin. Dyn. Syst. Ser. B* 6:835-866 (2006)
- [29] J. Kim: Invariant measures for a stochastic nonlinear Schrödinger equation, *Indiana Univ. Math. J.* 55:687-717 (2006)
- [30] N.V. Krylov, B.L. Rozovskii: Stochastic evolution equation, *J. Soviet Math.* 16:1233-1277 (1981)
- [31] B. Maslowski, J. Seidler: On sequentially weakly Feller solutions to SPDE's. *Atti della Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei Mat. Appl.* 10(2):69-78 (1999)
- [32] K.N. Le: Weak solutions of the Landau-Lifshitz-Bloch equation, *J. Differential Equations* 261:(12) 6699-6717 (2016)
- [33] D. Li, B. Wang, X. Wang: Limiting behavior of invariant measures of stochastic delay lattice systems, *J. Dyn. Differential Equations* 34:1453-1487 (2022)
- [34] O. Misiats, O. Stanzhytskyi, N. Yip: Existence and uniqueness of invariant measures for stochastic reaction-diffusion equations in unbounded domains, *J. Theor. Probab.* 29:996-1026 (2016)
- [35] L. Néel: Bases d'une nouvelle théorie générale du champ coercitif, *Ann. Univ. Grenoble* 22:299-343 (1946)
- [36] Z. Qiu, C. Sun: Stochastic Landau-Lifshitz-Bloch equation with transport noise: well-posedness, dissipation enhancement, *J. Stat. Phys.* 191:43 (2024)
- [37] B. Wang: Dynamics of fractional stochastic reaction-diffusion equations on unbounded domains driven by nonlinear noise, *J. Differential equations*, 268:1-59 (2019)
- [38] D. Wang, H. Wang: Global existence of martingale solutions to the three-dimensional stochastic compressible Navier-Stokes equations, *Diff. Int. Equ.* 28(11/12):1105-1154 (2015)

INSTITUTE OF APPLIED PHYSICS AND COMPUTATIONAL MATHEMATICS, BEIJING 100088, CHINA

Email address: `huang_daiwen@iapcm.ac.cn`

SCHOOL OF APPLIED MATHEMATICS, NANJING UNIVERSITY OF FINANCE AND ECONOMICS, NANJING, 210046, CHINA.

Email address: `zhqmath@163.com`

DEPARTMENT OF MATHEMATICS, NEW MEXICO INSTITUTE OF MINING AND TECHNOLOGY, SOCORRO, NM 87801, USA.

Email address: `bwang@nmt.edu`.