

Evaluation of tungsten influx rate using line emissions from W^{5+} ions in EAST Tokamak

Fengling Zhang^{1,2}, Darío Mitnik^{*3,1}, Ling Zhang^{†1}, Runjia Bao^{1,2}, Wenming Zhang^{1,2}, Yunxin Cheng¹, Ailan Hu¹, Shigeru Morita⁴, Xiaobin Ding⁵, Yinxian Jie¹, and Haiqing Liu¹

¹ Institute of Plasma Physics, Hefei Institutes of Physical Science, Chinese Academy of Sciences, Hefei 230031, China

² University of Science and Technology of China, Hefei 230026, China

³ Instituto de Astronomía y Física del Espacio (CONICET-Universidad de Buenos Aires), Buenos Aires 1428, Argentina

⁴ National Institute for Fusion Science, Toki 509-5292, Gifu, Japan

⁵ Key laboratory of Atomic and Molecular Physics & Functional Materials of Gansu Province, College of Physics and Electronic Engineering, Northwest Normal University, Lanzhou, 730070, China

E-mail: ^{*}dmitnik@df.uba.ar, [†]zhangling@ipp.ac.cn

September 2024

Abstract. The S/XB ratios (ionization per emitted photon) allow one to relate spectroscopic emissivity measurements to the impurity influx from a localized source. In this work, we determine the tungsten influx by examining two dominant EUV (Extreme Ultraviolet) line emissions at 282.13 Å and 394.07 Å, corresponding to the $4f^{14}5f \rightarrow 4f^{14}5d$ radiative transitions of the W^{5+} ion. The ground configuration of W^{5+} consists of the ground level and a metastable level, with the latter having a higher population than the ground state. Therefore, a simple approach assuming that the transitions are independent, i.e., only populated by a unique level source, requires correction.

To address this, we have developed a fully collisional–radiative modeling in which 430 levels contribute to the ionization. We have utilized three advanced computational codes – HULLAC (Hebrew University - Lawrence Livermore Atomic Code), AUTOSTRUCTURE, and FAC (Flexible Atomic Code) – for the atomic structure calculations. These codes provide the necessary information such as wavelengths, collisional and radiative transition rate coefficients. The FAC code was also used to calculate the direct electron-impact ionization under the distorted-wave approximation. We also included contributions to total ionization from excitation-autoionization processes up to $n = 15$ manifolds from the distorted-wave calculations.

Subsequently, we used these results to ascertain the tungsten impurity influx in a dedicated discharge of the EAST tokamak, which operates with full tungsten divertors. In our findings, we observed that for the density range relevant to the edge region of a tokamak reactor, the S/XB ratios are almost independent of electron density but exhibit significant variation with electron temperature.

Keywords: Impurity Influx in Tokamak, S/XB values, EUV Spectroscopy, Tungsten Spectroscopy

1. INTRODUCTION

Various intrinsic impurities are inevitably generated in tokamak discharges due to the interaction between plasma and wall. Impurities ionized in the core release a large number of electrons, severely diluting the concentration of the primary plasma and affecting the reaction power density of fusion, decreasing the overall performance of the plasma. Therefore, monitoring and controlling the entry of impurities into the core is an urgent problem for long-pulse discharge operations.

It is possible to relate spectroscopic measurements of emissivities from an impurity ion to the impurity influx into the plasma core. Behringer *et al.* [1] proposed the use of the “ionization per emitted photon” S/XB ratio, also known as the “inverse photon efficiency”, denoting the number of ionization events per observed photons (incoming ions over emitted photons). The S/XB ratios have been used to evaluate the influx in tokamak devices of common light impurities such as carbon, nitrogen, oxygen, and neon, as well as metal impurities such as chromium, iron, nickel, molybdenum, and others [1, 2, 3, 4, 5].

Tungsten has been selected as the divertor and first wall material for ITER due to its high melting point, high thermal conductivity, low sputtering rate, low fuel retention rate, and low neutron activation rate [6]. However, highly ionized tungsten ions in the core and edge emit intense radiation at various wavelength ranges, which can cause significant radiation power loss and severely reduce plasma confinement performance. Therefore, evaluating the amount of tungsten entering through the edge is crucial for functioning high confinement long pulse plasma. Many extensive studies have been dedicated to the evaluation of the tungsten influx using the 4009 Å spectroscopic line of W I [7, 8, 9]. Other lines from W I have been used by Beigman *et al.* [10], and W II was used by Pospieszczyk *et al.* [11]. Ballance *et al.* [12] conducted a detailed study on the S/XB ratios for the 1099.05 Å, 1119.7 Å, 1172.47 Å, and 1186.17 Å lines of W^{3+} , assuming these lines as independent. Dong *et al.* [13] calculated the influx rate of W^{6+} using the S/XB coefficients calculated with the ADAS code [14].

Recently, the spectrum of W^{5+} was observed during EAST discharge, which provides us with the necessary experimental spectrum for further studying the S/XB of low-ionized tungsten ions. As pointed out by Behringer *et al.* [1], data for any particular ionization stage of the atomic species of interest can be employed as long as no higher ionization stages of that atom emerge from the localized source. Thus, such spectroscopic measurements for specific radiative transitions in W^{5+} would allow for the determination of the influx of tungsten.

The most straightforward approach for the S/XB determination is to assume that only one ground (or metastable) level populates the upper level of the observed spectral line. Even if the effects of collisions to and cascades from higher lying levels are considered, this simple approximation only works for some cases, as shown in the following sections. In general, we must take allowance that the source of the level population can also arise from other metastables and not only from a unique level, the

so-called *metastable cross-coupling approach* [3].

The paper is organized as follows. In Section 3, we develop the main formulas needed for the evaluation of the S/XB coefficients. First, we show in 3.1 details about how to employ the rate coefficients for excitations and radiative decays into a quasistatic generalized collisional–radiative level population model. With this, we obtain the effective contribution to the population of the excited states involved in the radiative transitions to study. In 3.2, we show how to use these coefficients to determine S/XB ratios as a function of electron density and temperature. We realized that many works deal with cases in which only the ground level is populated or cases in which the emissivity in the metastable line could be considered independent (namely, the upper levels are only populated by direct excitation from a unique source). However, the discussion for cross-coupled metastables, in which the populations of the upper levels arise from different sources, is scarce and worth developing in detail. In Subsection 3.4, we briefly describe the three computational codes used in our calculations for the atomic structure. In Section 4, we show the results, first for the atomic structure calculations in 4.1. Then, we present the results obtained for a model consisting of only four levels, which accounts for most of the relevant physical processes involved, at least within the density and temperature range of interest for Tokamak plasmas. Following that, we solve a full collisional-radiative model with 430 levels, and present and discuss the final S/XB coefficients. We conclude with some final remarks in Section 5.

2. Experimental setup

EAST is a fully superconducting tokamak device equipped with ITER-like active water-cooled W/Cu monoblocks tungsten divertors, capable of high-power long pulse operation. The main parameters are: major radius $R \leq 1.9$ m, minor radius $a \leq 0.45$ m, plasma current $I_p \leq 1$ MA, and toroidal magnetic field $B_T \leq 3.5$ T [15, 16, 17]. Presently, various auxiliary heating and current driving have been installed, including two lower hybrid current drive (LHCD) systems, an electron cyclotron heating (ECH) system, an ion cyclotron resonant frequency (ICRF) system, and balanced neutral beam injection (NBI) systems [18]. A set of flat-field extreme ultraviolet (EUV) spectrometers, working in the 20-500 Å wavelength range with fast time response (5 ms/frame), named EUV_long [19], was developed to measure line emissions and to monitor the impurities present in the EAST plasmas. A laminar-type varied-line-spacing concave holographic grating 1200 grooves/mm is fixed at grazing incidence 87° with a narrow entrance slit width of 30 μm for increasing the spectral resolution. A back-illuminated charge-coupled device (CCD) with a total size of 26.6×6.6 mm² and pixel numbers of 1024×255 (26×26 μm^2 /pixel) is used for recording the spectrum. The wavelength calibration is performed by cubic polynomial fitting with many well-known spectral lines covering the observable range [20]. An absolute calibration, in which the raw measured spectral counts are converted into absolute spectral intensities, is further achieved by comparing the observed and calculated intensities of EUV and visible bremsstrahlung continua, a

procedure explained in previous works [19, 20, 21, 22].

Figure 1 shows the operating parameters of EAST plasma discharge #100300 as a function of time. The upper part (a) displays the plasma current I_p . Part (b) shows the electron temperature of the core plasma T_{e0} provided by the Thomson scattering system [23], and the line-averaged electron density n_e measured by the hydrogen cyanide (HCN) interferometer system [24]. Part (c) depicts the heating power of ECRH, LHW, and NBI. Part (d) shows the total radiation power loss measured by the AXUV (absolute extreme ultraviolet) photodiode array [25]. Finally, part (e) shows the emission intensity of the two dominant W^{5+} EUV spectral lines used here for determining the tungsten total flux. As shown in part (d), the radiation power loss starts to increase at time $t = 5.14$ s, as a consequence of core tungsten accumulation, which causes the plasma to cool down dramatically, as seen in part (b). Then, the plasma became unstable. The spectral lines from low-ionized tungsten ions (e) appear in the cool plasma after strong interactions between the unstable plasma and the wall at $t \approx 5.2$ s.

Figure 2 displays the 280-440 Å spectral range for the EAST discharge #100300 at two different times. The brown curve was obtained at time $t = 5.150$ s, when no trace of W was found in the plasma, and the black curve was obtained at time $t = 5.200$ s when the W intensity reached its maximum. Before the burst, only some lines emitted from HeII, Fe XVI, CIV, and CV are observable in this spectral range. The lines observed at the W peak intensity have been identified [26, 27] as belonging to different low-charged W ions, as indicated in the figure. The strong lines in the 280-320 Å range are associated with the $5p^6 \rightarrow 5p^55d$ transitions of W^{6+} , and from this ion, the $4f^{13}5s^25p^65d \rightarrow 4f^{13}5s^25p^65f$ lines are also observed. The line at 434.33Å, corresponds to the $5d^2 \rightarrow 5d7p$ transition of W^{4+} . In the region around 400 Å, two prominent lines are highlighted. We have identified these lines as corresponding to $5f \rightarrow 5d$ transitions of W^{5+} ; the line at 382.13 Å is emitted by the $(4f^{14}5f)_{\frac{5}{2}} \rightarrow (4f^{14}5d)_{\frac{3}{2}}$ transition, and the line at 394.07 Å arises from the $(4f^{14}5f)_{\frac{7}{2}} \rightarrow (4f^{14}5d)_{\frac{5}{2}}$ transition. As we shall see later, these lines will provide the spectroscopic information necessary to determine the amount of tungsten flux entering the plasma.




Figure 1. Evolution of specific parameters for the EAST discharge #100300. (a) Plasma current I_p . (b) Central electron temperature, T_{e0} , and line-averaged electron density, n_e . (c) Heating power of ECRH, LHW, and NBI. (d) Boundary radiation power loss. (e) W^{5+} emission line intensities at 382.13 Å (blue), and 394.07 Å (red).




Figure 2. EAST spectra obtained at discharge #100300 with the b_long spectrometer. Brown curve: Observed at $t = 5.150$, before the W burst. Black curve: Observed at $t = 5.200$, at the maximum of W intensity.

3. Theoretical methods

3.1. Collisional Radiative Model

To determine the impurity influx for a particular element using the photon emission from a particular ion, one needs to know the population of the excited levels of this ion. To calculate the population n_j of the excited levels j of a particular impurity ion, we solve for a given electronic density and temperature, the collisional-radiative equations:

$$\begin{aligned} \frac{dn_j}{dt} &= \sum_{k>j} n_k (A_{kj} + n_e Q_{kj}) + \sum_{i<j} n_i n_e Q_{ij} - \\ &\quad - n_j \left(\sum_{i<j} (A_{ji} + n_e Q_{ji}) + \sum_{k>j} n_e Q_{jk} \right) = \\ &= \sum_m C_{jm} n_m, \end{aligned} \quad (1)$$

where n_e is the electronic density, A_{ji} are the radiative rate coefficient for transitions from level j to level i . Q_{ji} are the electron-impact excitation rate coefficients if $i > j$, and the deexcitation rates if $i < j$. In matrixial form, Eq. (1) is

$$\frac{d\vec{n}}{dt} = \mathbf{C} \cdot \vec{n} \quad (2)$$

where the elements of the matrix \mathbf{C} are

$$C_{jm} \equiv \begin{cases} A_{mj} + n_e Q_{mj} & \text{for } m > j \\ n_e Q_{mj} & \text{for } m < j \\ - \left(\sum_{i<j} (A_{ji} + n_e Q_{ji}) + \sum_{k>j} n_e Q_{jk} \right) & \text{for } m = j \end{cases} \quad (3)$$

We partition the complete set of levels into metastables (denoted by greek letters σ, ρ, \dots, μ), and excited levels (denoted here with roman letters i, j, \dots, n)

$$\vec{n} = \begin{pmatrix} \vec{n}_\tau \\ \vec{n}_s \end{pmatrix} \quad (4)$$

and assume a quasi-static approximation in which the population of the excited levels are relaxed relative to the ground and metastable levels:

$$\frac{d\vec{n}_\tau}{dt} \neq 0 \quad \text{and} \quad \frac{d\vec{n}_s}{dt} = 0. \quad (5)$$

Under this approximation, Eq. (2) becomes

$$\begin{pmatrix} \frac{dn_\sigma}{dt} \\ \frac{dn_\rho}{dt} \\ \dots \\ \frac{dn_\mu}{dt} \\ 0 \\ 0 \\ \dots \\ 0 \end{pmatrix} = \begin{pmatrix} C_{\sigma\sigma} & C_{\sigma\rho} & \dots & C_{\sigma\mu} & C_{\sigma i} & C_{\sigma j} & \dots & C_{\sigma n} \\ C_{\rho\sigma} & C_{\rho\rho} & \dots & C_{\rho\mu} & C_{\rho i} & C_{\rho j} & \dots & C_{\rho n} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ C_{\mu\sigma} & C_{\mu\rho} & \dots & C_{\mu\mu} & C_{\mu i} & C_{\mu j} & \dots & C_{\mu n} \\ C_{i\sigma} & C_{i\rho} & \dots & C_{i\mu} & C_{ii} & C_{ij} & \dots & C_{in} \\ C_{j\sigma} & C_{j\rho} & \dots & C_{j\mu} & C_{ji} & C_{jj} & \dots & C_{jn} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ C_{n\sigma} & C_{n\rho} & \dots & C_{n\mu} & C_{ni} & C_{nj} & \dots & C_{nn} \end{pmatrix} \begin{pmatrix} n_\sigma \\ n_\rho \\ \dots \\ n_\mu \\ n_i \\ n_j \\ \dots \\ n_n \end{pmatrix} \quad (6)$$

To solve this problem, we take only the excited-levels part of the matrix, having the following equation

$$\begin{pmatrix} C_{i\sigma}n_\sigma + \cdots + C_{i\mu}n_\mu + C_{ii}n_i + \cdots + C_{in}n_n \\ C_{j\sigma}n_\sigma + \cdots + C_{j\mu}n_\mu + C_{ji}n_i + \cdots + C_{jn}n_n \\ \cdots \\ C_{n\sigma}n_\sigma + \cdots + C_{n\mu}n_\mu + C_{ni}n_i + \cdots + C_{nn}n_n \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \cdots \\ 0 \end{pmatrix} \quad (7)$$

which is equivalent to

$$\begin{pmatrix} C_{ii}n_i + \cdots + C_{in}n_n \\ C_{ji}n_i + \cdots + C_{jn}n_n \\ \cdots \\ C_{ni}n_i + \cdots + C_{nn}n_n \end{pmatrix} = \begin{pmatrix} -C_{i\sigma}n_\sigma + \cdots - C_{i\mu}n_\mu \\ -C_{j\sigma}n_\sigma + \cdots - C_{j\mu}n_\mu \\ \cdots \\ -C_{n\sigma}n_\sigma + \cdots - C_{n\mu}n_\mu \end{pmatrix} \quad (8)$$

The last expression can be written in a matricial form:

$$\mathbf{C}_s \cdot \vec{n}_s = \vec{C}_\sigma \cdot n_\sigma + \cdots + \vec{C}_\mu \cdot n_\mu \quad (9)$$

where the reduced matrix for the excited levels is

$$\mathbf{C}_s = \begin{pmatrix} C_{ii} & \cdots & C_{in} \\ C_{ji} & \cdots & C_{jn} \\ \cdots & \cdots & \cdots \\ C_{ni} & \cdots & C_{nn} \end{pmatrix} \quad (10)$$

and the column vectors are

$$\vec{C}_\tau = \begin{pmatrix} -C_{i\tau} \\ -C_{j\tau} \\ \cdots \\ -C_{n\tau} \end{pmatrix} \quad (11)$$

The solution for the excited levels is

$$\begin{aligned} \vec{n}_s &= [\mathbf{C}_s]^{-1} \cdot \vec{C}_\sigma \cdot n_\sigma + \cdots + [\mathbf{C}_s]^{-1} \cdot \vec{C}_\mu \cdot n_\mu = \\ &= \vec{\mathcal{F}}_\sigma \cdot n_\sigma + \cdots + \vec{\mathcal{F}}_\mu \cdot n_\mu \end{aligned} \quad (12)$$

in which the column vectors

$$\vec{\mathcal{F}}_\tau \equiv [\mathbf{C}_s]^{-1} \cdot \vec{C}_\tau \quad (13)$$

denote the effective contribution to the population of the excited levels via excitations from the metastable τ . From this point, we want to follow the usual notation used in the field, denoting the j component of the vector $\vec{\mathcal{F}}_\tau$ (i.e., the effective contribution to the population of j due to excitations from the metastable τ) as $\mathcal{F}_{j\tau}$.

3.2. S/XB – Ionization Events per Photon

The neutral impurity particles entering the Tokamak plasma from the wall are ionized in a narrow region around the surface. Neglecting recombination processes, the inward impurity influx density is equal to the ionization rate per unit surface area, integrated over the extension of the region towards the center of the plasma. Behringer *et al.* have shown [1] that for the calculation of the inward flux, it is enough to choose an ionization degree Z of the impurity ions sufficiently small to ensure that no ions emerge from the sputtering surface in higher ionization stages. In that case, the inward flux along the line-of-sight ξ up to stage Z becomes the overall impurity flux, which is related to the integral over the abundances of the metastables of stage Z alone:

$$\Gamma = \Gamma_\sigma = \int_0^\infty n_e \sum_\sigma S_\sigma(\xi) n_\sigma(\xi) d\xi, \quad (14)$$

where S_σ is the ionization rate coefficient from metastable level σ of the impurity ion of charge Z . Since a ionization event is followed by the emission of spectral lines, it is possible to relate the number of emitted photons with the flux. Considering the radiative transition from a level j to a lower level i and supposing that the upper-level j is populated only from levels excited from a unique metastable σ , the strength of a spectral line produced by the radiative transition $j \rightarrow i$ is proportional to its emissivity

$$\epsilon_{\sigma,i \rightarrow j} = A_{ij} n_i \quad (15)$$

and the corresponding line-of-sight emissivity is defined by

$$I_{\sigma,i \rightarrow j} = \int_0^\infty \epsilon_{\sigma,i \rightarrow j}(\xi) d\xi. \quad (16)$$

Substituting in the flux equation (14) and taking into account that for only one metastable

$$\mathcal{F}_{i\sigma} = \frac{1}{n_e} \frac{n_i}{n_\sigma} \quad (17)$$

the flux becomes [1]:

$$\begin{aligned} \Gamma_\sigma &= \int_0^\infty n_e S_\sigma(\xi) n_\sigma(\xi) d\xi \approx \frac{S_\sigma}{A_{ij} \mathcal{F}_{i\sigma}} \int_0^\infty A_{ij} n_i(\xi) d\xi = \\ &= \frac{S_\sigma}{A_{ij} \mathcal{F}_{i\sigma}} I_{\sigma,i \rightarrow j} = \mathcal{SXB}_{\sigma,i \rightarrow j} I_{\sigma,i \rightarrow j}. \end{aligned} \quad (18)$$

This equation allows us to identify the Ionization Events per Photon – coefficient S/XB (for only one metastable σ) as:

$$\mathcal{SXB}_{\sigma,i \rightarrow j} = \frac{S_\sigma}{A_{ij} \mathcal{F}_{i\sigma}}. \quad (19)$$

3.3. Derivation of S/XB formula for many metastables

Transitions from metastables can compete with the direct excitation from the ground state as the population processes for the upper transition levels. That happens even for low electronic densities and also even for small metastable populations. Therefore,

we need to reconsider Eq. (18) for *metastable cross-coupling*. First of all, if we have τ metastables, we need to measure the emissivity of ($n = \tau$) different lines

$$\begin{aligned} I_1 = I_i &\equiv I_{\sigma\rho\cdots\tau, i \rightarrow j} = \int_0^\infty A_{ij} n_i(\xi) d\xi \\ I_2 = I_l &\equiv I_{\sigma\rho\cdots\tau, l \rightarrow m} = \int_0^\infty A_{lm} n_l(\xi) d\xi \\ &\dots \\ I_n = I_s &\equiv I_{\sigma\rho\cdots\tau, s \rightarrow t} = \int_0^\infty A_{st} n_s(\xi) d\xi \end{aligned}$$

in which the subscripts $\sigma\rho\cdots\tau$ indicate that, in principle, all the upper levels i, l, \dots, s can be populated by excitations from all the metastables, namely,

$$\begin{aligned} n_i &= n_e n_\sigma \mathcal{F}_{i\sigma} + n_e n_\rho \mathcal{F}_{i\rho} + \dots + n_e n_\tau \mathcal{F}_{i\tau} \\ n_l &= n_e n_\sigma \mathcal{F}_{l\sigma} + n_e n_\rho \mathcal{F}_{l\rho} + \dots + n_e n_\tau \mathcal{F}_{l\tau} \\ &\dots \\ n_s &= n_e n_\sigma \mathcal{F}_{s\sigma} + n_e n_\rho \mathcal{F}_{s\rho} + \dots + n_e n_\tau \mathcal{F}_{s\tau} \end{aligned} \quad (20)$$

which, in matricial form is written as:

$$\frac{1}{n_e} (\vec{n}_k) = (\mathbf{F}) (\vec{n}_\mu), \quad (21)$$

where the reduced n -dimensional column vectors and the corresponding $n \times n$ operators are

$$(\vec{n}_k) \equiv \begin{pmatrix} n_i \\ n_l \\ \dots \\ n_s \end{pmatrix}; \quad (\vec{n}_\mu) \equiv \begin{pmatrix} n_\sigma \\ n_\rho \\ \dots \\ n_\tau \end{pmatrix}; \quad (\mathbf{F}) \equiv \begin{pmatrix} \mathcal{F}_{i\sigma} & \mathcal{F}_{i\rho} & \dots & \mathcal{F}_{i\tau} \\ \mathcal{F}_{l\sigma} & \mathcal{F}_{l\rho} & \dots & \mathcal{F}_{l\tau} \\ \dots & \dots & \dots & \dots \\ \mathcal{F}_{s\sigma} & \mathcal{F}_{s\rho} & \dots & \mathcal{F}_{s\tau} \end{pmatrix}. \quad (22)$$

The population numbers of the metastable levels n_σ and n_ρ are:

$$(\vec{n}_\mu) = \frac{1}{n_e} (\mathbf{F})^{-1} (\vec{n}_k) = \frac{1}{n_e} (\mathbf{R}) (\vec{n}_k), \quad (23)$$

in which the inverse of the reduced matrix (\mathbf{F}) is denoted as

$$(\mathbf{R}) \equiv (\mathbf{F})^{-1}$$

The total flux is

$$\begin{aligned} \Gamma &= \Gamma_\sigma + \Gamma_\rho + \dots + \Gamma_\tau = \\ &\approx n_e S_\sigma \int_0^\infty n_\sigma + n_e S_\rho \int_0^\infty n_\rho + \dots + n_e S_\tau \int_0^\infty n_\tau = \\ &= S_\sigma \int_0^\infty (\mathcal{R}_{\sigma i} n_i + \mathcal{R}_{\sigma l} n_l + \dots + \mathcal{R}_{\sigma s} n_s) + S_\rho \int_0^\infty (\mathcal{R}_{\rho i} n_i + \mathcal{R}_{\rho l} n_l + \dots + \mathcal{R}_{\rho s} n_s) + \\ &\quad + \dots + S_\tau \int_0^\infty (\mathcal{R}_{\tau i} n_i + \mathcal{R}_{\tau l} n_l + \dots + \mathcal{R}_{\tau n} n_n) = \\ &\approx \frac{1}{A_{ij}} (S_\sigma \mathcal{R}_{\sigma i} + S_\rho \mathcal{R}_{\rho i} + \dots + S_\tau \mathcal{R}_{\tau i}) \times \int_0^\infty A_{ij} n_i + \frac{1}{A_{lm}} \sum_{\mu=\sigma}^{\tau} S_\mu \mathcal{R}_{\mu l} \times I_l + \dots + \\ &\quad + \frac{1}{A_n} \sum_{\mu=\sigma}^{\tau} S_\mu \mathcal{R}_{\mu n} \times I_n = \sum_{k=1}^n \mathcal{SXB}_k \times I_k \end{aligned} \quad (24)$$

where $A_n \equiv A_{st}$ and

$$\mathcal{SXB}_k \equiv \sum_{\mu} \mathcal{SXB}_{\mu k} = \frac{1}{A_k} \sum_{\mu} S_{\mu} \mathcal{R}_{\mu k} = \frac{1}{A_k} \sum_{\mu} S_{\mu} (\mathbf{F})_{\mu k}^{-1}. \quad (25)$$

3.4. Computational Methods

We performed the evaluation of the atomic structure and the collisional and radiative rates needed to solve the collisional-radiative problem, with three separate independent calculations. We used the atomic structure computational codes HULLAC (Hebrew University - Lawrence Livermore Atomic Code) [28, 29], the AS (AutoStructure) code [30, 31, 32], and the FAC (Flexible Atomic Code) [33]. Since the three codes are well-known in astrophysics and fusion communities, we only briefly summarize their main features. HULLAC is a fully relativistic atomic structure package of codes that provides the transition energies and the corresponding rate coefficients, among other quantities. The code implements the relativistic parameter potential method to solve the Dirac Hamiltonian. Full multiconfiguration wavefunctions are used to compute the radiative transition rates; configuration mixing and, therefore, correlation effects are included in the calculations. The Breit interaction and quantum electrodynamic (vacuum polarization and the self-energy) corrections are treated as second-order perturbations. The AUTOSTRUCTURE code allows calculating energy levels, oscillator strengths, excitation, photoionization cross sections, and autoionization rates, among other quantities. These can be calculated with configuration resolution (configuration average CA), term resolution (LS coupling), or level resolution (intermediate coupling IC) using semi-relativistic kappa-averaged wavefunctions. It also allows the inclusion of mixing configurations. FAC employs a fully relativistic approach to solve the Dirac equation. Quantum-electrodynamic effects, mainly arising from Breit interaction, vacuum polarization, and electron self-energy, are all included in standard procedures in the code. FAC has been widely employed to interpret laboratory and astrophysical spectroscopic data like the other programs mentioned.

4. Results

4.1. Atomic Structure

The ion W^{5+} is isoelectronic to the Thulium, having a ground configuration [Xe] $4f^{14}5d$. This configuration has two levels separated by about 1 eV of energy. The lowest level of the ground configuration is $(4f^{14}5d)_{\frac{3}{2}}$ and the second level is $(4f^{14}5d)_{\frac{5}{2}}$. For most of the relevant electronic densities and temperatures, both of these two levels are significantly occupied and in particular, the population of the $J = \frac{5}{2}$ is higher than those of the ground level. We found two radiative transitions well separated and distinguishable in the experimental spectra, corresponding to the $(4f^{14}5f)_{\frac{5}{2}} \rightarrow (4f^{14}5d)_{\frac{3}{2}}$ transition at $\lambda_1 = 382.13 \text{ \AA}$, and $(4f^{14}5f)_{\frac{7}{2}} \rightarrow (4f^{14}5d)_{\frac{5}{2}}$ transition at $\lambda_1 = 394.07 \text{ \AA}$.

We include in our calculations, the $4f^{14}5l$ ($l = 2 - 4$), $4f^{14}6l$ ($l = 0 - 4$), and $4f^{14}7l$ ($l = 0 - 4$). We also include configuration-interaction (CI) between those configurations and $4f^{13}5d^2$, $4f^{13}5d6s$, $4f^{13}5d6p$, $4f^{13}6s^2$, $4f^{13}6s6p$, and also with the open $5p$ -shell $4f^{14}5p^55d^2$, $4f^{14}5p^55d6s$, $4f^{14}5p^55d6p$, $4f^{14}5p^56s^2$, $4f^{14}5p^56s6p$, and $4f^{14}5p^56s6p$. This structure conforms to a total of 436 levels, the model used in HULLAC calculations. Excluding the last $4f^{14}5p^56s6p$ configuration, the structure has 430 levels, and this is the model used for the AS and FAC calculations. We show in Table 1 the detailed results obtained with the three different calculations only for the four levels involved in the transitions.

Table 1. Atomic structure for the 4-levels of W^{5+} involved in the 382.13 Å and the 394.07 Å transitions.

Index	E (eV)			Level
	HULLAC	AS	FAC	
1	0.0000	0.0000	0.0000	$(4f^{14}5d)_{3/2}$
2	0.9718	1.3653	1.0142	$(4f^{14}5d)_{5/2}$
3	31.769	31.823	31.729	$(4f^{14}5f)_{5/2}$
4	31.841	31.973	31.814	$(4f^{14}5f)_{7/2}$

Figure 3 shows the spectra observed from EAST discharge (#100300) at a time $t = 5.200$ s, corresponding to the maximum of the W burst. The figure also includes our three independent W^{5+} synthetic spectra, which were obtained by solving the fully collisional-radiative model.

As the figure shows, the agreement between the three theoretical calculations is satisfactory, although the results are consistently higher than the experimental values by about 7 Å. Surprisingly, another simpler calculation, including only 20 levels from the $4f^{14}5p^65l$ ($l = 2 - 4$), $4f^{14}5p^66l$ ($l = 0 - 3$), and $4f^{14}5p^67l$ ($l = 0 - 3$) produces transition energies much closer than the experimental values, above the experimental for only 2 Å, in all three cases. However, these calculations do not take into account the configuration mixings between the $4f^{14}5f$ and the $4f^{13}5d^2$ configurations, which account for a total mixing of about 11% for both the $J = 5/2$ and $J = 7/2$ levels. This indicates that the latter model cannot be considered valid, and some serendipity is being introduced, which deserves to be investigated in the future.




Figure 3. EAST spectrum and theoretical collisional-radiative synthetic spectra for the $5f \rightarrow 5d$ transitions of the W^{5+} ion. (a) EAST Experimental spectra (#100300, at $t = 5.200$). (b) HULLAC calculation. (c) AS (d) FAC.

4.2. Excitation and Radiative Rates

The three codes have been employed for the calculation of the electron-impact excitation rate coefficients. The three codes use the distorted-waves approximation, which has been quite successful in the determination of excitation cross-sections for highly ionized species, where the effects of correlations are usually not significant. For W^{5+} , the coupling in the continuum is sufficiently small to allow the use of this approximation with reasonable accuracy. The overall agreement between the three codes is about 40%, for the range $20 < T_e < 100$ eV.


We also have calculated the radiative transition rate coefficients with the three codes, obtaining values that agree in a 20% for the dipole-allowed transitions. The values of the radiative transitions and the electron-impact excitation rate coefficients for the four levels involved in the $5f \rightarrow 5d$ lines are listed in Table 2, for HULLAC, AUTOSTRUCTURE, and FAC calculations.

Table 2. Radiative transitions A and electron-impact excitation rate coefficients (in cm^3s^{-1}) for the 4 levels involved in the W^{5+} lines. Note that $aE\pm b$ means $a \times 10^{\pm b}$.

HULLAC								
Lower	Upper	A (s^{-1})	T_e (eV)					
			20	40	60	80	90	100
1	2	5.20E+00	3.78E-09	2.47E-09	1.89E-09	1.56E-09	1.45E-09	1.35E-09
1	3	1.61E+10	3.36E-09	5.88E-09	6.84E-09	7.27E-09	7.40E-09	7.49E-09
2	3	1.16E+09	4.54E-10	6.56E-10	6.80E-10	6.68E-10	6.59E-10	6.49E-10
1	4	1.01E+01	3.29E-10	4.27E-10	4.01E-10	3.62E-10	3.44E-10	3.27E-10
2	4	1.83E+10	3.92E-09	6.66E-09	7.65E-09	8.09E-09	8.21E-09	8.30E-09
3	4	2.18E-03	4.91E-09	3.15E-09	2.42E-09	2.02E-09	1.87E-09	1.75E-09
AUTOSTRUCTURE								
Lower	Upper	A (s^{-1})	T_e (eV)					
			20	40	60	80	90	100
1	2	1.44E+01	8.62E-09	4.81E-09	3.36E-09	2.61E-09	2.35E-09	2.15E-09
1	3	1.64E+10	2.74E-09	4.94E-09	5.89E-09	6.37E-09	6.52E-09	6.63E-09
2	3	1.08E+09	4.50E-10	6.14E-10	6.27E-10	6.16E-10	6.08E-10	6.00E-10
1	4	1.04E+01	3.63E-10	4.36E-10	3.96E-10	3.53E-10	3.34E-10	3.17E-10
2	4	1.71E+10	3.34E-09	5.76E-09	6.74E-09	7.22E-09	7.36E-09	7.47E-09
3	4	2.02E-02	1.01E-08	5.46E-09	3.79E-09	2.95E-09	2.67E-09	2.44E-09
FAC								
Lower	Upper	A (s^{-1})	T_e (eV)					
			20	40	60	80	90	100
1	2	5.90E+00	5.07E-09	3.19E-09	2.40E-09	1.96E-09	1.81E-09	1.68E-09
1	3	1.28E+10	2.41E-09	4.27E-09	5.00E-09	5.35E-09	5.46E-09	5.54E-09
2	3	9.40E+08	4.31E-10	6.04E-10	6.16E-10	5.98E-10	5.87E-10	5.76E-10
1	4	9.29E+00	3.39E-10	4.31E-10	3.99E-10	3.56E-10	3.36E-10	3.18E-10
2	4	1.50E+10	3.11E-09	5.33E-09	6.16E-09	6.53E-09	6.64E-09	6.72E-09
3	4	3.72E-03	5.70E-09	3.39E-09	2.57E-09	2.14E-09	1.99E-09	1.86E-09

4.3. Electron-Impact Ionization Rates

For the calculation of the electron-impact ionization rate coefficients, we used the distorted-waves approximation. However, even by using this relatively simple approach, the calculations are cumbersome. First, the total direct ionization cross section σ^{DI} from the ground configuration levels involves the sum of the contribution of the ionization of the $4f$, $5s$, $5p$, and $5d$ subshells-electrons:




Second, the indirect pathway to ionization, through inner-shell excited intermediate levels must be taken into consideration since the cross-sections of these processes are much higher than the corresponding to the direct ionization, as indicated in the experimental ionization measurements by Stenke *et al.* [34] and by Spruck *et al.* [35]. The total cross-section σ_C^{EA} for excitation-autoionization from an initial level g , to any final level k of the following ion, through inner-shell excitation to any intermediate autoionizing level j within a given configuration or complex C is given by [36]

$$\sigma_C^{EA}(E) = \sum_{j \in C} \sigma_{gj}(E) \left[\frac{\sum_k A_{jk}^a + \sum_i A_{ji} B_i^a}{\sum_k A_{jk}^a + \sum_i A_{ji}} \right] \equiv \sum_{j \in C} \sigma_{gj}(E) B_j^a \quad (27)$$

where $\sigma_{gj}(E)$ is the cross section for electron-impact excitation from g to j as a function of the incident electron kinetic energy E . A_{jk}^a is the rate coefficient for autoionization from j to k , and A_{ji} is the Einstein coefficient for spontaneous emission from j to any lower-lying level i . B_j^a is the multiple branching ratio for autoionization from level j , defined by the bracket term. This term contains, in turn, the effective branching ratio B_i^a for further autoionization from level i , defined by a similar recursive expression [36]. The total excitation-autoionization cross-section is given by

$$\sigma^{EA}(E) = \sum_C \sigma_C^{EA}(E). \quad (28)$$

In our calculation, we include excitations from the $4f$, $5p$, and $5s$ subshells of the ground configuration. These excitations are graphically represented as follows:



where we take all the possibles $5 \leq n \leq 15$ and $l \leq 4$.

In a low-ionized species such as W^{5+} , the radiative rates from the doubly excited levels are quite small compared to the autoionizing rates. Therefore, we can neglect the multiple branching ratios in (27). In that way, we assume that once the excited electron reaches an autoionizing level, it autoionizes and does not follow further cascades.

The total electron-impact single ionization cross-section is

$$\sigma^{EIA}(E) = \sigma^{DI}(E) + \sigma_C^{EA}(E). \quad (30)$$

Since the calculations are demanding, we only use one computational code for the ionization cross-section results. Also, for comparisons with other calculations, we choose to use the FAC code. In Fig. 4, we show the contribution of the different channels to the electron-impact ionization cross-sections of W^{5+} . In the left part of the figure, we concentrate on the indirect excitation-autoionization channels σ^{EA} . The figure shows the accumulated contribution from the essential manifolds (dashed-color curves), together with the theoretical values reported by Jonauskas *et al.* [37]. The results are in excellent agreement and the slight differences can be attributed to the lack of some configuration interactions that we have not included in the calculations. On the other hand, Jonauskas *et al.* calculated the EA contribution up to the $n = 12$ manifold, whereas we also included the $n = (14, 15)$ configurations.

In the right part of the figure, the total ionization cross-sections σ^{EIA} are displayed. We intended to show the individual contribution of the ionization from the ground level $(4f^{14}5d)_{\frac{3}{2}}$ and from the metastable $(4f^{14}5d)_{\frac{5}{2}}$, but both cross-sections are very similar and are not easily distinguishable in the graph. The figure shows our results, together with the theoretical calculations reported by Jonauskas *et al.* [37] and also the results given by Zhang *et al.* [38]. As expected, the agreement is excellent. The differences can be attributed to the configurations used in the potential optimization and the configurations considered for the mixing interactions. To be able to assess the importance of the indirect channels, we include in the figure the total direct ionization σ^{DI} in dashed lines. The figure also shows the experimental results given by Stenke *et al.* [34], and more recent calculations from Spruck *et al.* [35]. The distorted-waves are known for generally overestimation of the cross-sections. Both experimental results show a high presence of metastable contributions, so the discrepancies between theoretical and experimental results could be even more significant.

Table 3. Total ionization rate coefficients from the metastable levels of W^{5+} , in cm^3s^{-1} . Note that aE-b means $a \times 10^{-b}$.

Metastable	T_e (eV)					
	20	40	60	80	90	100
1	2.01E-09	1.15E-08	2.04E-08	2.70E-08	2.96E-08	3.18E-08
2	2.07E-09	1.18E-08	2.08E-08	2.75E-08	3.01E-08	3.23E-08




Figure 4. Left: Excitation-Autoionization cross-sections from the ground state of W^{5+} . Color-dashed lines: Present calculations of the accumulated partial EA cross sections σ_C^{EA} for the principal manifolds C . Color squares: Theoretical results from Jonauskas *et al.* [37]. Red circles: Total theoretical results by Zhang *et al.* [38]. Black-solid lines: Present calculations for total σ^{EA} , considering $n \leq 15$.

Right: Total ionization cross-sections from the ground level. Black solid line: Present calculations of the total ionization cross-section σ^{EIA} (Eq. 30). Black-dashed line: Total direct-ionization cross-section σ^{DI} . Green-dashed line: Theoretical results from Jonauskas *et al.* [37]. Red-dot dashed line: Calculations by Zhang *et al.* [38]. Blue squares: Experimental results of Stenke *et al.* [34]; Green circles: Experimental results of Spruck *et al.* [35].

4.4. S/XB results

4.4.1. Simple Model (4-levels)

We found that a simple model having only four levels, namely, the ground level $(4f^{14}5d)_{3/2}$, the metastable $(4f^{14}5d)_{5/2}$, and the two upper-excited levels, $(4f^{14}5f)_{5/2}$ and $(4f^{14}5f)_{7/2}$, can be used to explain qualitatively many of the basic features in the determination of impurity flux. A detailed derivation of the S/XB partial coefficients is given in Appendix A. The final expressions for these coefficients are given in Eq. (A.18).

We show, in Figure 5, the S/XB coefficients corresponding to the transition $3 \rightarrow 1$ (upper part (a)), and in the lower part (b), the coefficients corresponding to the $4 \rightarrow 2$ transition. The coefficients for this table have been calculated using the rates obtained with FAC code for a typical electronic temperature $T_e = 60$ eV. As is expected, due to the high population of the metastable, the leading terms are the diagonal SXB_{11} and SXB_{21}




Figure 5. S/XB coefficients, according to the definition in Eq. (25), for a 4-level model. (a) S/XB for the $3 \rightarrow 1$ transition $((4f^{14}5f)_{5/2} \rightarrow (4f^{14}5d)_{3/2})$. (b) $4 \rightarrow 2$ transition $((4f^{14}5f)_{7/2} \rightarrow (4f^{14}5d)_{5/2})$, in W^{5+} . The electronic temperature is $T_e = 60$ eV. The low-density limit curves correspond to the approximation given in Eq. (31).

We can perform a further approximation, focusing on the low-density region, in order to understand the overall behavior of the S/XB coefficients. In this approximation, we neglect first the collisional decays that are proportional to the electron density and compete with stronger radiative decays. We also make use of the rates listed in Table 2, neglecting the relatively small terms, for example, $(A_{41} + A_{42} + A_{43}) \approx A_{42}$. Under

these assumptions, the low-density limit expressions for the S/XB coefficients are

$$\begin{aligned}
 S\mathcal{X}B_{11} &\approx \frac{S_1}{Q_{13} \frac{A_{31}}{A_{31}+A_{32}}} \\
 S\mathcal{X}B_{21} &\approx -\frac{S_2}{Q_{24} \frac{Q_{13}}{Q_{14}}} \\
 S\mathcal{X}B_{12} &\approx -\frac{S_1}{Q_{13} \frac{Q_{24}}{Q_{23}}} \\
 S\mathcal{X}B_{22} &\approx \frac{S_2}{Q_{24}}
 \end{aligned} \tag{31}$$

which are all density independent. As is shown in Figure 5, the approximated S/XB results are in excellent agreement with the exact expressions given in (A.18). The results of this approximation not only match the exact results at low electronic densities but this agreement is still maintained for densities that are above the standard operating values, even up to values such as $n_e \approx 10^{17} \text{ cm}^{-3}$.

We made comparisons between the S/XB coefficients resulting from the three independent calculations from HULLAC, AS, and FAC computational codes. The results are displayed in Figure 6 for an electronic temperature $T_e = 60 \text{ eV}$, and as before, the upper




Figure 6. Comparisons of the S/XB coefficients obtained with the three independent calculations using HULLAC, AS, and FAC codes for the 4-level model. (a) S/XB for the $3 \rightarrow 1$ transition $((4f^{14}5f)_{5/2} \rightarrow (4f^{14}5d)_{3/2})$. (b) $4 \rightarrow 2$ transition $((4f^{14}5f)_{7/2} \rightarrow (4f^{14}5d)_{5/2})$, in W^{5+} . The electronic temperature is $T_e = 60 \text{ eV}$.

The same results are shown in Figure 7 for the electronic density $n_e = 10^{13} \text{ cm}^{-3}$, which is within the normal range of operation in the EAST plasmas. The discrepancies between the different calculations are consistent with the differences obtained for the values




Figure 7. Comparisons of the S/XB coefficients obtained for the 4-level model. (a) S/XB for the $3 \rightarrow 1$ transition $((4f^{14}5f)_{5/2} \rightarrow (4f^{14}5d)_{3/2})$. (b) $4 \rightarrow 2$ transition $((4f^{14}5f)_{7/2} \rightarrow (4f^{14}5d)_{5/2})$, in W^{5+} . The electronic density is $n_e = 10^{13} \text{ cm}^{-3}$.

4.4.2. Full Model (430-levels)

We included all the configurations listed in Sec. 4.1 and solved the corresponding collisional-radiative equations, obtaining the S/XB ratios for the same transitions studied previously. The results are displayed in Fig. 8 for a fixed electronic temperature $T_e = 60$ eV and in Fig. 9, for a fixed electronic density $n_e = 10^{13}$ cm $^{-3}$. We note that for both lines, the S/XB ratios appear nearly constant around an electron density from 10^4 cm $^{-3}$, to a density of approximately 10^{14} cm $^{-3}$. This justifies the previous approach in which we considered only four levels and the possibility of using the low-density approximation to obtain a simple and quick determination of the incoming flux. At densities around 10^{15} cm $^{-3}$ the S/XB becomes sensitive to the collisional-radiative solutions. Therefore, we found substantial differences between the calculations, particularly between HULLAC results and the other two. Beyond $n_e > 10^{17}$ cm $^{-3}$, the plasma reaches the local thermodynamic equilibrium density regime, and then the S/XB ratios




Figure 8. Comparisons of the S/XB coefficients obtained with the three independent calculations using HULLAC, AS, and FAC codes, for the 430-level model, in W^{5+} . (a) S/XB for the $(4f^{14}5f)_{5/2} \rightarrow (4f^{14}5d)_{3/2}$ transition. (b) $(4f^{14}5f)_{7/2} \rightarrow (4f^{14}5d)_{5/2}$ transition. The electronic temperature is $T_e = 60$ eV.

The detailed results for electronic densities in the range 10^4 cm $^{-3} < n_e < 10^{18}$ cm $^{-3}$ and electronic temperatures in the range $20 < T_e < 100$ eV are presented in Tables 4, 5, and 6 for the calculations realized with HULLAC, AUTOSTRUCTURE, and FAC, respectively. The agreement between the calculations for the $(4f^{14}5f)_{5/2} \rightarrow (4f^{14}5d)_{3/2}$




Figure 9. Comparisons of the S/XB coefficients obtained for the 430-level model, in W^{5+} . (a) S/XB for the $(4f^{14}5f)_{5/2} \rightarrow (4f^{14}5d)_{3/2}$ transition. (b) $(4f^{14}5f)_{7/2} \rightarrow (4f^{14}5d)_{5/2}$ transition. The electronic density is $n_e = 10^{13} \text{ cm}^{-3}$.

transition (the $3 \rightarrow 1$ transition in the 4-level model) is of about 40% for $T_e = 50 \text{ eV}$ and densities up to $n_e = 10^{15} \text{ cm}^{-3}$ in the worst case (HULLAC vs. FAC). Fig. 9 shows that the agreement is much better at lower temperatures. For the $(4f^{14}5f)_{7/2} \rightarrow (4f^{14}5d)_{5/2}$ transition, the agreement at $T_e = 50 \text{ eV}$ is about 25%, it is much better for lower temperatures.

Table 4. SXB of W^{5+} for the radiative transitions (Line 1) $(4f^{14}5f)_{5/2} \rightarrow (4f^{14}5d)_{3/2}$, and (Line 2) $(4f^{14}5f)_{7/2} \rightarrow (4f^{14}5d)_{5/2}$, from a 436-level collisional-radiative model, calculated with the HULLAC computational suite.

Line	T_e (eV)	Electron density (cm^{-3})							
		10^4	10^6	10^8	10^{10}	10^{12}	10^{14}	10^{16}	10^{18}
1	20	0.543	0.532	0.525	0.519	0.502	0.504	0.804	1.710
	40	1.733	1.698	1.690	1.674	1.629	1.677	3.039	7.143
	60	2.626	2.577	2.569	2.549	2.492	2.579	4.798	11.90
	80	3.266	3.209	3.203	3.180	3.119	3.235	6.059	15.52
	90	3.516	3.458	3.452	3.429	3.367	3.494	6.543	16.92
	100	3.736	3.676	3.670	3.648	3.586	3.722	6.966	18.20
2	20	0.425	0.414	0.412	0.408	0.393	0.384	0.570	1.470
	40	1.391	1.354	1.367	1.355	1.318	1.331	2.258	5.758
	60	2.130	2.076	2.105	2.089	2.045	2.103	3.712	9.203
	80	2.661	2.598	2.636	2.620	2.575	2.678	4.805	11.70
	90	2.873	2.807	2.848	2.833	2.790	2.912	5.250	12.72
	100	3.056	2.988	3.032	3.017	2.975	3.116	5.639	13.60

Table 5. SXB of W^{5+} for the radiative transitions (Line 1) $(4f^{14}5f)_{5/2} \rightarrow (4f^{14}5d)_{3/2}$, and (Line 2) $(4f^{14}5f)_{7/2} \rightarrow (4f^{14}5d)_{5/2}$, from a 430-level collisional-radiative model, calculated with the AS computational code.

Line	T_e (eV)	Electron density (cm^{-3})							
		10^4	10^6	10^8	10^{10}	10^{12}	10^{14}	10^{16}	10^{18}
1	20	0.645	0.654	0.644	0.637	0.580	0.395	0.343	0.877
	40	2.076	2.107	2.099	2.080	1.921	1.352	1.136	2.457
	60	3.140	3.178	3.176	3.153	2.952	2.158	1.779	3.568
	80	3.889	3.927	3.928	3.905	3.694	2.787	2.272	4.356
	90	4.184	4.221	4.224	4.201	3.990	3.049	2.476	4.669
	100	4.442	4.476	4.480	4.457	4.249	3.285	2.657	4.929
2	20	0.469	0.480	0.479	0.474	0.430	0.293	0.249	0.674
	40	1.575	1.609	1.638	1.623	1.487	1.048	0.918	2.477
	60	2.431	2.474	2.526	2.507	2.326	1.699	1.487	3.913
	80	3.046	3.090	3.155	3.135	2.939	2.210	1.930	4.965
	90	3.293	3.336	3.404	3.385	3.186	2.425	2.115	5.390
	100	3.505	3.547	3.618	3.599	3.400	2.617	2.281	5.765

Table 6. SXB of W^{5+} for the radiative transitions (Line 1) $(4f^{14}5f)_{5/2} \rightarrow (4f^{14}5d)_{3/2}$, and (Line 2) $(4f^{14}5f)_{7/2} \rightarrow (4f^{14}5d)_{5/2}$, from a 430-level collisional-radiative model, calculated with the FAC computational code.

Line	T_e (eV)	Electron density (cm^{-3})							
		10^4	10^6	10^8	10^{10}	10^{12}	10^{14}	10^{16}	10^{18}
1	20	0.762	0.759	0.745	0.740	0.696	0.529	0.484	1.328
	40	2.484	2.479	2.448	2.436	2.326	1.828	1.670	4.094
	60	3.796	3.792	3.752	3.737	3.601	2.896	2.595	6.026
	80	4.730	4.725	4.681	4.666	4.525	3.710	3.273	7.351
	90	5.096	5.091	5.045	5.031	4.892	4.046	3.550	7.873
	100	5.414	5.409	5.361	5.348	5.211	4.343	3.790	8.314
2	20	0.520	0.518	0.512	0.510	0.482	0.371	0.323	0.830
	40	1.748	1.747	1.745	1.741	1.660	1.311	1.173	2.975
	60	2.707	2.709	2.714	2.710	2.603	2.098	1.866	4.619
	80	3.407	3.411	3.421	3.419	3.301	2.702	2.384	5.764
	90	3.685	3.690	3.702	3.701	3.580	2.951	2.595	6.203
	100	3.926	3.932	3.945	3.946	3.824	3.173	2.780	6.579

4.5. Tungsten Influx

The absolute calibration of the spectral line intensities [19, 21, 22, 20] results in values of the $5f \rightarrow 5d$ transitions about 3×10^{13} photons per cm^2 and per second at the emission peak corresponding to the tungsten burst. Assuming that the tungsten ions are sputtered from a region in which the electronic temperature is $T_e = 60$ eV and the electronic density is $n_e = 10^{13} \text{ cm}^{-3}$, the S/XB ratios are of the order of 3, we can infer a total tungsten flux of the order of 10^{14} particles per cm^2 and per second at the peak. Indeed, we show in Fig. 10 the calculated total tungsten flux from the plasma-facing surface towards the interior of the EAST device, which follows the behavior of the line intensity in time for discharge #100300, having a peak at $t = 5.2$ seconds. The fluxes are calculated using three independent calculations: HULLAC, AS, and FAC. The three calculations agree at the peak between 45% in the worst comparison case (HULLAC vs. FAC).




Figure 10. Total tungsten influx from the plasma-facing surface, calculated for the discharge #100300 at EAST, as a function of time.

5. Conclusions

We have developed the equations for calculating the S/XB ratios in a general case where many metastable levels may contribute to the ionization of the ion. We do not assume that the populations of the metastable terms are in quasistatic equilibrium with the population of the ground term. This approach eliminates the dependence of our results on the populations of the long-lived states.

Extensive theoretical calculations of atomic structure, radiative decay, electron-impact excitation, and electron-impact ionization have been carried out to generate the data necessary to model the W^{5+} ion in a plasma environment. We use three different atomic computational codes – FAC, HULLAC, and AUTOSTRUCTURE – which allow us to assess the accuracy of our theoretical results. For the electron-impact ionization, we included contributions from excitation-autoionization processes up to $n = 15$ manifolds and calculated the rate coefficients for the total ionization from both the two $4f^{14}5d$ configuration levels.

Using these data, we solved a collisional-radiative model to obtain the effective population coefficients needed in to determine the S/XB ratios. We selected the radiative transitions $(4f^{14}5f)_{5/2} \rightarrow (4f^{14}5d)_{3/2}$ and $(4f^{14}5f)_{5/2} \rightarrow (4f^{14}5d)_{3/2}$, which are prominent in the 320-460 Å range. The process of radiative cascade from higher energy terms than the upper terms involved in the radiative transition was found to have a negligible effect on the S/XB ratios for both types of transitions at low and intermediate energy densities. Therefore, by using a simple four-level collisional-radiative model consisting of the two upper levels and the two lower metastables, we can generate S/XB ratios that agree very well with the 430-level model. That is particularly useful because it allows us to derive simple analytical expressions that help us understand the principal mechanisms contributing to the ionization.

6. Acknowledgments

This work was supported by the National Magnetic Confinement Fusion Energy R & D Program of China (Grant Nos. 2022YFE03180400, 2022YFE03020004, 2019YFE030403), National Natural Science Foundation of China (Grant No. 12322512), and Chinese Academy of Sciences President’s International Fellowship Initiative (PIFI) (Grant Nos. 2024PVA0074). DM acknowledges partial support from CONICET by Project No. PIP11220200102421CO, and the ANPCyT by Project No. PICT- 2020-SERIE A-01931 in Argentina, and the Alliance of International Science Organizations (ANSO) Visiting Fellowship (ANSO-VF-2021-03), in China.

Appendix A. Derivation of S/XB coefficients for 4-levels and 2 metastables

The collisional-radiative equations (2) for 4-levels is:

$$\begin{pmatrix} \frac{dn_1}{dt} \\ \frac{dn_2}{dt} \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} C_{11} & C_{12} & C_{13} & C_{14} \\ C_{21} & C_{22} & C_{23} & C_{24} \\ C_{31} & C_{32} & C_{33} & C_{34} \\ C_{41} & C_{42} & C_{43} & C_{44} \end{pmatrix} \begin{pmatrix} n_1 \\ n_2 \\ n_3 \\ n_4 \end{pmatrix}, \quad (\text{A.1})$$

and for the excited-levels part of the matrix, the corresponding system of equations is

$$\begin{pmatrix} C_{31}n_1 + C_{32}n_2 + C_{33}n_3 + C_{34}n_4 \\ C_{41}n_1 + C_{42}n_2 + C_{43}n_3 + C_{44}n_4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (\text{A.2})$$

which is equivalent to

$$\begin{pmatrix} C_{33}n_3 + C_{34}n_4 \\ C_{43}n_3 + C_{44}n_4 \end{pmatrix} = \begin{pmatrix} -C_{31}n_1 - C_{32}n_2 \\ -C_{41}n_1 - C_{42}n_2 \end{pmatrix}. \quad (\text{A.3})$$

In matricial form:

$$\mathbf{C}_s \cdot \vec{n}_s = \vec{C}_1 \cdot n_1 + \vec{C}_2 \cdot n_2 \quad (\text{A.4})$$

where

$$\vec{n}_s = \begin{pmatrix} n_3 \\ n_4 \end{pmatrix}, \quad (\text{A.5})$$

the reduced matrix for the excited levels is

$$\mathbf{C}_s = \begin{pmatrix} C_{33} & C_{34} \\ C_{43} & C_{44} \end{pmatrix} \quad (\text{A.6})$$

and the column vectors are

$$\vec{C}_1 = \begin{pmatrix} -C_{31} \\ -C_{41} \end{pmatrix} \quad (\text{A.7})$$

and

$$\vec{C}_2 = \begin{pmatrix} -C_{32} \\ -C_{42} \end{pmatrix}. \quad (\text{A.8})$$

The solution for the excited levels is

$$\begin{aligned} \vec{n}_s &= [\mathbf{C}_s]^{-1} \cdot \vec{C}_1 \cdot n_1 + [\mathbf{C}_s]^{-1} \cdot \vec{C}_2 \cdot n_2 = \\ &= \vec{\mathcal{F}}_1 \cdot n_1 + \vec{\mathcal{F}}_2 \cdot n_2 \end{aligned} \quad (\text{A.9})$$

in which the column vectors are

$$\vec{\mathcal{F}}_1 \equiv [\mathbf{C}_s]^{-1} \cdot \vec{C}_1 \quad \text{and} \quad \vec{\mathcal{F}}_2 \equiv [\mathbf{C}_s]^{-1} \cdot \vec{C}_2. \quad (\text{A.10})$$

The inverse of \mathbf{C}_s is

$$[\mathbf{C}_s]^{-1} = \frac{1}{C_{33}C_{44} - C_{34}C_{43}} \begin{pmatrix} C_{44} & -C_{34} \\ -C_{43} & C_{33} \end{pmatrix} \quad (\text{A.11})$$

and

$$(\mathbf{F}) = \begin{pmatrix} \vec{\mathcal{F}}_1 & \vec{\mathcal{F}}_2 \end{pmatrix} \quad (\text{A.12})$$

becomes

$$(\mathbf{F}) = \frac{1}{C_{33}C_{44} - C_{34}C_{43}} \begin{pmatrix} -C_{31}C_{44} + C_{34}C_{41} & -C_{32}C_{44} + C_{34}C_{42} \\ C_{31}C_{43} - C_{33}C_{41} & C_{32}C_{43} - C_{33}C_{42} \end{pmatrix}. \quad (\text{A.13})$$

The inverse of the reduced matrix is

$$(\mathbf{R}) \equiv (\mathbf{F})^{-1} = \frac{1}{C_{31}C_{42} - C_{32}C_{41}} \begin{pmatrix} C_{32}C_{43} - C_{33}C_{42} & C_{32}C_{44} - C_{34}C_{42} \\ -C_{31}C_{43} + C_{33}C_{41} & -C_{31}C_{44} + C_{34}C_{41} \end{pmatrix}. \quad (\text{A.14})$$

From the definition (25) for the S/XB coefficients

$$\mathcal{SXB}_{\mu k} \equiv \frac{1}{A_k} S_\mu \mathcal{R}_{\mu k} \quad (\text{A.15})$$

we obtain

$$\mathcal{SXB} = \frac{1}{C_{31}C_{42} - C_{32}C_{41}} \begin{pmatrix} \frac{S_1}{A_{31}} (C_{32}C_{43} - C_{33}C_{42}) & \frac{S_2}{A_{31}} (-C_{31}C_{43} + C_{33}C_{41}) \\ \frac{S_1}{A_{42}} (C_{32}C_{44} - C_{34}C_{42}) & \frac{S_2}{A_{42}} (-C_{31}C_{44} + C_{34}C_{41}) \end{pmatrix} \quad (\text{A.16})$$

and replacing the corresponding elements of the matrix (Eq. (3)), we obtain finally the expressions for the S/XB coefficients

$$\begin{aligned} \mathcal{SXB}_{11} &= \frac{n_e S_1 n_e Q_{23} n_e Q_{34} - n_e Q_{24} (-A_{31} - A_{32} - n_e Q_{31} - n_e Q_{32} - n_e Q_{34})}{A_{31} n_e Q_{13} n_e Q_{24} - n_e Q_{14} n_e Q_{23}} \quad (\text{A.17}) \\ \mathcal{SXB}_{21} &= \frac{n_e S_2 - n_e Q_{13} n_e Q_{34} + n_e Q_{14} (-A_{31} - A_{32} - n_e Q_{31} - n_e Q_{32} - n_e Q_{34})}{A_{31} n_e Q_{13} n_e Q_{24} - n_e Q_{14} n_e Q_{23}} \\ \mathcal{SXB}_{12} &= \frac{n_e S_1 n_e Q_{23} (-A_{41} - A_{42} - A_{43} - n_e Q_{41} - n_e Q_{42} - n_e Q_{43}) - n_e Q_{24} (A_{43} + n_e Q_{43})}{A_{42} n_e Q_{13} n_e Q_{24} - n_e Q_{14} n_e Q_{23}} \\ \mathcal{SXB}_{22} &= \frac{n_e S_2 n_e Q_{13} (A_{41} + A_{42} + A_{43} + n_e Q_{41} + n_e Q_{42} + n_e Q_{43}) + n_e Q_{14} (A_{43} + n_e Q_{43})}{A_{42} n_e Q_{13} n_e Q_{24} - n_e Q_{14} n_e Q_{23}}. \end{aligned}$$

References

- [1] K. Behringer, H.P. Summers, B. Denne, M. Forrest, and M. Stamp, “Spectroscopic determination of impurity influx from localized surfaces”, *Plasma Phys. Control Fusion* **31**, 2059 (1989).
- [2] A.R. Field, C. Garcia-Rosales, G. Lieder, C.S. Pitcher, and R. Radtke, “Spectroscopic measurement of target plate erosion in the ASDEX Upgrade divertor”, *Nucl. Fusion* **36**, 119 (1996).
- [3] N.R. Badnell, T.W. Gorczyca, M.S. Pindzola, and H.P. Summers, “Excitation and ionization of neutral Cr and Mo, and the application to impurity influx”, *J. Phys. B* **29**, 3683 (1996).
- [4] D.C. Griffin, M.S. Pindzola, J.A. Shaw, N.R. Badnell, M. O’Mullane, and H.P. Summers, “Electron-impact excitation and ionization of Ar^+ for the determination of impurity influx in tokamaks”, *J. Phys. B* **30**, 3543 (1997).
- [5] B. Lipschultz, D.A. Pappas, B. LaBombard, J.E. Rice, D. Smith, and S.J. Wukitch, “A study of molybdenum influxes and transport in Alcator C-Mod”, *Nucl. Fusion* **41**, 585 (2001).
- [6] J.W. Coenen, S. Antusch, M. Aumann, W. Biel, J. Du, J. Engels, S. Heuer, A. Houben, T. Hoeschen, B. Jasper, *et al.*, “Materials for demo and reactor applications—boundary conditions and new concepts”, *Phys. Scrip.* **T167**, 014002 (2015).
- [7] A. Thoma, K. Asmussen, R. Dux, K. Krieger, A. Herrmann, B. Napiontek, R. Neu, J. Steinbrink, M. Weinlich, U. Wenzel, *et al.*, “Spectroscopic measurements of tungsten erosion in the ASDEX Upgrade divertor”, *Plasma Phys. Control. Fusion* **39**, 1487 (1997).
- [8] A. Geier, H. Maier, R. Neu, K. Krieger, and the ASDEX Upgrade Team, “Determination of the tungsten divertor retention at ASDEX Upgrade using a sublimation probe”, *Plasma Phys. and Contr. Fus.* **44**, 2091 (2002).
- [9] D. Nishijima, R.P. Doerner, M.J. Baldwin, A. Pospieszczyk, and A. Kreter, “Erratum: Experimental determination of S/XB values of W I visible lines”, *Phys. of Plasm.* **18**, 019901 (2011).
- [10] I. Beigman, A. Pospieszczyk, G. Sergienko, I. Yu. Tolstikhina, and L. Vainshtein, “Tungsten spectroscopy for the measurement of W-fluxes from plasma facing components”, *Plasma Phys. Control. Fusion* **49**, 1833 (2007).
- [11] A. Pospieszczyk, D. Borodin, S. Brezinsek, A. Huber, A. Kirschner, Ph. Mertens, G. Sergienko, B. Schweer, I.L. Beigman, and L. Vainshtein, “Determination of rate coefficients for fusion-relevant atoms and molecules by modelling and measurement in the boundary layer of textor”, *J. Phys. B* **43**, 144017 (2010).
- [12] C.P. Ballance, S.D. Loch, M.S. Pindzola, and D.C. Griffin, “Electron-impact excitation and ionization of W^{3+} for the determination of tungsten influx in a fusion plasma”, *J. Phys. B* **46**, 055202 (2013).
- [13] C.F. Dong, S. Morita, Z.Y. Cui, P. Sun, K. Zhang, I. Murakami, B.Y. Zhang, Z.C. Yang, D.L. Zheng, L. Feng, *et al.*, “Evaluation of tungsten influx rate and study of edge tungsten behavior based on the observation of EUV line emissions from W^{6+} ions in HL-2A”, *Nucl. Fus.* **59**, 016020 (2019).
- [14] “Open-ADAS, Atomic Data and Analysis Structure”, <https://open.adas.ac.uk>
- [15] B.N. Wan, Y.F. Liang, X.Z. Gong, J.G. Li, N. Xiang, G.S. Xu, Y.W. Sun, L. Wang, J.P. Qian, and H.Q. Liu, “Overview of EAST experiments on the development of high-performance steady-state scenario”, *Nucl. Fus.* **57**, 102019 (2017).
- [16] B.N. Wan, Y. Liang, X.Z. Gong, N. Xiang, G.S. Xu, Y. Sun, L. Wang, J.P. Qian, H.Q. Liu, and L. Zeng, “Recent advances in EAST physics experiments in support of steady-state operation for ITER and CFETR”, *Nucl. Fus.* **59**, 112003 (2019).
- [17] L. Wang, G.S. Xu, J.S. Hu, K.D. Li, Q.P. Yuan, J.B. Liu, F. Ding, Y.W. Yu, Z.P. Luo, J.C. Xu, *et al.*, “Progress of divertor heat and particle flux control in EAST for advanced steady-state operation in the last 10 years”, *Jour. of Fusion Energy* **40**, 1 (2021).
- [18] B.N. Wan, X.Z. Gong, Y. Liang, N.g Xiang, G. S. Xu, Y. Sun, L. Wang, J.P. Qian, H.Q. Liu, and B. Zhang, “Advances in the long-pulse steady-state high beta H-mode scenario with active

- controls of divertor heat and particle fluxes in EAST”, Nucl. Fus. **62**, 042010 (2022).
- [19] Z. Ling, S. Morita, X. Zong, Z. Wu, P. Zhang, C. Wu, W. Gao, T. Ohishi, M. Goto, and J. Shen, “A fast-time-response extreme ultraviolet spectrometer for measurement of impurity line emissions in the Experimental Advanced Superconducting Tokamak”, Rev. of Sci. Instr. **86**, 123509 (2015).
- [20] L. Li, L. Zhang, Z. Xu, S. Morita, Y. Cheng, F. Zhang, W. Zhang, Y. Duan, Q. Zang, S. Wang, S. Dai, G. Zuo, Z. Sun, L. Wang, X. Ding, J. Qian, H. Liu, and L. Hu, “Line identification of extreme ultraviolet (EUV) spectra from low-Z impurity ions in EAST Tokamak plasmas”, Plasma Sc. and Tech. **23**, 075102 (2021).
- [21] C. Dong, S. Morita, M. Goto, and E. Wang, “Absolute intensity calibration of flat-field space-resolved extreme ultraviolet spectrometer using radial profiles of visible and extreme ultraviolet bremsstrahlung continuum emitted from high-density plasmas in large helical device”, Rev. of Sci. Instr. **82**, 11 (2011).
- [22] C. Dong, S. Morita, M. Goto, and E. Wang, “Extension of wavelength range in absolute intensity calibration of space-resolved EUV spectrometer for LHD diagnostics”, Plasma and Fusion Research **7**, 2402139 (2012).
- [23] X. Han, A. Hu, D. Li, S. Xiao, B. Tian, Q. Zang, J. Zhao, C. Hsieh, X. Gong, L. Hu and G. Xu, “Multiple Laser System for High-Resolution Thomson Scattering Diagnostics on the EAST Tokamak”, IEEE Transactions on Plasma Science **46**, 406 (2018).
- [24] Q. Xu, X. Gao, Y. Jie, H. Liu, N. Shi, Y. Cheng and X. Tong, “HCN Laser Interferometer on the EAST Superconducting Tokamak”, Plasma Science and Technology, **10**, 519 (2008).
- [25] Y. Duan, L. Hu, S. Mao, P. Xu, K. Chen, S. Lin, G. Zhong, J. Zhang, L. Zhang and L. Wang, “Measurement of Radiated Power Loss on EAST”, Plasma Science and Technology, **13**, 546 (2011).
- [26] F. Ling Zhang, W. Zhang, Y. Cheng, A. Hu, X. Ding, S. Morita, Z. Li, Z. Zhou, Y. Cao, J. Ma, Z. Xu, L. Xu, C. Zhou, Y. Jie and D. Mitnik, “Line identification of extreme ultraviolet spectra from aluminum ions in EAST Tokamak plasmas”, Phys. Scr. **99**, 025615 (2024).
- [27] W. Zhang, L. Zhang, Y. Cheng, S. Morita, H. Sheng, D. Mitnik, Y. Sun, Z. Wang, Y. Chu, A. Hu, Y. Jie and H. Liu, “First observation of edge impurity behavior with $n = 1$ RMP application in EAST L-mode plasma”, Nucl. Fusion **64**, 086004 (2024).
- [28] M. Klapisch, J. L. Schwob, B. S. Fraenkel, and J. Oreg, “The $1s - 3p$ $K\beta$ -like X-ray spectrum of highly ionized iron”, J. Opt. Soc. Am. **67**, 148 (1977).
- [29] A. Bar-Shalom, M. Klapisch, and J. Oreg, “HULLAC, an integrated computer package for atomic processes in plasmas”, Jour. of Quant. Spectr. and Rad. Transf. **71**, 169 (2001).
- [30] N. R. Badnell, “Dielectronic recombination of Fe^{22+} and Fe^{21+} ”, J. Phys. B **19**, 3827 (1986).
- [31] N. R. Badnell, “On the effects of the two-body non-fine-structure operators of the Breit-Pauli Hamiltonian”, J. Phys. B **30**, 1 (1997).
- [32] N. R. Badnell, “A Breit-Pauli distorted wave implementation for AUTOSTRUCTURE”, Comp. Phys. Comm. **180**, 1528 (2011).
- [33] M. F. Gu, “The Flexible Atomic Code”, Can. J. Phys. **86**, 675 (2008).
- [34] M. Stenke, K. Aichele, D. Harthiramani, G. Hofmann, M. Steidl, R. Volpel, and E. Salzborn, “Electron-impact single-ionization of singly and multiply charged tungsten ions”, J. Phys. B **328**, 2711 (1995).
- [35] K. Spruck, A. Becker, A. Borovik, M.F. Gharaibeh, J. Rausch, S. Schippers, and A. Müller, “Electron-impact ionization of multiply charged tungsten ions”, Jour. of Phys. Conf. Ser. **488**, 062026 (2014).
- [36] D. Mitnik, P. Mandelbaum, J.L. Schwob, A. Bar-Shalom, and J. Oreg, “Excitation-autoionization cross sections and rate coefficients of Zn-like ions”, Phys. Rev A **55**, 307 (1997).
- [37] V. Jonauskas, A. Kynienė, S. Kučas, S. Pakalka, Š. Masys, A. Pranciukevičius, A. Borovik, M. F. Gharaibeh, S. Schippers, and A. Müller, “Electron-impact ionization of W^{5+} ”, Phys. Rev. A **100**, 062701 (2019).

- [38] D. Zhang, L. Xie, J. Jiang, Z. Wu, C. Dong, Y. Shi, and Y. Qu, “Electron-impact single ionization for W^{4+} and W^{5+} ”, *Chin. Phys. B* **27**, 053402 (2018).

