

Entangling power of spin- j systems: a geometrical approach

Eduardo Serrano-Ensástiga,^{1,*} Diego Morachis Galindo,^{1,2} Jesús A. Maytorena,² and Chryssomalis Chryssomalakos³

¹*Institut de Physique Nucléaire, Atomique et de Spectroscopie,
CESAM, University of Liège, B-4000 Liège, Belgium*

²*Departamento de Física, Centro de Nanociencias y Nanotecnología, Universidad Nacional Autónoma de México
Apartado Postal 14, 22800, Ensenada, Baja California, México*

³*Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México PO Box 70-543, 04510, CDMX, México*

(Dated: October 7, 2024)

Unitary gates with high entangling power are relevant for several quantum-enhanced technologies due to their entangling capabilities. For symmetric multiparticle systems, such as spin states or bosonic systems, the particle exchange symmetry restricts these gates and also the set of not-entangled states. In this work, we analyze the entangling power of spin systems by reformulating it as an inner product between vectors with components given by $SU(2)$ invariants. This approach allows us to study this quantity for small-spin systems including the detection of the unitary gate that maximizes it for small spins. We observe that extremal unitary gates exhibit entanglement distributions with high rotational symmetry, same that are linked to a convex combination of Husimi functions of certain states. Furthermore, we explore the connection between entangling power and the Schmidt numbers admissible in some spin state subspaces. Thus, the geometrical approach presented here suggests new paths for studying entangling power linked to other concepts in quantum information theory.

I. INTRODUCTION

Entanglement is a foundational concept in quantum theory and a vital resource for quantum technologies such as quantum computing, cryptography, metrology and simulation [1–3]. In multipartite quantum systems, it is generated through nonlocal unitary transformations, as local operations cannot alter the entanglement of a state [1]. It is therefore natural to investigate the entangling capacity of nonlocal unitary gates, and how to create highly entangled states via the evolution of non-local Hamiltonians or pulse sequences in physical systems [4–11]. On the theoretical side, several concepts have been proposed to assess the capability of unitary gates in generating quantum resources such as *gate typicality*, *disentangling power*, or *perfect entanglers* [12–15]. One of the most intuitively appealing quantities is the *entangling power* of a unitary gate [12] which is defined as the average entanglement generated by a unitary gate over the set of separable (non-entangled) states. Extensive studies of the entangling power have been carried out in bipartite systems [16–23] with generalizations to two-qubit global noise channels [24] and multipartite systems [25, 26]. The entangling power of a unitary operator is also connected to its entanglement in its own Schmidt decomposition [16, 27] or to invariant quantities under local transformations [8, 28, 29]. Furthermore, connections have been identified between highly entangling unitary gates and Quantum Error-Correcting Codes [25], absolutely maximally entangled (AME) states [26], unitary operators invariant under local actions of diagonal unitary and orthogonal groups [30], and quantum versions of chaotic systems [31–33].

When a multiqubit system is restricted to its symmetric subspace, the set of product states is significantly reduced to the set of Spin-Coherent (SC) states, which form a 2-sphere within the Hilbert space of quantum states [3, 34]. Additionally, the local unitary transformations are limited to the global rotations, specifically global $SU(2)$ transformations generated by the angular momentum operators. Symmetric multiqubit states arise in bosonic systems, such as two-mode multiphotonic systems or spin- j states. Moreover, the symmetric subspace of N qubit states is equivalent to the Hilbert space of spin- j ($j = N/2$) states, which makes the study of the entangling power of single spin states relevant in distinct physical platforms.

For the symmetric two-qubit case, which can be thought of as a spin-1 system, the entangling power of 3×3 unitary matrices has been studied in Ref. [35]. One key mathematical feature that arises in the study of those unitaries is the Cartan decomposition, which factorizes any quantum gate as a non-local operation multiplied on each side by local $SU(2)$ transformations [36]. The entangling power of the unitary is encoded into the non-local factor, which is obtained by exponentiating the maximally commuting (Cartan) subalgebra of $\mathfrak{su}(3)$. This allows to represent the unitaries with equivalent entangling properties in an euclidean two-dimensional space [35]. However, for higher spin systems the Cartan decomposition would not be very advantaging, since the number of relevant parameters grows quadratically with the dimension of the system.

In this work, we study the entangling power for spin- j states. We reformulate its original expression in terms of $SU(2)$ invariant quantities of the unitary operators. This geometrical approach connects the entangling power to other well-known quantities in quantum information theory. The structure of the paper is as follows: Sec. II

* ed.ensastiga@uliege.be

reviews the formal definitions of the entropy of entanglement and the entangling power for spin- j states. We derived its above mentioned geometric expression in Sec. III. We then calculate the entangling power and the entanglement distribution on the sphere for small spins in Sec. IV and V, respectively, including the search of unitary gates with the best entangling capabilities. Sec. VI contains the calculation of the average entangling power over all the unitary gates. The connection of the entangling power and the possible Schmidt numbers in states subspaces is explored in VII. Finally, we conclude with some last remarks in Sec. VIII.

II. BASIC CONCEPTS

We consider a single spin- j system with Hilbert space $\mathcal{H}^{(j)}$ and the Hilbert-Schmidt (HS) space of bounded operators $\mathcal{HS}(\mathcal{H}^{(j)})$ acting on $\mathcal{H}^{(j)}$. Any spin- j state is equivalent to a symmetric state of $2j$ spin-1/2 (qubits) constituents (see Ref. [37] for a detailed discussion of this equivalence). In this framework, we consider a bipartition of these $2j$ qubits $\mathcal{H}^{(j)} \subset \mathcal{H}_A \otimes \mathcal{H}_B$ and assume, without loss of generality, that $\dim(\mathcal{H}_A) \leq \dim(\mathcal{H}_B)$. Moreover, we can take as subsystems $\mathcal{H}_A \cong \mathcal{H}^{(q/2)}$ and $\mathcal{H}_B \cong \mathcal{H}^{(j-q/2)}$, *i.e.*, the bipartition involves states of spins $q/2$ and $j - q/2$.

The entanglement of a bipartite state $|\Psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$ can be quantified by the (normalized) linear entanglement entropy [12]

$$E(|\Psi\rangle) \equiv \frac{d}{d-1} [1 - \text{Tr}(\rho_A^2)], \quad (1)$$

where $E(|\Psi\rangle) \in [0, 1]$, $\rho_A = \text{Tr}_B(|\Psi\rangle\langle\Psi|)$ is the reduced mixed state after tracing out the subsystem B , and $d = \dim(\mathcal{H}_A)$. For a general bipartite pure state with Schmidt decomposition

$$|\Psi\rangle = \sum_{k=1}^d \sqrt{\Gamma_k} |\psi_{A,k}\rangle \otimes |\psi_{B,k}\rangle, \quad (2)$$

we have

$$E(|\Psi\rangle) = \frac{d}{d-1} \left[1 - \sum_{k=1}^d \Gamma_k^2 \right]. \quad (3)$$

In particular, $E(|\Psi\rangle) = 0$ for bipartite product states $|\Psi\rangle = |\psi_A\rangle \otimes |\psi_B\rangle$. On the other hand, the maximally entangled states, for which $E = 1$, have Schmidt decomposition

$$|\Psi\rangle = \frac{1}{\sqrt{d}} \sum_{k=1}^d |\psi_{A,k}\rangle \otimes |\psi_{B,k}\rangle, \quad (4)$$

where $\{|\psi_{A,k}\rangle\}_k \subset \mathcal{H}_A$ and $\{|\psi_{B,k}\rangle\}_k \subset \mathcal{H}_B$ are orthogonal sets of states. In our case, each value of q defines a different measure of entanglement E_q . Given that

$\text{Tr}(\rho_A^2) = \text{Tr}(\rho_B^2)$, we only need to consider $q = 1, \dots, \lfloor j \rfloor$, as the entanglements E_q for values of q higher than $\lfloor j \rfloor$ are linear combinations of those of lower q values.

The product (separable) states in $\mathcal{H}^{(j)}$ are the spin-coherent (SC) states $|j, \mathbf{n}\rangle \equiv D^{(j)}(\mathbf{n}) |j, j\rangle$ which constitute a 2-sphere in $\mathcal{H}^{(j)}$ [34]. A possible parametrization is given by the rotations $D^{(j)}(\mathbf{n}) = D^{(j)}(0, \theta, \phi)$ which align the \mathbf{z} axis to the direction \mathbf{n} with spherical angles (θ, ϕ) . $D^{(j)}(\alpha, \beta, \gamma)$ denotes the irreducible representation j (j -irrep) of a rotation matrix in the Euler angle parametrization [38]. The *entangling power* of a quantum unitary gate $U \in \mathcal{HS}(\mathcal{H}^{(j)})$, with respect to E_q , is defined as the average entanglement produced by U acting on SC states,

$$e_p(E_q, U) \equiv \overline{E_q(U|j, \mathbf{n}))} = \frac{1}{4\pi} \int_{S^2} E_q(U|j, \mathbf{n}) \, d\mathbf{n}. \quad (5)$$

It is easily seen that

$$e_p(E_q, R_1 U R_2) = e_p(E_q, U), \quad (6)$$

for all q , where $R_{1,2}$ are matrices representing arbitrary SU(2) elements. If U itself represents an SU(2) element, then $e_p(E_q, U) = 0$ for all q .

In what follows, we denote the coupled basis of two spins by

$$|j_1, j_2, L, M\rangle \equiv \sum_{m_2=-j_2}^{j_2} \sum_{m_1=-j_1}^{j_1} C_{j_1 m_1 j_2 m_2}^{LM} |j_1, m_1\rangle |j_2, m_2\rangle, \quad (7)$$

with $|j_1, m_1\rangle |j_2, m_2\rangle = |j_1, m_1\rangle \otimes |j_2, m_2\rangle$, and $C_{j_1 m_1 j_2 m_2}^{LM}$ denoting the Clebsch-Gordan coefficients.

A. Case $j = 1$

The entangling power $e_p(E_q, U)$ for $j = 1$ is only defined for $q = 1$, and it can be written as [35]

$$e_p(E_1, U) = \frac{3}{5} \left(1 - \frac{1}{9} |\text{Tr}(m)|^2 \right), \quad (8)$$

where

$$\text{Tr}(m) = \text{Tr}(U_B^T U_B), \quad (9)$$

and U_B is the unitary matrix U , transformed in the symmetric Bell states basis, $U_B = Q^\dagger U Q$ [39]

$$Q = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 0 & i \\ 0 & i\sqrt{2} & 0 \\ 1 & 0 & -i \end{pmatrix}. \quad (10)$$

After some algebra, we can also write Eq. (9) as

$$\text{Tr}(m) = \text{Tr}(\Phi U^T \Phi U), \quad \text{with } \Phi = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix}. \quad (11)$$




FIG. 1. Plot of the function $e_p(E_1, U)$, given in (14). One can clearly access a maximum on the $c_2 = 0$ curve (blue in the figure). Two particular maxima are shown (green points); the one on the right, corresponding to $c_1 = 2\pi/3$, gives the U_0 in (15).

The unitary transformations of spin-1 states can be parametrized using the Cartan decomposition [35, 36], with the $SU(2)$ -subgroup generated by the $j = 1$ angular momentum operators. A possible parametrization is $U = R_1 A R_2$ with $R_1, R_2 \in SU(2)$ and

$$A = \begin{pmatrix} \frac{\lambda_1 + \lambda_3}{2} & 0 & \frac{\lambda_1 - \lambda_3}{2} \\ 0 & \lambda_2 & 0 \\ \frac{\lambda_1 - \lambda_3}{2} & 0 & \frac{\lambda_1 + \lambda_3}{2} \end{pmatrix}, \quad (12)$$

with

$$\begin{aligned} \lambda_1 &= e^{\frac{i}{2}(c_1 - c_2 + c_3)}, \\ \lambda_2 &= e^{\frac{i}{2}(c_1 + c_2 - c_3)}, \\ \lambda_3 &= e^{\frac{i}{2}(-c_1 + c_2 + c_3)}, \end{aligned} \quad (13)$$

and where the real parameters c_k fulfill $c_1 + c_2 + c_3 = 0$ [35]. Given that the entangling power is invariant under left and right rotations (see Eq. (6)), $e_p(E_1, U)$ only depends on the c_i 's,

$$e_p(E_1, U) = \frac{4}{15} (\sin^2 c_{12} + \sin^2 c_{13} + \sin^2 c_{23}), \quad (14)$$

with $c_{ij} = c_i - c_j$. We plot $e_p(E_1, U)$ as a function of c_1 and c_2 in Fig. 1. For $c_2 = 0$ (blue curve), we observe two unitary matrices attaining the maximum at $c_1 = \pi/3$ and $2\pi/3$, respectively. The latter solution has corresponding unitary matrix (with $R_{1,2} = I$)

$$U_0 = \begin{pmatrix} \frac{1}{2}(1 + \omega^2) & 0 & \frac{1}{2}(1 - \omega^2) \\ 0 & -\omega & 0 \\ \frac{1}{2}(1 - \omega^2) & 0 & \frac{1}{2}(1 + \omega^2) \end{pmatrix}, \quad (15)$$

where $\omega = e^{i\pi/3}$ is the primitive cubic root of -1 and $e_p(E_1, U_0) = 3/5$.

III. ENTANGLING POWER FOR A GENERAL SPIN- j STATE

Our first result is the reformulation of $E_q(U |j, \mathbf{n})$ in the form

$$E_q(U |j, \mathbf{n}) = 1 - \langle 2j, \mathbf{n} | \mathcal{U}^\dagger \mathcal{M}_q \mathcal{U} | 2j, \mathbf{n} \rangle, \quad (16)$$

where $\mathcal{U} \equiv U \otimes U$ and

$$\mathcal{M}_q \equiv \frac{q+1}{q} \sum_{L=0}^{2j} (-1)^{2j+L} \chi(q, j, L) \mathcal{P}_L, \quad (17)$$

with

$$\chi(q, j, L) \equiv \sum_{\sigma=1}^q (2\sigma + 1) \frac{\left\{ \begin{matrix} q/2 & q/2 & q \\ q/2 & q/2 & \sigma \end{matrix} \right\} \left\{ \begin{matrix} j & j & L \\ j & j & \sigma \end{matrix} \right\}}{\left\{ \begin{matrix} j & j & 2j \\ j & j & \sigma \end{matrix} \right\}} \quad (18)$$

and

$$\mathcal{P}_L \equiv \sum_{M=-L}^L |j, j, L, M\rangle \langle j, j, L, M|. \quad (19)$$

Here, the curly bracket represents the Wigner 6- j symbol [38]. \mathcal{P}_L is the projector operator in the subspace of $\mathcal{H}^{(j) \otimes 2}$ defined with the states of coupled basis (7) with total angular momentum L . The derivation of Eq. (16) is given in Appendix A. We now use the resolution of unity of the SC states [40]

$$\frac{1}{4\pi} \int |2j, \mathbf{n}\rangle \langle 2j, \mathbf{n}| d\mathbf{n} = \frac{\mathcal{P}_{2j}}{4j+1} \equiv \mathcal{N}, \quad (20)$$

to calculate e_p , yielding

$$e_p(E_q, U) = 1 - \text{Tr}(\mathcal{U} \mathcal{N} \mathcal{U}^\dagger \mathcal{M}_q). \quad (21)$$

The fact that both operators \mathcal{N} and \mathcal{M}_q are linear combinations of \mathcal{P}_L 's, which are orthogonal among themselves, $\text{Tr}(\mathcal{P}_L \mathcal{P}_K) = (2L+1)\delta_{LK}$, suggests a further reformulation of the e_p . Indeed, to an operator $V \in \mathcal{HS}(\mathcal{H}^{(j) \otimes 2})$ we associate the $(2j+1)$ -dimensional vector \vec{V} , with $SU(2)$ -invariant components,

$$\vec{V} = \left(\text{Tr}(V \tilde{\mathcal{P}}_0), \dots, \text{Tr}(V \tilde{\mathcal{P}}_L), \dots, \text{Tr}(V \tilde{\mathcal{P}}_{2j}) \right), \quad (22)$$

where $\tilde{\mathcal{P}}_L = \mathcal{P}_L / \sqrt{2L+1}$. By direct calculation, we find

$$\langle \hat{\mathcal{U}} \vec{N}, \vec{\mathcal{M}}_q \rangle = \text{Tr}(\mathcal{U} \mathcal{N} \mathcal{U}^\dagger \mathcal{M}_q), \quad (23)$$

where $\langle \cdot, \cdot \rangle$ is the euclidean inner product,

$$\langle \vec{V}, \vec{W} \rangle \equiv \sum_{k=0}^{2j} V_k W_k, \quad (24)$$

and \widehat{U} is a $(2j+1) \times (2j+1)$ real matrix with entries

$$\left(\widehat{U}\right)_{mn} = \text{Tr}\left(\mathcal{U}\tilde{\mathcal{P}}_m\mathcal{U}^\dagger\tilde{\mathcal{P}}_n\right). \quad (25)$$

Therefore, the entangling power reads

$$e_p(E_q, U) = 1 - \left\langle \widehat{U}\vec{N}, \vec{\mathcal{M}}_q \right\rangle. \quad (26)$$

From Eqs. (24) and (25), we deduce that $\langle \widehat{U}\vec{N}, \vec{\mathcal{M}}_q \rangle = \langle \vec{N}, \widehat{U}^T \vec{\mathcal{M}}_q \rangle$. To summarize, we have expressed $e_p(E_q, U)$ in terms of the euclidean inner product of two vectors, with components given by SU(2)-invariant quantities of the operators \mathcal{N} and \mathcal{M}_q , one of them rotated by the unitary transformation \mathcal{U} . A similar formula for e_p , in terms of operators associated to the multipole operators [38], is given in Appendix B.

The projectors \mathcal{P}_L change by a $(-1)^{2j+L}$ sign under particle exchange in $\mathcal{H}^{(j)\otimes 2}$. Since the unitary operators $\mathcal{U} = U \otimes U$ preserve this symmetry, $\text{Tr}(\mathcal{U}\mathcal{P}_{2j}\mathcal{U}^\dagger\mathcal{P}_L)$ vanishes unless $L \equiv 2j \pmod{2}$. Thus, the vector $\vec{N} = (0, 0, \dots, 1/\sqrt{4j+1})$ after a \mathcal{U} transformation has components

$$\left(0, \text{Tr}(\mathcal{N}\tilde{\mathcal{P}}_1), 0, \text{Tr}(\mathcal{N}\tilde{\mathcal{P}}_3), 0, \dots, \text{Tr}(\mathcal{N}\tilde{\mathcal{P}}_{2j}) \right), \\ \left(\text{Tr}(\mathcal{N}\tilde{\mathcal{P}}_0), 0, \text{Tr}(\mathcal{N}\tilde{\mathcal{P}}_2), 0, \dots, \text{Tr}(\mathcal{N}\tilde{\mathcal{P}}_{2j}) \right), \quad (27)$$

for $2j$ odd or even, respectively. On the other hand, $\vec{\mathcal{M}}_q$ has components for L both odd and even. As an example, the components of $\vec{\mathcal{M}}_1$ are

$$\left(\vec{\mathcal{M}}_1\right)_L = \frac{L(L+1) - 2j(j+1)}{2j^2}. \quad (28)$$

Nevertheless, the relevant components of \mathcal{M}_q for e_p are the ones in common with $\widehat{U}\vec{N}$ (27), *i.e.*, with $2j \equiv L \pmod{2}$. These components of \vec{N} and $\vec{\mathcal{M}}_q$ lie in a hyperplane after any unitary transformation $\mathcal{U} = U \otimes U$ (see **Identity 3** of Appendix C)

$$\sum_{\substack{L=0 \\ 2j \equiv L \pmod{2}}}^{2j} \sqrt{2L+1} \left(\widehat{U}\vec{N}\right)_L = \sum_{\substack{L=0 \\ 2j \equiv L \pmod{2}}}^{2j} \text{Tr}(\mathcal{U}\mathcal{P}_L\mathcal{U}^\dagger\mathcal{N}) = 1, \\ \sum_{\substack{L=0 \\ 2j \equiv L \pmod{2}}}^{2j} \sqrt{2L+1} \left(\widehat{U}\vec{\mathcal{M}}_q\right)_L = \frac{(j+1)(2j+1)}{2j+1-q}. \quad (29)$$

The components of $\vec{\mathcal{M}}_q$ for several spin values are shown in Table I — they satisfy the inequalities

$$0 \leq \text{Tr}(\mathcal{U}\mathcal{P}_L\mathcal{U}^\dagger\mathcal{P}_K) \leq \min(2L+1, 2K+1), \quad (30)$$

for $L \equiv K \pmod{2}$, which become evident once the fact that $\mathcal{P}_L, \mathcal{P}_K$ project onto subspaces of dimensions $2L+1$

j	q	$\vec{\mathcal{M}}_q$
1	1	$(-2, -\sqrt{3}, \sqrt{5})$
3/2	1	$\left(-\frac{5}{3}, -\frac{11}{3\sqrt{3}}, -\frac{\sqrt{5}}{3}, \sqrt{7}\right)$
	2	$\left(-\frac{3}{2}, -\frac{5\sqrt{3}}{4}, -\frac{3\sqrt{5}}{4}, 0, 3\right)$
	2	$\left(-\frac{1}{4}, -\frac{\sqrt{3}}{2}, -\frac{3\sqrt{5}}{4}, -\frac{\sqrt{7}}{2}, 3\right)$
5/2	1	$\left(-\frac{7}{5}, -\frac{31\sqrt{3}}{25}, -\frac{23}{5\sqrt{5}}, -\frac{11\sqrt{7}}{25}, \frac{3}{5}, \sqrt{11}\right)$
	2	$\left(-\frac{7}{20}, -\frac{47\sqrt{3}}{100}, -\frac{31}{10\sqrt{5}}, -\frac{31\sqrt{7}}{50}, -\frac{3}{5}, \sqrt{11}\right)$

TABLE I. Vectors of SU(2) invariants of \mathcal{M}_q (17) for the spin values $j = 1, 3/2, 2, 5/2$ and $q = 1, \dots, [j]$.

and $2K+1$, respectively, is taken into account. The unitary operator rotates this subspace while preserving its dimension. The trace in (30) then is just the projection of one of the subspaces, rotated by \mathcal{U} , onto the other. These inequalities can be used to find bounds for the components of the rotated vector

$$0 \leq p_L \leq \frac{\sqrt{2L+1}}{4j+1}, \quad \text{with } p_L \equiv \left(\widehat{U}\vec{N}\right)_L. \quad (31)$$

A unitary operator \mathcal{U} that achieves a critical value of $e_p(E_q, U)$ must fulfill

$$\text{Tr}(\mathcal{U}\mathcal{N}\mathcal{U}^\dagger[\mathcal{M}_q, \mathcal{G}_a]) = 0, \quad \mathcal{G}_a = \mathbf{1} \otimes G_a + G_a \otimes \mathbf{1}, \quad (32)$$

for any generator of the Lie algebra $G_a \in \mathfrak{su}(2j+1)$. If it satisfies additionally that the Hessian of e_p evaluated at \mathcal{U} ,

$$H_{ab} \equiv -\text{Tr}(\mathcal{U}\mathcal{N}\mathcal{U}^\dagger[[\mathcal{M}_q, \mathcal{G}_a], \mathcal{G}_b]), \quad (33)$$

has only negative eigenvalues, then \mathcal{U} is a local maximum. Due to the invariance of $e_p(E_q, U)$ under left and right SU(2) operations (6), at least 6 eigenvalues of H are equal to zero.

IV. e_p FOR SMALL SPINS

A. Spin 1

We calculate again $e_p(E_1, U)$ for $j = 1$ using the formulation given in the previous section. In this case, $\widehat{U}\vec{N} = (p_0, 0, p_2)$. The restriction to the hyperplane (29), $p_0 + \sqrt{5}p_2 = 1$, lets us write e_p in terms of only p_0 ,

$$e_p(E_1, U) = 1 - \left\langle \widehat{U}\vec{N}, \vec{\mathcal{M}}_1 \right\rangle = 3p_0 \quad (34)$$

We recover (8) by calculating p_0 ,

$$5p_0 = \text{Tr}(\mathcal{U}\mathcal{P}_2\mathcal{U}^\dagger\mathcal{P}_0) \\ = \text{Tr}(\mathcal{U}(\mathbf{1} - \mathcal{P}_0 - \mathcal{P}_1)\mathcal{U}^\dagger\mathcal{P}_0) \\ = 1 - \text{Tr}(\mathcal{U}\mathcal{P}_0\mathcal{U}^\dagger\mathcal{P}_0). \quad (35)$$

Since \mathcal{P}_0 is a rank one operator, the last expression can be rewritten as

$$5p_0 = 1 - |\text{Tr}(\mathcal{P}_0\mathcal{U})|^2 = 1 - \frac{1}{9} |\text{Tr}(\Phi U^T \Phi U)|^2, \quad (36)$$

where Φ is defined in Eq. (11) and the last equality is derived as follows

$$\begin{aligned} \text{Tr}(\mathcal{P}_0\mathcal{U}) &= \sum_{\substack{m_1 m_2 \\ n_1 n_2}} C_{1m_1 1m_2}^{00} C_{1n_1 1n_2}^{00} U_{m_1 n_1} U_{m_2 n_2} \\ &= \frac{1}{3} \sum_{m_1 n_1} (-1)^{m_1+n_1} U_{m_1 n_1} U_{-m_1 -n_1} \\ &= \frac{1}{3} \text{Tr}(\Phi U^T \Phi U). \end{aligned} \quad (37)$$

We can obtain the upper bound $e_p(E_1, U) \leq 3/5$ from Eq. (31) — the inequality is in fact saturated by U_0 (15). In Fig. 2 (upper frame) we plot the vectors $\widehat{U}\vec{N}$ for random unitary operators, produced with the Haar measure, as well as the vector $\widehat{U}_0\vec{N}$ with U_0 given by (15). We also plot the orthogonal complement \vec{M}_q^\perp of \vec{M}_q with respect to the inner product (24). As expected, U_0 meets the criteria for a local maximum (32)-(33), with the Hessian there having two eigenvalues equal to -4 and six equal to 0.

B. Spin 3/2

The vector $\widehat{U}\vec{N} = (0, p_1, 0, p_3)$ is restricted in the hyperplane (29) $\sqrt{3}p_1 + \sqrt{7}p_3 = 1$. Hence, $e_p(E_1, U)$ is also a function of one SU(2)-invariant for $j = 3/2$,

$$e_p(E_1, U) = \frac{20\sqrt{3}}{9} p_1 \leq \frac{20}{21}. \quad (38)$$

The bound is saturated by the unitary operator

$$U_0 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & i \\ i & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad (39)$$

which we identified as outlined in Section VII. Note that in order to simplify the notation, we denote the optimal entangles by U_0 for all values of spin — which unitary operator is involved should be clear from the context. We plot the vectors $\widehat{U}\vec{N}$ for random unitary operators, as well as $\widehat{U}_0\vec{N}$ and \vec{M}_q^\perp in Fig. 2 (lower frame). Again, it is verified that U_0 is a critical value of e_p with Hessian having eigenvalues equal to -8 , $-8/5$ and 0, with degeneracies 4, 5 and 6, respectively.

C. Spin-2

Here, we have three different non-zero components in $\widehat{U}\vec{N} = (p_0, 0, p_2, 0, p_4)$ restricted to the plane $p_0 + \sqrt{5}p_2 +$




FIG. 2. Vectors of SU(2) invariants in the plane where $\widehat{U}\vec{N}$ has non-zero components (27) for $j = 1$ (upper frame) and $j = 3/2$ (lower frame). The black dots are the vectors corresponding to random, according to the Haar measure, unitary operators. The solid red segment is the part of the hyperplane (29) that also satisfies (31). The contour lines of $e_p(E_1, U)$ are parallel to the dashed blue line representing \vec{M}_q^\perp . The entangling power of U increases as the euclidean distance from $\widehat{U}\vec{N}$ to \vec{M}_q^\perp decreases. The upper bounds are saturated for the unitary transformations (15) (also (68)) for $j = 1$ and (39), for $j = 3/2$. To simplify the axes labels, we use the scaled variables $p'_\sigma = \sqrt{2\sigma + 1}p_\sigma$.

$3p_4 = 1$, and two different non-equivalent bipartite entanglements

$$\begin{aligned} e_p(E_1, U) &= \frac{10p_0 + 7\sqrt{5}p_2}{4}, \\ e_p(E_2, U) &= \frac{5p_0 + 7\sqrt{5}p_2}{4}. \end{aligned} \quad (40)$$

Unlike the previous spin values, the inequalities (31) provide a trivial bound for $e_p(E_q, U)$. Through numerical search, we identified unitary matrices $U_0^{(q)}$ that we conjecture are optimal entanglers for E_q , $q = 1, 2$. They read

$$U_0^{(1)} = \begin{pmatrix} \beta \cos \alpha & 0 & 0 & 0 & i\beta \sin \alpha \\ 0 & e^{i\frac{\pi}{4}} & 0 & 0 & 0 \\ 0 & 0 & \beta^2 & 0 & 0 \\ 0 & 0 & 0 & e^{-i\frac{\pi}{4}} & 0 \\ -i\beta \sin \alpha & 0 & 0 & 0 & -\beta \cos \alpha \end{pmatrix}, \quad (41)$$

with $\alpha = \arctan\left(\sqrt{83/53}\right)$ and $\beta = e^{-i \arctan\left(\sqrt{53/83}\right)}$, and

$$U_0^{(2)} = \begin{pmatrix} 0 & 0 & 0 & i & 0 \\ i & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & i \\ 0 & i & 0 & 0 & 0 \end{pmatrix}. \quad (42)$$

Both unitary matrices fulfill the criteria for local maximum (32) and (33) for their respective E_q , with entangling power

$$\begin{aligned} e_p(E_1, U_0^{(1)}) &= \frac{6889}{7140} \approx 0.9648, \\ e_p(E_2, U_0^{(2)}) &= \frac{25}{28} \approx 0.8929. \end{aligned} \quad (43)$$

As for the previous cases, we plot the vectors $\widehat{U}\vec{\mathcal{N}}$, $\widehat{U}_0^{(q)}\vec{\mathcal{N}}$ and $\vec{\mathcal{M}}_q^\perp$ in Fig. 3.

D. e_p of a unitary operator and its inverse

In this short section we comment on the relation between the entangling power of a unitary matrix U and its inverse. We find that, for $j = 1$, $e_p(E_1, U) = e_p(E_1, U^\dagger)$ for all $U \in \text{SU}(3)$. This result is derived from the expression for $e_p = 3[1 - \text{Tr}(U\mathcal{P}_0U^\dagger\mathcal{P}_0)]/5$ obtained from Eqs. (34)-(35). The same result holds for $j = 3/2$, with a similar proof starting from Eq. (38). On the other hand, numerical calculations show that, in general, $e_p(E_q, U) \neq e_p(E_q, U^\dagger)$ for $j \geq 2$.

V. ENTANGLEMENT DISTRIBUTION AND HUSIMI FUNCTIONS

Now, we would like to study the entanglement $E_q(U|j, \mathbf{n})$ as a function on the sphere where \mathbf{n} lives. We start with the expression

$$E_q(U|j, \mathbf{n}) = 1 - \langle 2j, \mathbf{n} | \mathcal{D}_q(U) | 2j, \mathbf{n} \rangle, \quad (44)$$

with

$$\mathcal{D}_q(U) \equiv \mathcal{P}_{2j} \mathcal{U}^\dagger \mathcal{M}_q \mathcal{U} \mathcal{P}_{2j}, \quad (45)$$

which follows from (16) since the state $|2j, \mathbf{n}\rangle$ lies in the subspace associated to \mathcal{P}_{2j} . The matrix $\mathcal{D}_q = \mathcal{D}_q(U)$ is Hermitian for any U and can be thought of as a $(4j+1) \times (4j+1)$ matrix when restricted to the image of \mathcal{P}_{2j} . Additionally, it rotates under $\text{SU}(2)$ transformations as $D^{(2j)}(R)\mathcal{D}_q D^{(2j)\dagger}(R)$ with $D^{(2j)}(R)$ the $(2j)$ -irrep of the rotation R . By Eq. (44), $E_q(U|j, \mathbf{n})$ inherits the rotational symmetries of \mathcal{D}_q . These symmetries can be scrutinized by the Majorana representation for Hermitian




FIG. 3. Vectors of $\text{SU}(2)$ invariants in the plane where $\widehat{U}\vec{\mathcal{N}}$ has non-zero components (27) for $j = 2$. The black dots are the vectors given by random unitary operators. The red plane is the restriction (29). The dashed blue and green lines denote the intersection between the red plane with $\vec{\mathcal{M}}_1^\perp$ and $\vec{\mathcal{M}}_2^\perp$, respectively. The yellow cubes represent the position of the vector $\vec{\mathcal{N}}$ and its corresponding rotated vector by the unitary operators $U_0^{(q)}$ given by Eqs. (41) and (42), respectively.

operators [41]. In particular, we use it to verify the rotational symmetries of $E_q(U|j, \mathbf{n})$ for the unitary gates mentioned below.

The eigendecomposition $\mathcal{D}_q = \sum_{k=1}^{4j+1} \sigma_k |\psi_k\rangle \langle \psi_k|$ also helps to recast the entanglement distribution on the sphere as

$$E_q(U|j, \mathbf{n}) = 1 - \sum_{k=1}^{4j+1} \sigma_k H_{|\psi_k\rangle}(\mathbf{n}), \quad (46)$$

with $H_{|\psi_k\rangle}(\mathbf{n}) = |\langle 2j, \mathbf{n} | \psi_k \rangle|^2$ being the Husimi function of $|\psi_k\rangle$ [40]. By averaging over the sphere, we find

$$\begin{aligned} e_p(E_q, U) &= 1 - \sum_{k=1}^{4j+1} \sigma_k \overline{H_{|\psi_k\rangle}(\mathbf{n})} \\ &= 1 - \frac{1}{4j+1} \sum_{k=1}^{4j+1} \sigma_k \\ &= 1 - \frac{\text{Tr}[\mathcal{D}_q(U)]}{4j+1}, \end{aligned} \quad (47)$$

which reduces to Eq. (26). Let us now study the entanglement distribution of the examples in Sec. IV.

A. $j=1$

We plot in Fig. 4a the entanglement distribution

$$\begin{aligned} E_1(U_0'|\mathbf{n}) &= 4\sqrt{2} \sin^5\left(\frac{\theta}{2}\right) \cos^3\left(\frac{\theta}{2}\right) \cos(3\phi) \\ &+ \frac{\sin^2\left(\frac{\theta}{2}\right)}{32} [90 + 105 \cos(\theta) + 54 \cos(2\theta) + 7 \cos(3\theta)], \end{aligned} \quad (48)$$




FIG. 4. Entanglement distribution $E_1(U|j, \mathbf{n})$ (Top) and its stereographic projection (Bottom) obtained by the SC states after the action of the unitary operator a) (68) b) (39) and c) (41) for the spin values $j = 1, 3/2, 2$, respectively. $E_q(U|\mathbf{n})$ has tetrahedral, icosahedral and octahedral symmetry, respectively. These unitary operators maximize e_p for $j = 1, 3/2$, and apparently for $j = 2$, respectively. The numerical values of the contour lines are marked in each color bar.

with U'_0 defined in (68). Its corresponding matrix \mathcal{D}_1 is with equal to

$$\mathcal{D}_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & \sqrt{2} \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & \sqrt{2} & 0 & 0 & 0 \end{pmatrix}, \quad (49)$$

which exhibits tetrahedral symmetry. The same point group can be observe in the entanglement distribution of U'_0 plotted in Fig. 4a. U'_0 does not create any entanglement when applied to four SC states pointing in the vertices of a regular tetrahedron, one of which is $|\mathbf{z}\rangle$. On the other hand, it transforms the SC states pointing along the vertices of the antipodal tetrahedron into maximally entangled states. For instance, $U'_0|-\mathbf{z}\rangle = |1, 0\rangle$. An alternative way to show the tetrahedral symmetry of the entanglement distribution is by direct calculation of the eigendecomposition of \mathcal{D}_1

$$\mathcal{D}_1 = \mathbb{1}_5 - 3|\psi_T\rangle\langle\psi_T| \quad (50)$$

$$|\psi_T\rangle = -\sqrt{\frac{2}{3}}|2, 1\rangle + \frac{1}{\sqrt{3}}|2, -2\rangle \quad (51)$$

a spin-2 state with tetrahedral symmetry [42]. By direct algebra, we obtain that

$$E_1(U'_0, |1, \mathbf{n}\rangle) = 3H_{|\psi_T\rangle}(\mathbf{n}), \quad (52)$$

i.e., the entanglement distribution of U'_0 is proportional to the Husimi function of the *tetrahedron* state $|\psi_T\rangle$.

Similar expressions for \mathcal{D}_1 and E_1 are obtained for a general $U \in \text{SU}(3)$ using the parametrization (12). By taking only the non-local term of the unitary gate $U = A$, we get that

$$\mathcal{D}_1 = \mathbb{1}_5 - \frac{4\Gamma_A}{3}|\psi_A\rangle\langle\psi_A|, \quad (53)$$

with

$$\Gamma_A = \sin^2 c_{12} + \sin^2 c_{13} + \sin^2 c_{23} \quad (54)$$

and

$$|\psi_A\rangle = \sqrt{\frac{3}{\Gamma_A}} \frac{\sin c_{12}}{2} (|2, 2\rangle + |2, -2\rangle) + i \left(\frac{\cos c_{12} - e^{-i(c_{13}+c_{23})}}{\sqrt{2\Gamma_A}} \right) |2, 0\rangle. \quad (55)$$

Thus,

$$E_1(U, |1, \mathbf{n}\rangle) = \frac{4\Gamma_A}{3} H_{|\psi_A\rangle}(\mathbf{n}), \quad (56)$$

with Γ_A proportional to $e_p(E_1, U)$ (14) [43].

B. $j=3/2$

We now plot $E_1(U_0 |3/2, \mathbf{n}\rangle)$, with U_0 the unitary transformation in (39), in Fig. 4b. The entanglement distribution

$$E_1(U_0 |3/2, \mathbf{n}\rangle) = \frac{1}{1152} [1090 + 192 \sin^5(\theta) \cos(\theta) \sin(5\phi) - 15 \cos(2\theta) - 18 \cos(4\theta) - 33 \cos(6\theta)] \quad (57)$$

has an icosahedral symmetry, and takes values in the interval [8/9, 80/81]. The rotational symmetries are also reflected in the configuration of the minima and maxima, which are arranged in an icosahedron and a dodecahedron, respectively. The corresponding \mathcal{D}_1 matrix has eigendecomposition given by

$$\mathcal{D}_1 = \mathbb{1}_7 - \frac{20}{9} \sum_{k=1}^3 |\psi_k\rangle \langle \psi_k|, \quad (58)$$

with states [44]

$$\begin{aligned} |\psi_1\rangle &= -i\sqrt{\frac{3}{5}} |3, 2\rangle + \sqrt{\frac{2}{5}} |3, -3\rangle, \\ |\psi_2\rangle &= i\sqrt{\frac{2}{5}} |3, 3\rangle + \sqrt{\frac{3}{5}} |3, -2\rangle, \\ |\psi_3\rangle &= |3, 0\rangle. \end{aligned} \quad (59)$$

The entanglement distribution is then given by

$$e_p(E_1, U_0) = \frac{20}{9} \sum_{k=1}^3 H_{|\psi_k\rangle}(\mathbf{n}). \quad (60)$$

C. $j=2$

We plot $E_1(U_0^{(1)}, |2, \mathbf{n}\rangle)$, with $U_0^{(1)}$ defined by (41), in Fig. 4c. We observe octahedral symmetry, confirmed by using the Majorana representation of mixed states [41]. In fact, the minima (maxima) of $E_1(U_0^{(1)}, |2, \mathbf{n}\rangle)$ lie on the vertices of a truncated octahedron (cuboctahedron) — the explicit expression of this function is rather long

and is not particularly enlightening. On the other hand, $E_2(U_0^{(2)}, |2, \mathbf{n}\rangle)$, with $U_0^{(2)}$ as in Eq. (42), is given by an affine transformation of the entanglement distribution plotted in Fig. 4b,

$$4E_2(U_0^{(2)}, |2, \mathbf{n}\rangle) = 9E_1(U_0^{\text{Eq. (39)}}, |3/2, \mathbf{n}\rangle) - 5. \quad (61)$$

Thus, $E_2(U_0^{(2)}, |2, \mathbf{n}\rangle)$ has icosahedral symmetry.

VI. AVERAGE OF e_p OVER THE UNITARY ORBIT

We calculate the average of $e_p(E_q, U)$ over the unitary operators $SU(d)$, with $d = 2j + 1$, and with respect to the normalized Haar measure ($\int d\mu(U) = 1$)

$$\overline{e_p(E_q, U)}^{\text{SU}(d)} \equiv \int e_p(E_q, U) d\mu(U) = 1 - \frac{1}{2j + 1 - q}, \quad (62)$$

the details can be found in Appendix D. The last equation can be written in terms of the dimensions of the initial and the reduced Hilbert spaces, $\mathcal{H}^{(q/2)} \otimes \mathcal{H}^{(j-q/2)}$ and $\mathcal{H}^{(q/2)}$ respectively,

$$\overline{e_p(E_q, U)}^{\text{SU}(d)} = 1 - \frac{\dim(\mathcal{H}^{(q/2)})}{\dim(\mathcal{H}^{(q/2)} \otimes \mathcal{H}^{(j-q/2)})}. \quad (63)$$

We observe that, similar to the non-symmetric case [25], the average of e_p increases as the dimension of the initial (resp. reduced) Hilbert space increases (resp. decreases). However, this dependence differs from that of the non-symmetric case. To corroborate this, let us briefly review e_p in the non-symmetric case.

We start with a system of n qudits $\mathcal{H}_D^{\otimes n}$. Then, we calculate its average linear entanglement entropy after we trace out $n - q$ constituents, Q_q (see [25] for the formal definition), where the average here means over all the possible bipartitions $q|n - q$ over the n constituents. In particular, if the state is symmetric, $Q_q = E_q$. Now, the entangling power with respect to a unitary matrix $\mathbb{U} \in SU(D^n)$, $e_p(Q_q, \mathbb{U})$, is equal to

$$e_p(Q_q, \mathbb{U}) = \int Q_q \left(\mathbb{U} (|\psi_1\rangle \otimes |\psi_2\rangle \otimes \cdots \otimes |\psi_n\rangle) \right) \otimes d\mu(\psi_1) \dots d\mu(\psi_n), \quad (64)$$

where $\int d\mu(\psi_k) = 1$. It turns out that its average over the unitary orbit $SU(D^n)$ is equal to [25]

$$\overline{e_p(Q_q, \mathbb{U})}^{\text{SU}(D^n)} = 1 - \frac{\dim(\mathcal{H}_D^{\otimes q}) + 1}{\dim(\mathcal{H}_D^{\otimes n}) + 1}. \quad (65)$$

where $\dim(\mathcal{H}_D^{\otimes \alpha}) = D^\alpha$. We can observe a difference between the previous equation and Eq. (62). However, a different nature between the average of e_p 's in the symmetric and non-symmetric case is expected because we integrated over different sets of product states (see Eqs. (5)

j	q	$\overline{e_p(E_q, U)}^{\text{SU}(2j+1)}$	$\max_U e_p(E_q, U)$
1	1	1/2	3/5
3/2	1	2/3	20/21
2	1	3/4	6889/7140
2	2	2/3	25/28

TABLE II. Average of $e_p(E_q, U)$ over the set of unitary gates $\text{SU}(2j+1)$ for several values of j and q . Additionally, we include the maximum possible value of e_p for $j=1$ and $3/2$, as well as the conjectured maximum values for $j=2$ obtained from numerical searches.

and (64) and over different sets of unitary gates (see Eqs. (62) and (65)). We tabulate $\overline{e_p(E_q, U)}$ for several values of j and q in Table II.

VII. e_p AND SCHMIDT NUMBERS

The reformulation of e_p in Eq. (26) shows that the entangling power increases as the subspace associated with the image of $\mathcal{N} = \mathcal{P}_{2j}/(4j+1)$, $\text{im}(\mathcal{N})$, is rotated to the orthogonal complement of \mathcal{M}_q^\perp . The Schmidt numbers of the states $|\psi\rangle \in \mathcal{H}^{(j)\otimes 2}$ are invariant under the action of \mathcal{U} . For instance, the states associated to coupled basis $|j, j, L, M\rangle$, or just $|L, M\rangle$ for short, have a Schmidt decomposition in terms of the decoupled basis $|j, m_1\rangle |j, m_2\rangle$. We write the explicit expressions for the states of $j=1$ and $3/2$ in Appendix E. In general, for a state $|\psi\rangle \in \text{im}(\mathcal{P}_{2j})$ to be connected by a \mathcal{U} transformation to a state $|\phi\rangle \in \text{im}(\mathcal{P}_{2j}^\perp)$, their Schmidt numbers have to be equal. We apply this idea to the search for the optimal unitary operators, for $j=1$ and $3/2$. In particular, this approach led us to the identification of U_0 for $j=3/2$, *i.e.*, Eq. (39).

A. Spin-1

Eq. (34) implies that the unitary transformations $\mathcal{U} = U \otimes U$ that maximize e_p are those that rotate one state from $\text{im}(\mathcal{P}_2)$ to $\text{im}(\mathcal{P}_0)$. Here, \mathcal{P}_0 contains only one state given by (see Appendix E)

$$|0, 0\rangle = \frac{1}{\sqrt{3}}(|1\rangle |-1\rangle - |0\rangle |0\rangle + |-1\rangle |1\rangle), \quad (66)$$

with Schmidt numbers $(1, 1, 1)/\sqrt{3}$. By direct inspection, we find that the state $|\Psi\rangle \in \text{im}(\mathcal{P}_2)$ given by

$$\begin{aligned} |\Psi\rangle &= \frac{1}{\sqrt{3}}(\sqrt{2}|2, 1\rangle - |2, -2\rangle) \\ &= \frac{1}{\sqrt{3}}(|1\rangle |0\rangle - |-1\rangle |-1\rangle + |0\rangle |1\rangle) \end{aligned} \quad (67)$$

can be transformed to $U'_0 \otimes U'_0 |\Psi\rangle = |0, 0\rangle \in \text{im}(\mathcal{P}_0)$ by

$$U'_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}. \quad (68)$$

U'_0 maximizes e_p and differs from U_0 in Eq. (15) by both left and right $\text{SU}(2)$ transformations.

B. Spin 3/2

Similarly, Eq. (38) suggests that \mathcal{U} must rotate three vectors from $\text{im}(\mathcal{P}_3)$ to $\text{im}(\mathcal{P}_1)$ in order to optimize e_p . The $|1, M\rangle$ states have Schmidt numbers (see Appendix E)

$$\begin{aligned} &\frac{1}{\sqrt{10}}(\sqrt{3}, \sqrt{3}, 2, 0) \quad \text{for } M = \pm 1, \\ &\frac{1}{2\sqrt{5}}(3, 3, 1, 1) \quad \text{for } M = 0. \end{aligned} \quad (69)$$

We find that certain states in $\text{im}(\mathcal{P}_3)$ have the same Schmidt numbers. Moreover, the \mathcal{U} defined by Eq. (39) effects the desired transformation

$$\begin{aligned} \mathcal{U} \left(\sqrt{\frac{3}{5}}|3, \mp 2\rangle - \sqrt{\frac{2}{5}}|3, \pm 3\rangle \right) &= |1, \pm 1\rangle, \\ \mathcal{U}|3, 0\rangle &= |1, 0\rangle. \end{aligned} \quad (70)$$

VIII. CONCLUSIONS AND PERSPECTIVES

In this work, we have studied the entangling power of unitary operators acting on pure spin- j states, which can be viewed as symmetric states of $N=2j$ qubits. The main differences with respect to the general non-symmetric case is that the set of product states are reduced to the SC states (see Eq. (5)), and the set of unitary gates is reduced from $\text{SU}(2^N)$ to $\text{SU}(2j+1)$ (See Sec. VI for more details). e_p is reformulated as the inner product of two $(2j+1)$ -vectors (26). One vector, \mathcal{M}_q , depends on the linear entanglement entropy E_q , while the other vector, \mathcal{N} , is transformed by the unitary matrix U . The components of both vectors are $\text{SU}(2)$ -invariant quantities, meaning that unitary transformations U, U' , differing only by left or right rotations, $U' = R_1 U R_2$, preserve them. Following this approach, several results and derivations are obtained, also raising new questions, which we summarize below.

We study in detail e_p for small spin values. Specifically, we identified the unitary gates that maximize $e_p(E_1, U)$ for $j=1$ and $j=3/2$. Through numerical calculations, we also discovered two unitary gates that are conjectured to maximize $e_p(E_q, U)$ for $q=1, 2$. These extremal unitaries possess highly symmetric entanglement distributions on the sphere (see Fig. 4) — note that a similar

characteristic is observed for the extremal spin states that maximize entanglement measures [45–48]. Additionally, the hermitian matrices $\mathcal{D}_q(U)$, associated with the entanglement distributions of these extremal unitaries, display peculiar characteristics such as high point group symmetries and multiple degeneracies in their eigenspectra. Also notably, these extremal unitaries can be represented as linear combinations of at most two permutations matrices with complex entries. The e_p reduces to an expression that contains the sum of the eigenvalues of \mathcal{D}_q . We also point out that $e_p(E_q, U) = e_p(E_q, U^\dagger)$ holds true for $j = 1$ and $3/2$, but fails, in general, for higher spins. This symmetry of e_p may be recast as time-reversal invariance, when U is generated by a hamiltonian, $U = e^{-itH}$.

We compute the average of $e_p(E_q, U)$ over unitary gates $SU(2j + 1)$ with respect to the Haar measure, Eq. (62), where we observe a difference with respect to the non-symmetric case Eq. (65). By sampling Haar-uniform random unitaries, we find that most of their associated invariant vectors cluster near the mean value (see Table II). These observations indicate that random unitary gates exhibit a statistical distribution with a narrow spread around the mean value. Further work could be done to derive the explicit statistical distribution of $e_p(E_q, U)$, as well as other variables such as the $SU(2)$ -invariant components p_L (31).

The vectors introduced in our geometrical approach to the calculation of e_p have components associated to the subspace projectors \mathcal{P}_L . Thus, finding the maximum of e_p involves searching for the unitary gate that rotates a subspace into another. Additionally, unitary transformations do not alter the Schmidt decomposition of the states in \mathcal{P}_L , suggesting that an alternative approach to maximizing e_p is to examine the possible Schmidt decomposition for the states spanning $\text{im}(\mathcal{P}_L)$. This approach proved useful in identifying the unitary gate that maximizes $e_p(E_1, U)$ for $j = 3/2$.

Lastly, we remark that, for spin- j pure states, the linear entanglement entropy E_q (1) coincides with the measure of anticoherent of order- q based on the purity (see Eq. (24) of Ref. [42]). Hence, the unitary gates with higher (symmetric) entangling power correspond to those with high capacities to generate anticoherence in the SC states. Anticoherence for pure states is known to be a measure of non-classicality [49–52]. It is also known that anticoherent states are useful in quantum-enhanced metrology of rotations [47, 53–55].

In summary, we have examined in detail the concept of entangling power for unitary matrices acting on spin states, introducing reformulations in terms of inner products between vectors associated to $SU(2)$ invariants and transformation of subspaces of bipartite states among themselves. Additionally, the entanglement distribution of a unitary gate is associated with a linear combinations of Husimi functions. These new perspectives could establish bridges between other quantities relevant for quantum information theory that initially appear unrelated, including those belonging to the non-symmetric case.

ACKNOWLEDGEMENTS

ESE acknowledges support from the postdoctoral fellowship of the IPD-STEMA program of the University of Liège (Belgium). DMG acknowledges support from the F.R.S.-FNRS under the Excellence of Science (EOS) programme. CC acknowledges support from the UNAM-PAPIIT project IN112224. JAM and DMG acknowledge support from the UNAM-PAPIIT project IN111122 (México).

Appendix A: Main equation

Here, we derive Eq. (16). But first, we introduce the basis of $\mathcal{HS}(\mathcal{H}^{(j)})$ defined by the *multipole operators* $\{T_{\sigma\mu}^{(j)}\}$, with $\sigma = 0, \dots, 2j$ and $\mu = -\sigma, \dots, \sigma$ [38, 56] and where we omit the superindex when there is no possible confusion. The $T_{\sigma\mu}$ operators can be written in terms of the Clebsch-Gordan coefficients $C_{j_1 m_1 j_2 m_2}^{j m}$ [38] and they are orthonormal with respect to the HS scalar product

$$\text{Tr}(T_{\sigma_1 \mu_1}^\dagger T_{\sigma_2 \mu_2}) = \delta_{\sigma_1 \sigma_2} \delta_{\mu_1 \mu_2}. \quad (\text{A1})$$

A density matrix $\rho \in \mathcal{HS}(\mathcal{H}^{(j)})$, representing a quantum state, can always be expanded in the $\{T_{\sigma\mu}\}$ basis

$$\rho = \sum_{\sigma=0}^{2j} \sum_{\mu=-\sigma}^{\sigma} \rho_{\sigma\mu} T_{\sigma\mu} = \sum_{\sigma=0}^{2j} \boldsymbol{\rho}_\sigma \cdot \mathbf{T}_\sigma, \quad (\text{A2})$$

where $\boldsymbol{\rho}_\sigma = (\rho_{\sigma\sigma}, \dots, \rho_{\sigma-\sigma})$ with $\rho_{\sigma\mu} = \text{Tr}(\rho T_{\sigma\mu}^\dagger)$, and $\mathbf{T}_\sigma = (T_{\sigma\sigma}, \dots, T_{\sigma-\sigma})$ is a vector of matrices. In particular, $T_{00} = (2j + 1)^{-1/2} \mathbb{1}$, and then $\rho_{00} = (2j + 1)^{-1/2}$ for any mixed state. In the picture of the spin- j states seeing as $2j$ spin-1/2 constituents, one can calculate the reduced density matrix $\rho_q = \text{Tr}_{2j-q} \rho$, after tracing out $2j - q$ spins-1/2. Notably, the multipole expansion (A2) of ρ_q has a simple expression in terms of the original state ρ [37, 41]

$$(\rho_q)_{\sigma\mu} = \frac{q!}{(2j)!} \sqrt{\frac{(2j - \sigma)!(2j + \sigma + 1)!}{(q - \sigma)!(q + \sigma + 1)!}} \rho_{\sigma\mu}, \quad (\text{A3})$$

with $\sigma = 0, \dots, q$ and $\mu = -\sigma, \dots, \sigma$.

Now, we calculate the entangling power (1) for a general spin- j system. First, we write the general expression of the SC states, transformed by U ,

$$U |j, \mathbf{n}\rangle \langle j, \mathbf{n}| U^\dagger = \frac{1}{\sqrt{2^j + 1}} T_{00}^{(j)} + \sum_{\sigma=1}^{2j} \boldsymbol{\rho}_\sigma \cdot \mathbf{T}_\sigma^{(j)}, \quad (\text{A4})$$

with

$$\rho_{\sigma\mu} = \langle j, \mathbf{n}| U^\dagger T_{\sigma\mu}^{(j)\dagger} U |j, \mathbf{n}\rangle. \quad (\text{A5})$$

By tracing out the subsystem B , *i.e.*, $2j - q$ spin-1/2 constituents, we obtain

$$\begin{aligned} \rho_A &= \text{Tr}_{2j-q} (U |j, \mathbf{n}\rangle \langle j, \mathbf{n}| U^\dagger) \\ &= \frac{1}{\sqrt{q+1}} T_{00}^{(q/2)} \\ &\quad + \sum_{\sigma=1}^q \frac{q!}{(2j)!} \sqrt{\frac{(2j-\sigma)!(2j+\sigma+1)!}{(q-\sigma)!(q+\sigma+1)!}} \boldsymbol{\rho}_\sigma \cdot \mathbf{T}_\sigma^{(q/2)}, \end{aligned} \quad (\text{A6})$$

where we use Eq. (A3). We then get that

$$\begin{aligned} E_q(U |j, \mathbf{n}\rangle) &= 1 - \frac{(q+1)(q!)^2}{q(2j)!^2} \times \\ &\quad \sum_{\sigma=1}^q \frac{(2j-\sigma)!(2j+\sigma+1)!}{(q-\sigma)!(q+\sigma+1)!} |\boldsymbol{\rho}_\sigma|^2, \\ &= 1 - \langle j, \mathbf{n} | \otimes \langle j, \mathbf{n} | \mathcal{U}^\dagger \mathcal{M}_q \mathcal{U} |j, \mathbf{n}\rangle \otimes |j, \mathbf{n}\rangle \end{aligned} \quad (\text{A7})$$

where $\mathcal{U} = U \otimes U$ as defined in the main text,

$$\mathcal{T}_\sigma = \sum_{\mu=-\sigma}^{\sigma} T_{\sigma\mu} \otimes T_{\sigma\mu}^\dagger \quad (\text{A8})$$

and

$$\begin{aligned} \mathcal{M}_q &= \frac{(q+1)(q!)^2}{q(2j)!^2} \sum_{\sigma=1}^q \frac{(2j-\sigma)!(2j+\sigma+1)!}{(q-\sigma)!(q+\sigma+1)!} \mathcal{T}_\sigma \\ &= \frac{q+1}{q} \sum_{\sigma=1}^q \frac{\begin{Bmatrix} q/2 & q/2 & q \\ q/2 & q/2 & \sigma \end{Bmatrix}}{\begin{Bmatrix} j & j & 2j \\ j & j & \sigma \end{Bmatrix}} \mathcal{T}_\sigma. \end{aligned} \quad (\text{A9})$$

The \mathcal{T}_σ 's can be written as a linear combination of the \mathcal{P}_L operators (see **Identity 1** of Appendix C)

$$\mathcal{T}_\sigma = (2\sigma+1) \sum_{L=0}^{2j} (-1)^{2j+L} \begin{Bmatrix} j & j & L \\ j & j & \sigma \end{Bmatrix} \mathcal{P}_L. \quad (\text{A10})$$

This result gives the expression (17) of \mathcal{M}_q . Lastly, we use that $|j, \mathbf{n}\rangle \otimes |j, \mathbf{n}\rangle = |2j, \mathbf{n}\rangle$ to obtain Eq. (16).

Appendix B: The T operator basis

Identity 1 of Appendix C lets us write e_p (16) as an inner product of vectors of SU(2) invariants associ-

ated to the \mathcal{T}_σ operators. The operator \mathcal{M}_q is written in Eq. (A9), while \mathcal{N} is expanded as

$$\mathcal{N} = \sum_{L=0}^{2j} \begin{Bmatrix} j & j & 2j \\ j & j & L \end{Bmatrix} \mathcal{T}_L. \quad (\text{B1})$$

The \mathcal{T}_σ operators are hermitian, orthogonal $\text{Tr}(\mathcal{T}_\sigma \mathcal{T}_{\sigma'}) = (2\sigma+1)\delta_{\sigma\sigma'}$, and invariant under diagonal SU(2) transformations in the double-copy space $\mathcal{H}^{(j)\otimes 2}$. Thus, and similarly as we did in Sec. III, we can associate to every operator $V \in \mathcal{HS}(\mathcal{H}^{(j)\otimes 2})$ a $(2j+1)$ -vector of SU(2) invariants

$$\vec{V}^\mathcal{T} = \left(\text{Tr}(V\tilde{\mathcal{T}}_0), \dots, \text{Tr}(V\tilde{\mathcal{T}}_\sigma), \dots, \text{Tr}(V\tilde{\mathcal{T}}_{2j}) \right), \quad (\text{B2})$$

with $\tilde{\mathcal{T}}_\sigma = \mathcal{T}_\sigma / \sqrt{2\sigma+1}$, and where the superscript denotes that the vector is written in the \mathcal{T} basis. We can follow the same procedure as in Sec. III to obtain the same reformulation of the e_p as an inner product of two vectors given in Eqs. (23)-(26) but now expressed in the \mathcal{T} basis. We also have that each vector, now written in the \mathcal{T} basis, lies in a hyperplane, even after the action of U ,

$$\begin{aligned} \sum_{\sigma=0}^{2j} \left(\hat{u} \vec{\mathcal{N}} \right)_\sigma^\mathcal{T} &= \sum_{\sigma=0}^{2j} \text{Tr}(u \mathcal{N} u^\dagger \mathcal{T}_\sigma) = 1. \\ \sum_{\sigma=0}^{2j} \left(\hat{u} \vec{\mathcal{M}}_q \right)_\sigma^\mathcal{T} &= \frac{2(j+1)(2j+1)}{(2j-q+1)}. \end{aligned} \quad (\text{B3})$$

We prove these results in Identity 3 of Appendix C. Since \mathcal{T}_0 is proportional to the identity, $\left(\hat{u} \vec{\mathcal{V}} \right)_0^\mathcal{T} = \left(\vec{\mathcal{V}} \right)_0^\mathcal{T}$. In particular $\left(\vec{\mathcal{N}} \right)_0^\mathcal{T} = 1/(2j+1)$ and $\left(\vec{\mathcal{M}}_q \right)_0^\mathcal{T} = 0$. We write \mathcal{N} and \mathcal{M}_q for $j = 1, 3/2, 2$ and $5/2$ in the \mathcal{T} basis in Table III. The properties of the 6j-symbols show that the components of $\vec{\mathcal{N}}$ and $\vec{\mathcal{M}}_q$ are always positive in the \mathcal{T} basis [38]. However, the disadvantage of the \mathcal{T} basis is that a \mathcal{U} transformation combines all the \mathcal{T} components, not only half of them as in the \mathcal{P} basis.

Appendix C: Useful identities

Here, we prove several identities used throughout the paper.

j	$\vec{\mathcal{N}}$ in \mathcal{T}_σ basis	$\vec{\mathcal{M}}_q$ in \mathcal{T}_σ basis
1	$\left(\frac{1}{3}, \frac{1}{2\sqrt{3}}, \frac{1}{6\sqrt{5}}\right)$	$(0, 2\sqrt{3}, 0)$
3/2	$\left(\frac{1}{4}, \frac{3\sqrt{3}}{20}, \frac{1}{4\sqrt{5}}, \frac{1}{20\sqrt{7}}\right)$	$\left(0, \frac{20}{3\sqrt{3}}, 0, 0\right)$
2	$\left(\frac{1}{5}, 0, \frac{2}{5\sqrt{3}}, \frac{2}{7\sqrt{5}}, \frac{1}{10\sqrt{7}}, \frac{1}{210}\right)$	$\begin{cases} \left(0, \frac{5\sqrt{3}}{2}, 0, 0, 0\right) \\ \left(0, \frac{15\sqrt{3}}{8}, \frac{7\sqrt{5}}{8}, 0, 0\right) \end{cases}$
5/2	$\left(\frac{1}{6}, 0, \frac{5}{14\sqrt{3}}, \frac{5\sqrt{5}}{84}, \frac{5}{36\sqrt{7}}, \frac{1}{84}, \frac{1}{252\sqrt{11}}\right)$	$\begin{cases} \left(0, \frac{14\sqrt{3}}{5}, 0, 0, 0, 0\right) \\ \left(0, \frac{21\sqrt{3}}{10}, \frac{21}{5\sqrt{5}}, 0, 0, 0\right) \end{cases}$

TABLE III. Vectors of SU(2) invariants of the operators \mathcal{N} and \mathcal{M}_q (A9) in the \mathcal{T}_σ basis for the spin values $j = 1, 3/2, 2, 5/2$ and $q = 1, \dots, \lfloor j \rfloor$. The vector $\vec{\mathcal{M}}_q$ has q nonzero entries.

Identity 1 *There is a linear transformation between the operators \mathcal{T}_σ and \mathcal{P}_L*

$$\mathcal{T}_\sigma = (2\sigma + 1) \sum_{L=0}^{2j} (-1)^{2j+L} \left\{ \begin{matrix} j & j & L \\ j & j & \sigma \end{matrix} \right\} \mathcal{P}_L, \quad \mathcal{P}_L = (-1)^{2j+L} (2L + 1) \sum_{\sigma=0}^{2j} \left\{ \begin{matrix} j & j & \sigma \\ j & j & L \end{matrix} \right\} \mathcal{T}_\sigma. \quad (\text{C1})$$

Proof.

$$\begin{aligned} \mathcal{T}_\sigma &= \sum_{\mu=-\sigma}^{\sigma} T_{\sigma\mu} \otimes T_{\sigma\mu}^\dagger \\ &= \left(\frac{2\sigma+1}{2j+1}\right) \sum_{\mu} \sum_{\substack{m, m' \\ n, n'}} C_{jm', \sigma\mu}^{jm} C_{jn', \sigma\mu}^{jn} |j, m\rangle \langle j, m'| \otimes |j, n'\rangle \langle j, n| \\ &= \left(\frac{2\sigma+1}{2j+1}\right) \sum_{\mu} \sum_{\substack{m, m' \\ n, n'}} \sum_{\substack{L_1, M_1 \\ L_2, M_2}} C_{jm', \sigma\mu}^{jm} C_{jn', \sigma\mu}^{jn} C_{jm, jn'}^{L_1 M_1} C_{jm', jn}^{L_2 M_2} |j, j, L_1, M_1\rangle \langle j, j, L_2, M_2| \\ &= \left(\frac{2\sigma+1}{2j+1}\right) (-1)^\sigma \sum_{\substack{L_1, M_1 \\ L_2, M_2}} \sum_M \sum_{\substack{m, m' \\ n, n'}} C_{jm, jn'}^{L_1 M_1} C_{jm', \sigma\mu}^{jm} C_{jm', jn}^{L_2 M_2} C_{\sigma\mu, jn'}^{jn} |j, j, L_1, M_1\rangle \langle j, j, L_2, M_2| \\ &= (2\sigma+1) \sum_{L_1, M_1} (-1)^{2j+L_1} \left\{ \begin{matrix} j & j & L_1 \\ j & j & \sigma \end{matrix} \right\} |L_1, M_1\rangle_C \langle L_1, M_1|_C \\ &= (2\sigma+1) \sum_{L=0}^{2j} (-1)^{2j+L} \left\{ \begin{matrix} j & j & L \\ j & j & \sigma \end{matrix} \right\} \mathcal{P}_L, \end{aligned} \quad (\text{C2})$$

where we use the decoupled and coupled bases of the angular momentum (7) and the definition of the 6j symbol (see p. 291, Eq. (8) of Ref. [38]). Lastly, we invert the previous equation with the orthogonality of the 6j symbols (See page 291 or Ref. [38] for the general formula)

$$(2\sigma+1) \sum_{L=0}^{2j} (2L+1) \left\{ \begin{matrix} j & j & L \\ j & j & \sigma \end{matrix} \right\} \left\{ \begin{matrix} j & j & L \\ j & j & \sigma' \end{matrix} \right\} = \delta_{\sigma\sigma'}, \quad (\text{C3})$$

to write \mathcal{P}_L in terms of the operators \mathcal{T}_σ . \square

Identity 2 *For any unitary transformation $\mathcal{U} = U \otimes U$ and operator $V = \sum_{K=0}^{2j} v_K \mathcal{P}_K$, it holds that*

$$\sum_{\substack{L=0 \\ 2j \equiv L + \delta \pmod{2}}}^{2j} \text{Tr}(\mathcal{U} V \mathcal{U}^\dagger \mathcal{P}_L) = \sum_{\substack{L=0 \\ 2j \equiv L + \delta \pmod{2}}}^{2j} \text{Tr}(V \mathcal{P}_L), \quad (\text{C4})$$

for $\delta = 0$ and 1, or when L runs over all the indices. In particular,

$$\sum_{\substack{L=0 \\ K \equiv L \pmod{2}}}^{2j} \text{Tr}(\mathcal{U}\mathcal{P}_L\mathcal{U}^\dagger\mathcal{P}_K) = 2K + 1, \quad \sum_{\substack{L=0 \\ 2j \equiv L \pmod{2}}}^{2j} \text{Tr}(\mathcal{U}\mathcal{P}_L\mathcal{U}^\dagger\mathcal{M}_q) = \frac{(j+1)(2j+1)}{2j+1-q}. \quad (\text{C5})$$

Proof. By resolution of the unity of the \mathcal{P}_L operators, we have that

$$\sum_{L=0}^{2j} \text{Tr}(\mathcal{U}\mathcal{P}_L\mathcal{U}^\dagger\mathcal{P}_K) = \text{Tr}(\mathcal{P}_K) = 2K + 1. \quad (\text{C6})$$

Now, the unitary transformations $\mathcal{U} = U \otimes U$ preserve the orthogonality between the projectors of different parity

$$\text{Tr}(\mathcal{U}\mathcal{P}_L\mathcal{U}^\dagger\mathcal{P}_K) = 0, \quad \text{for } L \equiv K + 1 \pmod{2}, \quad (\text{C7})$$

because they preserve the permutation exchange of the states $|\Psi\rangle \in \mathcal{H}^{(j)\otimes 2}$. Thus, any operator $V = \sum_{K=0}^{2j} v_K \mathcal{P}_K$ can be split in the components with K odd and even, and these components do not mixed after a transformation \mathcal{U} . This proves (C4). In particular,

$$\sum_{L=0}^{2j} \text{Tr}(\mathcal{U}\mathcal{P}_L\mathcal{U}^\dagger\mathcal{P}_K) = \sum_{\substack{L=0 \\ K \equiv L \pmod{2}}}^{2j} \text{Tr}(\mathcal{U}\mathcal{P}_L\mathcal{U}^\dagger\mathcal{P}_K) = \text{Tr}(\mathcal{P}_K) = 2K + 1. \quad (\text{C8})$$

For \mathcal{M}_q , we have that

$$\sum_{\substack{L=0 \\ 2j \equiv L + \delta \pmod{2}}}^{2j} \text{Tr}(\mathcal{U}\mathcal{M}_q\mathcal{U}^\dagger\mathcal{P}_L) = \frac{1}{2} \sum_{L=0}^{2j} [1 + (-1)^{2j+L+\delta}] \text{Tr}(\mathcal{M}_q\mathcal{P}_L) = (-1)^\delta \frac{(j+1)(2j+1)}{2j+1-q}, \quad (\text{C9})$$

where we use identities of the 6j symbols [38]. \square

Identity 3 Any hermitian operator $V = \sum_{\sigma=0}^{2j} v_\sigma \mathcal{T}_\sigma$ satisfies

$$\sum_{\sigma=0}^{2j} \text{Tr}(\mathcal{U}V\mathcal{U}^\dagger\mathcal{T}_\sigma) = \sum_{\sigma=0}^{2j} \text{Tr}(V\mathcal{T}_\sigma), \quad (\text{C10})$$

where $\mathcal{U} = U \otimes U$ and U a unitary operator. In particular,

$$\sum_{\sigma=0}^{2j} \text{Tr}(\mathcal{M}_q\mathcal{T}_\sigma) = \frac{2(j+1)(2j+1)}{(2j-q+1)}, \quad \sum_{\sigma=0}^{2j} \text{Tr}(\mathcal{N}\mathcal{T}_\sigma) = \frac{2j}{2j+1}. \quad (\text{C11})$$

Proof. The following calculation will prove convenient later on,

$$\begin{aligned} \sum_{\sigma=0}^{2j} \mathcal{T}_\sigma &= \sum_{\sigma=0}^{2j} \sum_{\mu=-\sigma}^{\sigma} \sum_{\substack{m_1, m_2 \\ n_1, n_2}} (-1)^{2j-m_2-n_2} C_{j m_1 j - m_2}^{\sigma \mu} C_{j n_1 j - n_2}^{\sigma \mu} |j, m_1, j, n_2\rangle \langle j, m_2, j, n_1| \\ &= \sum_{\substack{m_1, m_2 \\ n_1, n_2}} (-1)^{2j-m_2-n_2} \delta_{m_1, n_1} \delta_{m_2, n_2} |j, m_1, j, n_2\rangle \langle j, m_2, j, n_1| \\ &= \sum_{m_1, m_2} |j, m_1, j, m_2\rangle \langle j, m_2, j, m_1| \end{aligned} \quad (\text{C12})$$

where we use that $2(j-m)$ is always an even integer number. Now, let us prove the result for the \mathcal{T}_σ operators

$$\sum_{\sigma=0}^{2j} \text{Tr}(\mathcal{U}^\dagger\mathcal{T}_\tau\mathcal{U}\mathcal{T}_\sigma) = \sum_{\tau=-\sigma}^{\sigma} \sum_{m_1, m_2} \langle j, m_2 | U^\dagger T_{\tau\nu} U |j, m_1\rangle \langle j, m_1 | U^\dagger T_{\tau\nu}^\dagger U |j, m_2\rangle = 2\tau + 1 = \sum_{\sigma=0}^{2j} \text{Tr}(\mathcal{T}_\tau\mathcal{T}_\sigma), \quad (\text{C13})$$

The first equality will be valid for any V by linearity. In particular, we get for $V = \mathcal{N}$ that

$$\sum_{\sigma=0}^{2j} \text{Tr}(\mathcal{N}\mathcal{T}_\sigma) = \sum_{\sigma=0}^{2j} \sum_{L=1}^{2j} \left\{ \begin{matrix} j & j & 2j \\ j & j & L \end{matrix} \right\} \text{Tr}(\mathcal{T}_L\mathcal{T}_\sigma) = \sum_{L=1}^{2j} (2L+1) \left\{ \begin{matrix} j & j & 2j \\ j & j & L \end{matrix} \right\} = \frac{2j}{2j+1}, \quad (\text{C14})$$

where we use the following identities of the 6j symbol [38]

$$\sum_{L=0}^{2j} (2L+1) \left\{ \begin{matrix} j & j & 2j \\ j & j & L \end{matrix} \right\} = 1, \quad \left\{ \begin{matrix} j & j & 2j \\ j & j & 0 \end{matrix} \right\} = \frac{1}{2j+1}. \quad (\text{C15})$$

Similarly,

$$\sum_{\sigma=0}^{2j} \text{Tr}(\mathcal{M}_q\mathcal{T}_\sigma) = \frac{q+1}{q} \sum_{\tau=1}^q (2\tau+1) \frac{\left\{ \begin{matrix} q/2 & q/2 & q \\ q/2 & q/2 & \tau \end{matrix} \right\}}{\left\{ \begin{matrix} j & j & 2j \\ j & j & \tau \end{matrix} \right\}} = \frac{2(j+1)(2j+1)}{2j+1-q}. \quad \square \quad (\text{C16})$$

Appendix D: Proof of Eq. (62)

We start with the following identity regarding the integration over the $\text{SU}(d)$ unitary matrices and where A is a $d^2 \times d^2$ matrix (see **Proposition 3.9** in [57])

$$\int \mathcal{U} A \mathcal{U}^\dagger d\mu(U) = \left[\frac{\text{Tr}(A)}{d^2-1} - \frac{\text{Tr}(AF)}{d(d^2-1)} \right] \mathbb{1}_{d^2} - \left[\frac{\text{Tr}(A)}{d(d^2-1)} - \frac{\text{Tr}(AF)}{d^2-1} \right] F \quad (\text{D1})$$

where $F|\psi_1\rangle|\psi_2\rangle = |\psi_2\rangle|\psi_1\rangle$ is the swap operator. In particular $F\mathcal{P}_L = \mathcal{P}_L F = (-1)^{2j-L}\mathcal{P}_L$. Thus,

$$\int e_{\text{p}}(E_q, U) d\mu(U) = 1 - \text{Tr} \left(\left[\int \mathcal{U} \mathcal{N} \mathcal{U}^\dagger d\mu(U) \right] \mathcal{M}_q \right) = 1 - \frac{1}{d(d+1)} [\text{Tr}(\mathcal{M}_q) + \text{Tr}(F\mathcal{M}_q)]. \quad (\text{D2})$$

We add the resolution of the unity $\mathbb{1}_{d^2} = \sum_{L=0}^{2j} \mathcal{P}_L$ to use Eq. (C9) in the last equation

$$\int e_{\text{p}}(E_q, U) d\mu(U) = 1 - \frac{2(j+1)(2j+1)}{d(d+1)(2j+1-q)} = \frac{2j-q}{2j+1-q}. \quad (\text{D3})$$

Appendix E: Spin states in the coupled and decoupled bases

Here, we write the coupled basis $|L, M\rangle \equiv |j_1, j_2, L, M\rangle$ in terms of the decoupled basis $|m_1\rangle|m_2\rangle \equiv |j_1, m_1\rangle|j_2, m_2\rangle$. For $j = 1$, we have

$$\begin{aligned} |2, \pm 2\rangle &= |\pm 1\rangle|\pm 1\rangle, \\ |2, \pm 1\rangle &= \frac{1}{\sqrt{2}} (|\pm 1\rangle|0\rangle + |0\rangle|\pm 1\rangle), \\ |2, 0\rangle &= \frac{1}{\sqrt{6}} (|1\rangle|-1\rangle + 2|0\rangle|0\rangle + |-1\rangle|1\rangle), \\ |0, 0\rangle &= \frac{1}{\sqrt{3}} (|1\rangle|-1\rangle - |0\rangle|0\rangle + |-1\rangle|1\rangle), \end{aligned} \quad (\text{E1})$$

And for $j = 3/2$, it reads

$$\begin{aligned}
|3, \pm 3\rangle &= \left| \pm \frac{3}{2} \right\rangle \left| \pm \frac{3}{2} \right\rangle, \\
|3, \pm 2\rangle &= \frac{1}{\sqrt{2}} \left(\left| \pm \frac{3}{2} \right\rangle \left| \pm \frac{1}{2} \right\rangle + \left| \pm \frac{1}{2} \right\rangle \left| \pm \frac{3}{2} \right\rangle \right), \\
|3, \pm 1\rangle &= \frac{1}{\sqrt{5}} \left(\left| \pm \frac{3}{2} \right\rangle \left| \mp \frac{1}{2} \right\rangle + \left| \mp \frac{1}{2} \right\rangle \left| \pm \frac{3}{2} \right\rangle + \sqrt{3} \left| \pm \frac{1}{2} \right\rangle \left| \pm \frac{1}{2} \right\rangle \right), \\
|3, 0\rangle &= \frac{1}{2\sqrt{5}} \left(\left| \frac{3}{2} \right\rangle \left| -\frac{3}{2} \right\rangle + \left| -\frac{3}{2} \right\rangle \left| \frac{3}{2} \right\rangle + 3 \left| \frac{1}{2} \right\rangle \left| -\frac{1}{2} \right\rangle + 3 \left| -\frac{1}{2} \right\rangle \left| \frac{1}{2} \right\rangle \right),
\end{aligned} \tag{E2}$$

$$\begin{aligned}
|1, \pm 1\rangle &= \frac{1}{\sqrt{10}} \left(\sqrt{3} \left| \pm \frac{3}{2} \right\rangle \left| \mp \frac{1}{2} \right\rangle + \sqrt{3} \left| \mp \frac{1}{2} \right\rangle \left| \pm \frac{3}{2} \right\rangle - 2 \left| \pm \frac{1}{2} \right\rangle \left| \pm \frac{1}{2} \right\rangle \right), \\
|1, 0\rangle &= \frac{1}{2\sqrt{5}} \left(3 \left| \frac{3}{2} \right\rangle \left| -\frac{3}{2} \right\rangle + 3 \left| -\frac{3}{2} \right\rangle \left| \frac{3}{2} \right\rangle - \left| \frac{1}{2} \right\rangle \left| -\frac{1}{2} \right\rangle - \left| -\frac{1}{2} \right\rangle \left| \frac{1}{2} \right\rangle \right),
\end{aligned} \tag{E3}$$

-
- [1] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Rev. Mod. Phys.* **81**, 865 (2009).
- [2] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information: 10th Anniversary Edition*, 10th ed. (Cambridge University Press, New York, NY, USA, 2011).
- [3] I. Bengtsson and K. Życzkowski, *Geometry of Quantum States (2nd Ed.)* (Cambridge University Press, 2017).
- [4] W. Dür, G. Vidal, J. I. Cirac, N. Linden, and S. Popescu, *Phys. Rev. Lett.* **87**, 137901 (2001).
- [5] M. M. Wolf, J. Eisert, and M. B. Plenio, *Phys. Rev. Lett.* **90**, 047904 (2003).
- [6] H.-N. Xiong, X.-M. Lu, and X. Wang, *Journal of Physics B: Atomic, Molecular and Optical Physics* **45**, 015501 (2011).
- [7] B. Kraus and J. I. Cirac, *Phys. Rev. A* **63**, 062309 (2001).
- [8] J. Zhang, J. Vala, S. Sastry, and K. B. Whaley, *Phys. Rev. A* **67**, 042313 (2003).
- [9] T. Kalsi, A. Romito, and H. Schomerus, *Journal of Physics A: Mathematical and Theoretical* **55**, 264009 (2022).
- [10] H. L. Tang, K. Connelly, A. Warren, F. Zhuang, S. E. Economou, and E. Barnes, *Phys. Rev. Appl.* **19**, 044094 (2023).
- [11] F. Caruso, A. W. Chin, A. Datta, S. F. Huelga, and M. B. Plenio, *Phys. Rev. A* **81**, 062346 (2010).
- [12] P. Zanardi, C. Zalka, and L. Faoro, *Phys. Rev. A* **62**, 030301 (2000).
- [13] M. A. Nielsen, C. M. Dawson, J. L. Dodd, A. Gilchrist, D. Mortimer, T. J. Osborne, M. J. Bremner, A. W. Harrow, and A. Hines, *Phys. Rev. A* **67**, 052301 (2003).
- [14] B. Jonnadula, P. Mandayam, K. Życzkowski, and A. Lakshminarayan, *Phys. Rev. A* **95**, 040302 (2017).
- [15] L. Clarisse, S. Ghosh, S. Severini, and A. Sudbery, *Physics Letters A* **365**, 400 (2007).
- [16] P. Zanardi, *Phys. Rev. A* **63**, 040304 (2001).
- [17] A. T. Rezakhani, *Phys. Rev. A* **70**, 052313 (2004).
- [18] S. Balakrishnan and R. Sankaranarayanan, *Phys. Rev. A* **79**, 052339 (2009).
- [19] Y. Shen and L. Chen, *Journal of Physics A: Mathematical and Theoretical* **51**, 395303 (2018).
- [20] X. Wang, B. C. Sanders, and D. W. Berry, *Phys. Rev. A* **67**, 042323 (2003).
- [21] Y. Yang, X. Wang, and Z. Sun, *Physics Letters A* **372**, 4369 (2008).
- [22] M. Musz, M. Kuś, and K. Życzkowski, *Phys. Rev. A* **87**, 022111 (2013).
- [23] J. Chen, Z. Ji, D. W. Kribs, B. Zeng, and F. Zhang, *Journal of Physics A: Mathematical and Theoretical* **52**, 215302 (2019).
- [24] F.-Z. Kong and J.-L. Zhao, *Laser Physics Letters* **21**, 055206 (2024).
- [25] A. J. Scott, *Phys. Rev. A* **69**, 052330 (2004).
- [26] T. Linowski, G. Rajchel-Mieldzióć, and K. Życzkowski, *Journal of Physics A: Mathematical and Theoretical* **53**, 125303 (2020).
- [27] X. Wang and P. Zanardi, *Phys. Rev. A* **66**, 044303 (2002).
- [28] Y. Makhlin, *Quantum Information Processing* **1**, 243 (2002).
- [29] S. Balakrishnan and R. Sankaranarayanan, *Phys. Rev. A* **82**, 034301 (2010).
- [30] S. Singh and I. Nechita, *Journal of Physics A: Mathematical and Theoretical* **55**, 255302 (2022).
- [31] A. J. Scott and C. M. Caves, *Journal of Physics A: Mathematical and General* **36**, 9553 (2003).
- [32] R. Pal and A. Lakshminarayan, *Phys. Rev. B* **98**, 174304 (2018).
- [33] S. Aravinda, S. A. Rather, and A. Lakshminarayan, *Phys. Rev. Res.* **3**, 043034 (2021).
- [34] C. Chryssomalakos, E. Guzmán-González, and E. Serrano-Ensástiga, *J. Phys. A: Math. Theor.* **51**, 165202 (2018).
- [35] D. Morachis Galindo and J. A. Maytorena, *Phys. Rev. A* **105**, 012601 (2022).
- [36] M. Byrd, *Journal of Mathematical Physics* **39**, 6125 (1998).
- [37] J. Denis and J. Martin, *Phys. Rev. Res.* **4**, 013178 (2022).
- [38] D. Varshalovich, A. Moskalev, and V. Khersonskii, *Quan-*

- tum Theory of Angular Momentum* (World Scientific, 1988).
- [39] Here, we consider the Q matrix of [35] in the symmetric subspace of $\mathcal{H}_{1/2}^{\otimes 2}$.
- [40] G. S. Agarwal, *Phys. Rev. A* **24**, 2889 (1981).
- [41] E. Serrano-Ensástiga and D. Braun, *Physical Review A* **101**, 022332 (2020).
- [42] D. Baguette and J. Martin, *Phys. Rev. A* **96**, 032304 (2017).
- [43] Using the multipole expansion (A2), each spin-1 operator has associated to it a spin-2 state. Specifically, the components $\rho_2(U)$ can be expressed as $|\rho_2(U)\rangle = \sum_{m=-2}^2 \rho_{2m}(U) |2, m\rangle$. In particular, for $U = A \in \text{SU}(3)$, where A is defined in Eq. (12), we obtain that $|\rho_2(A^{*2})\rangle \propto |\psi_A\rangle$ (see Eq. (55)). However, this proportionality does not extend to higher spins.
- [44] The spin-3 states shown in Eq. (59) span a 3-dimensional 1-anticoherent subspace (see Ref. [55] for more details).
- [45] J. Martin, O. Giraud, P. A. Braun, D. Braun, and T. Bastin, *Phys. Rev. A* **81**, 062347 (2010).
- [46] O. Giraud, P. Braun, and D. Braun, *New Journal of Physics* **12**, 063005 (2010).
- [47] C. Chryssomalakos, L. Hanotel, E. Guzmán-González, D. Braun, E. Serrano-Ensástiga, and K. Życzkowski, *Phys. Rev. A* **104**, 012407 (2021).
- [48] A. Z. Goldberg, A. B. Klimov, M. Grassl, G. Leuchs, and L. L. Sánchez-Soto, *AVS Quantum Science* **2**, 044701 (2020).
- [49] J. Zimba, *Electronic Journal of Theoretical Physics* **3**, 143 (2006).
- [50] J. Crann, R. Pereira, and D. W. Kribs, *Journal of Physics A: Mathematical and Theoretical* **43**, 255307 (2010).
- [51] O. Giraud, D. Braun, D. Baguette, T. Bastin, and J. Martin, *Phys. Rev. Lett.* **114**, 080401 (2015).
- [52] D. Baguette, F. Damanet, O. Giraud, and J. Martin, *Phys. Rev. A* **92**, 052333 (2015).
- [53] C. Chryssomalakos and H. Hernández-Coronado, *Phys. Rev. A* **95** (2017).
- [54] A. Z. Goldberg and D. F. V. James, *Phys. Rev. A* **98**, 032113 (2018).
- [55] E. Serrano-Ensástiga, C. Chryssomalakos, and J. Martin, *Quantum metrology of rotations with mixed spin states* (2024), arXiv:2404.15548.
- [56] U. Fano, *Phys. Rev.* **90**, 577 (1953).
- [57] L. Zhang, *Matrix integrals over unitary groups: An application of schur-weyl duality* (2015), arXiv:1408.3782.