# Strongest quantum nonlocality in N-partite systems

Mengying Hu,<sup>1,2,3</sup> Ting Gao,<sup>1,2,3,\*</sup> and Fengli Yan<sup>4,†</sup>

<sup>1</sup>School of Mathematical Sciences, Hebei Normal University, Shijiazhuang 050024, China

<sup>2</sup>Hebei Mathematics Research Center, Hebei Normal University, Shijiazhuang 050024, China

<sup>3</sup>Hebei International Joint Research Center for Mathematics and Interdisciplinary Science,

Hebei Normal University, Shijiazhuang 050024, China

<sup>4</sup>College of Physics, Hebei Key Laboratory of Photophysics Research and Application,

Hebei Normal University, Shijiazhuang 050024, China

A set of orthogonal states possesses the strongest quantum nonlocality if only a trivial orthogonalitypreserving positive operator-valued measure (POVM) can be performed for each bipartition of the subsystems. This concept originated from the strong quantum nonlocality proposed by Halder *et al.* [Phy. Rev. Lett. **122**, 040403 (2019)], which is a stronger manifestation of nonlocality based on locally indistinguishability and finds more efficient applications in quantum information hiding. However, demonstrating the triviality of orthogonality-preserving local measurements (OPLMs) is not straightforward. In this paper, we present a sufficient and necessary condition for trivial OPLMs in *N*-partite systems under certain conditions. By using our proposed condition, we deduce the minimum size of set with the strongest nonlocality in system  $(\mathbb{C}^3)^{\otimes N}$ , where the genuinely entangled sets constructed in Ref. [Phys. Rev. A **109**, 022220 (2024)] achieve this value. As it is known that studying construction involving fewer states with strongest nonlocality contribute to reducing resource consumption in applications. Furthermore, we construct strongest nonlocal genuinely entangled sets in system  $(\mathbb{C}^d)^{\otimes N}$  ( $d \geq 4$ ), which have a smaller size than the existing strongest nonlocal genuinely unely entangled sets as *N* increases. Consequently, our results contribute to a better understanding of strongest nonlocality.

## I. INTRODUCTION

Locally indistinguishable states exhibit quantum nonlocality, which means that a set of orthogonal quantum states cannot be discriminated by local operations and classical communication (LOCC). This seminal concept, introduced by Bennett *et al.* [1], has been studied extensively [2–19]. This kind of state plays a crucial role in various quantum information processing tasks, such as accessing, transmitting, and storing quantum information. When information is encoded in locally indistinguishable states, it cannot be locally retrieved by spatially separated users who are restricted to using only LOCC, unless some (all) are together. As a consequence, these states have found practical applications in data hiding [20, 21] and quantum secret sharing [22, 23].

Local irreducibility, a stronger concept than locally indistinguishability, refers to a set of orthogonal states that cannot be eliminated one or more states by orthogonalitypreserving local measurements (OPLMs), and was proposed by Halder et al. [24]. It induces strong quantum nonlocality. A set of orthogonal states is strongly nonlocal if it is locally irreducible in every bipartition of the subsystems [24]. Naturally, information hidden in strongly nonlocal states enhances security since no information can be obtained even if N-1 users are collusive [25], indicating the higher practical value of such states. By definition, proving strong nonlocality bases on demonstrating local irreducibility. However, irreducibility is not easy to determine. At present, lots of efforts have been devoted to the construction of different types of strongly nonlocal sets [25-39]. The predominant approach employed by these papers [25-38] is to prove mathematically that the OPLMs can only be trivial, which is also a sufficient condition for local irreducibility. This is due to that a trivial measurement (a measurement  $\{E_x\}_x$  is called trivial if all the positive semidefinite operators are proportional to the identity operator) is orthogonality-preserving and gives us no information.

Wang et al. [33] defined that a set possesses the strongest nonlocality if only a trivial orthogonality-preserving positive operator-valued measure (POVM) can be performed for each bipartition of the subsystems. Therefore, the strongly nonlocal sets [25–37] constructed using the "trivial OPLMs" technique have the property of the strongest nonlocality. However, as the number and dimension of subsystems increase in multipartite systems, proving the triviality of OPLM becomes more intricate. Shi et al. [28] developed two lemmas, Block Zeros Lemma and Block Trivial Lemma, which contribute to simplifying the proof process. To further alleviate this difficulty, we firstly present several sufficient conditions for Block Trivial Lemma. Subsequently, under certain conditions, we present a sufficient and necessary condition for the triviality of OPLM in every bipartition in N-partite systems. By utilizing this result, we prove the minimum size of strongest nonlocal set in system  $(\mathbb{C}^3)^{\otimes N}$  under certain conditions, which is reached by the genuinely entangled sets constructed in Ref. [36].

Seeking fewer states that exhibit the strongest nonlocality contributes to resource conservation in quantum information processing tasks. Much efforts have been made on constructing smaller set for both orthogonal product states [26–31] and entangled states [25, 32–36]. Intuitively, entanglement gives rise to nonlocality as entangled states [40–48] exhibit Bell nonlocality [44] by violating Bell inequality. Bennett *et al.* [1] presented an indistinguishable orthogonal product basis (OPB) in Hilbert space  $\mathbb{C}^3 \otimes \mathbb{C}^3$  demonstrating that local indistinguishability is not directly related to entanglement. Subsequently, Walgate *et al.* [2] showed any pair of

<sup>\*</sup>Electronic address: gaoting@hebtu.edu.cn

<sup>&</sup>lt;sup>†</sup>Electronic address: flyan@hebtu.edu.cn

orthogonal states can always be perfectly distinguished locally, whether entangled or not. This indicates that entanglement is not a necessary condition for nonlocality.

Recently, a lower bound for the strongest nonlocal set on multipartite systems was proved by Li et al. [34]. In system  $\otimes_{i=1}^{N} \mathbb{C}^{d_i}$  (d<sub>i</sub> is the dimension of the *i*th subsystem), it has been proven that the size of the strongest nonlocal sets can not be smaller than  $\max_i \{\hat{d}_i + 1\}$ , where  $\hat{d}_i = (\prod_{i=1}^N d_i)/d_i$ . In tripartite systems, Zhen et al. [37] constructed the sets that reach the lower bound, one of which is product state and the rest is entangled state. So natural questions arise: What role do the entangled states play in the manifestation of the strongest nonlocality? How smaller can a strongest nonlocal entangled set could be in some given system? In the existing research on entangled states [25, 32-36], Li et al. [34] constructed the strongest nonlocal orthogonal genuinely entangled sets (OGESs) with size  $\prod_{i=1}^{N} d_i - \prod_{i=1}^{N} (d_i - 1) + 1$  in general multipartite systems  $\bigotimes_{i=1}^{N} \mathbb{C}^{d_i}$ . Then, in system  $(\mathbb{C}^3)^{\otimes N}$ , Hu et al. [36] presented a more less set of OGESs possessing the strongest nonlocality. Do the smaller set of OGESs exist in high-dimensional multipartite systems? In this paper, we generalize the set in Ref. [36] to system  $(\mathbb{C}^d)^{\otimes N}$  and demonstrate that the size in our construction is much smaller than that of Ref. [34] as N increases while keeping d fixed. It also provides an answer to the question given in Ref. [25], "How do we construct a strongly nonlocal orthogonal genuinely entangled set in  $(\mathbb{C}^d)^{\otimes N}$  for any  $d \geq 2$  and  $N \geq 5$ ?"

The paper is organized as follows. In Sec. II, some necessary notations and definitions are introduced. In Sec. III, we present a sufficient and necessary condition for the triviality of local orthogonality-preserving POVM under bipartition in *N*-partite systems. In this condition, we discuss the minimum size of set in system  $(\mathbb{C}^3)^{\otimes N}$ . In Sec. IV, we construct strongest nonlocal OGESs in system  $(\mathbb{C}^d)^{\otimes N}$  ( $d \ge 4, N \ge$ 3). Finally, we draw a conclusion in Sec. V.

#### **II. PRELIMINARIES**

We first introduce some definitions and lemmas needed in this paper. A POVM is a set of semidefinite operators  $\{E_m = M_m^{\dagger} M_m\}$  such that  $\sum_m E_m = \mathbb{I}$ , where  $\mathbb{I}$  is identity operation. Throughout this paper, we consider only pure state and POVM measurements, and do not normalize the states for simplicity.

Assume that  $E = (a_{i,j})_{i,j \in \mathbb{Z}_d}$  is the matrix representation of the operator  $E = M^{\dagger}M$  under the bases  $\mathcal{B} := \{|0\rangle, |1\rangle, \cdots, |d-1\rangle\}$ . Given two nonempty subsets S and  $\mathcal{T}$  of  $\mathcal{B}$ , where  $s = |\mathcal{S}|, t = |\mathcal{T}|$  are the cardinality of S and  $\mathcal{T}$ , respectively.  ${}_{\mathcal{S}}E_{\mathcal{T}} := \sum_{|i\rangle \in S} \sum_{|j\rangle \in \mathcal{T}} a_{i,j} |i\rangle\langle j|$  is a submatrix of E with row coordinates S and column coordinates

 $\mathcal{T}$ , specifically,  ${}_{\mathcal{S}}E_{\mathcal{S}}$  is denoted by  $E_{\mathcal{S}}$ . Ref. [28] gives two lemmas.

*Lemma* 1 (Block Zeros Lemma). Suppose that  $\{|\psi_i\rangle\}_{i=0}^{s-1}$ and  $\{|\phi_j\rangle\}_{j=0}^{t-1}$  are two orthogonal sets spanned by S and Trespectively, if  $\langle \psi_i | E | \phi_j \rangle = 0$  for any  $i \in \mathbb{Z}_s$ ,  $j \in \mathbb{Z}_t$ , then  $_{\mathcal{S}}E_{\mathcal{T}} = \mathbf{0} \text{ and } _{\mathcal{T}}E_{\mathcal{S}} = \mathbf{0}.$ 

Lemma 2 (Block Trivial Lemma). Let  $\{|\psi_j\rangle\}_{j=0}^{s-1}$  be an orthogonal set spanned by S, assume that  $\langle \psi_i | E | \psi_j \rangle = 0$  for any  $i \neq j \in \mathbb{Z}_s$ . If there exists a state  $|u_0\rangle$  in S such that  $\{|u_0\rangle\}_{ES \setminus \{|u_0\rangle\}} = 0$  and  $\langle u_0 | \psi_j \rangle \neq 0$  for any  $j \in \mathbb{Z}_s$ , then  $E_S \propto \mathbb{I}_S$ .

Lemma 3. (Shi et al. [28]) Let  $\{|\psi_j\rangle\}$  be a set of orthogonal states in multipartite system  $\bigotimes_{i=1}^{N} \mathbb{C}^{d_i}$ . For each  $i = 1, 2, \dots, N$ , define  $\bar{A}_i = \{A_1 A_2 \cdots A_N\} \setminus \{A_i\}$  is the joint party of all but the *i*th party. If any OPLM on  $\bar{A}_i$  is trivial, then the set  $\{|\psi_j\rangle\}$  possesses the strongest nonlocality.

Consider an N-partite quantum system  $\mathcal{H} = \mathcal{H}_{A_1} \otimes \mathcal{H}_{A_2} \otimes \cdots \otimes \mathcal{H}_{A_N}$  with dimensional  $d_k$  for the kth  $(1 \leq k \leq N)$  subsystem. The computational basis of the whole quantum system is  $\mathcal{B} = \{|i_1\rangle|i_2\rangle \cdots |i_N\rangle |i_k \in \mathbb{Z}_{d_k}\}$ . Let

$$\mathcal{B}_r := \{ |r_1^t\rangle | r_2^t\rangle \cdots | r_N^t\rangle | r_k^t \in \mathbb{Z}_{d_k} \}_{t \in \mathbb{Z}_{|\mathcal{B}_r|}}, \ r \in \mathcal{Q} \quad (1)$$

are the disjoint subsets of  $\mathcal{B}$ , where  $\mathcal{Q} = \{1, 2, \cdots, q\}$ . We define the set of orthogonal states spanned by  $\mathcal{B}_r$  as

$$\mathcal{S}_r := \left\{ |\Psi_{r,c}\rangle := \sum_{t \in \mathbb{Z}_{|\mathcal{B}_r|}} \omega_{|\mathcal{B}_r|}^{c \cdot t} |r_1^t\rangle |r_2^t\rangle \cdots |r_N^t\rangle \right\},$$
(2)

where  $\omega_{|\mathcal{B}_r|} = e^{\frac{2\pi\sqrt{-1}}{|\mathcal{B}_r|}}, c \in \mathbb{Z}_{|\mathcal{B}_r|}.$ 

The set of N-tuples that corresponding to the basis vectors in  $\mathcal{B}_r$  denotes as

$$\mathcal{G}_r^N = \{ (r_1^t, r_2^t, \cdots, r_N^t) \}_{t \in \mathbb{Z}_{|\mathcal{B}_r|}}.$$
(3)

We divide  $\mathbb{Z}_{|\mathcal{B}_r|}$  into  $m^r (0 < m^r \le |\mathcal{B}_r|)$  disjoint subsets  $R_{\tau} (\tau \in \mathbb{Z}_{m^r})$ , i.e.,  $\bigcup_{\tau \in \mathbb{Z}_{m^r}} R_{\tau} = \mathbb{Z}_{|\mathcal{B}_r|}$ , for any  $t \ne t^{'} \in R_{\tau}, r_1^t = r_1^{t'}$ . Let  $r_1^{R_{\tau}} = r_1^t (t \in R_{\tau})$ , the set of *N*-tuples in  $\mathcal{G}_r^N$  whose first component is same as  $r_1^{R_{\tau}}$  is represented as

$$\{r_1^{R_\tau}\} \times \{(r_2^t, \cdots, r_N^t)\}_{t \in R_\tau} = \{r_1^{R_\tau}\} \times \mathcal{G}_{[r, R_\tau]}^{N-1}, \quad (4)$$

where  $\mathcal{G}_{[r,R_{\tau}]}^{N-1} = \bigcup_{t \in R_{\tau}} \{ (r_2^t, \cdots, r_N^t) \}$ . Therefore

$$\mathcal{G}_{r}^{N} = \bigcup_{\tau \in \mathbb{Z}_{m}r} \left( \{ r_{1}^{R_{\tau}} \} \times \mathcal{G}_{[r,R_{\tau}]}^{N-1} \right).$$
<sup>(5)</sup>

Definition 1.  $\mathcal{I}_{[r,R_{\tau}]}^{N-1} = \bigcup_{v \in \mathcal{O}} \mathcal{G}_{[v,V_{\gamma}]}^{N-1}$   $(r \notin \mathcal{O} \subset \mathcal{Q}, \gamma \in \mathbb{Z}_{m^{v}})$  is called the block inclusion (BI) set of  $\mathcal{G}_{[r,R_{\tau}]}^{N-1}$ , if for any  $u \neq w \in \mathcal{O}$  there exist  $\alpha \in \mathbb{Z}_{m^{u}}, \beta \in \mathbb{Z}_{m^{w}}$  such that  $u_{1}^{U_{\alpha}} = w_{1}^{W_{\beta}} \neq r_{1}^{R_{\tau}}$  and  $\mathcal{G}_{[r,R_{\tau}]}^{N-1} \subset \mathcal{I}_{[r,R_{\tau}]}^{N-1}$ . Specifically,  $\mathcal{I}_{[r,R_{\tau}]}^{N-1}$  is called a tight block inclusion (TBI) set if there exists a subset  $\mathcal{G}_{[x,X_{\lambda}]}^{N-1} \subset \mathcal{I}_{[r,R_{\tau}]}^{N-1}$   $(x \in \mathcal{O}, \lambda \in \mathbb{Z}_{m^{x}})$  such that  $|\mathcal{G}_{[r,R_{\tau}]}^{N-1} \cap \mathcal{G}_{[x,X_{\lambda}]}^{N-1}| = 1$ , and  $\mathcal{G}_{[x,X_{\lambda}]}^{N-1}$  is called a tight subblock.

Definition 2. A family of sets  $\{\bigcup_{\tau} \mathcal{G}_{[r,R_{\tau}]}^{N-1}\}_{r \in \mathcal{Q}}$  is connected if it cannot be divided into two groups of sets  $\{\bigcup_{\gamma} \mathcal{G}_{[v,V_{\gamma}]}^{N-1}\}_{v \in \mathcal{O}} (\mathcal{O} \subseteq \mathcal{Q})$  and  $\{\bigcup_{\lambda} \mathcal{G}_{[x,X_{\lambda}]}^{N-1}\}_{x \in \mathcal{Q} \setminus \mathcal{O}}$  such

that

$$\left(\bigcup_{v\in\mathcal{O}}\bigcup_{\gamma\in\mathbb{Z}_{m^{v}}}\mathcal{G}^{N-1}_{[v,V_{\gamma}]}\right)\bigcap\left(\bigcup_{x\in\mathcal{Q}\setminus\mathcal{O}}\bigcup_{\lambda\in\mathbb{Z}_{m^{x}}}\mathcal{G}^{N-1}_{[x,X_{\lambda}]}\right)=\emptyset.$$
(6)

## **III. THE SUFFICIENT AND NECESSARY CONDITION** FOR THE TRIVIALITY OF **ORTHOGONALITY-PRESERVING LOCAL** MEASUREMENTS AND THE SMALLEST SIZE OF STATES UNDER THIS CONSTRAINTS

According to Lemma 3, to prove the strongest nonlocality, we need to show that the OPLM on  $\bar{A}_k$  =  $\{A_{k+1}\cdots A_N\cdots A_{k-1}\}$  can only be trivial. In this section, we first give a sufficient condition for Lemma 2, then we present a sufficient and necessary condition for the triviality of OPLM. Under this condition, we discuss the smallest size of set with strongest nonlocality in  $(\mathbb{C}^3)^{\otimes N}$   $(N \ge 3)$ .

Let  $\{\Pi\}$  be the orthogonality-preserving POVM on subsystem  $\overline{A}_k$ , and  $E = \mathbb{I} \otimes \Pi$ . For the states  $\{S_r\}$  in Eq. (2), we have

$$\langle \Psi_{r,c} | \mathbb{I} \otimes \Pi | \Psi_{r,c'} \rangle = 0, \quad \langle \Psi_{r,c} | \mathbb{I} \otimes \Pi | \Psi_{v,c''} \rangle = 0$$
 (7)

where  $c \neq c' \in \mathbb{Z}_{|\mathcal{B}_r|}, c'' \in \mathbb{Z}_{|\mathcal{B}_v|} \ (r \neq v \in \mathcal{Q}).$ 

*Theorem* 1. Consider the set  $\mathcal{B}_r$  (1), if one of the following conditions is true for the set given by Eq. (5), Block Trivial Lemma (Lemma 2) was satisfied, i.e.  $E_{\mathcal{B}_r} \propto \mathbb{I}_{\mathcal{B}_r}$ .

(i) There exists  $\tau_0 \in \mathbb{Z}_{m^r}$  such that  $|\mathcal{G}_{[r,R_{\tau_0}]}^{N-1}| = 1$ . (ii) There is a set  $\mathcal{G}_{[r,R_{\tau_1}]}^{N-1}$  ( $\tau_1 \in \mathbb{Z}_{m^r}$ ) has a TBI set.

(iii) There is a set  $\mathcal{G}_{[r,R_{\tau_2}]}^{N-\hat{1}}$   $(\tau_2 \in \mathbb{Z}_{m^r})$  which has a BI set  $\mathcal{I}^{N-1}_{[r,R_{\tau_2}]} = igcup_{v\in\mathcal{O}}\mathcal{G}^{N-1}_{[v,V_{\tau_2}]}(r\notin\mathcal{O}\subset\mathcal{Q}), ext{ for each } v\in\mathcal{O}, \mathcal{G}^N_v$ satisfies condition (i) or (ii).

N-*Proof.* Condition (i) : Consider the  $\begin{aligned} |\mathbf{r}\rangle &= |r_k^{R_{\tau_0}}, r_{k+1}^{t}, \cdots, r_N^{t}, \cdots, r_{k-1}^{t}\rangle \text{ belongs to } \mathcal{B}_r, \text{ for}\\ \text{any state } |\mathbf{r}'\rangle &= |r_k^{t'}, r_{k+1}^{t'}, \cdots, r_N^{t'}, \cdots, r_{k-1}^{t'}\rangle \in \mathcal{B}_r \setminus \{|\mathbf{r}\rangle\},\\ \text{there is } t' \notin R_{\tau_0} \text{ and } r_k^{t'} \neq r_k^{R_{\tau_0}}, \text{ thus} \end{aligned}$ 

$$\langle \boldsymbol{r}|E|\boldsymbol{r}'\rangle = \langle \boldsymbol{r}|\mathbb{I}\otimes\Pi|\boldsymbol{r}'\rangle = 0,$$
 (8)

we get  $\{|\mathbf{r}\rangle\} E_{\mathcal{B}_r \setminus \{|\mathbf{r}\rangle\}} = 0$ , since  $\langle \mathbf{r} | \Psi_{r,s} \rangle \neq 0$ , Lemma 2 is satisfied.

Condition (ii): For the TBI set  $\mathcal{I}_{[r,R_{\tau_1}]}^{N-1}$  $|r_k^{t'}, r_{k+1}^{t'}, \cdots, r_N^{t'}, \cdots, r_{k-1}^{t'}\rangle \in \mathcal{B}_r \setminus \{|\mathbf{r}\rangle\}, \text{ if } t' \notin R_{\tau_1}, \text{ i.e.,} \\ r_k^{t'} \neq r_k^{R_{\tau_1}}, \text{ we have } \langle \mathbf{r}|E|\mathbf{r}'\rangle = \langle \mathbf{r}|\mathbb{I} \otimes \Pi|\mathbf{r}'\rangle = 0. \text{ If } t' \in R_{\tau_1},$ 

we obtain that  $(r_{k+1}^{t'}, \cdots, r_N^{t'}, \cdots, r_{k-1}^{t'}) \in \mathcal{G}_{[r,R_{\tau_1}]}^{N-1}$ , since  $\mathcal{G}_{[x,X_{\lambda}]}^{N-1}$  is a tight sub-block, there must have  $u 
eq x \in \mathcal{O}$ such that  $(r_{k+1}^{t'}, \cdots, r_N^{t'}, \cdots, r_{k-1}^{t'}) \in \mathcal{G}_{[u, U_0]}^{N-1} \subset I_{[r, B_{\pi t}]}^{N-1}$ Because  $|\mathbf{x}\rangle = |x_k^{X_\lambda}, x_{k+1}, \cdots, x_N, \cdots, x_{k-1}\rangle \in \mathcal{B}_x$ ,  $|\mathbf{u}\rangle = |u_k^{U_\alpha}, r_{k+1}^{t'}, \cdots, r_N^{t'}, \cdots, r_{k-1}^{t'}\rangle \in \mathcal{B}_u$  and  $x_k^{X_\lambda} = u_k^{U_\alpha}$ , it follows from Eq. (7) that

$$\langle \mathbf{x} | E | \mathbf{u} \rangle$$
  
= $\langle x_{k+1}, \cdots, x_N, \cdots, x_{k-1} | \Pi | r_{k+1}^{t'}, \cdots, r_N^{t'}, \cdots, r_{k-1}^{t'} \rangle$   
=0.

Since  $r_k^{t'} = r_k^{R_{\tau_1}}$ , we have

$$\langle \boldsymbol{r} | E | \boldsymbol{r}' \rangle$$
  
= $\langle x_{k+1}, \cdots, x_N, \cdots, x_{k-1} | \Pi | r_{k+1}^{t'}, \cdots, r_N^{t'}, \cdots, r_{k-1}^{t'} \rangle$   
=0,

meanwhile,  $\langle \boldsymbol{r} | \Psi_{r,s} \rangle \neq 0$ , Lemma 2 is satisfied.

Condition (iii): For arbitrary two states  $|r\rangle$  $\begin{array}{l} |r_k^{R_{\tau_2}}, r_{k+1}^t, \cdots, r_N^t, \cdots, r_{k-1}^t\rangle \in \mathcal{B}_r \text{ and } |\mathbf{r}'\rangle = \\ |r_k^{t'}, r_{k+1}^{t'}, \cdots, r_N^{t'}, \cdots, r_{k-1}^{t'}\rangle \in \mathcal{B}_r \setminus \{|\mathbf{r}\rangle\}, \text{ if } t' \notin R_{\tau_2}, \\ \text{apparently, } \langle \mathbf{r}|E|\mathbf{r}'\rangle = 0. \quad \text{If } t' \in R_{\tau_2}, \text{ we get} \end{array}$  $(r_{k+1}^t, \cdots, r_k^t, \cdots, r_{k-1}^t) \neq (r_{k+1}^{t'}, \cdots, r_k^{t'}, \cdots, r_{k-1}^{t'}) \in \mathcal{G}_{[r, R_{\tau_2}]}^{N-1}.$  Since  $\mathcal{G}_{[r, R_{\tau_2}]}^{N-1}$  has a BI set  $\bigcup_{v \in \mathcal{O}} \mathcal{G}_{[v, V_{\gamma}]}^{N-1}$ , suppose  $\begin{array}{l} (r_{k+1}^t,\cdots,r_N^t,\cdots,r_{k-1}^t), \quad (r_{k+1}^{t'},\cdots,r_N^{t'},\cdots,r_{k-1}^{t'}) \in \\ \mathcal{G}_{[v,V_{\gamma}]}^{N-1} \ (v \in \mathcal{O}), \text{ as } \mathcal{G}_v^N \text{ satisfies condition (i) or (ii), we have} \end{array}$  $E_{\mathcal{B}_{v}} \propto \mathbb{I}_{\mathcal{B}_{v}}$ , i.e., for  $|\mathbf{v}\rangle = |v_{k}^{V_{\gamma}}, r_{k+1}^{t}, \cdots, r_{N}^{t}, \cdots, r_{k-1}^{t}\rangle$ and  $|\mathbf{v}'\rangle = |v_{k}^{V_{\gamma}}, r_{k+1}^{t'}, \cdots, r_{N}^{t'}, \cdots, r_{k-1}^{t'}\rangle$ , there is

$$\langle \boldsymbol{\nu} | E | \boldsymbol{\nu}' \rangle$$

$$= \langle r_{k+1}^t, \cdots, r_N^t, \cdots, r_{k-1}^t | \Pi | r_{k+1}^{t'}, \cdots, r_N^{t'}, \cdots, r_{k-1}^{t'} \rangle$$

$$= 0.$$

$$(9)$$

Thus

$$\langle \boldsymbol{r} | E | \boldsymbol{r}' \rangle$$
  
=  $\langle r_{k+1}^t, \cdots, r_N^t, \cdots, r_{k-1}^t | \Pi | r_{k+1}^{t'}, \cdots, r_N^{t'}, \cdots, r_{k-1}^{t'} \rangle$   
=0.

Suppose that  $(r_{k+1}^t, \cdots, r_N^t, \cdots, r_{k-1}^t) \in \mathcal{G}_{[v,V_{\gamma}]}^{N-1}$  and  $(r_{k+1}^t, \cdots, r_{k-1}^t) \in \mathcal{G}_{[v,V_{\gamma}]}^{N-1}$  $(r_{k+1}^{t'}, \cdots, r_{N}^{t'}, \cdots, r_{k-1}^{t'}) \in \mathcal{G}_{[w, W_{d}]}^{N-1}$ , where  $v \neq w \in \mathcal{O}$ , we have  $|\mathbf{v}\rangle = |v_k^{V_{\gamma}}, r_{k+1}^t, \cdots, r_N^t, \cdots, r_{k-1}^t\rangle \in \mathcal{B}_v, |\mathbf{w}\rangle = |w_k^{W_{\beta}}, r_{k+1}^{t'}, \cdots, r_N^{t'}, \cdots, r_{k-1}^t\rangle \in \mathcal{B}_w$  and  $v_k^{V_{\gamma}} = w_k^{W_{\beta}}$ , thus  $\langle \boldsymbol{r} | E | \boldsymbol{r}' \rangle$  $= \langle r_{k+1}^t, \cdots, r_N^t, \cdots, r_{k-1}^t | \Pi | r_{k+1}^{t'}, \cdots, r_N^{t'}, \cdots, r_{k-1}^{t'} \rangle$ (11)  $= \langle \mathbf{v} | \mathbb{I} \otimes \Pi | \mathbf{w} \rangle$ 

$$=0.$$

the last step is obtained by Eq. (7), so Lemma 2 is satisfied.■ Theorem 2. Consider the set  $S = \bigcup_{r \in Q} S_r$ , where  $S_r$  is given in Eq. (2). Suppose S is symmetric and  $\bigcup_{r \in Q} \bigcup_{\tau \in \mathbb{Z}_{m^r}} \mathcal{G}_{[r,\tau]}^{N-1} = \mathbb{Z}_{d_{k+1}} \times \cdots \times \mathbb{Z}_{d_N} \times \cdots \times \mathbb{Z}_{d_{k-1}}.$ When  $E_{\mathcal{B}_r}$  is proportional to the identity operator  $\mathbb{I}_r$ ,  $\Pi$  can only be trivial if and only if the set  $\bigcup_{r \in Q} \mathcal{G}_r^N$  (5) satisfy the following conditions:

(i) For any two different strings  $(i_{k+1}, \dots, i_N, \dots, i_{k-1}), (j_{k+1}, \dots, j_N, \dots, j_{k-1}) \in \mathbb{Z}_{d_{k+1}} \times \dots \times \mathbb{Z}_{d_N} \times \dots \times \mathbb{Z}_{d_{k-1}}$ , at least one element  $i_k$  in  $\mathbb{Z}_{d_k}$  such that  $(i_k, i_{k+1}, \dots, i_N, \dots, i_{k-1}), (i_k, j_{k+1}, \dots, j_N, \dots, j_{k-1}) \in \bigcup_{r \in \mathcal{Q}} \mathcal{G}_r^N$ .

(ii) The family of sets  $\{\bigcup_{\tau \in \mathbb{Z}_{m^r}} \mathcal{G}_{[r,R_{\tau}]}^{N-1}\}_{r \in \mathcal{Q}}$  are connected.

*Proof. Sufficient.* We first show that  $\Pi$  can only be trivial. Since  $E_{\mathcal{B}_r} \propto \mathbb{I}_{\mathcal{B}_r}$ , for any two different states  $|\mathbf{i}\rangle$ ,  $|\mathbf{i}'\rangle \in \mathcal{B}_r$ , we get  $\langle \mathbf{i}|E|\mathbf{i}'\rangle = \langle \mathbf{i}'|E|\mathbf{i}\rangle = 0$ . It follows from Lemma 1 that the states  $|\mathbf{i}\rangle \in \mathcal{B}_r$  and  $|\mathbf{j}\rangle \in \mathcal{B}_v$   $(r \neq v \in \mathcal{Q})$  satisfy  $\langle \mathbf{i}|E|\mathbf{j}\rangle = \langle \mathbf{j}|E|\mathbf{i}\rangle = 0$ .

So, for any two different states  $|i\rangle$ ,  $|j\rangle \in \bigcup_{r \in Q} \mathcal{B}_r$ , we have

$$\langle \boldsymbol{i}|E|\boldsymbol{j}\rangle = \langle \boldsymbol{j}|E|\boldsymbol{i}\rangle = 0.$$
 (12)

By condition (i), substituting  $|i\rangle$  $|i_k, i_{k+1}, \dots, i_N, \dots, i_{k-1}\rangle$  and  $|j\rangle$  $|i_k, j_{k+1}, \dots, j_N, \dots, j_{k-1}\rangle$  into Eq. (12), one gets

$$\langle i_k, i_{k+1}, \cdots, i_N, \cdots, i_{k-1} | E | i_k, j_{k+1}, \cdots, j_N, \cdots, j_{k-1} \rangle$$
  
= $\langle i_{k+1}, \cdots, i_N, \cdots, i_{k-1} | \Pi | j_{k+1}, \cdots, j_N, \cdots, j_{k-1} \rangle$   
=0,

thus the off-diagonal elements of  $\Pi$  are all zeros.

Furthermore, as  $E_{\mathcal{B}_r} \propto \mathbb{I}_{\mathcal{B}_r}$ , for two different vectors  $|\mathbf{i}\rangle = |i_k, i_{k+1}, \cdots, i_N, \cdots, i_{k-1}\rangle$  and  $|\mathbf{i}'\rangle = |i'_k, i'_{k+1}, \cdots, i'_N, \cdots, i'_{k-1}\rangle$  belong to  $\mathcal{B}_r$ , we have

$$\langle \boldsymbol{i}|E|\boldsymbol{i}\rangle = \langle \boldsymbol{i}'|E|\boldsymbol{i}'\rangle. \tag{14}$$

By condition (i), let  $i_k = i'_k$  in Eq. (14), we get

$$\langle i_{k+1}, \cdots, i_N, \cdots, i_{k-1} | \Pi | i_{k+1}, \cdots, i_N, \cdots, i_{k-1} \rangle = \langle i'_{k+1}, \cdots, i'_N, \cdots, i'_{k-1} | \Pi | i'_{k+1}, \cdots, i'_N, \cdots, i'_{k-1} \rangle$$
(15)

for any two different strings  $(i_{k+1}, \cdots, i_N, \cdots, i_{k-1})$ ,  $(i'_{k+1}, \cdots, i'_N, \cdots, i'_{k-1}) \in \bigcup_{\tau \in \mathbb{Z}_{m^r}} \mathcal{G}^{N-1}_{[r,R_{\tau}]}$ . Since  $\{\bigcup_{\tau \in \mathbb{Z}_{m^r}} \mathcal{G}^{N-1}_{[r,\tau]}\}_{r \in \mathcal{Q}}$  is connected, for any different (N - 1)-tuples  $(i_{k+1}, \cdots, i_N, \cdots, i_{k-1}) \neq (j_{k+1}, \cdots, j_N, \cdots, j_{k-1}) \in \{\bigcup_{\tau \in \mathbb{Z}_{m^r}} \mathcal{G}^{N-1}_{[r,\tau]}\}_{r \in \mathcal{Q}}$ , we have

$$\langle i_{k+1}, \cdots, i_N, \cdots, i_{k-1} | \Pi | i_{k+1}, \cdots, i_N, \cdots, i_{k-1} \rangle$$
  
= $\langle j_{k+1}, \cdots, j_N, \cdots, j_{k-1} | \Pi | j_{k+1}, \cdots, j_N, \cdots, j_{k-1} \rangle,$ (16)

thus the diagonal elements of  $\Pi$  are all equal. Accordingly, we get  $\Pi \propto \mathbb{I}.$ 

*Necessity.* By Eq. (2),  $S_r$  is an orthogonal set spanned by  $\mathcal{B}_r$ , Dim (span  $S_r$ ) =  $|\mathbb{Z}_{\mathcal{B}_r}|$ , then

$$\operatorname{span}\left\{|\Psi_{r,s}\right\rangle\right\}_{s\in\mathbb{Z}_{|\mathcal{B}_r|}} = \operatorname{span}\mathcal{B}_r.$$
(17)

Thus for  $|i\rangle \in \mathcal{B}_r$ , it has a linear combination of  $\{|\Psi_{r,s}\rangle\}_{s\in\mathbb{Z}_{|\mathcal{B}_r|}}$ , by Eq. (7), one obtains

$$\langle \boldsymbol{i} | \mathbb{I} \otimes \Pi | \boldsymbol{j} \rangle$$
  
= $\langle i_k | j_k \rangle \langle i_{k+1}, \cdots, i_{k-1} | \Pi | j_{k+1}, \cdots, j_{k-1} \rangle$  (18)  
=0

for any different vectors  $|\mathbf{i}\rangle = |i_k, i_{k+1}, \cdots, i_N, \cdots, i_{k-1}\rangle$ ,  $|j_k, j_{k+1}, \cdots, j_N, \cdots, j_{k-1}\rangle \in \bigcup_{r \in \mathcal{Q}} \mathcal{B}_r.$  $|\mathbf{j}\rangle =$ Since  $\Pi$  is proportional to the identity operator, when  $(i_{k+1}, \cdots, i_N, \cdots, i_{k-1}) = (j_{k+1}, \cdots, j_N, \cdots, j_{k-1})$ we get  $(i_{k+1}, \dots, i_{k-1} | \Pi | j_{k+1}, \dots, j_{k-1})$  $\neq$  0, then  $\langle i_k | j_k \rangle = 0$ , that is  $i_k \neq j_k$ . When  $(i_{k+1}, \dots, i_N, \dots, i_{k-1}) \neq (j_{k+1}, \dots, j_N, \dots, j_{k-1})$ there is  $\langle i_{k+1}, \cdots, i_{k-1} | \Pi | j_{k+1}, \cdots, j_{k-1} \rangle = 0$ , then  $\langle i_k | j_k \rangle$  equals 0 or 1. In this case, we will show that there must have  $\langle i_k | j_k \rangle = 1$  by contradiction. Assuming that  $\langle i_k | j_k \rangle$  is only equal to 0, we get  $i_k \neq j_k$ . By the symmetry of S, it follows that the vectors  $|i_{k+1}, \cdots, i_N, \cdots, i_{k-1}, i_k\rangle$ and  $|j_{k+1}, \cdots, j_N, \cdots, j_{k-1}, j_k\rangle$  belong to set  $\bigcup_{r \in \mathcal{Q}} \mathcal{B}_r$ . Substituting these two vectors to Eq. (18), we get  $i_{k+1} \neq j_{k+1}$  by the assumption. By the same token, one can deduce that

$$\begin{cases} i_{k+2} \neq j_{k+2}, \\ \vdots \\ i_N \neq j_N, \\ \vdots \\ i_{k-1} \neq j_{k-1}. \end{cases}$$

It follows that  $\bigcup_{r \in \mathcal{Q}} \bigcup_{\tau \in \mathbb{Z}_{m^r}} \mathcal{G}_{[r,\tau]}^{N-1} \neq \mathbb{Z}_{d_{k+1}} \times \cdots \times \mathbb{Z}_{d_N} \times \cdots \times \mathbb{Z}_{d_{k-1}}$ , this leads to a contradiction.

On the other hand, as  $E_{\mathcal{B}_r} \propto \mathbb{I}_{\mathcal{B}_r}$ , for any different vectors  $|\mathbf{i}\rangle = |i_k, i_{k+1}, \cdots, i_N, \cdots, i_{k-1}\rangle$ ,  $|\mathbf{i}'\rangle = |i'_k, i'_{k+1}, \cdots, i'_N, \cdots, i'_{k-1}\rangle \in \mathcal{B}_r$ , we have

$$\langle \boldsymbol{i}|E|\boldsymbol{i}\rangle = \langle \boldsymbol{i}'|E|\boldsymbol{i}'\rangle,$$
 (19)

then

=

\_

(13)

Since the diagonal elements of  $\Pi$  are all equal, one can deduce that  $\{\bigcup_{\tau \in \mathbb{Z}_{m^r}} \mathcal{G}_{[r,R_\tau]}^{N-1}\}_{r \in \mathcal{Q}}$  is connected.

Theorem 2 provides a sufficient and necessary condition for  $S = \bigcup_{r \in Q} S_r$  (2) which has the property of strongest nonlocality. Using the condition (i) of Theorem 2, in  $(\mathbb{C}^3)^{\otimes N}$ , we deduce the smallest size of S, what we need to do is to discuss the smallest size of  $\bigcup_{r \in Q} \mathcal{G}_r^N$ . Let  $\mathcal{C}(\mathbb{Z}_3^{N-1})$  be the set that satisfies the condition (i) of Theorem 2,  $\mathcal{SC}(\mathbb{Z}_3^{N-1})$  be all cyclic symmetric elements of  $\mathcal{C}(\mathbb{Z}_3^{N-1})$ , obviously,  $\mathcal{C}(\mathbb{Z}_3^{N-1}) \subset \mathcal{SC}(\mathbb{Z}_3^{N-1})$ . Theorem 3. For the set  $S = \bigcup_{r \in Q} \mathcal{S}_r$  (2) in

Theorem 3. For the set  $S = \bigcup_{r \in Q} S_r$  (2) in  $(\mathbb{C}^3)^{\otimes N}$   $(N \geq 3)$ , if S is symmetric and any orthogonalitypreserving POVM on  $\overline{A}_k$  can only be trivial, then S has the strongest nonlocality, the smallest size of this set is  $2 \times 3^{N-1}$ .

*Proof.* If set  $\bigcup_{r \in \mathcal{Q}} \mathcal{G}_r^N$  contains  $\mathcal{C}(\mathbb{Z}_3^{N-1})$ , due to the symmetry, it must contain  $\mathcal{SC}(\mathbb{Z}_3^{N-1})$ . In addition to the case that  $\mathcal{C}(\mathbb{Z}_3^{N-1}) = \mathbb{Z}_3^N$ , four cases of  $\mathcal{C}(\mathbb{Z}_3^{N-1})$  are as follows. Case I: For every string  $(i_2, \dots, i_N)$  in  $\mathbb{Z}_3^{N-1}$ , if only one element  $i_1 \in \mathbb{Z}_3$  such that  $(i_1, i_2, \dots, i_N) \in \mathcal{C}(\mathbb{Z}_3^{N-1})$ , we get  $\mathcal{C}(\mathbb{Z}_3^{N-1}) = \{i_1\} \times \mathbb{Z}_3^{N-1}$  and  $|\mathcal{C}(\mathbb{Z}_3^{N-1})| = 3^{N-1}$ . One can check that  $\mathcal{SC}(\mathbb{Z}_3^{N-1})$  is the collection of N-tuples that at least one component is  $i_n$ , thus  $\mathcal{SC}(\mathbb{Z}^{N-1})$  has size that at least one component is  $i_1$ , thus  $\mathcal{SC}(\mathbb{Z}_3^{N-1})$  has size  $3^N - 2^N$ . Therefore, the size of  $\bigcup_{r \in Q} \mathcal{G}_r^N$  is not less than  $3^N - 2^N$ .

Case II: For every string  $(i_2, \dots, i_N)$  in  $\mathbb{Z}_3^{N-1}$ , there are exactly two different elements  $i_1, i'_1 \in \mathbb{Z}_3$  such that are exactly two different elements  $i_1, i_1 \in \mathbb{Z}_3$  such that  $(i_1, i_2, \dots, i_N), (i'_1, i_2, \dots, i_N) \in \mathcal{C}(\mathbb{Z}_3^{N-1})$ . We get  $\mathcal{C}(\mathbb{Z}_3^{N-1}) = \{i_1, i'_1\} \times \mathbb{Z}_3^{N-1}$  and has size  $2 \times 3^{N-1}$ . Since  $\mathcal{C}(\mathbb{Z}_3^{N-1}) \subset \mathcal{SC}(\mathbb{Z}_3^{N-1})$ , then the size of  $\mathcal{SC}(\mathbb{Z}_3^{N-1})$  is not less than  $2 \times 3^{N-1}$ , thus the size of  $\bigcup_{r \in \mathcal{Q}} \mathcal{G}_r^N$  cannot be less than  $2 \times 3^{N-1}$ .

Let  $\mathbb{P}^{N-1}$  is a subset of  $\mathbb{Z}_3^{N-1}$  and  $0 < |\mathbb{P}^{N-1}| < 3^{N-1}$ . Case III: For every string  $(p_2, \cdots, p_N)$  in  $\mathbb{P}^{N-1}$ , there is only one element  $p_1 \in \mathbb{Z}_3$  such that  $(p_1, p_2, \cdots, p_N) \in$  $\mathcal{C}(\mathbb{Z}_3^{N-1}).$  In addition, for every string  $(q_2,\cdots,q_N)$  in  $\mathbb{C}(\mathbb{Z}_3^{N-1} \setminus \mathbb{P}^{N-1})$ , there is a subset  $\mathbb{Q}$  of  $\mathbb{Z}_3$  ( $|\mathbb{Q}| > 1$ ) such that  $\mathbb{Q} \times (q_2, \cdots, q_N) \subset \mathcal{C}(\mathbb{Z}_3^{N-1})$ . By condition (i) of Theorem 2, we get  $p_1 \in \mathbb{Q}$ , thus  $\{p_1\} \times \mathbb{Z}_3^{N-1} \subset \mathcal{C}(\mathbb{Z}_3^{N-1})$ . As stated in Case I, the size of  $\bigcup_{r \in \mathcal{Q}} \mathcal{G}_r^N$  is not less than  $3^N - 2^N$ .

Case IV: For every string  $(p_2, \dots, p_N)$  in  $\mathbb{P}^{N-1}$ , there are two different elements  $p_1, p'_1 \in \mathbb{Z}_3$  such that  $(p_1, p_2, \cdots, p_N), (p'_1, p_2, \cdots, p_N) \in \mathcal{C}(\mathbb{Z}_3^{N-1}).$  For the string  $(q_2, \cdots, q_N) \in \mathbb{Z}_3^{N-1} \setminus \mathbb{P}^{N-1}$  and the subset  $\mathbb{Q}$  of  $\mathbb{Z}_3$  ( $|\mathbb{Q}| > 1$ ), there is  $\mathbb{Q} \times (q_2, \dots, q_N) \subset \mathcal{C}(\mathbb{Z}_3^{N-1})$ . As  $|\mathbb{Q}| = 2$  is more likely than  $|\mathbb{Q}| = 3$  to get the smallest size of  $\bigcup_{r \in \mathcal{Q}} \mathcal{G}_r^N$ , we obtain  $|\mathcal{C}(\mathbb{Z}_3^{N-1})| = 2 \times 3^{N-1}$ , thus the size of  $\mathcal{CS}(\mathbb{Z}_3^{N-1})$  is not less than  $2 \times 3^{N-1}$ , then the size of  $\bigcup_{r \in \mathcal{Q}} \mathcal{G}_r^N$  cannot be less than  $2 \times 3^{N-1}$ .

Actually, in system  $(\mathbb{C}^3)^{\otimes N}$ , the set in Ref. [34] corresponds to Case I and has size  $3^N - 2^N$ . Obviously,  $2 \times 3^{N-1}$ is fewer to  $3^N - 2^N$ , meanwhile, the strongest nonlocal genuinely entangled set in Ref. [36] satisfies Theorem 2 and contains  $2 \times 3^{N-1}$  states. Consequently, in  $(\mathbb{C}^3)^{\otimes N}$ , the minimum size of the set  $\mathcal{S} = \bigcup_{r \in Q} \mathcal{S}_r$  is  $2 \times 3^{N-1}$ .

## IV. ORTHOGONAL GENUINELY ENTANGLED SETS WITH STRONGEST NONLOCALITY IN $(\mathbb{C}^d)^{\otimes N}$

In this section, we generalize the strongest nonlocal genuinely entangled set in Ref. [36] to  $(\mathbb{C}^d)^{\otimes N}$   $(d \ge 4)$  and show that there exist smaller strongest nonlocal OGESs in high-dimensional multipartite systems.

## A. An ORTHOGONAL GENUINELY ENTANGLED BASE in $(\mathbb{C}^d)^{\otimes N}$

For each element  $i \in \mathbb{Z}_d$ , let

$$\mathcal{G}_i^1 := \{i\},\tag{21}$$

then we construct d subsets of  $\mathbb{Z}_d^N$  for  $N \ge 2$ ,

$$\mathcal{G}_i^N := \bigcup_{j=0}^{d-1} \left( \{ i \oplus_d (d-j) \} \times \mathcal{G}_j^{N-1} \right), \qquad (22)$$

where  $i \in \mathbb{Z}_d$ ,  $i \oplus_d t = (i + t) \mod d$ .

Proposition 1. The sets in Eq. (22) are pairwise disjoint and the union of all is  $\mathbb{Z}_d^N$ , i.e.,

$$\bigcup_{i=0}^{d-1} \mathcal{G}_i^N = \mathbb{Z}_d^N \text{ and } \mathcal{G}_i^N \cap \mathcal{G}_j^N = \emptyset, \text{ where } i \neq j \in \mathbb{Z}_d.$$

The detailed proof is shown in Appendix A.

*Proposition* 2. The set  $\mathcal{G}_i^N$  in Eq. (22) is invariant under arbitrary permutation of the positions of the N components.

The detailed proof is shown in Appendix B.

Let  $\mathcal{H} := (\mathbb{C}^d)^{\otimes N}$ ,

$$\mathcal{S}_{i} := \{ |\Psi_{i,k}\rangle \in \mathcal{H} \mid k \in \mathbb{Z}_{s_{i}}, |\Psi_{i,k}\rangle := \sum_{\boldsymbol{j} \in \mathcal{G}_{i}^{N}} \omega_{s_{i}}^{kf_{i}(\boldsymbol{j})} |\boldsymbol{j}\rangle \}.$$
(23)

Here  $f_i : \mathcal{G}_i^N \longrightarrow \mathbb{Z}_{s_i}$  is any fixed bijection and  $\omega_{s_i} :=$  $e^{\frac{2\pi\sqrt{-1}}{s_i}}, s_i = |\mathcal{G}_i^N| \ (i \in \mathbb{Z}_d).$ 

Proposition 3. The set  $\bigcup_{i=0}^{d-1} S_i$  of states given by Eq. (23) is an orthogonal genuinely entangled base in  $(\mathbb{C}^d)^{\otimes N}$ .

*Proof.* By Proposition 1, it follows that  $\bigcup_{i=0}^{d-1} S_i$  is a base in system  $(\mathbb{C}^d)^{\otimes N}$ . The proof of genuine entanglement of  $|\Psi_{i,k}\rangle$  is similar to Ref. [[36], Theorem 1].

# **B.** STRONGEST NONLOCAL OGESs in $(\mathbb{C}^d)^{\otimes N}$ $(d \ge 4)$

Based on the above result, we present the strongest nonlocal genuinely entangled sets in system  $(\mathbb{C}^d)^{\otimes N}$ . We first consider which sets are sufficient to satisfy the condition (i) of Theorem 2. From Eq. (A4), the relationship be-tween  $\{\mathcal{G}_i^N\}_{i\in\mathbb{Z}_d}$  and  $\{\mathcal{G}_j^{N-1}\}_{j\in\mathbb{Z}_d}$  is described by  $\mathcal{M}_d =$  $\{m_{i,j}\}_{i,j\in\mathbb{Z}_d}$  (B6). For simplicity, we label row *i* by set  $\mathcal{G}_i^N$ and column j by set  $\mathcal{G}_i^{N-1}$ .

$$\mathcal{G}_{0}^{N-1} \quad \mathcal{G}_{1}^{N-1} \quad \mathcal{G}_{2}^{N-1} \quad \cdots \quad \mathcal{G}_{d-2}^{N-1} \quad \mathcal{G}_{d-1}^{N-1}$$

$$\mathcal{G}_{0}^{N} \begin{bmatrix} 0 & d-1 & d-2 & \cdots & 2 & 1 \\ 1 & 0 & d-1 & \cdots & 3 & 2 \\ 2 & 1 & 0 & \cdots & 4 & 3 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \mathcal{G}_{d-2}^{N} \\ \mathcal{G}_{d-1}^{N} \end{bmatrix} \begin{bmatrix} d-2 & d-3 & d-4 & \cdots & 0 & d-1 \\ d-1 & d-2 & d-3 & \cdots & 1 & 0 \end{bmatrix}$$

The distance of any two rows is defined as

$$r\left(\mathcal{G}_{i_1}^N, \mathcal{G}_{i_2}^N\right) = \min\{|i_1 - i_2|, \ d - |i_1 - i_2|\},\$$

where  $i_1 \neq i_2 \in \mathbb{Z}_d$ , then one gets

$$r\left(\mathcal{G}_{i_{1}}^{N},\mathcal{G}_{i_{2}}^{N}
ight) \in \left[1,\left\lfloor \frac{d}{2} 
ight],$$

with |.| denoting the floor function. The distance of any two columns is denoted as c  $(\mathcal{G}_{j_1}^{N-1},\mathcal{G}_{j_2}^{N-1})$  and has the same definition as  $r(\mathcal{G}_{i_1}^N, \mathcal{G}_{i_2}^N)$ . Then we get the following observations.

Observation 1. For any  $2 \times 2$  submatrix  $P_{2 \times 2}$  of  $\mathcal{M}_d$ ,

$$P_{2\times 2} = \begin{array}{cc} \mathcal{G}_{j_1}^{N-1} & \mathcal{G}_{j_2}^{N-1} \\ \mathcal{G}_{i_1}^{N} \begin{bmatrix} m_{i_1,j_1} & m_{i_1,j_2} \\ m_{i_2,j_1} & m_{i_2,j_2} \end{bmatrix}, \end{array}$$

if  $r(\mathcal{G}_{i_1}^N, \mathcal{G}_{i_2}^N) = c(\mathcal{G}_{j_1}^{N-1}, \mathcal{G}_{j_2}^{N-1})$  we get  $m_{i_1,j_1} = m_{i_2,j_2}$ or  $m_{i_1,j_2} = m_{i_2,j_1}$ , i.e., there are the same elements in both columns.

Observation 2. For any two different columns, since the union of all distances is  $\bigcup_{j_1 \neq j_2 \in \mathbb{Z}_d} \{ c \ (\mathcal{G}_{j_1}^{N-1}, \mathcal{G}_{j_2}^{N-1}) \} =$  $\begin{bmatrix} 1, \lfloor \frac{d}{2} \rfloor \end{bmatrix}, \text{ if we can find rows } \{\mathcal{G}_i^N\}_{i \in T \subset \mathbb{Z}_d} \text{ (c } (\mathcal{G}_{j_1}, \mathcal{G}_{j_2})\} = \begin{bmatrix} 1, \lfloor \frac{d}{2} \rfloor \end{bmatrix}, \text{ if owe can find rows } \{\mathcal{G}_i^N\}_{i \in T \subset \mathbb{Z}_d} \text{ such that } \bigcup_{i_1 \neq i_2 \in T} \{r(\mathcal{G}_{i_1}^N, \mathcal{G}_{i_2}^N)\} = \begin{bmatrix} 1, \lfloor \frac{d}{2} \rfloor \end{bmatrix}, \text{ it follows from Observation 1 that condition (i) of Theorem 2 is satisfied.}$ Proposition 4. The rows  $\{\mathcal{G}_t^N\}_{t \in T_1 \cup T_2}, \text{ where } T_1 = \{t_1 \mid 0 < t_1 < \lfloor \frac{d}{2} \rfloor, t_1 \text{ is odd}\} \text{ and } T_2 = \{0, \lfloor \frac{d}{2} \rfloor\}, \text{ satisfy the condition (i) of Theorem 2.}$ 

*Proof.* When  $\lfloor \frac{d}{2} \rfloor$  is odd, we have

$$\begin{cases} \bigcup_{t \in T_1} \{r(\mathcal{G}_0^N, \mathcal{G}_t^N)\} = \{1, 3, \cdots, \lfloor \frac{d}{2} \rfloor - 2\}, \\ \bigcup_{t \in T_1 \setminus \{1\}} \{r(\mathcal{G}_1^N, \mathcal{G}_t^N)\} = \{2, 4, \cdots, \lfloor \frac{d}{2} \rfloor - 3\}, \\ r(\mathcal{G}_1^N, \mathcal{G}_{\lfloor \frac{d}{2} \rfloor}^N) = \lfloor \frac{d}{2} \rfloor - 1, \\ r(\mathcal{G}_0^N, \mathcal{G}_{\lfloor \frac{d}{2} \rfloor}^N) = \lfloor \frac{d}{2} \rfloor. \end{cases}$$

When  $\lfloor \frac{d}{2} \rfloor$  is even, we have

$$\begin{cases} \bigcup_{t \in T_1} \{r(\mathcal{G}_0^N, \mathcal{G}_t^N)\} = \{1, 3, \cdots, \lfloor \frac{d}{2} \rfloor - 1\}, \\ \bigcup_{t \in T_1 \setminus \{1\}} \{r(\mathcal{G}_1^N, \mathcal{G}_t^N)\} = \{2, 4, \cdots, \lfloor \frac{d}{2} \rfloor - 2\}, \\ r(\mathcal{G}_0^N, \mathcal{G}_{\lfloor \frac{d}{2} \rfloor}^N) = \lfloor \frac{d}{2} \rfloor. \end{cases}$$

### By Observation 2, the condition (i) of Theorem 2 holds.

To complete the construction, we need to get the distribution of element  $(\xi)^{\times N}$  ( $\xi \in \mathbb{Z}_d$ ) in set  $\mathcal{G}_i^N$ , we first recall several knowledge of cyclic group.

Cyclic group. A group G is said to be cyclic if each element can be generated by  $q \in G$ , the element q is called a generator of G and denoted  $G = \langle g \rangle$ .

Lemma 4. Let  $G = \langle g \rangle$  is a cyclic group of order n, then the order of element  $g^s$  is  $\frac{n}{\gcd(s,n)}$ , where  $\gcd(s,n)$  is the greatest common divisor of s and n.

Theorem 4. Every set in  $\{\mathcal{G}_t^N\}_{t \in T_1 \cup T_2}$  contains *N*-tuples of the form  $(\xi')^{\times N}$   $(\xi' \in \mathbb{Z}_d \setminus \{0\})$ .

Proof. By Eq. (A3), we first obtain the distribution of element  $(\xi)^{\times N}$   $(\xi \in \mathbb{Z}_d)$  in  $\mathcal{G}_i^N$   $(j \in \mathbb{Z}_d)$ . Here we classify N by congruence of modulo d, let  $a \equiv N \pmod{d}$ , denoted as

TABLE I: The distribution of  $(\xi)^{\times N}$ , where  $\xi \in \mathbb{Z}_d$ .

System N	$(0)^{\times N}$	$(1)^{\times N}$	$(2)^{\times N}$	••• (6	$(d-2)^{\times N}$	$(d-1)^{\times N}$
[N] = [0]	$\mathcal{G}_0^N$	$\mathcal{G}_0^N$	$\mathcal{G}_0^N$	•••	$\mathcal{G}_0^N$	$\mathcal{G}_0^N$
[N] = [1]	$\mathcal{G}_0^N$	$\mathcal{G}_1^N$	$\mathcal{G}_2^N$	• • •	$\mathcal{G}_{d-2}^N$	$\mathcal{G}_{d-1}^N$
[N] = [2]	$\mathcal{G}_0^N$	$\mathcal{G}_2^N$	• • •			
÷						
[N] = [d-2]	$\mathcal{G}_0^N$	$\mathcal{G}_{d-2}^N$	•••			
[N] = [d-1]	$\mathcal{G}_0^N$	$\mathcal{G}_{d-1}^N$	$\mathcal{G}_{d-2}^N$		$\mathcal{G}_2^N$	$\mathcal{G}_1^N$

 $[N] = [a], a \in \mathbb{Z}_d$ . Then we find the relationship of component  $\xi$ , classification a and the subscript j of the set  $\mathcal{G}_i^N$ is

$$j = a \oplus_d a \oplus_d \dots \oplus_d a, \tag{24}$$

where a is repeated  $\xi$  times. The detailed distribution is shown in Table I. To complete the proof we need to show that each row in Table I from column  $(1)^{\times N}$  to column  $(d-1)^{\times N}$  contains the set  $\mathcal{G}_{j_0}^N$ , and the subscript  $j_0$  belongs to set  $T_1 \cup T_2$ .

From Eq. (24), we get that  $\{[j]\}_j = \langle [a] \rangle$  is a cyclic subgroup of group  $\{[0], [1], \dots, [d-1]\}$ . Using Lemma 4, we obtain the order of  $\langle [a] \rangle$  is  $k = \frac{d}{\gcd(s,d)}$ , where  $[a] = s[1] \ (0 \le s \le d-1)$ . We will discuss rows [N] = [a]in the following three cases.

When a = 0, we get j = 0 by Eq. (24), it follows that the sets in row [N] = [0] are all the same and are the  $\mathcal{G}_0^N$ .

When a and d are coprime, the order of  $\langle [a] \rangle$  is d, we get  $\{[j]\}_j = \{[0], [1], \cdots, [d-1]\}, \text{ which has an element } \lfloor \frac{d}{2} \rfloor$ belongs to  $T_2$ .

When a is a factor of d and 1 < a < d, we get the order k of  $\langle [a] \rangle$  satisfies k | d and 1 < k < d. Because there are d columns through column  $(0)^{\times N}$  to column  $(d-1)^{\times N}$ , the subscript  $0 \in T_2$  will appear at least twice. The proof is completed.

Now we give the construction of genuinely entangled states in  $(\mathbb{C}^d)^{\times N}$ , where N is divided into three cases as mentioned above.

Construction. Case I, [N] = [0]. Let

$$\widetilde{\mathcal{G}_{0}^{N}} = \mathcal{G}_{0}^{N} \setminus \{(0)^{\times N}, (\xi')^{\times N}\}, 
\{\widetilde{\mathcal{G}_{t}^{N}}\}_{t \in T_{1}} = \{\mathcal{G}_{t}^{N}\}_{t \in T_{1}}, 
\widetilde{\mathcal{G}_{\lfloor \frac{d}{2} \rfloor}^{N}} = \mathcal{G}_{\lfloor \frac{d}{2} \rfloor}^{N}, 
\widetilde{\mathcal{G}_{\lfloor \frac{d}{2} \rfloor+1}^{N}} = \{0\} \times \{(0)^{\times (N-1)}\} \bigcup \{\xi'\} \times \{(\xi')^{\times (N-1)}\},$$
(25)

where  $\xi' \in \mathbb{Z}_d \setminus \{0\}$  and satisfies  $[0] = \xi'[0]$ .

Case II, [N] = [a], where gcd(a, d) = 1. Let

$$\widetilde{\mathcal{G}_{0}^{N}} = \mathcal{G}_{0}^{N} \setminus \{(0)^{\times N}\}, 
\{\widetilde{\mathcal{G}_{t}^{N}}\}_{t \in T_{1}} = \{\mathcal{G}_{t}^{N}\}_{t \in T_{1}}, 
\widetilde{\mathcal{G}_{\lfloor \frac{d}{2} \rfloor}^{N}} = \mathcal{G}_{\lfloor \frac{d}{2} \rfloor}^{N} \setminus \{(\xi')^{\times N}\}, 
\widetilde{\mathcal{G}_{\lfloor \frac{d}{2} \rfloor+1}^{N}} = \{0\} \times \{(0)^{\times (N-1)}\} \bigcup \{\xi'\} \times \{(\xi')^{\times (N-1)}\}, 
(26)$$

where  $\xi' \in \mathbb{Z}_d \setminus \{0\}$  and satisfies  $\lfloor \lfloor \frac{d}{2} \rfloor = \xi'[a]$ . Case III, [N] = [a], where *a* divides *d*,  $a \neq 1, d$ . Let

$$\widetilde{\mathcal{G}_{0}^{N}} = \mathcal{G}_{0}^{N} \setminus \{(0)^{\times N}, (\xi')^{\times N}\}, \\ \{\widetilde{\mathcal{G}_{t}^{N}}\}_{t \in T_{1}} = \{\mathcal{G}_{t}^{N}\}_{t \in T_{1}}, \\ \widetilde{\mathcal{G}_{\lfloor \frac{d}{2} \rfloor}^{N}} = \mathcal{G}_{\lfloor \frac{d}{2} \rfloor}^{N}, \\ \widetilde{\mathcal{G}_{\lfloor \frac{d}{2} \rfloor+1}^{N}} = \{0\} \times \{(0)^{\times (N-1)}\} \bigcup \{\xi'\} \times \{(\xi')^{\times (N-1)}\},$$

$$(27)$$

where  $\xi' \in \mathbb{Z}_d \setminus \{0\}$  and satisfies  $[0] = \xi'[a]$ .

In each case, let  $T_3 = T_1 \cup T_2 \cup \{\lfloor \frac{d}{2} \rfloor + 1\}$ ,  $\widetilde{s_i}$  be the cardinality of the set  $\widetilde{\mathcal{G}_i^N}$ ,  $i \in T_3$ , denote

$$\widetilde{\mathcal{S}}_{i} := \{ |\widetilde{\Psi}_{i,k}\rangle \in \mathcal{H} \mid k \in \mathbb{Z}_{\widetilde{s}_{i}}, |\widetilde{\Psi}_{i,k}\rangle := \sum_{\boldsymbol{j} \in \widetilde{\mathcal{G}}_{i}^{\widetilde{N}}} \omega_{\widetilde{s}_{i}}^{kf_{i}(\boldsymbol{j})} |\boldsymbol{j}\rangle \}.$$
(28)

(28) Here  $f_i : \widetilde{\mathcal{G}_i^N} \longrightarrow \mathbb{Z}_{\widetilde{s_i}}$  is any fixed bijection and  $\omega_{\widetilde{s_i}} := e^{\frac{2\pi\sqrt{-1}}{s_i}}$ .

Even if we have moved  $(0)^{\times N}$  and  $(\xi')^{\times N}$ , each set in  $\{\widetilde{\mathcal{G}_t^N}\}_{t\in T_3}$  is still permutation invariance. Moreover, it follows from the proof of Ref. [[36], Theorem 1] that the states in Eq. (28) are still genuinely entangled.

*Theorem* 5. In  $(\mathbb{C}^d)^{\times N}$ , the states  $\widetilde{S} = \bigcup_{i \in T_3} \widetilde{S}_i$  given by Eq. (28) possess the strongest nonlocality, and

$$|\widetilde{\mathcal{S}}| = \begin{cases} (\frac{\lfloor \frac{d}{2} \rfloor + 1}{2} + 1) \times d^{(N-1)}, \lfloor \frac{d}{2} \rfloor \text{ is odd,} \\ (\frac{\lfloor \frac{d}{2} \rfloor}{2} + 2) \times d^{(N-1)}, \lfloor \frac{d}{2} \rfloor \text{ is even.} \end{cases}$$

*Proof.* To prove the strongest nonlocality, we need to show that the *N*-tuples in Eqs. (25-27) satisfy Theorem 1 and Theorem 2. Here we give a detailed proof for Case I, the remainder of the argument is analogous to Case I.

For Theorem 1, we first consider set  $\widetilde{\mathcal{G}_{\lfloor \frac{d}{2} \rfloor + 1}^{N}}$ , since  $| \{(0)^{\times (N-1)}\} |= 1$ , one gets that  $\widetilde{\mathcal{G}_{\lfloor \frac{d}{2} \rfloor + 1}^{N}}$  satisfies the condition (i). Then for any  $t \in T_1$ ,  $\widetilde{\mathcal{G}_t^N}$  has a subset  $\{t\} \times \mathcal{G}_0^{N-1}$  by Eq. (A3), we obtain  $\mathcal{G}_0^{N-1} = (\mathcal{G}_0^{N-1} \setminus \{(0)^{\times (N-1)}\}) \bigcup \{(0)^{\times (N-1)}\}$  and  $|\mathcal{G}_0^{N-1} \cap \{(0)^{\times (N-1)}\}| = 1$ , where the first term  $\mathcal{G}_0^{N-1} \setminus \{(0)^{\times (N-1)}\}|$  is taken from the subset  $\{0\} \times (\mathcal{G}_0^{N-1} \setminus \{(0)^{\times (N-1)}\})$  of  $\widetilde{\mathcal{G}_0^N}$  and the second term  $\{(0)^{\times (N-1)}\}$  is taken from the subset  $\{0\} \times \{(0)^{\times (N-1)}\}$  of  $\widetilde{\mathcal{G}_{\lfloor \frac{d}{2} \rfloor + 1}^N}$ . Thus  $\widetilde{\mathcal{G}_t^N}$  satisfy the condition (ii).  $\widetilde{\mathcal{G}_{\lfloor \frac{d}{2} \rfloor}^N}$  satisfies

condition (ii) for similar reason to  $\mathcal{G}_t^N$ .

For the set  $\widetilde{\mathcal{G}_0^N}$ ,  $\{\{0\} \times \mathcal{G}_0^{N-1}\} \setminus \{\{0\}^{\times N}\} = \{0\} \times (\mathcal{G}_0^{N-1} \setminus \{\{0\}^{\times (N-1)}\})$  is a subset of it. One can get  $\mathcal{G}_0^{N-1} \setminus \{\{0\}^{\times (N-1)}\} \subset \mathcal{G}_0^{N-1}$ , where  $\mathcal{G}_0^{N-1}$  on the right is taken from the subset  $\{t_0\} \times \mathcal{G}_0^{N-1}$  of  $\widetilde{\mathcal{G}_{t_0}^N}$  ( $t_0 \in T_1$ ). Since  $\widetilde{\mathcal{G}_{t_0}^N}$  satisfies the condition (ii), then  $\widetilde{\mathcal{G}_0^N}$  satisfies the condition (iii).

Now we provide an illustrative example of constructing d = 4, the distribution of element  $(\xi)^{\times N} (\xi \in \mathbb{Z}_4)$  in set  $\mathcal{G}_j^N (j \in \mathbb{Z}_4)$  is shown in Table II. For simplicity, we only present the set  $\bigcup_r \widetilde{\mathcal{G}_r^N}$  while omitting its corresponding set  $\widetilde{\mathcal{S}} = \bigcup_{i \in r} \widetilde{\mathcal{S}}_i$ .

TABLE II: The distribution of  $(\xi)^{\times N}$  when d = 4, where  $\xi \in \mathbb{Z}_4$ .

System $N$	$(0)^{\times N}$	$(1)^{\times N}$	$(2)^{\times N}$	$(3)^{\times N}$
[N] = [0]	$\mathcal{G}_0^N$	$\mathcal{G}_0^N$	$\mathcal{G}_0^N$	$\mathcal{G}_0^N$
[N] = [1]	$\mathcal{G}_0^N$	$\mathcal{G}_1^N$	$\mathcal{G}_2^N$	$\mathcal{G}_3^N$
[N] = [2]	$\mathcal{G}_0^N$	$\mathcal{G}_2^N$	$\mathcal{G}_0^N$	$\mathcal{G}_2^N$
[N] = [3]	$\mathcal{G}_0^N$	$\mathcal{G}_3^N$	$\mathcal{G}_2^N$	$\mathcal{G}_1^N$

*Example* 1. In  $(\mathbb{C}^4)^{\otimes N}$ , the following sets can be the strongest nonlocal OGESs.

$$\begin{split} [N] &= [0] \begin{cases} \widetilde{\mathcal{G}_{0}^{N}} &= \mathcal{G}_{0}^{N} \backslash \{(0)^{\times N}, \ (1)^{\times N}\}, \\ \widetilde{\mathcal{G}_{1}^{N}} &= \mathcal{G}_{1}^{N}, \\ \widetilde{\mathcal{G}_{2}^{N}} &= \mathcal{G}_{2}^{N}, \\ \widetilde{\mathcal{G}_{3}^{N}} &= \{(0)^{\times N}, (1)^{\times N}\}. \end{cases} \\ [N] &= [1] \begin{cases} \widetilde{\mathcal{G}_{0}^{N}} &= \mathcal{G}_{0}^{N} \backslash \{(0)^{\times N}\}, \\ \widetilde{\mathcal{G}_{1}^{N}} &= \mathcal{G}_{1}^{N}, \\ \widetilde{\mathcal{G}_{3}^{N}} &= \{(0)^{\times N}, (2)^{\times N}\}. \end{cases} \\ [N] &= [2] \begin{cases} \widetilde{\mathcal{G}_{0}^{N}} &= \mathcal{G}_{0}^{N} \backslash \{(0)^{\times N}, (2)^{\times N}\}, \\ \widetilde{\mathcal{G}_{1}^{N}} &= \mathcal{G}_{1}^{N}, \\ \widetilde{\mathcal{G}_{2}^{N}} &= \mathcal{G}_{2}^{N}, \\ \widetilde{\mathcal{G}_{3}^{N}} &= \{(0)^{\times N}, (2)^{\times N}\}. \end{cases} \\ [N] &= [3] \begin{cases} \widetilde{\mathcal{G}_{0}^{N}} &= \mathcal{G}_{0}^{N} \backslash \{(0)^{\times N}\}, \\ \widetilde{\mathcal{G}_{1}^{N}} &= \mathcal{G}_{1}^{N}, \\ \widetilde{\mathcal{G}_{3}^{N}} &= \{(0)^{\times N}, (2)^{\times N}\}. \end{cases} \end{cases} \\ [N] &= [3] \begin{cases} \widetilde{\mathcal{G}_{0}^{N}} &= \mathcal{G}_{2}^{N} \backslash \{(2)^{\times N}\}, \\ \widetilde{\mathcal{G}_{3}^{N}} &= \{(0)^{\times N}, (2)^{\times N}\}. \end{cases} \end{cases} \end{cases} \end{split}$$

It follows from the Theorem 4 Theorem 5 that the construction is not unique. In this regard, we give another example for d = 4 according to Table II.

*Example* 2. In  $(\mathbb{C}^4)^{\otimes N}$ , the following sets can also be the strongest nonlocal OGESs.

$$[N] = [0] \begin{cases} \widetilde{\mathcal{G}_0^N} = \mathcal{G}_0^N \backslash \{(0)^{\times N}, (2)^{\times N} \} \\ \widetilde{\mathcal{G}_1^N} = \mathcal{G}_1^N, \\ \widetilde{\mathcal{G}_2^N} = \mathcal{G}_2^N, \\ \widetilde{\mathcal{G}_3^N} = \{(0)^{\times N}, (2)^{\times N} \}. \end{cases}$$

$$\begin{split} [N] &= [1] \begin{cases} \widetilde{\mathcal{G}_{0}^{N}} = \mathcal{G}_{0}^{N} \backslash \{(0)^{\times N}\}, \\ \widetilde{\mathcal{G}_{1}^{N}} = \mathcal{G}_{1}^{N} \backslash \{(1)^{\times N}\}, \\ \widetilde{\mathcal{G}_{2}^{N}} = \mathcal{G}_{2}^{N}, \\ \widetilde{\mathcal{G}_{3}^{N}} = \{(0)^{\times N}, (1)^{\times N}\}. \end{cases} \\ [N] &= [2] \begin{cases} \widetilde{\mathcal{G}_{0}^{N}} = \mathcal{G}_{0}^{N} \backslash \{(0)^{\times N}\}, \\ \widetilde{\mathcal{G}_{1}^{N}} = \mathcal{G}_{1}^{N}, \\ \widetilde{\mathcal{G}_{3}^{N}} = \{(0)^{\times N}, (3)^{\times N}\}, \\ \widetilde{\mathcal{G}_{3}^{N}} = \{(0)^{\times N}, (3)^{\times N}\}. \end{cases} \\ [N] &= [3] \begin{cases} \widetilde{\mathcal{G}_{0}^{N}} = \mathcal{G}_{0}^{N} \backslash \{(0)^{\times N}\}, \\ \widetilde{\mathcal{G}_{1}^{N}} = \mathcal{G}_{1}^{N} \backslash \{(3)^{\times N}\}, \\ \widetilde{\mathcal{G}_{3}^{N}} = \{(0)^{\times N}, (3)^{\times N}\}, \\ \widetilde{\mathcal{G}_{3}^{N}} = \{(0)^{\times N}, (3)^{\times N}\}. \end{cases} \end{split}$$

The proof follows a similar approach as Theorem 5 and will not be repeated. Compared with Ref. [34], the set in our construction contains fewer states as N gradually increases. As shown in Tables III, IV, V, VI, for d = 4, 5 the size of states in our construction is significantly reduced when N > 4. For d = 6, 7 our size is much smaller when N > 3.

TABLE III: Comparisons of the sizes of strongest nonlocal OGESs in  $(\mathbb{C}^4)^{\otimes N}$ .

References	N=3	N = 4	N = 5	N = 6	N = 7	N = 8
Ref. [34]	38	176	782	3368	14198	58976
This work	48	192	768	3072	12288	49152

TABLE IV: Comparisons of the sizes of strongest nonlocal OGESs in  $(\mathbb{C}^5)^{\otimes N}$ .

References	N = 3	N = 4	N = 5	N = 6	N = 7	N = 8
Ref. [34]	62	370	2102	11530	61742	325090
This work	75	375	1875	9375	46875	234375

TABLE V: Comparisons of the sizes of strongest nonlocal OGESs in  $(\mathbb{C}^6)^{\otimes N}$ .

References	N = 3	N = 4	N = 5	N = 6	N = 7	N = 8
Ref. [34]	92	672	4652	31032	201812	1288992
This work	108	648	3888	23328	139968	839808

TABLE VI: Comparisons of the sizes of strongest nonlocal OGESs in  $(\mathbb{C}^7)^{\otimes N}$ .

References	N = 3	N = 4	N = 5	N = 6	N = 7	N = 8
Ref. [34]	128	1106	9032	70994	543608	4085186
This work	147	1029	7203	50421	352947	2470629

### V. CONCLUSION

In this paper, we investigate the strongest nonlocality in N-partite quantum system. First, we provide a sufficient and necessary condition for strongest nonlocal sets under some condition. Based on this condition, the minimum size of strongest nonlocal set in system  $(\mathbb{C}^3)^{\otimes N}$  is proven and supported by the OGESs constructed in Ref. [36]. Furthermore, we present a general construction in  $(\mathbb{C}^d)^{\otimes N}$  that demonstrates the existence of smaller strongest nonlocal set with genuine entanglement in high-dimensional multipartite systems. This construction also give an answer to the question proposed in Ref. [25], "How do we construct a strongly nonlocal orthogonal genuinely entangled set in  $(\mathbb{C}^d)^{\otimes N}$  for any  $d \geq 2$  and  $N \geq 5$ ?" Our work could enrich the understanding of the strongest nonlocality in multipartite systems.

### VI. ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China under Grants No. 12071110 and No. 62271189, the Hebei Central Guidance on Local Science and Technology Development Foundation of China under Grant No. 236Z7604G.

# Appendix A: PROOF OF PROPOSITION 1

Before proving the proposition, we introduce the following conception and notation.

*Circulant Matrix* [49]. A  $d \times d$  circulant matrix B is generated from the d-dimensional vector  $[b_0, \dots, b_{d-1}]$  by cyclically permuting its entries, and is

$$B = \begin{bmatrix} b_0 & b_1 & b_2 & \cdots & b_{d-2} & b_{d-1} \\ b_{d-1} & b_0 & b_1 & \cdots & b_{d-3} & b_{d-2} \\ b_{d-2} & b_{d-1} b_0 & \cdots & b_{d-4} & b_{d-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ b_2 & b_3 & b_4 & \cdots & b_0 & b_1 \\ b_1 & b_2 & b_3 & \cdots & b_{d-1} & b_0 \end{bmatrix}$$

In other word, a circulant matrix is cyclically shifted to the right by one position per row to form the subsequent rows.

Next, we use the circulant matrix to rewrite Eq. (22). Obviously,

$$\mathcal{G}_{i}^{N} = \left(\{i\} \times \mathcal{G}_{0}^{N-1}\right) \bigcup \left(\{i \oplus_{d} (d-1)\} \times \mathcal{G}_{1}^{N-1}\right) \bigcup \cdots \bigcup \left(\{i \oplus_{d} 2\} \times \mathcal{G}_{d-2}^{N-1}\right) \bigcup \left(\{i \oplus_{d} 1\} \times \mathcal{G}_{d-1}^{N-1}\right).$$
(A1)

We denote (A1) as

$$(\{i\} \times \mathcal{G}_0^{N-1}) \bigcup (\{i \oplus_d (d-1)\} \times \mathcal{G}_1^{N-1}) \bigcup \cdots \bigcup (\{i \oplus_d 2\} \times \mathcal{G}_{d-2}^{N-1}) \bigcup (\{i \oplus_d 1\} \times \mathcal{G}_{d-1}^{N-1})$$

$$= [i, i \oplus_d (d-1), \cdots, i \oplus_d 2, i \oplus_d 1] \times \begin{bmatrix} \mathcal{G}_0^{N-1} \\ \mathcal{G}_1^{N-1} \\ \vdots \\ \mathcal{G}_{d-2}^{N-1} \\ \mathcal{G}_{d-1}^{N-1} \end{bmatrix}.$$
(A2)

For each i in Eq. (A1), one gets

$$\begin{aligned}
\mathcal{G}_{0}^{N} &= \left(\{0\} \times \mathcal{G}_{0}^{N-1}\right) \bigcup \left(\{d-1\} \times \mathcal{G}_{1}^{N-1}\right) \bigcup \cdots \bigcup \left(\{2\} \times \mathcal{G}_{d-2}^{N-1}\right) \bigcup \left(\{1\} \times \mathcal{G}_{d-1}^{N-1}\right), \\
\mathcal{G}_{1}^{N} &= \left(\{1\} \times \mathcal{G}_{0}^{N-1}\right) \bigcup \left(\{0\} \times \mathcal{G}_{1}^{N-1}\right) \bigcup \cdots \bigcup \left(\{3\} \times \mathcal{G}_{d-2}^{N-1}\right) \bigcup \left(\{2\} \times \mathcal{G}_{d-1}^{N-1}\right), \\
\vdots \\
\mathcal{G}_{d-2}^{N} &= \left(\{d-2\} \times \mathcal{G}_{0}^{N-1}\right) \bigcup \left(\{d-3\} \times \mathcal{G}_{1}^{N-1}\right) \bigcup \cdots \bigcup \left(\{0\} \times \mathcal{G}_{d-2}^{N-1}\right) \bigcup \left(\{d-1\} \times \mathcal{G}_{d-1}^{N-1}\right), \\
\mathcal{G}_{d-1}^{N} &= \left(\{d-1\} \times \mathcal{G}_{0}^{N-1}\right) \bigcup \left(\{d-2\} \times \mathcal{G}_{1}^{N-1}\right) \bigcup \cdots \bigcup \left(\{1\} \times \mathcal{G}_{d-2}^{N-1}\right) \bigcup \left(\{0\} \times \mathcal{G}_{d-1}^{N-1}\right),
\end{aligned}$$
(A3)

which can be denoted as

$$\begin{bmatrix} \mathcal{G}_{0}^{N} \\ \mathcal{G}_{1}^{N} \\ \vdots \\ \mathcal{G}_{d-2}^{N} \\ \mathcal{G}_{d-2}^{N} \\ \mathcal{G}_{d-1}^{N} \end{bmatrix} = \begin{bmatrix} 0 & d-1 \cdots 2 & 1 \\ 1 & 0 & \cdots & 3 & 2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ d-2 & d-3 \cdots & 0 & d-1 \\ d-1 & d-2 \cdots & 1 & 0 \end{bmatrix} \times \begin{bmatrix} \mathcal{G}_{0}^{N-1} \\ \mathcal{G}_{1}^{N-1} \\ \vdots \\ \mathcal{G}_{d-2}^{N-1} \\ \mathcal{G}_{d-1}^{N-1} \end{bmatrix}.$$
(A4)  
$$\cdots \begin{array}{c} 2 & 1 \\ \cdots & 3 & 2 \end{bmatrix}$$

Here  $\begin{bmatrix} 0 & a-1 & \cdots & 2 & 1 \\ 1 & 0 & \cdots & 3 & 2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ d-2 & d-3 & \cdots & 0 & d-1 \\ d-1 & d-2 & \cdots & 1 & 0 \end{bmatrix}$  is a circulant matrix that generated from the *d*-dimensional vector  $[0, d-1, \cdots, 2, 1]$ .

Later on, we will give the proof of Proposition 1. Obviously, the sets  $\mathcal{G}_0^1, \dots, \mathcal{G}_{d-1}^1$  are pairwise disjoint and  $\bigcup_{i=0}^{d-1} \mathcal{G}_i^1 = \mathbb{Z}_d$ . We prove this proposition by induction. Assume that the claim is true for N = k, i.e.,  $\bigcup_{i=0}^{d-1} \mathcal{G}_i^k = \mathbb{Z}_d^k$  and  $\mathcal{G}_i^k \cap \mathcal{G}_j^k = \emptyset$ , if  $i \neq j \in \mathbb{Z}_d$ . Let l = k+1, for any two different sets  $\mathcal{G}_i^l = \bigcup_{x=0}^{d-1} (\{i_x\} \times \mathcal{G}_x^k)$  and  $\mathcal{G}_j^l = \bigcup_{y=0}^{d-1} (\{j_y\} \times \mathcal{G}_y^k)$ , according to Eq. (A4), if  $i_x = i_y$ , one gets  $x \neq y$ , from the induction hypothesis,  $\mathcal{G}_x^k \cap \mathcal{G}_y^k = \emptyset$  holds for k, thus  $(\{i_x\} \times \mathcal{G}_x^k) \cap (\{j_y\} \times \mathcal{G}_y^k) = \emptyset$ . If  $i_x \neq j_y$ , it definitely lead to  $(\{i_x\} \times \mathcal{G}_x^k) \cap (\{j_y\} \times \mathcal{G}_y^k) = \emptyset$ . In conclusion,  $\mathcal{G}_i^l \cap \mathcal{G}_j^l = \emptyset$ . On the other hand,

On the other hand,

$$\bigcup_{i=0}^{d-1} \mathcal{G}_i^l = \bigcup_{i=0}^{d-1} \bigcup_{j=0}^{d-1} \left( \{i \oplus_d (d-j)\} \times \mathcal{G}_j^k \right)$$
$$= \bigcup_{j=0}^{d-1} \left( \left( \bigcup_{i=0}^{d-1} \{i \oplus_d (d-j)\} \right) \times \mathcal{G}_j^k \right)$$
$$= \bigcup_{j=0}^{d-1} \left( \{0, 1, \cdots, d-1\} \times \mathcal{G}_j^k \right)$$
$$= \{0, 1, \cdots, d-1\} \times \bigcup_{j=0}^{d-1} \mathcal{G}_j^k$$
$$= \{0, 1, \cdots, d-1\} \times \mathbb{Z}_d^k$$
$$= \mathbb{Z}_d^l.$$

The proof is completed.

# **Appendix B: PROOF OF PROPOSITION 2**

Before the proof, we first give the following notation.

Assume that  $P_d = \{\sigma^r | r \in \mathbb{Z}_d\}$  is a cyclic permutation group of order d, and

$$\sigma[i_0, i_1, \cdots, i_{d-1}] = [i_{d-1}, i_0, \cdots, i_{d-2}],$$

where  $[i_0, i_1, \dots, i_{d-1}]$  is an arbitrary *d*-dimensional vector.

The proof of Proposition 2 is as follows. Observing Eq. (A2), we have that shifting the component of  $[i, i \oplus_d (d \begin{bmatrix} \mathcal{G}_0^{N-1} \\ \mathcal{G}_1^{N-1} \\ \vdots \\ \mathcal{G}_{d-2}^{N-1} \\ \mathcal{G}_{d-1}^{N-1} \end{bmatrix} \text{ to down by one position,}$ 

 $(1),\cdots,i\oplus_d 2,i\oplus_d 1]$  to the left by one position is equivalent to shifting the component of

that is,

$$\sigma^{-1}[i, i \oplus_{d} (d-1), \cdots, i \oplus_{d} 2, i \oplus_{d} 1] \times \begin{bmatrix} \mathcal{G}_{0}^{N-1} \\ \mathcal{G}_{1}^{N-1} \\ \vdots \\ \mathcal{G}_{d-2}^{N-1} \\ \mathcal{G}_{d-1}^{N-1} \end{bmatrix} = [i, i \oplus_{d} (d-1), \cdots, i \oplus_{d} 2, i \oplus_{d} 1] \times \sigma \begin{bmatrix} \mathcal{G}_{0}^{N-1} \\ \mathcal{G}_{1}^{N-1} \\ \vdots \\ \mathcal{G}_{d-2}^{N-1} \\ \mathcal{G}_{d-1}^{N-1} \end{bmatrix}.$$
(B1)

In Eq. (A4), performing a permutation  $\sigma^{-1}$  for each row of  $\begin{bmatrix} 0 & d-1 & \cdots & 2 & 1 \\ 1 & 0 & \cdots & 3 & 2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ d-2 & d-3 & \cdots & 0 & d-1 \\ d-1 & d-2 & \cdots & 1 & 0 \end{bmatrix}$ , one gets

$$\begin{bmatrix} d-1 \ d-2 \cdots & 1 & 0 \\ 0 \ d-1 \cdots & 2 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ d-3 \ d-4 \cdots & d-1 \ d-2 \\ d-2 \ d-3 \cdots & 0 \ d-1 \end{bmatrix} \times \begin{bmatrix} \mathcal{G}_0^{N-1} \\ \mathcal{G}_1^{N-1} \\ \vdots \\ \mathcal{G}_{d-2}^{N-1} \\ \mathcal{G}_{d-1}^{N-1} \end{bmatrix} = \begin{bmatrix} 0 \ d-1 \cdots & 2 & 1 \\ 1 \ 0 \ \cdots & 3 & 2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ d-2 \ d-3 \cdots & 0 \ d-1 \\ d-1 \ d-2 \cdots & 1 & 0 \end{bmatrix} \times \sigma \begin{bmatrix} \mathcal{G}_0^{N-1} \\ \mathcal{G}_1^{N-1} \\ \vdots \\ \mathcal{G}_{d-2}^{N-1} \\ \mathcal{G}_{d-1}^{N-1} \end{bmatrix}.$$
(B2)

On the other hand, one has

$$\begin{bmatrix} d-1 \ d-2 \cdots & 1 & 0 \\ 0 & d-1 \cdots & 2 & 1 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ d-3 \ d-4 \cdots & d-1 \ d-2 \\ d-2 \ d-3 \cdots & 0 & d-1 \end{bmatrix} \times \begin{bmatrix} \mathcal{G}_{0}^{N-1} \\ \mathcal{G}_{1}^{N-1} \\ \vdots \\ \mathcal{G}_{d-2}^{N-1} \\ \mathcal{G}_{d-1}^{N} \end{bmatrix} = \begin{bmatrix} \mathcal{G}_{d-1}^{N} \\ \mathcal{G}_{0}^{N} \\ \vdots \\ \mathcal{G}_{d-3}^{N} \\ \mathcal{G}_{d-2}^{N} \end{bmatrix} = \sigma \begin{bmatrix} \mathcal{G}_{0}^{N} \\ \mathcal{G}_{1}^{N} \\ \vdots \\ \mathcal{G}_{d-2}^{N} \\ \mathcal{G}_{d-1}^{N} \end{bmatrix}.$$
(B3)

By Eq. (B2) and Eq. (B3) we obtain

$$\sigma \begin{bmatrix} \mathcal{G}_{0}^{N} \\ \mathcal{G}_{1}^{N} \\ \vdots \\ \mathcal{G}_{d-2}^{N} \\ \mathcal{G}_{d-1}^{N} \end{bmatrix} = \begin{bmatrix} 0 & d-1 \cdots 2 & 1 \\ 1 & 0 & \cdots & 3 & 2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ d-2 & d-3 \cdots & 0 & d-1 \\ d-1 & d-2 \cdots & 1 & 0 \end{bmatrix} \times \sigma \begin{bmatrix} \mathcal{G}_{0}^{N-1} \\ \mathcal{G}_{1}^{N-1} \\ \vdots \\ \mathcal{G}_{d-2}^{N-1} \\ \mathcal{G}_{d-1}^{N-1} \end{bmatrix}.$$
(B4)

Let

$$\mathcal{V}_{N} := \begin{bmatrix} \mathcal{G}_{0}^{N} \\ \mathcal{G}_{1}^{N} \\ \vdots \\ \mathcal{G}_{d-2}^{N} \\ \mathcal{G}_{d-1}^{N} \end{bmatrix}$$
(B5)

and

$$\mathcal{M}_{d} := \begin{bmatrix} 0 & d-1 \cdots 2 & 1 \\ 1 & 0 & \cdots & 3 & 2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ d-2 & d-3 \cdots & 0 & d-1 \\ d-1 & d-2 \cdots & 1 & 0 \end{bmatrix},$$
(B6)

we get

$$\sigma \mathcal{V}_N = \mathcal{M}_d \times \sigma \mathcal{V}_{N-1}. \tag{B7}$$

Generating a circulant matrix of order d with column vectors  $\mathcal{V}_N, \sigma \mathcal{V}_N, \cdots, \sigma^{(d-1)} \mathcal{V}_N$ , by Eq. (B7), there is

$$\begin{bmatrix} \mathcal{V}_{N}, \sigma \mathcal{V}_{N}, \cdots, \sigma^{(d-2)} \mathcal{V}_{N}, \sigma^{(d-1)} \mathcal{V}_{N} \end{bmatrix}$$
  
= $\mathcal{M}_{d} \times \begin{bmatrix} \mathcal{V}_{N-1}, \sigma \mathcal{V}_{N-1}, \cdots, \sigma^{(d-2)} \mathcal{V}_{N-1}, \sigma^{(d-1)} \mathcal{V}_{N-1} \end{bmatrix}.$  (B8)

Repeating this argument, we obatin

$$\begin{bmatrix} \mathcal{V}_{N}, \sigma \mathcal{V}_{N}, \cdots, \sigma^{(d-2)} \mathcal{V}_{N}, \sigma^{(d-1)} \mathcal{V}_{N} \end{bmatrix}$$

$$= \underbrace{\mathcal{M}_{d} \times \cdots \times \mathcal{M}_{d}}_{N-1} \times \begin{bmatrix} \mathcal{V}_{1}, \sigma \mathcal{V}_{1}, \cdots, \sigma^{(d-2)} \mathcal{V}_{1}, \sigma^{(d-1)} \mathcal{V}_{1} \end{bmatrix}$$

$$= \underbrace{\mathcal{M}_{d} \times \cdots \times \mathcal{M}_{d}}_{N-1} \times \mathcal{M}_{d}.$$
(B9)

Here we use  $[\mathcal{V}_1, \sigma \mathcal{V}_1, \cdots, \sigma^{(d-2)} \mathcal{V}_1, \sigma^{(d-1)} \mathcal{V}_1] = \mathcal{M}_d$ . Substituting Eq. (B5) and Eq. (B6) into Eq. (B9), we have

[ 9 9	$egin{array}{ccc} \mathcal{G}_0^N & \mathcal{G}_{d-1}^N \ \mathcal{G}_1^N & \mathcal{G}_0^N \end{array}$	 	$\mathcal{G}_2^N \ \mathcal{G}_3^N$	$egin{array}{c} {\mathcal G}_1^N \ {\mathcal G}_2^N \end{array}$		$\begin{bmatrix} 0\\1 \end{bmatrix}$	d - 1 = 0	$1 \cdots 2$ $\cdots 3$	$\frac{1}{2}$	]	$\begin{bmatrix} 0\\ 1 \end{bmatrix}$	d-0	$1 \cdots 2$ $\cdots 3$	1 $2$			0 1	$d - 1 \\ 0$	· · · ·	$\frac{2}{3}$	$\begin{bmatrix} 1 \\ 2 \end{bmatrix}$
		۰.	:	:	=	:	÷	·.		×		÷	•	÷÷	×…	×	÷	÷	۰.	÷	:
$\mathcal{G}_{c}^{I}$	$\mathcal{G}_{d-2}^{N} \mathcal{G}_{d-3}^{N}$	• • •	$\mathcal{G}_0^N$	$\mathcal{G}_{d-1}^{N}$		d-2	2 d - 3	$3 \cdots 0$	d-1		d-1	2 d -	$3 \cdots 0$	d - 1			d-2	d-3	3	0 d	-1
$\left\lfloor \mathcal{G}_{c}^{r}\right\rfloor$	$\mathcal{G}_{d-1}^{\mathcal{H}}$		$\mathcal{G}_1^{n}$	$\mathcal{G}_0^{\prime\prime}$		d-1	1 d - 2	2 · · · 1	0	N	$\lfloor d - \rfloor$	1 d -	$2 \cdots 1$	0	N-1		d - 1	d-2	2	1 (B	$\begin{bmatrix} 0 \\ 10 \end{bmatrix}_1$

Beginning at the first square matrix on the right, like a snowball, the set of *N*-tuples is generated. As the right square matrices are the same, no matter how we change the position of the matrices, nothing is going to change on the left square matrix. Therefore the proof is now complete.

- C. H. Bennett, D. P. DiVincenzo, C. A. Fuchs, T. Mor, E. Rains, P. W. Shor, J. A. Smolin, and W. K. Wootters, Quantum nonlocality without entanglement, Phys. Rev. A 59, 1070 (1999).
- [2] J. Walgate, A. J. Short, L. Hardy, and V. Vedral, Local distinguishability of multipartite orthogonal quantum states, Phys. Rev. Lett. 85, 4972 (2000).
- [3] J. Walgate and L. Hardy, Nonlocality, asymmetry, and distinguishing bipartite states, Phys. Rev. Lett. 89, 147901 (2002).
- [4] H. Fan, Distinguishing bipartite states by local operations and classical communication, Phys. Rev. A 75, 014305 (2007).
- [5] S. M. Cohen, Local distinguishability with preservation of entanglement, Phys. Rev. A 75, 052313 (2007).
- [6] J. Niset and N. J. Cerf, Multipartite nonlocality without entanglement in many dimensions, Phys. Rev. A 74, 052103 (2006).
- [7] Y. Feng and Y. Shi, Characterizing locally indistinguishable orthogonal product states,

IEEE Trans. Inf. Theory 55, 2799 (2009).

- [8] N. Yu, R. Duan, and M. Ying, Any 2 ⊗ n subspace is locally distinguishable, Phys. Rev. A 84, 012304 (2011).
- [9] Z. C. Zhang, F. Gao, G. J. Tian, T. Q. Cao, and Q. Y. Wen, Nonlocality of orthogonal product basis quantum states, Phys. Rev. A 90, 022313 (2014).
- [10] Y. L. Wang, M. S. Li, Z. J. Zheng, and S. M. Fei, Nonlocality of orthogonal product-basis quantum states, Phys. Rev. A 92, 032313 (2015).
- [11] Y. L. Wang, M. S. Li, Z. J. Zheng, and S. M. Fei, The local indistinguishability of multipartite product states, Quantum Inf. Process. 16, 5 (2017).
- [12] S. Halder, Several nonlocal sets of multipartite pure orthogonal product states, Phys. Rev. A 98, 022303 (2018).
- [13] S. Ghosh, G. Kar, A. Roy, A. Sen (De), and U. Sen, Distinguishability of Bell states, Phys. Rev. Lett. 87, 277902 (2001).
- [14] S. Ghosh, G. Kar, A. Roy, and D. Sarkar, Distinguishability of maximally entangled states, Phys. Rev. A 70, 022304 (2004).
- [15] H. Fan, Distinguishability and indistinguishability by local operations and classical communication, Phys. Rev. Lett. 92, 177905 (2004).
- [16] S. Bandyopadhyay, S. Ghosh, and G. Kar, LOCC distinguishability of unilaterally transformable quantum states, New J. Phys. 13, 123013 (2011).
- [17] N. Yu, R. Duan, and M. Ying, Four locally indistinguishable ququad-ququad orthogonal maximally entangled states, Phys. Rev. Lett. **109**, 020506 (2012).
- [18] Y. L. Wang, M. S. Li, and Z. X. Xiong, One-way local distinguishability of generalized Bell states in arbitrary dimension, Phys. Rev. A 99, 022307 (2019).
- [19] M. Banik, T. Guha, M. Alimuddin, G. Kar, S. Halder, and S. S. Bhattacharya, Multicopy adaptive local discrimination: Strongest possible two-qubit nonlocal bases, Phys. Rev. Lett. **126**, 210505 (2021).
- [20] B. M. Terhal, D. P. DiVincenzo, and D. W. Leung, Hiding bits in Bell states, Phys. Rev. Lett. 86, 5807 (2001).
- [21] D. P. DiVincenzo, D. W. Leung, and B. M. Terhal, Quantum data hiding, IEEE Trans. Inf. Theory 48, 580 (2002).
- [22] R. Rahaman and M. G. Parker, Quantum scheme for secret sharing based on local distinguishability, Phys. Rev. A 91, 022330 (2015).
- [23] J. Wang, L. Li, H. Peng, and Y. Yang, Quantum-secret-sharing scheme based on local distinguishability of orthogonal multiqudit entangled states, Phys. Rev. A 95, 022320 (2017).
- [24] S. Halder, M. Banik, S. Agrawal, and S. Bandyopadhyay, Strong quantum nonlocality without entanglement, Phy. Rev. Lett. **122**, 040403 (2019).
- [25] F. Shi, Z. Ye, L. Chen, and X. Zhang, Strong quantum nonlocality in *N*-partite systems, Phys. Rev. A 105, 022209 (2022).
- [26] Z. C. Zhang and X. Zhang, Strong quantum nonlocality in multipartite quantum systems, Phys. Rev. A 99, 062108 (2019).
- [27] P. Yuan, G. J. Tian, and X. M. Sun, Strong quantum nonlocality without entanglement in multipartite quantum systems, Phys. Rev. A 102, 042228 (2020).
- [28] F. Shi, M. S. Li, M. Y. Hu, L. Chen, M. H. Yung, Y. L. Wang, and X. D. Zhang, Strongly nonlocal unextendible product bases do exist, Quantum 6, 619 (2022).
- [29] H. Q. Zhou, T. Gao, and F. L. Yan, Orthogonal product

sets with strong quantum nonlocality on a plane structure, Phys. Rev. A **106**, 052209 (2022).

- [30] H. Q. Zhou, T. Gao, and F. L. Yan, Strong quantum nonlocality without entanglement in an *n*-partite system with even *n*, Phys. Rev. A **107**, 042214 (2023).
- [31] Y. Y. He, F. Shi, and X. D. Zhang, Strong quantum nonlocality and unextendibility without entanglement in *N*-partite systems with odd *N*, Quantum **8**, 1349 (2024).
- [32] F. Shi, M. Y. Hu, L. Chen, and X. D. Zhang, Strong quantum nonlocality with entanglement, Phys. Rev. A 102, 042202 (2020).
- [33] Y. L. Wang, M. S. Li, and M. H. Yung, Graph-connectivitybased strong quantum nonlocality with genuine entanglement, Phys. Rev. A 104, 012424 (2021).
- [34] M. S. Li and Y. L. Wang, Bounds on the smallest sets of quantum states with special quantum nonlocality, Quantum 7, 1101 (2023).
- [35] J. Li, F. Shi, and X. D. Zhang, Strongest nonlocal sets with small sizes, Phys. Rev. A 108, 062407 (2023).
- [36] M. Y. Hu, T. Gao, and F. L. Yan, Strong quantum nonlocality with genuine entanglement in an *N*-qutrit system, Phys. Rev. A **109**, 022220 (2024).
- [37] X. F. Zhen, M. S. Li, and H. J. Zuo, Strongest nonlocal sets with minimum cardinality in tripartite systems, Phys. Rev. A 109, 052422 (2024).
- [38] A. Bhunia, S. Bera, I. Biswas, I. Chattopadhyay, and D. Sarkar, Strong quantum nonlocality: Unextendible biseparability beyond unextendible product basis, Phys. Rev. A 109, 052211 (2024).
- [39] Z. X, Xiong and M. S. Li, Existence of strongly nonlocal sets of three states in any N-partite system, arXiv: 2403.10969.
- [40] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Quantum entanglement, Rev. Mod. Phys. 81, 865 (2009).
- [41] J. S. Bell, On the Einstein Podolsky Rosen paradox, Physics 1, 195 (1964); On the problem of hidden variables in quantum mechanics, Rev. Mod. Phys. 38, 447 (1966).
- [42] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, Proposed experiment to test local hidden-variable theories, Phys. Rev. Lett. 23, 880 (1970).
- [43] F. L. Yan, T. Gao, and E. Chitambar, Two local observables are sufficient to characterize maximally entangled states of N qubits, Phys. Rev. A 83, 022319 (2011).
- [44] N. Brunner, D. Cavalcanti, S. Pironio, V. Scarani, and S. Wehner, Bell nonlocality, Rev. Mod. Phys. 86, 419 (2014); Erratum, Rev. Mod. Phys. 86, 839 (2014).
- [45] T. Gao, F. L. Yan, and S. J. van Enk, Permutationally invariant part of a density matrix and nonseparability of *N*-qubit states, Phys. Rev. Lett. **112**, 180501 (2014).
- [46] D. Ding, Y. Q. He, F. L. Yan, and T. Gao, Quantum nonlocality of generic family of four-qubit entangled pure states, Chin. Phys. B 24, 070301 (2015).
- [47] D. Ding, Y. Q. He, F. L. Yan, and T. Gao, Entanglement measure and quantum violation of Bell-type inequality, Int. J. Theor. Phys. 55, 4231 (2016).
- [48] H. X. Meng, J. Zhou, Z. P. Xu, H. Y. Su, T. Gao, F. L. Yan, and J. L. Chen, Hardy's paradox for multisetting high-dimensional systems, Phys. Rev. A 98, 062103 (2018).
- [49] D. S. Bernstein, Matrix Mathematics, Princeton University Press, Princeton, NJ, USA, 2005.