

Characterizing Quantum Codes via the Coefficients in Knill-Laflamme Conditions

Mengxin Du,¹ Chao Zhang,² Yiu Tung Poon,³ and Bei Zeng^{1,*}

¹*Department of Physics, The University of Texas at Dallas, Richardson, Texas 75080, USA*

²*Department of Physics, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong*

³*Department of Mathematics, Iowa State University, Ames, Iowa 50011, USA*

(Dated: October 11, 2024)

Quantum error correction (QEC) is essential for protecting quantum information against noise, yet understanding the structure of the Knill-Laflamme (KL) coefficients λ_{ij} from the condition $PE_i^\dagger E_j P = \lambda_{ij} P$ remains challenging, particularly for nonadditive codes. In this work, we introduce the signature vector $\vec{\lambda}(P)$, composed of the off-diagonal KL coefficients λ_{ij} , where each coefficient corresponds to equivalence classes of errors counted only once. We define its Euclidean norm $\lambda^*(P)$ as a scalar measure representing the total strength of error correlations within the code subspace defined by the projector P . We parameterize P on a Stiefel manifold and formulate an optimization problem based on the KL conditions to systematically explore possible values of λ^* . Moreover, we show that, for $((n, K, d))$ codes, λ^* is invariant under local unitary transformations. Applying our approach to the $((6, 2, 3))$ quantum code, we find that $\lambda_{\min}^* = \sqrt{0.6}$ and $\lambda_{\max}^* = 1$, with $\lambda^* = 1$ corresponding to a known degenerate stabilizer code. We construct continuous families of new nonadditive codes parameterized by vectors in \mathbb{R}^5 , with λ^* varying over the interval $[\sqrt{0.6}, 1]$. For the $((7, 2, 3))$ code, we identify $\lambda_{\min}^* = 0$ (corresponding to the non-degenerate Steane code) and $\lambda_{\max}^* = \sqrt{7}$ (corresponding to the permutation-invariant code by Pollatsek and Ruskai), and we demonstrate continuous paths connecting these extremes via cyclic codes characterized solely by λ^* . Our findings provide new insights into the structure of quantum codes, advance the theoretical foundations of QEC, and open new avenues for investigating intricate relationships between code subspaces and error correlations.

I. INTRODUCTION

Quantum error correction (QEC) is essential for protecting quantum information from the noise and errors that inevitably arise in quantum systems [1–6]. A deeper understanding of the structure of the set given by all possible coefficients λ_{ij} , which arise from the Knill-Laflamme (KL) conditions $PE_i^\dagger E_j P = \lambda_{ij} P$ [7], can provide valuable insights into the performance and underlying properties of quantum error-correcting codes. However, achieving this understanding is challenging. Nonadditive codes, which lie outside the stabilizer formalism, are particularly difficult to analyze, as relatively few examples have been systematically studied [8–10]. Moreover, degenerate codes—where multiple errors produce the same effect on the code space—exhibit inherently quantum phenomena, such as overlapping error syndromes, that lack classical analogues and remain poorly understood [11, 12]. These complexities make it difficult to systematically explore the structure of the set of all possible λ_{ij} values. As a result, there is currently no comprehensive framework for understanding the distribution of these coefficients, leaving important questions about their structure and implications for quantum error correction unanswered.

We analyze the structure defined by the set of all possible values of λ_{ij} that satisfy the KL conditions, which govern how pairs of errors interact within the code subspace defined by P . To encapsulate these interactions,

we introduce signature vector $\vec{\lambda}(P)$, composed of the off-diagonal elements λ_{ij} (with each coefficient corresponding to equivalent errors counted only once), which capture the non-trivial correlations between errors. The overall strength of these interactions is quantified by $\lambda^*(P)$, the Euclidean norm of the signature vector. This scalar value provides a measure of the total strength of error interactions within the code subspace, offering a new perspective on the role of these interactions in QEC. Crucially, for $((n, K, d))$ codes, λ^* is a function of the purity of the local reduced density matrices (RDMs) of the codewords, making it invariant under local unitary operations. This local unitary invariance allows λ^* to serve as a powerful tool for distinguishing locally unitary inequivalent quantum codes and identifying different codes based on their error interaction structures.

The focus of this paper is to study the range of λ^* : to understand the minimum and maximum values of λ^* (denoted by λ_{\min}^* and λ_{\max}^*), and to determine whether the range of λ^* is continuous between these extrema. We propose an algorithm to find λ_{\min}^* and λ_{\max}^* , as well as to identify possible intermediate values of λ^* between these limits. The core of our method involves characterizing the projector P on a Stiefel manifold, which provides a natural parameterization of the code subspace. We then formulate an optimization problem by constructing a loss function based on the KL conditions. This approach allows us to systematically explore the set of all possible λ_{ij} values and identify various values of λ^* corresponding to different quantum codes.

Using our method, we find that for the $((6, 2, 3))$ quantum code, $\lambda_{\min}^* = \sqrt{0.6}$ and $\lambda_{\max}^* = 1$. The value $\lambda^* = 1$

* bei.zeng@utdallas.edu

corresponds to the degenerate stabilizer code described in [11]. However, there are no known codes corresponding to $\lambda^* = \sqrt{0.6}$. We construct families of nonadditive codes, parameterized by four mutually orthogonal real vectors $a, b, c, d \in \mathbb{R}^5$, with λ^* parameterized by a vector $e \in \mathbb{R}^5$, orthogonal to a, b, c, d , which varies continuously over the interval $[\sqrt{0.6}, 1]$. This confirms that the range of λ^* for the $((6, 2, 3))$ code is indeed $[\sqrt{0.6}, 1]$. For each distinct value of $\lambda^* \in [\sqrt{0.6}, 1]$, our construction yields locally inequivalent codes, parameterized by the vector $e \in \mathbb{R}^5$.

For the $((7, 2, 3))$ code, we find that $\lambda_{\min}^* = 0$ and $\lambda_{\max}^* = \sqrt{7}$, where $\lambda^* = 0$ corresponds to the non-degenerate Steane code [13, 14], and $\lambda^* = \sqrt{7}$ corresponds to the permutation-invariant code proposed by Pollatsek and Ruskai [15]. We identify families of cyclic $((7, 2, 3))$ codes that trace continuous paths in the solution space. These paths, each characterized by a single parameter, which is simply λ^* , with λ^* varies continuously over the interval $[0, \sqrt{7}]$, directly connecting the Steane code and the permutation-invariant code. This finding demonstrates that it is possible to smoothly connect these two distinct codes while preserving cyclic symmetry, offering new insights into the relationship between different locally inequivalent quantum codes and their symmetry properties.

Our approach offers a systematic method to explore the range of λ^* , resulting in the construction of numerous new nonadditive codes for $((6, 2, 3))$ and $((7, 2, 3))$, with λ^* varying continuously from λ_{\min}^* to λ_{\max}^* . The ability to identify and quantify the range of λ^* provides novel insights into the structure of quantum codes, particularly in nonadditive cases. This framework opens new avenues for investigating the intricate relationships between code subspaces and error interactions, offering a deeper understanding of the mathematical structure underlying quantum error correction.

We organize our paper as follows. In Section II, we discuss preliminaries on quantum error correction and code parameters. In Section III, we define the signature vector and its norm λ^* , show that λ^* is invariant under local unitary operations by linking it to the purity of the RDMs of codewords, and develop an algorithm to find the maximum and minimum values of λ^* . In Sections IV and V, we apply our method to the $((6, 2, 3))$ and $((7, 2, 3))$ quantum codes, respectively, demonstrating how λ^* varies and constructing new nonadditive codes.

II. PRELIMINARY

In quantum error correction, the goal is to protect quantum information from errors caused by a noisy quantum channel. Quantum error-correcting codes (QECCs) are constructed to correct a specified set of errors. The Knill-Laflamme (KL) condition for quantum error correction can be expressed as:

$$PE_i^\dagger E_j P = \lambda_{ij} P, \quad \forall i, j, \quad (1)$$

where P denotes the projector onto the code subspace, E_i and E_j represent the Kraus operators corresponding to the possible errors, and λ_{ij} are complex scalars that characterize how the pair of errors E_i and E_j interact within the code subspace. This condition ensures that errors are correctable, provided that they act within the designated subspace and satisfy this equation.

The dimension of the code subspace is denoted as K , and if the logical information is encoded in a subspace of K -dimensional logical qubits within an n -dimensional physical qubit system, then P is an $n \times n$ matrix, and its rank equals K . The code subspace \mathcal{C} can be written as the span of orthonormal basis vectors $\{|\psi_1\rangle, |\psi_2\rangle, \dots, |\psi_K\rangle\}$, which span the logical space. The projector onto the code subspace is given by

$$P = \sum_{i=1}^K |\psi_i\rangle\langle\psi_i|. \quad (2)$$

The quantum error correction condition can then be expressed in terms of the basis vectors spanning the code subspace:

$$\langle\psi_k|E_i^\dagger E_j|\psi_l\rangle = \lambda_{ij}\delta_{kl}, \quad \forall i, j \quad (3)$$

where the scalars λ_{ij} describe how the errors E_i and E_j affect the code subspace.

A *non-degenerate QECC* is characterized by the Hermitian matrix λ_{ij} being non-singular (having full rank), which means that the determinant of λ_{ij} is non-zero and the matrix is invertible [1]. This implies that all errors have distinct effects on the code space and can be uniquely identified and corrected. In contrast, a *degenerate QECC* arises when the matrix λ_{ij} is singular (not of full rank), indicating that there are linear dependencies among the error operators when restricted to the code space [4]. Some errors or combinations of errors may have the same effect on the code space, making them indistinguishable. A *completely degenerate code*, or *decoherence-free subspace (DFS)*, represents an extreme case where all λ_{ij} elements are equal, resulting in a matrix of rank 1, meaning $\lambda_{ij} = \lambda$ for all i, j . In this scenario, the code space remains invariant under certain noise processes [11, 16].

An $((n, K, d))$ quantum error-correcting code is defined by three key parameters: n , the number of physical qubits used to encode the quantum information; K , the dimension of the code space, which corresponds to the number of logical qubits the code can protect (for example, if $K = 2^k$, the code protects k logical qubits); and d , the distance of the code, which determines the minimum number of physical qubit errors required to cause a logical error. The distance d indicates the code's ability

to detect and correct errors. Specifically, an $((n, K, d))$ code can detect up to $d - 1$ qubit errors and correct up to $t = \lfloor \frac{d-1}{2} \rfloor$ qubit errors [1, 4]. A well-known example is the Steane code, which is an $((7, 2, 3))$ code. This code encodes one logical qubit into seven physical qubits and can correct up to one qubit error and detect up to two qubit errors [4].

Furthermore, quantum error-correcting codes may either be *non-additive* or *additive*. Non-additive codes are a generalization of stabilizer (additive) codes and allow encoding of quantum information without adhering to the 2^k constraint for the dimension K . The code distance d , which is the minimum weight of an undetectable error, remains critical in both types of codes, as it determines how many errors can be detected and corrected.

Two quantum error-correcting codes P_1 and P_2 are locally equivalent if one can be transformed into the other by local unitary operations or local Clifford operations applied to individual qubits. Formally, P_1 and P_2 are locally equivalent if there exists a unitary transformation $U = U_1 \otimes U_2 \otimes \dots \otimes U_n$, where each U_i acts on a single qubit, such that $P_2 = UP_1U^\dagger$. This local equivalence ensures that the overall structure of the code and parameters n, K, d are preserved, even though individual states within the code space may change under the transformation [3].

In practice, to test whether two $((n, K, d))$ codes P_1 and P_2 are local unitary equivalent, we can use *Quantum weight enumerators* [17], which were defined by

$$A(z) = \sum_{j=0}^n A_j z^j, \quad B(z) = \sum_{j=0}^n B_j z^j \quad (4)$$

with coefficients

$$A_j = \frac{1}{K^2} \sum_{\text{wt}(O_\alpha)=j} \text{Tr}(O_\alpha P_c) \text{Tr}(O_\alpha^\dagger P_c), \quad (5)$$

$$B_j = \frac{1}{K} \sum_{\text{wt}(O_\alpha)=j} \text{Tr}(O_\alpha P_c O_\alpha^\dagger P_c). \quad (6)$$

Here

$$O_\alpha \in \{X, Y, Z, I\}^{\otimes n} \quad (7)$$

are n -fold Pauli tensor product. Denote the number of X factors, Y factors and Z factors in O_α as $\text{wt}_X(O_\alpha)$, $\text{wt}_Y(O_\alpha)$, and $\text{wt}_Z(O_\alpha)$. The weight of O_α is

$$\text{wt}(O_\alpha) = \text{wt}_X(O_\alpha) + \text{wt}_Y(O_\alpha) + \text{wt}_Z(O_\alpha). \quad (8)$$

And other related concepts including Rains' unitary and shadow quantum weight enumerators [18–20].

III. STRUCTURE OF λ_{ij}

Throughout this paper, we assume that the error operators E_i are Pauli operators for convenience. However,

our method naturally extends to non-Pauli errors as well. In this context, we simplify the analysis by focusing on the case where $i < j$. The quantum error correction criterion (KL) condition is given by $PE_i^\dagger E_j P = \lambda_{ij} P$, $\forall i, j$, where P is the projector onto the code subspace, and λ_{ij} encodes the interaction between errors E_i and E_j on the code subspace.

A. The signature vector

To capture the nature of these off-diagonal interactions, we define the signature vector as off-diagonal elements in matrix λ_{ij} in KL conditions. However, when dealing with Pauli errors, the product $E_i^\dagger E_j$ can be proportional to another Pauli operator that may already be included in our set of errors. This leads to a double counting problem, as the same operator can appear multiple times due to different pairs (i, j) . For example, with single-qubit errors such as X_i and Y_i , the product $X_i^\dagger Y_i$ is proportional to Z_i , which might already be included in the error set.

To resolve this issue and avoid double counting, we refine our definition by considering only unique error interactions. Specifically, we define an equivalence relation on the set of operator products $E_i^\dagger E_j$, where two operators are considered equivalent if they are proportional to a scalar multiple (including global phase). That is,

$$E_i^\dagger E_j \sim E_k^\dagger E_l \quad \text{if} \quad E_i^\dagger E_j = \alpha E_k^\dagger E_l,$$

for some non-zero α .

We then construct a set of representatives from each equivalence class of these operator products, ensuring that each unique operator (up to proportionality) is included only once. The *Signature Vector* $\vec{\lambda}(P)$ associated with a projector P is then defined using the corresponding λ_{ij} values for these representatives:

$$\vec{\lambda}(P) = (\lambda_{i_1 j_1}, \lambda_{i_2 j_2}, \dots, \lambda_{i_m j_m}), \quad (9)$$

where each $E_{i_k}^\dagger E_{j_k}$ is a distinct representative in \mathcal{S} , and $i_k < j_k$ for all k .

Each component of $\vec{\lambda}(P)$ represents the interaction between a pair of distinct errors E_{i_k} and E_{j_k} as captured by the quantum error correction criterion, without redundant counting. By focusing on these unique off-diagonal elements, the signature vector reflects the degree of correlation between different errors on the code space defined by P while avoiding double counting of equivalent error interactions.

Let $\mathcal{W}_{\text{error}}$ denote the set of all possible signature vectors corresponding to different projectors P that satisfy the quantum error correction criterion for a given error model $\{E_i\}$. Formally, we define:

$$\mathcal{W}_{\text{error}} = \{ \vec{\lambda}(P) : P \text{ satisfies the QEC criterion for error model } \{E_i\} \} \quad (10)$$

The set $\mathcal{W}_{\text{error}}$ represents all possible interactions between distinct errors under the given error model. Analyzing the structure of $\mathcal{W}_{\text{error}}$ is crucial for understanding the properties of quantum error-correcting codes.

The structure of $\mathcal{W}_{\text{error}}$ is closely related to the *rank- K joint numerical range* of the set of operators $\{E_i^\dagger E_j\}$. The rank- K joint numerical range for a set of operators $\{A_i\}_{i=1}^m$ is defined as:

$$\begin{aligned} W^{(K)}(A_1, A_2, \dots, A_m) = \\ \left\{ (\lambda_i) : \text{There exists a rank-}K \text{ projector } P \right. \\ \left. \text{such that } PA_iP = \lambda_i P \right\} \end{aligned} \quad (11)$$

This definition aligns with the quantum error correction conditions, therefore, studying $\mathcal{W}_{\text{error}}$ is equivalent to analyzing the rank- K joint numerical range of the operators $\{E_i^\dagger E_j\}$.

In particular, for $((n, K, d))$ codes, we have

$$\begin{aligned} \mathcal{W}_{\text{error}} &= W^{(K)}(\{E_{i_k}^\dagger E_{j_k}\}, i_k < j_k) \\ &= W^{(K)}(\{O_\alpha\}, 0 < \text{wt}(O_\alpha) < d). \end{aligned}$$

Here $O_\alpha \in \{X, Y, Z, I\}^{\otimes n}$ are n -fold Pauli tensor product.

The rank-1 joint numerical range is known to be always connected. It is convex for $m = 2$ Hermitian operators (as shown by the Toeplitz-Hausdorff theorem [21, 22]), so does the case for rank- K [23, 24]. But for $m > 2$ the properties of higher-rank joint numerical ranges are less well understood [25–29]. For $K \geq 2$, the rank- K joint numerical range of a set of operators is generally non-convex and can exhibit a complex structure, including disconnected components. As an example, consider a case of two qubits with $A_1 = X \otimes I$, $A_2 = X \otimes Z$, $A_3 = Y \otimes I$, $A_4 = Y \otimes Z$, $A_5 = Z \otimes I$. It can be shown that the rank-2 joint numerical range contains only two point

$$W^{(2)}(A_1, A_2, A_3, A_4, A_5) = \{(0, 0, 0, 0, 1), (0, 0, 0, 0, -1)\},$$

hence $W^{(2)}$ is disconnected.

In general, the connectedness of the rank- K joint numerical range depends on the specific operators involved. In this work, we study the connectedness of $\mathcal{W}_{\text{error}}$ for the case of the error-correcting code of interest.

B. The length of the signature vector $\lambda^*(P)$

We are particularly interested in the *length of the Signature Vector*, denoted by $\lambda^*(P)$, which is defined as:

$$\lambda^*(P) = \|\vec{\lambda}(P)\|_2 = \sqrt{\sum_{i_k < j_k} \lambda_{i_k j_k}^2} \quad (12)$$

This length provides a measure of the overall strength of the error interactions on the code subspace for the given

projector P [20]. A natural lower bound is $\lambda^* \geq 0$, where $\lambda^* = 0$ may be achieved when there exists codes with all $\lambda_{ij} = 0$, $i \neq j$ (e.g. nondegenerate stabilizer code). And for a given quantum channel a natural upper bound may be achieved when there exists codes with all $\lambda_{ij} = 1$ (i.e. the DFS).

Note that $\mathcal{W}_{\text{error}}$ may not be connected in general, implying that the set of possible values of $\lambda^*(P)$ may also not be continuous. In this study, we focus on the structure of the values of $\lambda^*(P)$, specifically investigating the minimum and maximum values of λ^* , denoted by λ_{\min}^* and λ_{\max}^* , respectively. Furthermore, we aim to determine whether the range of λ^* is continuous between these extrema. Although the continuity of the λ^* range does not imply the connectedness of $\mathcal{W}_{\text{error}}$, studying this range provides valuable insights into the structure of $\mathcal{W}_{\text{error}}$.

For $((n, K, d))$ codes, λ^* is the length of the vectors in $W^{(K)}(\{O_\alpha\}, 0 < \text{wt}(O_\alpha) < d)$. Furthermore, we show that λ^* , as defined above, is local unitary invariant (LUI), by linking λ^* to the purity of the reduced density matrices (RDMs) of the codewords (the purity of the reduced density matrix (RDM) is LUI).

Given a quantum state $|\psi\rangle$, the RDM for the i -th subsystem is defined as:

$$\rho^{(i)} = \text{Tr}_{(i)^c} [|\psi\rangle\langle\psi|],$$

where $\text{Tr}_{(i)^c}$ denotes the partial trace over all subsystems except the i -th one. The purity of this RDM is given by:

$$\mathcal{P}(\rho^{(i)}) = \text{Tr}[(\rho^{(i)})^2] \quad (13)$$

and since purity is invariant under local unitary transformations, the purity for 1-RDM, 2-RDM, ..., and up to $(d-1)$ -RDM is also LUI.

Next, consider the vector $\lambda^{(i)} = (\text{Tr}[\rho^{(i)} X_i], \text{Tr}[\rho^{(i)} Y_i], \text{Tr}[\rho^{(i)} Z_i])$, which captures how the i -th subsystem interacts with the Pauli operators. The length of this vector is LUI, and is expressed as:

$$\|\lambda^{(i)}\|_2 = \sqrt{\sum_{\alpha=1}^3 (\lambda_\alpha^{(i)})^2}.$$

This can be rewritten in terms of the purity as $\|\lambda^{(i)}\|_2^2 = 2\text{Tr}[\rho^{(i)}\rho^{(i)}] - 1$, demonstrating that $\|\lambda^{(i)}\|_2$ is LUI.

Now, let $\lambda^{(ij)}$ be a vector with 9 components, corresponding to the two-qubit interactions:

$$\lambda^{(ij)} = \left(\text{Tr}[\rho^{(ij)} X_i X_j], \text{Tr}[\rho^{(ij)} X_i Y_j], \dots, \text{Tr}[\rho^{(ij)} Z_i Z_j] \right).$$

The length of this vector is also LUI, and is given by:

$$\|\lambda^{(ij)}\|_2 = \sqrt{\sum_{\alpha=1}^9 (\lambda_\alpha^{(ij)})^2}.$$

We can express this as:

$$\|\lambda^{(ij)}\|_2^2 = 4\text{Tr}[\rho^{(ij)}\rho^{(ij)}] - 1 - \|\lambda^{(i)}\|_2^2 - \|\lambda^{(j)}\|_2^2,$$

where each term on the right-hand side has already been shown to be LUI.

With invariance of weight-1 and weight-2 vectors, in a similar fashion, the length of weight- $(d-1)$ vectors $\|\lambda^{(ij\dots)}\|_2$ can be proven to be LUI. Consequently, the length of the signature vector is given by:

$$(\lambda^*)^2 = \sum_i \|\lambda^{(i)}\|_2^2 + \sum_{ij} \|\lambda^{(ij)}\|_2^2 + \dots + (\text{weight-}(d-1) \text{ term}),$$

which is also LUI, as all terms involved are LUI. The LUI property can also be observed from the connection with quantum weight enumerators in Eq.(5). E.g., when $d=3$, $\lambda^{*2} = \sum_i \|\lambda^{(i)}\|_2^2 + \sum_{ij} \|\lambda^{(ij)}\|_2^2 = A_1 + A_2$.

Since λ^* is local unitary invariant (LUI), it follows that if two quantum codes P_1 and P_2 correspond to different values of λ^* , i.e., $\lambda^*(P_1) \neq \lambda^*(P_2)$, then the two codes must be local unitary inequivalent. This means that the distinct values of λ^* reflect different structures in the code subspaces that cannot be transformed into one another via local unitary operations. This shows that λ^* serves as a useful tool for distinguishing some local unitary inequivalent codes. However, the converse does not hold: two local unitary inequivalent codes may correspond to the same value of λ^* .

C. Algorithm for calculating range of λ^*

To parameterize the code space P , we use Stiefel manifold:

$$\text{St}(m, n) = \{x \in \mathbb{C}^{m \times n} : m \geq n, x^\dagger x = I_n\}.$$

Parametrization for Stiefel manifold is given by:

$$f(\theta) = \theta (\theta^\dagger \theta)^{-1/2} : \mathbb{C}^{m \times n} \rightarrow \text{St}(m, n)$$

Above is the polar decomposition which maps (full rank) complex matrix $\theta \in \mathbb{C}^{m \times n}$ to a Stiefel matrix and all Stiefel matrices can be generated in such a way [30]. We embed the code subspace into Stiefel manifold:

$$|\psi\rangle = \{|\psi_i\rangle : i = 1, \dots, K\} \in \text{St}(2^n, K) \subseteq \mathbb{C}^{2^n \times K}.$$

For the parametrized states $|\psi\rangle$ (not a valid code yet), we can calculate the tensor $\tilde{\lambda}_{\alpha, i, j} = \langle \psi_i | O_\alpha | \psi_j \rangle$. For the subspace to be a valid code, the following loss term \mathcal{L}_{KL} should be optimized to zero

$$\langle \tilde{\lambda}_{\alpha, i, i} \rangle_i = K^{-1} \sum_i \tilde{\lambda}_{\alpha, i, i}, \quad \|\tilde{\lambda}\|_2 = \sqrt{\sum_\alpha \langle \tilde{\lambda}_{\alpha, i, i} \rangle_i^2}$$

$$\mathcal{L}_{\text{KL}}(\theta) = \sum_{\alpha, i \neq j} |\tilde{\lambda}_{\alpha, i, j}|^2 + \sum_{\alpha, i} \left(\tilde{\lambda}_{\alpha, i, i} - \langle \tilde{\lambda}_{\alpha, i, i} \rangle_i \right)^2$$

To find the minimum length of λ vector, we can optimize the following loss

$$\mathcal{L}_\lambda(\theta; \mu) = \mu \mathcal{L}_{\text{KL}} + \|\tilde{\lambda}\|_2^2 \quad (14)$$

with \mathcal{L}_{KL} added as penalty and the hyper-parameter μ control the penalty strength. For a large enough μ , the optimal value of \mathcal{L}_λ should corresponds to λ with minimum length.

Similarly, to find the maximal length of λ , we can optimize the following loss function:

$$\mathcal{L}_{\tilde{\lambda}}(\theta; \mu) = \mu \mathcal{L}_{\text{KL}} - \|\tilde{\lambda}\|_2^2. \quad (15)$$

To find whether a code exists with length of λ equal to λ^* , we can define such a loss function:

$$\mathcal{L}(\theta; \mu, \lambda^*) = \mu \mathcal{L}_{\text{KL}} + \left(\|\tilde{\lambda}\|_2^2 - \lambda^{*2} \right)^2. \quad (16)$$

Notice that similarly one can also find the code with a predefined vector $\tilde{\lambda}$, just choose the loss function as:

$$\mathcal{L}(\theta; \mu, \tilde{\lambda}) = \mu \mathcal{L}_{\text{KL}} + \|\tilde{\lambda} - \tilde{\lambda}\|_2^2. \quad (17)$$

IV. THE ((6, 2, 3)) CODES

It is well known that ((5, 2, 3)) code is unique up to local unitary equivalence, with signature vector $\tilde{\lambda} = 0$, hence the range of λ^* is a single point 0. Much less is known about the range of λ^* for the case of ((6, 2, 3)). For stabilizer codes, there are only degenerate ones, for example the stabilizer code given in [31], with stabilizers given by

$$\begin{aligned} g_1 &: Y \ I \ Z \ X \ X \ Y \\ g_2 &: Z \ X \ I \ I \ X \ Z \\ g_3 &: I \ Z \ X \ X \ X \ X \\ g_4 &: I \ I \ I \ Z \ I \ Z \\ g_5 &: Z \ Z \ Z \ I \ Z \ I \end{aligned}$$

For this code, all components of signature vector are zero except the term $\langle 0_L | Z_4 Z_6 | 0_L \rangle = 1$, hence $\lambda^* = 1$. All the other ((6, 2, 3)) codes found in [32] also have $\lambda^* = 1$.

To find the range of λ^* , we sample $\lambda^* \in [0.5, 1.1]$, then calculate the optimal value for $\mathcal{L}(\theta; \mu, \lambda^*)$ in eq(16). The results are shown in Fig 1. For all optimizations, the violations of error-correcting conditions are less than $\mathcal{L}_{\text{KL}} \leq 10^{-15}$. From the figure, a sharp transition from almost zero to nonzero can be observed, which indicates $\|\lambda\|_2^2 \in [0.6, 1.0]$. This two boundaries are also found via optimizing $\mathcal{L}_{\tilde{\lambda}}$ and \mathcal{L}_λ .

A. Families of ((6, 2, 3)) codes with $\sqrt{0.6} \leq \lambda^* \leq 1$

To construct codes with $\sqrt{0.6} \leq \lambda^* \leq 1$, denote the six qubits by $q_1 q_2 q_3 q_4 q_5 q_6$, and choose the following five




FIG. 1: λ^{*2} range for $((6, 2, 3))$ code. Penalty factor is chosen $\mu = 1000$.

bases for qubits $q_2q_3q_4q_5q_6$

$$\begin{aligned} |S_1\rangle &= \frac{1}{\sqrt{2}} (|00001\rangle + |11110\rangle), \\ |S_2\rangle &= \frac{1}{\sqrt{2}} (|00010\rangle + |11101\rangle), \\ |S_3\rangle &= \frac{1}{\sqrt{2}} (|00100\rangle + |11011\rangle), \\ |S_4\rangle &= \frac{1}{\sqrt{2}} (|01000\rangle + |10111\rangle), \\ |S_5\rangle &= \frac{1}{\sqrt{2}} (|10000\rangle + |01111\rangle). \end{aligned}$$

Now choose logical states as:

$$|0_L\rangle = \sum_{i=1}^5 |x_i\rangle |S_i\rangle, \quad |1_L\rangle = \sum_{i=1}^5 |y_i\rangle |S_i\rangle.$$

Here

$$\begin{aligned} |x_i\rangle &= \gamma_i |0\rangle + \gamma_{i+5} |1\rangle, \quad i = 1, 2, 3, 4, 5 \\ |y_i\rangle &= \gamma_{i+5}^* |0\rangle - \gamma_i^* |1\rangle, \quad i = 1, 2, 3, 4, 5 \end{aligned}$$

The KL condition reduces to the following conditions on $|x_i\rangle$ and $|y_i\rangle$ (see Appendix A for details):

$$\sum_i |x_i\rangle \langle x_i| = \sum_i |y_i\rangle \langle y_i|, \quad \sum_i |y_i\rangle \langle x_i| = 0.$$

Notice that this is equivalent to require that the RDM of q_1 is $\frac{I}{2}$.

Now let

$$\gamma_j = a_j + ib_j, \quad \gamma_{5+j} = c_j + id_j,$$

for $j = 1, 2, 3, 4, 5$. Where $a_j, b_j, c_j, d_j \in \mathbb{R}$. Define the column vectors a, b, c , and d to be:

$$a = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \\ a_5 \end{pmatrix}, \quad b = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \\ b_5 \end{pmatrix}, \quad c = \begin{pmatrix} c_1 \\ c_2 \\ c_3 \\ c_4 \\ c_5 \end{pmatrix}, \quad d = \begin{pmatrix} d_1 \\ d_2 \\ d_3 \\ d_4 \\ d_5 \end{pmatrix}.$$

Then the KL condition will give the following conditions on a, b, c, d :

$$\begin{aligned} a \cdot a &= b \cdot b = c \cdot c = d \cdot d = \frac{1}{4}, \\ a \cdot b &= a \cdot c = a \cdot d = b \cdot c = b \cdot d = c \cdot d = 0. \end{aligned} \quad (18)$$

This means that a, b, c, d are orthogonal vectors in \mathbb{R}^5 .

We then choose e being the vector orthogonal to a, b, c, d , that is, the unnormalized orthogonal matrix composed from (a, b, c, d, e) as A

$$A = [a \ b \ c \ d \ e] = \begin{bmatrix} a_1 & b_1 & c_1 & d_1 & e_1 \\ a_2 & b_2 & c_2 & d_2 & e_2 \\ a_3 & b_3 & c_3 & d_3 & e_3 \\ a_4 & b_4 & c_4 & d_4 & e_4 \\ a_5 & b_5 & c_5 & d_5 & e_5 \end{bmatrix}$$

$$\rightarrow AA^T = A^T A = \frac{1}{4} I$$

which means each column (row) are orthogonal to each other. In other words, $2A$ is a 5×5 orthogonal matrix.

It turns out that the nonzero element of the signature vector, denoted as $PE_i^\dagger E_j P = \lambda_{ij} P$ of this code is given by the element of e (see Appendix A for details):

$$\begin{aligned} \lambda_{X_i X_j} &= \lambda_{Y_i Y_j} = -2e_{7-i} e_{7-j}, \\ \lambda_{Z_i Z_j} &= 2e_{7-i}^2 + 2e_{7-j}^2, \quad i, j \in \{2, 3, 4, 5, 6\}. \end{aligned}$$

And

$$\lambda^{*2} = \frac{1}{2} + 8 \sum_i e_i^4.$$

This means that λ^* is invariant with the rotation within the subspace spanned by (a, b, c, d) . To further understand this invariance, we can view $|0_L\rangle$ and $|1_L\rangle$ as bipartite states between q_1 (Party I) and $q_2q_3q_4q_5q_6$ (Party II). For Party I, an orthogonal transformation (i.e. $2A$, change of basis in the subspace spanned by $|S_i\rangle$) will correspond to a unitary transformation on Party II, hence will not change the RDM of Party I. This unitary in general cannot be realized by LU transformations on $q_2q_3q_4q_5q_6$, hence will lead to LU inequivalent codes.

It turns out (see Appendix A for details), however, when e is chosen, the freedom in the choice of (a, b, c, d) will lead to locally equivalent codes. This is due to the fact that, all such choices, given by

$$[a \ b \ c \ d] O,$$

where O is any 4×4 orthogonal matrix, can be generated by

1. local unitary transformations on party I (i.e. the first qubit) (leading to LU equivalent code), and
2. unitary transformations in the logical space spanned by $|0_L\rangle$ and $|1_L\rangle$ (leading to the same code).

In other words, the choice of e will in general lead to local inequivalent codes.

Specifically, the vector e for $\lambda_{\min}^{*2} = 0.6$ is

$$e = \frac{1}{2\sqrt{5}} [1 \ 1 \ 1 \ 1 \ 1],$$

and for $\lambda_{\max}^{*2} = 1$

$$e = \frac{1}{2} [0 \ 0 \ 0 \ 0 \ 1].$$

B. A single-parameter family

To have a single-parameter family of codes that connect λ_{\min}^* to $\lambda_{\max}^{*2} = 1$, since λ^* is only dependent on e , let us choose a single parameter family for e

$$e = \frac{1}{2} [\frac{1}{2} \sin \theta, \frac{1}{2} \sin \theta, \frac{1}{2} \sin \theta, \frac{1}{2} \sin \theta, \cos \theta],$$

for $\cos \theta \in [\frac{1}{\sqrt{5}}, 1]$.

Now we can choose the matrix A as

$$\mathbf{A} = \frac{1}{2} \begin{bmatrix} \frac{1}{2} & \frac{1}{2} & \frac{1}{2} & \frac{1}{2} \cos(\theta) & \frac{1}{2} \sin(\theta) \\ \frac{1}{2} & -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \cos(\theta) & \frac{1}{2} \sin(\theta) \\ -\frac{1}{2} & \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} \cos(\theta) & \frac{1}{2} \sin(\theta) \\ -\frac{1}{2} & -\frac{1}{2} & \frac{1}{2} & \frac{1}{2} \cos(\theta) & \frac{1}{2} \sin(\theta) \\ 0 & 0 & 0 & -\sin(\theta) & \cos(\theta) \end{bmatrix}$$

This gives a single parameter family of codes with the corresponding

$$\lambda^{*2} = \frac{1}{2} + \frac{1}{2} \left(\frac{1}{4} \sin^4(\theta) + \cos^4(\theta) \right)$$

runs continuously from 0.6 to 1.

For this family of codes, the matrix λ_{ij} will be block diagonal, and each block corresponding to $X_i X_j, Y_i Y_j, Z_i Z_j$ correlations, with the form (we only need to consider $i, j \in \{2, 3, 4, 5, 6\}$):

$$B = \begin{bmatrix} 1 & r & r & r & r \\ r & 1 & s & s & s \\ r & s & 1 & s & s \\ r & s & s & 1 & s \\ r & s & s & s & 1 \end{bmatrix}.$$

The eigenvalues of the matrix are:

$$\lambda_{B1} = \lambda_{B2} = \lambda_{B3} = 1 - s,$$

$$\lambda_{B4} = \frac{2 + 3s + \sqrt{9s^2 + 16r^2}}{2},$$

$$\lambda_{B5} = \frac{2 + 3s - \sqrt{9s^2 + 16r^2}}{2}.$$

Notice that

$$\begin{aligned} \lambda_{X_2 X_j} &= \lambda_{Y_2 Y_j} = -2e_5 e_{7-j} \\ &= -2 \left(\frac{1}{2} \cos \theta \frac{1}{4} \sin \theta \right), \quad j \in \{2, 3, 4, 5, 6\} \\ \lambda_{X_i X_j} &= \lambda_{Y_i Y_j} = -2e_{7-i} e_{7-j} \\ &= -2 \left(\frac{1}{4} \sin \theta \right)^2, \quad i, j \in \{3, 4, 5, 6\}. \end{aligned} \quad (19)$$

So the $X_i X_j, Y_i Y_j$ blocks are the same and correspond to

$$r = -\frac{1}{4} \sin \theta \cos \theta, \quad s = -2 \left(\frac{1}{4} \sin \theta \right)^2.$$

And

$$\begin{aligned} \lambda_{Z_2 Z_j} &= 2e_5^2 + 2e_{7-j}^2 \\ &= 2 \left(\frac{1}{4} \sin \theta \right)^2 + 2 \left(\frac{1}{2} \cos \theta \right)^2, \quad j \in \{2, 3, 4, 5, 6\} \\ \lambda_{Z_i Z_j} &= 2e_{7-i}^2 + 2e_{7-j}^2 \\ &= 4 \left(\frac{1}{4} \sin \theta \right)^2, \quad i, j \in \{3, 4, 5, 6\}. \end{aligned} \quad (20)$$

The $Z_i Z_j$ block corresponds to

$$r = 2 \left(\frac{1}{4} \sin \theta \right)^2 + 2 \left(\frac{1}{2} \cos \theta \right)^2, \quad s = 4 \left(\frac{1}{4} \sin \theta \right)^2.$$

So the matrix λ_{ij} will be full rank for $\cos \theta \in [\frac{1}{\sqrt{5}}, 1)$, i.e. $\lambda^* \in [\sqrt{0.6}, 1)$. For $\cos \theta = 1$, i.e. $\lambda^* = 1$, we have $\lambda_{B5} = 0$.

The enumerator is found to be:

$$\begin{aligned} A^{((6,2,3))} &= 1 + \left(\frac{3}{16} \cos(2\theta) + \frac{5}{64} \cos(4\theta) + \frac{47}{64} \right) z^2 \\ &\quad + \left(-\frac{3}{16} \cos(2\theta) - \frac{5}{64} \cos(4\theta) + \frac{17}{64} \right) z^3 \\ &\quad + \left(-\frac{3}{16} \cos(2\theta) - \frac{5}{64} \cos(4\theta) + \frac{721}{64} \right) z^4 \\ &\quad + \left(\frac{3}{16} \cos(2\theta) + \frac{5}{64} \cos(4\theta) + \frac{1007}{64} \right) z^5 + 3z^6 \end{aligned} \quad (21)$$

$$\begin{aligned} B^{((6,2,3))} &= 1 + \left(\frac{3}{16} \cos(2\theta) + \frac{5}{64} \cos(4\theta) + \frac{47}{64} \right) z^2 \\ &\quad + \left(\frac{3}{8} \cos(2\theta) + \frac{5}{32} \cos(4\theta) + \frac{751}{32} \right) z^3 \\ &\quad + \left(-\frac{3}{4} \cos(2\theta) - \frac{5}{16} \cos(4\theta) + \frac{577}{16} \right) z^4 \\ &\quad + \left(-\frac{3}{8} \cos(2\theta) - \frac{5}{32} \cos(4\theta) + \frac{1297}{32} \right) z^5 \\ &\quad + \left(\frac{9}{16} \cos(2\theta) + \frac{15}{64} \cos(4\theta) + \frac{1677}{64} \right) z^6 \end{aligned} \quad (22)$$

When $\cos(\theta) = 1$, we have $\lambda_{\max}^{*2} = 1$, and

$$e = \frac{1}{2} [0 \ 0 \ 0 \ 0 \ 1].$$

The code subspace, spanned by $(|0_L\rangle, |1_L\rangle)$, resides within the ground state space of the Hamiltonian

$$H = -2Z_2 \sum_{i \in \{3,4,5,6\}} Z_i + \frac{1}{2} \sum_{i \in \{3,4,5,6\}} \sum_{\substack{j \in \{3,4,5,6\} \\ j \neq i}} Z_i Z_j,$$

which is 16-dimensional degenerate, and is spanned by

$$|000001\rangle, |000010\rangle, |000100\rangle, |001000\rangle,$$

$$|011110\rangle, |011101\rangle, |011011\rangle, |010111\rangle,$$

$$|100001\rangle, |100010\rangle, |100100\rangle, |101000\rangle,$$

$$|111110\rangle, |111101\rangle, |111011\rangle, |110111\rangle.$$

This implies that the signature vector \vec{P} lies on the boundary of $W^{(1)}(\{O_\alpha\})$, where $\text{wt}(O_\alpha) = 1, 2$.

V. THE $((7, 2, 3))$ CODES

For the $((7, 2, 3))$ case, consider the Steane code with stabilizers

$$\begin{aligned} g_1 &: X \ I \ X \ I \ X \ I \ X \\ g_2 &: I \ X \ X \ I \ I \ X \ X \\ g_3 &: I \ I \ I \ X \ X \ X \ X \\ g_4 &: Z \ I \ Z \ I \ Z \ I \ Z \\ g_5 &: I \ Z \ Z \ I \ I \ Z \ Z \\ g_6 &: I \ I \ I \ Z \ Z \ Z \ Z \end{aligned}$$

This code has a signature vector $\vec{\lambda} = 0$, corresponding to $\lambda^* = 0$. To find the maximum value of λ^* , we run our algorithm and observe a sharp transition at $\lambda^* = \sqrt{7}$, as shown in Fig. 2.

It turns out that this maximum value $\lambda^* = \sqrt{7}$ corresponds to the permutation invariant code, which is constructed from the Dicke basis:

$$D_{n,k} = \binom{n}{k}^{-1/2} \sum_{\sigma \in \text{Sym}_n} \sigma |0\rangle^{\otimes n-k} \otimes |1\rangle^{\otimes k}$$

Two permutation invariant codes are given in [15] as:

$$\begin{aligned} 8|0_L\rangle &= \sqrt{15}D_{7,0} - \sqrt{7}D_{7,2} + \sqrt{21}D_{7,4} + \sqrt{21}D_{7,6} \\ |1_L\rangle &= X^{\otimes 7}|0_L\rangle \end{aligned} \quad (23)$$

and

$$\begin{aligned} 8|0_L\rangle &= \sqrt{15}D_{7,0} + \sqrt{7}D_{7,2} + \sqrt{21}D_{7,4} - \sqrt{21}D_{7,6} \\ |1_L\rangle &= X^{\otimes 7}|0_L\rangle \end{aligned} \quad (24)$$




FIG. 2: λ^{*2} range for $((7, 2, 3))$ code.

Notice that these two codes are local-unitary equivalent.

Now the key question is again, whether the set of all λ^* s is connected, i.e., whether the range of λ^* is indeed $[0, \sqrt{7}]$. Notice that by permuting the qubits, Steane code can have cyclic symmetry, with logical 0 and logical 1 given by

$$\begin{aligned} |0_L\rangle &= \frac{1}{\sqrt{8}} \left(|0000000\rangle + |1100101\rangle + |0101110\rangle + |0010111\rangle \right. \\ &\quad \left. + |1001011\rangle + |1110010\rangle + |0111001\rangle + |1011100\rangle \right) \\ &= \frac{1}{\sqrt{8}} |0000000\rangle + \sqrt{\frac{7}{8}} (|0010111\rangle + \text{cyc.}) \end{aligned} \quad (25)$$

and $|1_L\rangle = X^{\otimes 7}|0_L\rangle$. Here cyc. denotes all the other computational basis states with cyclic shift. Now we will explicitly construct families of cyclic codes with $\lambda^* \in [0, \sqrt{7}]$.

Let us choose the cyclic basis with even weights:

$$|\{0000000\}\rangle = |0000000\rangle$$

$$|\{0000011\}\rangle = \frac{1}{\sqrt{7}} (|0000011\rangle + \text{cyc.})$$

$$|\{0000101\}\rangle = \frac{1}{\sqrt{7}} (|0000101\rangle + \text{cyc.})$$

$$|\{0001001\}\rangle = \frac{1}{\sqrt{7}} (|0001001\rangle + \text{cyc.})$$

$$|\{0001111\}\rangle = \frac{1}{\sqrt{7}} (|0001111\rangle + \text{cyc.})$$

$$|\{0011011\}\rangle = \frac{1}{\sqrt{7}} (|0011011\rangle + \text{cyc.})$$

$$|\{0011101\}\rangle = \frac{1}{\sqrt{7}} (|0011101\rangle + \text{cyc.})$$

$$|\{0101011\}\rangle = \frac{1}{\sqrt{7}} (|0101011\rangle + \text{cyc.})$$

$$|\{0010111\}\rangle = \frac{1}{\sqrt{7}} (|0010111\rangle + \text{cyc.})$$

$$|\{0111111\}\rangle = \frac{1}{\sqrt{7}} (|0111111\rangle + \text{cyc.})$$

Using this basis, we parametrize $|0_L\rangle$ and $|1_L\rangle$ as follows:

$$\begin{aligned} |0_L\rangle &= c_0 |\{0000000\}\rangle \\ &+ \frac{c_1}{\sqrt{3}} \left(|\{0000011\}\rangle + |\{0000101\}\rangle + |\{0001001\}\rangle \right) \\ &+ c_2 |\{0010111\}\rangle + \frac{c_3}{2} \left(|\{0001111\}\rangle + |\{0011011\}\rangle \right. \\ &\left. + |\{0011101\}\rangle + |\{0101011\}\rangle \right) + c_4 |\{0111111\}\rangle, \\ |1_L\rangle &= X^{\otimes 7} |0_L\rangle \end{aligned} \quad (26)$$

Within the five-dimensional subspace given in Eq.(26), KL conditions will lead to three independent equations. Combined with normalization condition, the coefficients $(c_0, c_1, c_2, c_3, c_4)$ should satisfy the following four equations:

$$c_0^2 + c_1^2 + c_2^2 + c_3^2 + c_4^2 = 1 \quad (27)$$

$$\langle 0_L | Z_i | 0_L \rangle = 0 \rightarrow \quad (28)$$

$$7c_0^2 + 3c_1^2 - c_2^2 - c_3^2 - 5c_4^2 = 0$$

$$\langle 0_L | X_i X^{\otimes 7} | 0_L \rangle = 0 \rightarrow$$

$$2\sqrt{7}c_0c_4 + 2\sqrt{3}c_1c_2 + 4\sqrt{3}c_1c_3 \quad (29)$$

$$+ 4\sqrt{3}c_1c_4 + 4c_2c_3 + 3c_3^2 = 0$$

$$\langle 0_L | Y_i X^{\otimes 7} | 0_L \rangle = 0 \rightarrow$$

$$2\sqrt{7}c_0c_4 + 2\sqrt{3}c_1c_2 + 4\sqrt{3}c_1c_3 \quad (30)$$

$$- 4\sqrt{3}c_1c_4 - 4c_2c_3 - 3c_3^2 = 0$$

And for the signature vector, the following components are nonzero, satisfying (for $i \neq j$):

$$21\langle 0_L | X_i X_j | 0_L \rangle = 2\sqrt{21}c_0c_1 + 10c_1^2 + 4\sqrt{3}c_1c_2 + 8\sqrt{3}c_1c_3 + 12c_2c_3 + 6c_2c_4 + 9c_3^2 + 12c_3c_4 + 6c_4^2 \quad (31)$$

$$21\langle 0_L | Y_i Y_j | 0_L \rangle = -2\sqrt{21}c_0c_1 + 10c_1^2 - 4\sqrt{3}c_1c_2 - 8\sqrt{3}c_1c_3 + 12c_2c_3 - 6c_2c_4 + 9c_3^2 - 12c_3c_4 + 6c_4^2 \quad (32)$$

$$21\langle 0_L | Z_i Z_j | 0_L \rangle = 21c_0^2 + c_1^2 - 3c_2^2 - 3c_3^2 + 9c_4^2 \quad (33)$$

From Eq.(29) and Eq.(30) we obtain:

$$\sqrt{7}c_0c_4 + 2\sqrt{3}c_1c_2 + 2\sqrt{3}c_1c_3 = 0 \quad (34)$$

$$4\sqrt{3}c_1c_4 + 4c_2c_3 + 3c_3^2 = 0 \quad (35)$$

To solve these equations, we first find one solution for c_4 (see Appendix B for details):

$$c_4 = -\sqrt{3}c_1 \quad (36)$$

Then Eq.(36) and Eq.(34) derive another linear relation:

$$c_2 = -2c_3 + \sqrt{7}c_0 \quad (37)$$

Plug Eq.(36) and Eq.(37) into Eq.(27), (28), (31), (32), (33), (34) and (35), one finds f (for $i \neq j$):

$$\text{Normalization: } 8c_0^2 - 4\sqrt{7}c_0c_3 + 4c_1^2 + 5c_3^2 = 1 \quad (38)$$

$$\langle 0_L | Z_i | 0_L \rangle = 0 \rightarrow 4\sqrt{7}c_0c_3 - 12c_1^2 - 5c_3^2 = 0 \quad (39)$$

$$\langle 0_L | X_i X^{\otimes 7} | 0_L \rangle = 0 \rightarrow 4\sqrt{7}c_0c_3 - 12c_1^2 - 5c_3^2 = 0 \quad (40)$$

$$\langle 0_L | Y_i X^{\otimes 7} | 0_L \rangle = 0 \rightarrow -4\sqrt{7}c_0c_3 + 12c_1^2 + 5c_3^2 = 0 \quad (41)$$

$$21\langle 0_L | X_i X_j | 0_L \rangle = 12\sqrt{7}c_0c_3 + 28c_1^2 - 15c_3^2 \quad (42)$$

$$21\langle 0_L | Y_i Y_j | 0_L \rangle = 12\sqrt{7}c_0c_3 + 28c_1^2 - 15c_3^2 \quad (43)$$

$$21\langle 0_L | Z_i Z_j | 0_L \rangle = 12\sqrt{7}c_0c_3 + 28c_1^2 - 15c_3^2 \quad (44)$$

Since the signature vector components ((42), (43) and (44)) are equal, it is convenient to introduce λ^* as a parameter (for $i \neq j$):

$$21\langle 0_L | X_i X_j | 0_L \rangle = \sqrt{7}\lambda^* \quad (45)$$

$$21\langle 0_L | Y_i Y_j | 0_L \rangle = \sqrt{7}\lambda^* \quad (46)$$

$$21\langle 0_L | Z_i Z_j | 0_L \rangle = \sqrt{7}\lambda^* \quad (47)$$

By eliminating c_0 and c_3 through Eq.(39) and Eq.(42), we find $c_1 = \pm \frac{\sqrt{\sqrt{7}\lambda^*}}{8}$. With parameter $\lambda^* \in [0, \sqrt{7}]$, they become Steane code when $\lambda^* = 0$, and parametric code at $\lambda^* = \sqrt{7}$. The following two QECCs are related to QECC in eq(23):

$$\begin{aligned} c_0 &= \frac{\sqrt{\sqrt{7}\lambda^* + 8}}{8}, \\ c_1 &= -\frac{\sqrt{\sqrt{7}\lambda^*}}{8}, \\ c_4 &= -\sqrt{3}c_1, \end{aligned} \quad (48)$$

$$c_3 = \frac{2}{5} \left(\sqrt{7}c_0 \pm \sqrt{7c_0^2 - \frac{15\sqrt{7}\lambda^*}{64}} \right),$$

$$c_2 = -2c_3 + \sqrt{7}c_0$$

The following two correspond to QECC in Eq. (24)

$$\begin{aligned}
c_0 &= \frac{\sqrt{\sqrt{7}\lambda^* + 8}}{8}, \\
c_1 &= \frac{\sqrt{\sqrt{7}\lambda^*}}{8}, \\
c_4 &= -\sqrt{3}c_1, \\
c_3 &= \frac{2}{5} \left(\sqrt{7}c_0 \pm \sqrt{7c_0^2 - \frac{15\sqrt{7}\lambda^*}{64}} \right), \\
c_2 &= -2c_3 + \sqrt{7}c_0
\end{aligned} \tag{49}$$

For the quantity inside the square root non-negative, it requires $\lambda^* \leq \sqrt{7}$. The signature vector for these four codes are the same with the following nonzero components (for $i \neq j$):

$$\langle 0_L | X_i X_j | 0_L \rangle = \langle 0_L | Y_i Y_j | 0_L \rangle = \langle 0_L | Z_i Z_j | 0_L \rangle = \frac{\lambda^*}{3\sqrt{7}}$$

This means that all the 2-particle reduced density matrix of the code have the form

$$\rho^{(ij)} = \frac{1}{4}I + \frac{\lambda^*}{3\sqrt{7}}(X_i X_j + Y_i Y_j + Z_i Z_j).$$

Consequently, the matrix λ_{ij} will be block diagonal, and each block corresponding to $X_i X_j, Y_i Y_j, Z_i Z_j$ correlations, with the form

$$(1-s)I + sJ, \quad s = \frac{\lambda^*}{3\sqrt{7}} \in [0, \frac{1}{3}].$$

where:

- I is the 7×7 identity matrix.
- J is the 7×7 matrix with all entries equal to 1.

This matrix $(1-s)I + sJ$ has full rank and with one eigenvalues $6s + 1$ and six eigenvalues $1 - s$.

For the family given in Eq. (48), weight enumerators is given by

$$\begin{aligned}
A^{((7,2,3))} &= 1 + \lambda^{*2}z^2 + (21 - 2\lambda^{*2})z^4 + (42 + \lambda^{*2})z^6 \\
B^{((7,2,3))} &= 1 + \lambda^{*2}z^2 + 3(7 + \lambda^{*2})z^3 + (21 - 2\lambda^{*2})z^4 \\
&\quad + 6(21 - \lambda^{*2})z^5 + (42 + \lambda^{*2})z^6 + 3(15 + \lambda^{*2})z^7.
\end{aligned} \tag{50}$$

We have also explored all the local Clifford inequivalent $((7, 2, 3))$ stabilizer codes, and found that the only possible values of λ^* are $\{0, \sqrt{1}, \sqrt{2}, \sqrt{3}, \sqrt{5}\}$ (see Appendix C for details). For instance, the Bare code [33] corresponds to $\lambda^* = \sqrt{5}$.

When $\lambda^* = \lambda_{\max}^* = \sqrt{7}$, the code subspace, spanned by $(|0_L\rangle, |1_L\rangle)$, resides within the ground state space of the Hamiltonian

$$H = - \sum_{i \neq j} (X_i X_j + Y_i Y_j + Z_i Z_j).$$

This ground state space is 8-dimensional and corresponds to the symmetric subspace spanned by the Dicke basis. This implies that the signature vector \vec{P} lies on the boundary of $W^{(1)}(\{O_\alpha\})$, where $\text{wt}(O_\alpha) = 1, 2$.

ACKNOWLEDGMENTS

We gratefully acknowledge Chenfeng Cao for valuable discussions.

Appendix A: Details of the $((6,2,3))$ codes

1. The $|S_i\rangle$ basis

In the six-qubits system $q_1 q_2 q_3 q_4 q_5 q_6$, we choose a subspace for $q_2 q_3 q_4 q_5 q_6$ with basis $|S_i\rangle$:

$$\begin{aligned}
|S_1\rangle &= \frac{1}{\sqrt{2}} (|00001\rangle + |11110\rangle), \\
|S_2\rangle &= \frac{1}{\sqrt{2}} (|00010\rangle + |11101\rangle), \\
|S_3\rangle &= \frac{1}{\sqrt{2}} (|00100\rangle + |11011\rangle), \\
|S_4\rangle &= \frac{1}{\sqrt{2}} (|01000\rangle + |10111\rangle), \\
|S_5\rangle &= \frac{1}{\sqrt{2}} (|10000\rangle + |01111\rangle).
\end{aligned}$$

Their reduced density matrices (RDM) have clean forms. For example, 1-RDM

$$2\text{Tr}_{q_r q_s q_t q_\mu} [|S_i\rangle \langle S_j|] = \delta_{ij} I_2$$

2-RDM are

$$4\text{Tr}_{q_r q_s q_t} [|S_i\rangle \langle S_i|] = I_4 + (-1)^\delta Z \otimes Z$$

where the sign $\delta \in \{0, 1\}$ depends on how which qubits are chosen, for example:

$$\begin{aligned}
4\text{Tr}_{q_2 q_3 q_4} [|S_1\rangle \langle S_1|] &= I_4 - Z \otimes Z, \\
4\text{Tr}_{q_2 q_3 q_6} [|S_1\rangle \langle S_1|] &= I_4 + Z \otimes Z.
\end{aligned} \tag{A1}$$

Still, 2-RDM but with different basis

$$4\text{Tr}_{q_r q_s q_t} [|S_i\rangle \langle S_j|] = (X \otimes X + Y \otimes Y) \delta^{(i,j)},$$

$$4\text{Tr}_{q_2 q_3 q_4} [|S_1\rangle \langle S_2|] = (X \otimes X + Y \otimes Y),$$

$$\text{Tr}_{q_2 q_3 q_5} [|S_1\rangle \langle S_2|] = 0.$$

Our code are designed in the following 10-dimensional subspace

$$\{|0\rangle, |1\rangle\} \otimes \{|S_1\rangle, |S_2\rangle, |S_3\rangle, |S_4\rangle, |S_5\rangle\}$$

From above, all 2-RDM of pure states in this subspace will be in the form

$$\text{RDM}(q_i, q_j) = \frac{1}{4} I_4 + \alpha_{ij} (XX + YY) + \beta_{ij} ZZ.$$

The logical states are defined as

$$|x_i\rangle = \gamma_i |0\rangle + \gamma_{i+5} |1\rangle, \quad i = 1, 2, 3, 4, 5$$

$$|y_i\rangle = \gamma_{i+5}^* |0\rangle - \gamma_i^* |1\rangle, \quad i = 1, 2, 3, 4, 5$$

$$|0_L\rangle = \sum_{i=1}^5 |x_i\rangle |S_i\rangle, \quad |1_L\rangle = \sum_{i=1}^5 |y_i\rangle |S_i\rangle.$$

It is convenient to introduce the following shorthand notation for the 2-RDMs:

$$M_{ij}^{xx} = \langle x_i | x_j \rangle, \quad M_{ij}^{yx} = \langle y_i | x_j \rangle,$$

$$M_{ij}^{xy} = \langle x_i | y_j \rangle, \quad M_{ij}^{yy} = \langle y_i | y_j \rangle.$$

$$\begin{aligned} M(\square_1, \square_2, \square_3) &= \square_1 (|00\rangle \langle 00| + |11\rangle \langle 11|) \\ &\quad + \square_2 (|01\rangle \langle 01| + |10\rangle \langle 10|) \\ &\quad + \square_3 (|01\rangle \langle 10| + |10\rangle \langle 01|). \end{aligned}$$

$M(\square_1, \square_2, \square_3)$ is a 4-by-4 matrix with diagonal elements \square_1 and \square_2 , off-diagonal element \square_3 . Four important properties about the tensor M will be used later:

$$M_{ii}^{xy} = 0 \quad (\text{A2})$$

$$M_{ii}^{xx} = M_{ii}^{yy} \quad (\text{A3})$$

$$M_{ij}^{xx} = \gamma_i^* \gamma_j + \gamma_{i+5}^* \gamma_{j+5} = M_{ji}^{yy} \quad (\text{A4})$$

$$\begin{aligned} M_{ij}^{xy} + M_{ji}^{xy} &= (\gamma_i^* \gamma_{j+5}^* - \gamma_{i+5}^* \gamma_j^*) \\ &\quad + (\gamma_j^* \gamma_{i+5}^* - \gamma_{j+5}^* \gamma_i^*) = 0. \end{aligned} \quad (\text{A5})$$

Since $\langle 0_L | 0_L \rangle = \sum_i \langle x_i | x_i \rangle$, we have

$$\langle 0_L | 0_L \rangle = \sum_i M_{ii}^{xx} = \sum_i \gamma_i \gamma_i^* = 1.$$

With such a basis chosen, our code can already satisfy most of KL-conditions.

2. The 2-RDMs and KL conditions

a. RDM-(q_2, q_3) We first introduce RDM-(q_2, q_3):

$$\begin{aligned} \text{Tr}_{(23)^c} [|0_L\rangle \langle 0_L|] &= M(\square_1, \square_2, \square_3), \\ 2\square_1 &= M_{11}^{xx} + M_{22}^{xx} + M_{33}^{xx}, \\ 2\square_2 &= M_{44}^{xx} + M_{55}^{xx}, \\ 2\square_3 &= M_{45}^{xx} + M_{54}^{xx}, \end{aligned} \quad (\text{A6})$$

$$\begin{aligned} \text{Tr}_{(23)^c} [|1_L\rangle \langle 1_L|] &= M(\square'_1, \square'_2, \square'_3), \\ 2\square'_1 &= M_{11}^{yy} + M_{22}^{yy} + M_{33}^{yy}, \\ 2\square'_2 &= M_{44}^{yy} + M_{55}^{yy}, \\ 2\square'_3 &= M_{45}^{yy} + M_{54}^{yy}, \end{aligned} \quad (\text{A7})$$

$$\begin{aligned} \text{Tr}_{(23)^c} [|1_L\rangle \langle 0_L|] &= M(\square''_1, \square''_2, \square''_3), \\ 2\square''_1 &= M_{11}^{yx} + M_{22}^{yx} + M_{33}^{yx}, \\ 2\square''_2 &= M_{44}^{yx} + M_{55}^{yx}, \\ 2\square''_3 &= M_{45}^{yx} + M_{54}^{yx}. \end{aligned} \quad (\text{A8})$$

- From Eq.(A3), we have $\square_1 = \square'_1$ and $\square_2 = \square'_2$.
- From Eq.(A4), we have $\square_3 = \square'_3$.
- From Eq.(A2), we have $\square''_1 = \square''_2 = 0$.
- From Eq.(A5), we have $\square''_3 = 0$.

The $|0_L\rangle \langle 0_L|$ RDM is equal to $|1_L\rangle \langle 1_L|$ RDM and others are zero. It should be emphasized that this is true to all RDM(q_i, q_j), $i \neq 1, j \neq 1$.

b. RDM-($q_{i>1}, q_{j>1}$) Without loss of generality, only $|0_L\rangle \langle 0_L|$ is listed below:

$$\begin{aligned} \text{Tr}_{(24)^c} [|0_L\rangle \langle 0_L|] &= M(\square_1, \square_2, \square_3), \\ 2\square_1 &= M_{11}^{xx} + M_{22}^{xx} + M_{44}^{xx}, \\ 2\square_2 &= M_{33}^{xx} + M_{55}^{xx}, \\ 2\square_3 &= M_{35}^{xx} + M_{53}^{xx}, \end{aligned}$$

$$\begin{aligned} \text{Tr}_{(25)^c} [|0_L\rangle \langle 0_L|] &= M(\square_1, \square_2, \square_3), \\ 2\square_1 &= M_{11}^{xx} + M_{33}^{xx} + M_{44}^{xx}, \\ 2\square_2 &= M_{22}^{xx} + M_{55}^{xx}, \\ 2\square_3 &= M_{25}^{xx} + M_{52}^{xx}, \end{aligned}$$

$$\begin{aligned} \text{Tr}_{(26)^c} [|0_L\rangle \langle 0_L|] &= M(\square_1, \square_2, \square_3), \\ 2\square_1 &= M_{22}^{xx} + M_{33}^{xx} + M_{44}^{xx}, \\ 2\square_2 &= M_{11}^{xx} + M_{55}^{xx}, \\ 2\square_3 &= M_{15}^{xx} + M_{51}^{xx}, \end{aligned}$$

$$\begin{aligned} \text{Tr}_{(34)^c} [|0_L\rangle \langle 0_L|] &= M(\square_1, \square_2, \square_3), \\ 2\square_1 &= M_{11}^{xx} + M_{22}^{xx} + M_{55}^{xx}, \\ 2\square_2 &= M_{33}^{xx} + M_{44}^{xx}, \\ 2\square_3 &= M_{34}^{xx} + M_{43}^{xx}, \end{aligned}$$

$$\begin{aligned}
\text{Tr}_{(35)^c} [|0_L\rangle \langle 0_L|] &= M(\square_1, \square_2, \square_3), \\
2\square_1 &= M_{11}^{xx} + M_{33}^{xx} + M_{55}^{xx}, \\
2\square_2 &= M_{22}^{xx} + M_{44}^{xx}, \\
2\square_3 &= M_{24}^{xx} + M_{42}^{xx},
\end{aligned}$$

$$\begin{aligned}
\text{Tr}_{(36)^c} [|0_L\rangle \langle 0_L|] &= M(\square_1, \square_2, \square_3), \\
2\square_1 &= M_{22}^{xx} + M_{33}^{xx} + M_{55}^{xx}, \\
2\square_2 &= M_{11}^{xx} + M_{44}^{xx}, \\
2\square_3 &= M_{14}^{xx} + M_{41}^{xx},
\end{aligned}$$

$$\begin{aligned}
\text{Tr}_{(45)^c} [|0_L\rangle \langle 0_L|] &= M(\square_1, \square_2, \square_3), \\
2\square_1 &= M_{11}^{xx} + M_{44}^{xx} + M_{55}^{xx}, \\
2\square_2 &= M_{22}^{xx} + M_{33}^{xx}, \\
2\square_3 &= M_{23}^{xx} + M_{32}^{xx},
\end{aligned}$$

$$\begin{aligned}
\text{Tr}_{(46)^c} [|0_L\rangle \langle 0_L|] &= M(\square_1, \square_2, \square_3), \\
2\square_1 &= M_{22}^{xx} + M_{44}^{xx} + M_{55}^{xx}, \\
2\square_2 &= M_{11}^{xx} + M_{33}^{xx}, \\
2\square_3 &= M_{13}^{xx} + M_{31}^{xx},
\end{aligned}$$

$$\begin{aligned}
\text{Tr}_{(56)^c} [|0_L\rangle \langle 0_L|] &= M(\square_1, \square_2, \square_3), \\
2\square_1 &= M_{33}^{xx} + M_{44}^{xx} + M_{55}^{xx}, \\
2\square_2 &= M_{11}^{xx} + M_{22}^{xx}, \\
2\square_3 &= M_{12}^{xx} + M_{21}^{xx}. \tag{A9}
\end{aligned}$$

c. *RDM* ($q_1, q_{k>1}$) To satisfy KL-conditions, the RDMs

$$\text{Tr}_{(1k)^c} [|0_L\rangle \langle 0_L|] = \frac{1}{2} \sum_i |x_i\rangle \langle x_i| \otimes I_2,$$

$$\text{Tr}_{(1k)^c} [|1_L\rangle \langle 1_L|] = \frac{1}{2} \sum_i |y_i\rangle \langle y_i| \otimes I_2$$

$$\text{Tr}_{(1k)^c} [|1_L\rangle \langle 0_L|] = \frac{1}{2} \sum_i |y_i\rangle \langle x_i| \otimes I_2$$

must obey

$$\sum_i |x_i\rangle \langle x_i| = \sum_i |y_i\rangle \langle y_i|, \quad \sum_i |y_i\rangle \langle x_i| = 0.$$

From above, we can derive

$$\gamma = [a + ib \quad c + id] \in \mathbb{C}^{10}, \quad a, b, c, d \in \mathbb{R}^5,$$

$$\begin{aligned}
a \cdot a &= b \cdot b = c \cdot c = d \cdot d = \frac{1}{4}, \\
a \cdot b &= a \cdot c = a \cdot d = b \cdot c = b \cdot d = c \cdot d = 0.
\end{aligned}$$

$$\sum_i |x_i\rangle \langle x_i| = \frac{1}{2} I_2$$

Thus, vector (a, b, c, d) make four columns of orthogonal group $O(5)$.

3. Expression of λ^*

Let e being the vector orthogonal to a, b, c, d

$$e \cdot e = \frac{1}{4}, \quad e \cdot a = e \cdot b = e \cdot c = e \cdot d = 0, \quad e \in \mathbb{R}^5$$

Then we can prove

$$M_{ii}^{xx} = \frac{1}{4} - e_i^2, \quad M_{ij}^{xx} + M_{ji}^{xx} = -2e_i e_j.$$

Proof. Let's take $i = 1$ for example. Denote the unnormalized orthogonal matrix composed from (a, b, c, d, e) as:

$$\begin{aligned}
A &= [a \quad b \quad c \quad d \quad e] \\
&= \begin{bmatrix} a_1 & b_1 & c_1 & d_1 & e_1 \\ a_2 & b_2 & c_2 & d_2 & e_2 \\ a_3 & b_3 & c_3 & d_3 & e_3 \\ a_4 & b_4 & c_4 & d_4 & e_4 \\ a_5 & b_5 & c_5 & d_5 & e_5 \end{bmatrix} \tag{A10}
\end{aligned}$$

$$\rightarrow AA^T = A^T A = \frac{1}{4} I.$$

which means each column (row) are orthogonal to each other. Then

$$M_{11}^{xx} = a_1^2 + b_1^2 + c_1^2 + d_1^2 = \frac{1}{4} - e_1^2,$$

and similarly

$$\begin{aligned}
M_{12}^{xx} + M_{21}^{xx} &= a_1 a_2 + b_1 b_2 + c_1 c_2 + d_1 d_2 \\
&= (a_1 a_2 + b_1 b_2 + c_1 c_2 + d_1 d_2 + e_1 e_2) \\
&\quad - e_1 e_2 \\
&= -e_1 e_2. \tag{A11}
\end{aligned}$$

□

All nonzero components of signature vector can be written as

$$\begin{aligned}
\lambda_{X_i X_j} &= \lambda_{Y_i Y_j} = -2e_{7-i} e_{7-j}, \\
\lambda_{Z_i Z_j} &= 2e_{7-i}^2 + 2e_{7-j}^2, \quad i, j \in \{2, 3, 4, 5, 6\}. \tag{A12}
\end{aligned}$$

For example

$$\lambda_{X_2 X_3} = \lambda_{Y_2 Y_3} = -2e_4 e_5, \lambda_{Z_2 Z_3} = 2e_4^2 + 2e_5^2.$$

Then its square of length is

$$\lambda^{*2} = \frac{1}{2} + 8 \sum_i e_i^4$$

Proof. Two lemmas:

$$\sum_i \sum_{j \neq i} e_i^2 e_j^2 = \left(\sum_i e_i^2 \right)^2 - \sum_i e_i^4 = \frac{1}{16} - \sum_i e_i^4$$

$$\sum_i \sum_{j \neq i} e_i^4 = \sum_i \sum_j e_i^4 - \sum_i e_i^4 = 4 \sum_i e_i^4$$

Using lemmas above,

$$\begin{aligned} \lambda^{*2} &= \sum_i \sum_{j > i} 4e_i^2 e_j^2 + 4e_i^2 e_j^2 + 4(e_i^2 + e_j^2)^2 \\ &= \sum_i \sum_{j \neq i} 4e_i^2 e_j^2 + 2(e_i^2 + e_j^2)^2 \\ &= \sum_i \sum_{j \neq i} 8e_i^2 e_j^2 + 2e_i^4 + 2e_j^4 \\ &= \frac{1}{2} + 8 \sum_i e_i^4 \end{aligned}$$

□

4. Structure of SO(4) symmetry

The Lie algebra $\mathfrak{so}(4)$ comprises all 4×4 skew-symmetric matrices. A general skew-symmetric matrix $X \in \mathfrak{so}(4)$ satisfies $X^T = -X$. Such matrices have zeros on the diagonal and contain six independent off-diagonal elements. These non-zero elements correspond to infinitesimal rotations in the six independent planes of four-dimensional space ($x_i - x_j$ plane for $i \neq j$). Each generator can be represented by a matrix E_{ij} , where the (i, j) -th entry is +1, the (j, i) -th entry is -1, and all other entries are zero.

To obtain the Lie group SO(4), we exponentiate the Lie algebra elements. Specifically, for any skew-symmetric matrix $X \in \mathfrak{so}(4)$, the corresponding element in SO(4) can be obtained as:

$$\text{SO}(4) = \{e^X \mid X \in \mathfrak{so}(4)\}$$

Now choose the set

$$K_1 = E_{12} + E_{34}, K_2 = E_{12} - E_{34},$$

$$K_3 = E_{23} + E_{14}, K_4 = E_{23} - E_{14},$$

$$K_5 = E_{13} + E_{24}, K_6 = E_{13} - E_{24}.$$

Notice that

$$\exp(\theta K_1) = \begin{pmatrix} \cos \theta & \sin \theta & 0 & 0 \\ -\sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & \cos \theta & \sin \theta \\ 0 & 0 & -\sin \theta & \cos \theta \end{pmatrix}$$

$$\exp(\theta K_2) = \begin{pmatrix} \cos \theta & \sin \theta & 0 & 0 \\ -\sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & \cos \theta & -\sin \theta \\ 0 & 0 & \sin \theta & \cos \theta \end{pmatrix}$$

$$\exp(\theta K_3) = \begin{pmatrix} \cos \theta & 0 & 0 & \sin \theta \\ 0 & \cos \theta & \sin \theta & 0 \\ 0 & -\sin \theta & \cos \theta & 0 \\ -\sin \theta & 0 & 0 & \cos \theta \end{pmatrix}$$

$$\exp(\theta K_4) = \begin{pmatrix} \cos \theta & 0 & 0 & -\sin \theta \\ 0 & \cos \theta & \sin \theta & 0 \\ 0 & -\sin \theta & \cos \theta & 0 \\ \sin \theta & 0 & 0 & \cos \theta \end{pmatrix}$$

$$\exp(\theta K_5) = \begin{pmatrix} \cos \theta & 0 & \sin \theta & 0 \\ 0 & \cos \theta & 0 & \sin \theta \\ -\sin \theta & 0 & \cos \theta & 0 \\ 0 & -\sin \theta & 0 & \cos \theta \end{pmatrix}$$

$$\exp(\theta K_6) = \begin{pmatrix} \cos \theta & 0 & \sin \theta & 0 \\ 0 & \cos \theta & 0 & -\sin \theta \\ -\sin \theta & 0 & \cos \theta & 0 \\ 0 & \sin \theta & 0 & \cos \theta \end{pmatrix}$$

Notice that

$$\exp(-i\theta X) = \cos \theta I - i \sin \theta X = \begin{bmatrix} \cos \theta & -i \sin \theta \\ -i \sin \theta & \cos \theta \end{bmatrix}$$

$$\exp(-i\theta Y) = \cos \theta I - i \sin \theta Y = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

$$\exp(-i\theta Z) = \cos \theta I - i \sin \theta Z = \begin{bmatrix} e^{-i\theta} & 0 \\ 0 & e^{i\theta} \end{bmatrix}.$$

Write

$$|x_i\rangle = (a_i + ib_i) |0\rangle + (c_i + id_i) |1\rangle = \begin{bmatrix} a_i + ib_i \\ c_i + id_i \end{bmatrix}.$$

$$|y_i\rangle = (c_i - id_i) |0\rangle - (a_i - ib_i) |1\rangle = \begin{bmatrix} c_i - id_i \\ -a_i + ib_i \end{bmatrix}.$$

We now compare

$$\exp \theta K_i$$

with the representation of

- $\exp i\theta X_1, \exp i\theta Y_1, \exp i\theta Z_1$
- $\exp i\theta X_L, \exp i\theta Y_L, \exp i\theta Z_L$

as SO(4) rotations

a. $\exp(-i\theta X_1)$

$$\exp(-i\theta X_1) |x_i\rangle = \begin{bmatrix} \cos \theta & -i \sin \theta \\ -i \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} a_i + ib_i \\ c_i + id_i \end{bmatrix}$$

$$= \begin{bmatrix} (\cos \theta a_i + \sin \theta d_i) + i(\cos \theta b_i - \sin \theta c_i) \\ (\sin \theta b_i + \cos \theta c_i) + i(-\sin \theta a_i + \cos \theta d_i) \end{bmatrix}$$

As a SO(4) rotation, this is

$$[a \ b \ c \ d] \begin{bmatrix} \cos \theta & 0 & 0 & -\sin \theta \\ 0 & \cos \theta & \sin \theta & 0 \\ 0 & -\sin \theta & \cos \theta & 0 \\ \sin \theta & 0 & 0 & \cos \theta \end{bmatrix},$$

which corresponds to $\exp(\theta K_4)$.

b. $\exp(-i\theta Y_1)$

$$\exp(-i\theta Y_1) |x_i\rangle = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} a_i + ib_i \\ c_i + id_i \end{bmatrix}$$

$$= \begin{bmatrix} (\cos \theta a_i - \sin \theta c_i) + i(\cos \theta b_i - \sin \theta d_i) \\ (\sin \theta a_i + \cos \theta c_i) + i(\sin \theta b_i + \cos \theta d_i) \end{bmatrix}$$

As a SO(4) rotation, this is

$$[a \ b \ c \ d] \begin{bmatrix} \cos \theta & 0 & \sin \theta & 0 \\ 0 & \cos \theta & 0 & \sin \theta \\ -\sin \theta & 0 & \cos \theta & 0 \\ 0 & -\sin \theta & 0 & \cos \theta \end{bmatrix},$$

which corresponds to $\exp(\theta K_5)$.

c. $\exp(-i\theta Z_1)$

$$\exp(-i\theta Z_1) |x_i\rangle =$$

$$\begin{bmatrix} \cos \theta - i \sin \theta & 0 \\ 0 & \cos \theta + i \sin \theta \end{bmatrix} \begin{bmatrix} a_i + ib_i \\ c_i + id_i \end{bmatrix}$$

$$= \begin{bmatrix} (\cos \theta a_i + \sin \theta b_i) + i(-\sin \theta a_i + \cos \theta b_i) \\ (\cos \theta c_i - \sin \theta d_i) + i(\sin \theta c_i + \cos \theta d_i) \end{bmatrix}$$

As a SO(4) rotation, this is

$$[a \ b \ c \ d] \begin{bmatrix} \cos \theta & -\sin \theta & 0 & 0 \\ \sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & \cos \theta & \sin \theta \\ 0 & 0 & -\sin \theta & \cos \theta \end{bmatrix},$$

which corresponds to $\exp(-\theta K_2)$.

d. $\exp(-i\theta X_L)$

$$\exp(-i\theta X_L) |0_L\rangle = \cos \theta |0_L\rangle - i \sin \theta |1_L\rangle$$

$$\rightarrow \cos \theta \begin{bmatrix} a_i + ib_i \\ c_i + id_i \end{bmatrix} - i \sin \theta \begin{bmatrix} c_i - id_i \\ -a_i + ib_i \end{bmatrix}$$

$$= \begin{bmatrix} (\cos \theta a_i - \sin \theta d_i) + i(\cos \theta b_i - \sin \theta c_i) \\ (\sin \theta b_i + \cos \theta c_i) + i(\sin \theta a_i + \cos \theta d_i) \end{bmatrix}$$

As a SO(4) rotation, this is

$$[a \ b \ c \ d] \begin{bmatrix} \cos \theta & 0 & 0 & \sin \theta \\ 0 & \cos \theta & \sin \theta & 0 \\ 0 & -\sin \theta & \cos \theta & 0 \\ -\sin \theta & 0 & 0 & \cos \theta \end{bmatrix},$$

which corresponds to $\exp(\theta K_3)$.

e. $\exp(-i\theta Y_L)$

$$\exp(-i\theta Y_L) |0_L\rangle = \cos \theta |0_L\rangle + \sin \theta |1_L\rangle$$

$$\rightarrow \cos \theta \begin{bmatrix} a_i + ib_i \\ c_i + id_i \end{bmatrix} + \sin \theta \begin{bmatrix} c_i - id_i \\ -a_i + ib_i \end{bmatrix}$$

$$= \begin{bmatrix} (\cos \theta a_i + \sin \theta c_i) + i(\cos \theta b_i - \sin \theta d_i) \\ (-\sin \theta a_i + \cos \theta c_i) + i(\sin \theta b_i + \cos \theta d_i) \end{bmatrix}$$

As a SO(4) rotation, this is

$$[a \ b \ c \ d] \begin{bmatrix} \cos \theta & 0 & -\sin \theta & 0 \\ 0 & \cos \theta & 0 & \sin \theta \\ \sin \theta & 0 & \cos \theta & 0 \\ 0 & -\sin \theta & 0 & \cos \theta \end{bmatrix},$$

which corresponds to $\exp(-\theta K_6)$.

f. $\exp(-i\theta Z_L)$

$$\exp(-i\theta Z_L) |0_L\rangle = (\cos \theta - i \sin \theta) |0_L\rangle$$

$$\rightarrow (\cos \theta - i \sin \theta) \begin{bmatrix} a_i + ib_i \\ c_i + id_i \end{bmatrix}$$

$$= \begin{bmatrix} (\cos \theta a_i + \sin \theta b_i) + i(-\sin \theta a_i + \cos \theta b_i) \\ (\cos \theta c_i + \sin \theta d_i) + i(-\sin \theta c_i + \cos \theta d_i) \end{bmatrix}$$

As a SO(4) rotation, this is

$$\begin{bmatrix} a & b & c & d \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta & 0 & 0 \\ \sin \theta & \cos \theta & 0 & 0 \\ 0 & 0 & \cos \theta & -\sin \theta \\ 0 & 0 & \sin \theta & \cos \theta \end{bmatrix},$$

which corresponds to $\exp(-\theta K_1)$.

Appendix B: Details for the ((7,2,3) codes

We provide the details for solving the following system of equations:

$$c_0^2 + c_1^2 + c_2^2 + c_3^2 + c_4^2 = 1 \quad (\text{E1})$$

$$7c_0^2 + 3c_1^2 - c_2^2 - c_3^2 - 5c_4^2 = 0 \quad (\text{E2})$$

$$\begin{aligned} 2\sqrt{7} c_0 c_4 + 2\sqrt{3} c_1 c_2 + 4\sqrt{3} c_1 c_3 \\ + 4\sqrt{3} c_1 c_4 + 4c_2 c_3 + 3c_3^2 = 0 \end{aligned} \quad (\text{E3})$$

$$\begin{aligned} 2\sqrt{7} c_0 c_4 + 2\sqrt{3} c_1 c_2 + 4\sqrt{3} c_1 c_3 \\ - 4\sqrt{3} c_1 c_4 - 4c_2 c_3 - 3c_3^2 = 0 \end{aligned} \quad (\text{E4})$$

Our goal is to eliminate c_0 , c_2 , and c_3 to obtain an equation in terms of c_1 and c_4 .

Step 1: Subtract Eq.(E4) from Eq.(E3)

Subtracting Eq.(E4) from Eq.(E3):

$$\begin{aligned} & \left[2\sqrt{7} c_0 c_4 + 2\sqrt{3} c_1 c_2 + 4\sqrt{3} c_1 c_3 \right. \\ & \quad \left. + 4\sqrt{3} c_1 c_4 + 4c_2 c_3 + 3c_3^2 \right] \\ & - \left[2\sqrt{7} c_0 c_4 + 2\sqrt{3} c_1 c_2 + 4\sqrt{3} c_1 c_3 \right. \\ & \quad \left. - 4\sqrt{3} c_1 c_4 - 4c_2 c_3 - 3c_3^2 \right] = 0 \end{aligned}$$

Simplify:

$$\begin{aligned} 8\sqrt{3} c_1 c_4 + 8c_2 c_3 + 6c_3^2 = 0 \\ \Rightarrow 4\sqrt{3} c_1 c_4 + 4c_2 c_3 + 3c_3^2 = 0 \end{aligned} \quad (\text{E5})$$

Step 2: Add Equations (E3) and (E4)

Adding Equations (E3) and (E4):

$$\begin{aligned} & \left[2\sqrt{7} c_0 c_4 + 2\sqrt{3} c_1 c_2 + 4\sqrt{3} c_1 c_3 \right. \\ & \quad \left. + 4\sqrt{3} c_1 c_4 + 4c_2 c_3 + 3c_3^2 \right] \\ & + \left[2\sqrt{7} c_0 c_4 + 2\sqrt{3} c_1 c_2 + 4\sqrt{3} c_1 c_3 \right. \\ & \quad \left. - 4\sqrt{3} c_1 c_4 - 4c_2 c_3 - 3c_3^2 \right] = 0 \end{aligned}$$

Simplify:

$$\begin{aligned} 4\sqrt{7} c_0 c_4 + 4\sqrt{3} c_1 c_2 + 8\sqrt{3} c_1 c_3 = 0 \\ \Rightarrow \sqrt{7} c_0 c_4 + \sqrt{3} c_1 c_2 + 2\sqrt{3} c_1 c_3 = 0 \end{aligned} \quad (\text{E6})$$

Step 3: Eliminate c_0 Using Equations (E1) and (E2)

From Eq.(E1):

$$c_0^2 = 1 - c_1^2 - c_2^2 - c_3^2 - c_4^2 \quad (\text{E1a})$$

Substitute c_0^2 into Eq.(E2):

$$\begin{aligned} 7(1 - c_1^2 - c_2^2 - c_3^2 - c_4^2) + 3c_1^2 - c_2^2 - c_3^2 - 5c_4^2 = 0 \\ 7 - 7c_1^2 - 7c_2^2 - 7c_3^2 - 7c_4^2 + 3c_1^2 - c_2^2 - c_3^2 - 5c_4^2 = 0 \end{aligned}$$

Simplify:

$$-4c_1^2 - 8c_2^2 - 8c_3^2 - 12c_4^2 + 7 = 0 \quad (\text{E7})$$

Divide both sides by -1 :

$$4c_1^2 + 8c_2^2 + 8c_3^2 + 12c_4^2 = 7 \quad (\text{E7a})$$

Divide both sides by 4:

$$c_1^2 + 2c_2^2 + 2c_3^2 + 3c_4^2 = \frac{7}{4} \quad (\text{E8})$$

Step 4: Use Eq.(E6) to Express $c_0 c_4$

From Eq.(E6):

$$\sqrt{7} c_0 c_4 = -\sqrt{3} c_1 c_2 - 2\sqrt{3} c_1 c_3 \quad (\text{E9})$$

Step 5: Compute $(c_0c_4)^2$ from Eq.(E9)

Square both sides of Eq.(E9):

$$\begin{aligned} (\sqrt{7}c_0c_4)^2 &= \left(-\sqrt{3}c_1c_2 - 2\sqrt{3}c_1c_3\right)^2 \\ 7c_0^2c_4^2 &= 3c_1^2(c_2 + 2c_3)^2 \\ \Rightarrow c_0^2c_4^2 &= \frac{3}{7}c_1^2(c_2 + 2c_3)^2 \end{aligned} \quad (\text{E10})$$

Step 6: Express $c_0^2c_4^2$ in Terms of c_1 and c_4

From Eq.(E1a), we have:

$$c_0^2 = 1 - c_1^2 - c_2^2 - c_3^2 - c_4^2 \quad (\text{E1a})$$

Therefore:

$$c_0^2c_4^2 = (1 - c_1^2 - c_2^2 - c_3^2 - c_4^2)c_4^2 \quad (\text{E11})$$

Step 7: Equate the Two Expressions for $c_0^2c_4^2$

Set Eq.(E10) equal to Eq.(E11):

$$(1 - c_1^2 - c_2^2 - c_3^2 - c_4^2)c_4^2 = \frac{3}{7}c_1^2(c_2 + 2c_3)^2 \quad (\text{E12})$$

Step 8: Use Eq.(E8) to Express $c_2^2 + c_3^2$

From Eq.(E8):

$$c_2^2 + c_3^2 = \frac{1}{2} \left(\frac{7}{4} - c_1^2 - 3c_4^2 \right) \quad (\text{E13})$$

Simplify:

$$c_2^2 + c_3^2 = \frac{7}{8} - \frac{1}{2}c_1^2 - \frac{3}{2}c_4^2 \quad (\text{E14})$$

Step 9: Express $(c_2 + 2c_3)^2$ in Terms of Known Quantities

First, expand $(c_2 + 2c_3)^2$:

$$(c_2 + 2c_3)^2 = c_2^2 + 4c_2c_3 + 4c_3^2 \quad (\text{E15})$$

From Eq.(E5), rearranged:

$$c_2c_3 = -\sqrt{3}c_1c_4 - \frac{3}{4}c_3^2 \quad (\text{E16})$$

Substitute c_2c_3 into Eq.(E15):

$$\begin{aligned} (c_2 + 2c_3)^2 &= c_2^2 + 4 \left(-\sqrt{3}c_1c_4 - \frac{3}{4}c_3^2 \right) + 4c_3^2 \\ &= c_2^2 - 4\sqrt{3}c_1c_4 - 3c_3^2 + 4c_3^2 \\ &= c_2^2 - 4\sqrt{3}c_1c_4 + c_3^2 \end{aligned} \quad (\text{E17})$$

Now, using $c_2^2 + c_3^2$ from Eq.(E14):

$$(c_2 + 2c_3)^2 = \frac{7}{8} - \frac{1}{2}c_1^2 - \frac{3}{2}c_4^2 - 4\sqrt{3}c_1c_4 \quad (\text{E14})$$

Thus, from Eq.(E12), we have

$$\begin{aligned} \left(1 - c_1^2 - \left(\frac{7}{8} - \frac{1}{2}c_1^2 - \frac{3}{2}c_4^2 \right) - c_4^2 \right) c_4^2 \\ = \frac{3}{7}c_1^2 \left(\frac{7}{8} - \frac{1}{2}c_1^2 - \frac{3}{2}c_4^2 - 4\sqrt{3}c_1c_4 \right) \end{aligned} \quad (\text{E18})$$

Simplifying we have

$$\begin{aligned} 28c_4^4 + (7 + 8c_1^2)c_4^2 + 96\sqrt{3}c_1^3c_4 + (12c_1^4 - 21c_1^2) &= 0 \\ \Leftrightarrow \left(c_4 + \sqrt{3}c_1 \right) \left(28c_4^3 - 28\sqrt{3}c_1c_4^2 + (92c_1^2 + 7)c_4 \right. \\ \left. + \sqrt{3}(4c_1^3 - 7c_1) \right) &= 0 \end{aligned} \quad (\text{E19})$$

This then gives $c_4 = -\sqrt{3}c_1$ as a solution.

Appendix C: All ((7,2,3)) Stabilizer Codes

The codeword stabilized (CWS) formalism provides a systematic approach for enumerating all (7, 2, 3) stabilizer codes [10]. A CWS code is characterized by a graph and a classical binary code. We examined both connected and disconnected graphs corresponding to (7, 2, 3) stabilizer codes, taking into account graph isomorphisms and local unitary equivalence, as classified in [34]. This analysis results in 59 inequivalent graphs.

Assuming 0000000 is one of the classical codewords, we evaluated all 7-bit classical binary strings from 0000001 to 1111111 to identify the second codeword, selecting those that result in a CWS code with a minimum distance of 3. The resulting (7, 2, 3) stabilizer codes are summarized in Table I, which lists the corresponding lengths of the signature vectors λ^* .

The signature vector for a stabilizer code can only contain components of 0 or 1, which implies that the square of its norm, $(\lambda^*)^2$, must be an integer. An exhaustive search reveals that for (7, 2, 3) stabilizer codes, the only possible values of λ^* are $\{0, \sqrt{1}, \sqrt{2}, \sqrt{3}, \sqrt{5}\}$.

λ^*	Graph No.
0	40, 42, 43, 44
$\sqrt{1}$	38, 39, 41
$\sqrt{2}$	17, 30, 35, 37
$\sqrt{3}$	8, (1, 8), 28, 33
$\sqrt{5}$	(2, 4), 25, 31

TABLE I: Summary of all ((7,2,3)) stabilizer codes identified through the CWS formalism. Graph numbering follows [34]. For graphs with fewer than 7 vertices, isolated vertices were added, e.g., Graph No. 17. Notation (1, 8) refers to a disconnected graph composed of subgraphs No. 1 and No. 8.

-
- [1] A. R. Calderbank and P. W. Shor, Good quantum error-correcting codes exist, *Phys. Rev. A* **54**, 1098 (1996).
- [2] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information: 10th Anniversary Edition* (Cambridge University Press, 2010).
- [3] D. Gottesman, Stabilizer codes and quantum error correction, *arXiv: Quantum Physics* (1997).
- [4] A. M. Steane, Error correcting codes in quantum theory, *Phys. Rev. Lett.* **77**, 793 (1996).
- [5] D. A. Lidar and T. A. Brun, *Quantum Error Correction* (Cambridge University Press, 2013).
- [6] A. Y. Kitaev, Quantum computations: algorithms and error correction, *Russian Mathematical Surveys* **52**, 1191 (1997).
- [7] E. Knill and R. Laflamme, Theory of quantum error-correcting codes, *Phys. Rev. A* **55**, 900 (1997).
- [8] A. Calderbank, E. Rains, P. Shor, and N. Sloane, Quantum error correction via codes over $\text{GF}(4)$, *IEEE Transactions on Information Theory* **44**, 1369 (1998).
- [9] E. M. Rains, R. H. Hardin, P. W. Shor, and N. J. A. Sloane, A nonadditive quantum code, *Phys. Rev. Lett.* **79**, 953 (1997).
- [10] A. Cross, G. Smith, J. A. Smolin, and B. Zeng, Codeword stabilized quantum codes, in *2008 IEEE International Symposium on Information Theory* (2008) pp. 364–368.
- [11] D. A. Lidar and K. Birgitta Whaley, Decoherence-free subspaces and subsystems, in *Irreversible Quantum Dynamics*, edited by F. Benatti and R. Floreanini (Springer Berlin Heidelberg, Berlin, Heidelberg, 2003) pp. 83–120.
- [12] P. Sarvepalli and A. Klappenecker, Degenerate quantum codes and the quantum hamming bound, *Phys. Rev. A* **81**, 032318 (2010).
- [13] A. M. Steane, Simple quantum error-correcting codes, *Phys. Rev. A* **54**, 4741 (1996).
- [14] A. Steane, Multiple-particle interference and quantum error correction, *Proceedings of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences* **452**, 2551 (1996).
- [15] H. Pollatsek and M. B. Ruskai, Permutationally invariant codes for quantum error correction, *Linear Algebra and its Applications* **392**, 255 (2004).
- [16] D. A. Lidar, I. L. Chuang, and K. B. Whaley, Decoherence-free subspaces for quantum computation, *Phys. Rev. Lett.* **81**, 2594 (1998).
- [17] P. Shor and R. Laflamme, Quantum analog of the macwilliams identities for classical coding theory, *Phys. Rev. Lett.* **78**, 1600 (1997).
- [18] E. Rains, Quantum weight enumerators, *IEEE Transactions on Information Theory* **44**, 1388 (1998).
- [19] E. Rains, Quantum shadow enumerators, *IEEE Transactions on Information Theory* **45**, 2361 (1999).
- [20] D. Miller, K. Levi, L. Postler, A. Steiner, L. Bittel, G. A. White, Y. Tang, E. J. Kuehnke, A. A. Mele, S. Khatri, *et al.*, Experimental measurement and a physical interpretation of quantum shadow enumerators, *arXiv preprint arXiv:2408.16914* (2024).
- [21] O. Toeplitz, Das algebraische analogon zu einem satze von fejér, *Mathematische Zeitschrift* **2**, 187 (1918).
- [22] F. Hausdorff, Der wertvorrat einer bilinearform, *Mathematische Zeitschrift* **3**, 314 (1919).
- [23] H. J. Woerdeman, The higher rank numerical range is convex, *Linear and Multilinear Algebra* **56**, 65 (2008), <https://doi.org/10.1080/03081080701352211>.
- [24] C.-K. Li and N.-S. Sze, Canonical forms, higher rank numerical ranges, totally isotropic subspaces, and matrix equations, *Proc. Amer. Math. Soc.* **136**, 3013 (2008).
- [25] M.-D. Choi, D. W. Kribs, and K. Życzkowski, Higher-rank numerical ranges and compression problems, *Linear Algebra and its Applications* **418**, 828 (2006).
- [26] M.-D. Choi, M. Giesinger, J. A. Holbrook, and D. W. Kribs, Geometry of higher-rank numerical ranges, *Linear and Multilinear Algebra* **56**, 53 (2008), <https://doi.org/10.1080/03081080701336545>.
- [27] C.-K. Li, Y.-T. Poon, and N.-S. Sze, Higher rank numerical ranges and low rank perturbations of quantum channels, *Journal of Mathematical Analysis and Applications* **348**, 843 (2008).
- [28] C.-K. LI and Y.-T. POON, Generalized numerical ranges and quantum error correction, *Journal of Operator Theory* **66**, 335 (2011).
- [29] H.-L. Gau, C.-K. Li, Y.-T. Poon, and N.-S. Sze, Higher rank numerical ranges of normal matrices, *SIAM Journal on Matrix Analysis and Applications* **32**, 23 (2011), <https://doi.org/10.1137/09076430X>.
- [30] X. Zhu, C. Zhang, Z. An, and B. Zeng, *Unified framework for calculating convex roof resource measures* (2024), *arXiv:2406.19683 [quant-ph]*.
- [31] B. Shaw, M. M. Wilde, O. Oreshkov, I. Krensky, and

- D. A. Lidar, Encoding one logical qubit into six physical qubits, *Phys. Rev. A* **78**, 012337 (2008).
- [32] C. Cao, C. Zhang, Z. Wu, M. Grassl, and B. Zeng, Quantum variational learning for quantum error-correcting codes, *Quantum* **6**, 828 (2022).
- [33] M. Li, M. Gutiérrez, S. E. David, A. Hernandez, and K. R. Brown, Fault tolerance with bare ancillary qubits for a $[[7,1,3]]$ code, *Phys. Rev. A* **96**, 032341 (2017).
- [34] M. Hein, W. Dür, J. Eisert, R. Raussendorf, M. Van den Nest, and H.-J. Briegel, Entanglement in graph states and its applications, in *Quantum computers, algorithms and chaos* (IOS Press, 2006) pp. 115–218.