

# QUANTUM CELLULAR AUTOMATA AND CATEGORICAL DUALITIES OF SPIN CHAINS

COREY JONES<sup>1</sup>, KYLAN SCHATZ<sup>1</sup>, AND DOMINIC J. WILLIAMSON<sup>2,\*</sup>

ABSTRACT. Dualities play a central role in the study of quantum spin chains, providing insight into the structure of quantum phase diagrams and phase transitions. In this work we study categorical dualities, which are defined as bounded-spread isomorphisms between algebras of symmetry-respecting local operators on a spin chain. We consider generalized global symmetries that correspond to unitary fusion categories, which are represented by matrix-product operator algebras. A fundamental question about dualities is whether they can be extended to quantum cellular automata on the larger algebra generated by all local operators that respect the unit matrix-product operator. For conventional global symmetries, which are on-site representations of finite groups, this larger algebra is simply the tensor product of algebras associated to individual spins in the chain. We present a solution to the extension problem using the machinery of Doplicher-Haag-Roberts bimodules. Our solution provides a crisp categorical criterion for when an extension of a duality exists. We show that the set of possible extensions form a torsor over the invertible objects in the relevant symmetry category. Our approach recovers existing results from the literature when applied to quantum cellular automata that respect a conventional global symmetry.

## CONTENTS

1. Introduction	1
2. Dualities for spin chains with categorical symmetry	4
3. Abstract fusion spin chains	9
4. The algebra model for categorical inclusions	11
5. Solving the extension problem with DHR bimodules	13
6. Examples and Applications	16
Acknowledgments	18
References	18

## 1. INTRODUCTION

Quantum cellular automata (QCA) are a class of quantum operations that capture the fundamental properties of unitarity and locality [Wat95, Far20, Arr19]. The problem of classifying QCA up to finite depth unitary circuits has been the focus of much attention [GNVW09, SW04, FHH19, FH20, Haa21, Haa22], due to both the fundamental nature of QCA and their wide ranging applications. These include invertible quantum phases of matter [FHH20, JM21], floquet quantum dynamics [GNP21, PFM<sup>+</sup>16, PFVP17, PVF18, PM17, ZL21, AHLM23], many-body localization [LE24], unitary tensor network operators [CPGSV17, SSBC18, PC20, PTSC21, RWW20], and simulations of quantum field theories [ANW11, FS20, BM20].

<sup>1</sup> DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY, RALEIGH, NC 27695, USA

<sup>2</sup> SCHOOL OF PHYSICS, UNIVERSITY OF SYDNEY, SYDNEY, NEW SOUTH WALES 2006, AUSTRALIA

\* Current address: IBM Quantum, IBM Almaden Research Center, San Jose, CA 95120, USA.

Thus far, most work has focused on QCAs embodied by bounded-spread automorphisms on standard quasi-local operator algebras of spin systems  $A = \otimes_{\mathbb{Z}^n} M_d(\mathbb{C})$ , formed by taking the tensor product of finite qudit degrees of freedom on the sites of a lattice. This has led to a full classification of QCAs in one spatial dimension [GNVW09] and partial classifications in higher spatial dimensions [FH20, Haa21, Haa22]. There is a closely related problem of understanding *dualities*, which in this context are mathematically described by bounded-spread isomorphisms between *subalgebras* of the usual quasi-local algebra that interchange quantum phases of matter. Classification problems in this direction remain largely open (see [AMF16, TW19, CCH+21, KOZ22, LDOV23, Jon24, JL24, BDSNY24] for recent progress), and it is these dualities in one spatial dimension that form the primary focus of this work.

Dualities arise naturally in the presence of a global symmetry  $\mathcal{C} \curvearrowright A$ , which may correspond to a group, (weak) Hopf algebra, or fusion category. If a pair of Hamiltonians has symmetric local terms, then these lie in the algebra  $A^{\mathcal{C}} \subseteq A$  of symmetric operators. In this work, we use the term *duality* to formally denote a bounded-spread isomorphism between subalgebras of symmetric operators  $A^{\mathcal{C}}$  under some global symmetry which exchange Hamiltonians *but may not extend* to bounded-spread automorphisms of the whole quasi-local algebra  $A$ , also known as QCA. The motivating example for this class of dualities is due to Kramers and Wannier, originally introduced in the context of classical statistical mechanics [KW41]. The bounded-spread automorphism implementing KW duality acts on the symmetric subalgebra of operators on a chain of qubits that commute with the global  $\mathbb{Z}_2$  symmetry represented by  $\prod_i \sigma_i^x$ . The action of the Kramers-Wannier (KW) duality can be specified on a set of generators of this  $\mathbb{Z}_2$ -symmetric operator algebra via the map  $\sigma_i^x \mapsto \sigma_{i-1}^z \sigma_i^z$  and  $\sigma_i^z \sigma_{i+1}^z \mapsto \sigma_i^x$ .

In recent years, there has been significant interest in extending results from the setting of conventional global symmetries, which are represented by tensor products of single-site unitary operators, to categorical global symmetries, which are represented by matrix product operators (MPOs) [FFRS04, BMW+17, VBW+18, AMF16, TW19]. The resulting algebra generated by symmetric local operators is called a *fusion spin chain*. Recently, the idea of KW duality has been generalized in this direction to categorical symmetries, where matrix-product operators have been found to implement dualities based on any invertible bimodule categories between fusion categories [AMF16, WBV17, AFM20, LDOV23]. This suggests a deep connection between categorical dualities and the algebraic structure of fusion categories.

A key feature of non-trivial dualities following the KW paradigm is that bounded-spread isomorphisms of the symmetric operators *should not be* locally extendable to the whole spin chain (or in the case of non-unital MPOs, to the local edge-restricted sector). We call such an extension a *spatial implementation*. Spatial implementations are themselves ordinary 1+1D QCAs in the case of group or Hopf algebra symmetries. More generally, spatial implementations are “edge-restricted” QCA. These extensions lead naturally to the following question, whose answer sheds light on the relationship between QCA and categorical dualities:

**Question 1.1.** *When do bounded-spread isomorphisms of symmetric local operator algebras (i.e. categorical dualities) extend to QCA defined on the full (or edge-restricted) local operator algebras, and how are the possible extensions characterized?*

To answer this question, we first establish a precise reformulation in terms of *abstract fusion spin chains*  $A(\mathcal{E}, X)$ , built from a fusion category  $\mathcal{E}$  and an object  $X \in \mathcal{E}$ , by defining local algebras as endomorphism algebras of tensor powers of  $X$  (see Section 3). A fusion category  $\mathcal{C}$  acting on a spin chain via MPOs is specified by the data of an indecomposable right  $\mathcal{C}$ -module category  $\mathcal{M}$ , and an object  $X \in \mathcal{C}_{\mathcal{M}}^*$ . Then, the symmetric operator algebra is isomorphic to the fusion spin chain  $A(\mathcal{C}_{\mathcal{M}}^*, X)$ . If we view  $\mathcal{M}$  as a left  $\mathcal{C}_{\mathcal{M}}^*$ -module category, then rewriting  $\mathcal{E} := \mathcal{D}_{\mathcal{M}}^*$  we can reformulate the data of a categorical symmetry as an abstract fusion spin chain  $A(\mathcal{C}, X)$  together

with an indecomposable left  $\mathcal{C}$ -module category  $\mathcal{M}$ . The inclusion  $A(\mathcal{E}, X) \hookrightarrow A(\mathcal{E}, X)_{\mathcal{M}}$  (see Definition 3.3) corresponds to the inclusion of symmetric operators into all (edge-restricted) local operators, which we call a *spatial realization* of the chain  $A(\mathcal{E}, X)$ . Question 1.1 can be given a precise mathematical formulation as follows:

**Question 1.2.** *Given a bounded-spread isomorphism  $\alpha : A(\mathcal{C}, X) \rightarrow A(\mathcal{D}, Y)$  when does it extend to a QCA between spatial realizations  $A(\mathcal{C}, X)_{\mathcal{M}} \rightarrow A(\mathcal{D}, Y)_{\mathcal{N}}$ , and how are these extensions characterized?*

In this form, we can address the problem using the machinery of Doplicher-Haag-Roberts (DHR) bimodules, introduced in [Jon24] and generalizing the earlier work of [NS95]. Associated to an abstract spin system  $A$  is a braided  $C^*$ -tensor category  $\text{DHR}(A)$ , and associated to any bounded-spread isomorphism  $\alpha : A \rightarrow B$  is a braided equivalence  $\text{DHR}(\alpha) : \text{DHR}(A) \cong \text{DHR}(B)$ . Intuitively, if we view an abstract spin chain as operators living at a cut boundary of a 2+1D topologically ordered spin system (as in [JNPW23]) then DHR bimodules capture the structure of the bulk topological order (see also [CW23, IW23]). For a fusion spin chain  $A(\mathcal{C}, X)$ , there is a braided equivalence  $\text{DHR}(A(\mathcal{C}, X)) \cong \mathcal{Z}(\mathcal{C})$ , where the latter denotes the Drinfeld center of  $\mathcal{C}$ . We now set up the statement of our main theorem. For any indecomposable  $\mathcal{C}$ -module category  $\mathcal{M}$ , let  $Z(\mathcal{M}) \in \mathcal{Z}(\mathcal{C})$  denote the canonically associated Lagrangian algebra [Dav10]. Then we have the following (which is a special case of Theorem 5.3):

**Theorem 1.3.** *Let  $\alpha : A(\mathcal{C}, X) \rightarrow A(\mathcal{D}, Y)$  be a bounded-spread isomorphism between abstract fusion spin chains. Then for any indecomposable module categories  $\mathcal{M}$  and  $\mathcal{N}$  of  $\mathcal{C}$  and  $\mathcal{D}$  respectively, spatial implementations  $A(\mathcal{C}, X)_{\mathcal{M}} \rightarrow A(\mathcal{D}, Y)_{\mathcal{N}}$  of  $\alpha$  are in bijective correspondence with algebra isomorphisms  $\text{DHR}(\alpha)(Z(\mathcal{M})) \cong Z(\mathcal{N})$  in  $\mathcal{Z}(\mathcal{D})$ . As a consequence:*

- (1) *If  $\text{DHR}(\alpha)(Z(\mathcal{M}))$  is not isomorphic to  $Z(\mathcal{N})$ , then  $\alpha$  has no spatial implementation.*
- (2) *If  $\text{DHR}(\alpha)(Z(\mathcal{M})) \cong Z(\mathcal{N})$ , then the spatial implementations of  $\alpha$  form a torsor over  $\text{Aut}(Z(\mathcal{M})) \cong \text{Inv}(\mathcal{C}_{\mathcal{M}}^*)$*

For the Kramers-Wannier example described above, the fusion category is  $\mathcal{C} = \mathcal{D} = \text{Rep}(\mathbb{Z}_2)$ ,  $X$  is the left regular representation of  $\mathbb{Z}_2$ , and the DHR category of the symmetric spin chain is the toric code modular tensor category  $\mathcal{Z}(\text{Hilb}_{f.d.})$ . The module category in question is the standard fiber functor, which corresponds to the electric Lagrangian  $L = 1 \oplus e$ . The induced action of the Kramer-Wannier automorphism  $\alpha$  is the  $e \leftrightarrow m$  swap symmetry, which does not preserve  $L$ . By the above theorem,  $\alpha$  admits *no* bounded-spread extension to the whole quasi-local algebra  $A = \otimes_{\mathbb{Z}} M_2(\mathbb{C})$ .

Finally, we can use our formalism to make contact with the literature concerning symmetric QCA (sQCA). In this setting, we consider a global on-site finite group  $G$  acting by unitaries. We have the following (c.f. [GSSC20]) as a consequence of our machinery (see Theorem 6.3):

**Theorem 1.4.** *There is a homomorphism  $\pi : sQCA \rightarrow H^2(G, U(1))$ .*

Our results on the extension of dualities to QCAs point to a number of directions for future work. First, what is the full classification of bounded-spread isomorphisms on local algebras and similarly what is the classification of edge-restricted QCAs in one spatial dimension? Second, can our results be applied to characterize symmetries of topological orders in two dimensions via their action on the boundary? Third, can techniques from the theory of operator algebras, subfactors, and fusion categories be applied to classify dualities and QCAs in higher spatial dimensions?

The remainder of this work is presented as follows. In section 2 we introduce background material about categorical dualities of quantum spin chains. In section 3 we describe dualities in the setting of fusion spin chains. In section 4 we describe categorical inclusions defined by module categories

over algebra objects. In section 5 we describe how DHR bimodules can be applied to solve the extension problem for dualities of quantum spin chains. In section 6 we present examples that generalize the KW duality.

## 2. DUALITIES FOR SPIN CHAINS WITH CATEGORICAL SYMMETRY

In this section we introduce relevant background material. First, we briefly recall the algebraic approach to spin chains [BR97]. Consider  $\mathbb{Z} \subseteq \mathbb{R}$  as a metric space. A spin chain is specified by choosing an on-site Hilbert space  $\mathbb{C}^d$ ; for every interval  $I \subseteq \mathbb{Z}$ , we have the local Hilbert space  $\otimes_{x \in I} \mathbb{C}^d$  and the algebra of operators localized in  $I$ :

$$A_I := \otimes_{x \in I} M_d(\mathbb{C}).$$

For  $I \subseteq J$ , we have natural inclusions  $A_I \hookrightarrow A_J$  defined by  $a \mapsto 1^{J < I} \otimes a \otimes 1^{J > I}$ . The  $\mathbb{C}^*$ -completion of the union over all intervals (or more formally, the colimit in the category of  $\mathbb{C}^*$ -algebras) is called the *quasi-local algebra*. We call the *algebraic union* the local algebra. States of the system in the thermodynamic limit are normalized, positive linear functionals on  $A$ , while Hamiltonians can be expressed as (unbounded) derivations on  $A$  (see [BR87, BR97]).

A *local Hamiltonian* is an assignment of a self-adjoint element  $H_I \in A_I$  to every interval  $I \subseteq \mathbb{Z}$  for which there exists an  $R \geq 0$  such that if  $I$  has length greater than  $R$ , then  $H_I = 0$ . This defines an (unbounded) derivation on the local algebra:

$$\delta_H(a) := \sum_I [H_I, a],$$

which is well-defined since the above summation contains only finitely many non-zero terms.

The primary motivation for this paper is to understand a class of *categorical dualities* between spin systems. Broadly speaking, a duality between two physical theories is a mapping between the observables of the theories that intertwines the relevant physics (state, dynamics, etc.). However, interesting dualities should be equivalences that are in some salient sense “not obvious”. In the case of spin systems dualities are generally considered in the context of symmetries, whereby we have a locality-preserving equivalence on the symmetric subsystems that does not extend to an equivalence of the whole theory (for example, see [ECN11] or for the categorical context [LDOV23, AMF16]) We give a mathematical formulation of this in 1D in terms of fusion spin chains.

First we set up some basic terminology. We need a natural notion of a locality-preserving equivalence of spin systems, which is captured by the idea of bounded-spread isomorphisms (or quantum cellular automata).

**Definition 2.1.** If  $A, B$  are the local algebras of spin chains on  $\mathbb{Z}$ , a *bounded-spread isomorphism* is a  $\mathbb{C}^*$ -isomorphism  $\alpha : A \rightarrow B$  such that there exists an  $R > 0$  so that  $\alpha(A_I) \subseteq B_{I+R}$  for every interval  $I$ . A bounded-spread isomorphism  $\alpha : A \rightarrow A$  is called a *quantum cellular automaton* (QCA).

In the above definition  $I^{+R}$  denotes the  $R$ -neighborhood of the interval  $I$  in  $\mathbb{Z}$ . A bounded-spread isomorphism is a natural mapping between the observables of the two spin chains that *uniformly* preserves locality.

**Definition 2.2.** If  $H, K$  are local Hamiltonians on spin chains with local algebras  $A$  and  $B$  respectively, an equivalence between them is a bounded-spread isomorphism  $\alpha : A \rightarrow B$  such that  $\alpha(\delta_H(\alpha^{-1}(b))) = \delta_K(b)$  for all local operators  $b \in B$ . We denote this by  $\alpha(H) = K$ .

By the definition of bounded-spread isomorphism, for any local Hamiltonian  $H = \{H_I\}$  on  $A$ ,  $\alpha(H) := \{\alpha(H)_I := \alpha(H_{I-R})\}$  is a local Hamiltonian on  $B$  and  $\alpha$  implements an equivalence between these. Thus given any bounded-spread isomorphism  $\alpha : A \rightarrow B$  and any local Hamiltonian

$H$  on  $A$ ,  $\alpha$  always implements an equivalence between  $H$  and *some* local Hamiltonian on  $B$ . This suggests that it could be useful to shift our focus to the groupoid of local algebras of spin chains and bounded-spread isomorphisms between them.

**Example: Finite depth quantum circuits.** An obvious class of QCA are the *finite-depth quantum circuits*, (abbreviated FDQC). Recall a depth-one circuit is given by specifying a uniformly bounded partition  $\mathcal{P}$  of  $\mathbb{Z}$  into intervals, and for  $I \in \mathcal{P}$ , choosing a unitary  $u_I \in A_I$ . Then define a QCA acting on the local operator  $x$  by

$$\alpha(x) := \left( \prod_{I \in \mathcal{P}} u_I \right) x \left( \prod_{I \in \mathcal{P}} u_I^* \right).$$

Note that by locality, this infinite product is in fact finite since all but finite many of the  $u_I$  commute with  $x$ .

A *finite-depth quantum circuit* (FDQC) is a finite composition of depth one circuits. FDQC are QCA that can be locally implemented. It is easy to show that FDQC is a normal subgroup of QCA. FDQC equivalence for states is used to characterize long-range entanglement classes [CGW10], and thus we can view QCA/FDQC as long-range entanglement classes of uniformly locality-preserving equivalences [Haa22]. Since FDQC are locally implemented, they are often considered “trivial” as equivalences of Hamiltonians, which naturally leads us to consider the group  $QCA/FDQC$ , which has been frequently studied in the literature.

Before moving on to giving a mathematical definition for our Kramers-Wannier type dualities, it is useful to have in hand the notion of an *abstract spin chain*. This is strongly motivated by the Haag-Kastler axioms for an algebraic quantum field theory [HK64], and is a straightforward generalization of the properties satisfied by the quasi-local algebra of a spin system [BR97].

**Definition 2.3.** An abstract spin chain is a unital  $C^*$ -algebra  $A$  and for every interval  $I$ , a unital subalgebra  $A_I \subseteq A$  such that

- (1) (Isotony) If  $I \subseteq J$ ,  $A_I^\circ \subseteq A_J^\circ$
- (2) (Locality) If  $I \cap J = \emptyset$ , then  $[A_I^\circ, A_J^\circ] = 0$
- (3) (Density)  $(\cup A_I)^{\|\cdot\|} = A$
- (4) (Weak algebraic Haag duality) There is some  $R \geq 0$  such that for every interval  $I$ ,  $\{x \in A : [x, y] = 0 \text{ for all } y \in A_J, J \subseteq I^c\} \subseteq A_{I+R}$ .

If the last condition is satisfied with  $R = 0$ , it is called *algebraic Haag duality*. The abstract spin chains of interest in this paper always satisfy this stronger version of the last axiom. For two abstract spin chains, the definition of *bounded-spread isomorphism* used above makes perfect sense unchanged. We note that if  $\alpha$  is bounded spread, then it follows from weak algebraic Haag duality that  $\alpha^{-1}$  is as well.

**Actions of fusion categories on spin chains.** We are now ready to describe dualities, in the same sense as [LDOV23]. Suppose we have two spin systems with local algebras  $A$  and  $B$ , both with a global symmetry  $\mathcal{C} \curvearrowright A$  and  $\mathcal{D} \curvearrowright B$  (think of an on-site group symmetry, or more generally, a categorical symmetry determined by an MPO, see below). Suppose further that the local terms of the Hamiltonians  $H$  and  $K$  are both symmetric. Then we can restrict our attention to the symmetric operators, which consist of subalgebras  $A^{\mathcal{C}} \subseteq A$  and  $B^{\mathcal{D}} \subseteq B$  containing the terms of  $H$  and  $K$  respectively. We can then ask for bounded-spread isomorphisms *between the subalgebras*  $A^{\mathcal{C}}$  and  $B^{\mathcal{D}}$  which intertwine  $H$  and  $K$ . This is what we call a duality.

First, we formalize what we mean by a fusion categorical symmetry and the algebras of symmetric operators, resulting in the definition of fusion spin chains and duality. Recall a unitary fusion category is a unitary tensor category (in the sense of [ENO02]) with finitely many isomorphism

classes of simple objects. For further definitions concerning fusion categories, modules categories, and related concepts, see [EGNO15].

**Definition 2.4.** Let  $A$  be a spin system with local Hilbert space  $\mathbb{C}^d$  and let  $\mathcal{C}$  be a unitary fusion category. An action of  $\mathcal{C}$  on  $A$  consists of

- (1) An indecomposable, semi-simple right module category  $\mathcal{M}$  of  $\mathcal{C}$ .
- (2) An object  $X \in \mathcal{C}_{\mathcal{M}}^*$  which is *strongly tensor generating* for  $\mathcal{C}_{\mathcal{M}}^*$
- (3) A unitary isomorphism  $\bigoplus_{i,j \in \text{Irr}(\mathcal{M})} \mathcal{M}(i, X \triangleright j) \cong \mathbb{C}^d$ .

The action is called *unital* if  $\mathcal{M}$  is equivalent to  $\text{Hilb}_{f,d}$  as  $\mathbb{C}^*$ -categories.

From this data, we can construct *matrix product operators* which act by quantum channels on (a sector of) the spin system, implementing the fusion rules of  $\mathcal{C}$ .

In general a matrix product operator is defined from a 4-valent tensor

$$A_{\alpha\beta}^{ij} := \alpha \text{---} \boxed{M} \text{---} \beta \quad \begin{array}{c} i \\ | \\ \beta \\ | \\ j \end{array} \quad (2.5)$$

by

$$\sum_{\{i\}\{j\}} \text{tr}(A^{i_1 j_1} A^{i_2 j_2} \dots A^{i_n j_n}) |\{i\}\rangle \langle \{j\}| \quad (2.6)$$

where

$$\text{tr}(A^{i_1 j_1} A^{i_2 j_2} \dots A^{i_n j_n}) = \dots \text{---} \begin{array}{c} i_1 \\ | \\ \boxed{M} \\ | \\ j_1 \end{array} \text{---} \begin{array}{c} i_2 \\ | \\ \boxed{M} \\ | \\ j_2 \end{array} \text{---} \begin{array}{c} i_3 \\ | \\ \boxed{M} \\ | \\ j_3 \end{array} \text{---} \begin{array}{c} i_4 \\ | \\ \boxed{M} \\ | \\ j_4 \end{array} \text{---} \begin{array}{c} i_5 \\ | \\ \boxed{M} \\ | \\ j_5 \end{array} \text{---} \dots \quad (2.7)$$

where we treat  $A_{\alpha\beta}^{ij}$  as a matrix element of  $A^{ij}$  and take a matrix product over the  $\alpha, \beta$  indices of the tensors  $A_{\alpha\beta}^{ij}$ . From the data in Definition 2.4, one can construct for each simple object in  $\mathcal{C}$  a matrix product operator, satisfying a family of categorical relations [SWB<sup>+</sup>14, BMW<sup>+</sup>17]. It is in this sense that a fusion category is actually acting on the spin chain.

**Edge-restricted algebra.** For non-unital fusion categorical symmetries, it is natural to consider a non-unital “subalgebra” of  $A$ , called the *edge-restricted algebra*. If we pick an orthonormal basis  $E_{i,j}$  for the space  $\mathcal{M}(i, X \triangleleft j)$ , we can define a finite graph  $\mathcal{G}$  whose vertices are  $\text{Irr}(\mathcal{M})$ , and the edges from  $a \rightarrow b$  are defined to be  $E_{a,b}$ . Then using the isomorphism between the onsite Hilbert space  $H_i$  and  $\bigoplus_{i,j \in \text{Irr}(\mathcal{M})} \mathcal{M}(i, X \triangleleft j) \cong \mathbb{C}^d$ , we have a distinguished basis of  $H_m$  indexed by the edges of  $\mathcal{G}$ , for every  $m \in \mathbb{Z}$

Then for each pair of adjacent sites  $(i, i+1)$ , consider the local projection  $P_i \in A_{[i, i+1]}$  defined by  $P_i(e \otimes f) = \delta_{i(e)=s(f)} e \otimes f$ , which checks the adjacency of the edges  $e$  and  $f$  in the graph  $\mathcal{G}$ . Note that  $[P_i, P_j] = 0$  for all  $i$  and  $j$ , and thus we can (unambiguously) consider the projection

$$P := \prod_{i \in \mathbb{Z}} P_i.$$

We have two immediate observations:

- (1) This projection in general does not lie in the quasi-local C\*-algebra  $A$ . However, if we pick a state  $\phi$  on  $A$  such that  $\phi(P_i \cdot P_i) = \phi$  for all  $i$ , then  $P$  does define an operator on the GNS Hilbert space  $L^2(A, \phi)$  in the von Neumann completion of  $A$ . If we wish to avoid picking a state, we can view  $P$  as an element of the von Neumann algebra  $A^{**}$ , which is the von Neumann completion of  $A$  in the universal Hilbert space representation of  $A$  [Tak79].
- (2) If we consider a finite chain on an interval  $I$ , then the action of the restricted version of  $P$ ,  $P_I$  on  $H_I$  agrees with the action of the MPO assigned to the unit object  $\mathbb{1} \in \mathcal{C}$  away from the boundary. For this reason, it is reasonable to call the image of  $P$  in a Hilbert space representation the *unit MPO sector*. This is the sector on which the categorical MPO operators satisfy the required categorical relations [BMW<sup>+</sup>17].

We can consider the sector ‘‘corner subalgebra’’ of  $A$ , defined by  $A^\circ := PAP$  (in  $A^{**}$ ), with local subalgebras  $A_I^\circ := PA_I P$ . Technically speaking, the elements  $A_I^\circ$  are not properly elements of the algebra  $A$  or even its C\*-completion, but rather they are elements of the von Neumann completion of  $A$  (either in the double dual von Neumann algebra  $A^{**}$ , or in any faithful Hilbert space representation of  $A$  where  $P$  does not act by 0).

We can easily identify  $A_I^\circ$  as follows: If  $I = [i, j]$  with  $j - i = n$ , let  $L_I^{i,j}$  be the Hilbert space with orthonormal basis given by the paths of length  $n$  in  $\mathcal{G}$  starting at  $i$  and ending at  $j$ . Let  $d_{i,j}^n$  denote the number of length  $n$  paths from  $i$  to  $j$  so that  $d_n^{i,j} = \dim(L_I^{i,j})$ .

Then we have an isomorphism of Hilbert spaces to

$$\left( \prod_{a \leq i < b} P_i \right) (H_I) \cong \bigoplus_{i,j \in \mathcal{G}} L_I^{i,j}.$$

It is easy to see from the definitions that

$$A_I^\circ \cong \bigoplus_{i,j \in \text{Irr}(\mathcal{M})} \text{Mat}(L_I^{i,j})$$

where the summand  $\text{Mat}_{d_n^{i,j}}(\mathbb{C})$  is identified with all the linear operators on the Hilbert space  $L_I^{i,j}$ . In other words,  $A_I^\circ$  is isomorphic to the subalgebra of operators on  $L_I$  that satisfy the property that the coefficient of the matrix unit  $e_{p,q}$  is 0 if  $s(p) \neq s(q)$  or  $t(p) \neq t(q)$ .

Note that from this description,  $A_I^\circ$  is isomorphic to a (non-unital) subalgebra of  $A_I$ , but not coherently. Nevertheless,  $A^\circ$  has the structure of an abstract spin chain. All of the axioms are obvious expect for weak algebraic Haag duality, which in fact follows from Lemma 4.3 below.

**Symmetric algebra.** The operators of interest to us are the symmetric local operators. To define these, for each  $Y \in \text{Irr}(\mathcal{C})$ , and interval  $J$ , let  $M_J^Y$  denote the associated matrix product operator, acting on (a sufficiently large) edge-restricted Hilbert space with periodic boundary conditions

$$L_J^{per} := \bigoplus_{i,j \in \text{Irr}(\mathcal{M})} L_J^{i,i}$$

For an interval  $I \ll J$ , we have a unital, faithful representation  $A_I^\circ \subseteq \text{Mat}(L_J^{per})$ . Then we define

$$A_I^{\mathcal{C}} := \{x \in A_I^\circ : [M_J^Y, x] = 0\}.$$

Notice that since the MPO are defined locally, this doesn't depend on the choice of ambient interval for  $J \gg I$ . We also note that we need our graph  $\mathcal{G}$  to be sufficiently well-connected, which is guaranteed by the indecomposability and strongly tensor generating assumptions for  $X \in \mathcal{C}_{\mathcal{M}}^*$ .

For  $I \subseteq J$ , we have natural inclusions  $A_I^{\mathcal{C}} \hookrightarrow A_J^{\mathcal{C}}$ , and taking the colimit in the category of  $C^*$ -algebras, we obtain the algebra  $A^{\mathcal{C}}$ . Identifying  $A_I^{\mathcal{C}}$  with its image in the colimit gives an abstract spin system. We call the algebra  $A^{\mathcal{C}}$  the *symmetric algebra*.

We can now formally define a version of duality for Hamiltonians generalizing Kramers-Wannier duality to the categorical setting, following [LDOV23]. We can then state the problem that motivates this work.

**Definition 2.8.** Let  $H = \{H_I\}$  be a local Hamiltonian on the spin system  $A$ , and let  $\mathcal{C}$  be a fusion category acting on  $A$ . Then we say  $H$  is  $\mathcal{C}$ -symmetric if for every interval  $I$  with  $H_I \neq 0$ ,  $H_I P = P H_I \neq 0$  and  $H_I P \in A^{\mathcal{C}_I}$ .

In this case the collection  $H^\circ := \{H_I P\}$  actually defines a local Hamiltonian internally to the symmetric chain  $A^{\mathcal{C}}$ , which we denote  $H^{\mathcal{C}} := \{H_I^{\mathcal{C}}\}$

**Definition 2.9.** Let  $A$  and  $B$  be spin chains with local Hamiltonians  $H$  and  $K$  respectively. A *categorical duality* between  $(A, K)$  and  $(B, K)$  consists of:

- (1) Categorical symmetries  $\mathcal{C} \curvearrowright A$ ,  $\mathcal{D} \curvearrowright B$  such that  $H$  is  $\mathcal{C}$ -symmetric and  $K$  is  $\mathcal{D}$ -symmetric
- (2) A bounded-spread isomorphism  $\alpha : A^{\mathcal{C}} \rightarrow B^{\mathcal{D}}$  such that  $\alpha(H^{\mathcal{C}}) = K^{\mathcal{D}}$  (in the sense of Definition 2.2, but restricted to  $A^{\mathcal{C}}$ ).

**Motivating Example: Kramers-Wannier.** Here we spell out the motivating example of Kramers-Wannier duality [KW41]. In this case, the acting fusion category is  $\mathcal{C} = \text{Hilb}_{f.d.}(\mathbb{Z}_2)$ . The module category is  $\mathcal{M} = \text{Hilb}_{f.d.}$  equipped with the trivial  $\text{Hilb}_{f.d.}(\mathbb{Z}_2)$  action, and the dual object  $X = \mathbb{C}[\mathbb{Z}_2] \in \text{Rep}(\mathbb{Z}_2)$  is the left regular representation.

This translates into the standard on-site spin flip action of  $\mathbb{Z}_2$  a chain of qubits  $\otimes_{\mathbb{Z}} \mathbb{C}^2$ , so that  $A = \otimes_{\mathbb{Z}} M_2(\mathbb{C})$ . The non-trivial symmetry operator (or MPO) can be represented by the product of on-site operators (i.e. with trivial valence bonds) as

$$U_g := \prod \sigma_i^x.$$

In this case, since  $\mathcal{M}$  is rank one, the action is unital, so that the edge-restricted algebra and the quasi-local algebra of the entire spin system coincide,  $A^\circ = A$ .

The symmetric algebra is the  $C^*$ -algebra generated by the operators  $\sigma_i^x$  and  $\sigma_i^z \sigma_{i+1}^z$  as  $i$  ranges over  $\mathbb{Z}$ , or more formally

$$A^{\mathbb{Z}_2} = C^* - \text{alg}\{\sigma_i^x, \sigma_i^z \sigma_{i+1}^z : i \in \mathbb{Z}\}.$$

Then consider the Ising Hamiltonian

$$H = \sum_{i \in \mathbb{Z}} J \sigma_i^z \sigma_{i+1}^z + h \sigma_i^x,$$

which is  $\mathbb{Z}_2$ -symmetric according to our definition, since all local terms lie in the subalgebra  $A^{\mathbb{Z}_2}$ . Now, the Kramers-Wannier duality is defined by

$$\alpha(\sigma_i^x) = \sigma_{i-1}^z \sigma_i^z \quad \text{and} \quad \alpha(\sigma_i^z \sigma_{i+1}^z) = \sigma_i^x,$$

which extends to an isomorphism  $\alpha : A^{\mathbb{Z}_2} \rightarrow A^{\mathbb{Z}_2}$ . We see immediately from the definition

$$\alpha(H) = H.$$

A key feature of KW-duality is that the equivalence between theories *can only be witnessed* at the level of symmetric operators, and does not extend to a (uniformly locality-preserving) equivalence of the whole spin system [ECN11]. Alternatively, some authors say that any unitary extension does not preserve locality, though this is not rigorously defined in the algebraic formalism for the thermodynamic limit we are using here.

This leads us to the following definition.

**Definition 2.10.** A bounded-spread isomorphism  $\alpha : A^{\mathcal{C}} \rightarrow B^{\mathcal{D}}$  is called *spatially implemented* if it admits an extension to a bounded-spread isomorphism  $\tilde{\alpha} : A^{\circ} \rightarrow B^{\circ}$ .

The idea is that spatially implemented bounded-spread isomorphisms are not proper dualities in the sense of KW, and are actually more of a standard equivalence. This is not literally true on the nose in full generality, since the  $\tilde{\alpha} : A^{\circ} \rightarrow B^{\circ}$  generally does not extend to  $A \rightarrow B$ , and thus more properly we would say that  $\tilde{\alpha} : A^{\circ} \rightarrow B^{\circ}$  such that  $\tilde{\alpha}(H^{\mathcal{C}}) = K^{\mathcal{D}}$  is an *emergent equivalence* [ECN11]. Some may view an emergent equivalence as a kind of duality, but for us we regard it as closer to an equivalence than a KW-type duality. Indeed, if  $A^{\circ} = A$  and  $B^{\circ} = B$  (or in other words, the categorical action is unital, or equivalently, the module categories correspond to fiber functors), then the emergent equivalence reduces to an ordinary equivalence.

In any case, we can now give a precise mathematical statement of Question 1.1:

**Reformulation of Question 1.1.** Given a bounded-spread isomorphism  $\alpha : A^{\mathcal{C}} \rightarrow B^{\mathcal{D}}$ , can we characterize when it is spatially implemented?

In the sections that follow, we present a precise and abstract statement of this problem, and a solution using the DHR bimodule category of fusion spin chains.

### 3. ABSTRACT FUSION SPIN CHAINS

In the previous section, we set up the problem of studying bounded-spread isomorphisms between algebras of operators invariant under a categorical symmetry (i.e.  $A^{\mathcal{C}}$ ), and asked when this can be extended to the edge-restricted algebra (i.e.  $A^{\circ}$ ). In this section, we present an abstract formulation of this problem in terms of fusion spin chains. We refer the reader to [EGNO15] for definitions of fusion categories and related objects. For convenience, we assume all our fusion categories are strict.

**Fusion spin chains.** Given an indecomposable multi-fusion category  $\mathcal{E}$  and an object  $X \in \mathcal{E}$  (not necessarily simple) we construct an abstract spin chain  $A(\mathcal{E}, X)$  as follows:

- For each interval  $I$ , define  $A(\mathcal{C}, X)_I := \mathcal{E}(X^{\otimes |I|}, X^{\otimes |I|})$
- If  $I \subseteq J$ , then we define the inclusion  $A(\mathcal{E}, X)_I \hookrightarrow A(\mathcal{E}, X)_J$  by

$$f \mapsto (1_X)^{\otimes J < I} \otimes f \otimes (1_X)^{\otimes J > I}$$

- We set  $A(\mathcal{E}, X) = \text{colim}_I A(\mathcal{E}, X)_I$ , where the colimit is taken in the category of  $*$ -algebras.

The commutativity property follows from functoriality of the monoidal product. Abstract spin chains constructed as above are called *fusion spin chains* [Jon24, JL24].

It turns out the three examples of abstract spin chains we saw in the previous section can be realized as fusion spin chains:

- (1) **Concrete spin systems.** Choose the fusion category  $\mathcal{E} = \text{Hilb}_{f.d.}$  of finite-dimensional Hilbert spaces, and pick the object  $X := \mathbb{C}^d$ . Then  $A(\text{Hilb}_{f.d.}, X)$  can be identified with the ordinary spin chain  $A$  constructed from  $\mathbb{C}^d$ .
- (2) **Edge-restricted algebras.** Consider a fusion category action on an ordinary spin system  $\mathcal{C} \curvearrowright A$ , and let  $X \in \mathcal{C}_{\mathcal{M}}^*$ . This defines an object in the multi-fusion category  $\tilde{X} \in \mathcal{E} = \text{Mat}_m(\text{Hilb}_{f.d.})$ , where  $m$  is the rank of  $\mathcal{M}$ . This is the multi-fusion category consisting of  $m \times m$  matrices of finite-dimensional Hilbert spaces, with  $\otimes$  given by “matrix-multiplication” [FHP20]. The object  $\tilde{X}$  is defined by
$$\tilde{X}_{i,j} := \mathcal{M}(i, X \triangleleft j)$$
with Hilbert space structure on the components given by the composition inner product. It is easy to see that the edge-restricted abstract spin chain  $A^\circ$  associated with our action  $\mathcal{E} \curvearrowright A$  can be identified with the fusion spin chain  $A(\text{Mat}_m(\text{Hilb}_{f.d.}), \tilde{X})$ .
- (3) **Symmetric algebras** Again, suppose we have a fusion category  $\mathcal{C} \curvearrowright A$ . Then we can identify  $A^{\mathcal{C}}$  with the fusion spin chain  $A(\mathcal{C}_{\mathcal{M}}^*, X)$  [Kaw22, Kaw21].

This leads us to define the following groupoid:

**Definition 3.1.** Dua is the groupoid whose

- (1) Objects are fusion spin chains  $A(\mathcal{E}, X)$ .
- (2) Morphisms are bounded-spread isomorphisms with composition.

Our discussion above has motivated the definition of Dua, whose objects are *abstract* spin systems of symmetric operators. However, we are interested in studying spin systems *together with an embedding into an edge-restricted algebra* for some categorical symmetry. Here we give an abstract formulation.

**Definition 3.2.** Let  $\mathcal{E}$  be an indecomposable multi-fusion category. A *quotient* of  $\mathcal{E}$  is an indecomposable multifusion category  $\mathcal{D}$  and a dominant tensor functor  $F : \mathcal{E} \rightarrow \mathcal{D}$

We typically consider quotients up to (unitary) monoidal equivalence. Now, Given a dominant tensor functor  $F : \mathcal{E} \rightarrow \mathcal{D}$ , we have a natural inclusion of nets,

$$A(\mathcal{E}, X) \hookrightarrow_{\iota_F} A(\mathcal{D}, F(X))$$

where  $f \in A(\mathcal{E}, X)_I = \mathcal{E}(X^{\otimes n}, X^{\otimes n})$  is sent to the composition

$$F(X)^{\otimes n} \xrightarrow{\text{can}} F(X^{\otimes n}) \xrightarrow{F(f)} F(X^{\otimes n}) \xrightarrow{\text{can}} F(X)^{\otimes n}$$

in  $A(\mathcal{D}, F(X))_I$ . Here  $\text{can}$  denotes the canonical unitary isomorphisms built from the tensorators on  $F$ .  $\iota_F$  is a unital inclusion of the local  $C^*$ -algebras for each  $I$ , and is compatible with the connecting inclusions associated with  $I \subseteq J$ . Thus  $\iota_F$  extends to a unital inclusion of quasi-local algebras. If we have a quotient  $F : \mathcal{C} \rightarrow \mathcal{D}$  and a quotient  $G : \mathcal{D} \rightarrow \mathcal{E}$ , then the composition  $G \circ F : \mathcal{C} \rightarrow \mathcal{E}$  is a quotient. Furthermore, it is easy to see by construction that  $\iota_{G \circ F} := \iota_G \circ \iota_F$ . We call such an inclusion of nets a *categorical inclusion*.

Given a finitely semi-simple category  $\mathcal{M}$ , its category of endofunctor  $\text{End}(\mathcal{M})$  is an indecomposable multi-fusion category equivalent to  $\text{Mat}_n(\text{Hilb}_{f.d.})$ , where  $n = \text{rank}(\mathcal{M})$ . Recall that the data of a (left)  $\mathcal{E}$ -module category structure on a finitely semi-simple category  $\mathcal{M}$  can be expressed as a tensor functor  $F : \mathcal{E} \rightarrow \text{End}(\mathcal{M})$ , where  $F(Y) := Y \triangleright \cdot$ .

**Definition 3.3.** A *spatial realization* of a fusion categorical net  $A(\mathcal{E}, X)$  is a categorical inclusion  $A(\mathcal{E}, X) \hookrightarrow A(\text{End}(\mathcal{M}), X \triangleright \cdot)$  associated to an indecomposable left  $\mathcal{E}$ -module category  $\mathcal{M}$ . We denote  $A(\text{End}(\mathcal{M}), X \triangleright \cdot)$  by  $A(\mathcal{E}, X)_{\mathcal{M}}$ .

When  $\text{rank}(\mathcal{M}) = 1$ ,  $\text{End}(\mathcal{M}) \cong \text{Hilb}_{f.d.}$ , hence a spatial realization is an honest, unital inclusion of a fusion spin chain into the local operators of a concrete spin chain.

The motivation for this definition arises from our concept of fusion categorical symmetries. Above, we characterized these with the data of a fusion category  $\mathcal{C}$ , a right, indecomposable module category  $\mathcal{M}$ , and a (strongly tensor generating) object  $X \in \mathcal{C}_{\mathcal{M}}^*$ , correspond precisely (up to an obvious notion of equivalence) to spatial realizations of the fusion categorical net  $A(\mathcal{C}_{\mathcal{M}}^*, X)$ .

If we change our perspective and denote  $\mathcal{E} := \mathcal{C}_{\mathcal{M}}^*$ , then we can view  $\mathcal{M}$  as a left indecomposable  $\mathcal{E}$ -module category, and thus spatial realizations of the fusion spin chain  $A(\mathcal{E}, X)$ , parameterized by a left module category  $\mathcal{M}$ , correspond precisely the algebra symmetry operators inside  $A^\circ$  for an action of the dual category  $\mathcal{C}_{\mathcal{M}}^* \curvearrowright A$ . This leads us to reformulate Question 2 as follows:

**Reformulation of Question 1.1.** Let  $A(\mathcal{E}, X)$  and  $A(\mathcal{D}, Y)$  be fusion spin chains, with spatial realizations parameterized by module categories  $\mathcal{M}$  and  $\mathcal{N}$  respectively. If  $\alpha : A(\mathcal{E}, X) \rightarrow A(\mathcal{D}, Y)$  is a bounded-spread isomorphism, when does  $\alpha$  extend to a bounded-spread isomorphism  $\tilde{\alpha} : A(\mathcal{E}, X)_{\mathcal{M}} \rightarrow A(\mathcal{D}, Y)_{\mathcal{N}}$ ?

#### 4. THE ALGEBRA MODEL FOR CATEGORICAL INCLUSIONS

One way to build a categorical inclusion is to pick a connected, commutative Q-system object  $L \in \mathcal{Z}(\mathcal{C})$  and consider the *category  $\mathcal{C}_L$  of right  $A$ -modules in  $\mathcal{C}$*  (technically we should first apply the forgetful functor to  $L$  to obtain an object in  $\mathcal{C}$ ). The commutative central structure on  $L$  equips  $\mathcal{C}_L$  with the structure of a unitary multi-fusion category such that the free module functor  $F_L : \mathcal{C} \rightarrow \mathcal{C}_L$ ,  $x \mapsto x \otimes L$ , is a dominant tensor functor.

A theorem of [BN10] shows that any quotient  $F : \mathcal{C} \rightarrow \mathcal{D}$  is of the form described above, i.e. there exists a (unique up to isomorphism) connected commutative Q-system  $L \in \mathcal{Z}(\mathcal{C})$  such that the digram of monoidal functors commutes up to natural isomorphism

$$\begin{array}{ccc} \mathcal{C} & & \\ F \downarrow & \searrow^{F_L} & \\ \mathcal{D} & \xrightarrow{\cong} & \mathcal{C}_L \end{array}$$

This gives an “internal” description of a quotient as opposed to the a-priori “external” description requiring an indecomposable multi-fusion category  $\mathcal{D}$ . This is directly analogous to the first isomorphism theorem for groups.

Now we recall a concrete model for constructing  $\mathcal{C}_L$  and the free module functor  $F_L : \mathcal{C} \rightarrow \mathcal{C}_L$ . Let  $L \in \mathcal{Z}(\mathcal{C})$  be a connected, commutative Q-system. We denote the multiplication morphism  $m : L \otimes L \rightarrow L$  and the unit map  $\iota : \mathbb{1} \rightarrow L$ . We slightly abuse notation and identify  $L$  and its structure maps with their images in  $\mathcal{C}$  under the forgetful functor.

Define the category  $\mathcal{C}_L^\circ$  whose objects are objects of  $\mathcal{C}$  and whose morphisms are given by

$$\mathcal{C}_L^\circ(x, y) := \mathcal{C}(x, y \otimes A).$$

Composition of morphisms  $f \in \mathcal{C}_L^\circ(y, z)$  and  $g \in \mathcal{C}_L^\circ(x, y)$  is defined as

$$f \circ g := (1_z \otimes m) \circ (f \otimes 1_L) \circ g \in \mathcal{C}_L^\circ(x, z).$$

Following [JP17], it is not hard to see that  $\mathcal{C}_L^\circ$  becomes a  $\mathbb{C}^*$ -category with  $*$ -structure (denoted here with a  $\dagger$  to distinguish from the  $*$ -structure in  $\mathcal{C}$ )

$$g^\dagger := g^* \circ (1_y \otimes (m^* \circ \iota)).$$

To define the tensor structure on  $\mathcal{C}_L^\circ$ , the tensor product on objects (which we denote with  $\boxtimes$ ) is just the tensor product in  $\mathcal{C}$ ,  $x \boxtimes y = x \otimes y$  while the tensor product of morphisms is defined, for  $f \in \mathcal{C}_L^\circ(x, y)$  and  $g \in \mathcal{C}_L^\circ(z, w)$  by

$$f \boxtimes g := (1_{y \otimes w} \otimes m) \circ (1_y \otimes (\sigma_{L, w} \otimes 1_L)) \circ (f \otimes h).$$

This turns  $\mathcal{C}_L$  into a (strict) unitary multi-tensor category, and since  $L$  was connected in  $\mathcal{Z}(\mathcal{C})$ , it is in fact an indecomposable multi-fusion category. In fact, the functor  $\mathcal{C}_L^\circ$  to the category  $\text{FreeMod}_L$  of free (internal to  $\mathcal{C}$ ) right  $L$ -modules sending  $x \mapsto x \otimes L$  and  $f \mapsto (1_y \otimes m) \circ (f \otimes 1_L)$  is an equivalence of categories.

Thus the (unitary) Karoubian completion of  $\mathcal{C}_L^\circ$  is equivalent to the category of projective right  $L$ -modules, which is equivalent to the category of all right  $L$  modules since this category is semisimple. By a minor abuse of notation, we denote the unitary Karoubian completion of  $\mathcal{C}_L^\circ$  by  $\mathcal{C}_L$  and identify  $\mathcal{C}_L^\circ$  with its image in  $\mathcal{C}_L$ .

In this picture, the free module functor  $F_L : \mathcal{C} \rightarrow \mathcal{C}_L$  is just given by  $F_L(x) = x \in \mathcal{C}_L$  and  $F_L(f) := (1_y \otimes \iota) \circ f$ .

Now we have a nice way to describe the net obtained from categorical quotients  $F : \mathcal{C} \rightarrow \mathcal{D}$  directly in terms of  $\mathcal{C}$ . Let  $L$  be the commutative Q-system in the center corresponding to this quotient, and  $F_L$  the model for  $F$  as above. Then for an interval  $I \subseteq \mathbb{Z}$  with  $|I| = n$ ,

$$A(\mathcal{C}_L, X)_I = \mathcal{C}_L(X^{\otimes n}, X^{\otimes n}) \cong \mathcal{C}(X^{\otimes n}, X^{\otimes n} \otimes L).$$

Furthermore, for  $I \subseteq J$ , the inclusion  $A(\mathcal{C}_L, X)_I \hookrightarrow \mathfrak{A}(\mathcal{C}_L, X)_J$  is described for  $f \in \mathcal{C}(X^{\otimes n}, X^{\otimes n} \otimes L)$  by

$$f \mapsto 1_{X^{\otimes l+n}} \sigma_{L, X^{\otimes r}} (1_{X^{\otimes l}} \otimes f \otimes 1_{X^{\otimes r}}).$$

Here  $l$  is the number of points in  $J$  strictly less than  $I$ , and  $r$  is the number of points in  $J$  strictly greater than  $I$ .

The inclusion  $A(\mathcal{C}, X)_I \hookrightarrow A(\mathcal{C}_L, X)_I$  is simply given by

$$f \mapsto f \otimes \iota,$$

for  $f \in \mathcal{C}(X^{\otimes n}, X^{\otimes n})$ .

This generalizes to give a clean description of the lattice of intermediate categorical inclusions. For any sub Q-system  $K \leq L$ , the inclusion map  $j_K : K \rightarrow L$  gives the natural maps

$$A(\mathcal{C}_K, X)_I \cong \mathcal{C}(X^{\otimes n}, X^{\otimes n} \otimes K) \hookrightarrow \mathcal{C}(X^{\otimes n}, X^{\otimes n} \otimes L) \cong A(\mathcal{C}_L, X)_I,$$

$$f \mapsto (1_{X^{\otimes n}} \otimes j_K) \circ f.$$

This sets up an equivalence between the lattice of intermediate category inclusions between  $A(\mathcal{C}, X)$  and  $A(\mathcal{C}_L, X)$  and the lattice of sub Q-system objects  $K \leq L$ . We typically take the largest algebra  $L$  to be normalized so that  $m \circ m^* = 1_L$ , but for intermediate subalgebras, we only require that  $m \circ m^* = \lambda 1_K$  for some (necessarily positive) scalar  $\lambda$ .

**Remark 4.1. Canonical conditional expectation.** For any connected étale algebra  $L \in \mathcal{Z}(\mathcal{C})$ , identifying  $A(\mathcal{C}, X) \hookrightarrow A(\mathcal{C}_L, X)$ , there is a canonical, locality-preserving conditional expectation  $E : A(\mathcal{C}_L, X) \rightarrow A(\mathcal{C}, X)$  which is defined for  $f \in A(\mathcal{C}_L, X)_I = \mathcal{C}(X^{\otimes n}, X^{\otimes n} \otimes L)$  by

$$E(f) := \frac{1}{d_L} (1_{X^{\otimes n}} \otimes i^*) \circ f.$$

**Remark 4.2.** By the above discussion, spatial realizations of  $A(\mathcal{C}, X)$  are parameterized by choices of indecomposable  $\mathcal{C}$ -module categories. Every such module category canonically defines a Lagrangian algebra  $Z(\mathcal{M}) \in \mathcal{Z}(\mathcal{C})$ , its center [Dav10]. In particular, we can always view the spatial realizations  $A(\mathcal{C}, X) \hookrightarrow A(\mathcal{C}, X)_{\mathcal{M}}$  as  $A(\mathcal{C}, X) \hookrightarrow A(\mathcal{C}_{Z(\mathcal{M})}, X)$ .

In this section, we prove a lemma necessary for the proof of the main theorem, concerning relative commutants. As mentioned above, fusion spin chains satisfy weak algebraic Haag duality, and in fact, satisfy an even stronger version simply called algebraic Haag duality. In particular, if we choose  $n$  such that  $X^{\otimes n}$  contains a copy of every simple object in  $\mathcal{C}$  (the existence of such an  $n$  is the definition of strong tensor generating), then we have that for any interval  $I$  with  $|I| > n$ ,  $A(\mathcal{C}, X)_I = \{x \in \mathfrak{A} : [x, y] = 0 \text{ for all } y \in A(\mathcal{C}, X)_{I^c}\}$ . The following lemma shows that the nets  $\mathfrak{A}(\mathcal{C}_L, X)$  satisfy an even stronger “relative” version of this property.

**Lemma 4.3.** *Let  $A(\mathcal{C}, X)$  be a fusion spin chain and  $n$  as above. For any connected, commutative Q-system  $L \in \mathcal{Z}(\mathcal{C})$ , then for any interval  $I$  with  $|I| > n$ ,*

$$A(\mathcal{C}_L, X)_I = \{x \in A(\mathcal{C}_L, X) : [x, y] = 0 \text{ for all } y \in A(\mathcal{C}, X)_{I^c}\}.$$

*Proof.* Note that we can view the quasi-local algebra  $A_{\mathcal{C}_L}$  for  $A(\mathcal{C}, X)$  as a finite index extension of  $A_{I^c}$ , supported on the canonical copy of  $\mathcal{C} \boxtimes \mathcal{C}^{mp} \hookrightarrow \text{Bim}(A_{I^c})$  [Jon24]. Indeed,  $A(\mathcal{C}_L, X)$  is the Q-system realization of the Q-system  $K \in \mathcal{C} \boxtimes \mathcal{C}^{mp}$  obtained from considering  $\mathcal{C}_L$  as a  $\mathcal{C}$ - $\mathcal{C}$  bimodule category, choosing the object  $X^{\otimes n} \in \mathcal{C}_L$  and taking internal endomorphisms. In symbols,

$$A(\mathcal{C}_L, X) \cong |K|$$

Since the embedding of  $\mathcal{C} \boxtimes \mathcal{C}^{mp} \hookrightarrow \text{Bim}(\mathfrak{A}_{I^c})$  is fully faithful, by the usual Q-system arguments from subfactor theory [Jon24, CHPJP22],

$$A(\mathcal{C}, X)'_{I^c} \cap A(\mathcal{C}_L, X) \cong 1_{A(\mathcal{C}, X)_{I^c}} \otimes_{\mathbb{C}} \mathcal{C}(\mathbb{1}, K) \subseteq |K|.$$

By the definition of internal end and Q-system realization, under the identification of the realization subspace with our concrete net, we obtain

$$A(\mathcal{C}_L, X) \cong 1_{A(\mathcal{C}, X)_{I^c}} = \mathcal{C}(X^{\otimes n}, X^{\otimes n} \otimes L) = A(\mathcal{C}_L, X)_I.$$

□

## 5. SOLVING THE EXTENSION PROBLEM WITH DHR BIMODULES

Here, we recall the definition of a DHR bimodule on a net of algebras (over  $\mathbb{Z}$ ). First, if  $A$  is a (unital)  $C^*$ -algebra, then a *bimodule* is an algebraic  $A$ - $A$  bimodule  $X$ , together with a right  $A$ -valued inner product  $\langle \cdot | \cdot \rangle : X \times X \rightarrow A$  (making  $X$  into a Hilbert  $A$ -module) such that the left  $A$  action is by adjointable operators. What we call bimodules are more typically called *correspondences* in the  $C^*$ -algebra literature.

We denote the collection of all  $A$ - $A$  bimodules by  $\text{Bim}(A)$ . This forms a  $C^*$ -tensor category, whose objects are bimodules, morphisms are adjointable bimodule intertwiners, and tensor product is given by the relative tensor product of bimodules, denoted  $\boxtimes_A$  (see [CHPJP22]).

If  $X \in \text{Bim}(A)$ , a *projective basis* is a finite set  $\{b_i\} \subseteq X$  such that for all  $x \in X$

$$x = \sum_i b_i \langle b_i | x \rangle.$$

**Definition 5.1.** Let  $A$  be an abstract spin chain with quasi-local algebra  $A$ . A bimodule over the quasilocal algebra  $A$  is called a *DHR bimodule* if there exists an  $R$  such that for every interval with  $|I| > R$ , there exists a projective basis  $\{b_i\}$  for  $X$  such that  $ab_i = b_i a$  for all  $x \in A_{I^c}$ .

The collection of DHR bimodules is closed under  $\boxtimes_A$  and thus defines a full  $\otimes$  subcategory of  $\text{DHR}(A) \subseteq \text{Bim}(A)$ . If  $A$  also satisfies *weak algebraic Haag duality* (a property satisfied by fusion spin chains, see [Jon24]), then  $\text{DHR}(\mathfrak{A})$  admits a *canonical braiding*.

The clearest physical interpretation of DHR bimodules as a braided tensor category is in terms of bulk patch operators. One realization of a fusion spin chain is as the “boundary algebra” of of 2+1D topologically ordered spin system, in particular of Levin-Wen type models [LW05]. In this case, string operators associated to an anyon type  $Z \in \mathcal{Z}(\mathcal{C})$  in the bulk can be pushed onto the boundary, giving operators in the fusion spin chain  $A(\mathcal{C}, X)$  called *patch operators* [CW23, IW23]. If we push the right endpoint off to infinity, this operator no longer lives in  $A(\mathcal{C}, X)$ , but can be thought of formally as an element in a DHR-bimodule since it is localized in an interval around the left endpoint. Similarly to the GNS construction, we can take local perturbations and assemble this into a DHR-bimodule, corresponding to the anyon type  $Z \in \mathcal{Z}(\mathcal{C})$ . This suggests there should be a close relationship between  $\text{DHR}(A(\mathcal{C}, X))$  and  $\mathcal{Z}(\mathcal{C})$ , which is demonstrated in the following theorem.

**Theorem 5.2** ([Jon24], Theorem C). *If  $X$  is a strong, self-dual tensor generator of the unitary fusion category  $\mathcal{C}$ , then  $\text{DHR}(\mathfrak{A}(\mathcal{C}, X)) \cong \mathcal{Z}(\mathcal{C})$  as braided  $C^*$ -tensor categories.*

One of the main motivations of [Jon24] is that the DHR construction defines a functor  $\text{DHR} : \text{Net}_{\mathbb{Z}} \rightarrow C^*\text{-BrTens}$ , where  $\text{Net}_{\mathbb{Z}}$  is the groupoid of abstract spin chains and bounded-spread isomorphisms. This associates to each abstract spin chain its braided tensor category of DHR bimodules, while

$$\text{DHR}(\alpha) : \text{DHR}(A) \rightarrow \text{DHR}(B)$$

which acts on DHR bimodules by sending a DHR bimodule  $X$  of  $A$  to  $\text{DHR}(\alpha)(X) := X$  as a vector space, with left and right  $B$  actions given by

$$b \triangleright x \triangleleft c := \alpha^{-1}(b)x\alpha^{-1}(c),$$

with  $B$  valued inner product

$$\langle x | y \rangle_B := \alpha(\langle x | y \rangle_A).$$

In particular, for every bounded-spread isomorphism between fusion spin chains

$$\alpha : A(\mathcal{C}, X) \rightarrow A(\mathcal{D}, Y),$$

there is an induced unitary braided equivalence of categories

$$\text{DHR}(\alpha) : \mathcal{Z}(\mathcal{C}) \cong \text{DHR}(A(\mathcal{C}, X)) \rightarrow A(\mathcal{D}, Y) \cong \mathcal{Z}(\mathcal{D}).$$

**Theorem 5.3.** *Let  $\alpha : A(\mathcal{C}, X) \rightarrow A(\mathcal{D}, Y)$  be a bounded-spread isomorphism. Then for any commutative, connected  $Q$ -systems  $L \in \mathcal{Z}(\mathcal{C})$  and  $K \in \mathcal{Z}(\mathcal{D})$ , extensions of  $\alpha$  to bounded-spread isomorphisms  $\tilde{\alpha} : A(\mathcal{C}_L, X) \rightarrow A(\mathcal{D}_K, Y)$  are in bijective correspondence with algebra isomorphisms  $\gamma : \text{DHR}(\alpha)(L) \cong K$  in  $\mathcal{Z}(\mathcal{D})$ . As a consequence:*

- (1) *If  $\text{DHR}(\alpha)(L)$  is not isomorphic to  $Z(K)$ , then  $\alpha$  has no extension to a bounded-spread isomorphism  $\tilde{\alpha} : A(\mathcal{C}_L, X) \rightarrow A(\mathcal{D}_K, Y)$*
- (2) *If  $\text{DHR}(\alpha)(L) \cong K$ , spatial implementations form a torsor over  $\text{Aut}(L)$ .*

*Proof.* Let  $L \in \mathcal{Z}(\mathcal{C})$  be a commutative, connected  $Q$ -system. Then  $A(\mathcal{C}_L, X)$  is algebraically a bimodule over the quasi-local algebra  $A(\mathcal{C}, X)$  obtained by including the latter as a subalgebra of the former. The canonical condition expectation equips  $A(\mathcal{C}_L, X)$  with the structure of a right Hilbert  $A(\mathcal{C}, X)$  module, hence  $(A(\mathcal{C}_L, X), E) \in \text{Bim}(A(\mathcal{C}, X))$ . In fact, from the canonical embedding  $\mathcal{Z}(\mathcal{C}) \cong \text{DHR}(A(\mathcal{C}, X)) \subseteq \text{Bim}(A(\mathcal{C}, X))$ , we have that the quasi-local algebra with condition

expectation  $A(\mathcal{C}_L, X), E$  is simply the Q-system realization  $|L|$  of  $L \in \mathcal{Z}(\mathcal{C})$ , and in particular  $(A(\mathcal{C}_L, X), E) \in \text{DHR}(A(\mathcal{C}, X))$  is a connected, commutative Q-system.

With this picture in hand, to establish our bijection, we simply show that the data of one thing can simply be reinterpreted as the data of the other.

First, suppose  $\tilde{\alpha} : A(\mathcal{C}_L, X) \rightarrow A(\mathcal{D}_K, Y)$  is a bounded-spread isomorphism, restricting to  $\alpha$  on the subsystem  $A(\mathcal{C}, X)$ . We claim that  $\tilde{\alpha}$  is an algebra object isomorphism  $\text{DHR}(\alpha)(A(\mathcal{C}_L, X)) \cong A(\mathcal{D}_K, X)$ . Indeed,  $\text{DHR}(\alpha)(A(\mathcal{C}_L, X))$  is simply  $A(\mathcal{C}_K, X)$  with left and right action twisted by  $\alpha^{-1}$ . We check for  $a \in A(\mathcal{C}_L, X)$  and  $b, c \in A(\mathcal{D}, Y)$

$$\tilde{\alpha}(b \triangleright a \triangleleft c) = \tilde{\alpha}(\alpha^{-1}(b)a\alpha^{-1}(c)) = b\tilde{\alpha}(a)c$$

where we have used  $\tilde{\alpha}|_{A(\mathcal{C}, X)} = \alpha$ . Thus  $\tilde{\alpha} : \text{DHR}(\alpha)(A(\mathcal{C}_L, X)) \cong A(\mathcal{D}_K, X)$  is an isomorphism of  $A(\mathcal{D}, Y)$  bimodules. Since it is an algebra isomorphism, this bimodule isomorphism is an isomorphism of algebra objects.

Conversely, suppose we have an isomorphism of algebra objects  $\tilde{\alpha} : \text{DHR}(\alpha)(A(\mathcal{C}_L, X)) \cong A(\mathcal{D}_K, X)$  in the category of  $A(\mathcal{D}, Y)$  bimodules. Then reversing the above logic,  $\tilde{\alpha}$  is an algebra isomorphism  $A(\mathcal{C}_L, X) \rightarrow A(\mathcal{D}_K, Y)$  which restricts to  $\alpha$  on the algebra  $A(\mathcal{C}, X)$ . It remains to show that  $\tilde{\alpha}$  is bounded spread.

Suppose  $\alpha$  has spread at most  $R$ . We claim  $\tilde{\alpha}$  has spread at most  $R$ . Indeed, let  $a \in A(\mathcal{C}_L, X)_I$ . Note that  $A(\mathcal{D}, Y)_{(I+R)c} \subseteq \alpha(A(\mathcal{C}, X)_{Ic})$ , and since  $[a, b] = 0$  for all  $b \in A(\mathcal{C}, X)_{Ic}$ ,

$$0 = [\tilde{\alpha}(a), \tilde{\alpha}(b)] = [\tilde{\alpha}(a), \alpha(b)],$$

and hence

$$[\tilde{\alpha}(A(\mathcal{C}_L, X)_I), A(\mathcal{D}, Y)_{(I+R)c}] = 0.$$

By Lemma 4.3, this implies

$$\tilde{\alpha}(A(\mathcal{C}_L, X)_I) \subseteq A(\mathcal{C}_K, X)_{I+R},$$

so that  $\tilde{\alpha}$  is bounded spread. This concludes the proof of the main statement of the theorem.

The first consequence listed above follows immediately. The second follows from the fact that for any isomorphisms  $\gamma, \gamma' : \text{DHR}(\alpha(L)) \cong K$ ,  $\gamma^1 \circ \gamma' \in \text{Aut}(\text{DHR}(\alpha(L)) \cong \text{Aut}(L))$ . □

We recommend the reader to compare the above theorem with an analogous result concerning extending symmetries in 1+1D algebraic quantum field theory, [BJLP19, Section 6].

**Proof of Theorem 1.3.** The Theorem 1.3 stated in the introduction follows as a special case of the above, and we believe it provides a satisfactory answer to Question 2. Indeed, consider the case where  $L \in \mathcal{Z}(\mathcal{C})$  and  $K \in \mathcal{Z}(\mathcal{D})$  are  $Z(\mathcal{M})$  and  $Z(\mathcal{N})$  associated to the  $\mathcal{C}$  and  $\mathcal{D}$  module categories, respectively. Then we have identifications of the spatial realizations  $A(\mathcal{C}, X)_{\mathcal{M}} \cong A(\mathcal{C}_{Z(\mathcal{M})}, X)$ , whence the first part of the result follows from the previous theorem. That  $\text{Aut}(Z(\mathcal{M})) \cong (\mathcal{C}_{\mathcal{M}}^*)^\times$  follows, for example, from [BJ22]. □

For any algebra  $L \in \mathcal{Z}(\mathcal{C})$ , there is a group

$$\text{Aut}_{br}(\mathcal{Z}(\mathcal{C}) \mid L)$$

which consists of equivalence classes of pairs  $(\alpha, \gamma)$ , where  $\alpha$  is a braided tensor equivalence of  $\mathcal{Z}(\mathcal{C})$  and  $\gamma : \alpha(L) \cong L$  is an isomorphism of algebras. An equivalence between pairs  $(\alpha, \gamma)$  and  $(\beta, \delta)$  consists of a monoidal natural isomorphism  $\nu : \alpha \cong \beta$  such that  $\delta \circ \nu_L = \gamma$ . By [DMNO13, Sch24], for a Lagrangian algebra  $L$  corresponding to an indecomposable  $\mathcal{C}$ -module category  $\mathcal{M}$ , we have

$$\text{Aut}_{br}(\mathcal{Z}(\mathcal{C}) \mid L) \cong \text{Aut}_{\otimes}(\mathcal{C}_{\mathcal{M}}^*)$$

where the latter is the group of tensor autoequivalences up to monoidal natural isomorphism.

Let  $\text{QCA}(\mathcal{C}, X, \mathcal{M})$  be the group of bounded-spread autoequivalences of  $A(\mathcal{C}, X)_{\mathcal{M}}$  which preserve the subalgebra  $A(\mathcal{C}, X)$ . We can think of this as the group of spatially implemented dualities, or as the group of “edge-restricted” QCA that preserve the symmetric subalgebra. Then the above theorem yields the following corollary:

**Corollary 5.4.** *There is a homomorphism  $\pi : \text{QCA}(\mathcal{C}, X, \mathcal{M}) \rightarrow \text{Aut}_{\otimes}(\mathcal{C}_{\mathcal{M}}^*)$ .*

We can easily choose examples so that this is surjective. Indeed, consider the object  $X = \bigoplus_{Y \in \text{Irr}(\mathcal{C})} Y$ . Then  $\pi : \text{QCA}(\mathcal{C}, X, \mathcal{C}) \rightarrow \text{Aut}_{\otimes}(\mathcal{C})$  is surjective (see [Jon24]).

A family of examples of interest, which arise from onsite finite group symmetry, are when  $\mathcal{C} = \text{Rep}(G)$ ,  $\mathcal{M} = \text{Hilb}_{f.d.}$ , made into a  $\mathcal{C}$ -module category with the standard fiber functor structure on  $\text{Rep}(G)$ . In this case, we obtain a homomorphism

$$\pi : \text{QCA}(\text{Rep}(G), X, \text{Hilb}_{f.d.}) \rightarrow \text{Aut}_{\otimes}(\text{Hilb}_{f.d.}(G)) \cong H^2(G, U(1)) \rtimes \text{Aut}(G).$$

## 6. EXAMPLES AND APPLICATIONS

In this section, we consider a family of examples generalizing Kramers-Wannier (KW) dualities [LDOV23], and we also discuss applications of our previous result.

**Generalized shifts.** Let  $\mathcal{C}$  be a unitary fusion category, and  $\alpha \in \text{Aut}_{br}(\mathcal{Z}(\mathcal{C}))$ , such that  $\alpha$  generates a categorical action of  $\mathbb{Z}/n\mathbb{Z}$  on  $\mathcal{Z}(\mathcal{C})$  for some  $n$ . Then the  $o_4$  obstruction necessarily vanishes as  $H^4(\mathbb{Z}/n\mathbb{Z}, U(1))$  is trivial, and thus by [ENO10] there exists a  $\mathbb{Z}/n\mathbb{Z}$ -graded extension

$$\mathcal{D} := \bigoplus_{g \in \mathbb{Z}/n\mathbb{Z}} \mathcal{C}_g,$$

with  $\mathcal{C}_e = \mathcal{C}$ . For any  $Y \in \mathcal{C}_\alpha$ ,  $Y^{\otimes n} \in \mathcal{C}_e = \mathcal{C}$ . We can always choose  $Y$  such that  $X := Y^{\otimes n}$  is a strong tensor generator for  $\mathcal{C}$ , for example,  $Y = \bigoplus_{Z \in \text{Irr}(\mathcal{C}_\alpha)} Z$ .

Now, consider the fusion spin chain  $A(\mathcal{C}, X)$ . We define a “generalized translation” QCA for  $a \in A(\mathcal{C}, X)_{[i,j]} \cong \mathcal{C}(X^{j-i+1}, X^{j-i+1}) \cong \mathcal{D}(Y^{n(j-i+1)}, Y^{n(j-i+1)})$ ,

$$\tau_\alpha(a) := 1_{Y^{n-1}} \otimes a \otimes 1_Y \in A(\mathcal{C}, X)_{[i-1,j]}.$$

Then since for any  $Z \in \mathcal{Z}(\mathcal{C})$  we have natural isomorphisms  $Y \otimes Z \cong \alpha(Z) \otimes Y$ , it is easy to verify that

$$\text{DHR}(\tau_\alpha) \cong \alpha.$$

Thus, any braided autoequivalence of  $\mathcal{Z}(\mathcal{C})$  can be implemented by a QCA on the some fusion spin chain of the form  $A(\mathcal{C}, X)$ , although the choice of  $X$  depends on the autoequivalence.

**Dualities from Tambara-Yamagami.** The easiest example is the case  $\mathcal{C} := \text{Hilb}_{f.d.}(A)$ , where  $A$  is a finite abelian group.  $\mathcal{Z}(\mathcal{C}) \cong \text{Hilb}_{f.d.}(A \times \widehat{A})$ . Any symmetric, non-degenerate bicharacter  $\chi : A \times A \rightarrow U(1)$ , yields an isomorphism  $\tilde{\chi} : A \rightarrow \widehat{A}$ , given by  $\tilde{\chi}(x) = \chi(a, \cdot)$ . This results in a braided autoequivalence  $\alpha_\chi \in \text{Aut}_{br}(\text{Hilb}_{f.d.}(A \times \widehat{A}))$  defined by

$$\alpha_\chi(a, f) = (\tilde{\chi}^{-1}(f), \tilde{\chi}(a)).$$

Which further results in a  $\mathbb{Z}/2\mathbb{Z}$  extension of  $\text{Hilb}_{f.d.}(A)$  called a *Tambara-Yamagami* category [TY98], whose simple objects are simply  $A \cup \{\rho\}$  with fusion rules

$$\begin{aligned} a \otimes b &= ab, \\ a \otimes \rho &= \rho = \rho \otimes a, \end{aligned}$$

$$\rho \otimes \rho \cong \bigoplus_{a \in A} a.$$

If we set  $Y = \rho$ , then  $X := Y^2 = \bigoplus_{a \in A} a$  is a strong tensor generator for  $\mathcal{C}$ , and thus the above construction yields a QCA on the spin chain

$$\tau_{\alpha_\chi} : A(\mathcal{C}, X) \rightarrow A(\mathcal{C}, X)$$

with  $\text{DHR}(\tau_{\alpha_\chi}) \cong \alpha_\chi$ . For  $A = \mathbb{Z}_2$ , there is a unique symmetric, non-degenerate bicharacter  $\chi$ , and  $\tau_{\alpha_\chi}$  is exactly the Kramers-Wannier duality map described earlier.

There are two canonical spatial realizations of  $A(\mathcal{C}, X)$  described by the Lagrangian algebras  $L_1 := \mathbb{C}[A] \times 1$  and  $L_2 := 1 \times \mathbb{C}[\widehat{A}]$ , sometimes called electric and magnetic algebras. Then  $\alpha_\chi$  swaps  $L_1$  and  $L_2$ . In particular, by Theorem 1.3,  $\tau_{\alpha_\chi}$  cannot extend to a QCA on either spatial realization  $A(\mathcal{C}_{L_1}, X)$  or  $A(\mathcal{C}_{L_2}, X)$ .

**Q-system complete fusion categories.** A unitary fusion category  $\mathcal{C}$  is called *Q-system complete* (also called *torsion-free in the literature* if there is exactly one indecomposable module category (namely,  $\mathcal{C}$  itself) up to equivalence. This terminology comes from the fact that the fusion category  $\mathcal{C}$ , thought of as a 2-category with one object, is Q-system complete in the sense of [CHPJP22]. Examples include Fib and more generally  $\text{PSU}(2)_{2k+1}$ , and  $\text{Hilb}_{f.d.}(\mathbb{Z}/p\mathbb{Z}, \omega)$ , where  $[\omega] \in H^3(\mathbb{Z}/p\mathbb{Z})$  is non-trivial.

For Q-system complete fusion categories  $\mathcal{C}$ , any fusion spin chain of the form  $A(\mathcal{C}, X)$  has only one spatial realization  $A(\mathcal{C}, X)_{\mathcal{C}}$ , which in turn corresponds to a unique isomorphism class of Lagrangian algebra in  $\mathcal{Z}(\mathcal{C})$ . This results in the following corollary of Theorem 1.3.

**Proposition 6.1.** *Let  $\mathcal{C}$  be a Q-system complete fusion category. Suppose  $\alpha : A(\mathcal{C}, X) \rightarrow A(\mathcal{D}, Y)$  is a bounded-spread isomorphism between fusion spin chains. Then*

- (1)  $\mathcal{C} \cong \mathcal{D}$  as fusion categories and in particular  $\mathcal{D}$  is Q-system complete.
- (2) There exists a spatial implementation  $\tilde{\alpha} : A(\mathcal{C}, X)_{\mathcal{C}} \rightarrow A(\mathcal{D}, Y)_{\mathcal{D}}$  of  $\alpha$ .
- (3) The spatial implementations form a torsor over  $\text{Inv}(\mathcal{C})$ . In particular, if  $\mathcal{C}$  has no non-trivial invertible objects (e.g.  $\text{PSU}(2)_{2k+1}$ ), then there is a unique spatial implementation.

*Proof.* Let  $L$  denote the unique Lagrangian algebra object in  $\text{DHR}(A(\mathcal{C}, X)) \cong \mathcal{Z}(\mathcal{C})$ , so that  $\mathcal{C} \cong \mathcal{Z}(\mathcal{C})_L$ . Then  $\text{DHR}(\alpha)(L) \in \text{DHR}(A(\mathcal{D}, Y)) \cong \mathcal{Z}(\mathcal{D})$  is the unique Lagrangian algebra (since  $\text{DHR}(\alpha)$  is a braided equivalence). In particular, we have  $K := \text{DHR}(L)$  must be the canonical Lagrangian algebra in  $\mathcal{Z}(\mathcal{D})$ , hence  $\mathcal{Z}(\mathcal{D})_K \cong \mathcal{D}$ , and thus  $\text{DHR}(\alpha)$  induces a monoidal equivalence  $\mathcal{C} \cong \mathcal{Z}(\mathcal{C})_L \rightarrow \mathcal{Z}(\mathcal{D})_K \cong \mathcal{D}$ .

Since the spatial realizations must correspond to the Lagrangian algebras  $L$  and  $K$  respectively, the existence of a spatial implementation of  $\alpha$ , and the statement about the torsor, follows immediately from Theorem 1.3.  $\square$

**Symmetric QCA.** In this section, apply our above analysis to give a better understanding of QCA which commute with a group symmetry, and connect our perspective to the literature. Suppose we have a group  $G$  acting unitarily on the local Hilbert space  $V := \mathbb{C}^d$ , given by an assignment  $g \mapsto U_g \in U(d)$ . If we let  $A$  denote the spin system This extends to the automorphism of each  $A_I$  by  $\alpha_g(a) := (U_g)^{\otimes n} a (U_g^*)^{\otimes n}$ , which extends to a homomorphism  $G \rightarrow \text{Aut}(A)$ . We assume this homomorphism is injective.

In the literature, there have been several investigations into *symmetric QCA* (*sQCA*) [GSSC20], which in our language are bounded-spread isomorphisms  $\alpha : A \rightarrow A$  such that  $\alpha \circ \alpha_g = \alpha_g \circ \alpha$  for all  $g \in G$ . Our goal in this section is to make contact with these results from our perspective.

First, we recast this picture into the language of fusion spin chains by noting that  $A^G$  has the structure of an abstract spin chain with local algebras  $A_I^G := \{x \in A_I : \alpha_g(x) = x \text{ for all } g \in G\}$ .

Now if we view  $V$  as an object in  $\text{Rep}(G)$ , and we further assume that  $V$  is self-dual and strongly tensor generating (e.g.  $V \cong L^2(G)$  is the left regular representation), then

$$A^G \cong A(\text{Rep}(G), V).$$

The inclusion  $A^G \subseteq A$  is a spatial realization which corresponds to the standard fiber functor  $\text{Rep}(G) \rightarrow \text{Hilb}_{f.d.}$ . This leads to the following natural question:

**Question 6.2.** *Suppose  $\alpha : A^G \rightarrow A^G$  is a duality and  $\tilde{\alpha} : A \rightarrow A$  is a spatial implementation. Under what conditions the spatial implementation  $\tilde{\alpha}$  is a symmetric QCA?*

To answer this question, note that the Lagrangian algebra resulting from this spatial realization in  $\mathcal{Z}(\text{Rep}(G))$  can actually be viewed as an algebra object in  $\text{Rep}(G) \subseteq \mathcal{Z}(\text{Rep}(G))$ , where we identify  $\text{Rep}(G)$  with the full subcategory of its center using the standard symmetric braiding. The algebra is simply  $L := \text{Fun}(G)$ , the commutative algebra of  $\mathbb{C}$ -valued functions, viewed as a  $G$ -representation where  $G$  acts by translation [EGNO15].

**Theorem 6.3.** *Suppose  $\alpha : A^G \rightarrow A^G$  is a duality and  $\tilde{\alpha} : A \rightarrow A$  is a spatial implementation, and  $\gamma : \text{DHR}(\alpha)(L) \cong L$  is the associated algebra isomorphism. Then*

- (1)  $\tilde{\alpha}$  is a symmetric QCA if and only if for every  $\beta \in \text{Aut}(L)$ ,  $\gamma \circ \text{DHR}(\alpha)(\beta) = \beta \circ \gamma$ .
- (2) There is a homomorphism from the group of sQCA on  $A$  to  $H^2(G, U(1))$ .

*Proof.*  $A$ , as an algebra object (Q-system) in  $\text{Bim}(A^G)$ , is isomorphic to the Q-system realization  $|L|$  of the function algebra  $\text{Fun}(G) \in \text{Rep}(G) \subseteq \mathcal{Z}(\text{Rep}(G))$ , and thus we have that the group  $\text{Aut}(A|A^G) \leq \text{Aut}_{\text{Rep}(G)}(A)$  of automorphisms of  $A$  that are the identity on  $A^G$ , is isomorphic to  $\text{Aut}_{\text{Rep}(G)}(L) = G$ . But the group action  $g \mapsto \alpha_g$  is an injective homomorphism  $G \rightarrow \text{Aut}(A|A^G) \leq G$ , which therefore must also be surjective and we must have that  $\text{Aut}(A|A^G) = G$  since  $G$  is finite.

Furthermore, since  $\text{Hilb}_{f.d.}(G)$  is the dual category of  $\text{Rep}(G)$  associated to the Lagrangian  $L$ , we have that the linear span  $\mathbb{C}[\text{Aut}_{\text{Rep}(G)}(L)] = \text{End}_{\text{Rep}(G)}(L)$ , by [BJ22] This proves part (1). Part (2) follows from [Sch24].  $\square$

#### ACKNOWLEDGMENTS

CJ and KS were supported by NSF DMS-2247202. DJW was supported in part by the Australian Research Council Discovery Early Career Research Award (DE220100625). This paper grew out of discussions that took place during the ‘‘QCA group’’ at the workshop ‘‘Fusion categories and tensor networks’’ at the American Institute of Mathematics in 2022.

#### REFERENCES

- [AFM20] David Aasen, Paul Fendley, and Roger Mong. Topological defects on the lattice: Dualities and degeneracies, 2020.
- [AHLM23] David Aasen, Jeongwan Haah, Zhi Li, and Roger Mong. Measurement quantum cellular automata and anomalies in floquet codes, 2023.
- [AMF16] David Aasen, Roger Mong, and Paul Fendley. Topological defects on the lattice: I. the Ising model. *Journal of Physics A: Mathematical and Theoretical*, 49(35):354001, Aug 2016.
- [ANW11] Pablo Arrighi, Vincent Nesme, and Reinhard Werner. Unitarity plus causality implies localizability. *Journal of Computer and System Sciences*, 77(2):372–378, 2011. Adaptivity in Heterogeneous Environments.
- [Arr19] Pablo Arrighi. An overview of quantum cellular automata, 2019.

- [BDSNY24] Lakshya Bhardwaj, Thibault Décoppet, Sakura Schafer-Nameki, and Matthew Yu. Fusion 3-Categories for Duality Defects. aug 2024.
- [BJ22] Marcel Bischoff and Corey Jones. Computing fusion rules for spherical  $G$ -extensions of fusion categories. *Selecta Math. (N.S.)*, 28(2):Paper No. 26, 39, 2022.
- [BJLP19] Marcel Bischoff, Corey Jones, Yuan Ming Lu, and David Penneys. Spontaneous symmetry breaking from anyon condensation. *Journal of High Energy Physics*, 2019, 11 2019.
- [BM20] Todd A. Brun and Leonard Mlodinow. Quantum cellular automata and quantum field theory in two spatial dimensions. *Physical Review A*, 102(6), Dec 2020.
- [BMW<sup>+</sup>17] Nick Bultinck, Michaël Mariën, Dominic Williamson, Mehmet Burak Şahinoğlu, Jutho Haegeman, and Frank Verstraete. Anyons and matrix product operator algebras. *Ann. Physics*, 378:183–233, 2017.
- [BN10] Alain Bruguières and Sonia Natale. Exact sequences of tensor categories. *International Mathematics Research Notices*, 2011:5644–5705, 6 2010.
- [BR87] Ola Bratteli and Derek Robinson. *Operator algebras and quantum statistical mechanics. 1*. Texts and Monographs in Physics. Springer-Verlag, New York, second edition, 1987.  $C^*$ - and  $W^*$ -algebras, symmetry groups, decomposition of states.
- [BR97] Ola Bratteli and Derek Robinson. *Operator algebras and quantum statistical mechanics. 2*. Texts and Monographs in Physics. Springer-Verlag, Berlin, second edition, 1997. Equilibrium states. Models in quantum statistical mechanics.
- [CCH<sup>+</sup>21] Yichul Choi, Clay Córdova, Po Shen Hsin, Ho Tat Lam, and Shu Heng Shao. Non-Invertible Duality Defects in 3+1 Dimensions. *Physical Review D*, 105(12), nov 2021.
- [CGW10] Xie Chen, Zheng-Cheng Gu, and Xiao-Gang Wen. Local unitary transformation, long-range quantum entanglement, wave function renormalization, and topological order. *Phys. Rev. B*, 82:155138, Oct 2010.
- [CHPJP22] Quan Chen, Roberto Hernández Palomares, Corey Jones, and David Penneys. Q-system completion for  $C^*$  2-categories. *J. Funct. Anal.*, 283(3):Paper No. 109524, 59, 2022.
- [CPGSV17] J Ignacio Cirac, David Perez-Garcia, Norbert Schuch, and Frank Verstraete. Matrix product unitaries: structure, symmetries, and topological invariants. *Journal of Statistical Mechanics: Theory and Experiment*, 2017(8):083105, 2017.
- [CW23] Arkya Chatterjee and Xiao-Gang Wen. Symmetry as a shadow of topological order and a derivation of topological holographic principle. *Phys. Rev. B*, 107:155136, Apr 2023.
- [Dav10] Alexei Davydov. Centre of an algebra. *Adv. Math.*, 225(1):319–348, 2010.
- [DMNO13] Alexei Davydov, Michael Müger, Dmitri Nikshych, and Victor Ostrik. The witt group of non-degenerate braided fusion categories. *Journal für die Reine und Angewandte Mathematik*, pages 135–177, 9 2013.
- [ECN11] Gerardo Ortiz Emilio Cobanera and Zohar Nussinov. The bond-algebraic approach to dualities. *Advances in Physics*, 60(5):679–798, 2011.
- [EGNO15] Pavel Etingof, Shlomo Gelaki, Dmitri Nikshych, and Victor Ostrik. *Tensor categories*, volume 205 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2015.
- [ENO02] Pavel Etingof, Dmitri Nikshych, and Viktor Ostrik. On fusion categories. *Annals of Mathematics*, 162:581–642, 2002.
- [ENO10] Pavel Etingof, Dmitri Nikshych, and Victor Ostrik. Fusion categories and homotopy theory. *Quantum Topol.*, 1(3):209–273, 2010. With an appendix by Ehud Meir.
- [Far20] Terry Farrelly. A review of quantum cellular automata. *Quantum*, 4:368, Nov 2020.
- [FFRS04] Jürg Fröhlich, Jürgen Fuchs, Ingo Runkel, and Christoph Schweigert. Kramers-Wannier duality from conformal defects. *Physical Review Letters*, 93(7):70601, aug 2004.
- [FH20] Michael Freedman and Matthew Hastings. Classification of quantum cellular automata. *Communications in Mathematical Physics*, 376, 06 2020.
- [FHH19] Michael Freedman, Jeongwan Haah, and Matthew Hastings. The group structure of quantum cellular automata. *Communications in Mathematical Physics*, 389:1277 – 1302, 2019.
- [FHH20] Lukasz Fidkowski, Jeongwan Haah, and Matthew B. Hastings. Exactly solvable model for a 4 + 1D beyond-cohomology symmetry-protected topological phase. *Phys. Rev. B*, 101:155124, Apr 2020.
- [FHP20] Giovanni Ferrer and Roberto Hernández Palomares. Classifying module categories for generalized temperley–lieb–jones  $*$ -2-categories. *International Journal of Mathematics*, 31(04):2050027, 2020.
- [FS20] Terry Farrelly and Julien Streich. Discretizing quantum field theories for quantum simulation, 2020.
- [GNP21] Zongping Gong, Adam Nahum, and Lorenzo Piroli. Coarse-grained entanglement and operator growth in anomalous dynamics, 2021.
- [GNVW09] David Gross, Vincent Nesme, Holger Vogts, and Reinhard Werner. Index theory of one dimensional quantum walks and cellular automata. *Communications in Mathematical Physics*, 310, 10 2009.
- [GSSC20] Zongping Gong, Christoph Sünderhauf, Norbert Schuch, and J. Ignacio Cirac. Classification of matrix-product unitaries with symmetries. *Physical Review Letters*, 124, 03 2020.

- [Haa21] Jeongwan Haah. Clifford quantum cellular automata: Trivial group in 2D and Witt group in 3D. *J. Math. Phys.*, 62(9):092202, 2021.
- [Haa22] Jeongwan Haah. Topological phases of unitary dynamics: Classification in Clifford category. arXiv:2205.09141, 2022.
- [HK64] Rudolf Haag and Daniel Kastler. An algebraic approach to quantum field theory. *J. Mathematical Phys.*, 5:848–861, 1964.
- [IW23] Kansei Inamura and Xiao-Gang Wen. 2+1d symmetry-topological-order from local symmetric operators in 1+1d, 2023.
- [JL24] Corey Jones and Junhwi Lim. An index for quantum cellular automata on fusion spin chains. *Annales Henri Poincaré*, 25(10):4399–4422, 2024.
- [JM21] Robert A. Jones and Max A. Metlitski. One-dimensional lattice models for the boundary of two-dimensional majorana fermion symmetry-protected topological phases: Kramers-wannier duality as an exact  $Z_2$  symmetry. *Phys. Rev. B*, 104:245130, Dec 2021.
- [JNPW23] Corey Jones, Pieter Naaijken, David Penneys, and Daniel Wallick. Local topological order and boundary algebras. arXiv: 2307.12552, 2023.
- [Jon24] Corey Jones. DHR bimodules of quasi-local algebras and symmetric quantum cellular automata. *Quantum Topology*, 2024. arXiv: 2304.00068.
- [JP17] Corey Jones and David Penneys. Operator algebras in rigid  $C^*$ -tensor categories. *Comm. Math. Phys.*, 355(3):1121–1188, 2017.
- [Kaw21] Yasuyuki Kawahigashi. Two-dimensional topological order and operator algebras. *International Journal of Modern Physics B*, 35(8):2130003–2616, March 2021.
- [Kaw22] Yasuyuki Kawahigashi. Projector matrix product operators, anyons and higher relative commutants of subfactors. *Mathematische Annalen*, 2022.
- [KOZ22] Justin Kaidi, Kantaro Ohmori, and Yunqin Zheng. Symmetry TFTs for Non-Invertible Defects. *Communications in Mathematical Physics*, 404(2):1021–1124, sep 2022.
- [KW41] H. A. Kramers and G. H. Wannier. Statistics of the two-dimensional ferromagnet. Part i. *Physical Review*, 60(3):252–262, aug 1941.
- [LDOV23] Laurens Lootens, Clement Delcamp, Gerardo Ortiz, and Frank Verstraete. Dualities in one-dimensional quantum lattice models: Symmetric Hamiltonians and matrix product operator intertwiners. *PRX Quantum*, 4:020357, Jun 2023.
- [LE24] David M. Long and Dominic V. Else. Topological Phases of Many-Body Localized Systems: Beyond Eigenstate Order. aug 2024.
- [LW05] Michael A. Levin and Xiao Gang Wen. String-net condensation: A physical mechanism for topological phases. *Physical Review B - Condensed Matter and Materials Physics*, 71:045110, 1 2005. The OG.
- [NS95] Florian Nill and Kornél Szlachányi. Quantum chains of Hopf algebras with quantum double cosymmetry. *Communications in Mathematical Physics*, 187:159–200, 1995.
- [PC20] Lorenzo Piroli and J. Ignacio Cirac. Quantum cellular automata, tensor networks, and area laws. *Phys. Rev. Lett.*, 125:190402, Nov 2020.
- [PFM<sup>+</sup>16] Hoi Chun Po, Lukasz Fidkowski, Takahiro Morimoto, Andrew C. Potter, and Ashvin Vishwanath. Chiral floquet phases of many-body localized bosons. *Phys. Rev. X*, 6:041070, Dec 2016.
- [PFVP17] Hoi Chun Po, Lukasz Fidkowski, Ashvin Vishwanath, and Andrew C. Potter. Radical chiral floquet phases in a periodically driven kitaev model and beyond. *Phys. Rev. B*, 96:245116, Dec 2017.
- [PM17] Andrew C. Potter and Takahiro Morimoto. Dynamically enriched topological orders in driven two-dimensional systems. *Phys. Rev. B*, 95:155126, Apr 2017.
- [PTSC21] Lorenzo Piroli, Alex Turzillo, Sujeet K Shukla, and J Ignacio Cirac. Fermionic quantum cellular automata and generalized matrix-product unitaries. *Journal of Statistical Mechanics: Theory and Experiment*, 2021(1):013107, 2021.
- [PVF18] Andrew C. Potter, Ashvin Vishwanath, and Lukasz Fidkowski. Infinite family of three-dimensional floquet topological paramagnets. *Phys. Rev. B*, 97:245106, Jun 2018.
- [RWW20] Daniel Ranard, Michael Walter, and Freek Witteveen. A converse to lieb-robinson bounds in one dimension using index theory. *arXiv preprint arXiv:2012.00741*, 2020.
- [Sch24] Kylan Schatz. Boundary symmetries of (2+1)d topological orders. 2024.
- [SSBC18] M. Burak Sahinoglu, Sujeet K. Shukla, Feng Bi, and Xie Chen. Matrix product representation of locality preserving unitaries. *Phys. Rev. B*, 98:245122, Dec 2018.
- [SW04] Benjamin Schumacher and Reinhard Werner. Reversible quantum cellular automata. *arXiv: Quantum Physics*, 2004.

- [SWB<sup>+</sup>14] Mehmet Burak Sahinoglu, Dominic Williamson, Nick Bultinck, Michaël Mariën, Jutho Haegeman, Norbert Schuch, and Frank Verstraete. Characterizing Topological Order with Matrix Product Operators. *Annales Henri Poincaré*, 22(2):563–592, sep 2014.
- [Tak79] Masamichi Takesaki. *Theory of operator algebras. I*. Springer-Verlag, New York-Heidelberg, 1979.
- [TW19] Ryan Thorngren and Yifan Wang. Fusion Category Symmetry I: Anomaly In-Flow and Gapped Phases. dec 2019.
- [TY98] Daisuke Tambara and Shigeru Yamagami. Tensor categories with fusion rules of self-duality for finite abelian groups. *Journal of Algebra*, 209(2):692–707, 1998.
- [VBW<sup>+</sup>18] Robijn Vanhove, Matthias Bal, Dominic J. Williamson, Nick Bultinck, Jutho Haegeman, and Frank Verstraete. Mapping topological to conformal field theories through strange correlators. *Physical Review Letters*, 121:177203, 10 2018.
- [Wat95] J. Watrous. On one-dimensional quantum cellular automata. In *Proceedings of IEEE 36th Annual Foundations of Computer Science*, pages 528–537, 1995.
- [WBV17] Dominic J. Williamson, Nick Bultinck, and Frank Verstraete. Symmetry-enriched topological order in tensor networks: Defects, gauging and anyon condensation. nov 2017.
- [ZL21] Carolyn Zhang and Michael Levin. Classification of interacting floquet phases with  $u(1)$  symmetry in two dimensions. *Phys. Rev. B*, 103:064302, Feb 2021.