

# Robustness to Model Approximation, Empirical Model Learning, and Sample Complexity in Wasserstein Regular MDPs

Yichen Zhou, Yanglei Song and Serdar Yüksel

Department of Mathematics and Statistics, Queen’s University, (yichen.zhou,yanglei.song,yuksel)@queensu.ca

The paper studies the robustness properties of discrete-time stochastic optimal control under Wasserstein model approximation for both discounted cost and average cost criteria. Specifically, we study the performance loss when applying an optimal policy designed for an approximate model to the true dynamics compared with the optimal cost for the true model under the sup-norm-induced metric, and relate it to the Wasserstein-1 distance between the approximate and true transition kernels. A primary motivation of this analysis is empirical model learning, as well as empirical noise distribution learning, where Wasserstein convergence holds under mild conditions but stronger convergence criteria, such as total variation, may not. We discuss applications of the results to the disturbance estimation problem, where sample complexity bounds are given, and also to a general empirical model learning approach, obtained under either Markov or i.i.d. learning settings. Further applications regarding the continuity of invariant probability measures with respect to transition kernels are also discussed.

*Key words:* Markov decision processes, robustness, empirical model learning, sample complexity

*Funding:* This work was supported by the Natural Sciences and Engineering Research Council of Canada

---

**1. Introduction** In this paper, we study the robustness properties of discrete-time stochastic optimal control for model approximation under various performance criteria. Specifically, we characterize the relationship between the accuracy of the model approximation and the performance loss of applying an optimal control policy for an approximate model to the true dynamics. This robustness property is of practical importance, as learning algorithms are seldom implemented for the true models; see e.g. [30] and [35], the latter of which shows that quantized Q-learning algorithms for continuous spaces converge to the fixed point equation of an approximate model.

Robustness to model approximations, such as finite approximations, is a well-studied problem with rich literature. In such cases, the approximations are often designed, e.g. by quantization, allowing for a relatively more direct analysis; see [14, 4, 51, 49, 31] under a variety of models. In our paper, we will build on some of the findings and constructions in [51, 49, 31]

Another important problem is model learning from experimental data. In [17, 18, 10], models are approximated by replacing the distribution of the “disturbance process” with its empirical estimates. [32, 36] placed the empirical consistency problem in the context of robustness to weak convergence of kernels. Recently, further robustness questions related to the disturbance processes have been studied under the formulation of distributionally robust stochastic control [59, 56]. In this paper, we follow the general model approximation formulation considered in [20, 21, 32, 36, 6], and, in particular, focus on models with Lipschitz continuous costs and transition kernels that are continuous under the Wasserstein-1 distance. We study both discounted cost and average cost criteria and consider applications such as state discretization and disturbance process estimation through a unified lens. In some applications, we establish explicit sample complexity bounds. A detailed literature review is presented later in the paper.

**1.1. Preliminaries on Markov Decision Processes** We consider a discrete-time controlled Markov process with an underlying sample space  $(\Omega, \mathcal{F}, \mathbb{P})$ , a Polish (i.e. complete, separable, and metric) state space  $(\mathbb{X}, d_{\mathbb{X}})$  and an action space  $(\mathbb{U}, d_{\mathbb{U}})$ . We endow the two spaces with their Borel  $\sigma$ -algebras, denoted by  $\mathcal{F}_{\mathbb{X}}$  and  $\mathcal{F}_{\mathbb{U}}$  respectively. We denote by  $(\mathbb{X} \times \mathbb{U}, \sigma(\mathcal{F}_{\mathbb{X}} \times \mathcal{F}_{\mathbb{U}}))$  their product space.

**DEFINITION 1 (CONTROLLED TRANSITION KERNELS).** A map  $\mathcal{T} : \mathcal{F}_{\mathbb{X}} \times \mathbb{X} \times \mathbb{U} \rightarrow [0, 1]$  is a controlled transition kernel if a) for every fixed  $A \in \mathcal{F}_{\mathbb{X}}$ , the map  $(x, u) \mapsto \mathcal{T}(A|x, u)$  is  $\sigma(\mathcal{F}_{\mathbb{X}} \times \mathcal{F}_{\mathbb{U}})$ -measurable, and b) For every fixed  $(x, u) \in \mathbb{X} \times \mathbb{U}$ , the map  $A \mapsto \mathcal{T}(A|x, u)$  is a probability measure on  $(\mathbb{X}, \mathcal{F}_{\mathbb{X}})$ .

We define the history (or path) space at time  $t$  as  $(\mathbb{H}_t, \mathcal{F}_{\mathbb{H}_t}) := (\mathbb{X}^{t+1} \times \mathbb{U}^t, \sigma(\mathcal{F}_{\mathbb{X}}^{t+1} \times \mathcal{F}_{\mathbb{U}}^t))$ . An admissible policy at time  $t$  is a measurable probability-measure-valued function  $\gamma_t : \mathbb{H}_t \rightarrow \mathcal{P}(\mathbb{U})$ , where  $\mathcal{P}(\mathbb{U})$  is the space of probability measures on  $(\mathbb{U}, \mathcal{F}_{\mathbb{U}})$  endowed with the weak topology. The distribution of the controlled Markov process  $\{X_t, U_t : t \geq 0\}$  is then determined by the policies  $\{\gamma_t : t \geq 0\}$  and the distribution of  $X_0$ .

At each time  $t$ , a cost  $c(X_t, U_t)$  is incurred according to some per-step cost function  $c : \mathbb{X} \times \mathbb{U} \rightarrow \mathbb{R}$ . We refer to the quadruple  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  as a discrete-time Markov decision process, abbreviated as MDP. In this paper, we consider the fully observed optimal control problem: a decision maker (DM) employs a sequence of admissible policies  $\gamma := \{\gamma_t\}_{t=0}^{\infty}$ ; at time  $t$ , the DM observes the realized history  $h_t := \{X_{[0,t]}, U_{[0,t-1]}\} \in \mathbb{H}_t$ , and makes a decision  $U_t \in \mathbb{U}$  according to the probability measure  $\gamma_t(h_t)$ ; the system incurs a cost  $c(X_t, U_t)$  and transits to a new state according to the distribution  $\mathcal{T}(\cdot|X_t, U_t)$ ; this process is then repeated. The goal is to minimize the accumulated cost with respect to the control policy  $\{\gamma_t\}_{t \in \mathbb{N}}$  according to some performance criterion. Denote the set of admissible policies by

$$\Gamma_A := \{\{\gamma_t\}_{t=0}^{\infty} : \forall t \in \mathbb{N}, \gamma_t : \mathbb{H}_t \rightarrow \mathcal{P}(\mathbb{U}) \text{ measurable}\}.$$

**(a).** For the discounted cost criterion with discount factor  $\beta \in (0, 1)$ , cost function  $c$ , transition kernel  $\mathcal{T}$ , and policy  $\gamma$ , define:

$$\begin{aligned} \text{the value function under policy } \gamma : & \quad J_{\beta}(c, \mathcal{T}, \gamma)(x) := \mathbb{E}^{\mathcal{T}, \gamma} \left[ \sum_{t=0}^{\infty} \beta^t c(X_t, U_t) \mid X_0 = x \right], \\ \text{the optimal value function :} & \quad J_{\beta}^*(c, \mathcal{T})(x) := \inf_{\gamma \in \Gamma_A} J_{\beta}(c, \mathcal{T}, \gamma)(x). \end{aligned}$$

**(b).** For the average cost criterion with cost function  $c$ , transition kernel  $\mathcal{T}$ , and policy  $\gamma$ , define:

$$\begin{aligned} \text{the value function under policy } \gamma : & \quad J_{\infty}(c, \mathcal{T}, \gamma)(x) = \limsup_{T \rightarrow \infty} \frac{1}{T} \mathbb{E}^{\mathcal{T}, \gamma} \left[ \sum_{t=0}^{T-1} c(X_t, U_t) \mid X_0 = x \right], \\ \text{the optimal value function :} & \quad J_{\infty}^*(c, \mathcal{T})(x) := \inf_{\gamma \in \Gamma_A} J_{\infty}(c, \mathcal{T}, \gamma)(x). \end{aligned}$$

Here,  $\mathbb{E}^{\mathcal{T}, \gamma}$  denotes the expectation under the controlled transition kernel  $\mathcal{T}$  and the policy  $\gamma$ .

**1.2. Problem Statement and Contributions** Consider a reference MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and a decision maker who wants to perform optimal control. It is seldom the case that the decision maker has complete knowledge about the MDP, specifically  $c$  and  $\mathcal{T}$ . It is therefore natural to consider the scheme that the decision maker first learns an approximate MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$  using samples from the reference MDP, obtain an optimal policy in the approximate MDP,  $\gamma_{\hat{c}, \mathcal{S}}^*$ , and apply it to the true MDP. A critical question arises: how much performance do we lose by applying an optimal policy derived from some approximate model? In this paper, we provide upper bounds on the loss in terms of the estimation error of the cost function and transition kernel, and then apply

our findings to empirical model learning. We also study the implications of our findings regarding sample complexity bounds which relate the mismatch error to the number of samples.

### Contributions.

- (i) In Section 2.2.1, we show a condition under which Lipschitz continuity holds for the optimal discounted cost value function (Theorem 4). In Section 2.2.2, we show conditions under which Lipschitz continuity holds for the optimal average cost value function (Theorem 5 and 6).
- (ii) In Section 2.3.1, we show conditions under which the robustness performance loss holds for the optimal discounted cost value function (Theorem 7 and 8). In Section 2.3.2, we show conditions under which the robustness performance loss holds for the optimal average cost value function (Theorem 9 and 10, Corollary 3).
- (iii) In Section 3, we relate our results to the case where robustness is in terms of the distribution of the driving noise (Corollary 8). When the distribution is estimated by its empirical version, we provide explicit rates of convergence in terms of the sample size under various moment conditions (Theorem 14).
- (iv) In Section 4, as a primary contribution, we establish empirical consistency and robustness to empirical model learning, along with explicit rates of convergence results that lead to sample complexity bounds. Specifically, we propose learning algorithms based on data and establish empirical consistency and sample complexity results for near-optimal policies under two types of data used for model learning: data obtained along a controlled sample path (Corollary 11 and 12), and data achieved via a device simulating the Markov kernel (Corollary 13 and 14).
- (v) As a by-product of our analysis we show continuous dependence of invariant measures on transition kernels in Section 5, recovering a result of [47], which we achieve by viewing the problem as a control-free version of our analysis and framework.

**1.3. Literature Review** Stochastic control and reinforcement learning under model misspecification are fundamental to their applications and have been extensively studied. For a review of the literature concerning finite state and action spaces, we refer readers to Chapter 1 of [1] for a review. See also related studies such as [21, 20, 22, 32, 36, 6].

In [32, 36], the authors study robustness to incorrect transition kernels, focusing primarily on asymptotic convergence under weak convergence and setwise convergence, as well as on strong uniformity results with respect to total variation convergence. Specifically, a robustness result is obtained, in an asymptotic convergence sense, for the discounted cost criterion in [32] and for the average cost criterion in [36], respectively, under *continuous* weak convergence of transition kernels; uniformity results under total variation convergence are also established. A unified perspective is given in [37], which considers quantized approximations as a special case. This robustness analysis is further extended to continuous-time models in [45].

We also note that [38, Theorem 5.1] establishes continuity results for approximate models under discounted cost criterion and provides a set convergence result for sets of optimal control actions. However, this set convergence result is inconclusive for robustness without further assumptions on the true system model. Additionally, [42] is a related work which studies the continuity of the discounted cost value function for fully observed models under a general metric, known as the integral probability metric, which encompasses both the total variation metric and the Kantorovich metric in different configurations (which is not weaker than the metrics leading to weak convergence). An analogous result under the average cost criterion appeared in [24].

In [19, 17, 18, 16], the authors address the following disturbance distribution approximation problem: consider a stochastic dynamical system  $X_{t+1} = f(X_t, U_t, W_t)$ , where  $\{W_t\}_{t=0}^{\infty}$  is an i.i.d process with distribution  $\mu$ . If an alternative distribution  $\nu$  is proposed and an approximate optimal policy is computed under  $\nu$ , the objective is to characterize the relationship between the robustness performance loss and a certain distance between  $\mu$  and  $\nu$ . In [16], a robustness result is given

for the discounted cost criterion with the bounded-Lipschitz distance between  $\mu$  and  $\nu$ . In [19], similar results are obtained for the average cost criterion, using either the total variation distance or bounded-Lipschitz distance, via a minorization condition assumption.

Results on robustness to weak convergence in [32, 36] and to Wasserstein convergence in [17, 18, 10] imply empirical consistency with i.i.d. learning of models, as noted in these studies. In [17, 18], the authors consider a learning scenario where  $\nu = \frac{1}{n+1} \sum_{i=0}^n \delta_{w_i}$  is the empirical measure of  $\mu$  under samples  $\{w_i\}_{i=0}^n$ . The chosen distance between measures is Wasserstein-1 distance, with the discounted cost criterion considered in [17], and the average cost criterion in [18]. We show that the disturbance distribution approximation problem can be viewed as a special case of model approximation, and through Theorem 8 and 10, we obtain results analogous to those in [17, 18], with relaxed assumptions and improved bounds. A further related result on empirical consistency from a different perspective is presented in [23].

In [21], the authors focus on the discounted cost criterion, with a similar result established in the more recent work [6]. Notably, both papers use a weighted-norm-induced metric for the distance between cost functions to account for possibly unbounded costs. In this paper, we use the sup-norm-induced metric as the distance between cost functions and save discussions on unbounded costs for future work. We also acknowledge that Theorem 4 and 8 originally appeared in [6] as Corollary 1, and our Theorem 7 is a refinement of Theorem 5 of the same paper. In [20], the authors considered model approximation under the average cost criterion, utilizing either total variation or uniform bounded-Lipschitz distance. The authors obtained results similar to Theorem 9 by imposing contraction assumptions on Bellman operators of the reference and approximate models. In [22], the focus is restricted to approximating transition kernels, with an upper bound provided on the relative error of robustness performance loss.

A recently proposed approach motivated by robust stochastic optimization [43, 41, 15], referred to as distributional stochastic control [59, 56], considers a related, but distinct setting. The key difference is that in distributional robustness, the disturbance distribution is allowed to vary over time. In [59], the authors study the case of the Wasserstein ball, while in [56], both Wasserstein and f-divergence balls are considered.

In [10], the authors consider a setting where the transition kernel admits a  $(x, u)$ -dependent density with respect to some *known* reference measure  $\mu$  on  $(\mathbb{X}, \mathcal{F}_{\mathbb{X}})$ , and the decision maker uses an empirical estimate of  $\mu$  to approximate the model discretely. In [53, 60, 48, 50, 49], the authors consider static discretizations of state and action spaces and provide error bounds for compact state and action spaces and asymptotic analyses for more general settings. We refer readers to [49] for a review of related results, which also obtain several regularity results under the Wasserstein distance considered in this paper. Such discretization schemes have seen applications in analyzing the convergence of Q-learning [33, 31, 35]. In [54], the authors consider a generalization of the discretization scheme by introducing a class of approximate operators named averagers. However, their results are on the continuity of average cost value iteration, and require restrictive assumptions on transition kernels. In Section 4.1, we discuss how discretization in Wasserstein regular MDPs with compact spaces can be viewed as a special case of Wasserstein model approximation, and thus several results in [31] and [34] can be recovered from our analysis. This further motivates us to propose a sampling-based discretization algorithm in the spirit of, but with distinct assumptions and conditions from, [10], which we discuss in Section 4.2.

The key assumption in our paper is Lipschitz continuity in components of the MDPs. This has been a standard assumption in related works [49, 6]. To our knowledge, [27] is the first paper to study the assumption. In [46], the authors restrict their attention to Lipschitz continuous policies and show the Lipschitz continuity of the Q-factor under Lipschitz MDP assumptions. In [44], a policy gradient method is studied. In [2], the authors prove a continuity result on the value function (Theorem 2, [2]), although under restrictive assumptions.

**1.4. Space of Probability Measures and Transition Kernels** In this section, we collect some facts from probability theory to make the previous and following discussions precise.

**1.4.1. Convergence of Probability Measures** Given a complete separable metric space  $(\mathbb{X}, d_{\mathbb{X}})$  endowed with its Borel  $\sigma$ -algebra, denote by  $\mathcal{P}(\mathbb{X})$  the space of probability measures on it.

DEFINITION 2 (WEAK CONVERGENCE, P.292, [9]). A sequence of probability measures  $\{\mu_n\} \subset \mathcal{P}(\mathbb{X})$  converges weakly to  $\mu \in \mathcal{P}(\mathbb{X})$  if for all continuous and bounded  $f : \mathbb{X} \rightarrow \mathbb{R}$ ,

$$\int_{\mathbb{X}} f(x) \mu_n(dx) \rightarrow \int_{\mathbb{X}} f(x) \mu(dx).$$

DEFINITION 3 (TOTAL VARIATION DISTANCE, P.292, [9]). For two probability measures  $\mu, \nu \in \mathcal{P}(\mathbb{X})$ , the total variation distance between them is defined as

$$\|\mu - \nu\|_{TV} := 2 \sup_{A \in \mathcal{F}_{\mathbb{X}}} |\mu(A) - \nu(A)| = \sup_{\|f\|_{\infty} \leq 1} \left| \int_{\mathbb{X}} f(x) \mu(dx) - \int_{\mathbb{X}} f(x) \nu(dx) \right|,$$

where  $\|f\|_{\infty} := \sup_{x \in \mathbb{X}} |f(x)|$  is the sup-norm of a function  $f : \mathbb{X} \rightarrow \mathbb{R}$ .

DEFINITION 4 (WASSERSTEIN-P DISTANCE, DEFINITION 3.1.1, [12]). Let  $1 \leq p < \infty$ . For two probability measures  $\mu, \nu \in \mathcal{P}(\mathbb{X})$ , where  $\mathbb{X}$  is locally compact and separable, assume that for some  $x_0 \in \mathbb{X}$ ,  $\int_{\mathbb{X}} d_{\mathbb{X}}(x, x_0)^p \mu(dx) < \infty$  and  $\int_{\mathbb{X}} d_{\mathbb{X}}(x, x_0)^p \nu(dx) < \infty$ . The Wasserstein- $p$  distance is defined as follows:

$$\mathcal{W}_p(\mu, \nu) := \inf \left\{ \left( \int_{\mathbb{X} \times \mathbb{X}} d_{\mathbb{X}}(x, y)^p \pi(dx, dy) \right)^{\frac{1}{p}} : \pi \text{ is a coupling between } \mu, \nu \right\},$$

where a coupling  $\pi$  between two probability measures  $\mu, \nu \in \mathcal{P}(\mathbb{X})$  is defined as any probability measure on  $(\mathbb{X}^2, \sigma(\mathcal{F}_{\mathbb{X}} \times \mathcal{F}_{\mathbb{X}}))$  such that for any  $A \in \mathcal{F}_{\mathbb{X}}$ :  $\pi(A \times \mathbb{X}) = \mu(A)$ ,  $\pi(\mathbb{X} \times A) = \nu(A)$ .

REMARK 1. For  $p = 1$ , the Wasserstein distance admits a dual formulation (Kantorovich–Rubinstein formula, Chapter 6, [55]):

$$\mathcal{W}_1(\mu, \nu) = \sup_{\|f\|_{\text{Lip}} \leq 1} \left\{ \int_{\mathbb{X}} f(x) (\mu(dx) - \nu(dx)) \right\},$$

where  $\|f\|_{\text{Lip}} := \inf_{x \neq y \in \mathbb{X}} \frac{|f(x) - f(y)|}{d_{\mathbb{X}}(x, y)}$  is the Lipschitz norm of a function  $f : \mathbb{X} \rightarrow \mathbb{R}$ .

A key motivation for this paper is to show that if an MDP is regular in the Wasserstein sense, as detailed in Section 2.4, the MDP can be approximated via sampling, and the optimal policies obtained from the sampled MDPs are near-optimal. Specifically, the performance loss shares the same diminishing rate as the distance between the empirical and population-level distributions.

**1.4.2. On the Space of Controlled Transition Kernels** In this section, we discuss convergence criteria in the space of controlled probability kernels that are useful later on. Note that all controlled probability kernels can also be viewed as measure-valued functions:  $\mathcal{T} : \mathbb{X} \times \mathbb{U} \rightarrow \mathcal{P}(\mathbb{X})$ . Therefore, by equipping the space of probability measures  $\mathcal{P}(\mathbb{X})$  with a specific convergence criterion, we can extend the convergence criterion to the space of controlled probability kernels by combining it with a criterion for function convergence. Instead of listing every possible combination, we provide several examples to illustrate the ideas and naming conventions we use.

**Example 1.1 (Pointwise weak convergence)** Given a sequence of controlled probability kernels  $(\mathcal{T}_n)$ , we say  $\mathcal{T}_n$  converges pointwise weakly to a controlled probability kernel  $\mathcal{T}$  if,

$$\int_{\mathbb{X}} f(y) \mathcal{T}_n(dy|x, u) \rightarrow \int_{\mathbb{X}} f(y) \mathcal{T}(dy|x, u), \forall (x, u) \in \mathbb{X} \times \mathbb{U}, f \text{ Lipschitz and bounded.}$$

**Example 1.2 (Uniform convergence in Wasserstein- $p$  distance)** *Given a sequence of controlled probability kernels  $(\mathcal{T}_n)$  and  $p \geq 1$ , we say  $\mathcal{T}_n$  converges uniformly in Wasserstein- $p$  distance to a controlled probability kernel  $\mathcal{T}$  if  $\sup_{(x,u) \in \mathbb{X} \times \mathbb{U}} \mathcal{W}_p(\mathcal{T}_n(\cdot|x,u), \mathcal{T}(\cdot|x,u)) \rightarrow 0$ .*

REMARK 2 (METRICS ON SPACE OF CONTROLLED PROBABILITY KERNELS). We note here that  $\sup_{(x,u) \in \mathbb{X} \times \mathbb{U}} \mathcal{W}_p(\mathcal{T}(\cdot|x,u), \mathcal{S}(\cdot|x,u))$  defines a metric on the space of controlled probability kernels as a composition of  $\|\cdot\|_\infty$  norm induced metric and Wasserstein- $p$  distance. We refer to this metric as Uniform Wasserstein- $p$  distance and denote it by  $d_{\mathcal{W}_p}(\mathcal{T}, \mathcal{S})$  in later sections. We can similarly define, for example, uniform total variation distance as

$$d_{TV}(\mathcal{T}, \mathcal{S}) := \sup_{(x,u) \in (\mathbb{X}, \mathbb{U})} \|\mathcal{T}(\cdot|x,u) - \mathcal{S}(\cdot|x,u)\|_{TV}.$$

## 2. Regularity of Value Functions and Robustness to Model Approximation under Uniform Wasserstein Distance

**2.1. Optimality Equations** In this section, we present the basic assumptions that allow us to characterize solutions to stochastic optimal control problems by fixed point equations, and furthermore, allow us to restrict attention to deterministic stationary control policies  $\gamma: \mathbb{X} \rightarrow \mathbb{U}$ . We denote the set of continuous and bounded measurable functions mapping from  $(\mathbb{X}, \mathcal{F}_{\mathbb{X}})$  to  $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$  by  $C_b(\mathbb{X})$ .

ASSUMPTION 1. *Unless otherwise noted, we assume that any MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  in this paper satisfies the following:*

- (a).  $\mathbb{U}$  is compact.
- (b).  $c: \mathbb{X} \times \mathbb{U} \rightarrow \mathbb{R}$  is nonnegative, bounded ( $\|c\|_\infty < \infty$ ), and continuous on both  $\mathbb{X}$  and  $\mathbb{U}$ .
- (c). For any  $v \in C_b(\mathbb{X})$ ,  $\int_{\mathbb{X}} v(y) \mathcal{T}(dy|x, u)$  is a continuous on  $\mathbb{X}$  and  $\mathbb{U}$ , i.e.  $\mathcal{T}$  is weakly continuous on  $\mathbb{X} \times \mathbb{U}$ .

THEOREM 1 (DCOE, Theorem 5.2.1 and 5.5.2, [61]). *Consider an MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and a discount factor  $\beta \in (0, 1)$ . Suppose Assumption 1 holds.*

- (a). *The optimal discounted cost function  $J_\beta^*(c, \mathcal{T}) \in C_b(\mathbb{X})$  is the unique solution to the following fixed point equation in  $v \in C_b(\mathbb{X})$ , which we refer to as DCOE:*

$$v(x) = \inf_{u \in \mathbb{U}} \left\{ c(x, u) + \beta \int_{\mathbb{X}} v(y) \mathcal{T}(dy|x, u) \right\}, \forall x \in \mathbb{X}.$$

- (b). *There exists a deterministic stationary policy  $\gamma^*: \mathbb{X} \rightarrow \mathbb{U}$  such that*

$$v(x) = c(x, \gamma^*(x)) + \beta \int_{\mathbb{X}} v(y) \mathcal{T}(dy|x, \gamma^*(x)), \forall x \in \mathbb{X},$$

and  $J_\beta(c, \mathcal{T}, \gamma^*)(x) = J_\beta^*(c, \mathcal{T})(x)$ ,  $\forall x \in \mathbb{X}$ .

We note that part (a) follows from the fact that if we define the right-hand side of DCOE as an operator on  $C_b(\mathbb{X})$ , which we refer to as the Bellman operator  $\mathbb{T}$ , it can be shown that  $\mathbb{T}$  is a contraction on  $(C_b(\mathbb{X}), \|\cdot\|_\infty)$ , i.e. for any  $f_1, f_2 \in C_b(\mathbb{X})$ ,  $\mathbb{T}: C_b(\mathbb{X}) \rightarrow C_b(\mathbb{X})$  satisfies

$$\|\mathbb{T}f_1 - \mathbb{T}f_2\|_\infty \leq q \|f_1 - f_2\|_\infty,$$

where  $q \in [0, 1)$  is a constant. Thus, by Banach fixed-point theorem (see e.g. Theorem 3.1.7 of [61]), the DCOE admits a unique solution (see p.19 of [25]). As we discuss in Section 2.4, this fact is important for characterizing the Lipschitz continuity of value functions under suitable assumptions.

REMARK 3 (DISCOUNTED COST BELLMAN CONSISTENCY EQUATION). Given a fixed deterministic stationary policy  $\gamma$ , we consider the equation

$$v(x) = c(x, \gamma(x)) + \beta \int_{\mathbb{X}} v(y) \mathcal{T}(dy|x, \gamma(x)), \forall x \in \mathbb{X}.$$

Defining the right-hand side as an operator  $\mathbb{T}^\gamma$ , it can be shown that  $\mathbb{T}^\gamma$  is also a contraction, and thus admits a unique solution on the set of bounded measurable functions (as  $\gamma$  can be discontinuous). Furthermore, the solution is  $J_\beta(c, \mathcal{T}, \gamma)$ . We refer to this equation as the discounted cost Bellman consistency equation for  $\gamma$  (see Section 1.1.2, [1]).

For the average cost criterion, we present two additional sets of assumptions, each sufficient for the optimality equation. We apply these assumptions in different settings.

DEFINITION 5 (MINORIZATION CONDITION FOR KERNELS). The controlled transition kernel  $\mathcal{T} : \mathcal{F}_{\mathbb{X}} \times \mathbb{X} \times \mathbb{U} \rightarrow [0, 1]$  is said to satisfy the minorization condition if there exists a probability measure  $\rho$  on  $(\mathbb{X}, \mathcal{F}_{\mathbb{X}})$  and constant  $\epsilon > 0$  such that  $\forall x \in \mathbb{X}, u \in \mathbb{U}$ , and  $A \in \mathcal{F}_{\mathbb{X}}$ ,

$$\mathcal{T}(A|x, u) \geq \epsilon \rho(A).$$

THEOREM 2 (ACOE, Theorem 5.2.1 and 7.2.1, [61]). Consider an MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  such that  $\mathcal{T}$  satisfies the minorization condition, and suppose Assumption 1 holds. Then there exists a constant  $g$  and  $h \in C_b(\mathbb{X})$  such that the following fixed point equation, referred to as ACOE, holds:

$$g + h(x) = \inf_{u \in \mathbb{U}} \left\{ c(x, u) + \int_{\mathbb{X}} h(y) \mathcal{T}(dy|x, u) \right\}, \forall x \in \mathbb{X},$$

where  $g = \inf_{\gamma \in \Gamma_A} J_\infty(c, \mathcal{T}, \gamma)$ . Furthermore, there exists a deterministic stationary policy  $\gamma^* : \mathbb{X} \rightarrow \mathbb{U}$  such that

$$g + h(x) = c(x, \gamma^*(x)) + \int_{\mathbb{X}} h(y) \mathcal{T}(dy|x, \gamma^*(x)), \forall x \in \mathbb{X},$$

and  $J_\infty(c, \mathcal{T}, \gamma^*)(x) = g, \forall x \in \mathbb{X}$ .

REMARK 4. We refer to  $(g^*, h^*, \gamma^*)$  satisfying ACOE as a canonical triplet. To emphasize the dependence on the cost function  $c$  and kernel  $\mathcal{T}$ , we use the following notation for the canonical triplet:  $(g_{c, \mathcal{T}}^*, h_{c, \mathcal{T}}^*, \gamma_{c, \mathcal{T}}^*)$ . Note that  $g_{c, \mathcal{T}}^* = J_\infty^*(c, \mathcal{T})$ .

The above result relies on the fact that if the transition kernel satisfies the minorization condition with a scaling constant  $\epsilon > 0$  and minorizing probability measure  $\rho$ , the operator

$$\mathbb{T}v(x) := \inf_{u \in \mathbb{U}} \left\{ c(x, u) + \int_{\mathbb{X}} v(y) (\mathcal{T}(dy|x, u) - \epsilon \rho(dy)) \right\}, \forall x \in \mathbb{X}$$

is a contraction on  $C_b(\mathbb{X})$ , and thus admits a unique fixed point. We let  $h$  be the fixed point of  $\mathbb{T}$ , and let  $g := \epsilon \int_{\mathbb{X}} h(x) \rho(dx)$ . It is then clear that  $g$  and  $h$  satisfy ACOE.

REMARK 5 (AVERAGE COST BELLMAN CONSISTENCY EQUATION). Similar to the case of DCOE, under the minorization condition, for a fixed deterministic stationary policy  $\gamma$ , the operator

$$\mathbb{T}^\gamma v(x) := c(x, \gamma(x)) + \int_{\mathbb{X}} v(y) (\mathcal{T}(dy|x, \gamma(x)) - \epsilon \rho(dy)), \forall x \in \mathbb{X}$$

is a contraction operator. Following the above constructive approach, we have that there exists a constant  $g^\gamma$  and a bounded measurable function  $h^\gamma$  such that

$$g^\gamma + h^\gamma(x) = c(x, \gamma(x)) + \int_{\mathbb{X}} h^\gamma(y) \mathcal{T}(dy|x, \gamma(x)),$$

where  $g^\gamma = J_\infty(c, \mathcal{T}, \gamma)$ . We refer to this equation as the average cost Bellman consistency equation.

**ASSUMPTION 2 (Assumption 5.4.1 of [26]).** For an MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$ , there exists a state  $z \in \mathbb{X}$  and constants  $\alpha \in (0, 1)$ ,  $M > 0$  such that

$$(1 - \beta)J_{\beta}^*(c, \mathcal{T})(z) \leq M, \forall \beta \in [\alpha, 1).$$

Moreover, there exists a constant  $N \geq 0$  and a nonnegative function  $b: \mathbb{X} \rightarrow \mathbb{R}$  such that with  $h_{\beta}(c, \mathcal{T})(x) := J_{\beta}^*(c, \mathcal{T})(x) - J_{\beta}^*(c, \mathcal{T})(z)$ ,

$$-N \leq h_{\beta}(c, \mathcal{T})(x) \leq b(x), \forall x \in \mathbb{X}, \beta \in [\alpha, 1).$$

We note that the first part of the assumption is implied by Assumption 1, under which the per-step cost is bounded. A direct consequence of Assumption 2 is the following:

**LEMMA 1 (Lemma, p.88 of [26]).** Under Assumption 2 for the MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$ , there exists a sequence  $\beta(n) \rightarrow 1$  such that

$$\lim_{n \rightarrow \infty} (1 - \beta(n))J_{\beta(n)}^*(c, \mathcal{T})(x) = g_{c, \mathcal{T}}^* = J_{\infty}^*(c, \mathcal{T}), \forall x \in \mathbb{X}.$$

**THEOREM 3 (ACOE, Theorem 7.3.3, [61]).** Under Assumption 1 and 2 for the MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$ , there exists a triplet  $(g^*, h^*, \gamma^*)$ , where  $g^*$  is a constant,  $h^* \in C_b(\mathbb{X})$ , and  $\gamma^*$  is a deterministic stationary policy, such that  $g^* = \inf_{\gamma \in \Gamma_A} J_{\infty}(c, \mathcal{T}, \gamma)(x)$ , and  $J_{\infty}(c, \mathcal{T}, \gamma^*)(x) = g^*$ ,  $\forall x \in \mathbb{X}$ .

**2.2. Continuity of Optimal Value Functions** We begin with continuity results for optimal value functions, which serve as important intermediate steps in our robustness analysis in Subsection 2.3. We consider the discounted cost criterion in Subsection 2.2.1 and the average cost criterion in Subsection 2.2.2. Additionally, we note similar analyses in [36, 32].

### 2.2.1. Discounted Cost Criterion

**THEOREM 4.** Suppose that Assumption 1 holds for two MDPs, (reference)  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and (approximation)  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$ . If  $J_{\beta}^*(c, \mathcal{T})$  is Lipschitz continuous, then

$$\|J_{\beta}^*(c, \mathcal{T}) - J_{\beta}^*(\hat{c}, \mathcal{S})\|_{\infty} \leq \frac{1}{1 - \beta} \|c - \hat{c}\|_{\infty} + \frac{\beta}{1 - \beta} \|J_{\beta}^*(c, \mathcal{T})\|_{Lip} d_{W_1}(\mathcal{T}, \mathcal{S}).$$

*Proof.* For ease of notation, we use the following abbreviation:

$$v_{c, \mathcal{T}} := J_{\beta}^*(c, \mathcal{T}), \quad v_{\hat{c}, \mathcal{S}} := J_{\beta}^*(\hat{c}, \mathcal{S}). \quad (1)$$

By Theorem 1, we have that for any  $x \in \mathbb{X}$ :

$$v_{c, \mathcal{T}}(x) = \inf_{u \in \mathbb{U}} \left( c(x, u) + \beta \int v_{\mathcal{T}}(x') \mathcal{T}(dx'|x, u) \right), \quad v_{\hat{c}, \mathcal{S}}(x) = \inf_{u \in \mathbb{U}} \left( \hat{c}(x, u) + \beta \int v_{\mathcal{S}}(x') \mathcal{S}(dx'|x, u) \right).$$

Therefore, by triangular inequality, we have that for any  $x \in \mathbb{X}$ :

$$\begin{aligned} |v_{c, \mathcal{T}}(x) - v_{\hat{c}, \mathcal{S}}(x)| &\leq \sup_{u \in \mathbb{U}} |c(x, u) - \hat{c}(x, u)| + \beta \sup_{u \in \mathbb{U}} \left| \int v_{c, \mathcal{T}}(x') \mathcal{T}(dx'|x, u) - \int v_{\hat{c}, \mathcal{S}}(x') \mathcal{S}(dx'|x, u) \right| \\ &\leq \|c - \hat{c}\|_{\infty} + \beta \sup_{u \in \mathbb{U}} \left| \int v_{c, \mathcal{T}}(x') \mathcal{T}(dx'|x, u) - \int v_{c, \mathcal{T}}(x') \mathcal{S}(dx'|x, u) \right. \\ &\quad \left. + \int v_{c, \mathcal{T}}(x') \mathcal{S}(dx'|x, u) - \int v_{\hat{c}, \mathcal{S}}(x') \mathcal{S}(dx'|x, u) \right| \\ &\leq \|c - \hat{c}\|_{\infty} + \beta \sup_{u \in \mathbb{U}} \left| \int v_{c, \mathcal{T}}(x') \mathcal{T}(dx'|x, u) - \int v_{c, \mathcal{T}}(x') \mathcal{S}(dx'|x, u) \right| \end{aligned}$$



$$+ \beta \sup_{u \in \mathbb{U}} \left| \int v_{c,\mathcal{T}}(x') \mathcal{S}(dx'|x, u) - \int v_{\hat{c},\mathcal{S}}(x') \mathcal{S}(dx'|x, u) \right|.$$

Due to the dual formulation of  $\mathcal{W}_1$ , we have that for any  $x \in \mathbb{X}$ :

$$\begin{aligned} |v_{c,\mathcal{T}}(x) - v_{\hat{c},\mathcal{S}}(x)| &\leq \|c - \hat{c}\|_\infty + \beta \|v_{c,\mathcal{T}}\|_{\text{Lip}} \sup_{x,u} (\mathcal{W}_1(\mathcal{T}(\cdot|x, u), \mathcal{S}(\cdot|x, u))) \\ &\quad + \beta \sup_{u \in \mathbb{U}} \left( \int |v_{c,\mathcal{T}}(x') - v_{\hat{c},\mathcal{S}}(x')| \mathcal{S}(dx'|x, u) \right) \\ &\leq \|c - \hat{c}\|_\infty + \beta \|v_{c,\mathcal{T}}\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) + \beta \sup_{u \in \mathbb{U}} \left( \sup_{x \in \mathbb{X}} |v_{c,\mathcal{T}}(x) - v_{\hat{c},\mathcal{S}}(x)| \right) \\ &= \|c - \hat{c}\|_\infty + \beta \|v_{c,\mathcal{T}}\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) + \beta \sup_{x \in \mathbb{X}} |v_{c,\mathcal{T}}(x) - v_{\hat{c},\mathcal{S}}(x)|. \end{aligned}$$

Finally, if we take the supremum over all  $x \in \mathbb{X}$  on the left-hand side and by boundedness of  $v_{c,\mathcal{T}}$  and  $v_{\hat{c},\mathcal{S}}$ , which follows from  $\|c\|_\infty, \|\hat{c}\|_\infty < \infty$ , we can reorganize and obtain

$$\sup_{x \in \mathbb{X}} |v_{c,\mathcal{T}}(x) - v_{\hat{c},\mathcal{S}}(x)| \leq \frac{1}{1 - \beta} \|c - \hat{c}\|_\infty + \frac{\beta}{1 - \beta} \|v_{c,\mathcal{T}}\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).$$

□

**2.2.2. Average Cost Criterion** In the following, we first provide an extension of Theorem 4 to the average cost criterion under the minorization condition (Definition 5). Next, we present an alternative approach based on Theorem 4 that allows us to achieve a similar result without the minorization condition, utilizing Lemma 1.

**ASSUMPTION 3 (Minorization Condition for Weak Model Approximation).** *Given two controlled markov processes, (reference)  $(\mathbb{X}, \mathbb{U}, \mathcal{T})$  and (approximation)  $(\mathbb{X}, \mathbb{U}, \mathcal{S})$ , we assume there exists a constant  $\epsilon > 0$  and probability measures  $\rho, \tau \in \mathcal{P}((\mathbb{X}, \mathcal{F}_{\mathbb{X}}))$ , such that  $\forall x \in \mathbb{X}, u \in \mathbb{U}$ , and  $A \in \mathcal{F}_{\mathbb{X}}$ , the following hold  $\mathcal{T}(A|x, u) \geq \epsilon \rho(A)$ ,  $\mathcal{S}(A|x, u) \geq \epsilon \tau(A)$ .*

**THEOREM 5.** *Suppose that Assumption 1 holds for two MDPs, (reference)  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and (approximation)  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$ , both of which also satisfy Assumption 3 with some scaling constant  $\epsilon > 0$  and minorizing probability measures  $\rho$  and  $\tau$  respectively. Denote by the pairs  $(g_{c,\mathcal{T}}^*, h_{c,\mathcal{T}}^*)$  satisfying*

$$h_{c,\mathcal{T}}^*(x) = \inf_{u \in \mathbb{U}} \left( c(x, u) + \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') (\mathcal{T}(dx'|x, u) - \epsilon \rho(dx')) \right) \text{ for } x \in \mathbb{X}, \quad g_{c,\mathcal{T}}^* = \epsilon \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \rho(dx'),$$

and  $(g_{\hat{c},\mathcal{S}}^*, h_{\hat{c},\mathcal{S}}^*)$  satisfying

$$h_{\hat{c},\mathcal{S}}^*(x) = \inf_{u \in \mathbb{U}} \left( \hat{c}(x, u) + \int_{\mathbb{X}} h_{\hat{c},\mathcal{S}}^*(x') (\mathcal{S}(dx'|x, u) - \epsilon \tau(dx')) \right) \text{ for } x \in \mathbb{X}, \quad g_{\hat{c},\mathcal{S}}^* = \epsilon \int_{\mathbb{X}} h_{\hat{c},\mathcal{S}}^*(x') \tau(dx'),$$

whose existence is guaranteed by Theorem 2. If  $h_{c,\mathcal{T}}^*$  is Lipschitz continuous, we have

$$\begin{aligned} \|h_{c,\mathcal{T}}^* - h_{\hat{c},\mathcal{S}}^*\|_\infty &\leq \frac{1}{\epsilon} \left( \|c - \hat{c}\|_\infty + \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) \right) + \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} \mathcal{W}_1(\rho, \tau), \\ |g_{c,\mathcal{T}}^* - g_{\hat{c},\mathcal{S}}^*| &\leq \|c - \hat{c}\|_\infty + \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}). \end{aligned}$$

*Proof.* Define  $\mathcal{T}'(\cdot|x, u) := \mathcal{T}(\cdot|x, u) - \epsilon \rho(\cdot)$  and  $\mathcal{S}'(\cdot|x, u) = \mathcal{S}(\cdot|x, u) - \epsilon \tau(\cdot)$  for  $x \in \mathbb{X}, u \in \mathbb{U}$ . Applying the minorized fixed point equation, we have that for any  $x \in \mathbb{X}$ :

$$|h_{c,\mathcal{T}}^*(x) - h_{\hat{c},\mathcal{S}}^*(x)| \leq \sup_{u \in \mathbb{U}} |c(x, u) - \hat{c}(x, u)|$$

$$\begin{aligned}
& + \sup_{u \in \mathbb{U}} \left| \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') (\mathcal{T}(dx'|x, u) - \epsilon \rho(dx')) - \int_{\mathbb{X}} h_{\hat{c},\mathcal{S}}^*(x') (\mathcal{S}(dx'|x, u) - \epsilon \tau(dx')) \right| \\
& \leq \|c - \hat{c}\|_{\infty} + \sup_{u \in \mathbb{U}} \left| \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') (\mathcal{T}(dx'|x, u) - \epsilon \tau(dx')) - \int_{\mathbb{X}} h_{\hat{c},\mathcal{S}}^*(x') (\mathcal{S}(dx'|x, u) - \epsilon \tau(dx')) \right. \\
& \quad \left. + \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \epsilon \tau(dx') - \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \epsilon \rho(dx') \right| \\
& \leq \|c - \hat{c}\|_{\infty} + \sup_{u \in \mathbb{U}} \left| \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \mathcal{T}'(dx'|x, u) - \int_{\mathbb{X}} h_{\hat{c},\mathcal{S}}^*(x') \mathcal{S}'(dx'|x, u) \right| \\
& \quad + \epsilon \left| \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \tau(dx') - \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \rho(dx') \right| \\
& \leq \|c - \hat{c}\|_{\infty} + \epsilon \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} \mathcal{W}_1(\rho, \tau) + \sup_{u \in \mathbb{U}} \left| \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \mathcal{T}'(dx'|x, u) - \int_{\mathbb{X}} h_{\hat{c},\mathcal{S}}^*(x') \mathcal{S}'(dx'|x, u) \right|,
\end{aligned}$$

where in the last step we apply the dual formulation of Wasserstein-1 distance. We now focus on the last term. By the triangle inequality and the dual formulation of the Wasserstein-1 distance, we have that for any  $x \in \mathbb{X}$ :

$$\begin{aligned}
& \sup_{u \in \mathbb{U}} \left| \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \mathcal{T}'(dx'|x, u) - \int_{\mathbb{X}} h_{\hat{c},\mathcal{S}}^*(x') \mathcal{S}'(dx'|x, u) \right| \\
& \leq \sup_{u \in \mathbb{U}} \left| \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \mathcal{T}'(dx'|x, u) - \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \mathcal{S}'(dx'|x, u) + \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \mathcal{S}'(dx'|x, u) - \int_{\mathbb{X}} h_{\hat{c},\mathcal{S}}^*(x') \mathcal{S}'(dx'|x, u) \right| \\
& \leq \sup_{u \in \mathbb{U}} \left| \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \mathcal{T}'(dx'|x, u) - \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \mathcal{S}'(dx'|x, u) \right| + \sup_{u \in \mathbb{U}} \left| \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \mathcal{S}'(dx'|x, u) - \int_{\mathbb{X}} h_{\hat{c},\mathcal{S}}^*(x') \mathcal{S}'(dx'|x, u) \right| \\
& \leq \sup_{u \in \mathbb{U}} \left| \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') (\mathcal{T}'(dx'|x, u) + \epsilon \tau(dx')) - \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') (\mathcal{S}'(dx'|x, u) + \epsilon \tau(dx')) \right| \\
& \quad + \sup_{u \in \mathbb{U}} \left| \int_{\mathbb{X}} (h_{c,\mathcal{T}}^*(x') - h_{\hat{c},\mathcal{S}}^*(x')) (\mathcal{S}(dx'|x, u) - \epsilon \tau(dx')) \right| \\
& \leq \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} \sup_{(x,u) \in \mathbb{X} \times \mathbb{U}} (\mathcal{W}_1(\mathcal{T}(\cdot|x, u), \mathcal{S}(\cdot|x, u))) + \int_{\mathbb{X}} \sup_{u \in \mathbb{U}} |h_{c,\mathcal{T}}^*(x') - h_{\hat{c},\mathcal{S}}^*(x')| (\mathcal{S}(dx'|x, u) - \epsilon \tau(dx')) \\
& = \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) + (1 - \epsilon) \|h_{c,\mathcal{T}}^* - h_{\hat{c},\mathcal{S}}^*\|_{\infty}.
\end{aligned}$$

Thus,  $\|h_{c,\mathcal{T}}^* - h_{\hat{c},\mathcal{S}}^*\|_{\infty} \leq \frac{1}{\epsilon} (\|c - \hat{c}\|_{\infty} + \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S})) + \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} \mathcal{W}_1(\rho, \tau)$ . We proceed to bound the other term of concern:

$$\begin{aligned}
|g_{c,\mathcal{T}}^* - g_{\hat{c},\mathcal{S}}^*| & = \epsilon \left| \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \rho(dx') - \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \tau(dx') + \int_{\mathbb{X}} h_{c,\mathcal{T}}^*(x') \tau(dx') - \int_{\mathbb{X}} h_{\hat{c},\mathcal{S}}^*(x') \tau(dx') \right| \\
& \leq \epsilon \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} \mathcal{W}_1(\rho, \tau) + \epsilon \|h_{c,\mathcal{T}}^* - h_{\hat{c},\mathcal{S}}^*\|_{\infty} \\
& \leq \|c - \hat{c}\|_{\infty} + \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) + 2\epsilon \mathcal{W}_1(\rho, \tau).
\end{aligned}$$

We note that the second bound holds for any  $\epsilon > 0$ , and therefore

$$\begin{aligned}
|g_{c,\mathcal{T}}^* - g_{\hat{c},\mathcal{S}}^*| & \leq \inf_{\epsilon > 0} \left( \|c - \hat{c}\|_{\infty} + \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) + 2\epsilon \mathcal{W}_1(\rho, \tau) \right) \\
& = \|c - \hat{c}\|_{\infty} + \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).
\end{aligned}$$

□

Next, we present an approach to achieve similar results under different assumptions, utilizing Lemma 1.

**THEOREM 6.** *Suppose that Assumption 1 holds for two MDPs, (reference)  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and (approximation)  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$ . Further, assume that there exists a constant  $\alpha \in (0, 1)$  such that  $\forall \beta \in$*

$[\alpha, 1)$ ,  $\|J_\beta^*(c, \mathcal{T})\|_{Lip} < \infty$ ,  $\lim_{\beta \rightarrow 1} \|J_\beta^*(c, \mathcal{T})\|_{Lip}$  exists, and for any  $x \in \mathbb{X}$  there exists a sequence  $\beta_n(x) \rightarrow 1$  such that  $(1 - \beta_n(x))J_{\beta_n(x)}^*(\hat{c}, \mathcal{S})(x) \rightarrow g_{\hat{c}, \mathcal{S}}^*$ , then

$$\|J_\infty^*(c, \mathcal{T}) - J_\infty^*(\hat{c}, \mathcal{S})\|_\infty \leq \|c - \hat{c}\|_\infty + \left( \lim_{\beta \rightarrow 1} \|J_\beta^*(c, \mathcal{T})\|_{Lip} \right) d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).$$

*Proof.* By Theorem 4, for any  $\beta \in [\alpha, 1)$ , we have

$$\|(1 - \beta)(J_\beta^*(c, \mathcal{T}) - J_\beta^*(\hat{c}, \mathcal{S}))\|_\infty \leq \|c - \hat{c}\|_\infty + \beta \|J_\beta^*(c, \mathcal{T})\|_{Lip} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).$$

Consider the term  $(1 - \beta_n(x))J_{\beta_n(x)}^*(c, \mathcal{T})(x)$ . Since  $J_\beta^*(c, \mathcal{T})$  is Lipschitz  $\forall \beta \in [\alpha, 1)$ , we can pick an arbitrary  $z \in \mathbb{X}$  to satisfy Assumption 2, and thus by Lemma 1, we have that  $g_{c, \mathcal{T}}^*$  is a limit point of  $(1 - \beta_n(x))J_{\beta_n(x)}^*(c, \mathcal{T})(x)$  for any  $x \in \mathbb{X}$ . This implies that for each  $x \in \mathbb{X}$ , we can pick a subsequence  $\beta_{n_k}(x) \rightarrow 1$  such that

$$\begin{aligned} \lim_{k \rightarrow \infty} (1 - \beta_{n_k}(x))J_{\beta_{n_k}(x)}^*(c, \mathcal{T})(x) &\rightarrow g_{c, \mathcal{T}}^*, \quad \forall x \in \mathbb{X}, \\ \lim_{k \rightarrow \infty} (1 - \beta_{n_k}(x))J_{\beta_{n_k}(x)}^*(\hat{c}, \mathcal{S})(x) &\rightarrow g_{\hat{c}, \mathcal{S}}^*, \quad \forall x \in \mathbb{X}. \end{aligned}$$

Fixing this subsequence and by Theorem 4, we have

$$\begin{aligned} \sup_{x \in \mathbb{X}} \left| (1 - \beta_{n_k}(x)) \left( J_{\beta_{n_k}(x)}^*(c, \mathcal{T})(x) - J_{\beta_{n_k}(x)}^*(\hat{c}, \mathcal{S})(x) \right) \right| \\ \leq \|c - \hat{c}\|_\infty + \beta_{n_k}(x) \left\| J_{\beta_{n_k}(x)}^*(c, \mathcal{T}) \right\|_{Lip} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}). \end{aligned}$$

Now for each  $x \in \mathbb{X}$ , by continuity of the absolute value function, it is clear that

$$\begin{aligned} \lim_{k \rightarrow \infty} \left| (1 - \beta_{n_k}(x)) \left( J_{\beta_{n_k}(x)}^*(c, \mathcal{T})(x) - J_{\beta_{n_k}(x)}^*(\hat{c}, \mathcal{S})(x) \right) \right| \\ \leq \|c - \hat{c}\|_\infty + \left( \lim_{\beta \rightarrow 1} \|J_\beta^*(c, \mathcal{T})\|_{Lip} \right) d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}). \end{aligned}$$

Finally, by the definition of the subsequence, we have

$$|g_{c, \mathcal{T}}^* - g_{\hat{c}, \mathcal{S}}^*| \leq \|c - \hat{c}\|_\infty + \left( \lim_{\beta \rightarrow 1} \|J_\beta^*(c, \mathcal{T})\|_{Lip} \right) d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).$$

□

Regarding the existence of sequence  $\beta_n(x) \rightarrow 1$  such that  $(1 - \beta_n(x))J_{\beta_n(x)}^*(\hat{c}, \mathcal{S})(x) \rightarrow g_{\hat{c}, \mathcal{S}}^*$ , a sufficient condition is that  $J_\beta(\hat{c}, \mathcal{S})$  is also Lipschitz continuous for all  $\beta$  sufficiently close to 1. We discuss a particular application in which this assumption holds in Section 3.1. However, we do not impose such an assumption in the statement, as we hope to provide more relaxed conditions. We will save further discussions for future work.

**2.3. Robustness Performance Loss** We now use the continuity results discussed in Section 2.2 to establish our main results: an upper bound on the performance loss incurred by applying a policy that is optimal with respect to an *approximate* MDP. This bound is expressed in terms of the distance between cost functions and the uniform Wasserstein-1 distance between transition kernels.

### 2.3.1. Discounted Cost Criterion

**THEOREM 7.** *Suppose that Assumption 1 holds for two MDPs, (reference)  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and (approximation)  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$ . If  $J_\beta^*(c, \mathcal{T})$  is Lipschitz continuous, then*

$$\begin{aligned} \|J_\beta(c, \mathcal{T}, \gamma_{\hat{c}, \mathcal{S}}^*) - J_\beta^*(c, \mathcal{T})\|_\infty &\leq \frac{2 + (d_{TV}(\mathcal{T}, \mathcal{S}) - 2)\beta}{(1 - \beta)^2} \|c - \hat{c}\|_\infty \\ &\quad + \frac{2\beta - (2 - d_{TV}(\mathcal{T}, \mathcal{S}))\beta^2}{(1 - \beta)^2} \|J_\beta^*(c, \mathcal{T})\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}). \end{aligned}$$

*Proof.* For ease of notation, we adopt the abbreviations in (1), and let  $v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x) := J_\beta(c, \mathcal{T}, \gamma_{\hat{c}, \mathcal{S}}^*)(x)$  for  $x \in \mathbb{X}$ . By the triangle inequality and Theorem 4, we have:

$$\begin{aligned} \|v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - v_{c, \mathcal{T}}\|_\infty &= \|v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - v_{\hat{c}, \mathcal{S}} + v_{\hat{c}, \mathcal{S}} - v_{c, \mathcal{T}}\|_\infty \leq \|v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - v_{\hat{c}, \mathcal{S}}\|_\infty + \|v_{\hat{c}, \mathcal{S}} - v_{c, \mathcal{T}}\|_\infty \\ &\leq \|v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - v_{\hat{c}, \mathcal{S}}\|_\infty + \frac{1}{1 - \beta} \|c - \hat{c}\|_\infty + \frac{\beta}{1 - \beta} \|J_\beta^*(c, \mathcal{T})\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}). \end{aligned}$$

Thus, we only need to bound  $\|v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - v_{\hat{c}, \mathcal{S}}\|_\infty$ . By the discounted cost Bellman consistency equation, we have:

$$\begin{aligned} v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x) &= c(x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) + \beta \int v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x') \mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)), \text{ for } x \in \mathbb{X}, \\ v_{\hat{c}, \mathcal{S}}(x) &= \hat{c}(x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) + \beta \int v_{\hat{c}, \mathcal{S}}(x') \mathcal{S}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)), \text{ for } x \in \mathbb{X}. \end{aligned}$$

Therefore, for all  $x \in \mathbb{X}$ , we have

$$\begin{aligned} |v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x) - v_{\hat{c}, \mathcal{S}}(x)| &\leq |c(x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) - \hat{c}(x, \gamma_{\hat{c}, \mathcal{S}}^*(x))| + \beta \left| \int (v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x') - v_{\hat{c}, \mathcal{S}}(x')) \mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) \right. \\ &\quad \left. + \int v_{\hat{c}, \mathcal{S}}(x') (\mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) - \mathcal{S}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x))) \right| \\ &\leq \|c - \hat{c}\|_\infty + \beta \left| \int (v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x') - v_{\hat{c}, \mathcal{S}}(x')) \mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) \right| \\ &\quad + \beta \left| \int v_{\hat{c}, \mathcal{S}}(x') (\mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) - \mathcal{S}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x))) \right| \\ &\leq \beta \left| \int (v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x') - v_{\hat{c}, \mathcal{S}}(x')) \mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) \right| \\ &\quad + \beta \left| \int v_{c, \mathcal{T}}(x') (\mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) - \mathcal{S}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x))) \right| \\ &\quad + \beta \left| \int (v_{\hat{c}, \mathcal{S}}(x') - v_{c, \mathcal{T}}(x')) (\mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) - \mathcal{S}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x))) \right| + \|c - \hat{c}\|_\infty \\ &\leq \|c - \hat{c}\|_\infty + \beta \sup_{x \in \mathbb{X}} |v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x) - v_{\hat{c}, \mathcal{S}}(x)| + \beta \|v_{c, \mathcal{T}}\|_{\text{Lip}} \sup_{(x, u) \in \mathbb{X} \times \mathbb{U}} (\mathcal{W}_1(\mathcal{T}(\cdot|x, u), \mathcal{S}(\cdot|x, u))) \\ &\quad + \beta \|v_{\hat{c}, \mathcal{S}} - v_{c, \mathcal{T}}\|_\infty \sup_{\|f\|_\infty \leq 1} \left| \int f(x') (\mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) - \mathcal{S}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x))) \right| \\ &\leq \|c - \hat{c}\|_\infty + \beta \|v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - v_{\hat{c}, \mathcal{S}}\|_\infty + \beta \|v_{c, \mathcal{T}}\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) + \beta d_{TV}(\mathcal{T}, \mathcal{S}) \|v_{\hat{c}, \mathcal{S}} - v_{c, \mathcal{T}}\|_\infty \\ &\leq \beta \|v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - v_{\hat{c}, \mathcal{S}}\|_\infty + \frac{1 + (d_{TV}(\mathcal{T}, \mathcal{S}) - 1)\beta}{1 - \beta} \|c - \hat{c}\|_\infty + \frac{\beta + (d_{TV}(\mathcal{T}, \mathcal{S}) - 1)\beta^2}{1 - \beta} \|v_{c, \mathcal{T}}\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}). \end{aligned}$$

Then, taking the supremum over  $x \in \mathbb{X}$  on the left-hand side and rearranging terms, we have

$$\|v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - v_{\hat{c}, \mathcal{S}}\|_\infty \leq \frac{1 + (d_{TV}(\mathcal{T}, \mathcal{S}) - 1)\beta}{(1 - \beta)^2} \|c - \hat{c}\|_\infty + \frac{\beta + (d_{TV}(\mathcal{T}, \mathcal{S}) - 1)\beta^2}{(1 - \beta)^2} \|v_{c, \mathcal{T}}\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).$$

Therefore, we have

$$\begin{aligned} \|J_\beta(\mathcal{T}, \gamma_{\hat{c}, \mathcal{S}}^*) - J_\beta^*(c, \mathcal{T})\|_\infty &\leq \frac{2 + (d_{TV}(\mathcal{T}, \mathcal{S}) - 2)\beta}{(1 - \beta)^2} \|c - \hat{c}\|_\infty \\ &\quad + \frac{2\beta - (2 - d_{TV}(\mathcal{T}, \mathcal{S}))\beta^2}{(1 - \beta)^2} \|J_\beta^*(c, \mathcal{T})\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}). \end{aligned}$$

□

Specifically, using the fact that  $d_{TV}(\mathcal{T}, \mathcal{S}) \leq 2$ , we obtain the following:

COROLLARY 1. *Consider the same setup and conditions as in Theorem 7. Then we have*

$$\|J_\beta(c, \mathcal{T}, \gamma_{\hat{c}, \mathcal{S}}^*) - J_\beta^*(c, \mathcal{T})\|_\infty \leq \frac{2}{(1 - \beta)^2} \|c - \hat{c}\|_\infty + \frac{2\beta}{(1 - \beta)^2} \|J_\beta^*(c, \mathcal{T})\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).$$

We note that the above theorem addresses the case that the approximate value function may not be Lipschitz. However, if we multiply the above upper bound by  $1 - \beta$  and let  $\beta \rightarrow 1$ , it would diverge to  $\infty$ . Therefore, we cannot directly extend this result to the average cost criterion using the vanishing discount factor approach. Next, we present a theorem for the case where the approximate model is also Lipschitz, which allows for such an extension.

THEOREM 8. *Suppose that Assumption 1 holds for two MDPs, (reference)  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and (approximation)  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$ . If both  $J_\beta^*(c, \mathcal{T})$  and  $J_\beta^*(\hat{c}, \mathcal{S})$  are Lipschitz continuous, then*

$$\|J_\beta(c, \mathcal{T}, \gamma_{\hat{c}, \mathcal{S}}^*) - J_\beta^*(c, \mathcal{T})\|_\infty \leq \frac{1}{1 - \beta} \|c - \hat{c}\|_\infty + \frac{\beta}{1 - \beta} \left( \|J_\beta^*(c, \mathcal{T})\|_{\text{Lip}} + \|J_\beta^*(\hat{c}, \mathcal{S})\|_{\text{Lip}} \right) d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).$$

*Proof.* Recall the notations in the proof of Theorem 7. By a similar argument, after applying the triangle inequality, we only need to bound  $\|v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - v_{\hat{c}, \mathcal{S}}\|_\infty$ . Again due to the following fixed point equations

$$\begin{aligned} v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x) &= c(x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) + \beta \int v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x') \mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)), \text{ for } x \in \mathbb{X}, \\ v_{\hat{c}, \mathcal{S}}(x) &= c(x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) + \beta \int v_{\hat{c}, \mathcal{S}}(x') \mathcal{S}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)), \text{ for } x \in \mathbb{X}, \end{aligned}$$

we obtain that for any  $x \in \mathbb{X}$ ,

$$\begin{aligned} |v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x) - v_{\hat{c}, \mathcal{S}}(x)| &\leq \beta \left| \int (v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x') - v_{\hat{c}, \mathcal{S}}(x')) \mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) \right| \\ &\quad + \beta \left| \int v_{\hat{c}, \mathcal{S}}(x') (\mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) - \mathcal{S}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x))) \right| \\ &\leq \beta \|v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - v_{\hat{c}, \mathcal{S}}\|_\infty + \beta \left| \int v_{\hat{c}, \mathcal{S}}(x') (\mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) - \mathcal{S}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x))) \right| \\ &\leq \beta \|v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - v_{\hat{c}, \mathcal{S}}\|_\infty + \beta \|J_\beta^*(\hat{c}, \mathcal{S})\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}), \end{aligned}$$

which implies

$$\|v_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - v_{\hat{c}, \mathcal{S}}\|_\infty \leq \frac{\beta}{1 - \beta} \|J_\beta^*(\hat{c}, \mathcal{S})\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}). \quad (2)$$

Finally, together with Theorem 4, we have

$$\|J_\beta(c, \mathcal{T}, \gamma_{\hat{c}, \mathcal{S}}^*) - J_\beta^*(c, \mathcal{T})\|_\infty \leq \frac{1}{1 - \beta} \|c - \hat{c}\|_\infty + \frac{\beta}{1 - \beta} \left( \|J_\beta^*(c, \mathcal{T})\|_{\text{Lip}} + \|J_\beta^*(\hat{c}, \mathcal{S})\|_{\text{Lip}} \right) d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).$$

□

REMARK 6. By Theorem 4, if we assume the Lipschitz continuity of  $J_\beta^*(\hat{c}, \mathcal{S})$ , then

$$\|J_\beta^*(c, \mathcal{T}) - J_\beta^*(\hat{c}, \mathcal{S})\|_\infty \leq \frac{1}{1-\beta} \|c - \hat{c}\|_\infty + \frac{\beta}{1-\beta} \|J_\beta^*(\hat{c}, \mathcal{S})\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).$$

Furthermore, by the intermediate result (2) in the above proof, assuming only the Lipschitz continuity of  $J_\beta^*(\hat{c}, \mathcal{S})$ , we have:

$$\|J_\beta^*(c, \mathcal{T}, \gamma_S^*) - J_\beta^*(c, \mathcal{T})\|_\infty \leq \frac{1}{1-\beta} \|c - \hat{c}\|_\infty + \frac{2\beta}{1-\beta} \|J_\beta^*(\hat{c}, \mathcal{S})\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).$$

This recovers part (2) of Corollary 1 in [6].

**2.3.2. Average Cost Criterion** As discussed earlier, due to the  $(1-\beta)^2$  term in the denominators of the upper bounds in Theorem 7, we cannot directly extend Theorem 7 to the average cost criterion. In the following, we first provide an extension of Theorem 7 under the average cost criterion via the minorization approach, and then extend Theorem 8 using the vanishing discount factor approach.

**THEOREM 9.** *Suppose that Assumption 1 holds for two MDPs, (reference)  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and (approximation)  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$ , both of which satisfy Assumption 3 with a scaling constant  $\epsilon > 0$  and minorizing probability measures  $\rho$  and  $\tau$  respectively. We consider the pairs  $(g_{c, \mathcal{T}}^*, h_{c, \mathcal{T}}^*)$  satisfying*

$$h_{c, \mathcal{T}}^*(x) = \inf_{u \in \mathbb{U}} \left( c(x, u) + \int_{\mathbb{X}} h_{c, \mathcal{T}}^*(x') (\mathcal{T}(dx'|x, u) - \epsilon \rho(dx')) \right) \text{ for } x \in \mathbb{X}, \quad g_{c, \mathcal{T}}^* = \epsilon \int_{\mathbb{X}} h_{c, \mathcal{T}}^*(x') \rho(dx'),$$

and assume the existence of a deterministic stationary policy  $\gamma_{\hat{c}, \mathcal{S}}^*$  such that  $(g_{\hat{c}, \mathcal{S}}^*, h_{\hat{c}, \mathcal{S}}^*, \gamma_{\hat{c}, \mathcal{S}}^*)$  satisfies

$$\begin{aligned} h_{\hat{c}, \mathcal{S}}^*(x) &= \inf_{u \in \mathbb{U}} \left( \hat{c}(x, u) + \int_{\mathbb{X}} h_{\hat{c}, \mathcal{S}}^*(x') (\mathcal{S}(dx'|x, u) - \epsilon \tau(dx')) \right), \\ &= \hat{c}(x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) + \int_{\mathbb{X}} h_{\hat{c}, \mathcal{S}}^*(x') (\mathcal{S}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) - \epsilon \tau(dx')), \text{ for } x \in \mathbb{X}, \\ g_{\hat{c}, \mathcal{S}}^* &= \epsilon \int_{\mathbb{X}} h_{\hat{c}, \mathcal{S}}^*(x') \tau(dx'), \end{aligned}$$

and  $(g_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}, h_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}})$  satisfying

$$h_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x) = c(x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) + \int_{\mathbb{X}} h_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x') (\mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) - \epsilon \rho(dx')) \text{ for } x \in \mathbb{X}, \quad g_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} = \epsilon \int_{\mathbb{X}} h_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x') \rho(dx').$$

If  $h_{c, \mathcal{T}}^*$  is Lipschitz continuous, we have

$$\begin{aligned} |g_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - g_{c, \mathcal{T}}^*| &\leq \left( 1 + \frac{1}{\epsilon} d_{TV}(\mathcal{T}, \mathcal{S}) + \|\rho - \tau\|_{TV} \right) \left( \|c - \hat{c}\|_\infty + \|h_{c, \mathcal{T}}^*\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) \right) \\ &\quad + \left( d_{TV}(\mathcal{T}, \mathcal{S}) + \frac{1}{\epsilon} \|\rho - \tau\|_{TV} + \epsilon \|h_{c, \mathcal{T}}^*\|_{\text{Lip}} \right) \mathcal{W}_1(\rho, \tau). \end{aligned}$$

*Proof.* By the triangle inequality and Theorem 5, we have:

$$\begin{aligned} |g_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - g_{c, \mathcal{T}}^*| &= |g_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - g_{\hat{c}, \mathcal{S}}^* + g_{\hat{c}, \mathcal{S}}^* - g_{c, \mathcal{T}}^*| \leq |g_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - g_{\hat{c}, \mathcal{S}}^*| + |g_{\hat{c}, \mathcal{S}}^* - g_{c, \mathcal{T}}^*| \\ &\leq |g_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} - g_{\hat{c}, \mathcal{S}}^*| + \|c - \hat{c}\|_\infty + \|h_{c, \mathcal{T}}^*\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}). \end{aligned}$$

Applying the two fixed point equations, we have that for  $x \in \mathbb{X}$ ,

$$|h_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x) - h_{\hat{c}, \mathcal{S}}^*(x)| \leq |c(x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) - \hat{c}(x, \gamma_{\hat{c}, \mathcal{S}}^*(x))| + \left| \int_{\mathbb{X}} h_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}(x') (\mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) - \epsilon \rho(dx')) \right.$$

$$\begin{aligned}
 & - \int h_{\hat{c},\mathcal{S}}^*(x') (\mathcal{S}(dx'|x, \gamma_{\mathcal{S}}^*(x)) - \epsilon\tau(dx')) \Big| \\
 \leq & \|c - \hat{c}\|_{\infty} + \left| \int (h_{c,\mathcal{T}}^{\hat{c},\mathcal{S}}(x') - h_{\hat{c},\mathcal{S}}^*(x')) (\mathcal{T}(dx'|x, \gamma_{\mathcal{S}}^*(x)) - \epsilon\rho(dx')) \right. \\
 & \left. + \int h_{\hat{c},\mathcal{S}}^*(x') (\mathcal{T}(dx'|x, \gamma_{\mathcal{S}}^*(x)) - \mathcal{S}(dx'|x, \gamma_{\mathcal{S}}^*(x)) + \epsilon\tau(dx') - \epsilon\rho(dx')) \right| \\
 \leq & \|c - \hat{c}\|_{\infty} + (1 - \epsilon) \|h_{c,\mathcal{T}}^{\hat{c},\mathcal{S}} - h_{\hat{c},\mathcal{S}}^*\|_{\infty} \\
 & + \left| \int h_{c,\mathcal{T}}^*(x') (\mathcal{T}(dx'|x, \gamma_{\mathcal{S}}^*(x)) - \mathcal{S}(dx'|x, \gamma_{\mathcal{S}}^*(x)) + \epsilon\tau(dx') - \epsilon\rho(dx')) \right| \\
 & + \left| \int (h_{\hat{c},\mathcal{S}}^*(x') - h_{c,\mathcal{T}}^*(x')) (\mathcal{T}(dx'|x, \gamma_{\mathcal{S}}^*(x)) - \mathcal{S}(dx'|x, \gamma_{\mathcal{S}}^*(x))) \right| \\
 & + \epsilon \left| \int (h_{\hat{c},\mathcal{S}}^*(x') - h_{c,\mathcal{T}}^*(x')) (\tau(dx') - \rho(dx')) \right| \\
 \leq & \|c - \hat{c}\|_{\infty} + (1 - \epsilon) \|h_{c,\mathcal{T}}^{\hat{c},\mathcal{S}} - h_{\hat{c},\mathcal{S}}^*\|_{\infty} + \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} (d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) + \epsilon\mathcal{W}_1(\rho, \tau)) \\
 & + \|h_{c,\mathcal{T}}^* - h_{\hat{c},\mathcal{S}}^*\|_{\infty} (d_{TV}(\mathcal{T}, \mathcal{S}) + \epsilon\|\rho - \tau\|_{TV})
 \end{aligned}$$

After rearranging terms, we obtain

$$\|h_{c,\mathcal{T}}^{\hat{c},\mathcal{S}} - h_{\hat{c},\mathcal{S}}^*\|_{\infty} \leq \frac{1}{\epsilon} \left( \|c - \hat{c}\|_{\infty} + \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} (d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) + \epsilon\mathcal{W}_1(\rho, \tau)) \right. \\
 \left. + \|h_{c,\mathcal{T}}^* - h_{\hat{c},\mathcal{S}}^*\|_{\infty} (d_{TV}(\mathcal{T}, \mathcal{S}) + \epsilon\|\rho - \tau\|_{TV}) \right).$$

Finally, we have

$$\begin{aligned}
 |g_{c,\mathcal{T}}^{\hat{c},\mathcal{S}} - g_{c,\mathcal{T}}^*| & \leq \epsilon \|h_{c,\mathcal{T}}^{\hat{c},\mathcal{S}} - h_{\hat{c},\mathcal{S}}^*\|_{\infty} \\
 & \leq \|c - \hat{c}\|_{\infty} + \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} (d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) + \epsilon\mathcal{W}_1(\rho, \tau)) + \|h_{c,\mathcal{T}}^* - h_{\hat{c},\mathcal{S}}^*\|_{\infty} (d_{TV}(\mathcal{T}, \mathcal{S}) + \epsilon\|\rho - \tau\|_{TV}) \\
 & \leq \left( 1 + \frac{1}{\epsilon} d_{TV}(\mathcal{T}, \mathcal{S}) + \|\rho - \tau\|_{TV} \right) \left( \|c - \hat{c}\|_{\infty} + \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) \right) \\
 & \quad + \left( d_{TV}(\mathcal{T}, \mathcal{S}) + \frac{1}{\epsilon} \|\rho - \tau\|_{TV} + \epsilon \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} \right) \mathcal{W}_1(\rho, \tau),
 \end{aligned}$$

where at the second step we apply the inequality in Theorem 5.  $\square$

Using the fact that  $d_{TV}(\mathcal{T}, \mathcal{S}) \leq 2$ , we obtain the following corollary:

**COROLLARY 2.** *Consider the same setup and conditions as in Theorem 9. Then we have*

$$|g_{c,\mathcal{T}}^{\hat{c},\mathcal{S}} - g_{c,\mathcal{T}}^*| \leq \left( 3 + \frac{2}{\epsilon} \right) \left( \|c - \hat{c}\|_{\infty} + \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) \right) + \left( 2 + \frac{2}{\epsilon} + \epsilon \|h_{c,\mathcal{T}}^*\|_{\text{Lip}} \right) \mathcal{W}_1(\rho, \tau).$$

Next, we extend Theorem 8 to the average cost criterion using the vanishing discount factor approach.

**THEOREM 10.** *Suppose that Assumption 1 holds for two MDPs, (reference)  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and (approximation)  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$ . Suppose the ACOE for  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$  is satisfied with a deterministic stationary policy  $\gamma_{\hat{c},\mathcal{S}}^*$ . If there exists a constant  $\alpha \in (0, 1)$  such that  $\forall \beta \in [\alpha, 1)$ , both  $\|J_{\beta}^*(c, \mathcal{T})\|_{\text{Lip}} < \infty$ ,  $\lim_{\beta \rightarrow 1} \|J_{\beta}^*(c, \mathcal{T})\|_{\text{Lip}}$  and  $\lim_{\beta \rightarrow 1} \|J_{\beta}^*(\hat{c}, \mathcal{S})\|_{\text{Lip}}$  exist, and for any  $x \in \mathbb{X}$  there exists a sequence  $\beta_n(x) \rightarrow 1$  such that  $(1 - \beta_n(x))J_{\beta_n(x)}^*(c, \mathcal{T}, \gamma_{\hat{c},\mathcal{S}}^*)(x) \rightarrow g_{c,\mathcal{T}}^{\hat{c},\mathcal{S}}$ , then*

$$\begin{aligned}
 & \|J_{\infty}^*(c, \mathcal{T}) - J_{\infty}^*(c, \mathcal{T}, \gamma_{\hat{c},\mathcal{S}}^*)\|_{\infty} \\
 & \leq \|c - \hat{c}\|_{\infty} + \left( \lim_{\beta \rightarrow 1} \|J_{\beta}^*(c, \mathcal{T})\|_{\text{Lip}} + \lim_{\beta \rightarrow 1} \|J_{\beta}^*(\hat{c}, \mathcal{S})\|_{\text{Lip}} \right) d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).
 \end{aligned}$$

*Proof.* The proof proceeds in a similar manner as that of Theorem 6, with the modification that for the discounted cost result, we use Theorem 8 instead of Theorem 4.  $\square$

We note that, in general, no assumption necessarily ensures that  $J_\beta(c, \mathcal{T}, \gamma_{\hat{c}, \mathcal{S}}^*)$  is Lipschitz. Consequently, it is not immediately clear under what conditions there exists, for any  $x \in \mathbb{X}$ , a sequence  $\beta_n(x) \rightarrow 1$  such that  $(1 - \beta_n(x))J_{\beta_n(x)}(c, \mathcal{T}, \gamma_{\hat{c}, \mathcal{S}}^*)(x) \rightarrow g_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}$ . We now present an assumption concerning the ergodicity of approximate kernel  $\mathcal{S}$ , under which this result holds.

**ASSUMPTION 4.** *For the approximate model  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$ , assume that for any deterministic stationary policy  $\gamma$ , the induced Markov chain with transition kernel  $\mathcal{S}(\cdot|x, \gamma(x))$  is positive Harris recurrent and admits at least one invariant probability measure.*

**LEMMA 2 (Ergodicity for Positive Harris Recurrent Chains, Theorem 17.1.7 of [40]).** *For a Markov chain  $\{X_n : n \geq 0\}$ , which admits at least one invariant probability measure, the following statements are equivalent: a) The chain is positive Harris recurrent. b) There exists an invariant probability measure  $\pi$  such that for all  $f \in \mathcal{L}_1(\pi)$  and every initial distribution  $\mu$ ,*

$$\mathbb{P} \left( \lim_{n \rightarrow \infty} \frac{1}{n+1} \sum_{i=0}^n f(X_i) = \int_{\mathbb{X}} f(x) \pi(dx) \right) = 1.$$

**LEMMA 3 (Abelian Inequality, Lemma 5.3.1 of [26]).** *Let  $\{c_t\}_{t \in \mathbb{N}}$  be a sequence of non-negative numbers and  $\beta \in (0, 1)$ . Then*

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{t=0}^{n-1} c_t \leq \liminf_{\beta \rightarrow 1} (1 - \beta) \sum_{t=0}^{\infty} \beta^t c_t \leq \limsup_{\beta \rightarrow 1} (1 - \beta) \sum_{t=0}^{\infty} \beta^t c_t \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{t=0}^{n-1} c_t.$$

**COROLLARY 3.** *Suppose that Assumption 1 holds for two MDPs, (reference)  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and (approximation)  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$ , and that the approximation satisfies Assumption 4. Assume further that the ACOE for  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$  is satisfied with a deterministic stationary policy  $\gamma_{\hat{c}, \mathcal{S}}^*$ . If there exists a constant  $\alpha \in (0, 1)$  such that  $\forall \beta \in [\alpha, 1)$ ,  $\|J_\beta^*(c, \mathcal{T})\|_{Lip} < \infty$ ,  $\lim_{\beta \rightarrow 1} \|J_\beta^*(c, \mathcal{T})\|_{Lip}$  exists,  $\lim_{\beta \rightarrow 1} \|J_\beta^*(\hat{c}, \mathcal{S})\|_{Lip}$  exists, then*

$$\|J_\infty^*(c, \mathcal{T}) - J_\infty^*(c, \mathcal{T}, \gamma_{\hat{c}, \mathcal{S}}^*)\|_\infty \leq \|c - \hat{c}\|_\infty + \left( \lim_{\beta \rightarrow 1} \|J_\beta^*(c, \mathcal{T})\|_{Lip} + \lim_{\beta \rightarrow 1} \|J_\beta^*(\hat{c}, \mathcal{S})\|_{Lip} \right) d_{W_1}(\mathcal{T}, \mathcal{S}).$$

*Proof.* Due to Theorem 10, it suffices to show that for any  $x \in \mathbb{X}$ :  $\lim_{\beta \rightarrow 1} (1 - \beta)J_\beta(c, \mathcal{T}, \gamma_{\hat{c}, \mathcal{S}}^*)(x) = g_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}}$ . As the induced kernel  $\mathcal{T}^{\gamma_{\hat{c}, \mathcal{S}}^*}$  satisfies Lemma 2 (a), for any  $x \in \mathbb{X}$ :

$$\mathbb{P} \left( \lim_{n \rightarrow \infty} \frac{1}{n+1} \sum_{i=0}^n c(X_i, \gamma_{\hat{c}, \mathcal{S}}^*(X_i)) = \int_{\mathbb{X}} c(y, \gamma_{\hat{c}, \mathcal{S}}^*(y)) \pi(dy) \right) = 1,$$

where  $X_0 = x$  a.s. and  $\{X_i, i \geq 0\}$  is a Markov chain with kernel  $\mathcal{T}^{\gamma_{\hat{c}, \mathcal{S}}^*}$ . Furthermore, since  $c$  is assumed bounded, the family of time averages (i.e., the left-hand side) are uniformly bounded by  $\|c\|_\infty$ , and thus satisfy uniform integrability. Therefore, we also have

$$\lim_{n \rightarrow \infty} \frac{1}{n+1} \mathbb{E} \left[ \sum_{i=0}^n c(X_i, \gamma_{\hat{c}, \mathcal{S}}^*(X_i)) \right] = \int_{\mathbb{X}} c(y, \gamma_{\hat{c}, \mathcal{S}}^*(y)) \pi(dy).$$

Then, by the definition of the average cost criterion,

$$g_{c, \mathcal{T}}^{\hat{c}, \mathcal{S}} = J_\infty(c, \mathcal{T}, \gamma_{\hat{c}, \mathcal{S}}^*)(x) = \lim_{n \rightarrow \infty} \frac{1}{n+1} \mathbb{E} \left[ \sum_{i=0}^n c(X_i, \gamma_{\hat{c}, \mathcal{S}}^*(X_i)) \right] = \int_{\mathbb{X}} c(y, \gamma_{\hat{c}, \mathcal{S}}^*(y)) \pi(dy),$$



where  $\pi$  is an invariant probability measure of  $\mathcal{T}^{\gamma_{\hat{c},\mathcal{S}}^*}$ . Now, by the definition of the discounted cost criterion and Lemma 3, we further have

$$\begin{aligned} \limsup_{\beta \rightarrow 1} (1 - \beta) J_\beta(c, \mathcal{T}, \gamma_{\hat{c},\mathcal{S}}^*)(x) &= \limsup_{\beta \rightarrow 1} (1 - \beta) \sum_{i=0}^{\infty} \beta^i \int_{\mathbb{X}} c(y, \gamma_{\hat{c},\mathcal{S}}^*(y)) \mathcal{T}^{\gamma_{\hat{c},\mathcal{S}}^*,i}(dy|x) \\ &\leq \limsup_{n \rightarrow \infty} \frac{1}{n+1} \sum_{i=0}^n \int_{\mathbb{X}} c(y, \gamma_{\hat{c},\mathcal{S}}^*(y)) \mathcal{T}^{\gamma_{\hat{c},\mathcal{S}}^*,i}(dy|x) = g_{c,\mathcal{T}}^{\hat{c},\mathcal{S}}. \end{aligned}$$

Finally, since the limit of the right-hand side exists, we conclude the proof by Lemma 3.  $\square$

**2.4. Conditions Leading to Lipschitz Regularity** In previous subsections, we often assumed that for the reference MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$ ,  $J_\beta^*(c, \mathcal{T})$  is Lipschitz continuous. In this section, we present a sufficient condition for such Lipschitz continuity to hold. To our knowledge, these results are first introduced in [27], and one of the main results is also presented in [49]. For the case of average cost, see [7].

**ASSUMPTION 5 (Wasserstein Regular MDP).** *Consider an MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and a discount factor  $\beta < 1$ . In addition to Assumption 1, assume the following holds:*

- (a). *For any  $u \in \mathbb{U}$ ,  $c: \mathbb{X} \times \mathbb{U} \rightarrow \mathbb{R}$  is  $\|c\|_{Lip}$ -Lipschitz continuous as a function of  $x$ .*
- (b). *For any  $u \in \mathbb{U}$ ,  $\mathcal{T}: \mathbb{X} \times \mathbb{U} \rightarrow \mathcal{P}(\mathbb{X})$  is  $\|\mathcal{T}\|_{Lip}$ -Lipschitz continuous as function of  $x$ , with respect to Wasserstein-1 distance, i.e.*

$$\mathcal{W}_1(\mathcal{T}(\cdot|x, u), \mathcal{T}(\cdot|y, u)) \leq \|\mathcal{T}\|_{Lip} d_{\mathbb{X}}(x, y), \forall u \in \mathbb{U}, x, y \in \mathbb{X}.$$

- (c).  $\beta \|\mathcal{T}\|_{Lip} < 1$ .

**LEMMA 4 (Theorem 4.37 of [49]).** *For an MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and a discount factor  $\beta < 1$  satisfying Assumption 5,  $v_{c,\mathcal{T}} := J_\beta^*(c, \mathcal{T})$  is  $\frac{\|c\|_{Lip}}{1 - \beta \|\mathcal{T}\|_{Lip}}$ -Lipschitz.*

By combining Lemma 4 with Theorem 4, Theorem 7 and Theorem 8, we immediately have the following corollaries for the  $\beta$ -discounted cost case.

**COROLLARY 4.** *Consider two MDPs, (reference)  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and (approximation)  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$ . Suppose that the reference satisfies Assumption 1 and 5, and that the approximation satisfies Assumption 1. Then*

$$\begin{aligned} \|J_\beta^*(c, \mathcal{T}) - J_\beta^*(\hat{c}, \mathcal{S})\|_\infty &\leq \frac{1}{1 - \beta} \|c - \hat{c}\|_\infty + \frac{\beta \|c\|_{Lip}}{(1 - \beta)(1 - \beta \|\mathcal{T}\|_{Lip})} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}), \\ \|J_\beta(\mathcal{T}, \gamma_{\hat{c},\mathcal{S}}^*) - J_\beta^*(c, \mathcal{T})\|_\infty &\leq \frac{2}{(1 - \beta)^2} \|c - \hat{c}\|_\infty + \frac{2\beta \|c\|_{Lip}}{(1 - \beta)^2(1 - \beta \|\mathcal{T}\|_{Lip})} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}). \end{aligned}$$

**COROLLARY 5.** *Under Assumption 1 and 5, for two MDPs, (reference)  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and (approximation)  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$ , we have*

$$\|J_\beta(\mathcal{T}, \gamma_{\hat{c},\mathcal{S}}^*) - J_\beta^*(c, \mathcal{T})\|_\infty \leq \frac{1}{1 - \beta} \|c - \hat{c}\|_\infty + \frac{\beta}{1 - \beta} \left( \frac{\|c\|_{Lip}}{1 - \beta \|\mathcal{T}\|_{Lip}} + \frac{\|\hat{c}\|_{Lip}}{1 - \beta \|\mathcal{S}\|_{Lip}} \right) d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).$$

We now present results for the average cost criterion case.

**ASSUMPTION 6.** *In addition to Assumption 5, assume  $\|\mathcal{T}\|_{Lip} < 1$ .*

The following is an analog of Lemma 4 for the average cost criterion, and can be derived as a corollary of Lemma 4 and [7, Lemma 2.2].

LEMMA 5. *Consider an MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  satisfying Assumption 6. Let  $g \in \mathbb{R}$  and  $h \in C_b(\mathbb{X})$  satisfy:*

$$g + h(x) = \inf_{u \in \mathbb{U}} \left( c(x, u) + \int_{\mathbb{X}} h(y) \mathcal{T}(dy|x, u) \right) \quad \text{for } x \in \mathbb{X}.$$

Then,  $h$  is  $\frac{\|c\|_{Lip}}{1 - \|\mathcal{T}\|_{Lip}}$ -Lipschitz.

*Proof.* We first note that the existence of  $g \in \mathbb{R}$  and  $h \in C_b(\mathbb{X})$  is guaranteed by [7, Lemma 2.2], since Assumption 6 is stronger than [7, Assumption 9]. Furthermore, the proof of [7, Lemma 2.2] is constructive: for each  $\beta \in (0, 1)$  and some fixed  $x_0 \in \mathbb{X}$ , we can define

$$h_\beta(x) := J_\beta(x) - J_\beta(x_0).$$

For a sequence  $\{\beta(k)\}_{k \in \mathbb{N}}$  such that  $\lim_{k \rightarrow \infty} \beta(k) = 1$ , we define  $h(x) := \lim_{k \rightarrow \infty} h_{\beta(k)}(x)$ . Since Lemma 4 implies  $\{h_\beta(x) : \beta \in (0, 1)\}$  is a family of  $\frac{\|c\|_{Lip}}{1 - \|\mathcal{T}\|_{Lip}}$ -Lipschitz functions,  $\{h_\beta(x) : \beta \in (0, 1)\}$  is equicontinuous. Therefore the limit is well-defined for different sequences  $\{\beta(k)\}_{k \in \mathbb{N}}$ , and furthermore,  $h_{\beta(k)}$  converges to  $h$  uniformly. Since  $h$  is the uniform limit of a sequence of  $\frac{\|c\|_{Lip}}{1 - \|\mathcal{T}\|_{Lip}}$ -

Lipschitz functions, we have  $\|h\|_{Lip} \leq \frac{\|c\|_{Lip}}{1 - \|\mathcal{T}\|_{Lip}}$ .  $\square$

REMARK 7. Note that Lemma 5 is also proven in [34, Theorem 3.5] under the assumption that  $\mathcal{T}$  satisfies the minorization condition. The above proof demonstrates that this assumption is not necessary.

By combining Lemma 5 with Theorem 5, Theorem 6, Theorem 9 and Theorem 10, we immediately have the following corollaries for the average cost case.

COROLLARY 6. *Consider two MDPs, (reference)  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and (approximation)  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$ . Suppose that the reference satisfies Assumption 1 and 6, the approximation satisfies Assumption 1, and both satisfy Assumption 3 with some scaling constant  $\epsilon > 0$  and minorizing probability measures  $\rho$  and  $\tau$  respectively. Denote by the pairs  $(g_{c, \mathcal{T}}^*, h_{c, \mathcal{T}}^*)$  and  $(g_{\hat{c}, \mathcal{S}}^*, h_{\hat{c}, \mathcal{S}}^*)$  satisfying ACOE as in Theorem 2. We have*

$$\begin{aligned} \|h_{c, \mathcal{T}}^* - h_{\hat{c}, \mathcal{S}}^*\|_\infty &\leq \frac{1}{\epsilon} \left( \|c - \hat{c}\|_\infty + \frac{\|c\|_{Lip}}{1 - \|\mathcal{T}\|_{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) \right) + \frac{\|c\|_{Lip}}{1 - \|\mathcal{T}\|_{Lip}} \mathcal{W}_1(\rho, \tau), \\ |g_{c, \mathcal{T}}^* - g_{\hat{c}, \mathcal{S}}^*| &\leq \|c - \hat{c}\|_\infty + \frac{\|c\|_{Lip}}{1 - \|\mathcal{T}\|_{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}). \end{aligned}$$

Furthermore, if there exists  $\gamma_{\hat{c}, \mathcal{S}}^*$  such that  $(g_{\hat{c}, \mathcal{S}}^*, h_{\hat{c}, \mathcal{S}}^*, \gamma_{\hat{c}, \mathcal{S}}^*)$  forms a canonical triplet, and let  $(g_{\hat{c}, \mathcal{T}}^*, h_{\hat{c}, \mathcal{T}}^*)$  satisfies

$$h_{\hat{c}, \mathcal{T}}^*(x) = c(x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) + \int h_{\hat{c}, \mathcal{T}}^*(x') (\mathcal{T}(dx'|x, \gamma_{\hat{c}, \mathcal{S}}^*(x)) - \epsilon \rho(dx')), \quad g_{\hat{c}, \mathcal{T}}^* = \epsilon \int_{\mathbb{X}} h_{\hat{c}, \mathcal{T}}^*(x') \rho(dx'),$$

we have

$$\begin{aligned} |g_{\hat{c}, \mathcal{T}}^* - g_{c, \mathcal{T}}^*| &\leq \left( 1 + \frac{1}{\epsilon} d_{TV}(\mathcal{T}, \mathcal{S}) + \|\rho - \tau\|_{TV} \right) \left( \|c - \hat{c}\|_\infty + \frac{\|c\|_{Lip}}{1 - \|\mathcal{T}\|_{Lip}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}) \right) \\ &\quad + \left( d_{TV}(\mathcal{T}, \mathcal{S}) + \frac{1}{\epsilon} \|\rho - \tau\|_{TV} + \epsilon \frac{\|c\|_{Lip}}{1 - \|\mathcal{T}\|_{Lip}} \right) \mathcal{W}_1(\rho, \tau). \end{aligned}$$

Note that under Assumption 5, for any  $\beta \in (0, 1)$ ,  $J_\beta^*(c, \mathcal{T})$  is  $\frac{\|c\|_{\text{Lip}}}{1-\beta\|\mathcal{T}\|_{\text{Lip}}}$ -Lipschitz continuous, and thus  $\lim_{\beta \rightarrow 1} \|J_\beta^*(c, \mathcal{T})\|_{\text{Lip}} = \frac{\|c\|_{\text{Lip}}}{1-\|\mathcal{T}\|_{\text{Lip}}}$ . This allows us to use the vanishing discount factor results under Wasserstein regular MDPs. Specifically, for scenarios that both the reference and approximation model satisfy Assumption 6, we have the following:

**COROLLARY 7.** *Under Assumption 1 and 6, for two MDPs, (reference)  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$  and (approximation)  $(\mathbb{X}, \mathbb{U}, \mathcal{S}, \hat{c})$ ,*

$$\|J_\infty^*(c, \mathcal{T}) - J_\infty^*(\hat{c}, \mathcal{S})\|_\infty \leq \|c - \hat{c}\|_\infty + \frac{\|c\|_{\text{Lip}}}{1 - \|\mathcal{T}\|_{\text{Lip}}} d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).$$

*If further suppose that there exists deterministic stationary policy  $\gamma_{\hat{c}, \mathcal{S}}^*$  such that ACOE for approximate model is satisfied, then*

$$\|J_\infty^*(c, \mathcal{T}) - J_\infty(c, \mathcal{T}, \gamma_{\hat{c}, \mathcal{S}}^*)\|_\infty \leq \|c - \hat{c}\|_\infty + \left( \frac{\|c\|_{\text{Lip}}}{1 - \|\mathcal{T}\|_{\text{Lip}}} + \frac{\|\hat{c}\|_{\text{Lip}}}{1 - \|\mathcal{S}\|_{\text{Lip}}} \right) d_{\mathcal{W}_1}(\mathcal{T}, \mathcal{S}).$$

We now present a sufficient condition for the Wasserstein continuity of transition kernel.

**LEMMA 6.** *Consider the stochastic dynamical system*

$$X_{t+1} = f(X_t, U_t, W_t) \text{ for } t \geq 0, \text{ where } \{W_t\}_{t=0}^\infty \text{ is an i.i.d process with distribution } \mu \in \mathcal{P}(\mathbb{W}).$$

*For  $x \in \mathbb{X}$ ,  $u \in \mathbb{U}$  and  $A \subset \mathcal{F}_\mathbb{X}$ , define  $\mathcal{T}_\mu(A|x, u) := \mu(f_{x,u}^{-1}(A))$ , where  $f_{x,u}^{-1}(A) := \{w \in \mathbb{W} : f(x, u, w) \in A\}$ . Assume that for any  $u \in \mathbb{U}$ ,  $w \in \mathbb{W}$ ,  $f(\cdot, u, w) : \mathbb{X} \rightarrow \mathbb{X}$  is  $\|f\|_{\text{Lip}(\mathbb{X})}$ -Lipschitz continuous. Then for any  $u \in \mathbb{U}$ , the transition kernel  $\mathcal{T}(\cdot|x, u)$  is  $\|f\|_{\text{Lip}(\mathbb{X})}$ -Lipschitz continuous as a function of  $x$ , with respect to Wasserstein-1 distance, i.e.  $\forall x, y \in \mathbb{X}$ ,*

$$\mathcal{W}_1(\mathcal{T}(\cdot|x, u), \mathcal{T}(\cdot|y, u)) \leq \|f\|_{\text{Lip}(\mathbb{X})} d_\mathbb{X}(x, y).$$

*Proof.* By the dual formulation of  $\mathcal{W}_1$ , we have

$$\begin{aligned} \mathcal{W}_1(\mathcal{T}(\cdot|x, u), \mathcal{T}(\cdot|y, u)) &= \sup_{\|g\|_{\text{Lip}} \leq 1} \left( \int g(x) \mathcal{T}(dx|x, u) - \int g(x) \mathcal{T}(dx|y, u) \right) \\ &= \sup_{\|g\|_{\text{Lip}} \leq 1} \left( \int g(f(x, u, w)) \mu(dw) - \int g(f(y, u, w)) \mu(dw) \right) \\ &= \sup_{\|g\|_{\text{Lip}} \leq 1} \left( \int (g \circ f)(x, u, w) \mu(dw) - \int (g \circ f)(y, u, w) \mu(dw) \right) \\ &\leq \sup_{\|g\|_{\text{Lip}} \leq 1} \left( \int |(g \circ f)(x, u, w) - (g \circ f)(y, u, w)| \mu(dw) \right) \\ &\leq \sup_{\|g\|_{\text{Lip}} \leq 1} \left( \int \|f\|_{\text{Lip}(\mathbb{X})} d_\mathbb{X}(x, y) \mu(dw) \right) = \|f\|_{\text{Lip}(\mathbb{X})} d_\mathbb{X}(x, y). \end{aligned}$$

□

In the same spirit as the above proof, we note the following Lemma that shows Assumption 1, in general, holds for continuous stochastic dynamical systems.

**LEMMA 7.** *Consider the stochastic dynamical system as in Lemma 6. If for each  $w \in \mathbb{W}$ ,  $f$  is continuous as a function on  $\mathbb{X} \times \mathbb{U}$ , then the transition kernel  $\mathcal{T}(\cdot|x, u) := \mu(f_{x,u}^{-1}(\cdot))$  is weakly continuous on  $\mathbb{X} \times \mathbb{U}$ .*

*Proof.* It suffices to show for a sequence  $\{(x_n, u_n)\}_{n \in \mathbb{N}}$  and some  $(x, u) \in \mathbb{X} \times \mathbb{U}$  such that  $d_{\mathbb{X}}(x_n, x) \rightarrow 0$  and  $d_{\mathbb{U}}(u_n, u) \rightarrow 0$ , we have for any  $g \in C_b(\mathbb{X})$ ,  $\int_{\mathbb{X}} g(x) \mathcal{T}(dx|x_n, u_n) \rightarrow \int_{\mathbb{X}} g(x) \mathcal{T}(dx|x, u)$ . By the definition of  $\mathcal{T}$ , it suffices to show

$$\int_{\mathbb{X}} g(f(x_n, u_n, w)) \mu(dw) \rightarrow \int_{\mathbb{X}} g(f(x, u, w)) \mu(dw). \quad (3)$$

By continuity of  $g$  and the assumption that for each  $w \in \mathbb{W}$ ,  $f$  is continuous on  $\mathbb{X} \times \mathbb{U}$ , we have that for each  $w \in \mathbb{W}$ ,  $g(f(x_n, u_n, w)) \rightarrow g(f(x, u, w))$ . That is,  $g(f(x_n, u_n, w))$  converges to  $g(f(x, u, w))$  point-wise as a function on  $\mathbb{W}$ . Then by boundedness of  $g$  and the Dominated Convergence Theorem, we have that (3) holds.  $\square$

The applications of the above two lemmas are detailed in Section 3.1.

### 3. Application to Robustness to Noise Distribution and Empirical Noise Estimation

In this section, we consider the disturbance approximation and the associated robustness properties as discussed in [17, 18, 16, 10]. Specifically, we consider the following stochastic dynamical system:

$$X_{t+1} = f(X_t, U_t, W_t) \text{ for } t \geq 0, \text{ where } \{W_t\}_{t=0}^{\infty} \text{ is i.i.d. with some distribution } \mu. \quad (4)$$

Here,  $\{W_t\}_{t=0}^{\infty}$  is referred to as the disturbance process. Consider a decision maker who has knowledge of  $f$  and the true cost function  $c$ , but has no information about the distribution  $\mu$ . However, we assume  $\mu$  can be estimated, say, from realized samples  $\{w_t\}_{t=0}^n$ . The decision maker computes an optimal policy using an estimated distribution  $\nu$ , which may depend on the observed samples, and we want to upper bound the robustness performance loss by the distance between  $\mu$  and  $\nu$ . We show that this estimation scheme can be viewed as estimating the kernel, and thus the performance loss can be obtained via results from the previous section. We detail this discussion in Section 3.1.

**3.1. Robustness to Noise Distribution Approximations** To reduce the problem to the Wasserstein model approximation, we first note the following relationship between the distance of disturbance distributions and the distance between their corresponding transition kernels.

**LEMMA 8.** *Given two probability measures  $\mu, \nu \in \mathcal{P}(\mathbb{W})$  and a measurable function  $f: \mathbb{X} \times \mathbb{U} \times \mathbb{W} \rightarrow \mathbb{X}$ , we define the transition kernels as follows: for  $x \in \mathbb{X}$ ,  $u \in \mathbb{U}$  and  $A \subset \mathbb{F}_{\mathbb{X}}$ ,*

$$\mathcal{T}_{\mu}(A|x, u) := \mu(f_{x,u}^{-1}(A)), \quad \mathcal{T}_{\nu}(A|x, u) := \nu(f_{x,u}^{-1}(A)), \quad \text{where } f_{x,u}^{-1}(A) := \{w \in \mathbb{W} : f(x, u, w) \in A\}.$$

*Assume that for any  $(x, u) \in \mathbb{X} \times \mathbb{U}$ ,  $f(x, u, \cdot)$  is  $\|f\|_{\text{Lip}(\mathbb{W})}$ -Lipschitz continuous on  $\mathbb{W}$ . Then we have*

$$d_{\mathcal{W}_1}(\mathcal{T}_{\mu}, \mathcal{T}_{\nu}) \leq \|f\|_{\text{Lip}(\mathbb{W})} \mathcal{W}_1(\mu, \nu).$$

*Proof.* By the dual formulation of Wasserstein-1 distance,

$$\begin{aligned} \sup_{x,u} \mathcal{W}_1(\mathcal{T}_{\mu}(\cdot|x, u), \mathcal{T}_{\nu}(\cdot|x, u)) &= \sup_{x,u} \sup_{\|g\|_{\text{Lip}} \leq 1} \left| \int g(y) \mathcal{T}_{\mu}(dy|x, u) - \int g(y) \mathcal{T}_{\nu}(dy|x, u) \right| \\ &= \sup_{x,u} \sup_{\|g\|_{\text{Lip}} \leq 1} \left| \int g(f(x, u, w)) \mu(dw) - \int g(f(x, u, w)) \nu(dw) \right| \\ &\leq \sup_{x,u} \sup_{\|g\|_{\text{Lip}} \leq \|f\|_{\text{Lip}(\mathbb{W})}} \left| \int g(x, u, w) \mu(dw) - \int g(x, u, w) \nu(dw) \right| \\ &= \|f\|_{\text{Lip}(\mathbb{W})} \sup_{x,u} \sup_{\|g(x, u, \cdot)\|_{\text{Lip}} \leq 1} \left| \int g(x, u, w) \mu(dw) - \int g(x, u, w) \nu(dw) \right| \\ &= \|f\|_{\text{Lip}(\mathbb{W})} \sup_{x,u} \mathcal{W}_1(\mu, \nu) = \|f\|_{\text{Lip}(\mathbb{W})} \mathcal{W}_1(\mu, \nu). \end{aligned}$$

$\square$

Note that by Lemma 6, both  $\mathcal{T}_\mu$  and  $\mathcal{T}_\nu$  are  $\|f\|_{\text{Lip}(\mathbb{X})}$ -Lipschitz continuous. Therefore, if the cost  $c$  is Lipschitz continuous, we can conclude by Lemma 4 that for discount factor  $\beta$  such that  $\beta\|f\|_{\text{Lip}(\mathbb{X})} < 1$ , both  $J_\beta^*(c, \mathcal{T}_\mu)$  and  $J_\beta^*(c, \mathcal{T}_\nu)$  are Lipschitz continuous. This allows us to apply Corollary 5 and 7. We formalize the statement below.

**COROLLARY 8.** *Consider two stochastic dynamical systems (reference)  $X_{t+1} = f(X_t, U_t, W_t)$  and (approximation)  $\tilde{X}_{t+1} = f(\tilde{X}_t, \tilde{U}_t, \tilde{W}_t)$ , where  $\{W_t\}_{t=0}^\infty \sim \mu$ ,  $\{\tilde{W}_t\}_{t=0}^\infty \sim \nu$  are both i.i.d processes, and  $f$  is  $\|f\|_{\text{Lip}(\mathbb{X})}$ -Lipschitz continuous on  $\mathbb{X}$ , continuous on  $\mathbb{U}$ , and  $\|f\|_{\text{Lip}(\mathbb{W})}$ -Lipschitz continuous on  $\mathbb{W}$ . Denote their transition kernels by  $\mathcal{T}_\mu$  and  $\mathcal{T}_\nu$  respectively. Let  $\mathbb{U}$  be compact and  $c: \mathbb{X} \times \mathbb{U} \rightarrow \mathbb{R}$  be a common cost function such that  $c$  is  $\|c\|_{\text{Lip}}$ -Lipschitz continuous as a function of  $\mathbb{X}$ , and continuous as a function on  $\mathbb{U}$ . Let  $\beta \in (0, 1)$ , such that  $\beta\|f\|_{\text{Lip}(\mathbb{X})} < 1$ , be given, and denote the deterministic stationary optimal policy for the approximate system under discounted cost by  $\gamma_{\beta, \nu}^*$ . Then we have*

$$\|J_\beta^*(c, \mathcal{T}_\mu) - J_\beta(c, \mathcal{T}_\mu, \gamma_{\beta, \nu}^*)\|_\infty \leq \frac{2\beta\|c\|_{\text{Lip}}\|f\|_{\text{Lip}(\mathbb{W})}}{(1-\beta)(1-\beta\|f\|_{\text{Lip}(\mathbb{X})})} \mathcal{W}_1(\mu, \nu).$$

In addition, suppose that  $\|f\|_{\text{Lip}(\mathbb{X})} < 1$ , and that there exists a deterministic stationary policy  $\gamma_\nu^*$  such that the ACOE equation for the approximate model is satisfied. Then

$$\|J_\infty^*(c, \mathcal{T}_\mu) - J_\infty(c, \mathcal{T}_\mu, \gamma_\nu^*)\|_\infty \leq \frac{2\|c\|_{\text{Lip}}\|f\|_{\text{Lip}(\mathbb{W})}}{1-\|f\|_{\text{Lip}(\mathbb{X})}} \mathcal{W}_1(\mu, \nu).$$

### 3.2. General Sample Complexity Bounds

**3.2.1. Wasserstein Convergence of Empirical Measures** Suppose that we do not know the probability measure  $\mu \in \mathcal{P}(\mathbb{W})$ , but we can observe an i.i.d sequence  $\{W_t\}_{t=0}^\infty$  such that for any  $t \geq 0$ ,  $W_t \sim \mu$ . Then a natural way of estimating  $\mu$  is to use *empirical measures* (see e.g. [13]):

$$\mu_n(\cdot) := \frac{1}{n+1} \sum_{t=0}^n \delta_{W_t}(\cdot), \text{ for } n \geq 0, \quad (5)$$

where for any  $A \in \mathcal{F}_\mathbb{W}$  and  $w \in \mathbb{W}$ ,  $\delta_w(A) := \mathbb{1}\{w \in A\}$ . The rates at which empirical measures converge to the true probability measure in Wasserstein distance are well studied [13, 57, 8]. Here we collect several technical results from [13] that will be used later. The original results are stated for the general Wasserstein- $p$  distance, but we focus on the case  $p = 1$ .

**THEOREM 11 (Mean Convergence Rate of Empirical Measure, Theorem 1 [13]).** *Let  $\mu \in \mathcal{P}(\mathbb{R}^d)$ , and denote the  $q$ -th moment of  $\mu$  by  $M_q(\mu) := \int_{\mathbb{R}^d} |x|^q \mu(dx)$ . If there exists  $q > 1$  such that  $M_q(\mu) < \infty$ , then there exists a positive constant  $C$  depending on  $d$  and  $q$  such that for all  $n \geq 1$ ,*

$$\mathbb{E}(\mathcal{W}_1(\mu, \mu_n)) \leq CM_q^{1/q}(\mu) \begin{cases} n^{-1/2} + n^{-(q-1)/q}, & \text{if } d < 2, q \neq 2, \\ n^{-1/2} \log(1+n) + n^{-(q-1)/q}, & \text{if } d = 2, q \neq 2, \\ n^{-1/d} + n^{-(q-1)/q}, & \text{if } d > 2, q \neq d/(d-1). \end{cases}$$

For the concentration of empirical measure, we state the results under two sets of assumptions.

**THEOREM 12 (Concentration under Exponential Integrability, Theorem 2 [13]).** *Let  $\mu \in \mathcal{P}(\mathbb{R}^d)$ . Suppose there exists  $a > 1$  and  $b > 0$  such that  $\mathcal{E}_{a,b} := \int_{\mathbb{R}^d} e^{b|x|^a} \mu(dx) < \infty$ . There exists positive constants  $C$  and  $c$  that depends on  $d, a, b$ , and  $\mathcal{E}_{a,b}$ , such that for all  $n \geq 1$  and  $\epsilon \in (0, 1)$ ,*

$$\mathbb{P}(\mathcal{W}_1(\mu, \mu_n) > \epsilon) \leq C \begin{cases} \exp\{-c n \epsilon^2\}, & \text{if } d < 2, \\ \exp\{-c n (\epsilon / \log(2 + 1/\epsilon))^2\}, & \text{if } d = 2, \\ \exp\{-c n \epsilon^d\}, & \text{if } d > 2. \end{cases}$$

**THEOREM 13 (Concentration under Moment conditions, Theorem 2 [13]).** *Let  $\mu \in \mathcal{P}(\mathbb{R}^d)$ . Suppose there exists  $q > 2$  such that  $M_q(\mu) < \infty$ . There exists positive constants  $C$  and  $c$  that depends on  $q$  and  $M_q(\mu)$ , such that for all  $n \geq 1$  and  $\epsilon \in (0, 1)$ ,*

$$\mathbb{P}(\mathcal{W}_1(\mu, \mu_n) > \epsilon) \leq Cn(n\epsilon)^{-(q-\epsilon)} + C \begin{cases} \exp\{-cn\epsilon^2\}, & \text{if } d < 2, \\ \exp\{-cn(\epsilon/\log(2+1/\epsilon))^2\}, & \text{if } d = 2, \\ \exp\{-cn\epsilon^d\}, & \text{if } d > 2. \end{cases}$$

On the other hand, empirical measures do not, in general, converge in total variation if the true probability measure admits a density with respect to Lebesgue measure  $\lambda$ .

**3.2.2. Sample Complexity Bounds on Robustness to Empirical Noise Distribution Estimation** Recall the stochastic dynamic system in (4), where the function  $f$  is known but the disturbance distribution  $\mu$  is *unknown*. Denote by  $\mathcal{T}_\mu$  the associated controlled transition kernel.

Assume that we can observe an i.i.d sequence  $\{W_t\}_{t=0}^\infty$  such that for any  $t \geq 0$ ,  $W_t \sim \mu$ , and that we use the empirical measures  $\{\mu_n : n \geq 0\}$  in (5) to estimate  $\mu$ . For  $n \geq 0$ , given the estimate  $\mu_n$ , we consider the following stochastic dynamic system:

$$\tilde{X}_{t+1} = f(\tilde{X}_t, \tilde{U}_t, \tilde{W}_t) \text{ for } t \geq 0, \text{ where } \{\tilde{W}_t\}_{t=0}^\infty \text{ is i.i.d. with distribution } \mu_n,$$

and denote by  $\mathcal{T}_{\mu_n}$  the associated controlled transition kernel. Since the cost function  $c$  is assumed known, we can solve the  $\beta$ -discounted cost and the average cost MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}_{\mu_n}, c)$ , and denote the solutions by  $\gamma_{\beta, \mu_n}^*$  and  $\gamma_{\infty, \mu_n}^*$  respectively.

We now combine Corollary 8 with Theorem 11, 12, and 13 to establish a comprehensive result on the rate estimate of robustness performance loss resulting from empirical disturbance estimation.

**THEOREM 14 (Convergence Rate of Robustness Performance Loss).** *Assume  $f$  in (4) is  $\|f\|_{Lip(\mathbb{X})}$ -Lipschitz continuous on  $\mathbb{X}$ , continuous on  $\mathbb{U}$ , and  $\|f\|_{Lip(\mathbb{W})}$ -Lipschitz continuous on  $\mathbb{W}$ . Let  $\mathbb{U}$  be compact and  $c : \mathbb{X} \times \mathbb{U} \rightarrow \mathbb{R}$  be a known cost function such that  $c$  is  $\|c\|_{Lip}$ -Lipschitz continuous as a function of  $\mathbb{X}$ , and continuous as a function on  $\mathbb{U}$ . Let  $\beta \in (0, 1)$  such that  $\beta\|f\|_{Lip(\mathbb{X})} < 1$  be given.*

**(a).** *If there exists  $q > 1$  such that  $M_q(\mu) < \infty$ , then there exists a positive constant  $C$  depending on  $d, q, \mu, \beta, \|c\|_{Lip}, \|f\|_{Lip(\mathbb{X})}$ , and  $\|f\|_{Lip(\mathbb{W})}$ , such that for all  $n \geq 1$ ,*

$$\mathbb{E} \left[ \left\| J_\beta(c, \mathcal{T}_\mu, \gamma_{\beta, \mu_n}^*) - J_\beta^*(c, \mathcal{T}_\mu) \right\|_\infty \right] \leq C \begin{cases} n^{-1/2} + n^{-(q-1)/q}, & \text{if } d < 2, q \neq 2, \\ n^{-1/2} \log(1+n) + n^{-(q-1)/q}, & \text{if } d = 2, q \neq 2, \\ n^{-1/d} + n^{-(q-1)/q}, & \text{if } d > 2, q \neq d/(d-1). \end{cases}$$

**(b).** *If there exists  $a > 1$  and  $b > 0$  such that  $\mathcal{E}_{a,b} < \infty$ , then there exists a positive constant  $C$  that depends on  $d, a, b$ , and  $\mathcal{E}_{a,b}$ , and another positive constant  $c$ , which in addition depends on  $\beta, \|c\|_{Lip}, \|f\|_{Lip(\mathbb{X})}$ , and  $\|f\|_{Lip(\mathbb{W})}$ , such that for all  $n \geq 1$  and  $\epsilon \in (0, 1)$ ,*

$$\mathbb{P} \left( \left\| J_\beta(c, \mathcal{T}_\mu, \gamma_{\beta, \mu_n}^*) - J_\beta^*(c, \mathcal{T}_\mu) \right\|_\infty > \epsilon \right) \leq C \begin{cases} \exp\{-cn\epsilon^2\}, & \text{if } d < 2, \\ \exp\{-cn(\epsilon/\log(2+1/\epsilon))^2\}, & \text{if } d = 2, \\ \exp\{-cn\epsilon^d\}, & \text{if } d > 2. \end{cases}$$

**(c).** *If there exists  $q > 2$  such that  $M_q(\mu) < \infty$ , then there exists a positive constant  $C$  that depends on  $q$  and  $M_q(\mu)$ , and a positive constant  $c_1$  which in addition depends on  $\beta, \|c\|_{Lip}, \|f\|_{Lip(\mathbb{X})}$ , and  $\|f\|_{Lip(\mathbb{W})}$ , such that for all  $n \geq 1$  and  $\epsilon \in (0, 1)$ ,*

$$\mathbb{P} \left( \left\| J_\beta(c, \mathcal{T}_\mu, \gamma_{\beta, \mu_n}^*) - J_\beta^*(c, \mathcal{T}_\mu) \right\|_\infty > \epsilon \right) \leq$$

$$Cn(nc_2\epsilon)^{-(q-c_2\epsilon)} + C \begin{cases} \exp\{-c_1n\epsilon^2\}, & \text{if } d < 2, \\ \exp\{-c_1n(\epsilon/\log(2+1/\epsilon))^2\}, & \text{if } d = 2, \\ \exp\{-c_1n\epsilon^d\}, & \text{if } d > 2, \end{cases}$$

where  $c_2 := (1 - \beta) \left(1 - \beta \|f\|_{Lip(\mathbb{X})}\right) / \left(2\beta \|c\|_{Lip} \|f\|_{Lip(\mathbb{W})}\right)$ .

Additionally, assume  $\|f\|_{Lip(\mathbb{X})} < 1$ . Then the results in Part (a) and (b) hold under the average cost criterion if we replace  $J_\beta, J_\beta^*$ , and  $\gamma_{\beta,\mu_n}^*$  by  $J_\infty, J_\infty^*$ , and  $\gamma_{\infty,\mu_n}^*$ , respectively; in both cases,  $C$  no longer depends on  $\beta$ . Further, the result in Part (c) holds under the average cost criterion if we replace  $J_\beta, J_\beta^*, \gamma_{\beta,\mu_n}^*$ , and  $c_2$  by  $J_\infty, J_\infty^*, \gamma_{\infty,\mu_n}^*$ , and  $(1 - \|f\|_{Lip(\mathbb{X})}) / (2\|c\|_{Lip} \|f\|_{Lip(\mathbb{W})})$ , respectively; in this case,  $c_1$  no longer depends on  $\beta$ .

**Example 3.1** Let  $\mu$  be the standard Normal distribution on  $\mathbb{R}$ . It is known that all moments of  $\mu$  are finite, and with  $a = 2, b = \frac{1}{4}$ , we also know  $\mathcal{E}_{a,b} = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \exp\left\{-\frac{x^2}{4}\right\} \lambda(dx) < \infty$ , where  $\lambda$  is the Lebesgue measure on  $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ . Thus, both Theorem 11 and 12 apply to  $\mu$  and its empirical measures. By Theorem 14, we have that for any discount factor  $\beta \in (0, 1)$  such that  $\beta \|f\|_{Lip(\mathbb{X})} < 1$ ,

$$\begin{aligned} \mathbb{E} \left[ \left\| J_\beta^*(c, \mathcal{T}_\mu) - J_\beta(c, \mathcal{T}_\mu, \gamma_{\beta,\mu_n}^*) \right\|_\infty \right] &= \mathcal{O}(n^{-1/2}), \\ \mathbb{P} \left( \left\| J_\beta^*(c, \mathcal{T}_\mu) - J_\beta(c, \mathcal{T}_\mu, \gamma_{\beta,\mu_n}^*) \right\|_\infty > \epsilon \right) &= \mathcal{O}(\exp\{-c_n\epsilon^2\}), \end{aligned} \quad (6)$$

where  $c$  is a constant depending only on  $\mu, \beta, \|c\|_{Lip}, \|f\|_{Lip(\mathbb{W})}$ , and  $\|f\|_{Lip(\mathbb{X})}$ .

Assume, in addition,  $\|f\|_{Lip(\mathbb{X})} < 1$ . For the average cost problem, the inequalities in (6) continue to hold if we replace  $J_\beta, J_\beta^*$ , and  $\gamma_{\beta,\mu_n}^*$  by  $J_\infty, J_\infty^*$ , and  $\gamma_{\infty,\mu_n}^*$ , respectively. In this case,  $c$  does not depend on  $\beta$ .

Similarly, if  $\mu$  is the standard multivariate Normal distribution on  $\mathbb{R}^d$  with  $d > 2$ , the rates in (6) become  $\mathcal{O}(n^{-1/d})$  and  $\mathcal{O}(\exp\{-c_n\epsilon^d\})$ , respectively.

**3.3. Case where Model and Noise are Learned Simultaneously** In this section, we again consider the dynamic system in (4), but assume that *both* the function  $f$  and the distribution  $\mu$  are unknown.

We start with a lemma that bounds the Wasserstein-1 distance between two controlled transition kernels corresponding to different models with the same disturbance distribution. For a measurable function  $f: \mathbb{X} \times \mathbb{U} \times \mathbb{W} \rightarrow \mathbb{X}$ ,  $x \in \mathbb{X}, u \in \mathbb{U}$  and  $A \subset \mathcal{F}_{\mathbb{X}}$ , recall that  $f_{x,u}^{-1}(A) := \{w \in \mathbb{W} : f(x, u, w) \in A\}$ .

**LEMMA 9.** Let  $f, g: \mathbb{X} \times \mathbb{U} \times \mathbb{W} \rightarrow \mathbb{X}$  be measurable and  $\mu \in \mathcal{P}(\mathbb{W})$  be a probability measure. Define the controlled transition kernels as follows: for  $x \in \mathbb{X}, u \in \mathbb{U}$  and  $A \subset \mathcal{F}_{\mathbb{X}}$ ,

$$\mathcal{T}_{f,\mu}(A|x, u) := \mu(f_{x,u}^{-1}(A)), \quad \mathcal{T}_{g,\mu}(A|x, u) := \mu(g_{x,u}^{-1}(A)).$$

Then  $d_{\mathcal{W}_1}(\mathcal{T}_{f,\mu}, \mathcal{T}_{g,\mu}) \leq \|f - g\|_\infty$ .

*Proof.* By the dual formulation of the Wasserstein-1 distance,

$$\begin{aligned} \sup_{x,u} \mathcal{W}_1(\mathcal{T}_{f,\mu}(\cdot|x, u), \mathcal{T}_{g,\mu}(\cdot|x, u)) &= \sup_{x,u} \sup_{\|h\|_{Lip} \leq 1} \left| \int h(y) \mathcal{T}_{f,\mu}(dy|x, u) - \int h(y) \mathcal{T}_{g,\mu}(dy|x, u) \right| \\ &= \sup_{x,u} \sup_{\|h\|_{Lip} \leq 1} \left| \int h(f(x, u, w)) \mu(dw) - \int h(g(x, u, w)) \mu(dw) \right| \\ &\leq \sup_{x,u} \sup_{\|h\|_{Lip} \leq 1} \int |h(f(x, u, w)) - h(g(x, u, w))| \mu(dw) \end{aligned}$$

$$\begin{aligned} &\leq \sup_{x,u} \int |f(x,u,w) - g(x,u,w)| \mu(dw) \\ &\leq \sup_{x,u,w} |f(x,u,w) - g(x,u,w)| = \|f - g\|_\infty. \end{aligned}$$

□

A context in which Lemma 9 proves useful is the following additive noise model:

$$X_{t+1} = f(X_t, U_t) + W_t \text{ for } t \geq 0, \text{ where } \{W_t\}_{t=0}^\infty \text{ is i.i.d. with some distribution } \mu,$$

and  $f$  is assumed to be continuous on  $\mathbb{U}$  and  $\|f\|_{\text{Lip}(\mathbb{X})}$ -Lipschitz continuous on  $\mathbb{X}$ . Suppose the decision maker (DM) does not know the model  $f$  or the distribution  $\mu$ , but based on data, the DM estimates  $f$  by some estimator  $\hat{f}_n$ . Given an estimated model  $\hat{f}_n$ , the DM can then estimate the distribution  $\mu$  by the following “noisy” empirical measure

$$\tilde{\mu}_n(\cdot) := \frac{1}{n} \sum_{i=0}^{n-1} \delta_{(X_{i+1} - \hat{f}_n(X_i, U_i))}(\cdot).$$

For each realized sample path with outcome  $\omega \in \Omega$ , with a coupling that associates  $X_{i+1}(\omega) - \hat{f}_n(X_i(\omega), U_i(\omega))$  with  $W_i(\omega)$ , we can show that the noisy empirical measure is still a good estimator for  $\mu$  as long as  $\|f - \hat{f}_n\|_\infty$  is small:

$$\begin{aligned} \mathcal{W}_1(\tilde{\mu}_n(\omega), \mu) &\leq \mathcal{W}_1(\tilde{\mu}_n(\omega), \mu_n(\omega)) + \mathcal{W}_1(\mu_n(\omega), \mu) \\ &\leq \sum_{i=1}^{n-1} \frac{1}{n} \left| X_{i+1}(\omega) - \hat{f}_n(X_i(\omega), U_i(\omega)) - W_i(\omega) \right| + \mathcal{W}_1(\mu_n(\omega), \mu) \\ &= \sum_{i=1}^{n-1} \frac{1}{n} \left| (f(X_i(\omega), U_i(\omega)) + W_i(\omega)) - (\hat{f}_n(X_i(\omega), U_i(\omega)) + W_i(\omega)) \right| + \mathcal{W}_1(\mu_n(\omega), \mu) \\ &\leq \|f - \hat{f}_n\|_\infty + \mathcal{W}_1(\mu_n(\omega), \mu), \end{aligned}$$

where we recall the empirical measure  $\mu_n$  in (5). Under such an estimation scheme, we consider the approximate model:

$$\tilde{X}_{t+1} = \hat{f}_n(\tilde{X}_t, \tilde{U}_t) + \tilde{W}_t \text{ for } t \geq 0, \text{ where } \{\tilde{W}_t\}_{t=0}^\infty \text{ is i.i.d. with some distribution } \tilde{\mu}_n.$$

We assume that the DM knows the true cost function, and denote by  $\gamma_{\hat{f}_n, \tilde{\mu}_n, \beta}^*$  the optimal solution to the associated  $\beta$ -discounted cost MDP. If  $\hat{f}_n$  is continuous on  $\mathbb{X} \times \mathbb{U}$ , by Theorem 7 and Lemma 9, we have

$$\begin{aligned} &\mathbb{P} \left( \left\| J_\beta^*(c, \mathcal{T}_{f, \mu}) - J_\beta(c, \mathcal{T}_{f, \mu}, \gamma_{\hat{f}_n, \tilde{\mu}_n, \beta}^*) \right\|_\infty > \epsilon \right) = \mathcal{O} \left( \mathbb{P} \left( d_{\mathcal{W}_1}(\mathcal{T}_{f, \mu}, \mathcal{T}_{\hat{f}_n, \tilde{\mu}_n}) > \epsilon \right) \right) \\ &= \mathcal{O} \left( \mathbb{P} \left( d_{\mathcal{W}_1}(\mathcal{T}_{f, \mu}, \mathcal{T}_{f, \tilde{\mu}_n}) + d_{\mathcal{W}_1}(\mathcal{T}_{f, \tilde{\mu}_n}, \mathcal{T}_{\hat{f}_n, \tilde{\mu}_n}) > \epsilon \right) \right) = \mathcal{O} \left( \mathbb{P} \left( \mathcal{W}_1(\mu_n, \mu) > \frac{\epsilon}{2} \right) + \mathbb{P} \left( \|f - \hat{f}_n\|_\infty > \frac{\epsilon}{2} \right) \right), \end{aligned}$$

where the involved constants depend on  $\beta$ ,  $\|c\|_{\text{Lip}}$ ,  $\|f\|_{\text{Lip}(\mathbb{X})}$ , and  $\|f\|_{\text{Lip}(\mathbb{W})}$ . Note that the first term in the last expression above can be bounded using the concentration result in either Theorem 12 or Theorem 13. For bounding the uniform estimation error  $\|f - \hat{f}_n\|_\infty$ , we refer readers to [52, 39] for the rich nonparametric estimation literature. We note, however, that for the additive noise model, the classical theory requires i.i.d samples of triplets  $(X_i, U_i, X_{i+1})_{i=0}^{n-1}$ . One cannot obtain the i.i.d samples along a single sample path; instead, the system needs to be restarted independently for  $n$  times. We defer rigorous analysis and results on convergence rates to future work.



**4. Model Approximation, Empirical Model Learning, and Sample Complexity** In this section, we introduce a more general learning framework in which the model itself is learned from data. We begin by reviewing model approximation with finite model representations, noting that some cases fall within the scope of our general robustness results. We then propose learning algorithms to estimate the quantized model from data, establishing empirical consistency and sample complexity bounds. The generality of these results appears to be novel in the literature.

**4.1. Quantized Approximations** In this section, we briefly review the state quantization scheme, which was first introduced, to our knowledge, in [51, Section 3] (see also [31, Section 2.3]). The results will also be useful when we develop two general learning algorithms in the following section.

We demonstrate that the near-optimality of such quantization, as shown for the discounted cost problem in Theorem 6 of [31] (which slightly refines [49, Theorem 4.38]) and for the average cost problem in Theorem 3.5 of [34], can be considered as special cases of Theorem 7 and Theorem 9, respectively. For simplicity, we assume that  $\mathbb{U}$  is a finite set, which does not restrict generality if we consider a compact action space (see Chapter 3 of [49]).

Consider a state space  $\mathbb{X}$  and an  $M$ -partition of it,  $\{B_i\}_{i=1}^M$ . For each partition  $B_i$ , we pick some representative element  $y_i \in B_i$ . A quantizer is then a map  $q: \mathbb{X} \rightarrow \{y_i\}_{i=1}^M$ , where

$$q(x) = y_i \quad \text{if and only if} \quad x \in B_i.$$

Given a weighting measure  $\pi \in \mathcal{P}(\mathbb{X})$ , assuming  $\pi(B_i) > 0$  for  $1 \leq i \leq M$ , we denote its restriction on each quantization bin  $B_i$  by  $\hat{\pi}_i$ :

$$\forall A \subset B_i, \quad \hat{\pi}_i(A) = \pi(A)/\pi(B_i). \quad (7)$$

Then we define a new finite state MDP,  $(\{y_i\}_{i=1}^M, \mathbb{U}, \mathcal{T}^{\pi, M}, c^{\pi, M})$ , where  $c^{\pi, M}$  and  $\mathcal{T}^{\pi, M}$  are defined as follows: for  $1 \leq i, j \leq M$ ,

$$c^{\pi, M}(y_i, u) := \int_{B_i} c(x, u) \hat{\pi}_i(dx), \quad \mathcal{T}^{\pi, M}(y_j | y_i, u) := \int_{B_i} \mathcal{T}(B_j | x, u) \hat{\pi}_i(dx). \quad (8)$$

We denote by  $\gamma_\beta^{\pi, M}$  and  $\gamma_\infty^{\pi, M}$  the optimal policy for the discounted (by  $\beta$ ) and the average cost MDP  $(\{y_i\}_{i=1}^M, \mathbb{U}, \mathcal{T}^{\pi, M}, c^{\pi, M})$ , respectively. Next, we define the piecewise constant extension of such a finite model to the original state space as in [51, Section 3].

**DEFINITION 6 (PIECEWISE CONSTANT EXTENSION I).** For  $1 \leq i \leq M$ ,  $x \in B_i$ ,  $u \in \mathbb{U}$ , and  $A \in \mathcal{F}_{\mathbb{X}}$ , we define

$$\overline{c^{\pi, M}}(x) := c^{\pi, M}(y_i, u), \quad \overline{\mathcal{T}^{\pi, M}}(A | x, u) := \int_{B_i} \mathcal{T}(A | z, u) \hat{\pi}_i(dz),$$

and further

$$\overline{\gamma_\beta^{\pi, M}}(x) := \gamma_\beta^{\pi, M}(y_i), \quad \overline{\gamma_\infty^{\pi, M}}(x) := \gamma_\infty^{\pi, M}(y_i).$$

As discussed in [51, Section 3],  $\overline{\gamma_\beta^{\pi, M}}$  and  $\overline{\gamma_\infty^{\pi, M}}$  are the optimal policies for the discounted (by  $\beta$ ) and the average cost MDP  $(\mathbb{X}, \mathbb{U}, \overline{\mathcal{T}^{\pi, M}}, \overline{c^{\pi, M}})$ , respectively.

We note that the transition kernel under piecewise constant extension is, in general, not weakly continuous, so Assumption 1 does not apply to the approximate model. As a remedy, for the discounted problem, we adopt Assumption 2.1 of [25], which automatically holds for the piecewise constant model under Assumption 1 with a finite action set  $\mathbb{U}$ . For the average cost problem, in addition to Assumption 2.1 of [25], Assumption 3.1(4) of [25] also holds if the original kernel  $\mathcal{T}$  is minorized as in Definition 5. Therefore, we can invoke optimality equations for the piecewise

constant model for both problems, which means that all our results hold when using the piecewise constant model as an approximation. We further define the quantization error:

$$\delta_M := \max_{1 \leq i \leq M} \sup_{x, x' \in B_i} d_{\mathbb{X}}(x, x'). \quad (9)$$

The following approximation error is noted in [34]:

LEMMA 10 (**Lemma 3.3**, [34]). *Under Assumption 5(a) and (b),*

$$\|c - \overline{c^{\pi, M}}\|_{\infty} \leq \|c\|_{Lip} \delta_M, \quad d_{\mathcal{W}_1}(\mathcal{T}, \overline{\mathcal{T}^{\pi, M}}) \leq \|\mathcal{T}\|_{Lip} \delta_M.$$

Therefore instantiating Theorem 7 and 9 in view of Lemma 10, we have the following:

COROLLARY 9 (**Theorem 6**, [31]). *Under Assumption 1 and 5,*

$$\left\| J_{\beta}(c, \mathcal{T}, \overline{\gamma_{\beta}^{\pi, M}}) - J_{\beta}^*(c, \mathcal{T}) \right\|_{\infty} \leq \frac{2\|c\|_{Lip}}{(1-\beta)^2(1-\beta\|\mathcal{T}\|_{Lip})} \delta_M.$$

*Proof.* Combining Corollary 4 and Lemma 10:

$$\begin{aligned} \left\| J_{\beta}(c, \mathcal{T}, \overline{\gamma_{\beta}^{\pi, M}}) - J_{\beta}^*(c, \mathcal{T}) \right\|_{\infty} &\leq \frac{2}{(1-\beta)^2} \|c - \overline{c^{\pi, M}}\|_{\infty} + \frac{2\beta\|c\|_{Lip}}{(1-\beta)^2(1-\beta\|\mathcal{T}\|_{Lip})} d_{\mathcal{W}_1}(\mathcal{T}, \overline{\mathcal{T}^{\pi, M}}) \\ &\leq \frac{2\|c\|_{Lip}}{(1-\beta)^2} \delta_M + \frac{2\beta\|c\|_{Lip}\|\mathcal{T}\|_{Lip}}{(1-\beta)^2(1-\beta\|\mathcal{T}\|_{Lip})} \delta_M = \frac{2\|c\|_{Lip}}{(1-\beta)^2(1-\beta\|\mathcal{T}\|_{Lip})} \delta_M. \end{aligned}$$

□

The same also applies to the average cost problem; for corresponding results, see [34, Theorem 3.5], which can be viewed as a counterpart to Theorem 9 when the approximate model is obtained through finite model approximation.

COROLLARY 10 (**Theorem 3.5**, [34]). *Suppose that  $\mathcal{T}$  is minorized as in Definition 5. Under Assumption 1 and 6,*

$$\left\| J_{\infty}^*(c, \mathcal{T}, \overline{\gamma_{\infty}^{\pi, M}}) - J_{\infty}^*(c, \mathcal{T}) \right\|_{\infty} \leq \left(1 + \frac{2}{\epsilon}\right) \left(\frac{\|c\|_{Lip}}{1 - \|\mathcal{T}\|_{Lip}}\right) \delta_M.$$

*Proof.* By construction, as per Definition 6, if  $\mathcal{T}$  satisfies the minorization condition with a minorizing measure  $\rho$  and scaling constant  $\epsilon$ , then  $\overline{\mathcal{T}^{\pi, M}}$  also satisfies the minorization condition with the same minorizing measure and scaling constant. Consequently, the proof is complete by Corollary 6, where we upper bound the total variation distance between measures by 2. □

**4.2. Simultaneous Finite Model Approximation and Finite Model Learning** In this section, we focus on the case where the quantized model in (8) is unknown and needs to be learned from data. We consider two scenarios regarding data availability: (i) a single trajectory of a controlled Markov process and (ii) independent transitions where a simulation device is available for each initial state and action.

For either case, our goal is to show the robustness of the optimal policies derived from simultaneous finite model approximation and the learning of these approximate finite models obtained via empirical estimates. For the former, we present asymptotic empirical consistency results, whereas for the latter we provide explicit sample complexity bounds that relate performance loss to the number of samples.

As noted earlier, a related approach is presented in [10], which involves a different construction for scenarios where the model is known and satisfies an absolute continuity condition with respect to a reference measure. This construction aims to obtain a finite model by utilizing empirical data collected from the reference measure. Together with the known density, the empirical measure is used to construct a finite model.

---

**Algorithm 1** Sampling a Wasserstein Regular MDP along a Single Trajectory

---

**Require:** Simulation Length  $N$  and ergodic exploration policy  $\gamma$

- 1: **for all**  $n = 1, \dots, N - 1$  **do**
  - 2:     Observe  $(X_n, U_n, C_n, X_{n+1})$  that evolves according to the true kernel  $\mathcal{T}$  and policy  $\gamma$
  - 3: **end for**
  - 4: Set  $\hat{\mathcal{T}}_N(y_j|y_i, u) := \frac{\sum_{n=0}^{N-1} \mathbb{1}\{X_{n+1} \in B_j, X_n \in B_i, U_n = u\}}{\sum_{n=0}^{N-1} \mathbb{1}\{X_n \in B_i, U_n = u\}}$
  - 5: Set  $\hat{c}_N(y_i, u) := \frac{\sum_{n=0}^{N-1} C_n \mathbb{1}\{X_n \in B_i, U_n = u\}}{\sum_{n=0}^{N-1} \mathbb{1}\{X_n \in B_i, U_n = u\}}$
  - 6: **return**  $(\hat{\mathcal{T}}_N, \hat{c}_N)$
- 

**4.2.1. Empirical model learning from a single trajectory** Let  $\gamma \in \mathcal{P}(\mathbb{U})$  be a probability mass function on  $\mathbb{U}$  such that for  $u \in \mathbb{U}$ ,  $\gamma(u) > 0$ .  $\gamma$  may be viewed as a state-independent control policy. We observe a Markov chain  $(X_n, U_n, C_n), 0 \leq n \leq N$ , whose dynamics are determined by the kernel  $\mathcal{T}$  and policy  $\gamma$ , as shown in Algorithm 1. Specifically, for each  $x \in \mathbb{X}$ ,  $u, u' \in \mathbb{U}$ , and  $A \subset \mathbb{X}$ ,

$$\mathbb{P}(X_{n+1} \in A, U_{n+1} = u' | X_n = x, U_n = u) = \mathcal{T}(A|x, u)\gamma_{u'}, \quad C_n = c(X_n, U_n).$$

Define the following estimators for  $1 \leq i, j \leq M$  and  $u \in \mathbb{U}$ ,

$$\hat{\mathcal{T}}_N(y_j|y_i, u) := \frac{\sum_{n=0}^{N-1} \mathbb{1}\{X_{n+1} \in B_j, X_n \in B_i, U_n = u\}}{\sum_{n=0}^{N-1} \mathbb{1}\{X_n \in B_i, U_n = u\}}, \quad \hat{c}_N(y_i, u) := \frac{\sum_{n=0}^{N-1} C_n \mathbb{1}\{X_n \in B_i, U_n = u\}}{\sum_{n=0}^{N-1} \mathbb{1}\{X_n \in B_i, U_n = u\}}.$$

Given the learned MDP  $(\hat{\mathcal{T}}_N, \hat{c}_N)$ , we can solve the finite state space MDP for the optimal  $\beta$ -discounted cost policy  $\hat{\gamma}_{N,\beta}$  and the optimal average cost policy  $\hat{\gamma}_{N,\infty}$ . We then extend them to the original state space as in Definition 6, and denote the extensions by  $\overline{\hat{\gamma}_{N,\beta}}$  and  $\overline{\hat{\gamma}_{N,\infty}}$ .

Next, we show that the extended policies are robust. We assume  $(X_n, U_n), n \geq 0$  is Harris positive recurrent with a unique invariance distribution  $\pi$ , such that  $\pi(B_i) > 0$  for  $1 \leq i \leq M$ . It is important to note that we do not need to know  $\pi$  above (as the model is unknown); this is just used for analysis purposes in the following discussion. Then by Theorem 17.1.7 in [40] and Corollary 2.9 in [11], for  $1 \leq i, j \leq M$  and  $u \in \mathbb{U}$ ,

$$\begin{aligned} \hat{\mathcal{T}}_N(y_j|y_i, u) &\xrightarrow{a.s.} \frac{\int_{B_i} \mathcal{T}(B_j|x, u)\pi(dx)\gamma_u}{\int_{B_i} \pi(dx)\gamma_u} = \mathcal{T}^{\pi, M}(y_j|y_i, u), \\ \hat{c}_N(y_i, u) &\xrightarrow{a.s.} \frac{\int_{B_i} c(x, u)\pi(dx)\gamma_u}{\int_{B_i} \pi(dx)\gamma_u} = c^{\pi, M}(y_i, u), \end{aligned} \tag{10}$$

where  $\mathcal{T}^{\pi, M}(y_j|y_i, u)$  and  $c^{\pi, M}(y_i, u)$  are defined in (8).

We extend the cost function  $\hat{c}_N$  to the original state space as in Definition 6, and denote it as  $\overline{\hat{c}_N}$ . We adopt a different extension for the kernel for technical analysis. Denote by  $\mathbb{Q}_M := \{y_i\}_{i=1}^M$ .

**DEFINITION 7 (PIECEWISE CONSTANT EXTENSION II).** Let  $\mathcal{S} : \mathcal{F}_{\mathbb{Q}_M} \times \mathbb{Q}_M \times \mathbb{U} \rightarrow [0, 1]$  be a controlled transition kernel. Define its piecewise constant extension  $\tilde{\mathcal{S}} : \mathcal{F}_{\mathbb{X}} \times \mathbb{X} \times \mathbb{U} \rightarrow [0, 1]$  as follows: for  $i = 1, \dots, M$ ,  $x \in B_i$ ,  $u \in \mathbb{U}$ , and  $A \in \mathcal{F}_{\mathbb{X}}$ ,

$$\tilde{\mathcal{S}}(A|x, u) := \sum_{j=1}^M \mathcal{S}(y_j|y_i, u) \mathbb{1}\{y_j \in A\}.$$

It is clear that  $\overline{\hat{\gamma}_{N,\beta}}$  and  $\overline{\hat{\gamma}_{N,\infty}}$  are the optimal discounted cost and average cost MDP  $(\mathbb{X}, \mathbb{U}, \overline{\hat{\mathcal{T}}_N}, \overline{\hat{c}_N})$ .

We compare the original and learned MDP, i.e.,  $(\mathcal{T}, c)$  and  $(\tilde{\mathcal{T}}_N, \overline{\hat{c}_N})$ , through the finite model approximation  $(\overline{\hat{\mathcal{T}}_N}^{\pi, M}, c^{\pi, M})$ ; see (8). We start with the discounted problem. Recall  $\delta_M$  from (9).

LEMMA 11. Under Assumption 5(b),  $d_{\mathcal{W}_1}(\widetilde{\mathcal{T}}^{\pi, M}, \mathcal{T}) \leq (1 + \|\mathcal{T}\|_{\text{Lip}})\delta_M$ .

*Proof.* Fix an arbitrary  $f : \mathbb{X} \rightarrow \mathbb{R}$  with  $\|f\|_{\text{Lip}} \leq 1$ ,  $u \in \mathbb{U}$  and  $x \in B_i$  with  $1 \leq i \leq M$ . Note that

$$\int f(z) \widetilde{\mathcal{T}}^{\pi, M}(dz|x, u) = \int \sum_{B_j}^M f(y_j) \mathcal{T}(B_j|w, u) \hat{\pi}_i(dw).$$

For any  $w \in B_i$ , we have

$$\begin{aligned} & \left| \sum_{j=1}^M f(y_j) \mathcal{T}(B_j|w, u) - \int f(z) \mathcal{T}(dz|x, u) \right| \\ &= \left| \sum_{j=1}^M \int_{B_j} (f(y_j) - f(z)) \mathcal{T}(dz|w, u) + \int f(z) \mathcal{T}(dz|w, u) - \int f(z) \mathcal{T}(dz|x, u) \right| \leq (1 + \|\mathcal{T}\|_{\text{Lip}})\delta_M, \end{aligned}$$

which completes the proof.  $\square$

COROLLARY 11. Suppose  $\delta_M \rightarrow 0$  as  $M \rightarrow \infty$ . Under Assumption 1 and 5,

$$\mathbb{P} \left( \lim_{M \rightarrow \infty} \limsup_{N \rightarrow \infty} \|J_\beta(c, \mathcal{T}, \overline{\hat{\gamma}_{N, \beta}}) - J_\beta^*(c, \mathcal{T})\|_\infty = 0 \right) = 1.$$

*Proof.* Fix some outcome  $\omega \in \Omega$  such that (10) holds. Since  $M$  and  $\mathbb{U}$  is finite, we have that  $\limsup_{N \rightarrow \infty} \|c^{\pi, M} - \hat{c}_N\|_\infty = 0$ , and that

$$\limsup_{N \rightarrow \infty} d_{\mathcal{W}_1}(\mathcal{T}^{\pi, M}, \hat{\mathcal{T}}_N) = \limsup_{N \rightarrow \infty} \sup_{1 \leq i \leq M, u \in \mathbb{U}} d_{\mathcal{W}_1}(\mathcal{T}^{\pi, M}(\cdot|y_i, u), \hat{\mathcal{T}}_N(\cdot|y_i, u)) = 0.$$

By the triangle inequality, and due to the definition of extensions in (7),

$$d_{\mathcal{W}_1}(\mathcal{T}, \widetilde{\hat{\mathcal{T}}}_N) \leq d_{\mathcal{W}_1}(\mathcal{T}, \widetilde{\mathcal{T}}^{\pi, M}) + d_{\mathcal{W}_1}(\mathcal{T}^{\pi, M}, \hat{\mathcal{T}}_N). \quad (11)$$

Then due to Lemma 10 and Lemma 11,

$$\limsup_{N \rightarrow \infty} \|c - \overline{\hat{c}_N}\|_\infty \leq \|c\|_{\text{Lip}} \delta_M, \quad \limsup_{N \rightarrow \infty} d_{\mathcal{W}_1}(\mathcal{T}, \widetilde{\hat{\mathcal{T}}}_N) \leq (1 + \|\mathcal{T}\|_{\text{Lip}})\delta_M.$$

Then the proof is complete by Corollary 4.  $\square$

Finally, we focus on the average cost problem. Assume the reference kernel  $\mathcal{T}$  satisfies the minorization condition in Definition 5 with some probability measure  $\underline{\rho}$  and scaling constant  $\epsilon$ . We first find a minorization measure  $\tau^M$  for the approximating kernel  $\widetilde{\mathcal{T}}^{\pi, M}$ , and then bound the Wasserstein-1 distance between  $\rho$  and  $\tau^M$ .

LEMMA 12. Assume there exist a constant  $\epsilon > 0$  and a probability measures  $\rho \in \mathcal{P}(\mathbb{X})$ , such that  $\forall x \in \mathbb{X}, u \in \mathbb{U}$ , and  $A \in \mathcal{F}_{\mathbb{X}}$ ,  $\mathcal{T}(A|x, u) \geq \epsilon \rho(A)$ . Define the following probability measure  $\tau^M$ :

$$\tau^M(A) := \sum_{j=1}^M \rho(B_j) \mathbb{1}\{y_j \in A\}, \quad \text{for any } A \in \mathcal{F}_{\mathbb{X}}.$$

Then for any  $x \in \mathbb{X}$ ,  $u \in \mathbb{U}$  and  $A \in \mathcal{F}_{\mathbb{X}}$ ,

$$\widetilde{\mathcal{T}}^{\pi, M}(A|x, u) \geq \epsilon \tau^M(A) \quad \text{and} \quad \mathcal{W}_1(\rho, \tau^M) \leq \delta_M.$$

*Proof.* For any  $A \in \mathcal{F}_{\mathbb{X}}$ ,  $u \in \mathbb{U}$  and  $x \in B_i$  with  $1 \leq i \leq M$ , by definition,

$$\widetilde{\mathcal{T}}^{\pi, M}(A|x, u) = \sum_{j=1}^M \int_{B_j} \mathcal{T}(B_j|x, u) \hat{\pi}_i(dx) \mathbb{1}\{y_j \in A\} \geq \sum_{j=1}^M \int_{B_j} \rho(B_j) \hat{\pi}_i(dx) \mathbb{1}\{y_j \in A\}.$$

Since  $\hat{\pi}_i(B_i) = 1$  for  $1 \leq i \leq M$ , we have  $\widetilde{\mathcal{T}}^{\pi, M}(A|x, u) \geq \epsilon \tau^M(A)$ .

Further, for any  $f: \mathbb{X} \rightarrow \mathbb{R}$  with  $\|f\|_{\text{Lip}} \leq 1$ , by the triangle inequality,

$$\left| \int f(x) \rho(dx) - \int f(x) \tau^M(dx) \right| \leq \sum_{j=1}^M \left| \int_{B_j} f(x) \rho(dx) - \int_{B_j} f(y_j) \rho(B_j) \right| \leq \sum_{j=1}^M \int_{B_j} |f(x) - f(y_j)| \rho(dx),$$

which can be further bounded by  $\delta_M$ . The proof is complete.  $\square$

**COROLLARY 12.** *Suppose that assumption 6 and the minorization condition in Definition 5 hold for the reference MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$ , and that  $\delta_M \rightarrow 0$  as  $M \rightarrow \infty$ . Under Assumption 1 and 6,*

$$\mathbb{P} \left( \lim_{M \rightarrow \infty} \limsup_{N \rightarrow \infty} \|J_{\infty}(c, \mathcal{T}, \overline{\hat{\gamma}}_{N, \infty}) - J_{\infty}^*(c, \mathcal{T})\|_{\infty} = 0 \right) = 1.$$

*Proof.* Fix some outcome  $\omega \in \Omega$  such that (10) holds. First, fix some  $M$ . Due to Lemma 12, there exists some  $N^*$ , depending on  $\omega$  and  $M$ , such that for  $N \geq N^*$ ,

$$\widetilde{\mathcal{T}}_N(A|x, u) \geq \frac{\epsilon}{2} \tau^M(A), \text{ for } x \in \mathbb{X}, u \in \mathbb{U} \text{ and } A \in \mathcal{F}_{\mathbb{X}}.$$

The proof is completed by a similar argument as for Corollary 11, Corollary 6, and Lemma 12.  $\square$

**4.2.2. Empirical model learning from independently generated transition data** Let  $\pi \in \mathcal{P}(\mathbb{X})$  be a given weighting measure, such that  $\pi(B_i) > 0$  for  $1 \leq i \leq M$ . For  $1 \leq i \leq M$ , we denote its restriction on bin  $B_i$  by  $\hat{\pi}_i$  as defined in (7). We consider the data collected under the scheme described in Algorithm 2. Specifically, for each  $1 \leq i \leq M$  and action  $u \in \mathbb{U}$ , we have  $N$  independent triplets of observations,  $(X_{k,i,u}, Y_{k,i,u}, c(X_{k,i,u}, u))$ ,  $1 \leq k \leq N$ , where

$$X_{k,i,u} \sim \hat{\pi}_i, \quad Y_{k,i,u} | X_{k,i,u} \sim \mathcal{T}(\cdot | X_{k,i,u}, u).$$

Note that in the above  $\hat{\pi}_i$  has its full measure on  $B_i$  and that the total number of triplets is  $M \times |\mathbb{U}| \times N$ . Given the data, the estimated controlled kernel and cost function  $(\hat{\mathcal{T}}_N, \hat{c}_N)$  on the quantized space  $\{y_i\}_{i=1}^M$  are defined as follows. For each  $1 \leq i \leq M$  and action  $u \in \mathbb{U}$ ,

$$\hat{c}_N(y_i, u) := \frac{1}{N} \sum_{k=1}^N c(X_{k,i,u}, u), \quad \hat{\mathcal{T}}_N(y_j | y_i, u) := \frac{1}{N} \sum_{k=1}^N \mathbb{1}\{Y_{k,i,u} \in B_j\}, \text{ for } 1 \leq j \leq M. \quad (12)$$

As in the previous section, we can solve the finite state space MDP  $(\{y_i\}_{i=1}^M, \mathbb{U}, \hat{\mathcal{T}}_N, \hat{c}_N)$  for the optimal  $\beta$ -discounted cost policy  $\hat{\gamma}_{N, \beta}$  and the optimal average cost policy  $\hat{\gamma}_{N, \infty}$ . We then extend them to the original state space, denoted as  $\overline{\hat{\gamma}}_{N, \beta}$  and  $\overline{\hat{\gamma}}_{N, \infty}$ , which are the optimal policies for  $(\mathbb{X}, \mathbb{U}, \widetilde{\mathcal{T}}_N, \overline{\hat{c}}_N)$ , where we recall the extension of kernel  $\widetilde{\mathcal{T}}_N$  in Definition 7.

Next, we show the extended policies are robust with non-asymptotic upper bounds. For simplicity, we assume  $\mathbb{X} \subset \mathbb{R}^d$  for some  $d \geq 3$ ; the discussions for  $d = 1, 2$  are similar. Further, assume that for some constant  $L > 0$ ,

$$\sup_{x \in \mathbb{X}} \|x\| \leq L, \quad \|c\|_{\infty} \leq L. \quad (13)$$

We first bound the error due to quantization and learning. For a random variable  $Z$ , define its  $\psi_p$  norm for  $p \geq 1$  as follows:  $\|Z\|_{\psi_p} := \inf\{C > 0 : \exp(|Z|/C)^p \leq 2\}$ ; see Chapter 2.2 in [58]. As discussed in Chapter 2.2 of [58], for some constant  $C$ , that only depends on  $p$ ,  $\mathbb{E}|Z| \leq C \|Z\|_{\psi_p}$ .

**Algorithm 2** Sampling a Wasserstein Regular MDP with Restart

---

**Require:** number of repetitions  $N$  for each state and action pair

- 1: **for all**  $i = 1, \dots, M$  **do**
- 2:     **for all**  $u \in \mathbb{U}$  **do**
- 3:         **for all**  $k = 1, \dots, N$  **do**
- 4:             Sample  $X_{k,i,u} \sim \hat{\pi}_i$
- 5:             Sample i.i.d  $Y_{k,i,u} \sim \mathcal{T}(\cdot | X_{k,i,u}, u)$
- 6:             Obtain cost  $c(X_{k,i,u}, u)$
- 7:         **end for**
- 8:         Set  $\hat{c}_N(y_i, u) := \frac{1}{N} \sum_{k=1}^N c(X_{k,i,u}, u)$
- 9:         Set  $\hat{\mathcal{T}}_N(y_j | y_i, u) := \frac{1}{N} \sum_{k=1}^N \mathbb{1}\{Y_{k,i,u} \in B_j\}$  for  $j = 1, 2, \dots, M$
- 10:     **end for**
- 11: **end for**
- 12: **return**  $(\hat{\mathcal{T}}_N, \hat{c}_N)$

---

LEMMA 13. Under Assumption 5(a) and (b), for some constant  $C$  that depends only on  $d, L$ ,

$$\begin{aligned} \mathbb{E} [\|c - \bar{c}_N\|_\infty] &\leq \|c\|_{Lip} \delta_M + C \left( \frac{\log(M|\mathbb{U}|)}{N} \right)^{1/2}, \\ \mathbb{E} \left[ d_{W_1}(\mathcal{T}, \widetilde{\mathcal{T}}_N) \right] &\leq (1 + \|\mathcal{T}\|_{Lip}) \delta_M + C \left( \frac{\log(M|\mathbb{U}|)}{N} \right)^{1/d}. \end{aligned}$$

*Proof.* In this proof,  $C$  denotes constants, only depending on  $d$  and  $L$ , which may vary from line to line. For  $1 \leq i \leq M$  and  $u \in \mathbb{U}$ ,

$$\mathbb{E}[c(X_{k,i,i}, u)] = c^{\pi, M}(y_i, u), \text{ for } 1 \leq k \leq N,$$

where  $c^{\pi, M}$  is defined in (8). Due to independence, since  $\|c\|_\infty \leq L$ , by Hoeffding's inequality (see, e.g., Theorem 2.8 in [5]),  $\mathbb{P} \left( \sqrt{N} |\hat{c}_N(y_i, u) - c^{\pi, M}(y_i, u)| \geq t \right) \leq 2e^{-t^2/(2L^2)}$ , which, by Lemma 2.2.1 in [58], implies that  $\left\| \sqrt{N} |\hat{c}_N(y_i, u) - c^{\pi, M}(y_i, u)| \right\|_{\psi_2} \leq (6L^2)^{1/2}$ . Thus, by Lemma 2.2.2 in [58],

$$\mathbb{E} \left[ \sup_{1 \leq i \leq M, u \in \mathbb{U}} \sqrt{N} |\hat{c}_N(y_i, u) - c^{\pi, M}(y_i, u)| \right] \leq C \sqrt{\log(M|\mathbb{U}|)},$$

which implies the first statement due to Lemma 10 and the triangle inequality. Next, we focus on the second statement. For  $1 \leq i \leq M$  and  $u \in \mathbb{U}$ , we have

$$\mathbb{P}(Y_{k,i,u} \in B_j) = \mathcal{T}^{\pi, M}(y_j | y_i, u), \text{ for } 1 \leq j \leq M, 1 \leq k \leq N, \quad (14)$$

where  $\mathcal{T}^{\pi, M}$  is defined in (8). Thus,  $\hat{T}_N(\cdot | y_i, u)$  is the empirical distribution of a random sample of size  $N$  from  $\mathcal{T}^{\pi, M}(\cdot | y_i, u)$ , both of which are defined on  $\{y_j\}_{j=1}^M$ . By the definition of  $L$  in (13),

$$\sum_{j=1}^M e^{\|y_j\|^d} \mathcal{T}^{\pi, M}(y_j | y_i, u) \leq \exp(L^d).$$

Thus since  $d \geq 3$ , by Theorem 2 in [13], for all  $x > 0$ ,

$$\mathbb{P} \left( N^{1/d} d_{W_1} \left( \hat{T}_N(\cdot | y_i, u), \mathcal{T}^{\pi, M}(\cdot | y_i, u) \right) \geq x \right) \leq C \exp(-x^d/C),$$

which, by Lemma 2.2.1 in [58], implies that  $\left\| N^{1/d} d_{\mathcal{W}_1} \left( \hat{T}_N(\cdot|y_i, u), \mathcal{T}^{\pi, M}(\cdot|y_i, u) \right) \right\|_{\psi_d} \leq C$ . As a result, by Lemma 2.2.2 in [58],

$$\mathbb{E} \left[ d_{\mathcal{W}_1} \left( \hat{T}_N, \mathcal{T}^{\pi, M} \right) \right] = \mathbb{E} \left[ \sup_{1 \leq i \leq M, u \in \mathbb{U}} d_{\mathcal{W}_1} \left( \hat{T}_N(\cdot|y_i, u), \mathcal{T}^{\pi, M}(\cdot|y_i, u) \right) \right] \leq C (\log(M|\mathbb{U}|))^{1/d} N^{-1/d}.$$

Then the proof is complete due to Lemma 11 and the triangle inequality (11).  $\square$

COROLLARY 13. *Under Assumption 1 and 5,*

$$\mathbb{E} \left( \| J_{\beta}(c, \mathcal{T}, \overline{\hat{\gamma}_{N, \beta}}) - J_{\beta}^*(c, \mathcal{T}) \|_{\infty} \right) \leq C \left( \delta_M + \left( \frac{\log(M|\mathbb{U}|)}{N} \right)^{1/2} + \left( \frac{\log(M|\mathbb{U}|)}{N} \right)^{1/d} \right),$$

where  $C$  is a constant, depending only on  $\beta$ ,  $\|c\|_{Lip}$ ,  $\|\mathcal{T}\|_{Lip}$ ,  $d$  and  $L$ .

*Proof.* It follows from Corollary 4 and Lemma 13.  $\square$

Recall that  $\delta_M$  is the quantization error in (9). By a volume argument, due to the definition of  $L$  in (13), there exists a quantization scheme such that

$$\delta_M \leq (M/C)^{-1/d}, \tag{15}$$

for some constant  $C$ , depending only on  $d$  and  $L$ . Under this assumption, if

$$N = CM \log(M|\mathbb{U}|), \tag{16}$$

then we have

$$\mathbb{E} \left( \| J_{\beta}(c, \mathcal{T}, \overline{\hat{\gamma}_{N, \beta}}) - J_{\beta}^*(c, \mathcal{T}) \|_{\infty} \right) \leq C \left( M^{-1/d} + \left( \frac{\log(M|\mathbb{U}|)}{N} \right)^{1/d} \right) \leq CM^{-1/d}.$$

Note that the choice of  $N$  in (16) balances the error due to finite model approximation and due to finite model learning for a given, large  $M$ . The expected regret is of order  $M^{-1/d}$ .

Finally, we consider the average cost problem. Define

$$\kappa_M := \min \{ \mathcal{T}^{\pi, M}(y_j|y_i, u) : (i, j, u) \in \mathcal{I}_M \}, \tag{17}$$

where  $\mathcal{I}_M := \{ 1 \leq i, j \leq M, u \in \mathbb{U} : \mathcal{T}^{\pi, M}(y_j|y_i, u) > 0 \}$ . Thus,  $\kappa_M$  is the smallest non-zero element in  $\{ \mathcal{T}^{\pi, M}(y_j|y_i, u) : 1 \leq i, j \leq M, u \in \mathbb{U} \}$ , where  $\mathcal{T}^{\pi, M}$  is defined in (8). Further, define the event

$$\mathcal{E}_{N, M} := \bigcup_{1 \leq i, j \leq M, u \in \mathbb{U}} \left\{ \hat{T}_N(y_j|y_i, u) < \frac{1}{2} \mathcal{T}^{\pi, M}(y_j|y_i, u) \right\}. \tag{18}$$

We can upper bound the probability of  $\mathcal{E}_{N, M}$ , and on its complement,  $\widetilde{\hat{T}}_N$  is minorized. Recall the probability measure  $\tau^M$  in Lemma 12.

LEMMA 14. *The following upper bound holds:  $\mathbb{P}(\mathcal{E}_{N, M}) \leq 2M^2|\mathbb{U}| \exp(-N\kappa_M/10)$ . Further, suppose the minorization condition in Definition 5 hold for the reference MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$ . On the event  $(\mathcal{E}_{N, M})^c$ , for any  $x \in \mathbb{X}$ ,  $u \in \mathbb{U}$  and  $A \in \mathcal{F}_{\mathbb{X}}$ ,  $\widetilde{\hat{T}}_N(A|x, u) \geq (\epsilon/2)\tau^M(A)$ .*

*Proof.* For each  $(i, j, u) \in \mathcal{I}_M$ , in view of (14), by Bernstein's inequality [58, Lemma 2.2.9],

$$\mathbb{P} \left( \left| \hat{T}_N(y_j|y_i, u) - \mathcal{T}^{\pi, M}(y_j|y_i, u) \right| \geq 2^{-1} \mathcal{T}^{\pi, M}(y_j|y_i, u) \right) \leq 2 \exp(-N\kappa_M/10).$$

Then the first statement follows from the union bound. Further, the second statement follows from the definition of  $\mathcal{E}_{N, M}$  in (18) and Lemma 12.  $\square$

COROLLARY 14. *Suppose Assumption 6 and the minorization condition in Definition 5 hold for the reference MDP  $(\mathbb{X}, \mathbb{U}, \mathcal{T}, c)$ . Under Assumption 1 and 6,*

$$\mathbb{E} \left[ \left\| J_\infty(c, \mathcal{T}, \overline{\hat{\gamma}_{N,\infty}}) - J_\infty^*(c, \mathcal{T}) \right\|_\infty \right] \leq C \left( \delta_M + \left( \frac{\log(M|\mathbb{U}|)}{N} \right)^{1/2} + \left( \frac{\log(M|\mathbb{U}|)}{N} \right)^{1/d} \right) + 4LM^2|\mathbb{U}|\exp(-N\kappa_M/10).$$

where  $C$  is a constant, depending only on  $\epsilon$ ,  $\|c\|_{Lip}$ ,  $\|\mathcal{T}\|_{Lip}$ ,  $d$  and  $L$ .

*Proof.* By the definition of  $L$  in (13), due to Lemma 14,

$$\mathbb{E} \left[ \left\| J_\infty(c, \mathcal{T}, \overline{\hat{\gamma}_{N,\infty}}) - J_\infty^*(c, \mathcal{T}) \right\|_\infty ; \mathcal{E}_{N,M} \right] \leq 2L\mathbb{P}(\mathcal{E}_{N,M}) \leq 4LM^2|\mathbb{U}|\exp(-N\kappa_M/10).$$

On the event  $(\mathcal{E}_{N,M})^c$ , by Lemma 14,  $\tilde{T}_N$  is minorized by the probability measure  $\tau^M$  with constant  $\epsilon/2$ . Thus, by Corollary 6, Lemma 13, and Lemma 12,

$$\mathbb{E} \left[ \left\| J_\infty(c, \mathcal{T}, \overline{\hat{\gamma}_{N,\infty}}) - J_\infty^*(c, \mathcal{T}) \right\|_\infty ; (\mathcal{E}_{N,M})^c \right] \leq C \left( \delta_M + \left( \frac{\log(M|\mathbb{U}|)}{N} \right)^{1/2} + \left( \frac{\log(M|\mathbb{U}|)}{N} \right)^{1/d} \right).$$

Then the proof is complete.  $\square$

Assume  $\delta_M$  satisfies (15), and  $\kappa_M$  in (17) satisfies  $\kappa_M \geq 1/(CM)$  for some constant  $C > 0$ . Then if we choose  $N$  as in (16) for some large enough constant  $C$ , then again we have  $\mathbb{E} \left[ \left\| J_\infty(c, \mathcal{T}, \overline{\hat{\gamma}_{N,\infty}}) - J_\infty^*(c, \mathcal{T}) \right\|_\infty \right] \leq CM^{-1/d}$ .

**5. Concluding Remarks** In this paper, we consider the Wasserstein model approximation problem, where we upper bound the performance loss of applying an optimal policy from an approximate model to the true dynamics. This loss is bounded by the sup-norm-induced metric between the approximate and true costs, as well as the Wasserstein-1 distance between the approximate and true transition kernels. We study both the discounted cost and average cost criteria. Based on these results, we develop empirical model learning algorithms, establish their empirical consistency, and obtain sample complexity bounds. Additionally, we recover and generalize several existing results on continuous dependence, robustness and approximations.

An extension of the present work would be robustness of finite step value iteration, which generalizes [47], as clarified in the following observations. Consider applying Corollary 7 to the setup where  $c = \hat{c}$  and both the true and approximate system are control-free, i.e.

$$c(x, u) = c(x), \mathcal{T}(\cdot|x, u) = \mathcal{T}(\cdot|x), \mathcal{S}(\cdot|x, u) = \mathcal{S}(\cdot|x).$$

Further, suppose both probability kernels admit invariant probability measures, and denote them by  $\rho_\mathcal{T}$  and  $\rho_\mathcal{S}$ , respectively. By the result in Corollary 7, we have the following bound

$$\left| \int_{\mathbb{X}} c(x) \rho_\mathcal{T}(dx) - \int_{\mathbb{X}} c(x) \rho_\mathcal{S}(dx) \right| \leq \frac{\|c\|_{Lip}}{1 - \|\mathcal{T}\|_{Lip}} \sup_{x \in \mathbb{X}} (\mathcal{W}_1(\mathcal{T}(\cdot|x), \mathcal{S}(\cdot|x))).$$

We note that the choice of the cost function is arbitrary as long as it is Lipschitz. Restricting our attention to those with Lipschitz constant less than or equal to 1, and taking supremum over such cost functions on both sides, we have

$$\mathcal{W}_1(\rho_\mathcal{T}, \rho_\mathcal{S}) \leq \frac{1}{1 - \|\mathcal{T}\|_{Lip}} \sup_{x \in \mathbb{X}} (\mathcal{W}_1(\mathcal{T}(\cdot|x), \mathcal{S}(\cdot|x))).$$

This recovers Corollary 3.1 of [47] in an unweighted form. It is therefore natural to consider that a performance bound for finite-step approximate value iteration would be analogous to Theorem 3.1 in [47]. Additional future research directions include applying the approximation scheme within a parametric setting, as explored in a recent work [3], and non-asymptotic analog of Section 4.2.1, in view of recent works [29, 28].



## References

- [1] Agarwal A, Jiang N, Kakade SM, Sun W (2021) *Reinforcement Learning: Theory and Algorithms*. URL <https://rltheorybook.github.io/>.
- [2] Asadi K, Misra D, Littman M (2018) Lipschitz continuity in model-based reinforcement learning. Dy J, Krause A, eds., *Proceedings of the 35th International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, 264–273 (PMLR).
- [3] Bayraktar E, Kara AD (2024) Learning with linear function approximations in mean-field control. *arXiv preprint arXiv:2408.00991* .
- [4] Bertsekas D (1975) Convergence of discretization procedures in dynamic programming. *IEEE Transactions on Automatic Control* 20(3):415–419.
- [5] Boucheron S, Lugosi G, Massart P (2013) *Concentration Inequalities: A Nonasymptotic Theory of Independence* (Oxford University Press), ISBN 9780199535255.
- [6] Bozkurt B, Mahajan A, Nayyar A, Ouyang Y (2024) Model approximation in MDPs with unbounded per-step cost. *arXiv preprint arXiv:2402.08813* .
- [7] Demirci Y, Kara A, Yüksel S (2024) Average cost optimality of partially observed MDPs: Contraction of non-linear filters and existence of optimal solutions. *SIAM Journal on Control and Optimization*, *arXiv:2312.14111* .
- [8] Dudley RM (1969) The speed of mean Glivenko-Cantelli convergence. *The Annals of Mathematical Statistics* 40(1):40–50, ISSN 00034851.
- [9] Dudley RM (2002) *Real Analysis and Probability* (Cambridge University Press), ISBN 9780511755347.
- [10] Dufour F, Prieto-Rumeau T (2015) Approximation of average cost Markov decision processes using empirical distributions and concentration inequalities. *Stochastics* 87(2):273–307.
- [11] Eberle A (2023) Markov processes. <https://uni-bonn.sciebo.de/s/kzTUFff5FrWGAay>.
- [12] Figalli A, Glaudo F (2021) *An Invitation to Optimal Transport, Wasserstein Distances, and Gradient Flows* (EMS Press), ISBN 9783985475100.
- [13] Fournier N, Guillin A (2014) On the rate of convergence in Wasserstein distance of the empirical measure. *Probability Theory and Related Fields* 162(3–4):707–738, ISSN 1432-2064.
- [14] Fox B (1971) Finite-state approximations to denumerable-state dynamic programs. *Journal of Mathematical Analysis and Applications* 34(3):665–670, ISSN 0022-247X.
- [15] Gao R, Kleywegt A (2023) Distributionally robust stochastic optimization with Wasserstein distance. *Math. Oper. Res.* 48(2):603–655, ISSN 0364-765X.
- [16] Gordienko E, Chavez JRd (2022) Stability estimation of some Markov controlled processes. *Open Mathematics* 20(1):1509–1520, ISSN 2391-5455.
- [17] Gordienko E, Lemus-Rodríguez E, Montes-de Oca R (2007) Discounted cost optimality problem: stability with respect to weak metrics. *Mathematical Methods of Operations Research* 68(1):77–96, ISSN 1432-5217.
- [18] Gordienko E, Lemus-Rodríguez E, Montes-de Oca R (2008) Average cost Markov control processes: stability with respect to the Kantorovich metric. *Mathematical Methods of Operations Research* 70(1):13–33, ISSN 1432-5217.
- [19] Gordienko EI (1988) Stability estimates for controlled Markov chains with a minorant. *Journal of Soviet Mathematics* 40(4):481–486, ISSN 1573-8795.
- [20] Gordienko EI (1992) An estimate of the stability of optimal control of certain stochastic and deterministic systems. *Journal of Soviet Mathematics* 59(4):891–899, ISSN 1573-8795.
- [21] Gordienko EI, Salem FS (1998) Robustness inequality for Markov control processes with unbounded costs. *Systems and Control Letters* 33(2):125–130, ISSN 0167-6911.
- [22] Gordienko EI, Salem-Silva F (2000) Estimates of stability of Markov control processes with unbounded costs. *Kybernetika* 36(2):[195]–210.
- [23] Hanson J, Raginsky M, Sontag E (2021) Learning recurrent neural net models of nonlinear systems. *Learning for Dynamics and Control*, 425–435 (PMLR).

- [24] Hernández-Lerma O (1987) Approximation and adaptive control of markov processes: Average reward criterion. *Kybernetika* 23(4):265–288.
- [25] Hernández-Lerma O (1989) *Adaptive Markov Control Processes* (Springer New York), ISBN 9781441987143.
- [26] Hernández-Lerma O, Lasserre JB (1996) *Discrete-Time Markov Control Processes* (Springer New York), ISBN 9781461207290.
- [27] Hinderer K (2005) Lipschitz continuity of value functions in Markovian decision processes. *Mathematical Methods of Operations Research* 62(1):3–22, ISSN 1432-5217.
- [28] Huang D, Li X (2024) Bernstein-type inequalities for markov chains and markov processes: A simple and robust proof. *arXiv preprint arXiv:2408.04930* .
- [29] Huang D, Li X (2024) Non-asymptotic estimates for markov transition matrices with rigorous error bounds. *arXiv preprint arXiv:2408.05963* .
- [30] Johnson J (2018) Rethinking floating point for deep learning. *arXiv preprint arXiv:1811.01721* .
- [31] Kara A, Saldi N, Yüksel S (2023) Q-learning for MDPs with general spaces: Convergence and near optimality via quantization under weak continuity. *Journal of Machine Learning Research* 1–34.
- [32] Kara A, Yüksel S (2020) Robustness to incorrect system models in stochastic control. *SIAM Journal on Control and Optimization* 58(2):1144–1182.
- [33] Kara A, Yüksel S (2022) Near optimality of finite memory feedback policies in partially observed Markov decision processes. *Journal of Machine Learning Research* 23(11):1–46.
- [34] Kara A, Yüksel S (2023) Q-learning for continuous state and action MDPs under average cost criteria. *arXiv preprint arXiv:2308.07591* .
- [35] Kara A, Yüksel S (2024) Q-learning for stochastic control under general information structures and non-Markovian environments. *Transactions on Machine Learning Research (arXiv:2311.00123)* .
- [36] Kara AD, Raginsky M, Yüksel S (2022) Robustness to incorrect models and data-driven learning in average-cost optimal stochastic control. *Automatica* 139:110179.
- [37] Kara AD, Yüksel S (2021) Robustness to approximations and model learning in MDPs and POMDPs. Piunovskiy AB, Zhang Y, eds., *Modern Trends in Controlled Stochastic Processes: Theory and Applications, Volume III* (Luniver Press).
- [38] Langen H (1981) Convergence of dynamic programming models. *Mathematics of Operations Research* 6(4):493–512.
- [39] Mendelson S (2014) Learning without concentration. *arXiv preprint arXiv:1401.0304* .
- [40] Meyn SP, Tweedie R (1993) *Markov Chains and Stochastic Stability* (London: Springer-Verlag).
- [41] Mohajerin Esfahani P, Kuhn D (2017) Data-driven distributionally robust optimization using the Wasserstein metric: performance guarantees and tractable reformulations. *Mathematical Programming* 171(1–2):115–166, ISSN 1436-4646.
- [42] Müller A (1997) How does the value function of a markov decision process depend on the transition probabilities? *Mathematics of Operations Research* 22(4):872–885.
- [43] Namkoong H, Duchi JC (2016) Stochastic gradient methods for distributionally robust optimization with f-divergences. Lee D, Sugiyama M, Luxburg U, Guyon I, Garnett R, eds., *Advances in Neural Information Processing Systems*, volume 29 (Curran Associates, Inc.).
- [44] Pirodda M, Restelli M, Bascetta L (2015) Policy gradient in lipschitz markov decision processes. *Machine Learning* 100(2–3):255–283, ISSN 1573-0565.
- [45] Pradhan S, Yüksel S (2023) Robustness of stochastic optimal control to approximate diffusion models under several cost evaluation criteria. *Mathematics of Operations Research* .
- [46] Rachelson E, Lagoudakis MG (2010) On the locality of action domination in sequential decision making. *International Symposium on Artificial Intelligence and Mathematics*.
- [47] Rudolf D, Schweizer N (2018) Perturbation theory for Markov chains via Wasserstein distance. *Bernoulli* 24(4A), ISSN 1350-7265.

- [48] Saldi N, Linder T, Yüksel S (2015) Finite state approximations of Markov decision processes with general state and action spaces. *2015 American Control Conference (ACC)* (IEEE).
- [49] Saldi N, Linder T, Yüksel S (2018) *Finite Approximations in Discrete-Time Stochastic Control: Quantized Models and Asymptotic Optimality* (Cham: Springer).
- [50] Saldi N, Yüksel S, Linder T (2015) Finite-state approximation of Markov decision processes with unbounded costs and Borel spaces. *2015 54th IEEE Conference on Decision and Control (CDC)* (IEEE).
- [51] Saldi N, Yüksel S, Linder T (2017) On the asymptotic optimality of finite approximations to Markov decision processes with Borel spaces. *Mathematics of Operations Research* 42(4):945–978, ISSN 1526-5471.
- [52] Tsybakov AB (2009) *Introduction to Nonparametric Estimation* (Springer New York), ISBN 9780387790527.
- [53] Van Roy B (2006) Performance loss bounds for approximate value iteration with state aggregation. *Mathematics of Operations Research* 31(2):234–244, ISSN 1526-5471.
- [54] Vega-Amaya O, López-Borbón J (2019) A perturbation approach to approximate value iteration for average cost markov decision processes with borel spaces and bounded costs. *Kybernetika* 55(1):81–113.
- [55] Villani C (2009) *Optimal Transport* (Springer Berlin Heidelberg), ISBN 9783540710509.
- [56] Wang S, Si N, Blanchet J, Zhou Z (2024) Statistical learning of distributionally robust stochastic control in continuous state spaces. *arXiv preprint arXiv:2406.11281* .
- [57] Weed J, Bach F (2019) Sharp asymptotic and finite-sample rates of convergence of empirical measures in Wasserstein distance. *Bernoulli* 25(4A), ISSN 1350-7265.
- [58] Wellner J, et al. (2013) *Weak convergence and empirical processes: with applications to statistics* (Springer Science & Business Media).
- [59] Yang I (2021) Wasserstein distributionally robust stochastic control: A data-driven approach. *IEEE Transactions on Automatic Control* 66(8):3863–3870.
- [60] Yu H, Bertsekas D (2012) Discretized approximations for POMDP with average cost. *arXiv preprint arXiv:1207.4154* .
- [61] Yüksel S (2024) Optimization and control of stochastic systems. <https://mast.queensu.ca/~yukse1/LectureNotesOnStochasticOptControl.pdf/>.