

QRT MAP ON A BIELLIPTIC SURFACE

NALINI JOSHI, FRANK W. NIJHOFF, AND ALLAN STEEL

ABSTRACT. The family of mappings of the plane possessing a biquadratic invariant, which is known collectively as QRT maps, is composed of two involutions, one preserving a vertical shift and the other preserving a horizontal shift in the plane. In this paper, we extend the map by replacing each shift by the group operation on each of two families of elliptic curves, whose product forms a bielliptic surface.

1. INTRODUCTION

The Quispel-Roberts-Thompson (QRT) map is an 8-parameter family of mappings of the plane preserving a biquadratic invariant [11], cf. also [8]. It is the composition of two involutions, which can be expressed as the addition formula on a rational elliptic surface [15, 5]. The QRT map has played an important role in the development of the theory of discrete integrable systems, notably in the discovery of discrete Painlevé equations, see e.g. [8, Chapter 11] and [9]. In this paper, we generalise the QRT map to bielliptic surfaces¹.

Let Γ and Γ^* denote two elliptic curves with respective uniformizing variables ξ and η , given by

$$\Gamma : X^2 = 4x^3 - g_2x - g_3 = 4(x - e_1)(x - e_2)(x - e_3), \quad (1.1)$$

$$\Gamma^* : Y^2 = 4y^3 - G_2y - G_3 = 4(x - E_1)(x - E_2)(x - E_3), \quad (1.2)$$

with branch points given by coordinates e_1, e_2, e_3 and E_1, E_2, E_3 respectively, where we assume that, in general, the curves Γ and Γ^* are non-degenerate, i.e.,

$$27g_3^2 \neq g_2^3, \quad 27G_3^2 \neq G_2^3.$$

Furthermore, we do not impose any conditions on how the two curves Γ and Γ^* are related to each other; we leave the moduli g_2, g_3, G_2, G_3 undetermined (including their possible dependence on the variables of the other curve).

In this paper, we define two involutions ι_j , $j = 1, 2$, one acting on Γ and the other on Γ^* , by the group action on each respective curve. We will consider these involutions as a dynamical system on a bielliptic surface (see section 4 for the standard definition of a bielliptic surface). For this reason, we will call the composition of the two involutions the *bielliptic QRT map*. In remark 2.3 we show how the invariant of the bielliptic QRT map degenerates to that of the standard QRT map in a singular limit of the elliptic curves.

The involutions ι_j are introduced in section 2, where we also show that their composition is symplectic. Our focus lies primarily on the case when ι_1 acts on Γ while Γ^* is preserved and ι_2 acts on Γ^* while Γ is preserved. Under the condition that g_2, g_3 are independent of y and G_2, G_3 are independent of y , we provide the generic 10-parameter family of all such maps in Section 3, along with an explicit example.

2020 *Mathematics Subject Classification.* 37J70, 14H70 .

NJ's ORCID is 0000-0001-7504-4444.

¹Bielliptic surfaces are also known as hyperelliptic surfaces in earlier literature.

2. BIELLIPTIC QRT

In this section, we construct the bielliptic QRT map, show that it is symplectic and briefly describe how it reduces to the standard QRT map in the case when the elliptic curves Γ and Γ^* become degenerate.

Define an involution on the curve Γ by imposing the determinantal equation

$$\begin{vmatrix} 1 & x & X \\ 1 & \tilde{x} & \tilde{X} \\ \phi_1 & \phi_2 & \phi_3 \end{vmatrix} = (\tilde{x} - x)\phi_3 - (\tilde{X} - X)\phi_2 + (x\tilde{X} - \tilde{x}X)\phi_1 = 0, \quad (2.1)$$

where we assume that the vector $\phi = (\phi_1, \phi_2, \phi_3)$ is a constant (i.e., independent of (x, X)). The involution is given by the map

$$\iota_1 : (x, X) \mapsto (\tilde{x}, \tilde{X}), \quad (2.2)$$

where (x, X) denotes points on the Weierstrass curve Γ . For given $\phi = (\phi_1, \phi_2, \phi_3)$, we can solve eq. (2.1) for \tilde{x} , and substituting the result (in terms of x, X and \tilde{X}) into the curve relation for Γ we obtain a cubic for \tilde{X} , which has three solutions, one of which is $\tilde{X} = X$ (the trivial map) and the other is $\tilde{X} = \phi_3/\phi_1$, provided $(\phi_2/\phi_1, \phi_3/\phi_1)$ lies on the elliptic curve Γ as well. The condition $(\phi_2/\phi_1, \phi_3/\phi_1) \in \Gamma$ holds without loss of generality because neither the moduli g_2 and g_3 nor the corresponding uniformising variable are specified. The remaining solution of the cubic is the one we are interested in, and, as shown in the following lemma, it defines a single-valued involution ι_1 .

Lemma 2.1. *The map defined by (2.1) together with (1.1) is an involution, i.e., $\tilde{\tilde{\mathbf{X}}} = \mathbf{X}$.*

Proof. To prove the statement, apply the map ι_1 to the determinant on the left side of Equation (2.1). Then, in addition to (2.1), we obtain

$$\begin{vmatrix} 1 & \tilde{\tilde{x}} & \tilde{\tilde{X}} \\ 1 & \tilde{x} & \tilde{X} \\ \phi_1 & \phi_2 & \phi_3 \end{vmatrix} = 0, \quad (2.3)$$

given that the components of ϕ are constant. Subtracting (2.1) from this determinant, we see that a solution is given by $\tilde{\tilde{x}} = x$, $\tilde{\tilde{X}} = X$. However, there are other solutions. So, assuming that ϕ_1 does not identically vanish, and parametrising the curve in terms of Weierstrass elliptic functions, i.e. setting $(x, X) = (\wp(\xi), \wp'(\xi))$ for some dynamical variable ξ , and given (by the argument presented above) that ϕ_1, ϕ_2, ϕ_3 can be parametrised such that $(\frac{\phi_2}{\phi_1}, \frac{\phi_3}{\phi_1}) = (\wp(\delta), \wp'(\delta))$, we obtain from (2.1) the Frobenius-Stickelberger determinant, cf. [14],

$$\begin{vmatrix} 1 & \wp(\xi) & \wp'(\xi) \\ 1 & \wp(\tilde{\xi}) & \wp'(\tilde{\xi}) \\ 1 & \wp(\delta) & \wp'(\delta) \end{vmatrix} = 2 \frac{\sigma(\xi + \tilde{\xi} + \delta)}{\sigma^3(\xi) \sigma^3(\tilde{\xi}) \sigma^3(\delta)} \sigma(\tilde{\xi} - \xi) \sigma(\tilde{\xi} - \delta) \sigma(\xi - \delta), \quad (2.4)$$

and this vanishes if either $\tilde{\xi} = \xi$ (modulo the period lattice of the Weierstrass σ -function), which is a trivial solution, or if $\tilde{\xi} + \xi + \delta = 0 \pmod{\text{period lattice}}$. But the latter implies $\tilde{\tilde{\xi}} = \xi$, and hence the map $\mathbf{X} \rightarrow \tilde{\tilde{\mathbf{X}}}$ is an involution. \square

Remark 2.2. *Note that the proof of the lemma implies that*

$$\left(\frac{\phi_3}{\phi_1}\right)^2 = 4 \left(\frac{\phi_2}{\phi_1}\right)^3 - g_2 \left(\frac{\phi_2}{\phi_1}\right) - g_3. \quad (2.5)$$

Inspired by the QRT construction we now proceed by constructing an invariant for the map ι_1 . Introducing the vector $\mathbf{X} = (1, x, X)^T$, we can write (2.1) as

$$\phi \cdot (\mathbf{X} \times \widetilde{\mathbf{X}}) = 0 . \quad (2.6)$$

Assuming that ϕ depends on another set of variables y and Y such that the vector $\mathbf{Y} = (1, y, Y)^T$ is constant under the map: $\mathbf{Y} \rightarrow \widetilde{\mathbf{Y}} = \mathbf{Y}$, and taking ϕ of the form

$$\phi = \mathbf{A}_0 \mathbf{Y} \times \mathbf{A}_1 \mathbf{Y} , \quad (2.7)$$

we find that the ratio

$$I = \frac{\mathbf{X}^T \mathbf{A}_0 \mathbf{Y}}{\mathbf{X}^T \mathbf{A}_1 \mathbf{Y}} \quad (2.8)$$

is an invariant of the map $(x, X; y, Y) \rightarrow (\tilde{x}, \tilde{X}; \tilde{y}, \tilde{Y})$, i.e. $\tilde{I} = I$. This follows directly from inserting (2.7) into (2.6) leading to

$$(\mathbf{A}_0 \mathbf{Y} \times \mathbf{A}_1 \mathbf{Y}) \cdot (\mathbf{X} \times \widetilde{\mathbf{X}}) = (\mathbf{X}^T \mathbf{A}_0 \mathbf{Y})(\widetilde{\mathbf{X}}^T \mathbf{A}_1 \mathbf{Y}) - (\mathbf{X}^T \mathbf{A}_1 \mathbf{Y})(\widetilde{\mathbf{X}}^T \mathbf{A}_0 \mathbf{Y}) = 0,$$

and noting that $\widetilde{\mathbf{Y}} = \mathbf{Y}$. Importantly, we can allow for the moduli of the curve Γ to depend on the variables y and Y , i.e., $g_2 = g_2(\mathbf{Y})$ and $g_3 = g_3(\mathbf{Y})$.

Since the map ι_1 is an involution, i.e., a map of order 2, we need to extend it to create a map of infinite order. This is done by introducing a second involution ι_2 which does not commute with ι_1 , namely

$$\iota_2 : (y, Y) \mapsto (\hat{y}, \hat{Y}), \quad (2.9)$$

which leaves the variables x, X invariant, $\widehat{\mathbf{X}} = \mathbf{X}$, and such that

$$\begin{aligned} & \psi \cdot (\mathbf{Y} \times \widehat{\mathbf{Y}}) \\ &= \begin{vmatrix} 1 & y & Y \\ 1 & \hat{y} & \hat{Y} \\ \psi_1 & \psi_2 & \psi_3 \end{vmatrix} \\ &= (\hat{y} - y)\psi_3 - (\hat{Y} - Y)\psi_2 + (y\hat{Y} - \hat{y}Y)\psi_1 = 0 . \end{aligned} \quad (2.10)$$

This map shares the same invariant I of (2.8) provided we take the vector $\psi = (\psi_1, \psi_2, \psi_3)$ of the form

$$\psi = \mathbf{A}_0^T \mathbf{X} \times \mathbf{A}_1^T \mathbf{X} , \quad (2.11)$$

and where we assume that (y, Y) are points of the second Weierstrass curve Γ^* , with moduli $G_2 = G_2(\mathbf{X})$ and $G_3 = G_3(\mathbf{X})$ which may depend on the variables x, X . Similarly to (2.5) the functions $\psi_i(X)$ are subject to the curve relation on Γ^* , namely

$$\left(\frac{\psi_3}{\psi_1}\right)^2 = 4 \left(\frac{\psi_2}{\psi_1}\right)^3 - G_2 \left(\frac{\psi_2}{\psi_1}\right) - G_3. \quad (2.12)$$

The composition of the two non-commuting involutions $\iota_1 : (x, X; y, Y) \rightarrow (\tilde{x}, \tilde{X}; \tilde{y}, \tilde{Y})$ and $\iota_2 : (x, X; y, Y) \rightarrow (\hat{x}, \hat{X}; \hat{y}, \hat{Y})$, viewed as maps on the space of four variables $(x, X; y, Y)$, leads to a map $\mu = \iota_2 \circ \iota_1$. More explicitly we have the chain

$$(x, X; y, Y) \xrightarrow{\iota_1} (\tilde{x}, \tilde{X}; \tilde{y}, \tilde{Y}) \xrightarrow{\iota_2} (\hat{x}, \hat{X}; \hat{y}, \hat{Y}) \quad (2.13)$$

resulting in a combined map

$$\mu : (x, X; y, Y) \rightarrow (\bar{x}, \bar{X}; \bar{y}, \bar{Y}) \quad (2.14a)$$

given by

$$\left[\frac{1}{2}\phi_1(\bar{x} + x) - \phi_2\right] \frac{\bar{X} - X}{\bar{x} - x} = \frac{1}{2}\phi_1(\bar{X} + X) - \phi_3 , \quad (2.14b)$$

$$\left[\frac{1}{2} \bar{\psi}_1(\bar{y} + y) \right] - \bar{\psi}_2 \left] \frac{\bar{Y} - Y}{\bar{y} - y} = \frac{1}{2} \bar{\psi}_1(\bar{Y} + Y) - \bar{\psi}_3, \quad (2.14c)$$

subject to

$$X^2 = 4x^3 - g_2(Y)x - g_3(Y), \quad Y^2 = 4y^3 - G_2(X)y - G_3(X), \quad (2.14d)$$

and where $\phi = \phi(\mathbf{Y})$, and $\bar{\psi} = \psi(\bar{\mathbf{X}})$ given by (2.7) and (2.11) respectively.

The map μ defined by (2.14a) is what we will refer to as the bielliptic QRT map. This map is symplectic through the following argument (in contrast to the QRT map which in general position is only measure-preserving, cf. [12], also see [8, §6.3.3]). In fact, in terms of the uniformising variables on the curves Γ and Γ^* , the proof of measure-preservation is straightforward. Recalling that the proof of Lemma 2.1 gives the realization of the involution $\iota_1 : (\xi, \eta) \rightarrow (\tilde{\xi}, \tilde{\eta}) = (-\xi - \delta, \eta)$, we have

$$J_1 = \frac{\partial(\tilde{\xi}, \tilde{\eta})}{\partial(\xi, \eta)} = \begin{vmatrix} \frac{\partial \tilde{\xi}}{\partial \xi} & * \\ 0 & 1 \end{vmatrix} = -1.$$

Similarly, we have for $\iota_2 : (\xi, \eta) \rightarrow (\hat{\xi}, \hat{\eta}) = (\xi, -\eta - \varepsilon)$,

$$J_2 = \frac{\partial(\hat{\xi}, \hat{\eta})}{\partial(\xi, \eta)} = \begin{vmatrix} 1 & 0 \\ * & \frac{\partial \hat{\eta}}{\partial \eta} \end{vmatrix} = -1.$$

Hence, the Jacobian of the composition map $\iota_2 \circ \iota_1$ is given by $J = \tilde{J}_2 J_1 = 1$ which implies that in terms of the uniformising variables the map is not only measure-preserving, but actually symplectic. In terms of the variables on the bielliptic surface S , with affine coordinates $(x, X, y, Y) \in \mathbb{C}^4$, we define the symplectic form to be

$$\Omega = \frac{dx \wedge dy}{XY}.$$

As a consequence of the derivation above this symplectic form is preserved under both involutions ι_1 and ι_2 , i.e. $\tilde{\Omega} = \hat{\Omega} = \Omega$.

The following remark shows that the invariant of the bielliptic QRT map contains the the one of the standard QRT map, when the curves Γ and Γ^* become degenerate.

Remark 2.3. *Consider the following degeneration of the curve Γ (respectively Γ^*), which occurs when a pair of its branch points coalesce, with the other pair separating far apart.*

First, consider the change of variables $x = e_3 + u^2$, $X = U u$, which transforms Γ into a curve of the form

$$U^2 = 4(u^2 - \gamma_0^2)(u^2 - \gamma_1^2),$$

where $\gamma_0^2 = (e_1 - e_3)$ and $\gamma_1^2 = (e_2 - e_3)$. Assuming the dilation of one pair of branch points and coalescence of the remaining pair by taking $\gamma_0^2 = -1/\epsilon^2$, $\gamma_1^2 = \mu^2 \epsilon^2$, for sufficiently small ϵ , the resulting equation for (u, U) becomes

$$U^2 = \frac{4}{\epsilon^2} u^2 (1 + \mathcal{O}(\epsilon^2)),$$

i.e., $U \sim \pm 2u/\epsilon$. Putting these steps together leads to the transformation $x = e_3 + u^2$, $U = 2u^2/\epsilon$. The transformation of (y, Y) on Γ^ to (g, G) proceeds in a similar manner under the corresponding assumptions on $E_1 - E_3$ and $E_2 - E_3$.*

Substituting these transformations into the invariant I , while assuming $c_0 = c_0^\circ \epsilon$, $c_1 = c_1^\circ \epsilon$, $f_0 = f_0^\circ \epsilon$, $f_1 = f_1^\circ \epsilon$, $g_0 = g_0^\circ \epsilon$, $g_1 = g_1^\circ \epsilon$, $h_0 = h_0^\circ \epsilon$, $h_1 = h_1^\circ \epsilon$, and $i_0 = i_0^\circ \epsilon^2$, $i_1 = i_1^\circ \epsilon^2$, leads to a new invariant I° that is independent of ϵ as $\epsilon \rightarrow 0$. This invariant is biquadratic in two variables f and g and, so, we obtain the generic invariant of the standard QRT map.

3. CONSTANT MODULI CASE

The components of $\phi = (\phi_1, \phi_2, \phi_3)$ defined by equation (2.7) satisfy the cubic equation (2.5) and correspondingly the components of $\psi = (\psi_1, \psi_2, \psi_3)$ satisfy equation (2.12). These conditions place constraints on the entries of the matrices A_j . In this section, we provide the general solution of these constraints, under the condition that the moduli $g_2(\mathbf{Y}), g_3(\mathbf{Y})$ as well as $G_2(\mathbf{X}), G_3(\mathbf{X})$ are constants, i.e., independent of \mathbf{X} and \mathbf{Y} . Although the option that the moduli may depend in a nontrivial way on the variables of the map provides us with possibly additional freedom in constructing interesting dynamical systems, we will not consider this situation in the current paper, and restrict ourselves here to the constant moduli case.

The statement of finding the general solution for the latter case assumes that certain factors appearing in the denominators do not vanish. Their vanishing leads to 4 additional special solution branches, which are listed in Appendix A. In what follows we write the matrices A_0 and A_1 in terms of the following entries:

$$A_j = \begin{pmatrix} a_j & b_j & c_j \\ d_j & e_j & f_j \\ g_j & h_j & i_j \end{pmatrix}, \quad j = 0, 1.$$

(Note that the (3,1)-entry of each of A_0, A_1 is g_0 or g_1 respectively and should not be confused with g_2, g_3 in the definition of Γ) Taking the components of Equation (2.7) and substituting into (2.5), we get an equation in which we can use (1.2) to replace even powers of Y . The remaining equation is linear in Y . Since each coefficient in Y must vanish identically, we obtain two equations that are polynomial in y . The vanishing of each coefficient of a power of y leads to 18 polynomial equations for the entries of A_0 and A_1 .

The generic solution of these equations was found by using a variant of the primary decomposition algorithm in the MAGMA Computer Algebra System [4]. The algorithm computes a decomposition of the radical of the ideal generated by the polynomials (following the general approach of [6], combined with sparse interpolation techniques from [16]).

Each component of the decomposition gives a branch of the complete solution to the equations, and is described by a list of free parameters, a list of main variables v_1, \dots, v_l and a triangular system of polynomials in the v_i , so that the solutions for the branch can be obtained by setting the parameters to arbitrary values and then solving for the v_i in the triangular system by back-substitution, assuming the relevant denominators are non-zero.

The main branch of the solution is described with the free parameters $b_0, e_0, f_0, h_0, i_0, e_1, f_1, g_1, h_1, i_1, g_2, g_3, G_2, G_3$ and the main variables $a_0, c_0, d_0, g_0, a_1, b_1, c_1, d_1$. Write $D_0 = e_0 i_1 - h_0 f_1, D_1 = e_1 i_1 - f_1 h_1$. Then if $D_0 \neq 0, D_1 \neq 0, h_0 \neq 0, i_1 \neq 0$, the solution is described by the following triangular system in the main variables²:

$$a_0 = \frac{1}{D_0 D_1} \left((-e_0 i_0 h_1 + f_0 h_0 h_1) c_1 d_1 + (e_0 i_0 e_1 g_1 - f_0 h_0 e_1 g_1) c_1 \right. \\ \left. + (b_0 e_0 i_1^2 - b_0 f_0 h_1 i_1 - b_0 h_0 f_1 i_1 + b_0 i_0 f_1 h_1) d_1 \right. \\ \left. - b_0 e_0 f_1 g_1 i_1 + b_0 f_0 e_1 g_1 i_1 + b_0 h_0 f_1^2 g_1 - b_0 i_0 e_1 f_1 g_1 \right), \quad (3.1a)$$

$$c_0 = \frac{1}{D_0} \left((e_0 i_0 - f_0 h_0) c_1 + b_0 f_0 i_1 - b_0 i_0 f_1 \right), \quad (3.1b)$$

²In the list (3.1) the choice of the parameters for which we solve the constraint equations (2.5) and (2.12) are determined by the monomial ordering of the Groebner basis used in the algorithm.

$$d_0 = \frac{1}{D_1}((e_0 i_1 - f_0 h_1)d_1 - e_0 f_1 g_1 + f_0 e_1 g_1), \quad (3.1c)$$

$$g_0 = \frac{1}{D_1}((h_0 i_1 - i_0 h_1)d_1 - h_0 f_1 g_1 + i_0 e_1 g_1), \quad (3.1d)$$

$$a_1 = \frac{1}{D_0}((-h_0 d_1 + e_0 g_1)c_1 + b_0 i_1 d_1 - b_0 f_1 g_1), \quad (3.1e)$$

$$b_1 = \frac{1}{D_0}((e_0 h_1 - h_0 e_1)c_1 + b_0 e_1 i_1 - b_0 f_1 h_1), \quad (3.1f)$$

$$\begin{aligned} c_1^3 = \frac{1}{h_0^3} & \left((3b_0 h_0^2 i_1 + \frac{1}{4}e_0^3 i_1 - \frac{1}{4}e_0^2 h_0 f_1)c_1^2 \right. \\ & + (-3b_0^2 h_0 i_1^2 - \frac{1}{2}b_0 e_0^2 f_1 i_1 + \frac{1}{2}b_0 e_0 h_0 f_1^2 + \\ & \quad \frac{1}{4}e_0^2 h_0 i_1^2 g_2 - \frac{1}{2}e_0 h_0^2 f_1 i_1 g_2 + \frac{1}{4}h_0^3 f_1^2 g_2)c_1 \\ & + (b_0^3 i_1^3 + \frac{1}{4}b_0^2 e_0 f_1^2 i_1 - \frac{1}{4}b_0^2 h_0 f_1^3 - \frac{1}{4}b_0 e_0^2 i_1^3 g_2 + \\ & \quad \frac{1}{2}b_0 e_0 h_0 f_1 i_1^2 g_2 - \frac{1}{4}b_0 h_0^2 f_1^2 i_1 g_2 + \frac{1}{4}e_0^3 i_1^3 g_3 \\ & \quad \left. - \frac{3}{4}e_0^2 h_0 f_1 i_1^2 g_3 + \frac{3}{4}e_0 h_0^2 f_1^2 i_1 g_3 - \frac{1}{4}h_0^3 f_1^3 g_3) \right), \quad (3.1g) \end{aligned}$$

$$\begin{aligned} d_1^3 = \frac{1}{i_1^3} & \left((-\frac{1}{4}e_1 h_1^2 i_1 + 3f_1 g_1 i_1^2 + \frac{1}{4}f_1 h_1^3)d_1^2 \right. \\ & + (\frac{1}{2}e_1^2 g_1 h_1 i_1 + \frac{1}{4}e_1^2 i_1^3 G_2 - \frac{1}{2}e_1 f_1 g_1 h_1^2 \\ & \quad - \frac{1}{2}e_1 f_1 h_1 i_1^2 G_2 - 3f_1^2 g_1^2 i_1 + \frac{1}{4}f_1^2 h_1^2 i_1 G_2)d_1 \\ & + (-\frac{1}{4}e_1^3 g_1^2 i_1 - \frac{1}{4}e_1^3 i_1^3 G_3 + \frac{1}{4}e_1^2 f_1 g_1^2 h_1 - \frac{1}{4}e_1^2 f_1 g_1 i_1^2 G_2 \\ & \quad + \frac{3}{4}e_1^2 f_1 h_1 i_1^2 G_3 + \frac{1}{2}e_1 f_1^2 g_1 h_1 i_1 G_2 - \frac{3}{4}e_1 f_1^2 h_1^2 i_1 G_3 \\ & \quad \left. + f_1^3 g_1^3 - \frac{1}{4}f_1^3 g_1 h_1^2 G_2 + \frac{1}{4}f_1^3 h_1^3 G_3) \right). \quad (3.1h) \end{aligned}$$

Example 3.1. *The integer matrices*

$$A_0 = \begin{pmatrix} 2 & -16 & 2 \\ 3 & -2 & 3 \\ -2 & 5 & -2 \end{pmatrix}, \quad A_1 = \begin{pmatrix} -8 & -2 & -8 \\ 2 & 1 & 2 \\ 1 & 0 & 1 \end{pmatrix},$$

provide solutions of equations (3.1), under the conditions

$$2g_2 + g_3 = 16, \quad G_3 = -1.$$

There are 17 other branches of the solution for the cases where at least one of D_0 , D_1 , h_0 or i_0 is zero (see Appendix A).

4. GEOMETRY OF THE BIELLIPTIC QRT MAP

To make a connection to the standard terminology in the literature, we recall the construction of a bielliptic surface S of type I_a , as explained in [7]. Assuming that the period lattice of Γ^* is generated by $\{1, \tau\}$, we note that Γ^* has an automorphism of order 2, given by $\eta \mapsto -\eta$, or equivalently $Y \mapsto -Y$. The surface S is generated by quotienting $\Gamma \times \Gamma^*$ by the group of automorphisms generated by

$$\phi(\xi, \eta) = \left(\xi + \frac{\tau}{2}, -\eta \right).$$

While we do not make use of this construction explicitly, we rely on the fact that the resulting bielliptic surface S is a smooth algebraic surface.

In contrast, the geometric construction of the standard QRT map shows that it is a dynamical system on an elliptic surface [15, 5]. The bielliptic QRT map however can be considered as a higher dimensional generalization in 4D space of variables $(x, X; y, Y) \in \mathbb{P}^2 \times \mathbb{P}^2$ on the pencil

$$\{(x, X; y, Y) \mid \mathbf{X}^T \mathbf{A}_0 \mathbf{Y} - \lambda \mathbf{X}^T \mathbf{A}_1 \mathbf{Y} = 0\}_{\lambda \in \mathbb{P}^1} .$$

Rather than intersecting the corresponding surface with lines, as in [15], we intersect with the elliptic curves Γ and Γ^* . In fact, the pencil can be viewed as the solution space of a 10-parameter family of affine-linear equations:

$$Q_\lambda(x, X, y, Y; \mathbf{A}_0, \mathbf{A}_1) = 0 , \quad (4.1)$$

where the bielliptic QRT map comes from a reduction of the solution space to solutions coming from the intersection with the curves Γ and Γ^* .

Equation (4.1) resembles quadrilateral lattice equations (i.e., partial difference equations on a 2-dimensional grid), which are integrable in the sense that they satisfy the condition of consistency-around-the-cube (CAC) and were classified in [1]. This possible connection between the two subjects is tantalizing. In particular, we note that the most general equation in that class, known as Q4, has coefficients parametrised by an elliptic curve and possesses a class of N -soliton type solutions involving a pair of elliptic curves [3].

5. CONCLUSION

In this paper, we described an integrable, symplectic dynamical system on a bielliptic surface, which generalizes the QRT map. The situation we focused on is that where the moduli of the two curves Γ and Γ^* are constant (i.e., independent of the variables of the companion curve), in which case we find an 8-parameter family of two-dimensional symplectic maps possessing an algebraic invariant. The alternative option is to allow the moduli to depend on the companion variables, in which case we would have a more complicated system on a bielliptic surface. We will explore the latter situation in a future publication.

There are many remaining open questions. One concerns the special cases of matrices A_j , such as diagonal, symmetric, skew symmetric or upper/lower triangular matrices. These may be related to the classification of periodic fix points of the bielliptic QRT map, in the spirit of [15]. Another concerns the issue of what happens when the Weierstrass curves Γ and Γ^* are birationally transformed to elliptic curves in general position, and whether that would lift some of the conditions (3.1) on the matrices A_i . The geometric classification of bielliptic surfaces also raises interesting questions on how the bielliptic QRT map may be described geometrically.

ACKNOWLEDGMENTS

FWN is grateful for the support and hospitality of the Sydney Mathematical Research Institute (SMRI) during a visit where the current work was initiated. He was supported by the Foreign Expert Program of the Ministry of Sciences and Technology of China, grant number G2023013065L when this work was finalised.

APPENDIX A. SPECIAL BRANCHES OF SOLUTIONS

In this appendix, we list the special solutions that arise when one of i_1 , $e_1 i_1 - f_1 h_1$, h_0 , $e_0 i_1 - h_0 f_1$ vanishes.

A.1. Case 1: $i_1 = 0$.

A.1.1. *Case 1.1: $i_1 = 0, h_0 h_1 f_1 \neq 0$.*

$$\begin{aligned}
a_0 &= -\frac{(e_0 i_0 - f_0 h_0)}{h_0 f_1^2} c_1 d_1 - \frac{(-e_0 i_0 e_1 g_1 + f_0 h_0 e_1 g_1)}{h_0 f_1^2 h_1} c_1 \\
&\quad + \frac{b_0 i_0}{h_0 f_1} d_1 - \frac{(-b_0 h_0 f_1 g_1 + b_0 i_0 e_1 g_1)}{h_0 f_1 h_1}, \\
c_0 &= -\frac{(e_0 i_0 - f_0 h_0)}{h_0 f_1} c_1 + \frac{b_0 i_0}{h_0}, \\
d_0 &= \frac{f_0}{f_1} d_1 - \frac{(-e_0 f_1 g_1 + f_0 e_1 g_1)}{f_1 h_1}, \\
g_0 &= \frac{i_0}{f_1} d_1 - \frac{(-h_0 f_1 g_1 + i_0 e_1 g_1)}{f_1 h_1}, \\
a_1 &= \frac{1}{f_1} c_1 d_1 - \frac{e_0 g_1}{h_0 f_1} c_1 + \frac{b_0 g_1}{h_0}, \\
b_1 &= -\frac{(e_0 h_1 - h_0 e_1)}{h_0 f_1} c_1 + \frac{b_0 h_1}{h_0}, \\
c_1^3 &= -\frac{1}{4} \frac{e_0^2 f_1}{h_0^2} c_1^2 - \frac{(-\frac{1}{2} b_0 e_0 f_1^2 - \frac{1}{4} h_0^2 f_1^2 g_2)}{h_0^2} c_1 - \frac{(\frac{1}{4} b_0^2 f_1^3 + \frac{1}{4} h_0^2 f_1^3 g_3)}{h_0^2}, \\
d_1^2 &= \frac{2e_1 g_1}{h_1} d_1 - \frac{(e_1^2 g_1^2 h_1 + 4f_1^2 g_1^3 - f_1^2 g_1 h_1^2 G_2 + f_1^2 h_1^3 G_3)}{h_1^3}.
\end{aligned}$$

A.1.2. *Case 1.2: $i_1 = 0, h_0 = 0$, with $e_0 h_1 f_1 \neq 0$.*

$$\begin{aligned}
a_0 &= \frac{i_0}{f_1 h_1} b_1 d_1 - \frac{i_0 e_1 g_1}{f_1 h_1^2} b_1 + \frac{b_0 f_0 h_1 - b_0 i_0 e_1}{e_0 f_1 h_1} d_1 \\
&\quad + \frac{b_0 e_0 f_1 g_1 h_1 - b_0 f_0 e_1 g_1 h_1 + b_0 i_0 e_1^2 g_1}{e_0 f_1 h_1^2}, \\
c_0 &= \frac{i_0}{h_1} b_1 - \frac{b_0 f_0 h_1 - b_0 i_0 e_1}{e_0 h_1}, \\
d_0 &= \frac{f_0}{f_1} d_1 - \frac{-e_0 f_1 g_1 + f_0 e_1 g_1}{f_1 h_1}, \\
g_0 &= \frac{i_0}{f_1} d_1 - \frac{i_0 e_1 g_1}{f_1 h_1}, \\
a_1 &= \frac{g_1}{h_1} b_1 + \frac{b_0}{e_0} d_1 - \frac{b_0 e_1 g_1}{e_0 h_1}, \\
b_1^2 &= \frac{2b_0 e_1}{e_0} b_1 - \frac{4b_0^3 h_1^2 + b_0^2 e_0 e_1^2 - b_0 e_0^2 h_1^2 g_2 + e_0^3 h_1^2 g_3}{e_0^3}, \\
c_1 &= \frac{b_0 f_1}{e_0}, \\
d_1^2 &= \frac{2e_1 g_1}{h_1} d_1 - \frac{e_1^2 g_1^2 h_1 + 4f_1^2 g_1^3 - f_1^2 g_1 h_1^2 G_2 + f_1^2 h_1^3 G_3}{h_1^3}.
\end{aligned}$$

A.1.3. *Case 1.3: $i_1 = 0, h_0 = 0, f_1 = 0$, with $i_0 h_1 e_0 \neq 0$.*

$$\begin{aligned}
a_0 &= \frac{g_0 b_1}{h_1} + b_0 \frac{(f_0 h_1 - i_0 e_1)}{e_0 h_1 i_0} g_0 + \frac{b_0 g_1}{h_1}, \\
c_0 &= \frac{i_0 b_1}{h_1} + b_0 \frac{(f_0 h_1 - i_0 e_1)}{e_0 h_1}, \\
d_0 &= \frac{f_0 g_0}{i_0} + \frac{e_0 g_1}{h_1},
\end{aligned}$$

$$\begin{aligned}
g_0^2 &= -\frac{i_0^2}{h_1^3} \left(h_1^3 G_3 - h_1^2 g_1 G_2 + 4g_1^3 \right), \\
a_1 &= \frac{g_1 b_1}{h_1}, \\
b_1^2 &= \frac{2b_0 e_1}{e_0} b_1 - \frac{4b_0^3 h_1^2 + b_0^2 e_0 e_1^2 - b_0 e_0^2 h_1^2 g_2 + e_0^3 h_1^2 g_3}{e_0^3}, \\
c_1 &= 0, \\
d_1 &= \frac{e_1 g_1}{h_1}.
\end{aligned}$$

A.1.4. *Case 1.4.* $i_1 = 0, h_0 = 0, f_1 = 0, i_0 = 0$, with $h_1 e_0 \neq 0$

$$\begin{aligned}
a_0 &= \frac{b_0}{e_0} d_0, \\
c_0 &= \frac{b_0 f_0}{e_0}, \\
d_0^2 &= 2 \frac{e_0 g_1}{h_1} d_0 - \frac{e_0^2 h_1 g_1^2 + f_0^2 h_1^3 G_3 - f_0^2 h_1^2 g_1 G_2 + 4f_0^2 g_1^3}{h_1^3}, \\
g_0 &= 0, \\
a_1 &= \frac{g_1}{h_1} b_1, \\
b_1^2 &= 2 \frac{b_0 e_1}{e_0} b_1 - \frac{4b_0^3 h_1^2 + b_0^2 e_0 e_1^2 - b_0 e_0^2 h_1^2 g_2 + e_0^3 h_1^2 g_3}{e_0^3}, \\
c_1 &= 0, \\
d_1 &= \frac{e_1 g_1}{h_1}.
\end{aligned}$$

A.2. **Case 2:** $h_0 = 0$.

A.2.1. *Case 2.1:* $h_0 = 0, i_1 e_0 (e_1 i_1 - f_1 h_1) \neq 0$.

$$\begin{aligned}
a_0 &= -\frac{i_0 h_1 c_1 d_1}{e_1 i_1^2 - f_1 i_1 h_1} + \frac{i_0 e_1 g_1 c_1}{e_1 i_1^2 - f_1 i_1 h_1} - \frac{b_0 (-e_0 i_1^2 + f_0 i_1 h_1 - i_0 f_1 h_1)}{e_0 (e_1 i_1^2 - f_1 i_1 h_1)} d_1 \\
&\quad - \frac{b_0 g_1 (e_0 f_1 i_1 - f_0 e_1 i_1 + i_0 e_1 f_1)}{e_0 (e_1 i_1^2 - f_1 i_1 h_1)}, \\
c_0 &= \frac{i_0}{i_1} c_1 - \frac{-b_0 f_0 i_1 + b_0 i_0 f_1}{e_0 i_1}, \\
d_0 &= \frac{e_0 i_1 - f_0 h_1}{e_1 i_1 - f_1 h_1} d_1 - \frac{e_0 f_1 g_1 - f_0 e_1 g_1}{e_1 i_1 - f_1 h_1}, \\
g_0 &= -\frac{i_0 h_1}{e_1 i_1 - f_1 h_1} d_1 + \frac{i_0 e_1 g_1}{e_1 i_1 - f_1 h_1}, \\
a_1 &= \frac{g_1}{i_1} c_1 + \frac{b_0}{e_0} d_1 - \frac{b_0 f_1 g_1}{e_0 i_1}, \\
b_1 &= \frac{h_1}{i_1} c_1 - \frac{-b_0 e_1 i_1 + b_0 f_1 h_1}{e_0 i_1}, \\
c_1^2 &= \frac{2b_0 f_1}{e_0} c_1 - \frac{4b_0^3 i_1^2 + b_0^2 e_0 f_1^2 - b_0 e_0^2 i_1^2 g_2 + e_0^3 i_1^2 g_3}{e_0^3}, \\
d_1^3 &= -\frac{1}{i_1^3} \left(\frac{1}{4} (e_1 i_1 h_1^2 - 12 f_1 g_1 i_1^2 - f_1 h_1^3) d_1^2 \right. \\
&\quad \left. + \left(-\frac{1}{2} e_1^2 g_1 i_1 h_1 - \frac{1}{4} e_1^2 i_1^3 G_2 + \frac{1}{2} e_1 f_1 g_1 h_1^2 + \frac{1}{2} e_1 f_1 i_1^2 h_1 G_2 \right) \right)
\end{aligned}$$

$$\begin{aligned}
& + 3f_1^2g_1^2i_1 - \frac{1}{4}f_1^2i_1h_1^2G_2)d_1 \\
& + \left(\frac{1}{4}e_1^3g_1^2i_1 + \frac{1}{4}e_1^3i_1^3G_3 - \frac{1}{4}e_1^2f_1g_1^2h_1 + \frac{1}{4}e_1^2f_1g_1i_1^2G_2 \right. \\
& \quad - \frac{3}{4}e_1^2f_1i_1^2h_1G_3 - \frac{1}{2}e_1f_1^2g_1i_1h_1G_2 \\
& \quad \left. + \frac{3}{4}e_1f_1^2i_1h_1^2G_3 - f_1^3g_1^3 + \frac{1}{4}f_1^3g_1h_1^2G_2 - \frac{1}{4}f_1^3h_1^3G_3 \right).
\end{aligned}$$

A.2.2. *Case 2.2:* $h_0 = 0$, $(e_1i_1 - f_1h_1) = 0$, $i_1h_1i_0e_0 \neq 0$.

$$\begin{aligned}
a_0 &= \frac{g_0c_1}{i_1} - \frac{b_0(e_0i_1^2 - f_0h_1i_1 + i_0f_1h_1)}{e_0i_0h_1i_1}g_0 + \frac{b_0g_1}{h_1}, \\
c_0 &= \frac{i_0}{i_1}c_1 + \frac{b_0(f_0i_1 - i_0f_1)}{e_0i_1}, \\
d_0 &= -\frac{e_0i_1 - f_0h_1}{i_0h_1}g_0 + \frac{e_0g_1}{h_1}, \\
g_0^3 &= \frac{i_0(3g_1i_1^2 + \frac{1}{4}h_1^3)}{i_1^3}g_0^2 - \frac{i_0^2(3g_1^2 - \frac{1}{4}h_1^2G_2)}{i_1^2}g_0 \\
& \quad + \frac{i_0^3(g_1^3 - \frac{1}{4}g_1h_1^2G_2 + \frac{1}{4}h_1^3G_3)}{i_1^3}, \\
a_1 &= \frac{g_1}{i_1}c_1, \quad b_1 = \frac{h_1}{i_1}c_1, \\
c_1^2 &= \frac{2b_0f_1}{e_0}c_1 - \frac{4b_0^3i_1^2 + b_0^2e_0f_1^2 - b_0e_0^2i_1^2g_2 + e_0^3i_1^2g_3}{e_0^3}, \\
d_1 &= \frac{f_1g_1}{i_1}, \quad e_1 = \frac{f_1h_1}{i_1}.
\end{aligned}$$

A.2.3. *Case 2.3:* $h_0 = 0$, $(e_1i_1 - f_1h_1) = 0$, $h_1 = 0$, $i_1e_0 \neq 0$.

$$\begin{aligned}
a_0 &= \frac{b_0}{e_0}d_0 + \frac{i_0g_1}{i_1^2}c_1 - \frac{b_0i_0f_1g_1}{e_0i_1^2}, \\
c_0 &= \frac{i_0}{i_1}c_1 + \frac{b_0(f_0i_1 - i_0f_1)}{e_0i_1}, \\
d_0^3 &= 3\frac{f_0g_1}{i_1}d_0^2 + \frac{\frac{1}{4}e_0^2i_1^2G_2 - 3f_0^2g_1^2}{i_1^2}d_0 - \frac{1}{4}\frac{e_0^3g_1^2i_1 + e_0^3i_1^3G_3 + e_0^2f_0g_1i_1^2G_2 - 4f_0^3g_1^3}{i_1^3}, \\
g_0 &= \frac{i_0g_1}{i_1}, \quad a_1 = \frac{g_1}{i_1}c_1, \quad b_1 = 0, \\
c_1^2 &= \frac{2b_0f_1}{e_0}c_1 - \frac{4b_0^3i_1^2 + b_0^2e_0f_1^2 - b_0e_0^2i_1^2g_2 + e_0^3i_1^2g_3}{e_0^3}, \quad d_1 = \frac{f_1g_1}{i_1}, e_1 = 0.
\end{aligned}$$

A.2.4. *Case 2.4:* $h_0 = 0$, $(e_1i_1 - f_1h_1) = 0$, $i_0 = 0$, $i_1e_0 \neq 0$.

$$\begin{aligned}
a_0 &= \frac{b_0}{e_0}d_0, \quad c_0 = \frac{b_0f_0}{e_0}, \\
d_0^3 &= -\frac{1}{i_1^3}\left(-\frac{1}{4}(e_0h_1^2i_1 - 12f_0g_1i_1^2 - f_0h_1^3)d_0^2 \right. \\
& \quad - \left(-\frac{1}{2}e_0^2g_1h_1i_1 - \frac{1}{4}e_0^2i_1^3G_2 + \frac{1}{2}e_0f_0g_1h_1^2 + \frac{1}{2}e_0f_0h_1i_1^2G_2 + 3f_0^2g_1^2i_1 - \frac{1}{4}f_0^2h_1^2i_1G_2 \right)d_0 \\
& \quad \left. - \left(\frac{1}{4}e_0^3g_1^2i_1 + \frac{1}{4}e_0^3i_1^3G_3 - \frac{1}{4}e_0^2f_0g_1^2h_1 + \frac{1}{4}e_0^2f_0g_1i_1^2G_2 - \frac{3}{4}e_0^2f_0h_1i_1^2G_3 \right) \right)
\end{aligned}$$

$$\left. -\frac{1}{2}e_0f_0^2g_1h_1i_1G_2 + \frac{3}{4}e_0f_0^2h_1^2i_1G_3 - f_0^3g_1^3 + \frac{1}{4}f_0^3g_1h_1^2G_2 - \frac{1}{4}f_0^3h_1^3G_3 \right),$$

$$g_0 = 0, \quad a_1 = \frac{g_1}{i_1}c_1, \quad b_1 = \frac{h_1}{i_1}c_1,$$

$$c_1^2 = \frac{2b_0f_1}{e_0}c_1 - \frac{4b_0^3i_1^2 + b_0^2e_0f_1^2 - b_0e_0^2i_1^2g_2 + e_0^3i_1^2g_3}{e_0^3},$$

$$d_1 = \frac{f_1g_1}{i_1}, \quad e_1 = \frac{f_1h_1}{i_1}.$$

A.3. **Case** $e_0i_1 - h_0f_1 = 0$.

A.3.1. $e_0i_1 - h_0f_1 = 0$, $i_1(e_1i_1 - f_1h_1)h_0 \neq 0$.

$$\begin{aligned} a_0 = & -\frac{(f_0i_1 - i_0f_1)}{e_1^2i_1^2 - 2e_1f_1h_1i_1 + f_1^2h_1^2}b_1d_1h_1 + \frac{f_0i_1 - i_0f_1}{e_1^2i_1^2 - 2e_1f_1h_1i_1 + f_1^2h_1^2}b_1e_1g_1 \\ & - \frac{-b_0f_0h_1^2i_1 - b_0h_0e_1i_1^2 + b_0h_0f_1h_1i_1 + b_0i_0e_1h_1i_1}{h_0(e_1^2i_1^2 - 2e_1f_1h_1i_1 + f_1^2h_1^2)}d_1 \\ & - \frac{b_0f_0e_1g_1h_1i_1 + b_0h_0e_1f_1g_1i_1 - b_0h_0f_1^2g_1h_1 - b_0i_0e_1^2g_1i_1}{h_0(e_1^2i_1^2 - 2e_1f_1h_1i_1 + f_1^2h_1^2)}, \end{aligned}$$

$$c_0 = \frac{f_0i_1 - i_0f_1}{e_1i_1 - f_1h_1}b_1 - \frac{b_0f_0h_1i_1 - b_0i_0e_1i_1}{h_0(e_1i_1 - f_1h_1)},$$

$$d_0 = -\frac{f_0h_1 - h_0f_1}{e_1i_1 - f_1h_1}d_1 - \frac{-g_1(f_0e_1i_1 + h_0f_1^2)}{i_1(e_1i_1 - f_1h_1)},$$

$$e_0 = \frac{h_0f_1}{i_1},$$

$$g_0 = \frac{h_0i_1 - i_0h_1}{e_1i_1 - f_1h_1}d_1 + \frac{h_0f_1g_1 - i_0e_1g_1}{e_1i_1 - f_1h_1},$$

$$a_1 = \frac{i_1}{e_1i_1 - f_1h_1}b_1d_1 - \frac{f_1g_1}{e_1i_1 - f_1h_1}b_1 - \frac{b_0h_1i_1}{h_0(e_1i_1 - f_1h_1)}d_1 + \frac{b_0e_1g_1i_1}{h_0(e_1i_1 - f_1h_1)},$$

$$\begin{aligned} b_1^3 = & \frac{1}{h_0^3i_1^3}(3b_0h_1i_1^3 - \frac{1}{4}h_0e_1f_1^2i_1 + \frac{1}{4}h_0f_1^3h_1)b_1^2 \\ & - \frac{1}{h_0^2i_1^2}(3b_0^2h_1^2i_1^2 - \frac{1}{2}b_0h_0e_1^2f_1i_1 + \frac{1}{2}b_0h_0e_1f_1^2h_1 - \frac{1}{4}h_0^2e_1^2i_1^2g_2 \\ & \quad + \frac{1}{2}h_0^2e_1f_1h_1i_1g_2 - \frac{1}{4}h_0^2f_1^2h_1^2g_2)b_1 \\ & - \frac{1}{h_0^3i_1^3}(-b_0^3h_1^3i_1^3 + \frac{1}{4}b_0^2h_0e_1^3i_1^3 - \frac{1}{4}b_0^2h_0e_1^2f_1h_1i_1^2 \\ & \quad + \frac{1}{4}b_0h_0^2e_1^2h_1i_1^3g_2 - \frac{1}{2}b_0h_0^2e_1f_1h_1^2i_1^2g_2 + \frac{1}{4}b_0h_0^2f_1^2h_1^3i_1g_2 \\ & \quad + \frac{1}{4}h_0^3e_1^3i_1^3g_3 - \frac{3}{4}h_0^3e_1^2f_1h_1i_1^2g_3 + \frac{3}{4}h_0^3e_1f_1^2h_1^2i_1g_3 - \frac{1}{4}h_0^3f_1^3h_1^3g_3), \end{aligned}$$

$$c_1 = \frac{b_0i_1}{h_0},$$

$$\begin{aligned} d_1^3 = & -\frac{1}{i_1^3}\left(\frac{1}{4}(e_1h_1^2i_1 - 12f_1g_1i_1^2 - f_1h_1^3)d_1^2 \right. \\ & \quad + (-\frac{1}{2}e_1^2g_1h_1i_1 - \frac{1}{4}e_1^2i_1^3G_2 + \frac{1}{2}e_1f_1g_1h_1^2 + \frac{1}{2}e_1f_1h_1i_1^2G_2 + 3f_1^2g_1^2i_1 - \frac{1}{4}f_1^2h_1^2i_1G_2)d_1 \\ & \quad \left. + (\frac{1}{4}e_1^3g_1^2i_1 + \frac{1}{4}e_1^3i_1^3G_3 - \frac{1}{4}e_1^2f_1g_1^2h_1 + \frac{1}{4}e_1^2f_1g_1i_1^2G_2 - \frac{3}{4}e_1^2f_1h_1i_1^2G_3 \right) \end{aligned}$$

$$\left. - \frac{1}{2}e_1f_1^2g_1h_1i_1G_2 + \frac{3}{4}e_1f_1^2h_1^2i_1G_3 - f_1^3g_1^3 + \frac{1}{4}f_1^3g_1h_1^2G_2 - \frac{1}{4}f_1^3h_1^3G_3 \right).$$

A.3.2. $e_0i_1 - h_0f_1 = 0$, $e_1i_1 - f_1h_1 = 0$, $i_1h_0(h_0i_1 - i_0h_1) \neq 0$.

$$\begin{aligned} a_0 &= -\frac{(f_0h_1i_1 - i_0f_1h_1)}{(e_1i_1 - f_1h_1)^2}b_1d_1 - \frac{(-f_0e_1g_1i_1 + i_0e_1f_1g_1)}{(e_1i_1 - f_1h_1)^2}b_1 \\ &\quad - \frac{(-b_0f_0h_1^2i_1 - b_0h_0e_1i_1^2 + b_0h_0f_1h_1i_1 + b_0i_0e_1h_1i_1)}{h_0(e_1i_1 - f_1h_1)^2}d_1 \\ &\quad - \frac{(b_0f_0e_1g_1h_1i_1 + b_0h_0e_1f_1g_1i_1 - b_0h_0f_1^2g_1h_1 - b_0i_0e_1^2g_1i_1)}{h_0(e_1i_1 - f_1h_1)^2}, \\ c_0 &= \frac{(f_0i_1 - i_0f_1)}{e_1i_1 - f_1h_1}b_1 - \frac{(b_0f_0h_1i_1 - b_0i_0e_1i_1)}{h_0(e_1i_1 - f_1h_1)}, \\ d_0 &= -\frac{(f_0h_1 - h_0f_1)}{e_1i_1 - f_1h_1}d_1 + \frac{g_1(f_0e_1i_1 - h_0f_1^2)}{i_1(e_1i_1 - f_1h_1)}, \\ e_0 &= \frac{h_0f_1}{i_1}, \\ g_0 &= \frac{(h_0i_1 - i_0h_1)}{e_1i_1 - f_1h_1}d_1 + \frac{(h_0f_1g_1 - i_0e_1g_1)}{e_1i_1 - f_1h_1}, \\ a_1 &= \frac{i_1}{e_1i_1 - f_1h_1}b_1d_1 - \frac{f_1g_1}{e_1i_1 - f_1h_1}b_1 - \frac{b_0h_1i_1}{h_0(e_1i_1 - f_1h_1)}d_1 + \frac{b_0e_1g_1i_1}{h_0(e_1i_1 - f_1h_1)}, \\ b_1^3 &= \frac{3b_0h_1i_1^3 - \frac{1}{4}h_0e_1f_1^2i_1 + \frac{1}{4}h_0f_1^3h_1}{h_0i_1^3}b_1^2 \\ &\quad - \frac{3b_0^2h_1^2i_1^2 - \frac{1}{2}b_0h_0e_1^2f_1i_1 + \frac{1}{2}b_0h_0e_1f_1^2h_1 - \frac{1}{4}h_0^2e_1^2i_1^2g_2 + \frac{1}{2}h_0^2e_1f_1h_1i_1g_2 - \frac{1}{4}h_0^2f_1^2h_1^2g_2}{h_0^2i_1^2}b_1 \\ &\quad - \frac{1}{h_0^3i_1^3} \left(-b_0^3h_1^3i_1^3 + \frac{1}{4}b_0^2h_0e_1^3i_1^3 - \frac{1}{4}b_0^2h_0e_1^2f_1h_1i_1^2 + \frac{1}{4}b_0h_0^2e_1^2h_1i_1^3g_2 \right. \\ &\quad \left. - \frac{1}{2}b_0h_0^2e_1f_1h_1^2i_1^2g_2 + \frac{1}{4}b_0h_0^2f_1^2h_1^3i_1g_2 + \frac{1}{4}h_0^3e_1^3i_1^3g_3 \right. \\ &\quad \left. - \frac{3}{4}h_0^3e_1^2f_1h_1i_1^2g_3 + \frac{3}{4}h_0^3e_1f_1^2h_1^2i_1g_3 - \frac{1}{4}h_0^3f_1^3h_1^3g_3 \right), \\ c_1 &= \frac{b_0i_1}{h_0}, \\ d_1^3 &= \frac{1}{i_1^3} \left(-\frac{1}{4}(e_1h_1^2i_1 - 12f_1g_1i_1^2 - \frac{1}{4}f_1h_1^3)d_1^2 \right. \\ &\quad \left. - \left(-\frac{1}{2}e_1^2g_1h_1i_1 - \frac{1}{4}e_1^2i_1^3G_2 + \frac{1}{2}e_1f_1g_1h_1^2 + \frac{1}{2}e_1f_1h_1i_1^2G_2 + 3f_1^2g_1^2i_1 - \frac{1}{4}f_1^2h_1^2i_1G_2 \right) d_1 \right. \\ &\quad \left. - \left(\frac{1}{4}e_1^3g_1^2i_1 + \frac{1}{4}e_1^3i_1^3G_3 - \frac{1}{4}e_1^2f_1g_1^2h_1 + \frac{1}{4}e_1^2f_1g_1i_1^2G_2 - \frac{3}{4}e_1^2f_1h_1i_1^2G_3 \right. \right. \\ &\quad \left. \left. - \frac{1}{2}e_1f_1^2g_1h_1i_1G_2 + \frac{3}{4}e_1f_1^2h_1^2i_1G_3 - f_1^3g_1^3 + \frac{1}{4}f_1^3g_1h_1^2G_2 - \frac{1}{4}f_1^3h_1^3G_3 \right) \right). \end{aligned}$$

A.3.3. $e_0i_1 - h_0f_1 = 0$, $e_1i_1 - f_1h_1 = 0$, $h_0i_1 - i_0h_1 = 0$, $i_1h_1i_0(f_0i_1 - i_0f_1) \neq 0$.

$$\begin{aligned} a_0 &= -\frac{(f_0h_1i_1 - i_0f_1h_1)}{e_1^2i_1^2 - 2e_1f_1h_1i_1 + f_1^2h_1^2}b_1d_1 - \frac{-f_0e_1g_1i_1 + i_0e_1f_1g_1}{e_1^2i_1^2 - 2e_1f_1h_1i_1 + f_1^2h_1^2}b_1 \\ &\quad - \frac{-b_0f_0h_1^2i_1 - b_0h_0e_1i_1^2 + b_0h_0f_1h_1i_1 + b_0i_0e_1h_1i_1}{h_0e_1^2i_1^2 - 2h_0e_1f_1h_1i_1 + h_0f_1^2h_1^2}d_1 \end{aligned}$$

$$\begin{aligned}
& - \frac{b_0 f_0 e_1 g_1 h_1 i_1 + b_0 h_0 e_1 f_1 g_1 i_1 - b_0 h_0 f_1^2 g_1 h_1 - b_0 i_0 e_1^2 g_1 i_1}{h_0 e_1^2 i_1^2 - 2 h_0 e_1 f_1 h_1 i_1 + h_0 f_1^2 h_1^2}, \\
c_0 &= \frac{f_0 i_1 - i_0 f_1}{e_1 i_1 - f_1 h_1} b_1 - \frac{b_0 f_0 h_1 i_1 - b_0 i_0 e_1 i_1}{h_0 e_1 i_1 - h_0 f_1 h_1}, \\
d_0 &= - \frac{(f_0 h_1 - h_0 f_1)}{e_1 i_1 - f_1 h_1} d_1 - \frac{-f_0 e_1 g_1 i_1 + h_0 f_1^2 g_1}{e_1 i_1^2 - f_1 h_1 i_1}, \\
e_0 &= \frac{h_0 f_1}{i_1}, \\
g_0 &= \frac{h_0 i_1 - i_0 h_1}{e_1 i_1 - f_1 h_1} d_1 - \frac{h_0 f_1 g_1 - i_0 e_1 g_1}{e_1 i_1 - f_1 h_1}, \\
a_1 &= \frac{i_1}{e_1 i_1 - f_1 h_1} b_1 d_1 - \frac{f_1 g_1}{e_1 i_1 - f_1 h_1} b_1 - \frac{b_0 h_1 i_1}{h_0 e_1 i_1 - h_0 f_1 h_1} d_1 + \frac{b_0 e_1 g_1 i_1}{h_0 e_1 i_1 - h_0 f_1 h_1}, \\
b_1^3 &= \frac{b_1^2}{h_0 i_1^3} (3 b_0 h_1 i_1^3 - \frac{1}{4} h_0 e_1 f_1^2 i_1 + \frac{1}{4} h_0 f_1^3 h_1) \\
& - \frac{b_1}{h_0 i_1^2} (3 b_0^2 h_1^2 i_1^2 - \frac{1}{2} b_0 h_0 e_1^2 f_1 i_1 + \frac{1}{2} b_0 h_0 e_1 f_1^2 h_1 - \frac{1}{4} h_0^2 e_1^2 i_1^2 g_2 + \frac{1}{2} h_0^2 e_1 f_1 h_1 i_1 g_2 - \frac{1}{4} h_0^2 f_1^2 h_1^2 g_2) \\
& - \frac{1}{h_0^3 i_1^3} (-b_0^3 h_1^3 i_1^3 + \frac{1}{4} b_0^2 h_0 e_1^3 i_1^3 - \frac{1}{4} b_0^2 h_0 e_1^2 f_1 h_1 i_1^2 + \frac{1}{4} b_0 h_0^2 e_1^2 h_1 i_1^3 g_2 - \frac{1}{2} b_0 h_0^2 e_1 f_1 h_1^2 i_1^2 g_2 \\
& + \frac{1}{4} b_0 h_0^2 f_1^2 h_1^3 i_1 g_2 + \frac{1}{4} h_0^3 e_1^3 i_1^3 g_3 - \frac{3}{4} h_0^3 e_1^2 f_1 h_1 i_1^2 g_3 + \frac{3}{4} h_0^3 e_1 f_1^2 h_1^2 i_1 g_3 - \frac{1}{4} h_0^3 f_1^3 h_1^3 g_3), \\
c_1 &= \frac{b_0 i_1}{h_0}, \\
d_1^3 &= - \frac{d_1^2}{4 i_1^3} (e_1 h_1^2 i_1 - 12 f_1 g_1 i_1^2 - \frac{1}{4} f_1 h_1^3) \\
& - \frac{d_1}{i_1^3} (-\frac{1}{2} e_1^2 g_1 h_1 i_1 - \frac{1}{4} e_1^2 i_1^3 G_2 + \frac{1}{2} e_1 f_1 g_1 h_1^2 + \frac{1}{2} e_1 f_1 h_1 i_1^2 G_2 + 3 f_1^2 g_1^2 i_1 - \frac{1}{4} f_1^2 h_1^2 i_1 G_2) \\
& - \frac{1}{i_1^3} (\frac{1}{4} e_1^3 g_1^2 i_1 + \frac{1}{4} e_1^3 i_1^3 G_3 - \frac{1}{4} e_1^2 f_1 g_1^2 h_1 + \frac{1}{4} e_1^2 f_1 g_1 i_1^2 G_2 - \frac{3}{4} e_1^2 f_1 h_1 i_1^2 G_3 - \frac{1}{2} e_1 f_1^2 g_1 h_1 i_1 G_2 \\
& + \frac{3}{4} e_1 f_1^2 h_1^2 i_1 G_3 - f_1^3 g_1^3 + \frac{1}{4} f_1^3 g_1 h_1^2 G_2 - \frac{1}{4} f_1^3 h_1^3 G_3).
\end{aligned}$$

A.3.4. $e_0 i_1 - h_0 f_1 = 0$, $e_1 i_1 - f_1 h_1 = 0$, $h_0 i_1 - i_0 h_1 = 0$, $f_0 i_1 - i_0 f_1 = 0$. This is a degenerate branch resulting in a set of cross relations.

$$\begin{array}{lll}
f_0 h_0 - e_0 i_0 = 0, & b_0 a_1 - a_0 b_1 = 0, & c_0 a_1 - a_0 c_1 = 0, \\
d_0 a_1 - a_0 d_1 = 0, & e_0 a_1 - a_0 e_1 = 0, & f_0 a_1 - a_0 f_1 = 0, \\
g_0 a_1 - a_0 g_1 = 0, & h_0 a_1 - a_0 h_1 = 0, & i_0 a_1 - a_0 i_1 = 0, \\
c_0 b_1 - b_0 c_1 = 0, & d_0 b_1 - b_0 d_1 = 0, & e_0 b_1 - b_0 e_1 = 0, \\
f_0 b_1 - b_0 f_1 = 0, & g_0 b_1 - b_0 g_1 = 0, & h_0 b_1 - b_0 h_1 = 0, \\
i_0 b_1 - b_0 i_1 = 0, & d_0 c_1 - c_0 d_1 = 0, & e_0 c_1 - c_0 e_1 = 0, \\
f_0 c_1 - c_0 f_1 = 0, & g_0 c_1 - c_0 g_1 = 0, & h_0 c_1 - c_0 h_1 = 0, \\
i_0 c_1 - c_0 i_1 = 0, & e_0 d_1 - d_0 e_1 = 0, & f_0 d_1 - d_0 f_1 = 0, \\
g_0 d_1 - d_0 g_1 = 0, & h_0 d_1 - d_0 h_1 = 0, & i_0 d_1 - d_0 i_1 = 0, \\
f_0 e_1 - e_0 f_1 = 0, & g_0 e_1 - e_0 g_1 = 0, & h_0 e_1 - e_0 h_1 = 0, \\
i_0 e_1 - e_0 i_1 = 0, & g_0 f_1 - f_0 g_1 = 0, & h_0 f_1 - e_0 i_1 = 0, \\
i_0 f_1 - f_0 i_1 = 0, & h_0 g_1 - g_0 h_1 = 0, & i_0 g_1 - g_0 i_1 = 0, \\
f_0 h_1 - e_0 i_1 = 0, & i_0 h_1 - h_0 i_1 = 0, & f_1 h_1 - e_1 i_1 = 0.
\end{array}$$

A.4. Case $e_1 i_1 - f_1 h_1 = 0$.A.4.1. $e_1 i_1 - f_1 h_1 = 0$, $i_1 h_0 (h_0 i_1 - i_0 h_1) (e_0 i_1 - h_0 f_1) \neq 0$.

$$\begin{aligned}
a_0 &= -\frac{(e_0 i_0 h_1 - f_0 h_0 h_1)}{e_0 h_0 i_1^2 - e_0 i_0 h_1 i_1 - h_0^2 f_1 i_1 + h_0 i_0 f_1 h_1} g_0 c_1 \\
&\quad - \frac{-b_0 e_0 i_1^2 + b_0 f_0 h_1 i_1 + b_0 h_0 f_1 i_1 - b_0 i_0 f_1 h_1}{e_0 h_0 i_1^2 - e_0 i_0 h_1 i_1 - h_0^2 f_1 i_1 + h_0 i_0 f_1 h_1} g_0 \\
&\quad - \frac{-e_0 h_0 i_0 g_1 + f_0 h_0^2 g_1}{e_0 h_0 i_1^2 - e_0 i_0 h_1 i_1 - h_0^2 f_1 i_1 + h_0 i_0 f_1 h_1} c_1 \\
&\quad - \frac{b_0 e_0 i_0 g_1 i_1 - b_0 f_0 h_0 g_1 i_1}{e_0 h_0 i_1^2 - e_0 i_0 h_1 i_1 - h_0^2 f_1 i_1 + h_0 i_0 f_1 h_1}, \\
c_0 &= \frac{e_0 i_0 - f_0 h_0}{e_0 i_1 - h_0 f_1} c_1 - \frac{-b_0 f_0 i_1 + b_0 i_0 f_1}{e_0 i_1 - h_0 f_1}, \\
d_0 &= \frac{e_0 i_1 - f_0 h_1}{h_0 i_1 - i_0 h_1} g_0 - \frac{e_0 i_0 g_1 - f_0 h_0 g_1}{h_0 i_1 - i_0 h_1}, \\
g_0^3 &= -\frac{g_0^2}{4i_1^3} (h_0 h_1^2 i_1 - 12i_0 g_1 i_1^2 - i_0 h_1^3) \\
&\quad - \frac{g_0}{i_1^3} \left(-\frac{1}{2} h_0^2 g_1 h_1 i_1 - \frac{1}{4} h_0^2 i_1^3 G_2 + \frac{1}{2} h_0 i_0 g_1 h_1^2 + \frac{1}{2} h_0 i_0 h_1 i_1^2 G_2 + 3i_0^2 g_1^2 i_1 - \frac{1}{4} i_0^2 h_1^2 i_1 G_2 \right) \\
&\quad - \frac{1}{i_1^3} \left(\frac{1}{4} h_0^3 g_1^2 i_1 + \frac{1}{4} h_0^3 i_1^3 G_3 - \frac{1}{4} h_0^2 i_0 g_1^2 h_1 + \frac{1}{4} h_0^2 i_0 g_1 i_1^2 G_2 - \frac{3}{4} h_0^2 i_0 h_1 i_1^2 G_3 - \frac{1}{2} h_0 i_0^2 g_1 h_1 i_1 G_2 \right. \\
&\quad \left. + \frac{3}{4} h_0 i_0^2 h_1^2 i_1 G_3 - i_0^3 g_1^3 + \frac{1}{4} i_0^3 g_1 h_1^2 G_2 - \frac{1}{4} i_0^3 h_1^3 G_3 \right), \\
a_1 &= \frac{g_1}{i_1} c_1, \quad b_1 = \frac{h_1}{i_1} c_1, \\
c_1^3 &= \frac{c_1^2}{h_0^3} \left(3b_0 h_0^2 i_1 + \frac{1}{4} e_0^3 i_1 - \frac{1}{4} e_0^2 h_0 f_1 \right) \\
&\quad - \frac{c_1}{h_0^3} \left(3b_0^2 h_0 i_1^2 + \frac{1}{2} b_0 e_0^2 f_1 i_1 - \frac{1}{2} b_0 e_0 h_0 f_1^2 - \frac{1}{4} e_0^2 h_0 i_1^2 g_2 + \frac{1}{2} e_0 h_0^2 f_1 i_1 g_2 - \frac{1}{4} h_0^3 f_1^2 g_2 \right) \\
&\quad - \frac{1}{h_0^3} \left(-b_0^3 i_1^3 - \frac{1}{4} b_0^2 e_0 f_1^2 i_1 + \frac{1}{4} b_0^2 h_0 f_1^3 + \frac{1}{4} b_0 e_0^2 i_1^3 g_2 - \frac{1}{2} b_0 e_0 h_0 f_1 i_1^2 g_2 + \frac{1}{4} b_0 h_0^2 f_1^2 i_1 g_2 \right. \\
&\quad \left. - \frac{1}{4} e_0^3 i_1^3 g_3 + \frac{3}{4} e_0^2 h_0 f_1 i_1^2 g_3 - \frac{3}{4} e_0 h_0^2 f_1^2 i_1 g_3 + \frac{1}{4} h_0^3 f_1^3 g_3 \right), \\
d_1 &= \frac{f_1 g_1}{i_1}, \quad e_1 = \frac{f_1 h_1}{i_1}.
\end{aligned}$$

A.4.2. $e_1 i_1 - f_1 h_1 = 0$, $h_0 i_1 - i_0 h_1 = 0$, $i_1 h_1 i_0 (e_0 i_1^2 - i_0 f_1 h_1) \neq 0$.

$$\begin{aligned}
a_0 &= \frac{1}{e_0 i_1^2 - i_0 f_1 h_1} (i_0 h_1 d_0 c_1 + b_0 i_1^2 d_0 + e_0 i_0 g_1 c_1 - b_0 i_0 f_1 g_1), \\
c_0 &= \frac{1}{e_0 i_1^2 - i_0 f_1 h_1} ((e_0 i_0 i_1 - f_0 i_0 h_1) c_1 - (-b_0 f_0 i_1^2 + b_0 i_0 f_1 i_1)), \\
d_0^3 &= -\frac{d_0^2}{4i_1^3} (e_0 h_1^2 i_1 - 12f_0 g_1 i_1^2 - f_0 h_1^3) \\
&\quad - \frac{d_0}{i_1^3} \left(-\frac{1}{2} e_0^2 g_1 h_1 i_1 - \frac{1}{4} e_0^2 i_1^3 G_2 + \frac{1}{2} e_0 f_0 g_1 h_1^2 + \frac{1}{2} e_0 f_0 h_1 i_1^2 G_2 + 3f_0^2 g_1^2 i_1 - \frac{1}{4} f_0^2 h_1^2 i_1 G_2 \right) \\
&\quad - \frac{1}{i_1^3} \left(\frac{1}{4} e_0^3 g_1^2 i_1 + \frac{1}{4} e_0^3 i_1^3 G_3 - \frac{1}{4} e_0^2 f_0 g_1^2 h_1 + \frac{1}{4} e_0^2 f_0 g_1 i_1^2 G_2 - \frac{3}{4} e_0^2 f_0 h_1 i_1^2 G_3 \right)
\end{aligned}$$

$$\begin{aligned}
& -\frac{1}{2}e_0f_0^2g_1h_1i_1G_2 + \frac{3}{4}e_0f_0^2h_1^2i_1G_3 - f_0^3g_1^3 + \frac{1}{4}f_0^3g_1h_1^2G_2 - \frac{1}{4}f_0^3h_1^3G_3), \\
g_0 &= \frac{i_0g_1}{i_1}, \quad h_0 = \frac{i_0h_1}{i_1}, \\
a_1 &= \frac{g_1}{i_1}c_1, \quad b_1 = \frac{h_1}{i_1}c_1, \\
c_1^3 &= \frac{c_1^2}{i_0^3h_1^3} \left(3b_0i_0^2h_1^2i_1^2 + \frac{1}{4}e_0^3i_1^4 - \frac{1}{4}e_0^2i_0f_1h_1i_1^2 \right) \\
& - \frac{c_1}{i_0^3h_1^3} \left(3b_0^2i_0h_1i_1^4 + \frac{1}{2}b_0e_0^2f_1i_1^4 - \frac{1}{2}b_0e_0i_0f_1^2h_1i_1^2 - \frac{1}{4}e_0^2i_0h_1i_1^4g_2 + \frac{1}{2}e_0i_0^2f_1h_1^2i_1^2g_2 - \frac{1}{4}i_0^3f_1^2h_1^3g_2 \right) \\
& - \frac{1}{i_0^3h_1^3} \left(-b_0^3i_1^6 - \frac{1}{4}b_0^2e_0f_1^2i_1^4 + \frac{1}{4}b_0^2i_0f_1^3h_1i_1^2 + \frac{1}{4}b_0e_0^2i_1^6g_2 - \frac{1}{2}b_0e_0i_0f_1h_1i_1^4g_2 \right. \\
& \quad \left. + \frac{1}{4}b_0i_0^2f_1^2h_1^2i_1^2g_2 - \frac{1}{4}e_0^3i_1^6g_3 + \frac{3}{4}e_0^2i_0f_1h_1i_1^4g_3 - \frac{3}{4}e_0i_0^2f_1^2h_1^2i_1^2g_3 + \frac{1}{4}i_0^3f_1^3h_1^3g_3 \right), \\
d_1 &= \frac{f_1g_1}{i_1}, \quad e_1 = \frac{f_1h_1}{i_1}.
\end{aligned}$$

A.4.3. $e_1i_1 - f_1h_1 = 0$, $h_0i_1 - i_0h_1 = 0$, $h_1 = 0$, $i_1h_0 \neq 0$.

$$\begin{aligned}
a_0 &= \frac{b_0}{e_0}d_0 + \frac{i_0g_1}{i_1^2}c_1 - \frac{b_0i_0f_1g_1}{e_0i_1^2}, \quad c_0 = \frac{i_0}{i_1}c_1 - \frac{-b_0f_0i_1 + b_0i_0f_1}{e_0i_1}, \\
d_0^3 &= 3\frac{f_0g_1}{i_1}d_0^2 - \frac{-\frac{1}{4}e_0^2i_1^2G_2 + 3f_0^2g_1^2}{i_1^2}d_0 \\
& - \frac{\frac{1}{4}e_0^3g_1^2i_1 + \frac{1}{4}e_0^3i_1^3G_3 + \frac{1}{4}e_0^2f_0g_1i_1^2G_2 - f_0^3g_1^3}{i_1^3}, \\
g_0 &= \frac{i_0g_1}{i_1}, \quad h_0 = 0, \quad a_1 = \frac{g_1}{i_1}c_1, \quad b_1 = 0, \\
c_1^2 &= \frac{2b_0f_1}{e_0}c_1 - \frac{4b_0^3i_1^2 + b_0^2e_0f_1^2 - b_0e_0^2i_1^2g_2 + e_0^3i_1^2g_3}{e_0^3}, \\
d_1 &= \frac{f_1g_1}{i_1}, \quad e_1 = 0, \quad h_1 = 0.
\end{aligned}$$

A.4.4. $e_1i_1 - f_1h_1 = 0$, $h_0i_1 - i_0h_1 = 0$, $i_0 = 0$, $i_1e_0 \neq 0$.

$$\begin{aligned}
a_0 &= \frac{b_0}{e_0}d_0, \quad c_0 = \frac{b_0f_0}{e_0}, \\
d_0^3 &= -\frac{d_0^2}{4i_1^3} (e_0i_1h_1^2 - 12f_0g_1i_1^2 - f_0h_1^3) \\
& - \frac{d_0}{i_1^3} \left(-\frac{1}{2}e_0^2g_1i_1h_1 - \frac{1}{4}e_0^2i_1^3G_2 + \frac{1}{2}e_0f_0g_1h_1^2 + \frac{1}{2}e_0f_0i_1^2h_1G_2 + 3f_0^2g_1^2i_1 - \frac{1}{4}f_0^2i_1h_1^2G_2 \right) \\
& - \frac{1}{i_1^3} \left(\frac{1}{4}e_0^3g_1^2i_1 + \frac{1}{4}e_0^3i_1^3G_3 - \frac{1}{4}e_0^2f_0g_1^2h_1 + \frac{1}{4}e_0^2f_0g_1i_1^2G_2 - \frac{3}{4}e_0^2f_0i_1^2h_1G_3 - \frac{1}{2}e_0f_0^2g_1i_1h_1G_2 \right. \\
& \quad \left. + \frac{3}{4}e_0f_0^2i_1h_1^2G_3 - f_0^3g_1^3 + \frac{1}{4}f_0^3g_1h_1^2G_2 - \frac{1}{4}f_0^3h_1^3G_3 \right), \\
g_0 &= 0, \quad h_0, \\
a_1 &= \frac{g_1}{i_1}c_1, \quad b_1 = \frac{h_1}{i_1}c_1, \\
c_1^2 &= \frac{2b_0f_1}{e_0}c_1 - \frac{4b_0^3i_1^2 + b_0^2e_0f_1^2 - b_0e_0^2i_1^2g_2 + e_0^3i_1^2g_3}{e_0^3}, \\
d_1 &= \frac{f_1g_1}{i_1}, \quad e_1 = \frac{f_1h_1}{i_1}.
\end{aligned}$$

A.4.5. $e_1 i_1 - f_1 h_1 = 0$, $h_0 i_1 - i_0 h_1 = 0$, $e_0 i_1^2 - i_0 f_1 h_1 = 0$, $i_1 h_1 i_0 (f_0 i_1 - i_0 f_1) \neq 0$.

$$\begin{aligned}
a_0 &= \frac{1}{f_0 i_1 - i_0 f_1} (i_1 c_0 d_0 - i_0 f_1 g_1 c_0 - b_0 i_1^2 d_0 + b_0 f_0 g_1 i_1), \\
c_0^3 &= \frac{c_0^2}{h_1 i_1^3} (3b_0 i_1^4 - \frac{1}{4} f_0 f_1^2 h_1 i_1 + \frac{1}{4} i_0 f_1^3 h_1) \\
&\quad - \frac{c_0}{i_0 h_1^2 i_1^2} (3b_0^2 i_0 i_1^4 - \frac{1}{2} b_0 f_0^2 f_1 h_1 i_1^2 + \frac{1}{2} b_0 f_0 i_0 f_1^2 h_1 i_1 - \frac{1}{4} f_0^2 i_0 h_1^2 i_1^2 g_2 + \frac{1}{2} f_0 i_0^2 f_1 h_1^2 i_1 g_2 - \frac{1}{4} i_0^3 f_1^2 h_1^2 g_2) \\
&\quad - \frac{1}{i_0^2 h_1^3 i_1^3} (-b_0^3 i_0^2 i_1^6 + \frac{1}{4} b_0^2 f_0^3 h_1 i_1^5 - \frac{1}{4} b_0^2 f_0^2 i_0 f_1 h_1 i_1^4 + \frac{1}{4} b_0 f_0^2 i_0^2 h_1^2 i_1^4 g_2 - \frac{1}{2} b_0 f_0 i_0^3 f_1 h_1^2 i_1^3 g_2 \\
&\quad\quad + \frac{1}{4} b_0 i_0^4 f_1^2 h_1^2 i_1^2 g_2 + \frac{1}{4} f_0^3 i_0^2 h_1^3 i_1^3 g_3 - \frac{3}{4} f_0^2 i_0^3 f_1 h_1^3 i_1^2 g_3 + \frac{3}{4} f_0 i_0^4 f_1^2 h_1^3 i_1 g_3 - \frac{1}{4} i_0^5 f_1^3 h_1^3 g_3), \\
d_0^3 &= -\frac{d_0^2}{i_1^4} (3f_0 g_1 i_1^3 + \frac{1}{4} f_0 h_1^3 i_1 - \frac{1}{4} i_0 f_1 h_1^3) \\
&\quad - \frac{d_0}{i_1^6} (3f_0^2 g_1^2 i_1^4 - \frac{1}{4} f_0^2 h_1^2 i_1^4 G_2 + \frac{1}{2} f_0 i_0 f_1 g_1 h_1^3 i_1 + \frac{1}{2} f_0 i_0 f_1 h_1^2 i_1^3 G_2 - \frac{1}{2} i_0^2 f_1^2 g_1 h_1^3 - \frac{1}{4} i_0^2 f_1^2 h_1^2 i_1^2 G_2) \\
&\quad - \frac{1}{i_1^8} (-f_0^3 g_1^3 i_1^5 + \frac{1}{4} f_0^3 g_1 h_1^2 i_1^5 G_2 - \frac{1}{4} f_0^3 h_1^3 i_1^5 G_3 - \frac{1}{2} f_0^2 i_0 f_1 g_1 h_1^2 i_1^4 G_2 + \frac{3}{4} f_0^2 i_0 f_1 h_1^3 i_1^4 G_3 \\
&\quad\quad - \frac{1}{4} f_0 i_0^2 f_1^2 g_1^2 h_1^3 i_1 + \frac{1}{4} f_0 i_0^2 f_1^2 g_1 h_1^2 i_1^3 G_2 - \frac{3}{4} f_0 i_0^2 f_1^2 h_1^3 i_1^3 G_3 + \frac{1}{4} i_0^3 f_1^3 g_1^2 h_1^3 + \frac{1}{4} i_0^3 f_1^3 h_1^3 i_1^2 G_3), \\
e_0 &= \frac{i_0 f_1 h_1}{i_1^2}, \quad g_0 = \frac{i_0 g_1}{i_1}, \quad h_0 = \frac{i_0 h_1}{i_1}, \quad a_1 = \frac{b_0 g_1 i_1}{i_0 h_1}, \\
b_1 &= \frac{b_0 i_1}{i_0}, \quad c_1 = \frac{b_0 i_1^2}{i_0 h_1}, \quad d_1 = \frac{f_1 g_1}{i_1}, \quad e_1 = \frac{f_1 h_1}{i_1}.
\end{aligned}$$

REFERENCES

- [1] V.E. Adler, A.I. Bobenko and Yu.B. Suris, *Classification of integrable equations on quadrilaterals, the consistency approach*, Commun. Math. Phys. **233** (2003) 513–543.
- [2] J. Alonso, Yu.B. Suris and K. Wei, *A three-dimensional generalization of QRT maps*, arXiv: 2207.06051 [nlin.SI] (2023).
- [3] J. Atkinson and F.W. Nijhoff, *A constructive approach to the soliton solutions of integrable quadrilateral lattice equations*, Commun. Math. Phys. **299** (2010) 283–304.
- [4] W. Bosma, J. Cannon, and C. Playoust, *The Magma algebra system I: The user language*, Journal of Symbolic Computation **24**.3-4 (1997) 235–265.
- [5] J.H.H. Duistermaat, *Discrete Integrable Systems: the QRT Map and Elliptic Surfaces*, Springer, New York (2011).
- [6] P. Gianni, B. Trager, G. Zacharias, *Gröbner Bases and Primary Decomposition of Polynomial Ideals*. J. Symb. Comp. **6**, (1988), 149-167.
- [7] P. Griffiths and J. Harris, *Principles of Algebraic Geometry*, Wiley-Interscience, New York, 1978.
- [8] J. Hietarinta, N. Joshi and F.W. Nijhoff, *Discrete Systems and Integrability*, Cambridge University Press, Cambridge, UK (2016).
- [9] N. Joshi, *Discrete Painlevé equations*, CBMS Regional Conference Series in Mathematics, **131**, AMS, Providence, RI, (2019),
- [10] P. Kassotakis and N. Joshi, *Integrable non-QRT mappings of the plane*, Letters in Mathematical Physics **91** (2010) 71-81.
- [11] G.R.W. Quispel, J.A.G. Roberts and C. Thompson, *Integrable mappings and soliton equations. I & II*, Phys. Lett. **126A** (1988) 419–421; Physica **D34** (1989) 183–192.
- [12] J.A.G. Roberts, *Order and Chaos in reversible dynamical systems*, LaTrobe University PhD thesis, 1990.
- [13] J.A.G. Roberts and G.R.W. Quispel, *Creating and relating three-dimensional integrable maps*, J. Phys. A: Math. Gen. **39** (2006) L605–L615.
- [14] L. Stickelberger and G.F. Frobenius, “Ueber die Differentiation der elliptischen Functionen nach den Perioden und Invarianten.” Journal für die reine und angewandte Mathematik, **1882**, no. 92, 1882, pp. 311-327. <https://doi.org/10.1515/crll.1882.92.311>

- [15] T. Tsuda, *Integrable mappings via rational elliptic surfaces*, J. Phys. A: Math Gen **37** (2004) 2721–2730.
- [16] R. Zippel, *Probabilistic algorithms for sparse polynomials*, Symbolic and algebraic computation, LNCS **72** (1979), 216–226.

SCHOOL OF MATHEMATICS AND STATISTICS F07, THE UNIVERSITY OF SYDNEY, NSW 2006, AUSTRALIA.

Email address: `nalini.joshi@sydney.edu.au`

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9JT, UK

Email address: `frank.nijhoff@gmail.com`

SCHOOL OF MATHEMATICS AND STATISTICS F07, THE UNIVERSITY OF SYDNEY, NSW 2006, AUSTRALIA.

Email address: `allan.steel@sydney.edu.au`