Phenomenological quantum mechanics: deducing the formalism from experimental observations

Piotr Szańkowski¹, Davide Lonigro², Fattah Sakuldee^{3,4}, Łukasz Cywiński¹, and Dariusz Chruściński⁵

¹Institute of Physics, Polish Academy of Sciences, al. Lotników 32/46, PL 02-668 Warsaw, Poland

²Department Physik, Friedrich-Alexander-Universität Erlangen-Nürnberg, Staudtstraße 7, 91058 Erlangen, Germany

³Wilczek Quantum Center, School of Physics and Astronomy, Shanghai Jiao Tong University, 800 Dongchuan Road, Minhang, 200240 Shanghai, China

⁴The International Centre for Theory of Quantum Technologies, University of Gdańsk, Jana Bażyńskiego 1A, 80-309 Gdańsk, Poland

⁵Institute of Physics, Faculty of Physics, Astronomy and Informatics, Nicolaus Copernicus University, Grudziadzka 5/7, 87-100 Toruń, Poland

We propose an exercise in which one attempts to deduce the formalism of quantum mechanics from phenomenological observations only. Thus, the only assumed inputs are the multi-time probability distributions estimated from the results of sequential measurements of quantum observables; no presuppositions about the underlying mathematical structures are allowed. We show that it is indeed possible to derive in such a way a complete and fully functional formalism based on the structures of Hilbert spaces. However, the obtained formal description—the bi-trajectory formalism—turns out to be quite different from the standard state-focused formalism.

Contents

1	Introduction	2
2	Outline	4
3	Description of sequential measurement experiments	6
4	Phenomenology of sequential measurement experiments	8
	4.1 Quantum coarse-grained measurements	8
	4.2 Quantum interference	10
	4.3 Quantum interference between independent systems	12
	4.4 Markovianity and the initialization events	16
	4.5 Quantum Zeno effect	18
	4.6 Multiple observables and uncertainty relations	19
5	The bi-trajectory picture of quantum mechanics	21
	5.1 Basis for the deduced formalism	21
	5.2 Bi-probability distributions	23
	5.3 The bi-trajectory picture	25
6	The Hilbert space representation	26
	6.1 Bi-probability distributions as inner products	26
	6.2 The measurement–projector link	28
	6.3 Hilbert space representation for bi-probabilities	30
<u> </u>		

Piotr Szańkowski: piotr.szankowski@ifpan.edu.pl

7	Dynamical laws of the system7.1Unitary evolution operator7.2Generator of the evolution operator	32 32 33
8	Composite systems	35
9	The master object of the formalism	36
10	Summary of the deduced formalism	39
11	Conclusions	41

1 Introduction

One of the iconic features of quantum mechanics is that the standard formalism of the theory implements a dichotomous (two-fold) mode of description.

The first mode applies in *non-measurement contexts*, where the subject—a quantum system of interest—evolves freely without being disturbed by the probing with *measuring devices* deployed by an observer. In this case, a system is described by its time-dependent *state*, formally represented by a density matrix $\hat{\rho}_t$ in the form of a unit-trace positive semi-definite operator acting on a given Hilbert space. The system-specific *dynamical law* governing the state's time evolution is represented in the formalism by a unitary transformation

$$\hat{\rho}_{t_1} = \hat{U}_{t_1, t_0} \hat{\rho}_{t_0} \hat{U}_{t_1, t_0}^{\dagger}.$$
(1.1)

The second mode is intended for complementary measurement contexts, where it is used to describe the perceptions of an observer who intervenes at a given time t_1 by deploying a device capable of measuring a system observable. Accordingly, the outcome of such a measurement, as seen by the observer, is formally described by the probability distribution conditioned by the system state, given by the well-known *Born rule*,

$$P(f \mid \hat{\rho}_{t_1}) = \text{tr}[\hat{P}^F(f)\hat{\rho}_{t_1}].$$
(1.2)

Here, $\hat{F} = \sum_{f \in \Omega(F)} f \hat{P}^F(f)$ is a Hermitian operator representing the measured observable F, decomposed in terms of a complete family $\{\hat{P}^F(f) \mid f \in \Omega(F)\}$ of orthogonal projectors,

$$\sum_{f \in \Omega(F)} \hat{P}^F(f) = \hat{1} \quad \text{and} \quad \hat{P}^F(f) \hat{P}^F(f') = \delta_{f,f'} \hat{P}^F(f).$$
(1.3)

The *collapse rule* supplements the standard formalism by serving as an interface between the two modes. It states that, once the measurement is concluded and one transitions from the measurement context to the non-measurement context, the system is assigned a new state conditioned by the measurement result,

$$\hat{\rho}_{t_1} \xrightarrow{\text{state collapse}} \hat{\rho}_{t|t_1}(f_1) = \hat{U}_{t,t_1} \frac{\hat{P}^F(f_1)\hat{\rho}_{t_1}\hat{P}^F(f_1)}{\operatorname{tr}\left[\hat{P}^F(f_1)\hat{\rho}_{t_1}\right]} \hat{U}^{\dagger}_{t,t_1}, \qquad (1.4)$$

given that the result at time t_1 was $f_1 \in \Omega(F)$.

The addition of the collapse rule is necessary to construct the description of a measurement context that extends beyond the first deployment of the measuring device. Hence, in the standard formulation, the observed results of a measurement carried out with devices deployed in a chronological sequence are formally described by chaining the Born rule and the collapse rule: the system state at the time of the subsequent measurement is taken as the state collapsed due to previous measurement (and evolved for the duration in between the two measurements). That is, the probability that the observer measures the result f_1 at time t_1, f_2 at t_2, \ldots , and finally f_n at t_n , is given by

$$P_{t_n,\dots,t_1}(f_n,\dots,f_1) = P(f_1|\rho_{t_1}) \prod_{j=2}^n P(f_j|\hat{\rho}_{t_j|t_{j-1},\dots,t_1}(f_{j-1},\dots,f_1)),$$
(1.5)

where the conditional system states are found by a recursive application of the rule (1.4),

$$\hat{\rho}_{t|t_j...,t_1}(f_j,\ldots,f_1) = \hat{U}_{t,t_j} \frac{\hat{P}^F(f_j)\hat{\rho}_{t_j|t_{j-1},\ldots,t_1}(f_{j-1},\ldots,f_1)\hat{P}^F(f_j)}{P(f_j|\hat{\rho}_{t_j|t_{j-1},\ldots,t_1}(f_{j-1},\ldots,f_1))} \hat{U}_{t,t_j}^{\dagger}.$$
(1.6)

The choice to structure the formalism in this way has certain undesirable consequences. One of them is the necessity for the *Heisenberg cut* [1–3]. The issue is that the two description modes apply only in their respective domains, and so, whenever one wishes to employ the formalism, one has to decide whether a given situation counts as a measurement or not. But what are the quantitative criteria for making such a decision? The current understanding of the capabilities of the standard formalism is that it cannot provide a concrete answer to such a question [4–7]. Pinpointing the transition between the measurement and non-measurement context for a collection of interacting quantum systems remains an open problem. Therefore, the only available solution is to place the *cut* more or less arbitrarily; as long as one operates far away from the cut, the formalism can be used without further objections.

However, this is not—and never has been—a satisfactory resolution. On one hand, the friction due to the dichotomous description modes has been an inexhaustible source of inspiration for paradoxical thought experiments (perhaps best exemplified by the Wigner's friend scenarios, in which an observer observes another observer); on the other hand, modern quantum technologies also operate at scales that approach perilously close to the expected location of the Heisenberg cut. Thus, it seems inevitable that the standard formalism, being incapable of describing systems living in the admittedly fuzzy border separating the measurement and non-measurement contexts, will, sooner rather than later, reach the limits of its applicability in cutting-edge quantum physics.

Naturally, over the years, there have been a number of attempts at rectifying these problems. Examples include the de Broglie–Bohm pilot-wave theory [8, 9], Griffiths decoherent histories theory [10, 11], Everett many-worlds interpretation [12], and others. However, none of these proposed solutions has garnered universal acceptance as the *de facto* resolution of the tensions identified in the standard formalism. Perhaps the issue lies in the fact that these attempts attack the problem by *postulating* an often sweeping restructuring of the formalism. Even though the proposed modifications are usually motivated by some insightful intuition, in the end they are *ad hoc* propositions.

We concur with the diagnosis that the problems with the standard formalism cannot be reconciled within its framework, and the solution will most likely require a restructuring of the formalism. However, rather than postulating *ad hoc* modifications to the existing formalism, we propose a more systematic approach. The idea is to go back to basics, forget what we know about quantum theory, and attempt to *deduce* the formalism from the experimental observations only—*phenomenological* quantum mechanics.

We imagine that this route is not dissimilar to the one traversed a century ago by the pioneers of quantum theory, who invented the standard formalism in the first place. However, the breadth of the phenomenology of quantum systems has expanded tremendously over these 100 years; the experimental inputs available to us as a matter of course were considered, back then, as fanciful thought experiments at best and science-fiction at worst. Unsurprisingly, then, the theoreticians of quantum mechanics focused, at its inception, almost exclusively on the phenomenology of single-measurement experiments. This seems to be the main reason why the standard formalism has a dichotomous structure, and the interface between the description mode takes the form of the collapse rule. If this is a correct reading, then the collapse rule appears as an improvised *post hoc* solution that was "bolted" onto the formalism because it was just that—a solution to the problem of sequential measurements that, likely, was treated as an afterthought. Hence, to avoid the pitfalls encountered by the pioneers, we ought to treat experiments involving sequential measurements as the basic input for our deduction.

In this paper, we demonstrate that the phenomenology of sequential measurement experiments is indeed rich enough to deduce a fully functional and complete formalism without presupposing *anything* about its structure (not even a connection with Hilbert spaces has to be assumed). As expected, the so-deduced formalism—the *bi-trajectory formalism*, as we call it—turns out to be structured significantly differently from the standard formalism. In particular, the dichotomous mode of description is absent in the bi-trajectory formulation, allowing it to describe quantum, classical, and in-between systems without invoking the Heisenberg cut, state collapse, or other similar concepts.

Crucially, because the formalism is directly deduced from the phenomenology, its predictions are in full agreement with the empirically confirmed predictions that could also be made with the standard formalism. Additionally, the new formalism can be reduced to a form that is mathematically equivalent to the standard formulation.

The elements of the bi-trajectory formalism have been introduced in previous works [13–17]. In these works, the bi-trajectory-related concepts have been employed to parameterize open system dynamics [15], describe the origins of the classical noise limit in open systems [13], investigate the quantum-classical transition [14, 16], and in [17] all these elements were given solid mathematical foundations. However, only now, with the demonstration of a successful deduction from phenomenology, can the elements of the bi-trajectory formulation be brought together as a holistic formalism for quantum mechanics.

2 Outline

In the first part of the paper, we list the phenomenological observations constituting the inputs for the following deduction of the formalism. Section 3 introduces the formal description of the results of experiments involving sequential measurements using multi-time probability distributions. In Section 4, we use this description to quantify our phenomenological inputs:

We begin in Section 4.1 by providing evidence that sequential measurements of a quantum system cannot be interpreted as a sampling of a *trajectory* traced over time by the system's observable. This crucial observation establishes that such systems cannot be described using the classical theory. Furthermore, we recognize the fundamental distinction between classical and quantum systems: for classical systems all observables are essentially equivalent, whereas quantum systems support an infinite number of *inequivalent* observables. We explore some consequences of this peculiar feature of quantum systems by comparing measurements made with coarse- and fine-grained devices—an important example of inequivalent quantum observables.

Section 4.2 showcases the quantum interference effects in what amounts to a sequence of k-slit generalizations of the classic Young experiment, viewed as an example of sequential measurement with coarse-grained devices.

In Section 4.3, we discuss experiments involving multiple *independent* systems that exhibit quantum interference when measured with devices incapable of making a perfect "which-subsystem" distinction. To uphold the spirit of phenomenological quantum mechanics, we introduce, for the purposes of this analysis, the *phenomenological definition* of a composite quantum system.

Section 4.4 discusses experimental conditions under which the probability distributions describing sequential measurements exhibit *Markovianity*, enabling us to address the issue of experiment initialization.

Section 4.5 describes the quantum Zeno effect, consisting in the following phenomenon: sequences of measuring devices deployed in rapid succession become "frozen" in displaying the first measured result when the delay between the deployments tends to zero while their number goes to infinity. This observation is vital for probing the short-time nature of quantum dynamical laws.

Finally, Section 4.6 concludes the first part of the paper by discussing the *uncertainty relations* between measurements of inequivalent quantum observables.

The second part of the paper presents the process of the actual deduction of the formalism from the phenomenological data, including the detailed breakdown of the necessary mathematical inferences. Section 5 opens this part by showing how the phenomenological observations discussed in the first part lead to the conclusion that the emerging formalism is shaping into a *bi-trajectory picture*, a natural progression from the classical "(single) trajectory picture".

In Section 6, we find the Hilbert space representation for *bi-probabilities*, the complex-valued distributions on the space of trajectory pairs (the namesake of the bi-trajectory picture) and the basis of the bi-trajectory formalism.

In Section 6.1, we establish the critical link between bi-probabilities and Hilbert spaces via Gudder's theorem on inner product representation [18]. In Section 6.2, we exploit this connection along with the phenomenology of single-measurement experiments to infer the measurement-projector link, a foundational concept in the standard formalism. Finally, in Section 6.3, we derive how biprobabilities are represented in Hilbert space structures, such as projector operators and density matrices. This also reveals the limitations of the measurement-projector link.

In Section 7, we further refine the Hilbert space representation, drawing on the phenomenology of uncertainty relations and the Zeno effect, leading to the appearance of time-dependent unitary operators—and their generators—which are identified as the formal representation of the quantum dynamical laws.

In Section 8, we infer the form of the Hilbert space corresponding to a system composed of independent subsystems. We then deduce that independent systems become dependent when the dynamical law includes *interaction terms*. This establishes the correspondence between Hermitian operators and observables measurable by devices deployed by an observer.

Finally, Section 9 concludes the second part of the paper by identifying the *master object* of the bi-trajectory formalism—the fundamental element from which all other elements, including phenomenologically accessible multi-time probabilities, are derived. Final considerations and outlooks are gathered in Section 11.

Part I Phenomenology of quantum systems

In the first part of the paper, we analyze the outcomes of various experiments that can be reduced to the sequential deployment of devices measuring quantum observables. These outcomes are quantitatively described by phenomenological multi-time probability distributions, estimated using data obtained from many repetitions of each experiment. Our first conclusion is that the theory describing the analyzed experimental outcomes must be *non-classical* because we find that the measured sequences cannot be considered as sampling of a *trajectory* representing system's observables. The breakdown of this "trajectory picture" results from the violation of the *consistency condition* by the obtained multi-time probabilities—a feature fundamentally incompatible with any kind of classical theory.

The non-classical nature of the investigated systems manifests in various observed phenomena, including quantum interference, quantum Zeno effect, and uncertainty relations between the measured observables. Each of these phenomena corresponds to specific properties of the multi-time probabilities, providing clues for our deduction of the formalism carried out in the second part of the paper.

3 Description of sequential measurement experiments

Assume that the experimenter has access to a collection of measuring devices capable of probing observables of a quantum system. When a device measuring an observable F—an F-device for short—is deployed by the experimenter, one of the possible results $f \in \Omega(F)$,¹ corresponding to the measurement outcome obtained by the device at the time t of its deployment, is displayed.

Now, suppose that the experimenter carried out a vast number of sequential measurements with their devices, and diligently recorded the outcome of each experiment as a sequence (an *n*-tuple) of results,

$$\boldsymbol{f}_n = (f_n, \dots, f_1) \in \Omega(F_n) \times \dots \times \Omega(F_1) \equiv \Omega(\boldsymbol{F}_n), \tag{3.1}$$

displayed by the devices as they were deployed at the corresponding timings,

$$\boldsymbol{t}_n = (t_n, \dots, t_1) \qquad \text{such that } t_n > \dots > t_1. \tag{3.2}$$

These records, obtained for a wide spread of sequence lengths, timings, and device choices, were then used to estimate the corresponding probability distributions,

$$P_{t_n,\dots,t_1}^{F_n,\dots,F_1}(f_n,\dots,f_1) \equiv P_{t_n}^{F_n}(f_n).$$
(3.3)

Here, F_j at position j in the superscript indicates that the measurement at time t_j was performed with an F_j -device. Provided that the experimenter did an outstanding job at finding the most accurate estimations, the family of probabilities $P_{t_n}^{F_n}$ constitutes the complete quantitative description of the phenomenology of sequential measurements.

Consequently, the phenomenological inputs for our deduction of the formalism of quantum mechanics are described using the language of probability theory. Within this language, the set $\Omega(\mathbf{F}_n)$ is identified as the *sample space* for an experiment consisting of n measurements where the F_1 -device was deployed first, followed by the F_2 -device, ..., and terminating with the F_n -device. The sequence of results $\mathbf{f}_n \in \Omega(\mathbf{F}_n)$ recorded as described above is then a *singleton* event, so that

¹Here, we shall restrict our considerations to $\Omega(F)$ that are finite sets; we defer the case of (countable or uncountable) infinite sets of readouts for future projects.

the value of the probability distribution for a given f_n equals the *measure* of a set consisting of a single element,

$$P_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n) = P_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\{\boldsymbol{f}_n\}) \tag{3.4}$$

Note that, with a slight abuse of notation, we are using the same symbol here to denote the probability distribution and the probability measure it is associated with.

Any set of elements of the sample space $\Omega(\mathbf{F}_n)$, which represents an event, can always be decomposed into a finite disjoint union \sqcup of singletons. For example, take the event $A = \{\mathbf{f}_n \mid f_j \in A_j \subset \Omega(F_j)\}$ representing an *alternative* of singleton events; this set decomposes into the union of singletons as follows,

$$A = \bigsqcup_{\boldsymbol{f}_n \in A_n \times \dots \times A_1} \{\boldsymbol{f}_n\}.$$
(3.5)

Given that every probability measure $P_{t_n}^{F_n}$ is by definition additive,

$$P_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}\left(\bigsqcup_j A_j\right) = \sum_j P_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(A_j),\tag{3.6}$$

the singletons $\{f_n\}$ can therefore be considered as basis events: it suffices to specify the measure of each singleton sequence to completely characterize the probability of any event,

$$P_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(A) = \sum_{\boldsymbol{f}_n \in A} P_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n).$$
(3.7)

The probability of measuring any given sequence of results generally depends on the chronology of the performed measurements. Therefore, any given result $f_n \in \Omega(\mathbf{F}_n)$ obtained in a sequence of n measurements performed at the corresponding times \mathbf{t}_n and the same sequence of results but measured at different timings $\mathbf{t}'_n \neq \mathbf{t}_n$ are formally described as two distinct events. Consequently, the probabilities of those two events are given by two distinct probability measures $P_{\mathbf{t}_n}^{\mathbf{F}_n}$: Pow $(\Omega(\mathbf{F}_n)) \rightarrow [0,1]$ and $P_{\mathbf{t}'_n}^{\mathbf{F}_n}$: Pow $(\Omega(\mathbf{F}_n)) \rightarrow [0,1]$, each corresponding to a unique sequence of timings.²

To illustrate these points, consider the following example. From the additivity of probability measures, we obtain the equality

$$\sum_{f_j \in \Omega(F_j)} P_{t_n}^{F_n}(f_n, \dots, f_j, \dots, f_1) = P_{t_n}^{F_n} \Big(\Big\{ (f_n, \dots, f_j, \dots, f_1) \mid f_j \in \Omega(F_j) \Big\} \Big).$$
(3.8)

In words, this relation means that the probability obtained via marginalization over the result measured at time t_j equals the probability of measuring a sequence $(f_n, \ldots, f_j, \ldots, f_1)$ where all results are set to specific values, except for f_j , which can be anything displayed by the F_j -device. Intuitively, one might expect that such an alternative, where the sequence is not constrained by any particular value of f_j , should be equivalent to an event where the F_j -device was never deployed. If this were the case, then the marginalized probability would be equal to probability of a sequence of length n - 1, in which the entry corresponding to t_j has been struck out. However, since $P_{t_n,\ldots,t_j,\ldots,t_1}^{F_n,\ldots,F_1}$ and $P_{t_n,\ldots,t_j,\ldots,t_1}^{F_n,\ldots,F_1}$ are measures on distinct event spaces, there is no purely logical or mathematical reason for any relation between probability distributions pertaining to different sequences of timings. If any such relation does exist, it can only be established on the grounds of phenomenological observations of the physical processes they are used to describe. Therefore, whether the aforementioned intuitive postulate (or any other such postulate) is correct must be tested in experiments.

²The symbol Pow($\Omega(\mathbf{F}_n)$) indicates the power set of $\Omega(\mathbf{F}_n)$, that is, the set of all subsets of $\Omega(\mathbf{F}_n)$. This includes the empty set \emptyset , the whole set $\Omega(\mathbf{F}_n)$, all singletons $\{\mathbf{f}_n\}$, and all their unions.

4 Phenomenology of sequential measurement experiments

4.1 Quantum coarse-grained measurements

The first phenomenological observation concerns a relation between probability distributions $P_{t_n}^{F_n}$: $\Omega(F_n) \to [0, 1]$ describing sequences of different lengths:

Phenomenological observation 1 (Causality). When a probability distribution describing the sequential measurement is marginalized over the latest entry in the sequence, one obtains the distribution corresponding to the shorter time sequence,

$$\sum_{f_{n+1}\in\Omega(F_{n+1})} P_{t_{n+1}}^{F_{n+1}}(f_{n+1}) = P_{t_n}^{F_n}(f_n).$$

The follow-up conclusion is that, in general, this property does *not* extend to any other entry in the sequence:

Phenomenological observation 2 (Inconsistency). The marginalization over the readouts midsequence does not generally result in a distribution for a shorter sequence:

$$P_{t_n, \dots, t'_j, \dots, t_1}^{F_n, \dots, F_1}(f_n, \dots, f'_j, \dots, f_1) - \sum_{f_j \in \Omega(F_j)} P_{t_n}^{F_n}(f_n) \neq 0 \quad \text{for } j < n.$$

In terms of the mathematical theory of probability, this means that the family of distributions $P_{t_n}^F$ violates the Kolmogorov consistency condition. Consequently, the standard Kolmogorov extension theorem does not apply here, and the probabilities $P_{t_n}^F$ cannot be considered as discrete-time restrictions of a master measure $\mathcal{P}[f][\mathcal{D}f]$ on the space of trajectories $t \mapsto f(t) \in \Omega(F)$ [16, 17, 19], i.e., in general

$$P_{\boldsymbol{t}_n}^F(\boldsymbol{f}_n) \neq \int \Big(\prod_{j=1}^n \delta_{f(t_j), f_j}\Big) \mathcal{P}^F[f][\mathcal{D}f].$$
(4.1)

Physically, this means that, in general, a sequence of outcomes obtained by measuring quantum system *cannot* be interpreted as a sampling of the trajectory traced over time by the system's observable. This is the point where quantum and classical physics fundamentally diverge: in classical theories the Kolmogorov consistency is one of the defining assumptions, and the trajectory picture of physical quantities is a given.

The most salient consequence of the classical trajectory picture in regards to sequential measurements is that, at each instant of time, all classical observables have definite values—the corresponding trajectories unravel in time simultaneously. This statement is equivalent to the assumption that, in classical theories, all measurements are described by distributions satisfying Kolmogorv consistency condition. Consequently, for each observable, there exists a master measure on the space of trajectories describing its dynamics. Therefore, for every classical system, one can treat all mutually independent observables as components of a single vector trajectory—the *elementary* observable. In that case, the trajectory representing any other observable must be a function of the trajectory of the elementary observable, and thus, the probability distribution describing its measurements has the form

$$P_{\boldsymbol{t}_{n}}^{\boldsymbol{X}_{n}}(\boldsymbol{x}_{n}) = \int \Big(\prod_{j=1}^{n} p_{t_{j}}^{X_{j}}(x_{j}|e(t_{j}))\Big) \mathcal{P}^{E_{\text{cls}}}[e][\mathcal{D}e] = \sum_{\boldsymbol{e}_{n}} P_{\boldsymbol{t}_{n}}^{E_{\text{cls}}}(\boldsymbol{e}_{n}) \prod_{j=1}^{n} p_{t_{j}}^{X_{j}}(x_{j}|e_{j}), \qquad (4.2)$$

where E_{cls} is the elementary observable of the classical system.

When measurements are described by inconsistent distributions, like in quantum mechanics, observables do not have to have a definite value at each instant of time. Then, there is no reason why such a non-classical system would possess an elementary observable, and thus, not every device measuring its observables has to be described by a probability distribution of the form (4.2). This crucial categorical difference between quantum and classical observables, inevitably leads to some unintuitive (for a classical observer) experimental results.

One key example is observed in the following scenario. Consider an experiment in which, at some point in the sequence, instead of the device measuring the observable F_j , the experimenter deploys an \overline{F}_j -device that measures the same observable but with less definition, that is, a *coarse*grained variant of the fine-grained F_j -device. In more formal terms, the coarse-graining can be described as follows: given any $\overline{f} \in \Omega(\overline{F}_j)$, there exists a subset $\omega(\overline{f}) \subseteq \Omega(F_j)$ such that, if the F_j -device reads out any of the values $f \in \omega(\overline{f}) \subseteq \Omega(F_j)$, then, all else being equal, the \overline{F}_j -device deployed in place of the F_j -device displays the result $\overline{f} \in \Omega(\overline{F}_j)$. The relationship between the coarse- and fine-grained device is then summarized by the *resolution*,

$$\operatorname{Res}(\overline{F}_j|F_j) := \left\{ \omega(\overline{f}) \subseteq \Omega(F_j) \mid \overline{f} \in \Omega(\overline{F}_j), \bigsqcup_{\overline{f} \in \Omega(\overline{F}_j)} \omega(\overline{f}) = \Omega(F_j) \right\}.$$
(4.3)

As the concept of coarse-grained measurements will prove to be a crucial ingredient of various observations, we introduce here a short-hand notation that should make the subsequent analysis more transparent. Let $\omega(\bar{f}_j) = \{f'_j, f''_j, \dots, f^{(k_j)}_j\}$; then, for sequences of \bar{F}_j -device readouts, we write

$$\Omega(\overline{F}_n) \ni \overline{f}_n \equiv \left(f'_n \lor \dots \lor f_n^{(k_n)}, \dots, f'_1 \lor \dots \lor f_1^{(k_1)}\right) = \left(\bigvee_{\omega(\overline{f}_n)} f_n, \dots, \bigvee_{\omega(\overline{f}_1)} f_1\right) = \bigvee_{\omega(\overline{f}_n)} f_n. \quad (4.4)$$

Physically, the concept of coarse-grained device can be visualized with the following model of measurement. Suppose that the F-device operates by first analyzing the measured system into distinct components (e.g., magnetic field gradient splits the beam of spinful particles into spatially dislocated sub-beams), then detecting the presence of individual components (e.g., by scattering the light with short enough wavelength), and finally reversing the analysis step by recombining the components (e.g., inverse the magnetic field gradient). The coarse-grained \overline{F} -device would perform the same pre-detection analysis and post-detection synthesis, but its actual detection method would count a number of components as one individual (e.g., by using light with a longer wavelength).

An experimenter can exploit the classical-like causality property (observation 1) to test whether the given \overline{F} -device does indeed operate as a coarse-grained version of the *F*-device—i.e., whether the resolution (4.3) can be attributed to the device—by verifying if the corresponding probability distributions satisfy the condition

$$\sum_{f \in \omega(\bar{f})} P_t^F(f) = P_t^{\bar{F}}(\bar{f}) \quad \text{for every } \bar{f} \in \Omega(\bar{F}) \text{ and } t > 0.$$
(4.5)

However, this test only checks for a necessary, and not sufficient, condition. To understand this point, let us consider how coarse-grained measurements are achieved for classical observables. If we denote the classical coarse-grained observable as \tilde{F} , then, since the trajectory representing \tilde{F} is a function of the trajectory of F (which, in turn, is a function of the elementary trajectory), the effect of coarse-graining is obtained by setting the filter functions in Eq. (4.2) to $p_{\tilde{t}}^{\tilde{F}}(\bar{f}|f) = \sum_{f' \in \omega(\bar{f})} \delta_{f,f'}$, so that

$$P_{\boldsymbol{t}_n}^{\widetilde{F}}(\overline{\boldsymbol{f}}_n) = \sum_{\boldsymbol{f}_n} P_{\boldsymbol{t}_n}^F(\boldsymbol{f}_n) \prod_{j=1}^n p_{\boldsymbol{t}_j}^{\widetilde{F}}(\overline{\boldsymbol{f}}_j | \boldsymbol{f}_j) = \sum_{\boldsymbol{f}_n} P_{\boldsymbol{t}_n}^F(\boldsymbol{f}_n) \prod_{j=1}^n \sum_{\boldsymbol{f}_j' \in \omega(\overline{\boldsymbol{f}}_j)} \delta_{\boldsymbol{f}_j, \boldsymbol{f}_j'} = \sum_{\boldsymbol{f}_n \in \omega(\overline{\boldsymbol{f}}_n)} P_{\boldsymbol{t}_n}^F(\boldsymbol{f}_n).$$
(4.6)

Not only such a device satisfies the necessary condition (4.5), it is also the only *classical* device that does. Consequently, when one performs a sequential measurement of a classical observable, the readout of the coarse-grained \tilde{F} -device is equivalent to the *alternative* of readouts of the fine-grained variant of the device,

$$P_{t_n,\dots,t_j,\dots,t_1}^{F,\dots,F}\left(f_n,\dots,\bigvee_{\omega(\bar{f}_j)}f_j,\dots,f_1\right) = \sum_{f_j\in\omega(\bar{f}_j)}P_{t_n,\dots,t_j,\dots,t_1}^{F,\dots,F}(f_n,\dots,f_j,\dots,f_1).$$
 (4.7)

Such a *faux*-coarse-grained device can also be implemented for quantum observables: the quantum version of the \tilde{F} -device works by first performing the actual measurement with the fine-grained F-device, and then displaying the post-processed results to the experimenter; formally, this amounts

to using probability distribution of the form (4.2), in which the filter function describes the postprocessing. With the post-processing set to $p_t^{\widetilde{F}}$, the result of a sequential measurement using \widetilde{F} -device would be described by Eq. (4.6).

However, since quantum observables are not constrained by the classical trajectory picture, it is physically possible (or rather, it is not forbidden as a matter of principle) to construct a quantum coarse-graining \overline{F} -device which, on one hand, satisfies condition (4.5), and on the other hand, cannot be described by a distribution of the form (4.6). To phrase it less formally: A proper quantum coarse-grained \overline{F} -device must perform its measurement via physical processes that are distinct from those utilized by the *F*-device. Otherwise, the action of the \overline{F} -device would be physically equivalent to the action of the *F*-device, modulo some form of post-processing of the obtained data, which would render it a faux-coarse-grained device instead.

Phenomenological observation 3 (Quantum coarse-grained measurement). When a proper quantum coarse-graining \overline{F}_j -device, characterized by a resolution $\operatorname{Res}(\overline{F}_j|F_j)$, is deployed mid sequence of measurements at time $t_j < t_n$, we observe that

$$P_{t_n,\ldots,t_j,\ldots,t_1}^{F_n,\ldots,F_1}\left(f_n,\ldots,\bigvee_{\omega(\overline{f}_j)}f_j,\ldots,f_1\right)-\sum_{f_j\in\omega(\overline{f}_j)}P_{t_n}^{F_n}(f_n)\neq 0,$$

that is, the readout of a coarse-grained device is *not* equivalent to the alternative of fine-grained readouts. However, the classical picture always persists for the latest readout of the sequence due to causality (observation 1),

$$P_{t_n,t_{n-1},\ldots,t_1}^{\overline{F}_n,F_n,\ldots,F_1}\left(\bigvee_{\omega(\overline{f}_n)}f_n,f_{n-1},\ldots,f_1\right)=\sum_{f_n\in\omega(\overline{f}_n)}P_{t_n}^{F_n}(f_n).$$

We finally remark that, at least in some cases, it is possible to manipulate the physical construction of a quantum coarse-grained device to effectively adjust its resolution. In the example discussed previously, such a manipulation would amount to varying the wavelength of the light used for detection—the longer the wavelength, the more coarse-grained the device. This idea can be pushed to its logical extreme, where we observe the quantum analogue of the classical consistency relation:

Phenomenological observation 4 (Extreme quantum coarse-graining). Let the \overline{F}_j -device be an extremely coarse-grained variant of the F_j -device, so that it is incapable of resolving between any alternatives: $\operatorname{Res}(\overline{F}_j|F_j) = \{\Omega(F_j)\}$. Then, deploying this device at any time in the sequence yields the same results as if no device was deployed at that time,

$$P_{t_n,\dots,t_j,\dots,t_1}^{F_n,\dots,F_1}\left(f_n,\dots,\bigvee_{\Omega(F_j)}f_j,\dots,f_1\right) = P_{t_n,\dots,t_j,\dots,t_1}^{F_n,\dots,F_j,\dots,F_1}(f_n,\dots,f_j,\dots,f_1)$$

Given that quantum coarse-grained devices can be built with adjustable resolutions, we will generally use the symbol \overline{F} to represent a *class* of measuring devices rather than a specific device. Members of this class are all coarse-grained variants of the *F*-device, with each device characterized by a particular resolution. From this point onward, unless otherwise stated, when we write $P_{\dots, \overline{f_j}, \dots}^{\dots, \overline{F_j}, \dots}(\dots \bigvee_{\omega(\overline{f_j})} f_j \dots)$, we mean that at time t_j , the experimenter deployed one of the devices from the coarse-grained class, with the resolution partially defined by the argument $\bigvee_{\omega(\overline{f_j})} f_j$. Although this notation does not fully specify the resolution of the deployed device—since it provides only one element, $\omega(\overline{f_j}) \in \operatorname{Res}(\overline{F_j}|F_j)$ —in most cases, this will be sufficient for our purposes. If circumstances require an unambiguous determination of the resolution, it will be explicitly stated.

4.2 Quantum interference

The effect revealed by the coarse-grained measurements described in observation 3, consisting in quantum observable not conforming to the classical picture of a trajectory tracing through alternative routes, is an example of the emblematic phenomenon of *quantum interference*. Indeed, experiment in which a coarse-grained \overline{F} -device is deployed in place of the fine-grained F-device can be seen as an instance of a k-slit Young experiment analyzed as sequential measurements.

Traditionally, a k-slit experiment is presented as involving only a single measurement at its terminus [20]. The slits are then viewed as "alternative routes" the measured system can take to reach the measuring device that concludes the experiment; the quantum interference between those alternatives is pointed to as the cause of the measured interference pattern. The k-slit experiment is contrasted with a hypothetical scenario in which the experimenter knows—or could even potentially know—which slit the system has passed through on its route to the measuring device. It is then asserted that possessing such a knowledge (or even the potential to acquire it) would prevent the interference from occurring, allowing the results to be described as a sum of the probabilities of each alternative.

In the treatment presented here, this textbook summary of interference experiments can be reformulated with a more precise meaning. First, the notion of the experimenter "possessing the knowledge of which slit was taken" corresponds precisely to performing a fine-grained measurement with the F_j -device, with the result $f_j \in \omega(\bar{f}_j) \subset \Omega(F_j)$ indicating the slit taken. The "potential to know", on the other hand, corresponds to the measurement with a faux-coarse-grained \tilde{F}_j -device, which was discussed in Section 4.1. In both scenarios there is, indeed, no interference effect. In the latter case, this is obvious from Eq. (4.7). In the former case, the event corresponding to the experimenter performing measurements while knowing that f_j was one of the values in $\omega(\bar{f}_j)$ is represented by the set $\{(f_n, \ldots, f_j, \ldots, f_1) \mid f_j \in \omega(\bar{f}_j)\} = \bigsqcup_{f_j \in \omega(\bar{f}_j)} \{f_n\}$, and thus

$$P_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}\Big(\bigsqcup_{f_j\in\omega(\bar{f}_j)}\{\boldsymbol{f}_n\}\Big) = \sum_{f_j\in\omega(\bar{f}_j)} P_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n).$$
(4.8)

Finally, since the quantum coarse-grained \overline{F}_j -device, for which we observe interference, is neither an F_j -device nor a \widetilde{F}_j -device, deploying this device cannot count as "acquiring knowledge" or "having the potential to know" which slit the system passed through. This formulation also underscores the non-classical nature of the effect, as argued in Section 4.1, where it is demonstrated that the proper coarse-grained \overline{F}_j -device is a quintessentially quantum concept with no direct classical analogue.

Further experimentation with quantum coarse-grained devices reveals a crucial feature of quantum interference:

Phenomenological observation 5 (Pair-wise quantum interference). The results of a k-slit experiment are fully explained with the results of double-slit experiments, indicating that the quantum interference occurs only between *pairs* of alternatives. In particular, it is found that the results of experiments involving sequential measurements with quantum coarse-grained devices can be described using the results obtained with devices that coarse-grain, at most, pairs of fine-grained outcomes.

It is found that the explicit decomposition of an arbitrary probability distribution into a combination of distributions involving, at most, pair-wise coarse-graining is obtained by applying the following recurrence rule:

$$P_{\dots t_j \dots}^{\dots \overline{F}_j \dots} \left(\dots \bigvee_{\omega(\overline{f}_j)} f_j \dots \right) = \sum_{\substack{f_j \in \omega(\overline{f}_j)}} P_{\dots t_j \dots}^{\dots F_j \dots} (\dots f_j \dots)$$

+
$$\sum_{\substack{f_j^{\pm} \in \omega(\overline{f}_j) \\ f_j^{\pm} \neq f_j^{-}}} \frac{1}{2} \Big[P_{\dots t_j \dots}^{\dots \overline{F}_j \dots} (\dots f_j^{\pm} \vee f_j^{-} \dots) - P_{\dots t_j \dots}^{\dots F_j \dots} (\dots f_j^{\pm} \dots) - P_{\dots t_j \dots}^{\dots F_j \dots} (\dots f_j^{\pm} \dots) \Big],$$

which is applicable for j < n; the rule for j = n case is described in observation 3.

The decomposition obtained through the recurrence described in observation 5 becomes untraceable when considering sequences with more than a couple of coarse-grained measurements. Therefore, instead of insisting on expressing everything only in terms of the phenomenological probability distributions, it is convenient to parameterize those probabilities with a formal element that allows one to display the structure of the decomposition with more transparency. To this end, we shall posit that the probability distributions can be written as

$$P_{\boldsymbol{t}_n}^{\boldsymbol{F_n}}(\boldsymbol{f}_n) \equiv I_{\boldsymbol{t}_n}^{\boldsymbol{F_n}}(\boldsymbol{f}_n, \boldsymbol{f}_n), \tag{4.9}$$

where the functions $I_{t_n}^{F_n}: \Omega(F_n) \times \Omega(F_n) \to \mathbb{R}$ are defined to be symmetric with respect to exchange of their arguments $f_n^+ \leftrightarrow f_n^-$,

$$I_{t_n}^{F_n}(f_n^+, f_n^-) = I_{t_n}^{F_n}(f_n^-, f_n^+).$$
(4.10)

We assume nothing about observables in (4.9), therefore the parameterization works for both fineand coarse-grained devices. However, to successfully recreate the recurrence rule from observation 5, function $I_{t_n}^{F_n}$ must also exhibit additivity with respect to the coarse-graining \vee treated as a formal operation on sequences, so that

$$P_{\boldsymbol{t}_n}^{\overline{\boldsymbol{F}}_n}(\overline{\boldsymbol{f}}_n) = I_{\boldsymbol{t}_n}^{\overline{\boldsymbol{F}}_n} \left(\bigvee_{\omega(\overline{\boldsymbol{f}}_n)} \boldsymbol{f}_n^+, \bigvee_{\omega(\overline{\boldsymbol{f}}_n)} \boldsymbol{f}_n^-\right) \equiv \sum_{\boldsymbol{f}_n^\pm \in \omega(\overline{\boldsymbol{f}}_n)} I_{\boldsymbol{t}_n}^{\overline{\boldsymbol{F}}_n}(\boldsymbol{f}_n^+, \boldsymbol{f}_n^-).$$
(4.11)

To demonstrate its veracity, we now apply this parameterization and recover the recurrence rule:

$$\sum_{f_{j} \in \omega(\bar{f}_{j})} P_{\dots t_{j}\dots}^{\dots F_{j}\dots}(\dots f_{j}\dots) + \sum_{f_{j}^{+} \in \omega(\bar{f}_{j})} \frac{1}{2} \Big[P_{\dots t_{j}\dots}^{\dots F_{j}\dots}(\dots f_{j}^{+} \vee f_{j}^{-}\dots) - P_{\dots t_{j}\dots}^{\dots F_{j}\dots}(\dots f_{j}^{+}\dots) - P_{\dots t_{j}\dots}^{\dots F_{j}\dots}(\dots f_{j}^{-}\dots) \Big] \\ = \sum_{f_{j} \in \omega(\bar{f}_{j})} I_{\dots t_{j}\dots}^{\dots F_{j}\dots}(\dots f_{j}\dots; \dots f_{j}\dots) + \sum_{f_{j}^{+} \in \omega(\bar{f}_{j})} \frac{1}{2} \Big[I_{\dots t_{j}\dots}^{\dots F_{j}\dots}(\dots f_{j}^{+}\dots; \dots f_{j}^{-}\dots) + I_{\dots t_{j}\dots}^{\dots F_{j}\dots}(\dots f_{j}^{-}\dots; \dots f_{j}^{+}\dots) \Big] \\ + \sum_{f_{j}^{+} \in \omega(\bar{f}_{j})} \frac{1}{2} \Big[I_{\dots t_{j}\dots}^{\dots F_{j}\dots}(\dots f_{j}^{+}\dots; \dots f_{j}^{-}\dots) + I_{\dots t_{j}\dots}^{\dots F_{j}\dots}(\dots f_{j}^{-}\dots; \dots f_{j}^{+}\dots) \Big] \\ = \sum_{f_{j}^{+} \in \omega(\bar{f}_{j})} I_{\dots t_{j}\dots}^{\dots F_{j}\dots}(\dots f_{j}^{+}\dots; \dots f_{j}^{-}\dots) = I_{\dots t_{j}\dots}^{\dots F_{j}\dots}(\dots \bigvee_{\omega(\bar{f}_{j})} f_{j}^{+}\dots; \dots \bigvee_{\omega(\bar{f}_{j})} f_{j}^{-}\dots)$$

$$(4.12)$$

The added value in using the functions $I_{t_n}^{F_n}$ is the possibility to reformulate the content of observation 5 as an explicit decomposition that reduces all instances of the overt coarse-grained measurements in terms of relatively fine-grained measurements.

It has been pointed out [21, 22] that the pair-wise interference between routing alternatives could be considered as a defining feature of the quantum theory. The parameterization (4.11) reflects this fact via the structure of the *interference terms* $I_{t_n}^{F_n}$ as functions of *pairs* of sequences, specifically. In principle, one could imagine a non-classical theory (i.e., a physical theory not describable within the trajectory picture) in which the interference also occurs between k > 2 alternatives. As we shall demonstrate later down the line, the fact that k = 2 in quantum mechanics has profound consequences for the whole structure of its formalism. Undoubtedly, the same would be true for hypothetical theories with k > 2, which would likely be even more "exotic" than quantum mechanics itself, provided they could exist as physical theories.

4.3 Quantum interference between independent systems

The analysis of quantum interference suggests that it is possible to base the formal description of experimental results on (non-positive valued) distributions on the space of pairs of sequences of outcomes, associated with a given quantum system—or, at the very least, with observables of that

system. Indeed, the family of interference terms $I_{t_n}^{F_n} : \Omega(F_n) \times \Omega(F_n) \to \mathbb{R}$ are such distributions, and they seem to be capable to formally describe the sequential measurement outcomes, including the interference effects manifesting in (sequential) k-slit experiments. This is certainly the case as long as *one* system is probed by the experimenter. We will now consider certain type of quantum interference in experiments where *multiple* systems are measured at the same time—as it turns out, the description with interference terms alone will be incomplete.

Although the concept of a system composed of a number of subsystems is an intuitive one, we should start by establishing a *phenomenological* definition, that is, one that refers only to measurements: A system will be said to be *partitionable* (into subsystems) when it is possible to deploy at the same time the distinct measuring devices that could also be deployed individually to measure the corresponding observables one at the time. If this is the case, then each of those observables can be attributed to one of the subsystems the system is partitionable into.

Some clarifying comments are in order, in case one cannot fully immerse themselves in the role of someone completely ignorant of the standard formalism of quantum mechanics and take the above definition at the face value. The following points should be kept in mind when trying to compare the concept of partitionable system with the standard formal definition of composite system:

- (i) When deploying measuring devices, one has to take into account things like the physical size of the device and its structural integrity. In particular, the devices mentioned in the definition above must be built as separate constructs—otherwise they could not be deployed individually.
- (ii) The notion of "deployment of measuring devices at the same time" might seem to echo the notion of "simultaneous measurements of commuting observables" from the standard formulation; however, this is not the case. In accordance with the measurement-projector link in the standard formalism (we shall deduce such a link later in the paper), a measuring device corresponds to the partitioning of the Hilbert space of the system into the images of a complete set of orthonormal projectors. Therefore, when the device is deployed, it measures any observable whose eigenspaces (defined by the operator representing it in the standard formalism) overlap with the space partitioning of the device. Of course, "commuting observables" have the same eigenspaces, and thus, in this sense, one deployment measures all of them at the same time. However, performing the simultaneous measurement in the sense of the phenomenological definition means that two (or more) devices have to be deployed at the same time. In the case of commuting observables, this would mean deploying a number of identical copies of one device, which is impossible because they would have to occupy the same volume.
- (iii) A way to circumvent the issue of devices occupying the same physical space would be to augment them with a mechanism to spatially split and divert the system to make room for their deployment. However, the fact that one is able to split a system into parts that can then be simultaneously measured by a number of devices only enforces the idea that the system is partitionable.
- (iv) Coarse- and fine-grained observables are a special case of commuting observables: a device performing the simultaneous fine- and coarse-grained measurement is actually equivalent to fine-grained measurement, and thus, it counts as a faux-coarse-grained device.
- (v) Finally, it is possible that the proposed definition does not exactly match the notion of composite system from the standard formalism where it is defined as a system represented by a product of Hilbert spaces (whence the choice to use the name "partitionable" rather than "composite"). However, even if it were true, it would not be problematic, because this phenomenological definition leads to the definition of independent systems that does overlap with the notion from the standard formalism.

Now, suppose that the experimenter has access to a system partitionable into two subsystems, A and B. For brevity, we shall use the same symbols to denote the two observables that determine the system partitioning (as per the definition given above); and thus, we have an A- and a B-device that can be deployed simultaneously to measure observables belonging to the corresponding

subsystems. The results measured by those devices are then described by the sample spaces $\{a_n \mid a_j \in \Omega(A)\}$ and $\{b_n \mid b_j \in \Omega(B)\}$.

Phenomenological observation 6 (Simultaneous measurements of independent systems). Certain systems are found to be partitionable into *independent* subsystems. A trivial example would be a system for which the measurements defining the partitioning are performed in two remote laboratories.

The constituent subsystems of a partitionable system shall be considered as *independent* when the results of the measurements of any subsystem observable are indistinguishable from measurements of the same observable performed when the measured subsystem is the only present system. In terms of probability distributions, A and B are independent when

$$P_{\boldsymbol{t}_n}^{AB}(a_n \wedge b_n, \dots, a_1 \wedge b_1) = P_{\boldsymbol{t}_n}^{AB}(\boldsymbol{a}_n \wedge \boldsymbol{b}_n) = P_{\boldsymbol{t}_n}^{A}(\boldsymbol{a}_n)P_{\boldsymbol{t}_n}^{B}(\boldsymbol{b}_n),$$

where $a_j \wedge b_j$ indicate that the A-device and B-device were deployed simultaneously—an AB-device for short—at time t_j , and $P_{t_n}^{A/B}$ is the probability distribution describing the measurement with the A/B-device in an experiment where only one system (and thus one observable) is present.

As we can see, a distribution associated with a system composed of independent subsystems factorizes into distributions associated with individual subsystems. Nevertheless, the subsystemonly interference terms turn out to be insufficient to parameterize all probability distributions describing experiments involving quantum interference in systems partitionable into independent parts. To observe this, one has to deploy a coarse-grained \overline{AB} -device, a version of the device that measures A and B simultaneously but is incapable of making a perfect "which subsystem" distinction. Such a device must not be confused with the \overline{AB} -device, consisting in the deployment of the coarse-grained A- and B-only devices in tandem while the distinction between subsystems is always precise.

Phenomenological observation 7 (Quantum interference between independent systems). Given a quantum coarse-grained \overline{AB} -device with resolution

$$\operatorname{Res}(\overline{AB}|AB) = \Big\{ \omega(\overline{a \wedge b}) \subseteq \Omega(A) \times \Omega(B) \ \Big| \bigsqcup_{\overline{a \wedge b} \in \Omega(\overline{AB})} \omega(\overline{a \wedge b}) = \Omega(A) \times \Omega(B) \Big\},$$

one finds that quantum observables A and B of independent subsystems interfere with each other when the \overline{AB} -device, which is incapable of a perfect distinction between subsystems, is deployed in place of an AB-device that resolves subsystems perfectly,

$$\begin{split} I^{\dots AB\dots}_{\dots t_j\dots} &(\dots a_j^+ \wedge b_j^+ \dots; \dots a_j^- \wedge b_j^- \dots) \\ &= \frac{1}{2} \Big[P^{\dots \overline{AB}\dots}_{\dots t_j\dots} \Big(\dots (a_j^+ \wedge b_j^+) \vee (a_j^- \wedge b_j^-) \dots \Big) \\ &\quad - P^{\dots AB\dots}_{\dots t_j\dots} (\dots a_j^+ \wedge b_j^+ \dots) - P^{\dots AB\dots}_{\dots t_j\dots} (\dots a_j^- \wedge b_j^- \dots) \Big] \\ &= \frac{1}{2} \Big[P^{\dots \overline{AB}\dots}_{\dots t_j\dots} \Big(\dots (a_j^+ \wedge b_j^+) \vee (a_j^- \wedge b_j^-) \dots \Big) \\ &\quad - P^{\dots A\dots}_{\dots t_j\dots} (\dots a_j^+ \dots) P^{\dots B\dots}_{\dots t_j\dots} (\dots b_j^+ \dots) - P^{\dots A\dots}_{\dots t_j\dots} (\dots a_j^- \dots) P^{\dots B\dots}_{\dots t_j\dots} (\dots b_j^- \dots) \Big] \\ &\neq 0. \end{split}$$

However, even though subsystems A and B are independent, this interference cannot be fully explained exclusively in terms of subsystem interference terms. In formal terms, the *co-interference* term

$$\Phi_{\dots t_j \dots}^{\dots AB \dots} (\dots a_j^+ \wedge b_j^+ \dots ; \dots a_j^- \wedge b_j^- \dots) := I_{\dots t_j \dots}^{\dots AB \dots} (\dots a_j^+ \wedge b_j^+ \dots ; \dots a_j^- \wedge b_j^- \dots) - I_{\dots t_j \dots}^{\dots A\dots} (\dots a_j^+ \dots ; \dots a_j^- \dots) \times I_{\dots t_j \dots}^{\dots B\dots} (\dots b_j^+ \dots ; \dots b_j^- \dots),$$

is found to be non-zero in general.

The following observation confirms that the factorization rule has not been violated here. It also provides clues about the form of the actual distribution that can correctly parameterize the "which subsystem" interference experiments.

Phenomenological observation 8 (Interference between identical independent systems). Consider the "which subsystem" interference experiment carried out on a system partitionable into two independent and otherwise identical subsystems. Let the measurement causing the interference effect be performed with an \overline{FF} -device, which is a coarse-grained variant of the FF-device which simultaneously measures the same observable F in both subsystems (in particular, F = A or F = B).

Then, when the co-interference terms $\Phi_{\dots t_j \dots}^{\dots AA\dots}$ and $\Phi_{\dots t_j \dots}^{\dots BB\dots}$ obtained from such experiments are compared with the co-interference term $\Phi_{\dots t_j \dots}^{\dots AB\dots}$, it is found that they satisfy the following relation:

$$\begin{split} |\Phi_{\dots,a_j}^{\dots,AB\dots}(\dots a_j^+ \wedge b_j^+ \dots ; \dots a_j^- \wedge b_j^- \dots)| \\ &= \sqrt{|\Phi_{\dots,a_j}^{\dots,AA\dots}(\dots a_j^+ \wedge a_j^+ \dots ; \dots a_j^- \wedge a_j^- \dots)|} \sqrt{|\Phi_{\dots,a_j}^{\dots,BB\dots}(\dots b_j^+ \wedge b_j^+ \dots ; \dots b_j^- \wedge b_j^- \dots)|}. \end{split}$$

This shows that, for independent subsystems, the co-interference term indeed factorizes into a product of subsystem distributions.

Moreover, it is found that the co-interference term for a system composed of two identical subsystems satisfies the following property:

$$0 \ge \Phi_{\dots t_j \dots}^{\dots FF \dots} (\dots f_j^+ \wedge f_j^+ \dots ; \dots f_j^- \wedge f_j^- \dots) = -\Phi_{\dots t_j \dots}^{\dots FF \dots} (\dots f_j^+ \wedge f_j^- \dots ; \dots f_j^- \wedge f_j^+ \dots).$$

When the subsystems are identical, the co-interference term must have the form of a *square* of a subsystem distribution. Therefore, the above relation indicates that this distribution is purely imaginary (the square is negative) and anti-symmetric function of the sequence pairs.

To account for observations 7 and 8, the interference term-based parameterization of probability distributions postulated in the previous section, cf. Eq. (4.9), has to be further expanded. The most straightforward choice is to introduce a complex-valued distributions of sequence pairs, $Q_{t_n}^{F_n}$: $\Omega(F_n) \times \Omega(F_n) \to \mathbb{C}$, that are hermitian functions with respect to exchange of their arguments $f_n^+ \leftrightarrow f_n^-$

$$Q_{t_n}^{F_n}(f_n^-, f_n^+) = Q_{t_n}^{F_n}(f_n^+, f_n^-)^*,$$
(4.13)

and exhibits additivity with respect to the coarse-graining operation \lor :

$$Q_{\boldsymbol{t}_n}^{\boldsymbol{\overline{F}}_n} \left(\bigvee_{\omega(\boldsymbol{\overline{f}}_n)} \boldsymbol{f}_n^+, \bigvee_{\omega(\boldsymbol{\overline{f}}_n)} \boldsymbol{f}_n^-\right) = \sum_{\boldsymbol{f}_n^\pm \in \omega(\boldsymbol{\overline{f}}_n)} Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n^+, \boldsymbol{f}_n^-).$$
(4.14)

By definition, we set their real parts to be equal to the corresponding interference term,

$$I_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n^+, \boldsymbol{f}_n^-) \equiv \operatorname{Re} Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n^+, \boldsymbol{f}_n^-), \qquad (4.15)$$

and their imaginary part to be set by the factorization of the co-interference term:

$$\Phi_{\dots t_j \dots}^{\dots AB \dots} (\dots a_j^+ \wedge b_j^+ \dots; \dots a_j^- \wedge b_j^- \dots) \equiv \operatorname{i} \operatorname{Im} Q_{\dots t_j \dots}^{\dots A\dots} (\dots a_j^+ \dots; \dots a_j^- \dots) \times \operatorname{i} \operatorname{Im} Q_{\dots t_j \dots}^{\dots B\dots} (\dots b_j^+ \dots; \dots b_j^- \dots);$$
(4.16)

$$\sqrt{\left|\Phi_{\dots f_j \dots}^{\dots FF \dots} (\dots f_j^+ \wedge f_j^+ \dots; \dots f_j^- \wedge f_j^- \dots)\right|} = \left|\operatorname{Im} Q_{\dots f_j \dots}^{\dots F \dots} (\dots f_j^+ \dots; \dots f_j^- \dots)\right|.$$
(4.17)

As the function $Q_{t_n}^{F_n}$ is hermitian and its real part is symmetric, the imaginary part is thus antisymmetric, which is consistent with observation 8.

We can now use this parameterization to rewrite the "which subsystem" interference term as a combination of products of distributions associated with independent subsystems,

$$P_{\dots t_j \dots}^{\dots \overline{AB} \dots} \left(\dots \bigvee_{\omega \left((\overline{a \wedge b})_j \right)} a_j \wedge b_j \dots \right)$$

$$= I_{\dots t_{j}\dots}^{\dots \overline{AB}\dots} \left(\dots \bigvee_{\omega((\overline{a \wedge b})_{j})} a_{j}^{+} \wedge b_{j}^{+} \dots; \dots \bigvee_{\omega((\overline{a \wedge b})_{j})} a_{j}^{-} \wedge b_{j}^{-} \dots \right)$$

$$= \sum_{(a_{j}^{\pm}, b_{j}^{\pm}) \in \omega((\overline{a \wedge b})_{j})} I_{\dots t_{j}\dots}^{\dots AB\dots} (\dots a_{j}^{+} \wedge b_{j}^{+} \dots; \dots a_{j}^{-} \wedge b_{j}^{-} \dots)$$

$$= \sum_{(a_{j}^{\pm}, b_{j}^{\pm}) \in \omega((\overline{a \wedge b})_{j})} \left[I_{\dots t_{j}\dots}^{\dots A\dots} (\dots a_{j}^{+} \dots; \dots a_{j}^{-} \dots) I_{\dots t_{j}\dots}^{\dots B\dots} (\dots b_{j}^{+} \dots; \dots b_{j}^{-} \dots) + \Phi_{\dots t_{j}\dots}^{\dots AB\dots} (\dots a_{j}^{+} \wedge b_{j}^{+} \dots; \dots a_{j}^{-} \wedge b_{j}^{-} \dots) \right]$$

$$= \sum_{\substack{(a_j^{\pm}, b_j^{\pm}) \in \omega((\overline{a \wedge b})_j)}} \left[\operatorname{Re} Q_{\dots t_j \dots}^{\dots A \dots}(\dots a_j^{+} \dots; \dots a_j^{-} \dots) \operatorname{Re} Q_{\dots t_j \dots}^{\dots B \dots}(\dots b_j^{+} \dots; \dots b_j^{-} \dots) - \operatorname{Im} Q_{\dots t_j \dots}^{\dots A \dots}(\dots a_j^{+} \dots; \dots a_j^{-} \dots) \operatorname{Im} Q_{\dots t_j \dots}^{\dots B \dots}(\dots b_j^{+} \dots; \dots b_j^{-} \dots) \right]$$

$$= \sum_{(a_j^{\pm}, b_j^{\pm}) \in \omega \left((\overline{a \wedge b})_j \right)} \operatorname{Re} \left\{ Q_{\dots t_j \dots}^{\dots A \dots} (\dots a_j^{+} \dots ; \dots a_j^{-} \dots) Q_{\dots t_j \dots}^{\dots B \dots} (\dots b_j^{+} \dots ; \dots b_j^{-} \dots) \right\}.$$
(4.18)

On the other hand, since the imaginary part is anti-symmetric we have

$$Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n, \boldsymbol{f}_n) = \operatorname{Re} Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n, \boldsymbol{f}_n) = I_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n, \boldsymbol{f}_n) = P_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n),$$
(4.19)

and thus

=

=

$$P_{\dots,t_{j},\dots}^{\dots,\overline{AB},\dots}\left(\dots\bigvee_{\omega\left((\overline{a\wedge b})_{j}\right)}a_{j}\wedge b_{j}\dots\right)$$

$$=Q_{\dots,t_{j},\dots}^{\dots,\overline{AB},\dots}\left(\dots\bigvee_{\omega\left((\overline{a\wedge b})_{j}\right)}a_{j}^{+}\wedge b_{j}^{+}\dots;\dots\bigvee_{\omega\left((\overline{a\wedge b})_{j}\right)}a_{j}^{-}\wedge b_{j}^{-}\dots\right)$$

$$=\sum_{(a_{j}^{\pm},b_{j}^{\pm})\in\omega\left((\overline{a\wedge b})_{j}\right)}Q_{\dots,t_{j},\dots}^{\dots,AB,\dots}(\dots,a_{j}^{+}\wedge b_{j}^{+}\dots;\dots,a_{j}^{-}\wedge b_{j}^{-}\dots)$$

$$=\sum_{(a_{j}^{\pm},b_{j}^{\pm})\in\omega\left((\overline{a\wedge b})_{j}\right)}\operatorname{Re}Q_{\dots,t_{j},\dots}^{\dots,AB,\dots}(\dots,a_{j}^{+}\wedge b_{j}^{+}\dots;\dots,a_{j}^{-}\wedge b_{j}^{-}\dots),$$
(4.20)

and the last equality is, again, due to the anti-symmetry of the imaginary part. Comparing the two decompositions of $P_{\dots t_j \dots}^{\dots \overline{AB} \dots}$, we find the following crucial result: the distributions $Q_{t_n}^{AB}$ obey the factorization law when A and B are independent,

$$Q_{t_n}^{AB}(a_n^+ \wedge b_n^+, a_n^- \wedge b_n^-) = Q_{t_n}^A(a_n^+, a_n^-)Q_{t_n}^B(b_n^+, b_n^-).$$
(4.21)

In this way, we recover the factorization property for independent systems, while retaining the ability to parameterize the quantum interference effects in partitionable systems.

4.4 Markovianity and the initialization events

We shall now turn our attention to the special case of *perfectly fine-grained* devices—that is, devices that are not a coarse-grained variant of any other measuring device; we shall reserve symbols K, L, \ldots for such devices, while leaving F, G, \ldots to indicate generic devices that could be of any degree of graining.

Phenomenological observation 9 (Markovianity). The probability distributions describing measurements with a perfectly fine-grained K-device factorize as follows,

$$P_{t_n}^K(k_n) = P_{t_n|t_{n-1}}^{K|K}(k_n|k_{n-1}) \cdots P_{t_2|t_1}^{K|K}(k_2,|k_1) P_{t_1}^K(k_1), \qquad (4.22)$$

where the factors $P_{t_j|t_{j-1}}^{K|K}(k_j|k_{j-1})$ are *conditional probabilities*, as defined by the Bayes law,

$$P_{t|t_n,\dots,t_1}^{K|K}(k|k_n\dots,k_1) := \frac{P_{t,t_n,\dots,t_1}^K(k,k_n,\dots,k_1)}{P_{t_n,\dots,t_1}^K(k_n,\dots,k_1)}.$$
(4.23)

In probability theory, the property (4.22) is called *Markovianity* and is traditionally interpreted as a "lack of memory" because

$$P_{t|t_{n},\dots,t_{1}}^{K|K}(k|k_{n},\dots,k_{1}) = \frac{P_{t|t_{n}}^{K|K}(k|k_{n})P_{t_{n}|t_{n-1}}^{K|K}(k_{n}|k_{n-1})\cdots P_{t_{2}|t_{1}}^{K|K}(k_{2}|k_{1})P_{t_{1}}^{K}(k_{1})}{P_{t_{n}|t_{n-1}}^{K|K}(k_{n}|k_{n-1})\cdots P_{t_{2}|t_{1}}^{K|K}(k_{2}|k_{1})P_{t_{1}}^{K}(k_{1})} = P_{t|t_{n}}^{K|K}(k|k_{n}),$$
(4.24)

that is, the statistics of the result depend only on the latest prior readout, not on the whole history of all previous readouts.³

It is important to reiterate that Markovianity is inherently linked with the fine-graining of the measurement device. One can easily verify that the property is lost when the deployed devices have less than perfect definition, that is, when a coarse-grained \overline{K} -device is deployed in place of the K-device, e.g.,

$$P_{t_{2},t_{1}}^{K,K}(k_{2},k_{1}^{+}\vee k_{1}^{-}) = P_{t_{2},t_{1}}^{K}(k_{2},k_{1}^{+}) + P_{t_{2},t_{1}}^{K}(k_{2},k_{1}^{-}) + 2\operatorname{Re}Q_{t_{2},t_{1}}^{K}(k_{2},k_{2};k_{1}^{+},k_{1}^{-}) \\ = P_{t_{2}|t_{1}}^{K|K}(k_{2}|k_{1}^{+})P_{t_{1}}^{K}(k_{1}^{+}) + P_{t_{2}|t_{1}}^{K|K}(k_{2}|k_{1}^{-})P_{t_{1}}^{K}(k_{1}^{-}) + 2\operatorname{Re}Q_{t_{2},t_{1}}^{K}(k_{2},k_{2};k_{1}^{+},k_{1}^{-}) \\ \neq P_{t_{2}|t_{1}}^{K|\overline{K}}(k_{2}|k_{1}^{+}\vee k_{1}^{-})P_{t_{1}}^{\overline{K}}(k_{1}^{+}\vee k_{1}^{-}).$$

$$(4.25)$$

This is in line with the traditional notion of Markovianity: the property is known to be destroyed by coarse-graining [23].

So far we have avoided drawing the attention to the fact that the description in terms of probability distributions for the sequences of readouts characterized in Section 3 is not yet complete. The missing piece is the control over the *initial condition* of the experiment; this is, of course, a crucial element that allows the experimenter to reset the physical "state" of the system, so that the results of the subsequent measurements can be meaningfully compared with the results obtained in previous (and future) runs of the experiment. Therefore, all multitime probability distributions we have been considering so far should actually be interpreted as probabilities conditioned by some form of "initialization event",

$$P_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n) \to P_{\boldsymbol{t}_n|0}^{\boldsymbol{F}_n}(\boldsymbol{f}_n| \text{ initialization})$$
 (4.26)

One could imagine many different ways to realize this "initialization event" but, thanks to Markovianity, it is possible to define it in the terms of readouts of the measuring device; for example, if we consider the readout $k_0 \in \Omega(K)$ of a perfectly fine-grained K-device deployed at time t = 0 as the initialization, then we have

$$P_{\boldsymbol{t}_{n}|0}^{\boldsymbol{F}_{n}|\boldsymbol{K}}(\boldsymbol{f}_{n}|\boldsymbol{k}_{0}) = P_{\boldsymbol{t}_{n}|0,-s_{1},-s_{2},\ldots}^{\boldsymbol{F}_{n}|\boldsymbol{K},\boldsymbol{G},\boldsymbol{H},\ldots}(\boldsymbol{f}_{n}|\boldsymbol{k}_{0},\boldsymbol{g}_{-1},\boldsymbol{h}_{-2},\ldots) = P_{\boldsymbol{t}_{n}|0,-s_{1}',-s_{2}',\ldots}^{\boldsymbol{F}_{n}|\boldsymbol{K},\boldsymbol{G}',\boldsymbol{H}',\ldots}(\boldsymbol{f}_{n}|\boldsymbol{k}_{0},\boldsymbol{g}_{-1}',\boldsymbol{h}_{-2}',\ldots) = \dots$$

$$(4.27)$$

Therefore, this t = 0 readout of the K-device does indeed work as an initialization, since everything that happened to the system before that time cannot influence the subsequent measurements. Moreover, the initialization procedure defined in this way can be generalized to include a degree of uncertainty about the initializing measurement,

$$P_{\boldsymbol{t}_{n}|0}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n}|p) := \sum_{K} \sum_{k \in \Omega(K)} P_{\boldsymbol{t}_{n}|0}^{\boldsymbol{F}_{n}|K}(\boldsymbol{f}_{n}|k)p(K,k) \quad \text{such that } \sum_{K,k} p(K,k) = 1, \ p(K,k) \ge 0, \quad (4.28)$$

³We remark that the "lack of memory" should not be confused with "short memory". Indeed, "short memory" suggests that the statistics of the readout at t would depend only on previous events that occurred in the immediate past—i.e., the previous readouts have impact only when $|t - t_i|$ are in some sense small, while those readouts for which $|t - t_i|$ is large can be disregarded. This is not the case here, as the timings in (4.24) form an arbitrary chronological sequence, so it does not matter whether $t - t_n$ is large or small.

where p(K,k) denotes the probability that the experiment was initialized with a perfectly finegrained K-device that measured the result $k \in \Omega(K)$. From this point onward, when we write $P_{t_n}^{F_n}(f_n)$ without explicit indication of the initial condition, this has to be understood as a shorthand for $P_{t_n|0}^{F_n}(f_n|p)$ conditioned on a certain fixed distribution p(K,k).

4.5 Quantum Zeno effect

Consider an experiment in which some perfectly fine-grained K-devices are deployed one after the other with a small delay between the measurements. An intuitive expectation is that, in the limit in which the delay between consecutive measurements is negligible, the second readout should be the same as the first. This is indeed what is observed in experiments; one finds⁴

$$\lim_{\Delta t \to 0^+} P_{t+\Delta t|t}^{K|K}(k|k_0) = \delta_{k,k_0}.$$
(4.29)

Correlating the K-device with its coarse-grained variant—the \overline{K} -device—also yields an intuitive result,

$$\lim_{\Delta t \to 0^+} P_{t+\Delta t|t}^{\overline{K}|K}(\overline{k}|k_0) = \sum_{k \in \omega(\overline{k})} \delta_{k,k_0}.$$
(4.30)

However, these intuitions are based on the trajectory picture of classical theories: we assume that a trajectory traced by a classical observable is continuous, and thus, its samples taken within a small interval of time have to overlap. Therefore, it is nigh inevitable that quantum observables, for which the trajectory picture is invalid, will subvert expectations based on classical intuitions when one strings a frequent probing into sequences consisting of more than two measurements. Arguably, the most prominent example of such an intuition-defying result is the emergence of the *quantum Zeno effect* in experiments in which the measuring device is deployed with increasing frequency:

Phenomenological observation 10 (Quantum Zeno effect). Given a perfectly fine-grained K-device and a duration t, one observes

$$\lim_{n \to \infty} P_{\boldsymbol{s}_n|0}^{K|K}(k_0, \dots, k_0|k_0) = \lim_{n \to \infty} \prod_{j=0}^{n-1} P_{s_j + \frac{t}{n}|s_j}^{K|K}(k_0|k_0) = 1 \quad \text{where } s_j = jt/n.$$
(4.31)

That is, the device readouts becomes "frozen" in its initial value as the probing frequency tends to infinity. To put it differently, the probability that one measures any changes in the observable decreases as the frequency at which it is measured increases.

When combined with Markovianity, the occurrence of Zeno effect provides some insight regarding the laws that govern the dynamics of the measured observable. In particular, we can conclude from observation 10 that the *survival probability* must have the following short-time behavior,

$$P_{t+\Delta t|t}^{K|K}(k|k) \xrightarrow{\Delta t \to 0^+} 1 - v(k,t)^2 \Delta t^2 + O(\Delta t^3).$$

$$(4.32)$$

Indeed, if the linear term did not vanish, then the Zeno effect would be impossible,

$$\prod_{j=0}^{n-1} P_{s_j + \frac{t}{n} | s_j}^{K|K}(k_0 | k_0) = \prod_{j=0}^{n-1} \left(1 + L(k_0, s_j) \frac{t}{n} + O(n^{-2}) \right) \xrightarrow{n \to \infty} e^{\int_0^t L(k_0, s) ds} \neq 1.$$
(4.33)

⁴Phenomenological laws like (4.29) cannot be confirmed in an actual real-life experiment. The deployment of a physical measuring device must take some finite amount of time, so the delay Δt between consecutive measurements cannot ever be zero. Therefore, the limit $\Delta t \rightarrow 0^+$ is to be understood as an idealization where it is assumed that, in general, physical quantities have regular behavior, so that it is possible to extrapolate trends that were actually observed for small, but still finite, Δt . In formal terms, unless it is stated otherwise, we always assume that any phenomenological function $\varphi(t)$ possesses a well-behaved Taylor series expansion, $\varphi(t + \Delta t) = \varphi(t) + \varphi'(t)\Delta t + (1/2)\varphi''(t)\Delta t^2 + O(\Delta t^3)$.

It is important to remark that the Zeno effect is a quintessential quantum (or, in any case, nonclassical) effect, as it is incompatible with the trajectory picture. By definition, the measurements of a classical observable K_{cls} are described with consistent multi-time probabilities, i.e.,

$$P_{t_n}^{K_{\text{cls}}}(\boldsymbol{k}_n) = \sum_{\kappa \in \Omega(K_{\text{cls}})} P_{t_n,\dots,t_j,s,t_{j-1},\dots,t_1}^{K_{\text{cls}}}(k_n,\dots,k_j,\kappa,k_{j-1},\dots,k_1)$$

$$= \sum_{\kappa,\kappa' \in \Omega(K_{\text{cls}})} P_{\dots,t_j,s,t_{j-1},\dots,t_{j'},s',t_{j'-1},\dots}^{K_{\text{cls}}}(\dots,k_j,\kappa,k_{j-1},\dots,k_{j'},\kappa',k_{j'-1},\dots) = \dots$$
(4.34)

If we were to assume that K_{cls} exhibits the Zeno effect, then, by using consistency, we would conclude that, for any sequence of measurements, one gets

$$P_{\boldsymbol{t}_{n}|0}^{K_{\text{cls}}}(\boldsymbol{k}_{n}|k_{0}) = \lim_{N \to \infty} \sum_{\kappa_{j}: j \notin \iota_{N}(\{1,...,n\})} P_{\boldsymbol{s}_{N}|0}^{K_{\text{cls}}}(\boldsymbol{\kappa}_{N}|k_{0}) = \prod_{j=1}^{n} \delta_{k_{j},k_{0}}, \qquad (4.35)$$

with $\iota_N : \{1, \ldots, n\} \to \{1, \ldots, N\}$ such that $s_{\iota(j)} = t_j$ and $\kappa_{\iota(j)} = k_j$ for $j = 1, \ldots, n$. Therefore, in classical theories only trivial observables—that is, those that are "frozen" because they have no dynamics in the first place—can behave as if they exhibited Zeno effect.

4.6 Multiple observables and uncertainty relations

We previously noticed (see Section 4.1) that, due to the trajectory picture being in effect, any measurable quantity of a classical system can be formally represented as a function of a trajectory representing the elementary system observable. Generally, the elementary observable has the form of vector-valued trajectory of dimension equal to the number of degrees of freedom, e.g., a position in the three-dimensional space, $E_{cls} : t \mapsto (r_x(t), r_y(t), r_z(t)) \in \mathbb{R}^3$. Let us consider here a simple case of classical system whose elementary observable is one-dimensional and discretevalued, $E_{cls} : t \mapsto e(t) \in \Omega(E_{cls})$ where $\Omega(E_{cls})$ is a finite set. Then, an observable $K_{cls} : t \mapsto$ $k(t) = K(e(t)) \in \Omega(K_{cls})$ can be considered as a classical analogue of a perfectly fine-grained observable provided that the function $K : \Omega(E_{cls}) \to \Omega(K_{cls})$ is a bijection (one-to-one function); in that case, measurements of K_{cls} are essentially equivalent to measurements of the elementary observable itself,

$$P_{\boldsymbol{t}_{n}}^{K_{\text{cls}}}(\boldsymbol{k}_{n}) = \int \left(\prod_{j=1}^{n} p_{t_{j}}^{K_{\text{cls}}}(k_{j}|e(t_{j}))\right) \mathcal{P}^{E_{\text{cls}}}[e][\mathcal{D}e] = \int \left(\prod_{j=1}^{n} \delta_{k_{j},K(e(t_{j}))}\right) \mathcal{P}^{E_{\text{cls}}}[e][\mathcal{D}e]$$
$$= P_{t_{n},\dots,t_{1}}^{E_{\text{cls}}}(K^{-1}(k_{n}),\dots,K^{-1}(k_{1})) = P_{\boldsymbol{t}_{n}}^{E_{\text{cls}}}(K^{-1}(\boldsymbol{k}_{n})).$$
(4.36)

Therefore, in classical theories there is little incentive to ever make any distinctions between observables; ultimately, all fine-grained measurements yield essentially the same information. We can summarize this observation with the following *certainty relations*,

$$\delta_{e,e_{0}} = \lim_{\Delta t \to 0^{+}} P_{t+\Delta t|t}^{E_{cls}|E_{cls}}(e|e_{0})$$

$$= \lim_{\Delta t \to 0^{+}} P_{t+\Delta t|t}^{K_{cls}|K_{cls}}(K(e)|K(e_{0})) = \lim_{\Delta t \to 0^{+}} P_{t+\Delta t|t}^{L_{cls}|L_{cls}}(L(e)|L(e_{0}))$$

$$= \lim_{\Delta t \to 0^{+}} P_{t+\Delta t|t}^{K_{cls}|L_{cls}}(K(e)|L(e_{0})) = \lim_{\Delta t \to 0^{+}} P_{t+\Delta t|t}^{L_{cls}|K_{cls}}(L(e)|K(e_{0})), \quad (4.37)$$

where K_{cls} and L_{cls} are any two perfectly fine-grained observables, and E_{cls} is the elementary observable of the classical system.

Since the trajectory picture is invalid for quantum systems, it is possible to have physically inequivalent quantum observables. Consequently, quantum observables are not constrained by the classical certainty relations. So far, we have investigated only a special kind of such inequivalent observables, namely, an arbitrarily chosen observable F and its various (quantum) coarse-grained variants \overline{F} (cf. Section 4.2, Section 4.3). However, even observables measured with perfectly finegrained devices can be inequivalent: **Phenomenological observation 11** (Uncertainty relations). Given a pair of perfectly finegrained devices (K- and L-device), one observes the following relation when the two devices are deployed in a rapid succession:

$$\lim_{\Delta t \to 0^+} P_{t+\Delta t|t}^{K|L}(k|l) = \lim_{\Delta t \to 0^+} P_{t+\Delta t|t}^{L|K}(l|k) \equiv C_{k,l}^{K|L}, \quad \text{for any } t > 0,$$
(4.38)

where $C_{k,k'}^{K|K} = \delta_{k,k'}$ when L = K, but in general $C_{k,l}^{K|L} \neq \delta_{k,l}$, i.e., quantum observables can violate certainty relations.

Moreover, for every perfectly fine-grained K-device, it is always possible to find a perfectly fine-grained K^{\perp} -device such that

$$C_{k,k'}^{K|K^{\perp}} = |\Omega(K)|^{-1}, \text{ for every } k \in \Omega(K) \text{ and } k' \in \Omega(K^{\perp}).$$

$$(4.39)$$

Since $C_{k,l}^{K|L}$ does not depend on the time at which the measurements were taken, it can serve as a phenomenological measure of how "similar" the observables measured by the K- and L-devices are. On one extreme we have $C_{k,l}^{K|L} = \delta_{k,l}$, signifying that K and L are equivalent (analogously to all classical observables), while on the other end of the spectrum we have $C_{k,l}^{K|L} = |\Omega(K)|^{-1}$, meaning that there is no correlation whatsoever between observations of L and K.

It is important to underline that for classical observers there is no intuitive basis for how to interpret cases $C_{k,l}^{K|L} \neq \delta_{k,l}$. Indeed, since in classical physics there is essentially only one observable (and thus certainty relations are always true), it is impossible to extrapolate from the classical scenario to any situation in which there are two or more inequivalent observables. Any understanding of measurement, observation etc. acquired in classical setting has the existence of a single trajectory and the equivalence of all observables as foundational principles; there is simply no way to imagine how such a picture would be modified when observables are allowed to be inequivalent.

Part II Deduction of the formalism

In the second part of the paper, we demonstrate how to deduce a formalism of the theory from the phenomenology of quantum systems presented in the first part.

We begin by acknowledging that, in the case of quantum theory, it is impractical if not impossible—to construct a formalism that exclusively utilizes phenomenological quantities. Therefore, to advance the program, we base the emerging formalism on the complex-valued distributions on the space of sequence pairs. These distributions, initially introduced as a parameterization of the pair-wise interference decomposition of the phenomenological probability distributions, are now given a more fundamental role. Although these "bi-sequence" distributions are ostensibly an *ad hoc* inclusion, we argue that their addition is justified. This particular formal extension is both *minimal* and necessary, because its function cannot be reasonably fulfilled by the available phenomenological elements.

Consequently, the formalism takes shape as one that describes quantum system in terms of a complex-valued measure on the space of trajectory *pairs*. Thus, we arrive at an elegant and evocative explanation why quantum theory does not adhere to the classical trajectory picture: the *single*-trajectory picture is invalid in quantum mechanics, because, in reality, it is a *bi*-trajectory theory.

With the overall formal structure established, we then proceed to deduce the Hilbert space representation of the formalism. In this case, the connection to Hilbert space structures follows from the basic properties of bi-sequence distributions. However, the detailed form of the representation is found by referencing the previously analyzed phenomenological observations. Finding this representation is a crucial step, as it enables us to break down the formal elements into simple and comprehensible "building blocks." This is essential for model-building and, more broadly, for ensuring the theory's predictive power. A theory limited to describing experimental results without the capacity to propose and describe *potential* systems that have not yet been observed would be incomplete.

5 The bi-trajectory picture of quantum mechanics

5.1 Basis for the deduced formalism

The observations 5, 6 and 8 concerning various aspects of quantum interference described in Section 4 compelled us to consider a formal parameterization of phenomenological probability distributions based on a family of complex-valued distributions on the space of sequence pairs,

$$Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}: \Omega(\boldsymbol{F}_n) \times \Omega(\boldsymbol{F}_n) \to \mathbb{C}.$$
(5.1)

These distributions allowed us to succinctly describe the phenomenology of interference effects, and to conveniently encode the crucial observation that interference occurs only between pairs of alternatives within the very structure of the functions employed,

$$P_{\boldsymbol{t}_n}^{\overline{\boldsymbol{F}}_n}(\overline{\boldsymbol{f}}_n) = Q_{\boldsymbol{t}_n}^{\overline{\boldsymbol{F}}_n}(\overline{\boldsymbol{f}}_n, \overline{\boldsymbol{f}}_n) = \sum_{\boldsymbol{f}_n^{\pm} \in \omega(\overline{\boldsymbol{f}}_n)} Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n^+, \boldsymbol{f}_n^-) = \sum_{\boldsymbol{f}_n^{\pm} \in \omega(\overline{\boldsymbol{f}}_n)} \operatorname{Re} Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n^+, \boldsymbol{f}_n^-).$$
(5.2)

Furthermore, through a consistent application of the properties we have defined the functions $Q_{t_n}^{F_n}$ to possess, we proved that they obey the factorization rule for systems partitionable into independent subsystems [cf. Eq. (4.21)]. This is a key feature, without which it would be impossible to define a coherent formal description of isolated systems (i.e., a system independent of any other

system), and thus, any possibility of formulating a functioning formalism for the theory would be negated.

Consequently, the clear advantages of the distributions $Q_{t_n}^{F_n}$ makes them a prime candidate as the basis for the formalism that we strive to deduce from the phenomenological observations. However, concerns remain regarding whether the choice of $Q_{t_n}^{F_n}$ as the formal parameterization ingredient conflicts with the ethos of phenomenological quantum mechanics. To this regard we notice that, for the class of experiments analyzed so far, only a limited number of values $Q_{t_n}^{F_n}(f_n^+, f_n^-)$ can be considered as phenomenological quantities. These are (a) the diagonal values $Q_{t_n}^{F_n}(f_n, f_n)$, because they simply equal the measurable probabilities, and (b) the interference terms involved in the pair-wise decomposition in a *single k*-slit experiment,

$$\operatorname{Re} Q_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}}(f_{n},\ldots,f_{j}^{+},\ldots,f_{1};f_{n},\ldots,f_{j}^{-},\ldots,f_{1}) = \frac{1}{2} \Big[P_{t_{n},\ldots,t_{j},\ldots,t_{1}}^{F_{n},\ldots,F_{1}}(f_{n},\ldots,f_{j}^{+}\vee f_{j}^{-},\ldots,f_{1}) - P_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}}(\ldots,f_{j}^{+},\ldots) - P_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}}(\ldots,f_{j}^{-},\ldots) \Big].$$
(5.3)

On the other hand, as an alternative choice to distributions $Q_{t_n}^{F_n}$, which are generally nonphenomenological, we might consider functions defined as their symmetrized versions; namely,

$$\mathcal{S}[Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}](f_n^+, f_n^-; \dots; f_1^+, f_1^-) := \prod_{j=1}^n \left(\sum_{(\phi_j^+, \phi_j^-) \in S(\{f_j^+, f_j^-\})}\right) Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\phi_n^+, \dots, \phi_1^+; \phi_n^-, \dots, \phi_1^-), \quad (5.4)$$

where S(A) denotes the set of all permutations of elements of A, for example $S(\{f_j^+, f_j^-\}) = \{(f_j^+, f_j^-), (f_j^-, f_j^+)\}$ and $S(\{f_j, f_j\}) = \{(f_j, f_j)\}$. All such functions decompose into combinations of phenomenological probability distributions, thus making them phenomenological quantities; for example

$$\begin{split} \mathcal{S}[Q_{t_{n}}^{F_{n}}](f_{n}, f_{n}; \dots; f_{j}^{+}, f_{j}^{-}; \dots; f_{k}^{+}, f_{k}^{-}; \dots; f_{1}, f_{1}) \\ &= 2\operatorname{Re} Q_{t_{n}}^{F_{n}}(\dots, f_{j}^{+}, \dots, f_{j}^{+}, \dots; \dots, f_{j}^{-}, \dots, f_{k}^{-}, \dots) + 2\operatorname{Re} Q_{t_{n}}^{F_{n}}(\dots, f_{j}^{+}, \dots, f_{k}^{-}, \dots; \dots, f_{j}^{-}, \dots, f_{k}^{+}, \dots) \\ &= P_{\dots, t_{j} \dots t_{k} \dots}^{\dots, \overline{F}_{j} \dots, \overline{F}_{k} \dots}(\dots, f_{j}^{+} \vee f_{j}^{-}, \dots, f_{k}^{+} \vee f_{k}^{-}, \dots) \\ &- P_{\dots, t_{j} \dots t_{k} \dots}^{\dots, \overline{F}_{j} \dots, \overline{F}_{k} \dots}(\dots, f_{j}^{+} \vee f_{j}^{-}, \dots, f_{k}^{+}, \dots) - P_{\dots, t_{j} \dots, t_{k} \dots}^{\dots, \overline{F}_{j} \dots, \overline{F}_{k} \dots}(\dots, f_{j}^{+} \vee f_{j}^{-}, \dots, f_{k}^{-}, \dots) \\ &- P_{\dots, t_{j} \dots, t_{k} \dots}^{\dots, \overline{F}_{j} \dots, \overline{F}_{k} \dots}(\dots, f_{j}^{+} \dots, f_{j}^{+} \dots, f_{k}^{+} \vee f_{k}^{-}, \dots) - P_{\dots, t_{j} \dots, t_{k} \dots}^{\dots, \overline{F}_{j} \dots, \overline{F}_{k} \dots}(\dots, f_{j}^{-}, \dots, f_{k}^{+} \vee f_{k}^{-}, \dots) \\ &+ P_{\dots, t_{j} \dots, t_{k} \dots}^{\dots, F_{j} \dots, F_{k} \dots}(\dots, f_{j}^{+} \dots, f_{k}^{+} \dots) + P_{\dots, t_{j} \dots, t_{k} \dots}^{\dots, F_{j} \dots, F_{k} \dots}(\dots, f_{j}^{-}, \dots, f_{k}^{+} \dots) \\ &+ P_{\dots, t_{j} \dots, t_{k} \dots}^{\dots, F_{j} \dots, F_{k} \dots}(\dots, f_{j}^{+} \dots, f_{k}^{-} \dots) + P_{\dots, t_{j} \dots, t_{k} \dots}^{\dots, F_{j} \dots, F_{k} \dots}(\dots, f_{j}^{-} \dots, f_{k}^{+} \dots). \end{split}$$
(5.5)

This raises the following objection: perhaps, it would be more sensible to simply use $S[Q_{t_n}^{F_n}]$ as the formal parameterization, rather than the "bare" functions $Q_{t_n}^{F_n}$. After all, the pair-wise interference decomposition in observation 5 only consists of symmetric combinations anyway,

$$P_{t_n}^{\overline{F}_n} \left(\bigvee_{\omega(\overline{f}_n)} f_n\right) = \sum_{f_n^{\pm} \in \omega(\overline{f}_n)} Q_{t_n}^{F_n}(f_n^{+}, f_n^{-}) = \prod_{j=1}^n \left(\sum_{(f_j^{+}, f_j^{-}) \in \omega(\overline{f}_j) \times \omega(\overline{f}_j)} Q_{t_n}^{F_n}(f_n^{+}; f_n^{-}) \right)$$
$$= \prod_{j=1}^n \left(\sum_{\{f_j^{+}, f_j^{-}\} | f_j^{\pm} \in \omega(\overline{f}_j)} \sum_{(\phi_j^{+}, \phi_j^{-}) \in S(\{f_j^{+}, f_j^{-}\})} \right) Q_{t_n}^{F_n}(\phi_n^{+}, \dots, \phi_1^{+}; \phi_n^{-}, \dots, \phi_1^{-})$$
$$= \prod_{j=1}^n \left(\sum_{\{f_j^{+}, f_j^{-}\} | f_j^{\pm} \in \omega(\overline{f}_j)} \right) S[Q_{t_n}^{F_n}](f_n^{+}, f_n^{-}; \dots; f_1^{+}, f_1^{-}).$$
(5.6)

Hence, when debating the introduction of an *ad hoc* formal element, what standard should be used to determine whether distributions like $Q_{t_n}^{F_n}$ should be included over alternatives such as $\mathcal{S}[Q_{t_n}^{F_n}]$? In response, we propose the following two criteria:

- (i) There must be a compelling case demonstrating why the purpose of this new element cannot be fulfilled by phenomenological elements (the probability distributions $P_{t_n}^{F_n}$ and $\mathcal{S}[Q_{t_n}^{F_n}]$ in this case).
- (ii) Even if such a case is successfully made, minimal extensions of the phenomenological elements should be preferred over formulations that introduce additional, potentially excessive, complexity.

We believe that the distributions $Q_{t_n}^{F_n}$ comply with both requirements. First, the necessity for introducing an extension from single-sequence to sequence-pair distributions is evident when considering the difficulty of clearly expressing the crucial pair-wise interference decomposition by only using the phenomenological probability distributions. This issue alone does not present a strong enough case for adopting the "bare" $Q_{t_n}^{F_n}$, since this decomposition can also be accomplished using the phenomenological $\mathcal{S}[Q_{t_n}^{F_n}]$. However, the case for $Q_{t_n}^{F_n}$ is further strengthened by the demonstration of the factorization rule in "which subsystem" interference experiments-something that would remain obscured if only probabilities were used. Moreover, the case of "which subsystem" interference effects also disqualifies $\mathcal{S}[Q_{t_n}^{F_n}]$, as the symmetrization eliminates the anti-symmetric imaginary part of $Q_{t_n}^{F_n}$, which is crucial for parameterizing the observed co-interference term.

Second, the distributions $Q_{t_n}^{F_n}$ can be reasonably considered the simplest possible extension of phenomenological probabilities. Transitioning from $P_{t_n}^{F_n}$, which are (probability) distributions on the space of sequences $f_n \in \Omega(F_n)$, to $Q_{t_n}^{F_n}$, which are (non-probability) distributions on the space of sequence pairs $(f_n^+, f_n^-) \in \Omega(F_n) \times \Omega(F_n)$, is the simplest and most natural progression of the formal description. In contrast, due to their symmetries, the functions $\mathcal{S}[Q_{t_n}^{F_n}]$ should be viewed as distributions on the space of n pairs of alternatives. In fact, since $\mathcal{S}[Q_{t_n}^{F_n}]$ are defined as symmetrized interference terms (equal to the symmetric real part of $Q_{t_n}^{F_n}$), they should be viewed as functions of, up to, n different sequences—an extreme increase in complexity compared to $P_{t_n}^{F_n}$.

5.2 **Bi-probability distributions**

If we accept the distributions $Q_{t_n}^{F_n}$ as a valid extension of the formal description, the next step is to collate the data from the analyzed experiments and verify that the parameterization with these functions remains consistent with all phenomenological observations made thus far. In principle, the most straightforward approach would be to treat each measurement outcome as an criple, the most straightforward approach be to treat each measurement outcome as an equality constraint, where the values $Q_{t_n}^{F_n}(f_n^+, f_n^-)$ are the unknowns to be solved for, and the phenomenological probabilities $P_{t_n}^{F_n}(f_n)$ are the given parameters. For instance, constraints such as $\operatorname{Re} Q_{t_n}^{F_n}(f_n, f_n) - P_{t_n}^{F_n}(f_n) = 0$ and $\sum_{f_n^{\pm} \in \omega(\bar{f}_n)} Q_{t_n}^{F_n}(f_n^+, f_n^-) - P_{t_n}^{\bar{F}_n}(\bar{f}_n) = 0$, would be used. However, we know that such a system of equations does not have a single unique solution—were it so, the distributions $Q_{t_n}^{F_n}$ themselves would be phenomenological quantities! Consequently, in cuch an approach the constraints would constitute an infinite number of conditions defining the

It so, the distributions $Q_{t_n}^{F_n}$ themselves would be phenomenological quantities! Consequently, in such an approach, the constraints would constitute an infinite number of conditions defining the functions $Q_{t_n}^{F_n}$, which violates the requirement of minimizing the added complexity. Instead of attempting to solve every constraining equation, we should focus on identifying a minimal set of properties for the distributions $Q_{t_n}^{F_n}$ that would be sufficient for them to serve as a viable solutions. For example, if every combination of the type $\sum_{f_n^{\pm} \in \omega} Q_{t_n}^{F_n}(f_n^+, f_n^-)$ is a non-negative number not greater than 1, this would be sufficient for $Q_{t_n}^{F_n}$ to parameterize the pair-wise interference decomposition for any coarse grained measurements. interference decomposition for any coarse-grained measurements.

When composing such a list, aside from minding the contents of phenomenological observations, we can aid our efforts by considering how the properties of single-sequence probability distributions could be generalized for the case of complex-valued distributions of sequence pairs. The following properties can be identified in using this approach:

(Q1) Normalization

$$\sum_{\boldsymbol{f}_n^{\pm} \in \Omega(\boldsymbol{F}_n)} Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n^+, \boldsymbol{f}_n^-) = 1$$

(Q2) Bi-consistency:

$$\sum_{\substack{f_j^{\pm} \in \Omega(F_j) \\ f_n^{\pm} \in \Omega(F_j)}} Q_{t_n}^{F_n}(f_n^+, f_n^-) = Q_{t_n, \dots, t_j}^{F_n, \dots, F_j}(f_n^+, \dots, f_j^{\star}, \dots, f_1^+; f_n^- \dots, f_j^{\star}, \dots, f_1^-);$$

(Q3) Causality:

$$Q_{t_n,\dots,t_1}^{F_n,\dots,F_1}(f_n^+,\dots,f_1^+;f_n^-,\dots,f_1^-) \propto \delta_{f_n^+,f_n^-};$$

(Q4) Factorization rule for independent systems:

$$Q_{\boldsymbol{t}_n}^{AB}(\boldsymbol{a}_n^+ \wedge \boldsymbol{b}_n^+, \boldsymbol{a}_n^- \wedge \boldsymbol{b}_n^-) = Q_{\boldsymbol{t}_n}^A(\boldsymbol{a}_n^+, \boldsymbol{a}_n^-)Q_{\boldsymbol{t}_n}^B(\boldsymbol{b}_n^+, \boldsymbol{b}_n^-);$$

(Q5) Positive semi-definiteness:

$$\sum_{\boldsymbol{f}_n^{\pm} \in \Omega(\boldsymbol{F}_n)} Z(\boldsymbol{f}_n^+) Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n^+, \boldsymbol{f}_n^-) Z(\boldsymbol{f}_n^-)^* \geq 0 \quad \text{for any function } \boldsymbol{f}_n \mapsto Z(\boldsymbol{f}_n) \in \mathbb{C};$$

This property can be rephrased as follows: $\hat{Q}_{t_n}^{F_n} := [Q_{t_n}^{F_n}(f_n^+, f_n^-)]_{f_n^-, f_n^+}$, where sequences $f_n^{\pm} \in \Omega(F_n)$ are treated as indexes, is a positive semi-definite matrix.

Functions $Q_{t_n}^{F_n}$ satisfying the properties (Q1)–(Q5) are referred to as *bi-probability distributions* in [16, 17], and we shall stick to this convention; the name *decoherence functional* can also be found in the literature [18, 21, 22, 24, 25].

Examining the listed properties, we see that (Q1) is a straightforward generalization of the normalization condition for probability distributions. The property (Q3), for obvious reasons, does not correspond to any property of single-sequence distributions: instead, given the properties (Q1)and (Q2), it is a necessary and sufficient conditions for the bi-probabilities $Q_{t_n}^{F_n}$ to be compatible with observations 1 and 3. The factorization rule (Q4) simply reiterates the findings of Section 4.3, see Eq. (4.21). The positive semi-definiteness (Q5) can be seen as a generalization of the nonnegativity of single-sequence probability distributions. It is also the "workhorse" property: it enforces the Hermitianity (4.13), and more importantly, when combined with normalization (Q1), bi-consistency (Q2) and causality (Q3), it implies

$$Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n, \boldsymbol{f}_n) = \operatorname{Re} Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n, \boldsymbol{f}_n) \ge 0 \quad \text{and} \quad \sum_{\boldsymbol{f}_n \in \Omega(\boldsymbol{F}_n)} Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n, \boldsymbol{f}_n) = 1.$$
(5.7)

This indicates that the diagonal values of bi-probabilities are non-negative and normalized, thus consituting multi-time probability distributions,

$$Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n, \boldsymbol{f}_n) = P_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n),$$
(5.8)

which exhibit causality in agreement with observation 1,

$$\sum_{f_{n+1}} Q_{t_{n+1}}^{F_{n+1}}(f_{n+1}, f_{n+1}) = \sum_{f_{n+1}} \sum_{f_{n+1}} \delta_{f_{n+1}, f_{n+1}} Q_{t_{n+1}}^{F_{n+1}}(f_{n+1}, f_n; f_{n+1}', f_n) = Q_{t_n}^{F_n}(f_n, f_n)$$

$$\implies \sum_{f_{n+1}} P_{t_{n+1}}^{F_{n+1}}(f_{n+1}) = P_{t_n}^{F_n}(f_n),$$
(5.9)

and are inconsistent (thus compatible with observation 2),

$$\begin{aligned} Q_{t_{n},\dots,f_{j},\dots,f_{1}}^{F_{n},\dots,F_{j}}(f_{n},\dots,f_{j},\dots,f_{1};f_{n},\dots,f_{j},\dots,f_{1}) \\ &= \sum_{f_{j}^{\pm} \in \Omega(F_{j})} Q_{t_{n}}^{F_{n}}(f_{n},\dots,f_{j}^{+},\dots,f_{1};f_{n},\dots,f_{j}^{-},\dots,f_{1}) \\ &\implies P_{t_{n},\dots,f_{j},\dots,t_{1}}^{F_{n},\dots,F_{j}}(f_{n},\dots,f_{j},\dots,f_{1}) - \sum_{f_{j} \in \Omega(F_{j})} P_{t_{n}}^{F_{n}}(f_{n}) \end{aligned}$$

$$= \sum_{\substack{f_j^+ \neq f_j^- \\ f_j^+ \neq f_j^-}} Q_{t_n}^{F_n}(f_n, \dots, f_j^+, \dots, f_1; f_n, \dots, f_j^-, \dots, f_1)$$

$$= \sum_{\substack{f_j^+ \neq f_j^- \\ f_j^+ \neq f_j^-}} \frac{1}{2} \Big[P_{\dots t_j \dots}^{\dots \overline{F_j} \dots}(\dots, f_j^+ \vee f_j^-, \dots) - P_{\dots t_j \dots}^{\dots F_j \dots}(\dots, f_j^+, \dots) - P_{\dots t_j \dots}^{\dots F_j \dots}(\dots, f_j^-, \dots) \Big]$$

$$\neq 0.$$
(5.10)

Moreover, given a collection of resolutions $\operatorname{Res}(\overline{F}_j|F_j)$ for $j = 1, \ldots, n$ defined in accordance with Eq. (4.3), positive semi-definiteness also implies

$$\sum_{\boldsymbol{f}_{n}^{\pm} \in \omega(\bar{\boldsymbol{f}}_{n})} Q_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n}^{+}, \boldsymbol{f}_{n}^{-}) \geq 0 \quad \text{and} \quad \sum_{\bar{\boldsymbol{f}}_{n} \in \Omega(\bar{\boldsymbol{F}}_{n})} \sum_{\boldsymbol{f}_{n}^{\pm} \in \omega(\bar{\boldsymbol{f}}_{n})} Q_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n}^{+}, \boldsymbol{f}_{n}^{-}) = \sum_{\boldsymbol{f}_{n}^{\pm} \in \Omega(\boldsymbol{F}_{n})} Q_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n}^{+}, \boldsymbol{f}_{n}^{-}) = 1$$

$$(5.11)$$

thus explaining the compliance with the pair-wise interference decomposition (5.2).

Finally, we have the *bi-consistency* (Q2). This can be seen as a generalization of the classical Kolmogorov consistency condition (cf. Section 4.1) adopted for distributions on pairs of sequences. It is also a sufficient condition for functions $Q_{t_n}^{F_n}$ to be compliant with observation 4 of the effects of the extreme coarse-grained measurements.

5.3 The bi-trajectory picture

As we can see from Eq. (5.10), even though bi-probabilities are directly responsible for violating the consistency of the probability family $P_{t_n}^{F_n}$ through interference effects—and thus invalidating the trajectory picture for quantum observables—they still retain an analogous notion (Q2): rather than for a sequence of results, it is the *pair* of sequences of results that is consistent here. This begs the question if, in analogy to the extension theorem of classical probability theory, bi-consistency implies the existence of a master measure on the space of trajectory *pairs*.

As it was discussed in [17], from the mathematical point of view, when non-positive-valued distributions are involved, bi-consistency (Q2) alone is not sufficient for the master measure to exists. Nevertheless, we can show that the observations 9 (Markovianity) and 10 (Zeno effect)—the two phenomenological properties that give us a glimpse into the inner-workings of the quantum dynamical laws—imply that bi-probabilities *do* satisfy the additional necessary condition for the generalized Kolmogorov extension theorem. The formal proof of this assertion goes as follows.

Let $\mathbb{Q}_{[0,T]}^F = \{Q_{t_n}^F \mid n \in \mathbb{N}, t_1, \dots, t_n \in [0,T], 0 < t_1 < \dots < t_n\}$ be the family of bi-probabilities associated with observable F and the finite time window [0,T]. It was previously proven—see [17, Theorem 2.2]—that the family $\mathbb{Q}_{[0,T]}^F$ extends to the master measure $\mathcal{Q}_{[0,T]}^F[f^+, f^-][\mathcal{D}f^+][\mathcal{D}f^-]$ provided that (i) it is bi-consistent; and (ii) it is uniformly bounded,

$$\sup\left\{\|Q_{t_n}^F\|_1 \mid Q_{t_n}^F \in \mathbb{Q}_{[0,T]}^F\right\} < \infty \qquad \text{where} \quad \|Q_{t_n}^F\|_1 := \sum_{\boldsymbol{f}_n^{\pm}} |Q_{t_n}^F(\boldsymbol{f}_n^+, \boldsymbol{f}_n^-)|.$$
(5.12)

In this case, bi-consistency is automatically satisfied (Q2), but uniform boundedness has to be verified: (1) Due to bi-consistency we have (see also [17], Lemma 3.5 and Proposition 3.6)

$$\|Q_{t_n}^F\|_1 \le \lim_{n \to \infty} \|Q_{s_n}^F\|_1$$
 where $s_j = jT/n.$ (5.13)

(2) The property (Q5), i.e., $Q_{t_n}^F(f_n^+, f_n^-)$ is a positive *matrix* when one treats f_n^{\pm} as indexes, implies that

$$|Q_{\boldsymbol{s}_{n}}^{F}(\boldsymbol{f}_{n}^{+},\boldsymbol{f}_{n}^{-})| \leq \sqrt{Q_{\boldsymbol{s}_{n}}^{F}(\boldsymbol{f}_{n}^{+},\boldsymbol{f}_{n}^{+})Q_{\boldsymbol{s}_{n}}^{F}(\boldsymbol{f}_{n}^{-},\boldsymbol{f}_{n}^{-})} = \sqrt{P_{\boldsymbol{s}_{n}}^{F}(\boldsymbol{f}_{n}^{+})}\sqrt{P_{\boldsymbol{s}_{n}}^{F}(\boldsymbol{f}_{n}^{-})}.$$
(5.14)

(3) The short-time behavior of the survival probability (4.32), implied by the Markov property (4.22) and the Zeno effect, allows us to estimate the square root of probability,

$$\sqrt{P_{\boldsymbol{s}_n}^F(\boldsymbol{f}_n)} = \sqrt{P_{s_1|0}^F(f_1|p)} \prod_{j=1}^{n-1} \sqrt{P_{s_{j+1}|s_j}^{F|F}(f_{j+1}|f_j)}$$

$$\leq |\Omega(F)| \prod_{j=1}^{n-1} \left(\delta_{f_{j+1},f_j} + (1 - \delta_{f_{j+1},f_j}) \sqrt{1 - P_{s_j + \frac{T}{n}|s_j}^{F|F}(f_j|f_j)} \right) \\ \leq |\Omega(F)| \prod_{j=1}^{n-1} \left(\delta_{f_{j+1},f_j} + (1 - \delta_{f_{j+1},f_j}) \left(\sup_{f} |v(f,s_j)| \frac{T}{n} + O(n^{-2}) \right) \right),$$
(5.15)

which then leads to the required uniform bound,

$$\sup \|Q_{t_n}^F\|_1 \le |\Omega(F)|^2 \lim_{n \to \infty} \left[\prod_{j=1}^{n-1} \sum_{f_j} \left(\delta_{f_{j+1}, f_j} + (1 - \delta_{f_{j+1}, f_j}) \left(\sup_f |v(f, s_j)| \frac{T}{n} + O(n^{-2}) \right) \right) \right]^2$$

$$\le |\Omega(F)|^2 \lim_{n \to \infty} \left[\prod_{j=1}^{n-1} \left(1 + |\Omega(F)| \sup_f |v(f, s_j)| \frac{T}{n} + O(n^{-2}) \right) \right]^2$$

$$\le |\Omega(F)|^2 \exp\left[2|\Omega(F)| \int_0^T \sup_f |v(f, s)| ds \right] < \infty.$$
(5.16)

Therefore, $\mathcal{Q}_{[0,T]}^F$ extends the family $\mathbb{Q}_{[0,T]}^F$ for any choice of T. But this particular $\mathcal{Q}_{[0,T]}^F$ can also be seen as a restriction (to time window [0,T]) of a master measure \mathcal{Q}^F that extends $\mathbb{Q}^F = \lim_{n\to\infty} \mathbb{Q}_{[0,T_n]}^F$ such that $T_1 < T_2 < \cdots < T_n$; compare with [17], Section 4.1. This concludes the proof.

Therefore, it is possible to deduce at this point that the master bi-probability does, indeed, exist:

(Q6) The master bi-trajectory measure: The bi-probabilities $Q_{t_n}^F$ associated with a single observable F are discrete-time restrictions of a complex-valued measure $\mathcal{Q}^F[f^+, f^-][\mathcal{D}f^+][\mathcal{D}f^-]$ on the space of bi-trajectories, $t \mapsto (f^+(t), f^-(t)) \in \Omega(F) \times \Omega(F)$,

$$Q_{t_n}^F(f_n^+, f_n^-) = \iint \Big(\prod_{j=1}^n \delta_{f^+(t_j), f_j^+} \delta_{f^-(t_j), f_j^-} \Big) \mathcal{Q}^F[f^+, f^-][\mathcal{D}f^+][\mathcal{D}f^-].$$

In particular, due to the link (5.8) with probability distributions for sequential measurements, we also obtain that

$$P_{\boldsymbol{t}_n}^F(\boldsymbol{f}_n) = \iint \Big(\prod_{j=1}^n \delta_{f^+(t_j), f_j} \delta_{f^-(t_j), f_j} \Big) \mathcal{Q}^F[f^+, f^-][\mathcal{D}f^+][\mathcal{D}f^-],$$

showing that the observed probabilities $P_{t_n}^F$ can be interpreted as the result of a superposition of *interfering* pairs of elementary trajectories—the *bi-trajectory picture*.

Naturally, the master bi-trajectory measure Q^F is not a phenomenological quantity, as it cannot be directly or indirectly measured by a classical observer. Rather, it is a logically inferred *master object* of the formalism that, on the one hand, describes the dynamics of observable F in terms of bi-trajectories, and on the other hand, is a "source" of all observable probabilities regarding F. However, as we will discover in due time, Q^F turns out not to be the true master object: further deduction will lead us to the conclusion that, in this hierarchy of objects, the master bi-probability associated with single observable F is not on the very top. For now, we postpone the resolution of the true-master-object question to Section 9.

6 The Hilbert space representation

6.1 Bi-probability distributions as inner products

In his paper [18], S. Gudder shows that bi-probability distributions (there denoted as decoherence functionals) can be parameterized in terms of inner products on Hilbert spaces. This mathematical

result seems to be the simplest explanation of why the mathematical formalism of quantum theory should be based on the structure of Hilbert spaces. The construction itself is carried out in the following steps. First, define the $|\Omega(\mathbf{F}_n)| \times |\Omega(\mathbf{F}_n)|$ matrix,

$$\hat{Q}_{\boldsymbol{t}_n}^{\boldsymbol{F}_n} := [Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n^+, \boldsymbol{f}_n^-)]_{\boldsymbol{f}_n^+, \boldsymbol{f}_n^-}$$
(6.1)

by treating sequences \mathbf{f}_n^{\pm} as indexes. This matrix is positive semi-definite because the bi-probability satisfies (Q5), and tr $\hat{Q}_{\mathbf{t}_n}^{\mathbf{F}_n} = \sum_{\mathbf{f}_n} P_{\mathbf{t}_n}^{\mathbf{F}_n}(\mathbf{f}_n) = 1$ because of normalization. Second, take the vector space $\mathbb{C}^{|\Omega(\mathbf{F}_n)|}$, and pick the standard basis $\{|\psi_{\mathbf{t}_n}(\mathbf{f}_n)\rangle \mid \mathbf{f}_n \in \Omega(\mathbf{F}_n)\}$ which is orthonormal with respect to the standard inner product $\langle \psi_{\mathbf{t}_n}(\mathbf{f}_n^+)|\psi_{\mathbf{t}_n}(\mathbf{f}_n^-)\rangle = \prod_{j=1}^n \delta_{f_j^+, f_j^-}$; again, sequences \mathbf{f}_n are treated here as indexes. Third, define the following function on pairs of vectors:

$$\langle\!\langle \alpha | \beta \rangle\!\rangle_{\boldsymbol{t}_n} := \sum_{\boldsymbol{f}_n^{\pm} \in \Omega(\boldsymbol{F}_n)} A(\boldsymbol{f}_n^+) Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n^+, \boldsymbol{f}_n^-) B(\boldsymbol{f}_n^-)^*,$$
(6.2)

where $|\alpha\rangle = \sum_{\boldsymbol{f}_n} A(\boldsymbol{f}_n) |\psi_{\boldsymbol{t}_n}(\boldsymbol{f}_n)\rangle$ and $|\beta\rangle = \sum_{\boldsymbol{f}_n} B(\boldsymbol{f}_n) |\psi_{\boldsymbol{t}_n}(\boldsymbol{f}_n)\rangle$. The function $\langle\!\langle \cdot | \cdot \rangle\!\rangle_{\boldsymbol{t}_n} : \mathbb{C}^{|\Omega(\boldsymbol{F}_n)|} \times \mathbb{C}^{|\Omega(\boldsymbol{F}_n)|} \to \mathbb{C}$ defined this way is an inner product⁵, $\hat{Q}_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}$ is its metric, and thus, $(\mathbb{C}^{|\Omega(\boldsymbol{F}_n)|}, \langle\!\langle \cdot | \cdot \rangle\!\rangle_{\boldsymbol{t}_n}) = \mathcal{H}$ is a Hilbert space. Finally, when we evaluate the inner product between any two basis vectors, we arrive at the desired representation,

(Q7) Inner product representation

$$\begin{aligned} Q_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n}^{+},\boldsymbol{f}_{n}^{-}) &= \langle\!\langle \psi_{\boldsymbol{t}_{n}}(\boldsymbol{f}_{n}^{+}) | \psi_{\boldsymbol{t}_{n}}(\boldsymbol{f}_{n}^{-}) \rangle\!\rangle_{\boldsymbol{t}_{n}} &= \langle \psi_{\boldsymbol{t}_{n}}(\boldsymbol{f}_{n}^{+}) | \hat{Q}_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}} | \psi_{\boldsymbol{t}_{n}}(\boldsymbol{f}_{n}^{-}) \rangle \\ &= \operatorname{tr} \left[|\psi_{\boldsymbol{t}_{n}}(\boldsymbol{f}_{n}^{-}) \rangle \langle \psi_{\boldsymbol{t}_{n}}(\boldsymbol{f}_{n}^{+}) | \hat{Q}_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}} \right], \end{aligned}$$

where $\hat{Q}_{\boldsymbol{t}_n}^{\boldsymbol{F}_n} \geq 0$, tr $\hat{Q}_{\boldsymbol{t}_n}^{\boldsymbol{F}_n} = 1$, and

$$\left\{ |\psi_{\boldsymbol{t}_n}(\boldsymbol{f}_n)\rangle \mid \boldsymbol{f}_n \in \Omega(\boldsymbol{F}_n), \langle \psi_{\boldsymbol{t}_n}(\boldsymbol{f}_n^+) | \psi_{\boldsymbol{t}_n}(\boldsymbol{f}_n^-) \rangle = \delta_{\boldsymbol{f}_n^+, \boldsymbol{f}_n^-} \right\}$$

is an orthonormal basis.

The representation (Q7) is, however, not unique; it is always possible to use an appropriate unitary transformation to switch between Hilbert spaces, and thus, to switch between representations constructed with the corresponding inner products. For example, if the bi-probability is represented with the inner product of thespace $(W, ((\cdot|\cdot))_{t_n}) = \mathcal{K}$,

$$Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n^+, \boldsymbol{f}_n^-) = \left(\left(\phi_{\boldsymbol{t}_n}(\boldsymbol{f}_n^+) \middle| \phi_{\boldsymbol{t}_n}(\boldsymbol{f}_n^-) \right) \right)_{\boldsymbol{t}_n}, \tag{6.3}$$

where $\{ |\phi_{\boldsymbol{t}_n}(\boldsymbol{f}_n)\rangle \mid \boldsymbol{f}_n \in \Omega(F)^n \}$ spans \mathcal{K} , then there exist an unitary transformation $S: \mathcal{K} \to \mathcal{H}$ such that $\hat{S} |\phi_{\boldsymbol{t}_n}(\boldsymbol{f}_n)\rangle = |\psi_{\boldsymbol{t}_n}(\boldsymbol{f}_n)\rangle$, and thus

$$Q_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n^+, \boldsymbol{f}_n^-) = \left(\left(\phi_{\boldsymbol{t}_n}(\boldsymbol{f}_n^+) \middle| \phi_{\boldsymbol{t}_n}(\boldsymbol{f}_n^-) \right) \right)_{\boldsymbol{t}_n} = \left\langle \left\langle S \phi_{\boldsymbol{t}_n}(\boldsymbol{f}_n^+) \middle| S \phi_{\boldsymbol{t}_n}(\boldsymbol{f}_n^-) \right\rangle \right\rangle_{\boldsymbol{t}_n} = \left\langle \left\langle \psi_{\boldsymbol{t}_n}(\boldsymbol{f}_n^+) \middle| \psi_{\boldsymbol{t}_n}(\boldsymbol{f}_n^-) \right\rangle \right\rangle_{\boldsymbol{t}_n}.$$
(6.4)

Gudder's result provides us the recipe for a general purpose parameterization of an arbitrary bi-probability with the elements of Hilbert space formalism. However, here we are dealing with a specific kind of bi-probabilities whose various properties result from constraints imposed by the phenomenology of sequential measurements, rather than mathematics alone. Therefore, we shall now harness those constraints to narrow down the general parameterization to a *specific* form of Hilbert space representation adequate for the quantum theory.

⁵In fact, $\langle\!\langle \cdot | \cdot \rangle\!\rangle_{t_n}$ is "almost" an inner product. At this point it is not guaranteed that $\langle\!\langle \gamma | \gamma \rangle\!\rangle_{t_n} = 0$ only when $|\gamma\rangle = 0$; in other words, it is possible that $\operatorname{rank}(\hat{Q}_{t_n}^{F_n}) < \dim(\mathbb{C}^{|\Omega(F_n)|})$. This can be easily fixed by collecting all null vectors into the set $N = \{|\gamma\rangle \in \mathbb{C}^{|\Omega(F_n)|} \mid \langle\!\langle \gamma | \gamma \rangle\!\rangle_{t_n} = 0\}$ and switching to the quotient vector space $V = \mathbb{C}^{|\Omega(F_n)|}/N$, with elements in the form of equivalence classes $|[\alpha]\rangle = |\alpha\rangle + N$. Then, $\operatorname{rank}(\hat{Q}_{t_n}^{F_n}) = \dim(V)$ and the function on $V \times V$ defined by $\langle\!\langle [\alpha] | [\beta] \rangle\!\rangle_{t_n} := \langle\!\langle \alpha | \beta \rangle\!\rangle_{t_n}$ is a proper inner product. Henceforth it is assumed that this construction is employed when needed.

6.2 The measurement-projector link

As a first step, we wish to deduce the measurement-projector link that, in the standard formalism, is thought to relate the physical action of the measuring device with the action of a projector operator as a formal representation of that device. To this end we apply Gudder's construction in the simplest case of n = 1 measurement with an *F*-device, for which we have

$$P_t^F(f) = Q_t^F(f, f) = \delta_{f, f'} Q_t^F(f, f').$$
(6.5)

In this case, the inner product representation (Q7) is not complicated,

$$Q_t^F(f,f') = \langle\!\langle \psi_t(f) | \psi_t(f') \rangle\!\rangle_t^F = \langle\!\psi_t(f) | \hat{Q}^F | \psi_t(f') \rangle$$

= $\delta_{f,f'} \operatorname{tr}_{\mathcal{H}(F)} \left(|\psi_t(f) \rangle \langle\!\psi_t(f) | \hat{Q}^F \right) = \operatorname{tr}_{\mathcal{H}(F)} \left(|\psi_t(f) \rangle \langle\!\psi_t(f) | \hat{Q}^F | \psi_t(f') \rangle \langle\!\psi_t(f') | \right).$ (6.6)

Here, $(V(F), \langle\!\langle \cdot | \cdot \rangle\!\rangle^F)$ is the Hilbert space with the inner product used in Gudder's construction, and $\mathcal{H}(F) = (V(F), \langle \cdot | \cdot \rangle)$ denotes the Hilbert space corresponding to the same vector space equipped with the standard inner product. The basis $\{|\psi_t(f)\rangle \in V(F) \mid f \in \Omega(F)\}$ is chosen in such a way that its elements are mutually orthogonal in $\mathcal{H}(F)$, $\langle \psi_t(f) | \psi_t(f') \rangle = \delta_{f,f'}$; consequently, the operators $|\psi_t(f)\rangle \langle \psi_t(f)|$ are standard rank-1 projectors, and the metric satisfies the standard conditions: $\hat{Q}^F \geq 0$ and $\operatorname{tr}_{\mathcal{H}(F)} \hat{Q}^F = 1$.

When we repeat the construction for some coarse-grained variant of the *F*-device—an \overline{F} -device—we obtain analogous results,

$$Q_t^{\overline{F}}(\overline{f},\overline{f}') = \langle\!\langle \phi_t(\overline{f}) | \phi_t(\overline{f}') \rangle\!\rangle_t^{\overline{F}} = \delta_{\overline{f},\overline{f}'} \operatorname{tr}_{\mathcal{H}(\overline{F})} \left(|\phi_t(\overline{f})\rangle \langle \phi_t(\overline{f}) | \hat{Q}^{\overline{F}} \right) = \operatorname{tr}_{\mathcal{H}(\overline{F})} \left(|\phi_t(\overline{f})\rangle \langle \phi_t(\overline{f}) | \hat{Q}^{\overline{F}} | \phi_t(\overline{f}') \rangle \langle \phi_t(\overline{f}') | \right),$$
(6.7)

where, again, $\mathcal{H}(\overline{F}) = (V(\overline{F}), \langle \cdot | \cdot \rangle)$ is the Hilbert space obtained by endowing the vector space $V(\overline{F})$ with the standard inner product, spanned by an orthonormal basis $\{|\psi_t(\overline{f})\rangle \in V(\overline{F}) \mid \overline{f} \in \Omega(\overline{F})\}, |\psi_t(\overline{f})\rangle \langle \psi_t(\overline{f})|$ are rank-1 projector operators, and $\hat{Q}^{\overline{F}} \ge 0$, $\operatorname{tr}_{\mathcal{H}(\overline{F})} \hat{Q}^{\overline{F}} = 1$.

The measurements with F- and \overline{F} -devices, of course, satisfy the phenomenological condition (4.5),

$$Q_t^{\overline{F}}(\overline{f},\overline{f}) = P_t^{\overline{F}}(\overline{f}) = \sum_{f \in \omega(\overline{f})} P_t^F(f) = \sum_{f \in \omega(\overline{f})} Q_t^F(f,f) \quad \text{where} \quad \bigsqcup_{\overline{f} \in \Omega(\overline{F})} \omega(\overline{f}) = \Omega(\overline{F}), \quad (6.8)$$

which then allows us to establish the relation between their respective representations,

$$\operatorname{tr}_{\mathcal{H}(\overline{F})}\left(|\phi_t(\overline{f})\rangle\langle\phi_t(\overline{f})|\hat{Q}^{\overline{F}}\right) = \operatorname{tr}_{\mathcal{H}(F)}\left(\sum_{f\in\omega(\overline{f})}|\psi_t(f)\rangle\langle\psi_t(f)|\hat{Q}^{F}\right).$$
(6.9)

Now, there is no purely logical/mathematical reason for the Hilbert spaces $\mathcal{H}(F)$ and $\mathcal{H}(\overline{F})$ to be related to each other in any way, especially given the fact that $\dim \mathcal{H}(\overline{F}) \leq \dim \mathcal{H}(F)$. However, it seems grossly unnecessary that the respective formal representations "live" in different Hilbert spaces, even though, ostensibly, both describe a measurement of the same physical quantity (only with different degree of graining). Therefore, if we wish to be efficient with the "economy of entities" in our deduction, we ought to bring both representations to the same higher-dimensional space $\mathcal{H}(F)$.

To accomplish this, first we have to resolve a conundrum: we have to decide how to extend the projectors $|\phi_t(\bar{f})\rangle\langle\phi_t(\bar{f})|$ and the metric $\hat{Q}^{\bar{F}}$ from the lower-dimension space $\mathcal{H}(\bar{F})$ to the higherdimensional $\mathcal{H}(F)$. Generally, there are infinitely many ways to do it, but here we can rely on the phenomenological relation (6.9) as a guiding principle for our decision making. The simplest choice adhering to this principle is to identify the operators in $\mathcal{H}(\bar{F})$ with operators in $\mathcal{H}(F)$ that fulfill the same function, i.e., projectors correspond to projectors, and metric corresponds to metric,

$$|\phi_t(\bar{f})\rangle\langle\phi_t(\bar{f})| \to \hat{P}_t^{\bar{F}}(\bar{f}) := \sum_{f \in \omega(\bar{f})} |\psi_t(f)\rangle\langle\psi_t(f)| \quad \text{and} \quad \hat{Q}^{\bar{F}} \to \hat{Q}^{\bar{F}}.$$
(6.10)

We could imagine that the *F*-device itself is not perfectly fine-grained either, and that, in principle, there could exists a more fine-grained device measuring the observable *F*, i.e., $F = \overline{K}$ where *K* is perfectly fine-grained. We can account for that possibility by extending our projectors and metric to an even higher-dimensional *system* Hilbert space $\mathcal{H}_S = (V_S, \langle \cdot | \cdot \rangle)$,

$$|\psi_t(f)\rangle\langle\psi_t(f)| \to \hat{P}_t^F(f) : \hat{P}_t^F(f)\hat{P}_t^F(f') = \delta_{f,f'}\hat{P}_t^F(f), \sum_{f\in\Omega(F)}\hat{P}_t^F(f) = \hat{1}_S;$$
(6.11a)

$$\hat{Q}^F \to \hat{\rho}^p_{t_0} : \hat{\rho}^p_{t_0} \ge 0, \ \operatorname{tr}_S \hat{\rho}^p_{t_0} = 1,$$
(6.11b)

where the choice of the metric $\hat{\rho}_{t_0}^p$ has to be consistent with the phenomenology, i.e.,

$$P_{t|t_0}^F(f|p) = \operatorname{tr}_{\mathcal{H}(F)}\left(|\psi_t(f)\rangle\langle\psi_t(f)|\hat{Q}^F\right) = \operatorname{tr}_S[\hat{P}_t^F(f)\hat{\rho}_{t_0}^p],\tag{6.12}$$

where the distribution p(K,k) in $P_{t|t_0}^F(f|p)$ describes the initialization event [see Section 4.4, Eq. (4.28)]. The rank of $\hat{P}_t^F(f)$ could then be greater than 1 (i.e., $\hat{P}_t^F(f) \neq |\Psi_t^F(f)\rangle \langle \Psi_t^F(f)|$ for some $f \in \Omega(F)$), leaving an option for even more fine-grained measurements of F. Whatever the case, how fine the graining can go is ultimately determined by the dimension of the Hilbert space \mathcal{H}_S , which can be considered as a property of the measured quantum system. Moreover, the dimension of \mathcal{H}_S can be found out, in principle, by looking for an observable that exhibits Markovianity, which is the indicator of perfectly fine-grained measurements, as elaborated in Section 4.4.

Next, let us consider an observable G measured with a G-device that is not a coarse- or fine-grained variant of the F-device. In such a case, again we arrive at the corresponding parameterization with the basis $\{|\chi_t(g)\rangle \in V(G) \mid g \in \Omega(G)\}$ spanning the Hilbert space $\mathcal{H}(G) =$ $(V(G), \langle \cdot | \cdot \rangle)$, rank-1 projectors $|\chi_t(g)\rangle\langle\chi_t(g)|$, and the metric \hat{Q}^G . Since the measurements are still performed on the same physical system, we should extend this representation to the Hilbert space \mathcal{H}_S of the system. Therefore, the rank-1 projectors in $\mathcal{H}(G)$ are mapped to projectors in \mathcal{H}_S , $|\chi_t(g)\rangle\langle\chi_t(g)| \to \hat{P}^G_t(g)$ (where $\hat{P}^G_t(g)\hat{P}^G_t(g') = \delta_{g,g'}\hat{P}^G_t(g)$ and $\sum_{g\in\Omega(G)}\hat{P}^G_t(g) = \hat{1}_S$), and, since the choice of G is equally arbitrary as the choice of F before it, the metric should be mapped to the same operator in \mathcal{H}_S , $\hat{Q}^G \to \hat{\rho}^p_{t_0}$. Of course, just like it was with F, the mappings for G also have to be consistent with the phenomenology,

$$P_{t|t_0}^G(g|p) = \operatorname{tr}_{\mathcal{H}(G)}\left(|\chi_t(g)\rangle\langle\chi_t(g)|\hat{Q}^G\right) = \operatorname{tr}_S[\hat{P}_t^G(g)\hat{\rho}_{t_0}^p].$$
(6.13)

Then, it follows that the set of projectors representing the measurement of G cannot be the same projectors representing the action of F-device. If this were the case, then the probability distributions of both devices would be identical, thus contradicting the assumption that G-device is not a fine- or coarse-grained version of F-device. In this way, we have arrived naturally at the link between measuring devices and partitions of the Hilbert space into images of the set of orthogonal projectors representing the device.

By following the guiding principle that, for every system observable F, we extend the corresponding metric \hat{Q}^F to the same operator $\hat{\rho}^p_{t_0}$, we have forced this operator into the role of the Hilbert space representation of the initialization event. Indeed, the projectors encode which measuring device was deployed and when it was deployed, so they cannot be responsible for setting the initial conditions; this task therefore has to fall on the remaining object, $\hat{\rho}^p_{t_0}$. When discussing the phenomenology of the Markov property in Section 4.4 we had determined that the initialization is accomplished via the deployment of a perfectly fine-grained measuring device, see Eq. (4.28). We can now use this insight, together with the measurement-projector link, to deduce the specific form of the object $\hat{\rho}^p_{t_0}$ representing a given initialization event described phenomenologically by the distribution p. To this end, first consider an experiment in which the perfectly fine-grained K-device is deployed immediately after the initialization with the same device measuring result k_0 , i.e., $p(K', k') = \delta_{K',K} \delta_{k',k_0} \equiv \delta^K_{k_0}(K', k')$. In accordance with the phenomenological observations 9 and 11, the results of such a measurement are described by

$$\lim_{\Delta t \to 0^+} P_{t_0 + \Delta t \mid t_0}^K(k \mid \delta_{k_0}^K) = \lim_{\Delta t \to 0} P_{t_0 + \Delta t \mid t_0}^{K \mid K}(k \mid k_0) = C_{k,k_0}^{K \mid K} = \delta_{k,k_0}.$$
(6.14)

On the other hand, by using the Hilbert space representation of the measurement we have

$$\lim_{\Delta t \to 0^+} P_{t_0 + \Delta t \mid t_0}^K(k \mid \delta_{k_0}^K) = \operatorname{tr}_S \left[\hat{P}_{t_0}^K(k) \hat{\rho}_{t_0}^{\delta_{k_0}^K} \right] = \langle \Psi_{t_0}^K(k) \mid \hat{\rho}_{t_0}^{\delta_{k_0}^K} \mid \Psi_{t_0}^K(k) \rangle = \delta_{k,k_0}, \tag{6.15}$$

where $\{|\Psi_{t_0}^K(k)\rangle \in \mathcal{H}_S \mid k \in \Omega(K)\}$ form an orthonormal basis (the projectors representing perfectly fine-grained measurements are all rank-1). This, together with the requirements $\hat{\rho}_{t_0}^p \geq 0$ and $\operatorname{tr}_S \hat{\rho}_{t_0}^p = 1$ for any p(K,k), implies that the metric matrix representing the initialization event in the form of a fine-grained measurement is a rank-1 projection operator,

$$\hat{\rho}_{t_0}^{\delta_{k_0}^K} = \hat{P}_{t_0}^K(k_0) = |\Psi_{t_0}^K(k_0)\rangle \langle \Psi_{t_0}^K(k_0)|.$$
(6.16)

Therefore, the metric representing a general initialization event has the form of a convex combination of rank-1 projectors corresponding to the perfectly fine grained measurements,

$$\hat{\rho}_{t_0}^p = \sum_K \sum_{k \in \Omega(K)} p(K,k) \hat{\rho}_{t_0}^{\delta_k^K} = \sum_K \sum_{k \in \Omega(K)} p(K,k) \hat{P}_{t_0}^K(k) = \sum_K \sum_{k \in \Omega(K)} p(K,k) |\Psi_{t_0}^K(k)\rangle \langle \Psi_{t_0}^K(k)|.$$
(6.17)

We have deduced here that a (single) measurement with a given device formally corresponds to a set of orthogonal projectors. Then, it stands to reason that the reverse is also true: in principle, it should be possible to build a device such that its measurements can be represented by a given set of projectors. Bearing this observation in mind, consider the following formal statement,

$$P_{t|0}^{K}(k|\delta_{k_{0}}^{K}) = \operatorname{tr}_{S}\left[\hat{P}_{t}^{K}(k)\hat{P}_{0}^{K}(k_{0})\right] \equiv \operatorname{tr}_{S}\left[\hat{P}_{0}^{K_{t}}(k)\hat{P}_{0}^{K}(k_{0})\right] = \lim_{\Delta t \to 0^{+}} P_{\Delta t|0}^{K_{t}|K}(k|k_{0}) = C_{k,k_{0}}^{K_{t}|K}.$$
 (6.18)

Here, K_t denotes an observable such that the projectors corresponding to the deployment of K_t -device at time t = 0 are the same as the projectors corresponding to K-device deployed at time t > 0, i.e., $\hat{P}_0^{K_t}(k) = \hat{P}_t^K(k)$ —according to our previous observation, such a device should exist in principle. The interesting thing about the above "trick" is that we have exploited the formal measurement–projector link to essentially eliminate the passage of time from the picture. We have shown here that, in the context of single measurement experiments, the effects of the system dynamics unfolding during the time between the initialization and the terminating measurement are fully captured by the measurement with an appropriate device immediately following the initialization. In this sense, the single-measurement context can be considered as quantum statics, where the metric $\hat{\rho}_{t_0}^p$ is de facto the master object (i.e., all possible outcomes are derivable from it), and all phenomenologically answerable questions can be expressed in terms of a deployment of a certain measuring device in the instant following the initialization event.

We shall see soon that the static picture above does not hold anymore in contexts of sequential measurements, in which the dynamical laws of the system can no longer be simply "absorbed" by the freedom of choice of the measuring devices. We will further investigate the formal relations between representations of K_t and K in the upcoming Section 7, where we shall deduce the Hilbert space representation of the system dynamical laws.

6.3 Hilbert space representation for bi-probabilities

Consider a conditional probability distribution,

$$P_{t_{n+1}|t_n}^{F_{n+1}|F_n}(f_{n+1}|f_n) = \frac{P_{t_{n+1}}^{F_{n+1}}(f_{n+1})}{P_{t_n}^{F_n}(f_n)} = \frac{Q_{t_{n+1}}^{F_{n+1}}(f_{n+1}, f_{n+1})}{Q_{t_n}^{F_n}(f_n, f_n)}.$$
(6.19)

Formally, this is an n = 1 bi-probability, and so we can construct its representation in the Hilbert space of the system in the same fashion as it was previously done in Section 6.2,

$$P_{t_{n+1}|\mathbf{t}_n}^{F_{n+1}|\mathbf{F}_n}(f_{n+1}|\mathbf{f}_n) = \operatorname{tr}_S\left[\hat{P}_{t_{n+1}}^{F_{n+1}}(f_{n+1})\hat{\rho}_{\mathbf{t}_n}^{\mathbf{F}_n}(\mathbf{f}_n)\right] \equiv \operatorname{tr}_S\left[\hat{P}_{t_{n+1}}^{F_{n+1}}(f_{n+1})\frac{\hat{q}_{\mathbf{t}_n}^{F_n}(\mathbf{f}_n)}{Q_{\mathbf{t}_n}^{F_n}(\mathbf{f}_n,\mathbf{f}_n)}\right], \quad (6.20)$$

where we have introduced the operator $\hat{q}_{t_n}^{F_n}(f_n) \ge 0$ that plays the role of the (non-normalized) metric. Then, using the causality of bi-probabilities (Q3),

$$\sum_{f_{n+1}\in\Omega(F_{n+1})} P_{t_{n+1}|\mathbf{t}_n}^{F_{n+1}|\mathbf{F}_n}(f_{n+1}|\mathbf{f}_n) = \frac{\sum_{f_{n+1}\in\Omega(F_{n+1})} Q_{\mathbf{t}_{n+1}}^{F_{n+1}}(\mathbf{f}_{n+1}, \mathbf{f}_{n+1})}{Q_{\mathbf{t}_n}^{F_n}(\mathbf{f}_n, \mathbf{f}_n)} = \frac{Q_{\mathbf{t}_n}^{F_n}(\mathbf{f}_n, \mathbf{f}_n)}{Q_{\mathbf{t}_n}^{F_n}(\mathbf{f}_n, \mathbf{f}_n)} = 1, \quad (6.21)$$

and the completeness of the projector operators, $\sum_{f \in \Omega(F)} \hat{P}_t^F(f) = \hat{1}_S$ for any F, we find out that the bi-probability is given by the trace of the pseudo-metric $\hat{q}_{t_n}^{F_n}$,

$$Q_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n},\boldsymbol{f}_{n}) = \operatorname{tr}_{S}\left[\sum_{f_{n+1}\in\Omega(F_{n+1})}\hat{P}_{t_{n+1}}^{F_{n+1}}(f_{n+1})\hat{q}_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n})\right] = \operatorname{tr}_{S}\left[\hat{q}_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n})\right], \quad (6.22)$$

and thus,

$$\operatorname{tr}_{S}\left[\hat{q}_{\boldsymbol{t}_{n+1}}^{\boldsymbol{F}_{n+1}}(\boldsymbol{f}_{n+1})\right] = \operatorname{tr}_{S}\left[\hat{P}_{\boldsymbol{t}_{n+1}}^{\boldsymbol{F}_{n+1}}(\boldsymbol{f}_{n+1})\hat{q}_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n})\right] = \operatorname{tr}_{S}\left[\hat{P}_{\boldsymbol{t}_{n+1}}^{\boldsymbol{F}_{n+1}}(\boldsymbol{f}_{n+1})\hat{q}_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n})\hat{P}_{\boldsymbol{t}_{n+1}}^{\boldsymbol{F}_{n+1}}(\boldsymbol{f}_{n+1})\right].$$
(6.23)

This recurrence equation, with the initial condition $\hat{q}_{t_1}^{F_1}(f_1) = \hat{P}_{t_1}^{F_1}(f_1)\hat{\rho}_{t_0}^p\hat{P}_{t_1}^{F_1}(f_1)$, admits the following solution:

$$\hat{q}_{\boldsymbol{t}_n}^{\boldsymbol{F}_n}(\boldsymbol{f}_n) = \hat{P}_{t_n}^{F_n}(f_n) \cdots \hat{P}_{t_1}^{F_1}(f_1) \hat{\rho}_{t_0}^p \hat{P}_{t_1}^{F_1}(f_1) \cdots \hat{P}_{t_n}^{F_n}(f_n) \ge 0, \qquad (6.24)$$

which then leads to the parameterization of the diagonal part of the bi-probability,

$$P_{\boldsymbol{t}_{n}|t_{0}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n}|p) = \operatorname{tr}_{S}\left[\hat{P}_{t_{n}}^{F_{n}}(f_{n})\cdots\hat{P}_{t_{1}}^{F_{1}}(f_{1})\hat{\rho}_{t_{0}}^{p}\hat{P}_{t_{1}}^{F_{1}}(f_{1})\cdots\hat{P}_{t_{n}}^{F_{n}}(f_{n})\right] = Q_{\boldsymbol{t}_{n}|t_{0}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n},\boldsymbol{f}_{n}|\rho^{p}).$$
(6.25)

We can generalize it to the off-diagonal of $Q_{t_n|t_0}^{F_n}$ by making the use of the inconsistency relation (5.10). First, we have

$$\begin{aligned} P_{t_n,\dots,t'_j,\dots,t_1|t_0}^{F_n,\dots,F_1}(f_n,\dots,f'_j,\dots,f_1|p) \\ &= \operatorname{tr}_S \left[\hat{P}_{t_n}^{F_n}(f_n) \cdots \hat{P}_{t_{j+1}}^{F_{j+1}}(f_{j+1}) \sum_{f_j \in \Omega(F_j)} \hat{P}_{t_j}^{F_j}(f_j) \hat{P}_{t_{j-1}}^{F_{j-1}}(f_{j-1}) \cdots \hat{P}_{t_1}^{F_1}(f_1) \hat{\rho}_{t_0}^p \right. \\ & \times \hat{P}_{t_1}^{F_1}(f_1) \cdots \hat{P}_{t_{j-1}}^{F_{j-1}}(f_{j-1}) \sum_{f'_j \in \Omega(F_j)} \hat{P}_{t_j}^{F_j}(f'_j) \hat{P}_{t_{j+1}}^{F_{j+1}}(f_{j+1}) \cdots \hat{P}_{t_n}^{F_n}(f_n) \right], \end{aligned}$$

and so Eq. (5.10) can be rewritten as

$$\sum_{\substack{f_j^+ \neq f_j^- \\ f_j^- F_j^-$$

Since the inconsistency relation holds for all $n \in \mathbb{N}$, $1 \leq j \leq n$, $f_j \in \Omega(F_j)$, $t_n > \cdots > t_1 > t_0 > 0$, we arrive at the following conclusion:

(Q8) Hilbert space representation

$$Q_{\boldsymbol{t}_{n}|t_{0}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n}^{+},\boldsymbol{f}_{n}^{-}|\rho) = \operatorname{tr}_{S}\Big[\Big(\prod_{j=n}^{1}\hat{P}_{t_{j}}^{F_{j}}(f_{j}^{+})\Big)\hat{\rho}_{t_{0}}\Big(\prod_{j=1}^{n}\hat{P}_{t_{j}}^{F_{j}}(f_{j}^{-})\Big)\Big],$$

where $\prod_{j=n}^{1} \hat{A}_j$ $(\prod_{j=1}^{n} \hat{A}_j)$ denotes an ordered product of operators $\hat{A}_n \cdots \hat{A}_1$ $(\hat{A}_1 \cdots \hat{A}_n)$. In particular,

$$P_{t_n|t_0}^{F_n}(f_n|p) = I_{t_n|t_0}^{F_n}(f_n, f_n|p) = \operatorname{Re} Q_{t_n|t_0}^{F_n}(f_n, f_n|\rho) = Q_{t_n|t_0}^{F_n}(f_n, f_n|\rho).$$

Here, $\{\hat{P}_{t_j}^{F_j}(f_j) \mid f_j \in \Omega(F_j)\}$ are the orthogonal projectors associated with the observable measured by the F_j -device deployed at time t_j , the metric matrix $\hat{\rho}_{t_0}$ is given by

$$\hat{\rho}_{t_0} = \sum_K \sum_{k \in \Omega(K)} p(K,k) \hat{P}_{t_0}^K(k) = \sum_K \sum_{k \in \Omega(K)} p(K,k) |\Psi_{t_0}^K(k)\rangle \langle \Psi_{t_0}^K(k)|$$

and it represents the initialization event defined by the phenomenological distribution p(K, k) referring to measurements done with perfectly fine-grained K-devices deployed at time t_0 .

This result demonstrates the limitations of the measurement-projector link: the alluring idea that "the action of the *F*-device at time *t* is in one-to-one correspondence to the action of the projector operator $\hat{P}_t^F(f)$ ", which works for a single measurement, does not extend to sequential measurements. To see this, notice that, in the formal representation of a probability distribution for a sequence of measurements,

$$P_{\boldsymbol{t}_{n}|t_{0}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n}|p) = \operatorname{tr}_{S}\left[\left(\prod_{j=n}^{1}\hat{P}_{t_{j}}^{F_{j}}(f_{j})\right)^{\dagger}\left(\prod_{j=n}^{1}\hat{P}_{t_{j}}^{F_{j}}(f_{j})\right)\hat{\rho}_{t_{0}}\right] \equiv \operatorname{tr}_{S}\left[\hat{E}_{\boldsymbol{t}_{n}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n})\hat{\rho}_{t_{0}}\right],\tag{6.27}$$

the operator $\hat{E}_{t_n}^{F_n}(f_n)$ is generally *not* a projector. Therefore, given that $\hat{\rho}_{t_0}$ represents the initialization event, we cannot say that the action of a sequence of measuring devices is represented in an analogous way as the action of a single measuring device. Moreover, it is not true that the action of a sequence of devices [represented by $\hat{E}_{t_n,\dots,t_1}^{F_n,\dots,F_1}(f_n,\dots,f_1)$] corresponds to a sequence of actions of single devices [represented by $\prod_{j=n}^1 \hat{P}_{t_j}^{F_j}(f_j)$], even though the Hilbert-space representations of both consist of the same elementary building blocks—the projectors $\hat{P}_{t_j}^{F_j}(f_j)$. This leads us to the following conclusion: since, in general, the results of a sequence of measurements cannot be represented by a deployment of a single measuring device following immediately the initialization event, the quantum statics picture is incapable of describing the context of sequential measurements. Thus, the metric $\hat{\rho}_{t_0}$ cannot be identified as the master object of the formalism in non-static (dynamical) contexts.

7 Dynamical laws of the system

7.1 Unitary evolution operator

Consider an arbitrarily chosen observable K of the system S that is measured with a perfectly fine-grained K-device. Then, the projector operators associated with K are all rank-1, $\hat{P}_t^K(k) = |\Psi_t^K(k)\rangle\langle\Psi_t^K(k)|$, with $\mathbb{K}_t = \{|\Psi_t^K(k)\rangle \in \mathcal{H}_S \mid k \in \Omega(K)\}$ forming an orthonormal basis in the Hilbert space of the system. Since the operators $\{\hat{P}_t^K(k) \mid k \in \Omega(K)\}$ form a complete set of orthogonal projectors, they can be simultaneously diagonalized,

$$\hat{U}_t^K \hat{P}_t^K(k) (\hat{U}_t^K)^{\dagger} = \operatorname{diag}(\dots, 0, 1, 0 \dots), \quad \text{for all } k \in \Omega(K).$$

$$(7.1)$$

Here, the position of 1 in the diagonal corresponds to the value k of the projector, and we choose $\hat{U}_0^K = \hat{1}_S$ so that the matrices of $\hat{P}_0^K(k)$ are already diagonal. Then, formally, each unitary operator \hat{U}_t^K executes the basis transformation from \mathbb{K}_0 to \mathbb{K}_t —the basis in which the matrices representing each $\hat{P}_t^K(k)$ are all diagonal; alternatively, one can simply write

$$\hat{U}_{t}^{K}|\Psi_{t}^{K}(k)\rangle = |\Psi_{0}^{K}(k)\rangle, \quad (\hat{U}_{t}^{K})^{\dagger}\hat{P}_{0}^{K}(k)\hat{U}_{t}^{K} = \hat{P}_{t}^{K}(k), \quad \text{for all } k \in \Omega(K).$$
(7.2)

We can say that the operator-valued function $t \mapsto \hat{U}_t^K$ is the element of the Hilbert space representation responsible for the time-evolution of the projectors associated with the observable K. Since we associate any of its coarse-grained variants \overline{K} with projectors that are combinations of K-projectors, \hat{U}_t^K also works as an evolution operator for all \overline{K} ,

$$\hat{P}_{t}^{\overline{K}}(\overline{k}) = \sum_{k \in \omega(\overline{k})} \hat{P}_{t}^{K}(k) = \sum_{k \in \omega(\overline{k})} (\hat{U}_{t}^{K})^{\dagger} \hat{P}_{0}^{K}(k) \hat{U}_{t}^{K} = (\hat{U}_{t}^{K})^{\dagger} \Big(\sum_{f \in \omega(\overline{k})} \hat{P}_{0}^{K}(k) \Big) \hat{U}_{t}^{K} = (\hat{U}_{t}^{K})^{\dagger} \hat{P}_{0}^{\overline{K}}(\overline{k}) \hat{U}_{t}^{K}.$$
(7.3)

However, any fine-grained observable L inequivalent to K—and thus associated with projectors onto elements of bases $\mathbb{L}_t = \{|\Psi_t^L(l)\rangle \in S \mid l \in \Omega(L)\} \neq \mathbb{K}_t$ —will have its own evolution operator $t \mapsto \hat{U}_t^L$ that could be, in principle, distinct from \hat{U}_t^K . This begs the following question: what is the actual relation between the respective transformations \hat{U}_t^K and \hat{U}_t^L ?

We can answer this question by invoking the phenomenological observation 11: the correlation function with the following Hilbert space representation

$$P_{t+\Delta t|t}^{L|K}(l|k) = \frac{Q_{t+\Delta t,t|0}^{L,K}(l,k|\rho)}{Q_{t|0}^{K}(k,k|\rho)}$$

$$= \frac{\operatorname{tr}_{S}\left[|\Psi_{t+\Delta t}^{L}(l)\rangle\langle\Psi_{t+\Delta t}^{L}(l)|\Psi_{t}^{K}(k)\rangle\langle\Psi_{t}^{K}(k)|\hat{\rho}_{0}|\Psi_{t}^{K}(k)\rangle\langle\Psi_{t}^{K}(k)|\hat{\rho}_{0}\right]}{\operatorname{tr}_{S}\left[|\Psi_{t}^{K}(k)\rangle\langle\Psi_{t}^{K}(k)|\hat{\rho}_{0}\right]}$$

$$= \frac{|\langle\Psi_{t+\Delta t}^{L}(l)|\Psi_{t}^{K}(k)\rangle|^{2}\langle\Psi_{t}^{K}(k)|\hat{\rho}_{0}|\Psi_{t}^{K}(k)\rangle}{\langle\Psi_{t}^{K}(k)|\hat{\rho}_{0}|\Psi_{t}^{K}(k)\rangle} \xrightarrow{\Delta t \to 0^{+}} |\langle\Psi_{t}^{L}(l)|\Psi_{t}^{K}(k)\rangle|^{2}, \quad (7.4)$$

is found to be *independent* of the time t when the measurements were taken. Therefore, we obtain $|\langle \Psi_t^L(l)|\Psi_t^K(k)\rangle| = |\langle \Psi_0^L(l)|\Psi_0^K(k)\rangle|$ for all $l \in \Omega(L)$, $k \in \Omega(K)$ and $t \ge 0$, which then leads to the following system of equations,

$$\begin{split} |\langle \Psi_0^L(l) | \Psi_0^K(k) \rangle| &= |\langle \Psi_0^L(l) | \hat{U}_t^L(\hat{U}_t^K)^\dagger | \Psi_0^K(k) \rangle|; \quad |\langle \Psi_t^L(l) | \Psi_t^K(k) \rangle| = |\langle \Psi_t^L(l) | (\hat{U}_t^F)^\dagger \hat{U}_t^K | \Psi_t^K(k) \rangle|; \\ |\langle \Psi_0^K(k) | \Psi_0^L(l) \rangle| &= |\langle \Psi_0^K(k) | \hat{U}_t^K(\hat{U}_t^L)^\dagger | \Psi_0^L(l) \rangle|; \quad |\langle \Psi_t^K(k) | \Psi_t^L(l) \rangle| = |\langle \Psi_t^K(k) | (\hat{U}_t^K)^\dagger \hat{U}_t^L | \Psi_t^L(l) \rangle|, \end{split}$$

that admits a simple solution

$$\hat{U}_t^L = \mathrm{e}^{-\mathrm{i}\theta_t} \hat{U}_t^K. \tag{7.5}$$

The gauge phase θ_t can be set to zero without loss of generality because it does not influence the transformation of the projectors,

$$\hat{P}_{t}^{L}(l) = (\hat{U}_{t}^{L})^{\dagger} \hat{P}_{0}^{F}(l) \hat{U}_{t}^{L} = \mathrm{e}^{\mathrm{i}\theta_{t}} (\hat{U}_{t}^{K})^{\dagger} \hat{P}_{0}^{L}(l) \hat{U}_{t}^{K} \mathrm{e}^{-\mathrm{i}\theta_{t}} = (\hat{U}_{t}^{K})^{\dagger} \hat{P}_{0}^{L}(l) \hat{U}_{t}^{K}.$$
(7.6)

Crucially, since K and L were chosen arbitrarily, we can also drop the superscripts—in other words, there is only one operator-valued function $t \mapsto \hat{U}_t$ that describes the evolution for all observables.

Therefore, it stands to reason that, analogously to the Hilbert space of the system \mathcal{H}_S , the evolution operator \hat{U}_t can be ascribed to the quantum system itself; given what it does, it is fair to say that it represents the *dynamical law* of the system.

7.2 Generator of the evolution operator

The short-time behavior of the survival probability (4.32), which was implied by the phenomenology of the Zeno effect, inevitably constraints the possible forms of time-dependence in the evolution operator \hat{U}_t . In particular, we can immediately conclude that the function $t \mapsto \hat{U}_t$ has to be continuous and differentiable at each t, so that it possesses a well-behaved short-time series expansion,

$$\hat{U}_{t+\Delta t} = \hat{U}_t + \Delta t \frac{\mathrm{d}\hat{U}_t}{\mathrm{d}t} + \frac{\Delta t^2}{2} \frac{\mathrm{d}^2 \hat{U}_t}{\mathrm{d}t^2} + O(\Delta t^3), \tag{7.7}$$

which could then produce the corresponding expansion for the probability,

$$P_{t+\Delta t|t}^{K|K}(k|k) = |\langle \Psi_0^K(k)|\hat{U}_{t+\Delta t}\hat{U}_t^{\dagger}|\Psi_0^K(k)\rangle|^2$$

$$\begin{array}{l} \xrightarrow{\Delta t \to 0} 1 + \Delta t \langle \Psi_0^K(k) | \left(\frac{\mathrm{d}\hat{U}_t}{\mathrm{d}t} \hat{U}_t^{\dagger} + \hat{U}_t \frac{\mathrm{d}\hat{U}_t^{\dagger}}{\mathrm{d}t} \right) | \Psi_0^K(k) \rangle \\ + \frac{\Delta t^2}{2} \langle \Psi_0^K(k) | \left(\frac{\mathrm{d}^2 \hat{U}_t}{\mathrm{d}t^2} \hat{U}_t^{\dagger} + \hat{U}_t \frac{\mathrm{d}^2 \hat{U}_t^{\dagger}}{\mathrm{d}t^2} \right) | \Psi_0^K(k) \rangle \\ + \Delta t^2 \langle \Psi_0^K(k) | \frac{\mathrm{d}\hat{U}_t}{\mathrm{d}t} \hat{U}_t^{\dagger} | \Psi_0^K(k) \rangle \langle \Psi_0^K(k) | \hat{U}_t \frac{\mathrm{d}\hat{U}_t^{\dagger}}{\mathrm{d}t} | \Psi_0^K(k) \rangle + O(\Delta t^3).$$
(7.8)

We argued in Section 4.5 that, in the expansion above, the linear term vanishes and the quadratic term is negative; consequently, these phenomenological inputs constraint the form of \hat{U}_t even further:

$$\frac{\mathrm{d}\hat{U}_t}{\mathrm{d}t}\hat{U}_t^{\dagger} = -\hat{U}_t \frac{\mathrm{d}\hat{U}_t^{\dagger}}{\mathrm{d}t}, \quad \frac{\mathrm{d}^2\hat{U}_t}{\mathrm{d}t^2}\hat{U}_t^{\dagger} = \hat{U}_t \frac{\mathrm{d}^2\hat{U}_t^{\dagger}}{\mathrm{d}t^2}, \quad |\langle\phi|\frac{\mathrm{d}\hat{U}_t}{\mathrm{d}t}\hat{U}_t^{\dagger}|\phi\rangle|^2 \le -\langle\phi|\frac{\mathrm{d}^2\hat{U}_t}{\mathrm{d}t^2}\hat{U}_t^{\dagger}|\phi\rangle, \tag{7.9}$$

for all $|\phi\rangle$ and t > 0. Bearing these conditions in mind, we can now use the continuity (7.7) of \hat{U}_t to determine its exact form. Given t > 0, let $s_j = jt/n$: then

$$\hat{U}_{t} = \lim_{n \to \infty} \hat{U}_{t} \prod_{j=n-1}^{1} \hat{U}_{s_{j}}^{\dagger} \hat{U}_{s_{j}} = \lim_{n \to \infty} \hat{U}_{t} \hat{U}_{s_{n-1}}^{\dagger} \left(\prod_{j=n-2}^{1} \hat{U}_{s_{j}+\frac{t}{n}} \hat{U}_{s_{j}}^{\dagger} \right) \hat{U}_{s_{1}} = \lim_{n \to \infty} \prod_{j=n-1}^{0} \hat{U}_{s_{j}+\frac{t}{n}} \hat{U}_{s_{j}}^{\dagger}
= \lim_{n \to \infty} \prod_{j=n-1}^{0} \left(\hat{1} + \frac{t}{n} \frac{\mathrm{d}\hat{U}_{s_{j}}}{\mathrm{d}s_{j}} \hat{U}_{s_{j}}^{\dagger} + O(n^{-2}) \right) = \sum_{n=0}^{\infty} \int_{0}^{t} \mathrm{d}s_{n} \cdots \int_{0}^{s_{2}} \mathrm{d}s_{1} \prod_{j=n}^{1} \frac{\mathrm{d}\hat{U}_{s_{j}}}{\mathrm{d}s_{j}} \hat{U}_{s_{j}}^{\dagger}
= \mathcal{T} \exp\left[\int_{0}^{t} \frac{\mathrm{d}\hat{U}_{s}}{\mathrm{d}s} \hat{U}_{s}^{\dagger} \mathrm{d}s \right],$$
(7.10)

and thus, to satisfy all conditions (7.9) and $\hat{U}_t^{\dagger}\hat{U}_t = \hat{U}_t\hat{U}_t^{\dagger} = \hat{1}_S$ for all t > 0, we must have

$$\frac{\mathrm{d}\hat{U}_s}{\mathrm{d}s} = -\mathrm{i}\hat{H}(s)\hat{U}_s,\tag{7.11}$$

where $\hat{H}(s)$ is a Hermitian operator-valued function on [0, t]. Therefore, we have arrived at the expected exponential form of the evolution operator, generated by the *Hamiltonian* of the system,

$$\hat{U}_{t,t'} := \hat{U}_t \hat{U}_{t'}^{\dagger} = \mathcal{T} e^{-i \int_{t'}^{t} \hat{H}(s) ds}.$$
(7.12)

We conclude with a clarification: The method of "initialization through measurement" as the means of control over the initial conditions of the experiment, that was introduced back in Section 4.4, implicitly assumes the *stationarity* of probability distributions, i.e.,

$$P_{\boldsymbol{t}_n|0}^{\boldsymbol{F}_n}(\boldsymbol{f}_n|p) = P_{t_n+t_0,\dots,t_1+t_0|t_0}^{\boldsymbol{F}_n}(\boldsymbol{f}_n|p) \quad \text{for any } t_0 > 0.$$
(7.13)

Otherwise, even with Markovianity at our disposal, it would not be possible to reset the "state" of the system, as the results of any given experiment would depend on the absolute time it started. In other words, if the results were to depend on the *date* the experiment is taking place, then it would not be possible to compare the results of its past and future runs. But now that we have the detailed formal representation of the dynamics, we are obliged to determine how the requirement of stationarity translates into a constraint on the form of Hamiltonian.

When the stationarity property (7.13) is expressed within the Hilbert space representation,

$$\operatorname{tr}_{S}\left[\left(\prod_{j=n}^{1}\hat{P}_{0}^{F_{j}}(f_{j})\hat{U}_{t_{j}+t_{0},t_{j-1}+t_{0}}\right)\hat{\rho}_{0}\left(\prod_{j=1}^{n}\hat{U}_{t_{j}+t_{0},t_{j-1}+t_{0}}^{\dagger}\hat{P}_{0}^{F_{j}}(f_{j})\right)\right]$$
$$=\operatorname{tr}_{S}\left[\left(\prod_{j=n}^{1}\hat{P}_{0}^{F_{j}}(f_{j})\hat{U}_{t_{j},t_{j-1}}\right)\hat{\rho}_{0}\left(\prod_{j=1}^{n}\hat{U}_{t_{j},t_{j-1}}^{\dagger}\hat{P}_{0}^{F_{j}}(f_{j})\right)\right].$$
(7.14)

we conclude that the probabilities can be stationary only when the evolution operator satisfies the following condition,

$$\mathcal{T}e^{-i\int_{t_{j-1}+t_0}^{t_j+t_0}\hat{H}(s)ds} = \mathcal{T}e^{-i\int_{t_{j-1}}^{t_j}\hat{H}(s-t_0)ds} = \mathcal{T}e^{-i\int_{t_{j-1}}^{t_j}\hat{H}(s)ds} \quad \text{for all } t_{j-1} < t_j \text{ and } t_0 > 0.$$
(7.15)

which means that the Hamiltonian has to be time-independent, $\hat{H}(s) = \hat{H}$, in which case we obtain

$$\hat{U}_{t,t'} = \sum_{n=0}^{\infty} (-i)^n \frac{(t-t')^n}{n!} \hat{H}^n = e^{-i(t-t')\hat{H}}.$$
(7.16)

Once we deduce how one might introduce the concept of external fields into the formalism, we shall revisit the physical viability of time-dependent Hamiltonians.

8 Composite systems

Previously, we deduced that every quantum system is formally characterized by a Hilbert space \mathcal{H}_S and a Hamiltonian \hat{H} that generates the evolution operator $\hat{U}_{t,t'} = \exp[-i(t-t')\hat{H}]$ representing the system's dynamical law. We also established that bi-probabilities associated with observables of independent systems factorize in a manner analogous to probability distributions of independent stochastic variables. We can conclude from these facts that the Hilbert space representing a system composed of independent subsystems has the form of the tensor product of the Hilbert spaces of the constituents,

$$\mathcal{H}_{AB} = \mathcal{H}_A \otimes \mathcal{H}_B. \tag{8.1}$$

This means that the dynamical law of a system composed of independent subsystems is defined as the product of the evolution operators of its subsystems,

$$\hat{U}_{t,t'}^{AB} := \hat{U}_{t,t'}^A \otimes \hat{U}_{t,t'}^B, \tag{8.2}$$

which implies

$$\hat{H}_{AB} = \hat{H}_A \otimes \hat{1}_B + \hat{1}_A \otimes \hat{H}_B.$$
(8.3)

This tensor product structure extends to all operators involved in the Hilbert space representation: the density matrix of the total system factorizes,

$$\hat{\rho}_{AB} = \hat{\rho}_A \otimes \hat{\rho}_B, \tag{8.4}$$

and the projectors associated with subsystem observables also have the product form. For instance, $\{\hat{P}^{F_A}(f) \otimes \hat{1}_B \mid f \in \Omega(F_A)\}$ corresponds to the *A*-only observable measured by the F_A -device, $\{\hat{1}_A \otimes \hat{P}^{F_B}(f) \mid f \in \Omega(F_B)\}$ corresponds to the *B*-only F_B -device, and thus, $\{\hat{P}^{F_AF_B}(f_a \wedge f_b) = \hat{P}^{F_A}(f_a) \otimes \hat{P}^{F_B}(f_b) \mid f_a \wedge f_b \in \Omega(F_AF_B) = \Omega(F_A) \times \Omega(F_B)\}$ corresponds to the observable measured by an $F_A F_B$ -device, i.e., an F_A - an F_B -device deployed in tandem.

Therefore, subsystems cease to be independent when initialization is not in a product form, $\hat{\rho}_{AB} \neq \hat{\rho}_A \otimes \hat{\rho}_B$, or when the Hamiltonian includes a *coupling* operator, e.g.,

$$\hat{H}_{AB} = \hat{H}_A \otimes \hat{1}_B + \hat{1}_A \otimes \hat{H}_B + \lambda \hat{V}_A \otimes \hat{V}_B, \qquad (8.5)$$

so that $\hat{U}_{t,t'}^{AB} \neq \hat{U}_{t,t'}^{A} \otimes \hat{U}_{t,t'}^{B}$. When the subsystems are not independent because of interactions, the influence of one system onto another is compactly quantified by bi-probabilities associated with subsystem observables [15, 16],

$$Q_{t_{n}}^{F_{n}}(f_{n}^{+}, f_{n}^{-}|\rho_{A}) = \operatorname{tr}\left[\left(\prod_{j=n}^{1} \hat{U}_{0,t_{j}}^{AB} \hat{P}_{0}^{F_{j}}(f_{j}^{+}) \otimes \hat{1}_{B} \hat{U}_{t_{j},0}^{AB}\right) \hat{\rho}_{A} \otimes \hat{\rho}_{B}\left(\prod_{j=n}^{1} \hat{U}_{0,t_{j}}^{AB} \hat{P}_{0}^{F_{j}}(f_{j}^{-}) \otimes \hat{1}_{B} \hat{U}_{t_{j},0}^{AB}\right)\right] \\ = \iint \operatorname{tr}_{A}\left[\left(\prod_{j=n}^{1} \hat{P}_{t_{j}}^{F_{j}}[b^{+}](f_{j}^{+})\right) \hat{\rho}_{A}\left(\prod_{j=1}^{n} \hat{P}_{t_{j}}^{F_{j}}[b^{-}](f_{j}^{-})\right)\right] \mathcal{Q}^{B}[b^{+}, b^{-}][\mathcal{D}b^{+}][\mathcal{D}b^{-}],$$

$$(8.6)$$

where each $\{\hat{P}^{F_j}(f) \otimes \hat{1}_B \mid f \in \Omega(F_j)\}$ corresponds to the observable of the subsystem A measurable by the F_j -device, and

$$\hat{P}_{t}^{F_{j}}[h](f) = \left(\mathcal{T}\mathrm{e}^{-\mathrm{i}\int_{0}^{t}(\hat{H}_{A}+\lambda V_{B}(h(s))\hat{V}_{A})\mathrm{d}s}\right)^{\dagger}\hat{P}_{0}^{F_{j}}(f)\mathcal{T}\mathrm{e}^{-\mathrm{i}\int_{0}^{t}(\hat{H}_{A}+\lambda V_{B}(h(s))\hat{V}_{A})\mathrm{d}s},\tag{8.7}$$

where the projectors $\{\hat{P}_0^B(b) \mid b \in \Omega(B)\}$ and the function $V_B : \Omega(B) \to \mathbb{R}$ are the solution of the eigenproblem of the coupling operator \hat{V}_B ,

$$\hat{V}_B \hat{P}_0^B(b) = V_B(b) \hat{P}_0^B(b) \quad \text{for all } b \in \Omega(B).$$

$$(8.8)$$

We take note of two important features of Eq. (8.6). The first is that the unitary evolution operators in Eq. (8.7) are generated by the time-dependent Hamiltonian $\hat{H}[h](s) = \hat{H}_A + \lambda V_B(h(s))\hat{V}_A$. The real-valued function $V_B(h(s))$, which introduces a time-dependence into the Hamiltonian, can be interpreted as an external field driving the dynamics via coupling with the operator \hat{V}_A . Of course, in Eq. (8.6), the role of the external driver is played by the components of the bi-trajectory $(b^+(s), b^-(s))$: $b^+(s)$ is coupled to the sequence progressing forwards in time, while $b^-(s)$ drives the sequence going backwards in time, thereby facilitating the quantum interference between these two branches of the evolution.

In the special case where $\mathcal{Q}^B[b^+, b^-][\mathcal{D}b^+][\mathcal{D}b^-] = \delta[b^+ - b^0]\delta[b^- - b^0][\mathcal{D}b^+][\mathcal{D}b^-]$, meaning that both bi-trajectories overlap with the one trajectory $t \mapsto b^0(t)$, the influence from B takes on the form identical to the driving with the given external classical force field [13]. Therefore, the formally open system A can be described in this scenario as a closed system with the dynamical law generated by the time-dependent Hamiltonian $\hat{H}(s) = \hat{H}_A + \lambda V_B(b^0(s))\hat{V}_A$.

The second important feature is that the projectors $\{\hat{P}_t^B(b) \mid b \in \Omega(B)\}$ corresponding to a *B*-only observable, which we associate with the bi-probabilities $Q_{t_n}^B$,

$$Q_{t_n}^B(b_n^+, b_n^-) = \operatorname{tr}_B \left[\left(\prod_{j=n}^1 \hat{P}_{t_j}^B(b_j^+) \right) \hat{\rho}_B \left(\prod_{j=1}^n \hat{P}_{t_j}^B(b_j^-) \right) \right] \\ = \iint \left(\prod_{j=1}^n \delta_{b^+(t_j), b_j^+} \delta_{b^-(t_j), b_j^-} \right) \mathcal{Q}^B[b^+, b^-][\mathcal{D}b^+][\mathcal{D}b^-]$$
(8.9)

originate from the formal spectral decomposition of the coupling operator \hat{V}_B . Specifically, given the solution of the eigenproblem (8.8) we have

$$\hat{V}_B = \sum_{b \in \Omega(B)} V_B(b) \hat{P}_0^B(b) \equiv \sum_{v \in \Omega(V_B)} v \hat{P}_0^{V_B}(v).$$
(8.10)

Generalizing this observation, any Hermitian operator $\hat{F} = \sum_{f \in \Omega(F)} f \hat{P}^F(f)$ can be understood as the Hilbert space representation of the observable probed by a measuring device associated with the set of projectors $\{\hat{P}^F(f) \mid f \in \Omega(F)\}$ with images overlapping the corresponding eigenspaces of \hat{F} . Then, it follows that quantum systems influence each other by means of the dynamics generated by the coupling of their observables—the observables that could be in principle probed by the measuring device deployed by the classical observer. Thus, we establish the familiar notion, known from the standard formalism, that the observables of quantum systems correspond to Hermitian operators acting in the system Hilbert space.

9 The master object of the formalism

Back in Section 1, when reviewing the structure of the standard formalism of quantum mechanics, we observed that the density matrix $\hat{\rho}_t$ can be considered as a master object of this formalism: For as long as one restricts their interests to single-measurement experiments (ignoring, for the moment, the issue of initialization), then, indeed, $\hat{\rho}_t$ is all that is needed in order to fully describe all the observable facts. As it was noted in Section 6.2, even the passage of time can be reduced in this setting to observer choosing an appropriate measuring device. Hence, the single-measurement

context can be considered as quantum statics, analogous in its scope to, e.g., hydrostatics of classical fluid mechanics. It is then not surprising that for quantum statics, similarly to its classical analogues, it is a natural choice to designate the "system state" as the master object of the formalism.

Moving beyond "quantum statics" to "quantum dynamics" that involve scenarios such as sequential measurements, open quantum systems (cf. Section 8), etc., the state $\hat{\rho}$ is dethroned by bi-probabilities as the master object, and the role of density matrix is reduced to a convenient representation of the initial condition. In the setting where only a single observable F is relevant, as demonstrated in Section 5, all bi-probabilities $Q_{t_n}^F$ associated with F are simply the discretetime restrictions of the bi-trajectory measure $Q^F[f^+, f^-][\mathcal{D}f^+][\mathcal{D}f^-]$. It is thus revealed that, in this limited—but still dynamical—context, the status of master object belongs to the bi-trajectory measure, as it is the source of all other objects in the formalism.

However, unlike classical theories, quantum systems accommodate an infinite number of inequivalent observables, and in the general context, the formalism must utilize bi-probabilities $Q_{t_n|t_0}^{F_n}$ associated with multiple observables. Therefore, just like the density matrix before it, the master bi-trajectory measure $Q^F[f^+, f^-][\mathcal{D}f^+][\mathcal{D}f^-]$ surely has to be dethroned by something else. A natural candidate would be a bi-trajectory measure that extends *all* multi-observable bi-probabilities; the question is if such an object actually exists.

To answer this question, we must first introduce a higher level of abstraction to the formalism by defining *system* bi-probabilities that are not associated with any observable or initialization event. To begin with, note that any rank-1 projector in the system Hilbert space can be written as

$$\hat{P}_{t}^{K}(k) = |\Psi_{t}^{K}(k)\rangle\langle\Psi_{t}^{K}(k)| = e^{it\hat{H}}|\Psi_{0}^{K}(k)\rangle\langle\Psi_{0}^{K}(k)|e^{-it\hat{H}} \\
= e^{it\hat{H}}e^{i\sum_{\ell=1}^{d^{2}-1}\tau_{K}^{\ell}\hat{T}_{\ell}}|\eta(k)\rangle\langle\eta(k)|e^{-i\sum_{\ell=1}^{d^{2}-1}\tau_{K}^{\ell}\hat{T}_{\ell}}e^{-it\hat{H}} \\
\equiv \hat{U}_{t}^{\dagger}\hat{S}_{\vec{\tau}_{K}}^{\dagger}|\eta(k)\rangle\langle\eta(k)|\hat{S}_{\vec{\tau}_{K}}\hat{U}_{t},$$
(9.1)

where $d = \dim \mathcal{H}_S$, $\mathbb{E} = \{|\eta\rangle \in \mathcal{H}_S \mid \eta = 1, \dots, d, \langle \eta | \eta' \rangle = \delta_{\eta,\eta'} \}$ is an arbitrary reference orthonormal basis, $\{\hat{T}_\ell \mid \ell = 1, \dots, d^2 - 1\}$ is the set of Hermitian Lie algebra generators for unitary operators in \mathcal{H}_S , and $\vec{\tau}_K = (\tau_K^1, \dots, \tau_K^{d^2-1})$ is a vector of real coordinates generating the basis transformation between \mathbb{E} and $\mathbb{K}_0 = \{|\Psi_0^K(k)\rangle \in \mathcal{H}_S \mid k \in \Omega(K)\}$ (cf. Section 7). Then we define a family of bi-probabilities that, instead of time and observables, use the "space-time" d^2 -component coordinates $\tau_j = (\tau_j^0, \vec{\tau}_j) = (\tau_j^0, \tau_j^1, \dots, \tau_j^{d^2-1})$ as indexes,

$$Q_{\boldsymbol{\tau}_{n}|\tau_{0}}(\boldsymbol{\eta}_{n}^{+},\boldsymbol{\eta}_{n}^{-}|\boldsymbol{\eta}_{0}^{+},\boldsymbol{\eta}_{0}^{-}) := \operatorname{tr}\left[\left(\prod_{j=n}^{0}\hat{P}_{\tau_{j}}(\boldsymbol{\eta}_{j}^{+})\right)\left(\prod_{j=0}^{n}\hat{P}_{\tau_{j}}(\boldsymbol{\eta}_{j}^{-})\right)\right], \text{ where } \hat{P}_{\tau}(\boldsymbol{\eta}) := \hat{U}_{\tau^{0}}^{\dagger}\hat{S}_{\vec{\tau}}^{\dagger}|\boldsymbol{\eta}\rangle\langle\boldsymbol{\eta}|\hat{S}_{\vec{\tau}}\hat{U}_{\tau^{0}}.$$
(9.2)

At the moment these functions are purely abstract elements of the formalism, as we have not yet established any relations between them and the phenomenology. On the other hand, we see that, in the hierarchy of objects constituting the formalism we have deduced so far, the system bi-probabilities $Q_{\tau_n|\tau_0}$ are one step above their observable-associated cousins $Q_{t_n|t_0}^{F_n}$, as the latter decompose into convex combinations of the former,

$$Q_{\boldsymbol{t}_{n}|t_{0}}^{\boldsymbol{F}_{n}}(\boldsymbol{f}_{n}^{+},\boldsymbol{f}_{n}^{-}|\boldsymbol{\rho}) = \sum_{\boldsymbol{\eta}_{n}^{\pm},\eta_{0}} \Big(\prod_{j=1}^{n} \delta_{F_{j}(\boldsymbol{\eta}_{j}^{+}),f_{j}^{+}} \delta_{F_{j}(\boldsymbol{\eta}_{j}^{-}),f_{j}^{-}} \Big) Q_{(t_{n},\vec{\tau}_{n}),\dots,(t_{1},\vec{\tau}_{1})|(t_{0},\vec{\tau}_{0})}(\boldsymbol{\eta}_{n}^{+},\boldsymbol{\eta}_{n}^{-}|\eta_{0},\eta_{0})\boldsymbol{\rho}(\eta_{0}),$$
(9.3)

where the "spatial"-coordinates $\vec{\tau}_n, \ldots, \vec{\tau}_1$ and $\vec{\tau}_0$ are each found by solving the corresponding eigenproblem,

$$\hat{\rho}_{t_0}\hat{P}_{(t_0,\vec{\tau}_0)}(\eta) = \rho(\eta)\hat{P}_{(t_0,\vec{\tau}_0)}(\eta), \quad \left(\sum_{f\in\Omega(F_j)} f\hat{P}_{t_j}^{F_j}(f)\right)\hat{P}_{(t_j,\vec{\tau}_j)}(\eta) = F_j(\eta)\hat{P}_{(t_j,\vec{\tau}_j)}(\eta). \tag{9.4}$$

Therefore, the system bi-probabilities $Q_{\tau_n|\tau_0}$ are master objects with respect to observable-associated bi-probabilities, but it is only through $Q_{t_n|t_0}^{F_n}$ that they can be connected with the phenomenology.

It is easy to verify that the functions $Q_{\tau_n|\tau_0}$ satisfy all properties (Q1)–(Q5), including the crucial bi-consistency. Therefore, the natural next step would be to extend the family of system bi-probabilities to a bi-trajectory measure, as it was done previously for single-observable $Q_{t_n}^F$ (see Section 5). That is, we would like to say that there exists a bi-trajectory measure such that

$$Q_{\boldsymbol{\tau}_{n}|\tau_{0}}(\boldsymbol{\eta}_{n}^{+},\boldsymbol{\eta}_{n}^{-}|\boldsymbol{\eta}_{0}^{+},\boldsymbol{\eta}_{0}^{-}) = \iint \Big(\prod_{j=0}^{n} \delta_{\boldsymbol{\eta}^{+}(\tau_{j}),\boldsymbol{\eta}_{j}^{+}} \delta_{\boldsymbol{\eta}^{-}(\tau_{j}),\boldsymbol{\eta}_{j}^{-}} \Big) \mathcal{Q}[\boldsymbol{\eta}^{+},\boldsymbol{\eta}^{-}][\mathcal{D}\boldsymbol{\eta}^{+}][\mathcal{D}\boldsymbol{\eta}^{-}], \qquad (9.5)$$

where $\mathcal{Q}[\eta^+, \eta^-][\mathcal{D}\eta^+][\mathcal{D}\eta^-]$ is a complex measure on the space of *field* pairs,

$$\mathbb{R} \times S^{d^2 - 1} \ni \tau = (\tau^0, \vec{\tau}) \mapsto (\eta^+(\tau^0, \vec{\tau}), \eta^-(\tau^0, \vec{\tau})) \in \{1, \dots, d\} \times \{1, \dots, d\},$$
(9.6)

 $(S^n \text{ indicates here the hyper-sphere of dimension } n).$

Unfortunately, for several non-trivial reasons (see [17] for a detailed explanation), this time we cannot rely on the generalized Kolmogorov extension theorem to prove the existence of the master bi-trajectory measure. Nevertheless, even in the absence of a formal proof, we can still formulate a strong supporting argument, which goes as follows.

Assuming that the master measure exists, we can define the following super-operator:

$$\Lambda_{t} := \iint \left(\mathcal{T} e^{-i \int_{0}^{t} \hat{H}[\eta^{+}](s) ds} \right) \bullet \left(\mathcal{T} e^{-i \int_{0}^{t} \hat{H}[\eta^{-}](s) ds} \right)^{\dagger} \\ \times \sqrt{\rho(\eta^{+}(0, \vec{\tau}_{\rho}))\rho(\eta^{-}(0, \vec{\tau}_{\rho}))} \mathcal{Q}[\eta^{+}, \eta^{-}][\mathcal{D}\eta^{+}][\mathcal{D}\eta^{-}]$$
(9.7)

with $\hat{\rho}\hat{S}_{\vec{\tau}_{\rho}}|\eta\rangle = \rho(\eta)\hat{S}_{\vec{\tau}_{\rho}}|\eta\rangle$ an arbitrary density matrix and

$$\hat{H}[\eta](s) := \hat{H}_0 + \sum_{\alpha=1}^N F_\alpha \big(\eta(s, \vec{\tau}_\alpha) \big) \hat{H}_\alpha \quad : \quad \hat{F}_\alpha \hat{S}_{\vec{\tau}_\alpha} |\eta\rangle = F_\alpha(\eta) \hat{S}_{\vec{\tau}_\alpha} |\eta\rangle; \tag{9.8a}$$

where $\{\hat{H}_{\alpha} \mid \alpha = 0, 1, ..., N\}$ is a set of arbitrary Hermitian operators in some finite-dimensional Hilbert space \mathcal{H}_{O} , and $\{\hat{F}_{\alpha} = \sum_{f \in \Omega(F_{\alpha})} f\hat{P}_{0}^{F_{\alpha}}(f) \mid \alpha = 1, ..., N\}$ is an arbitrary set of observables in \mathcal{H}_{S} . Such Λ_{t} is a *characteristic function* of the measure $\mathcal{Q}[\eta^{+}, \eta^{-}][\mathcal{D}\eta^{+}][\mathcal{D}\eta^{-}]$, that generates moments in the form of bi-probabilities [15],

$$\begin{split} \Lambda_{t} &= \mathrm{e}^{-\mathrm{i}t\hat{H}_{0}} \Big(\sum_{n=0}^{\infty} (-\mathrm{i})^{n} \int_{0}^{t} \mathrm{d}s_{n} \cdots \int_{0}^{s_{2}} \mathrm{d}s_{1} \sum_{\alpha_{n},\dots,\alpha_{1}} \\ &\times \sum_{\eta_{n}^{\pm}} \sum_{\eta_{0}^{\pm}} Q_{(s_{n},\vec{\tau}_{\alpha_{n}}),\dots,(s_{1},\vec{\tau}_{\alpha_{1}})|(0,\vec{\tau}_{\rho})} (\eta_{n}^{+},\eta_{n}^{-}|\eta_{0}^{+},\eta_{0}^{-}) \sqrt{\rho(\eta_{0}^{+})\rho(\eta_{0}^{-})} \\ &\times \prod_{j=n}^{1} \left(F_{\alpha_{j}}(\eta_{j}^{+})\hat{H}_{\alpha_{j}}(s_{j}) \bullet - \bullet \hat{H}_{\alpha_{j}}(s_{j})F_{\alpha_{j}}(\eta_{j}^{-}) \right) \right) \mathrm{e}^{+\mathrm{i}t\hat{H}_{0}} \\ &= \mathrm{e}^{-\mathrm{i}t\hat{H}_{0}} \Big(\sum_{n=0}^{\infty} (-\mathrm{i})^{n} \int_{0}^{t} \mathrm{d}s_{n} \cdots \int_{0}^{s_{2}} \mathrm{d}s_{1} \sum_{\alpha_{n},\dots,\alpha_{1}} \\ &\times \sum_{\boldsymbol{f}_{n}^{\pm}} Q_{s_{n},\dots,s_{1}|0}^{F_{\alpha_{n}},\dots,F_{\alpha_{1}}} (\boldsymbol{f}_{n}^{+},\boldsymbol{f}_{n}^{-}|\rho) \prod_{j=n}^{1} \left(f_{j}^{+}\hat{H}_{\alpha_{j}}(s_{j}) \bullet - \bullet \hat{H}_{\alpha_{j}}(s_{j})f_{j}^{-} \right) \Big) \mathrm{e}^{+\mathrm{i}t\hat{H}_{0}} \end{split}$$

where $\hat{H}_{\alpha}(s) = e^{is\hat{H}_0}\hat{H}_{\alpha}e^{-is\hat{H}_0}$.

On the other hand, Λ_t is also a completely positive and trace-preserving (CPTP) dynamical map well known from the standard theory of open quantum systems,

$$\Lambda_t \hat{\rho}_O = \operatorname{tr}_S \left[\operatorname{e}^{-\operatorname{i} t \hat{H}_{OS}} \hat{\rho}_O \otimes \hat{\rho} \operatorname{e}^{+\operatorname{i} t \hat{H}_{OS}} \right] \quad : \quad \hat{H}_{OS} = \hat{H}_0 \otimes \hat{1}_S + \hat{1}_O \otimes \hat{H} + \sum_{\alpha=1}^N \hat{H}_\alpha \otimes \hat{F}_\alpha, \qquad (9.9)$$

where \hat{H} is the Hamiltonian of the system S that plays the role of the environment in this scenario. Therefore, the characteristic function of $\mathcal{Q}[\eta^+, \eta^-][\mathcal{D}\eta^+][\mathcal{D}\eta^-]$ is a well-behaved, regular CPTP map, and thus, the underlying bi-trajectory measure should also be well-behaved. This concludes the argument.

The system bi-trajectory measure $\mathcal{Q}[\eta^+, \eta^-][\mathcal{D}\eta^+][\mathcal{D}\eta^-]$ is the true master object of the formalism, as the relations (9.5) and (9.3) show that it is the source of all its elements. The master measure is not associated with any particular observable or initial condition, but rather it is associated with the quantum system itself. Indeed, the master measure is solely determined by the Hilbert space \mathcal{H}_S of the system and its Hamiltonian \hat{H} , and, as previously discussed, the system is also formally identified by the same pair. Thus, in the bi-trajectory formalism, the system is synonymous with $\mathcal{Q}[\eta^+, \eta^-][\mathcal{D}\eta^+][\mathcal{D}\eta^-]$.

10 Summary of the deduced formalism

In the framework of the bi-trajectory formalism, a quantum system is identified with a complexvalued measure on the space of trajectory pairs $\tau \mapsto (\eta^+(\tau), \eta^-(\tau))$ [cf. Eq. (9.6)] where the coordinate $\tau = (\tau^0, \vec{\tau})$ includes the time τ^0 and $(d^2 - 1)$ -component vector $\vec{\tau}$ parameterizing basis transformations in *d*-dimensional Hilbert space \mathcal{H} [cf. Eq. (9.1)]:

Quantum System
$$\longleftrightarrow \mathcal{Q}[\eta^+, \eta^-][\mathcal{D}\eta^+][\mathcal{D}\eta^-].$$
 (10.1)

For closed systems, the master bi-trajectory measure Q is generated by the system's Hamiltonian: given a hermitian operator \hat{H} , all discrete-time restrictions of the measure $Q_{\tau_n|\tau_0}(\eta_n^+, \eta_n^-|\eta_0^+, \eta_0^-)$ [cf. Eq. (9.5)] can be explicitly computed using the formula (9.2).

Using Q, one can obtain quantitative answer to questions about the system's dynamics, measurements outcomes, or multi-time correlations between observables. Depending on the question, this requires either a discrete-time restriction of the measure or computing an appropriate bi-trajectory average.

The probability of an outcome for a sequential measurement performed with an arbitrary collection of devices is given by the following restriction of Q:

$$P_{t_{n}|t_{0}}^{F_{n}}(f_{n}|\rho) = \iint \left(\sum_{\eta^{\pm}} \prod_{j=1}^{n} \left(\delta_{\eta^{+}(\tau_{j}),\eta_{j}^{+}} \delta_{F_{j}(\eta_{j}^{+}),f_{j}}\right) \left(\delta_{\eta^{-}(\tau_{j}),\eta_{j}^{-}} \delta_{F_{j}(\eta_{j}^{-}),f_{j}}\right) \right) \\ \times \sqrt{\rho(\eta^{+}(\tau_{0}))\rho(\eta^{-}(\tau_{0}))} \mathcal{Q}[\eta^{+},\eta^{-}][\mathcal{D}\eta^{+}][\mathcal{D}\eta^{-}] \\ = \sum_{\eta_{n}^{\pm},\eta_{0}} \left(\prod_{j=1}^{n} \delta_{F_{j}(\eta_{j}^{+}),f_{j}^{+}} \delta_{F_{j}(\eta_{j}^{-}),f_{j}^{-}}\right) \mathcal{Q}_{\tau_{n}|\tau_{0}}(\eta_{n}^{+},\eta_{n}^{-}|\eta_{0},\eta_{0})\rho(\eta_{0}) \\ = \operatorname{tr}\left[\left(\prod_{j=n}^{1} \hat{P}_{t_{j}}^{F_{j}}(f_{j})\right) \mathrm{e}^{-\mathrm{i}\hat{H}t_{0}}\hat{\rho} \,\mathrm{e}^{\mathrm{i}\hat{H}t_{0}}\left(\prod_{j=1}^{n} \hat{P}_{t_{j}}^{F_{j}}(f_{j})\right)\right] = \mathcal{Q}_{t_{n}|t_{0}}^{F_{n}}(f_{n},f_{n}|\rho), \qquad (10.2)$$

where $\tau_j^0 = t_j$ for each j, and $\vec{\tau}_j$ parameterize transformations to eigenbasis of the corresponding observables:

$$\hat{F}_j \hat{S}_{\vec{\tau}_j} |\eta\rangle = \Big(\sum_{f \in \Omega(F_j)} f \hat{P}_0^{F_j}(f) \Big) \hat{S}_{\vec{\tau}_j} |\eta\rangle = F_j(\eta) \hat{S}_{\vec{\tau}_j} |\eta\rangle,$$
(10.3)

and the initial condition is encoded in the probability distribution $\rho(\eta)$ and the coordinate $\tau_0 = (t_0, \vec{\tau}_0)$ such that

$$\hat{\rho}\hat{S}_{\vec{\tau}_0}|\eta\rangle = \rho(\eta)\hat{S}_{\vec{\tau}_0}|\eta\rangle. \tag{10.4}$$

Therefore, measurements are described by the "diagonal" part (i.e., $f_n^+ = f_n^-$) of the discrete-time and observable restriction, in which the bi-trajectory $(\eta^+(\tau), \eta^-(\tau))$ is filtered trough the sequence of coordinates τ_n corresponding to the measurement timings and choices of probed observables. Nonetheless, the "off-diagonal" parts of the master measure—where $\eta^+(\tau) \neq \eta^-(\tau)$ —always contribute to the overall result via interference between η^+ and η^- occurring during the period between measurements.

Arbitrary multi-time correlation functions—which might or might not be directly measurable —correspond to moments of the bi-trajectory measure. For example, the two-time correlation function, which is commonly used in standard quantum mechanics, is obtained in the bi-trajectory formalism as a second moment:

$$\operatorname{tr}\left([\hat{F}_{2}(t_{2}),\hat{F}_{1}(t_{1})]\hat{\rho}\right) = \iint \left[F_{2}\left(\eta^{+}(\tau_{2})\right)F_{1}\left(\eta^{+}(\tau_{1})\right) - F_{1}\left(\eta^{-}(\tau_{1})\right)F_{2}\left(\eta^{-}(\tau_{2})\right)\right] \\ \times \sqrt{\rho\left(\eta^{+}(\tau_{0})\right)\rho\left(\eta^{-}(\tau_{0})\right)} \mathcal{Q}[\eta^{+},\eta^{-}][\mathcal{D}\eta^{+}][\mathcal{D}\eta^{-}] \\ = \sum_{f_{2}^{\pm}}\sum_{f_{1}^{\pm}} (f_{2}^{\pm}f_{1}^{+} - f_{1}^{-}f_{2}^{-})Q_{t_{2},t_{1}|0}^{F_{2},F_{1}}\left(f_{2}^{\pm},f_{1}^{+};f_{2}^{-},f_{1}^{-}|\rho\right)$$
(10.5)

We can see in this example that certain types of correlation functions are determined solely by the "off-diagonal" parts of bi-trajectory measure restrictions (notice that the entire expression vanishes when $f_n^+ = f_n^-$).

One of the strengths of the bi-trajectory formalism is its utility in describing multi-time correlation functions for *open* systems. In Section 8, we introduced the bi-trajectory parameterization of dynamical maps, which can essentially be viewed as a combination of an infinite number of multi-time correlation functions of all orders. To understand what makes the bi-trajectory formulation particularly effective in this context, let us examine a relatively simple example of two-time correlation function with the system–environment coupling $\hat{H}_S + \hat{H}_E + \lambda \hat{V}_S \otimes \hat{V}_E$:

$$\operatorname{tr} \left(\left[\hat{U}_{0,t_{2}}^{SE}(\hat{F}_{2} \otimes \hat{1}) \hat{U}_{t_{2},0}^{SE}, \, \hat{U}_{0,t_{1}}^{SE}(\hat{F}_{1} \otimes \hat{1}) \hat{U}_{t_{1},0}^{SE} \right] \, \hat{\rho}_{S} \otimes \hat{\rho}_{E} \right)$$

$$= \iint \operatorname{tr}_{S} \left[\left(\hat{F}_{2}[v^{+}](t_{2}) \hat{F}_{1}[v^{+}](t_{1}) - \hat{F}_{1}[v^{-}](t_{1}) \hat{F}_{2}[v^{-}](t_{2}) \right) \hat{\rho}_{S} \right] \mathcal{Q}^{V_{E}}[v^{+}, v^{-}][\mathcal{D}v^{+}][\mathcal{D}v^{-}]$$

$$= \iint \left[F_{2}(\eta^{+}(\tau_{2})) F_{1}(\eta^{+}(\tau_{1})) - F_{1}(\eta^{-}(\tau_{1})) F_{2}(\eta^{-}(\tau_{2})) \right]$$

$$\times \sqrt{\rho_{S}(\eta^{+}(\tau_{0})) \rho_{S}(\eta^{-}(\tau_{0}))} \, \mathcal{Q}^{S|E}[\eta^{+}, \eta^{-}][\mathcal{D}\eta^{+}][\mathcal{D}\eta^{-}],$$

$$(10.6)$$

where, as per Eq. (8.7),

$$\hat{F}_{j}[h](t) = \sum_{f \in \Omega(F_{j})} f \hat{P}_{t}^{F_{j}}[h](f).$$
(10.7)

Crucially, the functional \mathcal{Q}^{V_E} is a bi-trajectory measure generated by \hat{H}_E , and thus represents the observable V_E in a *free* system E. In other words, \mathcal{Q}^{V_E} describes the dynamics of V_E as if S did not exist.

On the other hand, the measure $\mathcal{Q}^{S|E}$, that can formally be written as

$$\mathcal{Q}^{S|E}[\eta^{+},\eta^{-}][\mathcal{D}\eta^{+}][\mathcal{D}\eta^{-}] = \left(\iint \mathcal{K}^{S}[\eta^{+},\eta^{-}|v^{+},v^{-}]\mathcal{Q}^{V_{E}}[v^{+},v^{-}][\mathcal{D}v^{+}][\mathcal{D}v^{-}]\right)[\mathcal{D}\eta^{+}][\mathcal{D}\eta^{-}],$$
(10.8)

represents the bi-trajectory measure of system S that is coupled to its environment E via observable \hat{V}_E . This measure is conditioned by the free dynamics of system E, and as a result, is no longer generated by the Hamiltonian \hat{H}_S . However, since the influence of the environment is fully encapsulated by the free measure Q^{V_E} , the effects of E on the dynamics of S are determined only by the innate properties of E, even though the coupling is symmetric, with S also influencing E. Likewise, if we were to describe E in terms of its bi-trajectory measure while treating S as the environment, we would obtain $Q^{E|S}$ in a form of kernel averaged with Q^{V_S} , the bi-trajectory measure describing the free dynamics of observable V_S in system S. The fact that the description of the environment in isolation is sufficient to describe the dynamics of the open system, can be seen as a natural consequence of constructing the formal description based on trajectory distributions, whether for bi- or single-trajectory theories.

Finally, we arrive at the formal description of the classical limit and the associated quantumto-classical transition. We now recognize that quantum mechanics can be classified as a bitrajectory theory, meaning that the master object of its formalism is the bi-trajectory measure $\mathcal{Q}[\eta^+, \eta^-][\mathcal{D}\eta^+][\mathcal{D}\eta^-]$. This makes quantum mechanics explicitly non-classical, as the defining feature of any classical theory is its description in terms of a single-trajectory formalism, with the probability distribution $\mathcal{P}[e][\mathcal{D}e]$ as the master object. Nonetheless, it is possible to "force" the quantum bi-trajectory measure to behave as a classical single-trajectory probability distribution.

To achieved this, it is sufficient that a bi-trajectory measure associated with a given system observable, $Q^F[f^+, f^-][\mathcal{D}f^+][\mathcal{D}f^-]$ —that is, the measure extending the observable-restricted system's bi-probabilities

$$Q_{\boldsymbol{t}_{n}|t_{0}}^{F}(\boldsymbol{f}_{n}^{+},\boldsymbol{f}_{n}^{-}|\rho) = \sum_{\boldsymbol{\eta}_{n}^{\pm},\eta_{0}} \left(\prod_{j=1}^{n} \delta_{F(\boldsymbol{\eta}_{j}^{+}),f_{j}^{+}} \delta_{F(\boldsymbol{\eta}_{j}^{-}),f_{j}^{-}} \right) \rho(\eta_{0}) Q_{(\boldsymbol{t}_{n},\vec{\boldsymbol{\tau}}_{n})|(t_{0},\vec{\boldsymbol{\tau}}_{\rho})}(\boldsymbol{\eta}_{n}^{+},\boldsymbol{\eta}_{n}^{-}|\eta_{0},\eta_{0}) \big|_{\vec{\boldsymbol{\tau}}_{n},\dots,\vec{\boldsymbol{\tau}}_{1}=\vec{\boldsymbol{\tau}}_{F}}$$

$$(10.9)$$

—assigns non-zero values only when the components of its bi-trajectory overlap, i.e., $Q^F[f^+, f^-] \propto \delta[f^+ - f^-]$. Since bi-trajectory measures are positive semi-definite, such a functional becomes a proper probability measure on the space of single-trajectories,

$$\mathcal{Q}^{F}[f^{+}, f^{-}][\mathcal{D}f^{+}][\mathcal{D}f^{-}] \approx \delta[f^{+} - f]\delta[f^{-} - f]\mathcal{Q}^{F}[f, f][\mathcal{D}f^{+}][\mathcal{D}f^{-}] \equiv \mathcal{P}^{F}[f][\mathcal{D}f] \ge 0.$$
(10.10)

Identifying the conditions under which the system's dynamics cause this kind of collapse of bitrajectory representing its observable into a single trajectory is key to describing the classical limit within this formalism. A detailed analysis of various quantum models capable of achieving this limit can be found in [13, 15, 16].

In summary, the classical limit of the theory manifests in the following way: when an observable F of a quantum system undergoes the quantum-to-classical transition, it becomes possible to describe the observable as a stochastic process in any context. Firstly, the probabilities of measurement outcomes become consistent, as

$$P_{t_n,\dots,t_j}^F,\dots,t_1(f_n,\dots,f_j,\dots,f_1) - \sum_{f_j \in \Omega(F)} P_{t_n}^F(f_n) = \sum_{f_j^+ \neq f_j^-} Q_{t_n}^F(\dots,f_j^+\dots,\dots,f_j^-\dots) \to 0,$$
(10.11)

meaning that measurements deploying F-devices become indistinguishable from sampling trajectories of a classical observable [14]. Secondly, all "quantum" multi-time correlation functions for example the two-time auto-correlation $\operatorname{tr}([\hat{F}(t), \hat{F}(s)]\hat{\rho})$ —vanish as bi-probabilities are now purely "diagonal". Simultaneously, the remaining "classical-like" correlation functions, such as $\operatorname{tr}(\{\hat{F}(t), \hat{F}(s)\}\hat{\rho})$, coincide with the corresponding correlations of the stochastic process F(t), which can be identified with the single-trajectory the bi-trajectory has collapsed into. As a result, this stochastic process representation (or the surrogate field representation, as it is called in [13, 16]) must also extend to other systems coupled to observable \hat{F} . Indeed, since $f^{\pm}(t) = f(t)$ in this limit, the bi-trajectory measure of any open system O interacting with S via observable \hat{F} can be described as being generated by a stochastic Hamiltonian $\hat{H}_O + \lambda F(t)\hat{V}_O$, were F(t), with distribution $\mathcal{P}^F[f][\mathcal{D}f]$, can be interpreted as a process representing an external (possibly noisy) potential.

11 Conclusions

We have demonstrated how the formalism of quantum theory can be deduced from the phenomenology of experiments involving sequential deployment of measuring devices. By design, the formal description obtained this way is necessarily consistent with empirical observations it was derived from. However, the interpretation of the elements of this formalism does not have to (and usually does not) align with the interpretations of other formulations, including the standard formalism. This situation is analogous to classical mechanics in its various forms. Over the years, the formalism of the classical theory has been restructured in numerous ways: starting from the formulation based on Newton's Laws of Motion, and progressing into more elegant phase-space-based formalisms, exemplified most famously by Hamilton's formulation. Ostensibly, all these formulations describe the same *physical theory* and are in agreement regarding the predictions of empirical observations (within the domain of classical physics). Yet, the formulations are structured very differently, leading to different interpretations and, consequently, different *explanations*. For instance, Hamilton's reformulation was motivated by the quest to eliminate the problematic concept of *force* from the theory [26]. Hence, while Newton's formulation explains all motion through the application and balancing of forces, Hamilton's formalism explains the same motion without reference to such concepts.

The precedence set by classical mechanics shows that a specific formalism of a physical theory need not be its defining feature. In other words, a theory can retain its identity even when its formal description is replaced by a vastly different one. As the example of classical mechanics illustrates, though different formulations are often not equivalent (in terms of structure, interpretation, and explanations), it can be advantageous to switch between them depending on the context. Of course, conflicting interpretations remain an issue; in the case of classical mechanics, such conflicts can be sometimes resolved by, for example, treating forces as a part of an effective, rather than fundamental, description.

We believe that the bi-trajectory formalism stands in relation to the standard formalism of quantum mechanics much as Hamilton's formulation does to Newtonian mechanics, rather than as Einstein's theory of special relativity does to classical mechanics. In other words, we view the bi-trajectory formalism as a reformulation of quantum mechanics, not as the formalism of a new theory that would generalize, or be generalized by, the standard quantum theory. Nevertheless, the bi-trajectory formalism is not equivalent to the standard formalism; the two formulations are structured very differently, and thus, must be interpreted differently as well. Crucially, the bi-trajectory formulation operates with a *single mode of description*—unlike the standard formalism, there is no hard (Heisenberg) cut between measurement and non-measurement contexts. This is evident from the formulas (10.2), (10.5), (10.6), and (9.7) where the average over bi-trajectory measure is used in each context with no exceptions, be it measurement or otherwise. In fact, this feature has been explicitly used in [16] (and implicitly in [14]) to quantify, within the language of bi-trajectory picture, the emergence of objectivity in the quantum–classical transition.

Such a comprehensive restructuring of the formalism offers a relatively rare opportunity to reassess the physical and philosophical meaning of the theory from a fresh perspective. Moreover, it is possible that some fundamental problems with the theory that seem insurmountable within the standard formalism might have a solution—or cease to be problems—within a different formal framework. One such problematic aspect of the standard quantum mechanics could be the *measurement problem*. Most—if not all—conceptual and practical issues surrounding measurement in quantum mechanics remain unresolved because the standard formulation, with its dichotomous structure, makes modeling the process of measurement impossible. Indeed, for a model of measuring device to be successful it needs a formal description that does not break when transitioning between measurement and non-measurement contexts—due to the Heisenberg cut, such models are simply outside the scope of the standard formalism.

The bi-trajectory formalism, however, does away with the dichotomous mode of description, and thus, it has no need for neither the Heisenberg cut that separates the modes, nor the collapse rule that binds them. As such, we believe there is a good chance it can support viable models of measuring devices. These models would likely involve coupled quantum systems, where one represents the measured system and is largely arbitrary, and the other system, representing the device, is taken to the classical limit while still being strongly influenced by the interaction. Fortunately, the formal description of interacting systems and the quantum-to-classical transition are both strong suits of the bi-trajectory formulation. Overall, it is our belief that constructing the model of measuring device in the bi-trajectory picture is merely a *technical* problem, rather than the conceptual challenge it poses in the standard formalism. Given that the formalism has already been used with some success to describe coupled system and classical limit on separate occasions, we are optimistic that the technical challenge of combining the two into a single model is well within the reach.

Acknowledgments

D.L. acknowledges financial support by Friedrich-Alexander-Universität Erlangen-Nürnberg through the funding program "Emerging Talent Initiative" (ETI).

References

- H. Atmanspacher. "Cartesian cut, Heisenberg cut, and the concept of complexity". World Futures 49, 333–355 (1997).
- [2] M. Schlosshauer and K. Camilleri. "What classicality? Decoherence and Bohr's classical concepts". AIP Conference Proceedings 1327, 26–35 (2011).
- [3] L. Letertre. "The operational framework for quantum theories is both epistemologically and ontologically neutral". Studies in History and Philosophy of Science Part A 89, 129–137 (2021).
- [4] E. P. Wigner. "Remarks on the mind-body question". Pages 247–260. Springer Berlin Heidelberg. Berlin, Heidelberg (1995).
- [5] D. Frauchiger and R. Renner. "Quantum theory cannot consistently describe the use of itself". Nature Communications 9, 3711 (2018).
- [6] R. E. Kastner. "Unitary-Only Quantum Theory Cannot Consistently Describe the Use of Itself: On the Frauchiger–Renner Paradox". Foundations of Physics 50, 441–456 (2020).
- [7] M. Żukowski and M. Markiewicz. "Physics and Metaphysics of Wigner's Friends: Even Performed Premeasurements Have No Results". Phys. Rev. Lett. 126, 130402 (2021).
- [8] D. Bohm. "A Suggested Interpretation of the Quantum Theory in Terms of 'Hidden' Variables. I". Phys. Rev. 85, 166–179 (1952).
- [9] D. Bohm. "A Suggested Interpretation of the Quantum Theory in Terms of 'Hidden' Variables. II". Phys. Rev. 85, 180–193 (1952).
- [10] R. B. Griffiths. "Consistent histories and the interpretation of quantum mechanics". Journal of Statistical Physics 36, 219–272 (1984).
- [11] M. Gell-Mann and J. B. Hartle. "Classical equations for quantum systems". Phys. Rev. D 47, 3345–3382 (1993).
- [12] J. A. Barrett and P. Byrne, editors. "The Everett Interpretation of Quantum Mechanics: Collected Works 1955–1980 with Commentary". Princeton University Press. Princeton (2012).
- [13] P. Szańkowski and Ł. Cywiński. "Noise representations of open system dynamics". Scientific Reports 10, 22189 (2020).
- [14] P. Szańkowski. "Measuring trajectories of environmental noise". Phys. Rev. A 104, 022202 (2021).
- [15] P. Szańkowski. "Introduction to the theory of open quantum systems". SciPost Phys. Lect. Notes 68 (2023).
- [16] P. Szańkowski and Ł Cywiński. "Objectivity of classical quantum stochastic processes". Quantum 8, 1390 (2024).
- [17] D. Lonigro, F. Sakuldee, Ł. Cywiński, D. Chruśiński, and P. Szańkowski. "Double or nothing: a Kolmogorov extension theorem for multitime (bi)probabilities in quantum mechanics". Quantum 8, 1447 (2024).
- [18] S. Gudder. "Hilbert space representations of decoherence functionals and quantum measures". Mathematica Slovaca 62, 1209–1230 (2012).
- [19] A. N. Kolmogorov. "Foundations of the theory of probability". Chelsea Publishing Company. New York (1956).
- [20] R. P. Feynman, R. B. Leighton, and M. Sands. "The Feynman Lectures on Physics, Vol. III: The New Millennium Edition: Quantum Mechanics". Chapter 3: Probability Amplitudes, pages 65-76. Basic Books. New York (2011). url: https://www.feynmanlectures.caltech. edu/info.
- [21] R. D. Sorkin. "Quantum mechanics as quantum measure theory". Modern Physics Letters A 09, 3119–3127 (1994).
- [22] R. D. Sorkin. "Quantum dynamics without the wavefunction". Journal of Physics A: Mathematical and Theoretical 40, 3207 (2007).
- [23] N. G. Van Kampen. "Stochastic Processes in Physics and Chemistry". North-Holland Personal Library. Elsevier Science. (2011).

- [24] S. Gudder. "Quantum measure and integration theory". Journal of Mathematical Physics 50 (2009).
- [25] S. Gudder. "Finite quantum measure spaces". The American Mathematical Monthly 117, 512–527 (2010).
- [26] H. Poincaré. "Science and Hypothesis". Chapter VI: The Classical Mechanics, pages 89-139. The Walter Scott Publishing Co., Ltd. New York (1905). url: https://en.wikisource.org/ w/index.php?title=Science_and_Hypothesis/Chapter_6&oldid=6191544.