Surpassing the fundamental limits of distillation with catalysts

Kun Fang^{1, *} and Zi-Wen Liu^{2, †}

¹School of Data Science, The Chinese University of Hong Kong, Shenzhen, Guangdong, 518172, China

² Yau Mathematical Sciences Center, Tsinghua University, Beijing 100084, China

(Dated: October 21, 2024)

Quantum resource distillation is a fundamental task in quantum information science. Minimizing the distillation overhead, i.e., the amount of noisy source states required to produce some desired output state within target error ε , which typically scales as $O(\log^{\gamma}(1/\varepsilon))$, is crucial for the scalability of quantum computation and communication. A prior work [Phys. Rev. Lett. 125, 060405 (2020)] established a universal no-go theorem for resource distillation, indicating that no distillation protocol can achieve $\gamma < 1$ in the one-shot (finite-output) setting. Here, we show that this fundamental limit can be surpassed with the aid of quantum catalysts—an additional resource that facilitates the transformation but remains unchanged before and after the process. Specifically, we show that multishot distillation protocols can be converted into one-shot catalytic protocols, which hold significant practical benefits, while maintaining the distillation overhead. In particular, in the context of magic state distillation, our result indicates that the code-based low-overhead distillation protocols that rely on divergingly large batches can be promoted to the one-shot setting where the batch volume can be arbitrarily small for any accuracy. Combining with very recent results on asymptotically good quantum codes with transversal non-Clifford gates, we demonstrate that magic state distillation with constant overhead can be achieved with controllable output size using catalytic protocols. Furthermore, we demonstrate that catalysis enables a spacetime trade-off between overhead and success probability. Notably, we show that the optimal constant for constant-overhead catalytic magic state distillation can be reduced to 1 at the price of compromising the success probability by a constant factor. Finally, we present an illustrative example that extends the catalysis techniques to the study of dynamic quantum resources. This provides the channel mutual information with a one-shot operational interpretation, thereby addressing the open question posed by Wilming in [Phys. Rev. Lett. 127, 260402 (2021)].

Quantum technologies hold the potential of offering revolutionary advantages over classical methods in crucial technological tasks including computation and communication, driven by the precise manipulation of various kinds of quantum resources such as quantum entanglement [1], quantum coherence [2], and magic states [3]. However, the fragile nature of these quantum resources presents a formidable challenge, as these microscopic systems are highly susceptible to various sources of interference, including environmental noises and control imperfections [4]. These noise effects can significantly undermine the effectiveness and security of quantum computation and communication, thereby limiting the practical applicability of quantum technologies. To address these challenges, quantum resource distillation has emerged as a key strategy [3, 5], aiming to transform multiple noisy copies of a quantum resource into fewer, more purified copies, thereby enhancing their efficacy for subsequent applications. Notable examples include entanglement distillation [5-8] and coherence distillation [9-12]that play key roles in quantum networking and cryptography respectively, and in particular, magic state distillation [3, 13-19], which is of substantial interest as a leading scheme for the implementation of non-Clifford gates, which is considered a major bottleneck for fault-tolerant quantum computing.

Given the fundamental importance of resource distillation procedures to quantum information technology, minimizing the cost of distillation becomes a crucial problem. More specifically, we are interested in minimizing the *dis*tillation overhead, namely the amount of noisy inputs needed to produce certain desired outputs, for which a key figure of merit is the exponent γ in the $O(\log^{\gamma}(1/\varepsilon))$ scaling with respect to target error ε . A pressing problem in quantum information originated from the study of magic state distillation is whether the exponent γ can be reduced to below one (meaning that distillation can be performed using sublogarithmic resources) [13]. Recent studies [20, 21] addressed this problem, establishing a universal $\gamma > 1$ barrier, i.e., a logarithmic lower bound on the overhead in the one-shot setting that concerns a single output state, for virtually any type of resource. When considering the multi-shot setting, although surprising, it is recently found to be possible to achieve sublogarithmic and even constant average overhead per output in magic state distillation [14, 16–19, 22], but these protocols exhibit a common "batch size problem" that they need to involve a large number of states (e.g., the Hastings–Haah sublogarithmic protocol [14] is based on a code that entails $\geq 2^{58}$ input and $\geq 2^{44}$ output states) and the volume diverges rapidly as we take $\varepsilon \to 0$, which may not be suitable for practical needs.

In this work, we introduce a method to overcome both limitations by utilizing the idea of quantum catalysis—a phenomenon analogous to chemical catalysis where catalysts can facilitate reactions by providing alternative pathways with lower activation energies. In the realm of quantum information, the catalysis phenomenon involves an auxiliary quantum system, namely the catalyst, which enable transformations that would otherwise be unachievable but remains unaltered before and after the process and thus can be reused. Quantum catalysis was first discussed in the context of entanglement transformation [23] more than two decades ago and has since been extensively studied across various resource theories, including quantum coherence [24, 25], thermodynamics [26–28], and purity [29]. This area continues to be active, as evidenced by recent works [27, 30–33], with comprehensive reviews available in [34, 35]. While catalytic methods are known to be powerful in various scenarios, recent studies have also revealed "negative" results, such as demonstrating that catalysis *cannot* overcome bound entanglement [36] and *cannot* increase distillable entanglement [37]. These limitations highlight that whether catalysis can be helpful for a certain task is an nontrivial question. Furthermore, existing literature on distillation mostly focuses on the distillation rate, namely how much pure target resource can be extracted from a fixed amount of noisy input resource. This is conceptually dual to the overhead considered in this work, where the aim is to minimize the amount of input for some given target.

The key contribution of this work is to demonstrate the effectiveness of catalytic methods in enhancing distillation overhead, particularly their capability of breaking the logarithmic barrier in one-shot magic state distillation. Specifically, we establish a general and rigorous ordering of resource overheads for one-shot (unassisted) distillation, multi-shot distillation, and one-shot catalytic distillation, as illustrated in Fig. 1, by showing how any multi-shot distillation protocol can be converted into a one-shot catalytic protocol while maintaining the overhead by designing suitable catalysts with reusability guarantees. In particular, for magic state distillation,



FIG. 1. The relation of distillation overhead, $\log^{\gamma}(1/\varepsilon)$, across different settings. A fundamental no-go limit of $\gamma \geq 1$ exists for the one-shot (unassisted) distillation. We show in this work that this limit can be overcome with the aid of quantum catalysts.

the application of our catalytic approach to low-overhead protocols, such as the first sublogarithmic protocol by Hastings and Haah [14] and the recent constant-overhead protocols [16–19, 22] can yield protocols with arbitrary batch size (e.g., an n-to-m protocol can be converted into a $\lceil n/m \rceil$ -to-1 protocol) but no greater overhead for any accuracy, overcoming the aforementioned problems. Furthermore, we demonstrate that the failure probability and overhead, corresponding to time and space costs respectively, can be traded for each other with the help of catalysts, which is expected to have wide practical applications. Notably, we prove that the optimal constant for constant-overhead magic state distillation can be reduced to 1 at the price of sacrificing the success probability by a constant factor, pushing the efficiency of distillation to its ultimate limit. Lastly, by extending the catalytic technique to the channel (dynamical) setting, we provide a one-shot operational interpretation of channel mutual information, addressing an open question posed by Wilming in [28].

Preliminaries.— To present our results in a rigorous and unified manner, we shall adopt the language of resource theory, a framework that finds great success in studying different quantum resource features in recent years (see e.g. [38] for an introduction). A standard resource theory is defined by a set of *free operations* \mathcal{O} and a set of *free states* $\mathscr{F} \subseteq \mathscr{D}$ (the complement of which are resource states) where \mathscr{D} denotes the set of all density matrices. Taking the magic theory as an example, \mathscr{F} consists of stabilizer states, and Clifford operations are a standard choice of \mathcal{O} . Generally speaking, \mathcal{F} and \mathcal{O} can be adaptively defined, leading to a wide variety of meaningful resource theories, as long as they follow a *golden* rule: any free operation can only map a free state to another free state, i.e. $\mathcal{E}(\rho) \in \mathscr{F}, \forall \rho \in \mathscr{F}, \forall \mathcal{E} \in \mathscr{O}$. The general goal of distillation tasks is to transform noisy resources into pure ones by some protocol represented by a free operation. We are often also interested in protocols that only produce desired outputs with a certain probability (e.g. protocols based on error correction, which work upon passing the syndrome measurements). To encompass such probabilistic cases, consider the generalization of \mathscr{O} to the class $\mathscr{O}_{sub} := \{\mathcal{L} \mid \forall \rho \in \mathscr{F}, \exists p \geq 0, \psi \in \mathcal{F}\}$ \mathscr{F} , s.t. $\mathcal{L}(\rho) = p \cdot \psi$, which consists of subnormalized quantum operations, i.e., completely positive and tracenonincreasing maps. The case where \mathcal{L} is a completely positive trace-preserving (CPTP) map and thus p = 1corresponds to a deterministic protocol.

Denote the space for the source and target states as S, and the ancillary system that supports the catalyst as A. For any two quantum states $\rho, \sigma \in \mathscr{D}(S)$, define their trace distance by $\Delta(\rho, \sigma) := \|\rho - \sigma\|_1/2$ with $\|\cdot\|_1$ being the trace norm. Denote $\rho^{\otimes n} \xrightarrow{(\varepsilon,p)} \sigma^{\otimes m}$ if there exists a resource distillation protocol $\mathcal{L} \in \mathscr{O}_{\text{sub}}$ such that $\mathcal{L}(\rho^{\otimes n}) = p \cdot \eta^m$ with $\eta^m \in \mathscr{D}(S^m)$ and $\Delta(\eta^m_i, \sigma) \leq \varepsilon$ for all $i \in \{1, \cdots, m\}$, where η^m_i is the *i*-th marginal of η^m [39]. Then the one-shot (unassisted) distillation overhead $C_{\varepsilon,p}$ and the multi-shot average distillation overhead $\overline{C}_{\varepsilon,p}$ can be defined by $C_{\varepsilon,p}(\rho,\sigma) := \min\{n : \rho^{\otimes n} \xrightarrow{(\varepsilon,p)} \sigma\}$ and $\overline{C}_{\varepsilon,p}(\rho,\sigma) := \min\{\lceil n/m \rceil : \rho^{\otimes n} \xrightarrow{(\varepsilon,p)} \sigma^{\otimes m}\}$, respectively. Moreover, denote $\rho^{\otimes n} \otimes \omega \xrightarrow{(\varepsilon,p)} \sigma \otimes \omega$ if there exists a distillation protocol $\mathcal{L} \in \mathscr{O}_{\text{sub}}$ and a catalyst $\omega \in \mathscr{D}(A)$ such that $\mathcal{L}(\rho^{\otimes n} \otimes \omega) = p \cdot \nu$ with $\nu \in \mathscr{D}(SA), \ \Delta(\nu_S, \sigma_S) \leq \varepsilon$ and $\nu_A = \omega_A$, where ν_A, ν_S are two marginals of ν . Then the one-shot catalytic distillation overhead $\widetilde{C}_{\varepsilon,p}$ is defined by $\widetilde{C}_{\varepsilon,p}(\rho,\sigma) := \min\{n : \rho^{\otimes n} \otimes \omega \xrightarrow{(\varepsilon,p)} \sigma \otimes \omega, \omega \in \mathscr{D}\}$. Using these notations, the no-go theorem of [20] sets the fundamental limit $C_{\varepsilon,p}(\rho,\sigma) = \Omega(\log(1/\varepsilon)).$

General theory: one-shot catalytic distillation and theoretical guarantee for catalyst reuse. — Now we introduce the main results, with detailed proofs and extended discussions left to the Supplemental Materials. Here we are interested in the case where the target state is pure and $n \ge m$. Axiomatically, it is assumed that appending free systems, discarding and permuting subsystems, and classically controlled free operations are free, and no other properties of the resource theory are needed, rendering the framework highly flexible and applicable to most relevant theories.

Since a one-shot distillation protocol can be regarded as a multi-shot distillation protocol with further conditions, it follows directly that $\overline{C}_{\varepsilon,p}(\rho,\sigma) \leq C_{\varepsilon,p}(\rho,\sigma)$ for any quantum states ρ and σ . For the one-shot catalytic distillation, we adapt a catalysis technique originally proposed in [40] for LOCC transformations of quantum entanglement (further explored in recent works such as [25, 27, 28, 30-33]) to our distillation task, which involves an unequal number of source and target states. This is achieved by regrouping the source states into blocks and embedding the target state into a larger space. Particularly, we show that any multi-shot distillation protocol from $\rho^{\otimes n}$ to $\sigma^{\otimes m}$ can effectively be turned into a one-shot catalytic distillation protocol from $\rho^{\lceil n/m\rceil}\otimes\omega$ to $\sigma \otimes \omega$ with the same performance. Here, the catalyst ω plays a crucial role by enabling new transformation pathways that are otherwise unattainable and absorbing part of the complexity of the transformation process. Although the catalyst temporarily participates in the transformation, it is fully restored afterward. See Supplemental Materials for detailed descriptions of the procedure. As a result, we have the inequality $C_{\varepsilon,p}(\rho,\sigma) \leq$ $\overline{C}_{\varepsilon,p}(\rho,\sigma)$. In conclusion, the general relation of distillation overhead across different settings is as follows:

Theorem 1 For any quantum states ρ and σ , target error $\varepsilon \in [0, 1]$ and success probability $p \in [0, 1]$, the following relation holds: $\widetilde{C}_{\varepsilon,p}(\rho, \sigma) \leq \overline{C}_{\varepsilon,p}(\rho, \sigma) \leq C_{\varepsilon,p}(\rho, \sigma)$.

A key advantage of using a catalyst is its recoverability after the transformation, allowing for repeated reuse. We provide a theoretical guarantee that the catalyst can be reused indefinitely without any degradation in its expected performance.

Theorem 2 For a deterministic one-shot catalytic resource distillation procedure, after $l \geq 1$ repeated uses of the catalyst ω_A , we obtain a joint state $\nu_{S_1S_2...S_lA}$ such that the catalyst is exactly returned on its marginal $\nu_A = \omega_A$ and the target states $\nu_{S_1} = \nu_{S_2} = \cdots = \nu_{S_l}$ with error $\Delta(\nu_{S_i}, \sigma_S) \leq \varepsilon$ for all $i \in \{1, \dots, l\}$.

Note that the catalyst may exhibit correlations with the remaining systems. However, when addressing distillation tasks aimed at pure target states, which is the case of primary interest, the correlation between the obtained target state and the remaining systems is under control. This can be made explicit by applying Lemma 10 from [37], which states that an error threshold on the target state $\Delta(\nu_S, |\phi\rangle \langle \phi|_S) \leq \varepsilon$ ensures that the correlation between the target state and the catalyst remains small, with $\Delta(\nu_{SA}, |\phi\rangle \langle \phi|_S \otimes \nu_A) \leq \varepsilon + 3\sqrt{\varepsilon}$. Consequently, any intermediate usage of the target state and the catalyst will have a negligible impact on the other systems.

Application in magic state distillation. — As a major bottleneck of fault-tolerant quantum computing, magic state distillation has been a subject of extensive study since Bravyi and Kitaev's proposal in 2005 [3]. Significant effort has been devoted to reducing the overhead of magic state distillation, with numerous improvements achieved over the years [3, 13, 14, 16–19, 22, 41–44].

It was long believed that the overhead exponent γ cannot be reduced to below one. In the one-shot setting, this can indeed be shown by applying the universal no-go theorem for resource distillation in [20]. However, in the multi-shot setting, Hastings and Haah first found a code that achieves an average overhead of $n/m = O(\log^{\gamma}(1/\varepsilon))$ for $\gamma \approx 0.678$ [14], and very recent progress even further optimizes the exponent to $\gamma = 0$, namely achieving constant average overhead [16– 18]). As mentioned, despite these advances, the low overhead in these protocols comes with the batch size problem: the codes inherently involve a very large number of states, and furthermore, to achieve lower ε we need rapidly growing batch sizes. By applying the catalytic conversion established in Theorem 1, we can turn the constant-overhead distillation protocols [16, 22] into oneshot counterparts that achieves arbitrary accuracy with an arbitrary amount of (even only one) target states. This provides a way to bypass the fundamental limit of $\Omega(\log(1/\varepsilon))$ for one-shot distillation [20] and makes lowoverhead distillation more aligned with practical implementation.

Corollary 3 There exist one-shot catalytic magic state distillation protocols that achieve any given target error with constant success probability and constant overhead.

As discussed earlier, the correlation between the catalyst and the obtained target state can be made sufficiently small if the ideal target state is a pure state, as is the case in magic state distillation, where the target states are typically $|T\rangle$ and $|CCZ\rangle$. A related result by Rubboli and Tomamichel [45] indicates that making such residual correlations arbitrarily small requires a divergent amount of resources in the catalyst, given that the resource theory having multiplicative maximum fidelity of resource (i.e., $F(\rho_1 \otimes \rho_2) = F(\rho_1) \cdot F(\rho_2)$, where $F(\rho) := \max_{\sigma \in \mathscr{F}} F(\rho, \sigma)$, with F on the right-hand side referring to quantum fidelity). However, the maximum fidelity in the resource theory of magic (also named stabilizer fidelity) is known to be non-multiplicative [46, Section 6.2], hence the result in [45] does not apply, leaving open the possibility of discovering better catalysts with vanishing residual correlations.

Trading success probability for reduced overhead.— The resource overhead essentially represents the space required for distillation, characterizing the size of quantum systems that need to be coherently manipulated in a single experiment. On the other hand, the success probability corresponds to time cost, reflecting the number of experimental repetitions needed to achieve a sufficient number of successful outcomes. In the context of nearterm quantum computers where the available number of qubits is limited, it is often more feasible to repeat an experiment multiple times rather than to conduct it with a larger number of qubits in fewer trials. By leveraging quantum catalysis, we enable a trade-off between space and time, allowing the resource burden to be shifted from qubit overhead to time.

More specifically, we show that if a transformation from $\rho^{\otimes n}$ to $\sigma^{\otimes m}$ is achievable with success probability p, then a catalytic transformation from $\rho^{\otimes k}$ to σ can also be achieved with success probability $pm/\lceil n/k \rceil$, for any $1 \leq k \leq n/m$, effectively trading time for space. Notably, by setting k = 1, we obtain a one-shot catalytic distillation protocol with unit overhead and success probability pm/n, which represents the extreme scenario where the overhead is minimized to its absolute limit.

Theorem 4 Suppose there exists a distillation protocol transforming $\rho^{\otimes n}$ to $\sigma^{\otimes m}$ with success probability p and target error ε . Then it holds for any $1 \leq k \leq n/m$ that $\widetilde{C}_{\varepsilon,pm\lceil n/k\rceil^{-1}}(\rho,\sigma) \leq k$. In particular, $\widetilde{C}_{\varepsilon,pm/n}(\rho,\sigma) = 1$.

This result is nontrivial because trading success probability for overhead is not always possible without the use of a catalyst. Consider the example of entanglement distillation, where the goal is to transform a noisy entangled state ρ (e.g., an isotropic state) into a Bell state ψ with success probability p' < 1 and target error ε' . According to the no-go theorems established in [20, 47], there exist choices of (p', ε') within the forbidden regime, meaning the task cannot be accomplished with the desired performance. However, in the average case, this transformation is possible using the hashing protocol [48]: for some r and m, $\rho^{\otimes mr}$ can be transformed into $\psi^{\otimes m}$ with a target error and success probability of 1, where 1/r corresponds to the hashing bound. This demonstrates that achieving a multi-copy transformation does not necessarily guarantee the feasibility of a one-shot transformation by simply compromising the success probability, even when p' is chosen to be close to 0. Yet, our result indicates that this trade-off can always be achieved with the aid of a catalyst.

The proof of this result extends beyond existing catalysis techniques in the literature. To establish the result, we first consider $g = \lceil n/k \rceil$ groups of $\zeta = \rho^{\otimes k} \in \mathscr{D}(S^k)$ and aim to transform $\zeta^{\otimes g}$ into $\sigma^{\otimes m}$. This transformation requires embedding the state twice. In the first embedding, we append a free state $\pi^{\otimes (k-1)}$ such that the embedded state $\hat{\sigma}$ resides in the same space as $\zeta \in \mathscr{D}(S^k)$. For the second embedding, we append $\hat{\sigma}^{\otimes m}$ using a free state $\theta^{\otimes (g-m)}$. These embeddings allow us to apply the proof techniques from Theorem 1. It is crucial, however, to append the free state θ in the orthogonal subspace of S^k . This can be done by extending to a larger space, such as $S^k W$, where W is orthogonal to S^k , to prevent the free state from mixing with the target state. This ensures that the target state can be post-selected through projective measurement, preserving the fidelity in the final step. Furthermore, the catalyst used in the proof is not unique and can be made independent of the target state when m = 1. A complete proof is provided in the Supplemental Materials.

As mentioned, very recent works [16-18] demonstrated that magic state distillation can be achieved with constant overhead in the asymptotic limit but left the question of optimality of this constant open for future investigation. By applying Theorem 4, we find that the optimal constant can be ultimately reduced to 1 with the aid of a catalyst, at the expense of compromising the success probability by a constant factor.

Corollary 5 There exist one-shot catalytic magic state distillation protocols that achieve any given target error with constant success probability using only one copy of the source magic state.

Channel theory. — The concept of catalytic transformation can also be extended to the manipulation of quantum channels, which correspond to dynamic resources that play a crucial role in e.g. quantum communication and quantum error correction [21, 49–56]. By introducing the reduced channel of a multipartite quantum channel and replacing the trace norm with the diamond norm to characterize transformation errors, we can employ catalysis techniques to convert multi-shot channel coding strategies into one-shot catalytic channel coding strategies. As an illustrative example, consider quantum channel coding under entanglement-assisted or non-signalling codes. Leveraging the renowned quantum reverse Shannon theorem [49], we provide a one-shot operational interpretation of channel mutual information, thereby addressing the open question raised in [28]. See Supplemental Materials for detailed statements and proofs. More comprehensive studies of the channel setting will be left for follow-up works.

Discussion.— This work presents a theory of distillation overhead with the assistance of quantum catalvsis that is universally applicable to different quantum resources, underscoring the extraordinary potential of quantum catalysis in enhancing the performance of quantum information processing tasks. Most notably, we showed that it can indeed help overcome various kinds of fundamental limitations of distillation methods. In the particularly important context of magic state distillation, our approach can be combined with low-overhead protocols in the literature to obtain one-shot protocols with no greater overhead but fully controllable batch sizes for arbitrary accuracy. Furthermore, we demonstrated that catalysis enables a trade-off between overhead and success probability, especially allowing the optimal constant for magic state distillation overhead to be reduced to 1 by lowering the success probability by a constant factor, which optimizes the cost of the constant-overhead magic state distillation constructions.

We should point out that the catalysts used in this work generally have a size comparable to the system in the multi-shot protocols so that coherent manipulation of large quantum systems is still needed. However, the possibility for smaller catalysts is open, which is worth further exploration. More generally speaking, the current results highlight potential advantages in a theoretical setting, but we anticipate more detailed study of various issues that may arise in actual implementations, such as noise and correlation effects on the catalysts, to be an important step forward. Moreover, given the close connection between magic state distillation and quantum errorcorrecting (QEC) codes, our catalysis results may also provide new insights into catalytic QEC codes [57, 58]. Finally, extending these results to continuous variable systems and conducting a systematic study of channel theory remain promising directions for future work.

Acknowledgments. K.F. is supported by the University Development Fund (Grant No. UDF01003565) from the Chinese University of Hong Kong, Shenzhen. Z.-W.L. is supported in part by a startup funding from YMSC, Tsinghua University, and NSFC under Grant No. 12475023.

* kunfang@cuhk.edu.cn

Mod. Phys. 81, 865–942 (2009).

- [2] Alexander Streltsov, Gerardo Adesso, and Martin B. Plenio, "Colloquium: Quantum coherence as a resource," Rev. Mod. Phys. 89, 041003 (2017).
- [3] Sergey Bravyi and Alexei Kitaev, "Universal quantum computation with ideal clifford gates and noisy ancillas," Phys. Rev. A 71, 022316 (2005).
- [4] John Preskill, "Quantum Computing in the NISQ era and beyond," Quantum 2, 79 (2018).
- [5] Charles H Bennett, Gilles Brassard, Sandu Popescu, Benjamin Schumacher, John A Smolin, and William K Wootters, "Purification of noisy entanglement and faithful teleportation via noisy channels," Phys. Rev. Lett. 76, 722 (1996).
- [6] Charles H. Bennett, Herbert J. Bernstein, Sandu Popescu, and Benjamin Schumacher, "Concentrating partial entanglement by local operations," Phys. Rev. A 53, 2046–2052 (1996).
- [7] Charles H. Bennett, David P. DiVincenzo, John A. Smolin, and William K. Wootters, "Mixed-state entanglement and quantum error correction," Phys. Rev. A 54, 3824–3851 (1996).
- [8] K. Fang, X. Wang, M. Tomamichel, and R. Duan, "Nonasymptotic entanglement distillation," IEEE Trans. Inf. Theory, 1–1 (2019).
- [9] Andreas Winter and Dong Yang, "Operational resource theory of coherence," Phys. Rev. Lett. 116, 120404 (2016).
- [10] Kun Fang, Xin Wang, Ludovico Lami, Bartosz Regula, and Gerardo Adesso, "Probabilistic Distillation of Quantum Coherence," Phys. Rev. Lett. **121**, 070404 (2018), 1804.09500.
- [11] Bartosz Regula, Kun Fang, Xin Wang, and Gerardo Adesso, "One-shot coherence distillation," Phys. Rev. Lett. **121**, 010401 (2018).
- [12] Masahito Hayashi, Kun Fang, and Kun Wang, "Finite block length analysis on quantum coherence distillation and incoherent randomness extraction," IEEE Trans. Inf. Theory 67, 3926–3944 (2021).
- [13] Sergey Bravyi and Jeongwan Haah, "Magic-state distillation with low overhead," Phys. Rev. A 86, 052329 (2012).
- [14] Matthew B. Hastings and Jeongwan Haah, "Distillation with sublogarithmic overhead," Phys. Rev. Lett. 120, 050504 (2018).
- [15] Zi-Wen Liu, Kaifeng Bu, and Ryuji Takagi, "One-shot operational quantum resource theory," Phys. Rev. Lett. 123, 020401 (2019).
- [16] Adam Wills, Min-Hsiu Hsieh, and Hayata Yamasaki, "Constant-overhead magic state distillation," arXiv: 2408.07764 (2024).
- [17] Louis Golowich and Venkatesan Guruswami, "Asymptotically good quantum codes with transversal non-clifford gates," arXiv preprint arXiv:2408.09254 (2024).
- [18] Quynh T Nguyen, "Good binary quantum codes with transversal ccz gate," arXiv preprint arXiv:2408.10140 (2024).
- [19] Seok-Hyung Lee, Felix Thomsen, Nicholas Fazio, Benjamin J Brown, and Stephen D Bartlett, "Low-overhead magic state distillation with color codes," arXiv preprint arXiv:2409.07707 (2024).
- [20] Kun Fang and Zi-Wen Liu, "No-go theorems for quantum resource purification," Phys. Rev. Lett. 125, 060405 (2020).
- [21] Kun Fang and Zi-Wen Liu, "No-go theorems for quantum

[†] zwliu0@tsinghua.edu.cn

^[1] Ryszard Horodecki, Paweł Horodecki, Michał Horodecki, and Karol Horodecki, "Quantum entanglement," Rev.

resource purification: New approach and channel theory," PRX Quantum **3**, 010337 (2022).

- [22] Anirudh Krishna and Jean-Pierre Tillich, "Towards low overhead magic state distillation," Phys. Rev. Lett. 123, 070507 (2019).
- [23] Daniel Jonathan and Martin B Plenio, "Entanglementassisted local manipulation of pure quantum states," Phys. Rev. Lett. 83, 3566 (1999).
- [24] Johan Aberg, "Catalytic coherence," Phys. Rev. Lett. 113, 150402 (2014).
- [25] Ryuji Takagi and Naoto Shiraishi, "Correlation in catalysts enables arbitrary manipulation of quantum coherence," Phys. Rev. Lett. 128, 240501 (2022).
- [26] Markus P Müller, "Correlating thermal machines and the second law at the nanoscale," Phys. Rev. X 8, 041051 (2018).
- [27] Naoto Shiraishi and Takahiro Sagawa, "Quantum thermodynamics of correlated-catalytic state conversion at small scale," Phys. Rev. Lett. **126**, 150502 (2021).
- [28] H. Wilming, "Entropy and reversible catalysis," Phys. Rev. Lett. 127, 260402 (2021).
- [29] Paul Boes, Jens Eisert, Rodrigo Gallego, Markus P Müller, and Henrik Wilming, "Von neumann entropy from unitarity," Phys. Rev. Lett. **122**, 210402 (2019).
- [30] Priyabrata Char, Dipayan Chakraborty, Amit Bhar, Indrani Chattopadhyay, and Debasis Sarkar, "Catalytic transformations in coherence theory," Phys. Rev. A 107, 012404 (2023).
- [31] Chandan Datta, Ray Ganardi, Tulja Varun Kondra, and Alexander Streltsov, "Is there a finite complete set of monotones in any quantum resource theory?" Phys. Rev. Lett. 130, 240204 (2023).
- [32] Tulja Varun Kondra, Chandan Datta, and Alexander Streltsov, "Catalytic transformations of pure entangled states," Phys. Rev. Lett. 127, 150503 (2021).
- [33] Patryk Lipka-Bartosik and Paul Skrzypczyk, "Catalytic quantum teleportation," Phys. Rev. Lett. 127, 080502 (2021).
- [34] Chandan Datta, Tulja Varun Kondra, Marek Miller, and Alexander Streltsov, "Catalysis of entanglement and other quantum resources," Rep. Prog. Phys (2023).
- [35] Patryk Lipka-Bartosik, Henrik Wilming, and Nelly HY Ng, "Catalysis in quantum information theory," Rev. Mod. Phys. 96, 025005 (2024).
- [36] Ludovico Lami, Bartosz Regula, and Alexander Streltsov, "No-go theorem for entanglement distillation using catalysis," Phys. Rev. A 109 (2024).
- [37] Ray Ganardi, Tulja Varun Kondra, and Alexander Streltsov, "Catalytic and asymptotic equivalence for quantum entanglement," arXiv preprint arXiv:2305.03488 (2023).
- [38] Eric Chitambar and Gilad Gour, "Quantum resource theories," Rev. Mod. Phys. 91, 025001 (2019).
- [39] Here we follow the convention of magic state distillation that focuses on the error of each marginal states. Since the pure target is usually the scenario of interest, this is equivalent to define the error over the global states.
- [40] Runyao Duan, Yuan Feng, Xin Li, and Mingsheng Ying, "Multiple-copy entanglement transformation and entanglement catalysis," Phys. Rev. A 71, 042319 (2005).
- [41] Jeongwan Haah and Matthew B. Hastings, "Codes and Protocols for Distilling *T*, controlled-*S*, and Toffoli Gates," Quantum 2, 71 (2018).
- [42] Adam M Meier, Bryan Eastin, and Emanuel Knill,

"Magic-state distillation with the four-qubit code," Quantum Inf. Comput. **13**, 195–209 (2013).

- [43] Earl T Campbell, Hussain Anwar, and Dan E Browne, "Magic-state distillation in all prime dimensions using quantum reed-muller codes," Phys. Rev. X 2, 041021 (2012).
- [44] Cody Jones, "Multilevel distillation of magic states for quantum computing," Phys. Rev. A 87, 042305 (2013).
- [45] Roberto Rubboli and Marco Tomamichel, "Fundamental limits on correlated catalytic state transformations," Phys. Rev. Lett. **129**, 120506 (2022).
- [46] Sergey Bravyi, Dan Browne, Padraic Calpin, Earl Campbell, David Gosset, and Mark Howard, "Simulation of quantum circuits by low-rank stabilizer decompositions," Quantum 3, 181 (2019).
- [47] Bartosz Regula, "Probabilistic transformations of quantum resources," Phys. Rev. Lett. 128, 110505 (2022).
- [48] Igor Devetak and Andreas Winter, "Distillation of secret key and entanglement from quantum states," Proc. R. Soc. A. 461, 207–235 (2005).
- [49] Charles H. Bennett, Igor Devetak, Aram W. Harrow, Peter W. Shor, and Andreas Winter, "The quantum reverse shannon theorem and resource tradeoffs for simulating quantum channels," IEEE Trans. Inf. Theory 60, 2926–2959 (2014).
- [50] Gilad Gour, "Comparison of quantum channels by superchannels," IEEE Trans. Inf. Theory 65, 5880–5904 (2019).
- [51] Zi-Wen Liu and Andreas Winter, "Resource theories of quantum channels and the universal role of resource erasure," arXiv e-prints, arXiv:1904.04201 (2019), arXiv:1904.04201 [quant-ph].
- [52] Yunchao Liu and Xiao Yuan, "Operational resource theory of quantum channels," Phys. Rev. Res. 2, 012035 (2020).
- [53] Kun Fang, Xin Wang, Marco Tomamichel, and Mario Berta, "Quantum channel simulation and the channel's smooth max-information," in 2018 IEEE International Symposium on Information Theory (ISIT) (IEEE, 2018) pp. 2326–2330.
- [54] Kun Fang, Xin Wang, Marco Tomamichel, and Mario Berta, "Quantum channel simulation and the channel's smooth max-information," IEEE Trans. Inf. Theory 66, 2129–2140 (2019).
- [55] Gilad Gour and Carlo Maria Scandolo, "Dynamical resources," (2020), arXiv:2101.01552 [quant-ph].
- [56] Bartosz Regula and Ryuji Takagi, "Fundamental limitations on distillation of quantum channel resources," Nature Communications 12 (2021), 10.1038/s41467-021-24699-0.
- [57] Todd Brun, Igor Devetak, and Min-Hsiu Hsieh, "Correcting quantum errors with entanglement," Science 314, 436–439 (2006).
- [58] Todd A Brun, Igor Devetak, and Min-Hsiu Hsieh, "Catalytic quantum error correction," IEEE Trans. Inf. Theory 60, 3073–3089 (2014).
- [59] Mario Berta, Francesco Borderi, Omar Fawzi, and Volkher B Scholz, "Semidefinite programming hierarchies for constrained bilinear optimization," Mathematical Programming, 1–49 (2022).
- [60] A Yu Kitaev, "Quantum computations: algorithms and error correction," Russian Mathematical Surveys 52, 1191 (1997).

Supplemental Materials

In the Supplemental Materials, we provide more detailed expositions, proofs and discussions of the results in the main text. We may reiterate some of the steps to ensure that the Supplemental Materials are explicit and self-contained.

ONE-SHOT CATALYTIC RESOURCE DISTILLATION

Theorem S1 For any quantum states ρ and σ , target error $\varepsilon \in [0,1]$ and success probability $p \in [0,1]$, the following relation holds:

$$\widetilde{C}_{\varepsilon,p}(\rho,\sigma) \leq \overline{C}_{\varepsilon,p}(\rho,\sigma) \leq C_{\varepsilon,p}(\rho,\sigma).$$

Proof. The second inequality follows from their definitions. We now prove the first inequality by constructing an explicit catalytic transformation with the desired performance. The catalyst structure we use here was originally proposed in [40] in the context of LOCC transformations of quantum entanglement and it has also been investigated in various contexts in recent works [25, 27, 28, 30–33]. However, existing literature mostly considers the transformation of source states to an equal number of target states. To adapt to the distillation setting, we consider regrouping the source states into blocks. Consider any multi-shot distillation protocol $\mathcal{L} \in \mathscr{O}_{sub}$ such that $\mathcal{L}(\rho^{\otimes n}) = p\eta^m$ and $\Delta(\eta_i^m, \sigma) \leq \varepsilon$ for all $i \in \{1, \dots, m\}$, where $\eta^m \in \mathscr{D}(S^m)$ and η_i^m is the *i*-th marginal of η^m . Let $k = \lceil n/m \rceil$ be the number of copies in each block and denote $\zeta = \rho^{\otimes k}$. Since $mk = m\lceil n/m \rceil \geq n$, we have a free operation \mathcal{L}_1 such that

$$\mathcal{L}_1(\zeta^{\otimes m}) = \mathcal{L}_1(\rho^{\otimes mk}) = p\eta^m \text{ and } \Delta(\eta^m_i, \sigma) \le \varepsilon,$$

for all $i \in \{1, \dots, m\}$. This can be done by simply throwing away the residual mk - n copies of the source states and then performing the transformation \mathcal{L} on $\rho^{\otimes n}$. Note that η_m and $\sigma^{\otimes m}$ live in $\mathscr{D}(S^m)$, but we can embed them into a larger space $\mathscr{D}((S^k)^m)$ so that the individual subsystem matches the space of ζ . More precisely, consider embedding $\mathcal{E}(\cdot) = (\cdot) \otimes \pi_S^{\otimes (k-1)}$ where $\pi \in \mathscr{F}$ is a free state, and let $\hat{\eta}^m = \mathcal{E}^{\otimes m}(\eta^m)$ and $\hat{\sigma} = \mathcal{E}(\sigma)$. Consider $\mathcal{L}_2 = \mathcal{E} \circ \mathcal{L}_1$ we get a transformation

$$\mathcal{L}_2(\zeta^{\otimes m}) = p\hat{\eta}^m \quad \text{and} \quad \Delta(\hat{\eta}^m_i, \hat{\sigma}) \le \varepsilon_i$$

where $\hat{\eta}_i^m$ is the marginal state on the *i*-th system and the inequality follows since trace distance is invariant under our embedding.

Now, we construct a catalyst ω to transform $\zeta \otimes \omega$ to $\hat{\sigma} \otimes \omega$. Let $\hat{\eta}_{1:i}^m$ be the marginal state of $\hat{\eta}_m$ on the first *i* systems. Consider the catalyst

$$\omega := \frac{1}{m} \sum_{i=1}^{m} \zeta^{\otimes i-1} \otimes \hat{\eta}_{1:m-i}^{m} \otimes |i\rangle \langle i|.$$
(S1)

This gives an overall state

$$\zeta \otimes \omega = \frac{1}{m} \sum_{i=1}^{m} \zeta^{\otimes i} \otimes \hat{\eta}_{1:m-i}^{m} \otimes |i\rangle \langle i|.$$

We now aim to use this catalyst to complete the expected transformation. First, performing a classically controlled transformation on $\zeta \otimes \omega$, we get

$$\nu_{1} := \left(\mathcal{I} \otimes \sum_{i=1}^{m-1} |i\rangle \langle i| + \frac{1}{p} \mathcal{L}_{2} \otimes |m\rangle \langle m| \right) (\zeta \otimes \omega)$$
$$= \frac{1}{m} \sum_{i=1}^{m-1} \zeta^{\otimes i} \otimes \hat{\eta}_{1:m-i}^{m} \otimes |i\rangle \langle i| + \frac{1}{m} \hat{\eta}^{m} \otimes |m\rangle \langle m|,$$

where \mathcal{I} is an identity map. This succeeds with probability p. Second, by cyclically permuting the classical registers of ν_1 such that $i \to i + 1$ and $m \to 1$, we get

$$\nu_2 := \frac{1}{m} \sum_{i=1}^m \zeta^{\otimes i-1} \otimes \hat{\eta}^m_{1:m-i+1} \otimes |i\rangle \langle i|.$$
(S2)

Third, permuting the quantum registers in a similar way such that $i \to i+1$ and $m \to 1$, we get the resulting state ν_3 . Note that ν_3 and ν_2 are the same state but with different system orders. Finally, note that the embedding operation is reversible by removing the ancillary registers. Denote $\mathcal{E}^{-1}(\cdot) = \operatorname{Tr}_{S^{k-1}}(\cdot)$ to be the reverse embedding operation and $\nu := \mathcal{E}^{-1} \otimes \mathcal{I}^{\otimes m-1}(\nu_3)$ be the final state after reverse embedding.

After these transformations, we can show that (i) the marginal state on the first system of ν is our target state satisfying the target error; (ii) the marginal state on the last m-1 systems of ν is exactly the catalyst ω . For the first claim, the marginal state on the first system of ν (or equivalently, the marginal state on the last system of ν_2 after reverse embedding) is given by

$$\nu' := \mathcal{E}^{-1} \left(\frac{1}{m} \sum_{i=1}^{m} \hat{\eta}_i^m \right) = \frac{1}{m} \sum_{i=1}^{m} \eta_i^m.$$

Since $\Delta(\eta_i^m, \sigma) \leq \varepsilon$ we have $\Delta(\nu', \sigma) \leq \varepsilon$ by the convexity of trace distance. The second claim is evident by checking that the first m-1 systems of ν_2 gives ω in Eq. (S1). This gives a one-shot catalytic distillation protocol with the same performance in target error ε and success probability p, completing the proof.



FIG. 1. An illustration for the steps of the procedure used to prove Theorem S1, for the case where n = 15, m = 5, and therefore k = 3. Each dot represents a quantum state on the system S^3 . Red dots correspond to groups of states $\zeta = \rho^{\otimes 3} \in \mathcal{D}(S^3)$, while blue dots represent embedded states $\hat{\sigma} = \sigma \otimes \pi^{\otimes 2} \in \mathcal{D}(S^3)$ that match the system of ζ . The overall quantum state is a mixture of rows, each labeled with a classical register. In the first step, a classically controlled free operation is applied, transforming the last row from $\zeta^{\otimes 5} = \rho^{\otimes 15}$ to $\hat{\sigma}^{\otimes 5}$. The second step involves cyclically permuting the classical registers (i.e., the rows of dots), and the third step involves cyclically permuting the quantum registers (i.e., the columns of dots). The dashed boxes highlight the catalyst state, showing that it remains unchanged before and after the transformation.

Theorem S2 For a deterministic one-shot catalytic resource distillation, after $l \ge 1$ repeated uses of the catalyst ω_A , we obtain a joint state $\nu_{S_1S_2...S_lA}$ such that the catalyst is exactly returned on its marginal $\nu_A = \omega_A$ and the target states $\nu_{S_1} = \nu_{S_2} = \cdots = \nu_{S_l}$ with error $\Delta(\nu_{S_i}, \sigma_S) \le \varepsilon$ for all $i \in \{1, \dots, l\}$.

Proof. Let $T = S^k$ be the systems on the source state and $\zeta_T = \rho_S^{\otimes k}$. Let \mathcal{L} be the catalytic transformation. In the first round, we get $\mathcal{L}_{TA\to S_1A}(\zeta_T \otimes \omega_A) = \nu_{S_1A}$ with $\Delta(\nu_{S_1}, \sigma_{S_1}) \leq \varepsilon$, and $\nu_A = \omega_A$. Then we apply \mathcal{L} again to a fresh copy of the source state ζ_T together with the catalyst and leave the state on the system S_1 untouched. Denote the output systems as S_2A where S_2 is isomorphic to S_1 . This gives the global state $\mathcal{L}_{TA\to S_2A}(\zeta_T \otimes \nu_{S_1A})$. Then we can check the catalyst on the system A by $\operatorname{Tr}_{S_2S_1}\mathcal{L}_{TA\to S_2A}(\zeta_T \otimes \nu_{S_1A}) = \operatorname{Tr}_{S_2}\mathcal{L}_{TA\to S_2A}(\zeta_T \otimes \omega_A) = \omega_A$. So the marginal state on the catalytic system is unchanged. We can also check the target state on the system S_2 by $\operatorname{Tr}_{S_1A}\mathcal{L}_{TA\to S_2A}(\zeta_T \otimes \nu_{S_1A}) = \operatorname{Tr}_A\mathcal{L}_{TA\to S_2A}(\zeta_T \otimes \omega_A) = \operatorname{Tr}_A\mathcal{L}_{TA\to S_1A}(\zeta_T \otimes \omega_A) = \nu_{S_1}$, which means that the marginal state we distill remains exactly the same as in the first round. The stated result is obtained by applying the above process repeatedly l times.

Note that the catalyst is guaranteed to remain effective in deterministic protocols. In probabilistic transformations, however, there is a risk of losing the validity of the catalyst if the transformation fails, requiring its re-preparation. This risk arises from the probabilistic nature of multi-shot protocols. Nevertheless, if the multi-shot protocol fails with probability p, it is guaranteed that the probability of losing the catalyst in the converted one-shot catalytic protocol is no greater than p.

TRADING SUCCESS PROBABILITY FOR REDUCED OVERHEAD

Theorem S4 Suppose there exists a distillation protocol transforming $\rho^{\otimes n}$ to $\sigma^{\otimes m}$ with success probability p and target error ε . Then it holds for any $1 \leq k \leq n/m$ that $\widetilde{C}_{\varepsilon,pm\lceil n/k\rceil^{-1}}(\rho,\sigma) \leq k$. In particular, $\widetilde{C}_{\varepsilon,pm/n}(\rho,\sigma) = 1$.

Proof. Let $\mathcal{L} \in \mathcal{O}_{sub}$ be a resource distillation protocol that transforms n copies of the source state ρ_S to m $(m \leq n)$ copies of the target state σ_S with success probability p and within target error ε . That is, there exists a quantum state $\eta^m \in \mathscr{D}(S^m)$, such that $\mathcal{L}(\rho^{\otimes n}) = p\eta^m$ and $\Delta(\eta^m_i, \sigma) \leq \varepsilon$ for all $i \in \{1, \dots, m\}$, where η^m_i is the *i*-th marginal of η^m . Then we can consider $g = \lceil n/k \rceil \geq m$ groups of $\zeta = \rho^{\otimes k}$. Since $gk = k \lceil n/k \rceil \geq n$, there exists a free operation \mathcal{L}_1 such that $\mathcal{L}_1(\zeta^{\otimes g}) = \mathcal{L}_1(\rho^{\otimes gk}) = p\eta^m$ and $\Delta(\eta^m_i, \sigma) \leq \varepsilon$, for any $i \in \{1, \dots, m\}$. This can be done by simply abandoning the residual gk - n copies of the source states and then performing the transformation \mathcal{L} on $\rho^{\otimes n}$. Then we need to prove that there exists a catalytic transformation \mathcal{L}' that transforms one copy of ζ to one copy of σ with success probability pm/g and within target error ε .

Before the actual transformation, we need to do embedding twice to ensure that the systems match. First of all, as $\zeta \in \mathscr{D}(S^k)$ and $\sigma \in \mathscr{D}(S)$, we need to embed σ into a larger space by using the embedding $\mathcal{E}(\cdot) = (\cdot) \otimes \pi_S^{\otimes (k-1)}$ where $\pi \in \mathscr{F}$ is a free state. Let $\hat{\eta}^m = \mathcal{E}^{\otimes m}(\eta^m)$ and $\hat{\sigma} = \mathcal{E}(\sigma)$. Then, since $m \leq g$, we need to do a second embedding to compensate this system mismatching by using free states. Moreover, to avoid mixing the free state with our target state which could compromise the final fidelity, we need to append the free states on an orthogonal Hilbert space of $\mathscr{D}(S^k)$ so that we can distinguish them via projective measurements. More explicitly, let θ be a free states on the system W. We embed it into a joint Hilbert space $T = S^k W$, with the embedded state denoted as $\tilde{\theta}$. We then embed $\zeta, \hat{\sigma} \in \mathscr{D}(S^k)$ into the larger space T, with the embedded states denoted as $\tilde{\zeta} \in \mathscr{D}(T)$ and $\tilde{\sigma} \in \mathscr{D}(T)$, respectively. Similarly, we embed $\hat{\eta}^m$ into $\mathscr{D}(T^m)$ and denote it as $\tilde{\eta}^m$.

Then there exists a distillation protocol \mathcal{L}_2 that transforms $\tilde{\zeta}^{\otimes g} \in \mathscr{D}(T^g)$ to $\tilde{\eta}_m \otimes \tilde{\theta}^{\otimes g-m} \in \mathscr{D}(T^g)$. Let $\tilde{\eta}_{1:i}^m$ be the marginal state of $\tilde{\eta}_m$ on the first *i* systems and $\tilde{\eta}_i^m$ be the marginal state on the *i*-th system. Define the following catalyst state on $\mathscr{D}(T^{g-1}F)$ with classical register F:

$$\omega = \frac{1}{g} \sum_{i=1}^{g-m-1} \tilde{\zeta}^{\otimes i-1} \otimes \tilde{\eta}^m \otimes \tilde{\theta}^{\otimes g-m-i} \otimes |i\rangle \langle i| + \frac{1}{g} \sum_{i=g-m}^g \tilde{\zeta}^{\otimes i-1} \otimes \tilde{\eta}^m_{1:g-i} \otimes |i\rangle \langle i|.$$

Note that for m = g or m = g - 1 the first term vanishes. We have the overall state on $\mathcal{D}(T^g F)$

$$\tilde{\zeta} \otimes \omega = \frac{1}{g} \sum_{i=1}^{g-m-1} \tilde{\zeta}^{\otimes i} \otimes \tilde{\eta}^m \otimes \tilde{\theta}^{\otimes g-m-i} \otimes |i\rangle \langle i| + \frac{1}{g} \sum_{i=g-m}^g \tilde{\zeta}^{\otimes i} \otimes \tilde{\eta}^m_{1:g-i} \otimes |i\rangle \langle i|.$$

We then proceed with the following steps. First, performing a classically controlled operation, we get

$$\begin{split} \nu_{1} &:= \left(\mathcal{I} \otimes \sum_{i=1}^{g-1} |i\rangle \langle i| + \frac{1}{p} \mathcal{L}_{2} \otimes |g\rangle \langle g| \right) (\tilde{\zeta} \otimes \omega) \\ &= \frac{1}{g} \sum_{i=1}^{g-m-1} \tilde{\zeta}^{\otimes i} \otimes \tilde{\eta}^{m} \otimes \tilde{\theta}^{\otimes g-m-i} \otimes |i\rangle \langle i| + \frac{1}{g} \sum_{i=g-m}^{g-1} \tilde{\zeta}^{\otimes i} \otimes \tilde{\eta}^{m}_{1:g-i} \otimes |i\rangle \langle i| + \frac{1}{g} \tilde{\eta}^{m} \otimes \tilde{\theta}^{\otimes g-m} \otimes |g\rangle \langle g|. \end{split}$$

Second, by cyclically permuting the classical registers of ν_1 such that $i \to i+1$ and $g \to 1$, we get

$$\nu_2 := \frac{1}{g} \sum_{i=1}^{g-m} \tilde{\zeta}^{\otimes i-1} \otimes \tilde{\eta}^m \otimes \tilde{\theta}^{\otimes g-m-i+1} \otimes |i\rangle \langle i| + \frac{1}{g} \sum_{i=g-m+1}^{g} \tilde{\zeta}^{\otimes i-1} \otimes \tilde{\eta}^m_{1:g-i+1} \otimes |i\rangle \langle i|.$$

Third, permuting the quantum registers in a similar way such that $i \to i + 1$ and $g \to 1$, we get the resulting state ν_3 . Note that ν_3 and ν_2 are the same state but with different system orders. Then we can check that the last g - 1 quantum systems of ν_3 returns the catalyst, or equivalently, we can check the first g - 1 systems of ν_2 and get

$$\frac{1}{g}\sum_{i=1}^{g-m}\tilde{\zeta}^{\otimes i-1}\otimes\tilde{\eta}^m\otimes\tilde{\theta}^{\otimes g-m-i}\otimes|i\rangle\langle i|+\frac{1}{g}\sum_{i=g-m+1}^g\tilde{\zeta}^{\otimes i-1}\otimes\tilde{\eta}^m_{1:g-i}\otimes|i\rangle\langle i|=\omega.$$

The first quantum system of ν_3 or equivalently the last quantum system of ν_2 gives our target state:

$$\nu' = \frac{1}{g} \sum_{i=1}^{g-m} \tilde{\theta} + \frac{1}{g} \sum_{i=g-m+1}^{g} \tilde{\eta}_{g-i+1}^m = \frac{m}{g} \nu'' + \left(1 - \frac{m}{g}\right) \tilde{\theta}, \quad \text{with} \quad \nu'' = \frac{1}{m} \sum_{i=1}^{m} \tilde{\eta}_i^m$$

Since ν'' and $\tilde{\theta}$ are living effectively on $\mathscr{D}(S^k)$ and $\mathscr{D}(W)$ respectively, we can perform a projective measurement $\{P_{S^k}, P_W\}$ to distinguish these two states, where P_{S^k} and P_W are the projectors on the two subspaces. So we will have a probability m/g to obtain the final state ν'' . Since $\Delta(\eta_i^m, \sigma) \leq \varepsilon$, we have $\Delta(\nu'', \sigma) \leq \varepsilon$ by the invariance of trace distance under our embeddings and the convexity of trace distance. This concludes the proof.



FIG. 2. An illustration for the steps of the procedure used to prove Theorem S4, for the case where n = 5, m = 2, and k = 1. Each dot represents a quantum state on the system T. Red dots correspond to the embedded source state $\tilde{\rho} \in \mathscr{D}(T)$, while blue dots represent the embedded target state $\tilde{\sigma} \in \mathscr{D}(T)$. Grey dots represent the embedded free state $\tilde{\theta} \in \mathscr{D}(T)$, which is orthogonal to $\tilde{\sigma}$. The overall quantum state is a mixture of rows, each labeled with a classical register. In the first step, a classically controlled free operation is applied, transforming the last row from $\tilde{\rho}^{\otimes 5}$ to $\tilde{\sigma}^{\otimes 2} \otimes \tilde{\theta}^{\otimes 3}$. The second step involves cyclically permuting the classical registers (i.e., the rows of dots), and the third step involves cyclically permuting the quantum registers (i.e., the columns of dots). The dashed boxes highlight the catalyst state, showing that it remains unchanged before and after the transformation.

Remark 1 Some remarks concerning the above proof are in order:

- 1. Appending the free state θ on the orthogonal subspace is crucial, as it allows the target state to be post-selected after the projective measurement, ensuring that the fidelity is not compromised in the final step.
- 2. The catalyst used in the proof is not unique. Other options could also work in the same manner, such as

$$\omega = \frac{1}{g} \sum_{i=1}^{g-m} \tilde{\zeta}^{\otimes i-1} \otimes \tilde{\eta}^m_{1:m-1} \otimes \tilde{\theta}^{\otimes g-m-i+1} \otimes |i\rangle \langle i| + \frac{1}{g} \sum_{i=g-m+1}^g \tilde{\zeta}^{\otimes i-1} \otimes \tilde{\eta}^m_{1:g-i} \otimes |i\rangle \langle i|.$$

Such a catalyst is independent on the target state $\tilde{\eta}^m$ if m = 1.

CHANNEL MANIPULATION

In this section, we extend the catalysis technique to the channel setting. Before addressing catalytic channel manipulation, we first need to clarify the definition of a reduced channel. It is known that if a bipartite channel

satisfies the non-signaling condition $\operatorname{Tr}_{B_2} \circ \mathcal{N}_{A_1A_2 \to B_1B_2} = \operatorname{Tr}_{B_2} \circ \mathcal{N}_{A_1A_2 \to B_1B_2} \circ \mathcal{R}_{A_2}^{\pi}$, then we can define a unique channel such that $\operatorname{Tr}_{B_2} \circ \mathcal{N}_{A_1A_2 \to B_1B_2} = \mathcal{N}_{A_1 \to B_1} \circ \operatorname{Tr}_{A_2}$. This channel is given by

$$\mathcal{N}_{A_1 \to B_1}(\cdot) \coloneqq \operatorname{Tr}_{B_2} \mathcal{N}_{A_1 A_2 \to B_1 B_2}((\cdot) \otimes \pi_{A_2}), \tag{S3}$$

where π is a fixed state. By the non-signaling assumption, we can easily verify that the definition in (S3) is independent of the choice of π . Therefore, without loss of generality, we can always take π to be the maximally mixed state of appropriate dimension. See also discussions in [59, Section 2.2].

More generally, if the channel $\mathcal{N}_{A_1A_2 \to B_1B_2}$ does not satisfy the non-signaling condition, we can still define a linear map via (S3), which is clearly a quantum channel from $A_1 \to B_1$. The only difference is that the channel will depend on the choice of π . Hence, we denote this channel by $\mathcal{N}_{A_1 \to B_1}^{\pi}$ to indicate its dependence on the reference state π .

Let $\text{CPTP}(A \to B)$ be the set of all quantum channels (i.e., completely positive and trace-preserving maps) from A to B. Define $\Delta(\mathcal{N}, \mathcal{M}) := \frac{1}{2} ||\mathcal{N} - \mathcal{M}||_{\diamond}$ be the error between quantum channels \mathcal{N} and \mathcal{M} where $|| \cdot ||_{\diamond}$ is the diamond norm [60]. Now we define the catalytic channel transformation as follows.

Definition 1 Let $\mathcal{N} \in \operatorname{CPTP}(A \to B)$ and $\mathcal{M} \in \operatorname{CPTP}(A' \to B')$ be two quantum channels. We say \mathcal{N} can be catalytically transformed to \mathcal{M} with target error ε if there exists a free superchannel [50] Π and a channel $\mathcal{C} \in \operatorname{CPTP}(X \to Y)$ such that the output channel $\mathcal{P} := \Pi_n(\mathcal{N} \otimes \mathcal{C}_n) \in \operatorname{CPTP}(A' X \to B' Y)$ satisfying

$$\mathcal{P}_{X \to Y} = \mathcal{C}_{X \to Y}, \quad and \quad \Delta(\mathcal{P}_{A'X \to B'Y}, \mathcal{M}_{A' \to B'} \otimes \mathcal{C}_{X \to Y}) \le \varepsilon, \tag{S4}$$

where $\mathcal{P}_{X\to Y}(\cdot) := \operatorname{Tr}_{B'} \mathcal{P}_{A'X\to B'Y}((\cdot)\otimes \pi_{A'})$. Moreover, \mathcal{N} is said to be catalytically transformed to \mathcal{M} if there exsits a sequence of superchannels Π_n and a sequence of catalyst channel \mathcal{C}_n such that $\mathcal{N}^{\otimes n}$ can be catalytically transformed to $\mathcal{M}^{\otimes n}$ with error ε_n and $\lim_{n\to\infty} \varepsilon_n = 0$.

The first condition in (S4) implies that after the coding strategy Π the transmission scheme $\mathcal{P}_{X \to Y}(\cdot)$ functioning the same as the channel \mathcal{C} . So the functioning of \mathcal{C} is not violated by the coding and we can regard it as a catalyst channel. The second condition in (S4) implies that the correlation between the source channel and the catalyst channel is within a target error. This particularly implies that the reduced channel $\mathcal{P}_{A' \to B'}(\cdot) := \operatorname{Tr}_Y \mathcal{P}_{A'X \to B'Y}((\cdot) \otimes \pi_X)$ will function approximately as \mathcal{M} , in the sense that $\Delta(\mathcal{P}_{A' \to B'}, \mathcal{M}_{A' \to B}) \leq \varepsilon$. This can be checked as follows,

$$\begin{aligned} \|\mathcal{P}_{A'\to B'} - \mathcal{M}_{A'\to B'}\|_{\diamond} &= \sup_{\rho_{RA'}} \|\mathcal{P}_{A'\to B'}(\rho_{RA'}) - \mathcal{M}_{A'\to B'}(\rho_{RA'})\|_{1} \\ &= \sup_{\rho_{RA'}} \|\operatorname{Tr}_{Y} \mathcal{P}_{A'X\to B'Y}(\rho_{RA'}\otimes\pi_{X}) - \mathcal{M}_{A'\to B'}(\rho_{RA'})\|_{1} \\ &= \sup_{\rho_{RA'}} \|\operatorname{Tr}_{Y} \left[\mathcal{P}_{A'X\to B'Y}(\rho_{RA'}\otimes\pi_{X}) - \mathcal{M}_{A'\to B'}(\rho_{RA'})\otimes\mathcal{C}_{X\to Y}(\pi_{X})\right]\|_{1} \\ &\leq \sup_{\rho_{RA'}} \|\mathcal{P}_{A'X\to B'Y}(\rho_{RA'}\otimes\pi_{X}) - \mathcal{M}_{A'\to B'}(\rho_{RA'})\otimes\mathcal{C}_{X\to Y}(\pi_{X})\|_{1} \\ &\leq \sup_{\rho_{RA'_X}} \|\mathcal{P}_{A'X\to B'Y}(\rho_{RA'X}) - \mathcal{M}_{A'\to B'}\otimes\mathcal{C}_{X\to Y}(\rho_{RA'X})\|_{1} \\ &= \|\mathcal{P}_{A'X\to B'Y} - \mathcal{M}_{A'\to B'}\otimes\mathcal{C}_{X\to Y}\|_{\diamond}, \end{aligned}$$
(S5)

where the first inequality follows by the data-processing inequality of trace norm and the second inequality follows by relaxing the input state to all quantum states on RA'X.

With the above definition, we are now ready to discuss quantum channel simulation. By the well-known quantum reverse Shannon theorem [49], we know that channel manipulation under entanglement-assisted or nonsignalling (NS)-assisted codes is reversible. This, in particular, implies that \mathcal{N} can be asymptotically transformed to \mathcal{M} with unit rate if and only if $I(\mathcal{N}) \geq I(\mathcal{M})$, where

$$I(\mathcal{N}) := \sup_{\varphi_{RA}} I(B:R)_{\mathcal{N}_{A \to B}(\varphi_{RA})}$$

is the quantum channel mutual information and $I(B:R)_{\rho} = S(\rho_B) + S(\rho_R) - S(\rho_{BR})$, giving an operational meaning of channel mutual information in the conventional i.i.d. setting. Extending the catalysis technique for quantum states, the following result endows the channel mutual information an operational meaning at the one-shot level. It shows that given free entanglement and catalyst channels, \mathcal{N} can be catalytically transformed to one \mathcal{M} if and only if $I(\mathcal{N}) \geq I(\mathcal{M})$, thereby addressing the open question discussed in [28]. **Theorem 9** Let $\mathcal{N}, \mathcal{M} \in \operatorname{CPTP}(A \to B)$ be two quantum channels. Then \mathcal{N} can be catalytically transformed to \mathcal{M} under entanglement-assisted or NS-assisted codes if and only if $I(\mathcal{N}) \geq I(\mathcal{M})$.

The proof of this result requires the following lemmas.

Lemma 10 For any quantum channel \mathcal{N} and superchannel Π , the output channel $\Pi(\mathcal{N})$ satisfies $I(\Pi(\mathcal{N})) \leq I(\mathcal{N})$.

Proof. This can be understood from the existing operational interpretation of a channel's mutual information, which represents its entanglement-assisted quantum capacity. Since the channel $\Pi(\mathcal{N})$ is noisier than \mathcal{N} , it results in reduced communication capability. A similar argument is provided in [54, Remark 5].

Moreover, the quantum channel mutual information is continuous with respect to the diamond norm.

Lemma 11 Let $\mathcal{N}, \mathcal{M} \in \operatorname{CPTP}(A \to B)$ and d_{AB} be the dimension of $A \otimes B$. If $\|\mathcal{N} - \mathcal{M}\|_{\diamond} \leq \varepsilon$, it holds

$$|I(\mathcal{N}) - I(\mathcal{M})| \le 3\varepsilon \log d_{AB} + 3h_2(\varepsilon) := f(\varepsilon).$$

Proof. For any $\delta > 0$, choose φ_1 and φ_2 be such that $|I(\mathcal{N}) - I(B:R)_{\mathcal{N}(\varphi_1)}| \leq \delta$ and $|I(\mathcal{M}) - I(B:R)_{\mathcal{N}(\varphi_2)}| \leq \delta$ where $d_R = d_A$. Since $||\mathcal{N} - \mathcal{M}||_{\diamond} \leq \varepsilon$, we have $||\mathcal{N}(\varphi_1) - \mathcal{M}(\varphi_1)||_1 \leq \varepsilon$ and $||\mathcal{N}(\varphi_2) - \mathcal{M}(\varphi_2)||_1 \leq \varepsilon$. Recall the Fannes inequality $|S(\rho) - S(\psi)| \leq \varepsilon \log d + h_2(\varepsilon)$ if $||\rho - \psi||_1 \leq \varepsilon$. We have the continuity of the quantum mutual information as

$$|I(B:R)_{\rho} - I(B:R)_{\psi}| \le |S(\rho_B) - S(\psi_B)| + |S(\rho_R) - S(\psi_R)| + |S(\rho_{BR}) - S(\psi_{BR})|$$

$$\le 3\varepsilon \log d_{AB} + 3h_2(\varepsilon) := f(\varepsilon).$$

if $\|\rho_{BR} - \psi_{BR}\|_1 \leq \varepsilon$. Putting everything together, we have

$$I(\mathcal{N}) - \delta \le I(B:R)_{\mathcal{N}(\varphi_1)} \le I(B:R)_{\mathcal{M}(\varphi_1)} + f(\varepsilon) \le I(\mathcal{M}) + f(\varepsilon),$$

$$I(\mathcal{M}) - \delta \le I(B:R)_{\mathcal{M}(\varphi_2)} \le I(B:R)_{\mathcal{N}(\varphi_2)} + f(\varepsilon) \le I(\mathcal{N}) + f(\varepsilon).$$

So we get $|I(\mathcal{N}) - I(\mathcal{M})| \leq \delta + f(\varepsilon)$. Since δ can be made arbitrarily small, we get $|I(\mathcal{N}) - I(\mathcal{M})| \leq f(\varepsilon)$.

We are now ready to prove the result in Theorem 9.

Proof. We first show the "only if" direction. Suppose \mathcal{N} can be catalytically transformed to \mathcal{M} . For any $\varepsilon > 0$, there exists a catalyst channel \mathcal{C}_n and a code Π_n such that $\Delta(\Pi_n(\mathcal{N} \otimes \mathcal{C}_n), \mathcal{M} \otimes \mathcal{C}_n) \leq \varepsilon$. By the continuity of the channel mutual information and the monotonicity in Lemma 11 and Lemma 10, we get

$$I(\mathcal{M} \otimes \mathcal{C}_n) \le I(\Pi_n(\mathcal{N} \otimes \mathcal{C}_n)) + f(2\varepsilon) \le I(\mathcal{N} \otimes \mathcal{C}_n) + f(2\varepsilon).$$

Then by the additivity of channel mutual information, we get $I(\mathcal{M}) \leq I(\mathcal{N}) + f(2\varepsilon)$. Since the inequality holds for arbitrary ε , we get $I(\mathcal{M}) \leq I(\mathcal{N})$.

Now, we prove the "if" direction. That is, if $I(\mathcal{N}) \geq I(\mathcal{M})$ we can find a sequence of Π_n and \mathcal{C}_n that catalytically transform \mathcal{N} to \mathcal{M} . By the reversibility of the channel simulation, we know that for any given ε , there exists entanglement-assisted or NS-assisted transformations such that

$$\Pi_n(\mathcal{N}^{\otimes n}) = \mathcal{P}^n \quad \text{and} \quad \|\mathcal{P}^n - \mathcal{M}^{\otimes n}\|_{\diamond} \le \varepsilon.$$

Note that $\mathcal{P}^n \in \operatorname{CPTP}(A^n \to B^n)$. Define the reduced channel on the first *i* systems as

$$\mathcal{P}_{1:i}^{n}(\cdot) := \operatorname{Tr}_{B_{i+1:n}} \mathcal{P}_{A^{n} \to B^{n}}^{n}((\cdot) \otimes \pi_{A_{i+1:n}}),$$

where $\mathcal{P}_{1:n}^n := \mathcal{P}^n$. Now consider a catalyst channel

$$\mathcal{C}_n := \frac{1}{n} \sum_{k=1}^n \mathcal{N}^{\otimes k-1} \otimes \mathcal{P}_{1:n-k}^n \otimes |k\rangle \langle k|.$$

Then the overall channel is given by

$$\mathcal{N} \otimes \mathcal{C}_n = \frac{1}{n} \sum_{k=1}^n \mathcal{N}^{\otimes k} \otimes \mathcal{P}_{1:n-k}^n \otimes |k\rangle \langle k|.$$

Performing a classically controlled transformation $\mathcal{I} \otimes \sum_{k=1}^{n-1} + \Pi_n \otimes |n\rangle \langle n|$, we get

$$\mathcal{G}_1 = \frac{1}{n} \sum_{k=1}^{n-1} \mathcal{N}^{\otimes k} \otimes \mathcal{P}^n_{1:n-k} \otimes |k\rangle \langle k| + \frac{1}{n} \mathcal{P}^n \otimes |n\rangle \langle n|.$$

Next, we relabel the classical registers $i \to i+1$ and $n \to 1$ and get

$$\mathcal{G}_2 = \frac{1}{n} \sum_{k=1}^n \mathcal{N}^{\otimes k-1} \otimes \mathcal{P}_{1:n-k+1}^n \otimes |k\rangle \langle k|.$$

Then performing a cyclic permutation S_{A^n} and S_{B^n} on the input and the output, respectively, such that $i \to i + 1$ and $n \to 1$, we get

$$\mathcal{G}_3 = \frac{1}{n} \sum_{k=1}^n \mathcal{S}_{B^n} \circ (\mathcal{N}^{\otimes k-1} \otimes \mathcal{P}_{1:n-k+1}^n) \circ \mathcal{S}_{A^n} \otimes |k\rangle \langle k|.$$

Then we claim that \mathcal{G}_3 returns \mathcal{C}_n on its marginal and $\|\mathcal{G}_3 - \mathcal{M} \otimes \mathcal{C}_n\|_{\diamond} \leq 2\varepsilon$. The first claim is equivalent to check the first n-1 reduced channel of \mathcal{G}_2 ,

$$\operatorname{Tr}_{B_n} \circ \mathcal{G}_2((\cdot) \otimes \pi_{A_n}) = \frac{1}{n} \sum_{k=1}^n \operatorname{Tr}_{B_n} \mathcal{N}^{\otimes k-1} \otimes \mathcal{P}_{1:n-k+1}^n((\cdot) \otimes \pi_{A_n}) \otimes |k\rangle \langle k|$$
$$= \frac{1}{n} \sum_{k=1}^n \mathcal{N}^{\otimes k-1} \otimes \operatorname{Tr}_{B_n} \mathcal{P}_{1:n-k+1}^n((\cdot) \otimes \pi_{A_n}) \otimes |k\rangle \langle k|$$
$$= \frac{1}{n} \sum_{k=1}^n \mathcal{N}^{\otimes k-1} \otimes \mathcal{P}_{1:n-k}^n(\cdot) \otimes |k\rangle \langle k|$$
$$= \mathcal{C}_n.$$

Note that the second last equality holds by the definition of \mathcal{P}_i^n and here π_{A_n} and Tr_{B_n} are effectively acting on the last systems of the channel $\mathcal{P}_{1:n-k+1}^n$, which is also the n-k+1 system in the original definition of $\mathcal{P}_{1:n-k+1}^n$. Since the diamond norm is invariant under system permutation, the second claim is equivalent to check $\|\mathcal{G}_2 - \mathcal{C}_n \otimes \mathcal{M}\|_{\diamond} \leq 2\varepsilon$. This can be shown as follows:

$$\begin{aligned} \|\mathcal{G}_{2} - \mathcal{C}_{n} \otimes \mathcal{M}\|_{\diamond} &= \left\| \frac{1}{n} \sum_{k=1}^{n} \mathcal{N}^{\otimes k-1} \otimes \mathcal{P}_{1:n-k+1}^{n} \otimes |k\rangle \langle k| - \frac{1}{n} \sum_{k=1}^{n} \mathcal{N}^{\otimes k-1} \otimes \mathcal{P}_{1:n-k}^{n} \otimes \mathcal{M} \otimes |k\rangle \langle k| \right\|_{\diamond} \\ &= \frac{1}{n} \sum_{k=1}^{n} \left\| \mathcal{N}^{\otimes k-1} \otimes \mathcal{P}_{1:n-k+1}^{n} - \mathcal{N}^{\otimes k-1} \otimes \mathcal{P}_{1:n-k}^{n} \otimes \mathcal{M} \right\|_{\diamond} \\ &= \frac{1}{n} \sum_{k=1}^{n} \left\| \mathcal{P}_{1:n-k+1}^{n} - \mathcal{P}_{1:n-k}^{n} \otimes \mathcal{M} \right\|_{\diamond} \\ &\leq \frac{1}{n} \sum_{k=1}^{n} \left\| \mathcal{P}_{1:n-k+1}^{n} - \mathcal{M}^{\otimes n-k+1} \right\|_{\diamond} + \frac{1}{n} \sum_{k=1}^{n} \left\| \mathcal{P}_{1:n-k}^{n} \otimes \mathcal{M} - \mathcal{M}^{\otimes n-k+1} \right\|_{\diamond} \\ &= \frac{1}{n} \sum_{k=1}^{n} \left\| \mathcal{P}_{1:n-k+1}^{n} - \mathcal{M}^{\otimes n-k+1} \right\|_{\diamond} + \frac{1}{n} \sum_{k=1}^{n} \left\| \mathcal{P}_{1:n-k}^{n} - \mathcal{M}^{\otimes n-k+1} \right\|_{\diamond} \\ &\leq 2\varepsilon. \end{aligned}$$

where the first inequality follows by the triangle inequality of diamond norm and the last inequality follows from the assumption of $\|\mathcal{P}^n - \mathcal{M}^{\otimes n}\|_{\diamond} \leq \varepsilon$ and the monotonicity of diamond norm by taking reduced channel (same argument as Eq. (S5)). As this holds for arbitrary ε , we can take ε to be vanishingly small and conclude the proof.