

Thermodynamic functions of the Fermi gas at arbitrary temperatures

Yu.M. Poluektov, A.A. Soroka

National Science Center “Kharkov Institute of Physics and Technology”,
61108 Kharkov, Ukraine

E-mail: yuripoluektov@kipt.kharkov.ua (y.poluekt52@gmail.com)

Abstract

Thermodynamic functions of the ideal Fermi gas at arbitrary temperatures are calculated using the standard Fermi-Stoner functions. The properties of the Fermi-Stoner functions are analyzed. The limiting cases of low-temperature and classical limits with taking into account quantum corrections and the special case of zero chemical potential are considered.

Key words: fermion, electron, thermodynamic functions, equation of state, entropy, heat capacity, thermodynamic coefficients

PACS numbers: 05.30.Fk, 05.70.Ce, 64.10.+h, 67.10.Db, 67.10.Fj, 71.10.Ca

Contents

Introduction	2
1 Thermodynamic functions of the Fermi gas at arbitrary temperatures	3
1.1 Thermodynamic potential	4
1.2 Number of particles	4
1.3 Energy	4
1.4 Pressure	4
1.5 Entropy	4
1.6 Adiabatic equations	5
1.7 The degree of “quantumness” of the gas (“quantumness”)	6
1.8 Heat capacities	7
1.9 Isothermal and adiabatic speeds of sound	9
1.10 Number of particles collisions with a wall	9
1.11 Thermodynamic coefficients	10
2 Properties of the Fermi-Stoner functions	11
3 Thermodynamic quantities in the limiting and special cases	14
3.1 The classical limit and quantum corrections	14
3.2 Fermi gas close to degeneracy	15
3.3 Fermi gas at zero chemical potential	17
4 Conclusion	19
A appendix	20
B appendix	24
References	24
List of designations	26
Some fundamental constants	27

Introduction

The theory of ideal gases is presented in many textbooks and problem books on statistical mechanics [1 – 8]. However, as a rule, authors limit themselves to considering two limiting cases: a) high temperatures, at which the description of the motion of gas particles on the basis of the laws of classical mechanics is valid, and b) very low temperatures, when gases are close to degeneracy. Meanwhile, in low-temperature investigations, a situation is quite real when a system, which can be considered as an ideal gas (for example, quasiparticles in quantum liquids), is no longer classical, but is still far from strong degeneracy. In this intermediate case, for calculations of thermodynamic functions formulas relating to the above limiting cases cannot be used, but exact expressions

should be used. Since in the known to us literature on statistical physics the methodology of such calculations is not presented, the current work is devoted to the presentation of methods of calculation of thermodynamic functions for the Fermi gas at arbitrary temperatures. The ideal gas model allows to verify all fundamental thermodynamic relations on the basis of a microscopic description and to express all thermodynamic quantities through standard functions. In addition, a detailed study of the thermodynamic properties of ideal gases helps to better understand the general principles of statistical mechanics.

1 Thermodynamic functions of the Fermi gas at arbitrary temperatures

Thermodynamic functions of the Fermi gas at arbitrary temperatures are expressed in terms of the standard Fermi-Stoner functions.

In the ideal Fermi gas, the momentum distribution of particles with a half-integer spin is described by the Fermi-Dirac function

$$f_k = (\exp \beta \xi_k + 1)^{-1}, \quad (1.1)$$

where $\xi_k = \hbar^2 k^2 / 2m - \mu$, $\beta = 1/T$ – the inverse temperature, T – the temperature, μ – the chemical potential, k – the wave number of a particle, m – its mass. All thermodynamic functions of the ideal Fermi gas can be expressed through special functions defined by the formula

$$\Phi_s(t) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{z^{s-1} dz}{e^{z-t} + 1}, \quad (1.2)$$

where s is a positive integer or half-integer number. Similar functions were first introduced by Stoner [9] and therefore we will call them *the Fermi-Stoner functions*. The parameter t in (1.2) takes any real values from $-\infty$ to ∞ . The properties of these functions will be discussed in detail in the next section.

Further formulas will include the de Broglie thermal wavelength of a particle, defined by the formula

$$\Lambda \equiv \Lambda(T) = \left(\frac{2\pi\hbar^2}{mT} \right)^{1/2}. \quad (1.3)$$

As seen, the existence of the characteristic length Λ is conditioned by quantum laws, since the definition (1.3) contains the Planck's constant in the numerator. When passing to the classical description we should formally set $\hbar \rightarrow 0$, so that $\Lambda \rightarrow 0$. In reality, the decrease of the thermal wavelength, and hence the transition to the classical description, occurs as the temperature increases. The criterion of transition from the quantum to the classical description of a gas will be discussed in more detail below.

When finding the following relations, one should use the representation of the distribution function in the form of a series

$$\frac{1}{e^\xi + 1} = \begin{cases} \sum_{n=0}^{\infty} (-1)^n e^{n\xi}, & \xi < 0, \\ \sum_{n=0}^{\infty} (-1)^n e^{-(n+1)\xi}, & \xi > 0. \end{cases} \quad (1.4)$$

1.1 Thermodynamic potential

The thermodynamic potential of the ideal Fermi gas through the Fermi-Stoner function is expressed by the formula

$$\Omega = -gT \sum_k \ln(1 + e^{-\beta\xi_k}) = -\frac{gTV}{\Lambda^3} \Phi_{5/2}(t), \quad (1.5)$$

where $t = \beta\mu$, $g \equiv 2s + 1$, s – the particle spin. For the electron and the nucleons $s = 1/2$ and $g = 2$. The parameter $t = \beta\mu$, equal to the product of the chemical potential and the inverse temperature, is often encountered in the theory of quantum gases, so let us give this parameter the name – *the adiabaticity parameter*. The reason for this name will be explained below.

1.2 Number of particles

The total number of particles as a function of the chemical potential and temperature (or the parameter $t = \beta\mu$ and temperature) is determined by the formula

$$N = g \sum_k f_k = \frac{gV}{\Lambda^3} \Phi_{3/2}(t). \quad (1.6)$$

Note that the thermodynamic relation $N = -(\partial\Omega/\partial\mu)_T$ is fulfilled.

1.3 Energy

The energy is determined by the formula

$$E = g \sum_k \frac{\hbar^2 k^2}{2m} f_k = \frac{3}{2} \frac{gTV}{\Lambda^3} \Phi_{5/2}(t). \quad (1.7)$$

As seen, the known relation $\Omega = -\frac{2}{3}E$ is fulfilled.

1.4 Pressure

Since $\Omega = -pV$, the pressure is determined by the formula

$$p = \frac{gT}{\Lambda^3} \Phi_{5/2}(t). \quad (1.8)$$

Obviously, there holds the equation $pV = \frac{2}{3}E$.

1.5 Entropy

The entropy of the ideal Fermi gas can be found either from the combinatorial definition

$$S = -g \sum_k [f_k \ln f_k + (1 - f_k) \ln(1 - f_k)], \quad (1.9)$$

or using the thermodynamic relation $S = -(\partial\Omega/\partial T)_\mu$:

$$S = \frac{gV}{\Lambda^3} \left[\frac{5}{2}\Phi_{5/2}(t) - t\Phi_{3/2}(t) \right]. \quad (1.10)$$

Note that for the above thermodynamic quantities the known thermodynamic relation holds $E = \mu N + TS - pV$.

Formulas (1.6)–(1.8), (1.10) determine parametrically (the parameter t) the energy, pressure and entropy as functions of the temperature and particle number density. Practically, however, it is sometimes more convenient to use another dimensionless parameter θ . From formulas (1.6) and (1.8) it follows that

$$\frac{nT}{p} \equiv \theta = \frac{\Phi_{3/2}(t)}{\Phi_{5/2}(t)}, \quad (1.11)$$

where $n = N/V$ – the particle number density. Thus, formula (1.11) uniquely links the parameters θ and t . The dependence $\theta = \theta(t)$ is shown in Fig. 1. The parameter θ varies within the finite limits $0 < \theta < 1$. At $t \rightarrow -\infty$ $\theta \rightarrow 1$, and at $t \rightarrow +\infty$ $\theta \approx 5/2t$. By calculating the dependence of any thermodynamic quantity on θ , it is easy to obtain its dependence on temperature, density or pressure with the other two variables being fixed.




Figure 1: Dependence of the parameter θ on the adiabaticity parameter t .

1.6 Adiabatic equations

The entropy per one particle

$$\sigma \equiv \frac{S}{N} = \frac{5}{2} \frac{\Phi_{5/2}(t)}{\Phi_{3/2}(t)} - t \quad (1.12)$$

depends only on the parameter t . In adiabatic processes, when $\sigma = const$, the parameter $t = \mu/T$ is also constant. This is the reason for the adopted name of t – *the adiabaticity parameter*. The parameter θ (1.11) also does not change in adiabatic processes. The dependencies $\sigma = \sigma(t)$ and $\sigma = \sigma(\theta)$ are shown in Fig. 2. In the limit $t \rightarrow -\infty$ the dependence (1.12) becomes linear $\sigma \cong 5/2 - t$, and at $t \rightarrow \infty$ the entropy per one particle decreases as $\sigma \cong \pi^2/2t$. At $\theta \ll 1$ there is a linear dependence $\sigma(\theta) \approx \frac{\pi^2}{5} \theta$, and at $\theta \rightarrow 1$, when $t \approx (5/2) \ln 2 + \ln(1 - \theta)$, we have $\sigma(\theta) \approx (5/2)(1 - \ln 2) - \ln(1 - \theta)$.




Figure 2: Dependences of the entropy per one particle on: (a) the parameter $t - \sigma(t)$, (b) the parameter $\theta - \sigma(\theta)$.

Taking into account the constancy of t , from formulas (1.6), (1.8) there follow the adiabatic equations

$$\frac{n}{T^{3/2}} = \frac{g}{A} \Phi_{3/2}(t), \quad \frac{p}{n^{5/3}} = \frac{A^{2/3} \Phi_{5/2}(t)}{g^{2/3} \Phi_{3/2}^{5/3}(t)}, \quad \frac{p}{T^{5/2}} = \frac{g}{A} \Phi_{5/2}(t), \quad (1.13)$$

where $A = (2\pi\hbar^2/m)^{3/2}$. The values of the constant quantities on the right-hand side of equations (1.13) are completely determined by the adiabaticity parameter t or the parameter θ .

1.7 The degree of “quantumness” of the gas (“quantumness”)

Depending on the particle number density and temperature, the quantum mechanical properties of the gas will manifest themselves to a greater or lesser extent. As a measure of “quantumness” it is natural to use the ratio of the de Broglie thermal wavelength (1.3) to the average distance between particles $l = n^{-1/3}$. However, it is more convenient to take as such a measure the ratio of the volumes of spheres with radii Λ and l . Thus, the number q , characterizing the measure of “quantumness” of the gas, is defined as the ratio

$$q \equiv \left(\frac{\Lambda}{l}\right)^3 = \Lambda^3 n = g \Phi_{3/2}(t). \quad (1.14)$$

As seen, q depends only on the adiabaticity parameter t , i.e. the value of this parameter (or θ) uniquely determines the degree of “quantumness” of the gas. In the following, for brevity we will call the number q “quantumness”. With decreasing temperature at a fixed density the thermal length increases, and hence the quantumness also increases in this case. The dependences of the quantumness q on the parameters t and θ are presented in Fig. 3. The quantumness of the gas increases with increasing the parameter t and with decreasing θ . In the limit $t \rightarrow -\infty$ or $\theta \rightarrow 1$, the quantumness $q \rightarrow 0$ and the motion of gas particles obeys the laws of classical mechanics. The quantumness can be found based on the temperature and the particle number density, and the adiabaticity parameter t for a given state of the gas is determined based on the known quantumness from formula (1.14).




Figure 3: Dependences of the “quantumness” on: (a) the parameter $t - q(t)/g$, (b) the parameter $\theta - q(\theta)/g$.

1.8 Heat capacities

When finding heat capacities and thermodynamic coefficients, it is convenient to use the following expressions for differentials

$$dN = \frac{N}{V} dV + \frac{3N}{2T} dT + \frac{gV}{\Lambda^3} \Phi_{1/2} dt, \quad (1.15)$$

$$dp = \frac{5p}{2T} dT + \frac{gT}{\Lambda^3} \Phi_{3/2} dt, \quad (1.16)$$

$$dS = \frac{S}{V} dV + \frac{3S}{2T} dT + \frac{gV}{\Lambda^3} \left(\frac{3}{2} \Phi_{3/2} - t\Phi_{1/2} \right) dt. \quad (1.17)$$

Here, the relation is used $d\Lambda/\Lambda = -dT/2T$. The heat capacity for an arbitrary change of thermodynamic variables with temperature is defined by the expression

$$C \equiv T \frac{dS}{dT} = \frac{ST}{V} \frac{dV}{dT} + \frac{3S}{2} + \frac{gVT}{\Lambda^3} \left(\frac{3}{2} \Phi_{3/2} - t\Phi_{1/2} \right) \frac{dt}{dT}. \quad (1.18)$$

As a rule, systems with a fixed average number of particles are considered. In this case $dN = 0$, and from (1.15) it follows that

$$\frac{dV}{dT} = -\frac{3V}{2T} - \frac{gV^2}{N\Lambda^3} \Phi_{1/2} \frac{dt}{dT}. \quad (1.19)$$

For a constant volume from (1.19) we have

$$\frac{dt}{dT} = -\frac{3N\Lambda^3}{2gVT} \Phi_{1/2}^{-1}, \quad (1.20)$$

and for a constant pressure from (1.16) it follows

$$\frac{dt}{dT} = -\frac{5p\Lambda^3}{2gT^2} \Phi_{3/2}^{-1}. \quad (1.21)$$

Formulas (1.18)–(1.21) allow us to obtain formulas for the heat capacities of the gas under certain conditions. The heat capacity per one particle at a constant volume is determined by the formula

$$c_V \equiv \frac{C_V}{N} = \frac{15}{4} \frac{\Phi_{3/2}(t)}{\Phi_{1/2}(t)} \left[\frac{\Phi_{1/2}(t)\Phi_{5/2}(t)}{\Phi_{3/2}^2(t)} - \frac{3}{5} \right], \quad (1.22)$$

and the heat capacity at a constant pressure – by the formula

$$c_p \equiv \frac{C_p}{N} = \frac{25}{4} \frac{\Phi_{5/2}(t)}{\Phi_{3/2}(t)} \left[\frac{\Phi_{1/2}(t)\Phi_{5/2}(t)}{\Phi_{3/2}^2(t)} - \frac{3}{5} \right]. \quad (1.23)$$

Formulas (1.22), (1.23), together with (1.11), determine parametrically the dependences of the heat capacities on the positive parameter θ : $C_V/N = c_V(\theta)$, $C_p/N = c_p(\theta)$. The dependences of the heat capacities and their ratios on the parameters t and θ are shown in Fig. 4. In the classical limit $t \rightarrow -\infty$, the heat capacities tend to constant values $c_p = 5/2$, $c_V = 3/2$, and their ratio to $c_p/c_V = 5/3$. In the quantum limit $t \rightarrow \infty$, the heat capacities tend to zero according to the law $c_p \cong c_V \cong \pi^2/2t$, and their ratio tends to unity. For small $\theta \ll 1$ (the quantum limit) we have $c_V(\theta) = c_p(\theta) \approx \frac{\pi^2}{5} \theta$.




Figure 4: Dependences of the heat capacities per one particle and their ratios on the parameters t and θ : (a) 1 – $c_V(t)$, 2 – $c_p(t)$, 3 – $c_p(t)/c_V(t)$; (b) 1 – $c_V(\theta)$, 2 – $c_p(\theta)$, 3 – $c_p(\theta)/c_V(\theta)$.

Let us also give formulas for the ratio and the difference of heat capacities

$$\frac{C_p}{C_V} = \frac{5}{3} \frac{\Phi_{1/2}(t)\Phi_{5/2}(t)}{\Phi_{3/2}^2(t)}, \quad (1.24)$$

$$C_p - C_V = \frac{25}{4} \frac{gV}{\Lambda^3} \frac{\Phi_{3/2}^2(t)}{\Phi_{1/2}(t)} \left[\frac{\Phi_{1/2}(t)\Phi_{5/2}(t)}{\Phi_{3/2}^2(t)} - \frac{3}{5} \right]^2. \quad (1.25)$$

From (1.25) it is obvious that the condition $C_p - C_V > 0$ is satisfied. The heat capacities per one particle at constant values of volume and pressure (1.22), (1.23), as well as the ratio of heat capacities (1.24), are functions of only the adiabaticity parameter and remain constant in adiabatic processes.

1.9 Isothermal and adiabatic speeds of sound

The squares of isothermal and adiabatic speeds of sound are defined as derivatives of the pressure with respect to the density under the appropriate conditions and have the form

$$u_T^2 = \frac{1}{m} \left(\frac{\partial p}{\partial n} \right)_T = \frac{T}{m} \frac{\Phi_{3/2}(t)}{\Phi_{1/2}(t)}, \quad (1.26)$$

$$u_\sigma^2 = \frac{1}{m} \left(\frac{\partial p}{\partial n} \right)_\sigma = \frac{5T}{3m} \frac{\Phi_{5/2}(t)}{\Phi_{3/2}(t)}. \quad (1.27)$$

Their ratio is equal to the ratio of heat capacities (1.24)

$$\frac{u_\sigma^2}{u_T^2} = \frac{C_p}{C_V} = \frac{5}{3} \frac{\Phi_{1/2}(t)\Phi_{5/2}(t)}{\Phi_{3/2}^2(t)}. \quad (1.28)$$

We emphasize that this relation is valid at all temperatures.

It is of interest to determine the average of the square of the momentum of a gas particle. Based on the definition of the average, and taking into account relations (1.6), (1.7), we have:

$$\langle \hbar^2 k^2 \rangle = \frac{g \sum_k \hbar^2 k^2 f_k}{g \sum_k f_k} = \frac{2mE}{N} = 3mT \frac{\Phi_{5/2}(t)}{\Phi_{3/2}(t)}. \quad (1.29)$$

From here it follows that the average of the square of the velocity of a gas particle is proportional to the square of the adiabatic speed of sound (1.27) for all values of thermodynamic variables:

$$\langle v^2 \rangle = \frac{\langle \hbar^2 k^2 \rangle}{m^2} = \frac{9}{5} u_\sigma^2. \quad (1.30)$$

Since the functions $\Phi_s(t)$ take only positive values, the squares of the sound speeds are positive and *the conditions of thermodynamic stability*

$$\left(\frac{\partial p}{\partial n} \right)_T > 0, \quad C_V > 0, \quad (1.31)$$

or

$$\left(\frac{\partial p}{\partial n} \right)_\sigma > 0, \quad C_p > 0, \quad (1.32)$$

are satisfied for the ideal Fermi gas. In addition, the pressure is always positive $p > 0$.

1.10 Number of particles collisions with a wall

Above, the formula for pressure (1.8) was found using the thermodynamic relation. It is useful to determine the gas pressure on a wall directly through the number of particle collisions. Let the gas fill the space at $z > 0$. The number of impacts per unit area of a wall per unit time from particles with momenta in the range from $\hbar\mathbf{k}$ to $\hbar\mathbf{k} + \hbar d\mathbf{k}$ is equal to

$$d\nu = g|v_z|f_k \frac{2\pi\kappa d\kappa dk_z}{(2\pi)^3}, \quad (1.33)$$

where $|v_z| = \hbar|k_z|/m$, $\kappa = \sqrt{k_x^2 + k_y^2}$. The total number of impacts per unit area of a wall per unit time is obtained by integrating the relation (1.33):

$$\nu = \frac{gmT^2}{4\pi^2\hbar^3} \Phi_2(t). \quad (1.34)$$

Since at each collision a particle transfers momentum $2mv_z$ to a wall, then, multiplying (1.33) by this quantity and integrating, we obtain formula (1.8) for the pressure. Note that the quantities characterizing the volumetric properties of the gas are expressed through functions $\Phi_s(t)$ with a half-integer index s , while the number of impacts per unit area of a wall (1.34) is expressed through the function with an integer s .

1.11 Thermodynamic coefficients

Let us also present following from (1.15), (1.16) formulas for *the coefficient of volumetric expansion*

$$\alpha_p = \frac{1}{V} \left(\frac{\partial V}{\partial T} \right)_p = \frac{5}{2T} \left[\frac{\Phi_{1/2}(t)\Phi_{5/2}(t)}{\Phi_{3/2}^2(t)} - \frac{3}{5} \right], \quad (1.35)$$

the isothermal compressibility

$$\gamma_T = -\frac{1}{V} \left(\frac{\partial V}{\partial p} \right)_T = \frac{\Lambda^3 \Phi_{1/2}(t)}{gT \Phi_{3/2}^2(t)} \quad (1.36)$$

and *the isochoric thermal pressure coefficient*

$$\beta_V = \frac{1}{p} \left(\frac{\partial p}{\partial T} \right)_V = \frac{5}{2T} \frac{\Phi_{3/2}^2(t)}{\Phi_{1/2}(t)\Phi_{5/2}(t)} \left[\frac{\Phi_{1/2}(t)\Phi_{5/2}(t)}{\Phi_{3/2}^2(t)} - \frac{3}{5} \right]. \quad (1.37)$$

Comparing with (1.25), it is easy to verify that the known thermodynamic relation is fulfilled

$$C_p - C_V = TV \frac{\alpha_p^2}{\gamma_T}. \quad (1.38)$$

The remaining thermodynamic coefficients that can be constructed from the variables T, S and p, V are expressed through the above coefficients and the heat capacities [3]. The dependences of thermodynamic coefficients (1.35)–(1.37) on the parameters t and θ are shown in Fig. 5.




Figure 5: Dependencies of thermodynamic coefficients on:

(a) t : 1 – $\alpha_p(t)T$, 2 – $\beta_V(t)T$, 3 – $\gamma_T(t)gT\Lambda^{-3}$; (b) θ : 1 – $\alpha_p(\theta)T$, 2 – $\beta_V(\theta)T$, 3 – $\gamma_T(\theta)gT\Lambda^{-3}$.

2 Properties of the Fermi-Stoner functions

The general properties of the Fermi-Stoner functions and their calculation in the high-temperature and low-temperature limits are considered.

In this section we consider in more detail the properties of the *Fermi-Stoner* functions defined by the formula

$$\Phi_s(t) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{z^{s-1} dz}{e^{z-t} + 1}, \quad (2.1)$$

where s can be a positive integer or half-integer number, $\Gamma(s)$ is the gamma function, $t \equiv \mu/T$. As a rule, it is sufficient to use functions with indices $s = 1/2, 3/2, 5/2$, as well as the derivative $\Phi'_{1/2}(t)$. In such case: $\Gamma(1/2) = \sqrt{\pi}$, $\Gamma(3/2) = \sqrt{\pi}/2$, $\Gamma(5/2) = 3\sqrt{\pi}/4$. The form of these functions is shown in Fig. 6.




Figure 6: Graphs of the functions $\Phi_s(t)$ at $s = 1/2, 3/2, 5/2$ and the derivative $\Phi'_{1/2}(t)$; the maximum of the function $\Phi'_{1/2}(t)$ lies at $t_m = 1.11$.

At $t \leq 0$ the Stoner functions can be represented as a series

$$\Phi_s(t) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{e^{nt}}{n^s}. \quad (2.2)$$

At positive values of the parameter t the functions (2.1) can be written in the form

$$\Phi_s(t) = \frac{2t^s}{\Gamma(s)} \int_0^\infty \frac{x^{2s-1} dx}{e^{t(x^2-1)} + 1}. \quad (2.3)$$

In particular,

$$\Phi_0(t) = \frac{e^t}{1 + e^t}, \quad \Phi_1(t) = \ln(1 + e^t). \quad (2.4)$$

For $s \geq 1$, the derivative of the function of order s is determined by the function of order $s - 1$:

$$\frac{d\Phi_s(t)}{dt} = \Phi_{s-1}(t). \quad (2.5)$$

The functions $\Phi_s(t)$ are positive, and

$$\Phi_s(t) > \Phi_{s-1}(t). \quad (2.6)$$

At the characteristic point $t = 0$ the formula is valid

$$\Phi_s(0) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^s} = (1 - 2^{1-s})\zeta(s), \quad (2.7)$$

where $\zeta(s)$ is the Riemann zeta function [10]. Taking into account the values

$$\zeta(1/2) = -1.46035, \quad \zeta(3/2) = 2.61238, \quad \zeta(5/2) = 1.34149,$$

we obtain

$$\Phi_{1/2}(0) = 0.60490, \quad \Phi_{3/2}(0) = 0.76515, \quad \Phi_{5/2}(0) = 0.86720.$$

Since formula (2.5) is not valid for the derivative $\Phi'_{1/2}(t)$, this and higher derivatives of the function $\Phi_{1/2}(t)$ should be calculated separately. At $t < 0$, the derivative $\Phi'_{1/2}(t)$ can be calculated by differentiating the series (2.2), so that

$$\Phi'_{1/2}(t) = \sum_{n=1}^{\infty} (-1)^{n+1} \sqrt{n} e^{nt}. \quad (2.8)$$

At $t < 0$ the higher derivatives are calculated similarly. The series (2.8) converges the faster the smaller the value of t .

To calculate the derivatives of the function $\Phi_{1/2}(t)$ for arbitrary t , including $t \geq 0$, let us define the integrals

$$\Phi_{1/2}(t; n) \equiv \frac{1}{\sqrt{\pi}} \int_0^{\infty} \frac{z^{-1/2} dz}{(e^{z-t} + 1)^n} = \frac{2}{\sqrt{\pi}} \int_0^{\infty} \frac{du}{(e^{u^2-t} + 1)^n}. \quad (2.9)$$

The second form of recording the integral is more convenient for numerical calculations due to its faster convergence. Obviously, $\Phi_{1/2}(t; 1) = \Phi_{1/2}(t)$. The derivatives of the function $\Phi_{1/2}(t; n)$ with respect to t are expressed through the integrals (2.9):

$$\frac{d\Phi_{1/2}(t; n)}{dt} = n[\Phi_{1/2}(t; n) - \Phi_{1/2}(t; n+1)], \quad (2.10)$$

$$\frac{d^2\Phi_{1/2}(t; n)}{dt^2} = n[n\Phi_{1/2}(t; n) - (2n+1)\Phi_{1/2}(t; n+1) + (n+1)\Phi_{1/2}(t; n+2)], \quad (2.11)$$

$$\begin{aligned} \frac{d^3\Phi_{1/2}(t; n)}{dt^3} = n[n^2\Phi_{1/2}(t; n) - (3n^2 + 3n + 1)\Phi_{1/2}(t; n+1) + \\ + 3(n+1)^2\Phi_{1/2}(t; n+2) - (n+1)(n+2)\Phi_{1/2}(t; n+3)], \end{aligned} \quad (2.12)$$

$$\begin{aligned} \frac{d^4\Phi_{1/2}(t; n)}{dt^4} = n[n^3\Phi_{1/2}(t; n) - (4n^3 + 6n^2 + 4n + 1)\Phi_{1/2}(t; n+1) + \\ + (n+1)(6n^2 + 12n + 7)\Phi_{1/2}(t; n+2) - (n+1)(n+2)(4n+6)\Phi_{1/2}(t; n+3) + \\ + (n+1)(n+2)(n+3)\Phi_{1/2}(t; n+4)]. \end{aligned} \quad (2.13)$$

At $n = 1$, the relations (2.10)–(2.13) determine the derivatives of the function $\Phi_{1/2}(t)$. In particular, at $t = 0$ we have

$$\Phi'_{1/2}(0) = 0.38011, \quad \Phi''_{1/2}(0) = 0.11868, \quad \Phi'''_{1/2}(0) = -0.087841, \quad \Phi^{(IV)}_{1/2}(0) = -0.096048.$$

In the limit $t \rightarrow -\infty$, which corresponds to the classical case, it is sufficient to restrict ourselves to the first terms of the series (2.2)

$$\Phi_s(t) \approx e^t - \frac{e^{2t}}{2^s} + \frac{e^{3t}}{3^s} - \dots \quad (2.14)$$

In this approximation the relation of t with the parameter $\theta = nT/p$ (1.11) is given by the formulas

$$\theta - 1 = -\frac{e^t}{2^{5/2}} + \left(\frac{2}{3^{5/2}} - \frac{1}{32} \right) e^{2t}, \quad (2.15)$$

$$e^t = 2^{5/2}(1 - \theta) + 2^{17/2} \left(\frac{1}{3^{5/2}} - \frac{1}{64} \right) (1 - \theta)^2. \quad (2.16)$$

Another limiting case $t \rightarrow +\infty$ corresponds to the case of a quantum gas close to degeneracy near zero temperature. In this case, there holds an asymptotic expansion that is valid up to exponentially small terms

$$\Phi_s(t) = \frac{t^s}{\Gamma(s)} \left[\frac{1}{s} + 2 \sum_{k=0}^{\infty} C_{s-1}^{2k+1} \frac{(2k+1)!}{t^{2k+2}} (1 - 2^{-(2k+1)}) \zeta(2k+2) \right]. \quad (2.17)$$

Here for the binomial coefficients at integer $\nu \equiv n$ we have $C_\nu^r \equiv C_n^r = \frac{n!}{(n-r)!r!}$. If ν is not an integer, then $C_\nu^r = \frac{\nu(\nu-1)\dots(\nu-r+1)}{r!}$. The first terms of the expansion (2.17) have the form

$$\Phi_s(t) = \frac{t^s}{s\Gamma(s)} \left[1 + \frac{\pi^2}{6} \frac{s(s-1)}{t^2} + \frac{7\pi^4}{360} \frac{s(s-1)(s-2)(s-3)}{t^4} + \dots \right]. \quad (2.18)$$

In particular,

$$\Phi_{1/2}(t) = \frac{2t^{1/2}}{\sqrt{\pi}} \left[1 - \frac{\pi^2}{24t^2} - \frac{7\pi^4}{384t^4} \right], \quad (2.19)$$

$$\Phi_{3/2}(t) = \frac{4t^{3/2}}{3\sqrt{\pi}} \left[1 + \frac{\pi^2}{8t^2} + \frac{7\pi^4}{640t^4} \right], \quad (2.20)$$

$$\Phi_{5/2}(t) = \frac{8t^{5/2}}{15\sqrt{\pi}} \left[1 + \frac{5\pi^2}{8t^2} - \frac{7\pi^4}{384t^4} \right], \quad (2.21)$$

$$\Phi_2(t) = \frac{t^2}{2} \left[1 + \frac{\pi^2}{3t^2} \right]. \quad (2.22)$$

In this approximation the parameters t and θ are linked by the relations

$$\theta = \frac{5}{2t} \left[1 - \frac{\pi^2}{2t^2} + \frac{41\pi^4}{120t^4} \right], \quad (2.23)$$

$$\frac{1}{t} = \frac{2}{5}\theta + \frac{4\pi^2}{125}\theta^3. \quad (2.24)$$

The derivation of the asymptotic expansion (2.17) with account of exponential terms is given in Appendix A.

3 Thermodynamic quantities in the limiting and special cases

There is considered the behavior of thermodynamic quantities in the classical limit with account of the quantum correction, in the low-temperature limit and at zero chemical potential.

3.1 The classical limit and quantum corrections

Let us consider the limiting case $t \rightarrow -\infty$. In this case, it is sufficient to take into account some first terms in the series (2.2). We restrict ourselves to taking into account the first two terms, the first of which gives the classical limit, and the second one gives the main quantum correction to it. With a chosen accuracy the parameter t can be expressed from formulas (1.14), (2.14) through the parameter $q = \Lambda^3 n$, which was called the quantumness:

$$e^t = \frac{q}{g} \left(1 + \frac{q}{2^{3/2}g} \right). \quad (3.1)$$

This expansion is valid under condition $q \ll 1$, which is satisfied if the de Broglie thermal wavelength of a particle is much smaller than the average distance between gas particles.

The energy and pressure of the gas in this limit are given by the formulas

$$E = \frac{3NT}{2} \left(1 + \frac{q}{2^{5/2}g} \right), \quad (3.2)$$

$$p = nT \left(1 + \frac{q}{2^{5/2}g} \right). \quad (3.3)$$

As seen, the quantum correction to the ideal Fermi gas leads to an increase of the pressure, which corresponds to the effective repulsion of particles. This effect is conditioned by the antisymmetry of the wave function of a system of identical Fermi particles with respect to permutations.

The parameter t enters into the entropy linearly, so that

$$t = \ln \left(\frac{q}{g} \right) + \frac{q}{2^{3/2}g}. \quad (3.4)$$

The entropy takes the form $S = S_{cl} + S'$, where entropy in the classical limit is determined by the formula

$$S_{cl} = N \ln \left(\frac{ge^{5/2}}{q} \right). \quad (3.5)$$

This formula is called the Sakura-Tetrode formula [11, 12]. As seen, the entropy per one particle S_{cl}/N is determined only by the quantumness q . Note that this formula in the high-temperature limit includes the Planck's constant \hbar (through the thermal length). This indicates the special nature of this thermodynamic function, which even in the classical limit requires attracting the quantum-mechanical concepts for its correct definition. The mentioned circumstance is caused by the combinatorial nature of entropy, because of which its calculation requires taking into account the quantum-mechanical indistinguishability of particles at all temperatures. This also indicates that statistical physics can be built consistently only on the basis of quantum mechanics. The next quantum mechanical correction to the entropy has the form

$$S' = \frac{Nq}{2^{7/2}g}. \quad (3.6)$$

The heat capacities at constant volume and pressure with account of quantum corrections are expressed by the formulas

$$c_V \equiv \frac{C_V}{N} = \frac{3}{2} \left(1 - \frac{q}{2^{7/2}g} \right), \quad (3.7)$$

$$c_p \equiv \frac{C_p}{N} = \frac{5}{2} \left(1 - \frac{3q}{2^{7/2}g} \right). \quad (3.8)$$

The squares of the isothermal and adiabatic speeds of sound with account of quantum corrections have the form

$$u_T^2 = \frac{T}{m} \left(1 + \frac{q}{2^{3/2}g} \right), \quad (3.9)$$

$$u_\sigma^2 = \frac{5T}{3m} \left(1 + \frac{q}{2^{5/2}g} \right). \quad (3.10)$$

Quantum corrections result in an increase of the speed of sound. Let us also present the ratios of heat capacities and squares of speeds

$$\frac{C_p}{C_V} = \frac{u_\sigma^2}{u_T^2} = \frac{5}{3} \left(1 - \frac{q}{2^{5/2}g} \right), \quad (3.11)$$

as well as the formula for the difference of heat capacities

$$\frac{C_p - C_V}{N} = 1 - \frac{3q}{2^{5/2}g}. \quad (3.12)$$

Thermodynamic coefficients (1.35)–(1.37) with account of the first quantum correction have the form

$$\alpha_p = \frac{1}{T} \left(1 - \frac{5}{2^{7/2}} \frac{q}{g} \right), \quad (3.13)$$

$$\gamma_T = \frac{1}{nT} \left(1 - \frac{q}{2^{3/2}g} \right), \quad (3.14)$$

$$\beta_V = \frac{1}{T} \left(1 - \frac{3}{2^{7/2}} \frac{q}{g} \right). \quad (3.15)$$

3.2 Fermi gas close to degeneracy

Let us present formulas for thermodynamic functions near $T = 0$ with account of the first temperature correction. The thermodynamic potential in the case under consideration has the form

$$\Omega = -\frac{4g m^{3/2} \mu^{5/2} V}{15\sqrt{2} \pi^2 \hbar^3} \left(1 + \frac{5\pi^2 T^2}{8 \mu^2} \right). \quad (3.16)$$

The particle number density as a function of temperature and chemical potential is given by the formula

$$n(T, \mu) = \frac{\sqrt{2} g m^{3/2} \mu^{3/2}}{3\pi^2 \hbar^3} \left(1 + \frac{\pi^2 T^2}{8 \mu^2} \right). \quad (3.17)$$

From the latter formula, the chemical potential can be expressed through the temperature and particle number density

$$\mu = \mu_0 \left(1 - \frac{\pi^2 T^2}{12 \mu_0^2} \right), \quad (3.18)$$

where

$$\mu_0 \equiv \varepsilon_F = \left(\frac{6\pi^2}{g} \right)^{2/3} \frac{\hbar^2}{2m} n^{2/3} \quad (3.19)$$

is the value of the chemical potential at $T = 0$, coinciding with the Fermi energy. As seen, at $T = 0$ the chemical potential takes on a finite positive value, so that at $T \rightarrow 0$ the parameter $t = \mu/T$ increases indefinitely $t \rightarrow \infty$. The Fermi momentum is equal to

$$p_F \equiv \sqrt{2m\varepsilon_F} = \hbar \left(\frac{6\pi^2}{g} \right)^{1/3} n^{1/3}. \quad (3.20)$$

The energy of the gas with density n that is close to degeneracy is determined by the formula

$$E = E_0 \left(1 + \frac{5\pi^2 T^2}{12 \mu_0^2} \right), \quad (3.21)$$

where the energy at zero temperature

$$E_0 = \frac{3}{10} V \left(\frac{6\pi^2}{g} \right)^{2/3} \frac{\hbar^2}{m} n^{5/3} = \frac{3}{5} N \mu_0. \quad (3.22)$$

The pressure is given by the formula

$$p = p_0 \left(1 + \frac{5\pi^2 T^2}{12 \mu_0^2} \right), \quad (3.23)$$

where the pressure at zero temperature

$$p_0 = \frac{1}{5} \left(\frac{6\pi^2}{g} \right)^{2/3} \frac{\hbar^2}{m} n^{5/3} = \frac{2}{5} \mu_0 n. \quad (3.24)$$

The entropy at absolute zero turns to zero, so that the third law of thermodynamics in the Planck formulation holds. At low temperatures the entropy is proportional to temperature, and the next term of the expansion is proportional to the cube of temperature

$$S = V \frac{m}{\hbar^2} \left(\frac{g\pi}{6} \right)^{2/3} n^{1/3} \left(T - \frac{\pi^2 T^3}{10 \mu_0^2} \right) = \frac{\pi^2 N}{2} \left(\frac{T}{\mu_0} - \frac{\pi^2 T^3}{10 \mu_0^3} \right). \quad (3.25)$$

The heat capacities have the form

$$\frac{C_V}{N} = \frac{\pi^2}{2} \left(\frac{T}{\mu_0} - \frac{3\pi^2 T^3}{10 \mu_0^3} \right), \quad (3.26)$$

$$\frac{C_p}{N} = \frac{\pi^2}{2} \left(\frac{T}{\mu_0} + \frac{\pi^2 T^3}{30 \mu_0^3} \right), \quad (3.27)$$

and their difference is proportional to the cube of temperature

$$\frac{C_p - C_V}{N} = \frac{\pi^4 T^3}{6 \mu_0^3}. \quad (3.28)$$

The ratio of heat capacities tends to unity as the temperature tends to zero

$$\frac{C_p}{C_V} = 1 + \frac{\pi^2 T^2}{3 \mu_0^2}. \quad (3.29)$$

The squares of the sound speeds in the low-temperature limit have the form

$$u_T^2 = \frac{2 \mu_0}{3 m} \left(1 + \frac{\pi^2 T^2}{12 \mu_0^2} \right), \quad (3.30)$$

$$u_\sigma^2 = \frac{2 \mu_0}{3 m} \left(1 + \frac{5\pi^2 T^2}{12 \mu_0^2} \right). \quad (3.31)$$

Thermodynamic coefficients (1.35)–(1.37) in the low-temperature limit are

$$\alpha_p = \frac{\pi^2}{2\mu_0} \left(\frac{T}{\mu_0} - \frac{23\pi^2 T^3}{60 \mu_0^3} \right), \quad (3.32)$$

$$\gamma_T = \frac{3}{2} \frac{1}{n\mu_0} \left(1 - \frac{\pi^2 T^2}{12 \mu_0^2} \right), \quad (3.33)$$

$$\beta_V = \frac{5\pi^2}{6\mu_0} \left(\frac{T}{\mu_0} - \frac{43\pi^2 T^3}{60 \mu_0^3} \right). \quad (3.34)$$

3.3 Fermi gas at zero chemical potential

There occurs Bose-Einstein condensation in the ideal gas of bosons at zero chemical potential. In the gas of fermions the chemical potential at high temperatures is negative. As the temperature decreases, it also turns to zero at a certain temperature. However, this phenomenon is not associated with any phase transition. As the temperature further decreases down to zero, the chemical potential remains positive. Let us consider separately the state of the Fermi gas under conditions when its chemical potential is zero. In this case, quantum regularities remain essential. The quantumness (1.14) at $\mu = 0$ and $g = 2$ is not small $q_0 = g\Phi_{3/2}(0) = 1.53$. The particle number density, energy and pressure at $\mu = 0$ are given by

$$n = \frac{g}{\Lambda^3} \Phi_{3/2}(0), \quad \frac{E}{V} = \frac{3}{2} \frac{gT}{\Lambda^3} \Phi_{5/2}(0), \quad p = \frac{gT}{\Lambda^3} \Phi_{5/2}(0). \quad (3.35)$$

Here are the numerical values of $\Phi_s(0)$:

$$\Phi_{1/2}(0) = 0.60490, \quad \Phi_{3/2}(0) = 0.76515, \quad \Phi_{5/2}(0) = 0.86720.$$

Sometimes it is convenient to pass to a dimensionless notation of thermodynamic quantities. For this purpose, let us choose an arbitrary non-zero temperature T_0 . The corresponding de Broglie wavelength $\Lambda_0 \equiv \Lambda(T_0) = (2\pi\hbar^2/mT_0)^{1/2}$. At this temperature the particle number density, energy and pressure are

$$n_0 = \frac{g}{\Lambda_0^3} \Phi_{3/2}(0), \quad \frac{E_0}{V} = \frac{3}{2} \frac{gT_0}{\Lambda_0^3} \Phi_{5/2}(0), \quad p_0 = \frac{gT_0}{\Lambda_0^3} \Phi_{5/2}(0). \quad (3.36)$$

The dimensionless particle number density, energy and pressure at $\mu = 0$ are functions of only the dimensionless temperature $\tau \equiv T/T_0$:

$$\tilde{n} \equiv \frac{n}{n_0} = \tau^{3/2}, \quad \tilde{p} \equiv \frac{p}{p_0} = \tau^{5/2}, \quad \tilde{E} \equiv \frac{E}{E_0} = \tau^{5/2}. \quad (3.37)$$

These dependencies are presented in Fig. 7. The entropy and heat capacities per one particle are the following numbers

$$\sigma = \frac{5 \Phi_{5/2}(0)}{2 \Phi_{3/2}(0)} = 2.833, \quad c_V = 1.404, \quad c_p = 2.097. \quad (3.38)$$

The pressure, particle number density and temperature under condition $\mu = 0$ are linked by the relation

$$\frac{nT}{p} = \theta = \frac{\Phi_{3/2}(0)}{\Phi_{5/2}(0)} = 0.882. \quad (3.39)$$

The speeds of sound (1.26), (1.27) are proportional to temperature

$$u_T^2 = \frac{T \Phi_{3/2}(0)}{m \Phi_{1/2}(0)}, \quad u_\sigma^2 = \frac{5T \Phi_{5/2}(0)}{3m \Phi_{3/2}(0)}. \quad (3.40)$$

The coefficient of volumetric expansion α_p (1.35) and the isochoric thermal pressure coefficient β_V (1.37) are inversely proportional to temperature

$$\alpha_p = \frac{5}{2T} \left[\frac{\Phi_{5/2}(0)\Phi_{1/2}(0)}{\Phi_{3/2}^2(0)} - \frac{3}{5} \right] = \frac{0.740}{T}, \quad \beta_V = \frac{5}{2T} \left[1 - \frac{3}{5} \frac{\Phi_{3/2}^2(0)}{\Phi_{5/2}(0)\Phi_{1/2}(0)} \right] = \frac{0.826}{T}. \quad (3.41)$$

The isothermal compressibility γ_T (1.36) is given by the formula

$$\gamma_T = \frac{\Lambda^3 \Phi_{1/2}(0)}{gT \Phi_{3/2}^2(0)} = \frac{1}{g} \left(\frac{2\pi\hbar^2}{m} \right)^{3/2} \frac{\Phi_{1/2}(0)}{\Phi_{3/2}^2(0)} \frac{1}{T^{5/2}}, \quad (3.42)$$

so that $\gamma_T \sim T^{-5/2}$.




Figure 7: Dependences of the density $\tilde{n}(\tau) - 1$, the pressure and energy $\tilde{p}(\tau), \tilde{E}(\tau) - 2$ on the dimensionless temperature $\tau \equiv T/T_0$ at $\mu = 0$.

4 Conclusion

The paper provides a reference material containing formulas for the basic thermodynamic quantities of the ideal Fermi gas, expressed through the standard Fermi-Stoner functions, and considers the basic properties of these functions. The limiting cases of high and low temperatures and the special case of zero chemical potential are also considered. All given relations are valid for systems of large volume and large number of particles. Quantum distribution functions over discrete states in systems of non-interacting fermions and bosons in a finite volume with an arbitrary, including small, number of particles and their properties are considered by the authors in [13–15].

A appendix

The derivation of the low-temperature asymptotics with taking into account exponential corrections.

Let us write the initial formula (2.1), valid at $t > 0$, in the form $\Phi_s(t) \equiv \frac{2t^s}{\Gamma(s)} J_s(t)$, where

$$J_s(t) \equiv \int_0^\infty \frac{x^{2s-1} dx}{e^{t(x^2-1)} + 1} = J_s^{(1)}(t) + J_s^{(2)}(t), \text{ and}$$

$$J_s^{(1)}(t) \equiv \int_0^1 \frac{x^{2s-1} dx}{e^{t(x^2-1)} + 1}, \quad J_s^{(2)}(t) \equiv \int_1^\infty \frac{x^{2s-1} dx}{e^{t(x^2-1)} + 1}. \quad (\text{A.1})$$

Making a substitution $y = 1 - x^2$ in the first integral and $y = x^2 - 1$ in the second one, we obtain

$$J_s^{(1)}(t) = \frac{1}{2} \left[\frac{1}{s} + \sum_{n=1}^\infty (-1)^n \int_0^1 dy (1-y)^{s-1} e^{-nty} \right], \quad (\text{A.2})$$

$$J_s^{(2)}(t) = -\frac{1}{2} \sum_{n=1}^\infty (-1)^n \int_0^\infty dy (1+y)^{s-1} e^{-nty}.$$

Then we can write down $J_s(t) = J_s^{(3)}(t) + J_s^{(4)}(t)$, where

$$J_s^{(3)}(t) = \frac{1}{2} \left[\frac{1}{s} + \sum_{n=1}^\infty (-1)^n Z_{sn}(t) \right], \quad J_s^{(4)}(t) = -\frac{1}{2} \sum_{n=1}^\infty (-1)^n B_{sn}(t).$$

Here

$$Z_{sn}(t) \equiv \int_0^1 dy [(1-y)^{s-1} - (1+y)^{s-1}] e^{-nty}, \quad (\text{A.3})$$

$$B_{sn}(t) \equiv \int_1^\infty dy (1+y)^{s-1} e^{-nty}. \quad (\text{A.4})$$

To calculate the integral (A.3), one should use the expansions ($\nu > 0$) $(1 \pm x)^\nu = \sum_{r=0}^\infty (\pm 1)^r C_\nu^r x^r$. If $\nu \equiv n$ is an integer, then $C_\nu^r \equiv C_n^r = \frac{n!}{(n-r)!r!}$. If ν is a half-integer, then $C_\nu^r = \frac{\nu(\nu-1)\dots(\nu-r+1)}{r!}$. For a negative exponent $(1 \pm x)^{-\nu} = \sum_{r=0}^\infty (\pm 1)^r C_{-\nu}^r x^r$, where $C_{-\nu}^r \equiv (-1)^r \frac{\nu(\nu+1)\dots(\nu+r-1)}{r!}$. Then we obtain

$$Z_{sn}(t) = \sum_{r=1}^\infty C_{s-1}^r [(-1)^r - 1] \int_0^1 dy y^r e^{-nty}. \quad (\text{A.5})$$

Here

$$\int_0^1 dy y^r e^{-nty} = -e^{-nt} a_{nr}(t) + \frac{r!}{(nt)^{r+1}}$$

and

$$a_{nr}(t) \equiv \sum_{l=1}^{r+1} \frac{r!}{(nt)^l (r-l+1)!} = \frac{1}{nt} + \frac{r}{(nt)^2} + \frac{r(r-1)}{(nt)^3} + \dots + \frac{r!}{(nt)^r} + \frac{r!}{(nt)^{r+1}}. \quad (\text{A.6})$$

Therefore

$$Z_{sn}(t) = \sum_{r=1}^\infty C_{s-1}^r [(-1)^r - 1] \left[-e^{-nt} a_{nr}(t) + \frac{r!}{(nt)^{r+1}} \right].$$

As a result, taking into account that terms only with odd $r = 2k + 1$ remain in the sum, we have

$$J_s^{(3)}(t) = \frac{1}{2} \left\{ \frac{1}{s} + 2 \sum_{k=0}^{\infty} C_{s-1}^{2k+1} \left[\sum_{n=1}^{\infty} (-1)^n e^{-nt} a_{n,2k+1}(t) - \frac{(2k+1)!}{t^{2k+2}} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^{2k+2}} \right] \right\}. \quad (\text{A.7})$$

For the zeta function $\zeta(z) = \sum_{k=1}^{\infty} \frac{1}{k^z}$ ($\text{Re } z > 1$), there holds the relation $\sum_{k=1}^{\infty} \frac{(-1)^k}{k^z} = (2^{1-z} - 1)\zeta(z)$, so that

$$J_s^{(3)}(t) = \frac{1}{2} \left\{ \frac{1}{s} + 2 \sum_{k=0}^{\infty} C_{s-1}^{2k+1} \left[A_k(t) + \frac{(2k+1)!}{t^{2k+2}} (1 - 2^{-(2k+1)}) \zeta(2k+2) \right] \right\}, \quad (\text{A.8})$$

where the notation is used

$$\begin{aligned} A_k(t) &\equiv \sum_{n=1}^{\infty} (-1)^n e^{-nt} a_{n,2k+1}(t) = (2k+1)! \sum_{l=1}^{2k+2} \frac{1}{(2k+2-l)! t^l} \sum_{n=1}^{\infty} \frac{(-1)^n e^{-nt}}{n^l} = \\ &= -(2k+1)! \sum_{l=1}^{2k+2} \frac{1}{(2k+2-l)!} \frac{\Phi_l(-t)}{t^l}. \end{aligned}$$

Calculation of the quantity (A.4) gives $B_{sn}(t) = \frac{e^{nt}}{(nt)^s} I_s(2nt)$, where

$$I_s(p) = \int_p^{\infty} du u^{s-1} e^{-u} \equiv \Gamma(s, p), \quad (\text{A.9})$$

and $\Gamma(s, p)$ – the incomplete gamma function [10].

Two cases should be considered separately: 1) s – an integer, 2) s – a half-integer.

1) s – an integer:

$$I_s(p) = e^{-p} [p^{s-1} + (s-1)p^{s-2} + \dots + (s-1)!p + (s-1)!] = e^{-p} \sum_{r=0}^{s-1} \frac{(s-1)!}{(s-r-1)!} p^{s-1-r}. \quad (\text{A.10})$$

2) s – a half-integer.

If $s = 1/2$, then

$$I_{1/2}(p) \equiv \int_p^{\infty} du \frac{e^{-u}}{\sqrt{u}} = \sqrt{\pi} \operatorname{erfc}(\sqrt{p}), \quad (\text{A.11})$$

where the complementary probability integral:

$$\operatorname{erfc}(z) \equiv \frac{2}{\sqrt{\pi}} \int_z^{\infty} e^{-t^2} dt.$$

In particular, for $x \gg 1$ the asymptotics holds $\operatorname{erfc}(z) \approx \frac{e^{-z^2}}{\sqrt{\pi} z}$.

For $s > 1/2$,

$$I_s(p) = e^{-p} \sum_{r=0}^{s-3/2} \frac{(s-1)(s-2)\dots 1/2}{(s-r-1)(s-r-2)\dots 1/2} p^{s-1-r} + (s-1)(s-2)\dots \frac{3}{2} \cdot \frac{1}{2} \cdot I_{1/2}(p). \quad (\text{A.12})$$

In particular,

$$I_{3/2}(p) = e^{-p} \cdot p^{1/2} + \frac{1}{2} I_{1/2}(p), \quad I_{5/2}(p) = e^{-p} \left(p^{3/2} + \frac{3}{2} p^{1/2} \right) + \frac{3}{2} \cdot \frac{1}{2} \cdot I_{1/2}(p). \quad (\text{A.13})$$

Thus, the integral $J_s(t)$ is decomposed into the sum of the main $J_s^{(m)}(t)$ and exponential $J_s^{(\text{exp})}(t)$ parts

$$J_s(t) = J_s^{(m)}(t) + J_s^{(\text{exp})}(t). \quad (\text{A.14})$$

The main part of the decomposition:

$$J_s^{(m)}(t) = \frac{1}{2s} + \sum_{k=0}^{\infty} C_{s-1}^{2k+1} \frac{(2k+1)!}{t^{2k+2}} (1 - 2^{-(2k+1)}) \zeta(2k+2). \quad (\text{A.15})$$

The exponential part of the decomposition:

$$J_s^{(\text{exp})}(t) \equiv J_s^{(\text{exp1})}(t) + J_s^{(\text{exp2})}(t), \quad (\text{A.16})$$

where

$$J_s^{(\text{exp1})}(t) = \sum_{k=0}^{\infty} C_{s-1}^{2k+1} A_k(t) = - \sum_{k=0}^{\infty} C_{s-1}^{2k+1} (2k+1)! \sum_{l=1}^{2k+2} \frac{1}{(2k+2-l)!} \frac{\Phi_l(-t)}{t^l}, \quad (\text{A.17})$$

$$J_s^{(\text{exp2})}(t) = \frac{1}{2} \sum_{n=1}^{\infty} (-1)^{n+1} B_{sn}(t). \quad (\text{A.18})$$

At an integer s :

$$\begin{aligned} J_s^{(\text{exp2})}(t) &= -\frac{1}{2}(s-1)! \sum_{r=0}^{s-1} \frac{1}{2^{1-s+r}(s-1-r)! t^{r+1}} \sum_{n=1}^{\infty} (-1)^n \frac{e^{-nt}}{n^{r+1}} = \\ &= \frac{1}{2}(s-1)! \sum_{r=0}^{s-1} \frac{1}{2^{1-s+r}(s-1-r)!} \frac{\Phi_{r+1}(-t)}{t^{r+1}}. \end{aligned} \quad (\text{A.19})$$

At a half-integer $s = \frac{3}{2}, \frac{5}{2}, \dots$:

$$\begin{aligned} J_s^{(\text{exp2})}(t) &= \frac{1}{2} \sum_{r=0}^{s-3/2} \frac{(s-1)(s-2) \dots 1/2}{(s-r-1)(s-r-2) \dots 1/2} \cdot 2^{s-1-r} \cdot \frac{\Phi_{r+1}(-t)}{t^{r+1}} + \\ &+ \frac{1}{2}(s-1)(s-2) \dots \frac{3}{2} \cdot \frac{1}{2} \frac{1}{t^s} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{e^{nt}}{n^s} I_{1/2}(2nt). \end{aligned} \quad (\text{A.20})$$

For $s = \frac{1}{2}$,

$$J_{1/2}^{(\text{exp2})}(t) \equiv -\frac{1}{2} \frac{1}{t^{1/2}} \sum_{n=1}^{\infty} (-1)^n \frac{e^{nt}}{n^{1/2}} \int_{2nt}^{\infty} du \frac{e^{-u}}{\sqrt{u}}. \quad (\text{A.21})$$

The special cases for $\Phi_s(t) = \frac{2t^s}{\Gamma(s)} J_s(t)$:

$$1) s = \frac{1}{2}; \Phi_{1/2}(t) = \frac{2t^{1/2}}{\Gamma(1/2)} J_{1/2}(t).$$

$$J_{1/2}^{(m)} = 1 - \frac{\pi^2}{24} \cdot \frac{1}{t^2} - \frac{7\pi^4}{384} \cdot \frac{1}{t^4} - \frac{31\pi^6}{1024} \cdot \frac{1}{t^6},$$

$$J_{1/2}^{(\text{exp1})}(t) = \sum_{k=0}^{\infty} \frac{1 \cdot 3 \cdot \dots \cdot (4k+1)}{2^{2k+1}} \sum_{l=1}^{2k+2} \frac{1}{(2k+2-l)!} \frac{\Phi_l(-t)}{t^l},$$

$$J_{1/2}^{(\text{exp2})}(t) \equiv -\frac{1}{2} \frac{1}{t^{1/2}} \sum_{n=1}^{\infty} (-1)^n \frac{e^{nt}}{n^{1/2}} \int_{2nt}^{\infty} du \frac{e^{-u}}{\sqrt{u}}.$$

$$2) s = \frac{3}{2}; \Phi_{3/2}(t) = \frac{2t^{3/2}}{\Gamma(3/2)} J_{3/2}(t).$$

$$J_{3/2}^{(m)} = \frac{1}{3} \left(1 + \frac{\pi^2}{8} \cdot \frac{1}{t^2} + \frac{7\pi^4}{640} \cdot \frac{1}{t^4} + \frac{31\pi^6}{3072} \cdot \frac{1}{t^6} \right),$$

$$J_{3/2}^{(\text{exp1})}(t) = - \sum_{k=0}^{\infty} \frac{1 \cdot (-1) \cdot (-3) \cdot \dots \cdot (1-4k)}{2^{2k+1}} \sum_{l=1}^{2k+2} \frac{1}{(2k+2-l)!} \frac{\Phi_l(-t)}{t^l},$$

$$J_{3/2}^{(\text{exp2})}(t) = \frac{1}{2^{1/2}} \frac{\Phi_1(-t)}{t} + \frac{1}{2^2} \frac{1}{t^{3/2}} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{e^{nt}}{n^{3/2}} I_{1/2}(2nt).$$

$$3) s = \frac{5}{2}; \Phi_{5/2}(t) = \frac{2t^{5/2}}{\Gamma(5/2)} J_{5/2}(t).$$

$$J_{5/2}^{(m)} = \frac{1}{5} \left(1 + \frac{5\pi^2}{8} \cdot \frac{1}{t^2} - \frac{7\pi^4}{384} \cdot \frac{1}{t^4} - \frac{155\pi^6}{21504} \cdot \frac{1}{t^6} \right),$$

$$J_{5/2}^{(\text{exp1})}(t) = - \sum_{k=0}^{\infty} \frac{3 \cdot 1 \cdot (-1) \cdot \dots \cdot (3-4k)}{2^{2k+1}} \sum_{l=1}^{2k+2} \frac{1}{(2k+2-l)!} \frac{\Phi_l(-t)}{t^l},$$

$$J_{5/2}^{(\text{exp2})}(t) = \frac{1}{2} \left\{ 2^{3/2} \frac{\Phi_1(-t)}{t} + \frac{3}{2^{1/2}} \frac{\Phi_2(-t)}{t^2} \right\} + \frac{3}{2^3} \frac{1}{t^{5/2}} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{e^{nt}}{n^{5/2}} I_{1/2}(2nt).$$

B appendix

Some mathematical relations with the Riemann zeta function and the gamma function.

The Riemann zeta function at $\text{Re } z > 1$ is defined by the series

$$\zeta(z) = \sum_{n=1}^{\infty} \frac{1}{n^z}. \quad (\text{B.1})$$

At $\text{Re } z \leq 1$, $z \neq 1$, $\zeta(z)$ is defined as the analytic continuation of this series. At the point $z = 1$ it has a pole $\lim_{z \rightarrow 1} \left[\zeta(z) - \frac{1}{z-1} \right] = C = 0.577215$ – the Euler's constant.

At $\text{Re } z > 0$ there holds the representation

$$(1 - 2^{1-z})\zeta(z) = 1 - \frac{1}{2^z} + \frac{1}{3^z} - \frac{1}{4^z} + \dots = \frac{1}{\Gamma(z)} \int_0^{\infty} \frac{t^{z-1}}{e^t + 1} dt. \quad (\text{B.2})$$

Special values:

$$\begin{aligned} \zeta(2) &= \frac{\pi^2}{6} \approx 1.6449, & \zeta(4) &= \frac{\pi^4}{90} \approx 1.0823, \\ \zeta(6) &= \frac{\pi^6}{945} \approx 1.0173, & \zeta(8) &= \frac{\pi^8}{9450} \approx 1.0041; \\ \zeta(3) &\approx 1.202, & \zeta(5) &\approx 1.0369, & \zeta(7) &\approx 1.00835; \\ \zeta(0) &= -\frac{1}{2}, & \zeta'(0) &= -\frac{1}{2} \ln(2\pi), & \zeta(-2n) &= 0; \\ \zeta(-1) &= -\frac{1}{12} \approx -0.08333, & \zeta(-3) &= \frac{1}{120} \approx 0.008333, & \zeta(-5) &= -\frac{1}{252} \approx -0.003968. \end{aligned}$$

The gamma function is defined as the integral

$$\Gamma(n) = \int_0^{\infty} x^{n-1} e^{-x} dx = \int_0^1 \left(\ln \frac{1}{x} \right)^{n-1} dx, \quad (\text{B.3})$$

which has a finite value for real $n > 0$.

Some relations:

$$\Gamma(n+1) = n\Gamma(n),$$

$$\Gamma(n) = (n-1)! \quad (n - a \text{ positive integer}).$$

Special values:

$$\Gamma(1) = \Gamma(2) = 1,$$

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}, \quad \Gamma\left(\frac{3}{2}\right) = \frac{\sqrt{\pi}}{2}, \quad \Gamma\left(\frac{5}{2}\right) = \frac{3\sqrt{\pi}}{4},$$

$$\Gamma\left(n + \frac{1}{2}\right) = 1 \cdot 3 \cdot 5 \cdot \dots \cdot (2n-3)(2n-1) \frac{\sqrt{\pi}}{2^n}.$$

References

- [1] L.D. Landau, E.M. Lifshitz, *Statistical physics: Vol. 5 (Part 1)*, Butterworth-Heinemann, 544 p. (1980).
- [2] K. Huang, *Statistical mechanics*, J. Wiley & Sons, New York, 470 p. (1963).
- [3] Yu.B. Rumer, M.S. Ryvkin, *Thermodynamics, statistical physics, and kinetics*, Mir publishers, Moscow, 600 p. (1980).
- [4] A. Isihara, *Statistical physics*, Academic Press, New York, 438 p. (1971).
- [5] I.A. Kvasnikov, *Thermodynamics and statistical Physics: theory of equilibrium systems*, Izd. MGU, Moscow, 800 p. (1991).
- [6] G. Shilling, *Statistical physics in examples*, Mir, Moscow, 431 p. (1976).
- [7] P.T. Landsberg (Editor), *Problems in thermodynamics and statistical physics*, Pion, London, 573 p. (1971).
- [8] A.I. Akhiezer, S.V. Peletminskii, *Methods of statistical physics*, Pergamon Press, Oxford, 450 p. (1981).
- [9] J. McDougall, E.C. Stoner, *The computation of Fermi-Dirac functions*, Phil. Trans. Roy. Soc. A **237**, 67–104 (1938). doi:10.1098/rsta.1938.0004
- [10] M. Abramowitz, I. Stegun (Editors), *Handbook of mathematical functions*, National Bureau of Standards Applied mathematics Series **55**, 1046 p. (1964).
- [11] H. Tetrode, *Die chemische Konstante der Gase und das elementare Wirkungsquantum*, Annalen der Physik **343**, 434–442; **344**, 255–256 (1912). doi:10.1002/andp.19123430708; 10.1002/andp.19123441112
- [12] O. Sackur, *Die Anwendung der kinetischen Theorie der Gase auf chemische Probleme*, Annalen der Physik **341**, 958–980 (1911); *Die universelle Bedeutung des sog. elementaren Wirkungsquantums*, Annalen der Physik **345**, 67–86 (1913). doi:10.1002/andp.19113411505; 10.1002/andp.19133450103
- [13] Yu.M. Poluektov, A.A. Soroka, *Quantum distribution functions in systems with an arbitrary number of particles*. arXiv:2311.03003v2 [quant-ph]
- [14] Yu.M. Poluektov, A.A. Soroka, *On the thermodynamics of two-level Fermi and Bose nanosystems*, Optical and Quantum Electronics **56**, 1349 (2024). doi:10.1007/s11082-024-07266-x; arXiv:2405.02427v1 [quant-ph]
- [15] Yu.M. Poluektov, A.A. Soroka, *Thermodynamics of the Fermi gas in a cubic cavity of an arbitrary volume*. arXiv:2408.03667v1 [quant-ph]

List of designations

m – the mass of a gas particle,

k – the wave number of a particle, $\hbar k$ – its momentum,

T – the temperature in energy units,

$\beta = 1/T$ – the inverse temperature,

$g = 2s + 1$, s – the spin of a gas particle,

$\Lambda = (2\pi\hbar^2/mT)^{1/2}$ – the thermal wavelength of a particle,

$\Lambda_0 = (2\pi\hbar^2/mT_0)^{1/2}$ – the thermal wavelength at some characteristic temperature T_0 ,

μ – the chemical potential,

$f_k = (\exp \beta \xi_k + 1)^{-1}$ – the distribution function of Fermi particles in an ideal gas,

$\xi_k = \hbar^2 k^2 / 2m - \mu$,

$\Phi_s(t)$ – the special functions of order s (an integer or half-integer positive number), through which thermodynamic quantities of the ideal Fermi gas are expressed,

$t = \mu/T$ – the variable on which the functions $\Phi_s(t)$ depend – the adiabaticity parameter,

Ω – the thermodynamic potential,

N – the number of particles,

V – the volume,

$n = N/V$ – the particle number density,

p – the pressure,

E – the energy,

S – the entropy,

$\sigma = S/N$ – the entropy per one particle,

C_V, C_p – the heat capacities at constant volume and pressure,

$c_V = C_V/N, c_p = C_p/N$ – the heat capacities per one particle at constant volume and pressure,

u_T, u_σ – the isothermal and adiabatic speeds of sound,

$T_0 = \mu_0 = \varepsilon_F = \left(\frac{6\pi^2}{g}\right)^{2/3} \frac{\hbar^2 n^{2/3}}{2m}$, T_0 – the degeneracy temperature,

μ_0 – the chemical potential at zero temperature, ε_F – the Fermi energy,

$l = n^{-1/3}$ – the average distance between gas particles,

$q = (\Lambda/l)^3 = n\Lambda^3$ – the parameter, characterizing the degree of “quantumness” of the gas (quantumness),

$\theta \equiv nT/p$.

Some fundamental constants

The Planck constant (\hbar) – $1.055 \cdot 10^{-27}$ erg · s

The Boltzmann constant (k_B) – $1.381 \cdot 10^{-16}$ erg · K⁻¹

The electron mass (m_e) – $9.110 \cdot 10^{-28}$ g

The proton mass (m_p) – $1.673 \cdot 10^{-24}$ g

The mass of ³He atom (m_{3He}) – $5.01 \cdot 10^{-24}$ g

The electron charge (e) – $4.803 \cdot 10^{-10}$ cm^{3/2}g^{1/2}s⁻¹