Integral quantization based on the Heisenberg-Weyl group

Aleksandra Pędrak,^{1, [∗](#page-1-0)} Andrzej Góźdź,^{2,[†](#page-1-1)} Włodzimierz Piechocki,^{1,[‡](#page-1-2)} Patryk Mach,^{3, [§](#page-1-3)} and Adam Cieślik^{3, [¶](#page-1-4)}

¹Department of Fundamental Research, National Centre for Nuclear Research, Pasteura 7, 02-093 Warszawa, Poland ²Institute of Physics, Maria Curie-Skłodowska University, pl. Marii Curie-Skłodowskiej 1, 20-031 Lublin, Poland 3 Institute of Theoretical Physics, Jagiellonian University in Kraków, Łojasiewicza 11, 30-348 Kraków, Poland (Dated: November 1, 2024)

Abstract

We develop a relativistic framework of integral quantization applied to the motion of spinless particles in the four-dimensional Minkowski spacetime. The proposed scheme is based on coherent states generated by the action of the Heisenberg-Weyl group and has been motivated by the Hamiltonian description of the geodesic motion in General Relativity. We believe that this formulation should also allow for a generalization to the motion of test particles in curved spacetimes. A key element in our construction is the use of suitably defined positive operator-valued measures. We show that this approach can be used to quantize the one-dimensional nonrelativistic harmonic oscillator, recovering the standard Hamiltonian as obtained by the canonical quantization. Our formalism is then applied to the Hamiltonian associated with the motion of a relativistic particle in the Minkowski spacetime. A direct application of our model, including a computation of transition amplitudes between states characterized by fixed positions and momenta, is postponed to a forthcoming article.

CONTENTS

¶ adam.cieslik@doctoral.uj.edu.pl

I. INTRODUCTION

This work presents a generalization of the so-called integral quantization (IQ) method that is a special case of quantization based on deformations of quantum measures. The IQ has been used in quantizations of numerous physical systems. For a comprehensive review of applications we recommend references [\[1](#page-23-1)[–3\]](#page-23-2). We have already used the IQ approach based on the affine group. See, for instance, [\[4](#page-24-0)[–7\]](#page-24-1) for applications in astrophysics and cosmology $[8-11]$.

In this work we apply the IQ to the motion of a relativistic spinless particle in the Minkowski spacetime. We use the Hamiltonian description of the geodesic motion, known from General Relativity, and construct a scheme which, in principle, could be also applied to quantize the motion of a free particle in a curved spacetime. In the context of the motion in the Minkowski spacetime, this Hamiltonian approach suggests to construct the IQ procedure basing on the Heisenberg-Weyl group, as it allows for a one-to-one correspondence between the positions and momenta of the classical phase space and the group parameters.

The IQ method enables one to put forward the resolution of the time problem at the quantum level in a rather radical way. Traditionally, in quantum mechanics time is not considered to be a physical degree of freedom of the system, but rather a parameter. We propose that in the quantization of gravitational systems time should be treated on the same footing as space coordinates. The rationale for such dealing is that the distinction between time and space variables violates the general covariance of arbitrary transformations of temporal and spatial coordinates. The time should be considered to be a quantum observable. That is supported by series of experiments (see, e.g., [\[12,](#page-24-4) [13](#page-24-5)] and references therein). In fact, treating the time as an observable makes the quantization procedure more unique.

In this paper we describe the main elements of our model, focusing on formal details of the IQ. In particular, we pay special attention to a formulation in terms of the so-called positive operator valued measures (POVM), which leads to uniqueness in ascribing quantum operators to classical observables (see, e.g., $[14, 15]$ $[14, 15]$ and references therein). We introduce the eigenstates of the position and momentum operators and solve the eigenvalue problem of the quantum Hamiltonian associated with classical geodesics. As a test of the IQ based on the Heisenberg-Weyl group, we show that this method allows one to recover the results obtained by the canonical quantization for the standard nonrelativistic one-dimensional harmonic oscillator problem. Direct applications of our formalism to the motion of relativistic particles will be described in a forthcoming paper [\[16](#page-24-8)].

The paper is organized as follows: After specifying conventions concerning the Minkowski spacetime in Sec. II, we present the IQ based on the Heisenberg-Weyl group in Sec. III. The POVM approach is presented in Sec. IV with some applications. We conclude in Sec. V.

Throughout the paper we use geometric units with $c = 1$, where c denotes the speed of light. We assume the metric signature $(-, +, +, +)$.

II. PRELIMINARIES

Geodesic equations, describing the motion of free test particles in General Relativity, can be written in the Hamiltonian form

$$
\frac{dx^{\mu}}{d\tilde{s}} = \frac{\partial H}{\partial p_{\mu}}, \quad \frac{dp_{\mu}}{d\tilde{s}} = -\frac{\partial H}{\partial x^{\mu}}.
$$
\n(1)

Here x^{μ} , $\mu = 0, 1, 2, 3$, denote the coordinates along the geodesic. The corresponding covariant momentum components are defined as $p_{\mu} = g_{\mu\nu} p^{\nu} = g_{\mu\nu} dx^{\nu}/d\tilde{s}$. The Hamiltonian H can be written as

$$
H = \frac{1}{2}g^{\mu\nu}(x)p_{\mu}p_{\nu},\tag{2}
$$

where $g^{\mu\nu}$ denote the contravariant components of the metric tensor. The affine parameter \tilde{s} can be chosen in such a way that $g^{\mu\nu}p_{\mu}p_{\nu} = -m^2$, where m is the rest mass of the particle moving along the geodesic. Consequently $H = -\frac{1}{2}m^2$. For timelike geodesics, the proper time s is related to the affine parameter \tilde{s} by $\tilde{s} = s/m$.

We propose a semiclassical quantization scheme which, in principle, can be generalized to a quantization of the geodesic motion in curved spacetimes, described by the classical Hamiltonian [\(2\)](#page-3-1). In this paper we deal with the simplest, yet challenging, example of the motion in the flat Minkowski spacetime. In a sense, this renders the Hamiltonian description unnecessary (geodesics are simply straight lines in the Minkowski spacetime), but we adhere to this formalism to allow for future generalizations and as a guideline for the quantization procedure.

In the case of Minkowski spacetime the metric does not depend on x and is represented by the diagonal matrix $g = (-1, +1, +1, +1)$ so that we have (in Cartesian coordinates associated with an orthogonal frame) $g_{\mu\nu}x^{\mu}x^{\nu} = -x_0^2 + x_1^2 + x_2^2 + x_3^2$

and $g^{\mu\nu}p_{\mu}p_{\nu} = -p_0^2 + p_1^2 + p_2^2 + p_3^2$. The components of the four-momentum satisfy $-p_0 = p^0 = dx^0/d\tilde{s}, \ p_i = p^i = dx^i/d\tilde{s}, \ i = 1, 2, 3.$

The variables p_{μ} and x^{ν} are independent and they define the phase space $\{(p_{\mu},x^{\nu})\colon \mu,\nu=0,1,2,3\}\cong\mathbb{R}^{4}\times\mathbb{R}^{4}$, which can be identified with the cotangent bundle T^{\ast} *M* of the Minkowski spacetime (\mathcal{M}, g) .

III. INTEGRAL QUANTIZATION

The IQ procedure requires a specification of the group G that can be ascribed uniquely to the classical phase space of a given system. In this paper we choose the Heisenberg-Weyl group $\mathcal{HW}(4)$ to play that role, as this group can be identified with the cotangent bundle $T^*\mathcal{M}$ of the Minkowski spacetime. The IQ based on the Heisenberg-Weyl group has yet another advantage—the results obtained by this procedure remain, in many cases, consistent with the outcomes of the canonical quantization.

The group $\mathcal{HW}(4)$ has a unitary irreducible representation in the carrier Hilbert space $\mathcal{K} = L^2(\mathbb{R}^4, d^4 \xi)$, consisting of square integrable complex functions of four real variables ξ^{μ} , which enables us to construct the set of so-called coherent states in K .

A. The Heisenberg-Weyl group in four dimensions

Elements of the Heiseberg-Weyl group $\mathcal{HW}(4)$ are defined by 9 independent generators: four coordinate operators \hat{Q}^{μ} , four momentum operators \hat{P}_{μ} , and the unit operator 11ˆ, which satisfy the following commutation relations

$$
[\hat{Q}^{\mu}, \hat{P}_{\nu}] = i\hbar \delta^{\mu}_{\nu} \hat{\mathbb{1}} \,, \tag{3}
$$

$$
[\hat{Q}^{\mu}, \hat{\mathbf{1}}] = 0, \qquad (4)
$$

$$
[\hat{P}^{\mu}, \hat{\mathbb{1}}] = 0, \tag{5}
$$

$$
[\hat{Q}^{\mu}, \hat{Q}^{\nu}] = 0, \qquad (6)
$$

$$
[\hat{P}_{\mu}, \hat{P}_{\nu}] = 0, \qquad (7)
$$

where δ^{μ}_{ν} denotes the Kronecker delta, and $\mu, \nu = 0, 1, 2, 3$. The required realization of the above commutation relations in the space K can be defined by the following action of the operators \hat{Q}^{μ} and \hat{P}_{μ} :

$$
\hat{Q}^{\mu}\psi(\xi) = \xi^{\mu}\psi(\xi) , \qquad (8)
$$

$$
\hat{P}_{\mu}\psi(\xi) = -i\hbar \frac{\partial}{\partial \xi^{\mu}}\psi(\xi). \tag{9}
$$

Every element of the $H W(4)$ group can be written as the following unitary operator in ${\mathcal K}$

$$
g(\kappa; p, x) = g(\kappa; p_0, \dots, p_3, x^0, \dots, x^3) = \exp\left(i\kappa \hat{\mathbb{1}} + \frac{i}{\hbar}(p_\mu \hat{Q}^\mu - x^\mu \hat{P}_\mu)\right), \quad (10)
$$

where x^{μ} , p_{μ} , and κ denote group parameters. The multiplication law for the group reads^{[1](#page-5-0)}

$$
g(\kappa; p, x)g(\kappa'; p', x') = g\left(\kappa + \kappa' - \frac{1}{2\hbar}(p'_{\mu}x^{\mu} - p_{\mu}x'^{\mu}); \ p + p', \ x + x'\right). \tag{11}
$$

The Haar measure associated with the group $\mathcal{HW}(4)$ has the form $d\mu(\kappa, p, x) :=$ $d\kappa d\rho(p, x)$, where

$$
d\rho(p,x) = d^4p \, d^4x := dp_0 \, dp_1 dp_2 dp_3 \, dx^0 \, dx^1 dx^2 dx^3 \,. \tag{12}
$$

The action of unitary operators (10) in K is given by

$$
\hat{\mathcal{U}}(\kappa; p, x)\psi(\xi) = \exp(i\kappa)\exp\left(\frac{-ip_{\mu}x^{\mu}}{2\hbar}\right)\exp\left(\frac{ip_{\mu}\xi^{\mu}}{\hbar}\right)\psi(\xi - x), \quad (13)
$$

where we change the notation from $g(\kappa; p, x)$ to $\mathcal{U}(\kappa; p, x)$ to emphasise that in the latter case the action of operators \hat{Q}^{μ} and \hat{P}_{μ} is defined by Eqs. [\(8\)](#page-4-2) and [\(9\)](#page-4-2).

The subgroup parameterized by κ as

$$
g(\kappa; 0, 0) = \exp(i\kappa)\hat{\mathbb{1}},\tag{14}
$$

forms the unitary group $U(1)$, which is the center of the Heisenberg-Weyl group. This means that one can construct the homogeneous space $\mathcal{HW}(4)/\mathrm{U}(1) =: HW(4)$ to remove the redundant group parameter κ . According to the Stone-von Neumann theorem, any two unitary irreducible representations of the $H(W(4))$ group are equivalent, and the group parameter κ leads to the same states, in its action in K. The elements of the space $HW(4)$ are represented by the following unitary operators

$$
g(p,x) = \exp\left(\frac{i}{\hbar}(p_{\mu}\hat{Q}^{\mu} - x^{\mu}\hat{P}_{\mu})\right). \tag{15}
$$

 1 We use the identity

$$
\exp A \exp B = \exp \left(\frac{1}{2}[A, B]\right) \exp(A + B),
$$

which is valid if

$$
[A,[A,B]] = 0, \t [B,[A,B]] = 0.
$$

The multiplication law for operators [\(15\)](#page-5-2) reads

$$
g(p,x)g(\tilde{p},\tilde{x}) = \exp\left(-\frac{i}{2\hbar}(x^{\mu}\tilde{p}_{\mu} - p_{\mu}\tilde{x}^{\mu})\hat{I}\right)g(p+\tilde{p},x+\tilde{x}).
$$
 (16)

The unit operator can be identified with

$$
\hat{\mathbb{1}} = g(0,0). \tag{17}
$$

and the inverse operator reads

$$
g^{-1}(p,x) = g(-p, -x).
$$
 (18)

The unitary irreducible representation of the group $\mathcal{HW}(4)$ on the Hilbert space K is determined by the following action

$$
\hat{\mathcal{U}}(\kappa; p, x)\psi(\xi) = \exp(i\kappa)\,\hat{\mathcal{U}}(p, x)\psi(\xi)\,,\tag{19}
$$

where

$$
\hat{\mathcal{U}}(p,x)\psi(\xi) = \exp\left(\frac{-ip_{\mu}x^{\mu}}{2\hbar}\right)\exp\left(\frac{ip_{\mu}\xi^{\mu}}{\hbar}\right)\psi(\xi - x). \tag{20}
$$

B. Coherent states

The coherent states, $|p, x \rangle \in \mathcal{K} := L^2(\mathbb{R}^4, d^4 \xi)$, are constructed as follows

$$
|p,x\rangle = \hat{\mathcal{U}}(p,x)|\Phi_0\rangle, \quad \langle \xi|p,x\rangle = \hat{\mathcal{U}}(p,x)\langle \xi|\Phi_0\rangle = \hat{\mathcal{U}}(p,x)\Phi_0(\xi). \tag{21}
$$

where $\Phi_0(\xi) \colon \mathbb{R}^4 \to \mathbb{C}$ is the so-called fiducial vector and $|\Phi_0\rangle \in \mathcal{K}$ such that $\langle \Phi_0 | \Phi_0 \rangle = 1$. The freedom in the choice of the fiducial vector is a powerful feature of the IQ. In fact, the fiducial vector can be treated as a "parameter" of that quantization method.

In what follows we use the notation

$$
|\Phi_0\rangle = \hat{\mathcal{U}}(0,0)|\Phi_0\rangle = |0,0\rangle. \tag{22}
$$

Due to equation

$$
\hat{\mathcal{U}}^{-1}(p, x)\hat{\mathcal{U}}(\tilde{p}, \tilde{x})\psi(\xi) = \hat{\mathcal{U}}(-p, -x)\hat{\mathcal{U}}(\tilde{p}, \tilde{x})\psi(\xi)
$$
\n
$$
= \exp\left(-\frac{i}{2\hbar}(-x^{\mu}\tilde{p}_{\mu} + p_{\mu}\tilde{x}^{\mu})\right)\hat{\mathcal{U}}(\tilde{p} - p, \tilde{x} - x)\psi(\xi), \tag{23}
$$

we have

$$
\langle p, x | \tilde{p}, \tilde{x} \rangle = \langle \Phi_0 | \hat{\mathcal{U}}^{-1}(p, x) \hat{\mathcal{U}}(\tilde{p}, \tilde{x}) | \Phi_0 \rangle
$$

= $\exp \left(-\frac{i}{2\hbar} (-x^{\mu} \tilde{p}_{\mu} + p_{\mu} \tilde{x}^{\mu}) \right) \langle \Phi_0 | \hat{\mathcal{U}}(\tilde{p} - p, \tilde{x} - x) | \Phi_0 \rangle$
= $\exp \left(-\frac{i}{2\hbar} (-x^{\mu} \tilde{p}_{\mu} + p_{\mu} \tilde{x}^{\mu}) \right) \langle 0, 0 | \tilde{p} - p, \tilde{x} - x \rangle$. (24)

Making the group structure in the above formula explicit, we get

$$
\langle p, x | \tilde{p}, \tilde{x} \rangle = \langle \Phi_0 | \hat{\mathcal{U}} \left((p, x)^{-1} \circ (\tilde{p}, \tilde{x}) \right) | \Phi_0 \rangle = \langle 0, 0 | (p, x)^{-1} \circ (\tilde{p}, \tilde{x}) \rangle
$$

=
$$
\langle \Phi_0 | \hat{\mathcal{U}}^{-1} \left((\tilde{p}, \tilde{x})^{-1} \circ (p, x) \right) | \Phi_0 \rangle = \langle (\tilde{p}, \tilde{x})^{-1} \circ (p, x) | 0, 0 \rangle.
$$
 (25)

Using [\(25\)](#page-7-0) we obtain the following invariance property

$$
\langle (p', x') \circ (p, x) | (p', x') \circ (\tilde{p}, \tilde{x}) \rangle = \langle p, x | \tilde{p}, \tilde{x} \rangle.
$$
 (26)

Since the representation is irreducible, the operators $|p, x\rangle\langle p, x| : \mathcal{K} \to \mathcal{K}$ satisfy

$$
\frac{1}{A_{\Phi_0}} \int_{\mathbb{R}^8} d\rho(p, x) |p, x\rangle \langle p, x| = \hat{\mathbb{1}},
$$
\n(27)

where A_{Φ_0} is the normalization coefficient.

The Heisenberg-Weyl quantization consists in ascribing uniquely to each point of the phase space $T^{\ast} \mathcal{M}$ the projection operator

$$
\mathbb{R}^8 \ni (p, x) \longrightarrow |p, x\rangle\langle p, x| \,.
$$
 (28)

In the IQ Eq. (27) is used for mapping (quantization) of almost any classical observable $f: \mathbb{R}^8 \to \mathbb{R}$ onto an operator $\hat{f}: \mathcal{K} \to \mathcal{K}$ as follows

$$
f \longrightarrow \hat{f} := \frac{1}{A_{\Phi_0}} \int_{\mathbb{R}^8} d\rho(p, x) |p, x\rangle f(p, x) \langle p, x|.
$$
 (29)

Inserting the formulas

$$
\int_{\mathbb{R}^4} d^4 \xi \, |\xi\rangle\langle\xi| = \hat{\mathbb{1}}, \qquad \langle\xi''|\xi'\rangle = \delta^4(\xi'' - \xi'), \tag{30}
$$

where $\delta^4(\xi'' - \xi') = \delta(\xi''^0 - \xi'^0)\delta(\xi''^1 - \xi'^1)\delta(\xi''^2 - \xi'^2)\delta(\xi''^3 - \xi'^3)$, into [\(27\)](#page-7-1), and using (20) and (21) , as well as the expression

$$
\int_{\mathbb{R}} dp \exp\left(\frac{ipx}{\hbar}\right) = 2\pi\hbar \,\delta(x)\,,\tag{31}
$$

one can easily show that

$$
A_{\Phi_0} = (2\pi\hbar)^4. \tag{32}
$$

Thus, this coefficient does not depend on the choice of the fiducial vector $|\Phi_0\rangle \in \mathcal{K}$, which is not the case while applying the IQ based on other groups (see, e.g., $|4|$ -[\[11\]](#page-24-3) for more details).

Mapping [\(29\)](#page-7-2) leads to a symmetric operator which, in general, is not self-adjoint. In fact, a symmetric operator can have many self-adjoint extensions or none at all [\[17\]](#page-24-9). This feature makes the integral quantization non unique, which is undesirable. To solve this problem, we propose to use POVM type operators which make that mapping unique.

IV. POSITIVE OPERATOR VALUED MEASURES AND APPLICATIONS

First, let us define a general form of POVM operators. Let $\mathcal{L}(\mathcal{K})$ represent a set of bounded linear operators on the Hilbert space K. Let $\Omega \subset \mathbb{R}$ denote a set of allowed values of a quantum observable A and let F be the σ -algebra of subsets of Ω , with (Ω, \mathcal{F}) being a measurable space. The mapping $E \colon \mathcal{F} \to \mathcal{L}(\mathcal{K})$ is called the positive operator valued measure, POVM, if E fulfils the following conditions:

- $\forall_{X \in \mathcal{F}}, E(X)$ is positive semi-definite;
- if $\{X_k\}$ is a countable collection of disjoint sets, then

$$
E\left(\bigcup_{k} X_{k}\right) = \sum_{k} E(X_{k}),
$$

and this series converges in the weak topology;

• $E(\emptyset) = 0$ and $E(\Omega) = 1$.

The first condition ensures positivity of the quantum probability

$$
Prob(E(X); \hat{\gamma}) = Tr(E(X)\hat{\gamma}), \qquad (33)
$$

where $\hat{\gamma}$ denotes a density operator, i.e., the state of the quantum system under consideration. Prob $(E(X); \hat{\gamma})$ is interpreted as the probability that values of the observable E belong to the set X (see the so called "minimal interpretation of quantum mechanics" [\[14](#page-24-6)]). The second condition represents additivity of the measure for disjoint sets and the probability for mutually exclusive events. The last condition normalizes the probability to unity.

There is a well-known one-to-one correspondence between bounded sesquilinear forms and bounded linear operators defined on a Hilbert space [\[18](#page-24-10)], which may be understood as a precise formulation of the statement that any linear operator acting in a Hilbert space can be prescribed by its matrix elements. In fact, the assumption that the forms and operators should be bounded can be partially relaxed—see, for instance, Theorem 3.5.1 in [\[19](#page-24-11)]. In either case, sesquilinear forms can be used to study linear operators, but they can also be used to define quantum observables.

A sesquilinear form in a Hilbert space K is defined as a mappining $h: K \times K \to \mathbb{C}$, fulfilling the following conditions:

$$
\tilde{h}(\psi_2 + \psi'_2, \psi_1 + \psi'_1) = \tilde{h}(\psi_2, \psi_1) + \tilde{h}(\psi_2, \psi'_1) + \tilde{h}(\psi'_2, \psi_1) + \tilde{h}(\psi'_2, \psi'_1), \n\tilde{h}(\alpha\psi_2, \beta\psi_1) = \alpha^* \beta \tilde{h}(\psi_2, \psi_1),
$$
\n(34)

for all $\psi_1, \psi'_1, \psi_2, \psi'_2 \in \mathcal{K}$ and all $\alpha, \beta \in \mathbb{C}$. To construct the POVM operator corresponding to a classical observable $f(p, q)$, we define the following sesquilinear form

$$
\check{M}_f(U; \psi_2, \psi_1) := \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p, q) \langle \psi_2 | p, q \rangle \chi(f(p, x) \in U) \langle p, q | \psi_1 \rangle , \tag{35}
$$

where $\chi(S) = 1$, iff relation S is satisfied and $\chi(S) = 0$, otherwise. The condition $\chi(f(p,x) \in U) = 1$ restricts the part of the phase space transformed by the classical observable f to the set $U \subset \mathbb{R}$. Using the bra-ket notation, the corresponding operator can be written as

$$
\hat{M}_f(U) = \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p,q) \left| p, q \right\rangle \chi(f(p,x) \in U) \langle p, q \right|.
$$
 (36)

This operator is bounded and self-adjoint (see Appendix [A\)](#page-21-0). Operators of this type fulfil all conditions defining POVM operators, where U belongs to the σ -algebra \mathcal{F} .

We say that the full set of operators (36) represents the quantum observable f corresponding to a classical observable f. All required physical characteristics of the observable f can be obtained by means of operators constructed as functions of measures [\(36\)](#page-9-0). This includes the expectation value, the variance, and matrix elements of \ddot{f} .

In the traditional approach, an observable is described by a single operator. A synthesis of operator measures and values of a given observable is given on the basis of the spectral theorem [\[20](#page-24-12), [21\]](#page-24-13). Using this idea, in our case for non-orthogonal measures, one can write the approximate representation of the operator \hat{f} as

$$
\hat{f}(\epsilon; a, b) := \sum_{k} \bar{u}_k \,\hat{M}_f(Q_k) \,, \tag{37}
$$

where $(a < \cdots < u_k < u_{k+1} < u_{k+2} < \cdots \le b)$ is a partition of the interval $(a, b] \subset \mathbb{R}$, $Q_k = (u_k, u_{k+1}]$ and $\bar{u}_k \in Q_k$. $a < b$ determine the range and $\epsilon = \max_k |u_{k+1} - u_k|$ describes the resolution of this observable. In general, one needs to calculate the weak operator limit of self-adjoint operators $f(\epsilon; a, b)$ for $\epsilon \to 0$ and then take the limit $a \to -\infty$ and $b \to +\infty$. If this limit exists, it is represented by the integral

$$
\langle \psi_2 | \hat{f} | \psi_1 \rangle = \lim \langle \psi_2 | \hat{f}(\epsilon; a, b) | \psi_1 \rangle = \int_{\mathbb{R}} u d \langle \psi_2 | \hat{M}_f(u) | \psi_1 \rangle , \qquad (38)
$$

where

$$
\hat{M}_f(u) := \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p,q) \left| p, q \right\rangle \chi(f(p,q) \le u) \langle p, q \right|.
$$
 (39)

A link between measures and quantum physics is established by the fundamental formula

$$
\text{Prob}\left(M_f(U); \hat{\gamma}\right) = \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p, q) \,\chi(f(p, x) \in U) \text{Tr}(|p, q\rangle \langle p, q| \hat{\gamma}),\tag{40}
$$

which describes the probability that the observable \hat{f} has its value in the set U and the system is described by the density operator $\hat{\gamma}$.

Instead of dealing with symmetric sesquilinear forms, it is often more convenient to work with corresponding quadratic forms or, in the quantum context, the expectation values. We use the notation $\langle \hat{f}; \psi \rangle := \check{f}(\psi, \psi) = \langle \psi | \hat{f} | \psi \rangle$. The original sesquilinear form can be always recovered by the standard polarization identity [\[19](#page-24-11), [21\]](#page-24-13)

$$
\langle \psi_2 | \hat{f} | \psi_1 \rangle = \frac{1}{4} \left(\langle \hat{f}; \psi_1 + \psi_2 \rangle - \langle \hat{f}; \psi_1 - \psi_2 \rangle + i \langle \hat{f}; \psi_1 + i \psi_2 \rangle - i \langle \hat{f}; \psi_1 - i \psi_2 \rangle \right) \tag{41}
$$

In the following we consider only pure states $|\psi\rangle$, i.e., $\hat{\gamma} = |\psi\rangle\langle\psi|$. A generalization to mixed states is straighforward. Following definition [\(37\)](#page-9-1), the expectation value of the observable \hat{f} can be written as a limit of the following sum:

$$
\langle \hat{f}; \psi \rangle := \lim \langle \hat{f}(\epsilon; a, b); \psi \rangle = \lim \sum_{k} \bar{u}_{k} \langle \hat{M}_{f}(Q_{k}); \psi \rangle
$$

=
$$
\lim \sum_{k} \bar{u}_{k} [\text{Prob} (\hat{M}_{f}(u_{k+1}); \psi) - \text{Prob} (\hat{M}_{f}(u_{k}); \psi)]
$$

=
$$
\int_{\mathbb{R}} u d \langle \psi | \hat{M}_{f}(u) | \psi \rangle,
$$
 (42)

where $Q_k = (u_{k+1}, u_k]$ and $\bar{u}_k \in Q_k$. This limit means that the length of the largest subinterval $Q_k = (u_k, u_{k+1}]$ in the sum [\(42\)](#page-10-0) approaches zero, for every partition $(a < \cdots < u_k < u_{k+1} < u_{k+2} < \cdots \le b)$ of the interval $(a, b] \subset \mathbb{R}$. Subsequently, one needs to take limits $a \to -\infty$ and $b \to +\infty$.

A differentiation of the expectation value of operator (39) with respect to u gives the probability density that the observable \hat{f} has the value u:

$$
\frac{\partial}{\partial u} \langle \hat{M}_f(u); \psi \rangle = \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p, q) \, \delta(u - f(p, q)) |\langle p, q | \psi \rangle|^2, \tag{43}
$$

where the Dirac delta distribution is used as the derivative of the step function. Next, the condition $\bar{u}_k \in Q_k$ in [\(42\)](#page-10-0) implies that for every k there is a point $(p(k), q(k))$ for which $f(p(k), q(k)) = \bar{u}_k$, and formula [\(42\)](#page-10-0) can be rewritten as

$$
\langle \hat{f}; \psi \rangle := \lim \sum_{k} f(p(k), q(k)) \, |\langle p(k), q(k) | \psi \rangle|^2. \tag{44}
$$

If the above limit exists, the corresponding sesquilinear form can be written as

$$
\check{f}(\psi_2, \psi_1) = \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p, q) \langle \psi_2 | p, q \rangle f(p, q) \langle p, q | \psi_1 \rangle. \tag{45}
$$

Theorem 3.5.1 of $[19]$ allows one to associate with the sesquilinear form (45) a unique operator

$$
\hat{f} = \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p,q) \left| p, q \right\rangle f(p,q) \langle p, q \right|,\tag{46}
$$

defined on the domain $\mathcal{D} \subset \mathcal{K}$ consisting of all $|\psi_1\rangle$ for which there exists a state dependent finite constant $m(\psi_1) \geq 0$ such that for all $|\psi_2\rangle$ the following inequality is satisfied:

$$
|\check{f}(\psi_2, \psi_1)| \le m(\psi_1) || \psi_2 ||. \tag{47}
$$

This operator does not have to be bounded or self-adjoint, but it is symmetric in \mathcal{D} , defined by [\(47\)](#page-11-1). It can be shown that condition (47) is fulfilled if $f(p,q)\psi_1(p,q)$ is a square integrable function.

Finally, note that since symmetric sesquilinear forms and expectation values are related by the polarization identity (41) , a sesquilinar form associated with an expectation value (a quadratic form) is well defined, provided that the latter is also well defined (see Appendix [B\)](#page-22-0).

The above reasoning shows a limitation of the standard integral quantization formula [\(29\)](#page-7-2), which generally leads to a symmetric operator, which in turn may not always allow for a unique extension to a self-adjoint operator.

Matrix elements of any quantum observable correspond to special values of the associated sequilinear form. Every function of the quantum observable can be written in terms of these matrix elements. If a calculation based on operators [\(29\)](#page-7-2) fails, one can resort to POVM operators. Their matrix elements are well determined. In the following we use POVM operators to calculate appropriate matrix elements, even if we do not show this explicitly.

A. Elementary observables

The elementary observables are operators corresponding to group elements representing points of the configuration space. In practice one needs to construct operators corresponding to a given parametrization of the group $\mathcal{HW}(4)$.

In the canonical approach to quantization, the generators of the group $\mathcal{HW}(4)$, i.e., \hat{Q}^{μ} and \hat{P}_{μ} are considered to be momentum and position operators. In our approach to the $\mathcal{HW}(4)$ integral quantization the corresponding operators are defined by the appropriate measures [\(39\)](#page-10-1) as follows

$$
\hat{M}_{p_{\mu}}(u) := \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p, x) \left| p, x \right\rangle \chi(p_{\mu} \le u) \langle p, x \right|, \tag{48}
$$

$$
\hat{M}_{x^{\mu}}(v) := \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p, x) \left| p, x \right\rangle \chi(x^{\mu} \le v) \langle p, x \right|.
$$
\n(49)

According to [\(38\)](#page-10-3), the operators \hat{p}_{μ} and \hat{x}^{μ} are defined by the following matrix elements in the state space

$$
\langle e_n | \hat{p}_{\mu} | e_m \rangle := \int_{\mathbb{R}} u \, d \langle e_n | \hat{M}_{p_{\mu}} | e_m \rangle , \qquad (50)
$$

$$
\langle e_n|\hat{x}^{\mu}|e_m\rangle := \int_{\mathbb{R}} u \, d\langle e_n|\hat{M}_{x^{\mu}}|e_m\rangle ,\qquad(51)
$$

where vectors $|e_k\rangle$ form an arbitrary orthonormal basis in the state space.

It is interesting to compare generators of the group $\mathcal{HW}(4)$ and quantized operators of momenta and positions. For this purpose, it is easiest to compare corresponding matrix elements in a well chosen basis of the carrier space K .

In the case of momenta operators we choose the eigenbases of the generators \hat{P}_{μ} ,

$$
\langle \eta_{p''} | \hat{P}_{\mu} | \eta_{p'} \rangle = \delta^4 (p'' - p') p'_{\mu} . \tag{52}
$$

It is easy to show that

$$
\langle \xi | \eta_p \rangle =: \eta_p(\xi) = \left(\frac{1}{\sqrt{2\pi\hbar}}\right)^4 \exp\left(i\frac{p_\mu \xi^\mu}{\hbar}\right). \tag{53}
$$

In the case of position operators we can work in the eigenbases of the generators \hat{Q}^{μ} ,

$$
\langle \xi'' | \hat{Q}^{\mu} | \xi' \rangle = \delta^4 (\xi'' - \xi') \xi'^{\mu} . \tag{54}
$$

Straightforward calculations allow to obtain required matrix elements of the corresponding measures

$$
\langle \eta_{p''} | \hat{M}_{p\mu}(u) | \eta_{p'} \rangle = \delta^4(p'' - p') \int_{\mathbb{R}^4} d^4p \,\chi(p_\mu \le u) |\tilde{\Phi}_0(p' - p)|^2 \,, \tag{55}
$$

$$
\langle \xi'' | \hat{M}_{x^{\mu}}(u) | \xi' \rangle = \delta^4(\xi'' - \xi') \int_{\mathbb{R}^4} d^4x \, \chi(x^{\mu} \le u) |\Phi_0(\xi' - x)|^2 \,, \tag{56}
$$

where the Fourier transform of the fiducial vector is defined as

$$
\tilde{\Phi}_0(p) := \langle \eta_p | \Phi_0 \rangle = \frac{1}{(2\pi\hbar)^2} \int_{\mathbb{R}^4} d^4 \xi \exp\left(-i\frac{p_\mu \xi^\mu}{\hbar}\right) \Phi_0(\xi). \tag{57}
$$

To show (55) and (56) we use the expressions (see (21) and (20))

$$
\langle \xi | p, x \rangle = \exp\left(-i\frac{p_{\mu}x^{\mu}}{2\hbar}\right) \exp\left(i\frac{p_{\mu}\xi^{\mu}}{\hbar}\right) \Phi_{0}(\xi - x), \qquad (58)
$$

and

$$
\langle \eta_k | p, x \rangle = \int_{\mathbb{R}^4} d^4 \xi \langle \eta_k | \xi \rangle \langle \xi | p, x \rangle = \exp \left(i \frac{(p_\mu - 2k_\mu) x^\mu}{2\hbar} \right) \tilde{\Phi}_0(k - p), \tag{59}
$$

where in the last formula we use Eq. [\(53\)](#page-12-1). Finally, repeating the general calculation of Sec. IV, one obtains

$$
\langle \eta_{p''} | \hat{p}_{\mu} | \eta_{p'} \rangle = \delta^4(p'' - p') \left(p'_{\mu} - \int_{\mathbb{R}^4} d^4 p \, p_{\mu} | \tilde{\Phi}_0(p) |^2 \right) , \tag{60}
$$

$$
\langle \xi'' | \hat{x}^{\mu} | \xi' \rangle = \delta^4 (\xi'' - \xi') \left(\xi'^{\mu} - \int_{\mathbb{R}^4} d^4 \xi \, \xi^{\mu} | \Phi_0(\xi) |^2 \right) . \tag{61}
$$

In accordance with the condition [\(47\)](#page-11-1) and the subsequent commentary, the above operators are uniquely defined in the bases $|\eta_p\rangle$ and $|\xi\rangle$ if functions $p_\mu \tilde{\Phi}_0(p)$ and $\xi^{\mu}\Phi_0(\xi)$, respectively, are square integrable functions. We see that the generators of the group $\mathcal{HW}(4)$ and corresponding quantized momenta and positions coincide up to some constants. The constants depend only on the fiducial vector. Assuming additionally that the fiducial vector has good parity, i.e.,

$$
\Phi_0(-\xi) = \pm \Phi_0(\xi) \,, \tag{62}
$$

one can easily show that these constants are equal to zero. This means that the $HW(4)$ integral quantization method reproduces the canonical momentum and position operators.

B. Expectation values of the momentum and position operators

In our approach the coherent states are viewed as points of the quantum configuration space. This suggests a compatibility of their parametrization with expectation values of \hat{p}_{μ} and \hat{x}^{μ} operators, namely

$$
\langle \tilde{p}, \tilde{x} | \hat{p}_{\nu} | \tilde{p}, \tilde{x} \rangle = \tilde{p}_{\nu}, \qquad (63)
$$

$$
\langle \tilde{p}, \tilde{x} | \hat{x}^{\nu} | \tilde{p}, \tilde{x} \rangle = \tilde{x}^{\nu} . \tag{64}
$$

To fulfil the above compatibility conditions, one needs to choose an appropriate group parametrization and an appropriate fiducial vector. It turns out that in our case the good parity fiducial vector (62) allows to satisfy properties (63) . This is due to the fact that

$$
\langle \tilde{p}, \tilde{x} | \hat{p}_{\nu} | \tilde{p}, \tilde{x} \rangle = \tilde{p}_{\nu} + \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p, x) \, p_{\nu} \left| \int_{\mathbb{R}^4} d^4\xi \, \Phi_0^*(\xi) \exp\left(\frac{ip_{\mu}\xi^{\mu}}{\hbar}\right) \Phi_0(\xi - x) \right|^2 \tag{65}
$$

and

$$
\langle \tilde{p}, \tilde{x} | \hat{x}^{\nu} | \tilde{p}, \tilde{x} \rangle = \tilde{x}^{\nu} + \frac{1}{(2\pi\hbar)^{4}} \int_{\mathbb{R}^{8}} d\rho(p, x) x^{\nu} \left| \int_{\mathbb{R}^{4}} d^{4}\xi \Phi_{0}^{*}(\xi) \exp\left(\frac{ip_{\mu}\xi^{\mu}}{\hbar}\right) \Phi_{0}(\xi - x) \right|^{2}.
$$
\n(66)

Due to [\(62\)](#page-13-1) the r.h.s integrals are equal to zero.

To complete our derivation, one can check that the $H(W(4))$ generators have the same expectation values for the fiducial vectors satisfying [\(62\)](#page-13-1)

$$
\langle \tilde{p}, \tilde{x} | \hat{P}_{\nu} | \tilde{p}, \tilde{x} \rangle = \tilde{p}_{\nu} - i\hbar \int_{\mathbb{R}^4} d^4 \xi \, \Phi_0^*(\xi) \frac{\partial}{\partial \xi^{\nu}} \Phi_0(\xi) \tag{67}
$$

and

$$
\langle \tilde{p}, \tilde{x} | \hat{Q}^{\nu} | \tilde{p}, \tilde{x} \rangle = \tilde{x}^{\nu} + \int_{\mathbb{R}^4} d^4 \xi \, \xi^{\nu} \, |\Phi_0(\xi)|^2 \,. \tag{68}
$$

Again, the parity condition [\(62\)](#page-13-1) makes the r.h.s. integrals vanishing.

C. One dimensional harmonic oscillator

To compare the IQ based on the Heisenberg-Weyl group with the canonical quantization, we consider a quantization of a nonrelativistic one-dimensional harmonic oscillator. In this case the group $\mathcal{HW}(4)$ has to be replaced by $\mathcal{HW}(1)$. On the other hand, all formulas obtained in previous sections can be easily rewritten and applied in the present case.

The classical Hamiltonian of harmonic oscillations reads

$$
H(p,x) = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 x^2,
$$
\n(69)

where m represents the mass and ω is the frequency of the harmonic oscillator.

To quantize this Hamiltonian with the Heisenberg-Weyl group $H(W(1))$ we need two operators

$$
\widehat{p^2} := \int_{\mathbb{R}} u \, d\widehat{M}_{p^2}(u) \,, \tag{70}
$$

$$
\widehat{x}^2 := \int_{\mathbb{R}} v \, d\widehat{M}_{x^2}(v) \,, \tag{71}
$$

where the corresponding POVM operators are

$$
\hat{M}_{p^2}(u) = \frac{1}{2\pi\hbar} \int_{\mathbb{R}^2} dp \, dx \, |p, x\rangle \chi(p^2 \le u) \langle p, x| \,, \tag{72}
$$

$$
\hat{M}_{x^2}(u) = \frac{1}{2\pi\hbar} \int_{\mathbb{R}^2} dp \, dx \, |p, x\rangle \chi(x^2 \le u) \langle p, x| \,. \tag{73}
$$

In the following, we assume that the fiducial vector has good parity, i.e., $\Phi_0(-\xi)$ = $\pm\Phi(\xi)$. The matrix elements of operator [\(72\)](#page-15-0) within the "momentum" basis

$$
\langle \xi | \eta_p \rangle = \eta_p(\xi) := \frac{1}{\sqrt{2\pi\hbar}} \exp\left(\frac{ip\xi}{\hbar}\right) \tag{74}
$$

are

$$
\langle \eta_{p''} | \hat{M}_{p^2}(u) | \eta_{p'} \rangle = \delta(p'' - p') \int_{\mathbb{R}} dp \, \chi(p^2 \le u) | \tilde{\Phi}_0(p' - p) |^2.
$$
 (75)

Using these matrix elements and the decomposition of unity [\(89\)](#page-17-0), one gets required matrix elements for $\hat{M}_{p^2}(u)$ in the "position" basis

$$
\langle \xi'' | \hat{M}_{p^2}(u) | \xi' \rangle = \frac{1}{2\pi\hbar} \int_{\mathbb{R}} dp' \int_{\mathbb{R}} dp \exp\left(i\frac{p'(\xi'' - \xi')}{\hbar}\right) \chi\left((p' - p)^2 \le u\right) |\tilde{\Phi}_0(p)|^2.
$$
\n(76)

Similarly,

$$
\langle \xi'' | \hat{M}_{x^2}(v) | \xi' \rangle = \delta(\xi'' - \xi') \int_{\mathbb{R}} dx \, \chi(x^2 \le v) |\Phi_0(\xi' - x)|^2. \tag{77}
$$

Finally, the required matrix elements of operators [\(70\)](#page-15-1) are

$$
\langle \xi'' | \hat{p^2} | \xi' \rangle = -\hbar^2 \delta^{(2)} (\xi'' - \xi') + \delta(\xi'' - \xi') \int_{\mathbb{R}} dp \, p^2 | \tilde{\Phi}_0(p) |^2 \,, \tag{78}
$$

$$
\langle \xi''|\hat{x^2}|\xi'\rangle = \delta(\xi'' - \xi') \int_{\mathbb{R}} dx \, x^2 |\Phi_0(\xi' - x)|^2 , \qquad (79)
$$

where the parity of the fiducial vector and the formula for the second derivative of the delta distribution

$$
\frac{1}{2\pi\hbar} \int_{\mathbb{R}} dp \, p^2 \exp\left(i\frac{p(\xi'' - \xi')}{\hbar}\right) = -\hbar^2 \delta^{(2)}(\xi'' - \xi')\tag{80}
$$

were used.

Using these matrix element as the integral kernel of the quantized Hamiltonian, we get

$$
\int_{\mathbb{R}} d\xi' \langle \xi | \hat{H} | \xi' \rangle \psi(\xi') = \left\{ -\frac{\hbar^2}{2m} \frac{d^2}{d\xi^2} + \frac{1}{2} m \omega^2 \xi^2 \right\} \psi(\xi) + C \psi(\xi) ,\tag{81}
$$

where C is a constant dependent only on the fiducial vector

$$
C = \frac{1}{2m} \int_{\mathbb{R}} dp \, p^2 |\tilde{\Phi}_0(p)|^2 + \frac{1}{2} m \omega^2 \int_{\mathbb{R}} dx \, x^2 |\Phi_0(x)|^2.
$$
 (82)

Expression [\(81\)](#page-16-1) implies that the $H(W(1))$ quantization, that uses the POVM operators, reproduces the quantum harmonic oscillator Hamiltonian as obtained within the canonical quantization scheme:

$$
\hat{H} = -\frac{\hbar^2}{2m}\frac{d^2}{d\xi^2} + \frac{1}{2}m\omega^2\xi^2 + C.
$$
\n(83)

An open problem is related to the additional constant term C and its physical meaning. In principle, every classical Hamiltonian $H + C$ leads to the same Hamilton equations, independently of value of C . This suggests that such a constant term has no meaning. On the other hand, quantization of other observables can be sensitive to C, as it happens in the case of elementary observables p_{μ} and x^{μ} .

D. Quantum Hamiltonian of a test particle in the Minkowski spacetime

In what follows we examine the eigenvalue problem of the quantum Hamiltonian (2) describing the motion of a test particle with the rest mass m in the Minkowski spacetime. The classical Hamiltonian fulfils, due to [\(2\)](#page-3-1), the following equation

$$
H(p,x) = \frac{1}{2}g^{\mu\nu}p_{\mu}p_{\nu} = -\frac{1}{2}m^2.
$$
 (84)

The POVM operators of the quantum Hamiltonian \hat{H} have the form

$$
\hat{M}_H(u) = \left(\frac{1}{2\pi\hbar}\right)^4 \int_{\mathbb{R}^8} d\rho(p,x) \, |p,x\rangle \chi(H(p,x) \le u) \langle p,x| \,. \tag{85}
$$

The quantum Hamiltonian itself is determined by the following sesquilinear form

$$
\langle \psi_2 | \hat{H} | \psi_1 \rangle = \left(\frac{1}{2\pi\hbar}\right)^4 \int_{\mathbb{R}^8} d\rho(p, x) \langle \psi_2 | p, x \rangle H(p, x) \langle p, x | \psi_1 \rangle
$$

= $\frac{1}{2} g^{\mu\nu} \left(\frac{1}{2\pi\hbar}\right)^4 \int_{\mathbb{R}^8} d\rho(p, x) \langle \psi_2 | p, x \rangle \chi(p_\mu p_\nu \le u) \langle p, x | \psi_1 \rangle$. (86)

The last equality shows that the generalization to four dimensions of the harmonic oscillator matrix elements [\(78\)](#page-16-2) can be directly used in actual calculations.

In what follows we show that functions $\eta_p(\xi)$ defined in Eq. [\(53\)](#page-12-1) are generalized eigenstates of \hat{H} defined by [\(86\)](#page-17-1), if the coherent states $|p, x\rangle \in \mathcal{K}$ are generated from a suitably chosen fiducial vector $|\Phi_0\rangle \in \mathcal{K}$. For this purpose, we need to calculate matrix elements of the Hamiltonian [\(86\)](#page-17-1) within the states $\langle \eta_{k'} |$ and $|\eta_k \rangle$, i.e.,

$$
\langle \eta_{k'} | \hat{H} | \eta_k \rangle = \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d^4p \, d^4x \, \langle \eta_{k'} | p, x \rangle \frac{1}{2} \, g^{\mu\nu} p_\mu p_\nu \, \langle p, x | \eta_k \rangle \,. \tag{87}
$$

Inserting (59) into (87) we obtain

$$
\langle \eta_{k'}|\hat{H}|\eta_k\rangle = \frac{1}{2}\delta^4(k'-k)\int_{\mathbb{R}^4} d^4p\, g^{\alpha\beta}p_{\alpha}p_{\beta}\,\tilde{\Phi}_0^{\star}(k-p)\tilde{\Phi}_0(k'-p)\,. \tag{88}
$$

The key element of further procedure is to use the orthogonal decomposition of unity in the carrier space K in terms of the generalized states (53) , which reads

$$
\int_{\mathbb{R}^4} d^4p \, |\eta_p\rangle\langle\eta_p| = \hat{\mathbb{1}} \,. \tag{89}
$$

The validity of [\(89\)](#page-17-0) results from the theory of Fourier transforms in the context of distributions (see, e.g., [\[22\]](#page-25-0)) and is commonly used in quantum formalisms.

Using the matrix elements of the Hamiltonian [\(87\)](#page-17-2) and the decomposition of the unity [\(89\)](#page-17-0), we obtain the following:

$$
\hat{H}|\psi\rangle = \int_{\mathbb{R}^4} d^4k' |\eta_{k'}\rangle \langle \eta_{k'}|\hat{H}|\psi\rangle = \int_{\mathbb{R}^4} d^4k' |\eta_{k'}\rangle \int_{\mathbb{R}^4} d^4k'' \langle \eta_{k'}|\hat{H}|\eta_{k''}\rangle \langle \eta_{k''}|\psi\rangle
$$
\n
$$
= \int_{\mathbb{R}^4} d^4k' |\eta_{k'}\rangle \int_{\mathbb{R}^4} d^4k'' \frac{1}{2} \delta^4(k'-k'') \int_{\mathbb{R}^4} d^4p \, g^{\alpha\beta} p_{\alpha} p_{\beta} \, \tilde{\Phi}_0^{\star}(k''-p) \tilde{\Phi}_0(k'-p) \langle \eta_{k''}|\psi\rangle
$$
\n
$$
= \frac{1}{2} g^{\alpha\beta} \int_{\mathbb{R}^4} d^4k' |\eta_{k'}\rangle \int_{\mathbb{R}^4} d^4p \, p_{\alpha} p_{\beta} |\tilde{\Phi}_0(k'-p)|^2 \langle \eta_{k'}|\psi\rangle , \qquad (90)
$$

for any $|\psi\rangle \in \mathcal{K}$.

After the change of variables $p \to p + k'$, we get

$$
2\hat{H}|\psi\rangle = \left(\int_{\mathbb{R}^4} d^4p |\tilde{\Phi}_0(-p)|^2\right) \int_{\mathbb{R}^4} d^4k' g^{\alpha\beta} k'_{\alpha} k'_{\beta} \langle \eta_{k'}|\psi\rangle |\eta_{k'}\rangle + \left(\int_{\mathbb{R}^4} d^4p \, p_{\beta} |\tilde{\Phi}_0(-p)|^2\right) \int_{\mathbb{R}^4} d^4k' 2g^{\alpha\beta} k'_{\alpha} \langle \eta_{k'}|\psi\rangle |\eta_{k'}\rangle + \left(\int_{\mathbb{R}^4} d^4p \, g^{\alpha\beta} p_{\alpha} p_{\beta} |\tilde{\Phi}_0(-p)|^2\right) \int_{\mathbb{R}^4} d^4k' \langle \eta_{k'}|\psi\rangle |\eta_{k'}\rangle.
$$
 (91)

Now, assuming $|\psi\rangle = |\eta_k\rangle$, we obtain

$$
2\hat{H}|\eta_k\rangle = \tilde{\lambda}_k|\eta_k\rangle\,,\tag{92}
$$

where the generalized eigenvalues $\tilde{\lambda}_k$ are

$$
\tilde{\lambda}_k = \left(\int_{\mathbb{R}^4} d^4 p \, |\tilde{\Phi}_0(-p)|^2 \right) g^{\alpha \beta} k_\alpha k_\beta + \left(\int_{\mathbb{R}^4} d^4 p \, p_\beta |\tilde{\Phi}_0(-p)|^2 \right) 2g^{\alpha \beta} k_\alpha
$$
\n
$$
+ \int_{\mathbb{R}^4} d^4 p \, g^{\alpha \beta} p_\alpha p_\beta |\tilde{\Phi}_0(-p)|^2 \,. \tag{93}
$$

Quantization of the right hand side of [\(84\)](#page-16-3) gives the operator proportional to the unit operator, $-1/2 m^2 \hat{1}$. Taking into account that every function is an eigenfunction of the unit operator, one can identify the eigenvalues (93) with $-m^2$, so that we have

$$
\left(\int_{\mathbb{R}^4} d^4 p \, |\tilde{\Phi}_0(-p)|^2 \right) g^{\alpha\beta} k_\alpha k_\beta + \left(\int_{\mathbb{R}^4} d^4 p \, p_\beta |\tilde{\Phi}_0(-p)|^2 \right) 2g^{\alpha\beta} k_\alpha + \int_{\mathbb{R}^4} d^4 p \, g^{\alpha\beta} p_\alpha p_\beta |\tilde{\Phi}_0(-p)|^2 = -m^2.
$$
\n(94)

In what follows we try to simplify [\(94\)](#page-18-1). One can easily show that

$$
\int_{\mathbb{R}^4} d^4 p \, |\tilde{\Phi}_0(-p)|^2 = \int_{\mathbb{R}^4} d^4 \xi \, |\Phi_0(\xi)|^2 = 1 \,, \tag{95}
$$

as the fiducial vector Φ_0 is normalized.

If the fiducial vector is an even or odd function of each of its variables, it is not difficult to find that

$$
I_{\beta} := \int_{\mathbb{R}^4} d^4 p \, p_{\beta} |\tilde{\Phi}_0(p)|^2 = -I_{\beta}, \quad \beta = 0, 1, 2, 3,
$$
 (96)

so that $I_\beta = 0$.

Taking into account these two simplifications, we can write [\(94\)](#page-18-1) in the form

$$
-m^2 = g^{\alpha\beta}k_{\alpha}k_{\beta} + \int_{\mathbb{R}^4} d^4p \, g^{\alpha\beta}p_{\alpha}p_{\beta}|\tilde{\Phi}_0(p)|^2. \tag{97}
$$

Further simplifications are possible, depending on the choice of the fiducial vector in [\(97\)](#page-19-0). Assuming the fiducial vector as the vacuum state of the four dimensional harmonic oscillator, see App. [C,](#page-22-1)

$$
\Phi_0(\xi) = \prod_{\mu=0}^3 \left(\frac{\lambda_\mu}{\pi \hbar}\right)^{\frac{1}{4}} \exp\left(-\frac{\lambda_\mu (\xi^\mu)^2}{2\hbar}\right) \tag{98}
$$

with $\lambda_0, \lambda_1 = \lambda_2 = \lambda_3 > 0$ and $\lambda_0 = 3\lambda_3$, we obtain

$$
g^{\alpha\beta}k_{\alpha}k_{\beta} = -m^2.
$$
\n(99)

Therefore, a suitable choice of the fiducial vector leads to the result that the Hamiltonian eigenvalues satisfy the relationship quite similar to the relation among classical momenta. The quantum Hamiltonian H has a continuous spectrum, consisting of infinitely many eigenvalues $-\frac{1}{2}m^2$, each being infinitely many fold degenerate. In what follows we assume that the fiducial vector $|\Phi_0\rangle$ is chosen to be defined by [\(98\)](#page-19-1).

The assumed form of the fiducial vector has an additional advantage. In the context of the presented quantization method, coherent states are defined as representing the points of the quantum phase space of our physical system. In our model, coherent states are used to represent both four-momentum and four-position simultaneously. It is therefore important to find the smearing with which the positions and momenta are localised. This can be specified by the uncertainty principle, which states that the uncertainty in the position and momentum of a particle is related by the following equation [\[23](#page-25-1)]:

$$
\text{var}(\hat{p}_{\mu};|p,x\rangle)\text{var}(\hat{x}^{\nu};|p,x\rangle) \ge \frac{1}{4}|\langle p,x|[\hat{p}_{\mu},\hat{x}^{\nu}]|p,x\rangle|^2. \tag{100}
$$

For a real valued fiducial vector ($\Phi_0(\xi) \in \mathbb{R}$), the right hand side of inequality [\(100\)](#page-19-2) is equal to

$$
\frac{1}{4} |\langle p, x | [\hat{p}_{\mu}, \hat{x}^{\nu}] | p, x \rangle|^2 = \hbar^2 \left| \int_{\mathbb{R}^4} d^4 \xi \, \xi^{\nu} \Phi_0(\xi) \frac{d}{d\xi^{\mu}} \Phi_0(\xi) \right|^2. \tag{101}
$$

In case of the fiducial vector in form [\(98\)](#page-19-1), assuming $\lambda_0 = 3\lambda_3$, one gets

$$
\frac{1}{4} |\langle p, x | [\hat{p}_{\mu}, \hat{x}^{\nu}] | p, x \rangle|^2 = \begin{cases} 0 & \text{for } \nu \neq \mu, \\ \frac{\hbar^2}{4} & \text{for } \nu = \mu. \end{cases}
$$
 (102)

Therefore, the fiducial vector in the adopted form minimizes the uncertainty principle, independently of parameters λ_0 and λ_3 .

In Eq. (100) the variance, which describes a stochastic deviation from the expectation value of a quantum observable \hat{A} in the quantum state $|\Psi\rangle \in \mathcal{K}$, is defined as follows

$$
\text{var}(\hat{A}; |\Psi\rangle) := \langle (\hat{A} - \langle \hat{A}; \Psi \rangle)^2; \Psi \rangle = \langle \hat{A}^2; \Psi \rangle - \langle \hat{A}; \Psi \rangle^2, \tag{103}
$$

where $\langle \hat{B}; \Psi \rangle := \langle \Psi | \hat{B} | \Psi \rangle$.

V. SUMMARY

In this paper we use the Heisenberg-Weyl group to construct a space of coherent states. This group reproduces standard canonical commutation relations among positions and momenta. In addition, its natural parametrization is compatible with the Minkowski spacetime.

The carrier space of that representation is used as the Hilbert space of the considered quantum system. Since the representation is irreducible, there exists the decomposition of unity in the carrier space that can be used for mapping of almost any classical observable onto a symmetric operator in that Hilbert space.

The POVM approach is an extension of the standard approach with the formula [\(29\)](#page-7-2). It allows to overcome the problem that operators [\(29\)](#page-7-2) are usually only symmetric and may not have unique self-adjoint extensions. The idea is to construct first matrix elements of required POVM measure corresponding to a classical observable f. As the POVM operators are bounded and therefore self-adjoint, their matrix elements are defined on the entire state space K of the physical system. Using these matrix elements, one can construct required physical quantities, such as expectation values, variances, etc, directly from their definitions. Applying the POVM ideas, we show that our integral quantization reproduces the result obtained within the canonical quantization approach in the case of a commonly known harmonic oscillator.

The POVM approach renders the IQ method applicable to quantization of sophisticated gravitational systems. In particular, to the quantization of motion of test particles in curved spacetime [\[24\]](#page-25-2).

Appendix A: Selfadjoint measures

We show that every operator generated by the form (35) is bounded. To see this, let us consider the corresponding sesquilinear form

$$
\check{M}_f(U; \psi_2, \psi_1) = \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p, q) \langle \psi_2 | p, q \rangle \chi(f(p, x) \in U) \langle p, q | \psi_1 \rangle, \tag{A1}
$$

where ψ_1 and ψ_2 belong to the carrier space K. As $\chi(f(p,x) \in U) \leq 1$, this form can be bounded as follows:

$$
|\check{M}_f(\psi_2, \psi_1)| \le \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p, q) \left| \langle \psi_2 | p, q \rangle \langle p, q | \psi_1 \rangle \right|.
$$
 (A2)

Making use of the Hölder inequality

$$
\int_{\mathbb{R}^8} d\rho(p,q) \left| \langle \psi_2 | p, q \rangle \langle p, q | \psi_1 \rangle \right| \le \left(\int_{\mathbb{R}^8} d\rho(p,q) \left| \langle \psi_2 | p, q \rangle \right|^2 \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^8} d\rho(p,q) \left| \langle p, q | \psi_1 \rangle \right|^2 \right)^{\frac{1}{2}}, \tag{A3}
$$

we get

$$
|\check{M}_{f}(\psi_{2},\psi_{1})| \leq \frac{1}{A_{\phi}} \left(\int_{\mathbb{R}^{8}} d\rho(p,q) \left| \langle \psi_{2} | p, q \rangle \right|^{2} \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^{8}} d\rho(p,q) \left| \langle p, q | \psi_{1} \rangle \right|^{2} \right)^{\frac{1}{2}}.
$$
 (A4)

It can be shown directly, using the coherent states resolution of unity, that

$$
\langle \psi_2 | \psi_1 \rangle = \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p,q) \langle p,q | \psi_2 \rangle^* \langle p,q | \psi_2 \rangle < \infty \,. \tag{A5}
$$

This shows that every function $\langle p, q|\psi_k\rangle$, where $\psi_k \in \mathcal{K}$, belongs to the space of square integrable functions $L^2(\mathbb{R}^8, d\rho(p,q))$ and the right hand side of $(A4)$ is finite for every $\psi_1, \psi_2 \in \mathcal{K}$.

The inequality $(A4)$ is a required and sufficient condition that the operator generated by the sequilinear form (35) is bounded [\[18](#page-24-10)] (see also Theorem 3.5.2 of [\[19\]](#page-24-11)).

Since the sequilinear form (35) is symmetric, the generated operator is also symmetric, and because it is bounded, it is self-adjoint. In the bra-ket notation this operator can be shortly written as

$$
\hat{M}_f(U) = \frac{1}{(2\pi\hbar)^4} \int_{\mathbb{R}^8} d\rho(p,q) \, |p,q\rangle \chi(f(p,x) \in U) \langle p,q| \,. \tag{A6}
$$

Appendix B: Bounds of matrix elements

Matrix elements of the observable \hat{f} can be defined similarly to expectation value [\(38\)](#page-10-3). If the required limit does not exist, all matrix elements can be approximated by finite sums. In the case in which this limit leads to an integral form, matrix elements of the observable \hat{f} can be computed by the following sesquilinear form:

$$
\langle \psi_2 | \hat{f} | \psi_1 \rangle \equiv \check{f}(\psi_2, \psi_1) := \int_{\mathbb{R}} u \, d\langle \psi_2 | \hat{M}_f(u) | \psi_1 \rangle \, . \tag{B1}
$$

This form is bounded by the L[∞]-norm of the corresponding quadratic form $\check{f}(\psi,\psi)$, i.e., by the L^{∞} -norm of appropriate expectation values (see [\[19](#page-24-11)]):

$$
\|\langle \check{f} \rangle\|_{\infty} \le \sup_{\|\psi_1\| = \|\psi_2\| = 1} |\check{f}(\psi_2, \psi_1)| \le 2 \|\langle \check{f} \rangle\|_{\infty},
$$
 (B2)

where the L^{∞} -norm is defined as $\|\langle \check{f} \rangle\|_{\infty} := \sup_{\|\psi\|=1} \langle \check{f}, \psi \rangle$.

Appendix C: Choice of the fiducial vector

Let us try to choose Φ_0 in such a way that the integral

$$
J := \int_{\mathbb{R}^4} d^4 p \, g^{\alpha \beta} p_\alpha p_\beta |\tilde{\Phi}_0(p)|^2 \tag{C1}
$$

vanishes. For this purpose, we use the fiducial vector in the form of the fourdimensional harmonic oscillator ground state function

$$
\Phi_0(\xi) = \prod_{\mu=0}^3 \left(\frac{\lambda_\mu}{\pi \hbar}\right)^{\frac{1}{4}} \exp\left(-\frac{\lambda_\mu (\xi^\mu)^2}{2\hbar}\right) \tag{C2}
$$

with $\lambda_0, \lambda_1 = \lambda_2 = \lambda_3 > 0$, which is an even function of ξ^{μ} . Its Fourier transform reads

$$
\tilde{\Phi}_0(p) = \prod_{\mu=0}^3 (\pi \hbar \lambda_\mu)^{-1/4} \exp\left(-\frac{(p_\mu)^2}{2\hbar \lambda_\mu}\right) . \tag{C3}
$$

Therefore, the expression for J can be written as

$$
J = \int_{\mathbb{R}^4} dp_0 dp_1 dp_2 dp_3 \left(-p_0^2 + p_1^2 + p_2^2 + p_3^2 \right) (\pi \hbar \lambda_0)^{-1/2} \exp\left(-\frac{p_0^2}{\hbar \lambda_0} \right)
$$

$$
\times (\pi \hbar \lambda_1)^{-1/2} \exp\left(-\frac{p_1^2}{\hbar \lambda_1} \right) (\pi \hbar \lambda_2)^{-1/2} \exp\left(-\frac{p_2^2}{\hbar \lambda_2} \right)
$$

$$
\times (\pi \hbar \lambda_3)^{-1/2} \exp\left(-\frac{p_3^2}{\hbar \lambda_3} \right).
$$
 (C4)

For further discussion we need the following integrals

$$
K_{\mu} := \int_{\mathbb{R}} dp_{\mu} p_{\mu}^{2} \exp\left(-\frac{p_{\mu}^{2}}{\hbar \lambda_{\mu}}\right) = \frac{1}{2} \sqrt{\pi} (\hbar \lambda_{\mu})^{\frac{3}{2}}, \quad \mu = 0, 1, 2, 3 \quad (C5)
$$

and

$$
L_{\mu} := \int_{\mathbb{R}} dp_{\mu} \exp\left(-\frac{p_{\mu}^{2}}{\hbar \lambda_{\mu}}\right) = \sqrt{\pi \hbar \lambda_{\mu}}, \quad \mu = 0, 1, 2, 3. \tag{C6}
$$

Using the above integrals, we get

$$
J = \frac{1}{(\pi \hbar)^2 \sqrt{\lambda_0 \lambda_1 \lambda_2 \lambda_3}} (-K_0 L_1 L_2 L_3 + L_0 K_1 L_2 L_3 + L_0 L_1 K_2 L_3 + L_0 L_1 L_2 K_3), \quad (C7)
$$

and, as $\lambda_1 = \lambda_2 = \lambda_3$, we obtain

$$
J = \frac{1}{(\pi \hbar)^2 \sqrt{\lambda_0 \lambda_3^3}} (-K_0 L_3^3 + 3L_0 K_3 L_3^2) = \frac{1}{2} \hbar \left(-\lambda_0 + 3\lambda_3 \right) . \tag{C8}
$$

Thus, $J = 0$ if $\lambda_0 = 3\lambda_3$.

- [1] S. T. Ali, J.-P. Antoine, and J.-P. Gazeau, Coherent States, Wavelets and their Generalizations, (Theoretical and Mathematical Physics, Springer, New York, 2014), 2nd Edition. [I](#page-2-0)
- [2] J. R. Klauder, *Enhanced Quantization: Particles, Fields & Gravity* (World Scientfic, Singapore, 2015).
- [3] J.-P. Antoine, F. Bagarello, and J.-P. Gazeau, Coherent States and Their Applications: A Contemporary Panorama, Springer Proceedings in Physics, 2018. [I](#page-2-0)
- [4] A. Góźdź, M. Kisielowski, and W. Piechocki, "Quantum dynamics of gravitational mssive shell", Phys. Rev. D, 107, 046019 (20230. [I,](#page-2-0) [III B](#page-8-1)
- [5] W. Piechocki and T. Schmitz, "Quantum Oppenheimer-Snyder model", Phys. Rev. D, 102, 046004 (2020).
- [6] A. Góźdź, A. Pędrak, and W. Piechocki, "Ascribing quantum system to Schwarzschild spacetime with naked singularity", Class. Quantum Grav. 39, 145005 (2022).
- [7] A. Góźdź, A. Pędrak, and W. Piechocki, "Quantum system ascribed to the Oppenheimer-Snyder model of massive stars", Eur. Phys. J. C, 83:150 (2024). [I](#page-2-0)
- [8] A. Góźdź, W. Piechocki, and G. Plewa, "Quantum Belinski-Khalatnikov-Lifshitz scenario" Eur. Phys. J. C, 79:45 (2019). [I](#page-2-0)
- [9] A. Góźdź and W. Piechocki, "Robustnes of the BKL scenario", Eur. Phys. J. C, 80:142 (2020).
- [10] A. Góźdź, A. P¸edrak, and W. Piechocki, "Quantum dynamics corresponding to the chaotic BKL scenario", Eur. Phys. J. C, 83:150 (2023).
- [11] A. Góźdź, W. Piechocki, and T. Schmitz, "Dependence of the affine coherent states quantization on the parametrization of the affine group", Eur. Phys. J. Plus, 136:18 (2021). [I,](#page-2-0) [III B](#page-8-1)
- [12] A. Góźdź, M. Góźdź, and A. Pędrak, "Quantum time and quantum evolution", Universe, 9:256 (2023). [I](#page-2-0)
- [13] A. Góźdź and M. Góźdź, "Quantum clock in the projection evolution formalism", Universe, 10(3) (2024). [I](#page-2-0)
- [14] P. Busch, P. J. Lahti, and P. Mittelstaedt, The Quantum Theory of Measurement (Springer, 1996), 2-nd Edition. [I,](#page-2-0) [IV](#page-8-2)
- [15] J. P. Gazeau and B. Heller, "Positive-operator valued measure (povm) quantization", Axioms, 4:1–29 (215). [I](#page-2-0)
- [16] A. Cieślik, A. Góźdź, P. Mach, A. Pędrak, and W. Piechocki, "Semiclassical causal geodesics: I. Minkowski spacetime case", in progress. [I](#page-2-0)
- [17] M. Reed and B. Simon, Methods of Modern Mathematical Physics (San Diego, Academic Press, 1980). [III B](#page-8-1)
- [18] J. B. Conway, A Course in Functional Analysis, (Springer-Verlag, New York, 1990). [IV,](#page-8-2) [A](#page-21-2)
- [19] W. Mlak, Hilbert Spaces and Operator Theory, (PWN-Polish Scientific Publishers, Warszawa, Kluwer Academic Publishers, Dordrecht, 1991). [IV,](#page-8-2) [IV,](#page-10-4) [IV,](#page-11-0) [A,](#page-21-2) [B](#page-22-2)
- [20] F.W. Byron Jr. and R.W. Fuller, Mathematics of Classical and Quantum Mechanics, (Addison-Wesley, Don Mills, 1970). [IV](#page-9-0)
- [21] F. Riesz and B.Sz. Nagy, Functional analysis, translated from 2nd French edition by L.F. Baron, (Blackie & Son Limited, London and Glasgow, 1956). [IV,](#page-9-0) [IV](#page-10-4)
- [22] R. S. Strichartz, A Guide to Distribution Theory and Fourier Transforms (World Scientific Publishing, 2003). [IV D](#page-17-0)
- [23] H. P. Robertson, "The uncertainty principle", Phys. Rev., 34, 163 (1929). [IV D](#page-19-3)
- [24] A. Cieślik, A. Góźdź, P. Mach, A. Pędrak, and W. Piechocki, "Semiclassical causal geodesics: II. Schwarzschild spacetime case", in progress. [V](#page-20-0)