

FUNCTIONS OF SELF-ADJOINT OPERATORS UNDER RELATIVELY BOUNDED AND RELATIVELY TRACE CLASS PERTURBATIONS. RELATIVELY OPERATOR LIPSCHITZ FUNCTIONS

A.B. ALEKSANDROV AND V.V. PELLER

ABSTRACT. We study the behaviour of functions of self-adjoint operators under relatively bounded and relatively trace class perturbation. We introduce and study the class of relatively operator Lipschitz functions. An essential role is played by double operator integrals. We also consider study the class of resolvent Lipschitz functions. Then we obtain a trace formula in the case of relatively trace class perturbations and show that the maximal class of function for which the trace formula holds in the case of relatively trace class perturbations coincides with the class of relatively operator Lipschitz functions. Our methods also gives us a new approach to the inequality $\int |\xi(t)|(1+|t|)^{-1} dt < \infty$ for the spectral shift function ξ in the case of relatively trace class perturbations.

Contents

1.	Introduction	1
2.	Relatively bounded and relatively trace class perturbations	4
3.	The role of double operator integrals and Schur multipliers	7
4.	Schur multipliers originating from divided differences	10
5.	A representation of operator differences in terms of double operator integrals	13
6.	The necessity of the condition $\mathfrak{D}_I f \in \mathfrak{M}(\mathbb{R}^2)$	15
7.	Commutator estimates	17
8.	Differentiating in the strong operator topology	18
9.	Relatively trace class perturbations and the trace formula	21
	References	25

1. Introduction

In this paper we are going to study the behaviour of functions of self-adjoint operators under relatively bounded and relatively trace class perturbations.

Let us start, however, with the classical case of bounded and trace class perturbations. Let A be a (possibly unbounded) self-adjoint operator and let K be a bounded self-adjoint

The research on § 3-5 is supported by Russian Science Foundation [grant number 23-11-00153]. The research on the rest of the paper is supported by a grant of the Government of the Russian Federation for the state support of scientific research, carried out under the supervision of leading scientists, agreement 075-15-2024-631.

operator. The operator A can be interpreted as the initial (or unperturbed operator), K is considered as a perturbation while the operator $B = A + K$ is the perturbed operator.

For a function f continuous on the real line \mathbb{R} , the problem is to estimate the size (in one or another sense) of the operator $f(B) - f(A)$ in terms of the size of $B - A$. A function f is said to be an *operator Lipschitz function* if

$$\|f(B) - f(A)\| \leq \text{const} \|B - A\| \quad (1.1)$$

whenever $B - A$ is a bounded operator. Note that if inequality (1.1) holds for all bounded self-adjoint operator A and B , then it also holds for all not necessarily bounded operators.

The class of operator Lipschitz functions on \mathbb{R} will be denoted by $\text{OL}(\mathbb{R})$ or simply OL if it is clear that one deals with functions on \mathbb{R} .

Similarly, if instead of self-adjoint operators, we consider unitary operators, we can define the class of operator Lipschitz functions on the unit circle (see [AP1]).

We refer the reader to the survey article [AP1], which contains a lot of information about operator Lipschitz functions. Let us mention here that operator Lipschitz functions on the real line must be differentiable everywhere on \mathbb{R} and differentiable at infinity which means that the limit

$$\lim_{|t| \rightarrow \infty} \frac{f(t)}{t}$$

exists. However, operator Lipschitz functions do not have to be continuously differentiable, see [AP1].

It is well known (see [AP1]) that f is operator Lipschitz if and only if the following implication holds

$$B - A \in \mathbf{S}_1 \implies f(B) - f(A) \in \mathbf{S}_1, \quad (1.2)$$

where \mathbf{S}_1 is the trace class. We refer the reader to [GK] and [BS5] for the definition and basic properties of Schatten–von Neumann classes \mathbf{S}_p . Moreover (1.2) is equivalent to the fact that f is *trace class Lipschitz*, i.e.,

$$\|f(B) - f(A)\|_{\mathbf{S}_1} \leq \text{const} \|B - A\|_{\mathbf{S}_1},$$

whenever $B - A \in \mathbf{S}_1$.

In [L] physicist I.M. Lifshits when studying problems in quantum statistics and crystals theory arrived at the problem to evaluate the trace of the operator difference $f(B) - f(A)$ in the case when $B - A$ is a trace class perturbation of A . He discovered that there exists a real integrable function ξ on \mathbb{R} that is determined by A and B such that

$$\text{trace} (f(B) - f(A)) = \int_{\mathbb{R}} f'(t) \xi(t) dt \quad (1.3)$$

for sufficiently nice functions f . The function ξ is unique and is called the *spectral shift function* associated with A and B .

Later M.G. Krein has given in [Kr] a mathematically rigorous justification of the results of Lifshits and established that the spectral shift function ξ is integrable. In [Kr] a problem was posed to describe the maximal class of functions f , for which the above Lifshits–Krein trace formula holds for arbitrary self-adjoint operators A and B with trace class difference. The problem was solved in [Pe3]. It turned out that the maximal class of such functions coincides with the class of operator Lipschitz functions.

Another classical situation is when instead of measuring the size of the perturbation in terms of the differences $B - A$ we measure the size of the perturbation in terms of the difference of the resolvents $(B + iI)^{-1} - (A + iI)^{-1}$.

A function f on \mathbb{R} is called a *resolvent Lipschitz function* if

$$\|f(B) - f(A)\| \leq \text{const} \|(B + iI)^{-1} - (A + iI)^{-1}\|,$$

for arbitrary self-adjoint operators A and B .

We say that a self-adjoint operator K is a *resolvent trace class perturbation* of a self-adjoint operator A if

$$(A + K + iI)^{-1} - (A + iI)^{-1} \in \mathbf{S}_1.$$

It is also well known that a function f on \mathbb{R} is a resolvent Lipschitz function if and only if

$$f(B) - f(A) \in \mathbf{S}_1,$$

whenever $B - A$ is a resolvent trace class perturbation of A . Moreover, this is equivalent to the inequality

$$\|f(B) - f(A)\|_{\mathbf{S}_1} \leq \text{const} \|(B + iI)^{-1} - (A + iI)^{-1}\|_{\mathbf{S}_1}.$$

In the case of resolvent trace class perturbation there is an analog of the Lifshits–Krein trace formula. Indeed, suppose that K is a resolvent trace perturbation of A and $B = A + K$. Then there exists a real measurable function ξ that satisfies

$$\int_{\mathbb{R}} \frac{|\xi(t)|}{1 + t^2} dt < \infty$$

and such that trace formula (1.3) holds for sufficiently nice functions f . Note that unlike the case of trace class perturbations a spectral shift function ξ in the case of resolvent trace class perturbations is not unique. It is determined by A and B modulo an additive real constant.

The above facts are well known and can be obtained by applying the Cayley transform and reducing the situation to the case of unitary operator, see e.g., [MNP].

In this paper we mostly concentrate on the case of relatively bounded and relatively trace class perturbations; these notions will be introduced in § 2. We also introduce and study in § 2 the class of relatively operator Lipschitz functions, see § 2 for the definition

We obtain in § 4 various descriptions of the class of relatively operator Lipschitz functions. We also find in § 4 various new descriptions of the class of resolvent Lipschitz functions.

An important role will be played by double operator integrals. We give a brief introduction to double operator integrals in § 3 and introduce the notion of Schur multipliers with respect to spectral measures. We also discuss in § 3 how one can express the operator difference $f(B) - f(A)$ in terms of double operator integrals for various classes of operators and functions.

In Section 5 we obtain a representation of the operator difference $f(B) - f(A)$ in the case of a relatively bounded perturbation in terms of a double operator integral.

Then in § 2.7 we show that a function f on \mathbb{R} is relatively operator Lipschitz if and only if the function

$$(x, y) \mapsto \frac{f(x) - f(y)}{x - y}(x + i)$$

on \mathbb{R}^2 is a Schur multiplier with respect to arbitrary spectral measures.

Next, in Section § 7 we show that a function is relatively operator Lipschitz if and only if certain commutator estimates hold.

In § 8 we obtain a formula for the derivative of the function

$$t \mapsto f(A + t(B - A))$$

in the strong operator topology. This formula will be used in § 9 to obtain a trace formula in the case of relatively trace class perturbations. Our methods allows us to prove the inequality

$$\int_{\mathbb{R}} \frac{|\xi(t)|}{1 + |t|} dt < \infty.$$

for the spectral shift function ξ for the pair $\{A, B\}$, where $B - A$ is a relatively trace class perturbation of A . This inequality was obtained earlier in [CS]. However, our trace formula is valid for a considerably broader class of functions compared to the results of [CS]. Moreover, we describe the maximal class of functions, for which the trace formula holds in the case of relatively trace class perturbations and show that this class coincides with the class of relatively operator Lipschitz functions.

2. Relatively bounded and relatively trace class perturbations

Let us proceed now to relatively bounded perturbations and relatively trace class perturbations.

Let A be a not necessarily bounded self-adjoint operator. Suppose that K is a closed operator such that $\text{Dom}(A) \subset \text{Dom}(K)$, where the notation $\text{Dom}(T)$ stands for the domain of an operator T . It is well known (see [BS5], Theorem 1, Ch. 3, Sec. 4) that in this case K must be *dominated by* A , i.e., the following inequality must hold:

$$\|Kv\| \leq \text{const}(\|v\| + \|Av\|), \quad v \in \text{Dom}(A), \quad (2.1)$$

where $\text{Dom}(L)$ stands for the domain of an operator L .

If, in addition to this, K is a self-adjoint operator, we say that K is a *relatively bounded self-adjoint perturbation of* A .

It is easy to see that property (2.1) is equivalent to the fact that the operator

$$C \stackrel{\text{def}}{=} K(A + iI)^{-1} \quad (2.2)$$

is bounded which, in turn, is equivalent to the fact that the operator

$$G \stackrel{\text{def}}{=} K(A^2 + I)^{-1/2} \quad (2.3)$$

is bounded.

We say that a self-adjoint operator K is called a *relatively trace class perturbation* of A the operator C defined by (2.2) belongs to \mathbf{S}_1 which is, clearly, equivalent to the fact that the operator G defined by (2.3) belongs to \mathbf{S}_1 .

It is well known and easy to verify that if K is a relatively trace class perturbation of A , then it is a resolvent trace class perturbation of A . The converse is not true which can easily be seen from Theorems 4.1 and 4.2.

We are going to make three observations. All three are well known facts.

Lemma 2.1. *Suppose that K is a self-adjoint operator satisfying the inequality*

$$\|Kv\| \leq c\|v\| + d\|Av\|, \quad v \in \text{Dom}(A), \quad \text{for some } c > 0 \quad \text{and } d \in (0, 1) \quad (2.4)$$

(such operators are called strictly dominated by A). Then K is also dominated by $A + K$.

Proof. Indeed, let $v \in \text{Dom}(A)$. We have

$$\|Kv\| \leq c\|v\| + d\|Av\| = c\|v\| + d\|(A + K)v - Kv\| \leq c\|v\| + d\|(A + K)v\| + d\|Kv\|.$$

It follows that

$$\|Kv\| \leq (1 - d)^{-1}(c\|v\| + d\|(A + K)v\|), \quad (2.5)$$

and so K is dominated by $A + K$. ■

Lemma 2.2. *Suppose that K is a self-adjoint operator that is a relatively compact perturbation of A , i.e., the operator $K(A + iI)^{-1}$ compact. Then for any positive number d , there exists a positive number c such that*

$$\|Kv\| \leq c\|v\| + d\|Av\|, \quad v \in \text{Dom}(A),$$

and so, K is strictly dominated by $A + K$.

Proof. Let $\varepsilon > 0$. By the assumption, the operator $K(A + iI)^{-1}$ is compact, and so it can be represented in the form $K(A + iI)^{-1} = T + R$, where T is a finite rank operator and $\|R\| < \varepsilon$.

Consider the operators K_1 and K_2 defined on $\text{Dom}(A)$ by

$$K_1 = T(A + iI) \quad \text{and} \quad K_2 = R(A + iI).$$

Clearly,

$$\|K_2v\| = \|R(A + iI)v\| \leq \varepsilon\|(A + iI)v\| \leq \varepsilon\|Av\| + \varepsilon\|v\|, \quad v \in \text{Dom}(A),$$

and so, it suffices to show that for any positive number d , there exists a positive number c such that

$$\|K_1v\| \leq c\|v\| + d\|Av\|, \quad v \in \text{Dom}(A).$$

Since T is a linear combination of rank one operators, it suffices to consider the case when $\text{rank } T = \text{rank}(K_1(A + iI)^{-1}) = 1$. Suppose that

$$K_1v = ((A + iI)v, \varphi)\psi, \quad v \in \text{Dom}(A).$$

since A is densely defined, φ admit a representation $\varphi = \varphi_1 + \varphi_2$, where $\varphi_1 \in \text{Dom}(A)$ and $\|\varphi_2\| < \varepsilon\|\psi\|^{-1}$. We have

$$\begin{aligned} Kv &= ((A + iI)v, \varphi_1)\psi + ((A + iI)v, \varphi_2)\psi \\ &= (v, (A + iI)\varphi_1)\psi + ((A + iI)v, \varphi_2)\psi, \quad v \in \text{Dom}(A), \end{aligned}$$

and so

$$\begin{aligned} \|Kv\| &\leq \|(A + iI)\varphi_1\| \cdot \|\psi\| \cdot \|v\| + \|\varphi_2\| \cdot \|\psi\| \cdot \|v\| + \|\varphi_2\| \cdot \|\psi\| \cdot \|Av\| \\ &\leq (\|(A + iI)\varphi_1\| \cdot \|\psi\| + \|\varphi_2\| \cdot \|\psi\|)\|v\| + \varepsilon\|Av\|, \quad v \in \text{Dom}(A). \quad \blacksquare \end{aligned}$$

Lemma 2.3. *Suppose that a self-adjoint operator K is a relatively trace class perturbation of a self-adjoint operator A . Then K is a relatively trace class perturbation of $A + K$.*

Proof. It is easy to see that

$$K(A + K + iI)^{-1} - K(A + iI)^{-1} = -K(A + K + iI)^{-1}K(A + iI)^{-1} \in \mathbf{S}_1.$$

Indeed, by the assumption, $K(A + iI)^{-1} \in \mathbf{S}_1$. On the other hand, by Lemmata 2.1 and 2.2, the operator $K(A + K + iI)^{-1}$ is bounded. \blacksquare

Definition. A continuous function f on \mathbb{R} is called *relatively operator Lipschitz* if there is a positive number k such that

$$\|f(B) - f(A)\| \leq k\|(B - A)(A + iI)^{-1}\| \quad (2.6)$$

whenever A and B are self-adjoint operators such that $B - A$ is a relatively bounded perturbation of A . We denote by ROL the class of relatively operator Lipschitz function on \mathbb{R} .

Remark. Obviously, if f is a relatively operator Lipschitz function, then f is an operator Lipschitz function, and so, f is differentiable on \mathbb{R} . Indeed, (2.6) implies that

$$\|f(B) - f(A)\| \leq k\|(B - A)(A + iI)^{-1}\| \leq k\|A - B\| \cdot \|(A + iI)^{-1}\| \leq k\|A - B\|.$$

The following theorem is an elementary necessary condition for a function on \mathbb{R} to be relatively operator Lipschitz.

Theorem 2.4. *Let f be a relatively operator Lipschitz function on \mathbb{R} . Then the following equivalent condition holds:*

(a)

$$|f(s) - f(t)| \leq \text{const} |s - t| \cdot (1 + |t|)^{-1}, \quad s, t \in \mathbb{R}; \quad (2.7)$$

(b)

$$|f(s) - f(t)| \leq \text{const} \frac{|s - t|}{1 + |s| + |t|}, \quad s, t \in \mathbb{R};$$

(c) *the function*

$$s \mapsto (s + i)f(s), \quad s \in \mathbb{R},$$

is a Lipschitz function on \mathbb{R} .

Proof. Let us first show that (a) is a necessary condition. Indeed, it is clear that (2.7) exactly means that inequality (2.6) holds for all rank one self-adjoint operators A and B .

The implication (b) \implies (a) is trivial. To prove the implication (a) \implies (b), we observe that (a) implies that $|f(s) - f(t)| \leq \text{const} |s - t|(1 + |t|)^{-1}$ for $s, t \in \mathbb{R}$, and $\min((1 + |s|)^{-1}, (1 + |t|)^{-1}) \leq 2(1 + |s| + |t|)^{-1}$.

It remains to prove that (a) \iff (c). Substituting $t = 0$ we obtain that each of statements (a) and (c) implies that f is bounded.

First we prove that (a) \implies (c). We have

$$\begin{aligned} |(s + i)f(s) - (t + i)f(t)| &= |(s + i)(f(s) - f(t)) + f(t)(s - t)| \\ &\leq |(s + i)(f(s) - f(t))| + |f(t)(s - t)| \\ &\leq \text{const} \frac{|s + i|}{1 + |s|} |s - t| + \sup_{u \in \mathbb{R}} |f(u)| \cdot |s - t|. \end{aligned}$$

Similarly, we can prove that (c) \implies (a):

$$\begin{aligned} |(s + i)(f(s) - f(t))| &= |(s + i)f(s) - (t + i)f(t) - f(t)(s - t)| \\ &\leq |(s + i)f(s) - (t + i)f(t)| + |f(t)(s - t)| \\ &\leq \text{const} |s - t| + \sup_{u \in \mathbb{R}} |f(u)| \cdot |s - t|. \end{aligned}$$

It remains to observe that $|s + i|^{-1} \leq \frac{\sqrt{2}}{1 + |x|}$. ■

In § 6 we observe that condition (2.7) is not sufficient for a function to be relatively operator Lipschitz.

The following fact can be proved in the same way as the corresponding result for operator Lipschitz function, see Theorem 3.6.5 in [AP1].

Theorem 2.5. *Let f be a function on \mathbb{R} . The following statements are equivalent:*

- (a) $f \in \text{ROL}$;
- (b) if K is a self-adjoint relatively trace class perturbation of a self-adjoint operator A , then $f(A + K) - f(A) \in \mathbf{S}_1$;
- (c) f is a relatively trace class Lipschitz function, i.e.,

$$\|f(A + K) - f(A)\|_{\mathbf{S}_1} \leq \text{const} \|K(A + iI)^{-1}\|_{\mathbf{S}_1}$$

for an arbitrary relatively trace class perturbation K of a self-adjoint operator A .

3. The role of double operator integrals and Schur multipliers

Double operator integrals are expressions of the form

$$\iint \Phi(x, y) dE_1(x) Q dE_2(y). \tag{3.1}$$

Here E_1 and E_2 are spectral measures on Hilbert space, Φ is a bounded measurable function and Q is a bounded linear operator on Hilbert space.

Double operator integrals play a very important role in perturbation theory. They appeared first in the paper by Yu.L. Daletskii and S.G. Krein [DK]. Later M.S. Birman and M.Z. Solomyak created in [BS1, BS2] and [BS4] a beautiful rigorous theory of double operator integrals.

The starting point of the Birman–Solomyak theory is the case when Q belongs to the class \mathbf{S}_2 of Hilbert–Schmidt operators. Consider the set function \mathcal{E} that takes values in the set of orthogonal projections on the Hilbert Schmidt class \mathbf{S}_2 and is defined on the measurable rectangles by

$$\mathcal{E}(\mathbf{L})T \stackrel{\text{def}}{=} E_1(\mathbf{L})TE_2(\Delta), \quad T \in \mathbf{S}_2.$$

Birman and Solomyak showed in [BS6] that \mathcal{E} extends to a spectral measure on \mathbf{S}_2 . This allows one to define the double integral (3.1) in the case when $Q \in \mathbf{S}_2$ by

$$\iint \Phi dE_1 Q dE_2 \stackrel{\text{def}}{=} \left(\int \Phi d\mathcal{E} \right) Q.$$

Next, Φ is said to be a *Schur multiplier with respect to E_1 and E_2* if

$$Q \in \mathbf{S}_1 \implies \iint \Phi dE_1 Q dE_2 \in \mathbf{S}_1.$$

We use the notation \mathfrak{M}_{E_1, E_2} for the class of Schur multipliers with respect to E_1 and E_2 . The norm $\|\Phi\|_{\mathfrak{M}_{E_1, E_2}}$ in the space of Schur multipliers is defined as the norm of the transformer

$$Q \mapsto \iint \Phi dE_1 Q dE_2 \tag{3.2}$$

on \mathbf{S}_1 .

In the case when $\Phi \in \mathfrak{M}_{E_1, E_2}$, one can define by duality the double operator integral (3.1) for arbitrary bounded operator Q . Moreover, the norm of the transformer (3.2) on the space of bounded linear operators also coincides with $\|\Phi\|_{\mathfrak{M}_{E_1, E_2}}$.

There are several characterizations of the class \mathfrak{M}_{E_1, E_2} , see, e.g., [Pe1] and [AP3]. In particular, $\Phi \in \mathfrak{M}_{E_1, E_2}$ if and only if Φ belongs to the *Haagerup tensor product* $L_{E_1}^\infty \otimes_{\text{h}} L_{E_2}^\infty$ of L^∞ spaces $L_{E_1}^\infty$ and $L_{E_2}^\infty$, i.e., Φ admits a representation

$$\Phi(x, y) = \sum_n \varphi_n(x) \psi_n(y), \tag{3.3}$$

where $\varphi_n \in L_{E_1}^\infty$, $\psi_n \in L_{E_2}^\infty$, and

$$\{\varphi_n\}_{n \geq 0} \in L_{E_1}^\infty(\ell^2) \quad \text{and} \quad \{\psi_n\}_{n \geq 0} \in L_{E_2}^\infty(\ell^2).$$

By the *norm of Φ in $L^\infty(E_1) \otimes_{\text{h}} L^\infty(E_2)$* we mean the infimum of

$$\left\| \{\varphi_n\}_{n \geq 0} \right\|_{L_{E_1}^\infty(\ell^2)} \left\| \{\psi_n\}_{n \geq 0} \right\|_{L_{E_2}^\infty(\ell^2)} \tag{3.4}$$

over all representations in the form (3.3). Moreover, it was established recently in [AP3] that this description is *isometric*, i.e., the norm of Φ in \mathfrak{M}_{E_1, E_2} coincides with the norm of Φ in $L_{E_1}^\infty \otimes_{\text{h}} L_{E_2}^\infty$.

In the case when $\Phi \in L_{E_1}^\infty \otimes_{\text{h}} L_{E_2}^\infty$, the double operator integral (3.1) can be computed as follows:

$$\iint \Phi \, dE_1 Q \, dE_2 = \sum_n \left(\int \varphi_n \, dE_1 \right) Q \left(\int \psi_n \, dE_2 \right)$$

and the integral on the right converges in the weak operator topology.

We refer the reader to the survey article [AP1] for more detailed information on double operator integrals and Schur multipliers.

It turns out that a function f on \mathbb{R} is operator Lipschitz if and only if it is differentiable on \mathbb{R} and the divided difference $\mathfrak{D}f$ defined by

$$(\mathfrak{D}f)(x, y) \stackrel{\text{def}}{=} \begin{cases} \frac{f(x) - f(y)}{x - y}, & \text{if } x \neq y, \\ f'(x), & \text{if } x = y, \end{cases} \quad (3.5)$$

is a Schur multiplier with respect to arbitrary Borel spectral measures.¹ Moreover, for operator Lipschitz functions f the following representation holds for arbitrary self-adjoint operator A and B with bounded $B - A$:

$$f(B) - f(A) = \iint_{\mathbb{R} \times \mathbb{R}} (\mathfrak{D}f)(x, y) \, dE_B(x) (B - A) \, dE_A(y), \quad (3.6)$$

where E_A and E_B are the spectral measures of A and B .

We refer the reader to [AP1] for more detailed information.

The class of resolvent Lipschitz functions also admits a characterization in terms of Schur multipliers: a differentiable function f on \mathbb{R} is a resolvent Lipschitz function if and only if the function

$$(x, y) \mapsto (\mathfrak{D}f)(x, y)(x + i)(y + i)$$

on \mathbb{R}^2 is a Schur multiplier with respect to arbitrary Borel spectral measures. The following formula holds

$$f(B) - f(A) = \iint (\mathfrak{D}f)(x, y)(x + i)(y + i) \, dE_B(x) ((A + iI)^{-1} - (B + iI)^{-1}) \, dE_A(y)$$

for resolvent Lipschitz functions f . This can easily be reduced to an analogue of formula (3.6) for functions of unitary operators by passing to Cayley transform.

It is also well known (see e.g., [MNP]) that a function f on \mathbb{R} is resolvent Lipschitz if and only if it is the function φ on $\mathbb{T} \setminus \{1\}$ defined by

$$\varphi(\zeta) = f \left(i \frac{1 + \zeta}{1 - \zeta} \right), \quad (3.7)$$

then φ extends to an operator Lipschitz function on \mathbb{T} . This can also be proved by passing to Cayley transform.

¹We use the notation $\mathfrak{M}(\mathbb{R}^2)$ for the class of functions on \mathbb{R}^2 that are Schur multiplier with respect to arbitrary Borel spectral measure.

Suppose now that A and B are self-adjoint operators and $B - A$ is a resolvent trace class perturbation of A . Then the trace formula holds

$$\text{trace}(f(B) - f(A)) = \int_{\mathbb{R}} f'(t)\xi(t)$$

for an arbitrary resolvent Lipschitz function f , where ξ is a *spectral shift function* that corresponds to A and B . It is real valued, unique modulo a constant additive and satisfies the condition

$$\int_{\mathbb{R}} \frac{|\xi(t)|}{1+t^2} dt < \infty.$$

Moreover, the class of resolvent Lipschitz functions is the maximal class of functions, for which trace formula (3.6) holds as soon as $B - A$ is a resolvent trace class perturbation of A .

This again can be obtained by passing to the Cayley transforms of A and B . We refer the reader to [MNP] for details.

In this paper we characterize the class of relatively operator Lipschitz functions as the class of function f on \mathbb{R} , for which the function

$$(x, y) \mapsto (\mathfrak{D}f)(x, y)(x + i)$$

on \mathbb{R}^2 is a Schur multiplier with respect to arbitrary Borel spectral measures. In the case when $B - A$ is a relatively bounded perturbation of A and f is relatively operator Lipschitz, we represent $f(B) - f(A)$ in terms of the double operator integral

$$f(B) - f(A) = \iint (\mathfrak{D}f)(x, y)(y + i) dE_B(x)(B - A)(A + iI)^{-1} dE_A(y).$$

4. Schur multipliers originating from divided differences

Let f be a differentiable function on \mathbb{R} . As before, we use the symbol $\mathfrak{D}f$ the divided difference (3.5).

Definition. We say that a function φ on \mathbb{R} is a multiplier of the space OL of operator Lipschitz functions if

$$f \in \text{OL} \implies \varphi f \in \text{OL}.$$

We denote by \mathfrak{M}_{OL} the class of all multipliers of OL, By the norm $\|\varphi\|_{\mathfrak{M}_{\text{OL}}}$ of a function φ in \mathfrak{M}_{OL} we mean the norm of the multiplication operator $f \mapsto \varphi f$ on the space OL.

For convenience, throughout the paper we introduce the following notation. We denote by \mathbf{x} and \mathbf{y} the functions on \mathbb{R}^2 defined by

$$\mathbf{x}(s, t) \stackrel{\text{def}}{=} s, \quad (s, t) \in \mathbb{R}^2, \quad \text{and} \quad \mathbf{y}(s, t) \stackrel{\text{def}}{=} t, \quad (s, t) \in \mathbb{R}^2.$$

We are also going to use the notation \mathbf{x} for the function on \mathbb{R} defined by

$$\mathbf{x}(s) = s, \quad s \in \mathbb{R}.$$

We start with analyzing the condition

$$(\mathbf{x} + i)(\mathfrak{D}f) \in \mathfrak{M}(\mathbb{R}^2).$$

As we have mentioned in § 3, we are going to prove in this paper that this condition is equivalent to the fact that $f \in \text{ROL}$, i.e., f is a relatively operator Lipschitz function.

Theorem 4.1. *Let f be a differentiable function on \mathbb{R} . The following are equivalent:*

- (a) $(\mathbf{x} + \mathbf{i})(\mathfrak{D}f) \in \mathfrak{M}(\mathbb{R}^2)$;
- (b) $(\mathbf{x} + \mathbf{i})f \in \text{OL}(\mathbb{R})$;
- (c) $f \in \mathfrak{M}_{\text{OL}}$.

Proof. Let us first show that (a) \Rightarrow (b). Suppose that $(\mathbf{x} + \mathbf{i})(\mathfrak{D}f) \in \mathfrak{M}(\mathbb{R}^2)$. Analyzing $((\mathbf{x} + \mathbf{i})(\mathfrak{D}f))(x, y)$ for $y = 0$, we find that f is bounded. It remains to observe that

$$\frac{(x + \mathbf{i})f(x) - (y + \mathbf{i})f(y)}{x - y} = (x + \mathbf{i})(\mathfrak{D}f)(x, y) + f(y), \quad x, y \in \mathbb{R}. \quad (4.1)$$

Let us now prove that (b) \Rightarrow (a). Since $(\mathbf{x} + \mathbf{i})f \in \text{OL}(\mathbb{R})$, it follows that

$$|(t + \mathbf{i})f(t) - \mathbf{i}f(0)| \leq C|t| \quad \text{for every } t \in \mathbb{R}.$$

Consequently, f is bounded. It follows now from (4.1) that $(\mathbf{x} + \mathbf{i})(\mathfrak{D}f) \in \mathfrak{M}(\mathbb{R}^2)$.

Next, the implication (c) \Rightarrow (b) is trivial because $(\mathbf{x} + \mathbf{i}) \in \text{OL}(\mathbb{R})$. It remains to establish that (b) \Rightarrow (c).

Clearly, we may assume that $f(0) = 0$. Since $(\mathbf{x} + \mathbf{i})f \in \text{OL}(\mathbb{R})$, it is easy to see that

$$|(t + \mathbf{i})f(t)| \leq \|(\mathbf{x} + \mathbf{i})f\|_{\text{OL}(\mathbb{R})}|t| \quad \text{for every } t \in \mathbb{R}.$$

Consequently,

$$|f(t)| < \|(\mathbf{x} + \mathbf{i})f\|_{\text{OL}(\mathbb{R})} \quad \text{for every } t \in \mathbb{R}. \quad (4.2)$$

Let us show that $fh \in \text{OL}(\mathbb{R})$ for an arbitrary function h in $\text{OL}(\mathbb{R})$.

Let A and B be bounded self-adjoint operators.

Put $g = (\mathbf{x} + \mathbf{i})^{-1}h$. The identity

$$\begin{aligned} f(A)h(A) - f(B)h(B) &= f(A)(h(A) - h(B)) + (f(A) - f(B))h(B) \\ &= f(A)(h(A) - h(B)) + (f(A)(B + \mathbf{i}I) - f(B)(B + \mathbf{i}I))g(B) \\ &= f(A)(h(A) - h(B)) + (f(A)(A + \mathbf{i}I) - f(B)(B + \mathbf{i}I))g(B) \\ &\quad - f(A)(A - B)g(B) \end{aligned}$$

implies that

$$\begin{aligned} \|f(A)h(A) - f(B)h(B)\| &\leq \|f\|_{L^\infty(\mathbb{R})}\|h\|_{\text{OL}(\mathbb{R})}\|A - B\| \\ &\quad + \|(\mathbf{x} + \mathbf{i})f\|_{\text{OL}(\mathbb{R})}\|g\|_{L^\infty(\mathbb{R})}\|A - B\| + \|f\|_{L^\infty(\mathbb{R})}\|g\|_{L^\infty(\mathbb{R})}\|A - B\|. \end{aligned}$$

By (4.2), $\|f\|_{L^\infty(\mathbb{R})} \leq \|(\mathbf{x} + \mathbf{i})f\|_{\text{OL}(\mathbb{R})}$. It remains to observe that $g \in L^\infty(\mathbb{R})$. Indeed,

$$|g(t)| \leq |t + \mathbf{i}|^{-1}(|h(t) - h(0)| + |h(0)|) \leq \|h\|_{\text{OL}(\mathbb{R})} + |h(0)|, \quad t \in \mathbb{R}. \quad \blacksquare$$

Note that the equivalence of (b) and (c) in Theorem 4.1 is a special case of Theorem 4.7 in [A].

Remark. Obviously, condition (a) in the statement of the theorem is equivalent to the fact that $(\mathbf{y} + \mathbf{i})(\mathfrak{D}f) \in \mathfrak{M}(\mathbb{R}^2)$.

Let us proceed now to the condition

$$(\mathbf{x} + \mathbf{i})(\mathbf{y} + \mathbf{i})(\mathfrak{D}f) \in \mathfrak{M}(\mathbb{R}^2).$$

As we have already mentioned in § 3, this condition describes the class of resolvent Lipschitz functions. We have also mentioned in § 3 that this condition is equivalent to the fact that the function φ on $\mathbb{T} \setminus \{1\}$ defined by (3.7) extends to an operator Lipschitz function on \mathbb{T} .

Theorem 4.2. *Let f be a differentiable function on \mathbb{R} . The following are equivalent:*

- (a) $(\mathbf{x} + \mathbf{i})(\mathbf{y} + \mathbf{i})(\mathfrak{D}f) \in \mathfrak{M}(\mathbb{R}^2)$;
- (b) $(\mathbf{x} + \mathbf{i})^2(f - c) \in \text{OL}(\mathbb{R})$ for some $c \in \mathbb{C}$;
- (c) $(\mathbf{x} + \mathbf{i})(f - c) \in \mathfrak{M}_{\text{OL}}$ for some $c \in \mathbb{C}$.

If $c \in \mathbb{C}$ satisfies (b) or (c), then c must be equal to $\lim_{|x| \rightarrow \infty} f(x)$.

Proof. We need the following identity:

$$\begin{aligned} \frac{(x + \mathbf{i})^2 f(x) - (y + \mathbf{i})^2 f(y)}{x - y} &= (x + \mathbf{i})(y + \mathbf{i})(\mathfrak{D}f)(x, y) \\ &+ (x + \mathbf{i})f(x) + (y + \mathbf{i})f(y). \end{aligned} \quad (4.3)$$

Let us show that (a) \Rightarrow (b). Suppose that $(\mathbf{x} + \mathbf{i})(\mathbf{y} + \mathbf{i})(\mathfrak{D}f) \in \mathfrak{M}(\mathbb{R}^2)$. Then $(\mathbf{x} + \mathbf{i})(\mathfrak{D}f) \in \mathfrak{M}(\mathbb{R}^2)$. It follows from the previous theorem that $(\mathbf{x} + \mathbf{i})f \in \text{OL}(\mathbb{R})$. Thus, the limit $c \stackrel{\text{def}}{=} \lim_{|x| \rightarrow \infty} f(x)$ exists. Without loss of generality we may assume that $c = 0$. To deduce from (4.3) the inclusion $(\mathbf{x} + \mathbf{i})^2 f \in \text{OL}(\mathbb{R})$, it suffices to observe that the function $(\mathbf{x} + \mathbf{i})f$ is bounded. This follows from the fact that $\lim_{|x| \rightarrow +\infty} f(x) = 0$ and from the fact that the function $(\mathbf{x} + \mathbf{i})^2(\mathfrak{D}f)(\mathbf{x}, \mathbf{x}) = (\mathbf{x} + \mathbf{i})^2 f'$ is bounded.

Suppose now that $(\mathbf{x} + \mathbf{i})^2(f - c) \in \text{OL}(\mathbb{R})$ for some $c \in \mathbb{C}$. Let us show that $(\mathbf{x} + \mathbf{i})(\mathbf{y} + \mathbf{i})(\mathfrak{D}f) \in \mathfrak{M}(\mathbb{R}^2)$. Again, we may assume that $c = 0$. The result follows from the fact that the function $(\mathbf{x} + \mathbf{i})f$ is bounded and from identity (4.3).

The equivalence of (b) and (c) is a consequence of in the statement of Theorem 4.1.

Finally, it is obvious that (b) can hold only for one value of c . The same is true for (c). ■

Remark. As we have already observed, condition (a) in the statement of Theorem 4.2 is equivalent to the fact that the function φ on $\mathbb{T} \setminus \{1\}$ defined by (3.7) extends to an operator Lipschitz function on \mathbb{T} . This allows us to reduce the equivalence of statements (a) and (b) to Theorem 5.6 of the paper [A].

5. A representation of operator differences in terms of double operator integrals

Suppose that f is a differentiable function on \mathbb{R} . Consider the functions $\mathfrak{D}_I f$ and $\mathfrak{D}_{II} f$ on \mathbb{R}^2 defined by

$$(\mathfrak{D}_I f)(x, y) \stackrel{\text{def}}{=} \frac{f(x) - f(y)}{x - y}(y + i) \quad \text{and} \quad (\mathfrak{D}_{II} f)(x, y) \stackrel{\text{def}}{=} \frac{f(x) - f(y)}{x - y}(y^2 + 1)^{1/2}. \quad (5.1)$$

By the value of $(f(x) - f(y))(x - y)^{-1}$ in the case when $x = y$ we mean the derivative $f'(x)$.

It is easy to see that $\mathfrak{D}_{II} f$ is a Schur multiplier with respect to arbitrary Borel spectral measures on \mathbb{R} if and only if $\mathfrak{D}_I f$ has this property.

Theorem 5.1. *Suppose that A and B are self-adjoint operators such that $B - A$ is a relatively bounded perturbation of A . Let f be a differentiable function on \mathbb{R} such that $\mathfrak{D}_{II} f$ is a Schur multiplier with respect to arbitrary spectral measures on \mathbb{R} . Then f is a relatively operator Lipschitz function and*

$$\begin{aligned} f(B) - f(A) &= \iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y}(y + i) \, dE_B(x)(B - A)(A + iI)^{-1} \, dE_A(y) \\ &= \iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y}(y^2 + 1)^{1/2} \, dE_B(x)(B - A)(A^2 + I)^{-1/2} \, dE_A(y). \end{aligned} \quad (5.2)$$

Proof. Clearly, the double operator integral

$$\iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y}(y^2 + 1)^{1/2} \, dE_B(x)(B - A)(A^2 + I)^{-1/2} \, dE_A(y)$$

represents a bounded linear operator and

$$\begin{aligned} &\left\| \iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y}(y^2 + 1)^{1/2} \, dE_B(x)(B - A)(A^2 + I)^{-1/2} \, dE_A(y) \right\| \\ &\leq \|\mathfrak{D}_{II} f\|_{\mathfrak{M}(\mathbb{R}^2)} \|(B - A)(A^2 + I)^{-1/2}\|. \end{aligned} \quad (5.3)$$

For a positive number M , we consider the spectral projections, $P_M \stackrel{\text{def}}{=} E_A[-M, M]$ and $Q_M \stackrel{\text{def}}{=} E_B[-M, M]$. We are going to use the following agreement: by \int_{-M}^M we mean $\int_{[-M, M]}$, i.e., the integral over the interval $[-M, M]$.

We have

$$\begin{aligned}
Q_M & \left(\iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y} (y^2 + 1)^{1/2} dE_B(x) G(B - A)(A^2 + I)^{-1/2} dE_A(y) \right) P_M \\
&= \int_{-M}^M \int_{-M}^M \frac{f(x) - f(y)}{x - y} (y^2 + 1)^{1/2} dE_B(x) (B - A)(A^2 + I)^{-1/2} dE_A(y) \\
&= \int_{-M}^M \int_{-M}^M \frac{f(x) - f(y)}{x - y} (y^2 + 1)^{1/2} dE_B(x) (B - A)(A^2 + I)^{-1/2} dE_A(y) \\
&= \int_{-M}^M \int_{-M}^M \frac{f(x) - f(y)}{x - y} (y^2 + 1)^{1/2} (y^2 + 1)^{-1/2} dE_{A+K}(x) (B - A) dE_A(y) \\
&= \int_{-M}^M \int_{-M}^M \frac{f(x) - f(y)}{x - y} dE_B(x) (B - A) dE_A(y).
\end{aligned}$$

Note that the operator B is bounded on Range Q_M while A is bounded on Range P_M . Thus,

$$\begin{aligned}
& \int_{-M}^M \int_{-M}^M \frac{f(x) - f(y)}{x - y} dE_B(x) (B - A) dE_A(y) = \\
&= \int_{-M}^M \int_{-M}^M \frac{f(x) - f(y)}{x - y} dE_B(x) B dE_A(y) - \int_{-M}^M \int_{-M}^M \frac{f(x) - f(y)}{x - y} dE_B(x) A dE_A(y) \\
&= \int_{-M}^M \int_{-M}^M x \frac{f(x) - f(y)}{x - y} dE_B(x) dE_A(y) - \int_{-M}^M \int_{-M}^M y \frac{f(x) - f(y)}{x - y} dE_B(x) dE_A(y) \\
&= \int_{-M}^M \int_{-M}^M (x - y) \frac{f(x) - f(y)}{x - y} dE_B(x) dE_A(y) = \int_{-M}^M \int_{-M}^M (f(x) - f(y)) dE_B(x) dE_A(y) \\
&= \int_{-M}^M \int_{-M}^M f(x) dE_B(x) dE_A(y) - \int_{-M}^M \int_{-M}^M f(y) dE_B(x) dE_A(y) \\
&= Q_M(f(B) - f(A))P_M.
\end{aligned}$$

Since both Q_M and P_M converge as $M \rightarrow \infty$ to I in the strong operator topology, it follows that $f(A + K) - f(A)$ is a bounded operator and $f(A + K) - f(A)$ is equal to

the very last term in (5.2). The first equality in (5.2) can be proved in exactly the same way.

Finally, inequality (5.3) exactly means that f is relatively operator Lipschitz. ■

Below we prove Theorem 6.1, which asserts that the converse is also true, i.e., if f is relatively operator Lipschitz, then $\mathfrak{D}_I f$ and $\mathfrak{D}_{II} f$ belong to $\mathfrak{M}(\mathbb{R}^2)$.

Corollary 5.2. *Suppose that K is a self-adjoint relatively trace class perturbation of a self-adjoint operator A and let f be a differentiable function such that $\mathfrak{D}_I f$ is a Schur multiplier with respect to arbitrary Borel spectral measures on \mathbb{R} . Then $f(A+K) - f(A) \in \mathcal{S}_1$.*

Proof. Indeed, under the hypotheses of the corollary the transformer

$$D \mapsto \iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y} (y + i) dE_{A+K}(x) D dE_A(y)$$

is a bounded operator on \mathcal{S}_1 . The result follows from (5.2). ■

Recall that it was established in [Pe1] and [Pe2] that functions in the Besov class $B_{\infty,1}^1(\mathbb{R})$ on the real line must be operator Lipschitz. This together with Theorem 5.1 and Theorem 4.1 allows us to obtain the following sufficient condition for relative operator Lipschitzness.

Theorem 5.3. *Let f be a function on \mathbb{R} such that the function g defined by*

$$g(x) = f(x)(x + i), \quad x \in \mathbb{R},$$

belongs to the Besov class $B_{\infty,1}^1(\mathbb{R})$. Then f is a relatively operator Lipschitz functions.

6. The necessity of the condition $\mathfrak{D}_I f \in \mathfrak{M}(\mathbb{R}^2)$

To establish the necessity of the condition $\mathfrak{D}_I f \in \mathfrak{M}(\mathbb{R}^2)$ for f to be relatively operator Lipschitz, we consider the one-parametric family $A_t \stackrel{\text{def}}{=} A + tK$, $0 \leq t \leq 1$, and study the behaviour of the operators $f(A_t) - f(A)$. Here A and B are a self-adjoint operators on Hilbert space such that $K \stackrel{\text{def}}{=} B - A$ is a relatively bounded perturbation of A .

Theorem 6.1. *Let f be a relatively operator Lipschitz function on \mathbb{R} . Then the function*

$$\mathfrak{D}_I f = (\mathfrak{D}f)(\mathbf{y} + i)$$

is a Schur multiplier with respect to arbitrary Borel spectral measures;

Lemma 6.2. *Suppose that f is a relatively operator Lipschitz function. Let $M > 0$. Then the transformer*

$$T \mapsto \iint_{\mathbb{R} \times [-M, M]} \frac{f(x) - f(y)}{x - y} (y + i) dE_A(x) T dE_A(y), \quad T = T^* \in \mathcal{S}_2, \quad (6.1)$$

is bounded in the operator norm and the norm of the transformer is less than or equal to k , where k is the constant in (2.6).

Proof. Suppose that A and B are self-adjoint operators such that $K \stackrel{\text{def}}{=} B - A$ is a relatively bounded perturbation of A such that $T \stackrel{\text{def}}{=} K(A + iI)^{-1} \in \mathbf{S}_2$.

Put $A^{[M]} \stackrel{\text{def}}{=} AE_A([-M, M])$. Clearly, $A^{[M]}$ is a bounded linear operator. We have

$$\begin{aligned} \iint_{\mathbb{R} \times [-M, M]} \frac{f(x) - f(y)}{x - y} (y + i) dE_{A^{[M]}}(x) K (A^{[M]} + iI)^{-1} dE_{A^{[M]}}(y) \\ = \iint_{\mathbb{R} \times [-M, M]} \frac{f(x) - f(y)}{x - y} dE_{A^{[M]}}(x) K^{[M]} dE_{A^{[M]}}(y), \end{aligned}$$

where $K^{[M]} \stackrel{\text{def}}{=} KE_A([-M, M]) = T(A + iI)E_A([-M, M]) \in \mathbf{S}_2$.

Consider the one-parametric family $A_t^{[M]} = A^{[M]} + tK^{[M]}$, $t \in [0, 1]$. Then $A_0^{[M]} = A^{[M]}$.

Let f be a relatively operator Lipschitz function. Then f is operator Lipschitz (see the remark in § 2). We have

$$f(A_t^{[M]}) - f(A_0^{[M]}) = t \iint_{\mathbb{R} \times [-M, M]} \frac{f(x) - f(y)}{x - y} dE_{A_t^{[M]}}(x) K^{[M]} dE_{A^{[M]}}(y).$$

By Theorem 3.5.6 of [AP1],

$$\left. \frac{d}{dt} (f(A_t^{[M]}) - f(A_0^{[M]})) \right|_{t=0} = \iint_{\mathbb{R} \times [-M, M]} \frac{f(x) - f(y)}{x - y} dE_A(x) K^{[M]} dE_A(y)$$

in the strong operator topology.

Since f is relatively operator Lipschitz, we have

$$\|f(A_t^{[M]}) - f(A_0^{[M]})\| \leq kt \|K^{[M]}(A^{[M]} + iI)^{-1}\| \leq kt \|K(A + iI)^{-1}\|, \quad t > 0,$$

where k is the constant in (2.6). It follows that

$$\begin{aligned} \left\| \iint_{\mathbb{R} \times [-M, M]} \frac{f(x) - f(y)}{x - y} (y + i) dE_A(x) T dE_A(y) \right\| \\ = \left\| \iint_{\mathbb{R} \times [-M, M]} \frac{f(x) - f(y)}{x - y} (y + i) dE_A(x) K dE_A(y) \right\| \\ \leq k \|K(A + iI)^{-1}\| = k \|T\|. \quad \blacksquare \end{aligned}$$

Proof Theorem 6.1. Put

$$\mathcal{W}_M T \stackrel{\text{def}}{=} \iint_{\mathbb{R} \times [-M, M]} \frac{f(x) - f(y)}{x - y} (y + i) dE_A(x) T dE_A(y).$$

By Lemma 6.2,

$$\|\mathcal{W}_M T\| \leq k \|T\|.$$

It is easy to see now that the limit

$$\lim_{M \rightarrow \infty} \mathcal{W}_M T$$

exists in the strong operator topology for every T in \mathcal{S}_2 and equals

$$\iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y} (y + i) \, dE_A(x) T \, dE_A(y).$$

Thus,

$$\left\| \iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y} (y + i) \, dE_A(x) T \, dE_A(y) \right\| \leq k \|T\|$$

which proves the result. ■

Let us now observe that condition (2.7) does not imply that f is relatively operator Lipschitz. Indeed, it follows from Theorem 6.1 and Theorem 4.1 that if $f \in \text{ROL}$, then $(\mathbf{x} + i)f \in \text{OL}(\mathbb{R})$, and so f must satisfy the following necessary condition (see [AP1]):

Theorem 6.3. *If $f \in \text{ROL}$, then the function $(\mathbf{x} + i)f \in \text{OL}(\mathbb{R})$ must belong to the Besov space $B_1^1(\mathbb{R})$ locally.*

We refer the reader to [AP1] for other necessary conditions for a function to be operator Lipschitz.

7. Commutator estimates

In this section we are going to obtain a characterisation of the class of relatively operator Lipschitz functions in terms of commutator and quasi-commutator estimates.

The following result means, in particular, that a function is relatively operator Lipschitz if and only if it is *relatively commutator Lipschitz*, i.e., satisfies statement (b) of the following theorem.

Theorem 7.1. *Let $f : \mathbb{R} \rightarrow \mathbb{C}$ be a differentiable function. The following statements are equivalent:*

- (a) f is a relatively operator Lipschitz function;
- (b)

$$\|f(A)R - Rf(A)\| \leq \text{const} \|(AR - RA)(A + iI)^{-1}\|$$

for any bounded operator R and any self-adjoint operator A ;

- (c)

$$\|f(B)R - Rf(A)\| \leq \text{const} \|(BR - RA)(A + iI)^{-1}\|$$

for any bounded operator R and any self-adjoint operators A and B ;

- (d) The function $\mathfrak{D}_I f$ belongs to $\mathfrak{M}(\mathbb{R}^2)$.

Proof. First we prove that (a) \implies (c). Let φ be an operator Lipschitz function on \mathbb{T} satisfying (3.7) and such that $\varphi(1) = \lim_{|t| \rightarrow \infty} f(t)$. Clearly, φ is an operator Lipschitz on \mathbb{T} . Hence,

$$\|\varphi(V)R - R\varphi(U)\| \leq \text{const} \|VR - RU\| \tag{7.1}$$

for every bounded operator R and for every unitary operators U and V .

Let A and B be self-adjoint operators. Put $U = (A - iI)(A + iI)^{-1}$ and $V = (B - iI)(B + iI)^{-1}$. Applying (7.1) to unitary operators U and V we get

$$\begin{aligned}
\|f(B)R - Rf(A)\| &= \|\varphi(V)R - R\varphi(U)\| \leq \|\varphi\|_{\text{OL}(\mathbb{T})}\|VR - RU\| \\
&= \|\varphi\|_{\text{OL}(\mathbb{T})}\|(B + iI)^{-1}(B - iI)R - R(A - iI)(A + iI)^{-1}\| \\
&= \|\varphi\|_{\text{OL}(\mathbb{T})}\|(B + iI)^{-1}((B - iI)R(A + iI) - (B + iI)R(A - iI))(A + iI)^{-1}\| \\
&= 2\|\varphi\|_{\text{OL}(\mathbb{T})}\|(B + iI)^{-1}(BR - RA)(A + iI)^{-1}\| \\
&\leq 2\|\varphi\|_{\text{OL}(\mathbb{T})}\|(BR - RA)(A + iI)^{-1}\|.
\end{aligned}$$

The implication (c) \implies (a) is evident.

Let us prove that (b) \implies (c). Applying (a) to the self-adjoint operator $\mathbf{A} = \begin{pmatrix} A & \mathbf{0} \\ \mathbf{0} & B \end{pmatrix}$ and the bounded operator $\mathbf{R} = \begin{pmatrix} \mathbf{0} & \mathbf{0} \\ R & \mathbf{0} \end{pmatrix}$, we find that

$$\|f(\mathbf{A})\mathbf{R} - \mathbf{R}f(\mathbf{A})\| \leq \text{const} \|(\mathbf{A}\mathbf{R} - \mathbf{R}\mathbf{A})(\mathbf{A} + i\mathbf{I})^{-1}\|.$$

It remains to observe that

$$\|f(\mathbf{A})\mathbf{R} - \mathbf{R}f(\mathbf{A})\| = \|f(B)R - Rf(A)\|,$$

where $\mathbf{I} \stackrel{\text{def}}{=} \begin{pmatrix} I & \mathbf{0} \\ \mathbf{0} & I \end{pmatrix}$, and

$$\|(\mathbf{A}\mathbf{R} - \mathbf{R}\mathbf{A})(\mathbf{A} + i\mathbf{I})^{-1}\| = \|(BR - RA)(A + iI)^{-1}\|.$$

The implication (c) \implies (b) is evident.

It remains to observe that the implication (a) \implies (d) follows from Theorem 5.1 and the implication (d) \implies (a) follows from Theorem 6.1. \blacksquare

8. Differentiating in the strong operator topology

In this section we prove the differentiability of the function $t \mapsto f(A_t) - f(A)$ in the strong operator topology and obtain a formula for the derivative. This formula will be used in § 9 to study spectral shift functions in the case of relatively trace class perturbations.

Theorem 8.1. *Let f be a relatively operator Lipschitz function. Suppose that a self-adjoint operator K is a relatively bounded perturbation of a self-adjoint operator A . Then the function $t \mapsto f(A_t) - f(A)$ is differentiable at 0 in the strong operator topology and*

$$\begin{aligned}
\left. \frac{d}{dt}(f(A_t) - f(A)) \right|_{t=0} &= \iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y} (y + i) dE_A(x) C dE_A(y) \\
&= \iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y} (y^2 + 1)^{1/2} dE_A(x) G dE_A(y) \quad (8.1)
\end{aligned}$$

where the operator C and G are defined by (2.2) and (2.3).

To prove Theorem 8.1, we need the following result.

Theorem 8.2. *Let A be a self-adjoint operator and let K be a self-adjoint relatively bounded perturbation of A . Suppose that φ is a continuous function on \mathbb{R} such that the limit $\lim_{|t| \rightarrow \infty} \varphi(t)$ exists. Then the function*

$$s \mapsto \varphi(A + sK)$$

is continuous in the operator norm.

Proof. Without loss of generality we may assume that $\lim_{|t| \rightarrow \infty} \varphi(t) = 0$. Let $\varepsilon > 0$. Clearly, there exists an infinitely smooth function g on \mathbb{R} with compact support such that $\max_{t \in \mathbb{R}} |\varphi(t) - g(t)| < \varepsilon$. Clearly, g is a relatively operator Lipschitz function and it follows from Theorem 5.1 that

$$\|g(A + sK) - g(A + s_0K)\| \leq \text{const } |s - s_0| \cdot \|K(A + iI)^{-1}\|.$$

The result follows from the inequalities

$$\begin{aligned} \|\varphi(A + sK) - \varphi(A + s_0K)\| &\leq \|\varphi(A + sK) - g(A + sK)\| \\ &\quad + \|g(A + sK) - g(A + s_0K)\| + \|g(A + s_0K) - \varphi(A + s_0K)\| \\ &\leq 2\varepsilon + \text{const } |s - s_0| \cdot \|K(A + iI)^{-1}\|. \quad \blacksquare \end{aligned}$$

To prove Theorem 8.1, we need some preparation.

Let $\widehat{\mathbb{R}} \stackrel{\text{def}}{=} \mathbb{R} \cup \{\infty\}$ denote the one-point compactification of the real line \mathbb{R} . We recall that each function $f \in \text{OL}(\mathbb{R})$ is everywhere differentiable on $\widehat{\mathbb{R}}$ (see Theorem 3.3.3 in [AP1]). Recall that f is said to be differentiable at ∞ if there exists a finite limit $\lim_{|x| \rightarrow \infty} x^{-1}f(x) \stackrel{\text{def}}{=} f'(\infty)$.

Lemma 8.3. *Let f satisfy the equivalent conditions of Theorem 4.1. Then there are sequences $\{\varphi_n\}_{n \geq 0}$ and $\{\psi_n\}_{n \geq 0}$ of continuous functions on $\widehat{\mathbb{R}}$ and a positive number M such that the following statements hold:*

- (a) $\sum_{n \geq 0} |\varphi_n|^2 \leq M$ everywhere on $\widehat{\mathbb{R}}$;
- (b) $\sum_{n \geq 0} |\psi_n|^2 \leq M$ everywhere on $\widehat{\mathbb{R}}$;
- (c) $(x + i)(\mathfrak{D}f)(x, y) = \sum_{n \geq 0} \varphi_n(x)\psi_n(y)$ for all $x, y \in \mathbb{R}$.

Moreover, the constants in (a) and (b) depend only on f .

Proof. Note that there exists a finite limit $\lim_{|y| \rightarrow \infty} f(y)$. It suffices to observe that $\lim_{|y| \rightarrow \infty} f(y) = g'(\infty)$ for the operator Lipschitz function $g = (\mathbf{x} + i)f$.

Applying Theorem 3.5.7 in [AP1] to the operator Lipschitz function $(\mathbf{x} + \mathbf{i})f$, we find that

$$\frac{(x + \mathbf{i})f(x) - (y + \mathbf{i})f(y)}{x - y} = \sum_{n \geq 1} \varphi_n(x)\psi_n(y)$$

for some sequences $\{\varphi_n\}_{n \geq 1}$ and $\{\psi_n\}_{n \geq 1}$ of continuous functions on $\widehat{\mathbb{R}}$ such that

$$\sum_{n \geq 1} |\varphi_n|^2 \leq \|(\mathbf{x} + \mathbf{i})f\|_{\text{OL}(\mathbb{R})} \quad \text{and} \quad \sum_{n \geq 1} |\psi_n|^2 \leq \|(\mathbf{x} + \mathbf{i})f\|_{\text{OL}(\mathbb{R})}$$

everywhere on $\widehat{\mathbb{R}}$.

Applying equality (4.1), we obtain

$$(x + \mathbf{i})(\mathfrak{D}f)(x, y) = \sum_{n \geq 0} \varphi_n(x)\psi_n(y) - f(y).$$

To complete the proof, we put $\varphi_0(x) = 1$, $x \in \mathbb{R}$, and $\varphi_0(x) = -f(y)$, $y \in \mathbb{R}$. ■

Proof of Theorem 8.1. The proof is similar to the proof of Theorem 3.5.6 in [AP1]. We have

$$\begin{aligned} \frac{1}{t}(f(A_t) - f(A)) &= \iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y} (y + \mathbf{i}) \, dE_{A_t}(x) K(A + \mathbf{i}I)^{-1} \, dE_A(y) \\ &= \sum_{n \geq 0} \varphi_n(A_t) K(A + \mathbf{i}I)^{-1} \psi_n(A), \end{aligned}$$

where φ_n and ψ_n denote the same as in Lemma 8.3. It remains to prove that

$$\lim_{t \rightarrow 0} \sum_{n \geq 0} \varphi_n(A_t) K(A + \mathbf{i}I)^{-1} \psi_n(A) = \sum_{n \geq 0} \varphi_n(A) K(A + \mathbf{i}I)^{-1} \psi_n(A)$$

in the strong operator topology. Thus, we need to prove that for an arbitrary vector u ,

$$\lim_{t \rightarrow 0} \sum_{n \geq 0} (\varphi_n(A_t) - \varphi_n(A)) K(A + \mathbf{i}I)^{-1} \psi_n(A) u = \mathbf{0}$$

where the series is understood in the sense of the weak topology of \mathcal{H} while the limit is taken in the norm of \mathcal{H} . Assume that $\|u\| = 1$. Let $u_n \stackrel{\text{def}}{=} K(A + \mathbf{i}I)^{-1} \psi_n(A) u$. We have

$$\begin{aligned} \sum_{n \geq 0} \|u_n\|^2 &\leq \|K(A + \mathbf{i}I)^{-1}\|^2 \sum_{n \geq 0} \|(\psi_n(A)u)\|^2 \\ &= \|K(A + \mathbf{i}I)^{-1}\|^2 \sum_{n \geq 0} (|\psi_n|^2(A)u, u) \leq M \|K(A + \mathbf{i}I)^{-1}\|^2, \end{aligned}$$

where M is the number in the statement of Lemma 8.3. Let $\varepsilon > 0$ and choose a natural number N such that $\sum_{n>N} \|u_n\|^2 < \varepsilon^2$. Then it follows from Lemma 3.5.9 in [AP1] that

$$\begin{aligned} \left\| \sum_{n>N} (\varphi_n(A + tK) - \varphi_n(A))u_n \right\| &\leq \left\| \sum_{n>N} \varphi_n(A + tK)u_n \right\| \\ &+ \left\| \sum_{n>N} \varphi_n(A)u_n \right\| < 2M^{1/2}\varepsilon \end{aligned} \quad (8.2)$$

for all $t \in \mathbb{R}$. By Theorem 8.2,

$$\left\| \sum_{n=0}^N (\varphi_n(A + tK) - \varphi_n(A))u_n \right\| < \varepsilon$$

for all t sufficiently close to zero. Thus, in view of (8.2),

$$\left\| \sum_{n \geq 0} (\varphi_n(A + tK) - \varphi_n(A))u_n \right\| < (2M^{1/2} + 1)\varepsilon$$

for all t sufficiently close to zero.

The proof of the fact that the derivative on the left of (8.3) is equal to the second double operator integral in (8.3) is the same. ■

Corollary 8.4. *Let f be a relatively operator Lipschitz function. Suppose that a self-adjoint operator K is a relatively trace class perturbation of a self-adjoint operator A . Then the function $t \mapsto f(A_t) - f(A)$ is differentiable on \mathbb{R} in the strong operator topology and*

$$\begin{aligned} \frac{d}{ds}(f(A_s) - f(A_t)) \Big|_{s=t} &= \iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y} (y + i) dE_{A_t}(x) K(A_t + iI)^{-1} dE_{A_t}(y) \\ &= \iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y} (y^2 + 1)^{1/2} dE_{A_t}(x) K(A_t^2 + I)^{-1/2} dE_{A_t}(y) \end{aligned} \quad (8.3)$$

for any $t \in \mathbb{R}$.

Proof. The result follows from Theorem 8.1 (applied to the operators K and A_t) and from Lemma 2.3. ■

9. Relatively trace class perturbations and the trace formula

In this section we obtain an analogue of the Lifshits–Krein trace formula for relatively trace class perturbation. We also describe the maximal class of functions, for which the trace formula is applicable. The main results of this section are based on Theorem 8.1.

Let A be a self-adjoint operator and let K be a relatively trace class perturbation of A . Suppose that f is a relatively operator Lipschitz function on \mathbb{R} .

The purpose of this section is to prove that the trace formula in the case of relatively trace class perturbations holds for arbitrary relatively operator Lipschitz functions and the corresponding spectral shift function ξ satisfies inequality (9.1).

Theorem 9.1. *Let A be a self-adjoint operator and let K be a relatively trace class perturbation of A . Then there exists a unique measurable real-valued function ξ on \mathbb{R} satisfying*

$$\int_{\mathbb{R}} \frac{|\xi(t)|}{1+|t|} dt < \infty \quad (9.1)$$

and such that for an arbitrary relatively operator Lipschitz function f on \mathbb{R} the following trace formula holds

$$\text{trace} (f(A+K) - f(A)) = \int_{\mathbb{R}} f'(t)\xi(t) dt. \quad (9.2)$$

The function ξ satisfying (9.2) and (9.1) is called the *spectral shift function for the pair* $\{A, A+K\}$.

Note that inequality (9.1) was found earlier in [CS] by different methods. However, in [CS] trace formula (9.2) was obtained under considerably more restrictive assumptions on f .

Moreover, we describe in this section the maximal class of function f , for which the trace formula holds in the case of relatively trace class perturbations, see Theorem 9.3 below, and show that this maximal class coincides with the class of relatively operator Lipschitz functions.

The proof of Theorem 9.1 is close in spirit to the proof of Theorem 6.1 of [Pe3] and to the proof of Theorem 4.1 of [AP2].

Proof. Consider the one-parametric family of operators $A_t \stackrel{\text{def}}{=} A + tK$, $0 \leq t \leq 1$. By Theorem 8.1,

$$\frac{d}{ds} (f(A_s) - f(A_t)) \Big|_{s=t} = \iint_{\mathbb{R} \times \mathbb{R}} \frac{f(x) - f(y)}{x - y} (y + i) dE_{A_t}(x) K(A_t + iI)^{-1} dE_{A_t}(y), \quad (9.3)$$

where the derivative on the left is understood in the strong operator topology.

We need the following lemma:

Lemma 9.2. *The function $t \mapsto K(A_t + iI)^{-1}$ is continuous on $[0, 1]$ in the norm of \mathcal{S}_1 , and so $\|K(A_t + iI)^{-1}\|_{\mathcal{S}_1} \leq \text{const}$, $0 \leq t \leq 1$.*

Proof. We have

$$K(A_t + iI)^{-1} = K(A + iI)^{-1}(A + iI)(A_t + iI)^{-1}.$$

Clearly, it suffices to show that the function

$$t \mapsto ((A + iI)(A_t + iI)^{-1})^{-1} = (A_t + iI)(A + iI)^{-1}$$

is continuous. This follows immediately from the following obvious equality

$$(A_t + iI)(A + iI)^{-1} = I + tK(A + iI)^{-1}. \quad \blacksquare$$

Let us complete the proof of Theorem 9.1. Let Q_t be the operator on the right-hand side of (9.3). Since the function $\mathfrak{D}_I f$ is a Schur multiplier for arbitrary spectral measures, it follows that $Q_t \in \mathcal{S}_1$ for every t in $[0, 1]$ and $\sup_t \|Q_t\|_{\mathcal{S}_1} < \infty$.

Since the function $t \mapsto Q_t$ is continuous in the norm of \mathcal{S}_1 by Lemma 9.2, we obtain

$$f(A + K) - f(A) = \int_0^1 Q_t dt$$

in the sense of integration of continuous functions. Moreover,

$$\text{trace}(f(A + K) - f(A)) = \int_0^1 \text{trace} Q_t dt$$

We are going to use the following known result (see [Pe3], Theorem 5.1):

Let E be a Borel spectral measure on a locally compact topological space X . Suppose that Φ is a Schur multiplier with respect to E . If Φ is continuous in each variable, then

$$\text{trace} \left(\iint \Phi(x, y) dE(x) T dE(y) \right) = \int \Phi(x, x) d\mu(x) \quad (9.4)$$

for an arbitrary trace class operator T where the measure μ is defined by

$$\mu(\Delta) \stackrel{\text{def}}{=} \text{trace}(E(\Delta)T).$$

Applying (9.4) to the Schur multiplier $\mathfrak{D}_I f$ and the spectral measure E_{A_t} , we find that

$$\text{trace} Q_t = \int_{\mathbb{R}} f'(x)(x + i) d\nu_t(x), \quad (9.5)$$

where ν_t is the finite real Borel measure on \mathbb{R} defined by

$$\nu_t(\Delta) = \text{trace}(E_{A_t}(\Delta)K(A_t + iI)^{-1}).$$

As usual, we identify here the space of regular complex Borel measures on \mathbb{R} with the dual space to the Banach space of continuous functions on \mathbb{R} with zero limit at infinity. Then the function $t \mapsto \nu_t$ is continuous in the weak-* topology on the space of complex Borel measures. Indeed, if h is continuous on \mathbb{R} and $\lim_{|x| \rightarrow \infty} h(x) = 0$, then

$$\int h d\nu_t = \text{trace}(h(A_t)K(A_t + iI)^{-1}).$$

By Theorem 8.2, the function $t \mapsto h(A_t)$ is a continuous function on $[0, 1]$ in the operator norm, and so the function $t \mapsto \text{trace}(h(A_t)K(A_t + iI)^{-1})$ is continuous.

Consider the real Borel measure ν on \mathbb{R} defined by

$$\nu = \int_0^1 \nu_t dt.$$

Then by (9.5),

$$\text{trace}(f(A + K) - f(A)) = \int_0^1 f'(x)(x + i) d\nu(x).$$

We have mentioned in § 1 that the condition that K is a relatively trace class perturbation of A implies that

$$(A + K + iI)^{-1} - (A + iI)^{-1} \in \mathbf{S}_1.$$

Then the pair $\{A, A + K\}$ has a real-valued spectral shift function $\boldsymbol{\eta}$, i.e.,

$$\int_{\mathbb{R}} \frac{|\boldsymbol{\eta}(x)|}{1+x^2} dx < \infty$$

and

$$\text{trace} (f(A + K) - f(A)) = \int_{\mathbb{R}} f'(x)\boldsymbol{\eta}(x) dx$$

whenever f is a resolvent Lipschitz function on \mathbb{R} . It follows that

$$\int_{\mathbb{R}} f'(x)\boldsymbol{\eta}(x) dx = \int_{\mathbb{R}} f'(x)(x+i) d\nu(x)$$

for an arbitrary infinitely smooth function f with compact support. It follows that

$$\boldsymbol{\eta}(x) - (x+i) d\nu(x) = c dx$$

for some constant c . In particular, this means that ν is absolutely continuous with respect to Lebesgue measure. We can define now the function $\boldsymbol{\xi}$ by

$$\boldsymbol{\xi}(s) \stackrel{\text{def}}{=} \text{Re} \left((s+i) \frac{d\nu(x)}{dx}(s) \right).$$

Clearly, $\boldsymbol{\xi}$ satisfies (9.1) and equality (9.2) holds.

Let us show that such a spectral shift function is unique. Indeed, if $\boldsymbol{\xi}_1$ and $\boldsymbol{\xi}_2$ are spectral shift functions such that

$$\int_{\mathbb{R}} \frac{|\boldsymbol{\xi}_1(x)|}{1+|x|} dx < \infty \quad \text{and} \quad \int_{\mathbb{R}} \frac{|\boldsymbol{\xi}_2(x)|}{1+|x|} dx < \infty,$$

then $\boldsymbol{\xi}_1 - \boldsymbol{\xi}_2 = c$ is a constant function. Then

$$\int_{\mathbb{R}} \frac{|\boldsymbol{\xi}_1(x)|}{1+x^2} dx < \infty \quad \text{and} \quad \int_{\mathbb{R}} \frac{|\boldsymbol{\xi}_2(x)|}{1+x^2} dx < \infty.$$

Since $\boldsymbol{\xi}_1$ and $\boldsymbol{\xi}_2$ are spectral shift functions that corresponds to the resolvent trace class perturbation K , $\boldsymbol{\xi}_1 - \boldsymbol{\xi}_2 = c$ is a constant function. This is a well known fact. It follows from the fact that an integrable spectral shift function for a pair of unitary operators with trace class difference is unique modulo a constant additive.

If the constant c were nonzero, it would follow that

$$\int_{\mathbb{R}} \frac{|c|}{1+|x|} dx < \infty. \quad \blacksquare$$

The following result shows that the class ROL is the maximal class of functions on \mathbb{R} for which trace formula (9.2) holds.

Theorem 9.3. *Let f be a differentiable function on \mathbb{R} such that trace formula (9.2) holds for arbitrary self-adjoint operators A and K such that K is a relatively trace class perturbation of A . Then $f \in \text{ROL}$.*

Proof. Indeed, it follows from Theorem 2.5 that if $f \notin \text{ROL}$, then there are self-adjoint operators A and K such that K is a relatively trace class perturbation of A but $f(A + K) - f(A) \notin \mathcal{S}_1$. ■

REFERENCES

- [A] A. B. Aleksandrov, Operator Lipschitz functions and linear-fractional transformations, *Zapiski Nauchn. Semin. POMI* **401** (2012), 5–52 (Russian). English transl.: *J. Math. Sci. (N. Y.)* **194:6** (2013), 603–627.
- [AP1] A.B. ALEKSANDROV and V.V. PELLER, *Operator Lipschitz functions*, *Uspekhi Mat. Nauk* **71:4** (2016), 3–106 (Russian). English transl.: *Russian Math. Surveys* **71:4**, 605–702.
- [AP2] A.B. ALEKSANDROV and V.V. PELLER, *Krein’s trace formula for unitary operators and operator Lipschitz functions*, *Funkts. Anal. i Ego Pril.* **50:3** (2016), 1–11 (Russian). English transl.: *Funct. Anal. Appl.* **50:4** (2016), 167–175.
- [AP3] A.B. ALEKSANDROV and V.V. PELLER, *Haagerup tensor products and Schur multipliers*, *Algebra i Analiz* **36:5** (2024),
- [BS1] M.S. BIRMAN and M.Z. SOLOMYAK, *Double Stieltjes operator integrals*, *Problems of Math. Phys., Leningrad. Univ.* **1** (1966), 33–67 (Russian). English transl.: *Topics Math. Physics* **1** (1967), 25–54, Consultants Bureau Plenum Publishing Corporation, New York.
- [BS2] M.S. BIRMAN and M.Z. SOLOMYAK, *Double Stieltjes operator integrals. II*, *Problems of Math. Phys., Leningrad. Univ.* **2** (1967), 26–60 (Russian). English transl.: *Topics Math. Physics* **2** (1968), 19–46, Consultants Bureau Plenum Publishing Corporation, New York.
- [BS3] M.S. BIRMAN and M.Z. SOLOMYAK, *Remarks on the spectral shift function*, *Zapiski Nauchn. Semin. LOMI* **27** (1972), 33–46 (Russian). English transl.: *J. Soviet Math.* **3** (1975), 408–419.
- [BS4] M.S. BIRMAN and M.Z. SOLOMYAK, *Double Stieltjes operator integrals. III*, *Problems of Math. Phys., Leningrad. Univ.* **6** (1973), 27–53 (Russian).
- [BS5] M.S. BIRMAN and M.Z. SOLOMYAK, *Spectral theory of selfadjoint operators in Hilbert space, Mathematics and its Applications (Soviet Series)*, D. Reidel Publishing Co., Dordrecht, 1987.
- [BS6] M.S. BIRMAN and M.Z. SOLOMYAK, *Tensor product of a finite number of spectral measures is always a spectral measure*, *Integral Equations Operator Theory* **24** (1996), 179–187.
- [CS] A. CHATTOPADHYAY and A. SKRIPKA, *Trace formulas for relative Schatten class perturbations*, *J. Funct. Anal.* **274** (2018), 3377–3410.
- [DK] YU.L. DALETSKII and S.G. KREIN, *Integration and differentiation of functions of Hermitian operators and application to the theory of perturbations* (Russian), *Trudy Sem. Function. Anal., Voronezh. Gos. Univ.* **1** (1956), 81–105.
- [F] YU.B. FARFOROVSKAYA, *An example of a Lipschitzian function of selfadjoint operators that yields a nonnuclear increase under a nuclear perturbation*. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* **30** (1972), 146–153 (Russian).
- [GK] I.C. GOKHBERG and M.G. KREIN, *Introduction to the theory of linear nonselfadjoint operators in Hilbert space*, "Nauka", Moscow, 1965; English transl., Amer. Math. Soc., Providence, R.I., 1969.
- [Kr] M.G. KREIN, *On a trace formula in perturbation theory*, *Mat. Sbornik* **33** (1953), 597–626 (Russian).
- [L] I.M. LIFSHITS, *On a problem in perturbation theory connected with quantum statistics*, *Uspekhi Mat. Nauk* **7** (1952), 171–180 (Russian).
- [MNP] M.M. MALAMUD, H. NEIDHARDT and V.V. PELLER, *Absolute continuity of spectral shift*, *J. Funct. Anal.* **276** (2019), 1575–1621.
- [Pee] J. PEETRE, *New thoughts on Besov spaces*, Duke Univ. Press., Durham, NC, 1976.

- [Pe1] V.V. PELLER, *Hankel operators in the theory of perturbations of unitary and self-adjoint operators*, Funktsional. Anal. i Prilozhen. **19:2** (1985), 37–51 (Russian). English transl.: Funct. Anal. Appl. **19** (1985) , 111–123.
- [Pe2] V.V. PELLER, *Hankel operators in the perturbation theory of of unbounded self-adjoint operators*. Analysis and partial differential equations, 529–544, Lecture Notes in Pure and Appl. Math., **122**, Dekker, New York, 1990.
- [Pe3] V.V. PELLER, *The Lifshits–Krein trace formula and operator Lipschitz functions*, Proc. Amer. Math. Soc. **144** (2016), 5207–5215.

A.B. Aleksandrov
 St.Petersburg State University
 Universitetskaya nab., 7/9
 199034 St.Petersburg, Russia

V.V. Peller
 St.Petersburg State University
 Universitetskaya nab., 7/9
 199034 St.Petersburg, Russia

St.Petersburg Department
 Steklov Institute of Mathematics
 Russian Academy of Sciences
 Fontanka 27, 191023 St.Petersburg
 Russia
 email: alex@pdmi.ras.ru

St.Petersburg Department
 Steklov Institute of Mathematics
 Russian Academy of Sciences
 Fontanka 27, 191023 St.Petersburg
 Russia
 email: peller@math.msu.edu