

Viable Wormhole Structures and Energy Conditions in $f(Q, T)$ Theory

M. Zeeshan Gul ^{*}, M. Sharif [†], Shajee Shahid [‡] and Faisal Javed [§]
Department of Mathematics and Statistics, The University of Lahore,
1-KM Defence Road Lahore-54000, Pakistan.
Department of Physics, Zhejiang Normal University, Jinhua 321004,
People's Republic of China.

Abstract

This paper explores static wormhole solutions in $f(Q, T)$ theory, where Q is the non-metricity and T is the trace of energy-momentum tensor. We derive the field equations that describe gravitational phenomena in the existence of non-metricity and matter source terms. We examine different models of this theory to determine the explicit expressions of matter contents, which are useful for analyzing the wormhole structures. We investigate the existence of feasible traversable wormhole solutions for constant and variable redshift functions. To determine whether physically viable wormhole geometry exists, we examine the graphical interpretation of energy constraints for different values of model parameters. It is found that realistic traversable and stable wormhole solutions exist only for the first model of this gravity.

Keywords: $f(Q, T)$ theory; Wormhole; Energy conditions.

PACS: 98.62.Gq, 04.50.Kd, 83C15.

^{*}mzeeshangul.math@gmail.com

[†]msharif.math@pu.edu.pk

[‡]shajeeshaid0312@gmail.com

[§]faisaljaved.math@gmail.com

1 Introduction

The scientific community has been fascinated by the universe mystery and one of the most controversial subjects has been the theoretical idea of wormholes (*WHs*). Wormholes are fictitious tunnel-like formations that link different parts of spacetime [1]. The geometry of *WHs* was first formulated by using the Schwarzschild solution in [2]. Further, Einstein and Rosen [3] suggested that spacetime might be connected via a bridge between distinct regions, termed the Einstein-Rosen bridge. The formation of a *WH* structure requires a specific spacetime curvature induced by a sufficient amount of matter. However, exotic matter (which violates energy conditions (*ECs*)) is necessary for the stability of *WH* structures. Exotic matter is essential for stabilizing and making possible traversal of *WHs* according to theoretical analysis. Schwarzschild *WHs* are non-traversable because of the throat abrupt expansion and contraction [4]. Morris and Thorne [5] examined the first traversable *WH* solution that let matter flow through in both directions. It is important to note that the existence of exotic matter is necessary for traversable *WHs* to remain viable and stable. In theoretical physics, there is still discussion and interest in the study of *WHs* and their characteristics [6]-[8]. The maximal amount of exotic matter in the bridge raises concerns about the viability of the *WH* construction. Therefore, for a feasible *WH* geometry, there should be a sufficient amount of exotic matter in the bridge. Numerous techniques have been developed to examine the feasible *WH* structures [9]. Many researchers studied the viable *WH* geometry through different *WH* shape functions (*WSFs*) [10]. These functions enhance our understanding of the hypothetical configurations and attributes.

Traversable *WHs* are fascinating theoretical constructs that provide a unique window into understanding the interplay between gravity and quantum phenomena. In *GTR*, traversable *WHs* require exotic matter or matter with negative energy density to remain open and stable, which violates *ECs*. However, in the context of quantum gravity, these constraints may be relaxed or reinterpreted. In particular, quantum field theory in curved spacetime allows for the possibility of quantum fluctuations generating negative energy as seen in phenomena like the Casimir effect. These quantum fluctuations suggest that traversable *WHs* might exist without exotic matter in certain quantum gravity frameworks. In recent years, traversable *WHs* have also gained attention through their connections to quantum information theory. This provides a framework where *WHs* can be related to entangled quan-

tum states, offering a potential holographic description of quantum gravity phenomena. Furthermore, traversable *WHs* serve as theoretical laboratories for studying non-trivial spacetime topologies and the causal structure of spacetime in quantum gravity. They offer a promising avenue for investigating the nature of spacetime at the Planck scale and could lead to insights into fundamental issues such as the resolution of singularities, the unitarity problem in black hole evaporation and the possible existence of shortcuts through spacetime. This highlights how traversable *WHs* not only serve as speculative solutions in *GTR* but also play a significant role in advancing our understanding of quantum gravity, particularly in scenarios where classical constraints are reinterpreted through quantum mechanical principles.

Traversable *WHs* relate the geometry of spacetime (described by the metric) to the matter and energy present (described by the stress-energy tensor) have an interesting relationship with the *ECs*. For a *WH* to remain open and traversable, the geometry needs to avoid forming singularities or event horizons that would trap travelers inside. Various energy conditions are used to describe reasonable physical matter distributions. Energy conditions are sets of constraints on the stress-energy tensor that aim to describe normal matter, or exotic matter. When discussing traversable *WHs*, the focus is primarily on the violation of these *ECs*. For a *WH* to be traversable, the geometry must be such that the throat of the *WH* remains open and stable. In traversable *WHs*, it is generally required that the energy conditions are violated in regions around the throat. This implies the presence of exotic matter, allowing the throat to remain open for passage. The energy condition violations in traversable *WHs* are often associated with exotic matter, allowing the *WH* to maintain its open structure. This exotic matter counteracts the attractive nature of gravity, preventing the collapse of the *WH* throat. Thus, traversable *WHs* require violations of various *ECs* to ensure their stability and traversability. The presence of exotic matter plays a key role in this violation.

Einstein general theory of relativity serves as a cornerstone by providing a comprehensive description of gravitational field and matter on cosmic scales. In *GTR*, spacetime is described using mathematical structures defined by Riemann metric. This metric encodes information about distance and angles in spacetime which allow physicists to understand the curvature of spacetime caused by gravity. Weyl [11] introduced a more general geometric framework as an extension of *GTR*. Weyl theory incorporates the concept of a *length connection* which differs from the standard metric connection used

in Riemannian geometry (RG). Weyl theory focuses on gauging the conformal factor adjusting the scale of distances. Weyl introduced the concept of non-metricity which assures that covariant derivative of metric tensor exists. Non-metricity and torsion are two major concepts of non-Riemannian geometries. Alternative theories include torsion and non-metricity as additional geometrical characteristics of spacetime whereas Einstein formulation of GTR emphasizes curvature. There are two equivalent geometric representations of GTR , i.e., the curvature representation vanishes torsion and non-metricity, whereas the teleparallel representation vanishes curvature and non-metricity. However, one more comparable representation of the geometric features is the non-metricity of the metric.

In the framework of GTR , the gravitational interaction is described by a purely metric theory where the connection is assumed to be Levi-Civita, implying zero non-metricity. However, in alternative theories of gravity, especially those attempting to unify gravity with other forces or explain phenomena such as dark energy, dark matter, or early universe cosmology, the assumption of a purely metric theory is relaxed. Non-metricity arises in such scenarios and can be an essential feature of extended geometrical frameworks such as the metric-affine gravity or Teleparallel Gravity. Non-metricity describes how the length of vectors changes under parallel transport and it is closely linked to the concept of varying gravitational couplings or connections that include degrees of freedom beyond the metric. These modifications can provide a deep understanding of phenomena that GTR struggles to explain such as the nature of singularities, the behavior of gravity at quantum scales and potential resolutions to cosmological problems like the singularity in the Big Bang or cosmological bounce models. Thus, studying non-metricity offers a promising pathway to explore gravitational theories that extend beyond the well-established predictions of GTR , opening doors to resolving some of the fundamental issues in modern cosmology and gravitational physics.

Studying non-metricity is motivated by several key factors as

- Extensions to the Geometric Nature of Gravity

Non-metricity introduces an additional layer of geometry that departs from the constraints of GTR . While GTR is based on the curvature of spacetime, incorporating non-metricity allows for a more generalized theory of spacetime that includes non-metricity as fundamental geometric entity. This geometric extension provides alternative mechanisms for understanding gravitational interactions.

- Unification with Quantum Gravity

A persistent challenge in theoretical physics is reconciling *GTR* with quantum mechanics. Non-metricity may offer a pathway toward quantum gravity by introducing new symmetries or degrees of freedom that bridge the gap between classical gravity and quantum field theory. By modifying the fundamental structure of spacetime, theories with non-metricity have the potential to accommodate quantum effects that *GTR* cannot explain such as those predicted by string theory or loop quantum gravity.

- Exploring Dark Energy and Dark Matter

The introduction of non-metricity may also provide insight into the nature of dark energy and dark matter, which remain elusive in the context of *GTR*. Non-metricity could introduce additional fields or modifications to spacetime dynamics that mimic the effects of these unknown components of the universe. It also offers a possible explanation for the accelerated expansion of the universe and the missing mass inferred from galactic rotation curves without needing to invoke unknown particles.

- High-Energy Regimes and Early Universe Cosmology

In the early universe, conditions of extremely high energy and curvature likely deviate from the predictions of *GTR*. Theories that incorporate non-metricity such as teleparallel gravity or generalized affine theories could provide a more accurate description of the early universe, resolving issues like the big bang singularity and offering alternatives to cosmic inflation.

- The Role in Modified Gravity Theories

Non-metricity is a key feature in several modified gravity theories, including the metric-affine gravity framework and theories of gravity that generalize the Palatini formalism. These approaches offer solutions to long-standing issues in cosmology and astrophysics such as the cosmological constant problem and the need for new forms of energy to explain gravitational phenomena at large scales. Hence, the study of non-metricity extends the geometric framework of gravity, providing a fertile ground for new insights into gravitational phenomena that *GTR* cannot fully address. By incorporating non-metricity, researchers aim to uncover a more complete theory of gravity that not only explains known phenomena but also sheds light on the mysteries of dark matter, dark energy, quantum gravity and the early universe.

Teleparallel theory is one such alternative theory where torsion represents the gravitational interaction. In symmetric teleparallel theory, the gravitational interaction is represented by non-metricity. To characterize *GTR* in the context of torsion and non-metricity, the integral actions are expressed as $\int \sqrt{-g}\mathcal{T}$ [12] and $\int \sqrt{-g}Q$ [13] where \mathcal{T} represents torsion and Q represents non-metricity, respectively. Yixin et al [14] formulated the $f(Q, T)$ theory by assimilating the trace of stress-energy tensor in the functional action of $f(Q)$ gravity. The motivation behind this theory is to examine theoretical impacts by observational data and cosmological domains. Xu et al [15] developed the Weyl form of $f(Q, T)$ gravity and its implications in the background of cosmology. Arora et al [16] investigated the cosmic acceleration in the absence of dark energy. Bhattacharjee [17] explored the applications of $f(Q, T)$ gravity and found that this gravity alter the nature of tidal forces and equation of motion in Newtonian limit. Pati et al [18] established a mathematical framework in this theory in terms of the Hubble model. Agrawal et al [19] presented the dynamical features and matter bounce scenario in this background. Shiravand et al [20] explained the cosmic inflation in the same theory.

The $f(Q, T)$ theory of gravity is a modified theory that generalizes the symmetric teleparallel gravity, focusing on the non-metricity scalar and the trace of the energy-momentum tensor. In this framework, non-metricity represents a geometrical object that quantifies how the length of vectors changes during parallel transport. The trace of the energy-momentum tensor includes the contribution from matter fields. This theory departs from the *GTR* and other modified theories by introducing coupling between geometry (non-metricity) and matter field (trace of the energy-momentum tensor), explaining more accurately cosmological and astrophysical phenomena.

- Comparison with Other Modified Gravity Theories

Einstein general theory of relativity is based on the curvature of spacetime with the Ricci scalar as the fundamental geometrical object. The $f(R)$ gravity is one of the most well-known modifications to *GTR*, where the Ricci scalar in the Einstein-Hilbert action is replaced by a function $f(R)$, allowing more complex interactions between curvature and gravity. The field equations are fourth-order differential equations instead of the second-order equations in *GTR*. The $f(R, T)$ gravity extends $f(R)$ theory by introducing a dependence on both the Ricci scalar and the trace of the energy-momentum tensor, allowing a coupling between geometry and matter. The equations of

motion are more complex than $f(R)$ gravity and lead to non-conservation of the energy-momentum tensor. Teleparallel gravity reformulates GTR by using torsion rather than curvature to describe gravity. The gravitational action is constructed from the torsion scalar. In teleparallel gravity, the field equations involve the torsion tensor rather than the curvature of spacetime. Scalar-tensor theories introduce a scalar field with the tensor field (the metric) and gravity is described by both. The equations of motion involve both the metric and the scalar field, leading to a richer set of dynamics, including varying gravitational coupling. The higher-order theories extend GTR by adding terms involving higher powers of curvature such as the Gauss-Bonnet term, which is a specific combination of curvature invariants. The corresponding field equations lead to more complex field equations but retaining second-order derivatives in the Gauss-Bonnet case.

In $f(Q, T)$, the gravitational interaction is based on non-metricity rather than curvature with additional terms involving the matter trace, which introduces modifications to how gravity couples to matter. Both $f(Q, T)$ and $f(R)$ theories generalize GTR , but $f(Q, T)$ is based on non-metricity rather than curvature. Additionally, $f(Q, T)$ includes the matter-energy trace, creating a more direct interaction between matter and geometry. While $f(R, T)$ theory involves coupling between geometry and matter via the trace of energy-momentum tensor, the geometrical foundation differs as it is based on non-metricity in $f(Q, T)$ theory. This gives a distinct structure and different implications for cosmology and gravity. The $f(Q, T)$ does not involve an additional scalar field but modifies gravity directly through the non-metricity scalar. The coupling with T in $f(Q, T)$ introduces effective modifications similar to scalar fields but without introducing new fundamental fields. The $f(Q, T)$ theory can explain the late-time acceleration of the universe without invoking a cosmological constant or dark energy. The coupling between Q and T offers a novel approach to modifying gravity, allowing for an effective cosmological constant that varies with the matter content of the universe.

The study of WH geometry in modified theories has become subject of great interest for cosmologists in the recent years. Lobo et al [21] investigated traversable WH structures using various WSFs in $f(R)$ theory. Mazharimousavi and Halilsoy [22] discovered that WH solutions meet all required viability conditions near the WH throat in this theory. The traversable WH geometry through Noether symmetry in the context of scalar-tensor theory has been studied in [23]. The static WH solutions through ECs in $f(R, T)$ has been examined in [24]. In $f(R)$ theory, the viable WH solutions using

Noether symmetry have been studied in [25]. Mustafa et al [26] used the Karmarkar condition to examine the viable WH geometry in $f(Q)$ theory. Using the embedding class-I technique in $f(R)$ theory, Shamir and Fayyaz [27] created a WSF and found WH structure with a small amount of exotic matter. It was discovered that WH solutions which match the linear and exponential models of $f(Q)$ gravity models are stable and physically feasible [28]. The Karmarkar condition was applied in [29] to analyze traversable WH structure in $f(R)$ theory. Gul and his collaborators [30] developed WSF through Karmarkar constraint to examine the geometry of WH structures in different modified theories of gravity. Recently, the study of observational constraints in modified gravities discussed in [31]-[33].

This paper investigates viable traversable WH solutions using the embedding class-I technique in $f(R, G)$ theory. The analysis focuses on studying the behavior of shape function and ECs in this context. Wormholes are intriguing solutions to the Einstein field equations that have captured significant attention due to their implications in cosmology and interstellar travel. However, their viability and stability in the framework of alternative gravitational theories remain an open question. The motivation for exploring $f(R, G)$ gravity is twofold. First, this gravitational theory is an extension of GTR that allows for a more comprehensive description of gravitational phenomena. Second, WH solutions in $f(R, G)$ gravity offer new insights between gravity modifications and exotic structures like WHs . By investigating WHs in this modified gravity theory, we aim to contribute our understanding on the existence and stability of WH solutions. Furthermore, investigating WHs in this framework may shed light on the compatibility of WHs with modified gravity theories, which has implications for theoretical physics and observational cosmology.

To our knowledge, there has been limited exploration of WHs in the context of $f(Q, T)$ gravity. Our study takes a pioneering step in examining the existence and stability of WH solutions in this specific modified theory. Most previous research on WHs in modified theories focused on $f(Q)$ gravity. In contrast, our approach considers the joint effects of the non-metricity scalar and the trace of the stress-energy tensor in $f(Q, T)$ gravity, providing a more comprehensive analysis. We intend to perform a detailed viability analysis of the WH solutions in $f(Q, T)$ gravity, which is a novel aspect of our study. By conducting a comprehensive analysis of WH solutions in $f(Q, T)$ gravity, our work contributes to the broader understanding of gravitational theories and their astrophysical implications.

The literature mentioned above encourages us to investigate *WH* geometry in $f(Q, T)$ theory. This paper follows the following pattern. In section **2**, we provide the fundamental formulation of the field equations of $f(Q, T)$ theory. The field equations corresponding to Morris-Thorne spacetime in $f(Q, T)$ theory are developed in section **3**. The feasible *WH* geometry corresponding to several $f(Q, T)$ models with various shape functions and a constant redshift function is investigated in section **4**. Our results are summarized in the final section.

2 $f(Q, T)$ Theory-Basic Formalism

This section describes the basics of modified $f(Q, T)$ gravity and formulates the field equations using the variational method. Weyl [11] generalized *RG* based on the assumption that an arbitrary vector undergoes a change in length during parallel transport. Accordingly, the fundamental fields of Weyl space is represented by a new vector field (w^α) and the metric tensor. The expression $\delta l = l w_\alpha \delta x^\alpha$ [34] represents the change in length of a vector transported along an infinitesimal path in Weyl space. Additionally, the explanation for the variance in the vector length in Weyl space resulting from parallel transport is

$$\delta l = l(\nabla_\beta \omega_\alpha - \nabla_\alpha \omega_\beta) s^{\alpha\beta}. \quad (1)$$

If there is a local scaling length given by $\tilde{l} = \sigma(x)l$ then it changes the vector field w_α to $\tilde{w}_\alpha = w_\alpha + (ln\sigma)_{,\alpha}$. Moreover, the elements of the metric tensor transform under conformal transformations as $\tilde{g}_{\alpha\beta} = \sigma^2 g_{\alpha\beta}$ and $\tilde{g}^{\alpha\beta} = \sigma^{-2} g^{\alpha\beta}$ [35]. The Weyl geometry also contains a semi-metric connection, defined as

$$\bar{\Gamma}_{\alpha\beta}^\theta = \Gamma_{\alpha\beta}^\theta + g_{\alpha\beta} \omega^\theta - \delta_\alpha^\theta \omega_\beta - \delta_\beta^\theta \omega_\alpha, \quad (2)$$

where $\Gamma_{\alpha\beta}^\theta$ represents the Christoffel symbol. The gauge covariant derivative can be formulated using the fact that $\bar{\Gamma}_{\alpha\beta}^\theta$ is symmetric. The Weyl tensor is expressed as

$$\bar{R}_{\alpha\beta\gamma\eta} = \bar{R}_{(\alpha\beta)\gamma\eta} + \bar{R}_{[\alpha\beta]\gamma\eta}, \quad (3)$$

where

$$\begin{aligned} \bar{R}_{[\alpha\beta]\gamma\eta} &= R_{\alpha\beta\gamma\eta} + 2\nabla_\gamma \omega_{[\alpha} g_{\beta]\eta} + 2\nabla_\eta \omega_{[\beta} g_{\alpha]\gamma} + 2\omega_\gamma \omega_{[\alpha} g_{\beta]\eta} \\ &+ 2\omega_\eta \omega_{[\beta} g_{\alpha]\gamma} - 2\omega^2 g_\gamma [\alpha g_{\beta]}\eta, \end{aligned} \quad (4)$$

and

$$\bar{R}_{(\alpha\beta)\gamma\eta} = \frac{1}{2}(\bar{R}_{\alpha\beta\gamma\eta} + \bar{R}_{\beta\alpha\gamma\eta}) = g_{\alpha\beta}W_{\gamma\eta}. \quad (5)$$

The first contraction of the Weyl curvature tensor yields

$$\bar{R}^\alpha_\beta = \bar{R}^\gamma_\beta = R^\alpha_\beta + 2\omega^\alpha\omega_\beta + 3\nabla_\beta\omega^\alpha - \nabla_\alpha\omega^\beta + g^\alpha_\beta(\nabla_\gamma\omega^\gamma - 2\omega_\gamma\omega^\gamma). \quad (6)$$

Finally, the Weyl scalar is obtained as

$$\bar{R} = \bar{R}^\gamma_\gamma = R + 6(\nabla_\alpha\omega^\alpha - \omega_\alpha\omega^\alpha). \quad (7)$$

Compared to RG and Weyl geometry, Weyl-Cartan spaces with torsion provided a more generalized geometric framework. The term torsion describes the antisymmetric portion of the connection, which is a measurement of the connection twisting related to parallel transport. The law of parallel transport is given by $dv^\alpha = -v^\sigma\hat{\Gamma}^\alpha_{\sigma\beta}dx^\beta$ [36]. In Weyl-Cartan geometry, the connection is defined as

$$\hat{\Gamma}^\theta_{\alpha\beta} = \Gamma^\theta_{\alpha\beta} + C^\theta_{\alpha\beta} + L^\theta_{\alpha\beta}, \quad (8)$$

where

$$C^\theta_{\alpha\beta} = \hat{\Gamma}^\theta_{[\alpha\beta]} + g^{\theta\sigma}g_{\alpha\kappa}\hat{\Gamma}^\kappa_{[\beta\sigma]} + g^{\theta\sigma}g_{\beta\kappa}\hat{\Gamma}^\kappa_{[\alpha\sigma]}, \quad (9)$$

is the contortion tensor and the deformation tensor is expressed as

$$L^\theta_{\alpha\beta} = \frac{1}{2}g^{\theta\sigma}(Q_{\alpha\beta\sigma} + Q_{\beta\alpha\sigma} - Q_{\alpha\sigma\beta}), \quad (10)$$

with

$$Q_{\theta\alpha\beta} = -\frac{\partial g_{\alpha\beta}}{\partial\chi^\theta} + g_{\beta\sigma}\hat{\Gamma}^\sigma_{\alpha\theta} + g_{\sigma\alpha}\hat{\Gamma}^\sigma_{\beta\theta}. \quad (11)$$

Using Eq.(8) with $Q_{\theta\alpha\beta} = -2g_{\alpha\beta}\omega_\theta$, we have

$$\hat{\Gamma}^\theta_{\alpha\beta} = \Gamma^\theta_{\alpha\beta} + g_{\alpha\beta}\omega^\theta - \delta^\theta_\alpha\omega_\beta - \delta^\theta_\beta\omega_\alpha + C^\theta_{\alpha\beta}, \quad (12)$$

where

$$C^\theta_{\alpha\beta} = T^\theta_{\alpha\beta} - g^{\theta\eta}g_{\sigma\alpha}T^\sigma_{\eta\beta} - g^{\theta\eta}g_{\sigma\beta}T^\sigma_{\eta\alpha}. \quad (13)$$

The Weyl-Cartan tensor is given by

$$\hat{R}^\theta_{\alpha\beta\sigma} = \hat{\Gamma}^\theta_{\alpha\sigma,\beta} - \hat{\Gamma}^\theta_{\alpha\beta,\sigma} + \hat{\Gamma}^\gamma_{\alpha\sigma}\hat{\Gamma}^\theta_{\gamma\beta} - \hat{\Gamma}^\gamma_{\alpha\beta}\hat{\Gamma}^\theta_{\gamma\sigma}, \quad (14)$$

and

$$\begin{aligned}\hat{R} &= \hat{R}^{\alpha\beta}_{\alpha\beta} = R + 6\nabla_{\beta}\omega^{\beta} - 4\nabla_{\beta}T^{\beta} - 6\omega_{\beta}\omega^{\beta} + 8\omega_{\beta}T^{\beta} \\ &+ T^{\alpha\gamma\beta}T_{\alpha\gamma\beta} + 2T^{\alpha\gamma\beta}T_{\beta\gamma\alpha} - 4T_{\beta}T^{\beta}.\end{aligned}\quad (15)$$

By eliminating the boundary terms in the Ricci scalar, we can reformulate the gravitational action with coupling constant one as [37]

$$S = \frac{1}{2} \int g^{\alpha\beta} (\Gamma^{\gamma}_{\sigma\alpha} \Gamma^{\sigma}_{\beta\gamma} - \Gamma^{\gamma}_{\sigma\gamma} \Gamma^{\sigma}_{\alpha\beta}) \sqrt{-g} d^4x. \quad (16)$$

Using the assumption that the connection is symmetric, the gravitational action becomes

$$S = -\frac{1}{2} \int g^{\alpha\beta} (L^{\gamma}_{\sigma\alpha} L^{\sigma}_{\beta\gamma} - L^{\gamma}_{\sigma\gamma} \Gamma^{\sigma}_{\alpha\beta}) \sqrt{-g} d^4x. \quad (17)$$

This is the action of symmetric teleparallel gravity. However, these two gravitational theories differ in important ways. Because of the curvature tensor disappearance, the total spacetime geometry in the context of symmetric teleparallel gravity is flat. This leads to the adoption of the Weitzenbock configuration in the global geometry. Moreover, changes in the length of the vector during parallel transit cause gravitational effects rather than the rotation of the angle between the two vectors.

Now, we consider an extension of Eq.(17) as

$$S = \int \left(\frac{1}{2} f(Q, T) + L_M \right) \sqrt{-g} d^4x, \quad (18)$$

where

$$Q \equiv -g^{\alpha\beta} (L^{\gamma}_{\eta\alpha} L^{\eta}_{\beta\gamma} - L^{\gamma}_{\eta\gamma} L^{\eta}_{\alpha\beta}), \quad (19)$$

and

$$L^{\gamma}_{\eta\lambda} = -\frac{1}{2} g^{\gamma\theta} (\nabla_{\lambda} g_{\eta\theta} + \nabla_{\eta} g_{\theta\lambda} - \nabla_{\theta} g_{\eta\lambda}). \quad (20)$$

The superpotential is defined as

$$\begin{aligned}P^{\gamma}_{\alpha\beta} &\equiv \frac{1}{4} [-Q^{\gamma}_{\alpha\beta} + 2Q_{(\alpha}{}^{\gamma}{}_{\beta)} + Q^{\gamma} g_{\alpha\beta} - \hat{Q}^{\gamma} g_{\alpha\beta} - \delta^{\gamma}_{(\alpha} Q_{\beta)}] \\ &= -\frac{1}{2} L^{\gamma}_{\alpha\beta} + \frac{1}{4} (Q^{\gamma} - \tilde{Q}^{\gamma}) g_{\alpha\beta} - \frac{1}{4} \delta^{\gamma}_{(\alpha} Q_{\beta)}.\end{aligned}\quad (21)$$

We obtain the relation for non metricity as

$$Q = -Q_{\sigma\gamma\beta}P^{\sigma\gamma\beta} = -\frac{1}{4}(-Q^{\gamma\beta\rho}Q_{\gamma\beta\rho} + 2Q^{\gamma\beta\rho}Q_{\rho\gamma\beta} - 2Q^\rho\tilde{Q}_\rho + Q^\rho Q_\rho). \quad (22)$$

The calculation of this relation is given in [14]. The variation of Eq.(18) corresponding to the metric tensor yields

$$\begin{aligned} \delta S &= \int \frac{1}{2} \delta[f(Q, T)\sqrt{-g}] \delta[L_M \sqrt{-g}] d^4x, \\ &= \int \frac{1}{2} \left(-\frac{1}{2} f g_{\alpha\beta} \sqrt{-g} \delta g^{\alpha\beta} + f_Q \sqrt{-g} \delta Q + f_T \sqrt{-g} \delta T \right) \\ &\quad - \frac{1}{2} T_{\alpha\beta} \sqrt{-g} \delta g^{\alpha\beta} d^4x. \end{aligned} \quad (23)$$

The detailed variation of non metricity is provided in [14]. Furthermore, we define

$$T_{\alpha\beta} = -\frac{2}{\sqrt{-g}} \frac{(\delta\sqrt{-g}L_M)}{\delta g^{\alpha\nu}}, \quad \Theta_{\alpha\beta} \equiv g^{\gamma\eta} \frac{\delta T_{\gamma\eta}}{\delta g^{\alpha\beta}}, \quad (24)$$

which implies that $\delta T = \delta(T_{\alpha\beta}g^{\alpha\beta}) = (T_{\alpha\beta} + \Theta_{\alpha\beta})\delta g^{\alpha\beta}$. Thus, Eq.(24) becomes

$$\begin{aligned} \delta S &= \int \frac{1}{2} \left[-\frac{1}{2} f g_{\alpha\beta} \sqrt{-g} \delta g^{\alpha\beta} + f_T (T_{\alpha\beta} + \Theta_{\alpha\beta}) \sqrt{-g} \delta g^{\alpha\beta} \right. \\ &\quad \left. - f_Q \sqrt{-g} (P_{\alpha\gamma\eta} Q_\beta{}^{\gamma\eta} - 2Q^{\gamma\eta}{}_\alpha P_{\gamma\eta\beta}) \delta g^{\alpha\beta} + 2f_Q \sqrt{-g} P_{\gamma\alpha\beta} \nabla^\gamma \delta g^{\alpha\beta} \right] \\ &\quad - \frac{1}{2} T_{\alpha\beta} \sqrt{-g} \delta g^{\alpha\beta} d^4x. \end{aligned} \quad (25)$$

The resulting modified field equations are

$$\begin{aligned} T_{\alpha\beta} &= -\frac{2}{\sqrt{-g}} \nabla_\gamma (f_Q \sqrt{-g} P_{\alpha\beta}^\gamma) - \frac{1}{2} f g_{\alpha\beta} + f_T (T_{\alpha\beta} + \Theta_{\alpha\beta}) \\ &\quad - f_Q (P_{\alpha\gamma\eta} Q_\beta{}^{\gamma\eta} - 2Q^{\gamma\eta}{}_\alpha P_{\gamma\eta\beta}), \end{aligned} \quad (26)$$

where $f_T = \frac{\partial f}{\partial T}$ and $f_Q = \frac{\partial f}{\partial Q}$. This modified framework provides insights into the behavior of gravity through the solution of these field equations.

3 Wormhole and Energy Conditions

We consider the Morris-Thorne spacetime as [5]

$$ds^2 = dt^2 e^{2\mu(r)} - dr^2 \left(1 - \frac{\nu(r)}{r} \right)^{-1} - d\theta^2 r^2 - d\phi^2 r^2 \sin^2 \theta, \quad (27)$$

where $\mu(r)$ defines the redshift function and $\nu(r)$ represents the shape function. The given constraints must be fulfilled for a viable *WH* geometry.

- $\nu(r) < r$,
- $\nu(r_0) - r = 0$ at r_0 ,
- $\nu'(r_0) < 1$,
- $\frac{\nu(r)}{r} \rightarrow 0$ as $r \rightarrow \infty$,

where r_0 is the radius of *WH* throat. We assume fluid distribution as

$$T_{\alpha\beta} = (\rho + p_{\perp})u_{\alpha}u_{\beta} - p_{\perp}g_{\alpha\beta} + (p_r - p_{\perp})v_{\alpha}v_{\beta}, \quad (28)$$

where the four-velocity and four-vector are represented by the u_{α} and v_{α} , respectively. The matter-Lagrangian density is a fundamental concept in gravitational physics that describes the behavior and distribution of matter in a spacetime. When the matter distribution displays distinct characteristics along several spatial directions (anisotropic matter configuration), the important information can be gained by examining the particular matter-Lagrangian density. For anisotropic matter distribution, the well-known matter-Lagrangian density in the literature is considered as $L_M = -P = -\frac{p_r + 2p_{\perp}}{3}$ [38]-[40]. The expression of $\Theta_{\alpha\beta}$ can be expressed as

$$\Theta_{\alpha\beta} = -g_{\alpha\beta}P - 2T_{\alpha\beta}. \quad (29)$$

The non-metricity scalar corresponding to Moris-Throne spacetime turns out to be

$$Q = -\frac{\nu}{r^2} \left[\frac{r\nu' - \nu}{r(r - \nu)} + 2\mu' \right]. \quad (30)$$

Now, using Eqs.(26)-(30), the resulting field equations become

$$\begin{aligned} \rho &= \frac{1}{2r^3} \left[f_Q((2r - \nu)(r\nu' - \nu)(r - \nu)^{-1} + \nu(2r\mu' + 2)) + 2\nu r f_{QQ} f_Q \right. \\ &\quad \left. + f r^3 - 2r^3 f_T(P + \rho) \right], \\ p_r &= \frac{-1}{2r^3} \left[f_Q(\nu((r\nu' - \nu)(r - \nu)^{-1} + 2r\mu' + 2) - 4r\mu') + 2\nu r f_{QQ} f_Q \right] \end{aligned} \quad (31)$$

$$+ fr^3 + 2r^3 f_T(P - p_r) \Big], \quad (32)$$

$$p_\perp = \frac{-1}{4r^2} \left[f_Q \left\{ (r\nu' - \nu) \left(\frac{2r}{r - \nu} + 2r\mu' \right) (r)^{-1} + 4(2\nu - r)\mu' - 4r\mu'^2 \right. \right. \\ \left. \left. - 4r\mu'' \right\} - 4rf_{QQ}f_Q\mu' + 2fr^2r - \nu - 4r^2f_T(P - p_\perp) \right]. \quad (33)$$

The field equations are complicated because multivariate functions and their derivatives are present. We take a specific model with $f(T) = bT$ as

$$f(Q, T) = f(Q) + bT, \quad (34)$$

to simplify the field equations and obtain explicit expressions for the pressure components and energy density. Here, b is an arbitrary constant. The field equations (31)-(33) corresponding to this model become

$$\rho = \left[r^2 \{ 5br\mu'(2rQ'f_{QQ} - f_Q(\nu' - 4)) + r^2(10b\mu''f_Q - 3(b+1)f_Q) \right. \\ + 10br^2\mu'^2f_Q + 2f_Q(2b-3)\nu' \} + r\nu \{ -f_Q(20br^2\mu'' + 20br^2\mu'^2 \\ + r\mu'(-5b\nu' + 41b + 6) + (7b-3)\nu') - 2rQ'(10br\mu' - 2b + 3)f_{QQ} \\ + 3(b+1)r^2f_Q \} + \nu^2 \{ r(10br\mu''f_Q + \mu'(10br\mu' + 21b + 6)f_Q + 2Q' \\ \times (5br\mu' - 2b + 3)f_{QQ}) + 3(b+1)f_Q \} \Big] [6r^3(b+1)(2b-1) \\ \times (r - \nu)]^{-1}, \quad (35)$$

$$p_r = \left[r^2 \{ r\mu'(5b(f_Q\nu' - 2rQ'f_{QQ}) + 4f_Q(b-3)) + r^2(3(b+1)f_Q \right. \\ - 10b\mu''f_Q) - 10br^2f_Q\mu'^2 + 8b\nu'f_Q \} + r\nu \{ f_Q(20br^2\mu'' + 20br^2\mu'^2 \\ + r\mu'(-5b\nu' - 7b + 30) + (3-5b)\nu') + (6-12b)f_Q + 2rQ'f_{QQ} \\ \times (10br\mu' - 2b + 3) - 3(b+1)fr^2 \} - \nu^2 \{ f_Q(r(10br\mu'' + \mu'(10cr\mu' \\ - 3b + 18)) - 9b + 9) + 2rQ'f_{QQ}(5cr\mu' - 2b + 3) \} \Big] [6r^3(b+1) \\ \times (2b-1)(r - \nu)]^{-1}, \quad (36)$$

$$p_\perp = \left[r^2 \{ r\mu'(2f_{QQ}(b-3)rQ' - f_Q((b-3)\nu' + 8b + 6)) + r^2(2f_Q(b-3)\mu'' \right. \\ + 3(b+1)f) + 2f_Q(b-3)r^2\mu'^2 + f_Q(2b+3)\nu' \} + r\nu \{ f_Q(-4(b-3) \\ \times r^2\mu'' - 4(b-3)r^2\mu'^2 + r\mu'((b-3)\nu' + 23b + 15) + b\nu') + f_Q(6b-3) \\ + 4f_{QQ}rQ'(2b - (b-3)r\mu') - 3(b+1)nr^2 \} + \nu^2 \{ r(2f_Q(b-3)r\mu' \\ + f_Q\mu'(2(b-3)r\mu' - 3(5b+3)) + 2f_{QQ}Q'((b-3)r\mu' - 4b)) - 9f_Qb \} \Big] \\ \times [6r^3(b+1)(2b-1)(r - \nu)]^{-1}. \quad (37)$$

In the context of *WH* studies, the redshift function plays a crucial role in determining the gravitational redshift experienced by signals traveling through the *WH*. Physically, it represents how the gravitational potential changes along the radial coordinate of the *WH* throat and its surrounding geometry. When the redshift function is constant, it implies that there is no gravitational redshift experienced by signals passing through the *WH*. This is often a simplifying assumption in many *WH* models to avoid singularities at the throat and to ease the analysis of the spacetime geometry.

- Physical Meaning of a Constant Redshift Function

A constant redshift function ensures that the *WH* spacetime does not have an event horizon. Event horizons are typically associated with divergent redshift functions (as in the case of black holes), where signals cannot escape beyond a certain region. With constant redshift, signals can travel freely through the *WH*, making it a traversable *WH*. Since redshift function is related to the gravitational time dilation, a constant value indicates that there is no differential time dilation across the *WH* geometry. This can be important in constructing traversable *WHs* that are stable for interstellar travel, as travelers moving through the *WH* would not experience time distortion.

- Impact on the Modified Field Equations

When the redshift function is constant, it simplifies the Einstein field equations or any modified versions of these equations that govern the *WH* spacetime. The modified Einstein field equations often contain derivatives of redshift function. When redshift function is constant, terms involving derivatives of redshift vanish, significantly reducing the complexity of the equations. In theories of modified gravity, a constant redshift function might change the way curvature terms interact with the geometry, affecting the required modifications to the field equations. The lack of a varying potential could reduce the complexity of the additional terms introduced by these theories, leading to more manageable conditions for finding solutions. Thus, a constant redshift function in *WH* studies simplifies the field equations by eliminating time dilation effects and reducing the complexity of the spacetime metric. It affects the physical characteristics of the *WH*, such as the absence of an event horizon and simplified matter-energy conditions needed to sustain the *WH* structure.

Astrophysical objects are composed of a variety of materials. Determining the kind of substance (ordinary or exotic) that is contained in celestial objects

is essential. In order to comprehend the nature of matter in the cosmic objects, we look into certain inequalities, known as *ECs*. These constraints are useful in confirming the viability of *WHs*. The *ECs* for the anisotropic configuration are expressed as null energy bound ($0 \leq p_r + \rho$, $0 \leq p_\perp + \rho$), dominant energy bound ($0 \leq \rho \pm p_r$, $0 \leq \rho \pm p_\perp$), weak energy bound ($0 \leq p_r + \rho$, $0 \leq p_\perp + \rho$, $0 \leq \rho$) and strong energy bound ($0 \leq p_r + \rho$, $0 \leq p_\perp + \rho$, $0 \leq p_r + 2p_\perp + \rho$). The existence of traversable *WH* geometries and other hypothetical objects in spacetime requires a comprehension of these *ECs*. A feasible *WH* structure must diverge from these conditions.

3.1 Viable $f(Q, T)$ Models

Here, we examine the effects of several models of $f(Q, T)$ theory on the geometry of *WH*. Our work attempts to reveal obscure astrophysical and theoretical cosmological insights. The correction terms of this modified gravitational theory could produce insightful findings. The existence of feasible cosmic geometries could be significantly impacted by these modified terms. Therefore, exploring $f(Q, T)$ becomes crucial in identifying hypothetical objects. We examine three different models of $f(Q, T)$ in the following subsections.

Model 1

In this context, we examine a power-law $f(Q, T)$ model with arbitrary constant a_1 as

$$f(Q, T) = Q + a_1 Q^2 + bT. \quad (38)$$

This model has important cosmological implications, particularly in explaining early universe dynamics and late-time cosmic acceleration. It provides a framework for exploring alternatives to dark energy and possibly modifying gravitational wave predictions, making it a significant candidate for describing the cosmic evolution. The quadratic correction $a_1 Q^2$ introduces non-linearities in the gravitational field equations, which could have significant cosmological implications. Such quadratic terms are known to affect the early universe dynamics and they may lead to the possibility of a cosmological bounce, avoiding the singularity problem of the Big Bang. This term could also provide corrections to the late-time evolution of the universe, influencing cosmic acceleration or mimicking dark energy effects. The coupling of the trace of the energy-momentum tensor with gravity represents

a direct interaction between matter and geometry. This interaction implies that the energy content of the universe could influence the gravitational field in ways different from *GTR*. The term bT is expected to impact the equation of state of cosmological fluids, possibly leading to modifications in how matter and radiation evolve in the universe. It could offer explanations for cosmic acceleration without the need for dark energy by introducing effective pressure contributions arising from the coupling term. The combination of non-metricity and matter-geometry coupling introduces new dynamics in the cosmic evolution. Depending on the values of the parameters, this model could lead to deviations from the standard cosmological model, influencing the structure formation and the overall dynamics of cosmic expansion. We consider the model parameters values as $a_1 = 0.7$ and $b = 0.9$ to examine the viable *WH* solutions.

The field equations corresponding to this model are provided in Appendix **A**. We now look at an interesting case with a redshift function that is constant. The derivation of exact *WH* solutions is made possible by this simplification, which also significantly simplifies computations. In order to investigate the feasible *WH* geometry, we take into account the various shape functions in the following scenarios. In all the cases, we take *WH* throat at 0.5, i.e., $r_0 = 0.5$ for our convenience. It is important to mention that the all Morris-Thorne conditions are satisfied at $r_0 = 0.5$.

Case 1: $\nu(r) = \frac{r_0^2}{r}$

We consider the specific choice of shape function as $\nu(r) = \frac{r_0^2}{r}$ [41]. The field equations corresponding to this case turn out to be

$$\begin{aligned}\rho &= \frac{2a_1r_0^6((23b - 27)r_0^2 + 14(3 - 2b)r^2) - (2b - 3)r^4(r_0^3 - r_0r^2)^2}{3r^8(2b^2 + b - 1)(r_0^2 - r^2)^2}, \\ p_r &= \frac{2a_1r_0^6((b + 15)r_0^2 + 2(2b - 15)r^2) - (10b - 3)r^4(r_0^3 - r_0r^2)^2}{3r^8(2b^2 + b - 1)(r_0^2 - r^2)^2}, \\ p_\perp &= \frac{(2b - 3)r^4(r_0^3 - r_0r^2)^2 + 2a_1r_0^6((25b + 3)r_0^2 - 2(22b + 3)r^2)}{3(2b^2 + b - 1)r^8(r_0^2 - r^2)^2}.\end{aligned}$$

The graphical behavior of the *ECs* for different values of a_1 and b is shown in Figure 1. The behavior of *ECs* is examined in the upper panel for small values of the model parameters. The matter contents ρ , $\rho - p_r$ and $\rho - p_\perp$ in the left plot show negative behavior which violates the dominant and weak




Figure 1: Graphs of $\rho + p_r$ (blue), $\rho + p_\perp$ (green), $\rho - p_r$ (red), $\rho - p_\perp$ (black) and ρ (cyan) corresponding to Model 1 (Case 1) for different parametric values.

ECs, whereas all *ECs* are violated in the right plot. It is clear from the lower panel that for both large positive and negative parametric values, the dominant *EC* is violated. These graphs show that the fluid variables violate the *ECs*, which provides the feasible traversable *WH* structures in this case.

Case 2: $\nu(r) = r e^{r_0 - r}$

The field equations with respect to $\nu(r) = r e^{r_0 - r}$ [41] and constant redshift are

$$\rho = \frac{1}{6r^2(2b^2 + b - 1)(e^{r_0} - e^r)^2} \left[e^{r_0 - 2r} (a_1 e^{2r_0} ((19b - 21)e^{r_0} + (36 - 24$$

$$\begin{aligned}
& \times b)e^r) - 2e^r(2b - 3)(r - 1)(e^{r_0} - e^r)^2)], \\
p_r &= \frac{1}{6r^2(2b^2 + b - 1)(e^{r_0} - e^r)^2} [e^{r_0-2r} (a_1 e^{2r_0} ((5b + 9)e^{r_0} - 24e^r) - 2e^r \\
& \times (b(4r + 2) - 3)(e^{r_0} - e^r)^2)], \\
p_\perp &= \frac{1}{6r^2(2b^2 + b - 1)(e^{r_0} - e^r)^2} [e^{r_0-2r} (a_1 e^{2r_0} ((17b + 3)e^{r_0} - 6(6b + 1) \\
& \times e^r) - e^r(2b(r - 4) + 3r)(e^{r_0} - e^r)^2)].
\end{aligned}$$

The graphical representation of the *ECs* for various values of a_1 and b corresponding to case 2 is shown in Figure 2. It can be seen in the upper and lower panels that we obtain the identical behavior as found in case 1., i.e., the matter contents $(\rho - p_r, \rho - p_\perp, \rho)$ behave negatively in the upper panel and dominant *EC* is violated in the lower panel. Thus, we find that the viable traversable *WH* structures exist for both positive and negative values of the model parameters.

Case 3: $\nu(r) = r_0 e^{1-\frac{r}{r_0}}$

The corresponding field equations for $\nu(r) = r_0 e^{1-\frac{r}{r_0}}$ [42] are

$$\begin{aligned}
\rho &= \frac{1}{6(2b^2 + b - 1)r^6(e^{r_0} - re^{r/r_0})^2} [e^{1-\frac{2r}{r_0}} \{e^2 a_1 r_0 (e^{r_0} (3(9b - 11)r_0^2 + 2 \\
& \times (19b - 21)r_0 r + (19b - 21)r^2) - 4(2b - 3)re^{r/r_0}(4r_0^2 + 6r_0 r + 3r^2)) \\
& - 2(2b - 3)r^4 e^{r/r_0} (e^{r_0} - re^{r/r_0})^2 \}], \\
p_r &= \frac{1}{6(2b^2 + b - 1)r^6(e^{r_0} - re^{r/r_0})^2} [e^{1-\frac{2r}{r_0}} \{e^2 a_1 r_0 (e^{r_0} (-3(b - 7)r_0^2 + 2 \\
& \times (5b + 9)r_0 r + (5b + 9)r^2) - 4re^{r/r_0}((9 - 2b)r_0^2 + 12r_0 r + 6r^2)) - 2 \\
& \times r^3 e^{r/r_0} (e^{r_0} - re^{r/r_0})^2 ((6b - 3)r_0 + 4br) \}], \\
p_\perp &= \frac{1}{6(2b^2 + b - 1)r^6(e^{r_0} - re^{r/r_0})^2} [e^{1-\frac{2r}{r_0}} \{e^2 a_1 r_0 (e^{r_0} ((33b + 3)r_0^2 + 2 \\
& \times (17b + 3)r_0 r + (17b + 3)r^2) - 2re^{r/r_0}((26b + 3)r_0^2 + 6r_0(6br + r) \\
& + 3(6b + 1)r^2)) - r^3 e^{r/r_0} (e^{r_0} - re^{r/r_0})^2 ((3 - 6b)r_0 + (2b + 3)r) \}].
\end{aligned}$$

Figure 3 determines that the matter components $(\rho, \rho \pm p_r, \rho \pm p_\perp)$ exhibit negativity across all parametric values. This deviation from *ECs* suggests the presence of exotic matter, providing justification for the feasibility of a traversable *WH* geometry in this gravitational model.




Figure 2: Graphs of $\rho + p_r$ (blue), $\rho + p_\perp$ (green), $\rho - p_r$ (red), $\rho - p_\perp$ (black) and ρ (cyan) corresponding to Model 1 (Case 2) for different parametric values.




Figure 3: Graphs of $\rho + p_r$ (blue), $\rho + p_\perp$ (green), $\rho - p_r$ (red), $\rho - p_\perp$ (black) and ρ (cyan) corresponding to Model 1 (Case 3) for different parametric values.

Case 4: $\nu(r) = r \frac{\ln r + 1}{r_0 + 1}$

In this specific case [43], we obtain the field equations as

$$\begin{aligned}
\rho &= \frac{1}{6(2b^2 + b - 1)(r_0 + 1)^4(r + 1)^2 \left(r - \frac{r \ln(r+1)}{r_0+1}\right)^2} \left[a_1 \ln(r + 1)(\ln(r + 1)(8 \right. \\
&\times br_0 + (12 - 8b) \ln(r + 1) + 19b - 12r_0 - 21) - 8(2b - 3)(r_0 + 1)) + 2(2 \\
&\times b - 3)(r_0 + 1)(r + 1)(r + (r + 1) \ln(r + 1))(r_0 - \ln(r + 1) + 1)^2 \Big], \\
p_r &= \frac{1}{6(2b^2 + b - 1)(g + 1)^2 r^2 (r + 1)^2 (r_0 - \ln(r + 1) + 1)^2} \left[a_1 \ln(r + 1)(\ln \right. \\
&\times (r + 1)(-8br_0 + 4(2b - 3) \ln(r + 1) + 5b + 12r_0 + 9) - 4(2b + 3)(r_0 \\
&+ 1) - 2(r_0 + 1)(r + 1)((2b - 3)(r + 1) \ln(r + 1) - 4br)(r_0 - \ln(r + 1) \\
&+ 1)^2 \Big], \\
p_\perp &= \frac{1}{6(2b^2 + b - 1)(r_0 + 1)^2 r^2 (r + 1)^2 (r_0 - \ln(r + 1) + 1)^2} \left[(r_0 + 1)(r \right. \\
&+ 1)((2b + 3)r + 8b(r + 1) \ln(r + 1))(r_0 - \ln(r + 1) + 1)^2 - a_1 \ln(r \\
&+ 1)(\ln(r + 1)(-16br_0 + 16b \ln(r + 1) - 17b - 3) + 2(10b + 3)(r_0 \\
&+ 1)) \Big].
\end{aligned}$$

Figure 4 violates the *ECs* which manifests the existence of viable traversable *WH* geometry in this case.

Model 2

Here, we use another model with constant a_2 as [41]

$$f(Q, T) = Q + \frac{a_2}{Q} + bT, \quad (39)$$

This model has notable cosmological significance, particularly in addressing key issues like the early universe evolution, late-time cosmic acceleration and potential avoidance of singularities. Its impact on the early universe and the late-time accelerated expansion make it a candidate for explaining phenomena that the standard Λ CDM model might not fully capture. Additionally, observational signatures such as deviations in gravitational wave propagation or cosmic structure growth could help test the viability of this model. The term $\frac{a_2}{Q}$ introduces a non-linear modification to the non-metricity, which can have significant implications for the cosmological dynamics, particularly




Figure 4: Graphs of $\rho + p_r$ (blue), $\rho + p_\perp$ (green), $\rho - p_r$ (red), $\rho - p_\perp$ (black) and ρ (cyan) corresponding to Model 1 (Case 4) for different parametric values.

in the early universe and at large scales. The presence of this inverse term suggests that when Q becomes small (as it might near a bounce or at high-curvature regions), the dynamics become strongly modified. This term could contribute to avoiding cosmological singularities, leading to bouncing solutions or early-time inflation. The presence of the inverse term could also help in avoiding future singularities such as the Big Rip or other catastrophic fates, by altering the dynamics when Q becomes small. This could stabilize the cosmological evolution and lead to a more regular, non-singular universe. We consider $a_2 = 0.8$ and $b = 0.9$ to analyze the viable WH structures.

The modified field equations for this model are given in Appendix **B**. In this model, the graphical representations corresponding to cases **1-3** indicate the non-consistent behavior of the fluid parameters for both positive as well as negative values of the model parameters. This suggests that this model does not support the geometry of WH corresponding to specific shape functions given in the cases **1-3**.

Case 4: $\nu(r) = r \frac{\ln r + 1}{r_0 + 1}$

This case yields the field equations in the following form

$$\begin{aligned}
\rho &= -\frac{1}{6r^5(2b^2 + b - 1)\ln^2(r + 1)} \left[(r_0 + 1)(r + 1) \{ r^5(r_0 - \ln(r + 1) + 1) \right. \\
&\times (a_2(\ln(r + 1)(r(20br_0 + 13b - 30r_0 - 27) + 6(2b - 3)(r_0 + 1) - 2(2b - 3)(5r + 3)\ln(r + 1)) - 2(2b - 3)(r_0 + 1)r - 3a_2(b + 1)r\ln(r + 1)) \\
&\left. - \left(\frac{1}{(r_0 + 1)^2(r + 1)^2} (2(2b - 3)r^3\ln^2(r + 1)(r + (r + 1)\ln(r + 1))) \right) \right], \\
p_r &= -\frac{1}{6r^2(2b^2 + b - 1)(r_0 + 1)(r + 1)\ln^2(r + 1)} \left[2\ln^2(r + 1)((2b - 3) \right. \\
&\times (r + 1)\ln(r + 1) - 4br) - (r_0 + 1)^2r^2(r + 1)^2(r_0 - \ln(r + 1) + 1) \\
&\times (a_2(\ln(r + 1)(r(5b(4r_0 + 5) - 30r_0 - 33) + 6(2b - 3)(r_0 + 1) - 2(2b - 3)(5r + 3)\ln(r + 1)) - 4(4b - 3)(r_0 + 1)r - 3a_2(b + 1)r\ln(r + 1)) \left. \right], \\
p_\perp &= -\frac{1}{6r^2(2b^2 + b - 1)(r_0 + 1)(r + 1)\ln^2(r + 1)} \left[(r_0 + 1)^2r^2(r + 1)^2(r_0 \right. \\
&- \ln(r + 1) + 1)(a_2(b\ln(r + 1)(8r_0(5r + 3) + 41r - 8(5r + 3)\ln(r + 1) + 24) - (14b - 3)(r_0 + 1)r + 3a_2(b + 1)r\ln(r + 1)) + \ln^2(r + 1) \\
&\left. + 1)(-(2b + 3)r - 8b(r + 1)\ln(r + 1)) \right].
\end{aligned}$$




Figure 5: Graphs of $\rho+p_r$ (blue), $\rho+p_\perp$ (green), $\rho-p_r$ (red), $\rho-p_\perp$ (black) and ρ (cyan) corresponding to Model 2(Case 4) for different parametric values.

In Figure 8, the upper panels illustrate the negative trends of ρ , $\rho + p_r$ and $\rho - p_\perp$, indicating the violation of the *ECs*. Similarly, all *ECs* are observed to be violated in the lower panel. These graphical representations indicate that the fluid parameters violate the *ECs* across positive and negative values of model parameters. This suggests the presence of viable traversable *WH* structures in this framework.

Model 3

Finally, we consider [41]

$$f(Q, T) = Qe^{\frac{a_3}{Q}} + bT. \quad (40)$$

This mode has potential cosmological significance. Its non-linear modification to gravity through the non-metricity scalar can provide rich dynam-

ics that go beyond *GTR*. Additionally, the coupling to the trace of energy-momentum tensor lead to novel effects in the matter-gravity interaction. This model could be relevant for addressing key cosmological issues like the nature of dark energy, the origin of cosmic inflation, or the behavior of gravity at large scales. The term $Qe^{\frac{a_3}{Q}}$ introduces a non-linear interaction of Q , which might give rise to interesting cosmological effects. The exponential function suggests that the theory introduces corrections to *GTR* that become significant in regimes where Q is small, i.e., in the early or late universe when the geometry is evolving rapidly. The exponential term could play a role in the early universe. This might result in inflationary dynamics or provide an alternative to scalar field inflation models by driving an accelerated expansion. Depending on the values of the model might admit solutions that describe accelerating expansion, either in the early universe (inflation) or at late times (cosmic acceleration). For viable *WH* geometry, we assume the constant values as $a_3 = 0.8$ and $b = 0.9$.

The corresponding field equations for this model are bestowed in Appendix C. In this model, the graphical representation of the matter components $(\rho, \rho \pm p_r, \rho \pm p_\perp)$ indicate that *WH* does not exist corresponding to this model for all cases. Hence, it is found that this gravity model does not support the existence of a viable traversable *WH* structures for all cases. Thus, we conclude that this $f(Q, T)$ model is not suitable for *WH* solution.

4 Final Remarks

The exploration of *WH* structures holds paramount importance in the realm of astrophysics. Exotic matter is essential for physically viable *WH* geometries. Modified theories of gravity have drawn a lot of attention from the scientific community in recent decades as a potential replacement for *GTR*. These modified theories examine the viability of traversable *WH* geometries by the correction term ability to violate *ECs* and guarantee the existence of viable *WH* structures. Modified theories are not just important for theoretical conjecture, but they are essential for revealing the physical characteristics of celestial objects. In order to investigate the feasible *WH* configurations in the context of $f(Q, T)$ theory, we have considered the Morris-Throne space-time with anisotropic matter configuration. To get the explicit expressions of the pressure and energy density components, we have assumed the various models of this modified theory. The viability of *WH* structures was then

examined using *ECs* with various shape functions and a constant redshift function.

In this manuscript, we have investigated the feasibility of traversable *WH* solutions in the framework of $f(Q, T)$ theory. Our primary aim is to examine the theoretical aspects of *WH* solutions and their behavior under modified terms. Traversable *WHs* have garnered significant attention in theoretical physics due to their potential to connect different regions of spacetime. Identifying viable and stable traversable *WH* solutions in $f(Q, T)$ theory is not just a theoretical exercise rather, it holds profound implications for our understanding of the fundamental laws governing the universe. Firstly, our findings suggest that traversable *WHs* can exist in modified gravity theories such as $f(Q, T)$ theory. This challenges the concept that such structures are only theoretically feasible in the framework of *GTR*. By demonstrating their existence in this modified framework, we have expanded the scope for studying and investigating these intriguing phenomena with a wider range of theoretical frameworks than previously considered. Thus, our research advances the understanding of traversable *WHs* in a modified framework and lays a theoretical foundation for further exploration.

Traversable *WHs* represent a captivating subject in theoretical physics, particularly in the framework of $f(Q, T)$ theory. The exploration of *WHs* holds significant relevance and theoretical significance. Their existence introduces the intriguing prospect of time travel by enabling the formation of closed timelike curves. In the context of $f(Q, T)$ gravitational theory, the presence of traversable *WHs* challenges conventional notions of spacetime geometry. It implies that deviations from the standard Einstein-Hilbert action can yield exotic structures like *WHs*, thereby opening up new avenues for investigating the foundational principles of physics. The analysis of traversable *WHs* offers valuable insights into the impact of modified gravity on the large-scale structure of the universe. This contributes to refining our models of cosmic evolution and understanding the eventual fate of the universe. Moreover, the existence of viable traversable *WHs* in $f(Q, T)$ theory facilitates a deeper exploration of the *ECs* governing spacetime and their compatibility with exotic matter. Consequently, the practical implications of traversable *WHs* in $f(Q, T)$ theory have the potential to revolutionize both space travel and cosmology, while their theoretical ramifications reshape our comprehension of fundamental physics, encompassing gravity, spacetime and the essence of the cosmos.

For both positive and negative values of the model parameters, our analy-

sis in the first model shows the violation of *ECs* which indicates the presence of exotic matter at the *WH* throat. The dominate *EC* is found to be violated for all values of a_1 and b for the cases 1 and 2. This gives a traversable *WH* structure in this scenario (Figures 1 and 2). For the cases 3 and 4, we also obtain the feasible traversable *WH* geometry (Figures 3 and 4). The traversable *WH* geometry in the model 2 is only obtained for case 4 as the existence of exotic matter at the *WH* throat is ensured by the negative trends of fluid parameters, which indicate a violation of the *ECs* (Figures 5). We have noted that the *WH* geometry for model 2 is not supported by the behavior of matter variables corresponding to cases 1-3. Additionally, it is discovered that the *WH* geometry corresponding to all cases is not supported by model 3. Thus, the only model that supports the presence of a feasible traversable *WH* structure is $f(Q, T) = Q + a_1 Q^2 + bT$.

The geometric features of *WHs* in $f(R)$ gravity were studied by Lobo and Oliveira in [44], who found that there are no feasible *WH* solutions in vacuum. In $f(R)$ theory, Fayyaz and Shamir [45] investigated physically realistic traversable *WH* configurations using the Karmarkar condition. They found viable solutions with minimum exotic matter. In order to investigate *WHs* in $f(R, T)$ theory, Sharif and Fatima [46] found *WH* solutions for minimum radius. According to Banerjee et al [47], the power-law model $f(Q) = Q + a_1 Q^2$ does not have any *WH* solution. It is worthwhile to mention here that we have obtained viable *WH* solutions in $f(Q, T)$ theory corresponding to the functional form of $f(Q) = Q + a_1 Q^2$ in which Banerjee et al [47] found that the viable *WH* solution does not exist corresponding to this functional form.

The observational evidence for *WHs* presents a promising avenue for advancing our comprehension of the cosmos. To validate *WH* solutions, future studies and experiments could consider the following directions.

- Gravitational Lensing

Investigate the detection of distinct gravitational lensing signatures linked to traversable *WHs*. Analyzing lensing effects on background light sources might unveil characteristic patterns indicative of *WH* presence.

- Multi-Messenger Astronomy

Analyze gravitational wave observations with various forms of electromagnetic radiation such as gamma-ray bursts or neutrino detections.

Coordinated multi-messenger observations could furnish a comprehensive dataset for identifying viable *WH* events.

- Cosmic Microwave Background

Examine anomalies in the cosmic microwave background that may be attributed to the influence of nearby *WHs*. Deviations in temperature or polarization patterns could serve as indirect indicators of these exotic structures.

- Astrophysical Disk Dynamics

Investigate the dynamics of accretion disks around black holes to detect deviations that could be linked to the presence of *WHs*. Variations in disk behavior might offer indirect evidence of nearby traversable *WHs*.

- High-Energy Particle Colliders

Explore the viability of probing the microscopic scale for evidence of exotic matter or energy consistent with the theoretical requirements for stabilizing and sustaining *WHs*. Particle accelerator experiments may provide insights into the fundamental physics associated with these structures.

- Time-Delay Observations

Conduct precise timing observations of distant astrophysical events, such as gamma-ray bursts, and look for unexpected time delays that could be attributed to gravitational effects associated with traversable *WHs*.

These proposed avenues for future research aim to inspire the scientific community to explore diverse observational methods, fostering a viable approach to validating the existence of *WHs* and advancing our understanding of the fundamental nature of the universe.

Appendix A: Field Equations (Model 1)

Using Eqs.(35)-(37), we obtain the following field equations corresponding to model 1 as

$$\begin{aligned}
 \rho &= \frac{1}{6r^6(2b^2 + b - 1)(r - \nu)^2} \left[r^4(r - \nu)^2(10br(r - \nu)\mu'' - 5b\mu'(r(\nu' - 4) \right. \\
 &+ 3\nu) + 10br(r - \nu)\mu'^2 + (4b - 6)\nu') + a_1\{ - 20br^5\mu'\nu'(2r\mu' + \nu') - 2 \\
 &\times r^3\nu(20br^3\mu'^3 + 2\nu'(5br^2\mu'' + (4b - 6)\nu') - 60br^2\mu'^2\nu' + r\mu'(10br(4r \\
 &\times \mu'' + \nu'') - 15b\nu'^2 - 2(7b + 12)\nu')) + r^2\nu^2\{120br^3\mu'^3 + 8\nu'(5br^2\mu'' \\
 &+ 6b - 9) + 12r^2\mu'^2(-10b\nu' + b + 1) + 4r((b + 6)r\mu'' + (3 - 2b)\nu'') + 2 \\
 &\times r\mu'(20br(6r\mu'' + \nu'') - 5b\nu'^2 - 6(2b + 7)\nu' - 4(b + 6)) + (11b - 9)\nu'^2\} \\
 &- 2r\nu^3(60br^3\mu'^3 + \nu'(10br^2\mu'' + 19b - 21) + 4(b + 6)r^2\mu'' + 4r^2\mu'^2(-5b \\
 &\times \nu' + 3b + 3) + r\mu'(10br(12r\mu'' + \nu'') + 2(b - 9)\nu' + 3(b - 14)) + 2(3 \\
 &- 2b)r\nu'' + 8(2b - 3)) + \nu^4(2r(\mu'(2r\mu'(10br\mu' + 3b + 3) + 7b - 18) \\
 &+ 2r\mu''(20br\mu' + b + 6)) + 27b - 33)\} \Big], \\
 p_r &= \frac{1}{6r^6(2b^2 + b - 1)(r - \nu)^2} \left[(-10b\mu''r^3 + 8b\nu'r - 10b\mu'^2(r^2 - 4a_1\nu')r \right. \\
 &+ \mu'(4(b - 3)r^2 + 5b\nu'(r^2 + 4a_1\nu')))r^5 + \nu(40br^3a_1\mu'^3 + 6r^2(5br^2 - 4a_1 \\
 &\times (5\nu'b + 4b - 2))\mu'^2 + r(-30ba_1\nu'^2 - 2b(5r^2 + 62a_1)\nu' + r((36 - 17 \\
 &\times b)r + 20ba_1(4r\mu'' + \nu''))) \mu' - 4(2b + 3)a_1\nu'^2 + 6r^2(5b\mu''r^2 - 2b + 1) \\
 &+ 4br^2\nu'(5a_1\mu'' - 4)r^3 - \nu^2(120br^3a_1\mu'^3 + 6r^2(5br^2 + a_1(-20\nu'b - 46 \\
 &\times b + 26))\mu'^2 + r((-5\nu'b - 22b + 36)r^2 + 2a_1(-5b\nu'^2 + (6 - 108b)\nu' \\
 &+ 4b(5r(6r\mu'' + \nu'') - 13)))\mu' + (3 - 13b)a_1\nu'^2 + 8\nu'(a_1(5br^2\mu'' - 6) \\
 &- br^2) + 2r(6(1 - 2b)r + (15br^2 + 2(b + 6)a_1)\mu''r + 2(3 - 2b)a_1\nu''))r^2 \\
 &+ \nu^3((-12b + r(\mu'(10r\mu'b - 9b + 12) + 10br\mu'') + 6)r^2 + 2a_1(60br^3\mu'^3 \\
 &- 4r^2(5\nu'b + 33b - 21)\mu'^2 + r(10r(12r\mu'' + \nu'')b - 93b + (6 - 46b)\nu' \\
 &+ 6)\mu' + 4b + \nu'(10b\mu''r^2 - 5b - 9) + 2r(2(b + 6)r\mu'' + (3 - 2b)\nu'') \\
 &- 18))r + a_1\nu^4(-4(20r\mu'b + b + 6)\mu''r^2 - 2\mu'(-41b + 2r\mu'(10r\mu'b \\
 &- 21b + 15) + 6)r - 3(b - 7)) \Big],
 \end{aligned}$$

$$\begin{aligned}
p_{\perp} = & \frac{1}{6(2b^2 + b - 1)r^6(r - \nu)^2} \left[(2(b - 3)\mu''r^3 + (2b + 3)\nu'r + 2(b - 3)\mu'^2(r^2 \right. \\
& - 4a_1\nu')r - \mu'(2(4b + 3)r^2 + (b - 3)\nu'(r^2 + 4a_1\nu'))r^5 - \nu(8(b - 3)r^3a_1 \\
& \times \mu'^3 + 6r^2((b - 3)r^2 - 4a_1(2b + (b - 3)\nu' - 1))\mu'^2 + r(-6(b - 3)a_1\nu'^2 \\
& + (4(b + 15)a_1 - 2(b - 3)r^2)\nu' + r(4(b - 3)a_1(4r\mu'' + \nu'') - 5(5b + 3) \\
& \times r))\mu' + 2(10b + 3)a_1\nu'^2 + 3r^2(2(b - 3)\mu''r^2 - 2b + 1) + 2r^2\nu'(2b + 2 \\
& \times (b - 3)a_1\mu'' + 3))r^3 + \nu^2(24(b - 3)r^3a_1\mu'^3 + 6r^2((b - 3)r^2 + 2a_1(-13 \\
& \times b - 2(b - 3)\nu' + 5))\mu'^2 + r(2a_1(-(b - 3)\nu'(r)^2 + 48\nu' + 4(b + (b - 3) \\
& \times r(6r\mu'' + \nu'') + 6)) - r^2(26b + (b - 3)\nu' + 12))\mu' + (b + 3)a_1\nu'^2 + \nu' \\
& \times ((2b + 3)r^2 + 4a_1(2(b - 3)\mu''r^2 + 18b + 3)) + 2r(r(-6b + (3(b - 3) \\
& \times r^2 - 2(7b + 3)a_1)\mu'' + 3) - 8ba_1\nu''))r^2 + \nu^3(r^2(6b + r(\mu'(9b - 2(b \\
& - 3)r\mu' + 3) - 2(b - 3)r\mu'')) - 3) - 2a_1(12(b - 3)r^3\mu'^3 - 4r^2(21b + (b \\
& - 3)\nu' - 6)\mu'^2 + r(3(b + 13) - 2(b - 9)\nu' + 2(b - 3)r(12r\mu'' + \nu''))\mu' \\
& + 26b - 4(7b + 3)r^2\mu'' + \nu'(2(b - 3)\mu''r^2 + 17b + 3) - 8br\nu'' + 3))r \\
& + a_1\nu^4(33b + 2r(\mu'(-b + 2r\mu'(-15b + 2(b - 3)r\mu' + 3) + 15) + 2r \\
& \times (-7b + 4(b - 3)r\mu' - 3)\mu'')) + 3) \left. \right].
\end{aligned}$$

Appendix B: Field Equations (Model 2)

The resulting field equations corresponding to model 2 are

$$\begin{aligned}
\rho = & \frac{1}{6(2b^2 + b - 1)r^2\nu^3(\nu(2r\mu' + 1) - r(2r\mu' + \nu'))^3} \left[a_2r^5(r - \nu)(-20br^5 \right. \\
& \times \mu'\nu'(2r\mu' + \nu') + r^3\nu(20br^3\mu'^3 + 2\nu'(5br^2\mu'' + (3 - 2b)\nu') + 120br^2\mu'^2 \\
& \times (\nu' + 1) + r\mu'(-20br(r\mu'' + \nu'') + 15b\nu'^2 + 4(28b + 3)\nu')) - r^2\nu^2(60 \\
& \times br^3\mu'^3 + \nu'(20br^2\mu'' - 36b + 54) + 12r^2\mu'^2(10b\nu' + 31b + 1) + 2r((13b \\
& - 12)r\mu'' + 2(2b - 3)\nu'') + r\mu'(-20br(3r\mu'' + 2\nu'') - 5b\nu'^2 + 6(31b + 6) \\
& \times \nu' + 68b + 48) + (7b - 3)\nu'^2) + r\nu^3(60br^3\mu'^3 + 2(2((13b - 12)r^2\mu'' \\
& + (2b - 3)(r\nu'' - 4)) + \nu'(5br^2\mu'' - 7b + 18)) + 8r^2\mu'^2(5b\nu' + 48b + 3) \\
& + r\mu'(-20br(3r\mu'' + \nu'') + (74b + 24)\nu' + 3(41b + 36))) + \nu^4(r(2r\mu'' \\
& \times (10br\mu' - 13b + 12) - \mu'(4r\mu'(5br\mu' + 33b + 3) + 55b + 60)) + 21b \\
& - 39)) - \nu^2(\nu(2r\mu' + 1) - r(2r\mu' + \nu'))^2(3a_2(b + 1)r^5(r - \nu) + \nu(\nu(2r
\end{aligned}$$

$$\begin{aligned}
& \times (\mu' + 1) - r(2r\mu' + \nu'))(10br(\nu - r)\mu'' + 5b\mu'(r(\nu' - 4) + 3\nu) + 10br \\
& \times (\nu - r)\mu'^2 + (6 - 4b)\nu')) \Big], \\
p_r &= \frac{1}{6(2b^2 + b - 1)r^3\nu^3(\nu(2r\mu' + 1) - r(2r\mu' + \nu'))^3} \Big[a_2r^6(r - \nu)(-20 \\
& \times br^5\mu'\nu'(2r\mu' + \nu') + r^3\nu(20br^3\mu'^3 + 2\nu'(5br^2\mu'' + (6 - 8b)\nu') + 24r^2 \\
& \times \mu'^2(5b\nu' + 3b + 1) + r\mu'(-20br(r\mu'' + \nu'') + 15b\nu'^2 + 4(16b + 9)\nu')) \\
& - r^2\nu^2(6br^3\mu'^3 + \nu'(20br^2\mu'' - 60b + 66) + 12r^2\mu'^2(10b\nu' + 19b + 7) \\
& + 2r((13b - 12)r\mu'' + 2(2b - 3)\nu'') + r\mu'(-20br(3r\mu'' + 2\nu'') - 5b\nu'^2 \\
& + (90b + 84)\nu' + 20b + 72) + (3 - 5b)\nu'^2) + r\nu^3(60br^3\mu'^3 + 2(\nu'(5br^2\mu'' \\
& - 19b + 24) + 2r((13b - 12)r\mu'' + (2b - 3)\nu'') - 22b + 27) + 8r^2\mu'^2(5 \\
& \times b\nu' + 30b + 12) + r\mu'(-20br(3r\mu'' + \nu'') + (26b + 48)\nu' + 3(9b + 52))) \\
& + \nu^4(r(\mu'(-4r\mu'(5br\mu' + 21b + 9) - 7(b + 12)) + 2r\mu''(10br\mu' - 13b \\
& + 12)) + 33b - 45) - \nu^2(\nu(2r\mu' + 1) - r(2r\mu' + \nu'))^2(3a_2(b + 1)r^6(r \\
& - \nu) + \nu(\nu(2r\mu' + 1) - r(2r\mu' + \nu'))(br(r(\mu'(-10r\mu' + 5\nu' + 4) - 10r \\
& \times \mu'') + 8\nu') + \nu(r(10br\mu'' + \mu'(10br\mu' - 9b + 12)) - 12b + 6) - 12 \\
& \times r^2\mu')) \Big], \\
p_{\perp} &= \frac{1}{6(2b^2 + b - 1)r^3\nu^3(\nu(2r\mu' + 1) - r(2r\mu' + \nu'))^3} \Big[(a_2r^6(r - \nu)(-4 \\
& \times (b - 3)r^5\mu'\nu'(2r\mu' + \nu') + r^3\nu(4(b - 3)r^3\mu'^3 + \nu'(2(b - 3)r^2\mu'' + (3 \\
& - 14b)\nu') + 12r^2\mu'^2(2(b - 3)\nu' - 5) - r\mu'(4(b - 3)r(r\mu'' + \nu'') - 3(b \\
& - 3)\nu'^2 + 4(4b + 15)\nu')) + r^2\nu^2(-12(b - 3)r^3\mu'^3 + 2\nu'(-2(b - 3)r^2\mu'' \\
& + 42b - 3) + 12r^2\mu'^2(-2(b - 3)\nu' + b + 16) + 2r((3 - 17b)r\mu'' - 8b\nu'') \\
& + r\mu'(4((b - 3)r(3r\mu'' + 2\nu'') + 17b + 12) + (b - 3)\nu'^2 + 6(9b + 17)\nu') \\
& + b\nu'^2) + r\nu^3(12(b - 3)r^3\mu'^3 + 2\nu'((b - 3)r^2\mu'' - 29b) + 4r^2\mu'^2(2(b - 3) \\
& \times \nu' - 6b - 51) + 4r((17b - 3)r\mu'' + 4b\nu'') - r\mu'(4(b - 3)r(3r\mu'' + \nu'') \\
& + (38b + 42)\nu' + 153b + 93) - 70b + 3) + \nu^4(r(\mu'(4r\mu'(3(b + 6) - (b \\
& - 3)r\mu') + 85b + 45) + 2r\mu''(2(b - 3)r\mu' - 17b + 3)) + 57b)) + \nu^2(\nu(2 \\
& \times r\mu' + 1) - r(2r\mu' + \nu'))^2(3a_2(b + \mu' + 1) - r(2r\mu' + \nu'))(r(-2(b \\
& - 3)r^2\mu'' - 2(b - 3)r^2\mu'^2 + r\mu'((b - 3)\nu' + 8b + 6) - (2b + 3)\nu') + \nu(r
\end{aligned}$$

$$\times (2(b-3)r\mu'' + \mu'(2(b-3)r\mu' - 9b - 3) - 6b + 3)) \Big].$$

Appendix C: Field Equations (Model 3)

The corresponding $f(Q, T)$ field equations are

$$\begin{aligned} \rho = & \frac{1}{6(b+1)(2b-1)r^3(r-\nu)} \left[(10bJ\mu'^2r^2 + (10bJ\mu'' - 3(b+1)(k - \frac{1}{r^2} \right. \\ & \times (\nu(2\mu'r^2 \frac{r\nu' - \nu}{r(r-\nu)})))r^2 + 5b(\frac{1}{\nu^4(L)^4} (2kr^9a_3(a_3 - \frac{2\nu L}{r^2})\nu^4L)^4 (\frac{2L\nu}{r^3} \\ & - \frac{1}{r^2} ((-\frac{r\nu' - \nu}{r^2(r-\nu)} - \frac{(1-\nu')(r\nu' - \nu)}{r(r-\nu)^2} + 2\mu'' + \frac{\nu''}{r-\nu})\nu) - \frac{\nu'L}{r^2})) \\ & \times -(\nu' - 4)J)\mu'r + 2(2b-3)J\nu')r^2 + (-\frac{1}{\nu^4(L)^4} (2ka_3(10r\mu'b - 2b \\ & \times +3)(a_3 - \frac{2\nu L}{r^2}) (\frac{2L\nu}{r^3} - \frac{1}{r^2} ((-\frac{r\nu' - \nu}{r^2(r-\nu)} - \frac{(1-\nu')(r\nu' - \nu)}{r(r-\nu)^2} + 2\mu'' \\ & + \frac{\nu''}{r-\nu})\nu) - \frac{\nu'L}{r^2})r^9) + 3(b+1)(k - \frac{\nu L}{r^2})r^2 - J(20b\mu'^2r^2 + 20b\mu''r^2 \\ & + \mu'(-5\nu'b + 41b + 6)r + (7b-3)\nu'))\nu r + (3(b+1)J + r(10bJ\mu''r \\ & \times + J\mu'(10r\mu'b + 21b + 6) + \frac{1}{\nu^4(L)^4} (2kr^8a_3(5r\mu'b - 2b + 3)(a_3 - \frac{2\nu L}{r^2}) \\ & \times (\frac{2L\nu}{r^3} - \frac{1}{r^2} ((-\frac{r\nu' - \nu}{r^2(r-\nu)} - \frac{(1-\nu')(r\nu' - \nu)}{r(r-\nu)^2} + 2\mu'' + \frac{\nu''}{r-\nu})\nu) \\ & - \frac{\nu'L}{r^2})))\nu^2 \Big], \end{aligned}$$

$$\begin{aligned} p_r = & \frac{1}{6(b+1)(2b-1)r^3(r-\nu)} (r\nu(\frac{1}{L^4\nu^4} (2a_3K r^9(a_3 - \frac{2L\nu}{r^2}) (10br\mu' - 2b \\ & + 3)(\frac{2L\nu}{r^3} - \frac{L\nu'}{r^2} - \frac{1}{r^2} (\nu(-\frac{r\nu' - \nu}{r^2(r-\nu)} + 2\mu'' + \frac{\nu''}{r-\nu} - \frac{1}{r(r-\nu)^2} ((1 \\ & - \nu')(r\nu' - \nu)))))) + J(20br^2\mu'' + 20br^2\mu'^2 + r\mu'(-5b\nu' - 7b + 30) \\ & + (3-5b)\nu') + (6-12b)J - 3(b+1)r^2(K - \frac{L\nu}{r^2})) + r^2(r\mu'(5b(J\nu' \\ & - \frac{1}{L^4\nu^4} (2a_3K r^9(a_3 - \frac{2L\nu}{r^2}) (\frac{2L\nu}{r^3} - \frac{L\nu'}{r^2} - \frac{1}{r^2} (\nu(-\frac{r\nu' - \nu}{r^2(r-\nu)} + 2\mu'' \end{aligned}$$

$$\begin{aligned}
& + \frac{\nu''}{r-\nu} - \frac{(1-\nu')(r\nu' - \nu)}{r(r-\nu)^2} \Big) \Big) + 4(b-3)J + r^2(3(b+1)(K \\
& - \frac{L\nu}{r^2}) - 10bJ\mu'' - 10bJr^2\mu'^2 + 8bJ\nu') + (-\nu^2) \left(\frac{1}{L^4\nu^4} (2a_3Kr^9(a_3 \\
& - \frac{2L\nu}{r^2})(5br\mu' - 2b + 3) \left(\frac{2L\nu}{r^3} - \frac{L\nu'}{r^2} - \frac{1}{r^2} \left(\nu \left(-\frac{r\nu' - \nu}{r^2(r-\nu)} + 2\mu'' \right. \right. \right. \right. \\
& + \frac{\nu''}{r-\nu} - \frac{(1-\nu')(r\nu' - \nu)}{r(r-\nu)^2} \Big) \Big) \Big) + J(r(10br\mu'' + \mu'(r)(10br\mu' - 3b \\
& + 18)) - 9b + 9) \Big] , \\
p_{\perp} & = \frac{1}{6(b+1)(2b-1)r^3(r-\nu)} \Big[r^2(r\mu' \left(\frac{1}{L^4\nu^4} (2a_3(b-3)Kr^9(a_3 - \frac{2L\nu}{r^2}) \right. \right. \\
& \times \left. \left. \left(\frac{2L\nu}{r^3} - \frac{L\nu'}{r^2} - \frac{\nu \left(-\frac{r\nu' - \nu}{r^2(r-\nu)} + 2\mu'' + \frac{\nu''}{r-\nu} - \frac{(1-\nu')(r\nu' - \nu)}{r(r-\nu)^2} \right)}{r^2} \right) \right) - J((b \\
& - 3)\nu' + 8b + 6)) + r^2(2(b-3)J\mu'' + 3(b+1)(K - \frac{L\nu}{r^2})) + 2(b-3) \\
& \times Jr^2\mu'^2 + (2b+3)J\nu') + r\nu \left(\frac{1}{L^4\nu^4} (4a_3Kr^9(a_3 - \frac{2L\nu}{r^2})(2b - (b-3)r \\
& \times \mu') \left(\frac{2L\nu}{r^3} - \frac{L\nu'}{r^2} - \frac{1}{r^2} \left(\nu \left(-\frac{r\nu' - \nu}{r^2(r-\nu)} + 2\mu'' + \frac{\nu''}{r-\nu} - \frac{1}{r(r-\nu)^2} \left((1 \right. \right. \right. \right. \\
& - \nu')(r\nu' - \nu) \Big) \Big) \Big) \Big) + J(-4(b-3)r^2\mu'' - 4(b-3)r^2\mu'^2 + r\mu'((b-3) \\
& \times \nu' + 23b + 15) + b\nu') + (6b-3)J - 3(b+1)r^2(K - \frac{L\nu}{r^2})) + \nu^2 \left(r \left(\frac{1}{L^4\nu^4} \right. \right. \\
& \times \left. \left. (2a_3Kr^8(a_3 - \frac{2L\nu}{r^2})((b-3)r\mu' - 4b) \left(\frac{2L\nu}{r^3} - \frac{L\nu'}{r^2} - \frac{1}{r^2} \left(\nu \left(-\frac{r\nu' - \nu}{r^2(r-\nu)} \right. \right. \right. \right. \right. \\
& + 2\mu'' + \frac{\nu''}{r-\nu} - \frac{(1-\nu')(r\nu' - \nu)}{r(r-\nu)^2} \Big) \Big) \Big) \Big) + 2(b-3)Jr\mu'' + J\mu'(2(b-3)r\mu' \\
& - 3(5b+3)) - 9bJ \Big] ,
\end{aligned}$$

where

$$J = 1 - \frac{a_3r^4 \exp\left(-\frac{a_3r^2}{\nu\left(2\mu' + \frac{r\nu' - \nu}{r(r-\nu)}\right)}\right)}{\nu^2\left(2\mu' + \frac{r\nu' - \nu}{r(r-\nu)}\right)^2}, \quad k = \exp\left(-\frac{a_3r^2}{\nu\left(2\mu' + \frac{r\nu' - \nu}{r(r-\nu)}\right)}\right),$$

$$L = \left(2\mu' + \frac{r\nu' - \nu}{r(r - \nu)}\right).$$

Data Availability: No data was used for the research described in this paper.

References

- [1] Wheeler, J.A.: Ann. Phys. **2**(1957)604.
- [2] Flamm, L.: Phys. Z. **17**(1916)448.
- [3] Einstein, A. and Rosen, N.: Phys. Rev. **48**(1935)73.
- [4] Wheeler, J.A.: Phys. Rev. **97**(1955)511.
- [5] Morris, M.S. and Thorne, K.S.: Am. J. Phys. **56**(1988)395.
- [6] Visser, M., *Lorentzian Wormholes, From Einstein to Hawking* (American Institute of Physics, New York, 1995).
- [7] Dzhunushaliev, V.: Phys. Rev. **82**(2010)045032.
- [8] Visser, M., Kar, S. and Naresh, D.: Phys. Rev. Lett. **90**(2003)201102.
- [9] Lemos, J.P.S., Lobo, F.S.N and de Oliveira, S.Q.: Phys. Rev. D **68**(2003)064004.
- [10] Jahromi, A.S. and Moradpour, H.: Int. J. Mod. Phys. D **27**(2018)1850024; Godani, N. and Samanta, G.C.: Int. J. Mod. Phys. D **28**(2019)1950039; Sharif, M. and Gul, M.Z.: Symmetry **15**(2023)684.
- [11] Weyl, H.: Sitzungsber. Preuss. Akad. Wiss. **465**(1918)1.
- [12] Cai, Y.F., Capozziello, S., De Laurentis, M. and Saridakis, E.N.: Prog. Phys. **79**(2016)106901.
- [13] Jimenez, J.B., Heisenberg, I. and Koivisto, L.T.: Phys. Rev. D **98**(2018)044048.
- [14] Xu, Y. et al.: Eur. Phys. J. C **79**(2019)1.

- [15] Xu, Y. et al.: Eur. Phys. J. C **80**(2020)22.
- [16] Arora, S. et al.: Phys. Dark Universe **30**(2020)100664.
- [17] Bhattacharjee, S. and Sahoo, P.K.: Eur. Phys. J. C **80**(2020)289.
- [18] Pati, L., Mishra, B. and Tripathy, S.K.: Phys. Scr. **96**(2021)105003.
- [19] Agrawal, A.S. et al.: Phys. Dark Universe **33**(2021)100863.
- [20] Shiravand, M., Fakhry, S. and Farhoudi, M.: Phys. Dark Universe **37**(2022)101106.
- [21] Lobo, F.S.N. et al.: Phys. Rev. D **80**(2009)104012.
- [22] Mazharimousavi, S.H. and Halilsoy, M.: Mod. Phys. Lett. A **31**(2016)1650203.
- [23] Bahamonde, S. et al.: Phys. Rev. D **94**(2016)084042.
- [24] Zubair, M., Waheed, S. and Ahmed, Y.: Eur. Phys. J. C **76**(2016)444.
- [25] Sharif, M. and Nawazish, I.: Ann. Phys **389**(2018)283.
- [26] Mustafa, G. et al.: Int. J. Geom. Methods Mod. Phys. **17**(2020)2050103.
- [27] Shamir, M.F. and Fayyaz, I.: Eur. Phys. J. C **80**(2020)1102.
- [28] Hassan, Z., Mustafa, G. and Sahoo, P.K.: Symmetry **13**(2021)1260.
- [29] Malik, A. et al.: Chin. Phys. C **46**(2022)095104.
- [30] Gul, M.Z. and Sharif, M.: Chin. J. Phys. **88**(2024)388; New Astron. **106**(2024)102137; Gul, M.Z., Sharif, M. and Kanwal, I.: New Astron. **109**(2024)102204;
- [31] Mustafa, G. et al.: Phys. Rev. D **101**(2020)104013; Chin. J. Phys. **67**(2020)576; Phys. Dark Universe **30**(2020) 100652; Phys. Scr. **96**(2021)045009; Chin. J. Phys. **77**(2022)1742.
- [32] Mustafa, G. et al.: Phys. Dark Universe **31**(2021)100747; Phys. Scr. **96**(2021)105008; Ashraf, A. et al.: Ann. Phys. **422**(2020)168322; Shamir, M.F. and Mustafa, G. Ann. Phys. **418**(2020)168184.

- [33] Ditta, A. et al.: Eur. Phys. J. C **81**(2021)880; Mustafa, G., Maurya, S.K. and Ray, S.: Astrophys. J. **941**(2022)170; Javed, F. et al.: Nucl. Phys. B **990**(2023)116180; Mustafa, G. et al.: Eur. Phys. J. C **80**(2020)26; Phys. Scr. **96**(2021)045009.
- [34] Dirac, P.A.M.: Proc. R. Soc. Lond. A**333**(1973)403.
- [35] Novello, M. and Perez Bergliaffa, S.E.: Phys. Rep. **463**(2008)127.
- [36] Hehl, F.W. et al.: Rev. Mod. Phys. **48**(1976)393.
- [37] Landau, L.D. and Lifshitz, E.M., *The Classical Theory of Fields*, (Pergamon Press, Oxford 1970).
- [38] Lin, R.H., and Zhai, X.H.: Phys. Rev. D **103**(2021)124001.
- [39] Lobo, F.S.N.: Class. Quantum Grav. **25**(2008)175006.
- [40] Moradpour, H., Sadeghnezhad, N. and Hendi, S.H.: Can. J. Phys. **95**(2017)1257.
- [41] Banerjee, A. et al.: Eur. Phys. J. C **81**(2021)1031.
- [42] Chanda, A., Dey, S. and Paul, B.C.: Gen. Relativ. and Gravit. **53**(2021)78.
- [43] Samanta, G.C. and Godani, N.: Mod. Phys. Lett. A **34**(2019)1950224.
- [44] Lobo, F.S.N. and Oliveira, M.A.: Phys. Rev. D **80**(2009)104012.
- [45] Shamir, M.F. and Fayyaz, I.: Eur. Phys. J. C **80**(2020)1102.
- [46] Sharif, M. and Fatima, A.: Eur. Phys. J. Plus **138**(2023)196.
- [47] Banerjee, A. et al.: Eur. Phys. J. C **81**(2021)1031.