Investigating the Seebeck effect of the QGP medium using a novel relaxation time approximation model

Anowar Shaikh^{*1}, Shubhalaxmi Rath^{†2}, Sadhana Dash^{‡3}, and Binata Panda^{§1}

¹Department of Physics, Indian Institute of Technology (Indian School of Mines) Dhanbad, Jharkhand 826004, India

²Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica, Chile ³Department of Physics, Indian Institute of Technology Bombay, Mumbai 400076, India

Abstract

The highly energetic particle medium formed in the ultrarelativistic heavy ion collision displays a notable difference in temperatures between its central and peripheral regions. This temperature gradient can generate an electric field within the medium, a phenomenon referred to as the Seebeck effect. We have estimated the Seebeck coefficient for a dense quark-gluon plasma medium by using the relativistic Boltzmann transport equation in the recently developed novel relaxation time approximation (RTA) model within the kinetic theory framework. This study explores the Seebeck coefficient of individual quark flavors as well as the entire partonic medium, with the emphasis on its dependence on the temperature and the chemical potential. Our observation indicates that, for given current quark masses, the magnitude of the Seebeck coefficient for each quark flavor as well as for the partonic medium decreases as the temperature rises and increases as the chemical potential increases. Furthermore, we have investigated the Seebeck effect by considering the partonic interactions described in perturbative thermal QCD within the quasiparticle model. In addition, we have presented a comparison between our findings and the results of the standard RTA model.

^{*} anowar. 19 dr 0016 @ap.iitism.ac.in

[†]shubhalaxmi@academicos.uta.cl

[‡]sadhana@phy.iitb.ac.in

[§]binata@iitism.ac.in

1 Introduction

The primary objective in the study of ultrarelativistic heavy ion collisions is to detect and study the existence of a new form of matter as predicted by the quantum chromodynamics (QCD). These experiments at particle accelerators provide insight into the characteristics of fundamental particles like quarks and gluons. There is compelling evidence that the new form of matter, *i.e.* the quark-gluon plasma (QGP) matter has been detected at experimental facilities such as the Large Hadron Collider (LHC) [1-3] at CERN and the Relativistic Heavy Ion Collider (RHIC) [4-7] at BNL. There are also a number of observables which are regarded as the signatures of the presence of such a matter. These signatures include jet quenching [8,9], quarkonium suppression [10–12], elliptic flow [13, 14], photon and dilepton spectra [15, 16] etc. Since the discovery of QGP, numerous aspects of the strongly coupled QCD matter have been the focus of many investigations throughout the course of the last several decades. Quantitative data from the experiments on hadrons align with the theoretical results from ideal fluid dynamics, thus confirming the low viscosity of matter and its flow as a perfect fluid [17, 18]. Since then, it has been a pertinent and challenging task to comprehend the strongly interacting medium produced at high temperatures through studying its transport coefficients.

Different types of methodologies, such as the perturbative QCD and alternative effective models were used to determine the transport coefficients of quark matter [19–34]. The strongly coupled fluid behavior of QGP has made possible the use of theoretical approaches intended for systems of this kind. In kinetic theory, a simplified structure for the collision integral is often used, which is referred to as the relaxation time approximation (RTA) [35]. The Bhatnagar-Gross-Krook (BGK) model [36] and the Welander model [37] had previously accomplished progress in the nonrelativistic scenario. The relaxation time approximation has been used in several fields of physics. For the purpose of investigating the domain and microscopic basis of the relevance of hydrodynamics with relativistic effects, the relaxation time approximation has become an essential technique. The approximation given in ref. [35], although widely used, has some fundamental shortcomings and violates some macroscopic and microscopic conservation laws. These fundamental challenges are resolved and the essential characteristics of the linearized Boltzmann collision operator are preserved by using a novel collision integral [38]. Using the novel collision integral, the transport coefficients associated with the heat and the charge in a QGP medium have been derived in recent works [39, 40].

In the present work, we explore the thermoelectric effect by calculating the Seebeck coefficient for the strongly interacting matter created in heavy ion collisions with a temperature gradient between the fireball's core and its peripheral region. In the presence of a temperature gradient, a finite gradient of charge carriers may be produced, which ultimately results in an electric field. When there is a difference in the temperature inside a conducting material, the charge carriers in the hotter zone move towards the cooler zone, producing an electric field. When the generated electric field is strong enough to inhibit further charge flow, the diffusion ends. The Seebeck coefficient measures the magnitude of the electric field per unit temperature gradient in a medium. This thermoelectric transport coefficient is calculated by setting the electric current to zero [41, 42]. Unlike condensed matter systems, where only one kind of charge carrier participates in the transport process, in a medium with the strong interactions, both positive and negative charge carriers participate in the transport phenomena. The strongly interacting matter requires both a temperature gradient and finite baryon chemical potential in order to observe the thermoelectric effect. In the absence of the quark chemical potential, the numbers of particles and antiparticles are same, so no net thermoelectric effect is observed, thus resulting in no observable net Seebeck effect. The convention is to consider the sign of the Seebeck coefficient as positive when the thermoelectric current flows from the hotter end to the cooler end. Therefore, the sign of the Seebeck coefficient may be used to determine the sign of the dominating charge carriers in the system. It exhibits positive sign for the positive charge carriers and negative sign for the negative charge carriers. Recently, researchers have investigated the Seebeck effect in a hot hadron gas using the hadron resonance gas model [43]. Additionally, the Seebeck effect in the QGP phase has been examined within the framework of the Nambu-Jona-Lasinio (NJL) model in ref. [44]. In the background of the magnetic field, the thermoelectric response of the hot QCD medium has been investigated in references [45–49]. Our main objective in the present work is to investigate the impact of the novel RTA collision integral on the Seebeck coefficient of the QGP medium. In our approach, we incorporate the medium effects through the quasiparticle model [50], where the medium dependence is observed through the dispersion relations of the thermally massive quarks and gluons.

This work has been organized as follows. Section 2 describes the method for obtaining the Seebeck coefficient and addresses the relativistic Boltzmann transport equation (RBTE) in the novel relaxation time approximation model. Section 3 focuses on the determination of the Seebeck coefficient by solving the RBTE with the novel RTA collision integral. In section 4, we have discussed the results of the aforementioned thermoelectric coefficient by taking into account the current quark masses and the quasiparticle masses of the partons. Section 5 presents the summary of this work.

Notations and conventions: The covariant derivative ∂_{μ} represents the derivative with respect to x^{μ} , while $\partial_{\mu}^{(p)}$ represents the derivative with respect to p^{μ} . The fluid four velocity $u^{\mu} = (1, 0, 0, 0)$ is normalized to unity in the local rest frame $(u^{\mu}u_{\mu} = 1)$. In this work, the subscript f represents the flavor index, where f takes the values u, d and s. Furthermore, q_f , g_f , and δf_f $(\delta \bar{f}_f)$ represent the electric charge, the degeneracy factor,

and the infinitesimal change in the distribution function for the quark (antiquark) of the f^{th} flavor, respectively. The symbol m_f represents the current quark mass, with values of 3 MeV, 5 MeV, and 100 MeV for up, down, and strange quarks, respectively. In this work, $g_g = 2(N_c^2 - 1)$ represents the gluonic degrees of freedom, and $g_f(g_{\bar{f}}) = 2N_c$ denotes the quark (antiquark) degrees of freedom (for each flavor of quark), where $N_c = 3$ is the number of colors.

2 Relativistic Boltzmann transport equation in the context of the novel relaxation time approximation

In this section, we provide a comprehensive framework to analyze the thermoelectric effect in a hot partonic medium. We use this framework to calculate the Seebeck coefficient for each individual species as well as for the composite medium. We begin with the Boltzmann transport equation for a single particle, *i.e.*,

$$\frac{\partial f_f}{\partial t} + \frac{\mathbf{p}}{m} \cdot \nabla f_f + \mathbf{F} \cdot \frac{\partial f_f}{\partial \mathbf{p}} = \left(\frac{\partial f_f}{\partial t}\right)_{\mathbf{c}},\tag{1}$$

where $f_f = f_{eq,f} + \delta f_f$, **F** represents the force field that affects the particles in the medium and *m* represents the mass of the particle. The term on the right-hand side arises from particle collisions occurring inside the medium. Even for the most basic solution of eq. (1), the complex nature of the collision term poses some difficulties. Due to the complexity of quantum scattering theory, we use the relaxation time approximation to address the collision term which applies when the deviation (δf_f) is much less than the equilibrium particle distribution $(f_{eq,f})$, *i.e.*, $\delta f_f \ll f_{eq,f}$. To allow the linearization of the relativistic Boltzmann transport equation, we may assume that the quark distribution function is close to equilibrium.

When using the relaxation time approximation (RTA), a relaxation term of the following form replaces the collision term:

$$\left(\frac{\partial f_f}{\partial t}\right)_c = -\frac{1}{\tau_f} \Big(f_f(x, p, t) - f_{eq, f}(p) \Big).$$
⁽²⁾

The relaxation time τ_f is the time needed to restore the disturbed system to its original equilibrium condition. A damping frequency is represented by the expression $\frac{1}{\tau_f}$ in this context. In addition to the adaptability of the model, there is no instantaneous conservation of the particle number and it appears to be a noticeable drawback. This model is not compatible with the principles of microscopic and macroscopic conservation laws. As a result, it presents some challenges when attempting to describe relativistic gases using the energy-dependent relaxation time or general matching conditions. In the novel relaxation time approximation, the collision term is replaced by a relaxation term that has the following form:

$$\mathbf{C}[f_f] = -\frac{\omega_f}{\tau_{fp}} \bigg[\delta f_f - \frac{\langle (\omega_f/\tau_{fp})\delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\left\langle (\omega_f/\tau_{fp})P_1^{(0)}\delta f_f \right\rangle_0}{\left\langle (\omega_f/\tau_{fp})P_1^{(0)}P_1^{(0)} \right\rangle_0} + p^{\langle \mu \rangle} \frac{\left\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\delta f_f \right\rangle_0}{(1/3)\left\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \right\rangle_0} \bigg].$$

$$\tag{3}$$

Here,

$$P_1^{(0)} = 1 - \frac{\langle \omega_f / \tau_{fp} \rangle_0}{\langle \omega_f^2 / \tau_{fp} \rangle_0} \omega_f, \tag{4}$$

$$p^{\langle \mu \rangle} = \Delta^{\mu \nu} p_{\nu}, \tag{5}$$

$$\Delta^{\mu\nu} = g^{\mu\nu} - u^{\mu}u^{\nu}. \tag{6}$$

The momentum integrals are defined with respect to the local equilibrium distribution function $f_{eq,f}$ as follows,

$$\langle \dots \rangle_0 = \int \frac{d^3 p}{(2\pi)^3 p_0} f_{eq,f}(\dots)$$

$$= \int dP f_{eq,f}(\dots),$$
(7)

where τ_{fp} is the relaxation time and $\nu_f = \frac{1}{\tau_{fp}}$ denotes the collision frequency of the medium. The equilibrium distribution functions for f^{th} flavor of quark and antiquark are respectively given by

$$f_{eq,f} = \frac{1}{e^{\beta(\omega_f - \mu_f)} + 1},$$
(8)

$$\bar{f}_{eq,f} = \frac{1}{e^{\beta(\omega_f + \mu_f)} + 1},$$
(9)

where $\omega_f = \sqrt{\mathbf{p}^2 + m_f^2}$, $\beta = \frac{1}{T}$, and μ_f represents the chemical potential of f^{th} flavor of quark. The momentum-dependent relaxation time is given [51–53] by

$$\tau_{fp}(\omega_f) = (\beta \omega_f)^{\gamma} \tau_f, \tag{10}$$

where γ is an arbitrary constant which controls the energy dependence of the relaxation time and in the present article, we set the value of γ to 0.5. The expression for the τ_f is given [54] by

$$\tau_f(T) = \frac{1}{5.1T\alpha_s^2 \log(1/\alpha_s) \left[1 + 0.12(2N_f + 1)\right]}.$$
(11)

Here, α_s is the QCD running coupling constant, which is a function of both temperature and chemical potential and it has the following [55] form,

$$\alpha_s = \frac{g^2}{4\pi} = \frac{12\pi}{\left(11N_c - 2N_f\right)\ln\left(\Lambda^2/\Lambda_{\overline{\rm MS}}^2\right)} , \qquad (12)$$

where $\Lambda_{\overline{\text{MS}}} = 0.176 \text{ GeV}$, $\Lambda = 2\pi \sqrt{T^2 + \mu_f^2/\pi^2}$ for electrically charged particles (quarks and antiquarks) and $\Lambda = 2\pi T$ for gluons. The relativistic Boltzmann transport equation in its general relativistic covariant version for a QGP medium containing quarks, antiquarks and gluons is given by

$$p^{\mu}\partial_{\mu}f_f + q_f F^{\mu\nu}p_{\nu}\partial^{(p)}_{\mu}f_f = \mathbf{C}[f_f].$$
(13)

Here, $F^{\mu\nu}$ represents the electromagnetic field strength tensor, whose components describe the electric and magnetic fields. We consider the components $F^{i0} = \mathbf{E}$ and $F^{0i} = -\mathbf{E}$ to see the effect of electric field. The relativistic Boltzmann transport equation can further be expressed as

$$\mathbf{p} \cdot \frac{\partial f_{eq,f}}{\partial \mathbf{r}} + q_f \mathbf{E} \cdot \mathbf{p} \frac{\partial f_{eq,f}}{\partial p^0} + q_f p_0 \mathbf{E} \cdot \frac{\partial f_{eq,f}}{\partial \mathbf{p}} = \mathbf{C} [f_f].$$
(14)

3 Seebeck coefficient of the QGP medium

In the presence of the thermal gradient, the charge carriers move from the higher temperature zone to the lower temperature zone. Consequently, an electric current is generated in the medium and the corresponding induced current density is expressed as

$$\mathbf{J}^{i} = \sum_{f} g_{f} \int \frac{d^{3}p}{(2\pi)^{3}} \frac{p^{i}}{\omega_{f}} \left[q_{f} \delta f_{f}(x,p) + \bar{q}_{f} \delta \bar{f}_{f}(x,p) \right], \tag{15}$$

where J^i represents the spatial part of the current density vector. The symbols δf_f and $\delta \bar{f}_f$ respectively represent the infinitesimal changes in the phase space distribution functions of the f^{th} flavor of quark and antiquark. The relativistic Boltzmann transport equation (14) can be expressed in terms of the collision integral in the novel relaxation time approximation as

$$\mathbf{p}\frac{\partial f_{eq,f}}{\partial \mathbf{r}} + q_f \mathbf{E} \cdot \mathbf{p}\frac{\partial f_{eq,f}}{\partial p^0} + q_f p_0 \mathbf{E} \cdot \frac{\partial f_{eq,f}}{\partial \mathbf{p}} = -\frac{\omega_f}{\tau_{fp}} \bigg[\delta f_f - \frac{\langle (\omega_f/\tau_{fp}) \delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp}) P_1^{(0)} \delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp}) P_1^{(0)} P_1^{(0)} \rangle_0} + p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} \delta f_f \rangle_0}{(1/3) \langle (\omega_f/\tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \rangle_0} \bigg].$$
(16)

The partial derivatives appearing in the above equation are calculated as follows,

$$\frac{\partial f_{eq,f}}{\partial \mathbf{r}} = \beta^{2} (\omega_{f} - \mu_{f}) f_{eq,f} (1 - f_{eq,f}) \vec{\nabla} T(\mathbf{r})
\frac{\partial f_{eq,f}}{\partial p^{0}} = -\beta f_{eq,f} (1 - f_{eq,f})
\frac{\partial f_{eq,f}}{\partial \mathbf{p}} = -\beta \mathbf{v} f_{eq,f} (1 - f_{eq,f})$$
(17)

Using eq. (17), the left-hand side of eq. (16) can be written as

$$\mathbf{p}\frac{\partial f_{eq,f}}{\partial \mathbf{r}} + q_f \mathbf{E} \cdot \mathbf{p}\frac{\partial f_{eq,f}}{\partial p^0} + q_f p_0 \mathbf{E} \cdot \frac{\partial f_{eq,f}}{\partial \mathbf{p}} = \mathbf{p}\beta^2 (\omega_f - \mu_f) f_{eq,f} (1 - f_{eq,f}) \vec{\nabla} T(\mathbf{r}) - 2q_f \beta(\mathbf{E} \cdot \mathbf{p}) f_{eq,f} (1 - f_{eq,f}).$$
(18)

Based on the results of the calculations, the right-hand side of eq. (16) can be simplified (for a detailed calculation see appendix A) into

$$-\frac{\omega_f}{\tau_{fp}} \bigg[\delta f_f - \frac{\langle (\omega_f/\tau_{fp})\delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\left\langle (\omega_f/\tau_{fp})P_1^{(0)}\delta f_f \right\rangle_0}{\left\langle (\omega_f/\tau_{fp})P_1^{(0)}P_1^{(0)} \right\rangle_0} + p^{\langle \mu \rangle} \frac{\left\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\delta f_f \right\rangle_0}{(1/3)\left\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \right\rangle_0} \bigg] = -\frac{\omega_f A}{\tau_{fp}} \delta f_f. \quad (19)$$

Now, with the help of eq. (18) and eq. (19), one can solve eq. (16) to get the expression of δf_f as

$$\delta f_f = -\frac{\mathbf{p}\beta^2 \tau_{fp}}{\omega_f A} (\omega_f - \mu_f) f_{eq,f} (1 - f_{eq,f}) \vec{\nabla} T(\mathbf{r}) + \frac{2q_f \beta \tau_{fp} (\mathbf{E} \cdot \mathbf{p}) f_{eq,f} (1 - f_{eq,f})}{\omega_f A}, \quad (20)$$

where

$$A = \left[1 - \frac{\omega_f \int p^2 (f_{eq,f}/\tau_{fp}) dp}{\int p^2 \omega_f (f_{eq,f}/\tau_{fp}) dp}\right] \frac{\int p^2 (f_{eq,f}/\tau_{fp}) \left[1 - \left(\frac{\int p^2 (f_{eq,f}/\tau_{fp}) dp}{\int p^2 \omega_f (f_{eq,f}/\tau_{fp}) dp}\right) \omega_f\right] dp}{\int p^2 (f_{eq,f}/\tau_{fp}) \left[1 - \left(\frac{\int p^2 (f_{eq,f}/\tau_{fp}) dp}{\int p^2 \omega_f (f_{eq,f}/\tau_{fp}) dp}\right) \omega_f\right]^2 dp} + \frac{3p \int p^3 (f_{eq,f}/\tau_{fp}) dp}{\int p^4 (f_{eq,f}/\tau_{fp}) dp}.$$
 (21)

Similarly for antiquarks, we have

$$\delta \bar{f}_f = -\frac{\mathbf{p}\beta^2 \tau_{\bar{f}p}}{\omega_f \bar{A}} (\omega_f + \mu_f) \bar{f}_{eq,f} (1 - \bar{f}_{eq,f}) \vec{\nabla} T(\mathbf{r}) + \frac{2q_{\bar{f}}\beta \tau_{\bar{f}p} (\mathbf{E} \cdot \mathbf{p}) \bar{f}_{eq,f} (1 - \bar{f}_{eq,f})}{\omega_f \bar{A}}, \quad (22)$$

where

$$\bar{A} = \left[1 - \frac{\omega_f \int p^2 (\bar{f}_{eq,f}/\tau_{\bar{f}p}) dp}{\int p^2 \omega_f (\bar{f}_{eq,f}/\tau_{\bar{f}p}) dp}\right] \frac{\int p^2 (\bar{f}_{eq,f}/\tau_{\bar{f}p}) \left[1 - \left(\frac{\int p^2 (\bar{f}_{eq,f}/\tau_{\bar{f}p}) dp}{\int p^2 \omega_f (f_{eq,f}/\tau_{\bar{f}p}) dp}\right) \omega_f\right] dp}{\int p^2 (\bar{f}_{eq,f}/\tau_{\bar{f}p}) \left[1 - \left(\frac{\int p^2 (\bar{f}_{eq,f}/\tau_{\bar{f}p}) dp}{\int p^2 \omega_f (f_{eq,f}/\tau_{\bar{f}p}) dp}\right) \omega_f\right]^2 dp} + \frac{3p \int p^3 (\bar{f}_{eq,f}/\tau_{\bar{f}p}) dp}{\int p^4 (\bar{f}_{eq,f}/\tau_{\bar{f}p}) dp}.$$
 (23)

In order to get the spatial component of the induced current density, let us replace the values of δf_f and $\delta \bar{f}_f$ in eq. (15) for a single quark flavor as

$$\mathbf{J}^{i} = \frac{g_{f}q_{f}}{2\pi^{2}} \left[\int dp \frac{p^{4}}{\omega_{f}^{2}} \left[\frac{\tau_{fp}f_{eq,f}(1 - f_{eq,f})(\omega_{f} - \mu_{f})}{A} - \frac{\tau_{\bar{f}p}\bar{f}_{eq,f}(1 - \bar{f}_{eq,f})(\omega_{f} + \mu_{f})}{\bar{A}} \right] \left(-\beta^{2}\vec{\nabla}T(\mathbf{r}) \right) \right. \\ \left. + 2q_{f} \int dp \frac{p^{4}}{\omega_{f}^{2}} \left[\frac{\tau_{fp}f_{eq,f}(1 - f_{eq,f})}{A} + \frac{\tau_{\bar{f}p}\bar{f}_{eq,f}(1 - \bar{f}_{eq,f})}{\bar{A}} \right] \left(\beta \mathbf{E} \right) \right]. \quad (24)$$

When the system is in a condition of equilibrium, the net current resulting from the f^{th} quark flavor becomes zero. The current density mentioned above (eq. (24)) is assumed to be zero in order to determine the electric field resulting from the temperature gradient in the medium. This allows us to establish a relationship between the temperature gradient in the coordinate space and the induced electric field inside the medium as

$$\mathbf{E} = S\vec{\nabla}T(\mathbf{r}).\tag{25}$$

Here, the Seebeck coefficient (S) is defined as the electric field (magnitude) created in a conducting medium per unit temperature gradient when the electric current is set to zero and thus, we have

$$S = \frac{\beta}{2q_f} \frac{\int dp \frac{p^4}{\omega_f^2} \left[\frac{\tau_{fp} f_{eq,f}(1 - f_{eq,f})(\omega_f - \mu_f)}{A} - \frac{\tau_{\bar{fp}} \bar{f}_{eq,f}(1 - \bar{f}_{eq,f})(\omega_f + \mu_f)}{\bar{A}} \right]}{\int dp \frac{p^4}{\omega_f^2} \left[\frac{\tau_{fp} f_{eq,f}(1 - f_{eq,f})}{A} + \frac{\tau_{\bar{fp}} \bar{f}_{eq,f}(1 - \bar{f}_{eq,f})}{\bar{A}} \right]}{\bar{A}}.$$
 (26)

For a single species and degeneracy factor, eq. (26) may be recast into

$$S_f = \frac{\beta}{2q_f} \frac{Q_2}{Q_1},\tag{27}$$

where,

$$Q_{2} = \int dp \frac{p^{4}}{\omega_{f}^{2}} \Big[\frac{\tau_{fp} f_{eq,f} (1 - f_{eq,f}) (\omega_{f} - \mu_{f})}{A} - \frac{\tau_{\bar{f}p} \bar{f}_{eq,f} (1 - \bar{f}_{eq,f}) (\omega_{f} + \mu_{f})}{\bar{A}} \Big], \qquad (28)$$

$$Q_{1} = \int dp \frac{p^{4}}{\omega_{f}^{2}} \Big[\frac{\tau_{fp} f_{eq,f} (1 - f_{eq,f})}{A} + \frac{\tau_{\bar{f}p} \bar{f}_{eq,f} (1 - \bar{f}_{eq,f})}{\bar{A}} \Big].$$
(29)

Thus far, our analysis has included a single quark flavor. However, we will use the QGP medium that comprises of u, d and s quark flavors.

The overall spatial current in the medium is the resultant vector of the currents caused by the individual species and is written as

$$\mathbf{J} = \sum_{f} J^{f} = J^{u} + J^{d} + J^{s} \\
= \left[\frac{g_{u}q_{u}^{2}\beta}{\pi^{2}} (Q_{1})_{u} + \frac{g_{d}q_{d}^{2}\beta}{\pi^{2}} (Q_{1})_{d} + \frac{g_{s}q_{s}^{2}\beta}{\pi^{2}} (Q_{1})_{s} \right] \mathbf{E} \\
- \left[\frac{g_{u}q_{u}\beta^{2}}{2\pi^{2}} (Q_{2})_{u} + \frac{g_{d}q_{d}\beta^{2}}{2\pi^{2}} (Q_{2})_{d} + \frac{g_{s}q_{s}\beta^{2}}{2\pi^{2}} (Q_{2})_{s} \right] \vec{\nabla}T(\mathbf{r}).$$
(30)

The total induced electric field at steady state is obtained by setting the current density **J** equal to zero,

$$\mathbf{E} = \frac{\sum_{f} g_{f} q_{f}(Q_{2})_{f}}{\sum_{f} g_{f} q_{f}^{2}(Q_{1})_{f}} \frac{\beta}{2} \vec{\nabla} T(\mathbf{r}), \qquad (31)$$

which provides the total Seebeck coefficient for the whole medium as

$$S = \frac{\beta}{2} \frac{\sum_{f} g_{f} q_{f}(Q_{2})_{f}}{\sum_{f} g_{f} q_{f}^{2}(Q_{1})_{f}}.$$
(32)

The overall Seebeck coefficient of the medium can be expressed in terms of the Seebeck coefficient of each flavor (since each flavor has the same degeneracy factor (g_f)) as

$$S = \frac{\sum_{f} q_{f}^{2} S_{f}(Q_{1})_{f}}{\sum_{f} q_{f}^{2}(Q_{1})_{f}}.$$
(33)

4 Results and discussions

The quasiparticle description provides a phenomenological explanation of the behavior of quarks and gluons in a thermal QCD medium. This description takes into account the generation of thermal masses for the partons, in addition to their current masses. Thermal masses get generated when the partons interact with the medium and one may perceive QGP as a collection of the noninteracting quasiparticles inside the framework of the quasiparticle model. A model of this kind was first introduced by Goloviznin and Satz [56] to investigate the properties of the gluonic plasma. Subsequently, Peshier *et* al. [57, 58] used this model to examine the equation of state of the quark-gluon plasma derived from the lattice QCD calculations at finite temperature. In references [59,60], the quasiparticle model has been used to elucidate the lattice data, where a proper description of quasiparticles for the QGP medium has been utilized by taking into account the temperature and chemical potential dependent masses. The findings obtained from these models indicate that it is feasible to characterize the high temperature QGP phase by using a thermodynamically coordinated quasiparticle model. In this work, we have used the quasiparticle model, where the effective mass of the f^{th} quark flavor is written [50,61] as

$$m_{fT}^2 = m_f^2 + \sqrt{2}m_f m_{ft} + m_{ft}^2.$$
(34)

Here, m_f and m_{ft} represent the current mass and the thermal mass of the quark, respectively. The latter can be computed using the hard thermal loop (HTL) perturbation theory [58, 62] as

$$m_{fT}^2 = \frac{g^2 T^2}{6} \left(1 + \frac{\mu_f^2}{\pi^2 T^2} \right), \tag{35}$$

where, g is determined by the equation $g = (4\pi\alpha_s)^{1/2}$ and α_s is defined in equation (12). From eq. (12), it is evident that the value of α_s is affected by both temperature and chemical potential. The dispersion relation for the f^{th} flavor of quark is expressed as

$$\omega_f^2 = \mathbf{p}^2 + m_f^2 + \sqrt{2}m_f m_{ft} + m_{ft}^2.$$
(36)



Figure 1: The variation of Q_1 as a function of temperature (a) for u quark and (b) for d quark.



Figure 2: (a) The variation of Q_1 as a function of temperature for s quark and (b) the variation of Q_2 as a function of temperature for u quark.



Figure 3: The variation of Q_2 as a function of temperature (a) for d quark and (b) for s quark.

Using equations (27), (28), (29) and (33), we have calculated the Seebeck coefficient for each species as well as for the medium for quark chemical potential $\mu = 0.04$ GeV, 0.06 GeV, 0.08 GeV. We have used a temperature range from 0.16 GeV to 0.64 GeV for different chemical potential values. The variations of the integrals Q_1 and Q_2 with the temperature for u, d and s quarks at different finite chemical potentials are shown in figures 1a-2a and 2b-3b, respectively for the current quark mass scenario. The figures also compare our findings for the integrals Q_1 and Q_2 with the results obtained using the standard RTA model. It has been found that the modified RTA collision term leads to a decrease in these integrals for the QGP medium, regardless of the quark flavor. The closely comparable current masses of the u and d quarks result in similar values of the Q_1 and Q_2 integrals, whereas the values of these integrals are slightly different for the s quark (as seen in figures 2a and 3b) due of its large mass. However, the ratio Q_2/Q_1 for s quark produces values that are not significantly different from those of the u and dquarks. This observation on the integrals Q_1 and Q_2 facilitates the understanding of the Seebeck coefficient for individual quarks.



Figure 4: The variation of Seebeck coefficient as a function of temperature (a) for u quark and (b) for d quark.



Figure 5: (a) The variation of Seebeck coefficient as a function of temperature for s quark and (b) the variation of the total Seebeck coefficient as a function of temperature.

Figures 4a-5a and 5b demonstrate the changes in the Seebeck coefficient with an increase of the temperature for individual quarks and the composite system at various finite chemical potentials, respectively. Based on the information shown in figures 4a-5a, it is observed that the magnitude of the Seebeck coefficient for each quark decreases as the temperature increases, while keeping the chemical potential constant, for both collision terms. The reduction in the overall particle number density as the temperature increases, while the chemical potential remains constant, accounts for the decreasing value in the Seebeck coefficient. On the other hand, it has been observed that the Seebeck coefficient rises with the increasing chemical potential at a certain temperature. This is due to the

fact that a higher chemical potential is suggestive of a larger number of particles over antiparticles. The Seebeck coefficients for d and s quarks undergo a reversal in sign as a result of their negative charges. Similarly, a larger value of chemical potential for the d and s quarks would result in a greater number of negative charges (particles) over the number of positive charges (antiparticles), which would increase the negative value of the Seebeck coefficient. Regardless of the species of quark, we have noticed a significant decrease in the magnitude of the Seebeck coefficient in the novel RTA model as compared to the standard RTA model. The disparity in electric charges between u and d quarks is apparent in the substantial discrepancy in the Seebeck coefficient, with the Seebeck coefficient of the d quark being around twice that of the u quark. However, it is worth noting that the electric charge of the s quark is identical to that of the d quark. Thus, they show a significant similarity in the values of their respective individual Seebeck coefficients. Figure 5b illustrates the changes in the total Seebeck coefficient (S) in both the standard RTA and the novel RTA models as the temperature varies at different finite chemical potentials. A significant reduction in the quantity of S is seen in case of the novel RTA model. The rate of decline gets gradually smaller at higher temperatures as compared to that at lower temperatures. Our findings indicate that the magnitude of Sfalls as the temperature enhances and increases when we increase the chemical potential while keeping the temperature fixed, comparable to the behavior seen in individual quark cases. One notable observation when comparing to the standard RTA model is that, despite both S_d and S_s being negative, the magnitudes of S_u , S_d and S_s are such that the overall Seebeck coefficient of the medium is positive in the low temperature region and becomes slightly negative in the high temperature region. The sign of the Seebeck coefficient describes the direction of the generated field relative to the temperature gradient, often indicating a rising temperature. A positive Seebeck coefficient indicates that the induced field aligns with the temperature gradient in the system.

For a conducting medium, the presence of the thermoelectric effects will have an effect on both the electric current and the heat current. A finite value of the Seebeck coefficient can affect the electric current as well as the heat current in the medium. For instance, the electric current will be changed to $\mathbf{J} = \sigma_{el} \mathbf{E} - S \sigma_{el} \vec{\nabla} T(\mathbf{r})$, while the thermal conductivity can be modified to $\kappa = \kappa_0 - T \sigma_{el} S^2$. While the electrical and the thermal conductivities are always positive¹, the Seebeck coefficient can be negative for the temperature and chemical potential ranges considered in this study (fig. 5b). Thus, a negative Seebeck coefficient results in an increase in the net electric current within the medium. Similarly, the value of the Seebeck coefficient decreases the thermal conductivity. It would be intriguing to consider the thermoelectric effects while calculating entropy production in the field of

¹The manifestation of positivity of the electrical conductivity can be observed through the generation of entropy, which is governed by the second law of thermodynamics. In the presence of an electromagnetic field, the electrical conductivity and the thermal conductivity are positive by requiring that $T\partial_{\mu}\mathbf{s}^{\mu} \geq 0$ for the entropy current value [63, 64].

viscous hydrodynamics and viscous magnetohydrodynamics [63, 64].



Figure 6: The variation of Q_1 as a function of temperature (a) for u quark and (b) for d quark.



Figure 7: (a) The variation of Q_1 as a function of temperature for s quark and (b) the variation of Q_2 as a function of temperature for u quark.



Figure 8: The variation of Q_2 as a function of temperature (a) for d quark and (b) for s quark.



Figure 9: The variation of Seebeck coefficient as a function of temperature (a) for u quark and (b) for d quark.



Figure 10: (a) The variation of Seebeck coefficient as a function of temperature for s quark and (b) the variation of the total Seebeck coefficient as a function of temperature.

Figures 6a-7a and 7b-8b respectively illustrate the variations of integrals Q_1 and Q_2 with temperature for different quark flavors (u, d and s) at various finite chemical potentials using the quasiparticle framework. They have a trend that is similar to the current quark mass scenario, suggesting that they have positive values over the entire temperature range. According to the data shown in figures 9a-10a, it is found that the magnitude of the Seebeck coefficient for each quark decreases as the temperature increases and increases when the chemical potential increases. The introduction of the quasiparticle description leads to a small increase in the magnitudes of the Seebeck coefficients for all the quarks. Moreover, the total Seebeck coefficient of the QGP medium in the quasiparticle description, as seen in figure 10b, has been found to possess a slight positive value that decreases as the temperature range. This result is different from the current mass scenario (5b), where the Seebeck coefficient in the high temperature region exhibited a slight negative value. The small increase in magnitude of the Seebeck coefficient is the sole difference between the current quark mass scenario and the quasiparticle mass scenario.

5 Summary

This work focused on the novel RTA model to investigate the thermoelectric characteristics of a dense QGP medium by determining the Seebeck coefficient. We applied the kinetic theory method to solve the relativistic Boltzmann transport equation using a modified collision integral for precisely determining the Seebeck coefficient of the stated medium. This study examined both the current quark mass and quasiparticle mass contributions to the thermoelectric effect, where both the temperature and the chemical potential affect the individual component masses in the medium. The Seebeck coefficients were computed separately for the up, down and strange quarks. These distinct coefficients were then used to determine the Seebeck coefficient of the medium by a weighted average. The magnitude of the Seebeck coefficient decreases as the temperature of the medium increases and increases with increasing value of the chemical potential. The magnitude of the Seebeck coefficient was found to be increased for the quasiparticle mass scenario as compared to the current quark mass scenario.

6 Acknowledgments

One of the present authors (S. R.) acknowledges financial support from ANID Fondecyt postdoctoral grant 3240349 and S. D. acknowledges the SERB Power Fellowship, SPF/2022/000014 for the support on this work.

A Derivation of equation (19)

The collisional integral in the novel RTA model is given by

$$\mathbf{C}[f_f] = -\frac{\omega_f}{\tau_{fp}} \bigg[\delta f_f - \frac{\langle (\omega_f/\tau_{fp})\delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} + P_1^{(0)} \frac{\langle (\omega_f/\tau_{fp})P_1^{(0)}\delta f_f \rangle_0}{\langle (\omega_f/\tau_{fp})P_1^{(0)}P_1^{(0)} \rangle_0} + p^{\langle \mu \rangle} \frac{\langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}\delta f_f \rangle_0}{(1/3) \langle (\omega_f/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle} \rangle_0} \bigg].$$
(A.37)

The second term appearing in eq. (A.37) can be reduced to

$$\frac{\langle (\omega_f/\tau_{fp})\delta f_f \rangle_0}{\langle \omega_f/\tau_{fp} \rangle_0} = (1/\langle \omega_f/\tau_{fp} \rangle_0) \int dP \frac{\omega_f}{\tau_{fp}} f_{eq,f} \delta f_f = (1/\langle \omega_f/\tau_{fp} \rangle_0) \\ \times \left[\delta f_f \int dP \frac{\omega_f}{\tau_{fp}} f_{eq,f} - \int \left[\frac{d(\partial f_f)}{dP} \int dP \frac{\omega_f}{\tau_{fp}} f_{eq,f} \right] dP \right] \\ = (1/\langle \omega_f/\tau_{fp} \rangle_0) \delta f_f \left(\langle \omega_f/\tau_{fp} \rangle_0 \right) = \delta f_f. \quad (A.38)$$

Similarly, the 3rd and 4th terms appearing in eq. (A.37) can be respectively reduced to

$$P_{1}^{(0)} \frac{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \delta f_{f} \right\rangle_{0}}{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} P_{1}^{(0)} \right\rangle_{0}} = P_{1}^{(0)} \frac{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} \right\rangle_{0}}{\left\langle (\omega_{f}/\tau_{fp}) P_{1}^{(0)} P_{1}^{(0)} \right\rangle_{0}} \delta f_{f}, \qquad (A.39)$$

$$p^{\langle \mu \rangle} \frac{\left\langle (\omega_f / \tau_{fp}) p_{\langle \mu \rangle} \delta f_f \right\rangle_0}{(1/3) \left\langle (\omega_f / \tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \right\rangle_0} = p^{\langle \mu \rangle} \frac{\left\langle (\omega_f / \tau_{fp}) p_{\langle \mu \rangle} \right\rangle_0}{(1/3) \left\langle (\omega_f / \tau_{fp}) p_{\langle \mu \rangle} p^{\langle \mu \rangle} \right\rangle_0} \delta f_f.$$
(A.40)

Finally, eq. (A.37) takes the following form,

$$-\frac{\omega_{f}}{\tau_{fp}} \bigg[\delta f_{f} - \frac{\langle (\omega_{f}/\tau_{fp})\delta f_{f}\rangle_{0}}{\langle \omega_{f}/\tau_{fp}\rangle_{0}} + P_{1}^{(0)} \frac{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}\delta f_{f}\rangle_{0}}{\langle (\omega_{f}/\tau_{fp})P_{1}^{(0)}P_{1}^{(0)}\rangle_{0}} + p^{\langle \mu \rangle} \frac{\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}\delta f_{f}\rangle_{0}}{(1/3)\langle (\omega_{f}/\tau_{fp})p_{\langle \mu \rangle}p^{\langle \mu \rangle}\rangle_{0}} \bigg] = -\frac{\omega_{f}A}{\tau_{fp}} \delta f_{f}, \quad (A.41)$$

where

$$A = \left[1 - \frac{\omega_f \int p^2 (f_{eq,f}/\tau_{fp}) dp}{\int p^2 \omega_f (f_{eq,f}/\tau_{fp}) dp}\right] \frac{\int p^2 (f_{eq,f}/\tau_{fp}) \left[1 - \left(\frac{\int p^2 (f_{eq,f}/\tau_{fp}) dp}{\int p^2 \omega_f (f_{eq,f}/\tau_{fp}) dp}\right) \omega_f\right] dp}{\int p^2 (f_{eq,f}/\tau_{fp}) \left[1 - \left(\frac{\int p^2 (f_{eq,f}/\tau_{fp}) dp}{\int p^2 \omega_f (f_{eq,f}/\tau_{fp}) dp}\right) \omega_f\right]^2 dp} + \frac{3p \int p^3 (f_{eq,f}/\tau_{fp}) dp}{\int p^4 (f_{eq,f}/\tau_{fp}) dp}.$$
 (A.42)

References

- [1] A. Bzdak and V. Skokov, "Event-by-event fluctuations of magnetic and electric fields in heavy ion collisions", Phys. Lett. B **710** (2012), 171-174.
- [2] B. Alessandro *et al.* [ALICE], "ALICE: Physics Performance Report", J. Phys. G 32 (2006), 1295-2040.
- [3] F. Carminati *et al.* [ALICE], "ALICE: Physics performance report, volume I", J. Phys. G **30** (2004), 1517-1763.
- [4] V. Skokov, A. Y. Illarionov and V. Toneev, "Estimate of the magnetic field strength in heavy-ion collisions", Int. J. Mod. Phys. A 24 (2009), 5925-5932.
- [5] I. Arsene *et al.* [BRAHMS], "Quark gluon plasma and color glass condensate at RHIC? The Perspective from the BRAHMS experiment", Nucl. Phys. A **757** (2005), 1-27.
- [6] J. Adams *et al.* [STAR], "Experimental and theoretical challenges in the search for the quark gluon plasma: The STAR Collaboration's critical assessment of the evidence from RHIC collisions", Nucl. Phys. A 757 (2005), 102-183.
- [7] K. Adcox *et al.* [PHENIX], "Formation of dense partonic matter in relativistic nucleus-nucleus collisions at RHIC: Experimental evaluation by the PHENIX collaboration", Nucl. Phys. A **757** (2005), 184-283.
- [8] X. N. Wang and M. Gyulassy, "Gluon shadowing and jet quenching in A + A collisions at $s^{**}(1/2) = 200$ -GeV", Phys. Rev. Lett. **68** (1992), 1480-1483.

- [9] M. Gyulassy, P. Levai and I. Vitev, "Jet quenching in thin quark gluon plasmas. 1. Formalism", Nucl. Phys. B 571 (2000), 197-233.
- [10] J. P. Blaizot and J. Y. Ollitrault, "J/ ψ suppression in Pb-Pb collisions: A Hint of quark-gluon plasma production?", Phys. Rev. Lett. **77** (1996), 1703-1706.
- [11] H. Satz, "Quarkonium Binding and Dissociation: The Spectral Analysis of the QGP", Nucl. Phys. A 783 (2007), 249-260.
- [12] R. Rapp, D. Blaschke and P. Crochet, "Charmonium and bottomonium production in heavy-ion collisions", Prog. Part. Nucl. Phys. 65 (2010), 209-266.
- [13] R. S. Bhalerao and J. Y. Ollitrault, "Eccentricity fluctuations and elliptic flow at RHIC", Phys. Lett. B 641 (2006), 260-264.
- [14] S. A. Voloshin, A. M. Poskanzer, A. Tang and G. Wang, "Elliptic flow in the Gaussian model of eccentricity fluctuations", Phys. Lett. B 659 (2008), 537-541.
- [15] E. V. Shuryak, "Quark-Gluon Plasma and Hadronic Production of Leptons, Photons and Psions", Phys. Lett. B 78 (1978), 150.
- [16] J. I. Kapusta, P. Lichard and D. Seibert, "High-energy photons from quark gluon plasma versus hot hadronic gas", Phys. Rev. D 44 (1991), 2774-2788.
- [17] P. F. Kolb and U. W. Heinz, "Hydrodynamic description of ultrarelativistic heavy ion collisions", [arXiv:nucl-th/0305084 [nucl-th]].
- [18] P. Romatschke and U. Romatschke, "Viscosity Information from Relativistic Nuclear Collisions: How Perfect is the Fluid Observed at RHIC?", Phys. Rev. Lett. 99 (2007), 172301.
- [19] A. Muronga, "Relativistic Dynamics of Non-ideal Fluids: Viscous and heatconducting fluids. II. Transport properties and microscopic description of relativistic nuclear matter", Phys. Rev. C 76 (2007), 014910.
- [20] A. Puglisi, S. Plumari and V. Greco, "Electric Conductivity from the solution of the Relativistic Boltzmann Equation", Phys. Rev. D 90 (2014), 114009.
- [21] S. Mitra and V. Chandra, "Transport coefficients of a hot QCD medium and their relative significance in heavy-ion collisions", Phys. Rev. D 96 (2017) no.9, 094003.
- [22] S. Gupta, "The Electrical conductivity and soft photon emissivity of the QCD plasma", Phys. Lett. B 597 (2004), 57-62.
- [23] G. Aarts, C. Allton, A. Amato, P. Giudice, S. Hands and J. I. Skullerud, "Electrical conductivity and charge diffusion in thermal QCD from the lattice", JHEP 02 (2015), 186.

- [24] H. T. Ding, O. Kaczmarek and F. Meyer, "Thermal dilepton rates and electrical conductivity of the QGP from the lattice", Phys. Rev. D 94 (2016) no.3, 034504.
- [25] S. i. Nam, "Electrical conductivity of quark matter at finite T under external magnetic field", Phys. Rev. D 86 (2012), 033014.
- [26] M. Greif, I. Bouras, C. Greiner and Z. Xu, "Electric conductivity of the quark-gluon plasma investigated using a perturbative QCD based parton cascade", Phys. Rev. D 90 (2014) no.9, 094014.
- [27] B. Feng, "Electric conductivity and Hall conductivity of the QGP in a magnetic field", Phys. Rev. D 96 (2017) no.3, 036009.
- [28] P. V. Buividovich, M. N. Chernodub, D. E. Kharzeev, T. Kalaydzhyan, E. V. Luschevskaya and M. I. Polikarpov, "Magnetic-Field-Induced insulator-conductor transition in SU(2) quenched lattice gauge theory", Phys. Rev. Lett. **105** (2010), 132001.
- [29] M. Kurian and V. Chandra, "Effective description of hot QCD medium in strong magnetic field and longitudinal conductivity", Phys. Rev. D 96 (2017) no.11, 114026.
- [30] K. Hattori and D. Satow, "Electrical Conductivity of Quark-Gluon Plasma in Strong Magnetic Fields", Phys. Rev. D 94 (2016) no.11, 114032.
- [31] M. Kurian, S. Mitra, S. Ghosh and V. Chandra, "Transport coefficients of hot magnetized QCD matter beyond the lowest Landau level approximation", Eur. Phys. J. C 79 (2019) no.2, 134.
- [32] S. Rath and B. K. Patra, "Revisit to electrical and thermal conductivities, Lorenz and Knudsen numbers in thermal QCD in a strong magnetic field", Phys. Rev. D 100 (2019) no.1, 016009.
- [33] S. Rath and S. Dash, "Effects of weak magnetic field and finite chemical potential on the transport of charge and heat in hot QCD matter", Eur. Phys. J. A 59 (2023) no.2, 25.
- [34] A. Shaikh, S. Rath, S. Dash and B. Panda, "Flow of charge and heat in thermal QCD within the weak magnetic field limit: A Bhatnagar-Gross-Krook model approach", Phys. Rev. D 108 (2023) no.5, 056021.
- [35] J. Anderson and H. Witting, "A relativistic relaxation-time model for the Boltzmann equation", *Physica.* **74**, 466-488 (1974).
- [36] P. Bhatnagar, E. Gross and M. Krook, "A Model for Collision Processes in Gases 1. Small Amplitude Processes in Charged and Neutral One-Component Systems", *Phys. Rev.*. 94 pp. 511-525 (1954).

- [37] P. Welander, "On the temperature jump in a rarefied gas", Ark. Fys.. 7 pp. 507-553 (1954).
- [38] G. S. Rocha, G. S. Denicol and J. Noronha, "Novel Relaxation Time Approximation to the Relativistic Boltzmann Equation", Phys. Rev. Lett. **127** (2021) no.4, 042301.
- [39] S. Rath and S. Dash, "Analyzing the transport coefficients and observables of a rotating QGP medium in kinetic theory framework with a novel approach to the collision integral", [arXiv:2403.01240 [hep-ph]].
- [40] A. Shaikh, S. Rath, S. Dash and B. Panda, "Study of transport properties of a hot and dense QCD matter using a novel approximation method", [arXiv:2404.08919 [hep-ph]].
- [41] T. Scheidemantel, C. Ambrosch-Draxl, T. Thonhauser, J. Badding, and J. Sofo, "Transport coefficients from first-principles calculations", *Phys. Rev. B.* 68, (2003) no.9, 125210.
- [42] H. B. Callen, "Thermodynamics and an introduction to thermostatistics, second edition", chapter 14, John Wiley & sons, 1985.
- [43] J. R. Bhatt, A. Das and H. Mishra, "Thermoelectric effect and Seebeck coefficient for hot and dense hadronic matter", Phys. Rev. D 99 (2019) no.1, 014015.
- [44] A. Abhishek, A. Das, D. Kumar and H. Mishra, "Thermoelectric transport coefficients of quark matter", Eur. Phys. J. C 82 (2022) no.1, 71.
- [45] M. Kurian, "Thermoelectric behavior of hot collisional and magnetized QCD medium from an effective kinetic theory", Phys. Rev. D 103 (2021) no.5, 054024.
- [46] H. X. Zhang, J. W. Kang and B. W. Zhang, "Thermoelectric properties of the (an-)isotropic QGP in magnetic fields", Eur. Phys. J. C 81 (2021) no.7, 623.
- [47] D. Dey and B. K. Patra, "Seebeck effect in a thermal QCD medium in the presence of strong magnetic field", Phys. Rev. D 102 (2020) no.9, 096011.
- [48] D. Dey and B. K. Patra, "Thermoelectric response of a weakly magnetized thermal QCD medium", Phys. Rev. D 104 (2021), 076021.
- [49] S. A. Khan and B. K. Patra, "Seebeck and Nernst coefficients of a magnetized hot QCD medium with a number conserving kernel", Phys. Rev. D 107 (2023) no.7, 074034.
- [50] V. M. Bannur, "Quasiparticle model for QGP with nonzero densities", JHEP 09 (2007), 046.

- [51] K. Dusling, G. D. Moore and D. Teaney, "Radiative energy loss and v(2) spectra for viscous hydrodynamics", Phys. Rev. C 81 (2010), 034907.
- [52] K. Dusling and T. Schäfer, "Bulk viscosity, particle spectra and flow in heavy-ion collisions", Phys. Rev. C 85 (2012), 044909.
- [53] A. Kurkela and U. A. Wiedemann, "Analytic structure of nonhydrodynamic modes in kinetic theory", Eur. Phys. J. C 79 (2019) no.9, 776.
- [54] A. Hosoya and K. Kajantie, "Transport Coefficients of QCD Matter", Nucl. Phys. B 250 (1985), 666-688.
- [55] J. Kapusta and P. Landshoff, "Finite-temperature field theory", Journal Of Physics G: Nuclear And Particle Physics 15, 267 (1989).
- [56] V. Goloviznin and H. Satz, "The Refractive properties of the gluon plasma in SU(2) theory", Z. Phys. C 57 (1993), 671-676.
- [57] A. Peshier, B. Kampfer, O. P. Pavlenko and G. Soff, "A Massive quasiparticle model of the SU(3) gluon plasma", Phys. Rev. D 54 (1996), 2399-2402.
- [58] A. Peshier, B. Kampfer and G. Soff, "From QCD lattice calculations to the equation of state of quark matter", Phys. Rev. D 66 (2002), 094003.
- [59] M. Bluhm, B. Kampfer and G. Soff, "The QCD equation of state near T(c) within a quasiparticle model", Phys. Lett. B 620 (2005), 131-136.
- [60] M. Bluhm, B. Kampfer, R. Schulze, D. Seipt and U. Heinz, "A family of equations of state based on lattice QCD: Impact on flow in ultrarelativistic heavy-ion collisions", Phys. Rev. C 76 (2007), 034901.
- [61] V. M. Bannur, "Revisiting the quasiparticle model of the quark-gluon plasma", Eur. Phys. J. C 50 (2007), 629-634.
- [62] E. Braaten and R. D. Pisarski, "Simple effective Lagrangian for hard thermal loops", Phys. Rev. D 45 (1992) no.6, R1827.
- [63] X. G. Huang, M. Huang, D. H. Rischke and A. Sedrakian, "Anisotropic Hydrodynamics, Bulk Viscosities and R-Modes of Strange Quark Stars with Strong Magnetic Fields", Phys. Rev. D 81 (2010), 045015.
- [64] S. Gavin, "Transport coefficients in ultra-relativistic heavy-ion collisions", Nuclear Physics A 435 (1985), no.6, 826-843.