Morgane Austern<sup>\*</sup>

Lester Mackey<sup>†</sup>

#### Abstract

This work introduces a new, explicit bound on the Hellinger distance between a continuous random variable and a Gaussian with matching mean and variance. As example applications, we derive a quantitative Hellinger central limit theorem and efficient concentration inequalities for sums of dependent random variables.

**Keywords:** Hellinger distance; Stein's method; central limit theorem; efficient concentration inequalities; dependent random variables.

MSC2020 subject classifications: 60F05; 60F10.

# **1** Introduction

We introduce a new, explicit bound (Thm. 2.1) on the Hellinger distance [Hellinger, 1909] between a continuous random variable and a Gaussian with matching mean and variance. Thm. 2.1, based on Stein's method of exchangeable pairs [Stein, 1986], enables quantitative Hellinger central limit theorems (CLTs, Thm. 2.3) and efficient concentration inequalities (Thm. 2.5) for sums of locally dependent random variables. Notably, Thm. 2.3 also avoids the unknown constants, spectral gap conditions [Artstein et al., 2004], Stein kernel assumptions [Ledoux et al., 2015], and exponential dependence on Kullback-Leibler divergence [Bobkov et al., 2014, 2013] present in existing entropic CLTs for sums of independent random variables.

# **2** Bounding Hellinger Distance to a Gaussian

We begin by presenting a general upper bound on the Hellinger distance between a continuous random variable and a Gaussian with matching mean and variance. Throughout, we let Z denote a standard normal random variable with Lebesgue density  $\varphi$  and  $C_c^{\infty}$  denote the set of infinitely differentiable compact functions from  $\mathbb{R}$  to  $\mathbb{R}$ . Further, for a random variable S and  $p \geq 1$ , we write  $C^p$  for the set of p-times continuously differentiable functions from  $\mathbb{R}$  to  $\mathbb{R}$ , define  $||S||_p^p \triangleq \mathbb{E}[|S|^p]$ , and let  $L^p(S) \triangleq$  {measurable  $f : \mathbb{R} \to \mathbb{R} \mid \mathbb{E}[|f(S)|^p] < \infty$ }. Finally, we say that S admits a (first-order) score function  $\mathfrak{s}_S$  or a second-order score function  $I_S$  if  $\mathfrak{s}_S, I_S \in L^1(S)$  satisfy

 $\mathbb{E}(\mathfrak{s}_S(S)\Psi(S)) = -\mathbb{E}(\Psi'(S)) \quad \text{and} \quad \mathbb{E}(I_S(S)\Psi(S)) = \mathbb{E}(\Psi''(S)) \quad \text{for all} \quad \Psi \in C_c^{\infty}.$ (2.1)

<sup>\*</sup>Harvard University, United States of America. E-mail: morgane.austern@gmail.com

<sup>&</sup>lt;sup>†</sup>Microsoft Research New England, United States of America. E-mail: lmackey@microsoft.com

**Theorem 2.1** (Bounding Hellinger distance to a Gaussian). Let S be a random variable with convex support  $\Omega_S \subseteq \mathbb{R}$ ,  $\mathbb{E}(S) = 0$ ,  $\operatorname{Var}(S) = 1$ , Lebesgue density  $f_S \in C^1$ , and first and second-order score functions  $\mathfrak{s}_S, I_S \in L^2(S)$  satisfying  $\|\mathfrak{s}_S(S)S\|_2 < \infty$ . Fix any D > 0 and, for each  $t \ge -\frac{1}{2}\log(1-D)$ , let  $S^t$  be an exchangeable copy of S satisfying

$$\sum_{k=1}^{\infty} \frac{1}{\sqrt{k!}\sqrt{e^{2t}-1^k}} \mathbb{E}(|S^t - S|^k) < \infty.$$
(2.2)

Then, for any s > 0, we may bound the Hellinger distance

$$H(S,Z) \triangleq \left(\frac{1}{2} \int_{\mathbb{R}} \left(\sqrt{f_S(x)} - \sqrt{\varphi(x)}\right)^2 dx\right)^{1/2} \\ \leq \frac{1}{\sqrt{2}} \int_0^\infty \|I_t^S(X_t^S) - (X_t^S)^2 + 1 + X_t^S(\mathfrak{s}_{X_t^S}(X_t^S) + X_t^S)\|_2 dt \qquad (2.3) \\ \leq \frac{1}{\sqrt{2}} \left\{ -\frac{1}{2} \log(1-D)(1 + \|\mathfrak{s}_S(S)S\|_2) + \frac{D}{1-D} \|I_S(S)\|_2 \\ + \|\mathfrak{s}_S(S)\|_2 \left(\sqrt{\frac{D}{1-D}} - \tan^{-1}(\sqrt{\frac{D}{1-D}})\right) \\ + \int_{-\frac{1}{2}\log(1-D)}^\infty \|\frac{e^{-2t}}{1-e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} SZ + \tilde{\tau}_t^S \|_2 dt \right\}$$

in terms of the interpolants

$$X_t^S \triangleq e^{-t}S + \sqrt{1 - e^{-2t}}Z \text{ with } S \perp Z, \text{ density } f_t^S, \text{ and } (\mathfrak{s}_{X_t^S}, I_t^S) \triangleq (\frac{\partial_x f_t^S}{f_t^S}, \frac{\partial_x^2 f_t^S}{f_t^S})$$

for t > 0 and the Hermite sum

$$\tilde{\tau}_t^S \triangleq \frac{1}{s} \sum_{k=1}^{\infty} \frac{e^{-kt}}{k!\sqrt{1-e^{-2t}}} \mathbb{E}((S^t - S)^k | S) h_k(Z)$$

where  $h_k(x) \triangleq e^{x^2/2} \frac{\partial^k}{\partial x} e^{-x^2/2}$  for  $k \in \mathbb{N}$  designates the k-th Hermite polynomial.

**Remark 2.2** (Second-order score bound). The second-order score can often be bounded in terms of first-order score functions, as, by Lem. 6.2, if  $S \stackrel{d}{=} S_1 + S_2$  for independent  $S_1, S_2$  with score functions  $\mathfrak{s}_{S_1} \in L^2(S_1)$  and  $\mathfrak{s}_{S_2} \in L^2(S_2)$  then S admits a second-order score function  $I_S$  satisfying

$$||I_S(S)||_2 \le ||\mathfrak{s}_{S_1}(S_1)||_2 ||\mathfrak{s}_{S_2}(S_2)||_2.$$

Our proof of Thm. 2.1 in Sec. 3 relies on Stein's method of exchangeable pairs [Stein, 1986] and interpolation along the Ornstein-Uhlenbeck semigroup [Pavliotis, 2014, Chap. 4]. The first component takes inspiration from the arguments used by Bonis [2020] to bound Wasserstein distances.

#### 2.1 Application: Quantitative Hellinger CLT for Dependent Sums

As a first representative application of Thm. 2.1, we derive a quantitative Hellinger CLT for a sum of dependent random variables. We use the notation  $a_n \leq b_n$  to indicate that  $a_n \leq b_n + o(b_n)$  and defer the fully explicit Hellinger CLT bound to the proof in Sec. 4.

**Theorem 2.3** (Dependent Hellinger CLT). Let  $(V_i)_{i=1}^n$  be a sequence of i.i.d. random variables with n > 5,  $\mathbb{E}(V_1) = 0$ ,  $\operatorname{Var}(V_1) = 1$ ,  $|V_1| \stackrel{a.s.}{\leq} R$ , and a score function  $\mathfrak{s}_{V_1} \in L^2(V_1)$ . Fix any M > 2R, let  $\sigma \triangleq \sqrt{1 + M^2/2}$ , and define

$$S_n \stackrel{\Delta}{=} \frac{1}{\sqrt{n-1\sigma}} \sum_{i \le n-1} V_i V_{i+1} + \frac{1}{\sqrt{n-1\sigma}} \sum_{\substack{i \le n-1: \\ i \text{ is even}}} V_i M.$$
(2.5)

Then, for  $C_1 \triangleq \frac{3\sqrt{2}(R^2+1)(2R+M)^2}{(8+4M^2)}$  and  $C_2 \triangleq \frac{(1+M^2/2)(2\sqrt{2}+(2R+M)^2/2)}{(M-2R)^2}$ , we have

$$H(S_n, Z) \lessapprox \frac{C_1 \log(n)}{\sqrt{n}} + \frac{C_2}{\sqrt{n}} \|\mathfrak{s}_{V_1}(V_1)\|_2^2$$

Thm. 2.3 provides a quantitative Hellinger CLT bound that accommodates dependent summands, decays at an  $O(\frac{\log(n)}{\sqrt{n}})$  rate, and depends on  $V_1$  only through the boundedness parameter R and moments of the score function  $\mathfrak{s}_{V_1}$ . Prior entropic CLT bounds also bound the Hellinger distance but only apply to sums of independent observations and additionally depend on spectral gap conditions [Artstein et al., 2004]; exponentially on the KL divergence between each summand and Z [Bobkov et al., 2013, 2014]; or on moments of a Stein kernel [Ledoux et al., 2015], which need not exist and can be difficult to upper bound.

#### 2.2 Application: Efficient Concentration for Dependent Sums

As a second application of Thm. 2.1, we develop efficient concentration inequalities for sums of dependent random variables. Let  $\Phi$  represent the cumulative distribution function of a standard Gaussian and  $\Phi^c = 1 - \Phi$ . For an asymptotically normal sequence  $(S_n)_{n\geq 1}$  with  $\mathbb{E}(S_n) = 0$  and  $\operatorname{Var}(S_n) = 1$ , Austern and Mackey [2022] defined an efficient concentration inequality as an upper bound on  $\mathbb{P}(S_n > u)$  that converges to the asymptotically exact value  $\Phi^c(u)$  as  $n \to \infty$ . The authors also derive such inequalities for scaled sums of independent variables. Our Hellinger bounds enable efficient concentration for more general random variables via the following relation, proved in Sec. 5.

**Lemma 2.4** (Hellinger concentration). If *S* has a Lebesgue density, then, for all  $u \in \mathbb{R}$ ,

$$|\sqrt{\mathbb{P}(S>u)} - \sqrt{\Phi^c(u)}| \vee |\sqrt{\mathbb{P}(|S|>u)} - \sqrt{2\Phi^c(u)}| \le \sqrt{2}H(S,Z)$$

For example, Thm. 2.3 and Lem. 2.4 together directly imply the following efficient concentration inequality for the locally-dependent sum  $S_n$  (2.5).

Theorem 2.5 (Efficient dependent concentration). Under the assumptions of Thm. 2.3,

$$|\sqrt{\mathbb{P}(S_n > u)} - \sqrt{\Phi^c(u)}| \lesssim \frac{\sqrt{2}C_1 \log(n)}{\sqrt{n}} + \frac{\sqrt{2}C_2}{\sqrt{n}} \|\mathfrak{s}_{V_1}(V_1)\|_2^2$$

for all u in  $\mathbb{R}$  and  $C_1$  and  $C_2$  as defined in Thm. 2.3.

### **3 Proof of Thm. 2.1: Bounding Hellinger distance to a Gaussian**

We begin by establishing the inequality (2.3). Fix any  $T < \infty$ . By definition we have

$$H(S, X_T^S)^2 = 1 - \int_{\Omega_S} \sqrt{f_S(x)} \sqrt{f_T^S(x)} dx.$$

Fix any  $x \in \Omega_S$ . Since the function  $t \to \sqrt{f_t^S(x)}$  is continuous on  $[0, \infty)$  and differentiable on  $(0, \infty)$  by Lem. 6.3, the fundamental theorem of calculus [Knapp, 2007, Thm. 1.32] implies that  $\int_0^T \partial_t \sqrt{f_t^S(x)} dt = \sqrt{f_T^S(x)} - \sqrt{f_S(x)}$ . Moreover, since  $\int_0^T \int_{\Omega_S} \sqrt{f_S(x)} |\partial_t \sqrt{f_t^S(x)}| dx dt < \infty$  by Lem. 6.4, Fubini's theorem [Schilling, 2005, Cor. 13.9] implies that

$$H(S, X_T^S)^2 = -\int_0^T \int_{\Omega_S} \sqrt{f_S(x)} \partial_t \sqrt{f_s(x)} dx \, dt.$$

We will re-express this in terms of information-theoretic quantities. Write  $h_t^S \triangleq f_t^S / \varphi$ . By the Fokker-Planck equation [Pavliotis, 2014, Thm. 4.1], we have

$$\partial_{t}\sqrt{f_{t}^{S}(x)} = \frac{1}{2\sqrt{f_{t}^{S}(x)}}\partial_{t}f_{t}^{S}(x) = \frac{\varphi(x)}{2\sqrt{f_{t}^{S}(x)}}\partial_{t}\frac{f_{t}^{S}(x)}{\varphi(x)} = \frac{\varphi(x)}{2\sqrt{f_{t}^{S}(x)}}\partial_{t}h_{t}^{S}(x)$$
(3.1)  
$$= \frac{\varphi(x)}{2\sqrt{f_{t}^{S}(x)}}[\partial_{x}^{2}h_{t}^{S}(x) - x\partial_{x}h_{t}^{S}(x)] = \frac{f_{t}^{S}(x)}{2\sqrt{f_{t}^{S}(x)}} \left[\frac{\partial_{x}^{2}h_{t}^{S}(x)}{h_{t}^{S}(x)} - x\frac{\partial_{x}h_{t}^{S}(x)}{h_{t}^{S}(x)}\right].$$

We therefore obtain that

$$\begin{split} H(S, X_T^S)^2 &= -\int_0^T \int_{\Omega_S} \sqrt{f_S(x)} \frac{f_t^S(x)}{2\sqrt{f_t^S(x)}} \Big[ \frac{\partial_x^2 h_t^S(x)}{h_t^S(x)} - x \frac{\partial_x h_t^S(x)}{h_t^S(x)} \Big] dt \\ &= -\int_0^T \mathbb{E} \Big( \frac{\sqrt{f_S(X_t^S)}}{2\sqrt{f_t^S(X_t^S)}} \Big[ \frac{\partial_x^2 h_t^S(X_t^S)}{h_t^S(X_t^S)} - X_t^S \frac{\partial_x h_t^S(X_t^S)}{h_t^S(X_t^S)} \Big] \Big) dt \\ &\stackrel{(a)}{=} -\frac{1}{2} \int_0^T \mathbb{E} \Big( \Big[ \frac{\sqrt{f_S(X_t^S)}}{\sqrt{f_t^S(X_t^S)}} - 1 \Big] \Big[ \frac{\partial_x^2 h_t^S(X_t^S)}{h_t^S(X_t^S)} - X_t^S \frac{\partial_x h_t^S(X_t^S)}{h_t^S(X_t^S)} \Big] \Big) dt \\ &\stackrel{(b)}{\leq} \frac{1}{2} \sup_{t \le T} \Big\| 1 - \frac{\sqrt{f_S(X_t^S)}}{\sqrt{f_t^S(X_t^S)}} \Big\|_2 \int_0^T \| I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(-\mathfrak{s}_{X_t^S}(X_t^S) - X_t^S) \|_2 dt \\ &= \sup_{t \le T} \frac{1}{\sqrt{2}} H(S, X_t^S) \int_0^T \| I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(-\mathfrak{s}_{X_t^S}(X_t^S) - X_t^S) \|_2 dt, \end{split}$$

where (b) uses Cauchy-Schwarz and the identities

$$\begin{aligned} \frac{\partial_x h_t^S(x)}{h_t^S(x)} &= \frac{\partial_x f_t^S(x)}{f_t^S(x)} + x = \mathfrak{s}_{X_t^S}(x) + x \quad \text{and} \\ \frac{\partial_x^2 h_t^S(x)}{h_t^S(x)} &= \frac{\partial_x^2 f_t^S(x)}{f_t^S(x)} - x^2 + 1 + 2x \frac{\partial_x h_t^S(x)}{h_t^S(x)} \\ &= I_t^S(x) - x^2 + 1 + 2x (\mathfrak{s}_{X_t^S}(x) + x) = I_t^S(x) + x^2 + 1 + 2x \mathfrak{s}_{X_t^S}(x), \end{aligned}$$
(3.2)

and (a) follows from the identities

$$\mathbb{E}\left(\frac{\partial_x^2 h_t^S(X_t^S)}{h_t^S(X_t^S)} - X_t^S \frac{\partial_x h_t^S(X_t^S)}{h_t^S(X_t^S)}\right) \stackrel{(a_1)}{=} \mathbb{E}(I_t^S(X_t^S) + 1 + X_t^S \mathfrak{s}_{X_t^S}(X_t^S)) \stackrel{(a_2)}{=} 0$$

The equality  $(a_1)$  is a consequence of the identities (3.2) and (3.3), and  $(a_2)$  follows from the fact that, by our score assumption (2.1), we have  $\mathbb{E}(I_t^S(X_t^S)) = 0$  and  $\mathbb{E}(X_t^S\mathfrak{s}_{X_t^S}(X_t^S)) + 1 = 0$ . Since  $T < \infty$  was arbitrary and  $X_T^S \to Z$  surely as  $T \to \infty$ , we find that

$$H(S,Z) \le \sup_{T < \infty} H(S,X_T^S) \le \frac{1}{\sqrt{2}} \int_0^\infty \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(-\mathfrak{s}_{X_t^S}(X_t^S) - X_t^S)\|_2 dt.$$

To obtain the final bound (2.4), we fix any  $D \in (0,1)$  and s > 0, decompose the integral

$$\begin{split} &\int_0^\infty \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(-\mathfrak{s}_{X_t^S}(X_t^S) - X_t^S)\|_2 dt \\ &= \int_0^{-\frac{1}{2}\log(1-D)} \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(-\mathfrak{s}_{X_t^S}(X_t^S) - X_t^S)\|_2 dt \\ &+ \int_{-\frac{1}{2}\log(1-D)}^\infty \|I_t^S(X_t^S) - (X_t^S)^2 + 1 - X_t^S(-\mathfrak{s}_{X_t^S}(X_t^S) - X_t^S)\|_2 dt \triangleq (b_1) + (b_2), \end{split}$$

and bound each component separately.

By Lem. 6.1,  $\mathfrak{s}_{X_t^S}(X_t^S)$  is the almost-surely unique  $L^1$  random variable satisfying

$$\mathbb{E}(\mathfrak{s}_{X^S_t}(X^S_t)\Psi(X^S_t)) = -\mathbb{E}(\Psi'(X^S_t)) \quad \text{for all} \quad \Psi \in C^\infty_c.$$

Moreover, by the tower property and the independence of Z and S, we have

$$\mathbb{E}(\mathbb{E}(\mathfrak{s}_{S}(S) \mid X_{t}^{S})\Psi(X_{t}^{S})) = \mathbb{E}(\mathfrak{s}_{S}(S)\Psi(e^{-t}S + \sqrt{1 - e^{-2t}}Z)) = -e^{-t}\mathbb{E}(\Psi'(X_{t}^{S})), \quad \forall \Psi \in C_{c}^{\infty}.$$

Therefore we must have

$$\mathfrak{s}_{X_t^S}(X_t^S) = e^t \mathbb{E}(\mathfrak{s}_S(S) \mid X_t^S) \quad \text{almost surely.}$$
(3.4)

Similarly, by Lem. 6.1,  $I_t^S(X_t^S)$  is the almost-surely unique  $L^1$  random variable satisfying

$$\mathbb{E}(I_t^S(X_t^S)\Psi(X_t^S)) = \mathbb{E}(\Psi''(X_t^S)) \text{ for all } \Psi \in C_c^{\infty}.$$

Moreover, the tower property and independence of Z and S, imply that, for all  $\Psi \in C_c^{\infty}$ ,  $\mathbb{E}(\Psi(X_t^S)e^{2t}\mathbb{E}(I_S(S)|X_t^S)) = e^{2t}\mathbb{E}(\Psi(X_t^S)I_S(S)) = \mathbb{E}(\Psi''(e^{-t}S + \sqrt{1 - e^{-2t}}Z)) = \mathbb{E}(\Psi''(X_t^S)).$ 

Hence we obtain that, almost surely,

$$I_t^S(X_t^S) = e^{2t} \mathbb{E} \left( I_S(S) \mid X_t^S \right) \quad \text{and} \tag{3.5}$$
$$I_t^S(X_t^S) + X_t^S \mathfrak{s}_{X_t^S}(X_t^S) = e^t \mathbb{E} (e^t I_S(S) + \mathfrak{s}_S(S) X_t^S \mid X_t^S).$$

Therefore, the tower property, Jensen's inequality, and the triangle inequality yield

$$\begin{split} \|I_t^S(X_t^S) + 1 + X_t^S \mathfrak{s}_{X_t^S}(X_t^S)\|_2 &= \|\mathbb{E}(e^{2t}I_S(S) + e^t \mathfrak{s}_S(S)X_t^S + 1 \mid X_t^S)\|_2 \\ &\leq \|e^{2t}I_S(S) + e^t \mathfrak{s}_S(S)X_t^S + 1\|_2 \\ &\leq 1 + e^{2t}\|I_S(S)\|_2 + \|\mathfrak{s}_S(S)S\|_2 + e^t \sqrt{1 - e^{-2t}}\|\mathfrak{s}_S(S)\|_2 \end{split}$$

and hence

$$(b_1) \leq -\frac{1}{2}\log(1-D)(1+\|\mathfrak{s}_S(S)S\|_2) + \frac{D}{1-D}\|I_S(S)\|_2 + \|\mathfrak{s}_S(S)\|_2 \left(\sqrt{\frac{D}{1-D}} - \tan^{-1}(\sqrt{\frac{D}{1-D}})\right) \\ \leq -\frac{1}{2}\log(1-D)(1+\|\mathfrak{s}_S(S)S\|_2) + \frac{D}{1-D}\|I_S(S)\|_2 + \|\mathfrak{s}_S(S)\|_2 \frac{D}{\sqrt{1-D}}.$$

It remains to bound  $(b_2)$ . By Lem. 6.1 and the tower property, Z admits first and second-order score functions  $\mathfrak{s}_Z(Z) = -Z$  and  $I_Z(Z) = Z^2 - 1$  satisfying, for all  $\Psi \in C_c^{\infty}$ ,

$$\begin{split} & \mathbb{E}\big(\Psi(X^S_t)\frac{1}{\sqrt{1-e^{-2t}}}\mathbb{E}(Z|X^S_t)\big) = \mathbb{E}\big(\Psi(X^S_t)\frac{Z}{\sqrt{1-e^{-2t}}}\big) = \mathbb{E}\big(\Psi'(X^S_t)\big) \quad \text{and} \\ & \mathbb{E}\big(\Psi(X^S_t)\frac{1}{1-e^{-2t}}\mathbb{E}(Z^2-1|X^S_t)\big) = \mathbb{E}\big(\Psi(X^S_t)\frac{Z^2-1}{1-e^{-2t}}\big) = \mathbb{E}\big(\Psi''(X^S_t)\big). \end{split}$$

Using Lem. 6.1 once more we therefore obtain the almost sure identities

$$I_t^S(X_t^S) = \mathbb{E}(\frac{1}{1 - e^{-2t}}(Z^2 - 1) \mid X_t^S) \quad \text{and} \quad -\mathfrak{s}_{X_t^S}(X_t^S) - X_t^S = \mathbb{E}\Big[\frac{e^{-2t}}{\sqrt{1 - e^{-2t}}}Z - e^{-t}S \mid X_t^S\Big].$$

To conclude we will make use of the following lemma proved in Sec. 3.1.

Lemma 3.1 (Hermite sum conditional means). Under the assumptions of Thm. 2.1,

 $\mathbb{E}(\tilde{\tau}^S_t|X^S_t) = 0 \quad \text{almost surely for all} \quad t > 0.$ 

By Lem. 3.1 we have  $\mathbb{E}(\tilde{\tau}^S_t|X^S_t) = 0$  almost surely. Thus, with probability 1,

$$\begin{split} I_t^S(X_t^S) &- (X_t^S)^2 + 1 \\ &= \mathbb{E}(\frac{Z^2}{1 - e^{-2t}} - (X_t^S)^2 \mid X_t^S) - \frac{e^{-2t}}{1 - e^{-2t}} \\ &= \mathbb{E}\left(\frac{Z}{\sqrt{1 - e^{-2t}}} (\frac{Z}{\sqrt{1 - e^{-2t}}} - X_t^S) + X_t^S (\frac{Z}{\sqrt{1 - e^{-2t}}} - X_t^S) \mid X_t^S\right) - \frac{e^{-2t}}{1 - e^{-2t}} \\ &= \mathbb{E}\left(\frac{Z}{\sqrt{1 - e^{-2t}}} \left(\frac{e^{-2t}}{\sqrt{1 - e^{-2t}}} Z - e^{-t}S\right) \mid X_t^S\right) - \frac{e^{-2t}}{1 - e^{-2t}} + X_t^S \mathbb{E}\left[\left(\frac{e^{-2t}}{\sqrt{1 - e^{-2t}}} Z - e^{-t}S\right) \mid X_t^S\right] \\ &= \mathbb{E}\left(\frac{e^{-2t}}{1 - e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1 - e^{-2t}}} SZ + \tilde{\tau}_t^S \mid X_t^S\right) + X_t^S (-\mathfrak{s}_{X_t^S}(X_t^S) - X_t^S), \end{split}$$

and Jensen's inequality therefore implies

$$(b_2) = \int_{-\frac{1}{2}\log(1-D)}^{\infty} \|\mathbb{E}\left(\frac{e^{-2t}}{1-e^{-2t}}(Z^2-1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}}SZ \mid X_t^S\right)\|_2 dt \\ \leq \int_{-\frac{1}{2}\log(1-D)}^{\infty} \|\frac{e^{-2t}}{1-e^{-2t}}(Z^2-1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}}SZ + \tilde{\tau}_t^S\|_2 dt.$$

#### 3.1 Proof of Lem. 3.1: Hermite sum conditional means

Fix any  $\Psi \in C_c^{\infty}$ , define  $(P_t \Psi)(x) \triangleq \mathbb{E}(\Psi(e^{-t}x + \sqrt{1 - e^{-2t}}Z))$ , and, for each  $\psi \in C_c^{\infty}$ , let  $\psi^{(k)}$  denote its k-th derivative. Since  $\Psi \in C_c^{\infty}$ , we may write

$$(P_t \Psi)^{(k)}(x) = e^{-kt} \mathbb{E}(\Psi^{(k)}(e^{-t}x + \sqrt{1 - e^{-2t}}Z)).$$
(3.6)

By Bonis [2020, (16)], the Hermite polynomials satisfy  $\mathbb{E}(h_k(Z)\Psi(Z)) = \mathbb{E}(\Psi^{(k)}(Z))$  for each k. Therefore Hence by exploiting the independence of S and Z, we find that

$$\mathbb{E}(\Psi(X_t^S)h_k(Z) \mid S) = \mathbb{E}(\Psi(e^{-t}S + \sqrt{1 - e^{-2t}}Z)h_k(Z) \mid S)$$
  
=  $\sqrt{1 - e^{-2t}}^k \mathbb{E}(\Psi^{(k)}(e^{-t}S + \sqrt{1 - e^{-2t}}Z) \mid S) = \sqrt{1 - e^{-2t}}^k \mathbb{E}(\Psi^{(k)}(X_t^S) \mid S).$  (3.7)

We now invoke the definition of  $\tilde{\tau}_t^S$ , the tower property, the identities (3.7) and (3.6), the analyticity of  $P_t\Psi$  (by Bonis [2020, (24)] and Krantz and Parks [2002, Prop. 1.2.12]), and the fact that  $S \stackrel{d}{=} S^t$  in turn to conclude that

$$s \mathbb{E}(\mathbb{E}(\tilde{\tau}_{t}^{S} \mid X_{t}^{S})\Psi(X_{t}^{S})) \stackrel{(a)}{=} \sum_{k=1}^{\infty} \frac{e^{-kt}}{k!\sqrt{1-e^{-2t^{k}}}} \mathbb{E}\left(\mathbb{E}((S^{t}-S)^{k}|S)h_{k}(Z)\Psi(X_{t}^{S})\right)$$
$$= \sum_{k=1}^{\infty} \frac{e^{-kt}}{k!} \mathbb{E}\left(\mathbb{E}((S^{t}-S)^{k}|S)\Psi^{(k)}(X_{t}^{S})\right)$$
$$= \sum_{k=1}^{\infty} \frac{e^{-kt}}{k!} \mathbb{E}\left((S^{t}-S)^{k}\mathbb{E}(\Psi^{(k)}(X_{t}^{S})|S)\right)$$
$$= \sum_{k=1}^{\infty} \frac{1}{k!} \mathbb{E}\left((S^{t}-S)^{k}(P_{t}\Psi)^{(k)}(S)\right)$$
$$\stackrel{(b)}{=} \mathbb{E}\left((P_{t}\Psi)(S^{t}) - (P_{t}\Psi)(S)\right) = 0.$$

In (a) and (b) we used Fubini's theorem [Schilling, 2005, Cor. 13.9] to swap expectation and sum as

$$\sum_{k=1}^{\infty} \frac{e^{-kt}}{k!\sqrt{1-e^{-2t}}} \mathbb{E}\left(\mathbb{E}(|S^t - S|^k | S) | h_k(Z) | |\Psi(X_t^S)|\right) < \infty$$
(3.8)

by the boundedness of  $\Psi$ , the independence of Z and S, the Hermite polynomial inequality  $\mathbb{E}[|h_k(Z)|] \leq \sqrt{k!}$  [Bonis, 2020, Lem. 3], and the exchangeable pair assumption (2.2). Hence for all  $\Psi \in C_c^{\infty}$  we have  $\mathbb{E}(\mathbb{E}(\tilde{\tau}_t^S \mid X_t^S)\Psi(X_t^S)) = 0$ . Since  $\mathbb{E}(\tilde{\tau}_t^S \mid X_t^S) \in L^1$  by (3.8) with  $\Psi \equiv 1$ , Lem. 6.1 implies that  $\mathbb{E}(\tilde{\tau}_t^S \mid X_t^S) = 0$  almost surely, as otherwise  $\mathfrak{s}_{X_t^S}(X_t^S) + \mathbb{E}(\tilde{\tau}_t^S \mid X_t^S)$  would violate the almost sure uniqueness of the score function.

# 4 Proof of Thm. 2.3: Dependent Hellinger CLT

Thm. 2.3 follows from the following more detailed statement with D = 1 and  $\kappa = \frac{R^2(2R+M)^2\sqrt{n}}{n-1}$ .

Theorem 4.1 (Detailed dependent Hellinger CLT). Instantiate the assumptions of Thm. 2.3,

and fix any 
$$D \in (0, \sqrt{n})$$
 and  $\kappa > \frac{R^2(2R+M)^2\sqrt{n}}{(n-1)\min(D,(1+M^2/2))}$ . Then, for  $\alpha \triangleq e^{19/300}\pi^{1/4}$ ,  
 $\sqrt{2}H(S_n, Z) \leq \beta + \beta_0 + \beta_1$  for  
 $\beta \triangleq \frac{R^2}{\sqrt{n-1}\sigma} \sin^{-1}(\sqrt{1-\frac{D}{\sqrt{n}}}) + \frac{\sqrt{2}\log(n/D^2)}{4}(\frac{1}{n-1} + \frac{3(R^2+1)(2R+M)^2\sqrt{n}}{2\sigma^2(n-1)})$   
 $+ \frac{2R^2(2R+M)^2\alpha}{\sqrt{3(n-1)\sigma^2\kappa}}\sqrt{\frac{n}{n-1}}\int_0^{\sqrt{1-\frac{D}{\sqrt{n}}}} \frac{(e^{2x^2\kappa}-1)}{\sqrt{1-x^{2^3}}}dx$   
 $+ \frac{2R^2(2R+M)^2\alpha}{(n-1)\sigma^2\kappa}(\frac{1}{2} + \frac{2R(2R+M)\sqrt{n}}{\sigma(n-1)})\int_0^{\sqrt{1-\frac{D}{\sqrt{n}}}} \frac{(e^{2x^2\kappa}-1)}{\sqrt{1-x^{2^4}}}dx$   
 $\lesssim \frac{3\sqrt{2}(R^2+1)(2R+M)^2}{8\sigma^2}\frac{\log(n)}{\sqrt{n}},$   
 $\beta_0 \triangleq -\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})\left(1+\sigma^2\left\{1+\left(\frac{(2R+M)^2}{(M-2R)^2}+\frac{R^2}{\lfloor(n-1)/2\rfloor}\right)\|\mathfrak{s}_{V_1}(V_1)\|_2^2\right\}\right)$   
 $\lesssim \frac{D}{2\sqrt{n}}\left(1+\sigma^2+\frac{\sigma^2(2R+M)^2}{(M-2R)^2}\|\mathfrak{s}_{V_1}(V_1)\|_2^2\right),$   
 $\beta_1 \triangleq \frac{D}{\sqrt{n}}\left\{\frac{2\sqrt{2}\sigma^2}{(M-2R)^2(1-D/\sqrt{n})}\sqrt{\frac{n-1}{n-5}}\|\mathfrak{s}_{V_1}(V_1)\|_2^2+\frac{\sqrt{2}\sigma}{(M-2R)\sqrt{1-D/\sqrt{n}}}\|\mathfrak{s}_{V_1}(V_1)\|_2\right\},$ 

Proof of Thm. 4.1. Our result follows directly from three lemmas. The first, proved in Sec. 4.1, uses Thm. 2.1 to bound  $H(S_n, Z)$  in terms of an exchangeable copy of  $S_n$ . **Lemma 4.2** (Exchangeable pair bound on  $H(S_n, Z)$ ). Under the assumptions of Thm. 2.3, define

$$\tilde{\tau}_t \triangleq \sum_{k=1}^{\infty} \frac{e^{-kt}}{k! \sqrt{1 - e^{-2t^k}}} (n-1) \mathbb{E}((S'_n - S_n)^k | S_n) h_k(Z)$$
(4.1)

for fixed  $D \in (0, \sqrt{n})$ , all  $t \ge -\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})$ ,  $S'_n$  an exchangeable copy of  $S_n$  satisfying

$$\sum_{k=1}^{\infty} \frac{(\sqrt{n}-D)^{k/2}}{\sqrt{k!} D^{k/2}} \mathbb{E}(|S'_n - S_n|^k) < \infty,$$
(4.2)

and  $Z \perp (S'_n, S_n)$ . Then, for  $\beta_0$  and  $\beta_1$  as defined in Thm. 4.1,

$$\sqrt{2}H(S_n, Z) \le \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \left\| \frac{e^{-2t}}{1-e^{-2t}} (Z^2 - 1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} S_n Z + \tilde{\tau}_t \right\|_2 dt + \beta_0 + \beta_1.$$
(4.3)

The second, proved in Sec. 4.2, bounds the terms in (4.1) for a particular  $S'_n$  choice. **Lemma 4.3** (Exchangeable pair moment bounds). *Instantiate the assumptions of Thm. 2.3.* Then

$$S'_{n} \triangleq S_{n} + \frac{1}{\sqrt{n-1}\sigma} (V'_{J} - V_{J}) (V_{J+1} + M + V_{J-1}) \mathbb{I}(J \text{ is even}),$$
(4.4)

for  $(V'_i)_{i\geq 1} \stackrel{d}{=} (V_i)_{i\geq 1}$ ,  $J \sim \text{Uniform}(\{1, \cdots, n-1\})$ , and  $((V'_i)_{i\geq 1}, (V_i)_{i\geq 1}, J)$  independent is an exchangeable copy of  $S_n$  satisfying (4.2) for each  $D \in (0, \sqrt{n})$ ,

$$\|(n-1)\mathbb{E}(F|S_n) + S_n\|_2 \le \frac{R^2}{\sqrt{n-1}\sigma} \quad \text{for} \quad F \triangleq S'_n - S_n, \quad \text{and}$$

$$(4.5)$$

$$\|\frac{n-1}{2}\mathbb{E}(F^2|S_n) - 1\|_2 \le B_n \triangleq \frac{1}{n-1} + \frac{3(R^2+1)(2R+M)^2\sqrt{n}}{2\sigma^2(n-1)}.$$
(4.6)

 $\text{Moreover, for any } \kappa > \frac{R^2(2R+M)^2}{(n-1)\sigma^2} \text{ and } t \ge -\frac{1}{2}\log\left(1-\frac{R^2(2R+M)^2}{(n-1)\sigma^2\kappa}\right) \text{ with } c(t) \triangleq \sqrt{\kappa(1-e^{-2t})},$ 

$$\|\mathbb{E}(F^k|S_n)\|_2 \le \frac{2^{k+1}\sqrt{nc(t)^{k-3}R^2(2R+M)^2}}{(n-1)^{5/2}\sigma^2}, \ \forall \ odd \ k \ge 3 \quad and$$
(4.7)

$$\|\mathbb{E}(F^k|S_n)\|_2 \le \frac{2^{k+1}\sqrt{nc(t)^{k-4}R^3(2R+M)^3}}{(n-1)^3\sigma^3} + \frac{2^{k-1}c(t)^{k-4}R^2(2R+M)^2}{(n-1)^2\sigma^2}, \ \forall \ \text{even} \ k \ge 4.$$
(4.8)

The third, proved in Sec. 4.3, bounds the large-time component of (4.3) using Lem. 4.3. **Lemma 4.4** (Large-time Hellinger bound). Instantiate the assumptions and notation of Lem. 4.2 with  $S'_n$  defined in (4.4) and  $\kappa > \frac{R^2(2R+M)^2\sqrt{n}}{(n-1)\min(D,\sigma^2)}$ . Then, for  $\beta$  defined in Thm. 4.1,

$$\int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \left\| \frac{e^{-2t}}{1-e^{-2t}} (Z^2-1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} S_n Z + \tilde{\tau}_t \right\|_2 dt \le \beta.$$

## **4.1** Proof of Lem. 4.2: Exchangeable pair bound on $H(S_n, Z)$

We will first establish that  $S_n$  admits first and second-order score functions. Fix any  $\Psi \in C_c^{\infty}$  and, for each  $i \leq n$ , define  $V_{-i} \triangleq (V_1, \cdots, V_{i-1}, V_{i+1}, \cdots, V_n)$ . Then, for all even  $i \neq j \leq n-1$ ,

$$\begin{split} & \mathbb{E}\left(\frac{\mathfrak{s}_{V_{1}}(V_{i})}{(V_{i+1}+M+V_{i-1})}\Psi(S_{n})\right) = \mathbb{E}\left(\mathbb{E}\left(\frac{\mathfrak{s}_{V_{1}}(V_{i})}{(V_{i+1}+M+V_{i-1})}\Psi(S_{n})\big|V_{-i}\right)\right) = \frac{-\mathbb{E}(\Psi'(S_{n}))}{\sqrt{n-1}\sigma} \quad \text{and} \\ & \mathbb{E}\left(\frac{\mathfrak{s}_{V_{1}}(V_{i})\mathfrak{s}_{V_{1}}(V_{j})}{(V_{i+1}+M+V_{i-1})(V_{j+1}+M+V_{j-1})}\Psi(S_{n})\right) = \mathbb{E}\left(\frac{\mathfrak{s}_{V_{1}}(V_{j})}{(V_{j-1}+V_{j+1}+M)}\mathbb{E}\left(\frac{\mathfrak{s}_{V_{1}}(V_{i})}{(V_{i+1}+M+V_{i-1})}\Psi(S_{n})\big|V_{-i}\right)\right) \\ & = -\frac{1}{\sqrt{n-1}\sigma}\mathbb{E}\left(\frac{\mathfrak{s}_{V_{1}}(V_{j})}{(V_{j-1}+V_{j+1}+M)}\Psi'(S_{n})\right) \\ & = -\frac{1}{\sqrt{n-1}\sigma}\mathbb{E}\left(\mathbb{E}\left(\frac{\mathfrak{s}_{V_{1}}(V_{j})}{(V_{j+1}+M+V_{j-1})}\Psi'(S_{n})\big|V_{-j}\right)\right) = \frac{1}{(n-1)\sigma^{4}}\mathbb{E}\left(\Psi''(S_{n})\right). \end{split}$$

Therefore,  $S_n$  admits the first and second-order score functions

$$\begin{split} \mathfrak{s}_{S_n}(S_n) &= \frac{\sqrt{n-1}\sigma}{\lfloor (n-1)/2 \rfloor} \mathbb{E}(\sum_{i=2}^{n-1} \frac{\mathfrak{s}_{V_1}(V_i) \mathbb{I}(\text{i is even})}{(V_{i+1}+M+V_{i-1})} | S_n) \quad \text{and} \\ I_{S_n}(S_n) &= \frac{(n-1)\sigma^4}{\lfloor (n-1)/2 \rfloor (\lfloor (n-1)/2 \rfloor \rfloor - 1)} \sum_{\substack{i \neq j \leq n-1 \\ i, j \text{ even}}} \mathbb{E}\Big( \frac{\mathfrak{s}_{V_1}(V_i)\mathfrak{s}_{V_1}(V_j)}{(V_{i-1}+V_{i+1}+M)(V_{j-1}+V_{j+1}+M)} \Big| S_n \Big). \end{split}$$

We now turn our attention to bounding  $\|\mathfrak{s}_{S_n}(S_n)\|_2$ ,  $\|I_{S_n}(S_n)\|_2$ , and  $\|\mathfrak{s}_{S_n}(S_n)S_n\|_2$ . For the score bound, Jensen's inequality and the tower property yield

$$\|\mathfrak{s}_{S_n}(S_n)\|_2 \le \sigma \|\frac{\sqrt{n-1}}{\lfloor (n-1)/2 \rfloor} \sum_{i=2}^{n-1} \frac{\mathfrak{s}_{V_1}(V_i)\mathbb{I}(\text{i is even})}{(V_{i+1}+M+V_{i-1})}\|_2.$$

For each i, our independence, identical distribution, and boundedness assumptions imply that  $\mathbb{E}(\mathfrak{s}_{V_1}(V_i) \mid V_{-i}) = 0$  and  $\|\frac{\mathfrak{s}_{V_1}(V_i)\mathbb{I}(\text{i is even})}{(V_{i+1}+M+V_{i-1})}\|_2 \leq \frac{1}{M-2R}\|\mathfrak{s}_{V_1}(V_1)\|_2$ . Hence, we have

$$\|\mathfrak{s}_{S_n}(S_n)\|_2 \le \sigma \sqrt{\frac{(n-1)}{\lfloor (n-1)/2 \rfloor^2} \sum_{\substack{i \le n-1 \\ i \text{ even}}} \mathbb{E}\left(\frac{\mathfrak{s}_{V_1}(V_i)^2}{(V_{i+1}+M+V_{i-1})^2}\right)} \le \frac{\sqrt{2}\sigma}{M-2R} \|\mathfrak{s}_{V_1}(V_1)\|_2 \sqrt{\frac{n-1}{n-3}}.$$

Analogous reasoning yields

$$||I_{S_n}(S_n)||_2 \le \frac{2\sqrt{2}\sigma^2}{(M-2R)^2}\sqrt{\frac{n-1}{n-5}}||\mathfrak{s}_{V_1}(V_1)||_2^2.$$

Moreover, independence and the score function property (2.1) imply that, for all  $i \leq n-1$ ,  $\mathbb{E}(V_i|V_{-i}) = 0$  and  $\mathbb{E}(\mathfrak{s}_{V_1}(V_i)V_i|V_{-i}) = 1$ . Hence, Jensen's inequality and the tower property yield

$$\begin{split} & \mathbb{E}(\mathfrak{s}_{S_n}(S_n)^2 S_n^2) \\ & \leq \frac{\sigma^2}{\lfloor (n-1)/2 \rfloor^2} \mathbb{E}\Big(\Big(\sum_{\substack{i \leq n-1 \\ i \text{ even}}} \frac{\mathfrak{s}_{V_1}(V_i)}{V_{i+1} + M + V_{i-1}}\Big)^2 \Big(\sum_{\substack{i \leq n-1 \\ i \text{ even}}} V_i(V_{i+1} + M + V_{i-1})\Big)^2\Big) \\ & = \frac{\sigma^2}{\lfloor (n-1)/2 \rfloor^2} \sum_{\substack{i_{1:4} \leq n-1 \\ i_{1:4} \text{ even}}} \mathbb{E}\Big(\frac{\mathfrak{s}_{V_1}(V_{i_1})}{V_{i_1+1} + M + V_{i_1-1}} \frac{\mathfrak{s}_{V_1}(V_{i_2})}{(V_{i_2+1} + M + V_{i_2-1})} V_{i_3}(V_{i_3+1} + M + V_{i_3-1}) \\ & \quad V_{i_4}(V_{i_4+1} + M + V_{i_4-1})\Big) \\ & = \frac{\sigma^2}{\lfloor (n-1)/2 \rfloor^2} \sum_{\substack{i_1 \neq i_2 \leq n-1 \\ i_{1:2} \text{ even}}} \mathbb{E}\Big(\frac{\mathfrak{s}_{V_1}(V_{i_1})^2}{(V_{i_1+1} + M + V_{i_1-1})^2} V_{i_2}^2(V_{i_2+1} + M + V_{i_2-1})^2\Big) \\ & \quad + \frac{\sigma^2}{\lfloor (n-1)/2 \rfloor^2} \sum_{\substack{i_1 \neq i_2 \leq n-1 \\ i_{1:2} \text{ even}}} \mathbb{E}\Big(\mathfrak{s}_{V_1}(V_{i_1}) V_{i_1}\mathfrak{s}_{V_1}(V_{i_2}) V_{i_2}\Big) \\ & \quad + \frac{\sigma^2}{\lfloor (n-1)/2 \rfloor^2} \sum_{\substack{i_1 \neq i_2 \leq n-1 \\ i_{1:2} \text{ even}}} \mathbb{E}\Big(\mathfrak{s}_{V_1}(V_{i_1}) \|_{i_1}^2 \mathfrak{s}_{V_1}(V_{i_2}) V_{i_2}\Big) \\ & \quad \leq \sigma^2\Big\{\frac{(2R+M)^2}{(M-2R)^2} \|\mathfrak{s}_{V_1}(V_1)\|_2^2 + 1 + R^2 \|\mathfrak{s}_{V_1}(V_1)\|_2^2 \frac{1}{\lfloor (n-1)/2 \rfloor}\Big\}. \end{split}$$

The result now follows from these moment estimates and the Thm. 2.1 bound

$$\begin{split} \sqrt{2}H(S_n,Z) &\leq \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \left\| \frac{e^{-2t}}{1-e^{-2t}}(Z^2-1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}}S_nZ + \tilde{\tau}_t \right\|_2 dt \\ &- \frac{1}{2}\log(1-\frac{D}{\sqrt{n}})(1+\|\mathfrak{s}_{S_n}(S_n)S_n\|_2) + \frac{1}{\sqrt{n}}\frac{D}{1-D/\sqrt{n}}\|I_{S_n}(S_n)\|_2 + \|\mathfrak{s}_{S_n}(S_n)\|_2 \frac{D}{\sqrt{n}\sqrt{1-D/\sqrt{n}}} \|I_{S_n}(S_n)\|_2 + \|\mathfrak{s}_{S_n}(S_n)\|_2 + \|\mathfrak{s}_{S_n}(S_n)\|_2 \frac{D}{\sqrt{n}\sqrt{1-D/\sqrt{n}}} \|I_{S_n}(S_n)\|_2 + \|\mathfrak{s}_{S_n}(S_n)\|_2 \frac{D}{\sqrt{n}\sqrt{1-D/\sqrt{n}}} \|I_{S_n}(S_n)\|_2 + \|\mathfrak{s}_{S_n}(S_n)\|_2 + \|\mathfrak{s}_{S_$$

### 4.2 Proof of Lem. 4.3: Exchangeable pair moment bounds

Since  $F = \frac{(V'_J - V_J)(V_{J+1} + V_{J-1} + M)\mathbb{I}(J \text{ is even})}{\sqrt{n-1}\sigma}$ , J is uniform, and  $\mathbb{E}(V'_i|(V_i)_{i \le n}) = 0$ , we see

$$\begin{split} -(n-1)\sigma \ \mathbb{E}(F \mid (V_i)_{i \le n}) &= \frac{1}{\sqrt{n-1}} \sum_{\substack{i \le n-1 \\ i \text{ is even}}} V_i(V_{i+1} + V_{i-1} + M) \\ &= \frac{1}{\sqrt{n-1}} \sum_{\substack{i \le n-1 \\ i \text{ is even}}} V_i(V_{i+1} + M) + \frac{1}{\sqrt{n-1}} \sum_{\substack{i \le n-1 \\ i \text{ is even}}} V_iV_{i-1} \\ &= \frac{1}{\sqrt{n-1}} \sum_{\substack{i \le n-1 \\ i \text{ is even}}} V_i(V_{i+1} + M) + \frac{1}{\sqrt{n-1}} \sum_{\substack{i \le 2\lfloor \frac{n-1}{2} \rfloor - 1 \\ i \text{ is odd}}} V_iV_{i+1} \\ &= \sigma S_n + \frac{1}{\sqrt{n-1}} V_{n-1}V_n \ \mathbb{I}(n-1 \text{ is odd}). \end{split}$$

The inequality (4.5) now follows from the *R*-boundedness of each  $|V_i|$ . Similarly, since  $F^2 = \frac{(V'_J - V_J)^2 (V_{J+1} + V_{J-1} + M)^2 \mathbb{I}(J \text{ is even})}{(n-1)\sigma^2}$ , we have

$$\hat{\sigma}^2 \triangleq \frac{n-1}{2} \mathbb{E} \left( F^2 \mid (V_i)_{i \le n} \right) = \frac{1}{2(n-1)\sigma^2} \sum_{\substack{i \le n-1 \\ i \text{ even}}} (V_i^2 + 1) (V_{i+1} + V_{i-1} + M)^2.$$

The independence of variables  $(V_i)_{i \leq n}$  therefore implies that

$$\mathbb{E}(\hat{\sigma}^2) = \frac{1}{(n-1)\sigma^2} \sum_{\substack{i \le n-1 \\ i \text{ even}}} (2+M^2) = \frac{2\lfloor (n-1)/2 \rfloor}{n-1},$$

that  $|\mathbb{E}(\hat{\sigma}^2)-1| \leq \frac{1}{n-1},$  and hence, by Jensen's inequality and the triangle inequality, that

$$\|\frac{n-1}{2}\mathbb{E}(F^2 \mid S_n) - 1\|_2 \le \|\hat{\sigma}^2 - 1\|_2 \le \frac{1}{n-1} + \|\hat{\sigma}^2 - \mathbb{E}(\hat{\sigma}^2)\|_2.$$
(4.9)

Now consider the filtration  $(\mathcal{F}_i)_{i=1}^n$  with  $\mathcal{F}_i \triangleq \sigma(V_1, \ldots, V_i)$ . Boundedness and independence yield

$$\begin{aligned} |\mathbb{E}(\hat{\sigma}^2 \mid \mathcal{F}_i) - \mathbb{E}(\hat{\sigma}^2 \mid \mathcal{F}_{i-1})| &\leq \frac{1}{2(n-1)\sigma^2} \left[ (V_i^2 + 1)(V_{i+1} + V_{i-1} + M)^2 + (V_{i-1}^2 + 1)(V_{i-2} + V_i + M)^2 + (V_{i+1}^2 + 1)(V_{i+2} + V_i + M)^2 \right] &\leq \frac{3(R^2 + 1)(2R + M)^2}{2(n-1)\sigma^2} \end{aligned}$$

for each  $i \leq n$ . Since these martingale increments are uncorrelated,

$$\|\hat{\sigma}^2 - \mathbb{E}(\hat{\sigma}^2)\|_2 = \sqrt{\sum_{i=1}^n \mathbb{E}[(\mathbb{E}(\hat{\sigma}^2 \mid \mathcal{F}_i) - \mathbb{E}(\hat{\sigma}^2 \mid \mathcal{F}_{i-1}))^2]} \le \frac{3(R^2 + 1)(2R + M)^2 \sqrt{n}}{2(n-1)\sigma^2}.$$
 (4.10)

This estimate combined with (4.9) yields the second advertised inequality (4.6).

Next, fix  $k \ge 3$  and note that  $\|\mathbb{E}(F^k|S_n)\|_2 \le \|\mathbb{E}(F^k|(V_i)_{i\le n})\|_2 = \frac{\|\sum_{i\le n-1}\tilde{Z}_i^{k,i}\|_2}{n-1}$  for

$$\tilde{Z}_i^{k,t} \triangleq \frac{1}{\sqrt{n-1^k}\sigma^k} (V_i' - V_i)^k (V_{i+1} + V_{i-1} + M)^k \mathbb{I}(i \text{ is even})$$

Moreover, the triangle inequality, boundedness, and the fact that  $c(t) \geq \frac{R(2R+M)}{\sqrt{n-1}\sigma}$  imply

$$\begin{split} \|\mathbb{E}(\sum_{i \le n-1} \tilde{Z}_i^{t,k} \mid \mathcal{F}_l) - \mathbb{E}(\sum_{i \le n-1} \tilde{Z}_i^{t,k} \mid \mathcal{F}_{l-1})\|_2 \\ & \le \|\tilde{Z}_{l-1}^{t,k}\|_2 + \|\tilde{Z}_l^{t,k}\|_2 + \|\tilde{Z}_{l+1}^{t,k}\|_2 \le \frac{2^{k+1}\sigma}{\sqrt{n-1^k\sigma^k}} R^{k-1} (2R+M)^{k-1} \le \frac{2^{k+1}c(t)^{k-3}}{\sqrt{n-1^3\sigma^2}} R^2 (2R+M)^2 \end{split}$$

for each  $l \leq n$ . If k is odd, the inequality (4.7) then follows from the martingale variance decomposition as in (4.10). If k is even, we obtain (4.8) using the triangle inequality,

$$(n-1) \|\mathbb{E} (F^k \mid S_n)\|_2 \le \|\sum_{i \le n-1} \tilde{Z}_i^{k,t} - \mathbb{E} (\tilde{Z}_i^{k,t})\|_2 + \sum_{i \le n-1} \mathbb{E} (\tilde{Z}_i^{k,t}),$$

the martingale variance decomposition of  $\sum_{i \leq n-1} \tilde{Z}_i^{t,k} - \mathbb{E}(\tilde{Z}_i^{t,k})$ , and boundedness. Finally, we verify the condition (4.2). Fix any  $D \in (0, \sqrt{n})$ . Since n > 1, we have

$$|S'_n - S_n|^k = \frac{1}{\sqrt{n-1}^k \sigma^k} |V'_J - V_J| |V_{J+1} + M + V_{J-1}| \mathbb{I}(\mathbf{J} \text{ is even}) \le \frac{4^k R^k (2R+M)^k}{\sqrt{n}^k \sigma^k}, \quad \forall k \ge 1.$$

Since  $\sqrt{k!} \ge \lfloor k/2 \rfloor!$  for all  $k \ge 1$ , we find that, as advertised,

$$\begin{split} \sum_{k=1}^{\infty} \frac{(\sqrt{n}-D)^{k/2}}{\sqrt{k! D^{k/2}}} \mathbb{E}(|S'_n - S_n|^k) &\leq \sum_{k=1}^{\infty} \frac{(\sqrt{n}-D)^{k/2} 4^k R^k (2R+M)^k}{\sqrt{n^k \sigma^k} \sqrt{k! D^{k/2}}} &\leq \sum_{k=1}^{\infty} \frac{4^k R^k (2R+M)^k}{\sigma^k \sqrt{k! D^{k/2}}} \\ &\leq \sum_{k=1}^{\infty} \frac{4^k R^k (2R+M)^k}{\sigma^k \lfloor k/2 \rfloor! D^{k/2}} &\leq \sum_{k=0}^{\infty} \frac{1}{k!} \frac{4^{2k+1} R^{2k+1} (2R+M)^{2k+1}}{\sigma^{2k+1} D^{k+1/2}} + \sum_{k=0}^{\infty} \frac{1}{k!} \frac{4^{2k} R^{2k} (2R+M)^{2k}}{\sigma^{2k} D^k} \\ &= (1 + \frac{4R (2R+M)}{\sigma \sqrt{D}}) e^{\frac{16R^2 (2R+M)^2}{\sigma^2 D}} < \infty. \end{split}$$

### 4.3 Proof of Lem. 4.4: Large-time Hellinger bound

Using the definition of  $\tilde{\tau}_t$  and the triangle inequality we obtain the upper estimate

$$\begin{split} &\int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \left\| \frac{e^{-2t}}{1-e^{-2t}} (Z^2-1) - \frac{e^{-t}}{\sqrt{1-e^{-2t}}} S_n Z - \tilde{\tau}_t \right\|_2 \\ &\leq \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \frac{e^{-t} \|h_1(Z)\|_2}{\sqrt{1-e^{-2t}}} \|(n-1)\mathbb{E}(F|S_n) + S_n\|_2 dt \\ &+ \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \frac{e^{-2t} \|h_2(Z)\|_2}{1-e^{-2t}} \|\frac{n-1}{2}\mathbb{E}(F^2|S_n) - 1\|_2 dt \\ &+ \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \sum_{k\geq 3} \frac{e^{-tk} \|h_k(Z)\|_2}{k! (\sqrt{1-e^{-2t}})^k} (n-1) \|\mathbb{E}(F^k|S_n)\|_2 dt \triangleq (a_1) + (a_2) + (a_3). \end{split}$$

We will bound each of these terms in turn. First, by (4.5) of Lem. 4.3, we have

$$(a_1) \leq \frac{R^2}{\sqrt{n-1}\sigma} \sin^{-1}\left(\sqrt{1-\frac{D}{\sqrt{n}}}\right) \lessapprox \frac{R^2\pi}{2\sqrt{n-1}\sigma}.$$

Next, by (4.6) of Lem. 4.3 and the identity  $||h_2(Z)||_2^2 = \mathbb{E}[(Z^2 - 1)^2] = 2$ , we find that

$$(a_2) \le \|h_2(Z)\|_2 B_n \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \frac{e^{-2t}}{1-e^{-2t}} dt = \frac{\sqrt{2}B_n}{2} \int_0^{1-\frac{D}{\sqrt{n}}} \frac{1}{1-t} dt = \frac{\sqrt{2}B_n}{4} \log(n/D^2).$$

Now let  $k \ge 3$  be an odd integer and  $C_n \triangleq \frac{\sqrt{n}}{\sqrt{n-1}} \frac{2R^2(2R+M)^2}{\sigma^2}$ . By (4.7) of Lem. 4.3,

$$\begin{split} &\int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \frac{e^{-kt}}{\sqrt{1-e^{-2t^{k}}}} (n-1) \|\mathbb{E}\left(F^{k}|S_{n}\right)\|_{2} dt \leq \frac{C_{n}}{n-1} \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \frac{2^{k}e^{-tk}c(t)^{k-3}}{\sqrt{1-e^{-2t^{k}}}} dt \\ &= \frac{C_{n}2^{k}\sqrt{\kappa}^{k-3}}{n-1} \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \frac{e^{-tk}}{\sqrt{1-e^{-2t^{3}}}} dt = \frac{C_{n}\sqrt{\kappa}^{k-3}2^{k}}{n-1} \int_{0}^{\sqrt{1-\frac{D}{\sqrt{n}}}} \frac{x^{k-1}}{\sqrt{1-x^{2^{3}}}} dx. \end{split}$$

To further upper bound this, we will first propose a lower bound for  $\sqrt{2m+1!}$  for all  $m \in \mathbb{N} \setminus \{0\}$ . Indeed by using two applications of Stirling's approximation [Robbins, 1955] we observe that

$$\sqrt{(2m+1)!} \ge \sqrt{2m+1} \sqrt{\sqrt{2\pi(2m)} \cdot (2m/e)^{2m}} \cdot \exp\left(\frac{1}{12(2m)+1}\right)$$

$$\ge \sqrt{2m+1} e^{-19/300} 2^m m! / (\pi m)^{1/4}.$$
(4.11)

This directly implies that for  $\alpha \triangleq e^{19/300} \pi^{1/4}$  and all  $x \ge 1$ ,

$$\sum_{\substack{k \ge 3\\k \text{ odd}}} \frac{2^k x^{k-1} \sqrt{\kappa}^k}{\sqrt{k!}} \le \alpha \sqrt{\kappa} \sum_{k \ge 1} \frac{2^k x^{2k} \kappa^k k^{1/4}}{k! \sqrt{2k+1}} \le \frac{\alpha}{\sqrt{3}} \sqrt{\kappa} \left[ e^{2x^2 \kappa} - 1 \right].$$

Therefore, using Tonelli's theorem [Schilling, 2005, Thm. 13.8] to exchange the sum and integral and the Hermite inequality  $||h_k(Z)||_2 \leq \sqrt{k!}$  [Bonis, 2020, Lem. 3], we find that

$$\begin{split} &\sum_{k\geq 3} \frac{\|h_k(Z)\|_2}{k!} \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \frac{e^{-tk}}{\sqrt{1-e^{-2t^k}}} (n-1) \|\mathbb{E}(F^k|S_n)\|_2 dt \\ &\leq \frac{\alpha C_n}{\sqrt{3(n-1)\kappa}} \int_0^{\sqrt{1-\frac{D}{\sqrt{n}}}} \frac{(e^{2x^2\kappa}-1)}{\sqrt{1-x^{2^3}}} dx \lessapprox \sqrt{\frac{n}{n-1}} \frac{2R^2(2R+M)^2\alpha(e^{2\kappa}-1)n^{1/4}}{\sqrt{2D(n-1)\sigma^2\kappa}}. \end{split}$$

Next let  $k \ge 4$  be even and  $C'_n \triangleq \frac{\sqrt{n}}{\sqrt{n-1}} \frac{2R^3(2R+M)^3}{\sigma^3} + \frac{R^2(2R+M)^2\sqrt{n-1}}{2\sigma^2}$ . By (4.8),

$$\int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \frac{2^{k}e^{-tk}}{\sqrt{1-e^{-2t^{k}}}} (n-1) \|\mathbb{E}(F^{k}|S_{n})\|_{2} dt$$

$$\leq \frac{2^{k}\sqrt{\kappa}^{k-4}C_{n}'}{(n-1)^{3/2}} \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})}^{\infty} \frac{e^{-tk}}{\sqrt{1-e^{-2t^{4}}}} dt \leq \frac{2^{k}\sqrt{\kappa}^{k-4}C_{n}'}{(n-1)^{3/2}} \int_{0}^{\sqrt{1-\frac{D}{\sqrt{n}}}} \frac{x^{k-1}}{\sqrt{1-x^{2^{4}}}} dx.$$

The inequality (4.11) further implies that

$$\sum_{\substack{k \ge 4\\k \text{ even}}} \frac{2^k x^{k-1} \sqrt{\kappa^k}}{\sqrt{k!}} \le \sum_{k \ge 2} \frac{2^{2k} x^{2k-1} \kappa^k}{\sqrt{2k!}} \le 2^{1/4} \kappa \alpha (e^{2x^2 \kappa} - 1).$$

Hence, we find that, by Tonelli's theorem,

$$\begin{split} &\sum_{\substack{k \ge 4\\k \text{ is even}}} \frac{\|h_k(Z)\|_2}{k!} \int_{-\frac{1}{2}\log(1-\frac{D}{\sqrt{n}})} \frac{e^{-tk}}{\sqrt{1-e^{-2t^k}}} (n-1) \|\mathbb{E}(F^k|S_n)\|_2 dx \\ &\le 2\alpha \frac{C'_n}{\kappa\sqrt{n-1^3}} \int_0^{\sqrt{1-\frac{D}{\sqrt{n}}}} \frac{(e^{2x^2\kappa}-1)}{\sqrt{1-x^{2^4}}} dx. \end{split}$$

Combining our results, we find that  $(a_1) + (a_2) + (a_3) \le \beta$ , implying the result.  $\Box$ 

# 5 Proof of Lem. 2.4: Hellinger concentration

Write  $f_S$  for the Lebesgue density of S and define  $h \triangleq f_S/\varphi$ . The first claim follows from the triangle inequality and Cauchy-Schwarz as

$$\begin{split} |P(S>u) - \Phi^c(u)| &= |\mathbb{E}((h(Z) - 1)\mathbb{I}(Z>u))| = |\mathbb{E}((\sqrt{h(Z)} - 1)(\sqrt{h(Z)} + 1)\mathbb{I}(Z>u))| \\ &\leq |\mathbb{E}((\sqrt{h(Z)} - 1)\mathbb{I}(Z>u))| + |\mathbb{E}((\sqrt{h(Z)} - 1)\sqrt{h(Z)}\mathbb{I}(Z>u))| \\ &\leq \sqrt{\mathbb{E}((\sqrt{h(Z)} - 1)^2)} [\sqrt{\mathbb{E}(\mathbb{I}(Z>u))} + \sqrt{\mathbb{E}(h(Z)\mathbb{I}(Z>u))}] \\ &= \sqrt{\int_{-\infty}^{\infty} (\sqrt{f_S(x)} - \sqrt{\varphi(x)})^2 dx} [\sqrt{\Phi^c(u)} + \sqrt{\mathbb{P}(S>u)}] \\ &= \sqrt{2}H(S, Z) [\sqrt{\Phi^c(u)} + \sqrt{P(S>u)}]. \end{split}$$

The second claim follows analogously as

$$|P(|S| > u) - 2\Phi^{c}(u)| = |\mathbb{E}((h(Z) - 1)\mathbb{I}(|Z| > u))| \le \sqrt{2}H(S, Z)\left[\sqrt{2\Phi^{c}(u)} + \sqrt{P(|S| > u)}\right].$$

# 6 Additional Lemmas

**Lemma 6.1** (Score function properties). Let S be a random variable with convex support  $\Omega_S \subseteq \mathbb{R}$  and Lebesgue density  $f_S \in C^1$ . The following statements hold true.

- 1. Any first-order score function or second-order score function  $I_S$  for S is S-almost everywhere unique amongst elements of  $L^1(S)$  satisfying (2.1).
- 2. If  $f_S$  is twice differentiable with  $\mathfrak{s}_S \triangleq f'_S/f_S \in L^1(S), I_S \triangleq f''_S/f_S \in L^1(S)$ , and

$$\lim_{x \to \sup \Omega_S} \max(|f_S(x)|, |f'_S(x)|) = \lim_{x \to \inf \Omega_S} \max(|f_S(x)|, |f'_S(x)|) = 0,$$
(6.1)

then  $\mathfrak{s}_S$  and  $I_S$  are first and second-order score functions for S.

*Proof.* Suppose  $g, \tilde{g} \in L^1(S)$  are both score functions for S, fix  $\epsilon > 0$ , and choose  $\delta > 0$  so that  $\mathbb{E}(|g(S) - \tilde{g}(S)|\mathbb{I}(|S| > \delta)) < \epsilon$ . Since any measurable compactly supported indicator function can be arbitrarily approximated by  $C_c^{\infty}$  functions, select  $\Psi_{\epsilon,\delta} \in C_c^{\infty}$  satisfying

$$\|\Psi_{\epsilon,\delta}(S) - \mathbb{I}(g(S) > \tilde{g}(S), |S| \le \delta)\|_{\infty} \le \epsilon.$$

so that

$$\mathbb{E}(\max(g(S) - \tilde{g}(S), 0)) = \mathbb{E}([g(S) - \tilde{g}(S)] \mathbb{I}(g(S) > \tilde{g}(S)))$$
  
$$\leq \epsilon + \mathbb{E}([g(S) - \tilde{g}(S)] \Psi_{\epsilon,\delta}(S)) + \mathbb{E}(|g(S) - \tilde{g}(S)| \mathbb{I}(|S| > \delta)) \leq 2\epsilon.$$

Since  $\epsilon > 0$  was arbitrary we have  $g(S) \leq \tilde{g}(S)$  almost surely and, by symmetry,  $g(S) \geq \tilde{g}(S)$  almost surely, proving the first claim. Identical reasoning yields the claim for second-order scores.

Now suppose that  $f_S$  is twice differentiable with  $\mathfrak{s}_S, I_S \in L^1(S)$  and (6.1) holding. For all  $\Psi \in C_c^{\infty}$ , integration by parts, the boundedness of  $\Psi$ , and our vanishing assumption (6.1) imply that

$$\mathbb{E}(\frac{f'_{S}(S)}{f_{S}(S)}\Psi(S)) = \int_{\Omega_{S}} f'_{S}(x)\Psi(x)dx = f_{S}(x)\Psi(x)\Big|_{\inf\Omega_{S}}^{\sup\Omega_{S}} - \int_{\Omega_{S}} f_{S}(x)\Psi'(x)dx = -\mathbb{E}(\Psi'(S))$$
$$\mathbb{E}(\frac{f''_{S}(S)}{f_{S}(S)}\Psi(S)) = \int_{\Omega_{S}} f''_{S}(x)\Psi(x)dx = f'_{S}(x)\Psi(x)\Big|_{\inf\Omega_{S}}^{\sup\Omega_{S}} - \int_{\Omega_{S}} f'_{S}(x)\Psi'(x)dx$$
$$= -\mathbb{E}(\frac{f'_{S}(S)}{f_{S}(S)}\Psi'(S)) = \mathbb{E}(\Psi''(S)).$$

Hence  $f'_S/f_S$  and  $f''_S/f_S$  are first and second-order score functions.

**Lemma 6.2** (Second-order score bound). Suppose  $S \stackrel{d}{=} S_1 + S_2$  for independent random variables  $S_1, S_2$  with first-order score functions  $\mathfrak{s}_{S_1} \in L^2(S_1)$  and  $\mathfrak{s}_{S_2} \in L^2(S_2)$ . Then S admits a second-order score function  $I_S \in L^2(S)$  satisfying

$$||I_S(S)||_2 \le ||\mathfrak{s}_{S_1}(S_1)||_2 ||\mathfrak{s}_{S_2}(S_2)||_2$$

*Proof.* Fix any  $\Psi \in C_c^{\infty}$ . We invoke the tower property, the independence of  $S_1$  and  $S_2$ , and the first-order score function definition (2.1) in turn to discover that

$$\mathbb{E}(\mathbb{E}(\mathfrak{s}_{S_1}(S_1)\mathfrak{s}_{S_2}(S_2)|S_1 + S_2)\Psi(S_1 + S_2))$$
  
=  $\mathbb{E}(\mathfrak{s}_{S_1}(S_1)\mathfrak{s}_{S_2}(S_2)\Psi(S_1 + S_2))$   
=  $-\mathbb{E}(\mathfrak{s}_{S_1}(S_1)\Psi'(S_1 + S_2)) = \mathbb{E}(\Psi''(S_1 + S_2)).$ 

Hence  $I_S(S) = \mathbb{E}(\mathfrak{s}_{S_1}(S_1)\mathfrak{s}_{S_2}(S_2)|S)$  satisfies the second-order score definition (2.1). Finally, by Jensen's inequality and the independence of  $S_1$  and  $S_2$ , we have

$$||I_S(S)||_2 = ||\mathbb{E}(\mathfrak{s}_{S_1}(S_1)\mathfrak{s}_{S_2}(S_2)|S_1 + S_2)||_2 \le ||\mathfrak{s}_{S_1}(S_1)\mathfrak{s}_{S_2}(S_2)||_2 = ||\mathfrak{s}_{S_1}(S_1)||_2 ||\mathfrak{s}_{S_2}(S_2)||_2.$$

**Lemma 6.3** (Fundamental theorem precondition). Under the conditions of Thm. 2.1,  $t \to \sqrt{f_t^S(x)}$  is continuous on  $[0,\infty)$  and continuously differentiable on  $(0,\infty)$  for all  $x \in \Omega_S$ .

*Proof.* Fix any  $x \in \Omega_S$ . By Pavliotis [2014, Thm. 4.1],  $t \mapsto f_t^S(x)$  is continuous on  $[0, \infty)$  and continuously differentiable on  $(0, \infty)$ . Therefore,  $t \mapsto \sqrt{f_t^S(x)}$  is continuous on  $[0, \infty)$ .

Next, fix any t > 0. To conclude, it remains to show that  $f_t^S(x) > 0$  so that  $t \mapsto \partial_t \sqrt{f_t^S(x)}$  is also continuously differentiable on  $(0, \infty)$ . Since  $f_S$  is continuous, there exists a  $\delta > 0$  for which

$$\inf_{y \in [x-\delta, x+\delta]} f_S(y) > f_S(x)/2.$$

Now consider the interval  $M_x \triangleq x(1-e^{-t}) + \delta e^{-t}\sqrt{1-e^{-2t}}[-1,1]$ . Since  $f_S(x) > 0$ ,

$$\begin{split} f_t^S(x) &= \int_{\mathbb{R}} \frac{\varphi(y/\sqrt{1-e^{-2t}})}{\sqrt{1-e^{-2t}}} \frac{f_S((x-y)/e^{-t})}{e^{-t}} dy \ge \int_{M_x} \frac{\varphi(y/\sqrt{1-e^{-2t}})}{\sqrt{1-e^{-2t}}} \frac{f_S((x-y)/e^{-t})}{e^{-t}} dy \\ &\stackrel{(a)}{\ge} \frac{\varphi(|x|+\delta)}{\sqrt{1-e^{-2t}}} \int_{M_x} \frac{f_S((x-y)/e^{-t})}{e^{-t}} dy \stackrel{(b)}{\ge} \varphi(|x|+\delta) \delta f_S(x) > 0, \end{split}$$

as  $|y/\sqrt{1-e^{-2t}}| \le |x| + \delta$  for all  $y \in M_x$ , yielding (a), and  $|(x-y)/e^{-t} - x| \le \delta$  for all  $y \in M_x$  so that  $f_S((x-y)/e^{-t}) \ge f_S(x)/2$ , yielding (b).

Lemma 6.4 (Fubini precondition). Under the conditions of Thm. 2.1, we have

$$\int_0^T \int_{\Omega_S} \sqrt{f_S(x)} |\partial_t \sqrt{f_t^S(x)}| dx dt < \infty \quad \text{for all} \quad T > 0.$$
(6.2)

*Proof.* Fix any T > 0. The identities (3.1), (3.2), and (3.3), show that

$$|\partial_t \sqrt{f_t^S(x)}| = \frac{\sqrt{f_t^S(x)}}{2} |I_t^S(x) + x \mathfrak{s}_{X_t^S}(x) + 1| \quad \text{for all} \quad t \in [0, T] \quad \text{and} \quad x \in \mathbb{R}.$$

Hence we obtain that

$$\int_{0}^{T} \int_{\Omega_{S}} \sqrt{f_{S}(x)} |\partial_{t} \sqrt{f_{t}^{S}(x)}| \, dx \, dt = \int_{0}^{T} \mathbb{E} \Big( \frac{\sqrt{f_{S}(X_{t}^{S})}}{2\sqrt{f_{t}^{S}(X_{t}^{S})}} \Big| I_{t}^{S}(X_{t}^{S}) + X_{t}^{S} \mathfrak{s}_{X_{t}^{S}}(X_{t}^{S}) + 1 \Big| \Big) dt.$$

Moreover, the almost sure identities (3.4) and (3.5) and Cauchy-Schwarz imply

$$\mathbb{E}\left(\frac{\sqrt{f_{S}(X_{t}^{S})}}{2\sqrt{f_{t}^{S}(X_{t}^{S})}}\Big|I_{t}^{S}(X_{t}^{S}) + X_{t}^{S}\mathfrak{s}_{X_{t}^{S}}(X_{t}^{S}) + 1\Big|\right) \\
\leq \left\|\frac{\sqrt{f_{S}(X_{t}^{S})}}{2\sqrt{f_{t}^{S}(X_{t}^{S})}}\right\|_{2}\left[1 + e^{2T}\|I_{S}(S)\|_{2} + 2e^{T}\|\mathfrak{s}_{S}(S)\|_{2} + e^{T}\|\mathfrak{s}_{S}(S)S\|_{2}\right]$$

Since  $I_S, \mathfrak{s}_S \in L^2(S)$  and  $\|\mathfrak{s}_S(S)S\|_2 < \infty$  the result (6.2) follows from the observation

$$\left\|\frac{\sqrt{f_S(X_t^S)}}{2\sqrt{f_t^S(X_t^S)}}\right\|_2 = \frac{1}{2}\sqrt{\int_{\Omega_S} f_S(x)dx} = \frac{1}{2}.$$

## References

- S. Artstein, K. M. Ball, F. Barthe, and A. Naor. On the rate of convergence in the entropic central limit theorem. *Probability theory and related fields*, 129(3):381–390, 2004.
- M. Austern and L. Mackey. Efficient concentration with Gaussian approximation. arXiv preprint arXiv:2208.09922, 2022.
- S. G. Bobkov, G. P. Chistyakov, and F. Götze. Rate of convergence and edgeworth-type expansion in the entropic central limit theorem. *The Annals of Probability*, pages 2479–2512, 2013.

- S. G. Bobkov, G. P. Chistyakov, and F. Götze. Berry-Esseen bounds in the entropic central limit theorem. *Probability Theory and Related Fields*, 159(3-4):435–478, 2014.
- T. Bonis. Stein's method for normal approximation in wasserstein distances with application to the multivariate central limit theorem. *Probability Theory and Related Fields*, 178(3):827–860, 2020.
- E. Hellinger. Neue begründung der theorie quadratischer formen von unendlichvielen veränderlichen. *Journal für die reine und angewandte Mathematik*, 1909(136):210–271, 1909. doi: doi:10.1515/crll.1909.136.210. URL https://doi.org/10.1515/crll.1909.136.210.
- A. W. Knapp. Basic real analysis. Springer Science & Business Media, 2007.
- S. Krantz and H. Parks. A Primer of Real Analytic Functions. A Primer of Real Analytic Functions. Birkhäuser Boston, 2002. ISBN 9780817642648.
- M. Ledoux, I. Nourdin, and G. Peccati. Stein's method, logarithmic Sobolev and transport inequalities. *Geometric and Functional Analysis*, 25(1):256–306, 2015.
- G. A. Pavliotis. Stochastic processes and applications. Texts in applied mathematics, 60, 2014.
- H. Robbins. A remark on Stirling's formula. *The American Mathematical Monthly*, 62(1):26–29, 1955.
- R. L. Schilling. Measures, Integrals and Martingales. Cambridge University Press, 2005.
- C. Stein. Approximate computation of expectations. *Lecture Notes-Monograph Series*, 7:i–164, 1986. ISSN 07492170. URL http://www.jstor.org/stable/4355512.