Self-orthogonal codes from plateaued functions

Peng Wang^{a,b}, Ziling Heng^{a,b,∗}

^aSchool of Science, Chang'an University, Xi'an 710064, Chin a ^bNational Mobile Communications Research Laboratory, Southeast University, Nanjing 211111, China

Abstract

Self-orthogonal codes are of interest as they have important applications in quantum codes, lattices and many areas. In this paper, based on the weakly regular plateaued functions or plateaued Boolean functions, we construct a family of linear codes with four nonzero weights. This family of linear codes is proved to be not only self-orthogonal but also optimally or almost optimally extendable. Besides, we derive binary and ternary linearly complementary dual codes (LCD codes for short) with new parameters from this family of codes. Some families of self-dual codes are also obtained as byproducts.

Keywords: Weakly regular plateaued function, self-orthogonal code, LCD code, optimally extendable code *2000 MSC:* 94B05, 94A05

1. Introduction

Let \mathbb{F}_q be the finite field with q elements, where q is a power of a prime p. Let $\mathbb{F}_q^* := \mathbb{F}_q \setminus \{0\}$.

1.1. Linear codes and self-orthogonal codes

Let *n*, *k*, *d* be three positive integers. For a nonempty set $C \subseteq \mathbb{F}_q^n$, if *C* is a *k*-dimensional linear subspace of \mathbb{F}_q^n , then it is called an $[n, k, d]$ linear code over \mathbb{F}_q , where d denotes its minimum distance. Let \overrightarrow{A}_i be the number of codewords with weight *i* in a linear code C of length *n*, where $0 \le i \le n$. The sequence $(1, A_1, A_2, \dots, A_n)$ is called the weight distribution of C and the polynomial $A(z) = 1 + A_1 z + A_2 z^2 + \cdots + A_n z^n$ is referred to as the weight enumerator of C. Weight distribution is a significant research subject as it not only describes the error detection and error correction abilities of the code, but also can be used to calculate the error probabilities of error detection and correction of the code. The weight distributions of some linear codes were studied in the literature [\[11,](#page-22-0) [20,](#page-23-0) [32,](#page-23-1) [33\]](#page-23-2).

Optimal linear codes play an important role in both theory and practice. An $[n, k, d]$ linear code C over \mathbb{F}_q is said to be optimal if there does not exist $[n, k, d+1]$ linear code over \mathbb{F}_q and

1

[⋆]This research was supported in part by the National Natural Science Foundation of China under Grant 12271059, in part by the open research fund of National Mobile Communications Research Laboratory of Southeast University under Grant 2024D10 and in part by the Shaanxi Fundamental Science Research Project for Mathematics and Physics (Grant No. 23JSZ008).

[∗]Corresponding author

Email addresses: wp20201115@163.com (Peng Wang), zilingheng@chd.edu.cn (Ziling Heng)

almost optimal if there exists an $[n, k, d+1]$ optimal code over \mathbb{F}_q . The following is the well-known sphere-packing bound on the parameters of codes.

Lemma 1. *[\[18](#page-23-3), Sphere-packing bound] Let M be the maximum number of codewords in a code over* F*^q of length n and minimum distance d. Then*

$$
M \le \frac{q^m}{\sum\limits_{i=0}^t \binom{m}{i} (q-1)^i},
$$

where t = $\left| \frac{d-1}{2} \right|$ *and* $\left| \cdot \right|$ *denotes the floor function.*

Define the dual code of an $[n, k]$ linear code C by

$$
C^{\perp} = \left\{ \mathbf{u} \in \mathbb{F}_q^n : \langle \mathbf{u}, \mathbf{v} \rangle = \mathbf{0} \text{ for all } \mathbf{v} \in C \right\},\
$$

where $\langle \cdot \rangle$ denotes the standard inner product. Then C^{\perp} is a linear code with length *n* and dimension *n*−*k*. If $C \subseteq C^{\perp}$, then *C* is called a self-orthogonal code. In particular, if $C = C^{\perp}$, then *C* is said to be self-dual. Self-orthogonal codes have nice applications in lattices [\[37\]](#page-24-0), linear complementary dual codes (LCD codes for short) [\[26](#page-23-4)], quantum codes [\[6](#page-22-1)] and so on. For a given *q*-ary linear code *C*, it is natural to study whether it is self-orthogonal or not. If $q = 2,3$, the following provides simple conditions for a linear code to be self-orthogonal by the divisibility of its weights.

Lemma 2. [\[18](#page-23-3)*, Theorems 1.4.8 and 1.4.10] Let C be a linear code over* \mathbb{F}_p *. For* $p = 2$ *, if every codeword of C* has weight divisible by four, then *C* is self-orthogonal. For $p = 3$, *C* is self*orthogonal if and only if every codeword of C has weight divisible by three.*

Recently, Li and Heng established a sufficient condition for a *q*-ary linear code containing the all-1 vector to be self-orthogonal if *q* is a power of an odd prime [\[19](#page-23-5)].

Lemma 3. *[\[19](#page-23-5)] Let q be a power of p. where p is an odd prime. Let C be an* [*n*, *k*,*d*] *linear code over* \mathbb{F}_q *with* $1 \in \mathcal{C}$ *, where* 1 *is all-1 vector of length n. If* \mathcal{C} *is p-divisible, then* \mathcal{C} *is self-orthogonal.*

The following lemmas show that we can derive self-dual codes from self-orthogonal codes. These self-dual codes are the subcodes of the duals of the self-orthogonal codes.

Lemma 4. [\[4](#page-22-2), Proposition 4.5] Let q be a power of 2. A linear code $C \subseteq \mathbb{F}_q^n$ of dimension $k \geq \frac{n}{2}$ 2 *contains a self-dual subcode if and only if n is even and* C^{\perp} *is self-orthogonal, that is,* $C^{\perp} \subseteq C$ *.*

Lemma 5. [\[4,](#page-22-2) Proposition 4.6] Let $C \subseteq \mathbb{F}_q^n$ be a linear code of dimension $k \geq \frac{n}{2}$ $\frac{n}{2}$.

- *(1)* If $q \equiv 1 \pmod{4}$, then *C* contains a self-dual subcode if and only if n is even and C^{\perp} is self*orthogonal.*
- *(2)* If $q \equiv 3 \pmod{4}$, then *C* contains a self-dual subcode if and only if $n \equiv 0 \pmod{4}$ and C^{\perp} is *self-orthogonal.*

1.2. LCD codes

An $[n, k, d]$ linear code *C* is called an Euclidean LCD code if $C \cap C^{\perp} = \{0\}$, where 0 denotes the zero vector of length *n*. In [\[24](#page-23-6)], Massey introduced the definition and gave the algebraic characterization of LCD codes. He also proved the existence of asymptotically good LCD codes and provided an optimum linear coding solution for the two-user binary adder channel. In [\[31\]](#page-23-7), Sendrier showed that LCD codes satisfy the asymptotic Gilbert-Varshamov bound. LCD codes have nice applications in lattices, network coding, and multisecret-sharing schemes [\[1](#page-22-3), [3,](#page-22-4) [13\]](#page-22-5). Besides, LCD codes also have application in counter-measures to passive and active side channel analyses on embedded cryptosystems. Many families of LCD codes have been constructed in [\[34,](#page-24-1) [23](#page-23-8)]. The existence of *q*-ary (*q* > 3) Euclidean LCD codes has been studied in [\[8\]](#page-22-6). However, the existence of binary and ternary Euclidean LCD codes has not been totally investigated.

Self-orthogonal codes can be used to construct LCD codes. It is known that a generator matrix *G* of an [*n*, *k*, *d*] self-orthogonal code satisfies $GG^T = \mathbf{0}_{k,k}$, where $\mathbf{0}_{k,k}$ denotes the zero matrix of size $k \times k$. In other words, G is row-self-orthogonal. The following lemma shows how to construct leading-systematic codes from a row-self-orthogonal matrix, where a linear code is said to be leading-systematic if and only if it has a generator matrix of the form $G = [I : P]$ for an identity matrix *I*.

Lemma 6. [\[26](#page-23-4)] A leading-systematic linear code C with systematic generator matrix $G := [I : P]$ *is an LCD code if (but not only if) the matrix P is row-self-orthogonal, where I denotes an identity matrix.*

Note that a self-orthogonal code has many different generator matrices. If P_1 and P_2 are two different generator matrices of a self-orthogonal code *C*, then $G_1 := [I : P_1]$ and $G_2 := [I : P_2]$ may generate two inequivalent LCD codes. Finding a suitable generator matrix *P* such that $G = [I : P]$ generate an LCD code with large minimum distance, whose dual also have large minimum distance, is important.

1.3. Optimally and almost optimally extendable codes

In [\[2\]](#page-22-7), Bringer et al. introduced a direct sum masking (DSM) countermeasure to oppose side channel attack (SCA) and fault injection attack (FIA) which have brought great threats to the implementation of block cipher. It needs two linear codes *C* and *D* satisfying $C \oplus D = \mathbb{F}_q^n$, where C and D are used to encode the sensitive data and encode random data, respectively. In particular, if $\mathcal{D} = C^{\perp}$, then *C* is an LCD code. For a general pair (C, \mathcal{D}) satisfying $C \oplus \mathcal{D} = \mathbb{F}_q^n$, it is desirable to change C and D into C' and D' to protect the sensitive data stored in the register from SCA and FIA, where C' is obtained by adding k bits of zeros to the end of each codeword in *C* and \mathcal{D}' is derived by appending the identity matrix at the end of the generator matrix of \mathcal{D}' . If $d(\mathcal{D}^{\perp}) = d(\mathcal{D}^{\perp})$, then $\mathcal D$ is said to be *optimally extendable*. $\mathcal D$ is said to be *almost optimally extendable* if $d(D^{\perp}) = d(D^{\perp}) - 1$.

In the literature, there are only a few known constructions of optimally or almost optimally extendable codes. Carlet, Li and Mesnager gave two families of optimally extendable linear codes and two families of almost optimally extendable linear codes from the first-order Reed-Muller codes and irreducible cyclic codes of dimension two [\[7\]](#page-22-8). Quan, Yue and Hu studied (almost) optimally extendable linear codes from irreducible cyclic codes, MDS codes and NMDS codes [\[29\]](#page-23-9). In [\[36](#page-24-2)], Wang, Heng, Li and Yue constructed some families of almost optimally extendable

codes from self-orthogonal codes. In [\[12\]](#page-22-9), Heng, Li, Wu and Wang constructed two families of optimally or almost optimally extendable codes from some functions over finite fields.

1.4. The purpose of this paper

The purpose of this paper is to construct linear codes from weakly regular plateaued functions over finite fields. Let *p* be an odd prime and $f(x)$ be a function from \mathbb{F}_q to \mathbb{F}_p with $f(0) = 0$. We define a *p*-ary linear code as

$$
\overline{C_f} = \left\{ \mathbf{c}_{(a,b,c)} = (af(x) + \text{Tr}_{q/p}(bx) + c)_{x \in \mathbb{F}_q} : a \in \mathbb{F}_p, b \in \mathbb{F}_q, c \in \mathbb{F}_p \right\}.
$$
 (1)

We remark that $\overline{C_f}$ is the augmented code of the extended code of

$$
\mathcal{C}_f^* = \left\{ \mathbf{c}_{(a,b)} = (af(x) + \text{Tr}_{q/p}(bx))_{x \in \mathbb{F}_q^*} : a \in \mathbb{F}_p, b \in \mathbb{F}_q \right\}.
$$
 (2)

When *f* is a weakly regular bent function, the code $\overline{C_f}$ was studied in [\[12](#page-22-9)].

In this paper, we mainly study the parameters and weight distribution of $\overline{C_f}$ and prove the following if *f* is a weakly regular plateaued function:

- $\overline{C_f}$ is self-orthogonal;
- $\overline{C_f}$ is optimally or almost optimally extendable if we select a suitable generator matrix of $\overline{C_f}$.

Besides, we also derive ternary LCD codes from $\overline{C_f}$. Moreover, if f is a plateaued Boolean function, we also prove that $\overline{C_f}$ is self-orthogonal and determine their parameters and weight distribution. The binary code $\overline{C_f}$ is proved to be almost optimally extendable. We also use the binary code $\overline{C_f}$ to construct a family of binary LCD codes. Finally, we obtain some families of self-dual codes from the duals of the self-orthogonal codes in this paper.

2. Preliminaries

In this section, we present some mathematical foundations which will be used to give the main results in this paper.

2.1. Characters and Gaussian sums over finite fields

Let $q = p^m$ with p a prime. Let ζ_p denote the primitive p-th root of complex unity. Define an additive character χ of \mathbb{F}_q by the homomorphism from the additive group \mathbb{F}_q to the complex multiplicative group \mathbb{C}^* such that

$$
\chi(x+y) = \chi(x)\chi(y), \ x, y \in \mathbb{F}_q.
$$

For any $a \in \mathbb{F}_q$, an additive character of \mathbb{F}_q is defined by the function $\chi_a(x) = \zeta_p^{\text{Tr}_{q/p}(ax)}$, $x \in \mathbb{F}_q$, where $Tr_{q/p}(x)$ is the trace function from \mathbb{F}_q to \mathbb{F}_p . By definition, we have $\chi_a(x) = \chi_1(ax)$. Particularly, χ_0 is called the trivial additive character of \mathbb{F}_q and χ_1 is called the canonical additive character of \mathbb{F}_q . The following is the orthogonal relation of additive characters (see [\[21](#page-23-10)]):

$$
\sum_{x \in \mathbb{F}_q} \chi_a(x) = \begin{cases} q & \text{if } a = 0, \\ 0 & \text{if } a \in \mathbb{F}_q^*.\end{cases}
$$

Let α be a primitive element of \mathbb{F}_q . The multiplicative character ψ of \mathbb{F}_q is defined as the homomorphism from the multiplicative group \mathbb{F}_q^* to the complex multiplicative group \mathbb{C}^* such that

$$
\Psi(xy) = \Psi(x)\Psi(y), \ x, y \in \mathbb{F}_q^*.
$$

The function $\psi_j(\alpha^k) = \zeta_{a}^{jk}$ q_{q-1}^{k} , where $k = 0, 1, \dots, q-2$, is a multiplicative character for each $j = 1$ 0,1,...,*q*−2. In particular, ψ⁰ is called the trivial multiplicative character and η := ψ*^q*−¹ is referred to as the quadratic multiplicative character of \mathbb{F}_q if *q* is odd. The following is the orthogonal relation of multiplicative characters (see [\[21](#page-23-10)]):

$$
\sum_{x \in \mathbb{F}_q^*} \Psi_j(x) = \begin{cases} q - 1 & \text{if } j = 0, \\ 0 & \text{if } j \neq 0. \end{cases}
$$

For an additive character χ and a multiplicative character ψ of \mathbb{F}_q , the *Gaussian sum* $G(\psi, \chi)$ over \mathbb{F}_q is defined by

$$
G(\psi,\chi)=\sum_{x\in\mathbb{F}_q^*}\psi(x)\chi(x).
$$

Specially, $G(\eta, \chi)$ is called the *quadratic Gaussian sum* over \mathbb{F}_q for nontrivial χ . The explicit value of quadratic *Gaussian sum* is given in the following.

Lemma 7. [\[21,](#page-23-10) Theorem 5.15] Let $q = p^m$ with p odd. Then

$$
G(\eta, \chi_1) = (-1)^{m-1} (\sqrt{-1})^{(\frac{p-1}{2})^2 m} \sqrt{q}.
$$

2.2. Weakly regular plateaued functions and the cyclotomic fields

Let $f(x)$ be a *p*-ary function from \mathbb{F}_{p^m} to \mathbb{F}_p . Define the Walsh transform of $f(x)$ as follows:

$$
\mathrm{W}_f(\beta):=\sum_{x\in\mathbb{F}_{p^m}}\zeta_p^{f(x)-\mathrm{Tr}_{p^m/p}(\beta x)},\ \beta\in\mathbb{F}_{p^m}.
$$

Then *f* is said to be *balanced* over \mathbb{F}_p provided that *f* takes every value of \mathbb{F}_p exactly p^{m-1} times. Otherwise, *f* is said to be *unbalanced*. It is known that *f* is balanced if and only if $W_f(0) = 0$. A function *f* is referred to as a *bent* function if $|W_f(\beta)|^2 = p^m$ for any $\beta \in \mathbb{F}_{p^m}$. Besides, *f* is said to be *s-plateaued* provided that $| W_f(\beta) |^2 \in \{0, p^{m+s}\}$ for any $\beta \in \mathbb{F}_{p^m}$, with $0 \le s \le m$. For an *s*-plateaued function *f*, its *Walsh support* is defined by the set $S_f = \{ \beta \in \mathbb{F}_p^n : ||W_f(\beta)||^2 = p^{m+s} \}.$

Lemma 8. [\[27\]](#page-23-11) Let f be an s-plateaued function from \mathbb{F}_{p^m} to \mathbb{F}_p . Then for $\beta \in \mathbb{F}_{p^m}$, $|W_f(\beta)|^2$ *takes the values* p^{m+s} *and* 0 *for the times* p^{m+s} *and* $p^m - p^{m-s}$ *, respectively.*

Lemma 9. [\[27\]](#page-23-11) Let f be s-plateaued from \mathbb{F}_{p^m} to \mathbb{F}_p . Define the sets $Z(W_f) := \{(\alpha, \beta) \in \mathbb{F}_p^* \times \mathbb{F}_{p^m}$: $W_f(\alpha^{-1}\beta) = 0$ } *and* $S(W_f) := \{(\alpha, \beta) \in \mathbb{F}_p^* \times \mathbb{F}_{p^m} : W_f(\alpha^{-1}\beta) \neq 0\}$. Then, the sizes of $Z(W_f)$ and *S*(*W_f*) *are respectively equal to* $(p-1)(p^m-p^{m-s})$ *and* $(p-1)p^{m-s}$.

In particular, for $p = 2$, the Walsh distribution of plateaued Boolean functions is given in the following [\[9\]](#page-22-10),[\[10\]](#page-22-11).

Lemma 10. *Let m*+*s be an even integer with* 0 ≤ *s* ≤ *m and f be an s-plateaued Boolean function* with $f(0) = 0$ from \mathbb{F}_{2^m} to \mathbb{F}_2 . Then, for $\beta \in \mathbb{F}_{2^m}$, the Walsh distribution of f is given by

$$
W_f(\beta) = \begin{cases} 2^{\frac{m+s}{2}}, & 2^{m-s+1} + 2^{\frac{m-s-2}{2}} \text{ times,} \\ 0, & 2^m - 2^{m-s} \text{ times,} \\ -2^{\frac{m+s}{2}}, & 2^{m-s-1} - 2^{\frac{m-s-2}{2}} \text{ times.} \end{cases}
$$

Definition 11. *Let p be an odd prime and f be a p-ary s-plateaued function from* F*^p ^m to* F*p, where s* is an integer with $0 \le s \le m$. Then f is said to be weakly regular provided that there exists a *complex number u with unit magnitude (in fact,* | *u* |= 1 *and u does not depend on* β*) such that*

$$
W_f(\beta) \in \left\{ 0, u p^{\frac{m+s}{2}} \zeta_p^{f^*(\beta)} \right\},\tag{3}
$$

for all $\beta \in \mathbb{F}_{p^m}$, where f^* is a p-ary function over \mathbb{F}_{p^m} with $f^*(\beta) = 0$ for every $\beta \in \mathbb{F}_{p^m} \setminus S_f$. *Otherwise, f is said to be non-weakly regular.*

Lemma 12. Let f be a weakly regular s-plateaued function and $p^* = (-1)^{\frac{p-1}{2}}p$. Then, for all $\beta \in S_f$ *, we have*

$$
W_f(\beta) = \varepsilon \sqrt{p^*}^{m+s} \zeta_p^{f^*(\beta)},\tag{4}
$$

where $\varepsilon = \pm 1$ is the sign of W_f and f^* is a p-ary function over S_f .

Note that the weakly regular 0-plateaued function is the weakly regular bent function. Let *f* be a weakly regular *s*-plateaued function satisfying two homogenous conditions: $f(0) = 0$ and $f(ax) =$ $a^h f(x)$ for all $x \in \mathbb{F}_{p^m}$ and $a \in \mathbb{F}_p^*$, where *h* is an even positive integer with $gcd(h-1, p-1) = 1$. Let WRP and WRPB denote the sets of such weakly regular plateaued unbalanced and balanced functions, respectively.

Lemma 13. *[\[30\]](#page-23-12)* Let $f \in WRP$ or $f \in WRPB$. Then $f^*(0) = 0$ and $f^*(a\beta) = a^l f^*(\beta)$ for all $a \in \mathbb{F}_p^*$ and $\beta \in S_f$, where l is an even positive integer with $gcd(l-1, p-1) = 1$. For any $\beta \in S_f$ (s) (respectively, $\beta \in \mathbb{F}_{p^m} \backslash S_f$), we have $z\beta \in S_f$ (respectively, $z\beta \in \mathbb{F}_{p^m} \backslash S_f$) for every $z \in \mathbb{F}_p^*$.

To give the sufficient and necessary condition for a quadratic function to be *s*-plateaued, we now give a brief introduction on the quadratic functions (see [\[17](#page-23-13)]). Recall that any quadratic function from \mathbb{F}_{p^m} to \mathbb{F}_p with no linear term can be represented by

$$
Q(x) = \sum_{i=0}^{\lceil m/2 \rceil} \text{Tr}^m(a_i x^{p^i+1}),
$$

where $\lceil \cdot \rceil$ denotes ceiling function and $a_i \in \mathbb{F}_{p^m}$ for $0 \le i \le \lceil m/2 \rceil$. Then there exists an $m \times m$ symmetric matrix *A* such that $Q(x) = x^T A x$. Let *L* be the corresponding linearized polynomial over F*p ^m* defined as

$$
L(z) = \sum_{i=0}^{\lceil m/2 \rceil} (a_i z^{p^i} + a_i^{p^{m-i}} z^{p^{m-i}}).
$$

Then rank(*A*) = $m - s_L$ by [\[14,](#page-22-12) Proposition 2.1], where $0 \le s_L \le m$ is the dimension of the following \mathbb{F}_p -linear subspace of \mathbb{F}_{p^m} :

$$
\ker_{\mathbb{F}_p}(L) = \{ z \in \mathbb{F}_{p^m} : Q(z+y) = Q(z) + Q(y), \forall y \in \mathbb{F}_{p^m} \}.
$$

Proposition 14. *[\[28](#page-23-14)] Any quadratic function Q is s-plateaued if and only if* $rank(A) = m - s$, *i.e.* $s_L = s$.

By [\[17](#page-23-13), Proposition 1] and [\[5,](#page-22-13) Theorem 4.3], we have the following fact.

Proposition 15. *Any quadratic function is weakly regular plateaued. Namely, there is no quadratic non-weakly regular plateaued functions.*

2.3. Cyclotomic field

The following lemma gives some results on the cyclotomic field $\mathbb{Q}(\zeta_p)$.

Lemma 16. [\[27](#page-23-11)] Let $K = \mathbb{Q}(\zeta_p)$ denote the p-th cyclotomic field over the rational number field \mathbb{Q} , *wher p is an odd prime. Then the following hold.*

- *The ring of integers in K is* $O_K = \mathbb{Z}(\zeta_p)$ *and* $\{\zeta_p^i : 1 \le i \le p-1\}$ *is an integral basis of* O_K *, where* ζ*^p is the primitive p-th root of complex unity.*
- *The field extension K*/Q *is Galois of degree p*−1 *and Galois group Gal*(*K*/Q) = {σ*^a* : *a* ∈ \mathbb{F}_p^* , *where the automorphism* σ_a *of K is defined by* $\sigma_a(\zeta_p) = \zeta_p^a$.
- *The field K has a unique quadratic subfield* $L = \mathbb{Q}(\sqrt{p^*})$ *. For* $1 \le a \le p 1$ *,* $\sigma_a(\sqrt{p^*}) =$ $\eta_0(a)\sqrt{p^*}$, where $p^* = (-1)^{\frac{p-1}{2}}p$ and η_0 is the quadratic multiplicative character of \mathbb{F}_p . *Hence, the Galois group* $Gal(L/\mathbb{Q}) = \{1, \sigma_\gamma\}$ *, where* γ *is a nonsquare in* \mathbb{F}_p^* *.*

According to Lemma [16,](#page-6-0) we have

$$
\sigma_a(\zeta_p^b) = \zeta_p^{ab} \text{ and } \sigma_a(\sqrt{p^*}^m) = \eta_0^m(a)\sqrt{p^*}^m. \tag{5}
$$

3. Four-weight linear codes from *p*-ary weakly regular *s*-plateaued functions

In this section, if $f(x)$ is a weakly regular *s*-plateaued function, we will prove that the code $\overline{C_f}$ defined in Equation [\(1\)](#page-3-0) is a family of self-orthogonal codes and its parameters and weight distribution will be determined. Besides, we will construct a family of LCD codes from $\overline{C_f}$ and prove that $\overline{C_f}$ is an optimally extendable code.

3.1. The weight distribution of $\overline{C_f}$

From now on, we let λ_1 denote the canonical additive character of \mathbb{F}_p and ϕ_1 denote the canonical additive character of \mathbb{F}_{p^m} . Let $\text{wt}(\mathbf{c})$ denote the Hamming weight of a codeword \mathbf{c} in a linear code.

Lemma 17. Let $t \in \mathbb{F}_p^*$, $a \in \mathbb{F}_p$ and $b \in \mathbb{F}_{p^m}$, where p is an odd prime. Let $f(x)$ be a weakly regular *s-plateaued function with* $\varepsilon = \pm 1$ *the sign of its Walsh transform. Let* N_t *denote the number of solutions in* \mathbb{F}_{p^m} *of the equation af*(*x*) + $\text{Tr}_{p^m/p}(bx) = t$ *. If* $m + s$ *is even, then we have*

$$
N_t = \begin{cases} 0 & \text{if } a = 0, b = 0, \\ p^{m-1} & \text{if } a = 0, b \in \mathbb{F}_{p^m}^*, \text{ or } a \in \mathbb{F}_p^*, a^{-1}b \notin S_f, \\ p^{m-1} + \frac{\varepsilon(p-1)\sqrt{p^*}^{m+s}}{p} & \text{if } a \in \mathbb{F}_p^*, f^*(-_a^b) = a^{-1}t \text{ and } a^{-1}b \in S_f, \\ p^{m-1} - \frac{\varepsilon\sqrt{p^*}^{m+s}}{p} & \text{if } a \in \mathbb{F}_p^*, f^*(-_a^b) \neq a^{-1}t \text{ and } a^{-1}b \in S_f, \end{cases}
$$

where $p^* = (-1)^{\frac{p-1}{2}} p$. *If* $m + s$ *is odd, then we have*

$$
N_{t} = \begin{cases} 0 & \text{if } a = 0, b = 0, \\ p^{m-1} & \text{if } a = 0, b \in \mathbb{F}_{p^{m}}^{*}, or a \in \mathbb{F}_{p}^{*}, f^{*}(-\frac{b}{a}) = a^{-1}t, a^{-1}b \in S_{f}, \\ p^{m-1} + \frac{\varepsilon \sqrt{p^{*m+s}} G(\eta_{0}, \lambda_{1})}{p} & \text{if } a \in \mathbb{F}_{p}^{*}, f^{*}(-\frac{b}{a}) \neq a^{-1}t, \eta_{0}(f^{*}(-\frac{b}{a}) - a^{-1}t) = 1 \\ & \text{and } a^{-1}b \in S_{f}, \\ p^{m-1} - \frac{\varepsilon \sqrt{p^{*m+s}} G(\eta_{0}, \lambda_{1})}{p} & \text{if } a \in \mathbb{F}_{p}^{*}, f^{*}(-\frac{b}{a}) \neq a^{-1}t, \eta_{0}(f^{*}(-\frac{b}{a}) - a^{-1}t) = -1 \\ & \text{and } a^{-1}b \in S_{f}, \\ & & \text{and } a^{-1}b \in S_{f}, \end{cases}
$$

 $where G(\eta_0, \lambda_1) = (\sqrt{-1})^{(\frac{p-1}{2})^2} \sqrt{p}.$

Proof. Based on the orthogonal relation of additive characters, we have

$$
N_t = |\{x \in \mathbb{F}_{p^m} : af(x) + \text{Tr}_{p^m/p} (bx) = t\}|
$$

= $\frac{1}{p} \sum_{y \in \mathbb{F}_p} \sum_{x \in \mathbb{F}_{p^m}} \zeta_p^{yaf(x) + \text{Tr}_{p^m/p}(ybx) - yt}$
= $\frac{1}{p} \sum_{y \in \mathbb{F}_p^*} \lambda_1(-yt) \sum_{x \in \mathbb{F}_{p^m}} \zeta_p^{yaf(x) + \text{Tr}_{p^m/p}(ybx)} + p^{m-1}$

.

In order to calculate the value of N_t , we now consider the following cases. Case 1: If $a = 0$, then

$$
N_t = \frac{1}{p} \sum_{y \in \mathbb{F}_p^*} \lambda_1(-yt) \sum_{x \in \mathbb{F}_{p^m}} \phi_1(ybx) + p^{m-1} = \begin{cases} 0 & \text{if } b = 0, \\ p^{m-1} & \text{if } b \in \mathbb{F}_{p^m}^*.\end{cases}
$$

Case 2: If $a \neq 0$, then

$$
N_t = \frac{1}{p} \sum_{y \in \mathbb{F}_p^*} \lambda_1(-yt) \sum_{x \in \mathbb{F}_{p^m}} \zeta_p^{ya(f(x) - \text{Tr}_{p^m/p}(-\frac{b}{a}x))} + p^{m-1}
$$

$$
= \frac{1}{p} \sum_{y \in \mathbb{F}_p^*} \lambda_1(-yt) \sigma_{ya} \left(\mathbf{W}_f\left(-\frac{b}{a}\right)\right) + p^{m-1}
$$

$$
= \frac{1}{p} \sum_{y \in \mathbb{F}_p^*} \lambda_1(-ya^{-1}t) \sigma_y \left(\mathbf{W}_f\left(-\frac{b}{a}\right)\right) + p^{m-1}.
$$

where σ_y is the automorphism of $\mathbb{Q}(\zeta_p)$ defined by $\sigma_y(\zeta_p) = \zeta_p^y$. Subcase 2.1: If $W_f(-\frac{b}{a})$ $\frac{b}{a}$) = 0 (namely $a^{-1}b \notin S_f$), then

$$
N_t=p^{m-1}.
$$

Subcase 2.2: If $W_f(-\frac{b}{a})$ $\frac{b}{a}$) \neq 0 (namely $a^{-1}b \in S_f$), then we have

$$
\sigma_y\left(W_f\left(-\frac{b}{a}\right)\right) = \sigma_y\left(\epsilon\sqrt{p^*}^{m+s}\zeta_p^{f^*(-\frac{b}{a})}\right) = \epsilon\sqrt{p^*}^{m+s}\zeta_p^{yf^*(-\frac{b}{a})}\eta_0^{m+s}(y),
$$

by Lemma [12](#page-5-0) and Equation [5,](#page-6-1) where $p^* = (-1)^{\frac{p-1}{2}}p$. Then

$$
N_t = \frac{\varepsilon \sqrt{p^*}^{m+s}}{p} \sum_{y \in \mathbb{F}_p^*} \lambda_1(-ya^{-1}t) \zeta_p^{yf^*(-_a^b)} \eta_0^{m+s}(y) + p^{m-1}
$$

=
$$
\frac{\varepsilon \sqrt{p^*}^{m+s}}{p} \sum_{y \in \mathbb{F}_p^*} \lambda_1\left(y\left(f^*\left(-\frac{b}{a}\right) - a^{-1}t\right)\right) \eta_0^{m+s}(y) + p^{m-1}.
$$

Subcase 2.2.1: If $m + s$ is an even integer, by the orthogonality of additive characters, we have

$$
N_t = \frac{\varepsilon \sqrt{p^*}^{m+s}}{p} \sum_{y \in \mathbb{F}_p^*} \lambda_1 \left(y \left(f^* \left(-\frac{b}{a} \right) - a^{-1} t \right) \right) + p^{m-1}
$$

=
$$
\begin{cases} \frac{\varepsilon (p-1) \sqrt{p^*}^{m+s}}{p} + p^{m-1} & \text{if } f^* (-\frac{b}{a}) = a^{-1} t, \\ -\frac{\varepsilon \sqrt{p^*}^{m+s}}{p} + p^{m-1} & \text{if } f^* (-\frac{b}{a}) \neq a^{-1} t. \end{cases}
$$

Subcase 2.2.2: If $m + s$ is an odd integer, by the orthogonality of additive characters, we have

$$
N_t = \frac{\varepsilon \sqrt{p^{*}}^{m+s}}{p} \sum_{y \in \mathbb{F}_p^*} \lambda_1 \left(y \left(f^* \left(-\frac{b}{a} \right) - a^{-1} t \right) \right) \eta_0(y) + p^{m-1}
$$

$$
= \begin{cases} p^{m-1} & \text{if } f^* (-\frac{b}{a}) = a^{-1} t, \\ \frac{\varepsilon \sqrt{p^{*}}^{m+s}}{p} G(\eta_0, \lambda_1) + p^{m-1} & \text{if } \eta_0 (f^* (-\frac{b}{a}) - a^{-1} t) = 1, \\ -\frac{\varepsilon \sqrt{p^{*}}^{m+s}}{p} G(\eta_0, \lambda_1) + p^{m-1} & \text{if } \eta_0 (f^* (-\frac{b}{a}) - a^{-1} t) = -1. \end{cases}
$$

where $G(\eta_0, \lambda_1) = (\sqrt{-1})^{(\frac{p-1}{2})^2} \sqrt{p}$ by Lemma [7.](#page-4-0)

The desired conclusions follow from summarizing the above discussions.

Theorem 18. Let $q = p^m$, where p is an odd prime, s is an integer and m is a positive integer *with* $m \geq 2$ *. Let* $f(x)$ *be a weakly regular s-plateaued function with* $\varepsilon = \pm 1$ *the sign of its Walsh transform. Then* C_f *defined in Equation* [\(1\)](#page-3-0) *is a* [$q, m+2$] *linear code over* \mathbb{F}_p *and its weight distributions are respectively listed in Table [1](#page-9-0) for even m*+*s with* 0 ≤ *s* ≤ *m and Table [2](#page-9-1) for odd m* + *s* with 0 ≤ *s* ≤ *m* − 1*. Particularly,* $\overline{C_f}$ *is self-orthogonal if* $m + s$ ≥ 3*. Besides,* $\overline{C_f}^{\perp}$ *has parameters* $[q, q - m - 2, 3]$ *if* $m + s \geq 3$ *and is at least almost optimal according to the spherepacking bound.*

 \Box

Proof. Let $\mathbf{c}_{(a,b,c)} = (af(x) + \text{Tr}_{q/p}(bx) + c)_{x \in \mathbb{F}_q}$, $a \in \mathbb{F}_p$, $b \in \mathbb{F}_q$, $c \in \mathbb{F}_p$, be a codeword in $\overline{C_f}$. For $c = 0$, we have

$$
wt(c_{(a,b,0)}) = p^m - |\{x \in \mathbb{F}_q : af(x) + Tr_{q/p}(bx) = 0\}|
$$

$$
= p^m - \frac{1}{p} \sum_{y \in \mathbb{F}_p} \sum_{x \in \mathbb{F}_q} \phi_1(ybx) \lambda_1(yaf(x))
$$

Table 1: The weight distribution of $\overline{C_f}$ in Theorem [18](#page-8-0) for even $m + s$.

Weight	Frequency
$p^{m} - p^{m-1} - \frac{\varepsilon(p-1)(\sqrt{p^*})^{m+s}}{m}$	$(p-1)p^{m-s}$
$p^{m} - p^{m-1} + \frac{\varepsilon(\sqrt{p^*})^{m+s}}{n}$	$(p-1)^2 p^{m-s}$
$p^{m} - p^{m-1}$	$p(p^m-1)+p(p-1)(p^m-p^{m-s})$
n^m	$p-1$

Table 2: The weight distribution of $\overline{C_f}$ in Theorem [18](#page-8-0) for odd $m+s$.

$$
= pm - pm-1 - \frac{1}{p} \sum_{y \in \mathbb{F}_p^*} \sum_{x \in \mathbb{F}_q} \lambda_1(yaf(x) + yTr_{q/p}(bx))
$$

\n
$$
= \begin{cases} 0, & \text{if } a = 0, b = 0, \\ pm - pm-1, & \text{if } a = 0, b \neq 0, \\ pm - pm-1 - \frac{1}{p} \sum_{y \in \mathbb{F}_p^*} \sum_{x \in \mathbb{F}_q} \zeta_p^{ya(f(x) - Tr_{q/p}(-\frac{b}{a}x))}, & \text{if } a \neq 0. \end{cases}
$$

Denote by $N_1 = p^m - p^{m-1} - \frac{1}{p} \sum_{n \in \mathbb{N}} p^n$ *y*∈F[∗] *p* ∑ *x*∈F*^q* $\zeta_p^{ya(f(x)-\text{Tr}_{q/p}(-\frac{b}{a}x))}$. Similarly to the proof of Lemma [17,](#page-6-2) we have the following results.

(1) If $W_f(-\frac{b}{a})$ $\frac{b}{a}$) = 0 (i.e. $a^{-1}b \notin S_f$), then

$$
N_1=p^m-p^{m-1}.
$$

(2) If $W_f(-\frac{b}{a})$ $\frac{b}{a}$) \neq 0 (i.e. $a^{-1}b \in S_f$), we have

$$
N_1 = \begin{cases} p^m - p^{m-1} - \frac{\varepsilon(p-1)\sqrt{p^{*m+s}}}{p} & \text{if } f^*(-\frac{b}{a}) = 0, \\ p^m - p^{m-1} + \frac{\varepsilon\sqrt{p^{*m+s}}}{p} & \text{if } f^*(-\frac{b}{a}) \neq 0, \end{cases}
$$

for even $m + s$, and

$$
N_1 = \begin{cases} p^m - p^{m-1} & \text{if } f^*(-\frac{b}{a}) = 0, \\ p^m - p^{m-1} - \frac{\varepsilon \sqrt{p^{*m+s}} G(\eta_0, \lambda)}{p} & \text{if } f^*(-\frac{b}{a}) \neq 0 \text{ and } \eta_0(f^*(-\frac{b}{a})) = 1, \\ p^m - p^{m-1} + \frac{\varepsilon \sqrt{p^{*m+s}} G(\eta_0, \lambda)}{p} & \text{if } f^*(-\frac{b}{a}) \neq 0 \text{ and } \eta_0(f^*(-\frac{b}{a})) = -1, \end{cases}
$$

for odd $m + s$.

Then the value of $wt(c_{(a,b,0)})$ directly follows.

For $c \in \mathbb{F}_p^*$, denote by $N_{(a,b,c)} = |\{x \in \mathbb{F}_q : af(x) + \text{Tr}_{q/p}(bx) + c = 0\}|$. Then the Hamming weight of $\mathbf{c}_{(a,b,c)}$ is $\text{wt}(\mathbf{c}_{(a,b,c)}) = p^m - N_{(a,b,c)}$ for $a \in \mathbb{F}_p, b \in \mathbb{F}_{p^m}, c \in \mathbb{F}_p^*$. Besides, we can verify that $N_{(a,b,c)}$ is equal to N_t in Lemma [17](#page-6-2) for $t = -c$. Then by Lemma [17,](#page-6-2) if $m + s$ is an even integer, we have

$$
wt(c_{(a,b,c)}) = p^m - N_{(a,b,c)}
$$

\n
$$
= \begin{cases} p^m & \text{if } a = 0, b = 0, \\ p^m - p^{m-1} & \text{if } a = 0, b \in \mathbb{F}_{p^m}^*, \text{ or } a \in \mathbb{F}_{p^m}^*, a^{-1}b \notin S_f, \\ p^m - p^{m-1} - \frac{\varepsilon(p-1)\sqrt{p^m + s}}{p^m} & \text{if } a \in \mathbb{F}_{p^m}^*, f^*(-\frac{b}{a}) + a^{-1}c = 0, a^{-1}b \in S_f, \\ p^m - p^{m-1} + \frac{\varepsilon\sqrt{p^m + s}}{p} & \text{if } a \in \mathbb{F}_{p^m}^*, f^*(-\frac{b}{a}) + a^{-1}c \neq 0, a^{-1}b \in S_f; \end{cases}
$$

if $m + s$ is an odd integer, we have

$$
wt(c_{(a,b,c)}) = pm - N_{(a,b,c)}
$$

\n
$$
\begin{cases}\n pm & \text{if } a = 0, b = 0, \\
 pm - pm-1 & \text{if } a = 0, b \in \mathbb{F}_{pm}^*, or a \in \mathbb{F}_{pn}^*, f*(-\frac{b}{a}) + a-1c = 0, a-1b \in S_f, \\
 or a \in \mathbb{F}_{pn}^*, a-1b \notin S_f, \\
 pm - pm-1 - \frac{\varepsilon \sqrt{p*^{m+s}} G(\eta_0, \lambda)}{p} \\
 \text{if } a \in \mathbb{F}_{pn}^*, f*(-\frac{b}{a}) + a-1c \neq 0, \eta_0(f*(-\frac{b}{a}) + a-1c) = 1 \\
 \text{and } a-1b \in S_f, \\
 pm - pm-1 + \frac{\varepsilon \sqrt{p*^{m+s}} G(\eta_0, \lambda)}{p} \\
 \text{if } a \in \mathbb{F}_{pn}^*, f*(-\frac{b}{a}) + a-1c \neq 0, \eta_0(f*(-\frac{b}{a}) + a-1c) = -1 \\
 \text{and } a-1b \in S_f.\n\end{cases}
$$

Now we summarize the above discussions. If $m + s$ is an even integer, then

$$
\text{wt}(\mathbf{c}_{(a,b,c)}) = \begin{cases}\n0 & \text{if } a = 0, b = 0, c = 0, \\
p^m & \text{if } a = 0, b = 0, c \in \mathbb{F}_p^*, \\
p^m - p^{m-1} & \text{if } a = 0, b \in \mathbb{F}_{p^m}^*, c \in \mathbb{F}_p, \\
p^m - p^{m-1} - \frac{\varepsilon(p-1)\sqrt{p^{*m+s}}}{p^m} & \text{if } a \in \mathbb{F}_p^*, f^*(-\frac{b}{a}) + a^{-1}c = 0 \text{ and } a^{-1}b \in S_f, \\
p^m - p^{m-1} + \frac{\varepsilon\sqrt{p^{*m+s}}}{p} & \text{if } a \in \mathbb{F}_p^*, f^*(-\frac{b}{a}) + a^{-1}c \neq 0 \text{ and } a^{-1}b \in S_f;\n\end{cases}
$$

if $m + s$ is an odd integer, then

$$
\text{wt}(\mathbf{c}_{(a,b,c)}) = \begin{cases}\n0 & \text{if } a = 0, b = 0, c = 0, \\
p^m & \text{if } a = 0, b = 0, c \in \mathbb{F}_p^*, \\
p^m - p^{m-1} & \text{if } a = 0, b \in \mathbb{F}_{p^n}^*, c \in \mathbb{F}_p, \text{ or } a \in \mathbb{F}_p^*, a^{-1}b \in S_f, \\
f^*(-\frac{b}{a}) + a^{-1}c = 0, \text{ or } a \in \mathbb{F}_p^*, a^{-1}b \notin S_f, c \in \mathbb{F}_p, \\
p^m - p^{m-1} - \frac{\varepsilon \sqrt{p^{*m+s}} G(\eta_0, \lambda)}{p} & \text{if } a \in \mathbb{F}_p^*, f^*(-\frac{b}{a}) + a^{-1}c \neq 0, \\
\eta_0(f^*(-\frac{b}{a}) + a^{-1}c) = 1 \text{ and } a^{-1}b \in S_f, \\
p^m - p^{m-1} + \frac{\varepsilon \sqrt{p^{*m+s}} G(\eta_0, \lambda)}{p} & \text{if } a \in \mathbb{F}_p^*, f^*(-\frac{b}{a}) + a^{-1}c \neq 0, \\
\eta_0(f^*(-\frac{b}{a}) + a^{-1}c) = -1 \text{ and } a^{-1}b \in S_f.\n\end{cases}
$$

When $m + s$ is even, we denote by $w_1 = p^m$, $w_2 = p^m - p^{m-1}$, $w_3 = p^m - p^{m-1} - \frac{\varepsilon(p-1)\sqrt{p^{m}}}{p}$ $\frac{\sqrt{p}}{p}$, $w_4 = p^m - p^{m-1} + \frac{\varepsilon \sqrt{p^*}^{m+s}}{p}$ *p*. Next we will determine the frequency A_{w_i} , $1 \le i \le 4$. It is easy to deduce that $A_{w_1} = p - 1$ and $A_{w_2} = p(p^m - 1) + p(p - 1)(p^m - p^{m-s})$ by Lemma [8.](#page-4-1) Note that $A_{w_3} = |\{(a,b,c) \in \mathbb{F}_p^* \times \mathbb{F}_{p^m} \times \mathbb{F}_p : f^*(-\frac{b}{a})\}|$ $\left(\frac{b}{a}\right) + a^{-1}c = 0$ and $a^{-1}b \in S_f$ }. For fixed *a* and *b*, then the equation f^* $\left(-\frac{b}{a}\right)$ $\frac{b}{a}$) + $a^{-1}c$ = 0 with variable *c* has the unique solution. Hence, $A_{w_3} = (p-1)p^{m+s}$ by Lemma [8](#page-4-1) and $A_{w_4} = p(p-1)p^{m+s} - A_{w_3} = (p-1)^2p^{m+s}.$

When $m + s$ is odd, we denote by $w_1 = p^m$, $w_2 = p^m - p^{m-1}$, $w_3 = p^m - p^{m-1} - \frac{\varepsilon \sqrt{p^{m+1}} G(\eta_0, \lambda)}{p}$ $\frac{G(1|0,\mathcal{N})}{p}$, $w_4 = p^m - p^{m-1} + \frac{\varepsilon \sqrt{p^{*m+1}} G(\eta_0, \lambda)}{p}$ $\frac{G(\eta_0,\lambda)}{p}$. Next we will determine the frequency A_{w_i} , $1 \le i \le 4$. It is easy to deduce that $A_{w_1} = p - 1$ and $A_{w_2} = p(p^m - 1) + p(p - 1)(p^m - p^{m-s}) + p^{m-s}(p - 1)$ by Lemma [8.](#page-4-1) Note that $A_{w_3} = |\{(a, b, c) \in \mathbb{F}_p^* \times \mathbb{F}_{p^m} \times \mathbb{F}_p : f^* \left(-\frac{b}{a}\right)$ $\frac{b}{a}$) + $a^{-1}c \neq 0, \eta_0$ (f^* ($-\frac{b}{a}$ $\frac{b}{a}$ + $a^{-1}c$ = 1 and $a^{-1}b \in S_f$ }. For fixed $a \in \mathbb{F}_p^*$, $b \in \mathbb{F}_{p^m}$ and $\delta \in \langle \gamma^2 \rangle$, the solution of the equation $f^*(-\frac{b}{a})$ $\frac{b}{a}$) + $a^{-1}c = \delta$ with variable *c* is unique. Then $A_{w_3} = \frac{(p-1)^2 p^{m+s}}{2}$ $\frac{p}{2}$ by lemma [8.](#page-4-1) Similarly, we deduce $A_{w_4} = \frac{(p-1)^2 p^{m+s}}{2}$ $\frac{p}{2}$.

According to the weight distribution of $\overline{C_f}$, we deduce that $\overline{C_f}$ is *p*-divisible for $m + s \ge 3$. Then by Lemma [3,](#page-1-0) $\overline{C_f}$ is a self-orthogonal code.

Finally, we determine the parameters of $\overline{C_f}^{\perp}$. If $m + s$ is even, by the first four Pless power moments in [\[18,](#page-23-3) Page 260], we have

$$
A_1^{\perp} = A_2^{\perp} = 0, A_3^{\perp} = \frac{p^{m-1}(p-1)(p-2)\left(p^m - p + \epsilon \eta_0^{\frac{m+s}{2}}(-1)p^{\frac{m+s}{2}}(p-1)\right)}{6}.
$$

Note that $A_3^{\perp} > 0$ as $m + s \ge 4$ is an even integer. Then the parameters of $\overline{C_f}^{\perp}$ are $[q, q - m - 2, 3]$. If $m + s$ is odd, by the first four Pless power moments in [\[18,](#page-23-3) Page 260], we have

$$
A_1^{\perp} = A_2^{\perp} = 0, A_3^{\perp} = \frac{p^{m-1}(p-1)(p-2)(p^m - p)}{6}.
$$

Note that $A_3^{\perp} > 0$ as $m + s \geq 3$ is an odd integer. Then the parameters of $\overline{C_f}^{\perp}$ are $[q, q - m - 2, 3]$. The proof is completed by the sphere-packing bound in Lemma [1.](#page-1-1)

3.2. The optimally extendability of $\overline{C_f}$

In this subsection, we will prove that $\overline{C_f}$ is optimally extendable by selecting a suitable generator matrix of $\overline{C_f}$. Let $\mathbb{F}_q^* = \langle \alpha \rangle$ for $q = p^m$. Then $\{1, \alpha, \alpha^2, ..., \alpha^{m-1}\}$ is a \mathbb{F}_p -basis of \mathbb{F}_q . Let $d_1, d_2, \dots, d_{q-1}, d_q$ denote all elements of \mathbb{F}_q , where $d_q = 0$. A generator matrix *G* of $\overline{C_f}$ is given by

$$
G := \left[\begin{array}{cccc} 1 & 1 & \cdots & 1 \\ f(d_1) & f(d_2) & \cdots & f(d_q) \\ \mathrm{Tr}_{q/p}(\alpha^0 d_1) & \mathrm{Tr}_{q/p}(\alpha^0 d_2) & \cdots & \mathrm{Tr}_{q/p}(\alpha^0 d_q) \\ \mathrm{Tr}_{q/p}(\alpha^1 d_1) & \mathrm{Tr}_{q/p}(\alpha^1 d_2) & \cdots & \mathrm{Tr}_{q/p}(\alpha^1 d_q) \\ \vdots & \vdots & \ddots & \vdots \\ \mathrm{Tr}_{q/p}(\alpha^{m-1} d_1) & \mathrm{Tr}_{q/p}(\alpha^{m-1} d_2) & \cdots & \mathrm{Tr}_{q/p}(\alpha^{m-1} d_q) \end{array} \right]
$$

.

.

By an elementary row transformation, we obtain another generator matrix G_1 of $\overline{C_f}$ given as

$$
G_1 := \left[\begin{array}{cccc} 1 & 1 & \cdots & 1 \\ f(d_1) + 1 & f(d_2) + 1 & \cdots & f(d_q) + 1 \\ \mathrm{Tr}_{q/p}(\alpha^0 d_1) & \mathrm{Tr}_{q/p}(\alpha^0 d_2) & \cdots & \mathrm{Tr}_{q/p}(\alpha^0 d_q) \\ \mathrm{Tr}_{q/p}(\alpha^1 d_1) & \mathrm{Tr}_{q/p}(\alpha^1 d_2) & \cdots & \mathrm{Tr}_{q/p}(\alpha^1 d_q) \\ \vdots & \vdots & \ddots & \vdots \\ \mathrm{Tr}_{q/p}(\alpha^{m-1} d_1) & \mathrm{Tr}_{q/p}(\alpha^{m-1} d_2) & \cdots & \mathrm{Tr}_{q/p}(\alpha^{m-1} d_q) \end{array} \right]
$$

Let $G' := [I_{m+2,m+2} : G_1]$ generate a linear code $\overline{C_f}'$, where $I_{m+2,m+2}$ denotes the identity matrix of size $m+2$.

Theorem 19. Let $q = p^m$, where p is an odd prime, m is a positive integer, $0 \le s \le m$ is an integer *such that m*+*s is even and m*+*s* ≥ 4*. Let f*(*x*) *be a weakly regular s-plateaued function with* ε *the sign of the Walsh transform of* $f(x)$

1) When f(*x*) *is unbalanced, then we have the following results.*

1.1) If
$$
\varepsilon(\eta_0(-1))^{\frac{m+s}{2}} = 1
$$
, then $\overline{C_f}'$ has parameters
\n
$$
\left[p^m + m + 2, m + 2, p^m - p^{m-1} - (p-1)p^{\frac{m+s}{2}-1} + 2 \right].
$$

1.2) If $\epsilon(\eta_0(-1))^{\frac{m+s}{2}} = -1$ *, then* $\overline{C_f}$ ^{*'*} *has parameters*

$$
\[p^m+m+2,m+2,p^m-p^{m-1}-p^{\frac{m+s}{2}-1}+1\].
$$

2) When f(*x*) *is balanced, then we have the following results.*

2.1) If
$$
\varepsilon(\eta_0(-1))^{\frac{m+s}{2}} = 1
$$
, then $\overline{C_f}'$ has parameters
\n
$$
\left[p^m + m + 2, m + 2, \geq p^m - p^{m-1} - (p-1)p^{\frac{m+s}{2}-1} + 2 \right].
$$

2.2) If $\varepsilon(\eta_0(-1))^{\frac{m+s}{2}} = -1$, then $\overline{C_f}$ has parameters

$$
\[p^m + m + 2, m + 2, \geq p^m - p^{m-1} - p^{\frac{m+s}{2}-1} + 2\].
$$

Moreover, $\overline{C_f}^{\prime \perp}$ has parameters $[p^m + m + 2, p^m, 3]$ and is at least almost optimal with respect *to the sphere-packing bound.*

Proof. It is obvious that the length of $\overline{C_f}$ is $q + m + 2$ and the dimension of $\overline{C_f}$ is $m + 2$. It is clear that the minimum distance $d(\overline{C_f}')$ of $\overline{C_f}'$ satisfies $d(\overline{C_f}') \geq d(\overline{C_f})$, where $d(\overline{C_f})$ denotes the minimum distance of $\overline{C_f}$.

Next we prove that the minimum distance $d(\overline{C_f}')$ of $\overline{C_f}'$ satisfies that $d(\overline{C_f}') \geq d(\overline{C_f}) + 1$. For a codeword \mathbf{a}' in $\overline{C_f}'$, we have

$$
\mathbf{a}' = (c, a, a_0, a_1 \cdots, a_{m-1}) G'_1 \n= (c, a, a_0, \cdots, a_{m-1}, a f(d_1) + \sum_{i=0}^{m-1} a_i \text{Tr}_{q/p}(\alpha^i d_1) + c + a, \cdots, af(d_q) + \sum_{i=0}^{m-1} a_i \text{Tr}_{q/p}(\alpha^i d_q) + c + a),
$$

where $c, a, a_i \in \mathbb{F}_p$ for all $0 \le i \le m-1$. Then $\mathbf{a}' = (c, a, a_0, a_1 \cdots, a_{m-1}, \mathbf{a})$, where

$$
\mathbf{a} = (c, a, a_0, a_1 \cdots, a_{m-1}) G_1 = (af(x) + \text{Tr}_{p^m/p}(bx) + c + a)_{x \in \mathbb{F}_q}
$$

is a codeword in $\overline{C_f}$, where $b := \sum_{i=0}^{m-1} a_i \alpha^i$. Assume that **a**' has the minimum nonzero weight d'. If $d' = d(\overline{C_f})$, then we deduce that the first $m + 2$ locations are all zero. We have $c = a = a_i = 0$ (0 \leq *i* ≤ *m* − 1) and **a'** = **0**, which leads to a contradiction. Thus $d(\overline{C_f}') \ge d(\overline{C_f}) + 1$.

Next we consider the following cases:

Case 1: Let $f(x)$ be unbalanced. Then $W_f(0) = \varepsilon \sqrt{p^*}^{m+s}$ and $0 \in S_f$.

Subcase 1.1: Let $\varepsilon(\eta_0(-1))^{\frac{m+s}{2}} = 1$. By Theorem [18,](#page-8-0) we deduce that wt(**a**) = $d(\overline{C_f}) = p^m$ $p^{m-1} - (p-1)p^{\frac{m+s}{2}-1}$ if and only if $a \in \mathbb{F}_p^*$, $f^*(-\frac{b}{a})$ $\frac{b}{a}$) + $a^{-1}(c + a) = 0$ and $a^{-1}b \in S_f$. If $f^*(-\frac{b}{a}$ $\frac{b}{a}$) + $a^{-1}(c+a) = 0$, we have $(b, c) \neq (0, 0)$. Besides, $b = 0$ if and only if $(a_0, a_1, \dots, a_{m-1}) = (0, 0, \dots, 0)$. Thus $(c, a_0, a_1, \dots, a_{m-1}) \neq (0, 0, 0, \dots, 0)$ and $d(\overline{C_f}') \geq d(\overline{C_f}) + 2$. Specially, if $a \in \mathbb{F}_p^*, b = 0$ and $c = -a$, then $f^*(-\frac{b}{a})$ $\frac{b}{a}$) + $a^{-1}(c + a) = 0$ and wt(**a**[']) = wt(**a**) + 2 = $d(\overline{C_f})$ + 2. Hence $d(\overline{C_f}')$ = $d(\overline{C_f}) + 2.$

Subcase 1.2: Let $\mathcal{E}(\eta_0(-1))^{\frac{m+s}{2}} = -1$. By Theorem [18,](#page-8-0) we deduce that $wt(\mathbf{a}) = d(\overline{C_f}) =$ $p^m - p^{m-1} - p^{\frac{m+s}{2} - 1}$ if and only if $a \in \mathbb{F}_p^*$, $f^*(-\frac{b}{a})$ $\frac{b}{a}$) + $a^{-1}(c+a) \neq 0$ and $a^{-1}b \in S_f$. If $a \neq 0$, $b = 0$ and $c = 0$, then $f^*(-\frac{b}{a})$ $\frac{b}{a}$) + $a^{-1}(c + a) = 1$ and wt(**a**[']) = wt(**a**) + 1 = $d(\overline{C_f})$ + 1. Hence $d(\overline{C_f}') = d(\overline{C_f}) + 1.$

Case 2: Let $f(x)$ be balanced. Then $W_f(0) = 0$ and $0 \notin S_f$.

Subcase 2.1: Let $\varepsilon(\eta_0(-1))^{\frac{m+s}{2}} = 1$. By Theorem [18,](#page-8-0) we deduce wt(**a**) = $d(\overline{C_f}) = p^m - p^{m-1} (p-1)p^{\frac{m+s}{2}-1}$ if and only if $a \in \mathbb{F}_p^*$, $f^*(-\frac{b}{a})$ $(a^{\frac{b}{a}}) + a^{-1}(c + a) = 0$ and $a^{-1}b \in S_f$. Since $b \neq 0$, we have $wt(\mathbf{a}') \ge wt(\mathbf{a}) + 2$. Hence $d(\overline{C_f}') \ge d(\overline{C_f}) + 2$.

Subcase 2.2: Let $\mathcal{E}(\eta_0(-1))^{\frac{m+s}{2}} = -1$. By Theorem [18,](#page-8-0) we deduce wt(a) = $d(\overline{C_f}) = p^m$ $p^{m-1} - p^{\frac{m+s}{2}-1}$ if and only if $a \in \mathbb{F}_p^*$, $f^*(-\frac{b}{a})$ $(a^{\frac{b}{a}}) + a^{-1}(c + a) \neq 0$ and $a^{-1}b \in S_f$. Since $b \neq 0$, we also have $wt(\mathbf{a}') \ge wt(\mathbf{a}) + 2$. Hence $d(\overline{C_f}') \ge d(\overline{C_f}) + 2$.

It is obvious that that $\overline{C_f}'^{\perp}$ has length $q + m + 2$ and dimension q. We will prove that the minimum distance d^{\perp} of $\overline{C_f}'^{\perp}$ is 3 in the following. Note that G'_1 is a check matrix of $\overline{C_f}'^{\perp}$ and any column vector in G'_1 is nonzero. Hence, $d^{\perp} \geq 2$. Note that any two columns of the first $m+2$ columns of matrix G'_1 are \mathbb{F}_p -linearly independent and any two columns of the last *q* columns of matrix G'_1 are also \mathbb{F}_p -linearly independent. Select any column \mathbf{g}_1 in the first $m+2$ columns of matrix G'_1 and any column g_2 in the last *q* columns of matrix G'_1 . If $g_1 \neq (1,0,\dots,0)$, it is easy to deduce that g_1 and g_2 are \mathbb{F}_p -linearly independent. If $g_1 = (1,0,\dots,0)$, then g_1 and g_2 are \mathbb{F}_p -linearly dependent if and only if

$$
\begin{cases}\nf(d_i) + 1 = 0, \\
\operatorname{Tr}_{q/p}(\alpha^0 d_i) = 0, \\
\operatorname{Tr}_{q/p}(\alpha^1 d_i) = 0, \\
\vdots \\
\operatorname{Tr}_{q/p}(\alpha^{m-1} d_i) = 0,\n\end{cases}
$$
\n(6)

where $d_i \in \mathbb{F}_q$. This system of equations implies $d_i \in \mathbb{F}_q^*$. For any $x = \sum_{l=0}^{m-1} k_l \alpha^l \in \mathbb{F}_q$, $k_l \in \mathbb{F}_p$, the System [\(6\)](#page-14-0) implies

$$
\mathrm{Tr}_{q/p}(xd_i) = \sum_{l=0}^{m-1} k_l \mathrm{Tr}_{q/p}(\alpha^l d_i) = 0.
$$

This contradicts with the fact that $|\text{ker}(Tr_{q/p})| = p^{m-1}$. Hence, g_1 and g_2 are \mathbb{F}_p -linearly independent and $d^{\perp} \geq 3$. Besides, the first, second and last columns of G'_{1} are \mathbb{F}_p -linearly dependent. Then $d^{\perp} = 3.$ \Box

Next we will consider the case that $m + s$ is odd.

Theorem 20. Let $q = p^m$, where p is an odd prime, m is a positive integer, s is an integer and $m + s$ *is odd with m*+*s* ≥ 3*. Let f*(*x*) *be weakly regular s-plateaued function with* ε *the sign of the Walsh transform of* $f(x)$

1) When f(*x*) *is unbalanced, then we have the following results.*

1.1) If
$$
\varepsilon(\eta_0(-1))^{\frac{m+s+1}{2}} = 1
$$
, then the linear code $\overline{C_f}'$ has parameters
\n
$$
\left[p^m + m + 2, m + 2, p^m - p^{m-1} - p^{\frac{m+s-1}{2}} + 1 \right].
$$

1.2) If $\epsilon(\eta_0(-1))^{\frac{m+s+1}{2}} = -1$, then the linear code $\overline{C_f}$ has parameters

$$
\[p^m+m+2,m+2,p^m-p^{m-1}-p^{\frac{m+s-1}{2}}+2\].
$$

2) When f(*x*) *is balanced, then we have the following results.*

2.1) If
$$
\varepsilon(\eta_0(-1))^{\frac{m+s+1}{2}} = 1
$$
, then the linear code $\overline{C_f}'$ has parameters
\n
$$
\left[p^m + m + 2, m + 2, \ge p^m - p^{m-1} - p^{\frac{m+s-1}{2}} + 2 \right].
$$
\n2.2) If $\varepsilon(\eta_0(-1))^{\frac{m+s+1}{2}} = -1$, then the linear code $\overline{C_f}'$ has parameters
\n
$$
\left[p^m + m + 2, m + 2, \ge p^m - p^{m-1} - p^{\frac{m+s-1}{2}} + 2 \right].
$$

Moreover, the parameters of $\overline{C_f}'^{\perp}$ *is given by* $[p^m+m+2,p^m,3]$ *and* $\overline{C_f}'^{\perp}$ *is at least almost optimal with respect to the sphere-packing bound.*

 \Box

Proof. The proof is omitted as it is similar to that of Theorem [19.](#page-12-0)

The following gives the optimally extendability of $\overline{C_f}$.

Theorem 21. Let $\overline{C_f}$ be the linear code in Theorem [18](#page-8-0) and $\overline{C_f}'$ be the linear code with generator *matrix* $[I_{m+2,m+2} : G_1]$ *. Then* $\overline{C_f}$ *is an optimally extendable code with* $m+s \geq 3$ *.*

Proof. By Theorems [18,](#page-8-0)[19](#page-12-0) and [20,](#page-14-1) we have $d(\overline{C_f}^{\perp}) - d(\overline{C_f}^{\perp}) = 0$. Then the desired conclusion \Box follows.

Remark 22. Let $q = 3^m$, where $m \underline{\text{is}}_q$ a positive integer, s is an integer such that $m + s \geq 3$. *It is easy to verify that the codes* $\overline{C_f}'$ *in Theorems [19](#page-12-0) and [20](#page-14-1) are leading-systematic ternary LCD codes. We find that some ternary LCD codes are optimal according to the Code Tables at [http://www.codetables.de/.](http://www.codetables.de/) They are listed in Table [3.](#page-15-0)*

4. Binary self-orthogonal codes and LCD codes from *s*-plateaued Boolean functions

In this section, we study the code $\overline{C_f}$ defined in Equation [\(1\)](#page-3-0) if $f(x)$ is an *s*-plateaued Boolean function. We also construct binary LCD codes from $\overline{C_f}$.

Theorem 23. Let $q = 2^m$, where $0 \le s \le m-2$ is a nonnegative integer and m is a positive integer *with m* \geq 2 *such that m* $+$ *s is even. Let f*(*x*) *be an s-plateaued Boolean function. The* $\overline{C_f}$ *is a binary* $[q,m+2,2^{m-1}-2^{\frac{m+s-2}{2}}]$ *linear code and its weight distribution is listed in table [4.](#page-16-0) Particulary,* $\overline{C_f}$ is self-orthogonal if $m + s \geq 6$. Besides, $\overline{C_f}^{\perp}$ has parameters $[q, q-m-2, 4]$ and is optimal *according to the sphere-packing bound.*

Table 4: The weight distribution of $\overline{C_f}$ in Theorem [23](#page-15-1) when $p = 2$ and $m + s$ is even.

Weight	Frequency	
$\mathbf{\Omega}$		
$2^{m-1} - 2^{\frac{m+s-2}{2}}$	2^{m-s}	
$2^{m-1}+2^{\frac{m+s-2}{2}}$	2^{m-s}	
2^{m-1}	$2^{m+2} - 2^{m-s+1} - 2$	
2 ^m		

Proof. Let $\mathbf{c}_{(a,b,c)} = (af(x) + \text{Tr}_{q/p}(bx) + c)_{x \in \mathbb{F}_q}, a \in \mathbb{F}_2, b \in \mathbb{F}_q, c \in \mathbb{F}_2$, be a codeword in $\overline{C_f}$. Then its Hamming weight is given by

$$
wt(c_{a,b,c}) = 2m - \frac{1}{2} \sum_{y \in \mathbb{F}_2} \sum_{x \in \mathbb{F}_{2^m}} (-1)^{y(a f(x) + Tr_{2^m/2}(bx) + c}
$$

\n
$$
= 2m-1 - \frac{1}{2} \sum_{x \in \mathbb{F}_{2^m}} (-1)^{af(x) + Tr_{2^m/2}(bx) + c}
$$

\nif $a = 0, b = 0, c = 0$,
\nif $a = 0, b = 0, c = 1$,
\n
$$
2m-1 \qquad \text{if } a = 0, b \neq 0, c \in \mathbb{F}_2 \text{ or } a \neq 0, a^{-1}b \notin S_f, c \in \mathbb{F}_2,
$$

\n
$$
2m-1 - \frac{1}{2} W_f(a^{-1}b) \qquad \text{if } a \neq 0, b \in \mathbb{F}_{2^m}, c = 0, a^{-1}b \in S_f,
$$

\n
$$
2m-1 + \frac{1}{2} W_f(a^{-1}b) \qquad \text{if } a \neq 0, b \in \mathbb{F}_{2^m}, c = 1, a^{-1}b \in S_f.
$$

According to Lemma [10,](#page-5-1) we have $W_f(a^{-1}b) \in \{0, \pm 2^{(m+s)/2}\}\.$ Then the weight distribution directly follows where the frequency of each weight can be derived from Lemma [\[27\]](#page-23-11). By Lemma [2,](#page-1-2) we know $\overline{C_f}$ is self-orthogonal if $m + s \ge 6$. The parameters of $\overline{C_f}^{\perp}$ can be derived by the Pless power moments in [\[18,](#page-23-3) Page 260]. According to the sphere-packing bound, there exists no [q , q − m − 2,5] binary code. Then $\overline{C_f}^{\perp}$ is optimal with respect to the sphere-packing bound. \Box

In the following, we will construct a family of LCD codes from $\overline{C_f}$.

Let $I_{m+2,m+2}$ be the identity matrix of size $m+2$ and $\mathbb{F}_q^* = \langle \alpha \rangle$. Let $\{1, \alpha, \alpha^2, \dots, \alpha^{n-1}\}$ be a \mathbb{F}_2 -basis of \mathbb{F}_q . Let *d*₁, *d*₂, ···, *d*_{*q*−1}, *d*_{*q*} denote all elements of \mathbb{F}_q , where *d*_{*q*} = 0. By definition, the generator matrix *G* of $\overline{C_f}$ is given by

$$
G:=\left[\begin{array}{cccc} 1 & 1 & \cdots & 1 \\ f(d_1) & f(d_2) & \cdots & f(d_q) \\ \mathrm{Tr}_{q/2}(\alpha^0 d_1) & \mathrm{Tr}_{q/2}(\alpha^0 d_2) & \cdots & \mathrm{Tr}_{q/2}(\alpha^0 d_q) \\ \mathrm{Tr}_{q/2}(\alpha^1 d_1) & \mathrm{Tr}_{q/2}(\alpha^1 d_2) & \cdots & \mathrm{Tr}_{q/2}(\alpha^1 d_q) \\ \vdots & \vdots & \ddots & \vdots \\ \mathrm{Tr}_{q/2}(\alpha^{m-1} d_1) & \mathrm{Tr}_{q/2}(\alpha^{m-1} d_2) & \cdots & \mathrm{Tr}_{q/2}(\alpha^{m-1} d_q) \end{array}\right]
$$

However, the dual of the LCD code $\overline{C_f}$ generated by the matrix $G' := [I_{m+2,m+2} : G]$ has bad minimum distance 2. By an elementary row transformation on *G*, we obtain another generator

.

matrix G_2 of $\overline{C_f}$ given by

$$
G_2 := \begin{bmatrix} 1 & 1 & \cdots & 1 \\ f(d_1) + 1 & f(d_2) + 1 & \cdots & f(d_q) + 1 \\ \operatorname{Tr}_{q/2}(\alpha^0 d_1) & \operatorname{Tr}_{q/2}(\alpha^0 d_2) & \cdots & \operatorname{Tr}_{q/2}(\alpha^0 d_q) \\ \operatorname{Tr}_{q/2}(\alpha^1 d_1) & \operatorname{Tr}_{q/2}(\alpha^1 d_2) & \cdots & \operatorname{Tr}_{q/2}(\alpha^1 d_q) \\ \vdots & \vdots & \ddots & \vdots \\ \operatorname{Tr}_{q/2}(\alpha^{m-1} d_1) & \operatorname{Tr}_{q/2}(\alpha^{m-1} d_2) & \cdots & \operatorname{Tr}_{q/2}(\alpha^{m-1} d_q) \end{bmatrix} . \tag{7}
$$

Let $\overline{C_f}$ be the linear code generate by $G'_2 := [I_{m+2,m+2} : G_2]$.

Theorem 24. Let $q = 2^m$, where m is a positive integer and s is an integer such that $m + s$ is even with $m+s\geq 6$. The code $\overline{C_f'}$ generated by the matrix $G_2'=[I_{m+2,m+2}:G_2]$ is a leading-systematic *binary LCD code whose parameters are given as follows.*

(1) If $f(x)$ is balanced, then the binary LCD code $\overline{C_f}'$ has parameters $\left[2^m + m + 2, m + 2, \ge 2^{m-1} - 2^{\frac{m+s-2}{2}} + 2\right].$

(2) Let $f(x)$ be unbalanced.

(2.1) If
$$
W_f(0) = -2^{\frac{m+s}{2}}
$$
, then the binary LCD code $\overline{C_f}'$ has parameters
\n
$$
\left[2^m + m + 2, m + 2, 2^{m-1} - 2^{\frac{m+s-2}{2}} + 1\right].
$$
\n(2.2) If $W_f(0) = 2^{\frac{m+s}{2}}$, then $d = d + 1$; $\overline{D_f}(0) = 2^{\frac{m+s}{2}} + 1$

(2.2) If $W_f(0) = 2^{\frac{m+s}{2}}$, then the binary LCD code $\overline{C_f}'$ has parameters $\left[2^m + m + 2, m + 2, 2^{m-1} - 2^{\frac{m+s-2}{2}} + 2\right].$

Besides, $\overline{C_f}'^{\perp}$ *has parameters* $[2^m + m + 2, 2^m, 3]$ *and is at least almost optimal according to the sphere-packing bound.*

Proof. By Lemma [6,](#page-2-0) the linear code $\overline{C_f}$ is a leading-systematic LCD code. Obviously, the length of $\overline{C_f}$ is $q+m+2$ and the dimension of $\overline{C_f}$ is $m+2$. Similarly to the proof of Theorem [19,](#page-12-0) we can deduce that the minimum distance *d* of $\overline{C_f}$ satisfies that $d \geq d(\overline{C_f}) + 1$, where $d(\overline{C_f})$ denotes the minimum distance of $\overline{C_f}$. For a codeword \mathbf{a}' in $\overline{C_f}'$, we have

$$
\mathbf{a}' = (c, a, a_0, a_1 \cdots, a_{m-1})G'_2
$$

= $(c, a, a_0, \cdots, a_{m-1}, af(d_1) + \sum_{i=0}^{m-1} a_i \text{Tr}_{q/2}(\alpha^i d_1) + c + a, \cdots, af(d_q) + \sum_{i=0}^{m-1} a_i \text{Tr}_{q/2}(\alpha^i d_q) + c + a),$

where $c, a, a_i \in \mathbb{F}_2$ for all $0 \le i \le m-1$. Then $\mathbf{a}' = (c, a, a_0, a_1 \cdots, a_{m-1}, \mathbf{a})$, where

$$
\mathbf{a} = (c, a, a_0, a_1 \cdots, a_{m-1})G_2 = (af(x) + \text{Tr}_{q/2}(bx) + c + a)_{x \in \mathbb{F}_q}
$$

is a codeword in $\overline{C_f}$, where $b := \sum_{i=0}^{m-1} a_i \alpha^i$.

By Theorem [23,](#page-15-1) we deduce that $wt(a) = d(\overline{C_f}) = 2^{m-1} - 2^{\frac{m+s-2}{2}}$ if and only if $a \neq 0$, $c + a =$ $0, \mathbf{W}_f(a^{-1}b) = 2^{\frac{m+s}{2}}$ and $a^{-1}b \in S_f$, or $a \neq 0$, $c + a = 1$, $\mathbf{W}_f(a^{-1}b) = -2^{\frac{m+s}{2}}$ and $a^{-1}b \in S_f$.

- 1. If $f(x)$ is balanced, then $W_f(0) = 0$. Hence $W_f(a^{-1}b) \neq 0$ implies $b \neq 0$. Then we deduce $d \geq 2^{m-1} - 2^{\frac{m+s-2}{2}} + 2.$
- 2. Let $f(x)$ be unbalanced, then $W_f(0) \neq 0$.
	- (a) Let $W_f(0) = -2^{\frac{m+s}{2}}$. Then $wt(a) = d(\overline{C_f})$ if $a = 1, b = c = 0$. We then have $wt(a') =$ $d(\overline{C_f}) + 1$ and $d = 2^{m-1} - 2^{\frac{m+s-2}{2}} + 1$.
	- (b) Let $W_f(0) = 2^{\frac{m+s}{2}}$. Then we claim that $d \geq d(\overline{C_f}) + 2$. Otherwise, if there exists a codeword $\mathbf{a}' = (c, a, a_0, a_1 \cdots, a_{m-1}, \mathbf{a})$ of weight $d(\overline{C_f}) + 1$. Then $\text{wt}(\mathbf{a}) = d(\overline{C_f})$ and $wt(c, a, a_0, a_1 \cdots, a_{m-1}) = 1$. Note that $wt(a) = d(\overline{C_f})$ implies $a \neq 0$ by Theorem [23.](#page-15-1) Then $\text{wt}(c, a, a_0, a_1 \cdots, a_{m-1}) = 1$ implies $c = b = 0$. Since $W_f(0) = 2^{\frac{m+s}{2}}$, $\text{wt}(a) =$ $d(C_f)$ implies $a = c = 1$ which contradicts with $c = 0$. Then $d \geq d(C_f) + 2$. Let $a =$ $c = 1, b = 0$. Then there exists a codeword **a**' such that $wt(a') = d(\overline{C_f}) + 2$. Thus $d = 2^{m-1} - 2^{\frac{m+s-2}{2}} + 2.$

The parameters of $\overline{C_f}'^{\perp}$ can be derived by a similar method used in the proof of Theorem [19.](#page-12-0) \Box

Theorem 25. Let $m + s$ be even and $m + s \geq 6$. Let $\overline{C_f}$ be the linear code in Theorem [23](#page-15-1) with generator matrix G_2 and $\overline{C_f}$ be the linear code with generator matrix $[I_{m+2,m+2}:G_2]$. Then $\overline{C_f}$ is *an almost optimally extendable code.*

Proof. By Theorems [23](#page-15-1) and [24,](#page-17-0) we have $d(\overline{C_f}^{\perp}) - d(\overline{C_f}'^{\perp}) = 1$. The desired conclusion follows from the definition of almost optimally extendable code. \Box

5. Self-dual codes

In this section, we derive some families of self-dual codes from some self-orthogonal codes. The following is a simple result for self-orthogonal codes.

Lemma 26. Let *C* be a self-orthogonal code. Then for any $C' \subseteq C$, C' is also a self-orthogonal *code.*

 \Box

Proof. The desired conclusion directly follows from the definition of self-orthogonal code.

Lemma 27. Let G be a generator matrix of a linear code C' and $\begin{bmatrix} 1 \\ C \end{bmatrix}$ *G* 1 *be a generator matrix of a linear code C. If C is self-orthogonal, then C* ′ *is also self-orthogonal.*

Proof. Note that $C' \subseteq C$. Then the desired conclusion follows from Lemma [26.](#page-18-0) \Box

Theorem 28. Let $q = p^m$, where p is an odd prime, $m \ge 2$ is a positive integer and $0 \le s \le m$ is an *integer such that* $m+s \geq 3$ *. If* $p \equiv 1 \pmod{4}$ *and* $m \geq 2$ *<i>is any positive integer or* $p \equiv 3 \pmod{4}$ *and m is even, then there exists a* $\left[q-1, \frac{q-1}{2} \right]$ \int *self-dual code over* \mathbb{F}_p *which is a subcode of* $C_f^{*\perp}$ *.*

Proof. Define

$$
C_f = \left\{ (af(x) + \mathrm{Tr}_{q/p}(bx))_{x \in \mathbb{F}_q} : a \in \mathbb{F}_p, b \in \mathbb{F}_q \right\}.
$$

Then C_f is a subcode of $\overline{C_f}$ defined in Equation [\(1\)](#page-3-0). Since $\overline{C_f}$ is self-orthogonal by Theorem [18](#page-8-0) if $m + s \geq 3$, it follows from Lemma [27](#page-18-1) that C_f is also self-orthogonal. Note that C_f^* defined in Equation in [\(2\)](#page-3-1) can be derived by deleting the zero coordinate of each codeword of C_f . Hence C_f^* is self-orthogonal. Then the desired conclusion follows from Lemma [5.](#page-1-3)

Theorem 29. *Let* $q = 2^m$, where $0 ≤ s ≤ m − 2$ *is a nonnegative integer and m is a positive integer with* $m \geq 2$ such that $m+s \geq 6$ is even. Let $\overline{C_f}$ be defined in Equation [\(1\)](#page-3-0). Then there exists a $[q, \frac{q}{2}]$ $\frac{q}{2}$] binary self-dual code which is a subcode of $\overline{C_f}^\perp$.

Proof. By Theorem [23,](#page-15-1) $\overline{C_f}$ is self-orthogonal if $m + s \ge 6$ is even. Then the desired conclusion follows from Lemma [4.](#page-1-4)

In the following, we give an example to illustrate how to find self-dual codes by Theorems [28](#page-18-2) and [29.](#page-19-0)

Example 30. Let $\mathbb{F}_q^* = \langle \alpha \rangle$ and $q = p^m$. Let $p = 3$, $m = 2$ and $f(x) = \text{Tr}_{3^m/3}(\alpha x^4 + \alpha^8 x^2)$. Then $f(x)$ *is weakly regular ternary 1-plateaued function and* $\overline{C_f}$ *is a 3-weight ternary* $[9,4,3]$ *code with weight enumerator* $1 + 6z^3 + 66z^6 + 8z^9$. We know that $\overline{C_f}$ is self-orthogonal by Lemma [2.](#page-1-2) It is *easy to see that C* ∗ *f is also self-orthogonal with parameters* [8,3,3] *by Lemma [27.](#page-18-1) We will find a* $self\text{-}dual code \text{ }\widetilde{C} \text{ from } C_f^{\ast\perp}$. Select a parity check matrix of C_f^{\ast} as follows:

$$
H := \left[\begin{array}{cccccc} 2 & 0 & 0 & 0 & 0 & 1 & 2 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & 0 & 2 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 2 & 0 \end{array}\right].
$$
 (8)

Then H is a generator matrix of ${C_f}^{*\perp}$. Taking the first four rows of H yields the following matrix:

$$
\widetilde{H} := \left[\begin{array}{cccccc} 2 & 0 & 0 & 0 & 0 & 1 & 2 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & 0 & 2 \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 & 2 & 0 \end{array} \right].
$$
\n(9)

Then \widetilde{H} generate a [8,4,3] *ternary linear code* \widetilde{C} *which is a subcode of* $C_f^{*{\perp}}$ *.* \widetilde{C} *has weight enumerator* $1 + 16z^3 + 64z^6$ *. Hence* \tilde{C} *is self-orthogonal by Lemma* 2*.* Since $dim(\tilde{C}) = 4$, \tilde{C} *is self-dual.*

6. Summary and concluding remarks

In this paper, we mainly studied the linear code $\overline{C_f}$ if f is a weakly regular plateaued function or a plateaued Boolean function. The highlights of this family of codes are as follows:

• $\overline{C_f}$ is self-orthogonal under certain conditions (see Theorems [18](#page-8-0) and [23\)](#page-15-1);

- The dual of $\overline{C_f}$ is optimal (see Theorem [23\)](#page-15-1) or at least almost optimal (see Theorem [18\)](#page-8-0) according to the sphere-packing bound;
- $\overline{C_f}$ is an optimally extendable code (see Theorem [21\)](#page-15-2) or an almost optimally extendable code (see Theorem [25\)](#page-18-3) by selecting a suitable generator matrix.

Besides, we also derived some families of binary or ternary LCD codes from $\overline{C_f}$. In Table [5,](#page-21-0) we list known infinite families of binary or ternary LCD codes. From this table, our LCD codes have different parameters from known ones. In Table [3,](#page-15-0) these LCD codes are optimal in some cases. Furthermore, we also derived self-dual codes from some self-orthogonal codes in this paper. It is open to study the minimum distance of these self-dual codes.

$\, n$	[n,k,d]	Condition	Reference
$n = 3^m - 1$	$[n, 2m, \geq 2\delta_2]$	$\delta_2 = \frac{3^{m-1}-1}{4} + 3^{m-2}$, <i>m</i> odd	Th.4.5 in $[15]$
$n = 3^m - 1$	$[n, 3, \frac{3^m-1}{2}]$	m even	Th.4.6 in [15]
$n = 3^m - 1$	$[n, 2m, \geq 2\delta_3]$	$m \equiv 2 \pmod{4}$, $\delta_3 = \frac{3^{m-6}-1}{5}$ +3 ^{m-6} + 2 · 3 ^{m-5}	Th.4.7 in [15]
		$+2\cdot3^{m-3}+3^{m-2}$	
$n = 2^m - 1$	$[n, 2m, \geq 2\delta_1]$		Th.4.1 in $[16]$
$n = 2^m - 1$	$[n, 4m, \geq 2\delta_2]$	<i>m</i> > 3 odd, $\delta_1 = \frac{2^{m-1}-1}{3}$ <i>m</i> > 3 odd, $\delta_2 = \frac{2^{m-1}-7}{3}$ <i>m</i> > 3 odd, $\delta_3 = \frac{2^{m-1}-25}{3}$	Th.4.1 in $[16]$
$n = 2^m - 1$	$[n, 6m, \geq 2\delta_3]$		Th.4.1 in $[16]$
$n = 2^m - 1$	$[n, 2m + 2, > 2\delta_2]$		Th.4.4 in $[16]$
		$m \equiv 2 \pmod{4}, \delta_2 = \frac{2^{m-2}-1}{5}$	
$n = 2^m - 1$	$[n, 4m + 2, > 2\delta_3]$	$m \equiv 2 \pmod{4}$, $\delta_3 = \frac{2^{m-2}-31}{5}$ $+2^{m-3}$	Th.4.4 in $[16]$
$n = 2^m - 1$	$[n, 6, 2\delta_2]$	$m \equiv 0 \pmod{4}, \delta_2 = \frac{2^m - 1}{5}$	Th.4.5 in $[16]$
$n=q^{\ell}+1$	$[n,k,\geq 2(\delta-1)]$	$3 \le \delta \le q^{\lfloor(\ell-1)/2\rfloor} + 3$	Th.18 in [22]
		$k=q^{\ell}-2\ell(\delta-2-\lfloor\frac{\delta-2}{q}\rfloor)$	
$n = 2^m - 1$	$[n,k,\geq 2(2^{m-\ell}-1)]$	$m-2 \geq \ell \geq m - \lfloor (m-2)/2 \rfloor$ $k = 2^m - 2\sum_{j=0}^{m-1-\ell} {m \choose j}$	Th.26 in $[22]$
	$[n, n-1-2m\lceil \frac{(\delta-1)(q-1)}{q} \rceil, \geq 2\delta]$	$2 \leq \delta \leq q^{\lfloor (m-1)/2 \rfloor}$	Th.32 in [22]
<i>n</i> = $\frac{q^{m}-1}{q-1}$ <i>n</i> = $\frac{2^{m}+1}{3}$ <i>n</i> = $\frac{3m}{2}$ <i>n</i> = $\frac{2m}{2}$ <i>n</i> = $\binom{m}{t}$	$[n,k,\geq \delta]$	$m \ge 7$ odd and $3 \le \delta \le \frac{n}{2}$	Th.3.1 in [23]
	[n,m,2]	m even	Th. 10 in $[35]$
	$[n, \frac{m}{2}, 3]$	m even	Th.11 in [35]
	$[n, n-m+1, 3]$		Th.25 in [38]
		<i>t</i> even, $\binom{m-1}{t-1} \equiv 1 \pmod{2}$ and $\binom{m-2}{t-2} \equiv 0 \pmod{2}$; or	
		t even, $\overline{\binom{m-1}{t-1}} \equiv \binom{m-2}{t-2} + 1$	
		\equiv 0(mod 2) and tm \equiv 0(mod 2)	
$n = \binom{m}{t}$	$[n, n-m, 4]$	<i>t</i> odd, $\binom{m-1}{t-1} \equiv 1 \pmod{2}$ and $\binom{m-2}{t-2} \equiv 0 \pmod{2}$; or	Th.25 in [38]
		t odd, $\binom{m-1}{t-1} \equiv \binom{m-2}{t-2} + 1$	
		\equiv 0(mod 2) and $tm \equiv$ 0(mod 2)	
$n = \binom{m}{t} + 1$	$[n, n-m+1, 3]$	m, t even, $\binom{m-1}{t-1} \equiv 0 \pmod{2}$	Th.31 in [38]
		and $\binom{m-2}{t-2} \equiv 1 \pmod{2}$; or	
		<i>m</i> , <i>t</i> even $\binom{m-1}{t-1} \equiv \binom{m-2}{t-2}$	
		$+1 \equiv 1 \pmod{2}$	
$n = {m \choose t} + 1$ [n, n - m, 4]		t odd, $m \neq 2t$, $\binom{m-1}{t-1} \equiv 0 \pmod{2}$	Th.31 in [38]
		and $\binom{m-2}{t-2} \equiv 1 \pmod{2}$; or	
		t odd, $m \neq 2t$, $\binom{m-1}{t-1} \equiv \binom{m-2}{t-2}$	
		$+1 \equiv 1 \pmod{2}$ and <i>m</i> even	
	$n = n_f + m + 1$ $[n, m+1, \frac{n_f}{2} - 2^{\frac{m-4}{2}} + 2]$	$S = 0, m \ge 6$ even	Th.1 in $[36]$
		$n_f = 2^{m-1} \pm 2^{\frac{m-2}{2}}$	
	$n = n_f + m + 1$ $[n, m + 1, \frac{n_f}{2} - 2^{\frac{m-4}{2}} + 1]$	$S > 0, m \ge 6$ even	Th.1 in $[36]$
		$n_f = 2^{m-1} \pm 2^{\frac{m-2}{2}}$	
$n = n_f + m + 1$ [n, n, 3]		$m > 6$ even	Th.1 in $[36]$
		$n_f = 2^{m-1} \pm 2^{\frac{m-2}{2}}$	
	$n = q_0^{2r} + 3r + 1$ $[n, 3r + 1, (q_0 - 1)q_0^{2r-1} - q_0^{r-1} + 2]$	$r > 1, (q_0, r) \neq (2, 2)$	
$n = q_0^{2r} + 3r + 1$ [n, q_0^{2r} , 3]		$r > 1, (q_0, r) \neq (2, 2)$	Th.3 in [36]
	$n = p^m + m + 2$ $[n, m+2, p^m - p^{m-1} - (p-1)p^{\frac{m+s}{2}-1} + 2]$	$m+s \ge 4$ even, $f(x)$ unbalanced, $\epsilon(\eta_0(-1))^{\frac{m+s}{2}} = 1$	Th.3 in $[36]$
			Theorem 19
	$n = p^m + m + 2$ $[n, m+2, p^m - p^{m-1} - (p-1)p^{\frac{m+s}{2}-1} + 2]$	$m+s \geq 4$ even, $f(x)$ unbalanced, $\mathcal{E}(\eta_0(-1))^{\frac{m+s}{2}} = -1$	Theorem 19
	$n = p^m + m + 2$ $[n, m + 2] \geq p^m - p^{m-1} - (p-1)p^{\frac{m+s}{2}-1} + 2$	$m+s \geq 4$ even, $f(x)$ balanced, $\epsilon(\eta_0(-1))^{\frac{m+s}{2}} = 1$	Theorem 19
	$n = p^m + m + 2$ $[n, m + 2, \geq p^m - p^{m-1} - p^{\frac{m+s}{2}-1} + 2]$	$m+s \geq 4$ even, $f(x)$ balanced, $\mathcal{E}(\eta_0(-1))^{\frac{m+s}{2}} = -1$	Theorem 19
	$n = p^m + m + 2 \quad [n, m+2, p^m - p^{m-1} - p^{\frac{m+s-1}{2}} + 1]$	$m+s \geq 3$ odd, $f(x)$ unbalanced, $\epsilon(\eta_0(-1))^{\frac{m+s}{2}} = 1$	Theorem 20
	$n = p^m + m + 2$ $[n, m + 2, p^m - p^{m-1} - p^{\frac{m+s-1}{2}} + 2]$	$m+s \geq 3$ odd, $f(x)$ unbalanced, $\mathcal{E}(\eta_0(-1))^{\frac{m+s}{2}} = -1$	Theorem 20
	$n = p^m + m + 2$ $[n, m + 2, \geq p^m - p^{m-1} - p^{\frac{m+s-1}{2}} + 2]$	$m + s \geq 3$ odd, $f(x)$ balanced	Theorem 20
$n = p^m + m + 2 \quad [n, p^m, 3]$		$m + s > 3$	Theorem 19, Theorem 20
	$n = 2^m + m + 2$ $[n, m + 2, \geq 2^{m-1} - 2^{\frac{m+s-2}{2}} + 2]$	$m+s \geq 6$ even, $f(x)$ balanced	Theorem 24
$n = 2^m + m + 2$	$[n,m+2,2^{m-1}-2^{\frac{m+s-2}{2}}+1]$	$m+s \ge 6$ even, $f(x)$ unbalanced, $W_f(0) = -2^{\frac{m+s}{2}}$	Theorem 24
$n = 2^m + m + 2$	$[n,m+2,2^{m-1}-2^{\frac{m+s-2}{2}}+2]$	$m+s \ge 6$ even, $f(x)$ unbalanced, $W_f(0) = 2^{\frac{m+s}{2}}$	Theorem 24
$n = 2^m + m + 2$	$[n, 2^m, 3]$	$m+s \geq 6$ even	Theorem 24

Table 5: The known infinite families of binary or ternary LCD codes.

References

References

- [1] A. Alahmadi, A. Altassan, A. AlKenani, S. Çalkavur, H. Shoaib, P. Solé, A multisecretsharing scheme based on LCD codes, Mathematics 8 (2) (2020) 272.
- [2] J. Bringer, C. Carlet, H. Chabanne, S. Guilley, H. Maghrebi, Orthogonal direct sum masking - a smartcard friendly computation paradigm in a code with builtin protection against sidechannel and fault attacks, in: WISTP, Springer, Heraklion, 2014, pp. 40-56.
- [3] M. Braun, T. Etzion, A. Vardy, Linearity and complementsin projective space, Linear Algebra Appl. 430 (1) (2013) 57-70.
- [4] E. Berardini, A. Caminata, A. Ravagnani, Structure of CSS and CSS-T quantum codes. Des. Codes Cryptogr. (2024). online at https://doi.org/10.1007/s10623-024-01415-9.
- [5] A. Cesmelioğlu, W. Meidl, A construction of bent functions from plateaued functions, Des. Codes Cryptogr. 66 (2013) 231-242.
- [6] A.R. Calderbank, E.M. Rains, P.W. Shor, N.J.A. Sloane, Quantum error correction and orthogonal geometry, Phys. Rev. Lett. 78 (3) (1997) 405-408.
- [7] C. Carlet, C. Li, S. Mesnager, Some (almost) optimally extendable linear codes, Designs Codes Cryptogr. 87 (2019) 2813-2834.
- [8] C. Carlet, S. Mesnager, C. Tang, Y. Qi, R. Pellikaan, Linear codes over \mathbb{F}_q are equivalent to LCD codes for *q* > 3, IEEE Trans. Inf. Theory 64 (4) (2018) 3010-3017.
- [9] A. Canteaut, P. Charpin, Decomposing bent functions, IEEE Trans. Inform. Theory 49 (8) (2003) 2004-2019.
- [10] A. Canteaut, P. Charpin, G. Kyureghyan, A new class of monomial bent functions, Finite Fields Appl. 14 (1) (2008) 221-241.
- [11] K. Ding, C. Ding, A class of two-weight and three-weight codes and their applications in secret sharing, IEEE Trans. Inf. Theory 61 (2015) 5835-5842.
- [12] Z. Heng, X. Li, Y. Wu, Q. Wang, Two families of linear codes with desirable properties from some functions over finite fields, IEEE Trans. Inf. Theory, DOI: 10.1109/TIT.2024.3439408.
- [13] X. Hou, F. Oggier, On LCD codes and lattices, In Proc. IEEE Int. Symp. Inform. Theory (2016) 1501-1505.
- [14] X. Hou, Solution to a problem of S. Payne, Proc. Am. Math. Soc. 132 (2004) 1-6.
- [15] X. Huang, Q. Yue, Y. Wu, X. Shi, Ternary primitive LCD BCH codes, Advances in Mathematics of Communications 17 (3) (2023) 644-659.
- [16] X. Huang, Q. Yue, Y. Wu, X. Shi, J. Michel, Binary primitive LCD BCH codes, Des. Codes Cryptogr. 88 (2020) 2453-2473.
- [17] T. Helleseth, A. Kholosha, Monomial and quadratic bent functions over the finite fields of odd characteristic, IEEE Trans. Inform. Theory 52 (5) (2006) 2018-2032.
- [18] W.C. Huffman, V. Pless, Fundamentals of Error-Correcting Codes, Cambridge University Press, Cambridge, 2003.
- [19] X. Li, Z. Heng, Self-orthogonal codes from *p*-divisible codes, IEEE Trans. Inf. Theory, Doi: 10.1109/TIT.2024.3449921.
- [20] X. Li, Z. Heng, Constructions of near MDS codes which are optimal locally recoverable codes, Finite Fields Appl. 88 (2023) 102184.
- [21] R. Lidl, H. Niederreiter, Finite Fields, Cambridge University Press, Cambridge, 1997.
- [22] C. Li, C, Ding, S. Li, LCD cyclic codes over finite fields, IEEE Trans. Inf. Theory 63 (7) (2017) 4344-4356.
- [23] Y. Liu, X. Kai, S. Zhu, On binary LCD BCH codes of length $\frac{2^m+1}{3}$ $\frac{1}{3}$, Adv. Math. Commun. (2022), online at https://www.aimsciences.org/article/doi/10.3934/amc.2022053.
- [24] J. L. Massey, Reversible codes, Inf. Control 7 (3) (1964) 369-380.
- [25] J. L. Massey, Linear codes with complementary duals, Discrete Math. 106-107 (1992) 337- 342.
- [26] J. L. Massey, Orthogonal, antiorthogonal and self-orthogonal matrices and their codes, In Communications and Coding, Somerset, England: Research Studies Press, (1998) 3-9.
- [27] S. Mesnager, F. Ozbudak, A. Sinak, Linear codes from weakly regular plateaued functions and their secret sharing schemes, Des. Codes Cryptogr. 87 (2-3) (2019) 463-480.
- [28] S. Mesnager, F. Özbudak, A. Sinak, Results on characterizations of plateaued functions in arbitrary characteristic, in: International Conference on Cryptography and Information Security in the Balkans, Springer, 2015, pp. 17-30.
- [29] X. Quan, Q. Yue, L. Hu, Some constructions of (almost) optimally extendable linear codes. Adv. Math. Commun. (2022). online at https://doi.org/10.3934/amc.2022027.
- [30] A. Sınak, Construction of minimal linear codes with few weights from weakly regular plateaued functions, Turkish Journal of Math. 46 (3) (2022) 953-972.
- [31] N. Sendrier, Linear codes with complementary duals meet the Gilbert-Varshamov bound, Discrete Math. 285 (1-3) (2004) 345-347.
- [32] J. Sui, Q. Yue, X. Li and D. Huang, MDS, near-MDS or 2-MDS self-dual codes via twisted generalized Reed-Solomon codes, IEEE Trans. Inf. Theory 68(12) (2022) 7832-7841.
- [33] Y. Wu, C. Li, F. Xiao, Quaternary linear codes and related binary subfield codes, IEEE Trans. Inf. Theory 68 (5) (2022) 3070-3080.
- [34] Y. Wu, Q. Yue, Factorizations of binomial polynomials and enumerations of LCD and selfdual constacyclic codes, IEEE Trans. Inf. Theory 65 (3) (2019) 1740-1751.
- [35] Y. Wu, Y. Lee, Binary LCD codes and self-orthogonal codes via simplicial complexes, IEEE Commun. Lett. 24(6), 1159-1162, (2020).
- [36] X. Wang, Z. Heng, Li, F. et al. LCD codes and almost optimally extendable codes from selforthogonal codes. Des. CodesCryptogr. (2024). https://doi.org/10.1007/s10623-024-01420-y.
- [37] Z. Wan, A characteristic property of self-orthogonal codes and its application to lattices, Bulletin of the Belgian Mathematical Society-Simon Stevin 5 (2/3) (1998) 477-482.
- [38] Z. Zhou, X. Li, C. Tang, C. Ding, Binary LCD codes and self-orthogonal codes from a generic construction, IEEE Trans. Inform. Theory 65 (1) (2019) 16-27.