# MORSE INEQUALITIES FOR NONCOMPACT MANIFOLDS

TSUYOSHI KATO, DAISUKE KISHIMOTO, AND MITSUNOBU TSUTAYA

Abstract. We establish Morse inequalities for a noncompact manifold with a cocompact and properly discontinuous action of a discrete group, where Morse functions are not necessarily invariant under the group action. The inequalities are given in terms of the  $L^2$ -Betti numbers and functions on the acting group which describe rough configurations of critical points of a Morse function.

#### 1. INTRODUCTION

Morse inequalities relate the number of critical points of a Morse function on a compact manifold to its Betti numbers. Here, compactness of a manifold is essential as it guarantees finiteness of the number of critical points and the Betti numbers. Then it is challenging to generalize Morse inequalities to noncompact manifolds. There should have been attempts for it, but there is no result except for the very special cases that critical points are finitely many  $[4]$  and have strong symmetry  $[16]$ .

Throughout the paper, let  $M$  be a connected *n*-dimensional manifold without boundary, possibly noncompact, equipped with a cocompact and properly discontinuous action of a discrete group G such that the orbit manifold  $M/G$  is oriented. Namely, we will consider a Galois covering  $G \to M \to M/G$  such that M is connected and  $M/G$  is a closed oriented n-dimensional manifold. We fix any  $G$ -invariant metric on  $M$ . If  $G$ is amenable, then a manifold  $M$  is a typical example of a manifold having bounded geometry and a regular exhaustion, for which the index theory is developed in [18, 19]. The purpose of this paper is to establish Morse inequalities for a certain Morse function on  $M$  which is essentially irrelevant to the action of G.

Roe [18] crudely classified generalizations of the index theorem to noncompact manifolds into three types. In particular, for the type II theorems, including Atiyah's Γ-index theorem [1] and Roe's index theorem [18, 19], the index is interpreted through some kind of averaging or renormalization procedure. Our approach takes such procedure. To generalize Morse inequalities to noncompact manifolds, we need to replace Betti numbers by other invariants because the Betti numbers of a noncompcat manifold may not be defined. For a manifold  $M$ , there is a nice variant of Betti numbers,  $L^2-Betti \ numbers,$  and we employ them for our purpose. As in [13, Example 1.37],  $L^2$ -Betti numbers may be thought of as a kind of average of Betti

Date: November 8, 2024.

<sup>2010</sup> Mathematics Subject Classification. 57R19, 58E05.

Key words and phrases. Morse inequality, noncompact manifold,  $L^2$ -Betti number, piecewise trace, Witten deformation.

numbers, so they should match our approach. On the other hand, Gromov [6] proved that vanishing of  $L^2$ -Betti numbers is a quasi-isometry invariant, and as in  $[17]$ , for a large class of groups,  $L^2$ -Betti numbers themselves are proved to be quasi-isometry invariants. Here, quasi-isometry invariants are the object of study in large scale geometry. However, we only use some results in large scale geometry, and do not work in it because quasi-isometries identify a compact set with a single point so that the number of critical points does not make sense.

Let us introduce Morse functions that we are going to consider. It is shown in [8] that every connected noncompact manifold admits a function without critical points. To exclude such a trivial situation, we assume the following boundedness of Morse functions (cf. [12]).

**Definition 1.1.** A smooth function  $f: M \to \mathbb{R}$  is bounded if  $\nabla^i f$  is bounded for any  $0 \leq i \leq \max\{3, \lceil \frac{n}{2} \rceil\}$  $\frac{n}{2}$ ].

Actually, boundedness of the gradient vector field  $\nabla f$  is essential, and that of f and its higher derivatives may be a technical condition. We note that the boundedness of a function is independent of the choice of a G-invariant metric on  $M$  because  $G$ -invariant metrics on  $M$  are mutually equivalent. By (1.2) below, we describe a rough configuration of critical points of a Morse function on  $M$  by a function on  $G$ , which may be thought of as counting of critical points. In order to make this function controllable, we assume the following uniformness of Morse functions. We fix a G-invariant triangulation of  $M$ . Then as in Section 2, we can construct a fundamental domain  $K$  as a finite subcomplex of M such that

$$
(1.1)\t\t\t\t M = \bigcup_{g \in G} gK
$$

and  $gK \cap hK$  has dimension at most  $n-1$  for  $g \neq h \in G$ .

**Definition 1.2.** A Morse function  $f: M \to \mathbb{R}$  is uniform if there is  $\epsilon > 0$ satisfying that

- (1) any two critical points of f are at least  $2\epsilon$  distant, and
- (2) for each critical point p of f, there is  $g \in G$  such that the  $\epsilon$ neighborhood of  $p$  is contained in  $gK$ .

In Section 5, we define a *strongly uniform* Morse function on M as a uniform Morse function on M satisfying additional mild conditions on critical points. Let  $f: M \to \mathbb{R}$  be a Morse function, and let  $\mathrm{Crit}_k(f)$  denote the set of critical points of  $f$  with index  $k$ . Define a function

(1.2) 
$$
c_k \colon G \to \mathbb{R}, \quad g \mapsto |\operatorname{Crit}_k(f) \cap gK|.
$$

Then this is a function describing a rough configuration of critical points of f with index  $k$ , mentioned above. This way of counting of discrete points on M was discovered in [12]. Observe that  $c_k$  is a bounded function for each  $k \geq 0$  whenever f is uniform, because a fundamental domain K is compact.

We set notation. Let  $b_k^{(2)}$  $k<sup>(2)</sup>$  denote the k-th  $L<sup>2</sup>$ -Betti number of M. Let  $\ell^{\infty}(G)$  denote the Banach space of bounded functions on G. Then G acts on  $\ell^{\infty}(G)$  by  $(g \cdot \phi)(x) = \phi(xg)$  for  $g, x \in G$  and  $\phi \in \ell^{\infty}(G)$ . Let I denote the subspace of  $\ell^{\infty}(G)$  generated by  $\phi - g \cdot \phi$  for  $g \in G$  and  $\phi \in \ell^{\infty}(G)$ , and

let  $\Im$  denote the closure of  $\Im$  by the weak topology. Taking modulo  $\Im$  may be thought of as averaging process mentioned above. For  $\phi_1, \phi_2 \in \ell^{\infty}(G)$ , we write

- $\phi_1 \ge \phi_2 \mod \overline{\mathcal{I}}$  if there is  $\phi_3 \in \overline{\mathcal{I}}$  such that the pointwise inequality  $\phi_1 \ge \phi_2 + \phi_3$  holds, and
- $\phi_1 \approx \phi_2 \mod \overline{J}$  if  $\phi_1 \geq \phi_2 \mod \overline{J}$  and  $\phi_2 \geq \phi_1 \mod \overline{J}$ .

The first relation is a preorder on  $\ell^{\infty}(G)$ , not necessarily a partial order, so that we need the second relation. The quotient  $\ell^{\infty}(G)/\mathcal{I}$  is isomorphic to the 0-th uniformly finite homology of  $M$  defined by Block and Weinberger  $[2]$ (see also [3]), and is related with finite propagation operators that play an important role in this paper [9, 10, 11]. Let  $\mathbb{1} \in \ell^{\infty}(G)$  denote the constant function on G with values 1.

Now we state the main theorem.

**Theorem 1.3.** For a strongly uniform bounded Morse function  $f: M \to \mathbb{R}$ , we have

$$
c_k - c_{k-1} + \dots + (-1)^k c_0 \ge (b_k^{(2)} - b_{k-1}^{(2)} + \dots + (-1)^k b_0^{(2)}) \mathbb{1} \mod \overline{\mathfrak{I}}
$$
  
for  $k = 0, 1, \dots, n-1$ , and  

$$
c_n - c_{n-1} + \dots + (-1)^n c_0 \approx (-1)^n \chi(M/G) \mathbb{1} \mod \overline{\mathfrak{I}}.
$$

We remark that Theorem  $1.3$  makes sense only when  $G$  is amenable because  $\ell^{\infty}(G) \neq \overline{\ell}$  if and only if G is amenable (Proposition 2.12). By definition, G is amenable if there is a G-invariant positive linear map  $\mu: \ell^{\infty}(G) \to$ R with  $\mu(1) = 1$ , which is called a G-invariant mean. As a corollary to Theorem 1.3, we have the following mean value Morse inequalities.

**Corollary 1.4.** Let  $f: M \to \mathbb{R}$  be a strongly uniform bounded Morse function. If G is amenable, then for any G-invariant mean  $\mu: \ell^{\infty}(G) \to \mathbb{R}$ ,

 $\mu(c_k) - \mu(c_{k-1}) + \cdots + (-1)^k \mu(c_0) \ge b_k^{(2)} - b_{k-1}^{(2)} + \cdots + (-1)^k b_0^{(2)}$  $\overline{0}$ 

for  $k = 0, 1, ..., n - 1$  and

$$
\mu(c_n) - \mu(c_{n-1}) + \cdots + (-1)^n \mu(c_0) = (-1)^n \chi(M/G).
$$

We can deduce from Theorem 1.3 the weak Morse inequalities

$$
c_k \ge b_k^{(2)} \mod \bar{\mathcal{I}}
$$

for  $k = 0, 1, \ldots, n$ , and by applying a property of the module of coinvariants  $\ell^{\infty}(G)/\ell$  proved in [12] to it, we get the following property of the  $L^2$ -Betti numbers, which has its own interest.

**Corollary 1.5.** Let  $f: M \to \mathbb{R}$  be a strongly uniform bounded Morse function. If G is an infinite amenable group and  $b_k^{(2)}$  $\binom{2}{k} \neq 0$ , then f must have infinitely many critical points with index k.

We also see that Theorem 1.3 recovers the result of Novikov and Shubin [16] on Morse inequalities for a Morse function on  $M/G$  and the  $L^2$ -Betti numbers of  $M$  when  $G$  is amenable (Corollary 5.9).

We prove Theorem 1.3 by employing the Witten deformation by a Morse function. Then we need to disassemble the trace of an operator on the Hilbert space of  $L^2$ -sections of a vector bundle over M with respect to the

cover (1.1). In Section 2, we introduce a piecewise trace-class operator and its piecewise trace which is the desired disassembly of the usual (or Roe's) trace. In Section 3, we introduce an operator whose kernel function has propagation subordinate to a Gaussian function centered at the diagonal set, which we call an operator with Gaussian propagation. We prove the trace property of the piecewise trace of a product of operators with Gaussian propagation. We also consider the functional calculus of a generalized Dirac operators such that the resulting operator has Gaussian propagation. In Section 4, we introduce a disassembly of the normalized Betti number of Roe [19, 20], which is related to the  $L^2$ -Betti number of M, with respect to the cover  $(1.1)$ , and prove its invariance under a certain deformation, including the Witten deformation by a bounded function. In Section 5, we collect all results obtained so far, and apply the Witten deformation to prove Theorem 1.3 and its corollaries. The boundedness of a Morse function itself is used only in the proof of the invariance of the normalized Betti number, though the boundedness of the derivative is essential as mentioned above. Then we pose some question on the boundedness of a Morse function itself.

Acknowledgement. The authors are partially supported by JSPS KAK-ENHI Grant numbers JP23K22394 (Kato), JP22K03284 (Kishimoto), and JP22K03317 (Tsutaya)

# 2. Piecewise trace

In [18], Roe defined the trace of a uniform operator over a noncompact manifold of bounded geometry with regular exhaustion, and use it to build the index theory for such a noncompact manifold. In this section, we introduce the piecewise trace of an operator on the space of  $L^2$ -sections of a G-invariant vector bundle over  $M$ , which may be thought of as disassembly of Roe's trace with respect to the action of G.

2.1. Fundamental domain. First of all, we recall from [12] the construction of a fundamental domain. Choose a G-invariant triangulation  $L$  of  $M$ , that is, the lift of a triangulation of  $M/G$ . Then  $L/G$  is a triangulation of  $M/G$ . For each *n*-simplex of  $L/G$ , we choose one of its lifts to L, and define the subcomplex  $K$  of  $L$ , called a *fundamental domain*, by the union of those *n*-simplices of  $L$ . Remark that a fundamental domain  $K$  needs not be connected. By definition, we have

$$
M = \bigcup_{g \in G} gK
$$

and  $\dim(qK \cap hK) \leq n-1$  for  $q \neq h \in G$ , implying  $\text{Int}(qK) \cap \text{Int}(hK) = \emptyset$ for  $q \neq h$ . In the sequel, we fix a fundamental domain K.

2.2. Piecewise Hilbert-Schmidt operator. Let  $E \to M$  be a G-invariant vector bundle with  $G$ -invariant metric, that is,  $E$  is the lift of a vector bundle over  $M/G$  with metric. Let  $L^2(E)$  denote the Hilbert space of  $L^2$ -sections of E. Then the map  $g^{-1}: gK \to K$  induces an isometry  $(g^{-1})^*: L^2(E|_K) \to$  $L^2(E|_{gK})$ . We choose an orthonormal basis  $\{e_i \mid i \in I\}$  of  $L^2(E|_K)$ , and set  $e_i^g = (g^{-1})^*(e_i)$  for  $g \in G$ . Then  $\{e_i^g\}$  $_i^g \mid i \in I$  is an orthonormal basis of  $L^2(E|_{gK})$ . Moreover, if we consider  $e_i^{\hat{g}}$  $_i^g$  as an  $L^2$ -section of the entire vector

bundle E in the obvious way, then  $\{e_i^g\}$  $_i^g \mid i \in I, g \in G$  is an orthonormal basis of  $L^2(E)$ . For  $u \in L^2(E)$ , let  $||u||$  denote the  $L^2$ -norm.

**Definition 2.1.** A bounded operator  $A: L^2(E) \to L^2(E)$  is called *piecewise* Hilbert-Schmidt if

$$
\rho_2(A)\colon G\to\mathbb{R},\quad g\mapsto \left(\sum_{i\in I}\|Ae_i^g\|^2\right)^{\frac{1}{2}}.
$$

is a well-defined bounded function.

Let  $\mathbf{1}_S : M \to \{0,1\}$  denote the characteristic function of a subset  $S \subset M$ . An operator  $A: L^2(E) \to L^2(E)$  is a piecewise Hilbert-Schmidt operator if and only if  $A1_{gK}$  is a Hilbert-Schmidt operator for each  $g \in G$  such that  $\sup_{g \in G} ||A1_{gK}||_{HS}$  is bounded, where  $|| \cdot ||_{HS}$  denotes the Hilbert-Schmidt norm. Observe that

(2.1) 
$$
\rho_2(A)(g) = \|A\mathbf{1}_{gK}\|_{\text{HS}}.
$$

Then  $\rho_2(A)$  is independent of the choice of an orthonormal basis of  $L^2(E|_K)$ , and we may think of  $\rho_2$  as disassembly of the Hilbert-Schmidt norm with respect to the cover  $(1.1)$ . The following lemma is immediate from the definition of  $\rho_2$ .

**Lemma 2.2.** Let  $A, B: L^2(E) \to L^2(E)$  be piecewise Hilbert-Schmidt operators.

- (1)  $\rho_2(A + B) \leq \rho_2(A) + \rho_2(B)$ .
- (2)  $\rho_2(cA) = |c|\rho_2(A)$  for  $c \in \mathbb{R}$ .
- (3)  $\rho_2(A) = 0$  if and only if  $A = 0$ .

For a bounded operator  $A: L^2(E) \to L^2(E)$ , let ||A|| denote its operator norm.

**Proposition 2.3.** Let  $A: L^2(E) \to L^2(E)$  be a bounded operator, and let  $B: L<sup>2</sup>(E) \to L<sup>2</sup>(E)$  be a piecewise Hilbert-Schmidt operator. Then AB is a piecewise Hilbert-Schmidt operator such that

$$
\rho_2(AB) \le \|A\|\rho_2(B).
$$

*Proof.* By  $(2.1)$ ,  $\rho_2(AB)(g) = ||AB\mathbf{1}_{gK}||_{\text{HS}} \le ||A|| ||B\mathbf{1}_{gK}||_{\text{HS}} = ||A|| \rho_2(B)(g)$ for any  $q \in G$ . Then the statement follows.

Let  $A: L^2(E) \to L^2(E)$  be a Hilbert-Schmidt operator, and let  $B: L^2(E) \to$  $L^2(E)$  be a bounded operator. Then we have

$$
||AB||_{\text{HS}} \le ||A||_{\text{HS}}||B||
$$
 and  $||A^*||_{\text{HS}} = ||A||_{\text{HS}}.$ 

Hence we may expect that  $\rho_2$  has analogous properties, but it fails. Choose any  $e \in L^2(E|_K)$  with  $||e|| = 1$  and  $g \in G$  with  $g \neq 1$ . We define a piecewise Hilbert-Schmidt operator

$$
A = \langle -, e \rangle (g^{-1})^*(e)
$$

where  $\langle -, - \rangle$  denotes the L<sup>2</sup>-inner product. Then  $A^* = \langle -, (g^{-1})^*(e) \rangle e$  is a piecewise Hilbert-Schmidt operator. Observe that  $\rho_2(A^*A)(1) = 1 \geq 0$  $\rho_2(A^*)(1)$ ||A|| and  $\rho_2(A)(1) = 1 \neq 0 = \rho_2(A^*)(1)$ . Then

$$
\rho_2(A^*A) \nleq \rho_2(A^*) \|A\|
$$
 and  $\rho_2(A^*) \neq \rho_2(A)$ .

Let  $A: L^2(E) \to L^2(E)$  be an operator. We say that A is represented by a kernel function  $k_A$  if  $k_A$  is a section of the vector bundle  $E^* \boxtimes E \to M \times M$ such that for any  $u \in L^2(E)$ ,

$$
(Au)(x) = \int_M k_A(x, y)u(y) \text{ vol}(y)
$$

where vol denotes the pullback of the volume form of  $M/G$  to M. We also say that an operator A is smoothing if it is represented by a smooth kernel function.

**Lemma 2.4.** If  $A: L^2(E) \to L^2(E)$  is a piecewise Hilbert-Schmidt smoothing operator, then for each  $q \in G$ ,

$$
\rho_2(A)(g) = \left(\int_M \int_{gK} |k_A(x, y)|^2 \operatorname{vol}(y) \operatorname{vol}(x)\right)^{\frac{1}{2}}.
$$

Proof. Observe that

$$
(\rho_2(A)(g))^2 = \sum_{i \in I} (||Ae_i^g||^2)^{\frac{1}{2}}
$$
  
= 
$$
\sum_{i \in I} \int_M \left| \int_M k_A(x, y) e_i^g(y) \operatorname{vol}(y) \right|^2 \operatorname{vol}(x)
$$
  
= 
$$
\int_M \sum_{\substack{i \in I \\ h \in G}} \left| \int_M k_A(x, y) \mathbf{1}_{gK} e_i^h(y) \operatorname{vol}(y) \right|^2 \operatorname{vol}(x)
$$
  
= 
$$
\int_M \int_M |k_A(x, y) \mathbf{1}_{gK}|^2 \operatorname{vol}(y) \operatorname{vol}(x)
$$
  
= 
$$
\int_M \int_M |k_A(x, y)|^2 \operatorname{vol}(y) \operatorname{vol}(x).
$$

where we use Parseval's identity for the fourth equality. Then the statement follows.  $\Box$ 

**Lemma 2.5.** If  $A, B: L^2(E) \to L^2(E)$  are piecewise Hilbert-Schmidt operators, then

$$
\rho_2(A, B): G \to \mathbb{R}, \quad g \mapsto \sum_{i \in I} \langle Ae_i^g, Be_i^g \rangle.
$$

is a well-defined bounded function which is independent of the choice of an orthonormal basis of  $L^2(E|_K)$ .

Proof. Define an operator

(2.2) 
$$
P = \sum_{i \in I, g \in G} \text{sgn}(\langle Ae_i^g, Be_i^g \rangle) \langle -, e_i^g \rangle e_i^g
$$

where we put  $sgn(0) = 0$ . Then BP is a piecewise Hilbert-Schmidt operator such that  $\rho_2(BP) = \rho_2(B)$  and  $\rho_2(A, BP) = \sum_{i \in I} |\langle Ae_i^g, Be_i^g \rangle|$ . Observe that for any piecewise Hilbert-Schmidt operators  $C$  and  $D$ , the Schwartz inequality

$$
(2.3) \qquad \qquad \rho_2(C,D) \le \rho_2(C)\rho_2(D)
$$

holds. Thus  $\rho_2(A, BP)$  is a bounded function, and therefore  $\rho_2(A, B)$  is a well-defined bounded function. The independence of the choice of an orthonormal basis of  $L^2(E|_K)$  follows from the fact that

$$
\rho_2(A, B)(g) = \langle A\mathbf{1}_{gK}, B\mathbf{1}_{gK} \rangle
$$

for  $g \in G$ .

2.3. Piecewise trace-class operator. We define a piecewise trace-class operator as an analogy to a trace-class operator.

**Definition 2.6.** An operator  $A: L^2(E) \rightarrow L^2(E)$  is a piecewise traceclass operator if  $A = B^*C$  for some piecewise Hilbert-Schmidt operators  $B, C: L<sup>2</sup>(E) \rightarrow L<sup>2</sup>(E)$ . Its *piecewise trace* is defined by

$$
\text{Tr}(A) = \rho_2(B, C).
$$

By Lemma 2.5, the piecewise trace is a well-defined bounded function on G which is independent of the choice of an orthonormal basis of  $L^2(E|_K)$ . It is easy to see that  $\text{Tr}(A)$  is independent of the choice of a decomposition  $A = B^*C$ . We show the basic properties of a piecewise trace.

**Lemma 2.7.** If  $A = B^*C$  for piecewise Hilbert-Schmidt smoothing operators  $B, C: L<sup>2</sup>(E) \rightarrow L<sup>2</sup>(E)$ , then A is a trace-class smoothing operator such that

$$
\operatorname{Tr}(A)(g) = \int_{gK} \operatorname{tr}(k_A(x, x)) \operatorname{vol}(x).
$$

Proof. By definition, A is of piecewise trace-class, and since the composite of smoothing operators is smoothing, A is a smoothing operator. Since  $\text{Tr}(A)(g)$  is the trace of the trace-class operator  $\mathbf{1}_q A \mathbf{1}_q$ , the equality in the statement follows quite similarly to [20, Theorems 8.12].  $\Box$ 

**Proposition 2.8.** Let  $A: L^2(E) \to L^2(E)$  be a piecewise trace-class smoothing operator. Then the piecewise trace of A modulo  $\mathfrak I$  is independent of the choice of a fundamental domain K.

*Proof.* By Lemma 2.7,  $Tr(A)$  is given by the integral of the bounded *n*-form  $tr(k_A(x, x))$  vol(x) in the sense of [12]. It is shown in [14] that such an integral modulo J is independent of the choice of  $K$ .

For a piecewise trace-class operator  $A: L^2(E) \to L^2(E)$ , we can define a bounded function

$$
\rho_1(A) \colon G \to \mathbb{R}, \quad g \mapsto \sum_{i \in I} |\langle e_i^g, Ae_i^g \rangle|.
$$

Indeed, the proof of Lemma 2.5 implies that  $\rho_1(A)$  is a well-defined bounded function. Here, we remark that  $\rho_1$  depends on the choice of an orthonormal basis of  $L^2(E|_K)$ , so when we consider  $\rho_1$ , we always fix an orthonormal basis of  $L^2(E|_K)$ . We record properties of  $\rho_1$  which are immediate from the definition.

**Lemma 2.9.** Let  $A, B: L^2(E) \to L^2(E)$  be piecewise trace-class operators. (1)  $\rho_1(A + B) \leq \rho_1(A) + \rho_1(B)$ .

(2)  $\rho_1(cA) = |c| \rho_1(A)$  for  $c \in \mathbb{R}$ .

**Proposition 2.10.** Let  $A: L^2(E) \to L^2(E)$  be a piecewise trace-class operator. If A is positive, then

$$
\text{Tr}(A) = \rho_1(A) \ge 0.
$$

*Proof.* If A is positive, then  $\langle e_i^g \rangle$  $\{g}_i^g, Ae_i^g \rangle \ge 0$  for any  $g \in G$  and  $i \in I$ , implying  $\text{Tr}(A) = \rho_1(A) \geq 0.$ 

**Proposition 2.11.** If  $A, B: L^2(E) \to L^2(E)$  are piecewise Hilbert-Schmidt operators, then

$$
\rho_1(A^*B) \le \rho_2(A)\rho_2(B).
$$

*Proof.* Let P be as in  $(2.2)$ . Then  $\rho_1(A^*B) = \rho_2(A, BP)$ , so by  $(2.3)$ ,  $\rho_1(A^*B) \leq \rho_2(A)\rho_2(BP) = \rho_2(A)\rho_2(B).$ 

2.4. Roe's trace. Recall that a discrete group  $G$  is amenable if there is a G-invariant positive linear map  $\mu: \ell^{\infty}(G) \to \mathbb{R}$  satisfying  $\mu(1) = 1$ , called a G-invariant mean. We have the following characterization of amenability.

Proposition 2.12. The following conditions are equivalent.

- (1) G is amenable.
- $(2) \ell^{\infty}(G) \neq \overline{\mathcal{I}}.$

*Proof.* Let  $\mu: \ell^{\infty}(G) \to \mathbb{R}$  be a *G*-invariant mean. Then for  $\phi \in \ell^{\infty}(G)$ ,

$$
|\mu(\phi)| \le |\mu(\sup_{x \in G} |\phi(x)| \mathbb{1})| = \sup_{x \in G} |\phi(x)|
$$

and then  $\mu$  is bounded, implying  $\mu(\bar{J}) = 0$ . Thus by  $\mu(1) = 1$ , (1) implies (2). By [2, 3], if G is not amenable then  $\ell^{\infty}(G) = \mathcal{I}$ . Hence (2) implies  $(1).$ 

We compare the piecewise trace with Roe's trace  $[18]$  when G is amenable. Recall that Følner's theorem also characterizes amenability, which states that for any  $\epsilon > 0$  and a nonempty finite subset  $S \subset G$ , there is a finite subset  $F \subset G$  satisfying

$$
\frac{|F \triangle gF|}{|F|} < \epsilon
$$

for any  $g \in S$ . For  $k \geq 1$ , we define  $S_k = \{ g \in G \mid d(K, gK) \leq k \}$ . Then we get a sequence  $S_1 \subset S_2 \subset \cdots$  of finite subsets of G which exhausts G. By Følner's theorem, there is a finite subset  $F_k$  of G satisfying

$$
\frac{|F_k \triangle g F_k|}{|F_k|} < \frac{1}{k}
$$

for any  $g \in S_k$ . Let  $M_k$  be the union of all  $gK$  for  $g \in F_k$ . Then as is shown in [18, p.106],  $\{M_k\}_{k>1}$  is a regular exhaustion of M, hence by [20, (3.5)], Roe's trace of a suitable smoothing operator  $A: L^2(E) \to L^2(E)$  is given by

$$
\tau(A) = \lim_{k \to \infty} \frac{1}{\text{vol}(M_k)} \int_{M_k} \text{tr}(k_A(x, x)) \text{vol}(x).
$$

On the other hand, we can define a G-invariant finitely additive probability measure on  $G$  by

$$
\mu(S) = \lim_{\omega} \frac{|F_k \cap S|}{|F_k|}
$$

for  $S \subset G$ , where  $\omega$  is a nonprincipal ultrafilter on N, which yields a Ginvariant mean  $\mu: \ell^{\infty}(G) \to \mathbb{R}$ . Thus we can easily deduce the following, where  $vol(K) = vol(M/G)$ .

**Proposition 2.13.** Let  $A: L^2(E) \to L^2(E)$  be a piecewise trace-class operator, and let  $\mu: \ell^{\infty}(G) \to \mathbb{R}$  be the above G-invariant mean. Then

$$
\mu(\text{Tr}(A)) = \frac{\tau(A)}{\text{vol}(M/G)}.
$$

#### 3. Operator with Gaussian propagation

In this section, we define an operator with Gaussian propagation, and show the trace property of the piecewise trace of the product of operators with Gaussian propagation. We also investigate functional calculus of a generalized Dirac operator that produces operators with Gaussian propagation.

3.1. Trace property. Let  $E \to M$  be a G-invariant vector bundle with G-invariant metric.

**Definition 3.1.** A smoothing operator  $A: L^2(E) \to L^2(E)$  has *Gaussian* propagation if there are  $C_1, C_2 > 0$  such that

$$
|k_A(x,y)| \le C_1 e^{-C_2 d(x,y)^2}.
$$

Since the group  $G$  is the quotient of the fundamental group of a compact manifold  $M/G$ , it is finitely generated. We choose a finite generating set of G, and consider the associated word metric on G, where those metrics are mutually quasi-isometric. Now we take any  $q, h \in G$ . By the fundamental observation in geometric group theory, an inclusion  $G \to M$  is a quasiisometry, so there are  $C_1, C_2, C_3, C_4 > 0$  which are independent of the choice of g, h such that for any  $x \in gK$ ,  $y \in hK$ , we have

(3.1) 
$$
C_1 e^{-C_2 d(g,h)^2} \le e^{-d(x,y)^2} \le C_3 e^{-C_4 d(g,h)^2}
$$

For  $m \geq 0$ , let  $G_m = \{g \in G \mid d(g, 1) \leq m\}$ . Then we get a sequence of subsets of G

$$
G_0 \subset \cdots \subset G_m \subset G_{m+1} \subset \cdots
$$

which exhausts G. As in [15], there is  $C > 0$  such that

$$
(3.2) \t\t |G_m| \le e^{Cm}.
$$

Let  $\text{Gau}(E)$  denote the set of smoothing operators on  $L^2(E)$  with Gaussian propagation.

# **Proposition 3.2.** The set of operators  $\text{Gau}(E)$  is a  $*$ -algebra.

*Proof.* If A is a smoothing operator, then  $A^*$  is also a smoothing operator with  $k_{A^*}(x, y) = k_A(y, x)^*$ . Hence  $\text{Gau}(E)$  is closed under  $*$ . Clearly,  $Gau(E)$  is closed under linear combinations, so it remains to show that  $Gau(E)$  is closed under products. Let  $A, B \in Gau(E)$ , and let  $q_1, q_2 \in G$  with  $d(g_1, g_2) = l$ . Take any  $x_1 \in g_1 K$  and  $x_2 \in g_2 K$ . Then

$$
|k_{AB}(x_1, x_2)| \leq \int_M |k_A(x_1, y)k_B(y, x_2)| \operatorname{vol}(y)
$$
  
\n
$$
\leq \sum_{m=0}^{\infty} \sum_{g \in G_m} \int_{gg_1K} |k_A(x_1, y)k_B(y, x_2)| \operatorname{vol}(y)
$$
  
\n
$$
\leq \sum_{m=0}^{\infty} \sum_{g \in G_m} \int_{gg_1K} C_1 e^{-C_2(d(x_1, y)^2 + d(y, x_2)^2)} \operatorname{vol}(y)
$$

for some  $C_1, C_2 > 0$  which are independent of the choice of  $g_1, g_2, x_1, x_2$ . By (3.1) and (3.2),

$$
\sum_{m=0}^{\infty} \sum_{g \in G_m} \int_{gg_1K} C_1 e^{-C_2(d(x_1,y)^2 + d(y,x_2)^2)} \operatorname{vol}(y)
$$
  
\n
$$
\leq \sum_{m=0}^{\infty} \sum_{g \in G_m} C_1 e^{-C_3(m^2 + (m-l)^2)}
$$
  
\n
$$
\leq e^{-C_3 l^2} \sum_{m=0}^{\infty} C_1 e^{-2C_3(m^2 - l_m) + C_4 m}
$$
  
\n
$$
\leq C_5 e^{-C_3 l^2}
$$

for some  $C_3, C_4, C_5 > 0$  which are independent of the choice of  $g_1, g_2, x_1, x_2$ . Moreover, by  $(3.1)$ ,

$$
C_5 e^{-C_3 l^2} \le C_6 e^{-C_7 d(x_1, x_2)^2}.
$$

for some  $C_6, C_7 > 0$  which are independent of the choice of  $g_1, g_2, x_1, x_2$ . Then we obtain  $|k_{AB}(x_1,x_2)| \leq C_6 e^{-\tilde{C}_7 d(x_1,x_2)^2}$ . Thus AB belongs to  $\text{Gau}(E)$ , completing the proof.

**Proposition 3.3.** Elements of  $\text{Gau}(E)$  are piecewise Hilbert-Schmidt.

*Proof.* The proof of Lemma 2.4 implies that a smoothing operator  $A: L^2(E) \to$  $L^2(E)$  is piecewise Hilbert-Schmidt if and only if

(3.3) 
$$
\sup_{g\in G} \int_M \int_{gK} |k_A(x,y)|^2 \operatorname{vol}(y) \operatorname{vol}(x) < \infty.
$$

Now we let  $A \in \text{Gau}(E)$ , and take any  $h \in G$  and  $y \in hK$ . By (3.1),

$$
\int_M |k_A(x,y)|^2 \operatorname{vol}(x) = \sum_{g \in G} \int_{gK} |k_A(x,y)|^2 \operatorname{vol}(x) \le \sum_{g \in G} C_1 e^{-C_2 d(g,h)^2}
$$

for some  $C_1, C_2 > 0$  which are independent of the choice of h and y. By  $(3.2)$ , we have

$$
\sum_{g \in G} C_1 e^{-C_2 d(g,h)^2} \le \sum_{m=0}^{\infty} \sum_{g \in G_m} C_1 e^{-C_2 d(gh,h)^2} \le \sum_{m=0}^{\infty} C_1 e^{-C_2 m^2 + C_3 m} < C_4.
$$

for some  $C_3, C_4 > 0$  which are independent of the choice of h and y. Then  $(3.3)$  holds, hence A is piecewise Hilbert-Schmidt as stated.

**Corollary 3.4.** If  $A, B \in \text{Gau}(E)$ , then AB is of piecewise trace-class.

*Proof.* If  $A, B \in \text{Gau}(E)$ , then by Propositions 3.2 and 3.3,  $A^*$  and B are piecewise Hilbert-Schmidt operators. Hence  $AB = (A^*)^*B$  is of piecewise trace-class.

We consider a function  $F: G \times G \to \mathbb{R}$  such that there are  $C_1, C_2 > 0$ satisfying

(3.4) 
$$
|F(g,h)| \leq C_1 e^{-C_2 d(g,h)^2}
$$

for all  $q, h \in G$ .

**Lemma 3.5.** There is  $C > 0$  such that for any  $q \in G$ , we have

$$
\sum_{h \in G} |F(h,g)| \le C \quad and \quad \sum_{h \in G} |F(g,h)| \le C.
$$

*Proof.* By (3.2), there is  $C_3 > 0$  such that for any  $m \geq 0$  and  $g \in G$ ,

$$
\sum_{h \in G_m} |F(hg, g)| = \sum_{k=0}^m \sum_{h \in G_k - G_{k-1}} |F(hg, g)| \le \sum_{k=0}^m C_1 e^{-C_2 k^2 + C_3 k}.
$$

Clearly, the last term is bounded, and then by taking  $m \to \infty$ , the first inequality is obtained. By symmetry, the second inequality is also obtained.  $\Box$ 

By Lemma 3.5, we can define bounded functions  $F_1, F_2: G \to \mathbb{R}$  by

$$
F_1(g) = \sum_{h \in G} F(h, g)
$$
 and  $F_2(g) = \sum_{h \in G} F(g, h)$ .

Let  $\widehat{\mathcal{I}}$  denote the closure of  $\mathcal{I}$  by the  $\ell^{\infty}$ -norm topology. Note that  $\widehat{\mathcal{I}} \subset \overline{\mathcal{I}}$ .

**Lemma 3.6.**  $F_1 \equiv F_2 \mod \widehat{\mathcal{I}}$ .

*Proof.* For  $m \geq 0$  and  $q \in G$ , let

$$
F_1^m(g) = \sum_{h \in G_m} F(g, hg) \text{ and } F_2^m(g) = \sum_{h \in G_m} F(h^{-1}g, g).
$$

By Lemma 3.5,  $F_1^m$  and  $F_2^m$  are bounded functions on G. By  $(3.2)$ ,

$$
|F_1(g) - F_1^m(g)| \le \sum_{k=m}^{\infty} \sum_{h \in G_k - G_{k-1}} |F(g, hg)| \le \sum_{k=m}^{\infty} C_1 e^{-C_2 k^2 + C_3 k} \le C_4 e^{-C_5 m^2}
$$

for some  $C_3, C_4, C_5 > 0$  which are independet of the choice of m and g. Then  $F_1^m$  converges uniformly to  $F_1$  as  $m \to \infty$ . By symmetry,  $F_2^m$  also converges uniformly to  $F_2$  as  $m \to \infty$ . For  $g \in G$ , we define a function  $\phi_g : G \to \mathbb{R}$  by  $\phi_g(h) = F(g, gh)$ . By (3.4),  $\phi_g$  is bounded. We set

$$
\Phi_m(h) = \sum_{g \in G_m} (1 - h^{-1}) \phi_g(h).
$$

Then  $\Phi_m \in \mathcal{I}$  and  $F_2^m - F_1^m = \Phi_m$ . Then  $\Phi_m$  converges uniformly to  $F_2 - F_1$ as  $m \to \infty$ , implying  $F_1 - F_2 \in \overline{J}$ .

**Proposition 3.7.** For  $A, B \in \text{Gau}(S)$ , we have

$$
\text{Tr}(AB) \equiv \text{Tr}(BA) \mod 1.
$$

Proof. By Corollary 3.4, AB and BA are piecewise trace-class operators, so their piecewise traces are defined. We consider a function  $F: G \times G \to \mathbb{R}$ given by

$$
F(g, h) = \int_{gK \times hK} \text{tr}(k_A(x, y)k_B(y, x)) \text{ vol}(x) \text{ vol}(y).
$$

Since A and B have Gaussian propagation, there are  $C_1, C_2 > 0$  such that for any  $g, h \in G$ ,

$$
|F(g,h)| \leq \int_{gK \times hK} |\text{tr}(k_A(x,y)k_B(y,x))| \text{vol}(x) \text{vol}(y)
$$
  

$$
\leq \int_{gK \times hK} C_1 e^{-C_2 d(x,y)^2} \text{vol}(x) \text{vol}(y).
$$

Then by  $(3.1)$ , the function F satisfies  $(3.4)$ . By Lemma 2.7, we have

$$
F_1(g) = \sum_{h \in G} F(h, g)
$$
  
= 
$$
\int_{gK} tr \left( \int_M k_A(x, y) k_B(y, x) vol(y) \right) vol(x)
$$
  
= 
$$
\int_{gK} tr (k_{AB}(x, x)) vol(x)
$$
  
= Tr
$$
Tr(AB)(g)
$$

for any  $g \in G$ , implying  $F_1 = \text{Tr}(AB)$ . Quite similarly, we can see that  $F_2 =$ Tr(BA). Thus by Lemma 3.6, we obtain  $\text{Tr}(AB) \equiv \text{Tr}(BA) \mod \hat{J}$ .  $\Box$ 

3.2. Functional calculus. In the sequel, let  $S \to M$  be a G-invariant Clifford bundle with Dirac operator  $D$ . Then  $S$  has bounded geometry in the sense of  $[18, p. 93]$ , and the Dirac operator D is G-invariant. We consider a generalized Dirac operator

$$
\widetilde{D} = D + A
$$

where  $A$  is a self-adjoint endomorphism of  $S$ , not necessarily  $G$ -invariant. For the rest of this section, we assume that  $\nabla^i A$  is bounded for  $i = 0, 1, \ldots, \lceil \frac{n}{2} \rceil$  $\frac{n}{2}$ ] — 1. Let  $C_c^{\infty}(S)$  denote the set of compactly supported smooth sections of S.

**Lemma 3.8.** There is  $C_k > 0$  satisfying an ineuality

$$
\sum_{i=0}^{k} \|D^i u\|^2 \le C_k \sum_{i=0}^{k} \|\widetilde{D}^i u\|^2
$$

for  $k = 0, 1, \ldots \lceil \frac{n}{2} \rceil$  $\frac{n}{2}$  and any  $u \in C_c^{\infty}(S)$ .

*Proof.* We induct on k. The  $k = 0$  case is trivial. We assume that the statement holds for  $k \leq m-1$ . Then

$$
||D^m u|| = ||D^{m-1}(\tilde{D} - A)u||
$$
  
\n
$$
\leq ||D^{m-1}\tilde{D}u|| + ||D^{m-1}Au||
$$
  
\n
$$
\leq C_1 \left(\sum_{i=0}^{m-1} ||\tilde{D}^{i+1}u||^2\right)^{\frac{1}{2}} + ||D^{m-1}Au||
$$

for some  $C_1 > 0$  which is independent of the choice of u. We also have

$$
||D^{m-1}Au||^2 \leq C_2 \sum_{i=0}^{m-1} ||\nabla^i(Au)||^2 \leq C_3 \sum_{i=0}^{m-1} ||\nabla^i u||^2 \leq C_4 \sum_{i=0}^{m-1} ||D^i u||^2
$$

for some  $C_2, C_3, C_4 > 0$  which are independent of the choice of u. Here, the first inequality follows as in  $[21, \text{ page } 73]$ , the second inequality follows from  $\nabla(Au) = (\nabla A - A\nabla)u$  and the assumption that  $\nabla^i A$  is bounded for  $i=0,1,\ldots\lceil\frac{n}{2}\rceil$  $\frac{n}{2}$  – 1, and the third inequality follows from the elliptic estimate (cf.  $[21,$  Proposition 5.16]). Then by the induction hypothesis, we obtain

$$
||D^{m-1}Au||^2 \leq C_5 \sum_{i=0}^{m-1} ||\widetilde{D}^i u||^2
$$

for some  $C_5 > 0$  which is independent of the choice of u, and thus the statement follows.  $\Box$ 

Let G denote the set of Schwartz-class functions  $\phi: \mathbb{R} \to \mathbb{R}$  such that there are  $C_1, C_2 > 0$  satisfying

$$
|\widehat{\phi}^{(i)}(x)| \le C_1 e^{-C_2 x^2}
$$

for  $i = 0, 1, ..., 2\lceil \frac{n}{2} \rceil$  $\frac{n}{2}$ , where  $\widehat{\phi}$  denotes the Fourier transform of  $\phi$  and  $f^{(i)}$ stands for the *i*-th derivative of a function  $f: \mathbb{R} \to \mathbb{R}$ .

Proposition 3.9. The set of functions  $\mathcal G$  is an algebra.

*Proof.* The only nontrivial property is that  $\mathcal{G}$  is closed under products. For Schwartz-class functions  $\phi$  and  $\psi$ , we have  $(\phi \psi)^{(i)} = \hat{\phi}^{(i)} * \hat{\psi}$ . Then since the convolution of Gaussian functions is a Gaussian function, G is closed under  $\Box$ 

**Lemma 3.10.** A generalized Dirac operator  $\widetilde{D}$  is essentially self-adjoint.

*Proof.* By [7, Theorem 1.17], D is essentially self-adjoint. Then as A is self-adjoint and continuous,  $\widetilde{D}$  is essentially self-adjoint too.

Lemma 3.10 enables us to perform functional calculus for a generalized Dirac operator  $\overline{D}$ . We remark that all results in [21, Chapter 9] holds not only for the Dirac operator D but also for a generalized Dirac opertor  $\widetilde{D}$  as mentioned in  $[21, p. 95]$ . In particular, we have:

**Lemma 3.11** (cf. [21, Proposition 7.20]). For any  $u \in C_c^{\infty}(S)$ ,  $e^{\sqrt{-1}t\widetilde{D}}u$  is supported within  $N(\text{supp}(u); |t|)$ .

For  $\phi \in \mathcal{G}$ , we can define the operator  $\phi(\widetilde{D})$  as in [21, Chapter 9], that is,

$$
\langle u, \phi(\tilde{D})v \rangle = \frac{1}{2\pi} \int_{\mathbb{R}} \widehat{\phi}(t) \langle u, e^{\sqrt{-1}t\tilde{D}}v \rangle dt
$$

for  $u, v \in C_c^{\infty}(S)$ .

Proposition 3.12. The map

 $\mathcal{G} \to \text{Gau}(S), \quad \phi \mapsto \phi(\widetilde{D})$ 

is a well-defined homomorphism.

*Proof.* Let  $\widetilde{K}$  denote the  $\epsilon$ -neighborhood of K for some  $\epsilon > 0$ . We take any  $g, h \in G$ , and let  $\delta$  be the distance between  $g\widetilde{K}$  and  $h\widetilde{K}$ . We also take any  $u \in L^2(S|_{g\widetilde{K}})$  and  $v \in C_c^{\infty}(S)$  with supp $(v) \subset h\widetilde{K}$ . Let  $\phi \in \mathcal{G}$ . Then by Lemma  $3.\overline{1}1$ , we have

$$
\begin{split}\n&\|\langle u, \widetilde{D}^i(\phi(\widetilde{D})\widetilde{D}^j v)|_{g\widetilde{K}}\rangle_{L^2(S|_{g\widetilde{K}})}| \\
&= |\langle u, \widetilde{D}^i(\phi(\widetilde{D})\widetilde{D}^j v)\rangle_{L^2(S)}| \\
&= \frac{1}{2\pi} \left| \int_{\mathbb{R}} \widehat{\phi}(t) \langle u, e^{\sqrt{-1}t\widetilde{D}} \widetilde{D}^{i+j} v \rangle_{L^2(S)} dt \right| \\
&= \frac{1}{2\pi} \left| \int_{\mathbb{R}} \widehat{\phi}^{(i+j)}(t) \langle u, e^{\sqrt{-1}t\widetilde{D}} v \rangle_{L^2(S)} dt \right| \\
&\leq \frac{1}{2\pi} \int_{|t| \geq \delta} C_1 e^{-C_2 t^2} dt \|u\|_{L^2(S|_{g\widetilde{K}})} \|v\|_{L^2(S|_{h\widetilde{K}})} \\
&\leq C_3 e^{-C_2 \delta^2} \|u\|_{L^2(S|_{g\widetilde{K}})} \|v\|_{L^2(S|_{h\widetilde{K}})}\n\end{split}
$$

for some  $C_1, C_2, C_3 > 0$  which are independent of the choice of  $u, v$ , where  $i, j = 0, 1, \ldots, \lceil \frac{n}{2} \rceil$  $\frac{n}{2}$ . Hence as u is an arbitrary element of  $L^2(S|_{g\widetilde{K}})$ , we get

(3.5) 
$$
\|\widetilde{D}^{i}((\varphi(\widetilde{D})\widetilde{D}^{j}v)|_{g\widetilde{K}})\|_{L^{2}(S|_{g\widetilde{K}})} \leq C_{3}e^{-C_{2}\delta^{2}}\|v\|_{L^{2}(S|_{h\widetilde{K}})}
$$

for  $i, j = 0, 1, \ldots, \lceil \frac{n}{2} \rceil$  $\frac{n}{2}$ . By Lemma 3.8, we have the Sobolev inequality

$$
\sup_{x\in g\widetilde{K}}|(\varphi(\widetilde{D})\widetilde{D}^{j}v)(x)|\leq C_4\left(\sum_{i=0}^{\lceil\frac{n}{2}\rceil}\|\widetilde{D}^{i}(\varphi(\widetilde{D})\widetilde{D}^{j}v)|_{g\widetilde{K}}\|_{L^2(S|_{g\widetilde{K}})}^2\right)
$$

 $\frac{1}{2}$ 

 $\begin{array}{c} \hline \end{array}$  $\overline{\phantom{a}}$ I  $\overline{\phantom{a}}$ 

for some  $C_4 > 0$  which is independent of the choice of v and g, where  $j = 0, 1, \ldots, \lceil \frac{n}{2} \rceil$  $\frac{n}{2}$ . Then by  $(3.5)$ , we obtain

(3.6) 
$$
\sup_{x \in g\widetilde{K}} |(\varphi(\widetilde{D})\widetilde{D}^j v)(x)| \leq \lceil \frac{n}{2} \rceil C_3 C_4 e^{-C_2 \delta^2} ||v||_{L^2(S|_{h\widetilde{K}})}
$$

for  $j = 0, 1, ..., \lceil \frac{n}{2} \rceil$  $\frac{n}{2}$ . Now we take any  $x \in gK$  and  $u_x \in S_x$ . Let  $u =$  $\mathbf{1}_{h\widetilde{K}}(y)k_{\phi(\widetilde{D})}(x,-)^*u_x \in L^2(S|_{h\widetilde{K}})$ . It follows from  $(3.6)$  that

$$
\langle \widetilde{D}^j u, v \rangle_{L^2(S|_{h\widetilde{K}})} = \langle u, \widetilde{D}^j v \rangle_{L^2(S)}
$$
  
= 
$$
\left| \int_M \langle k_{\phi(\widetilde{D})}(x, y)^* u_x, v(y) \rangle \operatorname{vol}(y) \right|
$$
  
= 
$$
\left| \int_M \langle u_x, k_{\phi(\widetilde{D})}(x, y) v(y) \rangle \operatorname{vol}(y) \right|
$$
  
= 
$$
|\langle u_x, (\varphi(\widetilde{D}) \widetilde{D}^j v)(x) \rangle|
$$
  

$$
\leq \lceil \frac{n}{2} \rceil C_3 C_4 e^{-C_2 \delta} |u_x| \|v\|_{L^2(S|_{h\widetilde{K}})}
$$

for  $j = 0, 1, ..., \lceil \frac{n}{2} \rceil$  $\frac{n}{2}$ . Then since v is any element of  $C_c^{\infty}(S|_{h\widetilde{K}})$ , we get

$$
\|\widetilde{D}^{j}u\|_{L^{2}(S|_{h\widetilde{K}})}\leq C_{5}e^{-C_{2}\delta^{2}}|u_{x}|.
$$

for some  $C_5 > 0$  which is independent of the choice of x and  $u_x$ , where  $j = 0, 1, \ldots, \lceil \frac{n}{2} \rceil$  $\frac{n}{2}$ . Thus by arguing as above, we obtain

$$
|k(x,y)^*u_x| \le C_7 e^{-C_2 \delta^2} |u_x|
$$

for some  $C_{\overline{z}} > 0$  which is independent of the choice of g and h, where  $x \in g\widetilde{K}$ and  $y \in hK$ . As  $u_x$  is an arbitrary element of  $S_x$ , this implies that

2

$$
|k(x, y)| = |k(x, y)^*| \le C_7 e^{-C_2 \delta}
$$

for any  $x \in g\widetilde{K}$  and  $y \in h\widetilde{K}$ . Thus by (3.1), we obtain  $\phi(\widetilde{D}) \in \mathcal{G}$ . Clearly, the map in the statement is a homomorphism completing the proof the map in the statement is a homomorphism, completing the proof.

# 4. Piecewise normalized Betti number

In this section, we define piecewise normalized Betti numbers as an analogy to normalized Betti numbers defined by Roe [19, 20], and prove its invariance under a certain deformation.

4.1. **Abstract differential.** Let  $\Lambda^k = \Lambda^k T^* M$  for  $k = 0, 1, ..., n$ , and let  $\Lambda = \bigoplus_{k=0}^n \Lambda^k$ . Then  $\Lambda$  is a *G*-invariant Clifford bundle with Dirac operator  $D = d + d^*$ . Roe [19] defined an abstract differential on  $\Lambda$  and the normalized Betti number of it. Here, we define a slightly different operator adopted to our situation, but we ambiguously call it an abstract differential.

**Definition 4.1.** An operator  $\tilde{d}: C_c^{\infty}(\Lambda^k) \to C_c^{\infty}(\Lambda^{k+1})$  is an abstract differential if it satisfies that

- (1)  $\tilde{d}^2 = 0$ , and
- (2)  $\tilde{d} + \tilde{d}^*$  is a generalized Dirac operator, that is,  $D + A$  for some self-adjoint endomorphism A of  $\Lambda$ , such that  $\nabla^i A$  is bounded for  $i=0,1,\ldots\lceil\frac{n}{2}\rceil$  $\frac{n}{2}$ ] – 1.

We give the most important example of an abstract differential.

**Example 4.2.** Let  $f: M \to \mathbb{R}$  be a function. The Witten deformation of the differential d on  $\Lambda$  is defined by  $d_t = e^{-tf} de^{tf}$ . Clearly,  $d_t^2 = 0$  and

$$
d_t + d_t^* = D + tR
$$

where  $R = (df \wedge) - (df \wedge)$ . If f is bounded, that is,  $\nabla^i f$  is bounded for  $i=0,1,\ldots,\lceil\frac{n}{2}\rceil$  $\frac{n}{2}$ , then  $\nabla^i R$  is bounded for  $i = 0, 1, \ldots, \lceil \frac{n}{2} \rceil$  $\frac{n}{2}$ ] – 1. Thus  $d_t$  is an abstract differential.

We define piecewise normalized Betti numbers of an abstract differential by replacing trace with piecewise trace in the definition of normalized Betti numbers [20, p.766]. However, compactly supported nonzero functions do not belong to the algebra G. Then, instead of showing that  $\phi(D)$  is of piecewise trace-class for a compactly supported function  $\phi \colon \mathbb{R} \to \mathbb{R}$ , we extend the definition of piecewise trace.

**Definition 4.3.** Let  $A: L^2(\Lambda) \to L^2(\Lambda)$  be an operator with bounded kernel function. We define a bounded function

$$
\widetilde{\text{Tr}}(A) \colon G \to \mathbb{R}, \quad g \mapsto \int_{gK} \text{tr}(k_A(x, x)) \text{vol}(x).
$$

By Lemma 2.7, we have:

**Proposition 4.4.** If  $A: L^2(\Lambda) \to L^2(\Lambda)$  is of piecewise trace-class, then

$$
\operatorname{Tr}(A) = \operatorname{Tr}(A).
$$

Let  $\tilde{d}$ :  $\Lambda \to \Lambda$  be an abstract differential, and let  $\tilde{D} = \tilde{d} + \tilde{d}^*$ .

**Lemma 4.5.** For any Schwartz-class function  $\phi: \mathbb{R} \to \mathbb{R}$ ,  $\phi(\widetilde{D})$  has a bounded kernel function.

*Proof.* By Lemma 3.8, there is an embedding  $\widetilde{W}^k \to W^k$ , where  $\widetilde{W}^k$  and  $W^k$  denote the k-th Sobolev spaces with respect to  $\widetilde{D}$  and D, respectively. Then the statement can be verified quite similarly to Proposition 3.12.  $\Box$ 

Let C denote the set of compactly supported functions  $\phi \colon \mathbb{R}_{\geq 0} \to \mathbb{R}$  such that  $\phi(x^2)$  is smooth,  $0 \le \phi \le 1$  and  $\phi(0) = 1$ .

Definition 4.6. We define the *piecewise normalized* k-th Betti number associated to an abstract differential  $d$  as a bounded function

$$
b_k(\tilde{d})
$$
:  $G \to \mathbb{R}$ ,  $g \mapsto \inf_{\phi \in \mathcal{C}} \widetilde{\text{Tr}}(\phi(\tilde{D}^2|_{\Lambda^k}))(g)$ .

Clearly,  $\text{Tr}(\phi(\tilde{D}^2|_{\Lambda^k})) \geq 0$  for  $\phi \in \mathcal{C}$ , so the infimum in the above definition exists. We choose  $C > 0$  satisfying  $|\text{Tr}(e^{-\tilde{D}^2})| \leq C1$ . We say that a subset  $\mathcal{C}_0 \subset \mathcal{C}$  is *cofinal* if for any  $\tilde{\phi} \in \mathcal{C}$  and  $\epsilon > 0$ , there is  $\phi \in \mathcal{C}_0$  satisfying

(4.1) 
$$
\tilde{\phi}(x^2) + C^{-1} \epsilon e^{-x^2} \ge \phi(x^2).
$$

**Lemma 4.7.** If  $\mathcal{C}_0$  is a cofinal subset of  $\mathcal{C}$ , then for each  $g \in G$ ,

$$
b_k(\tilde{d})(g) = \inf_{\phi \in \mathfrak{C}_0} \widetilde{\text{Tr}}(\phi(\widetilde{D}^2|_{\Lambda_k}))(g).
$$

*Proof.* Take any  $\tilde{\phi} \in \mathcal{C}$  and  $\epsilon > 0$ . Then there is  $\phi \in \mathcal{C}_0$  satisfying (4.1), so we get

$$
\widetilde{\text{Tr}}(\tilde{\phi}(\widetilde{D}^2|_{\Lambda_k})) + \epsilon \geq \widetilde{\text{Tr}}(\phi(\widetilde{D}^2|_{\Lambda_k})).
$$

Thus as  $\epsilon$  can be arbitrarily small, we obtain

$$
b_k(\tilde{d})(g) \ge \inf_{\phi \in \mathcal{C}_0} \widetilde{\text{Tr}}(\phi(\tilde{D}^2|_{\Lambda_k}))(g)
$$

for each  $g \in G$ . The reverse inequality is obvious by the definition of  $b_k(\tilde{d})$ , and therefore the statement follows.

Any function on  $[0, 1]$  may be considered as a function on  $\mathbb{R}$  by extending it by zero outside of [0, 1]. Then we can define  $\text{Tr}(\phi(\tilde{D}^2|_{\Lambda^k}))$  for any  $\phi \in$  $C^{\infty}[0,1].$ 

**Lemma 4.8.** For any bounded linear function  $\alpha: \ell^{\infty}(G) \to \mathbb{R}$ , there is a signed regular Borel measure  $\mu$  on [0, 1] such that for any  $\phi \in C^{\infty}[0,1],$ 

$$
\alpha(\widetilde{\text{Tr}}(\phi(\widetilde{D}^2|_{\Lambda^k}))) = \int_0^1 \phi d\mu.
$$

*Proof.* There is  $C_0 > 0$  such that

$$
(4.2) \quad |\widetilde{\text{Tr}}(\phi(\widetilde{D}^2|_{\Lambda^k}))| \le \sup_{x \in [0,1]} |\phi(x)| |\widetilde{\text{Tr}}(\mathbf{1}_{[0,1]}(\widetilde{D}^2|_{\Lambda^k}))| \le C_0 \sup_{x \in [0,1]} |\phi(x)| \mathbb{1}
$$

for any  $\phi \in C^{\infty}[0,1]$ . Then the map

$$
C^{\infty}[0,1] \to \mathbb{R}, \quad \phi \mapsto \widetilde{\text{Tr}}(\phi(\widetilde{D}^2|_{\Lambda^k}))
$$

is a bounded linear function. Thus the statement follows from the Riesz representation theorem  $[22, 6.19$  Theorem.

**Proposition 4.9.** The piecewise trace  $\text{Tr}(e^{-t\widetilde{D}^2|_{\Lambda^k}})$  converges weakly to  $b_k(\tilde{d})$  as  $t \to \infty$ .

*Proof.* Clearly, there is  $\phi \in \mathcal{C}$  satisfying  $e^{-x^2} \geq \phi(x^2)$ . Let  $\mathcal{C}_0 = \{\phi(tx) \mid$  $t > 0$ . Then it is easy to see that  $\mathcal{C}_0$  is a cofinal subset of  $\mathcal{C}$ . Hence by Lemma 4.8 and Lebesgue's monotone convergence theorem,  $\text{Tr}(\phi(t\bar{D}^2|_{\Lambda^k}))$ converges weakly to  $b_k(\tilde{d})$  as  $t \to \infty$ . Observe that for any  $\epsilon, s > 0$ , there is  $t > 0$  satisfying

$$
\phi(sx^2) + C^{-1} \epsilon e^{-x^2} \ge e^{-tx^2}
$$

where  $C > 0$  is the constant in the definition of a cofinal subset of C. Then

$$
\widetilde{\text{Tr}}(\phi(t\widetilde{D}^2|_{\Lambda^k})) \le \text{Tr}(e^{-t\widetilde{D}^2|_{\Lambda^k}}) \le \widetilde{\text{Tr}}(\phi(s\widetilde{D}^2|_{\Lambda^k})) + \epsilon \mathbb{1}.
$$

Let  $\alpha: \ell^{\infty}(G) \to \mathbb{R}$  be a positive bounded linear function. Then we have

$$
\alpha(\widetilde{\text{Tr}}(\phi(t\widetilde{D}^2|_{\Lambda^k}))) \leq \alpha(\text{Tr}(e^{-t\widetilde{D}^2|_{\Lambda^k}})) \leq \alpha(\widetilde{\text{Tr}}(\phi(s\widetilde{D}^2|_{\Lambda^k}))) + \epsilon \alpha(\mathbb{1}).
$$

Note that  $\epsilon$  can be arbitrarily small, and accordingly, s and t can be arbitrarily large. Then we get

(4.3) 
$$
\lim_{t \to \infty} \alpha(\text{Tr}(e^{-t\widetilde{D}^2|_{\Lambda^k}})) \to \alpha(b_k(\widetilde{d})).
$$

As in [5, Chapter IV, 8.16], any bounded linear function on  $\ell^{\infty}(G)$  is given by the integral with a bounded finitely additive signed measure, so by [5, Chapter III, 1.8], it admits the Jordan decomposition. Then (4.3) holds for any bounded linear function  $\alpha$ :  $\ell^{\infty}(G) \to \mathbb{R}$ , completing the proof.  $\Box$ 

4.2. Invariance. The rest of this section is devoted to prove the following theorem which is an analogy to  $[20, (2.6)$  Theorem.

**Theorem 4.10.** Let  $F: \Lambda \to \Lambda$  be a bounded self-adjoint automorphism, and let  $\tilde{d}_1, \tilde{d}_2$  be abstract differentials satisfying

$$
\tilde{d}_1 F = F \tilde{d}_2.
$$

Then for  $k = 0, 1, \ldots, n$ , we have

$$
b_k(\tilde{d}_1) \approx b_k(\tilde{d}_2) \mod \overline{\mathfrak{I}}.
$$

The proof of this theorem is basically a modification of that of  $[20, (2.6)]$ Theorem].

**Lemma 4.11** (cf. [20, Theorem 4.1]). Let  $E: L^2(\Lambda^k) \to L^2(\Lambda^k)$  be a selfadjoint piecewise trace-class operator. Then

$$
\text{Tr}(E) \ge -\rho_1(E - E^2).
$$

*Proof.* By Proposition 2.3,  $E^2$  is of piecewise trace-class, so  $\rho_1(E - E^2)$  and the piecewise traces of  $E^2$  and  $E - E^2$  are defined. As E is self-adjoint,  $E^2$ is positive, so by Proposition 2.10,  $\text{Tr}(E^2) \geq 0$ . Then we get

$$
\text{Tr}(E) = \text{Tr}(E - E^2) + \text{Tr}(E^2) \ge \text{Tr}(E - E^2) \ge -\rho_1(E - E^2)
$$

and hence the statement follows.

For positive integers  $p$  and  $q$ , we set

$$
P_p = e^{-p(\tilde{d}_1 + \tilde{d}_1^*)^2|_{\Lambda^k}} \quad \text{and} \quad Q_q = F^{-1}e^{-q(\tilde{d}_2 + \tilde{d}_2^*)^2|_{\Lambda^k}F}
$$

where  $\tilde{d}_1, \tilde{d}_2$  and F are as in Theorem 4.10.

**Lemma 4.12.** The operator  $Q_q$  is of piecewise trace-class such that

$$
\operatorname{Tr}(Q_q) = \operatorname{Tr}(FQ_q F^{-1}).
$$

Proof. Observe that

$$
k_{AF^{-1}}(x, y) = k_A(x, y)F^{-1}
$$
 and  $k_{AF}(x, y) = k_A(x, y)F$ 

where  $A = e^{-\frac{q}{2}(\tilde{d}_2 + \tilde{d}_2^*)^2|_{\Lambda^k}}$ . Then since F is bounded and A is piecewise Hilbert-Schmidt, the proof of Lemma 2.4 implies that  $AF^{-1}$  and  $AF$  are piecewise Hilbert-Schmidt, implying that  $Q_q = (AF^{-1})^*(AF)$  is of piecewise trace-class. Moreover, we have

$$
\text{tr}(k_{FQ_qF^{-1}}(x,y)) = \text{tr}(Fk_{Q_q}(x,y)F^{-1}) = \text{tr}(k_{Q_q}(x,y)).
$$

Thus by Lemma 2.7, the statement follows.  $\square$ 

**Lemma 4.13** (cf.  $[20, (5.4)$  Corollary). There is an inequality

$$
\rho_2(P_{2p}-P_pQ_qP_p)\leq C\sqrt{\frac{q}{p-1}}\mathbb{1}.
$$

for some  $C > 0$  which is independent of the choice of p and q. *Proof.* By  $[20, (5.3)$  Proposition, we have

$$
\|P_{2p-2}-P_{p-1}Q_qP_{p-1}\|\leq 4\|F\|\|F^{-1}\|\sqrt{\frac{q}{p-1}}
$$

so by Propositions 2.3, we get

$$
\rho_2(P_{2p} - P_p Q_q P_p) \le ||P_1|| ||P_{2p-2} - P_{p-1} Q_q P_{p-1} || \rho_2(P_1) \le C \sqrt{\frac{q}{p-1}} 1
$$

for some  $C > 0$  which is independent of the choice of p and q.

**Lemma 4.14** (cf. [20, (6.2) Proposition]). Let  $f : \mathbb{N} \to \mathbb{N}$  be an increasing function. Then  $\rho_1(P_p - P_{f(p)})$  and  $\rho_1(Q_q - Q_{f(q)})$  converge weakly to zero as  $p \to \infty$  and  $q \to \infty$ .

*Proof.* By Lemma 4.9,  $Tr(P_p)$  is weakly convergent as  $p \to \infty$ , so  $\rho_1(P_p - p)$  $P_{f(p)}) = \text{Tr}(P_p - P_{f(p)})$  converges weakly to zero as  $p \to \infty$ . By Lemma 4.12, we also get the result for  $\rho_1(Q_q - Q_{f(q)})$ .  $\Box$ 

**Lemma 4.15.** Let  $A: L^2(\Lambda) \to L^2(\Lambda)$  be a bounded operator, and let  $B: L^2(\Lambda) \to$  $L^2(\Lambda)$  be a piecewise Hilbert-Schmidt operator. Then for  $p + q \geq 1$ ,

$$
\rho_1(ABP_pQ_q) \le C||A||\rho_2(B)
$$

where  $C > 0$  is independent of the choice of  $A, B, p, q$ .

Proof. By Propositions 2.3 and 2.11, we have

$$
\rho_1(ABP_pQ_q) \le \rho_2(AB)\rho_2(P_pQ_q) \le ||A||\rho_2(B)\rho_2(P_pQ_q).
$$

If  $q > 0$ , then by Proposition 2.3,

$$
\rho_2(P_p Q_q) \leq ||P_p Q_{q-1}|| \rho_2(Q_1) = \rho_2(Q_1).
$$

If  $q = 0$ , then we can see quite similarly that  $\rho_2(P_pQ_q) \leq \rho_2(P_1)$ . Since  $\rho_2(P_1)$  and  $\rho_2(Q_1)$  are bounded functions, the statement follows.

**Lemma 4.16.** Let  $\alpha: \ell^{\infty}(G) \to \mathbb{R}$  be a positive bounded linear function. Suppose that for  $0 < \epsilon < 1$  and  $i = 0, 1$ ,

$$
\alpha(\rho_1(P_p - P_{2p+i})) < \epsilon \qquad \alpha(\rho_1(Q_q - Q_{2q+i})) < \epsilon
$$
  
\n
$$
\alpha(\rho_1(P_p - P_{p+1})) < \epsilon \qquad \rho_2(P_{2p} - P_pQ_qP_p) < \epsilon \mathbb{1}.
$$

Then there is an inequality

$$
\alpha(\text{Tr}(P_p)) < \alpha(\text{Tr}(Q_q)) + \alpha(\text{Tr}([P_pQ_q, P_{p+1}])) \\
 &+ \alpha(\text{Tr}([P_pQ_{q+1}, Q_q])) + C\epsilon^{\frac{1}{2}}
$$

for some  $C > 0$  which is independent of the choice of  $p, q, \epsilon$ .

Proof. By Lemma 4.15, we have

$$
\begin{aligned} \text{Tr}(P_p) &\le \rho_1(P_p - P_{2p+1}) + \rho_1((P_{2p} - P_p Q_q P_p) P_1) + \text{Tr}(P_p Q_q P_{p+1}) \\ &\le \rho_1(P_p - P_{2p+1}) + C_1 \rho_2(P_{2p} - P_p Q_q P_p) + \text{Tr}(P_p Q_q P_{p+1}) \end{aligned}
$$

for some  $C_1 > 0$  which is independent of the choice of  $p, q, \epsilon$ . Then we get

 $\alpha(\text{Tr}(P_p)) \leq \alpha(\text{Tr}(P_pQ_qP_{p+1})) + (1+C_1)\epsilon.$ 

By Lemma 4.15, we also have

$$
\begin{aligned} \text{Tr}(P_p Q_q P_{p+1}) &\leq \text{Tr}(P_p Q_q) + \rho_1 ((P_{2p+1} - P_p) Q_q) + \text{Tr}([P_p Q_q, P_{p+1}]) \\ &\leq \text{Tr}(P_p Q_q) + C_2 \rho_2 (P_{2p+1} - P_p) + \text{Tr}([P_p Q_q, P_{p+1}]) \end{aligned}
$$

for some  $C_2 > 0$  which is independent of the choice of  $p, q, \epsilon$ . Observe that by Proposition 2.3, 1

(4.4) 
$$
\rho_2(P_{p_1} - P_{p_2}) = \rho_1((P_{p_1} - P_{p_2})^2)^{\frac{1}{2}} \le \rho_1(P_{p_1} - P_{p_2})^{\frac{1}{2}}
$$
  
for  $p_1 < p_2$  because  $(e^{-p_1x} - e^{-p_2x})^2 \le e^{-p_1x} - e^{-p_2x}$ . Then we get

$$
\alpha(\text{Tr}(P_p Q_q P_{p+1})) \leq \alpha(\text{Tr}(P_p Q_q)) + \alpha(\text{Tr}([P_p Q_q, P_{p+1}])) + C_2 \epsilon^{\frac{1}{2}}.
$$

Since  $\rho_2(P_p) \leq ||P_{p-1}|| \rho_2(P_1) = \rho_2(P_1)$  by Proposition 2.3, we have  $\text{Tr}(P_pQ_q) \leq \text{Tr}(P_pQ_{2q+1}) + \rho_1(P_p(Q_q - Q_{2q+1}))$ 

$$
\leq \text{Tr}(P_p Q_{2q+1}) + \rho_2(P_p)\rho_2(Q_q - Q_{2q+1})
$$
  
\n
$$
\leq \text{Tr}(P_p Q_{2q+1}) + \rho_2(P_1)\rho_2(Q_q - Q_{2q+1})
$$
  
\n
$$
= \text{Tr}(Q_q P_p Q_{q+1}) + \text{Tr}([P_q Q_{q+1}, Q_q]) + \rho_2(P_1)\rho_2(Q_q - Q_{2q+1}).
$$

Quite similarly to (4.4), we can see that

(4.5) 
$$
\rho_2(Q_{q_1} - Q_{q_2}) \leq ||F||^2 ||F^{-1}||^2 \rho_1 (Q_{q_1} - Q_{q_2})^{\frac{1}{2}}
$$

for  $q_1 < q_2$ . Then we get

$$
\alpha(\text{Tr}(P_pQ_q))\leq \alpha(\text{Tr}(Q_qP_pQ_{q+1}))+\alpha(\text{Tr}([P_qQ_{q+1},Q_q]))+C_3\epsilon^{\frac{1}{2}}
$$

for some  $C_3 > 0$  which is independent of the choice of  $p, q, \epsilon$ . Summarizing, we obtain

$$
\alpha(\text{Tr}(P_p)) \leq \alpha(\text{Tr}(Q_q P_p Q_{q+1})) + \alpha(\text{Tr}([P_p Q_q, P_{p+1}]))
$$

$$
+ \alpha(\text{Tr}([P_p Q_{q+1}, Q_q])) + C_4 \epsilon^{\frac{1}{2}}
$$

for some  $C_4 > 0$  which is independent of the choice of  $p, q, \epsilon$ . Now we set  $E = Q_q - Q_q P_p Q_{q+1}$ . Then

$$
E - E2 = (Qq - Q2q) + (Q2q - Qq)PpQq+1 + QqPpQ1(Q2q - Qq)+ Qq(Pp - P2p)Qq+1 + Qq(P2p - PpQqPp)Qq+1+ QqPp(Qq - Q2q+1)PpQq+1.
$$

and so by Lemma  $4.15$  together with  $(4.4)$  and  $(4.5)$ ,

$$
\alpha(\rho_1(E - E^2)) \le C_5 \epsilon^{\frac{1}{2}}
$$

for some  $C_5 > 0$  which is independent of the choice of  $p, q, \epsilon$ . Hence by Lemma 4.11, we get

$$
\alpha(\text{Tr}(Q_q P_p Q_{q+1})) = \alpha(\text{Tr}(Q_q)) - \alpha(\text{Tr}(E)) \leq \alpha(\text{Tr}(Q_q)) + C_5 \epsilon^{\frac{1}{2}}.
$$

Thus the statement follows.

Now we are ready to prove Theorem 4.10.

*Proof of Theorem 4.10.* Let  $\alpha$ :  $\ell^{\infty}(G) \to \mathbb{R}$  be any bounded positive linear function. By Lemmas 4.13 and 4.14, there are integer sequences  $p_1 < p_2$  $\cdots$  and  $q_1 < q_2 < \cdots$  such that

$$
\alpha(\rho_1(P_{p_i} - P_{2p_i + j})) < \frac{1}{i} \qquad \alpha(\rho_1(Q_{q_i} - Q_{2q_i + j})) < \frac{1}{i}
$$
  

$$
\alpha(\rho_1(P_{p_i} - P_{p_i + 1})) < \frac{1}{i} \qquad \rho_2(P_{2p_i} - P_{p_i}Q_{q_i}P_{p_i}) < \frac{1}{i}1
$$

where  $j = 0, 1$ . By Lemmas 4.9 and 4.15 together with  $(4.4)$  and  $(4.5)$ , it is straightforward to see that  $\alpha(\text{Tr}([P_{p_i}Q_{q_i}, P_{p_i+1}])$  and  $\alpha(\text{Tr}([P_{p_i}Q_{q_i+1}, Q_{q_i}]))$ are Cauchy sequences. Then they converges as  $i \to \infty$ , and so by Proposition 3.7 and the Jordan decomposition mentioned in the proof of Lemma 4.9, as  $i \to \infty$ , Tr( $[P_{p_i}Q_{q_i+1}, Q_{q_i}]$ ) and Tr( $[P_{p_i}Q_{q_i+1}, Q_{q_i}]$ ) converge weakly to some elements of  $\overline{J}$  which are independent of the choice of  $\alpha$ . Thus by Lemmas 4.9, 4.12 and 4.16, we get

$$
\alpha(b_k(\tilde{d}_1)) \ge \alpha(b_k(\tilde{d}_2)) + \alpha(\delta)
$$

for some  $\delta \in \bar{\mathcal{I}}$  which is independent of the choice of  $\alpha$ . Thus since the substitution of an element of  $G$  is a positive bounded linear function on  $\ell^{\infty}(G)$ , we obtain

$$
b_k(\tilde{d}_1) \ge b_k(\tilde{d}_2) \mod \bar{\mathfrak{I}}.
$$

By symmetry, we have the reverse inequality, and therefore the proof is finished.  $\Box$ 

### 5. Morse inequality

In this section, we prove Theorem 1.3 and its corollaries by collecting the results obtained in the previous sections. Then we pose some questions on the boundedness of a Morse function that we anticipate will be fruitful.

$$
\Box
$$

5.1. Strongly uniform Morse function. The key technical ingredient of Morse theory is the Morse lemma which states that near a critical point, a Morse function is identified with a nondegenerate quadratic function by choosing an appropriate coordinate. Such a coordinate is not generally a normal coordinate, which is inconvenient for our purpose. Then, instead of Morse lemma, we consider replacement of a given Morse function by a nice Morse function with the same critical points. We show that this is possible for a strongly uniform Morse function, and so we only need to prove Theorem 1.3 for such a nice Morse function.

**Definition 5.1.** A bounded Morse function  $f: M \to \mathbb{R}$  is called *strongly* uniform if it is uniform with respect to  $\epsilon > 0$  and satisfies

$$
\inf_{\lambda} |\lambda| = \delta_1 \quad \text{and} \quad \inf_{x \in M - N(\text{Crit}(f); \epsilon_0)} |\nabla f(x)| \ge \delta_2
$$

for some  $\delta_1, \delta_2 > 0$ , where  $\lambda$  ranges over all eigenvalues of the Hessian  $\nabla^2 f(p)$ for all  $p \in \text{Crit}(f)$  and

$$
\epsilon_0 = \frac{\delta_2 + 8(\sup |\nabla^3 f|)\epsilon^2}{\delta_1}.
$$

Since  $\delta_2$  and  $\epsilon$  can be arbitrarily small, we may assume that

$$
\epsilon_0 < \frac{\epsilon}{2}.
$$

Let g denote the fixed G-invariant metric on  $M$ . We introduce a new  $G$ invariant metric on  $M$ . Let  $L$  denote the fixed  $G$ -invariant triangulation of  $M$ , which is used to construct a fundamental domain  $K$ . Take any small  $\epsilon > 0$ . We define a G-invariant metric  $q(\epsilon)$  on M such that  $q(\epsilon) = q$  on  $N(L_{n-1};\frac{\epsilon}{2})$  $\frac{\epsilon}{2}$  and  $g(\epsilon)$  is flat on the complement of  $N(L_{n-1}; \epsilon)$ , where  $L_{n-1}$ denotes the  $(n-1)$ -skeleton of L. Indeed, such a G-invariant metric  $g(\epsilon)$ exists. Let  $f: M \to \mathbb{R}$  be a uniform Morse function with respect to  $\epsilon > 0$ . Then for any critical point p of f,  $N(p; \epsilon)$  is in the complement of  $N(L_{n-1}; \epsilon)$ , and so the metric  $g(\epsilon)$  is flat on  $N(p;\epsilon)$ . We remark that all results in the previous sections hold for any G-invariant metric on M, hence for  $q(\epsilon)$ . As mentioned above, boundedness of a function is independent of the choice of a G-invariant metric on  $M$ . We will use the fixed metric  $q$  for considering upper and lower bounds for a Morse function.

Let us deform a given strongly uniform Morse function on  $M$  to the one which is quadratic near each critical point. Let  $f: M \to \mathbb{R}$  be a uniform Morse function with respect to  $\epsilon > 0$ , and consider the metric  $g(\epsilon)$ . Take any  $p \in \text{Crit}(f)$ . Then there is a flat normal coordinate  $(x_1, \ldots, x_n)$  around p such that  $(0, \ldots, 0)$  corresponds to p. By rotating the flat normal coordinate if necessary, we have the Taylor expansion of  $f$  at  $p$  as

(5.2) 
$$
f(x) = f(p) + \frac{1}{2}(\lambda_1 x_1^2 + \dots + \lambda_n x_n^2) + r_p(x)
$$

where  $\lambda_1, \ldots, \lambda_n$  are eigenvalues of the Hessian  $\nabla^2 f(p)$  and  $r_p(x) = O(d(x, p)^3)$ as  $x \to p$ .

**Lemma 5.2.** If  $f: M \to \mathbb{R}$  is a strongly uniform bounded Morse function with respect to  $\epsilon > 0$ , then for any  $p \in \text{Crit}(f)$  and  $x \in N(p; \epsilon_0) - N(p; \frac{\epsilon_0}{2})$  $\frac{\epsilon_0}{2}),$ 

the Taylor expansion  $(5.2)$  satisfies

$$
((\lambda_1 x_1)^2 + \dots + (\lambda_n x_n)^2)^{\frac{1}{2}} - |\nabla r_p(x)| - \frac{3}{\epsilon} |r_p(x)| > \delta_2
$$

where  $\epsilon_0$  and  $\delta_2$  are as in Definition 5.1.

*Proof.* Let  $C = \sup |\nabla^3 f|$ , and take any  $x \in N(p; \epsilon)$ . Then by the mean value theorem, we have

$$
|\nabla^2 r_p(x)| \le |\nabla^3 r_p(y)|\epsilon
$$

for some  $y \in N(p; \epsilon)$  because  $\nabla r_p(p) = \nabla^2 r_p(p) = 0$ . Hence  $|\nabla^2 r_p(x)| \leq C\epsilon$ as  $\nabla^3 r_p(x) = \nabla^3 f(x)$  on  $N(p; \epsilon)$ . Quite similarly, we can get  $|\nabla r_p(x)| \leq C\epsilon^2$ and  $|r_p(x)| \leq C\epsilon^3$ . Thus by (5.1)

$$
((\lambda_1x_1)^2+\cdots+(\lambda_nx_n)^2)^{\frac{1}{2}}-|\nabla r_p(x)|-\frac{3}{\epsilon}|r_p(x)|\geq \delta_1|x|-4C\epsilon^2>\delta_2.
$$

Therefore the statement is proved.

**Proposition 5.3.** Let  $f: M \to \mathbb{R}$  be a strongly uniform bounded Morse function. Then there is a strongly uniform bounded Morse function  $\hat{f} : M \rightarrow$ R satisfying that

(1)  $\operatorname{Crit}_k(\tilde{f}) = \operatorname{Crit}_k(f)$  for  $k = 0, 1, ..., n$ , and

(2) for any 
$$
p \in \text{Crit}(f)
$$
,

$$
\tilde{f}(x) = \tilde{f}(p) + \frac{1}{2}(\lambda_1 x_1^2 + \dots + \lambda_n x_n^2)
$$

on  $N(p; \frac{\epsilon_0}{2})$  $\frac{z_0}{2}$ , where  $\lambda_1, \ldots, \lambda_n$  and  $(x_1, \ldots, x_n)$  are as in  $(5.2)$ .

*Proof.* Let  $\rho: \mathbb{R} \to \mathbb{R}$  be an increasing smooth function such that

$$
\rho(t) = \begin{cases} 0 & t \le \frac{\epsilon_0}{2} \\ 1 & t \ge \epsilon_0. \end{cases} \quad \text{and} \quad 0 \le \frac{d\rho}{dt} \le \frac{3}{\epsilon_0}.
$$

Now we define

$$
\tilde{f}(x) = f(p) + \frac{1}{2}(\lambda_1 x_1^2 + \dots + \lambda_n x_n^2) + \rho(d(x, p))r_p(x)
$$

for  $p \in \text{Crit}(f)$  and  $x \in N(p; \epsilon_0)$ , and  $\tilde{f} = f$  on  $M - N(\text{Crit}(f); \epsilon_0)$ . Clearly,  $\mathrm{Crit}_k(\tilde{f}) = \mathrm{Crit}_k(f)$  for  $k = 0, 1, \ldots, n$ . By Lemma 5.2,  $|\nabla \tilde{f}| \geq \delta_2$ on  $N(\text{Crit}(f); \epsilon_0) - N(\text{Crit}(f); \frac{\epsilon_0}{2})$ , and it is straightforward to check that  $|\nabla \tilde{f}| \geq \delta_2$  on  $M - N(\text{Crit}(f); \epsilon_0)$ . Thus the statement follows.

5.2. **Proof of Theorem 1.3.** Let  $f: M \to \mathbb{R}$  be a bounded function, and for  $t > 0$ , let  $d_t = e^{-tf}de^{tf}$  for the differential  $d: C_c^{\infty}(\Lambda) \to C_c^{\infty}(\Lambda)$ . Then as in Example 4.2,  $d_t$  is an abstract differential. Let  $D_t = d_t + d_t^*$ .

**Proposition 5.4.** Let  $\phi: \mathbb{R} \to \mathbb{R}$  be a function such that  $\phi(x^2) \in \mathcal{G}$ . and let  $\psi(x) = x\phi(x)^2$ . Then for  $t > 0$ , we have

$$
\sum_{i=0}^k (-1)^{k-i} \operatorname{Tr}(\psi(D_t^2|_{\Lambda^i})) \ge 0 \mod \widehat{\mathfrak{I}}.
$$

for  $k = 0, 1, ..., n - 1$  and

$$
\sum_{i=0}^{n} (-1)^{n-i} \operatorname{Tr}(\psi(D_t^2|\Lambda_i)) \equiv 0 \mod \widehat{\mathfrak{I}}.
$$

$$
\qquad \qquad \Box
$$

Proof. Quite similarly to the proof of Proposition 3.2, we can show that the operator  $d_t \phi(D_t^2|_{\Lambda^i})$  is piecewise Hilbert-Schmidt, so by Proposition 3.2,  $d_t^* \phi(D_t^2|_{\Lambda^i}) = (d_t \phi(D_t^2|_{\Lambda^i}))^*$  is piecewise Hilbert-Schmidt too. Then by Proposition 3.7, we have

$$
\begin{aligned} \text{Tr}(d_t d_t^* \phi(D_t^2 |_{\Lambda^i})^2) &= \text{Tr}(d_t \phi(D_t^2 |_{\Lambda^{i-1}}) d_t^* \phi(D_t^2 |_{\Lambda^i})) \\ &\equiv \text{Tr}(d_t^* \phi(D_t^2 |_{\Lambda^i}) d_t \phi(D_t^2 |_{\Lambda^{i-1}})) \mod \widehat{\mathcal{I}} \\ &= \text{Tr}(d_t^* d_t \phi(D_t^2 |_{\Lambda^{i-1}})^2). \end{aligned}
$$

Hence we get

$$
\sum_{i=0}^{k} (-1)^{k-i} \text{Tr}(\psi(D_t^2|\Lambda_i))
$$
\n
$$
= \sum_{i=0}^{k} (-1)^{k-i} (\text{Tr}(d_t d_t^* \phi(D_t^2|\Lambda_i)^2) + \text{Tr}(d_t^* d_t \phi(D_t^2|\Lambda_i)^2))
$$
\n
$$
\equiv \sum_{i=0}^{k} (-1)^{k-i} (\text{Tr}(d_t^* d_t \phi(D_t^2|\Lambda_i)^2) + \text{Tr}(d_t^* d_t \phi(D_t^2|\Lambda_i)^2)) \mod \hat{\mathcal{I}}
$$
\n
$$
= \text{Tr}(d_t^* d_t \phi(D_t^2|\Lambda_i)^2).
$$

Since  $d_t^* d_t \phi(D_t^2|_{\Lambda^k})^2 = (d\phi(D_t^2|_{\Lambda^k})) (d\phi(D_t^2|_{\Lambda^k}))^*$ , the operator  $d_t^* d_t \phi(D_t^2|_{\Lambda^k})^2$ is positive, hence by Proposition 2.10,  $\text{Tr}(d_t^* d_t \phi(D_t^2|_{\Lambda^k})^2) \geq 0$ . Clearly,  $d_t^* d_t \phi(D_t^2 |_{\Lambda^n})^2 = 0$ , and thus the proof is finished.

**Corollary 5.5.** Let  $\phi$  and  $\psi$  be as in Proposition 5.4. If  $\psi(x^2) \geq e^{-x^2}$ , then for any  $t > 0$ , we have

$$
\sum_{i=0}^{k} (-1)^{k-i} \operatorname{Tr}(\psi(D_t^2 |_{\Lambda^i})) \ge \sum_{i=0}^{k} (-1)^{k-i} b_i^{(2)} \mod \overline{\mathcal{I}}
$$

 $for k = 0, 1, ..., n - 1, and$ 

$$
\sum_{i=0}^{n} (-1)^{i} \operatorname{Tr}(\psi(D_t^2|_{\Lambda^i})) \approx (-1)^n \chi(M/G) \mod \overline{\mathfrak{I}}.
$$

*Proof.* By Proposition 5.4, for any  $s > 2$ , we have

$$
\sum_{i=0}^{k} (-1)^{k-i} \operatorname{Tr}(\psi(D_t^2|_{\Lambda^i})) \ge \sum_{i=0}^{k} (-1)^{k-i} \operatorname{Tr}(e^{-sD_t^2|_{\Lambda^i}}) \mod \widehat{\mathcal{I}}.
$$

Then by Lemma 4.9, we take  $s \to \infty$  to get

$$
\sum_{i=0}^{k} (-1)^{k-i} \operatorname{Tr}(\psi(D_t^2|_{\Lambda^i})) \ge \sum_{i=0}^{k} (-1)^{k-i} b_i(d_t) \mod \bar{\mathcal{I}}.
$$

Now we can apply Theorem 4.10 to  $F = e^{-tf}$ ,  $\tilde{d}_1 = d$ ,  $\tilde{d}_2 = d_t$ , so that  $b_i(d_t) \approx b_i(d) = b_i^{(2)} \mod \overline{J}$ . Thus the inequalities in the statement follow. Quite similarly, we can get

$$
\sum_{i=0}^{n} (-1)^{i} \operatorname{Tr}(\psi(D_{t}^{2} |_{\Lambda^{i}})) \approx \sum_{i=0}^{n} (-1)^{n-i} b_{i}^{(2)} \mod \bar{\mathcal{I}}.
$$

Thus the proof is finished by the fact that  $\sum_{i=0}^{n}(-1)^{i}b_{i}^{(2)} = \chi(M/G)$ .  $\Box$ 

Now we suppose that  $f: M \to \mathbb{R}$  is a strongly uniform bounded Morse function, and let  $\tilde{f}$ :  $M \to \mathbb{R}$  be as in Proposition 5.3. We set  $d_t = e^{-t\tilde{f}}de^{t\tilde{f}}$ and  $D_t = d_t + d_t^*$ .

**Lemma 5.6.** Let  $\phi: \mathbb{R} \to \mathbb{R}$  be a smooth function such that  $\widehat{\phi}$  is supported within  $(-\frac{\epsilon_0}{4})$  $\frac{\epsilon_0}{4}, \frac{\epsilon_0}{4}$  $\frac{\varepsilon_0}{4}$ . Then on the complement of  $N(\mathrm{Crit}(\tilde{f}); \frac{\varepsilon_0}{2}), \phi(D_t^2)$  converges uniformly to zero as  $t \to \infty$ .

*Proof.* The proof is a modification of that for  $[21, \text{Lemma } 14.6]$ . It is easy to see that  $|\nabla \tilde{f}| \geq \delta$  for some  $\delta > 0$  on the complement of  $N(\mathrm{Crit}(\tilde{f}); \frac{\epsilon_0}{2})$ . Let  $u$  be any differential form compactly supported within the complement of  $N(\text{Crit}(\tilde{f}); \frac{\epsilon_0}{2})$ . Then as in the proof of [21, Lemma 14.6], there is an inequality

(5.3) 
$$
\|\phi(D_t^2)u\| \le q(t)\|u\|
$$

where  $q(t)$  decays rapidly as  $t \to \infty$  and is independent of the choice of u. As mentioned in the proof of  $[18, (2.8)$  Proposition, the Sobolev inequality

$$
||u||_{L^{\infty}}^{2} \leq C_{1} \sum_{i=0}^{\lceil \frac{n}{2} \rceil} ||D^{i}u||^{2}
$$

holds, where  $C_1 > 0$  is independent of the choice of u. Then by Lemma 3.8, the Sobolev inequality

$$
\|u\|_{L^\infty}^2 \leq C_2(t) \sum_{i=0}^{\lceil \frac{n}{2} \rceil} \|D_t^i u\|^2
$$

also holds, where  $C_2(t) > 0$  is independent of the choice of u (but possibly dependent on t). On the other hand, since  $D_t^2 = D^2 + t(\nabla^2 \tilde{f}) + t^2 |\nabla \tilde{f}|^2$  and we are assuming that  $\nabla \tilde{f}$  and  $\nabla^2 \tilde{f}$  are bounded, we have

$$
||(1+D_t^2)^{-1}u||_{k+2} \le p_k(t)||u||_k
$$

for some polynomial  $p_k(t)$  which is independent of the choice of u, where  $\|\cdot\|_k$  denotes the k-th Sobolev norm with respect to D. Then we can argue as in  $[21, \text{Proof of Lemma } 14.6]$  to improve  $(5.3)$  as

$$
\|\phi(D_t^2)u\|_{L^{\infty}} \le p(t)\|u\|_{L^1}
$$

where  $p(t)$  decays rapidly as  $t \to \infty$  and is independent of the choice of u. Thus the statement follows.  $\hfill \square$ 

Let  $\tilde{\rho} \colon \mathbb{R} \to \mathbb{R}$  be a smooth function supported within  $\left[-\frac{\epsilon_0}{2}\right]$  $\frac{\epsilon_0}{2}, \frac{\epsilon_0}{2}$  $\frac{[0]}{2}$  such that  $\tilde{\rho}(0) = 1$ . We define a smooth function  $\rho \colon M \to \mathbb{R}$  by

$$
\rho(x) = \sum_{p \in \text{Crit}(f)} \tilde{\rho}(d(x, p)).
$$

**Lemma 5.7.** Let  $\phi$  be as in Proposition 5.6. Then  $\text{Tr}(\rho \phi(D_t^2|_{\Lambda_k}))$  converges uniformly to  $c_k$  as  $t \to \infty$ .

*Proof.* Take any  $p \in \mathrm{Crit}_k(\tilde{f})$ . Then  $p \in g(\mathrm{Int}(K))$  for some  $g \in G$ , so g is unique. Let  $\rho_p(x) = \tilde{\rho}(d(x, p))$ . Recall that we are working with the metric  $g(\epsilon)$  on M, which is flat on  $N(p;\epsilon)$  for each  $p \in \mathrm{Crit}(\tilde{f})$ . Let  $\lambda_1, \ldots, \lambda_n$  be the eigenvalue of the Hessian  $\nabla^2 \tilde{f}(p)$ , and let

$$
\lambda(a,b) = \sum_{i=0}^{n} (|\lambda_i|(1+2a_i) + \lambda_i b_i)
$$

for  $a = \{a_1, \ldots, a_n\}$  with nonnegative integers  $a_i$  and  $b = \{b_1, \ldots, b_n\}$  with  $b_i = \pm 1$  such that exactly k of  $b_i$  are  $+1$ . Then as in the proof of [21, Lemma 14.11],  $\phi(t\lambda(a,b)^{\frac{1}{2}})$  is a spectrum of  $\phi(D_t^2|_{\Lambda^k})$  such that

$$
\text{Tr}(\rho_p \phi(D_t^2|_{\Lambda^k}))(g) = \sum_{a,b} \phi(t\lambda(a,b)^{\frac{1}{2}}) \langle \rho_p e(a,b), e(a,b) \rangle
$$

where  $e(a, b)$  is a normalized eigenvector corresponding to  $\lambda(a, b)$ . Thus since  $\phi$  is rapidly decaying, we get

$$
\lim_{t \to \infty} \sum_{a,b} \phi(t\lambda(a_i, b_i)^{\frac{1}{2}}) = \begin{cases} 1 & p \text{ has index } k \\ 0 & \text{otherwise} \end{cases}
$$

where only  $\lambda(a, b) = 0$  contributes to the first equality. As in the proof of [21, Lemma 14.11], if  $\lambda(a, b) = 0$ , then

$$
e(a,b)=(\pi^{-\frac{n}{4}}t^{\frac{n}{2}}(\lambda_1\cdots\lambda_n)^{\frac{n}{2}})e^{-t\frac{1}{2}(\lambda_1x_1^2+\cdots+\lambda_nx_n^2)}dx_1\cdots dx_n
$$

hence  $\langle \rho_p e(a, b), e(a, b) \rangle \rightarrow \text{as } t \rightarrow 0$ . On the other hand, we have

$$
\text{Tr}(\rho_p \varphi(D_t^2|_{\Lambda^k}))(h) = 0
$$

for  $h \neq g$ . Thus  $\text{Tr}(\rho_p \phi(D_t^2|_{\Lambda_k}))$  converges uniformly to a function  $c_k^p$  $k^p: G \to$ R given by

$$
c_k^p(h) = \begin{cases} 1 & p \text{ has index } k \text{ and } h = g \\ 0 & \text{otherwise.} \end{cases}
$$

Since the Morse function  $f$  is strongly uniform, we have

$$
\inf_{p \in \text{Crit}(f)} |\lambda| = \delta_1 > 0
$$

where  $\lambda$  ranges over all eigenvalues of the Hessian  $\nabla^2 \tilde{f}(p)$ . Then the uniformness of the above convergence is also uniform as p ranges over all  $p \in \text{Crit}(f)$ . Note that  $\sum_{p \in \text{Crit}(\tilde{f})} c_k^p = c_k$  and

$$
\text{Tr}(\rho\varphi(D_t^2|_{\Lambda^k})) = \sum_{p \in \text{Crit}(f)} \text{Tr}(\rho_p\varphi(D_t^2|_{\Lambda^k}))
$$

where  $\text{supp}(\rho_p) \cap \text{supp}(\rho_q) = \emptyset$  for  $p \neq q \in \text{Crit}(f)$ . Thus the statement follows.  $\square$ 

Now we are ready to prove Theorem 1.3

*Proof of Theorem 1.3.* Combine Corollary 5.5 and Lemmas 5.6 and 5.7.  $\Box$ 

We prove Corollaries 1.4 and 1.5.

*Proof of Corollary 1.4.* As in the proof of Proposition 2.12, we have  $\mu(\mathcal{I}) =$ 0. Then the statement follows from Theorem 1.3 and the positivity of  $\mu$ .  $\Box$ 

To prove Corollary 1.5, we need the following proposition [12, Proposition 2.3].

**Proposition 5.8.** Let G be a finitely generated infinite group. If  $\phi \in \ell^{\infty}(G)$ satisfies  $\phi(q) = 0$  for all but finitely many  $q \in G$ , then  $\phi \equiv 0 \mod 1$ .

Proof of Corollary 1.5. We can easily deduce from Theorem 1.3 the weak Morse inequality

$$
c_k \ge b_k^{(2)} \mathbb{1} \mod \bar{\mathfrak{I}}.
$$

Then we get  $\mu(c_k) \geq b_k^{(2)}$ <sup>(2)</sup> for any *G*-invariant mean  $\mu$ :  $\ell^{\infty}(G) \to \mathbb{R}$ . Hence by Proposition 5.8,  $c_k(g) \neq 0$  for infinitely many  $g \in G$ .

Novikov and Shubin [16] proved Morse inequalities for a Morse function on  $M/G$  and the  $L^2$ -Betti numbers of M. We recover this result when G is amenable. For a Morse function  $\bar{f}: M/G \to \mathbb{R}$ , let  $\bar{c}_k$  denote the number of critical points of  $f$  having index  $k$ .

**Corollary 5.9.** If G is amenable and  $\bar{f}$ :  $M/G \rightarrow \mathbb{R}$  is a Morse function, then we have

$$
\bar{c}_k - \bar{c}_{k-1} + \dots + (-1)^k \bar{c}_0 \ge b_k^{(2)} - b_{k-1}^{(2)} + \dots + (-1)^k b_0^{(2)}
$$

for  $k = 0, 1, ..., n - 1$ , and

$$
\bar{c}_n - \bar{c}_{n-1} + \cdots + (-1)^n \bar{c}_0 = (-1)^n \chi(M/G).
$$

*Proof.* Let  $f: M \to \mathbb{R}$  be the lift of  $\overline{f}$ . Then as  $M/G$  is compact, f is a bounded Morse function. Moreover, by deforming a fundamental domain K slightly if necessary, we may assume that  $f$  is strongly uniform. Since  $f$ is G-invariant, we have  $c_k = \bar{c}_k \mathbb{1}$  for  $k = 1, 2, \ldots, n$ . Then the statement follows from Corollary 1.4.

5.3. Questions on boundedness. As mentioned in Section 1, the boundedness of the derivative  $\nabla f$  of a Morse function  $f: M \to \mathbb{R}$  is an essential assumption, and the boundedness of  $\nabla^i f$  for  $i \geq 2$  and f itself are assumed by technical reasons. Here, we consider the boundedness of  $f$  itself. It is used to apply Theorem 4.10 to the  $F = e^{-tf}$  case. However, the normalized Betti number  $b_k(d_t)$  for the Witten defrmation  $d_t = e^{-tf}de^{tf}$  is defined in terms of the derivatives of  $f$ , so its properties may not depend on the boundedness of f. Then we ask:

#### Question 5.10. Does Theorem 1.3 hold for an unbounded Morse function?

Corollary 1.5 has its own interest, and so we also ask:

## Question 5.11. Does Corollary 1.5 hold for an unbounded Morse functioin?

Observe that Corollary 1.5 makes sense for any infinite group  $G$ , though Theorem 1.3 makes sense only for an amenable group  $G$  as in Proposition 2.12. Then we can further ask whether or not Corollary 1.5 holds for an unbounded Morse function and a nonamenable group. However, we can get a negative answer to this further question as follows. Suppose that  $M/G$ has nonpositive sectional curvature, and that M is simply-connected. In this case, the group  $G$  is nonamenable. Fix a basepoint  $p$  of  $M$ . Then by the Cartan-Hadamard theorem, for each point  $x \in M$ , there is a unique

geodesic  $\gamma_x: [0, t_x] \to M$  from p to x, where  $t_x = d(p, x)$ . Observe that for any small  $\epsilon > 0$ ,  $\left|\frac{d\gamma_x}{dt}\right| = 1$  on  $[\epsilon, t_x]$ . Then a function  $f: M \to \mathbb{R}$  which is a small perturbation of a function

$$
M \to \mathbb{R}, \quad x \mapsto d(p, x)
$$

has finitely many nondegenerate critical points near  $p$ . Namely,  $f$  is a Morse function on  $M$  with finitely many critical points. Thus if  $G$  is infinite, e.g.  $M/G$  is a surface of genus  $\geq 2$ , this example is a negative answer to the above further question.

#### **REFERENCES**

- [1] M. Atiyah, Elliptic operators, discrete groups and von Neumann algebras, Astérisque 32 (1976), 43-72. 1
- [2] J. Block and S. Weinberger, Aperiodic tilings, positive scalar curvature, and amenability of spaces, J. Amer. Math. Soc. 5 (1992), no. 4, 907-918. 3, 8
- [3] J. Brodzki, G.A. Niblo, and N. Wright, Pairings, duality, amenability and bounded cohomology, J. Eur. Math. Soc. 14 (2012), 1513-1518. 3, 8
- [4] X. Dai and J. Yan, Witten deformation for noncompact manifolds with bounded geometry, J. Inst. Math. Jussieu 22 (2023), no. 2, 643-680. 1
- [5] N. Dunford and J.T. Schwartz, Linear Operators, Part I: General Theory, Wiley, New York, 1958. 17
- [6] M. Gromov, Asymptotic invariants of infinite groups, in Geometric group theory, vol. 2 (Sussex, 1991), 1-295, London Math. Soc. Lecture Note Ser. 182, Cambridge Univ. Press, Cambridge. 2
- [7] M. Gromov and H.B. Lawson, Jr., Positive scalar curvature and the Dirac operator on complete Riemannian manifolds, Publ. Math. I.H.E.S. 58 (1983), 83-196. 13
- [8] M.W. Hirsch, On imbedding differentiable manifolds in euclidean space, Ann. of Math. (2) 73 (1961), 566-571. 2
- [9] T. Kato, D. Kishimoto, and M. Tsutaya, Homotopy type of the space of finite propagation unitary operators on Z, Homology Homotopy Appl. 25 (2023), no. 1, 375-400. 3
- [10] T. Kato, D. Kishimoto, and M. Tsutaya, Homotopy type of the unitary group of the uniform Roe algebra on  $\mathbb{Z}^n$ , J. Topol. Anal. 15 (2023), no. 2, 495-512. 3
- [11] T. Kato, D. Kishimoto, and M. Tsutaya, Hilbert bundles with ends, J. Topol. Anal. 16 (2024), no. 2, 291-322. 3
- [12] T. Kato, D. Kishimoto, and M. Tsutaya, Vector fields on non-compact manifolds, accepted by Algebr. Geom. Topol. 2, 3, 4, 7, 26
- [13] W. Lück,  $L^2$ -invariants: Theory and Applications to Geometry and K-theory, Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge Bd. 44, Springer-Verlag, Berlin, 2002. 1
- [14] F. Milizia, Bounded differential forms and coinvariants of bounded functions, arVix:2311.07731. 7
- [15] J. Milnor, A note on curvature and fundamental group, J. Differential Geometry 2 (1968) 1-7. 9
- [16] S.P. Novikov and M.A. Shubin, Morse inequalities and von Neumann II1-factors, Doklady Akad. Nauk SSSR 289 (1986), 289-292. 1, 3, 26
- [17] P. Pansu, Cohomologie  $L^p$ : invariance sous quasiisométries, preprint, 1995. 2
- [18] J. Roe, An index theorem on open manifolds. I, J. Differential Geom. 27 (1988), no. 1, 87-113. 1, 4, 8, 12, 24
- [19] J. Roe, An index theorem on open manifolds. II, J. Differential Geom. 27 (1988), no. 1, 115-136. 1, 4, 15
- [20] J. Roe, On the quasi-isometry invariance of  $L^2$  Betti numnbers, Duke Math. J. 59 (1989), no. 3, 765-783. 4, 7, 8, 15, 17, 18
- [21] J. Roe, Elliptic operators, Topology and Asymptotic Methods, Pitman Res. Notes Math. Ser. 179, Longman Scientific & Technical, Harlow; copublished in the United States with John Wiley & Sons, Inc., New York, 1988. 13, 24, 25
- [22] W. Rudin, Real and Complex Analysis, McGraw-Hill, 1966.

17

Department of Mathematics, Kyoto University, Kyoto 606-8502, Japan Email address: tkato@math.kyoto-u.ac.jp

Faculty of Mathematics, Kyushu University, Fukuoka 819-0395, Japan  $\emph{Email address: }$ kishimoto@math.kyushu-u.ac.jp

Faculty of Mathematics, Kyushu University, Fukuoka 819-0395, Japan Email address: tsutaya@math.kyushu-u.ac.jp