Conjugate gradient methods for high-dimensional GLMMs

Andrea Pandolfi^{*}, Omiros Papaspiliopoulos[†], and Giacomo Zanella[‡]

November 8, 2024

Generalized linear mixed models (GLMMs) are a widely used tool in statistical analysis. The main bottleneck of many computational approaches lies in the inversion of the high dimensional precision matrices associated with the random effects. Such matrices are typically sparse; however, the sparsity pattern resembles a multi partite random graph, which does not lend itself well to default sparse linear algebra techniques. Notably, we show that, for typical GLMMs, the Cholesky factor is dense even when the original precision is sparse. We thus turn to approximate iterative techniques, in particular to the conjugate gradient (CG) method. We combine a detailed analysis of the spectrum of said precision matrices with results from random graph theory to show that CG-based methods applied to high-dimensional GLMMs typically achieve a fixed approximation error with a total cost that scales linearly with the number of parameters and observations. Numerical illustrations with both real and simulated data confirm the theoretical findings, while at the same time illustrating situations, such as nested structures, where CG-based methods struggle.

Keywords: Bayesian computation; High-dimensional Gaussians; Conjugate gradient samplers; Cholesky factorization; Random graphs.

^{*}Bocconi University, Department of Decision Sciences

[†]Bocconi University, Department of Decision Sciences and BIDSA

[‡]Bocconi University, Department of Decision Sciences and BIDSA

GZ acknowledges support from the European Research Council (ERC), through StG "PrSc-HDBayLe" grant ID 101076564.

1. Introduction

Generalized linear mixed models (GLMMs) are a foundational statistical tool, widely used across multiple disciplines (Gelman and Hill, 2007; Wood, 2017). GLMMs extend the framework of Generalized Linear Models by incorporating both fixed effects and random effects. Fixed effects capture population-level trends and relationships, while random effects account for individual deviations from these trends. GLMMs utilize a link function to establish a relationship between the mean of the response variable and the linear predictor, which is given by a linear combination of both fixed and random effects (see Section 3).

In various settings, the factors associated with the random effects may include numerous categories, known as *levels*, resulting in models with a large number of parameters p and a large number of observations N, both potentially in the order of several thousands. This scenario often arises in contemporary applications. For instance, within the political sciences, one may consider a geographic factor with numerous units, or one may include factors encoding so-called deep interactions (Ghitza and Gelman, 2013). Another context where such models find utility is in recommendation systems, where categorical variables represent customers and products (Perry, 2016; Gao and Owen, 2017).

Our work is relevant for various computational algorithms used to fit GLMMs, such as Gibbs sampling, variational inferences, algorithms for restricted maximum likelihood estimation, and Laplace approximations (see Section 8 for a discussion). The primary computational bottleneck of these methodologies lies in the factorization of a sparse high-dimensional matrix, denoted by Q. This matrix decomposes as

$$\boldsymbol{Q} = \boldsymbol{T} + \boldsymbol{V}^T \boldsymbol{\Omega} \boldsymbol{V},\tag{1}$$

where T is the prior component (or, equivalently, a regularization term) and is usually diagonal or easily factorizable; Ω is a diagonal matrix and adjusts for the variances of the response variables; and $V \in \mathbb{R}^{N \times p}$ denotes the design matrix. In particular, we mostly focus on performing Bayesian inferences using so-called *blocked Gibbs samplers*, that alternate the updates of regression parameters, variance hyperparameters and potential additional latent variables (see Section 6.1). The update of the regression parameters requires sampling from a multivariate Gaussian distribution with precision matrix Q as in (1), and it is usually the most computationally intensive steps in those algorithms.

The standard procedure to sample from a multivariate normal distribution involves the computation of the Cholesky factor of Q (see Section 2.1). It is well established that the Cholesky factorization can be computed efficiently with nested factors (see e.g. Papaspiliopoulos and Zanella, 2017 and references therein). On the contrary, we will show that, for general crossed factors, under standard random design assumptions, the cost of computing the Cholesky factor scales as $\mathcal{O}(p^3)$, even when Q is sparse (see Section 4 for further details). Thus, since exact factorization of Q is too expensive, we consider an alternative strategy to sample from the desired Gaussian distribution, which involves the solution of a properly perturbed linear system $Q\theta = b$ (see Section 2.2). Here, instead of finding the exact solution of the linear system, we employ the well-known conjugate gradient (CG) algorithm to find an approximate solution or, equivalently, produce an approximate sample. We will demonstrate that, in the general case where the Cholesky factorization becomes inefficient, the CG sampler only requires a constant (not growing with N and p) number of matrix-vector multiplications Qb, which results in a total of $\mathcal{O}(\max(N, p))$ cost to produce an approximate sample.

The paper is organized as follows. In Section 2, we briefly review the algorithms for the Cholesky factorization and the CG method, with a particular focus on their time complexity. In Section 3, we describe the specific instance of GLMMs, which we refer to as *random-intercept crossed effects models*, that will serve as the reference model for the theoretical analysis in Sections 4 and 5. In Section 4, we show that exact Cholesky factorization is expensive for such models; while, in Section 5, we show that the CG algorithm converges fast for such models. The bulk of our technical contribution lies in the analysis of the spectrum of Q performed in Section 5. Finally, in Section 6, we describe the proposed methodologies for the case of general GLMMs, and illustrate them with simulated and real data. Specifically, we consider an application to a survey data for the American presidential elections of 2004 (Ghitza and Gelman, 2013) and to a data set for Instructor Evaluations by Students at ETH (from 1me4 R-library). The proofs of the presented results are given in the supplementary material and code to reproduce the experiments is available at github.com/AndreaPandolfi/ASLA.jl.

2. Review of high-dimensional Gaussian sampling

In this section, we review some well-known algorithms to sample from a high dimensional Gaussian distribution. Specifically, we consider the problem of sampling $\boldsymbol{\theta} \sim \mathcal{N}_p(\boldsymbol{Q}^{-1}\boldsymbol{m}, \boldsymbol{Q}^{-1})$. Such structure arises in several contexts, such as Bayesian regression (Nishimura and Suchard, 2023), spatial models with GMRFs (Rue et al., 2009), and GLMMs, as in this case (see Section 3). We are mostly interested in the complexity of these algorithms when p is large and \boldsymbol{Q} is sparse. We say that a matrix $\boldsymbol{Q} \in \mathbb{R}^{p \times p}$ is sparse, if the number of non-zero entries grows slower than the number of total possible entries, namely if $n_{\boldsymbol{Q}} = o(p^2)$ as $p \to +\infty$, where $n_{\boldsymbol{Q}}$ denotes the number of non-zero entries of \boldsymbol{Q} .

2.1. Cholesky factorization

The Cholesky factorization of a positive-definite matrix $\boldsymbol{Q} \in \mathbb{R}^{p \times p}$ is a factorization of the form $\boldsymbol{Q} = \boldsymbol{L}\boldsymbol{L}^T$, where $\boldsymbol{L} \in \mathbb{R}^{p \times p}$ is a lower triangular matrix with real and positive diagonal entries. Such factorization can be computed with the following column-wise recursion

$$L_{mm} = \sqrt{Q_{mm} - \sum_{\ell=1}^{m-1} L_{m\ell}^2}, \qquad L_{jm} = \frac{1}{L_{mm}} \left(Q_{jm} - \sum_{\ell=1}^{m-1} L_{m\ell} L_{j\ell} \right), \quad j > m. \quad (2)$$

Given the Cholesky factor L, one can sample from $\mathcal{N}(Q^{-1}m, Q^{-1})$ by solving linear systems in L and L^T , which can be done efficiently in $\mathcal{O}(n_L)$ time via forward and

backward substitution. A description of the algorithm can be found in the supplementary material.

For dense matrices, the exact computation of the Cholesky factor requires $\mathcal{O}(p^3)$ time, and the storage of $\mathcal{O}(p^2)$ entries. For sparse matrices with specific structures, the computational requirements can be drastically reduced. If Q can be turned into a banded matrix via row and column permutations, the cost of computing L can be reduced to $\mathcal{O}(pb^2)$, where b is the bandwidth. When considering spatial Gaussian Markov random fields (GMRFs) or nested hierarchical models, such reordering can be done efficiently, which makes sampling via Cholesky factor efficient in such cases (see Papaspiliopoulos and Zanella, 2017, and extensive references therein). On the other hand, there is little work in the literature that studies the computational cost of the Cholesky factorization for general GLMMs with non-nested designs.

2.1.1. Conditional Independence Graphs and Cholesky complexity analysis

The conditional independence structure of a given Gaussian vector $\boldsymbol{\theta} \sim \mathcal{N}_p(\boldsymbol{Q}^{-1}\boldsymbol{m}, \boldsymbol{Q}^{-1})$ is described by \boldsymbol{Q} through the relation

$$\theta_j \perp \theta_m \mid \boldsymbol{\theta}_{-jm} \Longleftrightarrow Q_{jm} = 0, \qquad (3)$$

where θ_{-jm} denotes θ after the removal of the entries j and m. Hence, the *conditional* independence (CI) graph of θ is entirely described by $Supp(\mathbf{Q})$, the support of \mathbf{Q} (Rue and Held, 2005, Ch. 2.1.5). We denote the CI graph of θ by $G_{\mathbf{Q}}$, whose vertices are the variables $\{\theta_j, j = 1, \ldots, p\}$ and the edges are those (θ_j, θ_m) s.t. $Q_{jm} \neq 0$ for $j \neq m$.

The Cholesky factor \boldsymbol{L} of \boldsymbol{Q} has also a probabilistic interpretation: for m < j, $L_{jm} = 0$ if and only if θ_j and θ_m are independent given the *future set* of θ_m excluding θ_j , i.e. $\theta_m \perp \theta_j \mid \boldsymbol{\theta}_{\{(m+1):p\}\setminus j}$ (Rue and Held, 2005, Theorem 2.8). Therefore, even when \boldsymbol{Q} is sparse, \boldsymbol{L} might not be and the order we assign to elements of $\boldsymbol{\theta}$ influences the sparsity of \boldsymbol{L} . The possibly non-zero entries of \boldsymbol{L} can be deduced from $G_{\boldsymbol{Q}}$ as follows: a sufficient condition to ensure $L_{jm} = 0$ for m < j is that the future set of θ_m separates it from θ_j in $G_{\boldsymbol{Q}}$. This motivates defining the number of possible non-zero entries in \boldsymbol{L} as $n_{\boldsymbol{L}} = \sum_{m=1}^{p} n_{\boldsymbol{L},m}$ where

$$n_{L,m} = |\{j \ge m : \text{the future set of } \theta_m \text{ does not separate it from } \theta_j \text{ in } G_Q\}|.$$
 (4)

Thus, the Cholesky factor L involves $n_L - n_Q \ge 0$ additional potential non-zero entries compared to the original matrix Q. Such additional non-zeros terms are commonly referred to as *fill-ins*. Since n_L depends on the ordering of the variables in θ , standard algorithms for Cholesky factorizations of sparse matrices proceed in two steps: first they try to find an ordering of variables that reduces n_L as much as possible, and then compute the corresponding Cholesky factor using (2). Finding the ordering that minimizes n_L is NP-hard, but various heuristic strategies to find good orderings are available (Golub and van Loan, 2013, Ch. 11).

Sparsity in L has direct consequences on the computational cost required to compute it - although it has to be appreciated that a sparse Cholesky is not necessarily computable efficiently. The following theorem quantifies this connection.

Theorem 1. Denoting with Cost(Chol) the number of floating point operations (flops) needed to compute the Cholesky factor L of a positive definite matrix Q, we have

$$\mathcal{O}\left(n_{\boldsymbol{L}}^{2}/p\right) \leq \operatorname{Cost}(\operatorname{Chol}) = \mathcal{O}\left(\sum_{m=1}^{p} n_{\boldsymbol{L},m}^{2}\right) \leq \mathcal{O}\left(n_{\boldsymbol{L}}^{1.5}\right).$$
(5)

The equality and lower bound in (5) are well-known, while the upper bound is more involved, and we have not been able to find it in the literature. Note that, trivially, the result also implies $\text{Cost}(\text{Chol}) \geq \mathcal{O}\left(n_Q^2/p\right)$.

When Q is a dense matrix we have $n_L = \mathcal{O}(p^2)$ and thus the lower and upper bounds in (5) coincide, being both cubic in p, and they are both tight. For sparse matrices, instead, the two bounds can differ up to a $\mathcal{O}(p^{0.5})$ multiplicative factor and each can be tight depending on the sparsity pattern. For example, if Q is a banded matrix with bandwidth b, we have $n_L = \mathcal{O}(pb)$ and $\operatorname{Cost}(\operatorname{Chol}) = \mathcal{O}(pb^2)$, hence the lower bound is tight while the upper bound is off by a $\mathcal{O}((p/b)^{0.5})$ factor. On the contrary, for a matrix Q with a dense $p^{0.5} \times p^{0.5}$ sub-matrix and diagonal elsewhere we have $n_L = \mathcal{O}(p)$ and $\operatorname{Cost}(\operatorname{Chol}) = \mathcal{O}(p^{1.5})$, meaning that the upper bound is tight while the lower bound is off by a $\mathcal{O}(p^{0.5})$ factor.

2.2. Conjugate Gradient

Conjugate Gradient (CG) is a widely-used iterative optimization algorithm employed for solving large systems of linear equations $Q\theta = b$, for which Q is positive-definite (Golub and van Loan, 2013; Saad, 2003). CG iteratively minimizes the quadratic form associated to the linear system. However, instead of optimizing along the gradient direction, it restricts to the component of the gradient which is conjugate (i.e. Q-orthogonal) to the previous search directions.

Each CG iteration only requires evaluation of matrix-vector products $Q\theta$ and scalar products of *p*-dimensional vectors. This feature makes CG methods very appealing for solving sparse linear systems, as each iteration only requires $\mathcal{O}(n_Q)$ operations. The algorithm is also optimal in terms of memory efficiency, as it only requires storing $\mathcal{O}(p)$ values: basically only the approximate solution, which gets updated in place.

Several strategies have been proposed to use CG algorithm to sample from high dimensional Gaussian distributions. In this paper, we will refer to the *perturbation optimization* sampler (Papandreou and Yuille, 2010; Nishimura and Suchard, 2023), which requires solving the following linear system

$$Q\theta = m + z, \qquad z \sim \mathcal{N}(0, Q)$$
 (6)

Simple computations show that, when the linear system in (6) is solved exactly, $\boldsymbol{\theta}$ is an exact sample from $\mathcal{N}_p(\boldsymbol{Q}^{-1}\boldsymbol{m}, \boldsymbol{Q}^{-1})$. If the linear system is solved via CG method, an approximate sample will be returned. A potential limitation for the application of this algorithm is given by the necessity to sample efficiently from $\mathcal{N}_p(\boldsymbol{0}, \boldsymbol{Q})$, which is however feasible for matrices as in (1) (see end of Section 3). Notice that in this case, CG is only

used to solve the linear system in (6), while other CG samplers (Parker and Fox, 2012; Vono et al., 2022) exploit CG algorithms, actually Lanczos tri-diagonalization (Lanczos, 1950), to build an approximate low-rank square root of Q^{-1} which is then used to sample. Notice, however, that, if Q is full-rank but has only k < p distinct eigenvalues, then (6) would return an exact sample in k-steps (see Theorem 2), while other methodologies based on low-rank approximations would not.

2.2.1. Rate of convergence

When studying the complexity of CG algorithm, one needs to quantify how fast the solution at iteration k approaches the exact solution of the linear system. The convergence behavior of CG algorithm has been extensively studied in the literature (Golub and van Loan, 2013, Section 11). We report the most well-known result in the following Theorem.

Theorem 2. Consider the linear system $Q\theta = b$, with Q positive definite. Denote the starting vector with θ^0 , then the Q-norm distance between the k-th CG iterate θ^k and the solution θ satisfies the inequality

$$\frac{||\boldsymbol{\theta}^{k} - \boldsymbol{\theta}||_{\boldsymbol{Q}}}{||\boldsymbol{\theta}^{0} - \boldsymbol{\theta}||_{\boldsymbol{Q}}} \le 2\left(\frac{\sqrt{\kappa(\boldsymbol{Q})} - 1}{\sqrt{\kappa(\boldsymbol{Q})} + 1}\right)^{k},\tag{7}$$

where $\kappa(\mathbf{Q}) = \lambda_{max}(\mathbf{Q})/\lambda_{min}(\mathbf{Q})$ denotes the condition number of \mathbf{Q} . Moreover, if \mathbf{Q} has only k < p distinct eigenvalues, CG returns the exact solution after k iterations.

See Trefethen and Bau (1997) for a proof of these results. Theorem 2 shows that CG has a fast convergence rate when either Q has a small condition number or when it has few distinct eigenvalues.

It is well-known in the CG literature that the convergence rate presented in Theorem 2 is very conservative. Indeed, CG is also fast when most of the eigenvalues, except few outlying ones, are clustered in a small interval $[\mu_s, \mu_{p-r}]$, with μ_s not close to 0. In this case, one observes that, after a small number of iterations, CG behaves as if the components corresponding to the outlying eigenvalues have been removed, and the CG convergence rate changes as if the condition number in (7) is replaced by the effective value μ_{p-r}/μ_s (Concus et al., 1976; van der Vorst, 2003). This behavior is known as superlinear convergence of conjugate gradients. For this reason, instead of considering $\kappa(\mathbf{Q})$, we will focus on the effective condition number (Van der Sluis and Van der Vorst, 1986)

$$\kappa_{s+1,p-r}(\boldsymbol{Q}) = \frac{\mu_{p-r}}{\mu_{s+1}},\tag{8}$$

obtained by removing the smallest s eigenvalues of Q and the r largest ones. In Section 5, we will provide upper bounds on $\kappa_{s+1,p-r}(Q)$ in the GLMM context, for small s and r.

3. Random-intercept crossed effects models

We now describe the model we will refer to in the theoretical analysis of Section 4 and 5. Specifically, we present the class of *random-intercept crossed effects models*.

Model 1 (Random-intercept crossed effects models). For each observation $i \in \{1, ..., N\}$, consider a univariate continuous response $y_i \in \mathbb{R}$ distributed as

$$y_i \mid \eta_i \sim \mathcal{N}\left(\eta_i, \ \tau^{-1}\right), \qquad \eta_i = \theta_0 + \sum_{k=1}^K \boldsymbol{z}_{i,k}^T \boldsymbol{\theta}_k,$$
(9)

where θ_0 is a global intercept, $\boldsymbol{\theta}_k = (\theta_{k,1}, \dots, \theta_{k,G_k})^T$ is the vector of random effects for factor k, and $\boldsymbol{z}_{i,k} \in \{0,1\}^{G_k}$ with $\sum_{g=1}^{G_k} z_{i,k,g} = 1$ is a "one-hot" vector which encodes the level of factor k for the observation i. Here K is the number of factors, and G_k the number of levels in factor K. We assume independent priors $\theta_{k,g} \stackrel{ind.}{\sim} \mathcal{N}(0, T_k^{-1})$, for each factor k and level g. The fixed effect is assigned either a normal or improper flat prior, and the precision parameters are assigned conjugate gamma distributions. Though, neither of these choices are directly relevant for our analysis.

If we define $\boldsymbol{\theta} = (\theta_0, \boldsymbol{\theta}_1^T, \dots, \boldsymbol{\theta}_K^T)^T \in \mathbb{R}^p$ (with $p = 1 + \sum_k G_k$) and $\boldsymbol{v}_i = (1, \boldsymbol{z}_{i,1}^T, \dots, \boldsymbol{z}_{i,K}^T)^T \in \mathbb{R}^p$, we can write η_i as $\boldsymbol{v}_i^T \boldsymbol{\theta}$. Under Model 1, the posterior conditional distribution of $\boldsymbol{\theta}$ is

$$\boldsymbol{\theta} \mid \boldsymbol{y}, \boldsymbol{V}, \boldsymbol{T}, \tau \sim \mathcal{N}_p(\boldsymbol{Q}^{-1}(\boldsymbol{T}\boldsymbol{m}_0 + \tau \boldsymbol{V}^T \boldsymbol{y}), \ \boldsymbol{Q}^{-1}), \qquad (10)$$

where \boldsymbol{m}_0 is the prior mean of $\boldsymbol{\theta}, \boldsymbol{T}$ its prior precision, $\boldsymbol{V} = [\boldsymbol{v}_1 | \dots | \boldsymbol{v}_N]^T \in \mathbb{R}^{N \times p}$ and

$$\boldsymbol{Q} = \boldsymbol{T} + \tau \boldsymbol{V}^T \boldsymbol{V} \,. \tag{11}$$

In Sections 4 and 5, we study the time complexity of sampling from (10) with the algorithms presented in Section 2. Regarding the CG sampler, we will only focus on the complexity of the CG step. Indeed, sampling $\boldsymbol{z} \sim \mathcal{N}_p(\boldsymbol{0}, \boldsymbol{Q})$ with \boldsymbol{Q} as in (11), can be done in $\mathcal{O}(N+p)$ time, by sampling $\boldsymbol{\zeta} \sim \mathcal{N}_p(\boldsymbol{0}, \boldsymbol{I}_p)$ and $\boldsymbol{\eta} \sim \mathcal{N}_N(\boldsymbol{0}, \boldsymbol{I}_N)$ independently, and setting $\boldsymbol{z} = \boldsymbol{T}^{1/2} \boldsymbol{\zeta} + \sqrt{\tau} \boldsymbol{V}^T \boldsymbol{\eta}$.

Random-intercept models defined as in Model 1 are a specific instance of the more general class of GLMMs considered in Section 6, which motivates our work. Throughout the paper, Model 1 will serve as a study case to develop theoretical results. We restrict our attention to the random-intercept case as it allows for an exhaustive theoretical tractability, while still preserving the crossed sparsity structure that constitute the main computational challenge for general GLMMs with non-nested designs (see e.g. Section 4). Notice also that various recent works provided methodologies and theoretical results for this class of model (Papaspiliopoulos et al., 2019; Ghosh et al., 2022). Note that, while we develop theoretical results in the context of Model 1, our methodology and numerical results apply to the more general class of GLMMs described in Section 6, which in particular includes random slopes and interaction terms.

For the sake of simplicity, the only fixed effect included in Model 1 is the intercept $\theta_0 \in \mathbb{R}$. Nonetheless, all the results of the following sections could be restated to include

multivariate fixed effects, $\boldsymbol{\theta}_0 \in \mathbb{R}^{D_0}$, leading to the same conclusions, provided that D_0 is sufficiently small relative to the dimensionality of the random effects $(\boldsymbol{\theta}_1, \ldots, \boldsymbol{\theta}_K)$ (see Section B.1 in the supplementary material for further details).

3.1. Design assumptions

Consider the precision matrix defined in (11). Since T is a diagonal matrix, the offdiagonal support of Q is entirely characterized by the likelihood term, which we will denote with $U = V^T V = \sum_{i=1}^n v_i v_i^T$. In particular, simple computations show that

$$U[\theta_0, \theta_0] = N, \qquad U[\theta_0, \theta_{k,g}] = |\{i = 1, \dots, N : z_{i,k,g} = 1\}|, \\ U[\theta_{k,g}, \theta_{k,g}] = |\{i : z_{i,k,g} = 1\}|, \qquad U[\theta_{k,g}, \theta_{k',g'}] = |\{i : z_{i,k,g} = 1, z_{i,k',g'} = 1\}|,$$
(12)

where $U[\theta_{k,g}, \theta_{k',g'}]$ refer to the entry relative to the pair $(\theta_{k,g}, \theta_{k',g'})$. We assume without loss of generality that each level of each factor is observed at least once, which is equivalent to say that $U[\theta_{k,g}, \theta_{k,g}] \ge 1$, for all factors $k = 1, \ldots, K$ and levels $g = 1 \ldots, G_k$.

In the analogy with conditional independence graphs described in Section 2.1.1, the last equality in (12) implies that $\theta_{k,g}$ is connected to $\theta_{k',g'}$ in $G_{\mathbf{Q}}$ if and only if $k \neq k'$ and there exists at least one observation where the two levels are observed together. Thus, the resulting CI graph is a (K+1)-partite graph, with one trivial block containing the vertex θ_0 , and a block of size G_k for each factor $k = 1, \ldots, K$. Since $U[\theta_0, \theta_{k,g}] = U[\theta_{k,g}, \theta_{k,g}] > 0$ for each k, g, then θ_0 is connected to all the other vertices.

Throughout our analysis, we will consider a regime where $N, p \to \infty$, while K is fixed. Notice that $N \ge p/(K+1)$, since we assume that each $\theta_{k,g}$ appears in at least one observation. Finally, we assume that $n_{\mathbf{Q}}$ or, equivalently, the number of edges is $G_{\mathbf{Q}}$ is $\mathcal{O}(N)$.

4. Cholesky factorization with sparse crossed designs

In this section, we will provide a simple theoretical example as well as numerical results with random designs showing that the Cholesky factorization is not suitable to efficiently factorize the posterior precision matrices that arise from Model 1.

Proposition 1. Under Model 1, n_L is a non-increasing function of the position of θ_0 in the ordering of θ . Hence, placing θ_0 last in the ordering always minimizes n_L .

Proposition 1 implies that we can place θ_0 last without loss of generality. This helps to reduce n_L but it does not solve the problem. The following example describes a simple sparse design that is catastrophic in this sense, where $n_Q = \mathcal{O}(p)$, but $n_L = \mathcal{O}(p^2)$.

Example 1. Consider a random intercept model with K = 2 and $G_1 = G_2 = G$. Fix an integer $d \ge 2$, and for each $g = 1, \ldots, G-1$, connect the vertex $\theta_{1,g}$ to $\theta_{2,j}$ (this is equivalent to adding an observation *i*, s.t. $g_1[i] = g$ and $g_2[i] = j$) for all *j*'s for which at least one of the following conditions hold: (a) g = j; (b) $d(g-1) \le j-2 < dg$ mod (G-1) and g < j; (c) $d(g-1) \le j-1 < dg \mod (G-1)$ and g > j. For g = G connect $\theta_{1,G}$ to all the $\theta_{2,j}$'s that have degree less or equal than d. A graphical representation of the resulting precision matrix Q can be found in Section D of the supplementary material.

Proposition 2. Under the design of Example 1 with d fixed and $G \to \infty$, it holds that

$$n_{\boldsymbol{Q}} = \mathcal{O}(pd), \ n_{\boldsymbol{L}} = \mathcal{O}(p^2) \ and \ \operatorname{Cost}(\operatorname{Chol}) = \mathcal{O}(p^3).$$
 (13)

Proposition 2 shows that sparse crossed designs can lead to poor performances of sparse Cholesky solvers. We now provide a numerical study that shows that this is indeed the case for most sparse crossed designs. We consider missing completely at random designs where each cell in the K-dimensional data contingency table contains an observation with probability $\pi \in (0, 1)$ and is empty otherwise. We take $G_1 = \cdots = G_K = G$, hence $N \sim \text{Binomial}(G^K, \pi)$, and we consider, for increasing values of G, different values of K and different ways that π relates to G. Specifically, we examine the following designs:

(a)
$$K = 2, \ \pi = 20G^{-1}$$
; (b) $K = 2, \ \pi = G^{-1/2}$; (c) $K = 5, \ \pi = G^{-K+3/2}$.

Notice that the expected degree is constant in scenario (a), but it increases with G in scenarios (b) and (c). Moreover, notice that by picking levels independently among all the possible G^K combinations, we guarantee $n_{\boldsymbol{Q}} = \mathcal{O}(N)$. Figure 1 reports the results, plotting the number of flops required to compute \boldsymbol{L} , versus p = KG+1 in a log-log scale. In all three regimes, despite \boldsymbol{Q} being sparse, the Cholesky factor \boldsymbol{L} is dense, leading to the worst case scenario $\text{Cost}(\text{Chol}) = \mathcal{O}(p^3)$. Notably, the same results were obtained regardless of the ordering strategy. In Figure 1, we also include the number of flops required for sampling with CG algorithm, which we will analyze in Section 5.5.



Figure 1: Cost(Chol) (red dots) as a function of p (x-axis) in a log-log scale, for scenarios (a)–(c) described above. The purple triangles represent the cost of the CG sampler, which is discussed in Section 5.5. Q is obtained by setting $\tau = 1$ and $T = I_p$. The results are obtained by averaging over 30 independent simulations.

5. Spectral analysis and CG convergence

In this section, we provide a series of spectral results about the posterior precision matrix for Model 1. Namely, we show that all but a fixed number of the eigenvalues of Q, if appropriately preconditioned, concentrate in a small interval, as $p \to +\infty$. In particular, Theorem 4 and 6 will show that the effective condition number remains bounded for increasing p. This suggests that the number of CG iterations required to reach a desired level of accuracy is independent of the size of the problem, leading to dimension-free convergence of the algorithm, which is consistent with numerical experiments (see Section 5.5).

5.1. Notation

We denote with $Diag(\mathbf{Q})$, the diagonal matrix whose diagonal coincide with the one of \mathbf{Q} and, for a given vector \mathbf{v} , we denote with $Diag(\mathbf{v})$ the diagonal matrix whose diagonal is \mathbf{v} . Recalling that $\mathbf{U} = \mathbf{V}^T \mathbf{V}$, we start by decomposing \mathbf{Q} in (11) as $\mathbf{Q} =$ $\mathbf{T} + \tau \cdot Diag(\mathbf{U}) + \tau \mathbf{A}$, where $\mathbf{A} = \mathbf{U} - Diag(\mathbf{U})$ can be seen as a weighted adjacency matrix, which counts the number of times that two levels $\theta_{k,g}$ and $\theta_{k',g'}$ are observed together, see (12). We also define $\mathbf{D} = Diag(\mathbf{A}\mathbb{1}_p)$, where $\mathbb{1}_p \in \mathbb{R}^p$ is a vector of ones, so that \mathbf{D} is a diagonal matrix whose elements are the row-wise sums of \mathbf{A} . Notice that, for crossed effect models, $\mathbf{D} = KDiag(\mathbf{U})$, indeed $U[\theta_{k,g}, \theta_{k,g}]$ counts the number of times that the level $\theta_{k,g}$ is observed, see (12), however, each time it is observed it is also "connected" to other K - 1 levels from the other factors and to the global parameter θ_0 , hence the equality.

When we consider the sub-matrix relative to the random effects only, we use the notation $\mathbf{Q}^{(r)} \in \mathbb{R}^{p-1 \times p-1}$, which is obtained by removing the row and column relative to the global effect θ_0 . We use the same notation for the adjacency matrix $\mathbf{A}^{(r)}$. We let $\mathbf{D}^{(r)} = Diag(\mathbf{A}^{(r)} \mathbb{1}_{p-1}) \in \mathbb{R}^{(p-1) \times (p-1)}$, so that $\mathbf{D}^{(r)} = (K-1) \cdot Diag(\mathbf{U}^{(r)})$.

Denote with \bar{Q} the Jacobi preconditioned precision matrix (Golub and van Loan, 2013, Sec. 11.5.3), i.e.

$$\bar{\boldsymbol{Q}} = Diag(\boldsymbol{Q})^{-1/2} \boldsymbol{Q} Diag(\boldsymbol{Q})^{-1/2} \,. \tag{14}$$

In Section 5.2, we expand on the importance of such preconditioning. We define $\bar{A}^{(r)} = (D^{(r)})^{-1/2} A^{(r)} (D^{(r)})^{-1/2}$, as one would usually normalize the adjacency matrix of a graph.

5.2. Jacobi preconditioning

In this section, we briefly investigate the choice of Jacobi preconditioning. The general idea of preconditioning (Golub and van Loan, 2013, Section 11.5) is to find a non-singular and easy to factorize matrix M s.t. the preconditioned matrix $M^{-1/2}QM^{-1/2}$ is better conditioned than the original Q. A simple modification of the standard CG algorithm allows working with the preconditioned matrix instead of Q (Golub and van Loan, 2013, Algorithm 11.5.1) at the additional cost of computing $x \to M^{-1}x$ at each iteration.

For positive definite matrices, the Jacobi preconditioning is often a natural choice, since $x \to M^{-1}x$ is very cheap to evaluate.

For the specific case of Model 1, the Jacobi preconditioning allows rescaling Q so that its eigenvalues lie within a bounded interval independently of the design (see Theorem 3 for more details). To illustrate this point, we consider the following setting. There are Kfactors of size G_1, \ldots, G_K and, for a given N, we sample uniformly without replacement N out of the possible $\prod_{k=1}^{K} G_k$ combinations of levels. This generates a specific random design V. Notice that this is equivalent to the random design of Figure 1, but for general G_k and for fixed N. In this case, simple computations show that the expected value of the diagonal element $Q[\theta_{k,g}, \theta_{k,g}]$ is $T_k + \tau N/G_k$. When looking at the spectrum of the resulting Q, one can observe, for each factor k, a bulk of G_k eigenvalues centered around the expected value $T_k + \tau N/G_k$, which is a manifestation of spectral concentration for high-dimensional random matrices. The left panel in Figure 2 shows this phenomenon in the case of K = 3 factors. On the other hand, when applying Jacobi preconditioning to Q, the bulks relative to different factors are grouped into a single one centered around 1 (see the right panel of Figure 2). These two different spectral behaviors are reflected in



Figure 2: The two histograms show the spectrum of Q (left) and Q (right). They are obtained by setting $T = I_p$, $\tau = 1$, $G_1 = 300$, $G_2 = 1000$, $G_3 = 2000$. The dashed lines on the left panel correspond to the values $T_k + \tau N/G_k$, for k = 1, 2, 3.

the number of iterations needed for convergence of CG. See for example Table 1, which considers a sequence of designs with 2 factors of increasingly different sizes. There, the effective condition number and the number of CG iteration increase as the difference between factor sizes increases, while they remain constant when considering the Jacobi preconditioned version \bar{Q} . Theorems 4 and 6 provide theoretical results supporting the empirical behavior of the spectrum of \bar{Q} . Given the better performances, we will focus our analysis on the spectrum of \bar{Q} .

Note that, in the right panel of Figure 2, one can observe a cluster of 2 small eigenvalues separated from the remaining part of the spectrum. This shows why we need to consider

C.	Ga	N	No preconditioning		Jacobi preconditioning	
		1 V	$\kappa_{3,p-2}$	n. iters	$\kappa_{3,p-2}$	n. iters
100	100	2828	10.61	31	5.37	25
100	1000	36482	33.08	50	4.39	24
100	10000	1015037	166.91	64	3.33	19

effective condition numbers defined as in (8), and it will be made formal in Theorem 3.

Table 1: Effective condition number and number of CG iterations for Q and \bar{Q} . The design is the same as in Figure 1, with $\pi = p^{-9/10}$. Q is obtained with $T = I_p$ and $\tau = 1$. In the linear system $Q\theta = b$, b has independent entries with uniform distribution on (-0.5, 0.5).

Remark 1 (CG preconditioners). There is a vast literature that studies the choice of the optimal CG preconditioner for different linear systems (Saad, 2003, Ch.10). In our analysis, we have also explored other preconditioning techniques, such as so-called incomplete Cholesky factorizations (see the supplementary material for a more detailed discussion). Nevertheless, we observed results comparable, or slightly worse, to the ones obtained with Jacobi preconditioning. Nishimura and Suchard (2023) also proposed a prior-preconditioned version of conjugate gradient method. However, they focused on Bayesian sparse regression with a large number of covariates, identifying the relevant ones through shrinkage priors. Since we do not consider sparsity-inducing priors, the diagonal elements of the prior term **T** are much smaller than those of the likelihood term $\tau \mathbf{V}^T \mathbf{V}$. This is why, in our case, using **T** as a preconditioner does not improve much the spectrum of **Q**, leading to the same issues observed in Table 1.

5.3. Outlying eigenvalues of \bar{Q}

In this section, we identify a set of outlying eigenvalues of the matrix \bar{Q} . Namely, Theorem 3 shows that, independently of the data design, \bar{Q} always has K small eigenvalues and a large one, close to K + 1.

Theorem 3. Let $U = V^T V$ be the likelihood term of Q in (11). Denote by \overline{U} the matrix $Diag(U)^{-1/2}UDiag(U)^{-1/2}$ and let $\lambda_1(\overline{U}) \leq \cdots \leq \lambda_p(\overline{U})$ be its eigenvalues. Then:

$$\lambda_1(\bar{U}) = \cdots = \lambda_K(\bar{U}) = 0, \qquad \qquad \lambda_p(\bar{U}) = K + 1$$

The previous equalities extend to the eigenvalues $\bar{\mu}_1 \leq \cdots \leq \bar{\mu}_p$ of Q as follows:

$$0 < \bar{\mu}_1 \le \dots \le \bar{\mu}_K \le \frac{\max_{k=0,\dots,K} T_k}{\tau + \max_{k=0,\dots,K} T_k}, \qquad K+1 \le \bar{\mu}_p \le K+2$$

The eigenvalues of Theorem 3 corresponds to those directions where the likelihood is highly or poorly informative. This is a well-known behavior due to the additive structure of the linear predictor η_i in (9). The predictor is very informative about the sum of fixed and random effects, whose direction corresponds to the eigenvalue $\bar{\mu}_p$. However, it provides little information about their relative differences, whose directions correspond to the eigenvalues $\lambda_1(\bar{U}) = \cdots = \lambda_K(\bar{U})$. Notice also that the upper bounds on the small eigenvalues $\bar{\mu}_1, \ldots, \bar{\mu}_K$ in Theorem 3 are quite conservative. In all the numerical examples we have explored, the K smallest eigenvalues of \bar{Q} are very close to zero.

5.4. Connection with graphs and adjacency matrices

The purpose of this section is to draw a connection between the spectrum of \bar{Q} and the one of the adjacency matrix $\bar{A}^{(r)}$ of a *K*-partite graph. We will give sufficient conditions that guarantee that the eigenvalues of \bar{Q} , except for those identified by Theorem 3, concentrate around 1 as $p \to +\infty$. The following corollary extends Theorem 3 to the matrix $\bar{A}^{(r)}$.

Corollary 1. Let $\bar{A}^{(r)} = (D^{(r)})^{-1/2} A^{(r)} (D^{(r)})^{-1/2} \in \mathbb{R}^{(p-1) \times (p-1)}$ be as in Section 5.1, and denote with $\bar{\nu}_1 \leq \cdots \leq \bar{\nu}_{p-1}$ its eigenvalues. Then,

$$\bar{\nu}_1 = \dots = \bar{\nu}_{K-1} = -(K-1)^{-1}, \qquad \bar{\nu}_{p-1} = 1.$$
 (15)

We now focus on the remaining part of the spectrum of $\bar{A}^{(r)}$, which lies in the interval $[\bar{\nu}_K, \bar{\nu}_{p-2}]$. If this interval becomes smaller, we would also expect the spectrum of \bar{Q} to concentrate in a smaller interval, therefore improving the effective condition number. The following lemma quantifies this connection.

Lemma 1. Under Model 1, consider $\bar{Q} \in \mathbb{R}^{p \times p}$ and $\bar{A}^{(r)}$ as above, with eigenvalues $\bar{\mu}_1 \leq \cdots \leq \bar{\mu}_p$ and $\bar{\nu}_1 \leq \cdots \leq \bar{\nu}_{p-1}$ respectively. Let $\bar{\nu}_K \leq 0 \leq \bar{\nu}_{p-2}$, then,

$$\kappa_{K+1,p-2}(\bar{\boldsymbol{Q}}) \le \frac{1 + (K-1)\bar{\nu}_{p-2}}{1 + (K-1)\bar{\nu}_K}.$$
(16)

Lemma 1 provides an interesting connection between the effective condition number $\kappa_{K+1,p-2}(\bar{Q})$, and the concentration of eigenvalues of $\bar{A}^{(r)}$ around 0. This allows to focus on the spectrum of $\bar{A}^{(r)}$ and to leverage results from random graph theory, in order to bound the condition number of \bar{Q} using (16).

Remark 2. The upper-bound in (16) is quite tight in the regimes that we consider. When the bound becomes uninformative for $\bar{\nu}_K = -1/(K-1)$, we numerically observed that $\kappa_{K+1,p-2}(\bar{\mathbf{Q}})$ is very large, in the order of several hundreds. Theoretically, if $\bar{\nu}_K = -1/(K-1)$, we can prove that $\kappa_{K+1,p-2}(\bar{\mathbf{Q}})$ is lower bounded by $\tau (\max_k T_k)^{-1}$, hence the effective condition number can be arbitrarily large depending on the value of $\max_k T_k$. Heuristically, if we want $\kappa_{K+1,p-2}(\bar{\mathbf{Q}})$ to be small, we need $\bar{\nu}_K$ to be bounded away from -1/(K-1).

However, it is important to remark that the condition $\bar{\nu}_K > -1/(K-1)$ is only sufficient for the fast convergence of CG. Clearly, it can happen that, $\bar{\nu}_r > -1/(K-1)$, for some small r > K, even if $\bar{\nu}_K = -1/(K-1)$. In such case, we can extend Lemma 1 to bound $\kappa_{r+1,p-2}(\bar{\mathbf{Q}})$, and CG would still converge fast. In the following analysis, we will focus on $\kappa_{K+1,p-2}(\mathbf{Q})$ for mathematical convenience, in order to obtain a sufficient condition for CG to converge fast. Though, we will also study necessary conditions, as in Theorem 5.

In order for the upper-bound in (16) to be effective, we would like $\bar{\nu}_{p-2}$ to be small, or equivalently $1 - \bar{\nu}_{p-2}$ to be large; but mostly we need $\bar{\nu}_K$ to be bounded away from $-\frac{1}{K-1}$.

The spectral gap $1 - \bar{\nu}_{p-2}$ is a well-studied quantity since it is directly related to the global connectivity of a graph. One may wonder whether global connectivity is also associated to the gap $\bar{\nu}_K + \frac{1}{K-1}$. In Section 5.4.1, we will see that, in the simple case with two factors, this association is straightforward. Indeed, we will prove that a sufficient condition to bound the effective condition number is that the associated graph is made of a unique connected component. On the other hand, in Section 5.4.2, we will show that, when considering more than two factors, a different notion of connectivity is needed.

5.4.1. The case K = 2

In this section, we consider Model 1 with only two factors. In this case, $\bar{A}^{(r)}$ represents a bipartite graph and its spectrum has a very peculiar structure. Indeed, its eigenvalues are symmetric around zero, and, in modulo, less or equal than 1 (Brito et al., 2022). With such property, the link with connectivity of a graph expressed in the previous section becomes straightforward. Indeed, $1 - \bar{\nu}_{p-2} = \bar{\nu}_2 + 1$, which simplifies the bound in Lemma 1 as follows.

Corollary 2. Consider Model 1 with K = 2, then

$$\kappa_{3,p-2}(\bar{\boldsymbol{Q}}) \le \frac{1 + \bar{\nu}_{p-2}}{1 - \bar{\nu}_{p-2}}.$$
(17)

Assume now, that $\mathbf{A}^{(r)}$ is a proper adjacency matrix with binary entries in $\{0, 1\}$. In this case, one can show that $\bar{\nu}_{p-2} < 1$ if and only if the graph associated to the adjacency matrix $\bar{\mathbf{A}}^{(r)}$ consists of a unique connected component (Godsil and Royle, 2001). More in general, the higher is its connectivity, the larger is the gap $1 - \bar{\nu}_{p-2}$ (Godsil and Royle, 2001, Ch. 13.5). With this in mind, Corollary 2 states that CG converges fast, when the graph associated to $\bar{\mathbf{A}}^{(r)}$ is well-connected.

Clearly, higher connectivity comes at the price of having a denser support for Q and consequently a larger cost per iteration of CG. Ideally, CG would be most efficient when the CI graph is well-connected while being very sparse. There is a vast literature of random graph theory that studies graphs with this property. For example, the wellknown *Alon-Boppana bound* (Alon, 1986) gives an upper bound on the spectral gap of regular graphs. Graphs that attain this bound are called *Ramanujan*. Random regular graphs are also known for having great connectivity property, indeed, they can get arbitrarily close to the Alon-Boppana bound, with high probability, as the number of vertices goes to infinity (Friedman, 2003). Brito et al. (2022) extends this result to the case of *random biregular bipartite graph*, i.e. a bipartite graph sampled uniformly at random among all the possible bipartite graphs, where the G_1 vertices of the first part has constant degree d_1 , and the remaining G_2 vertices has degree d_2 . This is why one would expect CG algorithm to have a fast convergence rate under this random design. The following result formalizes this intuition.

Theorem 4. Consider Model 1 with $\mathbf{A}^{(r)}$ being the adjacency matrix of a bipartite, biregular random graph. Then, for any sequence $\epsilon_p \to 0$ as $p \to \infty$, it holds asymptotically almost surely that

$$\kappa_{3,p-2}(\bar{\boldsymbol{Q}}) \le \frac{1 + (1/\sqrt{d_1} + 1/\sqrt{d_2}) + \epsilon_p}{1 - (1/\sqrt{d_1} + 1/\sqrt{d_2}) - \epsilon_p}.$$
(18)

Theorem 4 shows that, even with relatively small degrees d_1 and d_2 , the effective condition number of \bar{Q} becomes close to 1. An analogous, but weaker result can be stated also for Erdős-Rényi random bipartite graphs and can be found in the supplementary material.

5.4.2. The case K > 2

When considering more than two factors, the symmetry of the spectrum of $\bar{A}^{(r)}$ around 0 is lost. One may wonder whether the direct link established by Corollary 2, between the connectivity of the graph associated to $\bar{A}^{(r)}$, expressed by the spectral gap $1 - \bar{\nu}_{p-2}$, and the good conditioning of the matrix \bar{Q} holds true also for K > 2. Unfortunately, this is not the case. Example 2 describes a 3-partite graph with excellent connectivity, whose associated precision matrix \bar{Q} has $\mathcal{O}(p)$ eigenvalues close to zero.

Example 2. Consider the case with K = 3 factors and $G_1 = G_2 = G_3 = G$. We consider $N = G^2$ observations, s.t. for each i = 1, ..., N, the first two factors appear with the level $g_1[i] = g_2[i] = \lceil i/G \rceil$; while the third factor has level $g_3[i] = (i \mod G) + 1$. An example of the resulting conditional independence graph is given in the supplementary material. With such design, the first and third factors, as well as the second and third, are fully connected; however, the first and second factor are very poorly connected. Indeed, each level $\theta_{1,g}$ of the first factor only connects to $\theta_{2,g}$ of the second one, and the sub-graph restricted to the first two factors has G disconnected components.

In this case, the spectral gap of $\bar{\mathbf{A}}^{(r)}$ is $1 - \bar{\nu}_{p-2} = 0.5$ and the CI graph has great connectivity, as each vertex is at most at distance 2 from any other one. Nonetheless, G+1 eigenvalues of $\bar{\mathbf{A}}^{(r)}$ are equal to the lower-bound of -1/2 in (15). As a consequence, CG would need to remove many small eigenvalues before getting to a faster rate of convergence. For instance, if we take G = 200, $\mathbf{T} = \mathbf{I}_p$ (the identity matrix) and $\tau = 1$, we have $\kappa_{G+3,p-2}(\bar{\mathbf{Q}}) \approx 2$, but $\kappa_{G+2,p-2}(\bar{\mathbf{Q}}) \approx 400$.

The intuition behind Example 2 is made more general in the following theorem, which shows that *pairwise connectivity* between all factors is a necessary condition for a well conditioned precision matrix.

Theorem 5. Consider Model 1. Fix any permutation π of the first K integers. For each $\ell = 1, \ldots, K - 1$, let C_{ℓ} be the number of connected components of the sub-graph

restricted to the factors $(\pi(\ell), \pi(\ell+1))$. Then $\bar{\mathbf{A}}^{(r)}$ has at least $\sum_{\ell=1}^{K-1} C_{\ell}$ eigenvalues equal to $-\frac{1}{K-1}$.

Theorem 5 shows that, if we want $\bar{\nu}_K$ to be bounded away from $-\frac{1}{K-1}$, one needs each pair of bipartite sub-graphs to have only one connected component, that is $C_{\ell} = 1$ for each ℓ and π . This is clearly a stronger property than global connectivity of the graph of $\bar{A}^{(r)}$, and in several cases it may not be satisfied. For instance, some datasets may have one factor which is nested into another one, i.e. when each level of the former factor can be observed only with one level of the latter one (see Section 6.2 for an example). In this case, the sub-graph associated to these two factors would present as many connected components as the number of levels in the first factor, and $\bar{A}^{(r)}$ will have as many eigenvalues equal to -1/(K-1). This issue becomes even more relevant when considering interaction terms between factors. Consider two factors of sizes G_1 and G_2 respectively, the interaction term between them is given by a third factor of size $G_1 \cdot G_2$ that consider all the possible pairwise combinations among the levels of the two factors. One can notice that, by construction, the interaction term is nested inside both of the original factors. Hence, the corresponding $\bar{A}^{(r)}$ will have at least $G_1 + G_2$ eigenvalues equal to -1/(K-1). As we will see in Section 6.2, including nested factors and interaction terms adds to the spectrum of \bar{Q} several "problematic" eigenvalues close to zero, slowing down the convergence of CG algorithm.

Unfortunately, neither pairwise connectivity between all factors guarantees $\bar{\nu}_K > -\frac{1}{K-1}$, an example is given in the supplementary material. Indeed, a stronger notion of pairwise connectivity is needed to lower bound $\bar{\nu}_K$. The following result describes such condition, and shows that the effective condition number $\kappa_{K+1,p-2}(\bar{Q})$ is bounded independently of p.

Theorem 6 (Strong pairwise connectivity). Consider Model 1. For any pair of factors $k \neq h$, let $\mathbf{A}^{(k,h)} \in \mathbb{R}^{(G_k+G_h)\times(G_k+G_h)}$ be the adjacency matrix of the bipartite graph restricted to the pair of factors (k,h). Denote with $\bar{\mathbf{A}}^{(k,h)}$ the normalized version obtained as $\mathbf{M}^{-1/2}\mathbf{A}^{(k,h)}\mathbf{M}^{-1/2}$, with $\mathbf{M} = Diag(\mathbf{A}^{(k,h)}\cdot 1)$. Assume that

$$\lambda^* = \sqrt{K-1} \cdot \max\left\{ |\lambda| \colon \lambda \in \sigma\left(\bar{\boldsymbol{A}}^{(k,h)}\right), \, |\lambda| \neq 1, \, k \neq h \right\} < 1 \;. \tag{19}$$

Then, the effective condition number is bounded by

$$\kappa_{K+1,p-2}(\bar{\boldsymbol{Q}}) \le \frac{1+\lambda^*}{1-\lambda^*} .$$
(20)

Condition (19) states that, if the second-largest eigenvalue of $\bar{A}^{(k,h)}$ is bounded away from $\frac{1}{\sqrt{K-1}}$ for all pairs (k, h), then \bar{Q} has a bounded condition number after the removal of few extreme eigenvalues. For example, if we consider a random K-partite d-regular graph, the condition $\frac{2}{\sqrt{d}} < \frac{1}{\sqrt{K-1}}$ i.e. d > 4(K-1), would guarantee (19) asymptotically almost surely, as $p \to \infty$ (see Theorem 4).

5.5. Numerical experiments

The above bounds on the effective condition numbers of \bar{Q} suggest that CG samplers are highly effective for large unstructured designs, which resembles random graphs. In this section, we verify these conclusions through numerical simulations. Specifically, we consider the same numerical experiment as in the end of Section 4 and compute the cost of obtaining an approximate sample by solving (6) with CG, which is given by the product of the cost per iteration and the number of iterations needed for convergence. The former is of order $\mathcal{O}(n_Q)$, while the latter can be quantified as

N. iterations = inf
$$\left\{ k \ge 0 : || \boldsymbol{Q} \boldsymbol{\theta}^k - \boldsymbol{b} ||_2 < \epsilon || \boldsymbol{b} ||_2 \right\}$$
, (21)

where θ^k is the approximate CG solution of the linear system $Q\theta = b$ at iteration k, and Q is the precision matrix of the conditional distribution of θ . Monitoring the relative norm of the residuals is a standard stopping criterion in libraries that implements CG algorithm. In this case, it also allows for a fair comparison between problems with different sizes. In our numerical experiments, we will set the tolerance to $\epsilon = 10^{-8}$.

We consider the three scenarios presented in Figure 1. Table 2 reports the number of iterations defined in (21) in the three different designs, while the plots in Figure 1 show, in purple, the total number of floating points operations in logarithmic scale.

In the first design, we have $N \sim Binomial(G^2, 20/G)$, so that $n_{\mathbf{Q}} = \mathcal{O}(p)$. Here the number of CG iterations is basically constant with p (see the first column of Table 2) and, as a result, the slope for CG in Figure 1 is roughly equal to 1. To be more precise, the small increase in the number of CG iterations is due to the fact that the number of disconnected components in Erdős-Rényi random graphs with a constant degree also increases as the graph size grows (Erdős et al., 1960). Instead, for designs (b) and (c) we have $n_{\mathbf{Q}} = \mathcal{O}(p^{3/2})$ and expected degree and connectivity of the graph that increase with p. This leads to an increasingly better effective condition number, which explains the decrease in the number of iterations observed in Table 2 (second and third column) as well as why the slopes in Figure 1 are below 1.5 even though the cost per iteration is of order $\mathcal{O}(p^{3/2})$.

m	N. iterations			
P	Case (a)	Case (b)	Case (c)	
100	17	22	45	
205	18	21	47	
435	19	20	48	
910	19	21	43	
1910	19	18	39	
4000	19	17	34	

Table 2: Number of CG iterations (21) for different values of p, in scenarios (a)-(c).

As these simulations illustrate, CG excels when it comes to large sparse unstructured random designs. Indeed, for p in the order of 10^3 , CG converges in less than 40 iterations

in all three designs. Therefore, while this framework represents a worst-case scenario for sampling via Cholesky factorization, as its computation requires $\mathcal{O}(p^3)$ time, it is optimal for sampling via CG, since the number of iterations is independent of the dimension p.

6. Application to Generalized Linear Mixed Models

In this section, we discuss and illustrate the application of the conjugate gradient sampler to the more general class of GLMMs. In particular, relative to the simplified framework of Model 1, we allow for the inclusion of multivariate fixed effects, random slopes and interaction terms, as well as non-Gaussian likelihoods with data augmentation. We numerically explore the computational performances of the CG sampler when applied to synthetic and real data in this general framework, considering applications on a recommendation system type data and on a survey data. This comparison will illustrate that CG sampler is efficient in the former scenario, but loses efficiency in the latter one.

6.1. Model and algorithms

We consider GLMMs of the form

$$y_{i} \mid \eta_{i} \sim f\left(y_{i} \mid \eta_{i}\right), \quad \eta_{i} = \boldsymbol{x}_{i,0}^{T} \boldsymbol{\theta}_{0} + \sum_{k=1}^{K} \boldsymbol{x}_{i,k}^{T} \boldsymbol{\theta}_{k}, \qquad i = 1, \dots, N,$$

$$\boldsymbol{\theta}_{k,g} \sim \mathcal{N}\left(\mathbf{0}, \boldsymbol{T}_{k}^{-1}\right), \quad \boldsymbol{\theta}_{k} = \left(\boldsymbol{\theta}_{k,1}^{T}, \cdots, \boldsymbol{\theta}_{k,G_{k}}^{T}\right)^{T}, \quad g = 1, \dots, G_{k}; \quad k = 1, \dots, K.$$

$$(22)$$

Here $\boldsymbol{\theta}_0 \in \mathbb{R}^{D_0}$ denotes the vector of fixed effects and $\boldsymbol{x}_{i,0} \in \mathbb{R}^{D_0}$ denotes the corresponding covariates. For each factor k and level g, we consider the random effect $\boldsymbol{\theta}_{k,g} \in \mathbb{R}^{D_k}$, for $g = 1, \ldots, G_k$, with corresponding covariates $\boldsymbol{w}_{i,k} \in \mathbb{R}^{D_k}$. Up to now, we only considered random intercept models with $D_k = 1$ and $\boldsymbol{w}_{i,k} = 1$, while we now also consider random slopes models with $D_k > 1$. As before, the vector $\boldsymbol{z}_{i,k} \in \{0,1\}^{G_k}$ encodes the level of the factor k to which observation y_i is assigned to. With such notation, $\boldsymbol{\theta}_k$ has covariates $\boldsymbol{x}_{i,k} = \boldsymbol{z}_{i,k} \otimes \boldsymbol{w}_{i,k} \in \mathbb{R}^{G_k D_k}$, where \otimes denotes the Kronecker product.

In previous sections, we discussed the case of Gaussian likelihood function with identity link, namely $f(y_i | \eta_i) = \mathcal{N}(\eta_i, \tau^{-1})$. We consider also binomial likelihood with logistic link function $f(y_i | \eta_i) = \text{Binomial}(n_i, (1 + \exp(-\eta_i))^{-1})$, using the Polya-Gamma data augmentation (Polson et al., 2013) defined as

$$p(y_i, \omega_i \mid \eta_i) = \frac{1}{2^{n_i}} \exp\left(\left[y_i - n_i/2 \right] \eta_i - \omega_i \frac{\eta_i^2}{2} \right) f_{PG}(\omega_i \mid n_i, 0) \,,$$

where $f_{PG}(\omega|b,c)$ indicates a Polya-Gamma variable with parameters b and c. We assign improper flat priors to the fixed effects parameters $\boldsymbol{\theta}_0$ and we assign Wishart priors, $W(\alpha_k, \boldsymbol{\Phi}_k^{-1})$ to the matrices \boldsymbol{T}_k , which simplify to gamma distributions when $D_k = 1$.

Denote with T and m_0 the prior precision and prior mean of θ respectively, with $\kappa = (y_1 - n_1/2, \dots, y_N - n_N/2)^T$ the vector of "centered" observations, and with $\Omega =$

 $(\omega_1, \ldots, \omega_N)^T$ the vector of PG latent variables. With such notation, simple Bayesian computation leads to the following conditional posterior updates

$$p(\boldsymbol{\theta} \mid \boldsymbol{y}, \boldsymbol{V}, \boldsymbol{\Omega}, \{\boldsymbol{T}_k\}_{k=1}^K) \sim \mathcal{N}_p\left(\boldsymbol{Q}^{-1}(\boldsymbol{T}\boldsymbol{m}_0 + \boldsymbol{V}^T\boldsymbol{\kappa}), \ \boldsymbol{Q}^{-1}\right),$$
(23)

$$p(\omega_i \mid \boldsymbol{y}, \boldsymbol{V}, \boldsymbol{\theta}) \stackrel{ind.}{\sim} PG(n_i, \ \boldsymbol{v}_i^T \boldsymbol{\theta}), \qquad \forall i = 1, \dots, N, \quad (24)$$

$$p(\boldsymbol{T}_k \mid \boldsymbol{y}, \boldsymbol{V}, \boldsymbol{\theta}) \stackrel{ind.}{\sim} W\left(\alpha_k + G_k, \ (\boldsymbol{\Phi}_k + \sum_{g=1}^{G_k} \boldsymbol{\theta}_{k,g} \boldsymbol{\theta}_{k,g}^T)^{-1}\right), \quad \forall k = 1, \dots, K, \ (25)$$

where $\boldsymbol{Q} = \boldsymbol{T} + \boldsymbol{V}^T \boldsymbol{\Omega} \boldsymbol{V}$. The resulting Gibbs sampler is described in the supplementary material. The computational bottleneck is the $\mathcal{O}(p^3) \gg \mathcal{O}(N)$ cost for exact sampling from (23), which we seek to reduce with CG samplers.

6.2. Application to voter turnout and student evaluations

We explore the performances of the CG sampler when used to sample from the distribution in (23) in high-dimensional contexts. We consider the following two data sets:

- A survey data of the 2004 American political elections (Ghitza and Gelman, 2013; Goplerud et al., 2024). It collects the vote (Democrat or Republican) of 79148 American citizens. The data set also contains information about their *income* (5 levels), *ethnic group* (4 levels), *age* (4 levels), *region of the country* (5 levels) and *state* (51 levels).
- 2. Instructor Evaluations by Students at ETH (Papaspiliopoulos et al., 2019). A data set with 73421 observations for the following variables: students Id (2972 levels), a factor denoting individual professors (1128 levels), student's age (4 levels), the department of the lecture (14 levels) and a factor describing when the lecture rated had taken place (6 levels). The response is the ratings of lectures by the students (discrete rating from poor to very good), which will be modelled as $y_i \sim Binom(4, \eta_i)$, according to the notation of (22).

We monitor the number of iterations needed for CG to converge, defined as in (21). We also explored other stopping criterion, such as $||\theta^k - \theta||_2 < \epsilon_1 ||\theta||_2$ or $||\theta^k - \theta||_{\infty} < \epsilon_2$, but decided not to include them in the analysis since they yielded similar results. We set the relative accuracy threshold to 10^{-8} , which in our simulations led to practically negligible error introduced by the CG solver. We also tested different accuracy levels (from 10^{-4} to 10^{-8}), but opted for a more conservative one. Indeed, in our experiments once CG reached a higher convergence rate after the "removal" of the extreme eigenvalues, it often reaches very low error with only a few additional iterations (see end of Section 2.2).

We investigate how the number of CG iterations changes depending on:

• **Design complexity**. So far, we have considered missing completely at random designs, which have proven very beneficial for the convergence of the CG method (see Section 5.5). Now, we aim to study how the number of CG iterations varies

when moving away from this favorable random design. Moreover, we also construct two simulated datasets with the same number of factors and levels as in the voter turnout and student evaluations data, but with a random design matrix simulated as described in Section 5.5. We will compare the simulated datasets with the real ones, in order to isolate the contribution of the complexity in the sparsity structure of Q.

- Model complexity. We consider several variations of the model in (22), which we report below in order of increasing size and complexity.
 - (a) *Random intercept*. When we associate to each factor a one-dimensional random effect.
 - (b) Random intercept with nested factors. This scenario often arises where certain levels of a given factor are divided into groups and the group label is also included as another factor, e.g. students who are divided in different classes. For the voter turnout, we consider the factor *state*, which is nested inside *region of the country*. For the student evaluations, the factor denoting the professor is nested inside *department*.
 - (c) *Random slopes.* We include interactions between a continuous predictor and a factor, which "modifies the slope" of the predictor depending on the level of the factor.
 - (d) Interactions. We add interaction terms to the original factors in each data set, e.g. if we have two factors of sizes G_1 and G_2 respectively, the twoway interaction term is a third factor with G_1G_2 levels, which encodes the information about the pairwise interactions of the first two factors. Notice that each interaction term between two factors is, by construction, nested in both factors. We will also include three-way interactions.
- Sample size. We can analyze the impact of sample size by considering, for each data set, a subsample with a smaller size, such as N = 7000, and a larger one with N = 70000. However, instead of subsampling uniformly at random, we enforce each level of each factor to appear in at least one observation. This guarantee to have the same p, as long as we don't include interaction terms, whose size doesn't depend only on which levels are observed, but also on how they interact.
- Factor size. The two data sets we consider differ significantly when it comes to factor size. E.g. the student evaluations data set involves factors with thousands of levels, compared to tens of levels for the voter turnout. We will see that factor size plays a crucial role when considering the number of CG iterations.

Table 3 summarizes the results of our numerical study. Each row shows the number of CG iterations for the different designs described above. We use the simple additive model (a) as a benchmark and consider only one additional term at a time to isolate the contribution of each component. E.g., if we include random slopes, we remove the nested factor and interaction terms. The only exception is the three-way interactions, as it would be unusual to consider them without the pairwise interactions. The last row presents results for the full model, which includes all the effects mentioned. Each entry in Table 3 reports the number of iterations for different sample sizes. Additionally, for each data set, we include a column displaying the number of iterations obtained using simulated data.

Casa	Voter Turnout		Students Evaluations	
Case	Real	Simulated	Real	Simulated
Random intercents	30(68)	25~(68)	26 (4101)	18 (4101)
Random intercepts	36~(68)	24 (68)	35~(4101)	16(4101)
Nested factor	42(73)	28(73)	63 (4115)	25 (4115)
Nested factor	54(73)	27(73)	94~(4115)	22 (4115)
Bandom slopes	48(127)	38(127)	84 (5229)	34 (5229)
Random slopes	68(127)	40(127)	150 (5229)	34 (5229)
2 way interactions	$134\ (763)$	99~(787)	91~(19523)	31 (23585)
	343~(786)	190(787)	$122 \ (89385)$	$35 \ (97897)$
3 way interactions	166 (2952)	111 (3720)	98~(27588)	32 (31650)
	442 (3569)	258 (3723)	$128 \ (159633)$	36~(168028)
Full	189(3016)	147 (3784)	122 (28730)	58 (32792)
1 un	532(3633)	351 (3787)	266 (160775)	65(169170)

Table 3: Average number of CG iterations. The average is obtained over 200 Gibbs sampler iterations after an initial burn-in. The number in parentheses represents p. In each entry, we report results obtained with $N = 7\,000$ (above) and $N = 70\,000$.

Based on the results in Table 3, we draw the following conclusions:

- In accordance with Theorem 5, we observe that introducing factors with poor pairwise connectivity results in an increase in CG iterations. This increase occurs when we include nested factors (e.g., *region* for voter turnout and *department* for student evaluations) or interaction terms, which introduce nesting by construction. For the voter turnout case, the number of CG iterations increases significantly when multiway interactions are added, moving from 36 to 343 iterations by simply introducing 2 way interactions. For the student evaluations data, adding interaction terms appears less problematic relative to the problem size, although including interactions between such large factors is usually uncommon. Analogous patterns are observed in the simulated data sets.
- Table 3 shows, for the real data, an overall increase in the number of CG iterations when the sample size increases. This might seem counter-intuitive since, for larger N, the CI graph becomes denser and better connected. However, when examining the spectrum of \bar{Q} for the random intercept only case, we observe that, for small N, the interval $[\bar{\mu}_{K+1}, \ \bar{\mu}_{p-2}]$ is larger but most of the eigenvalues therein are

tightly concentrated in its center, around 1. In contrast, for larger N, the interval $[\bar{\mu}_{K+1}, \bar{\mu}_{p-2}]$ is smaller, consistently with the CI graph being better connected, but the spectrum is more diffuse. In other words, for smaller N, the effective condition number $\kappa_{K+1,p-2}(\bar{Q})$ is worse in accordance with the theoretical analysis of Section 5. However, in this example, $\kappa_{s+1,p-r}$ is better for larger s, r, resulting in an overall faster convergence rate.

• The comparison between these two data sets highlights the significant impact of factor size on the number of CG iterations. For the voter turnout data, which presents relatively small factors, CG can require several hundred iterations, even for solving relatively small problems on the order of a few thousand. Moreover, this issue is not specific to the details of the voter turnout design, as a similarly large number of iterations is needed also in the simulated case with a missing completely at random design but same factor sizes. On the other hand, when considering the student evaluations case, where some factors have numerous levels, CG proves to be very efficient, requiring only a few hundred iterations to solve significantly large linear systems in the order of 10⁵. This difference becomes even more significant in the simulated case, where approximately 50 iterations are sufficient for convergence, even for the largest problems.

7. Comparison with existing works

GLMMs present significant computational challenges, especially in high-dimensional settings. Efficient computation for these models has been extensively studied in scenarios where the random effects have a nested dependence structure (Tan and Nott, 2013, 2018; Tan, 2020; Papaspiliopoulos and Zanella, 2017). Successful strategies have also been proposed for more complex dependence structures, arising e.g. from time-series models (Saul and Jordan, 1995; Salimans and Knowles, 2012; Tan and Nott, 2018; Frazier et al., 2023) or spatial data analysis (Rue and Held, 2005; Katzfuss and Guinness, 2021; Schäfer et al., 2021). However, the complexity of the crossed dependence structure hinder the applicability of these methodologies for GLMMs with more than one factor.

It's known that standard techniques become inefficient when applied to high dimensional crossed effects models (Gao and Owen, 2020; Papaspiliopoulos et al., 2019). For this reason, various computational approaches have been explored to address this issue, including backfitting with centering (Ghosh et al., 2022), method of moments (Gao and Owen, 2020), collapsed Gibbs sampler (Papaspiliopoulos et al., 2019; Ceriani and Zanella, 2024; Papaspiliopoulos et al., 2023), and variational inference methods (Menictas et al., 2023; Goplerud et al., 2024).

The CG method is widely recognized for its effectiveness in solving large, sparse linear systems, and it has been extensively applied also for sampling from high dimensional Gaussian distributions (Papandreou and Yuille, 2010; Simpson et al., 2013; Vono et al., 2022). However, there is a lack of theoretical studies analyzing the effectiveness of CG for GLMMs. The recent work Nishimura and Suchard (2023) provides a detailed study

of the use of CG in the context of large Bayesian sparse regression models. Differently from their analysis, we consider the case where the design matrix itself V is sparse.

8. Discussion

We remark that the applicability of the methodologies presented in this article goes beyond Gibbs sampling. Standard inferential methodologies such as restricted maximum likelihood estimation, Laplace approximation and variational inference require the computation of the Cholesky factorization of a matrix as in (1), and would suffer from the same limitations presented in Section 4. On the other hand, some of these methodologies can benefit from the use of CG and similar iterative methods. For example, in the context of ML estimation, one could employ the method of moments proposed in Gao and Owen (2020), to obtain consistent estimates of the variance components and then use conjugate gradient to compute the GLS estimator. Also, in the context of partially factorized or unfactorized variational inference (Goplerud et al., 2024), one is required to invert matrices with the same structure of (1), which can be computed via CG. In all these cases, the spectral analysis of Section 5 may be of interest to predict theoretical performances. However, the applicability of CG algorithm is hindered in those cases where the determinant of large matrices is also needed, e.g., for the computation of marginal quantities. Several strategies have been proposed to approximate determinants of matrices via Lanczos algorithm (Ubaru et al., 2017), though they are computationally more expensive than CG, and it is harder to control for the error induced by such approximations.

This paper raises both theoretical and methodological questions. For example, it would be interesting to provide an explicit result, which proves the cubic complexity of the Cholesky factorization under generic (K-partite) random designs. As for the analysis of CG algorithm, Theorem 3 and 5 shows how linear dependence among the columns of the design matrix V relates to small eigenvalues of Q, which, in general, significantly slows down the convergence of CG. A standard technique to address this issue involves imposing linear constraints on the vector of parameters θ (Wood, 2017; Zanella and Roberts, 2020). Perhaps surprisingly, an initial numerical exploration of such procedure suggests that it actually does not lead to a significant improvement of the rate of convergence of CG algorithm, and, in some cases, may even slow it down. We leave a more detailed investigation to future work.

References

Alon, N. (1986). Eigenvalues and expanders. Combinatorica, 6(2):83–96.

Ashcroft, C. J. (2021). On the eigenvalues of Erdős-Rényi random bipartite graphs.

Brito, G., Dumitriu, I., and Harris, K. D. (2022). Spectral gap in random bipartite biregular graphs and applications. *Combinatorics, Probability and Computing*.

- Ceriani, P. M. and Zanella, G. (2024). Linear-cost unbiased posterior estimates for crossed effects and matrix factorization models via couplings. *preprint arXiv:2410.08939*.
- Chung, F. R. (1997). Spectral graph theory, volume 92. American Mathematical Soc.
- Concus, P., Golub, G. H., and O'Leary, D. P. (1976). A generalized conjugate gradient method for the numerical solution of elliptic partial differential equations. In *Sparse* matrix computations, pages 309–332. Elsevier.
- Erdős, P., Rényi, A., et al. (1960). On the evolution of random graphs. Publ. math. inst. hung. acad. sci, 5(1):17–60.
- Frazier, D. T., Loaiza-Maya, R., and Martin, G. M. (2023). Variational Bayes in state space models: Inferential and predictive accuracy. *Journal of Computational and Graphical Statistics*, 32(3):793–804.
- Friedman, J. (2003). Relative expanders or weakly relatively Ramanujan graphs. Duke Mathematical Journal, 118(1).
- Gao, K. and Owen, A. (2017). Efficient moment calculations for variance components in large unbalanced crossed random effects models. *Electronic Journal of Statistics*, 11(1):1235–1296.
- Gao, K. and Owen, A. B. (2020). Estimation and inference for very large linear mixed effects models. *Statistica Sinica*, 30:1741–1771.
- Gelman, A. and Hill, J. (2007). Data analysis using regression and multilevel/hierarchical models, volume 3. Cambridge University Press New York, New York, USA.
- Ghitza, Y. and Gelman, A. (2013). Deep interactions with MRP: Election turnout and voting patterns among small electoral subgroups. *American Journal of Political Science*, 57(3):762–776.
- Ghosh, S., Hastie, T., and Owen, A. B. (2022). Backfitting for large scale crossed random effects regressions. *The Annals of Statistics*, 50(1):560 583.
- Godsil, C. and Royle, G. F. (2001). *Algebraic graph theory*, volume 207. Springer Science & Business Media.
- Golub, G. H. and van Loan, C. F. (2013). *Matrix Computations*. JHU Press, fourth edition.
- Goplerud, M., Papaspiliopoulos, O., and Zanella, G. (2024). Partially factorized variational inference for high-dimensional mixed models. *Biometrika, in press.*
- Horn, R. A. and Johnson, C. R. (2012). *Matrix Analysis*. Cambridge University Press, 2 edition.

- Katzfuss, M. and Guinness, J. (2021). A general framework for Vecchia approximations of Gaussian processes. *Statistical Science*, 36(1):124 141.
- Lanczos, C. (1950). An iteration method for the solution of the eigenvalue problem of linear differential and integral operators. *Journal of Research of the National Bureau* of Standards.
- Lin, C.-J. and Moré, J. J. (1999). Incomplete Cholesky factorizations with limited memory. SIAM Journal on Scientific Computing, 21(1):24–45.
- Menictas, M., Di Credico, G., and Wand, M. P. (2023). Streamlined variational inference for linear mixed models with crossed random effects. *Journal of Computational and Graphical Statistics*, 32(1):99–115.
- Nishimura, A. and Suchard, M. A. (2023). Prior-preconditioned conjugate gradient method for accelerated Gibbs sampling in "large n, large p" Bayesian sparse regression. *Journal of the American Statistical Association*, 118(544):2468–2481.
- Papandreou, G. and Yuille, A. L. (2010). Gaussian sampling by local perturbations. Advances in Neural Information Processing Systems, 23.
- Papaspiliopoulos, O., Roberts, G. O., and Zanella, G. (2019). Scalable inference for crossed random effects models. *Biometrika*, 107(1):25–40.
- Papaspiliopoulos, O., Stumpf-Fétizon, T., and Zanella, G. (2023). Scalable Bayesian computation for crossed and nested hierarchical models. *Electronic Journal of Statis*tics, 17(2):3575 – 3612.
- Papaspiliopoulos, O. and Zanella, G. (2017). A note on MCMC for nested multilevel regression models via belief propagation. Arxiv 1704.06064.
- Parker, A. and Fox, C. (2012). Sampling Gaussian distributions in Krylov spaces with conjugate gradients. SIAM Journal on Scientific Computing, 34(3):B312–B334.
- Perry, P. O. (2016). Fast moment-based estimation for hierarchical models. Journal of the Royal Statistical Society: Series B (Statistical Methodology), 79(1):267–291.
- Polson, N. G., Scott, J. G., and Windle, J. (2013). Bayesian inference for logistic models using Pólya–Gamma latent variables. *Journal of the American Statistical Association*, 108(504):1339–1349.
- Rivin, I. (2002). Counting cycles and finite dimensional Lp norms. Advances in Applied Mathematics, 29(4):647 – 662.
- Rue, H. and Held, L. (2005). *Gaussian Markov random fields: theory and applications*. Chapman & Hall.
- Rue, H., Martino, S., and Chopin, N. (2009). Approximate Bayesian inference for latent Gaussian models by using integrated nested Laplace approximations. *Journal of the* royal statistical society: Series b (statistical methodology), 71(2):319–392.

- Saad, Y. (2003). Iterative Methods for Sparse Linear Systems. Other Titles in Applied Mathematics. SIAM, second edition.
- Salimans, T. and Knowles, D. (2012). Fixed-form variational posterior approximation through stochastic linear regression. *Bayesian Analysis*, 8.
- Saul, L. and Jordan, M. (1995). Exploiting tractable substructures in intractable networks. In Advances in Neural Information Processing Systems, volume 8. MIT Press.
- Schäfer, F., Katzfuss, M., and Owhadi, H. (2021). Sparse Cholesky factorization by Kullback–Leibler minimization. SIAM Journal on Scientific Computing, 43(3):A2019– A2046.
- Simpson, D. P., Turner, I. W., Strickland, C. M., and Pettitt, A. N. (2013). Scalable iterative methods for sampling from massive Gaussian random vectors. *preprint at* arXiv:1312.1476.
- Tan, L. S. L. (2020). Use of model reparametrization to improve variational Bayes. Journal of the Royal Statistical Society Series B: Statistical Methodology, 83(1):30–57.
- Tan, L. S. L. and Nott, D. J. (2013). Variational inference for generalized linear mixed models using partially noncentered parametrizations. *Statistical Science*, 28(2):168– 188.
- Tan, L. S. L. and Nott, D. J. (2018). Gaussian variational approximation with sparse precision matrices. *Statistics and Computing*, 28:259–275.
- Trefethen, L. and Bau, D. (1997). Numerical Linear Algebra. SIAM.
- Ubaru, S., Chen, J., and Saad, Y. (2017). Fast estimation of tr(f(a)) via stochastic Lanczos quadrature. SIAM Journal on Matrix Analysis and Applications.
- Van der Sluis, A. and Van der Vorst, H. A. (1986). The rate of convergence of conjugate gradients. Numerische Mathematik, 48:543–560.
- van der Vorst, H. A. (2003). *Iterative Krylov methods for large linear systems*. Cambridge University Press, Cambridge; New York.
- Vono, M., Dobigeon, N., and Chainais, P. (2022). High-dimensional Gaussian sampling: A review and a unifying approach based on a stochastic proximal point algorithm. SIAM Review, 64(1):3–56.
- Wood, S. N. (2017). Generalized additive models: an introduction with R. CRC press.
- Zanella, G. and Roberts, G. (2020). Multilevel linear models, Gibbs samplers and multigrid decompositions. *Bayesian Analysis*.

A. Proofs

Proof of Theorem 1. The number of operations required by the recursive equation in (2) is

$$Cost(Chol) = \sum_{m=1}^{p} 1 + n_{L,m} + n_{L,m} (1 + n_{L,m}), \qquad (26)$$

see Theorem 2.2 of https://www.tau.ac.il/~stoledo/Support/chapter-direct.pdf for a step-by-step derivation of (26). In both cases, we have $\text{Cost}(\text{Chol}) = \mathcal{O}\left(\sum_{m=1}^{p} n_{\boldsymbol{L},m}^{2}\right)$ as stated in (5). Then, using Jensen's inequality and $\sum_{m=1}^{p} n_{\boldsymbol{L},m} = n_{\boldsymbol{L}}$, we have

$$\sum_{m=1}^{p} n_{\boldsymbol{L},m}^{2} = p\left(p^{-1} \sum_{m=1}^{p} n_{\boldsymbol{L},m}^{2}\right) \ge p\left(p^{-1} \sum_{m=1}^{p} n_{\boldsymbol{L},m}\right)^{2} = n_{\boldsymbol{L}}^{2}/p,$$

which proves the lower bound in (5).

The upper bound in (5), instead, cannot be deduced purely from $\operatorname{Cost}(\operatorname{Chol}) = \mathcal{O}\left(\sum_{m=1}^{p} n_{L,m}^{2}\right)$, but rather it follows from a characterization of $\operatorname{Cost}(\operatorname{Chol})$ in terms of 3-cycles of an undirected graph associated to L. More precisely, define G_{L} as the undirected graph with nodes $\{1, \ldots, p\}$ and an edge between vertices j > m if and only if the future set of θ_{m} does not separate it from θ_{j} in G_{Q} . For any $j, m \in \{1, \ldots, p\}$ with $j \neq m$, we write $\{j, m\} \in G_{L}$ if there is an edge between j and m in G_{L} . By the arguments in Section 2.1, $L_{j\ell}$ is (a potential) non-zero if and only if $\{j, \ell\} \in G_{L}$. The dominating cost of the recursion (2) to obtain L is the computation of $\sum_{\ell=1}^{m-1} L_{m\ell}L_{j\ell}$ for $m = 2, \ldots, p$ and $j = m + 1, \ldots, p$. Ignoring multiplications by zero and summation of zeros, this corresponds to

$$\mathcal{O}(|\{(\ell, m, j) : \{m, \ell\} \in G_L, \{j, \ell\} \in G_L \text{ and } 1 \le \ell < m < j \le p\}|)$$
(27)

operations. By definition of G_L , if $\{\ell, m\} \in G_L$ and $\{\ell, j\} \in G_L$ for $\ell < m < j$, then also $\{\ell, m\} \in G_L$. Thus, the cost in (27) coincides with the number of 3-cycles in G_L . The upper bound in (5) then follows, noting that the number of 3-cycles in an undirected graph with n_L edges is less or equal than $n_L^{1.5}$, see e.g. (Rivin, 2002, Theorem 4).

Proof of Proposition 1. Let $\boldsymbol{\theta} \in \mathbb{R}^p$ be an arbitrary permutation of $(\theta_0, \theta_1, \ldots, \theta_K)$ and denote by ℓ the position of θ_0 in $\boldsymbol{\theta}$. We show that, whenever $\ell < p$, switching the positions of θ_0 and the variable immediately after it in $\boldsymbol{\theta}$ does not increase n_L . Such switching of positions does not change the future set of θ_m for any $m \notin \{\ell, \ell+1\}$ and thus leaves also $n_{L,m}$ unchanged by its definition in (4). Moreover, since θ_0 is connected to all other variables in G_Q , it follows that $n_{L,m}$ equals the maximum value of p-m+1for all θ_m located after θ_0 in $\boldsymbol{\theta}$ and for θ_0 itself. Thus, when θ_0 is in position ℓ both $n_{L,\ell}$ and $n_{L,\ell+1}$ take their maximal values and moving θ_0 to position $\ell + 1$ cannot increase the value of $(n_{L,\ell} + n_{L,\ell+1})$.

Proof of Proposition 2. Denote with $Q[\theta_{k,g}; \theta_{k',g'}]$ the entry of Q relative to the pair $(\theta_{k,g}, \theta_{k',g'})$, and recall that, for $k \neq k'$, its value coincides with the number of observations $i = 1, \ldots, N$ s.t. $z_{i,k,g} = 1$ and $z_{i,k',g'} = 1$. Finally, $\theta_{k,g}$ is adjacent to $\theta_{k',g'}$ in G_Q if and only if $Q[\theta_{k,g}; \theta_{k',g'}] > 0$.

Define the function $r : \{1, \ldots, G\} \mapsto \{1, \ldots, G\}$ as r(1) = 1 and $r(j) = \lceil d^{-1}(j-1) \rceil$ for $j = 2, \ldots, G$, so that $Q[\theta_{1,r(j)}; \theta_{2,j}] = 1$ for all $j \ge 1$. Then, for every couple j and m such that $r(j) \le m \le j$ we now show that the future set of $\theta_{2,m}$ does not separate it from $\theta_{2,j}$ in $G_{\mathbf{Q}}$. We construct a path in $G_{\mathbf{Q}}$ between $\theta_{2,m}$ and $\theta_{2,j}$ that goes through $\theta_{2,1}$ as follows. Since both $Q[\theta_{1,j}; \theta_{2,j}]$ and $Q[\theta_{1,r(j)}; \theta_{2,j}]$ are positive for all $j \ge 1$, the path going from $\theta_{2,j}$ to $\theta_{1,r(j)}$ to $\theta_{2,r(j)}$ to $\theta_{1,r(r(j))}$ to $\theta_{2,r(r(j))}$ etc. is supported on $G_{\mathbf{Q}}$. Also, since $r(\ell) < \ell$ for all $\ell \ge 2$ and r(1) = 1, the above path eventually reaches $\theta_{2,1}$. The same strategy can be applied to construct a path in $G_{\mathbf{Q}}$ from $\theta_{2,m}$ to $\theta_{2,1}$. Joining the two above paths at $\theta_{2,1}$, we obtain a path from $\theta_{2,m}$ to $\theta_{2,j}$ in $G_{\mathbf{Q}}$. The same strategy can be applied to construct a path in $G_{\mathbf{Q}}$ from $\theta_{2,m}$ to $\theta_{2,1}$. Joining the two above paths at $\theta_{2,1}$, we obtain a path from $\theta_{2,m}$ to $\theta_{2,j}$ in $G_{\mathbf{Q}}$. The same strategy can be applied to construct a path in $G_{\mathbf{Q}}$ from $\theta_{2,m}$ to $\theta_{2,m}$ is a potential non-zero whenever $r(j) \le m \le j$, meaning that the row of \mathbf{L} corresponding to $\theta_{2,j}$ contains at least j - r(j) + 1 potential non-zeros. Summing over j we obtain

$$n_{L} \ge \sum_{j=2}^{G} (j - r(j) + 1) \ge \sum_{j=2}^{G} \left(j - \frac{j-1}{d} \right) \ge \frac{d-1}{d} \sum_{j=2}^{G} j = \frac{d-1}{d} \left(\frac{G(G+1)}{2} - 1 \right) \,.$$

Thus, $n_{L} = \mathcal{O}(G^{2})$ which also implies $\operatorname{Cost}(\operatorname{Chol}) = \mathcal{O}(G^{3})$ by the lower bound in Theorem 1. The statements about n_{L} and $\operatorname{Cost}(\operatorname{Chol})$ follow from the above equalities and p = 1 + 2G.

Proof of Theorem 3. We start by proving the results relative to the likelihood term $\boldsymbol{U} = \boldsymbol{V}^T \boldsymbol{V} = \sum_{i=1}^N \boldsymbol{v}_i \boldsymbol{v}_i^T$.

Proof of part 1. First recall, from Section 3, that $\mathbf{v}_i = (1, \mathbf{z}_{i,1}^T, \dots, \mathbf{z}_{i,K}^T)^T \in \mathbb{R}^p$, where $\mathbf{z}_{i,k}$ have only one entry with unitary value and the remaining ones are zero. Consider the sets of vector $W = \{\mathbf{w}_k; k = 1, \dots, K\}$, where each $\mathbf{w}_k = (-1, \mathbf{0}_{G_1}^T, \dots, \mathbf{1}_{G_k}^T, \dots, \mathbf{0}_{G_K}^T)^T$, and the vector of ones $\mathbb{1}_{G_k}$ is in the position relative to the k-th factor. Then, for all $i = 1, \dots, N$ and $k = 1, \dots, K$, we have $\mathbf{v}_i^T \mathbf{w}_k = \mathbf{0}_p$. Thus, $\mathbf{U}\mathbf{w}_k = \mathbf{0}_p$ for each k. Since the vectors in W are linearly independent, we can deduce that $dim(Null(\mathbf{U})) \geq K$. Also, since U is positive semidefinite, then at least the first K eigenvalues must be equal to zero. The same holds for $\bar{\mathbf{U}}$, since the matrix $Diag(\mathbf{U})$ is positive definite by hypothesis $(U_{ii} \geq 1 \text{ for each } i)$. We have thus shown $\lambda_1(\bar{\mathbf{U}}) = \dots = \lambda_K(\bar{\mathbf{U}}) = 0$, as desired.

Proof of part 2. Recall that U = Diag(U) + A, where A can be interpreted as a weighted adjacency matrix representing a (K + 1)-partite graph (we can think of the global effect θ_0 as an additional part with one single vertex). We define \bar{A} as the adjacency matrix A normalized by the row-wise sum of its elements, i.e. $\bar{A} = D^{-1/2}AD^{-1/2}$. As we showed in Section 5, it holds that

$$\boldsymbol{D}_{ii} = \sum_{j=1}^{p} A_{ij} = K U_{ii} \,.$$

It follows that \bar{U} and \bar{A} satisfy $\bar{U} = I_p + K\bar{A}$ and, in particular, they share the same eigenvectors. Finally, recall that the largest eigenvalue of any normalized adjacency matrix \bar{A} is $\lambda_p(\bar{A}) = 1$ (Chung, 1997). As a consequence, the largest eigenvalue relative to \bar{U} is $\lambda_p(\bar{U}) = 1 + K\lambda_p(\bar{A}) = K + 1$, as desired.

Proof of the bounds on $\bar{\mu}_1, \ldots, \bar{\mu}_p$. Finally, we translate the spectral results on \bar{U} into bound on the eigenvalues of \bar{Q} . Notice that $\bar{Q} = E + \bar{U}$ where E is a diagonal matrix with elements $E_{ii} = (T_{ii} + \tau U_{ii})/T_{ii}$. Since $U_{ii} \ge 1$, we thus can bound the eigenvalues of E as

$$\lambda_p(\boldsymbol{E}) = \max_i E_{ii} \leq \frac{T_{ii}}{T_{ii} + \tau} \leq \frac{T}{\tau + \bar{T}}$$

where $\overline{T} = \max_{k=0,...K} T_k$. We can then obtain the inequalities of Theorem 3 combining the above inequality with Weyl's inequality (Horn and Johnson, 2012, Theorem 4.3.1). Indeed we have

$$0 < \bar{\mu}_K \leq \underbrace{\lambda_K(\bar{\boldsymbol{U}})}_{=0} + \lambda_p(\boldsymbol{E}) \leq \frac{T}{\tau + \bar{T}},$$

as well as

$$\bar{\mu}_p \le \lambda_p(\bar{U}) + \underbrace{\lambda_p(E)}_{\le 1} \le K + 2$$

and

$$\bar{\mu}_p \ge \lambda_p(\bar{U}) + \underbrace{\lambda_1(E)}_{\ge 0} \ge K + 1.$$

Proof of Corollary 1. We begin by defining $\bar{\boldsymbol{U}}^{(r)} = Diag(\boldsymbol{U}^{(r)})^{-1/2}\boldsymbol{U}^{(r)}Diag(\boldsymbol{U}^{(r)})^{-1/2}$. Theorem 3 applied to such matrix guarantees that its spectrum is concentrated in the interval [0, K] and, more specifically:

$$\lambda_1(\bar{\boldsymbol{U}}^{(r)}) = \dots = \lambda_1(\bar{\boldsymbol{U}}^{(r)})_{K-1} = 0; \qquad \lambda_1(\bar{\boldsymbol{U}}^{(r)})_{p-1} = K.$$
 (28)

Recalling from Section 5.1 that $\boldsymbol{D}^{(r)} = (K-1)Diag(\boldsymbol{U}^{(r)})$, we have

$$\bar{\boldsymbol{U}}^{(r)} = \boldsymbol{I}_{p-1} + (K-1)\bar{\boldsymbol{A}}^{(r)}$$

This implies that for each eigenvalue λ of $\bar{U}^{(r)}$, $(\lambda - 1)/(K - 1)$ is an eigenvalue for $\bar{A}^{(r)}$, from which follows (15).

Proof of Lemma 1. Since $\bar{\boldsymbol{Q}}^{(r)}$ is obtained from $\bar{\boldsymbol{Q}}$ by removing the row and column relative to θ_0 , we can apply Cauchy interlacing theorem (Horn and Johnson, 2012, Theorem 4.3.17), which implies

$$\bar{\mu}_1 \le \lambda_1(\bar{\boldsymbol{Q}}^{(r)}) \le \bar{\mu}_2 \le \dots \bar{\mu}_{p-1} \le \lambda_{p-1}(\bar{\boldsymbol{Q}}^{(r)}) \le \bar{\mu}_p .$$
⁽²⁹⁾

Note that, with the notation introduced in Section 5.1, it holds that

$$\bar{\boldsymbol{Q}}^{(r)} = \boldsymbol{I}_{p-1} + (\boldsymbol{C}^{(r)})^{1/2} \bar{\boldsymbol{A}}^{(r)} (\boldsymbol{C}^{(r)})^{1/2},$$

where $C^{(r)}$ is the diagonal matrix with elements

$$C_{ii}^{(r)} = \frac{\tau D_{ii}^{(r)}}{T_{ii}^{(r)} + \tau U_{ii}^{(r)}} = \frac{\tau (K-1)U_{ii}^{(r)}}{T_{ii}^{(r)} + \tau U_{ii}^{(r)}}$$

Since $U_{ii}^{(r)} > 0$ by assumption and $T_{ii}^{(r)} \ge 0$, then also $0 < C_{ii}^{(r)} \le (K-1)$. Thus, $\mathbf{C}^{(r)}$ is a positive definite diagonal matrix with the largest diagonal element $\gamma = \max_i C_{ii}^{(r)} \le K-1$. Now, if $\bar{\nu}_{p-r} \ge 0$ also $\lambda_{p-r}((\mathbf{C}^{(r)})^{1/2}\bar{\mathbf{A}}^{(r)}(\mathbf{C}^{(r)})^{1/2}) \ge 0$, and it can be upperbounded by $\gamma \bar{\nu}_{p-1}$. An analogous lower bound can be obtained for all those $\bar{\nu}_s \le 0$. To conclude, we have

$$\kappa_{s+1,p-r}(\bar{\boldsymbol{Q}}) = \frac{\mu_{p-r}}{\bar{\mu}_{s+1}}$$

$$\leq \frac{\lambda_{p-r}(\bar{\boldsymbol{Q}}^{(r)})}{\lambda_s(\bar{\boldsymbol{Q}}^{(r)})}$$

$$= \frac{1 + \lambda_{p-r}((\boldsymbol{C}^{(r)})^{1/2}\bar{\boldsymbol{A}}^{(r)}(\boldsymbol{C}^{(r)})^{1/2})}{1 + \lambda_s((\boldsymbol{C}^{(r)})^{1/2}\bar{\boldsymbol{A}}^{(r)}(\boldsymbol{C}^{(r)})^{1/2})}$$

$$\leq \frac{1 + (K-1)\bar{\nu}_{p-r}}{1 + (K-1)\bar{\nu}_s}.$$

		I
		I
		I
		I

Proof of Corollary 2. We just need to prove that the spectrum of $\bar{A}^{(r)}$ is symmetric around zero. This together with Lemma 1 gives the result. Recall that the matrix $\bar{A}^{(r)}$ can be represented as

$$ar{m{A}}^{(r)} = egin{bmatrix} m{0} & m{B} \ m{B}^T & m{0} \end{bmatrix}$$
 ,

with $\boldsymbol{B} \in \mathbb{R}^{G_1 \times G_2}$. Consider w.l.o.g. any positive eigenvalue $\lambda > 0$ of $\bar{\boldsymbol{A}}^{(r)}$. Let \boldsymbol{x} be one of its eigenvector and split it into $\boldsymbol{x} = [\boldsymbol{x}_1, \boldsymbol{x}_2]$, where $\boldsymbol{x}_1 \in \mathbb{R}^{G_1}$ and $\boldsymbol{x}_2 \in \mathbb{R}^{G_2}$. Then $\boldsymbol{x}^* = [\boldsymbol{x}_1, -\boldsymbol{x}_2]$ is an eigenvalue relative to $-\lambda$. Indeed:

$$ar{oldsymbol{A}}^{(r)}oldsymbol{x}^* = egin{bmatrix} -oldsymbol{B}^Toldsymbol{x}_2\ Boldsymbol{x}_1\end{bmatrix} = egin{bmatrix} -\lambdaoldsymbol{x}_1\ \lambdaoldsymbol{x}_2\end{bmatrix} = -\lambdaoldsymbol{x}^*.$$

Since this operation also preserve the multiplicity of the eigenvalues, the result holds true. $\hfill \Box$

Proof of Theorem 4. By Theorem 3.2 of Brito et al. (2022), the adjacency matrix $A^{(r)}$ satisfies

$$\lambda_{p-2}(\mathbf{A}^{(r)}) \le \sqrt{d_1 - 2} + \sqrt{d_2 - 1} + \epsilon_p,$$

$$\lambda_2(\mathbf{A}^{(r)}) \ge -(\sqrt{d_1 - 2} + \sqrt{d_2 - 1}) - \epsilon_p,$$

asymptotically almost surely, with $\epsilon_p \to 0$, as $p \to \infty$. For a bipartite biregular graph, it holds that $\bar{\mathbf{A}}^{(r)} = \mathbf{A}^{(r)}/\sqrt{d_1 d_2}$, then the bound above extends to the eigenvalues $\bar{\nu}_2$ and $\bar{\nu}_{p-2}$ of $\bar{\mathbf{A}}^{(r)}$ as follows

$$\bar{\nu}_{p-2} \le \frac{1}{\sqrt{d_1}} + \frac{1}{\sqrt{d_2}} + \epsilon_p , \\ \bar{\nu}_2 \ge -\frac{1}{\sqrt{d_1}} - \frac{1}{\sqrt{d_2}} - \epsilon_p .$$

Finally, we apply Lemma 1 with s = q = 2 so that, when K = 2, equation (16) becomes

$$\kappa_{3,p-2} \le \frac{1+\bar{\nu}_{p-2}}{1+\bar{\nu}_2}$$

Combining the above bounds we obtain the desired statement.

Proof of Theorem 5. Notice that

$$\bar{U}^{(r)} = I_{p-1} + (K-1) \cdot \bar{A}^{(r)}$$

Then, the eigenspace relative to -1/(K-1) of $\bar{\boldsymbol{A}}^{(r)}$ coincides with the null space of $\bar{\boldsymbol{U}}^{(r)}$. For simplicity, we will refer to the latter. Notice also, that $dim(Null(\bar{\boldsymbol{U}}^{(r)})) = dim(Null(\boldsymbol{U}^{(r)}))$, since $\bar{\boldsymbol{U}}^{(r)}$ is the product of $\boldsymbol{U}^{(r)}$ and full-rank matrices. In the remaining part of the proof, we will find $\sum_{\ell=1}^{K-1} C_{\ell}$ linearly independent eigenvalues in the null space of $\boldsymbol{U}^{(r)}$, hence proving Proposition 5. We split the proof in two parts.

1. Construction of $\{\boldsymbol{x}_{m}^{(\ell)}, \ell = 1, ..., K-1, m = 1, ..., C_{\ell}\} \subset Null(\boldsymbol{U}^{(r)})$. Consider w.l.o.g. the trivial permutation π equal to the identity. For each observation i and factor k, denote with $g_{k}[i]$ the unique non-null entry of the one-hot vector $\boldsymbol{z}_{i,k}$. Then, for each $\boldsymbol{x} \in \mathbb{R}^{p-1}, \boldsymbol{V}^{(r)}\boldsymbol{x} = \sum_{k=1}^{K} x_{g_{k}[i]}$. Thus

$$\boldsymbol{x}^{T} \boldsymbol{U}^{(r)} \boldsymbol{x} = || \boldsymbol{V}^{(r)} \boldsymbol{x} ||^{2} = \sum_{i=1}^{n} \left(\sum_{k=1}^{K} x_{g_{k}[i]} \right)^{2} = 0$$

happens if and only if $\sum_{k=1}^{K} x_{g_k[i]} = 0$ for each $i = 1, \ldots, N$. Hence, we have the equivalence

$$\boldsymbol{x} \in Null(\boldsymbol{U}^{(r)}) \qquad \Leftrightarrow \qquad \sum_{k=1}^{K} x_{g_k[i]} = 0, \ \forall i = 1, \dots, N.$$
 (30)

For any $\ell = 1, \ldots, K - 1$, denote with $P_1^{(\ell)}, \ldots, P_{C_{\ell}}^{(\ell)}$ and $Q_1^{(\ell)}, \ldots, Q_{C_{\ell}}^{(\ell)}$ the disconnected components in the bipartite graph restricted to the pair $(\ell, \ell + 1)$. In particular, the

 $P_m^{(\ell)}$'s are disjoint subsets of the levels in factor ℓ , and the same goes for $Q_m^{(\ell)}$'s. For any $\ell = 1, \ldots, K - 1$ and $m = 1, \ldots, C_1$, we define the vector $\boldsymbol{x}_m^{(\ell)} \in Null(\boldsymbol{U}^{(r)})$ as

$$\begin{cases} x_{m,j}^{(\ell)} = 1, & \text{if j belongs to } P_m \\ x_{m,j}^{(\ell)} = -1, & \text{if j belongs to } Q_m \\ x_{m,j}^{(\ell)} = 0, & \text{otherwise}. \end{cases}$$

Note that $\boldsymbol{x}_{m}^{(\ell)}$ satisfies (30) because, if an observation *i* does not involve $P_{m}^{(\ell)}$, then $x_{m,g_{k}^{k}[i]}^{(\ell)} = 0$ for each *k*. On the other hand, if $x_{m,g_{\ell}[i]}^{(\ell)} = 1$, then $x_{m,g_{\ell+1}[i]}^{(\ell)} = -1$ (by definition of connected components, an edge from $P_{m}^{(\ell)}$ can only connect to $Q_{m}^{(\ell)}$), while $x_{m,g_{k}[i]} = 0$ for each $k \neq \ell, \ell + 1$ (by construction).

2. $\{\boldsymbol{x}_m^{(\ell)}\}_{\ell,m}$ are linearly independent. Consider the linear combination

$$\sum_{\ell=1}^{K-1}\sum_{m=1}^{C_\ell}\beta_m^{(\ell)}\boldsymbol{x}_m^{(\ell)}=\boldsymbol{0}.$$

If we consider the first G_1 components (i.e. we restrict to the levels of the first factor), only $\{\boldsymbol{x}_m^{(1)}\}_{m=1}^{C_1}$ have non-zero entries on such components. Moreover, since they are supported on the disjoint sets $(P_m^{(1)})_{m=1}^{C_1}$, it implies that $\beta_1^{(1)} = \cdots = \beta_{C_1}^{(1)} = 0$. We now consider the G_2 components relative to the second factor. If we don't consider the previous eigenvectors, only $\{\boldsymbol{x}_m^{(2)}\}_{m=1}^{C_2}$ have non-zero entries on such components. Analogously to the previous case, we can conclude $\beta_1^{(2)} = \cdots = \beta_{C_2}^{(2)} = 0$. We can further extend this procedure, proving that $\beta_m^{(\ell)} = 0$ for each ℓ, m .

Proof of Theorem 6. Consider the matrix $\bar{A}^{(r)} \in \mathbb{R}^{p-1}$, Theorem 3 applied to it implies that $\bar{\nu}_1 = \ldots \bar{\nu}_{K-1} = -1/(K-1)$ and $\bar{\nu}_{p-1} = 1$. In particular, the invariant subspace associated to these eigenvalues is the K-dimensional space $W = \{x \in \mathbb{R}^{p-1}: x \text{ is constant factor-wise}\} = Span(w_1, \ldots, w_K)$, where $w_k = (\mathbf{0}_{G_1}^T, \ldots, \mathbf{1}_{G_k}^T, \ldots, \mathbf{0}_{G_K}^T)^T$. We denote its orthogonal with $\delta W = W^{\perp}$. With this notation, Theorem 6 becomes a straightforward consequence of Lemma 1 and 2 below.

Lemma 2. Under assumption (19), it holds that:

$$\max(|\bar{\nu}_K|, \bar{\nu}_{p-2}) = \max_{\boldsymbol{x} \in \delta W} \frac{||\bar{\boldsymbol{A}}^{(r)} \boldsymbol{x}||}{||\boldsymbol{x}||} < \frac{\lambda^*}{K-1}, \qquad (31)$$

where $||\mathbf{x}||$ denotes the standard Euclidean norm.

Proof. Denote for simplicity $\eta = \frac{\lambda^*}{\sqrt{K-1}}$. Denote instead with $\bar{A}^{(r)}[k,h] \in \mathbb{R}^{G_k \times G_h}$, the block of $\bar{A}^{(r)}$ relative to the pair of factors (k,h). It holds that:

$$\bar{\boldsymbol{A}}^{(k,h)} = (K-1) \cdot \begin{pmatrix} \boldsymbol{0} & \bar{\boldsymbol{A}}^{(r)}[k,h] \\ \bar{\boldsymbol{A}}^{(r)}[h,k] & \boldsymbol{0} \end{pmatrix},$$

where the multiplicative factor K - 1 comes from the fact that $\bar{A}^{(r)}$ is made of K cliques, hence each edge in the clique is connected to K - 1 levels in each other factor; however when restricting to the pair (k, h), it's connected to only one level. Moreover, by properties of bipartite adjacency matrix (Brito et al., 2022), it holds that

$$\max_{\lambda \in \sigma\left(\bar{\boldsymbol{A}}^{(k,h)}\right): |\lambda| \neq 1} |\lambda| = (K-1) \max_{\boldsymbol{y} \in \mathbb{R}^{G_h}: \; \boldsymbol{y}^T \mathbb{1}_{G_h} = 0} \frac{||\bar{\boldsymbol{A}}^{(r)}[k,h]\boldsymbol{y}||}{||\boldsymbol{y}||}.$$

Hence, (19) is equivalent to

$$\max_{\boldsymbol{y} \in \mathbb{R}^{G_h}: \ \boldsymbol{y}^T \mathbb{1}_{G_h} = 0} \frac{||\bar{\boldsymbol{A}}^{(r)}[k,h]\boldsymbol{y}||}{||\boldsymbol{y}||} < \frac{\eta}{K-1}$$

For a given vector $\boldsymbol{x} \in \delta W$, denote with $\boldsymbol{x}[k] \in \mathbb{R}^{G_k}$ the vector restricted to the factor k; by definition of δW , it holds that $\boldsymbol{x}[k]^T \mathbb{1}_{G_k} = 0$. Finally, taking $\boldsymbol{x} \in \delta W$, we have

$$\begin{split} |\bar{\boldsymbol{A}}^{(r)}\boldsymbol{x}||^{2} &= \sum_{k=1}^{K} \left\| \sum_{h \neq k} \bar{\boldsymbol{A}}^{(r)}[k,h]\boldsymbol{x}[h] \right\|^{2} \\ &\leq \sum_{k=1}^{K} \sum_{h \neq k} \left\| \bar{\boldsymbol{A}}^{(r)}[k,h]\boldsymbol{x}[h] \right\|^{2} \\ &\leq \frac{\eta^{2}}{(K-1)^{2}} \sum_{k=1}^{K} \sum_{h \neq k} \|\boldsymbol{x}[h]\|^{2} \\ &= \frac{\eta^{2}}{(K-1)^{2}} (K-1) ||\boldsymbol{x}||^{2} \\ &= \frac{\eta^{2}}{K-1} ||\boldsymbol{x}||^{2} \\ &= \frac{(\lambda^{*})^{2} ||\boldsymbol{x}||^{2}}{(K-1)^{2}} \,. \end{split}$$

B. Supplementary material for Section 5

B.1. Including multivariate fixed effects

Lemma 1 provides an upper bound on $\kappa_{K+1,p-2}(\bar{\boldsymbol{Q}})$, by interlacing the spectrum of $\bar{\boldsymbol{Q}}$ with the one of $\bar{\boldsymbol{Q}}^{(r)}$ (see proof of the lemma). A similar result can also be obtained for $\boldsymbol{\theta}_0 \in \mathbb{R}^{D_0}$, with $D_0 > 1$, i.e. when we include multivariate fixed effects. In which case, the upper bound would become

$$\kappa_{K+D_0,p-D_0-1}(\bar{\boldsymbol{Q}}) \le \frac{1+(K-1)\bar{\nu}_{p-D_0-1}}{1+(K-1)\bar{\nu}_K},$$

where $\bar{\nu}_K$ and $\bar{\nu}_{p-D_0-1}$ would be respectively the K-th smallest and second-largest eigenvalues of $\bar{A}^{(r)} \in \mathbb{R}^{(p-D_0)\times(p-D_0)}$. Provided that θ_0 is low-dimensional, the above inequality would give an upper bound on the effective condition number after the removal of few extreme eigenvalues.

B.2. Conjugate Gradient with Incomplete Cholesky preconditioner

In this section, we explore alternative preconditioning techniques to lower the number of CG iterations. In particular, we will refer to the Incomplete Cholesky (IC) preconditioner (Saad, 2003, Ch. 10). IC algorithm computes an approximate factorization $\boldsymbol{L}\boldsymbol{L}^T \approx \boldsymbol{Q}$, by restraining the support of \boldsymbol{L} to a subset S, which can be chosen a priori or on the run (see Algorithm 2). A standard choice for S is the lower-triangular support of \boldsymbol{Q} itself. Algorithm 2 with such choice of S is also referred to as zero fill-in incomplete Cholesky, which is a widely used CG preconditioner.

An important limitation of Algorithm 2 is that it may fail for general positive definite matrices. Indeed, by restraining the support of L, it may happen that $Q_{jj} - \sum_{k: (j,k) \in S} L_{jk}^2 < 0$. Several modifications have been proposed to prevent this issue, usually by scaling or shifting Q. Nevertheless, in the following numerical examples, we obtained the best results (in terms of rate of convergence of CG) by taking the absolute value of the previous quantity, hence forcing the continuation of Algorithm 2.

A lot of work has been done to refine the choice of S in order to improve the quality of the approximation, while preserving the computational efficiency of Algorithm 2. For example, Lin and Moré (1999) proposed a variation of the IC algorithm that selects Sdepending on the numerical entries of the matrix and satisfies $|S| \leq p \cdot n_Q$. Here, the choice of p allows controlling the density of the factorization, obtaining a more accurate solution at a price of additional memory and computational time. The results obtained with such preconditioner were comparable to the ones obtained with the standard IC preconditioner, hence we did not include them in this article.

Table 4 reports the comparison between the Jacobi (see Section 5.2) and the Incomplete Cholesky preconditioner. For both the American political elections survey data and the Instructor Evaluations data set, we display the average number of CG iterations (after an initial burn-in), obtained with Jacobi and IC preconditioner, respectively. While the IC preconditioned CG requires a smaller number of iterations, such reduction is not significant (at most a factor of 4) but comes at the additional price, in terms of time and memory complexity, of computing and inverting L. Finally, the overall cost of the two algorithms is basically equivalent in the case we have considered. However, the Jacobi preconditioning is easier to implement and does not incur in the problems of Algorithm 2.

B.3. Erdős-Rényi random bipartite graph

In this section, we extend Theorem 4 to the with K = 2 factors case and design given by an Erdős-Rényi bipartite random graph (i.e. each edge is missing independently at random with probability $1 - \pi$). The resulting bound on $\kappa_{3,p-2}(\bar{Q})$ is very similar to

Caso	Voter T	urnout	Students Evaluations	
Case	Jacobi	IChol	Jacobi	IChol
Random intercepts	36	19	35	16
Nested effect	53	31	94	37
Random slopes	70	27	150	17
2 way interactions	338	104	121	53
3 way interactions	445	108	126	54
Everything	517	140	262	67

Table 4: Average number of CG iterations with Jacobi and Incomplete Cholesky preconditioning for the data sets considered in Section 6. We considered $N = 70\,000$ observations.

the one of Theorem 4, however, in this case, we need an asymptotic lower bound on the expected degree $G_1\pi$ (and $G_2\pi$) to guarantee sufficient connectivity in the limit.

Theorem 7. Let $\mathbf{Q} = Diag(\mathbf{Q}) + \tau \mathbf{A}$ be the posterior precision matrix in (11). Let $\mathbf{A}^{(r)}$ be the adjacency matrix of an Erdős-Rényi bipartite random graph. Let $G_2 \geq G_1 \geq 1$ and $\pi = \pi(G_1)$ be such that

$$G_1 \pi = \Omega(\sqrt{G_1 \pi} \log^3(G_1)), \quad G_2 \pi = \Omega(\sqrt{G_2 \pi} \log^3(G_2)).$$

Then, with probability tending to 1 as G_1 tends to infinity

$$\kappa_{3,p-2}(\bar{\boldsymbol{Q}}) = \frac{\bar{\mu}_{p-2}}{\bar{\mu}_3} \le \frac{1+\epsilon}{1-\epsilon},$$

with

$$\epsilon = 2[2 + o(1)] \left(\frac{1}{\sqrt{G_1 \pi}} + \frac{1}{\sqrt{G_2 \pi}} + \sqrt{\frac{1}{G_1 \pi} + \frac{1}{G_2 \pi}} \right),$$

as $G_1 \to \infty$.

Proof. The proof of this result follows the same steps of the proof of Theorem 4, with the additional result from Ashcroft (2021). \Box

B.4. Pairwise connected components is not sufficient

Example 3. Consider a simple crossed random effects model with K = 3 factors with size $G_1 = G_2 = G_3 = 2$. Let N = 3, and let the observed levels be

$$g_1[1] = 1, \quad g_2[1] = 1, \quad g_3[1] = 2,$$

$$g_1[2] = 2, \quad g_2[2] = 2, \quad g_3[2] = 2,$$

$$g_1[3] = 2, \quad g_2[3] = 1, \quad g_3[3] = 1.$$

The corresponding $U^{(r)}$ will be

$$\boldsymbol{U}^{(r)} = \begin{bmatrix} 1 & 0 & 1 & 0 & 0 & 1 \\ 0 & 2 & 1 & 1 & 1 & 1 \\ 1 & 1 & 2 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 & 0 & 2 \end{bmatrix}$$

Notice that $dim(Null(U^{(r)})) = 3$ and that the following vectors generate the null-space of $U^{(r)}$:

In particular, the first two vectors are the directions identified by Theorem 3, while the third one is specific to the symmetry of this particular example. By construction also $\bar{\boldsymbol{U}}^{(r)}$ has a null-space of dimension 3. Finally, because of the relationship in (15), the corresponding eigenvalues of $\bar{\boldsymbol{A}}^{(r)}$ will satisfy $\bar{\nu}_1 = \bar{\nu}_2 = \bar{\nu}_3 = -1/2$.

To conclude, this example provides a design where the bipartite subgraph of any pair of factors is made of a unique connected component, but $\bar{\nu}_K = -\frac{1}{K-1}$.

C. Pseudocode of the algorithms

C.1. Cholesky factorization

Algorithm 1 shows a pseudocode of the algorithm that allows computing the Cholesky factorization of a given positive-definite matrix $\boldsymbol{Q} \in \mathbb{R}^{p \times p}$. A simple modification of Algorithm 1 allows computing an approximate factorization $\hat{\boldsymbol{L}}$, by restricting the support of \boldsymbol{L} to a given subset $S \subseteq \{(i, j): i \geq j, i = 1, \ldots, p, j = 1, \ldots, p\}$. The resulting algorithm is called incomplete Cholesky factorization (Golub and van Loan, 2013) and it is outlined in Algorithm 2.

Algorithm 1 Cholesky factorization. Algorithm 2 Incomplete Cholesky **Input:** $Q \in \mathbb{R}^{p \times p}$ symmetric and positive **Input:** $Q \in \mathbb{R}^{p \times p}$ symmetric, **S** sparsity set definite **Output:** $L \in \mathbb{R}^{p \times p}$ lower triangular s.t. **Output:** $L \in \mathbb{R}^{p \times p}$ lower triangular with $Q = LL^T$ $Supp(\mathbf{L}) \subset S$ 1: for $j \in \{1, ..., p\}$ do 1: for $j \in \{1, ..., p\}$ do $L_{jj} \leftarrow \sqrt{Q_{jj} - \sum_{k: (j,k) \in S} L_{jk}^2}$ for $i > j: (i, j) \in S$ do $L_{jj} \leftarrow \sqrt{Q_{jj} - \sum_{k: (j,k) \in S} L_{jk}^2}$ for $i \in \{j + 1, \dots, N\}$ do 2: 2: 3: 3: $L_{ij} \leftarrow Q_{ij}/L_{jj}$ $L_{ij} \leftarrow Q_{ij}/L_{jj}$ 4: 4: for $k \in \{1, ..., j-1\}$ do for k < j: $(i, k), (j, k) \in S$ do 5: 5: $L_{ij} \leftarrow L_{ij} - (L_{ik} \cdot L_{jk}/L_{jj})$ $L_{ij} \leftarrow L_{ij} - (L_{ik} \cdot L_{jk}/L_{jj})$ 6: 6: end for end for 7: 7: end for end for 8: 8: 9: end for 9: end for 10: return L10: return L

C.2. Sampling with Cholesky factorization

For a given Cholesky factor \boldsymbol{L} , sampling from a Gaussian distribution with precision matrix $\boldsymbol{Q} = \boldsymbol{L}\boldsymbol{L}^T$ is straightforward. E.g. one can obtain a sample $\boldsymbol{\theta} \sim \mathcal{N}(\boldsymbol{Q}^{-1}\boldsymbol{m}, \boldsymbol{Q}^{-1})$, by first sampling $\boldsymbol{z} \sim \mathcal{N}(\boldsymbol{0}, \boldsymbol{I}_p)$, and then solving the following two linear system

$$Lw = m$$
, $L^T \theta = w + z$.

Specifically, one needs to solve a lower and an upper-triangular linear system respectively, which can be done with $\mathcal{O}(n_L)$ cost via forward and backward substitution.

C.3. Conjugate gradient

For the conjugate gradient algorithm, we refer to the formulation of Saad (2003). Namely, we refer to Algorithm 6.17 and to Algorithm 9.1, for the preconditioned version. For simplicity, we report them in Algorithm 3 and 4 respectively.

If we only consider vectorial operations, Algorithm 3 requires the computation of three scalar products, three linear combinations of vectors and a matrix-vector multiplication, accounting for a total of $4p + 2n_{\boldsymbol{Q}}$ flops. The preconditioned variant, only requires the additional cost of evaluating $\boldsymbol{M}^{-1}\boldsymbol{z}$, which, for Jacobi preconditioning, accounts for p flops.

Algorithm 3 Conjugate Gradient	Algorithm 4 Preconditioned CG
Input: $\boldsymbol{Q} \in \mathbb{R}^{p \times p}$ symmetric positive	Input: $\boldsymbol{Q} \in \mathbb{R}^{p \times p}$ symmetric positive
definite, $\boldsymbol{b} \in \mathbb{R}^p$, $\epsilon > 0$ desired accuracy.	definite, $\boldsymbol{b} \in \mathbb{R}^p$, $\epsilon > 0$, \boldsymbol{M} preconditioner.
Output: Approx. solution of $Qx = b$.	Output: Approx. solution of $Qx = b$.

1: $x_0 = 0, r_0 = b, z_0 = M^{-1}r_0, p_0 = z_0.$ 1: $x_0 = 0, r_0 = b, p_0 = r_0.$ 2: for $j = 0, 1, \ldots$ until $||\boldsymbol{r}_j||_2 < \epsilon$, do 2: for j = 0, 1, ... until $||r_j||_2 < \epsilon$, do $\mathbf{r} \ j = 0, 1, \dots \text{ until } || i$ $\alpha_j = \frac{\mathbf{r}_j^T \mathbf{r}_j}{\mathbf{p}_j^T \mathbf{Q} \mathbf{p}_j}$ $\mathbf{x}_{j+1} = \mathbf{x}_j + \alpha_j \mathbf{p}_j$ $\mathbf{r}_{j+1} = \mathbf{r}_j - \alpha_j \mathbf{Q} \mathbf{p}_j$ $\beta_j = \frac{\mathbf{r}_{j+1}^T \mathbf{r}_{j+1}}{\mathbf{r}_j^T \mathbf{r}_j}$ $\mathbf{p}_{j+1} = \mathbf{r}_{j+1} + \beta_j \mathbf{p}_j$ and for $lpha_j = rac{oldsymbol{r}_j^T oldsymbol{z}_j}{oldsymbol{p}_j^T oldsymbol{Q} oldsymbol{p}_j} \ oldsymbol{x}_{j+1} = oldsymbol{x}_j + lpha_j oldsymbol{p}_j$ 3: 3: 4: 4: $\boldsymbol{r}_{j+1} = \boldsymbol{r}_j - \alpha_j \boldsymbol{Q} \boldsymbol{p}_j$ 5: 5: $egin{aligned} egin{aligned} egi$ 6: 6: 7:7:8: end for $\boldsymbol{p}_{j+1} = \boldsymbol{z}_{j+1} + \beta_j \boldsymbol{p}_j$ 8: 9: end for 9: return x_{j+1} 10: return x_{j+1}

C.4. Polya-Gamma augmented Gibbs sampler

Algorithm 5 PG augmented Gibbs sampler
Input: hyperparameters m_0 , α_k , Φ_k ; vector of observations y ; design matrix V ;
initial values $\boldsymbol{\theta}^{(0)}, \ \boldsymbol{\Omega}^{(0)}, \ \boldsymbol{T}_k^{(0)}$
for $t = 1 : T$ do
Sample $\boldsymbol{\theta}^{(t)} \sim p(\boldsymbol{\theta} \mid \boldsymbol{y}, \boldsymbol{V}, \boldsymbol{\Omega}^{(t-1)}, \{\boldsymbol{T}_k^{(t-1)}\}_{k=1}^K)$ according to (23)
Sample $\omega_i^{(t)} \sim p(\omega_i \mid \boldsymbol{y}, \boldsymbol{V}, \boldsymbol{\theta}^{(t)})$, for each $i = 1, \dots, N$, according to (24)
Sample $\boldsymbol{T}_{k}^{(t)} \sim p(\boldsymbol{T}_{k} \mid \boldsymbol{y}, \boldsymbol{V}, \boldsymbol{\theta}^{(t)})$, for each $k = 1, \dots, K$, according to (25)
end for
Output: MCMC samples $\{(\boldsymbol{\theta}^{(t)}, \boldsymbol{\Omega}^{(t)}, , \{\boldsymbol{T}_k^{(t)}\}_{k=1}^K); t = 0, \dots, T\}.$

D. Additional Figures

D.1. Sparsity structure of Q resulting from Example 1



Figure 3: Precision matrix \boldsymbol{Q} of $\mathcal{L}(\boldsymbol{\theta} \mid \boldsymbol{y}, \boldsymbol{g})$ induced by the structured design of Example 1 with G = 20, d = 3 and default ordering $(\theta_{1,1}, \ldots, \theta_{1,G}, \theta_{2,1}, \ldots, \theta_{2,G}, \theta_0)$.

D.2. Graphical representation of the graph in Example 3



Figure 4: Conditional independence graph of Example 3. Obtained for $G_1 = G_2 = G_3 = 3$, and N = 9.