

# SHIFTED QUASI-SYMMETRIC FUNCTIONS AND THE HOPF ALGEBRA OF PEAK FUNCTIONS

NANTEL BERGERON, STEFAN MYKYTIUK, FRANK SOTTILE,  
AND STEPHANIE VAN WILLIGENBURG

**SUMMARY.** In his work on  $P$ -partitions, Stembridge defined the algebra of peak functions  $\Pi$ , which is both a subalgebra and a retraction of the algebra of quasi-symmetric functions. We show that  $\Pi$  is closed under coproduct, and therefore a Hopf algebra, and describe the kernel of the retraction. Billey and Haiman, in their work on Schubert polynomials, also defined a new class of quasi-symmetric functions — shifted quasi-symmetric functions — and we show that  $\Pi$  is strictly contained in the linear span  $\Xi$  of shifted quasi-symmetric functions. We show that  $\Xi$  is a coalgebra, and compute the rank of the  $n$ th graded component.

**RÉSUMÉ.** Dans ses travaux sur les  $P$ -partitions, Stembridge définit l'algèbre  $\Pi$  des fonctions de pics. Cette algèbre peut être vue comme une sous-algèbre ou un quotient de l'algèbre des fonctions quasi-symétriques. Nous montrons ici que  $\Pi$  est fermée sous le coproduit, et est donc une algèbre de Hopf. Nous décrivons aussi le noyau du quotient ci-dessus. D'autre part, dans leurs travaux sur les polynômes de Schubert, Billey et Haiman ont défini une nouvelle classe de fonctions quasi-symétriques: les fonctions quasi-symétrique décalé. Nous montrons que  $\Pi$  est strictement contenue dans l'espace linéaire  $\Xi$  des fonctions quasi-symétrique gauchis. Puis nous montrons que  $\Xi$  est une coalgèbre et calculons les dimensions des composantes de degré  $n$ .

## 1. INTRODUCTION

Schur  $Q$  functions first arose in the study of projective representations of  $S_n$  [8]. Since then they have appeared in variety of contexts including the representations of Lie superalgebras [9] and cohomology classes dual to Schubert cycles in isotropic Grassmanians [4, 7]. While studying the duality between skew Schur  $P$  and  $Q$  functions and their connection to the Schubert calculus of isotropic flag manifolds, we were led to their quasi-symmetric analogues: the *peak functions* of Stembridge [10]. We show that *the linear span of peak functions is a Hopf algebra* (Theorem 2.2). We also show that these peak functions are contained in the strictly larger set of *shifted quasi-symmetric functions* (Theorem 3.6) introduced by Billey and Haiman [1]. We remark that the quasi-symmetric functions here are not any apparent specialization of the quasi-symmetric  $q$ -analogues of Hivert [3].

---

*Date:* March 10, 2018.

*1991 Mathematics Subject Classification.* 05E15, 05E05, 05A15, 16W30.

*Key words and phrases.* Pieri, graded operation, poset, quasi-symmetric functions.

Bergeron supported in part by NSERC.

Sottile supported in part by NSF grant DMS-9701755 and NSERC grant OGP0170279.

van Willigenburg is supported in part by the Leverhulme Trust.

From extensive calculations, we believe that the set of all shifted quasi-symmetric functions form a Hopf algebra, but at present we can only show that:

*The set of all shifted quasi-symmetric functions forms a graded coalgebra whose  $n$ th graded component has rank  $\pi_n$ , where  $\pi_n$  is given by the recurrence*

$$\pi_n = \pi_{n-1} + \pi_{n-2} + \pi_{n-4},$$

*with initial conditions  $\pi_1 = 1$ ,  $\pi_2 = 1$ ,  $\pi_3 = 2$ ,  $\pi_4 = 4$ .*

We shall prove this result (Theorems 3.2 and 4.3) and in addition shall establish some other properties of these functions.

A composition  $\alpha = [\alpha_1, \alpha_2, \dots, \alpha_k]$  of a positive integer  $n$  is an ordered list of positive integers whose sum is  $n$ . We denote this by  $\alpha \vDash n$ . We call the integers  $\alpha_i$  the *components* of  $\alpha$ , and denote the number of components in  $\alpha$  by  $k(\alpha)$ . There exists a natural one-to-one correspondence between compositions of  $n$  and subsets of  $[n-1]$ . If  $A = \{a_1, a_2, \dots, a_{k-1}\} \subset [n-1]$ , where  $a_1 < a_2 < \dots < a_{k-1}$ , then  $A$  corresponds to the composition,  $\alpha = [a_1 - a_0, a_2 - a_1, \dots, a_k - a_{k-1}]$ , where  $a_0 = 0$  and  $a_k = n$ . For ease of notation, we shall denote the set corresponding to a given composition  $\alpha$  by  $I(\alpha)$ . For compositions  $\alpha$  and  $\beta$  we say that  $\alpha$  is a *refinement* of  $\beta$  if  $I(\beta) \subset I(\alpha)$ , and denote this by  $\alpha \preceq \beta$ .

For any composition  $\alpha = [\alpha_1, \alpha_2, \dots, \alpha_k]$  we denote by  $M_\alpha$  the *monomial quasi-symmetric function* [2]

$$M_\alpha = \sum_{i_1 < i_2 < \dots < i_k} x_{i_1}^{\alpha_1} \dots x_{i_k}^{\alpha_k}.$$

We define  $M_0 = 1$ , where 0 denotes the unique empty composition of 0. We denote by  $F_\alpha$  the *fundamental quasi-symmetric function* [2]

$$F_\alpha = \sum_{\alpha \preceq \beta} M_\beta.$$

**Definition 1.1.** *For any subset  $A \subset [n-1]$ , let  $A+1$  be the subset of  $\{2, \dots, n\}$  formed from  $A$  by adding 1 to each element of  $A$ . Let  $\alpha \vDash n$ . Then we define*

$$\theta_\alpha = \sum_{\substack{\beta \vDash n \\ I(\alpha) \subset I(\beta) \cup I(\beta)+1}} 2^{k(\beta)} M_\beta.$$

This is the natural extension of the definition of peak functions given in [10].

**Example 1.2.** *We shall often omit the brackets that surround the components of a composition.*

*If  $\alpha = 21$ , then  $I(\alpha) = \{2\}$ , and  $I(\alpha) + 1 = \{3\}$ . Hence*

$$\theta_{21} = 4M_{21} + 4M_{12} + 8M_{111}.$$

Let  $\Sigma^n$  be the  $\mathbb{Z}$ -module of quasi-symmetric functions spanned by  $\{M_\alpha\}_{\alpha \vDash n}$  and let  $\Sigma = \bigoplus_{n \geq 0} \Sigma^n$  be the graded  $\mathbb{Z}$ -algebra of quasi-symmetric functions. This is a Hopf algebra [5] with coproduct given by

$$\Delta(M_\alpha) = \sum_{\alpha = \beta \cdot \gamma} M_\beta \otimes M_\gamma,$$

where  $\beta \cdot \gamma$  is the concatenation of compositions  $\beta$  and  $\gamma$ .

**Example 1.3.**  $\Delta(M_{32}) = 1 \otimes M_{32} + M_3 \otimes M_2 + M_{32} \otimes 1$ .

We compute the coproduct of the functions  $\theta_\alpha$ .

**Lemma 1.4.** *For any composition  $\alpha \vDash n$  we have that*

$$(1) \quad \Delta(\theta_\alpha) = \sum \theta_{\varepsilon \cdot a} \otimes \theta_{\phi(b \cdot \zeta)}$$

where the sum is over all ways of writing  $\alpha$  as  $\varepsilon \cdot (a+b) \cdot \zeta$ , that is, the concatenation of compositions  $\varepsilon$  and  $\zeta$ , and a component of  $\alpha$  written as the sum of numbers  $a, b \geq 0$ . Also  $\phi(b \cdot \zeta) = [1 + \zeta_1, \zeta_2, \dots]$  if  $b = 1$  and  $b \cdot \zeta$  otherwise.

We shall use this result to show that certain subsets of functions  $\theta_\alpha$  span coalgebras (Theorems 2.2 and 3.2).

*Proof.* Definition 1.1 is equivalent to

$$\theta_\alpha = \sum_{\substack{\beta \vDash n \\ \beta^* \preceq \alpha}} 2^{k(\beta)} M_\beta,$$

where  $\beta^*$  is the refinement of  $\beta$  obtained by replacing all components  $\beta_i > 1$ , for  $i > 1$ , by  $[1, \beta_i - 1]$ . Thus the LHS of equation (1) is equal to

$$(2) \quad \sum_{\substack{\beta \vDash n \\ \beta^* \preceq \alpha \\ \beta = \gamma \cdot \delta}} 2^{k(\beta)} M_\gamma \otimes M_\delta = \sum_{\substack{\gamma \cdot \delta \vDash n \\ (\gamma \cdot \delta)^* \preceq \alpha}} 2^{k(\gamma)} M_\gamma \otimes 2^{k(\delta)} M_\delta.$$

Let  $2^{k(\gamma)} M_\gamma \otimes 2^{k(\delta)} M_\delta$  be a term of this sum, with  $\gamma \vDash m$ . This term can only appear in one summand on the RHS of equation (1), namely  $\theta_{\varepsilon \cdot a} \otimes \theta_{\phi(b \cdot \zeta)}$  with  $\varepsilon \cdot a \vDash m$ . To show that it does indeed appear, we need to prove that  $\gamma^* \preceq \varepsilon \cdot a$  and  $\delta^* \preceq \phi(b \cdot \zeta)$ . Let  $\delta^{**}$  be the refinement of  $\delta^*$  obtained by replacing the part  $\delta_1$  by  $[1, \delta_1 - 1]$  if  $\delta_1 > 1$ . We have that

$$\gamma^* \cdot \delta^{**} = (\gamma \cdot \delta)^* \preceq \varepsilon \cdot (a+b) \cdot \zeta,$$

which implies that  $\gamma^* \preceq \varepsilon \cdot a$ , and  $\delta^{**} \preceq b \cdot \zeta \preceq \phi(b \cdot \zeta)$ .

If  $\delta_1 = 1$  then  $\delta^* = \delta^{**} \preceq \phi(b \cdot \zeta)$ . However, if  $\delta_1 > 1$  then there are two possible cases: either  $\delta_1 \leq b$ , or  $b = 1$  and  $\delta_1 - 1 \leq \zeta_1$ . In the former case  $\delta^* \preceq b \cdot \zeta = \phi(b \cdot \zeta)$ , while in the latter,  $\delta_1 \preceq 1 + \zeta_1$ , whence  $\delta^* \preceq [1 + \zeta_1, \zeta_2, \dots] = \phi(b \cdot \zeta)$ .

Conversely, let  $2^{k(\gamma)}M_\gamma \otimes 2^{k(\delta)}M_\delta$  be a term belonging to a tensor  $\theta_{\varepsilon \cdot a} \otimes \theta_{\phi(b \cdot \zeta)}$  on the RHS of equation (1). To show that it appears in equation (2) we must prove that  $(\gamma \cdot \delta)^* \preceq \varepsilon \cdot (a + b) \cdot \zeta$ . We have that  $\gamma^* \preceq \varepsilon \cdot a$  and  $\delta^* \preceq \phi(b \cdot \zeta)$ , which imply that

$$(\gamma \cdot \delta)^* = \gamma^* \cdot \delta^{**} \preceq \gamma^* \cdot \delta^* \preceq \varepsilon \cdot a \cdot \phi(b \cdot \zeta).$$

If  $b > 1$  then

$$(\gamma \cdot \delta)^* \preceq \varepsilon \cdot a \cdot \phi(b \cdot \zeta) = \varepsilon \cdot a \cdot b \cdot \zeta \preceq \varepsilon \cdot (a + b) \cdot \zeta.$$

If  $b = 1$  then  $\delta^* \preceq \phi(b \cdot \zeta) = [1 + \zeta_1, \zeta_2, \dots]$  implies that

$$\delta^{**} = [1, \dots] \preceq [1, \zeta_1, \dots] = b \cdot \zeta.$$

Therefore,

$$(\gamma \cdot \delta)^* = \gamma^* \cdot \delta^{**} \preceq \varepsilon \cdot a \cdot b \cdot \zeta \preceq \varepsilon \cdot (a + b) \cdot \zeta$$

as desired.  $\square$

## 2. THE PEAK HOPF ALGEBRA

**Definition 2.1.** For any composition  $\alpha = [\alpha_1, \alpha_2, \dots, \alpha_k]$  we say that  $\theta_\alpha$  is a peak function if  $\alpha_i = 1 \Rightarrow i = k$ .

Observe that if  $\theta_\alpha$  is a peak function and  $\alpha \vDash n$ , then  $I(\alpha) \subset \{2, \dots, n-1\}$  such that no two  $i$  in  $I(\alpha)$  are consecutive.

Let  $\Pi^n$  be the  $\mathbb{Z}$ -module spanned by all peak functions  $\theta_\alpha$ ,  $\alpha \vDash n$ , and let  $\Pi = \bigoplus_{n \geq 0} \Pi^n$ . This was studied by Stembridge [10] who showed that the peak functions are F-positive, are closed under product, and form a basis for  $\Pi$ , and so the rank of  $\Pi^n$  is the  $n$ th Fibonacci number. In addition we also know the following about the algebra of peaks,  $\Pi$ .

**Theorem 2.2.**  $\Pi$  is closed under coproduct.

*Proof.* If all components of a composition  $\alpha$ , except perhaps the last, are greater than 1, then the same is true for all compositions  $\varepsilon \cdot a$  and  $\phi(b \cdot \zeta)$  appearing in the RHS of equation (1).  $\square$

Let  $\Theta$  be the  $\mathbb{Z}$ -linear map from  $\Sigma$  to  $\Pi$  defined by  $\Theta(F_\alpha) = \theta_{\Lambda(\alpha)}$ , where  $\Lambda(\alpha)$  is the composition formed from  $\alpha = [\alpha_1, \alpha_2, \dots, \alpha_k]$  by adding together adjacent components  $\alpha_i, \alpha_{i+1}, \dots, \alpha_{i+j}$  where  $\alpha_{i+l} = 1$  for  $l = 0, \dots, j-1$ , and either  $\alpha_{i+j} \neq 1$ , or  $i + j = k$ .

**Example 2.3.** If  $\alpha = 31125111$  then  $\Lambda(\alpha) = 3453$ .

Stembridge [10] showed that  $\Theta : \Sigma \rightarrow \Pi$  is a graded surjective ring homomorphism, and was an analogue of the retraction from the algebra of symmetric functions to Schur  $Q$  functions. It is clear from our proof above that this morphism is in fact a Hopf homomorphism. We can describe the kernel of  $\Theta$  as follows.

**Lemma 2.4.** *The non-zero differences  $F_\alpha - F_{\Lambda(\alpha)}$  form a basis of the kernel of  $\Theta$ .*

*Proof.* Each difference  $F_\alpha - F_{\Lambda(\alpha)}$  is in the kernel of  $\Theta$  as  $\Theta(F_\alpha - F_{\Lambda(\alpha)}) = 0$  since  $\Lambda(\Lambda(\alpha)) = \Lambda(\alpha)$ . In addition, the non-zero differences are linearly independent as they have different leading terms. Letting  $f_n$  denote the  $n$ th Fibonacci number, there are  $2^{n-1} - f_n$  such differences, and since

$$\begin{aligned} \dim \ker \Theta &= \dim \Sigma^n - \dim \Pi^n \\ &= 2^{n-1} - f_n, \end{aligned}$$

our result follows. □

### 3. THE COALGEBRA OF SHIFTED QUASI-SYMMETRIC FUNCTIONS

**Definition 3.1.** *For any composition  $\alpha = [\alpha_1, \alpha_2, \dots, \alpha_k] \vDash n$  we say that  $\theta_\alpha$  is a shifted quasi-symmetric function (sqs-function) if  $n \leq 1$  or  $\alpha_1 > 1$ .*

Observe that if  $\theta_\alpha$  is an sqs-function and  $\alpha \vDash n$ , then  $I(\alpha) \subset \{2, \dots, n-1\}$ .

For integers  $n \geq 0$ , let  $\Xi^n$  be the  $\mathbb{Z}$ -module spanned by all sqs-functions  $\theta_\alpha$ ,  $\alpha \vDash n$ , and let  $\Xi = \bigoplus_{n \geq 0} \Xi^n$ .


**Theorem 3.2.**  *$\Xi$  is closed under coproduct.*

*Proof.* If the first component of a composition  $\alpha$  is greater than 1, then the same is true for all compositions  $\varepsilon \cdot a$  and  $\phi(b \cdot \zeta)$  appearing in the RHS of equation (1). □

Unlike peak functions [10], sqs-functions are not  $F$ -positive since

$$\theta_{211} = F_{22} + F_{112} + 2F_{121} + F_{211} - F_{1111}.$$

**Definition 3.3.** *For any composition,  $\alpha \vDash n$ , we define the complement  $\alpha^c$  of  $\alpha$  to be the composition for which  $I(\alpha^c) = (I(\alpha))^c$ , the set complement of  $I(\alpha)$  in  $[n-1]$ . We define the graph  $G(\alpha)$  of  $\alpha$  to be the graph obtained from*




by removing the edge  $(i, i+1)$  if and only if  $i \in I(\alpha)$ .

Observe that  $G(\alpha^c)$  contains the edge  $(i, i+1)$  if and only if this edge is not contained in  $G(\alpha)$ . These graphs will be used later to simplify the proof of Theorem 3.6.


Let a *word* of length  $n$  be any  $n$ -tuple,  $w_1 w_2 \dots w_n$ , and let a *binary word* of length  $n$  be a word  $w_1 w_2 \dots w_n$  such that  $w_i \in \{0, 1\}$  for all  $i$ . For  $2 \leq i \leq n-1$ , let us denote by  $3^{(i)}$  the composition  $[1^{i-2}, 3, 1^{n-i-1}]$  of  $n$ . For some subset  $S \subset \{2, \dots, n-1\}$ , let us denote by  $\bigwedge_{i \in S} 3^{(i)}$  the composition of  $n$  for which  $G(\bigwedge_{i \in S} 3^{(i)})$  has an edge

between vertices  $i$  and  $i + 1$  if and only if an edge exists between vertices  $i$  and  $i + 1$  in  $G(3^{(i)})$  for some  $i \in S$ .


**Example 3.4.** Let  $S = \{2, 3\} \subset [3]$ . Then  $G(3^{(2)})$  is



and  $G(3^{(3)})$  is



hence  $G(\bigwedge_{i \in S} 3^{(i)})$  is



so  $\bigwedge_{i \in S} 3^{(i)}$  is the composition 4.

**Definition 3.5.** [1] Let  $\alpha$  be a composition of  $n$ . Let  $\mathcal{A}(I(\alpha))$  denote the set of all sequences  $j_1 \leq j_2 \leq \dots \leq j_n$  in  $\mathbb{N}$  such that we do not have  $j_{i-1} = j_i = j_{i+1}$  for any  $i \in I(\alpha)$ . The shifted quasi-symmetric function  $\theta_\alpha^{BH}$  is given by

$$\theta_\alpha^{BH} = \sum_{\substack{J = (j_1, \dots, j_n) \\ j_1 \leq \dots \leq j_n \\ J \in \mathcal{A}(I(\alpha))}} 2^{|j|} x_{j_1} \dots x_{j_n},$$

where  $|j|$  denotes the number of distinct values  $j_i$  in  $J$ .

**Theorem 3.6.** For any sqs-function  $\theta_\alpha$  we have that  $\theta_\alpha = \theta_\alpha^{BH}$ .

*Proof.* For each  $i \in I(\alpha) \subset [n - 1]$ ,  $j_{i-1} = j_i = j_{i+1}$  is forbidden in any monomial

$$x_{j_1} x_{j_2} \dots x_{j_i} \dots x_{j_n}$$

appearing as a summand of the function  $\theta_\alpha^{BH}$ . This is equivalent to saying that  $M_\beta$  is a summand of  $\theta_\alpha^{BH}$  if and only if  $G(3^{(i)}) \not\subset G(\beta)$  for all  $i \in I(\alpha)$ . Therefore at least one of  $i - 1$  or  $i$  must be the largest label of a vertex in a connected component in  $G(\beta)$ .

Now when going from compositions of  $n$  to subsets of  $[n - 1]$  we can do so using our graphs,  $G$ . All we have to do is list the label of the vertex that is the largest in each connect component, not listing  $n$ . We call these vertices the *end-points*. We are now in a position to prove the equivalence of Definitions 1.1 and 3.5 for sqs-functions.

The powers of 2 agree so we need only show that the indices of summation do too. To see this, take any sqs-function  $\theta_\alpha$  and let  $i \in I(\alpha)$ . Then  $M_\beta$  is a summand in  $\theta_\alpha^{BH}$  if at least one of  $i - 1$  or  $i$  is an end-point in  $G(\beta)$ . Therefore  $i$  or  $i - 1$  belongs to  $I(\beta)$ , and  $M_\beta$  is a summand of  $\theta_\alpha$ . Conversely, if  $M_\beta$  is a summand of  $\theta_\alpha$ , then this implies that for each  $i \in I(\alpha)$ , we have that  $i - 1$  or  $i$  belongs to  $I(\beta)$ , so one of  $i - 1$  or  $i$  is an end-point in  $G(\beta)$ , so  $M_\beta$  is a summand of  $\theta_\alpha^{BH}$ .  $\square$

4. A BASIS FOR  $\Xi$ 

**Definition 4.1.** Let  $\theta_\alpha$  be an sqs-function and  $\alpha \vDash n$ . We define an internal peak  $i \in I(\alpha)$  such that  $i - 1, i + 1 \notin I(\alpha)$ , and  $i \in \{3, \dots, n - 2\}$ .

**Remark** Observe that the occurrence of an internal peak in the  $i$ th position in  $I(\alpha) = \{w_1, w_2, \dots\}$ , where  $w_1 < w_2 < \dots$ , is equivalent to having two components of  $\alpha$ , say  $\alpha_i, \alpha_{i+1}$  such that  $\alpha_{i+1} \geq 2$ , and  $\alpha_i \geq 2$  if  $i \neq 1$ , or  $\alpha_i \geq 3$  if  $i = 1$ .

We can now describe the basis of  $\Xi$  as follows.

**Theorem 4.2.** The coalgebra  $\Xi$  has a basis consisting of all sqs-functions  $\theta_\alpha$  where  $I(\alpha)$  contains no internal peak.

We sketch the proof of Theorem 4.2 later.

**Theorem 4.3.** The rank of  $\Xi^n$  is given by the recurrence

$$\pi_n = \pi_{n-1} + \pi_{n-2} + \pi_{n-4},$$

with initial conditions  $\pi_1 = 1, \pi_2 = 1, \pi_3 = 2, \pi_4 = 4$ .

This recurrence was suggested by a superseeker query [6].

*Proof.* By direct calculation we obtain that  $\pi_1 = 1, \pi_2 = 1, \pi_3 = 2$ , and  $\pi_4 = 4$ .

To obtain our recurrence, we observe that for each sqs-function,  $\theta_\alpha$  where  $\alpha \vDash n$ , we can encode  $I(\alpha)$  as a binary word of length  $n - 2$ , by placing a 1 in position  $i - 1$  if  $i$  is contained in  $I(\alpha)$ , and 0 otherwise. By this one-to-one correspondence we see that  $I(\alpha)$  contains no internal peak if its corresponding binary word does not contain 010 as a subword.

We therefore count binary words of length  $n$  that avoid the subword 010. Appending either 1 or 0 to such a binary word of length  $n - 1$  gives one of length  $n$ , provided that we have not created the subword 010 in the last three positions. Let  $a_n, b_n, c_n$ , and  $d_n$  enumerate those binary words of length  $n - 2$  that avoid the subword 010 and end in, respectively 00, 01, 10, and 11. We then obtain the following 4 simultaneous recursions.

$$a_n = a_{n-1} + c_{n-1}, \quad b_n = a_{n-1} + c_{n-1}, \quad c_n = d_{n-1}, \quad d_n = b_{n-1} + d_{n-1}.$$

Clearly the number of  $I(\alpha)$ s in  $[n - 1]$  with no internal peaks is given by

$$\pi_n = a_n + b_n + c_n + d_n,$$

However by substituting in our recurrences we obtain

$$\begin{aligned}
\pi_n &= a_n + b_n + c_n + d_n \\
&= 2a_{n-1} + b_{n-1} + 2c_{n-1} + 2d_{n-1} \\
&= \pi_{n-1} + a_{n-1} + c_{n-1} + d_{n-1} \\
&= \pi_{n-1} + a_{n-2} + b_{n-2} + c_{n-2} + 2d_{n-2} \\
&= \pi_{n-1} + \pi_{n-2} + d_{n-2} \\
&= \pi_{n-1} + \pi_{n-2} + b_{n-3} + d_{n-3} \\
&= \pi_{n-1} + \pi_{n-2} + a_{n-4} + b_{n-4} + c_{n-4} + d_{n-4} \\
&= \pi_{n-1} + \pi_{n-2} + \pi_{n-4}.
\end{aligned}$$

□

We say that  $M_\beta$  is a maximal term of  $\theta_\alpha$  if for any  $\gamma$  higher in the partial order of compositions  $M_\gamma$  is not a summand of  $\theta_\alpha$ . The following lemma is stated without proof.

**Lemma 4.4.** *Let  $\theta_\alpha$  be an sqs-function. Consider the collection  $S$  of all possible sets derived from  $I(\alpha)$  by adding either  $i-1$  or  $i+1$  to  $I(\alpha)$  for all internal peaks  $i \in I(\alpha)$ . If  $M_\beta$  is a maximal term of  $\theta_\alpha$ , then  $\beta$  is derived from*

$$\bigwedge_{\substack{i \in (I(\tilde{\alpha}))^c \\ I(\tilde{\alpha}) \in S}} 3^{(i)}$$

by adding adjacent components equal to 1 together to give a component equal to 2 as often as possible.

**Lemma 4.5.** *Let  $\theta_\alpha$  be an sqs-function, and let  $I(\alpha)$  have an internal peak in the  $j$ th position, then we have the following linear relation*

$$\begin{aligned}
\theta_\alpha &= \theta_{[\alpha_1, \dots, \alpha_j-1, 1, \alpha_{j+1}, \dots, \alpha_k]} + \theta_{[\alpha_1, \dots, \alpha_j, 1, \alpha_{j+1}-1, \dots, \alpha_k]} \\
&\quad - \theta_{[\alpha_1, \dots, \alpha_j-1, 1, 1, \alpha_{j+1}-1, \dots, \alpha_k]}.
\end{aligned}$$

*Proof.* By Definition 3.5 we have that the leading terms of  $\theta_\alpha$  determine the other summands that belong to  $\theta_\alpha$ . Hence by Lemma 4.4 it follows that the summands of  $\theta_\alpha$  will be the union of the summands of  $\theta_{[\alpha_1, \dots, \alpha_j-1, 1, \alpha_{j+1}, \dots, \alpha_k]}$  and  $\theta_{[\alpha_1, \dots, \alpha_j, 1, \alpha_{j+1}-1, \dots, \alpha_k]}$ . However, those summands that appear in both will be duplicated. By definition these will be the summands of  $\theta_{[\alpha_1, \dots, \alpha_j-1, 1, 1, \alpha_{j+1}-1, \dots, \alpha_k]}$ , and the result follows. □

*Sketch of proof of Theorem 4.2.* From our relation in Lemma 4.5, it follows that any  $\theta_\alpha$  can be rewritten as a linear combination of functions  $\theta_{\tilde{\alpha}}$ , where  $I(\tilde{\alpha})$  contains no internal peaks. In addition, by Lemma 4.4 and definition 3.5 we have that the set



of all sqs-functions  $\theta_\alpha$  where  $I(\alpha)$  contains no internal peaks is linearly independent and thus form a basis for  $\Xi$ .  $\square$

## REFERENCES

- [1] S. Billey, and M. Haiman, *Schubert polynomials for the classical groups*, Journal of AMS, **8** (1995) 443–482.
- [2] I. Gessel, *Multipartite  $P$ -partitions and inner products of skew Schur functions*, Contemporary Mathematics, **34** (1984) 289–301.
- [3] F. Hivert, *Quasi-symmetric functions and Hecke algebra action*, Proceeding of the 10th FPSAC, Toronto (1998) 377–388.
- [4] T. Józefiak, *Schur  $Q$  functions and cohomology of isotropic Grassmanians*, Math. Proc. Camb. Phil. Soc., **109** (1991) 471–478.
- [5] C. Malvenuto and C. Reutenauer, *Duality between quasi-symmetric functions and the Solomon descent algebra*, Journal of Algebra, **177** (1995) 967–982.
- [6] N.J.A. Sloane, *An on-line version of the Encyclopedia of integer sequences*, Elect. J. Combin., **1** No.1, (1994) F1.  
<http://akpublic.research.att.com/~njas/sequences/ol.html>.
- [7] P. Pragacz, *Algebro-geometric applications of Schur  $S$ - and  $Q$ - polynomials*, Topics in Invariant Thoery, Séminaire d'Algèbre Dubreil-Malliavin 1989-90, Springer-Verlag, (1991) 130–191.
- [8] I. Schur, *Über die Darstellung der symmetrischen und der alternierenden Gruppe durch gebrochene lineare Substitutionen*, J. reine angew. Math. , **139** (1911) 155–250.
- [9] A.N. Sergeev, *The tensor algebra of the identity representation as a module over the Lie superalgebras  $gl(n, m)$  and  $Q(n)$* , Math. USSR Sbornik, **51** (1985) 419–427.
- [10] J. Stembridge, *Enriched  $P$ -partitions*, Trans. AMS, **349** (1997) 763–788.

(Nantel Bergeron and Stefan Mykytiuk and Stephanie van Willigenburg) DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, TORONTO, ONTARIO M3J 1P3, CANADA

(Frank Sottile) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, VAN VLECK HALL, 480 LINCOLN DRIVE, MADISON, WISCONSIN 53706-1388, USA

*E-mail address*, Nantel Bergeron: [bergeron@mathstat.yorku.ca](mailto:bergeron@mathstat.yorku.ca)

*URL*, Nantel Bergeron: <http://www.math.yorku.ca/bergeron>

*E-mail address*, Stefan Mykytiuk: [Mykytiuk@pascal.math.yorku.ca](mailto:Mykytiuk@pascal.math.yorku.ca)

*E-mail address*, Frank Sottile: [sottile@math.wisc.edu](mailto:sottile@math.wisc.edu)

*URL*, Frank Sottile: <http://www.math.wisc.edu/~sottile>

*E-mail address*, Stephanie van Willigenburg: [steph@pascal.math.yorku.ca](mailto:steph@pascal.math.yorku.ca)

*URL*, Stephanie van Willigenburg: <http://www-theory.dcs.st-and.ac.uk/~stephvw>