Peccei-Quinn mechanism and axion interactions in the 3-3-1 model with Cosmological Inflation

H. N. Long^{a,b1,2,*} and L. T. Hue^{a,b1,2,[†](#page-0-1)}

^{1 a} Subatomic Physics Research Group, Science and Technology Advanced Institute,

Van Lang University, Ho Chi Minh City 700000, Vietnam

 2^b Faculty of Applied Technology, School of Technology,

Van Lang University, Ho Chi Minh City 700000, Vietnam

(Dated: October 10, 2023)

Based on the Peccei-Quinn (PQ) assignment, the PQ charge operator in the 3-3-1 model with Cosmological Inflation is constructed in terms of diagonal generators T_3 and T_8 of the $SU(3)_L$ subgroup. The formula shows that difference of PQ charges of up and down quarks is 2, i.e., $\Delta Q_A = 2$, while for electric charge, are assumed to be equal, while the (Q_A) are opposite. PQ charge of neutral scalars equal ± 2 , while for charged scalars, it vanishes. The couplings of axion with fermions are presented. To have correct kinetic term for the axion, the PQ scale f_a is to be VEV of the singlet scalar boson, namely, $f_a = v_\phi$. The photon couples to charged particles only, while in this model, the axion does not couple to charged scalar/gauge bosons. The point is worth emphasizing that the axion has doubly derivative coupling with scalar playing the role of inflaton. The new effects mainly happen in the energy region from 10^7 GeV to 10^{11} GeV. The chiral effective Lagrangian as usually provides axion mass consistent with model-independent prediction.

I. INTRODUCTION

At present, axion is one of the hottest issues in particle physics from both theoretical and experimental aspects [\[1](#page-14-0)[–4\]](#page-15-0). As shown in Refs. [\[5](#page-15-1), [6](#page-15-2)], the $\tilde{\theta}$ term: $\mathcal{L}_{\theta} \sim \tilde{\theta} \tilde{G}.G$ where \tilde{G} being the dual gluon strength tensor is responsible for CP violation. It is well-known that the absence of CP violation in strong sector of the Standard Model (SM) leads to upper bound of the value

$$
\tilde{\theta} \le 10^{-10} \,. \tag{1}
$$

Question is why this value so small called strong CP problem.

To solve the puzzle, Peccei and Quinn [\[7,](#page-15-3) [8\]](#page-15-4) have represented a singlet field $\Phi(x)$ in which an axion is complex phase in polar coordinate i.e., $\Phi(x) = R(x)e^{i\theta(x)}$. Here θ is responsible for neutron electric diplole moment (EDM). Next point in PQ recipe is to make $\theta = 0$ energetically favourable. This means that if one writes potential $V = V(\Phi)$ then it will have a minimum at desired value, for example $R = v \gg 0$. To solve the strong CP problem, the potential also favour $\theta = 0$. Setting neutron EDM to zero, the simplest way is harmonic form of potential $V(\theta) \sim \theta^2$.

Combining the above discussions, one comes to redefinition of the singlet field as follows [\[9](#page-15-5)]

$$
\Phi = R(x) e^{i\frac{a(x)}{f_a}x_{\phi}},\tag{2}
$$

where x_{ϕ} and f_a are PQ charge of ϕ and axion decay constant, respectively. It is worth noting that to satisfy the above upper, the axion decay constant f_a must be very large: $f_a \ge 10^{10}$ GeV. The factor a/f_a ensures a solution to the strong CP puzzle.

The PQ formalism is considered in many versions of the beyond Standard Models (BSMs). Among BSMs, the models based on the $SU(3)_C \times SU(3)_L \times U(1)_N$ (3-3-1) gauge group [\[10](#page-15-6)[–19](#page-15-7)] have some intriguing properties such as: i) Due to fact that one generation of quarks transforms differently from two other ones, the anomaly free condition leads to number of generation is multiple of color number. Combining with QCD asymptotic free requiring number of quarks is not bigger than five, one gets number of generation is equal to three. By the same reason, it explains why top quark is so heavy (175 GeV). ii) The PQ symmetry is automatically fulfilled in the 3-3-1 models [\[20\]](#page-15-8). iii) The models provide electric charge quantization [\[21](#page-15-9), [22\]](#page-15-10).

[∗]Electronic address: hoangngoclong@vlu.edu.vn

[†]Electronic address: lethohue@vlu.edu.vn

In the frameworks of the 3-3-1 models, the PQ formalism has been considered for almost two decades ago [\[23](#page-15-11)[–25\]](#page-15-12). However, in the above mentioned works, the main ingredient, namely the singlet ϕ is expanded as ordinary complex scalar field, i.e., in the sum of CP even and CP odd components. In that cases, the axion mixs with other CP odd scalars. The axion is the pure imaginary part of ϕ only in the limit $v_{\phi} \gg v_{\chi}$, which is the vacuum expectation value (VEV) responsible for breaking from $SU(3)_L$ to SM subgroup. In Ref. [\[25](#page-15-12)], the PQ symmetry was considered for two main versions: the minimal 3-3-1 model [\[10](#page-15-6)[–13\]](#page-15-13) and the version with right-handed neutrinos [\[14](#page-15-14)[–19](#page-15-7)].

New development in this direction was done five years ago in Ref. [\[26](#page-15-15)], where the discrete symmetry $Z_{11} \times Z_2$ is imposed, and Majorana right-handed neutrinos are introduced. To provide their masses, a complex scalar transforming as a $SU(3)_{L_2}$ singlet is added. As a consequence, the model also contains a heavy CP even scalar with mass in the range of 10^{10} GeV called by inflaton. However, Ref. [\[26\]](#page-15-15) still contains some flawed points in the assignment of the Z_2 symmetry, incorrect mixing matrix of the CP odd scalars, and still an absence of identification of the Standard Model-like Higgs boson. The mentioned problems have been solved in Ref. [\[27](#page-15-16)]. In the above work, using Euler rotation method, the correct mixing matrix of CP odd sector has been performed. For the CP even sector, applying the Hatree-Fock approximation, the 4×4 matrix has been diagonalized. As a result, the model contains the expected inflaton with mass around 10^{11} GeV, one heavy scalar with mass at TeV scale labeled by H_x , one scalar with mass at the EW scale (h_5) , and of course the SM-like Higgs boson (h) .

In summary, the 3-3-1 model under consideration has some interesting properties such as: i) The existence of Majorana right-handed neutrino with masses produced at the very high PQ scale around 10^{11} GeV, ii) The CP-even component of the singlet plays a role of the inflaton with very large mass too. ii) Lastly, the scalar sector is quite simple and very interesting beside the scalar with mass 125 GeV, there is also a partner with mass at the EW scale 150/96 GeV.

Note that in Refs. [\[26,](#page-15-15) [27](#page-15-16)], the axion becomes the pure imaginary part of the singlet ϕ , only if $\tan \theta_{\phi} \approx v_{\chi}/v_{\phi} \to 0$. In addition, the axion does not explicitly present in the law of the PQ transformation. For example Ref. [\[26](#page-15-15)], PQ transformation of the single is $\phi \to e^{-i2\alpha X_d} \phi$, which is quite different from that given in Eq. [\(2\)](#page-0-2).

In this work, the singlet field ϕ takes complex value everywhere, and axion is complex phase in polar coordinates. Then the axion also appears as a phase of PQ transformations. To solve the strong CP puzzles in QCD, the axion will associate factor 10^{-10} and in this case $1/f_a$ where $f_a \sim 10^{10}$ GeV.

The layout of the reminder of this work is as follows. In section [II](#page-1-0) we briefly present fermion and scalar with their quantum numbers of the model. Section [III](#page-3-0) is devoted to the PQ symmetry, where the PQ charge operator is constructed in terms of diagonal generators. The couplings of axion are presented in section [IV.](#page-5-0) The anomalous and derivative axion's couplings are discussed. The axion couplings to scalar are discussed in detail in section [V.](#page-11-0) We show that to have the correct axion's kinetic form: $f_a = v_\phi$. We make outlines and conclusions in the final section - section [VI.](#page-13-0)The PQ transformation and its charge operator are presented in sections [A](#page-13-1) and [B,](#page-14-1) respectively.

II. BRIEF REVIEW OF THE MODEL

To have Dirac and Majorana mass terms for ν_L and N_R , the particles and their transforms [\[26](#page-15-15), [27](#page-15-16)] under SU(3)_C × $\text{SU(3)}_L \times \text{U(1)}_N \times Z_2 \times Z_{11}$ group are presented in Table [I](#page-1-1) (Reason why we have changed the label for U(1) subgroup will be cleared in subsection [III A\)](#page-3-1).

	Q_{nL}					Q_{3L} u_{aR} d_{aR} T_{3R} D_{nR} ψ_{aL} l_{aR} N_{aR}				η	X	Ω	Φ
$SU(3)_C$	3	3	3	3	3	3							
$SU(3)_L$	$\overline{\mathbf{3}}$	3					3			3	3	3	
$U(1)_N$	0	$\frac{1}{2}$	$\frac{2}{3}$	$\overline{ }$	$\frac{2}{3}$	\sim	$\mathbf{\Omega}$		$\overline{0}$	÷	-	$\frac{2}{3}$	0
Z_2													
Z_{11}	\sim ω_4	ω_0	ω_5	ω_2	ω_3	ω_4	ω_1	ω_3	ω_{5}	ω_{5}	$\omega_{\rm o}$	ω_{2}	ω .

Table I: $SU(3)_C \times SU(3)_L \times U(1)_N \times Z_{11} \times Z_2$ charge assignments of the particle content of the model. Here $w_k = e^{ik2\pi/11}$, $a = 1, 2, 3$ and $\alpha = 1, 2$.

Masses of fermions and gauge bosons request VEVs of three triplets and one singlet

$$
\langle \eta \rangle = \frac{1}{\sqrt{2}} (v_{\eta}, 0, 0)^T, \langle \chi \rangle = \frac{1}{\sqrt{2}} (0, 0, v_{\chi})^T,
$$

$$
\langle \rho \rangle = \frac{1}{\sqrt{2}} (0, v_{\rho}, 0)^T, \quad \langle \phi \rangle = \frac{1}{\sqrt{2}} v_{\phi}.
$$
 (3)

The VEV v_x triggers the spontaneous breaking of the $SU(3)_L \times U(1)_N$ gauge symmetry down to the SM electroweak gauge

group, while VEVs of η and ρ break the SM electroweak gauge group. The VEV of the singlet v_{ϕ} breaks the PQ symmetry and provides Majoran mass for N_R .

A. Neutrino masses

With the above specified particle content, the following Yukawa interactions invariant under the $SU(3)_C \times SU(3)_L \times U(1)_N \times$ $Z_2 \times Z_{11}$ symmetry, arise [\[26](#page-15-15)]: With the above specified particle content, the following Yukawa couplings for quarks invariant under the above mentioned group, arise [\[26,](#page-15-15) [27\]](#page-15-16):

$$
-\mathcal{L}_{q}^{Y} = y_{1}\bar{Q}_{3L}T_{R}\chi + \sum_{n,m=1}^{2} (y_{2})_{n,m} \bar{Q}_{nL}D_{mR}\chi^{*} +\sum_{a=1}^{3} (y_{3})_{3a} \bar{Q}_{3L}u_{aR}\eta + \sum_{n=1}^{2} \sum_{a=1}^{3} (y_{4})_{na} \bar{Q}_{nL}d_{aR}\eta^{*} +\sum_{a=1}^{3} (y_{5})_{3a} \bar{Q}_{3L}d_{aR}\rho + \sum_{n=1}^{2} \sum_{a=1}^{3} (y_{6})_{na} \bar{Q}_{nL}u_{aR}\rho^{*} + \text{H.c.}
$$
 (4)

and for the lepton [\[27\]](#page-15-16)

$$
-\mathcal{L}_l^Y = \sum_{a=1}^3 \sum_{b=1}^3 g_{ab} \bar{\psi}_{aL} l_{bR} \rho + \sum_{a=1}^3 \sum_{b=1}^3 \left(y_\nu^D \right)_{ab} \bar{\psi}_{aL} \eta N_{bR} + \sum_{a=1}^3 \sum_{b=1}^3 \left(y_N \right)_{ab} \phi \bar{N}_{aR}^C N_{bR} + \text{H.c.}.
$$
 (5)

The last two terms in [\(5\)](#page-2-0) contains related to neutrino mass

$$
-\mathcal{L}_l^Y \supset \left(y_\nu^D\right)_{ab} \bar{\psi}_{aL} \eta N_{bR} + (y_N)_{ab} \bar{N}_{aR}^C N_{bR} \phi + \text{H.c.} \tag{6}
$$

The Dirac neutrino mass term arises from v_{η} , while the Majorana mass term arises from v_{ϕ} .

From [\(6\)](#page-2-1), it follows that the light active neutrinos are generated from a type I seesaw mechanism mediated by right handed Majorana neutrinos. Thus implying that the resulting light active neutrino mass matrix has the form [\[27](#page-15-16)]

$$
M_{\nu} = M_{\nu}^{D} M_{N}^{-1} \left(M_{\nu}^{D} \right)^{T}, \quad M_{\nu}^{D} = \frac{v_{\eta}}{\sqrt{2}} y_{\nu}^{D}, \quad M_{N} = \sqrt{2} v_{\phi} y_{N}. \tag{7}
$$

The standard neutrinos get mass at eV scale, while $M_R \sim 10^7$ GeV [\[26\]](#page-15-15). Note that the similar script containing axion, inflaton and seesaw exists in Ref. [\[28\]](#page-15-17).

B. The Higgs sector

The full potential of the model under consideration has the form [\[26](#page-15-15)]

$$
V_{tot} = \mu_{\chi}^2 \chi^{\dagger} \chi + \mu_{\rho}^2 \rho^{\dagger} \rho + \mu_{\eta}^2 \eta^{\dagger} \eta + \mu_{\phi}^2 \phi^* \phi + \lambda_1 (\chi^{\dagger} \chi)^2 + \lambda_2 (\eta^{\dagger} \eta)^2 + \lambda_3 (\rho^{\dagger} \rho)^2 + \lambda_4 (\chi^{\dagger} \chi)(\eta^{\dagger} \eta) + \lambda_5 (\chi^{\dagger} \chi)(\rho^{\dagger} \rho) + \lambda_6 (\eta^{\dagger} \eta)(\rho^{\dagger} \rho) + \lambda_7 (\chi^{\dagger} \eta)(\eta^{\dagger} \chi) + \lambda_8 (\chi^{\dagger} \rho)(\rho^{\dagger} \chi) + \lambda_9 (\eta^{\dagger} \rho)(\rho^{\dagger} \eta) + \lambda_{10} (\phi^* \phi)^2 + \lambda_{11} (\phi^* \phi)(\chi^{\dagger} \chi) + \lambda_{12} (\phi^* \phi)(\rho^{\dagger} \rho) + \lambda_{13} (\phi^* \phi)(\eta^{\dagger} \eta) + \left(\lambda_{\phi} \epsilon^{ijk} \eta_i \rho_j \chi_k \phi^* + \text{H.c.} \right)
$$
\n(8)

For future presentation, here we write explicitly the scalar fields

$$
\chi^{T} = (\chi_{1}^{0}, \chi_{2}^{-}, \chi_{3}^{0}) \sim (1, 3, -\frac{1}{3}), \eta^{T} = (\eta_{1}^{0}, \eta_{2}^{-}, \eta_{3}^{0}) \sim (1, 3, -\frac{1}{3}),
$$

\n
$$
\rho^{T} = (\rho_{1}^{+}, \rho_{2}^{0}, \rho_{3}^{+}) \sim (1, 3, \frac{2}{3}), \quad \phi = \frac{1}{2} (v_{\phi} + R_{\phi}) e^{i \frac{\sigma}{2f_{a}} x_{\phi}} | \sim (1, 1, 0),
$$
\n(9)

where $\tan \theta_{PQ} = \frac{I_{\phi}}{n+1}$ $v_\phi+R_\phi$

In this case, the axion is decoupled from the scalar sector, and this case corresponds to the limit $v_{\phi} \gg v_{\chi}$ in Ref. [\[27](#page-15-16)]. However, it does not touch to other sectors such as CP even scalars.

The VEV v_{ϕ} is responsible for the PQ symmetry breaking resulting (see below). Then VEV v_{χ} breaks SU(3)_L × U(1)_N to the SM group. Two others v_{ρ}, v_{η} are needed for the usual $U(1)_{Q}$ symmetry. Hence, it follows $v_{\phi} \gg v_{\chi} \gg v_{\rho}, v_{\eta}$. The constraint conditions at tree level were analyzed in Ref. [\[24\]](#page-15-18). From [\(8\)](#page-2-2), it is quite reasonable to accept : $\lambda_2 \approx \lambda_3$, $\lambda_4 \approx \lambda_5$, $\lambda_7 \approx \lambda_8$, $\lambda_{12} \approx$ λ_{13} . From result in Ref. [\[29](#page-15-19)], it follows $v_{\chi} \ge 10357$ GeV for $M_{Z'} \ge 4.1$ TeV.

The result of Ref. [\[27](#page-15-16)] showed that in the CP-even scalar sector, there are six fields. One massless field is part of G_{X0} , another massive in TeV scale is associated to χ_1^0 . One heavy field with mass in the range of 10^{11} GeV and associated with singlet ϕ is identified to inflaton Φ. One SM-like Higgs boson h with mass ∼ 125 GeV. Two remain fields include one heavy with mass at TeV scale (H_x) and another with mass at EW scale (h_5) .

In the limit $v_{\phi} \gg v_{\chi} \gg v_{\rho} \gg v_{\eta}$, one has [\[27](#page-15-16)]

$$
\eta \simeq \begin{pmatrix} \frac{1}{\sqrt{2}} (u + h_5 + iA_5) \\ H_1^- \\ G_{X^0} \end{pmatrix}, \ \chi \simeq \begin{pmatrix} \chi_1^0 \\ G_{Y^-} \\ \frac{1}{\sqrt{2}} (v_X + H_X + iG_{Z'}) \end{pmatrix}, \ \rho \simeq \begin{pmatrix} G_{W^+} \\ \frac{1}{\sqrt{2}} (v + h + iG_Z) \\ H_2^+ \end{pmatrix},
$$
\n
$$
\phi = \frac{1}{\sqrt{2}} (v_{\phi} + \Phi) e^{-i\frac{\theta}{f_a}}.
$$
\n(10)

Combination of table [I](#page-1-1) and Eqs. (4) , (5) , (10) leads to some interesting consequences: Firstly, the SM-like Higgs boson h has Yukawa couplings with only SM fermions. Secondly, the pseudoscalar A_5 and H_X can have Yukawa couplings with not only exotic quarks but also SM quarks and leptons.

III. PECCEI - QUINN MECHANISM

A. PQ charge in the 3-3-1 model with Inflation

The new format of writing PQ transforms is given in Refs. [\[1](#page-14-0), [2](#page-14-2), [9](#page-15-5)] and in Refs. [\[30,](#page-15-20) [31\]](#page-15-21). According to Ref. [\[24](#page-15-18)], and using notations in Refs. [\[30](#page-15-20), [31\]](#page-15-21) for an arbitrary fermion and scalar boson, the PQ transformations are as follows

$$
f \rightarrow f' = e^{i\left(\frac{x_f}{2f_a}\right)\gamma_5 a} f, \quad \bar{f} \rightarrow \bar{f}' = \bar{f} e^{i\left(\frac{x_f}{2f_a}\right)\gamma_5 a}, \quad \varphi \rightarrow \varphi' = e^{i\left(\frac{x_\varphi}{2f_a}\right)a} \varphi,
$$
\n
$$
(11)
$$

$$
f_L \rightarrow f'_L = e^{-i\left(\frac{x_f}{2f_a}\right)a} f_L, \quad \bar{f}_L \rightarrow \bar{f}'_L = \bar{f}_L e^{i\left(\frac{x_f}{2f_a}\right)a},
$$

\n
$$
f_R \rightarrow f'_R = e^{i\left(\frac{x_f}{2f_a}\right)a} f_R, \quad \bar{f}_R \rightarrow \bar{f}'_R = \bar{f}_R e^{-i\left(\frac{x_f}{2f_a}\right)a},
$$
\n(12)

where x_f is PQ charge of fermion and $f_a \sim 10^{11}$ GeV is axion decay constant relating to the scale of symmetry breaking of $U(1)_{PQ}$ global group. The PQ charges of fermions are given as [\[26](#page-15-15)]

$$
x_u = x_T = -x_d = -x_D = x_l = -x_{lR} = -x_{\nu} = x_{\nu_R} = -x_N \equiv R. \tag{13}
$$

Note that the parameter R in (13) in principle, is any non-zero integer. However, as seen later, within notations in (11) , the absolute value of R should be equal to the unit, i.e., $|R| = 1$. Here, we choose the sign is plus, i.e., $R = 1$.

For scalars, from Yukawa couplings, it follows that charged scalars have vanishing PQ charge since they connect up and down particles with opposite values, while electrically neutral scalar has PQ charge duplicate charge (with opposite sign) of fermion to which it provides mass, because it connects to both up or down particles:

$$
\eta_1^0 \rightarrow e^{i\frac{a}{f_a}} \eta_1^0, \quad \chi_1^0 \rightarrow e^{i\frac{a}{f_a}} \chi_1^0, \quad \rho_1^+ \rightarrow \rho_1^+, \n\phi \rightarrow e^{i\frac{a}{f_a}} \phi, \quad \rho_2^0 \rightarrow e^{-i\frac{a}{f_a}} \rho_2^0, \quad \chi_2^- \rightarrow \chi_2^-.
$$
\n(14)

The last term in [\(8\)](#page-2-2) shows that ρ has the same value and opposite to that of η , χ and ϕ . This is explicitly clarified in Table [II.](#page-3-5)

			$d \left T\right D_{\alpha} \left l\right \left \nu\right \left \nu_R\right N_R \left \eta_1^0\right \eta_3^0 \left \chi_1^0\right \chi_3^0 \left \rho^0\right \phi \left \eta_2^-\right \chi_2^-\left \rho_1^+\right \rho_3^+$					
$U(1)_{PQ}$ 1 -1 1 -1 -1 -1 +1 1 -1 2 2 2 2 -2 0 0 0 0								

Table II: $U(1)_{PQ}$ charge assignments of the particle content of the model. Here $x_F = x_{F_R} = -x_{F_L}$

For singlets of right-handed fermions and scalar ϕ , the generators T_3 and T_8 produce the zero, but the general PQ charge \mathcal{X}_{pq} for right handed fermions takes opposite value of left handed partner, while for ϕ , as usually, it is followed from Yukawa couplings, i.e., $\mathcal{X}_{pq}(f_R) = -Q_A(f_L)$ and $\mathcal{X}_{pq}(\phi) = 2$, $\Rightarrow Q_A(\phi) = 2$. To be noted that there exists the same situation when one deals with electric charges of right-handed fermions.

The PQ charges of the model particle content can be clearly drawn in Table [III.](#page-4-0)

Table III: Multiplets and Q_A charges in the model

From Tables [II,](#page-3-5) [III](#page-4-0) and [\(14\)](#page-3-6), we can formulate PQ charge assignment for triplets in the model under consideration as follows (for details, see Appendix [B\)](#page-14-1)

$$
Q_A = 2T_3 - \frac{2}{\sqrt{3}}T_8 + \mathcal{X}_{pq} \,. \tag{15}
$$

Therefore, the PQ transformation in [\(12\)](#page-3-4) and [\(14\)](#page-3-6) can be written in form of PQ charge operator labeled by Q_A as follows

$$
U(1)_{PQ}: f \to f' = e^{i\left(\frac{a}{2f_a}\right)\gamma_5 Q_A} f, \quad \bar{f} \to \bar{f}' = \bar{f} e^{i\left(\frac{a}{2f_a}\right)\gamma_5 Q_A}, \tag{16}
$$

$$
f_L \rightarrow f'_L = e^{-i\left(\frac{a}{2f_a}\right)Q_A} f_L, \quad \bar{f}_L \rightarrow \bar{f}'_L = \bar{f}_L e^{i\left(\frac{a}{2f_a}\right)Q_A},
$$
\n
$$
f \rightarrow f' = e^{i\left(\frac{a}{2f_a}\right)Q_A} f_L, \quad \bar{f} \rightarrow \bar{f}'_L = \bar{f}_L e^{-i\left(\frac{a}{2f_a}\right)Q_A},
$$
\n
$$
(17)
$$

$$
f_R \rightarrow f'_R = e^{i\left(\frac{a}{2f_a}\right)Q_A}f_R, \quad \bar{f}_R \rightarrow \bar{f}'_R = \bar{f}_Re^{-i\left(\frac{a}{2f_a}\right)Q_A},\tag{17}
$$

$$
\varphi \ \to \ \varphi' = e^{+i\left(\frac{a}{2f_a}\right)Q_A}\varphi \,. \tag{18}
$$

It is worth noting that in the realization the Georgi-Kaplan-Randall (GKR) field basis, all fields except the axion, transform by additive constant.

The following remarks are in order. The formula [\(15\)](#page-4-1) shows that difference of electric charges of up and down quarks/leptons is 1, i.e., $\Delta Q = 1$, while for PQ charge $\Delta Q_A = 2$. For right handed fermions, electric charges (Q) of left-handed and right-handed are assumed to be equal, while for (Q_A) is opposite. For scalar bosons, only neutral scalar have a PQ charge equal to ± 2 , where the sign plus for scalar in top of doublet and minus for bottom one. The charged scalar does not have a PQ charge. So the electric charge (Q) and (Q_A) have again opposite property. This is illustrated in Table [IV.](#page-4-2)

	$U(1)_Q$	$U(1)_{Q_A}$
f_u f_d	$\Delta Q=1$	$\Delta Q_A = 2$
φ^0	$Q(\varphi^0) = 0$	$Q_A(\varphi^0) = 2$ $Q_A(\varphi^-) = 0$
Chiral fermion f	$Q(f_R) = Q(f_L)$	$ Q_A(f_R) = -Q_A(f_L)$

Table IV: Some characteristic properties between $U(1)_Q$ and $U(1)_{Q_A}$.

To solve the strong CP problem, the $U(1)_{pq}$ symmetry is spontaneously breaking by VEV of the singlet ϕ as follows

$$
Q_A \langle \phi \rangle = \frac{2}{\sqrt{2}} v_{\phi} \neq 0. \tag{19}
$$

B. Yukawa couplings and quark masses

From Table [II,](#page-3-5) we see that PQ charge of neutral scalar equals twice of PQ charges of fermions receiving for which scalar provides masses, so all Yukawa couplings given in (4) and (5) are invariant. In addition, exotic quark carrying lepton number 2, while ordinary SM quarks do not, so mass eigenstates of exotic quarks are their original states, while ordinary quarks are with mass mixing. Hence, we just deal with ordinary quarks. Note that top (t) quark gets mass from VEV of scalar η while c and u quarks get mass from VEV of ρ . For down quarks, the situation is converse. Therefore, for up quarks, one has [\[27\]](#page-15-16)

$$
M_u = \frac{v_{\eta}}{\sqrt{2}} \begin{pmatrix} (y_6)_{11} \frac{v_{\rho}}{v_{\eta}} & (y_6)_{12} \frac{v_{\rho}}{v_{\eta}} & (y_6)_{13} \frac{v_{\rho}}{v_{\eta}} \\ (y_6)_{21} \frac{v_{\rho}}{v_{\eta}} & (y_6)_{22} \frac{v_{\rho}}{v_{\eta}} & (y_6)_{23} \frac{v_{\rho}}{v_{\eta}} \end{pmatrix} = V_{uL} \widetilde{M}_u V_{uR}^{\dagger},
$$
\n
$$
(20)
$$

with

$$
\widetilde{M}_u = \text{diag}(m_u, m_c, m_t) \tag{21}
$$

In a similar way, one gets for down quarks

$$
M_d = \frac{v_\rho}{\sqrt{2}} \begin{pmatrix} (y_4)_{11} \frac{v_\eta}{v_\rho} & (y_4)_{12} \frac{v_\eta}{v_\rho} & (y_4)_{13} \frac{v_\eta}{v_\rho} \\ (y_4)_{21} \frac{v_\eta}{v_\rho} & (y_4)_{22} \frac{v_\eta}{v_\rho} & (y_4)_{23} \frac{v_\eta}{v_\rho} \\ (y_5)_{31} & (y_5)_{32} & (y_5)_{33} \end{pmatrix} = V_{dL} \widetilde{M}_d V_{dR}^{\dagger},
$$
\n(22)

with

$$
M_d = \text{diag}(m_d, m_s, m_b) \tag{23}
$$

In the above matrices, all Yukawa couplings of the form $(y_i)_{ab}$ $a, b = 1, 2, 3; i = 3, 4, 5, 6$ are positive and real. With $\alpha = 1, 2$ and $a = \alpha, 3$, these couplings can be determined by the following equations:

$$
(y_6)_{na} = \frac{\sqrt{2}}{v_\rho} \left(V_{uL} \widetilde{M}_u V_{uR}^\dagger \right)_{na}, \qquad (y_3)_{3a} = \frac{\sqrt{2}}{v_\eta} \left(V_{uL} \widetilde{M}_u V_{uR}^\dagger \right)_{3a} \tag{24}
$$

$$
(y_4)_{na} = \frac{\sqrt{2}}{v_\eta} \left(V_{dL} \widetilde{M}_d V_{dR}^\dagger \right)_{na}, \qquad (y_5)_{3a} = \frac{\sqrt{2}}{v_\rho} \left(V_{dL} \widetilde{M}_d V_{dR}^\dagger \right)_{3a} . \tag{25}
$$

From (20) and (22) , the diagonalized mass matrix of ordinary quarks are determined as below:

$$
\widetilde{M}_{u,d} = \left(V_L^{(u,d)}\right)^\dagger M_{u,d} V_R^{(u,d)}.
$$
\n(26)

In general, one gets:

$$
M_f = (M_f)_{diag} = V_{fL}^{\dagger} M_f V_{fR}, \qquad f_{(R,L)} = V_{f(R,L)} \hat{f}_{(R,L)},
$$

$$
\overline{f}_{aL} (M_f)_{ab} f_{bR} = \overline{\hat{f}}_{kL} (V_{fL}^{\dagger})_{ka} (M_f)_{ab} (V_{fR})_{bl} \tilde{f}_{lR} = \overline{\hat{f}}_{kL} (V_{fL}^{\dagger} M_f V_{fR})_{kl} \tilde{f}_{lR} = \overline{\hat{f}}_{kL} (\widetilde{M}_f)_{kl} \tilde{f}_{lR} = m_{f_k} \overline{\hat{f}}_{kL} \tilde{f}_{kR},
$$

$$
k = 1, 2, 3.
$$
 (27)

Here, $f_{k(R,L)}$ and $f_{k(R,L)}$ ($k = 1, 2, 3$) are the SM fermionic fields in the mass and interaction bases, respectively. From [\(5\)](#page-2-0), it follows charged leptons acquire masses from VEV of ρ , while the Majorana right handed neutrinos N_R get mass from $\langle \phi \rangle$. Since we focus on the strong CP problem, we leave lepton part for this moment.

Now we turn into coupling of axion with matter fields and scalar bosons

IV. AXION INTERACTIONS

A. Axion gauge boson couplings

Note that the gauge boson self-couplings in the 3-3-1 models were calculated in [\[32](#page-15-22), [33\]](#page-15-23). To understand these couplings, let us denote W to be gauge boson of the group $SU(3)$. Then the *electric* and *magnetic* fields are:

$$
W_{0\nu}^{a} \equiv E_{\nu}^{a}, \quad \nu = 1, 2, 3 = i \; \text{(space index)}; a = 1, 2, \dots 8
$$
\n
$$
B_{k}^{a} \equiv \epsilon_{ijk} W_{ij}^{a} \quad i \neq j \neq k, (j, k = 1, 2, 3)
$$
\n
$$
(28)
$$

Then the Lagrangian of gauge bosons are follows

$$
-4\mathcal{L}_{gauge} \supset a(x)W_{a\mu\nu}W^{a\mu\nu} = 2a(x)(W_{a0\nu}W^{a0\nu} + W_{aij}W^{aij}) = 2a(x)(E_{ai}E^{ai} + B_{ak}B^{ak}) = 2a(x)(\mathbf{E}^2 + \mathbf{B}^2). \tag{29}
$$

In the 3-3-1 models, this part contains triple and quartic gauge boson couplings which have been presented in Refs. [\[32](#page-15-22), [33](#page-15-23)]. Now we consider a part with the dual tensor

$$
\mathcal{L}_{\theta} \supset \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} W_{a\mu\nu} W_{\alpha\beta}^{a} = 4! \epsilon^{0i\alpha\beta} W_{a0i} W_{\alpha\beta}^{a} = 4! \epsilon^{0ijk} W_{a0i} W_{jk}^{a} = 4! (E_{ai} B^{ai}) = 4! \mathbf{E} \cdot \mathbf{B}
$$
\n(30)

$$
= 4! \epsilon^{0ijk} [\partial_0 W_i^a - \partial_i W_0^a + gf_{abc} W_0^b W_i^c] [\partial_j W_k^a - \partial_k W_j^a + gf_{ade} W_j^d W_k^e]. \tag{31}
$$

Here

$$
W_{\mu\nu}^a = \partial_{\mu} W_{\nu}^a - \partial_{\nu} W_{\mu}^a + g f_{abc} W_{\mu}^b W_{\nu}^c \tag{32}
$$

Note that in [\(32\)](#page-6-0), four space-time indexes are different. Then $G\tilde{G}$ is violated by $x_o \to -x_0$ (T inverse) or $x_i \to -x_i$ (parity). Since CPT invariance, one has CP violation of CT violation.

According to Ref. [\[8](#page-15-4)], for Lagrangian below

$$
\mathcal{L} = -\frac{1}{4} F^a_{\mu\nu} F^{a\mu\nu} + i \bar{\psi} \gamma_\mu D^\mu \psi + L_{Yukawa} - V(\phi) , \qquad (33)
$$

with transformation

$$
\to e^{i\sigma\gamma_5}\psi \,, \quad \bar{\psi} \to \bar{\psi}e^{i\sigma\gamma_5} \quad \phi \to e^{-2i\sigma}\phi \,. \tag{34}
$$

then one has

$$
\partial^{\mu} J_{\mu}^{5} = \frac{g^2}{16\pi^2} F_{\mu\nu}^a \tilde{F}^{a\mu\nu}
$$
\n(35)

The the 3-3-1 model with three gauge subgroups, the anomalous couplings have the form [\[35\]](#page-15-24)

 ψ

$$
\mathcal{L}_{ag} = c_G \frac{\alpha_s}{4\pi} \frac{a}{2f_a} G\tilde{G} + c_{WW} \frac{\alpha_2}{4\pi} \frac{a}{2f_a} W\tilde{W} + c_{BB} \frac{\alpha_1}{4\pi} \frac{a}{2f_a} B\tilde{B}.
$$
\n(36)

Using Eq. [\(32\)](#page-6-0) and replace W^a by physical fields in given in [\(62\)](#page-11-1), one will have coupling of axion (a) with gauge bosons with four external indexes (space-time) different.

B. Axion-fermion couplings

Let us start from kinematic term of fermion as follows

$$
\mathcal{L}_f^0 = \frac{i}{2} \bar{f} \gamma^\mu \stackrel{\leftrightarrow}{\partial_\mu} f = \frac{i}{2} (\bar{f} \gamma^\mu \partial_\mu f - \partial_\mu \bar{f} \gamma^\mu f). \tag{37}
$$

For the gauge version, where ordinary derivative replaced by covariant one, we have

$$
\mathcal{L}_f = \frac{i}{2} (\bar{f}\gamma^\mu D_\mu f - D_\mu \bar{f}\gamma^\mu f)
$$

\n
$$
= \frac{i}{2} [\bar{f}\gamma^\mu (\partial_\mu - iP_\mu^f) f - (\partial_\mu \bar{f} + i \bar{f} P_\mu^f) \gamma^\mu f]
$$

\n
$$
= \frac{i}{2} \bar{f}\gamma^\mu \stackrel{\leftrightarrow}{\partial_\mu} f + \bar{f} P_\mu^f \gamma^\mu f \equiv I + II
$$
\n(38)

Here $P_{\mu} = g_S G_{\mu} - g A_{\mu} - g' B_{\mu}$ and $G_{\mu}, A_{\mu}, B_{\mu}$ are matrices of gluon, $SU(3)_L$ and $U(1)_X$ bosons, respectively. We next deal with the first term

$$
I = \frac{i}{2} \bar{f}' \gamma^{\mu} \stackrel{\leftrightarrow}{\partial_{\mu}} f' = \frac{i}{2} \bar{f} e^{i \left(\frac{a}{2f_a} \right) \gamma_5 Q_A} \gamma^{\mu} \stackrel{\leftrightarrow}{\partial_{\mu}} e^{i \left(\frac{a}{2f_a} \right) \gamma_5 Q_A} f
$$

\n
$$
= \frac{i}{2} \left[\bar{f} e^{i \left(\frac{x_f}{2f_a} \right) \gamma_5 a} \gamma^{\mu} \partial_{\mu} (e^{i \left(\frac{x_f}{2f_a} \right) \gamma_5 a} f) - \partial_{\mu} (\bar{f} e^{i \left(\frac{x_f}{2f_a} \right) \gamma_5 a} \gamma^{\mu} e^{i \left(\frac{x_f}{2f_a} \right) \gamma_5 a} f \right]
$$

\n
$$
= \frac{1}{2} \left\{ \bar{f} e^{i \left(\frac{x_f}{2f_a} \right) \gamma_5 a} i \gamma^{\mu} \left[i \left(\frac{x_f}{2f_a} \right) e^{i \left(\frac{x_f}{2f_a} \right) \gamma_5 a} \gamma_5 \partial_{\mu} a f + e^{i \left(\frac{x_f}{2f_a} \right) \gamma_5 a} \partial_{\mu} f \right]
$$

\n
$$
- \left[(\partial_{\mu} \bar{f}) e^{i \left(\frac{x_f}{2f_a} \right) \gamma_5 a} + i \bar{f} \left(\frac{x_f}{2f_a} \right) e^{i \left(\frac{x_f}{2f_a} \right) \gamma_5 a} \gamma_5 \partial_{\mu} a \right] i \gamma^{\mu} e^{i \left(\frac{x_f}{2f_a} \right) \gamma_5 a} f \right\}
$$

\n
$$
= \frac{i}{2} \bar{f} \gamma^{\mu} \stackrel{\leftrightarrow}{\partial_{\mu}} f - \frac{i^2}{2} \times 2 \left(\frac{x_f}{2f_a} \right) (\partial_{\mu} a) \bar{f} \gamma_5 \gamma^{\mu} f.
$$

\n
$$
= \frac{i}{2} \bar{f} \gamma^{\mu} \stackrel{\leftrightarrow}{\partial_{\mu}} f + \left(\frac{x_f}{2f_a} \right) (\partial_{\mu} a) \bar{f} \gamma_5 \gamma^{\mu} f.
$$

\n(39)

The first term in [\(39\)](#page-6-1) is kinematic term of fermion: $I \supset \frac{1}{2} \bar{f} i \gamma^{\mu} \stackrel{\leftrightarrow}{\partial_{\mu}} f$. The second term is th famous derivative coupling of axion to fermion:

$$
\mathcal{L}_{(f-a)} = \left(\frac{x_f}{2f_a}\right) \partial_\mu a \bar{f} \gamma_5 \gamma^\mu f. \tag{40}
$$

For up quarks with $x_f = 1$, one has

$$
\mathcal{L}_{(u-a)} = \left(\frac{1}{2f_a}\right) \partial_\mu a \,\bar{u} \gamma^\mu \gamma_5 u \,. \tag{41}
$$

We continue to the second term

$$
II' = \bar{f}e^{-i(\frac{x_f}{2f_a})\gamma_5 a}\gamma^{\mu}P_{\mu}^f e^{-i(\frac{x_f}{2f_a})\gamma_5 a}f = \bar{f}\gamma^{\mu}P_{\mu}^f f = II
$$
\n(42)

This means that there are not new couplings, and this term just gives couplings between gauge boson and two fermions. Thus the axion - fermion derivative couplings have the form

$$
\mathcal{L}_{(f-a)} = + \left(\frac{1}{f_a}\right) \partial_\mu a \left[\bar{d} \mathbf{c}_d \gamma^\mu \gamma_5 d + \bar{u} \mathbf{c}_u \gamma^\mu \gamma_5 u + \bar{T} \mathbf{c}_T \gamma^\mu \gamma_5 T + \bar{D}_\alpha \mathbf{c}_{D_\alpha} \gamma^\mu \gamma_5 D_\alpha \right]
$$
(43)

$$
+ \bar{l} \mathbf{c}_l \gamma^\mu \gamma_5 l + I_\nu \bar{\nu}_a \mathbf{c}_\nu \gamma^\mu \gamma_5 \nu_a + \frac{1}{2} \bar{N}_a \mathbf{c}_{N_a} \gamma^\mu P_R N_a \bigg] \ . \tag{44}
$$

In [\(44\)](#page-7-0), for the coefficients $(c_f, f = d, u, \cdots N_R)$, one has to count the number of color, flavor indexes and PQ charge $Q_A(f)$. In the model under consideration, these coefficients have the form

Our result coincides with that in Ref. [\[34](#page-15-25)].

Note that the above derivative couplings are completely different from usual Yukawa ones in Ref. [\[27](#page-15-16)].

C. Axion-scalar couplings

We continue to scalar sector. For complex scalar φ , its Lagrangian is

$$
\mathcal{L}_{\varphi} = (D^{\mu} \varphi)^{\dagger} D_{\mu} \varphi = [(\partial_{\mu} - i P^{\varphi}_{\mu}) \varphi]^{\dagger} (\partial^{\mu} - i P^{\varphi \mu}) \varphi \n= \partial^{\mu} \varphi^{\dagger} \partial_{\mu} \varphi - i \partial_{\mu} \varphi^{\dagger} P^{\varphi \mu} \varphi + i \varphi^{\dagger} P^{\varphi \mu} \partial_{\mu} \varphi + \varphi^{\dagger} P^{\varphi} P^{\varphi \mu} \varphi \equiv \mathbf{A} \mathbf{1} + \mathbf{A} \mathbf{2} + \mathbf{A} \mathbf{3}.
$$
\n(45)

Within PQ transformation in Eq. [\(14\)](#page-3-6), one has

$$
\mathbf{A1} = \partial^{\mu} \left(\varphi^{\dagger} e^{i \left(\frac{a}{2f_a} \right) Q_A} \right) \partial_{\mu} \left(e^{-i \left(\frac{a}{2f_a} \right) Q_A} \varphi \right) = \partial^{\mu} \left(\varphi^{\dagger} e^{i \left(\frac{x \varphi}{2f_a} \right) a} \right) \partial_{\mu} \left(e^{-i \left(\frac{x \varphi}{2f_a} \right) a} \varphi \right)
$$

\n
$$
= \left[\partial_{\mu} \varphi^{\dagger} . e^{i \left(\frac{x \varphi}{2f_a} \right) a} + \varphi^{\dagger} e^{i \left(\frac{x \varphi}{2f_a} \right) a} (i) \left(\frac{x \varphi}{2f_a} \right) \partial_{\mu} a \right] \left[i \left(\frac{x \varphi}{2f_a} \right) e^{-i \left(\frac{x \varphi}{2f_a} \right) a} \partial^{\mu} a \varphi + e^{-i \left(\frac{x \varphi}{2f_a} \right) a} \partial^{\mu} \varphi \right]
$$

\n
$$
= \partial_{\mu} \varphi^{\dagger} \partial^{\mu} \varphi + \left(\frac{x \varphi}{2f_a} \right)^2 \partial_{\mu} a \partial^{\mu} a \varphi^{\dagger} \varphi - i \left(\frac{x \varphi}{2f_a} \right) \partial_{\mu} a (\partial^{\mu} \varphi^{\dagger} \varphi - \varphi^{\dagger} \partial^{\mu} \varphi)
$$

\n
$$
= \text{kinetic term of } \varphi + \left(\frac{x \varphi}{2f_a} \right)^2 \partial_{\mu} a \partial^{\mu} a \varphi^{\dagger} \varphi - i \left(\frac{x \varphi}{2f_a} \right) \partial_{\mu} a (\partial^{\mu} \varphi^{\dagger} \varphi - \varphi^{\dagger} \partial^{\mu} \varphi) \tag{46}
$$

Applying Eq. (46) for the model under consideration, one gets the first term in (46) is kinematic for scalars. There are quartic couplings of two axions with two neutral scalars and triple couplings of one axion to two neutral scalars. Namely

$$
\mathbf{A1} = \text{kinetic term of } H + \left(\frac{1}{f_a}\right)^2 \partial_\mu a \partial^\mu a \left(\sum_{H=\eta_1^0, \eta_3^0, \rho_2^0}^{x_1^0, x_3^0, \phi} H^* H\right) - i \left(\frac{x_\varphi}{2f_a}\right) \partial^\mu a \sum_{D=\eta, \chi}^{K=\rho^0 \phi^0} \left[D^* \stackrel{\leftrightarrow}{\partial_\mu} D - K^* \stackrel{\leftrightarrow}{\partial_\mu} K\right]
$$
(47)

The explicit results are

$$
\mathbf{A1331} \equiv \mathbf{A1}^{331} = \text{kinetic term of scalars} + \left(\frac{1}{f_a}\right)^2 \partial_\mu a \partial^\mu a \left(\eta_1^{0*} \eta_1^0 + \eta_3^{0*} \eta_3^0 + \chi_1^{0*} \chi_1^0 + \chi_3^{0*} \chi_3^0 + \rho_2^{0*} \rho_2^0 + \phi^{0*} \phi^0\right) - i \left(\frac{1}{f_a}\right) \partial^\mu a \left[\eta_1^{0*} \stackrel{\leftrightarrow}{\partial_\mu} \eta_1^0 + \eta_3^{0*} \stackrel{\leftrightarrow}{\partial_\mu} \eta_3^0 + \chi_1^{0*} \stackrel{\leftrightarrow}{\partial_\mu} \chi_1^0 + \chi_3^{0*} \stackrel{\leftrightarrow}{\partial_\mu} \chi_3^0 - \rho_2^{0*} \stackrel{\leftrightarrow}{\partial_\mu} \rho_2^0 - \phi^{0*} \stackrel{\leftrightarrow}{\partial_\mu} \phi^0\right].
$$
\n(48)

Next, for the second term in Eq. [\(45\)](#page-7-2)

$$
\mathbf{A2} = -i\partial_{\mu} \left[\varphi^{\dagger} e^{i\left(\frac{x_{\varphi}}{2f_{a}}\right)a} \right] P^{\varphi\mu} e^{-i\left(\frac{x_{\varphi}}{2f_{a}}\right)a} \varphi + i\varphi^{\dagger} e^{i\left(\frac{x_{\varphi}}{2f_{a}}\right)a} P^{\varphi\mu} \partial_{\mu} \left[e^{-i\left(\frac{x_{\varphi}}{2f_{a}}\right)a} \varphi \right]
$$

\n
$$
= -i \left[(\partial_{\mu} \varphi^{\dagger}) e^{+i\left(\frac{x_{\varphi}}{2f_{a}}\right)a} + i\varphi^{\dagger} \left(\frac{x_{\varphi}}{2f_{a}}\right) \partial_{\mu} a e^{i\left(\frac{x_{\varphi}}{2f_{a}}\right)a} \right] P^{\varphi\mu} e^{-i\left(\frac{x_{\varphi}}{2f_{a}}\right)a} \varphi
$$

\n
$$
+ i\varphi^{\dagger} e^{i\left(\frac{x_{\varphi}}{2f_{a}}\right)a} P^{\varphi\mu} \left[-i\left(\frac{x_{\varphi}}{2f_{a}}\right) \partial_{\mu} a e^{-i\left(\frac{x_{\varphi}}{2f_{a}}\right)a} \varphi + e^{-i\left(\frac{x_{\varphi}}{2f_{a}}\right)a} \partial_{\mu} \varphi \right]
$$

\n
$$
= -i \left[\partial^{\mu} \varphi^{\dagger} + i\varphi^{\dagger} \left(\frac{x_{\varphi}}{2f_{a}}\right) \partial^{\mu} a \right] P^{\varphi}_{\mu} \varphi
$$

\n
$$
- i\varphi^{\dagger} P^{\varphi}_{\mu} \left[i\left(\frac{x_{\varphi}}{2f_{a}}\right) \partial^{\mu} a \varphi - \partial^{\mu} \varphi \right]
$$

\n
$$
= -i \left(\partial^{\mu} \varphi^{\dagger} P^{\varphi}_{\mu} \varphi - \varphi^{\dagger} P^{\varphi}_{\mu} \partial^{\mu} \varphi \right) + 2 \left(\frac{x_{\varphi}}{2f_{a}} \right) (\partial^{\mu} a) \left[\varphi^{\dagger} P^{\varphi}_{\mu} \varphi \right].
$$

\n(4

The first term in Eq. [\(49\)](#page-8-0) is ordinary one in the 3-3-1 model (without axion's participation). The second term is quartic couplings of axion, gauge and two scalar bosons.

There are no new coupling in A3, or other word speaking, A3 contains ordinary couplings of scalars to gauge bosons. In summary, new couplings of axion with neutral scalars are (including operators of up to dimension 5)

$$
\mathcal{L}_{(aS)} = \left(\frac{1}{f_a}\right)^2 \partial_\mu a \partial^\mu a \left(\sum_{H=\eta_1^0, \eta_3^0, \rho_2^0}^{x_1^0, x_3^0, \rho_3^0} H^*H\right)
$$

\n
$$
- i \left(\frac{x_\varphi}{2f_a}\right) (\partial^\mu a) \sum_{D=\eta, \chi}^{K=\rho^0 \phi^0} \left[D^* \stackrel{\leftrightarrow}{\partial_\mu} D - K^* \stackrel{\leftrightarrow}{\partial_\mu} K\right]
$$

\n
$$
+ 2 \left(\frac{x_\varphi}{2f_a}\right) (\partial^\mu a) H^\dagger P_\mu^H H
$$

\n
$$
\equiv \mathcal{L}(a a H H) + \mathcal{L}(a H H) + \mathcal{L}(a G H H), \qquad (50)
$$

where G is labeled for gauge bosons. For more detail of axion couplings, the reader is referred to Refs. [\[2](#page-14-2), [31,](#page-15-21) [35,](#page-15-24) [36](#page-15-26)].

D. Interactions of axion to scalar and gauge bosons

We proceed now explicit form of the above terms. Note that in Eq. (50) , there are not only couplings of axion to scalar bosons, but also to gauge bosons. Let us consider quartic couplings

$$
\mathcal{L}(aaHH) = \left(\frac{1}{f_a}\right)^2 \partial_\mu a \partial^\mu a \left(\sum_{H=\eta_1^0,\eta_3^0,\rho_2^0}^{x_1^0,\phi_3^0,\phi_4^0} H^*H\right)
$$
\n
$$
= \left(\frac{1}{f_a}\right)^2 \partial_\mu a \partial^\mu a \left(\eta_1^{0*}\eta_1^0 + \eta_3^{0*}\eta_3^0 + \chi_1^{0*}\chi_1^0 + \chi_3^{0*}\chi_3^0 + \rho_2^{0*}\rho_2^0 + \phi^*\phi\right)
$$
\n
$$
= \frac{1}{2} \left(\frac{1}{f_a}\right)^2 \partial_\mu a \partial^\mu a \left\{ \left[(v_\eta + R_\eta^1)^2 + (I_\eta^1)^2 + (v_\chi + R_\chi^3)^2 + (I_\chi^3)^2 + (v_\rho + R_\rho)^2 + (I_\rho)^2 + (v_\phi + R_\phi)^2 \right] \right.
$$
\n
$$
+ 2(\eta_3^{0*}\eta_3^0 + \chi_1^{0*}\chi_1^0) \left.\right\}
$$
\n
$$
= \frac{1}{2} \left(\frac{1}{f_a}\right)^2 \partial_\mu a \partial^\mu a \left[(v_\eta^2 + v_\rho^2 + v_\chi^2 + v_\phi^2) + (I_\chi^3 + v_\phi R_\phi) \right.
$$
\n
$$
+ 2(v_\eta R_\eta^1 + v_\rho R_\rho + v_\chi R_\chi^3 + v_\phi R_\phi) + (R_\eta^1)^2 + (I_\eta^1)^2 + (R_\rho^1)^2 + (I_\rho^2)^2 + (I_\chi^3)^2 + (I_\chi^3)^2 + (I_\chi^3)^2 + 2(\eta_3^{0*}\eta_3^0 + \chi_1^{0*}\chi_1^0) \right]. \quad (52)
$$

From Eq. [\(51\)](#page-9-0), it follows that to get a right form of axion kinetic term, the following condition is

$$
f_a^2 = v_\eta^2 + v_\rho^2 + v_\chi^2 + v_\phi^2. \tag{53}
$$

Thus one gets finally

$$
\mathcal{L}(aaHH) = \frac{1}{2}\partial_{\mu}a\partial^{\mu}a + \left(\frac{1}{f_a}\right)^2\partial_{\mu}a\partial^{\mu}a \left\{v_{\eta}R_{\eta}^1 + v_{\rho}R_{\rho} + v_{\chi}R_{\chi}^3 + v_{\phi}R_{\phi} + \eta_3^{0*}\eta_3^0 + \chi_1^{0*}\chi_1^0\right\}
$$
(54)

$$
+\frac{1}{2}\left[(R_{\eta}^{1})^{2} + (I_{\eta}^{1})^{2} + (R_{\rho})^{2} + (I_{\rho})^{2} + (R_{\chi}^{3})^{2} + (I_{\chi}^{3})^{2} + (R_{\phi})^{2} \right] \} \tag{55}
$$

In Eq. [\(54\)](#page-9-1), there exist triple couplings of two axions to one CP-even scalar. One of them with the strength of coupling is $\partial_{\mu}a\partial^{\mu}a R_{\phi} = v_{\phi}(f_a)^{-2} \sim (f_a)^{-1}$ - the coupling of two axions to inflaton. We should add this interaction for the future study, since $\propto (f_a)^{-1}$, while terms in the second line can be neglected.

We next consider triple couplings of an axion to two scalars.

$$
\mathcal{L}(aHH) = -i\left(\frac{1}{f_a}\right)\partial^{\mu}a\left(\eta_1^{0*} \stackrel{\leftrightarrow}{\partial_{\mu}} \eta_1^{0} + \eta_3^{0*} \stackrel{\leftrightarrow}{\partial_{\mu}} \eta_3^{0} + \chi_1^{0*} \stackrel{\leftrightarrow}{\partial_{\mu}} \chi_1^{0} + \chi_3^{0*} \stackrel{\leftrightarrow}{\partial_{\mu}} \chi_3^{0}
$$

\n
$$
- \rho_2^{0*} \stackrel{\leftrightarrow}{\partial_{\mu}} \rho_2^{0} - \phi^{0*} \stackrel{\leftrightarrow}{\partial_{\mu}} \phi^{0}\right)
$$

\n
$$
= -i\left(\frac{1}{f_a}\right)\partial^{\mu}a\left(-iv_{\eta}\partial_{\mu}I_{\eta}^{1} + iI_{\eta}^{1}\partial_{\mu}R_{\eta}^{1} - iR_{\eta}^{1}\partial_{\mu}I_{\eta}^{1} - iv_{\chi}\partial_{\mu}I_{\chi}^{3} + iI_{\chi}^{3}\partial_{\mu}R_{\chi}^{3} - iR_{\chi}^{3}\partial_{\mu}I_{\chi}^{3}
$$

\n
$$
-iv_{\rho}\partial_{\mu}I_{\rho} + iI_{\rho}\partial_{\mu}R_{\rho} - iR_{\rho}\partial_{\mu}I_{\rho}
$$

\n
$$
+ \eta_3^{0*} \stackrel{\leftrightarrow}{\partial_{\mu}} \eta_3^{0} + \chi_1^{0*} \stackrel{\leftrightarrow}{\partial_{\mu}} \chi_1^{0}\right)
$$

\n
$$
-i\left(\frac{1}{f_a}\right)\partial^{\mu}a\left(v_{\eta}\partial_{\mu}I_{\eta}^{1} + v_{\chi}\partial_{\mu}I_{\chi}^{3} + v_{\rho}\partial_{\mu}I_{\rho}\right)
$$

\n
$$
-i\left(\frac{1}{f_a}\right)\partial^{\mu}a\left(iI_{\eta}^{1}\partial_{\mu}R_{\eta}^{1} + iI_{\rho}\partial_{\mu}R_{\rho} + iI_{\chi}^{3}\partial_{\mu}R_{\chi}^{3}
$$

\n
$$
-iR_{\eta}^{1}\partial_{\mu}I_{\eta}^{1} - iR_{\chi}^{3}\partial_{\mu}I_{\chi}^{3} - iR_{\rho}\partial_{\mu}I_{\rho}
$$

where the mixing of axion with CP-odd scalars is determined as

$$
\mathcal{L}_{(aA)} = -\left(\frac{1}{f_a}\right)\partial^\mu a \left(v_\eta \partial_\mu I_\eta^1 + v_\chi \partial_\mu I_\chi^3 + v_\rho \partial_\mu I_\rho\right). \tag{57}
$$

11

The terms in Eq. [\(57\)](#page-9-2) are some kind of mixing between axion and Goldstone bosons eaten by neutral gauge bosons. This unwanted mixing could be rotated away by making $U(1)$ rotation [\[9](#page-15-5)]. Hence one has coupling of axion to two scalar fields given below

$$
\mathcal{L}_{(aSS)} = -i \left(\frac{1}{f_a} \right) \partial^{\mu} a \left(i I_{\eta}^1 \partial_{\mu} R_{\eta}^1 + i I_{\rho} \partial_{\mu} R_{\rho} + i I_{\chi}^3 \partial_{\mu} R_{\chi}^3 \n- i R_{\eta}^1 \partial_{\mu} I_{\eta}^1 - i R_{\chi}^3 \partial_{\mu} I_{\chi}^3 - i R_{\rho} \partial_{\mu} I_{\rho} \n+ \eta_3^{0*} \stackrel{\leftrightarrow}{\partial_{\mu}} \eta_3^0 + \chi_1^{0*} \stackrel{\leftrightarrow}{\partial_{\mu}} \chi_1^0
$$
\n(58)

We close this part by considering

$$
\mathcal{L}(aGHH) = 2\left(\frac{x_{\varphi}}{2f_a}\right)\partial^{\mu}a\,\varphi^{\dagger}P_{\mu}^{\varphi}\varphi
$$

\n
$$
= 2\left(\frac{x_{\varphi}}{2f_a}\right)\partial^{\mu}a\left[\eta^{\dagger}P_{\mu}^{\eta}\eta + \chi^{\dagger}P_{\mu}^{\chi}\chi + \rho^{\dagger}P_{\mu}^{\rho}\rho\right]
$$

\n
$$
= 2\left(\frac{g}{f_a}\right)\partial^{\mu}a\left\{\left(W_{3\mu} + \frac{1}{\sqrt{3}}W_{8\mu} - \frac{1}{3}t\sqrt{\frac{2}{3}}B_{\mu}\right)v_{\eta}^{2} - \left(W_{3\mu} - \frac{1}{\sqrt{3}}W_{8\mu} - \frac{2}{3}t\sqrt{\frac{2}{3}}B_{\mu}\right)v_{\rho}^{2} - \left(\frac{2}{\sqrt{3}}W_{8\mu} + \frac{1}{3}t\sqrt{\frac{2}{3}}B_{\mu}\right)v_{\chi}^{2}\right\}
$$

\n
$$
+ 2\left(\frac{g}{f_a}\right)\partial^{\mu}a\left\{\left(W_{3\mu} + \frac{1}{\sqrt{3}}W_{8\mu} - \frac{1}{3}t\sqrt{\frac{2}{3}}B_{\mu}\right)[2v_{\eta}R_{\eta}^{1} + (R_{\eta}^{1})^{2} + (I_{\eta}^{1})^{2}] + \left(-W_{3\mu} + \frac{1}{\sqrt{3}}W_{8\mu} + \frac{2}{3}t\sqrt{\frac{2}{3}}B_{\mu}\right)[2v_{\rho}R_{\rho}^{2} + (R_{\rho})^{2} + (I_{\rho})^{2}] + \left(-\frac{2}{\sqrt{3}}W_{8\mu} - \frac{1}{3}t\sqrt{\frac{2}{3}}B_{\mu}\right)[2v_{\chi}R_{\chi}^{3} + (R_{\chi}^{3})^{2} + (I_{\chi}^{3})^{2}] + \left(-\frac{2}{\sqrt{3}}W_{8\mu} - \frac{1}{3}t\sqrt{\frac{2}{3}}B_{\mu}\right)\eta_{3}^{0*}\eta_{3}^{0} + \left(W_{3\mu} + \frac{1}{\sqrt{3}}W_{8\mu} - \frac{1}{3}t\sqrt{\frac{2}{3}}B_{\
$$

where the first term in Eq. [\(59\)](#page-10-0) is term mix axion with weak gauge bosons

$$
\mathcal{L}_{(aW)} = 2\left(\frac{g}{f_a}\right)\partial^{\mu}a \left[\left(\frac{Z_{\mu}}{cw} + Z_{\mu}' \frac{1 - t_W^2}{\sqrt{3 - t_W^2}}\right) v_{\eta}^2 + \left(\frac{Z_{2\mu}'}{c_W^2 \sqrt{3 - t_W^2}} - \frac{Z_{\mu}}{cw}\right) v_{\rho}^2 + \left(-\frac{2Z_{\mu}'}{\sqrt{3 - t_W^2}}\right) v_{\chi}^2 \right] \n= vev^2 \cdot \partial_{\mu}a Z^{\mu},
$$
\n(60)

and

$$
\mathcal{L}_{aGHH} = 2\left(\frac{g}{f_a}\right)\partial^{\mu}a \left[\left(\frac{Z_{\mu}}{cw} + Z_{\mu}'\frac{1-t_W^2}{\sqrt{3-t_W^2}}\right)[2v_{\eta}R_{\eta}^1 + (R_{\eta}^1)^2 + (I_{\eta}^1)^2] + \left(\frac{Z_{2\mu}'}{c_W^2\sqrt{3-t_W^2}} - \frac{Z_{\mu}}{cw}\right)[2v_{\rho}R_{\rho}^2 + (R_{\rho})^2 + (I_{\rho})^2] + \left(-\frac{2Z_{\mu}'}{\sqrt{3-t_W^2}}\right)[2v_{\chi}R_{\chi}^3 + (R_{\chi}^3)^2 + (I_{\chi}^3)^2] + \left(-\frac{2Z_{\mu}'}{\sqrt{3-t_W^2}}\right)\eta_3^{0*}\eta_3^0 + \left(\frac{Z_{\mu}}{cw} + Z_{\mu}'\frac{1-t_W^2}{\sqrt{3-t_W^2}}\right)\chi_1^{0*}\chi_1^0\right] = \text{Couplings}(aGHH), \tag{61}
$$

where $t = \frac{3\sqrt{2}tw}{\sqrt{3-t_W^2}}$. We also use the limit $v_x^2 \gg v^2$, which result in the following relations between physical basis (A_μ, Z_μ, Z'_μ)

and the original gauge boson states [\[17](#page-15-27)]:

$$
W_{3\mu} = A_{\mu} s_W + c_W Z_{\mu},
$$

\n
$$
W_{8\mu} = (-A_{\mu} c_W + s_W Z_{\mu}) \frac{t_W}{\sqrt{3}} + \frac{\sqrt{3 - t_W^2} Z_{\mu}'}{\sqrt{3}},
$$

\n
$$
B_{\mu} = -(-A_{\mu} c_W + s_W Z_{\mu}) \frac{\sqrt{3 - t_W^2}}{\sqrt{3}} + \frac{t_W Z_{\mu}'}{\sqrt{3}},
$$
\n(62)

In Eq. [\(61\)](#page-10-1), there are couplings, up to dimension 5, between axion, gauge boson with one scalar or two scalars. To close this section, it is worth noting that unlike photon, the axion does not couple to charged scalar and gauge bosons.

V. TOTAL AXION LAGRANGIAN

The total part concerned to axion is given below

$$
\mathcal{L}_a = \frac{1}{2} \partial_\mu a \partial^\mu a - \frac{1}{2} m_{a\sigma}^2 a^2 + c_{GG} \frac{\alpha_s}{4\pi} \frac{a}{2f_a} G\tilde{G} + c_{WW} \frac{\alpha_2}{4\pi} \frac{a}{2f_a} W \tilde{W} + c_{BB} \frac{\alpha_1}{4\pi} \frac{a}{2f_a} B\tilde{B}
$$
(63)

$$
+ \frac{\partial^{\mu} a}{2f_a} \left(\sum_{f=u,d,T,D}^{l,\nu} \bar{\psi}_f c_f \gamma_{\mu} \gamma_5 \psi + \frac{1}{2} \bar{N}_a \mathbf{c}_{N_a} \gamma^{\mu} P_R N_a \right) - (\bar{q}_L M_a q_R + \text{H.c.}) \tag{64}
$$

$$
+\left(\frac{1}{f_a}\right)^2 \partial_\mu a \partial^\mu a \left(\sum_{H=\eta_1^0,\eta_3^0,\rho_2^0}^{x_1^0,x_3^0,\phi} H^*H\right) \qquad ;\tag{65}
$$

$$
- i \left(\frac{x_{\varphi}}{2f_a}\right) \partial^{\mu} a \sum_{D=\eta,\chi}^{K=\rho^0 \phi^0} \left[D^* \stackrel{\leftrightarrow}{\partial_{\mu}} D - K^* \stackrel{\leftrightarrow}{\partial_{\mu}} K \right]
$$
(66)

$$
+ 2\left(\frac{x_{\varphi}}{2f_a}\right)\partial^{\mu}a \sum_{H=\eta_1^0,\eta_3^0,\rho_2^0}^{\chi_1^0,\chi_3^0,\phi} H^{\dagger}P_{\mu}^H H,
$$
\n
$$
\tag{67}
$$

where $H^*H = \frac{1}{2}[(v_H + R_H)^2 + I_H^2]$.

The mass of xion m_{ao} in the first line is acquired from mixing of the particle with π^0 and η pseudoscalars. Term in Eq. [\(65\)](#page-11-2) provides kinematic term of axion with f_a^2 is sum over all VEVs of scalars. Beside ordinary dimension 6 terms, there are dimension 5 terms associated with VEVs in the form $\left(\frac{1}{f_a}\right)$ $\int_{0}^{2} v_{H} \partial_{\mu} a \partial^{\mu} a H$. When $H = \phi$, then we have a key term proportional to $1/f_a$ only: $\left(\frac{1}{f_a}\right)^2 v_\phi \partial_\mu a \partial^\mu a \Phi \sim \left(\frac{1}{f_a}\right) \Phi \partial_\mu a \partial^\mu a$. The last two terms in the above expression give couplings of axion with scalar and gauge fields. The last term in [\(67\)](#page-11-2) contains unwanted mixing of axion with Goldstone bosons eaten by massive Z and Z' bosons. Fortunately, this mixing can be rotated away by the $U(1)$ rotation. It is emphasized that the last term in the second line, the right handed Majorana neutrinos are very heavy with mass around 10^7 GeV.

The up to $1/f_a$ couplings of axion field are presented in Table [V.](#page-11-3) However, the coupling of two axions with inflation is included in bottom of the table.

Coupling	Expression	Vertex	Figure	Note
$a \, F \, F$	$\left(\frac{x_f}{f_a}\right) \partial_\mu\, a\, \bar{f} \gamma^\mu \gamma_5 f$	$I_{(c,f)}\frac{x_f}{f_a}k\gamma_5$		$2\frac{3}{2f_c}P_R\rlap{/}k$ for N_R
$a\,a\,R_\phi$	$\left(\frac{x_{\varphi}}{f_a}\right)^2\,v_{\phi}\partial^{\mu}a\partial_{\mu}a\,R_{\phi}\,\left.-2i\left(\frac{x_{\varphi}}{f_a}\right)^2\,v_{\phi}k.q$			$-2i\left(\frac{x_{\varphi}^2}{f_a}\right) k.q$ large

Table V: k, q are ingoing momentum of axion and scalar, A, S and Z are CP -odd, CP -even and neural gauge bosons, respectively

A. Elimination of mixing between axion and Goldstone bosons G_Z and $G_{Z'}$

As seen in sub-section [IV D,](#page-9-3) there are terms of mixing axion with Goldstone bosons G_Z and $G_{Z'}$. To avoid this trouble [\[9](#page-15-5)], we rotate scalar triplets by opposite $U(1)_{PQ}$ as follows

$$
\chi \rightarrow \chi' = e^{-i\left(\frac{a}{2f_a}\right)} \chi,
$$

\n
$$
\eta \rightarrow \eta' = e^{-i\left(\frac{a}{2f_a}\right)} \eta,
$$

\n
$$
\rho \rightarrow \rho' = e^{i\left(\frac{a}{2f_a}\right)} \rho.
$$
\n(68)

Of course, the quarks have to be changed to keep Yukawa interactions invariant. Note that the singlet ϕ in unchanged. Then, Eq. [\(67\)](#page-11-2) becomes final Lagrangian

$$
\mathcal{L}_{a} = \frac{1}{2} \partial_{\mu} a \partial^{\mu} a - \frac{1}{2} m_{a\sigma}^{2} a^{2} + c_{GG} \frac{\alpha_{s}}{4\pi} \frac{a}{2f_{a}} G\tilde{G} + c_{WW} \frac{\alpha_{2}}{4\pi} \frac{a}{2f_{a}} W\tilde{W} + c_{BB} \frac{\alpha_{1}}{4\pi} \frac{a}{2f_{a}} B\tilde{B}
$$

$$
+ \frac{1}{2} \frac{\partial^{\mu} a}{f_{a}} \left(\sum_{f=u,d,T,D}^{l,\nu} \bar{\psi}_{f} c_{f} \gamma_{\mu} \gamma_{5} \psi + \frac{1}{2} \bar{N}_{a} \mathbf{c}_{N_{a}} \gamma^{\mu} P_{R} N_{a} \right) - (\bar{q}_{L} M_{a} q_{R} + \text{H.c.}) \tag{69}
$$

$$
+ \frac{1}{2} \left(\frac{1}{f_a}\right)^2 \partial_\mu a \partial^\mu a \left(v_\phi^2 + 2v_\phi R_\phi + R_\phi^2\right). \tag{70}
$$

To have correct axion's kinetic term, Eq. [\(70\)](#page-12-0) leads to condition

$$
f_a = v_\phi. \tag{71}
$$

The point is worth noting that the terms in last line (doubly $(\partial_\mu a \partial^\mu a)$ derivative coupling of axion to inflaton) are characteristic for the model under consideration. Note that this coupling is dimension 5 too. As mentioned in Ref. [\[9](#page-15-5)], the axion has only two kinds of couplings: derivative couplings to fermions and anomalous couplings to gauge bosons $aG\tilde{G}$. It is worth noting that the new effects mainly happen in the energy region from 10^7 GeV to 10^{11} GeV, namely in the region from mass of Majorana right-handed neutrino (N_R) to mass of inflaton Φ .

B. Axion mass from mixing with π^0

In the frameworks of chiral perturbative Lagrangian in the energy scale ≤ 1 GeV [\[49](#page-15-28), [50](#page-15-29)], the axion's mass was derived [\[2](#page-14-2), [9](#page-15-5), [37](#page-15-30)[–40](#page-15-31)]. For two quark flavors, the leading order chiral Lagrangian is given by [\[2](#page-14-2)]

$$
\mathcal{L}_{a}^{\chi (LO)} = +\frac{f_{\pi}^{2}}{4} \text{Tr} \left[(D_{\mu} U U)^{\dagger} D_{\mu} U \right] + \frac{\partial_{\mu} a}{2 f_{a}} \frac{1}{2} \text{Tr} \left[c_{q} \sigma^{a} \right] J_{A,\,\mu}^{a} + \frac{f_{\pi}^{2}}{2} B_{0} \text{Tr} \left[M_{a} U^{\dagger} + U M_{a}^{\dagger} \right]. \tag{72}
$$

The terms in the first line connected with kinetic of fields, while the term in the second line provides mass of the fields. Hence we continue with this terms.

$$
UM_a^{\dagger} + M_a U^{\dagger} = UM_q + M_q U^{\dagger} + i \frac{a}{2f_a} \{Q_A, M_q\} U^{\dagger} - 2 \left(\frac{a}{2f_a}\right)^2 U(Q_A^2 M_q + 2Q_A M_q Q_A + M_q Q_A^2) + \cdots,
$$
 (73)

where $M_q = \text{diag}(m_u, m_d)$. Using

$$
U = e^{i\frac{\pi^a \sigma^a}{f_\pi}} = \mathbf{1} \cos\left(\frac{\pi}{f_\pi}\right) + i\frac{\pi^a \sigma^a}{\pi} \sin\left(\frac{\pi}{f_\pi}\right),\tag{74}
$$

where $\pi = \sqrt{(\pi^0)^2 + 2\pi^+ \pi^-}$ and $Q_A = \frac{M_q^{-1}}{\text{Tr}(M_q^{-1})}$, one gets

$$
\mathcal{L}_{amass} = \frac{f_{\pi}^2}{2} B_0 \text{Tr}[M_a U^{\dagger} + U M_a^{\dagger}] = f_{\pi}^2 m_{\pi}^2 \cos\left(\frac{\pi}{f_{\pi}}\right) - \frac{m_a^2}{2} \,. \tag{75}
$$

Here, the axion mass is given by

$$
m_a^2 = \frac{m_\pi^2 f_\pi^2}{f_a^2} \frac{m_u m_d}{(m_u + m_d)^2} \cos\left(\frac{\pi}{f_\pi}\right) \simeq \frac{m_\pi^2 f_\pi^2}{f_a^2} \frac{m_u m_d}{(m_u + m_d)^2} \,. \tag{76}
$$

A consequence is as follows

$$
m_a \simeq 5.7 \left(\frac{10^{12} \text{ GeV}}{f_a}\right) \mu \text{eV} \,. \tag{77}
$$

In a similar way, we can deal with three flavour quark (with s quark). The result is given by

$$
m_a^2 = \frac{1}{2} m_\pi^2 \left(\frac{f_\pi}{f_a}\right)^2 \frac{1}{(m_u + m_d)} \left(\frac{m_u m_d m_s}{m_u m_d + m_u m_s + m_d m_s}\right). \tag{78}
$$

This result coincides with one given in Ref. [\[41](#page-15-32)]

$$
m_a = 4 \frac{f_{\pi} m_{\pi}}{f_a/N} \left[\frac{m_u m_d m_s}{(m_u m_d + m_u m_s + m_d m_s)(m_u + m_d)} \right]^{\frac{1}{2}}
$$

\n
$$
\simeq (1.2 \times 10^{-5} \text{ eV}) \left(\frac{10^{12} \text{ GeV}}{f_a/N} \right), \tag{79}
$$

VI. CONCLUSIONS

In this paper, we have realized PQ formalism of the 3-3-1 model with Cosmological Inflation. The singlet field ϕ takes complex value everywhere, and axion is complex phase in polar coordinates. Then the axion also appears as a phase of PQ transformations. Using the GKS formation, we have constructed PQ charge operator Q_A in terms of diagonal generators of $SU(3)_L$ subgroup. The formula shows that the difference of PQ charges (Q_A) of up and down quarks equal to 2, i.e., $\Delta Q_A = 2$, while for electric charge, as usually $\Delta Q = 1$. For right-handed fermions, electric charges (Q) of left-handed and right-handed are assumed to be equal, while for (Q_A) is the opposite. PQ charges of neutral scalars equal ± 2 , while for charged scalars, it vanishes. In the model under consideration, in contract to photon which couples to charged particles only, while the axion does not couple to charged scalar/gauge bosons. To have correct kinetic term for the axion, the PQ scale f_a is to equal VEV of the singlet scalar boson, namely $f_a = v_\phi$. The derivative couplings of axion to fermions are presented. The point is worth emphasizing that the axion has doubly derivative coupling with scalar playing the role of inflaton. This coupling increases magnitude of coupling to $1/f_a$. It is worth emphasizing that the new effects mainly happen in the energy region from 10^7 GeV to 10^{11} GeV, namely in the region from mass of Majorana right-handed neutrino (N_R) to mass of inflaton Φ . The chiral effective Lagrangian as usually provides axion mass consistent with model-independent prediction.

Acknowledgments

The authors thank D. T. Nhung for useful discussion.

Appendix A: PQ transformation for chiral fermions

Proof of Eq. (11)

$$
f \rightarrow f' = e^{i\left(\frac{x_f}{2f_a}\right)\gamma_5 a} f, \quad \bar{f} \rightarrow \bar{f}' = \bar{f} e^{i\left(\frac{x_f}{2f_a}\right)\gamma_5 a}, \quad \varphi \rightarrow \varphi' = e^{i\left(\frac{x_\varphi}{2f_a}\right)a} \varphi \tag{A1}
$$

$$
f_L \rightarrow f'_L = e^{-i\left(\frac{x_f}{2f_a}\right)a} f_L, \quad \bar{f}_L \rightarrow \bar{f}'_L = \bar{f}_L e^{+i\left(\frac{x_f}{2f_a}\right)a} \tag{A2}
$$

Noting that

$$
\gamma_5 P_L = P_L \gamma_5 = -P_L \rightarrow (\gamma_5)^k P_L = P_L (\gamma_5)^k = (-1)^k P_L ,
$$

$$
\gamma_5 P_R = P_R \gamma_5 = P_R \rightarrow (\gamma_5)^k P_R = P_R (\gamma_5)^k = P_R .
$$
 (A3)

As a consequence, we have

$$
P_{L}e^{i\left(\frac{x_{f}}{2f_{a}}\right)\gamma_{5}a} = \left[\sum_{k=0}^{\infty} \frac{P_{L}(\gamma_{5})^{k}}{k!} \times \left(\frac{ix_{f}a}{2f_{a}}\right)^{k}\right] = \left[\sum_{k=0}^{\infty} \frac{(-1)^{k}P_{L}}{k!} \times \left(\frac{ix_{f}a}{2f_{a}}\right)^{k}\right] = e^{-i\left(\frac{x_{f}}{2f_{a}}\right)a}P_{L} = e^{i\left(\frac{x_{f}}{2f_{a}}\right)\gamma_{5}a}P_{L}
$$

\n
$$
\Rightarrow f_{L} \rightarrow f'_{L} = P_{L}f' = P_{L}\left(e^{i\left(\frac{x_{f}}{2f_{a}}\right)\gamma_{5}a}f\right) = e^{-i\left(\frac{x_{f}}{2f_{a}}\right)a}P_{L}f = e^{-i\left(\frac{x_{f}}{2f_{a}}\right)a}f_{L},
$$

\n
$$
P_{R}e^{i\left(\frac{x_{f}}{2f_{a}}\right)\gamma_{5}a} = e^{i\left(\frac{x_{f}}{2f_{a}}\right)\gamma_{5}a}P_{R} = e^{i\left(\frac{x_{f}}{2f_{a}}\right)a}P_{R} \Rightarrow f_{R} \rightarrow f'_{R} = e^{i\left(\frac{x_{f}}{2f_{a}}\right)a}f_{R}.
$$
\n(A4)

The Dirac conjugation:

$$
\overline{f} \rightarrow \overline{f'} = (f')^{\dagger} \gamma_0 = \left(e^{i\left(\frac{x_f}{2f_a}\right)\gamma_5 a} f \right)^{\dagger} \gamma_0 = f^{\dagger} e^{-i\left(\frac{x_f}{2f_a}\right)\gamma_5 a} \gamma_0 = f^{\dagger} \gamma_0 e^{i\left(\frac{x_f}{2f_a}\right)\gamma_5 a} = \overline{f} e^{i\left(\frac{x_f}{2f_a}\right)\gamma_5 a},
$$
\n
$$
\overline{f_L} \rightarrow \overline{f'_L} = \left(e^{i\left(\frac{x_f}{2f_a}\right)\gamma_5 a} f \right)^{\dagger} P_L \gamma_0 = f^{\dagger} e^{-i\left(\frac{x_f}{2f_a}\right)\gamma_5 a} P_L \gamma_0 = f^{\dagger} P_L \gamma_0 \times e^{i\left(\frac{x_f}{2f_a}\right)a} = \overline{f_L} e^{i\left(\frac{x_f}{2f_a}\right)a}
$$
\n
$$
\overline{f_R} \rightarrow \overline{f'_R} = \left(e^{i\left(\frac{x_f}{2f_a}\right)\gamma_5 a} f \right)^{\dagger} P_R \gamma_0 = f^{\dagger} e^{-i\left(\frac{x_f}{2f_a}\right)\gamma_5 a} P_R \gamma_0 = f^{\dagger} P_R \gamma_0 \times e^{-i\left(\frac{x_f}{2f_a}\right)a} = \overline{f_R} e^{-i\left(\frac{x_f}{2f_a}\right)a}.
$$
\n(A5)

This means that in vector fermion, PQ transformation has γ_5 , but in chiral form without γ_5 . Therefore, the mass matrices have usual form as in ordinary in the 3-3-1 model [\[27\]](#page-15-16).

Appendix B: Formula of PQ charge operator

The PQ charges given in Table [II](#page-3-5) allows us to write some nice formula as generalized lepton number in [\[45](#page-15-33), [46](#page-15-34)]. Let us write PQ charge operator in diagonal operators as follows (for left-handed fermions sitting in non-singlets).

$$
Q_A = \alpha T_3 + \beta T_8 + \delta X_{pq}, \qquad (B1)
$$

Applying for Q3, one gets

$$
\alpha = +2, \quad \beta = -\frac{2}{\sqrt{3}}, \quad \delta \mathcal{X}_{pq}(Q_3) = +\frac{1}{3}
$$
\n(B2)

Assuming $\delta = 1$, one gets $\mathcal{X}_{pq}(Q_3) = +\frac{1}{3}$. Hence

$$
Q_A = 2T_3 - \frac{2}{\sqrt{3}}T_8 + \mathcal{X}_{pq} \,. \tag{B3}
$$

For all fermion triplets, one has $\mathcal{X}_{pq}(3) = \frac{1}{3}$.

Note that the above formula is applicable for left-handed fermions, for right-handed fermions just take opposite For scalars

$$
\mathcal{X}_{pq}(\chi,\eta) = \frac{4}{3}, \quad \mathcal{X}_{pq}(\rho) = -\frac{2}{3}.
$$
 (B4)

The PQ charge of the singlet ϕ follows from Yukawa coupling and equal 2. Let us connect PQ charge with electric one being as follows

$$
Q = T_3 - \frac{1}{\sqrt{3}} T_8 + N \,,\tag{B5}
$$

From Eqs. $(B3)$ and $(B5)$, it follows

$$
Q_A = 2Q + \mathcal{X}_{pq} - 2N. \tag{B6}
$$

For singlets, their values are given from Yukawa interactions as follows

$$
\mathcal{X}_{pq}(N_R) = -1 \,, \quad \mathcal{X}_{pq}(f_R) = -Q_A(f_L) \,, \quad \mathcal{X}_{pq}(\phi) = 2 \,. \tag{B7}
$$

For minimal 3-3-1 model, PQ symmetry may be not suitable because of the Landau pole around 5 TeV. Note that electric charges of up and down elements differ by one unit, while PQ charges of the above elements differ by two. That is why the factor 2 in Eq. $(B4)$.

Therefore we can write PQ transformation for left/right handed fermion and scalar as follows

$$
\Psi(x) \rightarrow e^{i\frac{a}{2f_a}Q_A}\Psi(x), \quad \Psi(x) = f_L, \phi(\text{scalar fields}), \tag{B8}
$$

$$
\overline{\Psi(x)} \rightarrow \overline{\Psi(x)} e^{-i\frac{a}{2f_a}Q_A} \tag{B9}
$$

- [3] C. B. Adams et al, Snowmass 2021 White Paper, Axion Dark Matter, arXiv:2203.14923 [hep-ex].
- [4] Bartram C, et al. (ADMX), Phys Rev Lett. 2021;127(26):261803.
- [5] G.'tHooft, Phys. Rev. Lett. **37**, 8(1976).
- [6] G.'tHooft, Phys. Rev. D **14**, 3432 (1976).
- [7] R. D. Peccei and H. Quinn, Phys. Rev. Lett. 38,1440(1977).
- [8] R. D. Peccei and H. Quinn, Phys.Rev. D.16,1791(1977).
- [9] H. Georgi, D. B. Kaplan and L. Randall, Phys. Lett. B 169 (1986) 73.
- [10] F. Pisano and V. Pleitez, Phys. Rev. D 46, 410 (1992), arXiv:hep-ph/9206242.
- [11] P. H. Frampton, Phys. Rev. Lett. **69**, 2889 (1992), DOI: 10.1103/PhysRevLett.69.2889.
- [12] R. Foot, O.F. Hernandez, F. Pisano, and V. Pleitez, Phys. Rev. D 47, 4158 (1993), arXiv:hep-ph/9207264.
- [13] D. Ng, Phys. Rev. D 49 (1994) 4805, e-Print: hep-ph/9212284 [hep-ph]
- [14] M. Singer, J. W. F. Valle and J. Schechter, Phys. Rev. D **22**, 738 (1980).
- [15] R. Foot, H. N. Long and Tuan A. Tran, Phys. Rev. D 50, 34 $(R)(1994)$ [arXiv:hep-ph/9402243].
- [16] J. C. Montero, F. Pisano and V. Pleitez, Phys. Rev. D 47, 2918 (1993).
- [17] H. N. Long, Phys. Rev. D 54, 4691 (1996), arXiv: hep-ph/9607439.
- [18] H. N. Long, Phys. Rev. D 53, 437 (1996), arXiv: hep-ph/9504274.
- [19] M. Ozer, Phys.Rev.D 54 (1996) 1143-1149, DOI: 10.1103/PhysRevD.54.1143.
- [20] P. B. Pal, Phys. Rev. D **52** (1995) 1659. [hep-ph/9411406].
- [21] de S. Pires, C. Antonio and O. P. Ravinez, Phys. Rev. D 58, 03500 (1998), eprint: hep-ph/9803409.
- [22] P. V. Dong, H. N. Long, Int. J. Mod. Phys. A 21, 6677 (2006), eprint: hep-ph/0507155.
- [23] A. G. Dias, V. Pleitez and M. D. Tonasse, Phys. Rev. D 67, 095008 (2003) arXiv:hep-ph/0211107
- [24] A. G. Dias, C. A. de S. Pires and P. S. Rodrigues da Silva, Phys. Rev. D 68 (2003) 115009, arXiv:0309058 [hep-ph],
- [25] A. G. Dias and V. Pleitez, Phys.Rev. D 69 (2004) 077702, arXiv:hep-ph/0308037.
- [26] J. G. Ferreira, C. A. de S. Pires, J. G. Rodrigues and P.S. Rodrigues da Silva, Phys. Lett. B 771 (2017) 199.
- [27] V. H. Binh, D. T. Binh, A. E. Cárcamo Hernández, D. T. Huong, D. V. Soa, and H. N. Long, Phys. Rev. D 107 , 095030 (2023), arXiv:2007.05004[hep-ph].
- [28] G. Ballesteros, J. Redondo, A. Ringwald, and C. Tamarit, JCAP 08 (2017) 001 , arXiv: 1610.01639 [hep-ph]
- [29] V. Oliveira and C. A. de S. Pires, Phys. Rev. D, 106, no. 1, p. 015031, 2022.
- [30] Pei-Hong Gu, JCAP 2016, arXiv:1603.05070.
- [31] K. Choi, S. H. Im, H. J. Kim and H. Seong, JHEP 08 (2021) 058, arXiv: 2106.05816.
- [32] H. N. Long and D. V. Soa, Nucl. Phys. B 601, (2001) 361, arXiv: hep-ph/0104150.
- [33] D. T. Binh, D. T. Huong, Tr. T. Huong, H. N. Long, and D. V. Soa, J. Phys. G: Nucl. Part. Phys. 29 (2003) 1213, arXiv: hep-ph/0211072.
- [34] A. M. Galda, M. Neubert, $ALP-LEFT$ Interference and the Muon $(g-2)$, arXiv:2308.01338
- [35] M. Bauer, M Neubert, S. Renner, M. Schnubelb and A. Thammg, JHEP 04 (2021) 063, arXiv:2012.12272.
- [36] I. G. Irastorza, J. Redondo, Progress in Particle and Nuclear Physics, 102 (2018) 89, arXiv:1801.08127.
- [37] J. Stern, R. Zaoui, Regge pole and scattering lengths, Nucl. Phys. B 17 (1970) 253 - 266.
- [38] L. Di Luzio and G. Piazza, JHEP 12 (2022) 041, arXiv:2206. 04061.
- [39] Y. Giraldo, R. Martinez, E. Rojas, Juan C. Salazar, Eur. Phys. J. C 82 (2022) 1131, arXiv:2007.05653 [hep-ph]
- [40] S. Weinberg, Phys. Rev. Lett. 40 (1978) 223-226.
- [41] M. Srednicki, Nucl. Phys. B 260 (1985) 689.
- [42] M. Gorghetto, G. Villadoro, JHEP 03 (2019) 033, arXiv:1812.01008 .
- [43] G. Grilli di Cortona, E. Hard, J. P. Vega and G. Villadoro, JHEP 01 (2016) 034, arXiv: 1511.02867.
- [44] Mingqiu Li, Sichun Sun, Qi-Shu Yan, Zhijie Zhao, Gravitational waves from axion wave production, arXiv:2309.08407.
- [45] D. Chang and H. N. Long, Phys. Rev. D 73, (2006) 053006, arXiv:hep-ph/0603098.
- [46] A. E. Cárcamo Hernández, Sergey Kovalenko, H. N. Long, and Ivan Schmidt, J. High Energy Phys. 07 (2018) 144, arXiv:1705.09169 [hep-ph].
- [47] H. N. Long and T. Inami, Phys. Rev. D 61, (2000) 075002, arXiv: hep-ph/9902475.
- [48] M. B. Tully and G. C. Joshi, Phys. Rev. D 64, (2001) 011301(R).
- [49] S. Scherer, Introduction to Chiral Perturbation Theory, Adv. Nucl. Phys. 27 (2003) 277, arXiv:hep-ph/0210398
- [50] L. Di Luzio, G. Martinelli, G. Piazza, Phys. Rev. Lett. 126, 241801 (2021), arXiv:2101.10330 [hep-ph].