

# Objective Features Extracted from Motor Activity Time Series for Food Addiction Analysis Using Machine Learning

Borisenkov M.,<sup>1\*</sup> Velichko A.,<sup>2</sup> Belyaev M.,<sup>2</sup> Korzun D.,<sup>3</sup> Tserne T.,<sup>4,5</sup> Bakutova L.,<sup>1</sup> Gubin D.<sup>6,7,8</sup>

<sup>1</sup>Department of Molecular Immunology and Biotechnology, Institute of Physiology of Komi Science Centre of the Ural Branch of the Russian Academy of Sciences, 167982 Syktyvkar, Russia

(BM: [borisenkov@physiol.komisc.ru](mailto:borisenkov@physiol.komisc.ru)), (BL: [foksmars14@yandex.ru](mailto:foksmars14@yandex.ru))

<sup>2</sup>Institute of Physics and Technology, Petrozavodsk State University, 185910 Petrozavodsk, Russia

(VA: [velichkogf@gmail.com](mailto:velichkogf@gmail.com)), (BM: [biomax89@yandex.ru](mailto:biomax89@yandex.ru))

<sup>3</sup>Institute of Mathematics and Information Technology, Petrozavodsk State University, 185910 Petrozavodsk, Russia (KD: [dkorzun@cs.karelia.ru](mailto:dkorzun@cs.karelia.ru))

<sup>4</sup>Pitirim Sorokin Syktyvkar State University, 167000 Syktyvkar, Russia (TT: [cerne\\_tatyana@mail.ru](mailto:cerne_tatyana@mail.ru))

<sup>5</sup>Municipal Institution of Additional Education "Center for Psychological, Pedagogical, Medical and Social Assistance", 167031 Syktyvkar, Russia

<sup>6</sup>Laboratory for Chronobiology and Chronomedicine, Research Institute of Biomedicine and Biomedical Technologies, Medical University, 625023 Tyumen, Russia (GD: [dgubin@mail.ru](mailto:dgubin@mail.ru))

<sup>7</sup>Department of Biology, Medical University, 625023 Tyumen, Russia

<sup>8</sup>Tyumen Cardiology Research Center, Tomsk National Research Medical Center, Russian Academy of Sciences, 634009 Tomsk, Russia

\* Correspondence: [borisenkov@physiol.komisc.ru](mailto:borisenkov@physiol.komisc.ru)

## Abstract

This study investigates machine learning algorithms to identify objective features for diagnosing food addiction (FA) and assessing confirmed symptoms (SC). Data were collected from 81 participants (mean age:  $21.5 \pm 9.6$  years, range: 18-61 years, women: 77.8%) whose FA and SC were measured using the Yale Food Addiction Scale (YFAS). Participants provided demographic and anthropometric data, completed the YFAS, the Zung Self-Rating Depression Scale, and the Dutch Eating Behavior Questionnaire, and wore an actimeter on the non-dominant wrist for a week to record motor activity. Analysis of the actimetric data identified significant statistical and entropy-based features that accurately predicted FA and SC using ML. The Matthews correlation coefficient (MCC) was the primary metric. Activity-related features were more effective for FA prediction (MCC=0.88) than rest-related features (MCC=0.68). For SC, activity segments yielded MCC=0.47, rest segments MCC=0.38, and their combination MCC=0.51. Significant correlations were also found between actimetric features related to FA, emotional, and restrained eating behaviors, supporting the model's validity. Our results support the concept of a human bionic suite composed of IoT devices and ML sensors, which implements health digital assistance with real-time monitoring and analysis of physiological indicators related to FA and SC.

**Keywords:** food addiction, actimetry, machine learning, eating behavior, depression.

## Highlights:

- The analysis of actimetric data revealed significant statistical and entropy-based features that accurately predicted FA and SC. This finding makes a significant contribution to advancing the concept of smart IoT devices for enhanced health monitoring.
- Actimetric features derived from activity segments carry more information about FA than features obtained from rest segments.
- A significant correlation exists between actimetric indicators and emotional and restrained eating behavior. This finding underscores the relevance of actimetric data in understanding different eating behaviors.

## 1. Introduction

Obesity is considered a "disease of civilization", with its growth rates recently acquiring an epidemic character (James et al., 2001). There is a steady increase in the number of individuals with overweight and obesity in countries with different levels of economic development and among various social and age groups (Chooi et al., 2019). Obesity is a multifactorial disease, with significant contributions from genetic, environmental, and social factors (Hampl et al., 2023). Disordered eating behaviors are also a notable risk factor for obesity (Lister et al., 2023). Obesity is associated with challenges in human resilience to stress and

mental health (Meigal et al., 2023). These challenges can be addressed through advances in Machine Learning (ML) and smart sensor technology within the Internet of Things (IoT) (Korzun, 2017).

Currently, the DSM-V (Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition) identifies three primary eating disorders: anorexia nervosa (AN), bulimia nervosa (BN), and binge eating disorder (BED) (American Psychiatric Association, 2013). The latest edition also proposes the inclusion of food addiction (FA) as a distinct disorder. The Yale Food Addiction Scale (YFAS; Gearhardt et al., 2009, 2013) and its modified version (YFAS 2.0; Schulte & Gearhardt, 2017) are used to assess FA.

The YFAS measures addiction-like eating of palatable foods based on the seven diagnostic criteria for substance dependence in the fourth revision of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) (Meule et al., 2017). Numerous studies conducted in both general and clinical populations have noted a connection between FA and BMI (Gearhardt et al., 2013) and between FA and depression (Borisenkov et al., 2020). Experimental studies have identified objective neuroimaging correlates of FA (Gearhardt et al., 2011).

However, some authors highlight difficulties in interpreting FA test results. Several studies have found no significant association between FA and BMI (Gearhardt et al., 2011). In some cases, FA symptoms are identified in individuals without signs of weight disorders (Borisenkov et al., 2018), as well as in underweight individuals (Schulte & Gearhardt, 2018). It has been suggested (Gearhardt et al., 2011) that YFAS can identify disordered eating behaviors before the onset of morphological signs of obesity. However, longitudinal studies are needed to confirm this hypothesis.


FA has been shown to have a close relationship with AN, BN, and BED (Meule & Gearhardt, 2019), and there is also considerable overlap between the criteria for FA and BED (Burrows et al., 2017), complicating their differential diagnosis. Consequently, some experts question the necessity of recognizing FA as a separate eating disorder (Ziauddeen et al., 2012). Other researchers (Meule et al., 2014) argue that further development of the FA concept requires improving the differential diagnosis system for eating disorders based on longitudinal study results.

The diagnosis of eating disorders such as FA (Figure 1a) is often limited to questionnaire-based methods, which significantly reduce diagnostic accuracy (Meule & Gearhardt, 2014). Introducing simple and objective assessment methods for these disorders represents a crucial step in developing the FA concept. Direct and objective evaluation of the brain centers responsible for eating behavior requires costly equipment like functional magnetic resonance imaging, making it unsuitable for long-term studies. Instead, chronobiology methods, such as continuous monitoring of physiological indicators (body temperature and motor activity), are more effective, following the concept of human bionic suite composed of smart IoT devices and sensors (Korzun, 2017; Korzun & Meigal, 2019). Although these methods assess parameters not directly related to eating behavior, they have already been applied in this field.

The actimetry method (Figure 1b) has been extensively used to study the 24-hour behavioral rhythms of patients with eating disorders, such as BED (Jones et al., 2005; Roveda et al., 2018). Literature suggests that dysfunction of the circadian system plays a key role in the etiology of BED (Romo-Nava et al., 2022). Given the strong connections between the circadian system and eating behavior (Mistlberger & Antle, 2011; Wehrens et al., 2017), it is expected that other forms of eating disorders are also associated with circadian system dysfunction.

In our previous study, we identified a direct link between the number of FA symptoms, the mean level (MESOR), and fragmentation (intradaily variability, I.V.) of circadian motor activity (Borisenkov et al., 2022). These indicators could potentially differentiate between FA and BED diagnoses (Borisenkov et al., 2022).

Currently, actimetry is mainly used in chronobiology and sleep medicine, including detecting sleep disorders. For example, early-stage Alzheimer's disease shows increased fragmentation and decreased stability in daily activity-rest rhythms (Musiek et al., 2018). Recent advancements include applying ML to automate actimetry data analysis in sleep medicine and chronobiology (Li et al., 2020; Meteier et al., 2024; Rani et al., 2022). ML methods have been used for assessing driver drowsiness (Meteier et al., 2024), detecting chronic insomnia (Rani et al., 2022), and analyzing rest-activity rhythms (Li et al., 2020). However, literature lacks ML applications for assessing eating behavior based on actimetry data.



**Figure 1.** A schematic illustration showing the diagnosis of eating disorders (a), such as food addiction, using actimetric data measurement (b).

The current subjective FA assessment methods lack sensitivity. There is a need to develop objective diagnostic criteria for FA using ML for automated data collection and analysis.

This study aims to test the hypothesis that analyzing motor activity time series with an entropy-based ML algorithm can provide new criteria for FA assessment. Our team has experience with effective features, including entropy-based ones, for EEG signal analysis (Belyaev et al., 2023), and has developed entropy features based on NNetEn (Velichko et al., 2023; Heidari et al., 2023). Entropy-based ML algorithms offer automated analysis of complex motor activity time series, revealing patterns and correlations not accessible through traditional methods (Conejero et al., 2024). This approach provides objective criteria for FA assessment and improves diagnostic accuracy by enhancing the understanding of eating behavior and neurophysiological processes.

Integrating wearable IoT devices and ML sensors further enhances this approach by enabling real-time monitoring, digital assistance, and personalized feedback, which aids in managing eating behaviors through tailored interventions. This combination of advanced ML algorithms and IoT technology offers a dynamic and comprehensive method for diagnosing and treating food addiction, ultimately contributing to improved mental health outcomes.

## 2. Materials and Methods

### 2.1 Study Design

This publication presents the results of the third phase of a study focused on investigating external and internal factors associated with FA. In the first phase, climatic, socio-demographic, anthropometric, and physiological factors associated with FA were examined. In the second phase, we analyzed the relationship between metrics characterizing the daily rhythm of motor activity and FA symptoms to identify the most significant circadian rhythm indicators suitable for the differential diagnosis of FA. The results of the first and second phases were published in previous works (Borisenkov et al., 2020, 2020a, 2022). This study used data from questionnaires conducted in the first phase and actimetry data collected from participants in the second phase.

### 2.2 Study Participants

The characteristics of the study participants are detailed in our previous article (Borisenkov et al., 2022). Briefly, data were collected from February 2019 to March 2020 in Syktyvkar and Tyumen (Lat./Long.: 61.7/50.9 and 57.2/65.5, respectively). A total of 81 participants were examined (average age:  $21.5 \pm 9.6$ , range: 18-61 years, women: 77.8%). Exclusion criteria included night/shift work and sleep disorders.

### 2.3 Instruments

Each participant provided personal data, including residence/study place, gender, age, height, and weight, and completed the YFAS questionnaire (Gearhardt et al., 2009) and the Dutch Eating Behavior Questionnaire (DEBQ) (van Strien et al., 1986). They wore a Daqtometer v2.4 actimeter (Daqtix, Germany) on their non-dominant wrist for a week to record motor activity levels. Height and weight were used to calculate body mass index (BMI), with percentiles adjusted for gender and age using growth charts (de Onis et al., 2007). BMI categories were defined as (1) underweight, (2) normal weight, (3) overweight, and (4) obese, with categories 3 and 4 combined into a group for overweight and obesity (ov/ob).

### 2.3.1 Yale Food Addiction Scale

The YFAS (Gearhardt et al., 2009) was used to identify food addictions (e.g., high-fat, high-sugar, and processed foods). The Russian-translated version (YFAS-Rus) was used, with its validity confirmed by strong internal consistency and a significant correlation with the emotional eating subscale of the DEBQ (Borisenkov et al., 2020), similar to earlier studies (Gearhardt et al., 2013; Manzoni et al., 2018). The scale has 25 items and includes 7 diagnostic criteria for substance dependence and eating behavior impairments (Gearhardt et al., 2009). Results are given as (1) the number of FA symptoms (SC), ranging from 0 to 7, and (2) a dichotomous measure where three or more symptoms and significant impairments indicate FA. Cronbach's alpha for this sample was 0.87.

### 2.3.2 Dutch Eating Behavior Questionnaire

The DEBQ (van Strien et al., 1986) was used to assess eating behavior and translated into Russian (DEBQ-Rus). As shown previously (Borisenkov et al., 2020), DEBQ-Rus has satisfactory internal consistency. The test has 33 questions divided into three subscales: restrained (DEBQrestr, 10 items), external (DEBQextern, 10 items), and emotional (DEBQemo, 13 items). A Likert scale from 1 (never) to 5 (very often) is used. Scores are averaged for each subscale. DEBQ categories (DEBQc) were assigned based on whether scores on the subscales exceeded the sample average thresholds (DEBQrestrc: 2.20, DEBQexternc: 2.93, DEBQemoc: 1.96). Cronbach's  $\alpha$  values were 0.88 for DEBQ, 0.92 for DEBQrestr, 0.52 for DEBQextern, and 0.94 for DEBQemo.

### 2.3.3 Zung Self-Rating Depression Scale

The ZSDS which consists of 20 items was used to assess the level of depression (Zung, 1965). Raw scores were transformed into ZSDS indices (ZSDSIs) as described earlier (Zung 1972; Passik et al. 2000). The ZSDSI varying from 25 to 100 scores was used as quantitative measure of the depression. In addition, ZSDSI categories (ZSDSIc; threshold value – 60 scores) were used as qualitative measure of depression. Cronbach's  $\alpha$  for this sample was 0.857.

### 2.3.4 Wrist Actimetry

Each participant wore a Daqtometer v2.4 actimeter (Daqtix, Germany) continuously on their non-dominant wrist for one week. Activity was measured at 1 Hz, with values summed per minute and expressed in arbitrary units (a.u.). The actimeter recorded dynamic (movement) and static (positional change) acceleration using a two-axis accelerometer. To quantify activity, values for each axis ( $x_i$  and  $y_i$ ) were read every second. The linear difference between consecutive readings was summed for each 1-minute interval (*bin*). This value was stored for each cell and computed as follows:

$$bin = \sum_{i=1}^{60} ((x_i - x_{i-1}) + (y_i - y_{i-1})) \quad (1)$$

## 2.4 Dataset

The dataset under investigation contains data from 78 participants, including 13 anthropometric, psychoemotional, and behavioral features for each individual, collected over a period of 6 to 7 days. The dataset contained actimetric data and statistical features (original dataset). The statistical features in original dataset are presented in Tables 1 and 2.

The first target variable is FA, where 10 subjects were diagnosed with FA=1 (class 1), and 68 patients had no FA detected, FA=0 (class 0). The second target variable is SC. Four categories were identified based on the number of confirmed symptoms:

1. Class 1 (0-1 symptoms), number of records: 33;
2. Class 2 (2 symptoms), number of records: 16;

3. Class 3 (3 symptoms), number of records: 15;
4. Class 4 (4-7 symptoms), number of records: 14.

Since the number of elements in each class varies, classification metrics designed for imbalanced datasets were used (see section 2.7).

**Table 1.** General characteristics of quantitative variables

#	Variables	Min	Max	M	SD	Missing data
1	Age	15	62	21.65	9.64	0
2	BMI%	5	97	47.75	25.20	3
3	ZSDSI	28	90	47.27	12.24	0
4	DEBQrestr	1	4.8	2.20	1.00	0
5	DEBQextern	1.3	4	2.93	0.58	0
6	DEBQemo	1	4.2	1.96	0.50	0

BMI%: BMI percentiles, ZSDSI: Total score on the Zung Self-Rating Depression Scale, DEBQrestr: Total score on the DEBQ restrained eating subscale, DEBQextern: Total score on the DEBQ external eating subscale, DEBQemo: Total score on the DEBQ emotional eating subscale.

**Table 2.** General characteristics of qualitative variables

#	Variables	Categories	Codes	N	%	Missing data
1	Sex	Female	1	59	73.8	0
		Male	2	21	26.2	
2	BMIC	Underweight	1	5	6.5	3
		Normal weight	2	61	79.2	
		Overweight	3	7	9.1	
		Obesity	4	4	5.2	
3	ov/ob	No	0	66	85.7	3
		Yes	1	11	14.3	
4	ZSDSIc	No	0	68	85.0	0
		Yes	1	12	15.0	
5	DEBQrestrc	No	0	51	63.7	0
		Yes	1	29	36.3	
6	DEBQexternc	No	0	33	41.2	0
		Yes	1	47	58.8	
7	DEBQemoc	No	0	46	57.5	0
		Yes	1	34	42.5	

BMIC: BMI categories, ov/ob: prevalence of overweight/obesity, ZSDSIc: prevalence of depression, DEBQrestrc: prevalence of restrained eating behavior, DEBQexternc: prevalence of external eating behavior, DEBQemoc: prevalence of emotional eating behavior.

## 2.5 Segmentation of Actimetric Data

To extract features from actimetric data, the original time series were segmented into activity and rest periods for separate analysis. The process included:


1. Cleaning the data of empty values and removing inactivity periods longer than 12 hours.
2. Calculating a moving average curve with a 1-hour window.
3. Segmenting the time series using a change point detection algorithm based on the moving average.
4. Merging adjacent activity and rest segments with a threshold method.
5. Combining rest periods shorter than 3 hours with adjacent activity periods.
6. Combining activity periods shorter than 4 hours with adjacent rest periods.

To reduce data volatility, the actimetric curve was averaged. Segmentation used the ruptures library (Truong et al., 2020) and kernel change point detection algorithms (Arlot et al., 2019; Celisse et al., 2018), resulting in segmentations that could exceed the number of activity and rest periods. Change points identified transitions between different types of activity or rest phases.

The threshold method, with a global threshold set at three-quarters of the median of the actimetric curve, was used to classify segments. Segments above the threshold were classified as activity, while those below were classified as rest. Adjacent segments were merged according to specified criteria.

## 2.6 Calculation of Actimetric Features

For each participant, actimetric features were divided into two groups: activity segments (Group 1) and rest segments (Group 2) (Figure 2). Statistical methods and entropy calculation methods were employed to compute these features.



**Figure 2.** An example of actimetric data segmented into activity (A) and rest (R) segments. From a physiological point of view, the A-segment coincides with the period of daytime activity, the R-segment – with the period of nighttime sleep.

The statistical methods used included the mean, maximum, minimum, range (the difference between the maximum and minimum), standard deviation, variance, coefficient of variation, and percentile values (1st, 5th, 25th, 50th, 75th, 95th, 99th percentiles). The entropy calculation methods included fuzzy entropy (FuzzyEn) (Chen et al., 2007), distribution entropy (DistEn) (Li et al., 2015), singular value decomposition entropy (SVDEn) (Roberts et al., 1999), permutation entropy (PermEn) (Bandt & Pompe, 2002), and phase entropy (PhaseEn) (Rohila & Sharma, 2019). Each entropy calculation method was applied using ten different sets of parameters.

Let's examine the steps for calculating feature values using the activity segments (Group 1) as an example:

1. All activity segments from the actimetric data were selected ( $A_1, A_2, A_3 \dots A_N$ ), as shown in Figure 2. The A segments are time series of varying lengths.
2. For each time series segment, a statistical or entropy feature was calculated, resulting in another time series composed of these metrics. For example,  $TS = \text{FuzzyEn}(A_1), \text{FuzzyEn}(A_2), \text{FuzzyEn}(A_3) \dots \text{FuzzyEn}(A_N)$ . Thus, a set of  $TS$  values was formed for each method and segment for each participant.
3. The obtained  $TS$  values for each method were aggregated using the mean ( $\text{Mean}(TS)$ ) and standard deviation ( $\text{Std}(TS)$ ) functions.

A similar procedure was conducted for the rest segments (Group 2). Consequently, each statistical and entropy method produced four features: mean and standard deviation for the activity segments, as well as mean and standard deviation for the rest segments.

It is also important to note that various parameter sets were used for the entropy calculation methods to accurately characterize the dynamics of changes in participants' activity. Using this algorithm, 256 actimetric features were calculated for each record in the dataset.

## 2.7 Machine Learning Methods

For classifying FA and SC based on the extracted features, a ML model was implemented using the scikit-learn library in Python. The model was structured as a data processing pipeline consisting of six main stages: feature selection, handling missing values, feature scaling, classification using the chosen algorithm, cross-validation, and classification accuracy assessment.

**Feature Selection.** The primary goal of this study was to identify the most significant actimetric features and their combinations for classifying FA and SC. The features were categorized into three groups: actimetric features from activity segments, actimetric features from rest segments, and subjective features. A sequential search was conducted across various combinations of these feature groups to determine their impact on the classification results.

**Handling Missing Values.** The original dataset contained missing values in the subjective features (see Tables 1-2). These gaps were filled using the mean values for each feature to ensure consistency across the dataset.

**Feature Scaling.** Scaling of feature values was necessary when the model included more than one feature. The min-max scaling method was applied, normalizing feature values to a range between 0 and 1 to ensure comparability.

**Classification Algorithm.** The K-Nearest Neighbors (KNN) algorithm (Mack, 1981) was selected for classification due to its simplicity and efficiency. KNN classifies data points based on their proximity to other samples in the training set.

**Cross-Validation.** To assess the model's accuracy, leave-one-out cross-validation (LOOCV) was employed (Vehtari & Ojanen, 2012). This method involves iterating the training and testing process  $K=78$  times, where each iteration leaves out one sample from the training data to serve as the test sample. LOOCV is particularly useful for small datasets, as it maximizes the use of available data (Bradshaw et al., 2023). The Matthews correlation coefficient (MCC) (Matthews, 1975) for each iteration was calculated, and the overall MCC served as the criterion for selecting the most significant features and their combinations. The influence of hyperparameters on model accuracy was also evaluated, with optimal values selected based on the MCC for each feature set.

**Classification Accuracy Assessment.** The MCC was used to evaluate the classification accuracy. Additionally, a confusion matrix was generated to derive parameters such as Accuracy, Sensitivity, Specificity, and F1-score.

In binary classification, there are four possible outcomes: correct predictions of positive samples (True Positive, TP) and negative samples (True Negative, TN), as well as incorrect predictions of negative samples (False Positive, FP) and positive samples (False Negative, FN). The MCC is calculated using the following formula:

$$MCC = \frac{TP \cdot TN - FP \cdot FN}{\sqrt{(TP+FP) \cdot (TP+FN) \cdot (TN+FP) \cdot (TN+FN)}} \quad (2)$$

This metric is particularly valuable for evaluating model accuracy with imbalanced datasets because it considers all four outcomes, including True Negatives, unlike the F1-score. MCC was calculated using standard libraries from scikit-learn for both binary and multiclass tasks, providing a comprehensive assessment of model performance.

## 3. Results

### 3.1 Segmentation Results of Actimetric Data

Figure 3 shows examples of the segmentation of actimetric data from the studied dataset, illustrating various levels of FA and the number of confirmed SC. The visualization is presented in the form of three columns, each corresponding to different levels of FA and SC:

(a) FA=0, SC=1: This chart shows data from a participant with no food addiction and minimal symptoms. It reveals regular activity and rest cycles with clear segmentation, indicating a balanced daily routine.

(b) FA=0, SC=6: This chart depicts a participant without food addiction but with many symptoms. Increased fragmentation and instability in activity rhythm suggest possible circadian disturbances despite no FA.

(c) FA=1, SC=3: This example features a participant with food addiction and moderate symptoms. The chart shows significant activity fluctuations and prolonged periods, reflecting impulsive or irregular behavior typical of FA.

Data segmentation involved several steps: cleaning gaps (e.g., for water procedures), removing erroneous inactivity periods over 12 hours, and using a change point detection algorithm to identify state changes. Adjacent activity and rest segments were then merged, smoothed, and stabilized.

A median-based threshold method combined segments effectively, distinguishing phases of activity and rest even in imbalanced datasets, as described in section 2.7.


Overall, the charts demonstrate the proposed method's effectiveness in distinguishing and visualizing different levels of FA and related SC, facilitating detailed analysis of actimetric data in eating behavior studies.

### 3.2 FA Classification Results

### 3.2.1 Classification Results of Individual Features

After segmenting the data and calculating the actimetric features, the entire dataset was analyzed using a classification algorithm to assess the impact of each individual feature. Tables 3, 4, and 5 present the five most important features from the original dataset of subjective features (see Tables 1 and 2) and actimetric features, divided into activity and rest segments.

Analyzing individual features shows that actimetric features (Tables 4 and 5) have higher MCC values than features from the original dataset (Table 3). The highest MCC value of 0.62 is for activity segments (standard deviation of the maximum value). Other features from activity segments also show higher MCC values, highlighting their greater importance for FA classification compared to rest segments.



**Figure 3.** Examples of actimetry data segmentation from the study dataset with varying levels of FA and SC a) FA=0, SC=1, b) FA=0, SC=6, c) FA=1, SC=3; FA: food addiction, SC: symptom counts of FA.

**Table 3.** The most significant subjective features for determining FA from the original dataset

#	Feature Name	MCC
7	ZSDSIc	0.42
8	DEBQrestr	0.34
9	DEBQrestrc	0.34
10	DEBQextern	0.34



The analysis used variables listed in Tables 1&2; MCC: Matthew's correlation coefficient, other abbreviations as in Tables 1&2.

**Table 4.** The most significant actimetric features for determining FA based on activity segments

Aggregation Function	Feature	MCC
Standard Deviation	Maximum Value	0.62
Standard Deviation	FuzzyEn	0.58
Mean Value	SVDEn	0.53
Mean Value	DistEn	0.42
Mean Value	99th Percentile	0.37
Mean Value	Standard Deviation	0.36

FuzzyEn: fuzzy entropy, SVDEn: singular value decomposition entropy, DistEn: distribution entropy, MCC: Matthew's correlation coefficient.

**Table 5.** The most significant actimetric features for determining FA based on rest segments

Aggregation Function	Feature	MCC
Mean Value	FuzzyEn	0.61
Standard Deviation	75th Percentile	0.43
Standard Deviation	FuzzyEn	0.42
Standard Deviation	DistEn	0.38
Standard Deviation	Standard Deviation	0.36

Abbreviations as in Table 4.

Notably, the highest accuracy for activity segments corresponds to the standard deviation as the aggregation function, indicating that variation in daily activity indicators is crucial for determining FA. For rest segments, the most important feature is the mean value of FuzzyEn.

### 3.2.2 Classification Results of Feature Combinations

In the next stage of the work, a search was conducted for feature combinations that provide the highest classification accuracy. Table 6 presents the results for all four groups.

**Table 6.** The most significant feature combinations for determining FA

Group	Features	MCC
Objective and Subjective Features	ZSDSIc, DEBQrestrc, DEBQexternc, DEBQemo, BMI%,	0.75
Actimetric Features, Activity Segments Only	Standard Deviation of FuzzyEn, Mean Value of the 99th Percentile, Standard Deviation of SVDEn, Mean Value of the 5th Percentile,	0.88
Actimetric Features, Rest Segments Only	Mean Value of FuzzyEn, Standard Deviation of the Maximum Value,	0.68
Actimetric Features, Both Rest and Activity Segments	Standard Deviation of FuzzyEn (Activity), Standard Deviation of FuzzyEn (Rest), Mean Value of the 99th Percentile (Activity)	0.88

Abbreviations as in Tables 1, 2&4.

Table 6 shows that the highest MCC value (0.88) is achieved using features from activity segments, offering better classification accuracy than the original dataset features (Tables 1 and 2). The lowest MCC value (0.68) is from features derived from rest segments, highlighting the stronger link between FA and activity segments. Using all actimetric features does not improve accuracy but reduces the number of features from four to three.

In cases with an MCC of 0.88, errors involved incorrectly identifying two subjects with FA=1. The confusion matrix is shown in Table 7, with a classification accuracy of 97.4%, sensitivity of 0.8, specificity of 1, and an F1-score of 0.89.

**Table 7.** Confusion matrix based on classification results

	Predicted Values	
	FA=0	FA=1

<b>Actual Values</b>	FA=0	68	0
	FA=1	2	8

FA: food addiction.

### 3.3 SC Classification Results

#### 3.3.1 Classification Results of Individual Features

Tables 8, 9, and 10 present the five most significant features (based on MCC values) from the original dataset (see Table 1), as well as actimetric features divided into activity and rest segments.

**Table 8.** The most significant features for determining SC from the original dataset

#	Feature Name	MCC
1	Age	0.25
6	ZSDSI	0.25
10	DEBQextern	0.21
4	BMIc	0.2
5	ov/ob	0.2

The analysis used variables listed in Tables 1 and 2.

**Table 9.** The most significant actimetric features for determining SC based on activity segments

Aggregation Function	Feature	MCC
Standard Deviation	95th Percentile	0.29
Mean Value	DistEn	0.26
Standard Deviation	FuzzyEn	0.24
Mean Value	PermEn	0.23
Mean Value	Mean Value	0.22
Mean Value	Standard Deviation	0.14

**Table 10.** The most significant actimetric features for determining SC based on rest segments

Aggregation Function	Feature	MCC
Standard Deviation	FuzzyEn	0.28
Mean Value	FuzzyEn	0.28
Standard Deviation	PhaseEn	0.20
Mean Value	Variance	0.20
Mean Value	PhaseEn	0.19

Abbreviations as in Table 4.

#### 3.3.2 Classification Results of Feature Combinations

In the next stage of the work, a search was conducted for feature combinations that provide the highest classification accuracy. Table 11 presents the results for all four groups.

Table 11 shows that the highest MCC value (0.51) is achieved with features from both activity and rest segments, offering better classification accuracy than the objective and subjective (OaS) dataset features (MCC=0.46). Comparable values are seen with actimetric features from activity segments alone (MCC=0.47). The lowest MCC value (0.38) is for features from rest segments, highlighting the greater significance of activity segments in determining SC. The confusion matrix for classification using both activity and rest segments is shown in Table 12.

**Table 11.** The most significant feature combinations for determining SC

Group	Features	MCC
Objective and Subjective Features	Age, BMIc, ZSDSI, DEBQrestrc, DEBQextern, DEBQemoc	0.46
Actimetric Features, Activity Segments Only	Mean Value of PermEn, Standard Deviation of FuzzyEn, Standard Deviation of PhaseEn,	0.47

Mean Value of the 99th Percentile,

Actimetric Features, Rest Segments Only	Mean Value of FuzzyEn, Mean Value of the Coefficient of Variation, Standard Deviation of FuzzyEn, Standard Deviation of the Range of Values,	0.38
Actimetric Features, Both Activity and Rest Segments	Mean Value of the Coefficient of Variation (Activity), Standard Deviation of the 95th Percentile (Activity), Standard Deviation of the Maximum Value (Rest), Standard Deviation of the 95th Percentile (Rest), Standard Deviation of the Range of Values (Rest)	0.51

Abbreviations as in Tables 1, 2 & 4

**Table 12.** Confusion matrix for SC classification using actimetric features from activity and rest segments

		Predicted Values				Sensitivity	Specificity	F1-score
		SC=1	SC=2	SC=3	SC=4			
Actual Values	SC=1	27	3	0	3	0.818	0.644	0.710
	SC=2	4	11	0	1	0.687	0.887	0.647
	SC=3	9	3	3	0	0.200	1.000	0.333
	SC=4	3	1	0	10	0.714	0.937	0.714

SC: symptom counts of food addiction.

**Table 13.** Confusion matrix for SC classification using objective and subjective features

		Predicted Values				Sensitivity	Specificity	F1-score
		SC=1	SC=2	SC=3	SC=4			
Actual Values	SC=1	31	0	1	1	0.939	0.466	0.704
	SC=2	9	6	0	1	0.375	0.983	0.521
	SC=3	11	1	1	2	0.066	0.984	0.117
	SC=4	4	0	0	10	0.714	0.937	0.714

The analysis used variables listed in Tables 1&2. SC: symptom counts of food addiction.

The model combining features from both activity and rest segments shows higher accuracy than the one using only OaS features (Table 13). The classification accuracy was 65% for actimetric features and 62% for OaS features. The OaS-based model better identifies SC=1, with a sensitivity of 0.939 and an F1-score of 0.704, but performs worse for other classes (SC={2, 3}). The actimetric model has lower sensitivity for SC=1 at 0.818 but a slightly higher F1-score of 0.710.

For SC=2, the actimetric model has a sensitivity of 0.687, specificity of 0.887, and an F1-score of 0.647, outperforming the OaS model with a sensitivity of 0.375, specificity of 0.983, and an F1-score of 0.521. However, for SC=3, the actimetric model struggles with a sensitivity of 0.200 and an F1-score of 0.333, compared to the OaS model's sensitivity of 0.066 and F1-score of 0.117. For SC=4, both models show comparable results with the same sensitivity of 0.714, specificity of 0.937, and F1-score of 0.714.

### 3.4 Relationship Between Actimetry and Subjective Features

To analyze the relationship between the OaS features (BMI%, ZSDSI, DEBQrestr, DEBQextern, DEBQemo) from Table 1 and the most significant actimetry features for determining FA or SC (Tables 4, 5, 9, 10), Pearson's correlation analysis was used (Table 14).

**Table 14.** Results of the correlation analysis of the relationships between the studied indicators

Aggregation Function	Feature	Subjective Features				
		BMI%	ZSDSI	DEBQrestr	DEBQextern	DEBQemo
Segments of activity						
<i>SD</i>	Max	-	-	-	-	0.263*
<i>SD</i>	FuzzyEn	-	-	-	0.265*	-
<i>M</i>	SVDEn	-0.296**	-	-	-	-
<i>M</i>	99th percentile	-	-	-	-	0.303**
<i>M</i>	<i>SD</i>	-	-	0.289*	-	0.382**

<i>M</i>	<i>M</i>	-	-	0.288*	-	-
<b>Segments of rest</b>						
<i>M</i>	Variance	-	-	0.244*	-	0.259*

The table presents Pearson correlation coefficients for the features BMI%, ZSDSI, DEBQrestr, DEBQextern, DEBQemo, and the most significant actimetry features taken from Tables 4, 5, 9, and 10, segmented by activity and rest. \* -  $P < 0.05$ , \*\* -  $P < 0.01$ ; the rest of the abbreviations are as in Tables 1, 2&4.

Table 14 demonstrates significant correlations between certain actimetry and subjective features, highlighting the importance of some metrics for assessing behavioral and physiological parameters. Among the actimetry features, significant correlations are observed in the activity segments for DEBQemo, BMI%, DEBQrestr, and DEBQextern. For DEBQemo and DEBQrestr, the correlation is positive, with the highest values observed for the mean (*M*) of the standard deviation (*SD*). A positive correlation is also observed for DEBQextern and the *SD* of FuzzyEn. For BMI%, a negative correlation is observed with the *M* of SVDeN. In the rest segments, significant correlations were found only between DEBQemo, DEBQrestr, and the *M* of variance. It can be noted that the strongest linear correlations are between the actimetry features of the activity segments and the initial features.

## 4. Discussion

### 4.1 Relationship Between Actimetric Features of FA and Psychophysiological Characteristics

The results of our study highlight the value of combining actimetric data with ML methods for diagnosing FA. Activity segments proved more informative for assessing FA and SC than rest segments, consistent with our previous research (Borisenkov et al., 2022), which identified daytime motor activity predictors like MESOR and I.V. as important for SC.

The most significant correlations were found between actimetric features and emotional eating behavior, confirming its well-documented link to FA (Meule et al., 2017; Lacroix & von Ranson, 2021). Notably, there were also significant correlations with restrained eating behavior, emphasizing its relevance to FA diagnosis (Rios et al., 2023).

We observed only one significant correlation between an actimetric feature for FA and BMI%, related to activity segments, which aligns with prior studies showing either a lack of or nonlinear relationship between FA and anthropometric characteristics (Schulte & Gearhardt, 2018).

Our study did not find significant correlations between FA and depression, contrary to earlier research suggesting a strong link (Borisenkov et al., 2020). This may indicate a need for further investigation into the complex relationship between FA and depression.

### 4.2 Analysis of FA and SC Classification Accuracy

The analysis of data from Tables 4, 5, 9, and 10 reveals that MCC values for features from activity segments are slightly higher than for rest segments, indicating a stronger connection between FA and SC and motor activity characteristics during the day. This is further evident in the optimal combinations of actimetric features (Tables 6 and 11). Combining features from activity and rest segments either does not improve the MCC metric or results in only slight gains. Thus, daytime activity alone is sufficient for classifying FA and SC.

Comparing OaS (Tables 1 and 2) and actimetric features for FA (Tables 3-6) and SC (Tables 8-11) classification shows that actimetric features enable more accurate classification. For SC, the maximum MCC values are close: MCC=0.29 (actimetry) vs. MCC=0.25 (OaS features) for individual features, and MCC=0.51 (actimetry) vs. MCC=0.46 (OaS features) for optimal combinations. For FA, the difference is more pronounced: MCC=0.62 (actimetry) vs. MCC=0.42 (OaS features) for individual features, and MCC=0.88 (actimetry) vs. MCC=0.75 (OaS features) for optimal combinations.

The maximum MCC values for SC are significantly lower than for FA, likely due to more classes (4 for SC vs. 2 for FA) and difficulty in classifying SC classes {2,3} (Tables 12 and 13). Accurate SC classification is challenging with both OaS and actimetric data, as classification errors are similar. For instance, errors where SC=3 samples are misclassified as SC=1 exceed the correctly classified samples.

### 4.3 Strengths and Limitations

The proposed method accurately identifies individuals with food addiction with high specificity and sensitivity. A key advantage of the study is its inclusion of participants from two geographically distant

regions. Analysis results show that while basic demographic and anthropometric features (such as age, depression level, and body mass index) have moderate significance, actimetric features related to activity segments offer more precise indicators for assessing FA symptoms. This highlights the need for further research and the integration of objective motor activity data into diagnostic algorithms for better assessment of eating disorders. However, the study has limitations, including a small sample size, gender imbalance, and a cross-sectional design that prevents conclusions about causal relationships.

## 5. Conclusion and Future Directions

This study demonstrated that combining actimetric data with ML enables the creation of a model that accurately reflects the relationship between food addiction and psychophysiological characteristics. Our findings suggest that the identified actimetric features can serve as objective criteria for FA, highlighting the potential of using actimetric data for developing objective diagnostic methods for eating disorders. Actimetric features also allow for more accurate classification of FA and SC compared to traditional features, with daily activity analysis being sufficient for classification.

Future research should combine subjective and objective methods to enhance actimetric data analysis, aiming to uncover hidden patterns related to eating disorders. Key future steps include evaluating the sensitivity, specificity, and accuracy of FA diagnosis in a double-blind experiment, testing the methodology with a larger sample size, assessing effectiveness in small ethnic groups, and conducting longitudinal studies to evaluate the method's potential for mass screening. Recent advancements in wearable technologies and actigraphy have incorporated light sensors (Danilenko et al., 2022), enabling the expansion of data analyses to circadian patterns of light exposure that associate with metabolic issues (Kim et al., 2023; Gubin et al., 2024) and via mechanisms of circadian disruption can be related to food addiction and eating disorders including binge eating and night eating (Plano et al., 2022).

In line with the concept of a human bionic suite, IoT technology can enhance monitoring and analysis capabilities by providing real-time data on human motor activity. Real-time data processing and ML sensors could improve the reliability and accuracy of diagnoses, particularly in cases where traditional methods fall short. These promising results indicate that wearable IoT devices are effective in analyzing physiological indicators related to FA and SC within digital health assistance, and in supporting human resilience to stress and mental health challenges. Future research will focus on IoT integration and the refinement of ML algorithms to develop advanced tools for diagnosing and managing eating disorders, aiming for a more precise and personalized diagnostic approach.

**Funding Statement:** The research (Introduction, Methods 2.5 – 2.7, Results, Discussion) is implemented with financial support by Russian Science Foundation, project no. 22-11-20040 (<https://rscf.ru/en/project/22-11-20040/> (accessed on 20 October 2023)), jointly with Republic of Karelia and funding from Venture Investment Fund of Republic of Karelia (VIF RK) (Head: KD) and (Methods 2.1-2.4) by the West-Siberian Science and Education Center, Government of Tyumen District, Decree of 20.11.2020, No. 928-rp (Head: GD).

**Conflict of Interest Disclosure:** The authors declare no conflict of interest.

**Ethics Approval Statement:** The study adhered to the tenets of the Declaration of Helsinki and research program was approved by the Ethics Committee of the Institute of Physiology of the Komi Scientific Center of the Ural Branch of the Russian Academy of Sciences. Each participant signed an informed consent form.

## References

- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5)*. Washington, DC: American Psychiatric Association; 2013.
- Arlot, S., Celisse, A., Harchaoui, Z.A. (2019). Kernel multiple change-point algorithm via model selection. *Journal of Machine Learning Research*, 20(162), 1-56. <http://jmlr.org/papers/v20/16-155.html>.
- Bandt, C., Pompe, B. (2002). Permutation entropy: A natural complexity measure for time series. *Physical Review Letters*, 88(17), 174102. <https://doi.org/10.1103/PhysRevLett.88.174102>.
- Belyaev, M., Murugappan, M., Velichko, A., Korzun, D. (2023). Entropy-based machine learning model for fast diagnosis and monitoring of Parkinson's disease. *Sensors*, 23, 8609. <https://doi.org/10.3390/s23208609>.
- Borisenkov, M.F., Tserne, T.A., Bakutova, L.A. (2018). Food addiction in Russian adolescents: Associations with age, sex, weight, and depression. *European Eating Disorders Review*, 26(6), 671-676. <https://doi.org/10.1002/erv.2644>.

- Borisenkov, M.F., Popov, S.V., Pecherkina, A.A., Dorogina, O.I., Martinson, E.A., Vetosheva, V.I., ... Symaniuk, E.E. (2020). Food addiction in young adult residents of Russia: Associations with emotional and anthropometric characteristics. *European Eating Disorders Review*, 28(4), 465-472. <https://doi.org/10.1002/erv.2731>.
- Borisenkov, M.F., Popov, S.V., Tserne, T.A., Bakutova, L.A., Pecherkina, A.A., Dorogina, O.I., ... Symanyuk, E.E. (2020a). Food addiction and symptoms of depression among inhabitants of the European North of Russia: Associations with sleep characteristics and photoperiod. *European Eating Disorders Review*, 28(3), 332-342. <https://doi.org/10.1002/erv.2728>.
- Borisenkov, M.F., Tserne, T.A., Bakutova, L.A., Gubin, D.G. (2022). Food addiction and emotional eating are associated with intradaily rest-activity rhythm variability. *Eating and Weight Disorders*, 27(8), 3309-3316. <https://doi.org/10.1007/s40519-022-01461-z>.
- Bradshaw, T.J., Huemann, Z., Hu, J., Rahmim, A.A (2023). Guide to cross-validation for artificial intelligence in medical imaging. *Radiology: Artificial Intelligence*, 5(4), e220232. <https://doi.org/10.1148/ryai.220232>.
- Burrows, T., Skinner, J., McKenna, R., Rollo, M. (2017). Food addiction, binge eating disorder, and obesity: Is there a relationship? *Behavioral Sciences (Basel)*, 7(3), 54. <https://doi.org/10.3390/bs7030054>.
- Celisse, A., Marot, G., Pierre-Jean, M., Rigail, G.J. (2018). New efficient algorithms for multiple change-point detection with reproducing kernels. *Computational Statistics & Data Analysis*, 128, 200-220. <https://doi.org/10.1016/j.csda.2018.07.002>.
- Chen, W., Wang, Z., Xie, H., Yu, W. (2007). Characterization of surface EMG signal based on fuzzy entropy. *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, 15(2), 266-272. <https://doi.org/10.1109/TNSRE.2007.897025>.
- Chooi, Y.C., Ding, C., Magkos, F. (2019). The epidemiology of obesity. *Metabolism*, 92, 6-10. <https://doi.org/10.1016/j.metabol.2018.09.005>.
- Conejero, J.A., Velichko, A., Garibo-i-Orts, Ò., Izotov, Y., Pham, V.-T. (2024). Exploring the entropy-based classification of time series using visibility graphs from chaotic maps. *Mathematics*, 12, 938. <https://doi.org/10.3390/math12070938>.
- Danilenko, K.V., Stefani, O., Voronin, K.A., Mezhakova, M.S., Petrov, I.M., Borisenkov, M.F., Markov, A.A., Gubin, D.G. (2022). Wearable light-and-motion dataloggers for sleep/wake research: A review. *Applied Sciences*, 12, 11794. <https://doi.org/10.3390/app122211794>
- de Onis, M., Onyango, A.W., Borghi, E., Siyam, A., Nishida, C., Siekmann, J. (2007). Development of a WHO growth reference for school-aged children and adolescents. *Bulletin of the World Health Organization*, 85(9), 660-667. <https://doi.org/10.2471/blt.07.043497>.
- Gearhardt, A.N., Corbin, W.R., Brownell, K.D. (2009). Preliminary validation of the Yale Food Addiction Scale. *Appetite*, 52(2), 430-436. <https://doi.org/10.1016/j.appet.2008.12.003>.
- Gearhardt, A.N., Yokum, S., Orr, P.T., Stice, E., Corbin, W.R., Brownell, K.D. (2011). Neural correlates of food addiction. *Archives of General Psychiatry*, 68(8), 808-816. <https://doi.org/10.1001/archgenpsychiatry.2011.32>.
- Gearhardt, A.N., Roberto, C.A., Seamans, M.J., Corbin, W.R., Brownell, K.D. (2013). Preliminary validation of the Yale Food Addiction Scale for children. *Eating Behavior*, 14(4), 508-512. <https://doi.org/10.1016/j.eatbeh.2013.07.002>.
- Gubin, D., Danilenko, K., Stefani, O., Kolomeichuk, S., Markov, A., Petrov, I., ... Cornelissen, G. (2024). Blue light and temperature actigraphy measures predicting metabolic health are linked to melatonin receptor polymorphism. *Biology (Basel)*, 3(1), 22. doi: 10.3390/biology13010022.
- Hampl, S.E., Hassink, S.G., Skinner, A.C., Armstrong, S.C., Barlow, S.E., Bolling, C.F., ... Okechukwu, K. (2023). Clinical practice guideline for the evaluation and treatment of children and adolescents with obesity. *Pediatrics*, 151(2), e2022060640. <https://doi.org/10.1542/peds.2022-060640>.
- Heidari, H., Velichko, A., Murugappan, M., Chowdhury, M.E. (2023). Novel techniques for improving NNetEn entropy calculation for short and noisy time series. *Nonlinear Dynamics*, 111(10), 9305-9326. <https://doi.org/10.1007/s11071-023-08298-w>.
- James, P.T., Leach, R., Kalamara, E., Shayeghi, M. (2001). The worldwide obesity epidemic. *Obesity Research*, 9, Suppl. 4, 228S-233S. <https://doi.org/10.1038/oby.2001.123>.
- Jones, S.H., Hare, D.J., Evershed, K. (2005). Actigraphic assessment of circadian activity and sleep patterns in bipolar disorder. *Bipolar Disorders*, 7(2), 176-186. <https://doi.org/10.1111/j.1399-5618.2005.00187.x>.
- Kim, M., Vu, T.H., Maas, M.B., Braun, R.I., Wolf, M.S., Roenneberg, T., ... Zee, P.C. (2023). Light at night in older age is associated with obesity, diabetes, and hypertension. *Sleep*, 46(3), zsa130. doi: 10.1093/sleep/zsa130.
- Korzun, D. (2017). Internet of things meets mobile health systems in smart spaces: An overview. In: Bhatt, C., Dey, N., Ashour, A. (eds) *Internet of Things and Big Data Technologies for Next Generation Healthcare. Studies in Big Data*, 23, Springer, Cham. [https://doi.org/10.1007/978-3-319-49736-5\\_6](https://doi.org/10.1007/978-3-319-49736-5_6)
- Korzun D., Meigal A. (2019). Multi-source data sensing in mobile personalized healthcare systems: Semantic linking and data mining. *Proc. 24th Conference of Open Innovations Association (FRUCT)*, 187-192, doi: 10.23919/FRUCT.2019.8711950.
- Lacroix, E., von Ranson, K.M. (2021). Prevalence of social, cognitive, and emotional impairment among individuals with food addiction. *Eating and Weight Disorders*, 26(4), 1253-1258. <https://doi.org/10.1007/s40519-020-01014-2>.

- Li, P., Liu, C., Li, K., Zheng, D., Liu., C, Hou, Y. (2015). Assessing the complexity of short-term heartbeat interval series by distribution entropy. *Medical & Biological Engineering & Computing*, 53(1), 77-87. <https://doi.org/10.1007/s11517-014-1216-0>.
- Li, X., Zhang, Y., Jiang, F., Zhao, H. (2020). A novel machine learning unsupervised algorithm for sleep/wake identification using actigraphy. *Chronobiology International*, 37(7), 1002-1015. <https://doi.org/10.1080/07420528.2020.1754848>.
- Lister, N.B., Baur, L.A., Felix, J.F., Hill, A.J., Marcus, C., Reinehr, T., ... Wabitsch, M. (2023). Child and adolescent obesity. *Nature Reviews Disease Primers*, 9(1), 24. <https://doi.org/10.1038/s41572-023-00435-4>.
- Mack, Y.P. (1981). Local properties of k-NN regression estimates. *SIAM Journal of Algebraic Discrete Methods*, 2(3), 311-323. <https://doi.org/10.1137/0602035>.
- Manzoni, G.M., Rossi, A., Pietrabissa, G., Varallo, G., Molinari, E., Poggiogalle, E., ... Castelnuovo, G. (2018). Validation of the Italian Yale Food Addiction Scale in postgraduate university students. *Eating and Weight Disorders*, 23(2), 167-176. <https://doi.org/10.1007/s40519-018-0495-0>.
- Matthews, B.W. (1975). Comparison of the predicted and observed secondary structure of T4 phage lysozyme. *Biochimical et Biophysical Acta (BBA) - Protein Structure*, 405(2), 442-451. [https://doi.org/10.1016/0005-2795\(75\)90109-9](https://doi.org/10.1016/0005-2795(75)90109-9).
- Meigal, A. L. Gerasimova-Meigal and D. Korzun (2023). A concept model of mHealth sensorics for digital assistance of human cognitive resilience. *Proc. 34th Conference of Open Innovations Association (FRUCT)*, 100-107, doi: 10.23919/FRUCT60429.2023.10328167.
- Meteier, Q., Favre, R., Viola, S., Capallera, M., Angelini, L., Mugellini, E., Sonderegger, A. (2024). Classification of driver fatigue in conditionally automated driving using physiological signals and machine learning. *Transportation Research Interdisciplinary Perspectives*, 26, 101148. <https://doi.org/10.1016/j.trip.2024.101148>.
- Meule, A., Gearhardt, A.N. (2014). Five years of the Yale Food Addiction Scale: Taking stock and moving forward. *Current Addiction Reports*, 1, 193-205. <https://doi.org/10.1007/s40429-014-0021-z>.
- Meule, A., Gearhardt, A.N. (2019). Ten years of the Yale Food Addiction Scale: A review of version 2.0. *Current Addiction Reports*, 6, 218-228. <https://doi.org/10.1007/s40429-019-00261-3>.
- Meule, A., von Rezori, V., Blechert, J. (2014). Food addiction and bulimia nervosa. *European Eating Disorders Review*, 22(5), 331-337. <https://doi.org/10.1002/erv.2306>.
- Meule, A., Müller, A., Gearhardt, A.N., Blechert, J. (2017). German version of the Yale Food Addiction Scale 2.0: Prevalence and correlates of 'food addiction' in students and obese individuals. *Appetite*, 115, 54-61. <https://doi.org/10.1016/j.appet.2016.10.003>.
- Mistlberger, R.E., Antle, M.C. (2011). Entrainment of circadian clocks in mammals by arousal and food. *Essays in Biochemistry*, 49(1), 119-136. <https://doi.org/10.1042/bse0490119>.
- Musiek, E.S., Bhimasani, M., Zangrilli, M.A., Morris, J.C., Holtzman, D.M., Ju, Y.-E.S. (2018). Circadian rest-activity pattern changes in aging and preclinical Alzheimer disease. *JAMA Neurology*, 75(5), 582. <https://doi.org/10.1001/jamaneurol.2017.4719>.
- Passik, S.D., Lundberg, J.C., Rosenfeld, B., Kirsh, K.L., Donaghy, K., Theobald, D., ... Dugan, W. (2000). Factor analysis of the Zung Self-Rating Depression Scale in a large ambulatory oncology sample. *Psychosomatics*, 41, 121-127. <https://doi.org/10.1176/appi.psy.41.2.121>.
- Plano, S.A., Soneira, S., Tortello, C., Golombek, D.A. (2022). Is the binge-eating disorder a circadian disorder? *Frontiers in Nutrition*, 9, 964491. doi: 10.3389/fnut.2022.964491
- Rani, S., Shelyag, S., Karmakar, C., Zhu, Y., Fossion, R., Ellis, J.G., ... Angelova, M. (2022). Differentiating acute from chronic insomnia with machine learning from actigraphy time series data. *Frontiers in Network Physiology*, 2, 1036832. <https://doi.org/10.3389/fnetp.2022.1036832>.
- Rios, J.M., Berg, M.K., Gearhardt, A.N. (2023). Evaluating bidirectional predictive pathways between dietary restraint and food addiction in adolescents. *Nutrients*, 15, 2977. <https://doi.org/10.3390/nu15132977>.
- Roberts, S.J., Penny, W., Rezek, I. (1999). Temporal and spatial complexity measures for electroencephalogram based brain-computer interfacing. *Medical & Biological Engineering & Computing*, 37(1), 93-98. <https://doi.org/10.1007/BF02513272>.
- Rohila, A., Sharma, A. (2019). Phase entropy: a new complexity measure for heart rate variability. *Physiological Measurements*, 40(10), 105006. <https://doi.org/10.1088/1361-6579/ab499e>.
- Romo-Nava, F., Guerdjikova, A.I., Mori, N.N., Scheer, F.A.J.L., Burgess, H.J., McNamara, R.K., ... McElroy, S.L. (2022). A matter of time: A systematic scoping review on a potential role of the circadian system in binge eating behavior. *Frontiers in Nutrition*, 9, 978412. <https://doi.org/10.3389/fnut.2022.978412>.
- Roveda, E., Montaruli, A., Galasso, L., Pesenti, C., Bruno, E., Pisanisi, P., ... Esposito, F. (2018). Rest-activity circadian rhythm and sleep quality in patients with binge eating disorder. *Chronobiology International*, 35(2), 198-207. <https://doi.org/10.1080/07420528.2017.1392549>.
- Schulte, E.M., Gearhardt, A.N. (2017). Development of the modified Yale Food Addiction Scale Version 2.0. *European Eating Disorders Review*, 25(4), 302-308. <https://doi.org/10.1002/erv.2515>.

Schulte, E.M., Gearhardt, A.N. (2018). Associations of food addiction in a sample recruited to be nationally representative of the United States. *European Eating Disorders Review*, 26(2), 112-119. <https://doi.org/10.1002/erv.2575>.

Truong, C., Oudre, L., Vayatis, N. (2020). Selective review of offline change point detection methods. *Signal Processing*, 167, 107299. <https://doi.org/10.1016/j.sigpro.2019.107299>.

van Strien, T., Frijters, J.E., Bergers, G.P., Defares, P.B. (1986). The Dutch Eating Behavior Questionnaire (DEBQ) for assessment of restrained, emotional, and external eating behavior. *International Journal of Eating Disorders*, 5(2), 295-315. [https://doi.org/10.1002/1098-108X\(198602\)5:2<295::AID-EAT2260050209>3.0.CO;2-T](https://doi.org/10.1002/1098-108X(198602)5:2<295::AID-EAT2260050209>3.0.CO;2-T).

Vehtari, A., Ojanen, J. (2012). A survey of Bayesian predictive methods for model assessment, selection and comparison. *Statistics Surveys*, 6, 142-228. <https://doi.org/10.1214/12-SS102>.

Velichko, A., Belyaev, M., Izotov, Y., Murugappan, M., Heidari, H. (2023). Neural network entropy (NNetEn): Entropy-based EEG signal and chaotic time series classification, python package for NNetEn calculation. *Algorithms*, 16, 255. <https://doi.org/10.3390/a16050255>.

Wehrens, S.M.T., Christou, S., Isherwood, C., Middleton, B., Gibbs, M.A., Archer, S.N., ... Johnston J.D. (2017). Meal timing regulates the human circadian system. *Current Biology*, 27(12), 1768-1775.e3. <https://doi.org/10.1016/j.cub.2017.04.059>.

Ziauddeen, H., Farooqi, I.S., Fletcher, P.C. (2012). Obesity and the brain: how convincing is the addiction model? *Nature Review Neuroscience*, 13(4), 279-286. <https://doi.org/10.1038/nrn3212>.

Zung, W.W. (1965). A self-rating depression scale. *Archives of General Psychiatry*, 12, 63-70. <https://doi.org/10.1001/archpsyc.1965.01720310065008>.

Zung, W.W. (1972). The depression status inventory: An adjunct to the self-rating depression scale. *Journal of Clinical Psychology*, 28, 539-543. [https://doi.org/10.1002/1097-4679\(197210\)28:4<539::aid-jclp2270280427>3.0.co;2-s](https://doi.org/10.1002/1097-4679(197210)28:4<539::aid-jclp2270280427>3.0.co;2-s).