# Dynamical homogenization in effective loop quantum cosmology

Edward Wilson-Ewing\*

Department of Mathematics and Statistics, University of New Brunswick, Fredericton, NB, Canada E3B 5A3

The effective dynamics of loop quantum gravity for marginally bound Lemaître-Tolman-Bondi spacetimes predict that the big-bang singularity is resolved and replaced by a cosmic bounce. Numerics show that these effective dynamics also homogenize small regions soon after the bounce when inhomogeneities before the bounce are sufficiently large. These homogeneous regions typically have a width of a few Planck lengths where relative perturbations in the energy density remain less than 15%. If the bounce is followed by an inflationary period, the homogeneous region can reach cosmic scales and the amplitude of relative perturbations can be suppressed to a value compatible with observations.

### I. INTRODUCTION

The small amplitude of the temperature fluctuations  $\delta T/T \sim 10^{-5}$  observed in the cosmic microwave background (CMB) [1], combined with the assumption of the Copernican principle that our location in the universe is a typical one, lead to the conclusion that the early universe was extremely homogeneous. This homogeneity provides initial conditions for the later evolution of the universe, that matches observations but begs the question as to why the early universe is homogeneous to such a high degree in the first place. Further, given the expectation that gravity will dominate the dynamics at the largest scales and, being attractive, can be expected to cause inhomogeneities to grow, it would then seem that an even higher degree of homogeneity is required at times well before the formation of the CMB.

This is a serious fine-tuning problem if there is no dynamical mechanism for homogenizing, or smoothing, the early pre-CMB universe. Interestingly, there do exist a few proposals for such a process, notably through inflation [2–9] (although see [10]) or ekpyrosis [11, 12]; this letter points out the existence of a another possibility where homogenization occurs during a cosmic bounce, studied here in the context of loop quantum cosmology but possibly holding more generally.

Inflation can alleviate the homogeneity problem in two ways. First, if there is a small region that is moderately homogeneous, then inflationary expansion can blow this small region up to cosmological scales and further homogenize it [3]; and second, under some assumptions, it has been shown that large homogeneous regions will necessarily have formed at sufficiently late times in an inflationary spacetime [2, 4, 7–9]. Note however that in the first case, it is necessary to already have small regions that are moderately homogeneous before inflation starts, while for the second family of results extremely long periods of inflation may be required for sufficient smoothing to occur. As shall be shown below, the cosmic bounce arising in loop quantum cosmology can generate the small and moderately homogeneous regions needed for the first scenario.

A phase of ekpyrotic contraction is an alternative mechanism that can efficiently smooth out inhomogeneities. This smoothing is successful even if the initial departures from homogeneity are of order one, and it can homogenize large regions of space [11, 12]. The drawback of ekpyrosis is that multiple fields are needed to generate (nearly) scale invariant scalar perturbations [13–15] and the perturbations are sensitive to anisotropies [16], which introduces a separate fine-tuning problem (this is in contrast to the background ekpyrotic dynamics which are robust to the presence of anisotropies).

As mentioned above, another possibility is that quantum gravity could help to alleviate the inhomogeneity problem. This was first suggested based on the observation that in loop quantum cosmology, the big-bang singularity is resolved and replaced by a non-singular bounce; heuristically, this can be understood as occurring due to quantum gravity becoming repulsive when the spacetime curvature is Planckian. Could this effective repulsion during a cosmic bounce perhaps also smooth out inhomogeneities in the early universe [17]?

Although this is an intriguing idea, perturbative calculations give no sign of such an effect. Indeed, in classical general relativity, linearized cosmological perturbations will grow in a contracting spacetime if the equation of state  $w = p/\rho$  relating the energy density  $\rho$  and pressure p of the matter field satisfies  $-1 < w \leq 1$ . Assuming the strong energy condition w > -1, the only remaining possibility consists of ekpyrotic fields satisfying w > 1 for which the amplitude of the cosmological perturbations remains constant. In this perturbative analysis, therefore, a contracting pre-bounce phase will not homogenize the spacetime, rather inhomogeneities will either grow or at best remain at a constant amplitude.

What about the bounce itself? A simple modelindependent way to check this possibility is by approximating the bounce as a discontinuous change in the Hubble rate H from  $H = -H_{\star}$  to  $H = H_{\star}$  and require that the cosmological perturbations satisfy appropriate matching conditions [18]; the result of this calculation is

<sup>\*</sup> edward.wilson-ewing@unb.ca

that the amplitude of the perturbations remains of the same order of magnitude as before the bounce [19–21]. Alternatively, it is possible to consider specific theories that predict a cosmic bounce (including loop quantum cosmology) and explicitly calculate the evolution of the perturbations through the resulting bounce: the result is the same and there is no suppression of the amplitude of cosmological perturbations across the bounce; for reviews of these calculations in general bouncing cosmological scenarios as well as specifically in loop quantum cosmology, see, respectively, [22, 23].

Nonetheless, it is important to keep in mind that these previous analyses are entirely based on the truncation of inhomogeneities to the approximation of linear perturbation theory. As shall be shown in the following, a non-perturbative analysis of this problem in Lemaître-Tolman-Bondi (LTB) spacetimes gives the opposite result: in effective loop quantum gravity, an inhomogeneous cosmic bounce can indeed homogenize small regions. Although these regions are small, they can later be expanded to cosmological scales, and also further homogenized, by inflation, as argued in [3].

#### II. NON-PERTURBATIVE ANALYSIS IN LTB SPACETIMES

The Lemaître-Tolman-Bondi (LTB) spacetime, corresponding to spherically symmetric dust, is an exact solution to the Einstein equations that allows arbitrarily large inhomogeneities in the radial direction. It will be convenient to write the metric in generalized Painlevé-Gullstrand coordinates,

$$ds^{2} = -dt^{2} + \frac{1}{1 + \mathcal{E}} (dr + N^{r} dt)^{2} + r^{2} d\Omega^{2}, \quad (1)$$

where the shift vector  $N^r$  depends on the radius r and time t coordinates, and  $d\Omega^2 = d\theta^2 + \sin^2 \theta \, d\phi^2$  is the metric on the unit 2-sphere. For simplicity, in the following I will focus on the  $\mathcal{E} = 0$  marginally bound configurations.

As a first step in exploring loop quantum gravity (LQG) effects, it is possible to derive effective equations that include corrections due to the discrete nature of quantum geometry as predicted in loop quantum gravity, but that neglect quantum fluctuations; specifically, the discrete nature of quantum geometry is captured by evaluating holonomies along paths of a minimal Planckian length. Such effective equations were first derived in the context of homogeneous cosmology [24, 25], and for sharply-peaked quantum states they provide an excellent approximation to the dynamics of expectation values, so long as the volume of the spatial region considered is much larger than the Planck volume [26, 27].

Following the same process, effective equations can also be derived from the loop quantization of the LTB spacetime [28–31]. These equations are expected to give a good approximation to the quantum dynamics at large length scales, but quantum fluctuations could become important at shorter scales. As a starting point, here I will focus on the effective dynamics, and leave the inclusion of quantum fluctuations for future work.

For marginally bound LTB spacetimes expressed in generalized Painlevé-Gullstrand coordinates, there is a single (local) degree of freedom corresponding to the angular component b of the Ashtekar-Barbero connection. In the effective LQG theory [28–31],

$$\partial_t b = -\frac{1}{2\Delta r} \partial_r \left( r^3 \sin^2 \frac{\sqrt{\Delta b}}{r} \right), \qquad (2)$$

where  $\Delta \sim \ell_{\rm Pl}^2$  is the area gap, namely the smallest nonzero eigenvalue of the area operator in loop quantum gravity, and the Barbero-Immirzi parameter is set to 1. Note that the velocity  $v_b$  of the field (i.e., the prefactor to  $\partial_r b$  after expanding the equation of motion) is

$$v_b = -\frac{r}{\sqrt{\Delta}} \sin \frac{\sqrt{\Delta}b}{r} \cos \frac{\sqrt{\Delta}b}{r}, \qquad (3)$$

and depends on both r and b itself.

Then, b determines the shift vector  $N^r$  through [30–33]

$$N^{r} = -\frac{r}{\sqrt{\Delta}} \sin \frac{\sqrt{\Delta}b}{r} \cos \frac{\sqrt{\Delta}b}{r}, \qquad (4)$$

which is identically equal to  $v_b$ , and the dust energy density  $\rho$  by [28–30]

$$\rho = \frac{\rho_c}{3r^2} \,\partial_r \left( r^3 \sin^2 \frac{\sqrt{\Delta}b}{r} \right),\tag{5}$$

where  $\rho_c = 3/8\pi G\Delta$  is the critical energy density of loop quantum cosmology. This can be used to rewrite (2) as  $\partial_t b = -4\pi G r \rho$ , showing that b is monotonically decreasing since  $\rho \ge 0$ .

In homogeneous cosmology, the effective dynamics resolve the big-bang singularity, and replace it with a nonsingular bounce that occurs at  $v_b(t) = 0$  [24]. As shall be shown below, the dynamics are to some extent qualitatively similar for marginally bound LTB spacetimes with the difference that the bounce is inhomogeneous since  $v_b$ is now a function also of r, and the bounce at each radius occurs at different times, when  $v_b(r, t) = 0$ .

The fact that the bounce can occur at different times at different radii has important consequences. In particular, if there is a region  $r \in [r_1, r_2]$  with a large over-density, it may bounce before the region  $r < r_1$  lying immediately inside. In such a situation, the outer region that bounces first starts to expand with  $v_b^{(out)} > 0$  while the inner region continues to contract with  $v_b^{(in)} < 0$ . This causes a rarefaction wave (located around  $v_b = 0$ ) to form between the two regions, with the rarefaction wave moving inwards, following the inner region that continues to contract. In the meantime, the outer region is expanding, and the new intermediate region located between the

ingoing rarefaction wave and the outermost part of the now-expanding outer region will be quite homogeneous, because all of these points will have had  $v_b = 0$  nearly simultaneously, which from (3) and (5) implies that  $\rho$  will be nearly constant in this region.

Note that these regions are nearly, but not exactly, homogeneous. The reason for this is that the rarefaction wave (where  $v_b = 0$ , and therefore the location where the bounce occurs) moves inwards, and as a result at larger r the bounce happens at slightly earlier times, giving  $\rho$  more time to dilute during the post-bounce expanding phase:  $\rho$  will then be monotonically decreasing as a function of r in these nearly homogeneous regions. This effect is confirmed by the numerics (discussed below) that clearly show this (small) slope.

The size of the homogeneous region can be maximized by increasing the longevity of the rarefaction wave (that vanishes when the less dense inner region bounces). This can be done by increasing the difference in  $\rho$  in the inner and outer regions: ironically, larger inhomogeneities in  $\rho$ before the bounce will lead to larger homogeneous regions after the bounce.

To see the dynamical formation of such homogeneous regions, it is necessary to solve the dynamics numerically. It is convenient to introduce B = rb, as the dynamics for B are given by a conservation law  $\partial_t B + \partial_r f(B; r) = 0$ with  $f = (r^3/2\Delta) \cdot \sin^2(\sqrt{\Delta}B/r^2)$ . Then, the Godunov algorithm for non-linear wave equations can be directly applied to numerically solve the dynamics [34]; this has been done for the LQG effective dynamics of LTB spacetimes for configurations of  $\rho$  that have compact support [29, 30, 35]. For the cosmological configurations of interest here, the difference is that one of the boundary conditions changes: while the inner boundary condition B(r=0) = 0 remains the same (this is due to B = rbvanishing at r = 0, now the outer boundary condition at  $r = r_{max}$  is fixed by assuming a cosmological exterior with homogeneous energy density which implies  $B(r = r_{max}) \propto r^2$ . Choosing a discretization in the radial direction of constant width  $\delta r$ , this boundary condition is imposed at each time step through the condition  $B(r_{max}) = r_{max}^2 B(r_{max} - \delta r) / (r_{max} - \delta r)^2$  used to determine  $B(r_{max})$ .

Initial conditions are given for the interval  $r \in [0, r_{max}]$ that correspond to a contracting spacetime with inhomogeneities in  $\rho$  of order 1, and the Godunov algorithm is then used to solve the dynamics numerically. The initial conditions are given in terms of  $\rho$  as

$$\rho(t_i) = \rho_{avg} + \sum_{n=1}^{m} \alpha_n \cos\left(\frac{n\pi r}{r_{max}}\right), \qquad (6)$$

where  $\rho_{avg} > 0$ , the  $\alpha_n$  are randomly selected from a uniform distribution in the same interval [-A, A], with A chosen so that perturbations in  $\rho(t_i)$  are large while keeping  $\rho > 0$ . The sum over n ends at  $m = r_{max}/\sqrt{\Delta}$ , corresponding to a Planck-scale cutoff on the Fourier expansion for  $\rho(t_i)$ ; note however that despite this cutoff



(d) Post-bounce expansion, t = 8.40.

FIG. 1. These frames from a representative example show  $\rho(r)$  at four instants, using units where  $\hbar = \Delta = 1$ . The top frame shows the initial conditions, the second shows a later time during contraction, while the third frame shows part of the bounce, with the region located approximately between  $r \approx 7$  and  $r \approx 11$  having already bounced and the region located approximately between  $r \approx 11$  and  $r \approx 13$  in the process of bouncing, and the fourth frame shows the time by which R expanded by a factor of 1.5 after the bounce. The critical energy density  $\rho_c$  is shown as a dotted line in the bottom two frames. The formation of small homogeneous regions can be seen in the last frame: there are four regions with a radial extent of at least  $\sqrt{\Delta}$  where  $|\partial_r \rho|/\rho < 0.15$ , located around r = 9, r = 12, r = 15, and r = 29. Note that the range of r differs for each frame, in order to follow the geometry as it initially contracts and later expands.

there are no homogeneous regions in the initial profile for  $\rho(t_i)$ , see Fig. 1a for an explicit example. Note that  $\sin(n\pi r/r_{max})$  terms are excluded from the series to ensure that  $\rho$  is smooth at the origin. (Also, the Fourier expansion is used only for setting initial conditions—as the dynamics are non-linear, this expansion does not help in solving the dynamics.)  $\rho(t_i)$  is then used to determine  $B(t_i)$  by inverting (5), and then the Godunov algorithm is used to evolve B. The lattice spacing  $\delta r$  in the radial direction is fixed, while the time steps  $\delta t$  are determined dynamically using the Courant-Friedrichs-Lewy condition [34]. Once b = B/r is known, the energy density  $\rho$  at later times can then be calculated from (5). The code is available online [36].

The numerical solutions depend of course on the initial conditions, but importantly there are certain key features that appear quite generically: (i) there is a bounce in the cosmological dynamics also in the presence of large inhomogeneities, in agreement with what has been found for homogeneous cosmology [24]; (ii) shock waves form during contraction, as is guaranteed to happen if inhomogeneities are sufficiently large [37, 38]; and (iii) small homogeneous regions emerge during the bounce, generated by sufficiently over-dense regions during the bounce. Due to the formation of shocks, it is necessary to allow for weak solutions, as is done by the Godunov algorithm used here; for more on weak solutions in effective LQG see [29, 30, 38]. As explained above, the homogeneous regions form due to an outer over-dense region bouncing before a neighbouring inner under-dense region, causing the formation of a rarefaction wave moving inwards so that a continuous interval of r bounces nearly simultaneously.

To quantitatively characterize the homogeneous regions formed during the bounce, a reasonable definition that I will use here is that they are intervals in r of width greater than  $\sqrt{\Delta}$  where for each r in that interval,  $|\partial_r \rho|/\rho < 0.15$  (results are similar for other possible definitions of homogeneous regions).

A typical example, with  $r_{max} = 100, \delta r = 10^{-4}, \rho_{avg} =$  $10^{-3}, A = 0.08$  (using units where  $c = \hbar = \Delta = 1$ ), is shown in Fig. 1. Note that in terms of the coordinate r, during contraction the dust will initially move inwards, and then outwards after the bounce; the results are depicted using the range for r from the origin to R(t), the outer edge of the inhomogeneous region (that is initially  $R(t_i) = r_{max}$ ). For the example shown in Fig. 1, R decreased by a factor of  $\sim 4.9$  during contraction, and increased by a factor of  $\sim 1.5$  after the bounce, at which time the numerics were stopped. For this run, at the final time there are four homogeneous regions where  $|\partial_r \rho / \rho| < 0.15$  with widths of 1.93, 3.04, 1.54, and 2.12, and located around r = 9, 12, 15 and 29 respectively; note that these cover over a quarter of the range of the numeric results. It is worth pointing out that the gradient in  $\rho$  is for the most part due to a linear and quadratic dependence on r in the homogeneous regions, with the linear best fit accounting for approximately one third of  $(\partial_r \rho)/\rho$ , and the quadratic best fit accounting for another third in each of the four homogeneous regions.

As can be seen in Fig. 1d, there are typically no bumps or other types of features in the homogenized regions, except for a small decrease in  $\rho$ , due to the outer region bouncing first, and therefore expanding and diluting more, as explained above.

On its own, a smoothing to  $|\partial_r \rho / \rho| < 0.15$  over a length scale of a few  $\sqrt{\Delta}$  is not sufficient to explain perturbations of amplitude ~ 10<sup>-5</sup> at cosmological scales, but if this is combined with a later sufficiently long-lasting era of inflation then the size of the smoothed region will be exponentially enlarged, and also further homogenized, by an amount depending on the number of e-folds of inflation. In particular, the amplitude of the relative perturbations over the observable universe can be made arbitrarily small by choosing a sufficiently large number of e-folds.

Note however that if there is a limited amount of inflation, the remaining small near-linear change in  $\rho$  could potentially have an observational impact, for example by introducing a preferred direction in the CMB corresponding to the direction where  $\rho$  increases. More work is needed to determine whether this could cause the hemispherical power asymmetry observed in the CMB [39] (in particular, such a study will require going beyond spherical symmetry as well as including more realistic matter fields); a detailed investigation of this possibility is left for future work.

#### **III. DISCUSSION**

In marginally bound LTB spacetimes, the cosmic bounce of loop quantum cosmology can provide a mechanism to smooth out inhomogeneities in the very early universe, forming small moderately homogeneous regions. These regions can then be blown up to cosmological size and further homogenized by inflation, as argued in [3], thereby potentially solving the homogeneity problem of the early universe. Although this analysis focused on the loop quantum cosmology bounce, it is plausible that a similar homogenization mechanism could occur in other bouncing cosmological scenarios as well. Note that in this scenario, the Copernican principle fails at the largest (super-horizon) scales, where there exist significant inhomogeneities.

The physics underlying the homogenization process is not as simple as quantum gravity becoming repulsive when the spacetime curvature is sufficiently large (indeed, some regions are not homogenized during the bounce, contrary to what would be expected if quantum gravity were to become 'universally repulsive' at Planck scales). Rather, if inhomogeneities are sufficiently large, then a rarefaction wave will form between an outer sufficiently over-dense region adjacent to an inner sufficiently under-dense region. It is the new intermediate region between the outer region (that bounces first) and the rarefaction wave (that moves inwards following the inner region that contracts for a longer time), that is approximately homogeneous after the bounce.

The specific model considered here, namely the marginally bound Lemaître-Tolman-Bondi spacetime with LQG effective dynamics, provides a comparatively simple framework to study this problem. This has the important advantage of making calculations tractable, but it is important in future work to consider richer models to check the robustness of these results.

A first step in this direction is to include non-vanishing pressure—pressure may help strengthen the homogenization process, but this needs to be verified; alternatively, it may decrease inhomogeneities during contraction and suppress the formation of homogeneous regions during the bounce. A more challenging next step is to move beyond spherical symmetry and allow general inhomogeneities; this however has two major difficulties: first, the general dynamics for effective LQG are not presently known beyond spherical symmetry, so considerable work will be required simply to derive the equations of motion; and second, more powerful analytic and numeric methods will be required, although recent progress on this problem in classical general relativity (see, e.g., [8, 9, 12]) is encouraging.

It is already known that loop quantum cosmology can provide suitable initial conditions for the onset of inflation in a homogeneous and isotropic cosmology [40–43]. Building on this work, it would be interesting to couple an inflaton field to this model to compute the size the homogeneous regions expand to by the end of inflation, for a given inflationary potential, and to what degree the remaining inhomogeneities in these regions are further smoothed out.

Finally, the homogenization that is found to occur in these models comes from an effective description that includes leading order quantum gravity effects for sharplypeaked states, but neglects the effect of quantum fluctuations. Since quantum gravity effects can homogenize the expectation values of such states in this approximation, it is natural to ask what may happen when quantum fluctuations are included. Do quantum fluctuations affect (either enhancing or destroying) the smoothing mechanism observed here? And this also raises the complementary question of whether quantum gravity could also 'smooth out' and dynamically minimize quantum fluctuations (at least in some regions of space). In other words, can quantum gravity dynamically drive the wave function of the universe to a state that is well approximated (at least in some region of space) by the tensor product of a homogeneous background and something close to an appropriate vacuum state for linear perturbations? These questions are left for future work.

# ACKNOWLEDGMENTS

I thank Viqar Husain for helpful discussions. This work was supported in part by the Natural Sciences and Engineering Research Council of Canada.

- Planck, N. Aghanim *et al.*, "Planck 2018 results. I. Overview and the cosmological legacy of Planck," Astron. Astrophys. 641 (2020) A1, arXiv:1807.06205.
- [2] A. A. Starobinsky, "Isotropization of arbitrary cosmological expansion given an effective cosmological constant," JETP Lett. **37** (1983) 66–69.
- [3] A. D. Linde, "Initial Conditions for Inflation," Phys. Lett. B 162 (1985) 281–286.
- [4] L. G. Jensen and J. A. Stein-Schabes, "Is Inflation Natural?," Phys. Rev. D 35 (1987) 1146.
- [5] W. E. East, M. Kleban, A. Linde, and L. Senatore, "Beginning inflation in an inhomogeneous universe," JCAP 09 (2016) 010, arXiv:1511.05143.
- [6] K. Clough, E. A. Lim, B. S. DiNunno, W. Fischler, R. Flauger, and S. Paban, "Robustness of Inflation to Inhomogeneous Initial Conditions," JCAP 09 (2017) 025, arXiv:1608.04408.
- [7] M. Kleban and L. Senatore, "Inhomogeneous Anisotropic Cosmology," JCAP 10 (2016) 022, arXiv:1602.03520.
- [8] P. Creminelli, O. Hershkovits, L. Senatore, and A. Vasy, "A de Sitter no-hair theorem for 3+1d cosmologies with isometry group forming 2-dimensional orbits," Adv. Math. 434 (2023) 109296, arXiv:2004.10754.
- J. Wang and L. Senatore, "On the asymptotics of 3+1D cosmologies with bounded scalar potential and isometry group forming 2-dimensional orbits," arXiv:2111.09257.
- [10] D. Garfinkle, A. Ijjas, and P. J. Steinhardt, "Initial conditions problem in cosmological inflation revisited," Phys. Lett. B 843 (2023) 138028, arXiv:2304.12150.

- [11] D. Garfinkle, W. C. Lim, F. Pretorius, and P. J. Steinhardt, "Evolution to a smooth universe in an ekpyrotic contracting phase with w > 1," Phys. Rev. D 78 (2008) 083537, arXiv:0808.0542.
- [12] W. G. Cook, I. A. Glushchenko, A. Ijjas, F. Pretorius, and P. J. Steinhardt, "Supersmoothing through Slow Contraction," Phys. Lett. B 808 (2020) 135690, arXiv:2006.01172.
- [13] F. Finelli, "Assisted contraction," Phys. Lett. B 545 (2002) 1–7, arXiv:hep-th/0206112.
- [14] F. Di Marco, F. Finelli, and R. Brandenberger, "Adiabatic and isocurvature perturbations for multifield generalized Einstein models," Phys. Rev. D 67 (2003) 063512, arXiv:astro-ph/0211276.
- [15] J.-L. Lehners, P. McFadden, N. Turok, and P. J. Steinhardt, "Generating ekpyrotic curvature perturbations before the big bang," Phys. Rev. D 76 (2007) 103501, arXiv:hep-th/0702153.
- [16] I. Agullo, J. Olmedo, and E. Wilson-Ewing, "Observational constraints on anisotropies for bouncing alternatives to inflation," JCAP 10 (2022) 045, arXiv:2206.04037.
- [17] I. Agullo, A. Ashtekar, and W. Nelson, "A Quantum Gravity Extension of the Inflationary Scenario," Phys. Rev. Lett. **109** (2012) 251301, arXiv:1209.1609.
- [18] N. Deruelle and V. F. Mukhanov, "On matching conditions for cosmological perturbations," Phys. Rev. D 52 (1995) 5549–5555, arXiv:gr-qc/9503050.
- [19] R. Brandenberger and F. Finelli, "On the spectrum of fluctuations in an effective field theory of the Ekpyrotic universe," JHEP **11** (2001) 056,

arXiv:hep-th/0109004.

- [20] J.-c. Hwang, "Cosmological structure problem in the ekpyrotic scenario," Phys. Rev. D 65 (2002) 063514, arXiv:astro-ph/0109045.
- [21] E. J. Copeland and D. Wands, "Cosmological matching conditions," JCAP 06 (2007) 014, arXiv:hep-th/0609183.
- [22] R. Brandenberger and P. Peter, "Bouncing Cosmologies: Progress and Problems," Found. Phys. 47 (2017) 797–850, arXiv:1603.05834.
- [23] E. Wilson-Ewing, "Separate universes in loop quantum cosmology: framework and applications," Int. J. Mod. Phys. D 25 (2016) 1642002, arXiv:1512.05743.
- [24] A. Ashtekar, T. Pawlowski, and P. Singh, "Quantum Nature of the Big Bang: Improved dynamics," Phys. Rev. D74 (2006) 084003, arXiv:gr-qc/0607039.
- [25] V. Taveras, "Corrections to the Friedmann Equations from LQG for a Universe with a Free Scalar Field," Phys. Rev. D 78 (2008) 064072, arXiv:0807.3325.
- [26] C. Rovelli and E. Wilson-Ewing, "Why are the effective equations of loop quantum cosmology so accurate?," Phys. Rev. D 90 (2014) 023538, arXiv:1310.8654.
- [27] M. Bojowald and S. Brahma, "Minisuperspace models as infrared contributions," Phys. Rev. D 93 (2016) 125001, arXiv:1509.00640.
- [28] J. G. Kelly, R. Santacruz, and E. Wilson-Ewing, "Black hole collapse and bounce in effective loop quantum gravity," Class. Quant. Grav. 38 (2021) 04LT01, arXiv:2006.09325.
- [29] V. Husain, J. G. Kelly, R. Santacruz, and
  E. Wilson-Ewing, "Quantum gravity of dust collapse: Shock waves from black holes," Phys. Rev. Lett. 128 (2022) 121301, arXiv:2109.08667.
- [30] V. Husain, J. G. Kelly, R. Santacruz, and
   E. Wilson-Ewing, "Fate of quantum black holes," Phys. Rev. D 106 (2022) 024014, arXiv:2203.04238.
- [31] K. Giesel, H. Liu, P. Singh, and S. A. Weigl, "Generalized analysis of a dust collapse in effective loop quantum gravity: fate of shocks and covariance," arXiv:2308.10953.

- [32] R. Gambini, J. Olmedo, and J. Pullin, "Spherically symmetric loop quantum gravity: analysis of improved dynamics," Class. Quant. Grav. 37 (2020) 205012, arXiv:2006.01513.
- [33] J. G. Kelly, R. Santacruz, and E. Wilson-Ewing, "Effective loop quantum gravity framework for vacuum spherically symmetric spacetimes," Phys. Rev. D 102 (2020) 106024, arXiv:2006.09302.
- [34] R. J. LeVeque, Finite Volume Methods for Hyperbolic Problems. Cambridge University Press, 2002.
- [35] L. Cipriani, F. Fazzini, and E. Wilson-Ewing, "Gravitational collapse in effective loop quantum gravity: beyond marginally bound configurations," arXiv:2404.04192.
- [36] The code is available with the source files on the arXiv.org page.
- [37] C. Hellaby and K. Lake, "Shell crossings and the Tolman model," Astrophys. J. 290 (1985) 381.
- [38] F. Fazzini, V. Husain, and E. Wilson-Ewing, "Shell-crossings and shock formation during gravitational collapse in effective loop quantum gravity," Phys. Rev. D 109 (2024) 084052, arXiv:2312.02032.
- [39] Planck, P. A. R. Ade *et al.*, "Planck 2015 results. XVI. Isotropy and statistics of the CMB," Astron. Astrophys. 594 (2016) A16, arXiv:1506.07135.
- [40] A. Ashtekar and D. Sloan, "Loop quantum cosmology and slow roll inflation," Phys. Lett. B 694 (2011) 108–112, arXiv:0912.4093.
- [41] A. Corichi and A. Karami, "On the measure problem in slow roll inflation and loop quantum cosmology," Phys. Rev. D 83 (2011) 104006, arXiv:1011.4249.
- [42] L. Linsefors and A. Barrau, "Duration of inflation and conditions at the bounce as a prediction of effective isotropic loop quantum cosmology," Phys. Rev. D 87 (2013) 123509, arXiv:1301.1264.
- [43] B. Bonga and B. Gupt, "Phenomenological investigation of a quantum gravity extension of inflation with the Starobinsky potential," Phys. Rev. D 93 (2016) 063513, arXiv:1510.04896.