

Amplitude-driven nonreciprocity for energy guiding

Mathieu Padlewski* and Hervé Lissek†

Signal Processing Laboratory 2, Ecole Polytechnique Fédérale de Lausanne, 1015 Lausanne, Switzerland

Romain Fleury‡

Laboratory of Wave Engineering, Ecole Polytechnique Fédérale de Lausanne, 1015 Lausanne, Switzerland

(Dated: October 1, 2024)

The non-Hermitian skin effect is an intriguing physical phenomenon, in which all eigen-modes of a non-Hermitian lattice become localized at boundary regions. While such an exotic behavior has been demonstrated in various physical platforms, these realizations have been so far restricted to the linear regime. Here, we explore the interplay between nonlinearity and the non-Hermitian skin effect which can lead to extraordinary amplitude-driven skin localization dynamics that are unfeasible in linear systems. We show that nonlinear non-Hermitian systems can exhibit new kinds of non-Hermitian skin modes, capable of concentrating the energy of a source at any point of space, depending on the power level. This allows the realization of an energy guide that adapts spontaneously to the amount of energy previously stored in the system: it transfers the energy stored in bulk modes to a point only after a sufficient level of energy has been accumulated. Our theoretical, numerical and experimental findings open up exciting paths for a new generation of non-reciprocal systems, in which nonlinearities serve as a strategic tuning knob to manipulate the guiding process.

MAIN

Non-Hermitian systems [1, 2] have recently attracted significant attention because of their unique physical properties. Unlike their Hermitian counterparts, non-Hermitian systems exhibit intriguing features, such as complex eigenvalue spectra and the non-orthogonality of eigenmodes. These characteristics fundamentally challenge the conventional constraints imposed by Hermitian systems and enable phenomena such as non-reciprocal transport [3–6], exceptional-point-based sensing [7–9] and most notably the non-Hermitian skin effect (NHSE) [10–16] to be realized. This latter effect arises in tight-binding models with asymmetric hoppings, rendering the Hamiltonian inherently non-Hermitian. As a consequence, bulk modes become exponentially localized near specific boundaries, aligning with the dominant direction of non-reciprocal energy flow. Remarkably, the sensitivity of these non-Hermitian skin modes to boundary modifications opens up exciting possibilities for applications such as sensing, ultra-sharp filtering, and interferometry [17].

While the NHSE has been experimentally realized in various physical platforms ranging from photonics [13, 14] to mechanics [5, 18, 19] and electronics [12, 20], previous investigations have primarily focused on linear systems. The interplay between the skin effect and nonlinearity, however, remains largely unexplored.

In this work, we extend the non-Hermitian skin effect into the nonlinear regime. Specifically, we explore the intrinsic nonlinearities of a non-Hermitian lattice, revealing a new class of amplitude-driven non-Hermitian skin




FIG. 1. Nonlinear non-Hermitian skin effect. Consider a tight-binding lattice composed of resonators coupled to each other with specified hopping strengths. The cell hoppings (t_+ and t_-) are non-reciprocal and intensity-dependent, which are key ingredients to nucleate a nonlinear non-Hermitian skin effect. The system is partitioned such as to yield an interface between two finite chains (50 unit cells) of inverted coupling lattices. The eigen-mode distributions are plotted for increasing non-Hermiticity ($I \approx 0 \rightarrow I = I_{max}$). At low intensity, the eigen-mode spectrum consists of many delocalized states, spreading over the bulk of the lattice. As the intensity level is increased, nonlinear dynamics are triggered causing all eigen-modes to localize at the interface.

modes. These modes selectively localize above a critical intensity threshold. Leveraging these findings, we demonstrate their relevance for selectively guiding energy from high-power sources. Moreover, by dynamically tuning the system's nonlinearity level, we can control the characteristics of the skin modes, enabling efficient energy focusing. Our study bridges the gap between nonlinearity and the NHSE, offering novel insights into the

* mathieu.padlewski@epfl.ch

† herve.lissek@epfl.ch

‡ romain.fleuryk@epfl.ch

rich physics of non-Hermitian systems. The emergence of spontaneous non-Hermitian guiding systems promises innovative applications, where energy naturally flows toward arbitrary locations upon exceeding the threshold.

Inspired from the Hatano-Nelson model[10, 11], let's define two tight-binding Hamiltonians, namely \hat{H}^+ and \hat{H}^- , defined as:

$$\hat{H}^\pm = \sum_i U_i |\psi_i\rangle \langle \psi_i| + \sum_{i,j} (t_{i,j}^\pm |\psi_i\rangle \langle \psi_j| + t_{i,j}^\mp |\psi_j\rangle \langle \psi_i|) \quad (1)$$

in which U_i corresponds to an onsite interaction at site i and $t_{i,j}^\pm$ to a coupling interaction site j . $t_{i,j}^\pm$ can be specified to manifest nonlinear character - i.e. have some form of dependency on the eigen-state $|\psi_j\rangle$. More specifically, we assume the form $t_{i,j}^\pm = t \pm \alpha_{i,j} I$, in which t is site-independent, $\alpha_{i,j}$ is the Kerr-like coefficient and $I := \langle \psi_j | \psi_j \rangle$ is the local intensity of the field. The top and bottom insets of Fig. 1 represent the tight-binding lattices corresponding to H^+ and H^- , which are related through spatial inversion. The non-reciprocal character of the coupling interactions $t_{i,j}^+ \neq t_{i,j}^-$ combined with non-linearity can trigger a nonlinear non-Hermitian skin effect, as we demonstrate in the following.

To start, consider a finite one-dimensional chain partitioned in one part described by H^+ and another by H^- such as to form an interface. In other words, the system is configured such that the nonlinear couplings are favored towards the interface. Since the effective Hamiltonians H^\pm are amplitude-dependent, the eigen-modes of the corresponding system are investigated for different excitation energies I and presented in Figure. 1 (see Methods for the details on the simulations). For $I \approx 0$, the eigen-mode distribution across the chain is uniform which is reminiscent of a Hermitian and reciprocal system. Indeed, the latter is simply a consequence of a vanishingly small nonlinear coupling, i.e. $t_{i,j}^\pm \approx t$. Increasing the intensity however, unequivocally drives all the modes towards the interface where they localize.

The remarkable amplitude-dependence of the non-Hermitian skin effect offers a unique opportunity to design exotic systems capable of selectively guiding broadband wave packets to any location in space. Picture a fictional table top composed of atoms that are nonlinearly and nonreciprocally coupled towards its center. A light tap on the surface would evoke usual non-dispersive acoustic waves across the plane whereas, on the other hand, a hard hit would spark nonlinear nonreciprocity and guide the energy toward its center - regardless of the location of the impact. Using iterative numerical techniques (see Methods), this scenario was executed using Eq.(1) and is presented in Fig. 1 (animations can be found in the Supplementary Material). In the following, we turn this fictional system into reality by presenting a state of the art experimental platform capable of producing these nonlinear and non-Hermitian skin effects.




FIG. 2. Nonlinear non-Hermitian skin effect in two dimensions. **a,b** Two dimensional version of the nonlinear model discussed in Fig. 1. The unit cell consists of evanescently coupled resonators with specified coupling coefficients that are arranged such as to guide the energy to the center (red cross-hair). Furthermore, the non-reciprocal extra-cell coupling coefficients t^+ , and t^- are nonlinear, enabling an intensity-dependent non-Hermiticity. **a** Low energy pulse case - typical non-dispersive wave-like behavior. **b** High energy pulse case - energy converges towards the center.

Amplitude-dependant interface skin modes are experimentally realized, closely matching theoretical predictions. A finite metacrystal composed of 16 individually controlled active electroacoustic resonators (AERs) was successfully developed and manufactured - the details of which are found in the Methods. A comprehensive overview of the experimental apparatus is presented detailing the electronic coupling scheme, the unit cell and the coupling configuration of the metacrystal (Figs 3a-c). The first half of the metacrystal is electronically configured such as to favor nonlinear coupling to the right neighbors and the other to the left. In other words, the metacrystal can host amplitude-dependant skin modes at its center (at site $n^{\circ}9$). Note that the interface can be arbitrarily shifted to any site along the chain (See Fig.S1 in the Supplementary Material). Amplitude-dependent third order nonlinear couplings are achieved by means of an active feedback loop as explained in Figure 3a. Amplitude-dependency is demonstrated by the introduction of a broadband acoustic pulse of low intensity at one end of the chain. As shown in Fig. 3d, the signal propagates unimpeded throughout the system until it reaches the anechoic termination. Conversely, for a high intensity pulse, the behavior is strikingly different. With the activation of the amplitude-driven nonreciprocal coupling, the signal increases in amplitude up to the skin interface and sharply drops after passing the crest.




FIG. 3. Experimental broadband nonlinear non-Hermitian sound guiding. **a** Electronic coupling scheme of an active electroacoustic resonator (AER) that depends on the pressure amplitude of the nearest neighbor. The non-reciprocal extra-cell coupling coefficients t^+ , and t^- are nonlinear, enabling an intensity-dependent non-Hermiticity. **b** Close-up of an active cell. **c** Cross-section of the entire metacrystal composed of 16 individually controlled AERs. The waveguide is capped by a source speaker on one end and an anechoic termination on the other. **d,e** Spatio-temporal plots of a system configured such as to have nonlinear couplings guiding the energy to the center (highlighted in grey) for low and high energy pulse excitations respectively.

To summarize, the intrinsic nonlinearity of a non-Hermitian lattice was studied, leveraging it to induce the non-Hermitian skin effect. Amplitude-driven skin modes allow for a novel spontaneous energy guiding system, which collects energy from the pump during high-power operation and conserves it when the energy level is low. We demonstrate that the proposed system can be dynamically tuned by adjusting the excitation intensity. While our investigation focuses on a simple coupling scheme, future work should explore the interplay between nonlinearity, NHSE and topology by configuring the system such as to obtain a nonlinear non-hermitian SSH chain. Overall, our theoretical and experimental findings contribute to our understanding of the physics behind the non-Hermitian skin effect and hold promise for engineering applications such as lasing, sensing, and signal processing.

METHODS

A. Numerical details

The nonlinearity of our proposed model is related to the fact that the associated Hamiltonian is a function of its eigen-modes. Therefore, the associated eigen-vectors should be calculated throughout a recursive process. In this process, an initial guess is assumed for the eigen-vectors. This initial guess is used to generate a new Hamiltonian, based on which a new set of eigen-vectors is obtained. This procedure is repeated until achieving a converged solution. In our eigen-mode calculations, we used the relative error of 1 % as the convergence criterion. Our initial guess for the eigen-vector $|\psi\rangle$ was the solution of Eq. (1) in the absence of nonlinearity (note that for the linear case the eigen vectors of Eq. (1) can be easily obtained by setting the determinant of the Hamiltonian to zero). The results of Fig. 2 of the manuscript were obtained by solving the matrix equation $i\partial_t |\psi\rangle = H |\psi\rangle$, in which H is the Hamiltonian of the system. A Runge-

Kutta iterative method was employed to solve the associated temporal equation using a weighted Kronecker-Delta as the initial condition.

B. Active Acoustic Crystal Design and Control Scheme

The acoustic waveguide is composed of CNC-milled 20mm thick PVC slabs lined with 16 off-the-shelf Visaton FRWS 5 - 8 Ohm drivers, 3D-printed back-cavities, and quarter-inch PCB microphones. The pressure microphones placed at the AERs (Fig. 3b) serve both for interaction control and mode probing. Active control is carried out by a commercial Speedgoat machine equipped with the IO135 module. This Direct Memory Access (DMA) module enables both on-site tunability

and low-latency driver control over all 16 channels. COMSOL Multiphysics finite element simulations were performed enabling design optimization and experimental validation for linear coupling configurations. In order to account for manufacturing inconsistencies, the mechanical mass, resistance and compliance of each speaker was characterized following methods described by E. Rivet [21] and actively tuned to match one another. Finally, the quality factor of the resonators where all enhanced by synthetically reducing AER acoustic resistance by 80%. (- details in Supplementary Material). As a last note, both the tuning and reduced resistance are captured by the Φ gate in Figure 3.

Acknowledgements This research was supported by the Swiss National Science Foundation under Grant No. 200020.200498.

-
- [1] Carl M. Bender and Stefan Boettcher. Real spectra in non-hermitian hamiltonians having PT-symmetry. 80(24):5243–5246. Publisher: American Physical Society.
- [2] Carl M. Bender. Making sense of non-hermitian hamiltonians. 70(6):947.
- [3] Zin Lin, Hamidreza Ramezani, Toni Eichelkraut, Tsampikos Kottos, Hui Cao, and Demetrios N. Christodoulides. Unidirectional invisibility induced by PT-symmetric periodic structures. 106(21):213901. Publisher: American Physical Society.
- [4] Romain Fleury, Dimitrios L. Sounas, Caleb F. Sieck, Michael R. Haberman, and Andrea Alù. Sound isolation and giant linear nonreciprocity in a compact acoustic circulator. 343(6170):516–519. Publisher: American Association for the Advancement of Science.
- [5] Martin Brandenbourger, Xander Locsin, Edan Lerner, and Corentin Coulais. Non-reciprocal robotic metamaterials. 10(1):4608. Publisher: Nature Publishing Group.
- [6] Xinxin Guo, Hervé Lissek, and Romain Fleury. Observation of non-reciprocal harmonic conversion in real sounds. 6(1):1–6. Publisher: Nature Publishing Group.
- [7] Zhicheng Xiao, Huanan Li, Tsampikos Kottos, and Andrea Alù. Enhanced sensing and non-degraded thermal noise performance based on $\mathcal{P}\mathcal{T}$ -symmetric electronic circuits with a sixth-order exceptional point. 123(21):213901. Publisher: American Physical Society.
- [8] Jan Wiersig. Prospects and fundamental limits in exceptional point-based sensing. 11(1):2454. Publisher: Nature Publishing Group.
- [9] Jan Wiersig. Review of exceptional point-based sensors. 8(9):1457–1467. Publisher: Optica Publishing Group.
- [10] Naomichi Hatano and David R. Nelson. Localization transitions in non-hermitian quantum mechanics. 77(3):570–573. Publisher: American Physical Society.
- [11] Naomichi Hatano and David R. Nelson. Non-hermitian delocalization and eigenfunctions. 58(13):8384–8390. Publisher: American Physical Society.
- [12] Hui Jiang, Li-Jun Lang, Chao Yang, Shi-Liang Zhu, and Shu Chen. Interplay of non-hermitian skin effects and anderson localization in nonreciprocal quasiperiodic lattices. 100(5):054301. Publisher: American Physical Society.
- [13] Xueyi Zhu, Huaiqiang Wang, Samit Kumar Gupta, Haijun Zhang, Biye Xie, Minghui Lu, and Yanfeng Chen. Photonic non-hermitian skin effect and non-bloch bulk-boundary correspondence. 2(1):013280. Publisher: American Physical Society.
- [14] Sebastian Weidemann, Mark Kremer, Tobias Helbig, Tobias Hofmann, Alexander Stegmaier, Martin Greiter, Ronny Thomale, and Alexander Szameit. Topological funneling of light. 368(6488):311–314. Publisher: American Association for the Advancement of Science.
- [15] Xiujuan Zhang, Yuan Tian, Jian-Hua Jiang, Ming-Hui Lu, and Yan-Feng Chen. Observation of higher-order non-hermitian skin effect. 12(1):5377. Publisher: Nature Publishing Group.
- [16] Xiujuan Zhang, Tian Zhang, Ming-Hui Lu, and Yan-Feng Chen. A review on non-hermitian skin effect. 7(1):2109431. Publisher: Taylor & Francis.
- [17] Linhu Li, Ching Hua Lee, and Jiangbin Gong. Topological switch for non-hermitian skin effect in cold-atom systems with loss. 124(25):250402. Publisher: American Physical Society.
- [18] Ananya Ghatak, Martin Brandenbourger, Jasper van Wezel, and Corentin Coulais. Observation of non-hermitian topology and its bulk-edge correspondence in an active mechanical metamaterial. 117(47):29561–29568. Publisher: Proceedings of the National Academy of Sciences.
- [19] Colin Scheibner, William T.M. Irvine, and Vincenzo Vitelli. Non-hermitian band topology and skin modes in active elastic media. 125(11):118001. Publisher: American Physical Society.
- [20] Li-Jun Lang, Yijiao Weng, Yunhui Zhang, Enhong Cheng, and Qixia Liang. Dynamical robustness of topological end states in nonreciprocal su-schrieffer-heeger models with open boundary conditions. 103(1):014302. Publisher: American Physical Society.
- [21] Etienne Rivet, Sami Karkar, and Herve Lissek. Broadband low-frequency electroacoustic absorbers through

hybrid sensor-/shunt-based impedance control. 25(1):10.
63–72. Publisher: Institute of Electrical and Electronics

Engineers Place: Piscataway.