

# THE EXACT QUANTUM CHROMATIC NUMBER OF HADAMARD GRAPHS

A. MEENAKSHI MCNAMARA

ABSTRACT. We compute the exact value of the quantum chromatic numbers of Hadamard graphs of order  $n = 2^N$  for  $N$  a multiple of 4 using the upper bound derived by Avis, Hasegawa, Kikuchi, and Sasaki, as well as an application of the Hoffman-like lower bound of Clive and Wocjan that was generalized by Ganesan for quantum graphs.

## INTRODUCTION

Quantum coloring of graphs is a modification of the usual notion of graph coloring using non-local games. In a non-local game, players Alice and Bob collaborate to answer pairs of questions without communication using some prior agreed-upon strategy. The possible strategies used by the players have many classifications (see, for instance, [CMN<sup>+</sup>07] and [OP16]), including classical strategies and “quantum” strategies where a shared entangled resource is available. The use of entanglement often allows quantum strategies to outperform classical ones, leading to a “quantum advantage.” In fact, the famous CHSH inequality may be described as such a quantum advantage in a non-local game. There has been especially great interest in non-local games in recent years due to their importance in quantum information theory. Further, the resolution of the Connes Embedding Conjecture through showing that  $\text{MIP}^* = \text{RE}$  in [JNV<sup>+</sup>22] relied upon non-local games. Hence, studying non-local games is important to understanding the broader impacts of this result.

In the graph coloring game, the players wish to convince a referee that they have a coloring for a graph, i.e., an assignment of colors to vertices of the graph such that adjacent vertices are different colors. Graph coloring games are of particular interest because they provide explicit examples of non-local games, and often naturally tie into other problems in quantum information theory and communication as seen in [AHKS06]. Additionally, there are numerous modifications of these games allowing for the concrete study of different models of non-local games, as has been done in [OP16] where the equivalence of graph coloring questions to the Connes Embedding Conjecture through Tsirelson’s Problem were studied. Recent work by Harris in [Har24] showed that all non-local games may be reformulated as a kind of graph coloring games; furthering the importance of graph coloring in the study of non-local games.

In particular, when studying graph coloring games we are interested in the minimum number of colors needed to win. Restricting to quantum strategies, we call this the quantum chromatic number of the graph,  $\chi_q$ , and it is known that determining this number is an NP hard problem in general [Ji13]. Numerous examples of graphs  $G$  have been found exhibiting a quantum advantage such that  $\chi_q(G) < \chi(G)$ , where  $\chi(G)$  is the usual (i.e., classical) chromatic number of  $G$ . In particular [AHKS06] found the first such examples to be Hadamard graphs on  $2^N$  vertices with  $N > 3$  a multiple of 4,  $H_N$ . In particular for  $N = 12$  we have  $\chi(H_{12}) = 13$  while  $\chi_q(H_{12}) = 12$ . These upper bounds on the quantum chromatic number of the Hadamard graph can be understood by viewing the Hadamard graph as an orthogonality graph, and

similar results for general orthogonality graphs were developed in [CMN<sup>+</sup>07]. However, in these examples, the computations of the quantum chromatic of graphs give only upper bounds. Further, many of the lower bounds are given by other non-local invariants such as the tracial rank defined in [PSS<sup>+</sup>16], and thus are similarly difficult to study. Hence, except for a few cases of graphs without any quantum advantage (such as the graphs studied in [EW19] and the subfamily of Kneser graphs studied in [WED20]) or with few colors needed in their classical coloring (for example, see [CMN<sup>+</sup>07]), there are very few examples of the precise quantum chromatic numbers of graphs being known.

In this paper, we will study the quantum chromatic number of the Hadamard graphs; giving precise results for an infinite family of graphs. Note that throughout this work, we consider Hadamard graphs as defined by Ito in [Ito85a] and [Ito85b] (see Sections 1.2 and 1.3). This is in contrast to another, perhaps more common, definition of Hadamard graphs that comes more directly from the notion of Hadamard matrices, such as those considered in [CM24]. However, our notion of Hadamard graphs is still related to Hadamard matrices. In particular, as is discussed in [Ito85a], the conjecture of the existence of Hadamard matrices for every  $N$  a multiple of 4 can be rephrased as conjecturing that the clique number of  $H_N$  is  $N$ . These graphs have been studied in such depth in the quantum coloring setting due to their connection to quantum information theory problems. For instance, the first results concerning quantum coloring Hadamard graphs are contained within [BCT99] for  $N = 2^n$ , in the process of studying the problem of simulating entangled systems using classical hidden variable models and communication. In these studies, an upper bound for the quantum chromatic number of Hadamard graphs has been established in [AHKS06]. In fact, using results of [FR87], it was shown in [AHKS06] that Hadamard graphs provide examples of an exponential gap between the quantum and classical chromatic numbers. These results on quantum coloring Hadamard graphs were further applied to the problem of simulating quantum entanglement via classical communication in [GTW13]. However, despite the success of developing upper bounds, no lower bounds on the quantum chromatic number of these graphs have been established.

In general, very few graphs have had concrete lower bounds on their quantum chromatic numbers established. In this work, we will make use of the spectral lower bound derived in [EW19]. These results were generalized to quantum graphs in [Gan23], and in fact the proof of these results was simplified using the notions of quantum coloring on quantum graphs. As classical graphs are special cases of quantum graphs, we will emphasize the applicability of our approach to quantum graphs as well by phrasing results using the language of quantum graphs when convenient. This is especially relevant as interest in quantum graphs has been growing in recent years due to their connections with quantum information theory. In fact, quantum graphs were developed to generalize the role of graphs in classical information theory to the quantum setting [DSW13]. Quantum graphs are a non-commutative topology generalization of graphs that have been independently developed several times leading to numerous equivalent descriptions (see [EKS98, DSW13, Wea21, MRV18, BCE<sup>+</sup>20]). The notion of quantum chromatic numbers has been extended to quantum graphs, and the quantum adjacency matrix approach of [MRV18] lends itself to spectral lower bounds as found in Ganesan's work in [Gan23] where she generalized the results of [EW19]. In our

work, these lower bounds will prove particularly useful in the study of Hadamard graphs due to an equivalent description of these graphs as conjugacy class graphs, allowing for the spectral results to be found using character theory as in [Ito85a].

Using these lower bounds in addition to the more usual approach to obtaining upper bounds on quantum chromatic numbers, we obtain our main result:

**Theorem 0.1** (Main Theorem). *Let  $H_N$  be the Hadamard graph on  $2^N$  vertices,  $N$  a multiple of 4. Then,*

$$\chi_q(H_N) = N.$$

The structure of this paper is as follows. In Section 1 we provide the necessary background on quantum coloring and quantum graphs, and the various viewpoints on Hadamard graphs. Then, Section 2 is divided into two main subsections: First, in Section 2.1 we review the proof of the upper bound for the quantum chromatic number of Hadamard graphs, reformulating it in the language of positive operator valued measures. Then, in Section 2.2 we review results of Ito on Hadamard graphs and develop lower bounds, leading to our result.

**Acknowledgements.** The author would like to thank Rolando de Santiago for his guidance and help throughout this project. The author would also like to thank Priyanga Ganesan for thoughtful discussions about quantum coloring and non-local games. Research at Perimeter Institute is supported in part by the Government of Canada through the Department of Innovation, Science and Economic Development and by the Province of Ontario through the Ministry of Colleges and Universities.

## 1. BACKGROUND

**1.1. Quantum Colorings of Graphs.** In a non-local game, cooperating players Alice and Bob play against a referee by answering questions while separated such that no communication is allowed (hence, non-local). The referee will ask both players questions, and determine if they win a given (independent) round based on their answers. Alice and Bob may agree on a strategy ahead of time, and we call a strategy winning if they win with probability 1. There are numerous strategies Alice and Bob may be allowed to use, we are primarily interested in classical (i.e. local) and quantum strategies. Other strategies often studied include quantum approximate, quantum commuting, and non-signaling.

A quantum strategy allows Alice and Bob to share an entangled state  $\psi \in \mathbb{C}^a \otimes \mathbb{C}^b$  that they measure with positive operator valued measures (POVMs)  $\{E_{x_A, y_A}\} \in M_a$  and  $\{F_{x_B, y_B}\} \in M_b$  corresponding to the referee asking questions  $x_A$  and  $x_B$  to Alice and Bob, respectively, and Alice and Bob responding with  $y_A$  and  $y_B$  respectively. Using this setup, the chance that Alice and Bob respond with  $y_A$  and  $y_B$  given questions  $x_A$  and  $x_B$  is

$$p(y_A, y_B | x_A, x_B) = \langle \psi | E_{x_A, y_A} \otimes F_{x_B, y_B} | \psi \rangle.$$

Alternatively, in a local strategy no entanglement is present. This corresponds to requiring that  $\psi$  is a product state in the strategy above. Equivalently, local strategies satisfy  $p(y_A, y_B | x_A, x_B) = p_1(y_A | x_A)p_2(y_B | x_B)$  for some probabilities  $p_1$  and  $p_2$  [PSS<sup>+</sup>16].

The non-local game we will consider in this paper is the graph coloring game. Classically, an  $m$ -coloring of a graph  $G$  is an assignment of a list of colors  $m$  colors to  $V(G)$  such that adjacent vertices are given different colors. The chromatic number of a graph  $\chi(G)$  is the smallest value of  $m$  such that the graph has an  $m$ -coloring. In the **graph coloring game** Alice and Bob want to convince a referee that they have a coloring for a graph. To do so, the referee gives Alice and Bob vertices of the graph  $x_A, x_B \in V(G)$ , and the players respond with colors  $y_A, y_B \in \{1, 2, \dots, c\}$  for the vertices. Alice and Bob have a winning strategy if their responses follow the rules of graph coloring, i.e., if (1)  $p(y_A \neq y_B | x_A = x_B) = 0$ , and (2) if  $p(y_A = y_B | x_A \sim x_B) = 0$ . Condition (1) is precisely the definition of a synchronous game. Thus, Bob's POVMs can be obtained from Alice's as described in [CMN<sup>+</sup>07]. Classically, a winning strategy must use at least  $\chi(G)$  colors. However, if Alice and Bob use a quantum strategy it may be possible to win using fewer colors. We call the number of colors needed for a winning quantum strategy the **quantum chromatic number** of a graph,  $\chi_q(G)$ .

These types of coloring games have also been generalized to quantum graphs in [BGH22]. One approach to studying coloring on quantum graphs by [BGH22] uses the operator space notion: an (irreflexive) **quantum graph** is a tuple  $(\mathcal{S}, \mathcal{M}, \mathcal{B}(H))$  where  $\mathcal{M} \subset \mathcal{B}(H)$  is a finite dimensional, von Neumann algebra and  $\mathcal{S}$  is an operator space that is a bimodule over  $\mathcal{M}'$  that satisfies  $\mathcal{S} \perp \mathcal{M}'$ . Here, we consider a **quantum coloring** to be a projection valued measure (PVM)  $\{P_a\}_{a \in [c]} \in \mathcal{M} \bar{\otimes} \mathcal{N}$  where  $[c]$  are the colors,  $\mathcal{N}$  is a finite dimensional von Neumann algebra, and the projections satisfy the coloring relation

$$P_a(S \otimes I_{\mathcal{N}})P_a = 0 \forall a \in \{1, 2, \dots, c\}.$$

We may modify the definition to generalize other strategies by changing the conditions on  $\mathcal{N}$  (see [BGH22]).

Another approach to quantum graphs that we will use is the **quantum adjacency matrix**. We consider a quantum set to be a pair  $(\mathcal{M}, \psi)$  where  $\mathcal{M}$  is a finite dimensional  $C^*$ -algebra and  $\psi : \mathcal{M} \rightarrow \mathbb{C}$  is a faithful state. Note that we may identify  $\mathcal{M}$  with  $L^2(\mathcal{M})$ , where  $L^2(\mathcal{M}) = L^2(\mathcal{M}, \psi)$  is the GNS completion of  $\mathcal{M}$  with respect to  $\psi$ . Hence, given  $m : \mathcal{M} \otimes \mathcal{M} \rightarrow \mathcal{M}$  via multiplication we obtain  $m^*$  its adjoint on  $L^2(\mathcal{M})$ . We say  $\psi$  is a  $\delta$ -form if there exists  $\delta > 0$  such that  $mm^* = \delta^2 I$ . Then, the tuple  $(\mathcal{M}, \psi, A)$  is a quantum graph with quantum adjacency matrix  $A : L^2(\mathcal{M}) \rightarrow L^2(\mathcal{M})$  if  $A$  is schur idempotent ( $m(A \otimes A)m^* = \delta^2 A$ ) and satisfies  $(I \otimes \eta^* m)(I \otimes A \otimes I)(m^* \eta \otimes I) = A$  where  $\eta : \mathbb{C} \rightarrow \mathcal{M}$  via  $\lambda \mapsto \lambda I$ . An irreflexive quantum graph further satisfies  $m(A \otimes I)m^* = 0$ . Throughout the remainder of this paper we will assume that all (quantum) graphs are irreflexive as coloring ideas are best applied in this setting. These two equivalent definitions are described and compared in [Gan23].

The quantum adjacency matrix approach has allowed many spectral lower bounds to be extended to quantum coloring as was done in [Gan23]. In particular, we will make use of the bound **Hoffman bound** for quantum coloring of classical graphs that was found in [EW19], phrased here for its generalization to quantum graphs given by [Gan23]: Let  $\mathcal{G} = (\mathcal{M}, \psi, A)$  be a quantum graph, and let  $\lambda_{max} = \lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_{\dim \mathcal{M}} = \lambda_{min}$  be all the eigenvalues of  $A$  (with multiplicity). Then  $1 + \frac{\lambda_{max}}{|\lambda_{min}|} \leq \chi_q(\mathcal{G})$ .

Both notions of quantum graphs generalize classical graphs, and hence results for quantum graphs specialize to classical graphs. In the operator space approach a classical graph corresponds to a quantum graph where  $\mathcal{M}$  is abelian. Likewise, in the quantum adjacency matrix approach a classical graph corresponds to

the usual notion of adjacency matrices, i.e., a  $|V| \times |V|$  matrix with 1 at  $ij$ th entry if  $i \sim j$  in  $G$  and zeros elsewhere.

**1.2. Hadamard Graphs as Orthogonality Graphs.** Let  $N \in \mathbb{N}$  be a multiple of 4. The Hadamard graph of size  $n = 2^N$  is a graph defined as follows: the vertices are vectors of size  $N$  with entries  $\pm 1$ ; two vertices are adjacent iff the inner product of the vectors is 0. We will denote the vertex  $x$  as the vector  $(x_i)_{i=0}^{N-1}$  where  $x_i$  is the  $i$ -th component of the vector. In this notation, the vertices  $x$  and  $y$  are adjacent iff the vectors  $(x_i)_{i=0}^{N-1}$  and  $(y_i)_{i=0}^{N-1}$  differ in exactly half of the entries. Note that this makes the Hadamard graph an orthogonality graph as we are representing the vertices as vectors and edges are determined by the orthogonality relation.

*Remark 1.* Note that the Hadamard graph defined above is disconnected. In particular, vertices with an even number of  $+1$ s and  $-1$ s and those with an odd number form identical connected components. Hence, we may restrict the Hadamard graph to the graph on  $2^{N-1}$  vertices corresponding to the connected component containing the all  $1$ s vector (the component with even numbers of  $+1$ s and  $-1$ s).

*Remark 2.* It is convenient to define the POVMs for coloring using a slightly different notation for the Hadamard graph. Namely, the Hadamard graph of size  $n = 2^N$  is defined as follows: the vertices are vectors of size  $N$  with entries  $0$ s and  $1$ s; two vertices are adjacent iff exactly half of the vector entries differ. Note that this is equivalent to the above definition with  $-1$ s replaced by  $0$ s. This convention will be applied for notational convenience.

**1.3. Hadamard Graphs as Conjugacy Class Graphs.** In [Ito85a] it was shown that the Hadamard graph can also be viewed as a conjugacy class graph in the sense of [Ito84]. In a conjugacy class graph, the vertices may be identified with elements of a group  $\Gamma$ . Further, there is some  $C \subset \Gamma$  that is a union of conjugacy classes, satisfies  $C = C^{-1}$  and generates  $\Gamma$  such that vertices  $x$  and  $y$  are adjacent iff  $x = cy$  for some  $c \in C$ . Characterizing the Hadamard graph as a conjugacy class graph will allow for the use of irreducible characters in order to calculate the spectrum of the adjacency matrix of the Hadamard graph, as will be further developed in Section 2.2.

To view the Hadamard graph as a conjugacy class graph we will consider the Hadamard graph as described in Remark 1. Following [Ito85a], we see that the vertices form an abelian group with group multiplication given by componentwise multiplication of the vectors and unit the all  $1$ s vector  $h$ . If we let  $C$  be the set of vertices adjacent to  $h$ , then vertices  $x$  and  $y$  are adjacent iff  $x = cy$  for some  $c \in C$ . We note that  $C$  is the union of conjugacy classes since in an abelian group conjugacy classes are single elements. Additionally,  $C = C^{-1}$  as each element is its own inverse. To see that  $C$  generates the group, let  $w(i, j)$  be the vector of all  $+1$ s except at the  $i$ th and  $j$ th spot where there are  $-1$ s. Observe  $\{w(0, 1), w(0, 2), \dots, w(0, N)\}$  is a generating set for the group. Further, the elements  $w(0, i_1) \cdot w(0, i_2) \cdots w(0, i_{N/2})$  and  $w(0, i_2) \cdot w(0, i_3) \cdots w(0, i_{N/2})$ , for  $0 < i_k < N$  and  $i_j \neq i_k$  for all  $j, k$ , are both in  $C$ , and we notice that  $\binom{N-1}{N/2} + \binom{N-1}{N/2-1} = \binom{N}{N/2}$  so all elements of  $C$  are of this form. Then we notice that the product of these elements is  $w(0, i_1)$  for any  $i_1 \in \{0, \dots, N-1\}$ , so  $C$  generates the group.

## 2. COMPUTING THE CHROMATIC NUMBER

**2.1. Upper Bounds.** In this section, we will describe the work of [AHKS06] which shows that the quantum chromatic number of the Hadamard graph on  $2^N$  vertices with  $N$  a multiple of 4 is less than or equal to  $N$ . Note that, as is noted in [AHKS06], this result along with [FR87] implies an exponential gap between the chromatic number and quantum chromatic number of Hadamard graphs with  $N = 4p^q$ . Hence, this work provided an early example of quantum advantage in non-local games.

In the work of [AHKS06], the authors described a winning strategy to color the Hadamard graph on  $2^N$  vertices using  $N$  colors. Given vertices  $(a_i)$  and  $(b_i)$  in the convention used in Remark 2, Alice and Bob win using the following protocol:

- (1) Prepare an initial state  $|\psi\rangle := \frac{1}{\sqrt{N}} \sum_{j=0}^{N-1} |j\rangle \otimes |j\rangle$
- (2) Alice and Bob apply phase shifts using  $P_{a_i}$  and  $P_{b_i}$ , where  $P_{l_i}|i\rangle = (-1)^{l_i}|i\rangle$ . This results in  $(P_{a_i} \otimes P_{b_i})|\psi\rangle$ .
- (3) Apply the general quantum Fourier transform to get

$$(QFT_N \otimes QFT_N^{-1})(P_{a_i} \otimes P_{b_i})|\psi\rangle = \frac{1}{N^{3/2}} \sum_{j=0}^{N-1} \sum_{k=0}^{N-1} \sum_{\ell=0}^{N-1} (\omega)^{\ell(j-k)} (-1)^{a_i \oplus b_i} |j\rangle \otimes |k\rangle.$$

- (4) Measure the state using this computational basis. This will obtain one of  $N$  basis states, and they return the corresponding color to the referee.

We wish to reframe this strategy into the familiar POVM framework of section 1.1. Recall that as this is a synchronous game we may assume that Alice and Bob use the same POVM, and also that they are projections. Hence, we must find a set of projections  $\{E_{x,\alpha}\}_{x \in V(H_n), \alpha \in [n]}$  such that  $\sum_{\alpha \in [n]} E_{x,\alpha} = I$ , and such that using these projections and their conjugates for Alice and Bob is a winning strategy (namely, we must check that given adjacent vertices they return different colors).

**Theorem 2.1** (Theorem 3.1 in [AHKS06]). *Let  $G$  be the Hadamard graph on  $2^N$  vertices where  $N$  is a multiple of 4. Then  $\chi_q(G) \leq N$*

*Proof.* We will adopt the convention for Hadamard graphs used in Remark 2. Fix  $\omega$ , a primitive root of unity. For each  $\alpha \in [N]$   $x \in V(G)$ , define

$$\langle \tilde{E}_x^\alpha | = \frac{1}{\sqrt{N}} \sum_{j=0}^{N-1} \omega^{j\alpha} (-1)^{x_i} \langle j |$$

and set  $E_x^\alpha = |\tilde{E}_x^\alpha\rangle\langle \tilde{E}_x^\alpha|$ . Note that since  $\tilde{E}_x^\alpha$  is a unit vector, it is readily checked that  $E_x^\alpha$  is a projection for each  $x \in V(G)$  and  $\alpha \in [N]$ . Consider the shared entangled state  $|\psi\rangle = \frac{1}{\sqrt{N}} \sum_{j=0}^{N-1} |j\rangle \otimes |j\rangle$  We now verify

that  $\{E_x^\alpha\}_{\alpha \in [N]}$  forms a PVM for each  $x \in V(G)$ . To this end,

$$\begin{aligned} \sum_{\alpha=1}^N E_x^\alpha &= \sum_{\alpha=1}^N \frac{1}{N} \sum_{j,k=0}^{N-1} \omega^{(j-k)\alpha} (-1)^{x_j+x_k} |k\rangle\langle j| \\ &= \left( \frac{1}{N} \sum_{j,k=0}^{N-1} (-1)^{x_j+x_k} |k\rangle\langle j| \right) \sum_{\alpha=1}^N \omega^{(j-k)\alpha} \\ &= \sum_{j=0}^{N-1} |j\rangle\langle j| = I \end{aligned}$$

where in the second to the last line, we used the following facts: if  $j = k$  then  $(-1)^{x_j+x_k} = 1$  and  $\sum_{\alpha=1}^N \omega^0 = N$ ; and if  $j \neq k$  then we get  $\sum_{\alpha=1}^N \omega^{(j-k)\alpha} = 0$ .

Next, we verify the coloring condition. First we calculate,

$$\begin{aligned} \langle \psi | E_x^\alpha \otimes \overline{E_y^\beta} | \psi \rangle &= \frac{1}{N^3} \sum_{\ell=0}^{N-1} \langle \ell | \otimes \langle \ell | \left( \sum_{j,k=0}^{N-1} \omega^{(j-k)\alpha} (-1)^{x_j+x_k} |k\rangle\langle j| \right) \otimes \\ &\quad \left( \sum_{j',k'=0}^{N-1} \omega^{(k'-j')\beta} (-1)^{y_{j'}+y_{k'}} |k'\rangle\langle j'| \right) \sum_{\ell'=0}^{N-1} |\ell'\rangle \otimes |\ell'\rangle \\ &= \frac{1}{N^3} \sum_{\ell,\ell'=0}^{N-1} \left( \omega^{(\ell'-\ell)\alpha} (-1)^{x_{\ell'}+x_\ell} \right) \left( \omega^{(\ell'-\ell)\beta} (-1)^{y_{\ell'}+y_\ell} \right) \\ &= \frac{1}{N^3} \sum_{\ell,\ell'=0}^{N-1} \omega^{(\ell'-\ell)(\alpha+\beta)} (-1)^{x_\ell+x_{\ell'}+y_\ell+y_{\ell'}} \end{aligned}$$

Thus we have

$$(1) \quad \langle \psi | E_x^\alpha \otimes \overline{E_y^\beta} | \psi \rangle = \frac{1}{N^3} \sum_{\ell=0}^{N-1} \omega^{\ell(\alpha-\beta)} (-1)^{x_\ell+y_\ell} \sum_{\ell'=0}^{N-1} \omega^{\ell'(\beta-\alpha)} (-1)^{x_{\ell'}+y_{\ell'}}$$

Now we wish to verify that if  $x \sim y$  and  $\alpha = \beta$  then  $\langle \psi | E_x^\alpha \otimes \overline{E_y^\beta} | \psi \rangle$  evaluates to 0. Using these assumptions in Equation (1), we find

$$\langle \psi | E_x^\alpha \otimes \overline{E_y^\beta} | \psi \rangle = \frac{1}{N^3} \sum_{\ell=0}^{N-1} (-1)^{x_\ell+y_\ell} \sum_{\ell'=0}^{N-1} (-1)^{x_{\ell'}+y_{\ell'}} = 0$$

since exactly half of the  $x_\ell + y_\ell$  are even and exactly half are odd when  $x \sim y$ .

Finally, since Bob's POVMs are found from Alice's POVMs as described in [CMN<sup>+</sup>07] this must be a synchronous game. Hence, it is easily checked that if  $\alpha \neq \beta$  then

$$\langle \psi | E_x^\alpha \otimes \overline{E_x^\beta} | \psi \rangle = 0.$$

□

*Remark 3.* We obtain the PVM corresponding to colors used in coloring quantum graphs as in [BGH22] by considering the following block diagonal matrix,

$$P_\alpha = \begin{pmatrix} E_1^\alpha & 0 & \cdots & 0 \\ 0 & E_2^\alpha & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & E_n^\alpha \end{pmatrix},$$

where the vertices are labeled using the set  $[n]$ . Notice that we have  $P_\alpha \in D_n \otimes M_N(\mathbb{C})$  as  $E_x^\alpha \in M_N(\mathbb{C})$ , so as  $M_N(\mathbb{C})$  is finite dimensional this is a quantum coloring in the above description as expected.

The verification that  $\{P_\alpha\}_\alpha$  forms a PVM is immediate from the result for  $\{E_x^\alpha\}_\alpha$ , and the coloring condition may be checked a quick calculation verifying that

$$P_\alpha(E_{x,y} \otimes I_N)P_\alpha = 0$$

whenever  $x \sim y$ , where  $E_{x,y} \in M_n(\mathbb{C})$  is the elementary matrix with a 1 at the  $xy$  entry and zero elsewhere.

**2.2. Lower Bounds.** We will now derive a lower bound for the quantum chromatic number of the Hadamard graph. In particular, we will use the Hoffman bound that was shown in [Gan23] using eigenvalues of the quantum adjacency matrix. As this is a classical graph, the quantum adjacency matrix is precisely the usual concept of an adjacency matrix. Hence, we will use the results of Ito in [Ito84] and [Ito85a] in order to compute the eigenvalues by viewing the Hadamard graph as a conjugacy class graph.

We begin by noting that conjugacy class graphs are determined by a group and a (union of) conjugacy classes of the group. Hence, representation theory is useful in studying such graphs, and we shall review some of the relevant definitions. In particular, a representation of a group  $\Gamma$  on a vector space  $V$  is a group homomorphism  $\rho : \Gamma \rightarrow GL(V)$ . A representation is *irreducible* if there are no invariant subspaces of  $\rho(\Gamma)$  except for  $\{0\}$  and  $V$  itself. If  $V$  is finite dimensional (as will be considered here), then the *character* corresponding to  $\rho$  is  $\chi_\rho : \Gamma \rightarrow \mathbb{C}$  via  $\chi_\rho(g) = \text{Tr}(\rho(g))$  for any  $g \in \Gamma$ . A character  $\chi_\rho$  is called irreducible if  $\rho$  is irreducible. Characters possess several properties that will be used in the following:

- Characters are constant on conjugacy classes of  $\Gamma$  since  $\text{Tr}(ab) = \text{Tr}(ba) \implies \text{Tr}(aba^{-1}) = \text{Tr}(b)$ .
- The set of class functions (functions that take a constant value on conjugacy classes) has a natural inner product:  $\langle \alpha, \beta \rangle = \frac{1}{|\Gamma|} \sum_{g \in \Gamma} \alpha(g) \overline{\beta(g)}$ . Further, if  $\alpha$  and  $\beta$  are characters corresponding to a finite dimensional representation, and  $\Gamma$  is of finite order then

$$(2) \quad \langle \alpha, \beta \rangle = \frac{1}{|\Gamma|} \sum_{g \in \Gamma} \alpha(g) \beta(g^{-1}),$$

because finite order invertible linear transformations have the trace of their inverse is the complex conjugate of the trace.

- The set of irreducible characters forms an orthonormal basis for the class functions. In particular, if  $\rho = \rho_1 \oplus \rho_2$  is a reducible representation, then  $\chi_\rho = \chi_{\rho_1} + \chi_{\rho_2}$ .
- For  $h \in \Gamma$  and  $\Gamma$  of finite order, we will also use

$$(3) \quad \sum_{g \in \Gamma} \chi_s(g^{-1}) \chi_r(gh) = |\Gamma| \frac{\chi_r(h)}{\chi_r(e)} \delta_{r,s}$$



for any two irreducible characters  $\chi_r$  and  $\chi_s$ .

- An important example of a representation is the regular representation, defined by  $R(g) : \ell^2(\Gamma) \rightarrow \ell^2(\Gamma)$  via  $R(g)e_h = e_{gh}$  for all  $g, h \in \Gamma$  and  $\{e_g\}$  the basis for  $\ell^2(\Gamma)$ . Note that  $\ell^2(\Gamma) \cong \mathbb{C}^{|\Gamma|}$  when  $\Gamma$  is of finite order. Then if we let  $\gamma$  be the character corresponding to  $R$ , we find  $\gamma(g) = |\Gamma|\delta_{g,e}$ , since  $R(g)$  is a permutation matrix with zero trace unless  $g = e$ . Further,  $\langle \gamma, \chi \rangle = \frac{1}{|\Gamma|} \sum_{g \in \Gamma} \gamma(g) \overline{\chi(g)} = \frac{1}{|\Gamma|} \gamma(e) \chi(e) = \chi(e)$ , so if  $\{\chi_i\}_{i=1}^k$  is the set of irreducible representations of  $\Gamma$  then

$$(4) \quad \gamma = \sum_{i=1}^k \chi_i(e) \chi_i.$$

Hence, we find  $|\Gamma| = \sum_{i=1}^k \chi_i(e)^2$

The proof of the following theorem is contained within [Ito84]. We include it for completeness.

**Theorem 2.2** ([Ito84]). *Suppose  $G$  is a conjugacy class graph defined using group  $\Gamma$  and conjugacy class (or inverse closed union of conjugacy classes)  $C$ . Further, suppose that  $\chi$  is an irreducible character of  $\Gamma$ . Then  $\chi$  contributes to the spectrum of  $G$  an eigenvalue  $\lambda = |C|\chi(c)/\chi(e)$  (or  $\lambda = \sum_{c \in C} \chi(c)/\chi(e)$ ), where  $c \in C$ , with multiplicity  $\chi(e)^2$ . (Note that distinct characters may add to the multiplicity of a single eigenvalue).*

*Proof.* In this proof we shall assume that  $C$  is a single conjugacy class for ease of notation. If  $C$  is an inverse closed union of conjugacy classes then the proof follows identically but with  $|C|$  replaced by  $\sum_{c \in C}$  throughout.

Let  $V(G) = \{x_1, \dots, x_g\}$  with  $g = |\Gamma|$  as vertices are identified with elements of  $\Gamma$ . Further, let  $\delta_C(z) = 1$  or 0 depending on if  $z \in C$  or not. Then we see that the adjacency matrix of  $G$  is  $A = (\delta_C(x_i x_j^{-1}))_{i,j \leq 1}^g$ .

Since  $\delta_C$  is a class function it is a  $\mathbb{C}$  linear combination of irreducible characters of  $\Gamma$ , i.e.  $\delta_C = \sum_{i=1}^k a_i \chi_i$ , where  $a_i \in \mathbb{C}$  and  $\chi_i$ ,  $1 \leq i \leq k$ , are the irreducible characters. Then orthogonality of irreducible characters using the inner products in Equation (2) gives

$$g \langle \delta_C, \chi_i \rangle = a_i g = \sum_{j=1}^g \delta_C(x_j) \chi_i(x_j^{-1}) = \sum_{c \in C} \chi_i(c^{-1}) = |C| \chi_i(c)$$

for  $c \in C$ . Hence,  $a_i = |C| \chi_i(c) / g$ .

Now, letting  $D_\ell = (\chi_\ell(x_i x_j^{-1}))_{i,j=1}^g$  we get  $A = \frac{|C|}{g} \sum_{\ell=1}^k \chi_\ell(c) D_\ell$ . Additionally, let  $X_{s,1} = (\chi_s(x_1^{-1}), \dots, \chi_s(x_g^{-1}))$  for  $1 \leq s \leq k$ . Then  $X_{s,1} A = \frac{|C|}{g} \sum_{\ell=1}^k \chi_\ell(c) X_{s,1} D_\ell$ .

The relation of group characters given in Equation (3) gives us the  $j$ th component of  $X_{s,1} D_\ell$  equals

$$\sum_{i=1}^g \chi_s(x_i^{-1}) \chi_\ell(x_i x_j^{-1}) = (\chi_s(x_j^{-1}) / \chi_s(e)) g \delta_{s,\ell}.$$

Hence, the  $j$ th component of  $X_{s,1} A$  is  $|C| (\chi_s(c) / \chi_s(e)) \chi_s(x_j^{-1})$ , and we see that  $X_{s,1}$  is an eigenvector of  $A$  with eigenvalue  $|C| \chi_s(c) / \chi_s(e)$ .

We may further notice that  $X_{s,1}$  is the first row vector of  $D_s$ , so let  $X_{s,m}$  be the  $m$ th row vector of  $D_s$ . Similarly, we find these are eigenvectors of  $A$  with eigenvalues  $|C| \chi_s(c) / \chi_s(e)$  as well. Additionally, we have

$$X_{s,l} \cdot X_{t,m} = \sum_{i=1}^g \chi_s(x_l x_i^{-1}) \chi_t(x_i x_m^{-1}) = \sum_{i=1}^g \chi_s(x_i^{-1}) \chi_t(x_i x_l x_m^{-1}) = 0$$

if  $s \neq t$ . Hence, we complete the proof if we can show that  $D_s$  has rank  $\chi_s(e)^2$  for  $1 \leq s \leq k$ .

Let  $R$  be the regular representation of  $\Gamma$ , and  $\gamma$  the character of  $R$ . Using Equation (4) we get  $(\gamma(x_i x_j^{-1})) = \sum_{\ell=1}^k \chi_\ell(e) D_\ell$ . If we let  $\{e_i\}_{i=1}^g$  be the standard basis vectors for  $\mathbb{C}^g$ , then this gives  $ge_i = \sum_{\ell=1}^k \chi_\ell(e) X_{i,\ell}$ . Hence,  $\{X_{s,i}\}_{s,i=1}^{s=k,i=g}$  generates  $\mathbb{C}^g$ , and since  $g = \sum_{\ell=1}^k \chi_\ell(e)^2$  it suffices to show that the rank of  $D_\ell$  does not exceed  $\chi_\ell(e)^2$ .

Let  $R_\ell(x) = (a_{rs}^\ell(x))^{\chi_\ell(e)}$  for  $x \in \Gamma$  be an irreducible representation of  $\Gamma$  corresponding to the character  $\chi_\ell$ . Further, let

$$A_i^{(\ell)} = \begin{pmatrix} a_{i1}^\ell(x_1) & \cdots & a_{i\chi_\ell(e)}^\ell(x_1) \\ \vdots & & \vdots \\ a_{i1}^\ell(x_g) & \vdots & a_{i\chi_\ell(e)}^\ell(x_g) \end{pmatrix} \begin{pmatrix} a_{1i}^\ell(x_1^{-1}) & \cdots & a_{1i}^\ell(x_g^{-1}) \\ \vdots & & \vdots \\ a_{\chi_\ell(e)i}^\ell(x_1^{-1}) & \vdots & a_{\chi_\ell(e)i}^\ell(x_g^{-1}) \end{pmatrix}$$

for  $1 \leq i \leq \chi_\ell(e)$ . We have  $\chi_\ell(x_j x_i^{-1}) = \sum_{r,t=1}^{\chi_\ell(e)} a_{rt}^\ell(x_i) a_{ir}^\ell(x_j^{-1})$ . Hence, we find  $D_\ell = A_1^{(\ell)} + \cdots + A_{\chi_\ell(e)}^{(\ell)}$ . Since the rank of  $A_i^{(\ell)}$  cannot exceed  $\chi_\ell(e)$  we get the rank of  $D_\ell$  cannot exceed  $\chi_\ell(e)^2$ .  $\square$

Applying the above theorem to the Hadamard graph, we obtain the following result. Note that this result is contained in [Ito85a] Proposition 5. However, we restrict our attention to only the largest and smallest eigenvalues, as only these are needed to apply the Hoffman bound. The proof of the following is also found in [Ito85a]. We include a sketch of the relevant portions of the proof for completeness.

**Theorem 2.3** ([Ito85a], Part of Proposition 5). *Let  $A$  be the adjacency matrix for the Hadamard graph on  $2^N$  vertices where  $N$  is a multiple of 4. Then the largest eigenvalue of  $A$  is  $\lambda_{max} = \binom{N}{N/2}$  and the smallest eigenvalue is  $\lambda_{min} = -\frac{\lambda_{max}}{N-1}$ .*

*Proof.* As was shown in Section 1.2, we may view the Hadamard graph as a conjugacy class graph for an abelian group  $\Gamma$  and picking the union of conjugacy classes to be the set of vertices adjacent to the all 1s vertex  $h$ ,  $D_1(h)$ . Hence, we need only understand the characters of  $\Gamma$  and their values on the  $D_1(h)$  by Theorem 2.2. Further, as  $\Gamma$  is abelian the characters are precisely 1-dimensional representations, and for any character  $\chi$ ,  $\chi(e) = 1$ .

Recall that we denote the vector of all ones except at the  $i$ th and  $j$ th spots by  $w(i, j)$ , and that the set  $\{w(0, 1), \dots, w(0, N-1)\}$  forms a generating set for  $\Gamma$ . Then define the character  $\chi_{0,i}$ ,  $1 \leq i \leq N-1$ , by  $\chi_{0,i}(w(0, i)) = -1$  and  $\chi_{0,i}(w(0, j)) = 1$  for  $i \neq j$ . Then  $\{\chi_{0,1}, \dots, \chi_{0,N-1}\}$  forms a generating system for the character group of  $\Gamma$ .

Let  $\chi_0$  be the identity character. Then  $\chi_0(c) = 1$  for any  $c \in D_1(h)$ , so we get the eigenvalue associated to  $\chi_0$  is  $|D_1(h)| = \binom{N}{N/2}$ . This is the maximal eigenvalue as it is clear from the generating set of characters that no character will evaluate to value larger than 1.

It can be shown that any product of  $r$  distinct characters in the generating set, say  $\chi = \chi_{0,i_1} \cdots \chi_{0,i_r}$ , will contribute 0 to the spectrum of  $A$  if  $r$  is odd. Thus, we now see the case when  $2 \leq r \leq N$  even. Let the product

of  $r$  distinct characters in the generating set be  $\chi = \chi_{0,i_1} \cdots \chi_{0,i_r}$ . This character contributes the eigenvalue

$$\begin{aligned} \lambda(r) &= \sum_{v \in D_1(h)} \chi(v) \\ &= \binom{N}{N/2} - 2 \sum_{i \text{ odd}} \binom{r}{i} \binom{n-r}{(N/2)-i} \\ &= a_{N,r}(N/2) = (-1)^{r/2} \frac{r!(N-r)!}{(N/2)!(r/2)!((N-r)/2)!} \end{aligned}$$

where  $a_{N,r}(N/2)$  is the coefficient of  $x^{N/2}$  in  $(1-x)^r(1+x)^{N-r}$ , and the last equality is a result of K. Nomura as described in [Ito84]. Moreover  $\lambda(r+4)/\lambda(r) = (r+3)(r+1)/(n-r-1)(n-r-3)$ , so a quick calculations shows that picking  $r = N - 2$  gives  $\lambda(r) = -\frac{\lambda(N)}{N-1}$  is the minimum eigenvalue.  $\square$

Then combining these spectral results for Hadamard graphs in Theorem 2.3 with the Hoffman lower bound from [EW19] and [Gan23] gives the main result.

**Corollary 2.4.** *For  $N$  a multiple of 4, let  $H_N$  be the Hadamard graph on  $2^N$  vertices. Then  $\chi_q(H_N) = N$ .*

#### REFERENCES

- [AHKS06] David Avis, Jun Hasegawa, Yosuke Kikuchi, and Yuuya Sasaki, *A quantum protocol to win the graph colouring game on all hadamard graphs*, IEICE Trans. Fundam. Electron. Commun. Comput. Sci. **E89-A** (2006), no. 5, 1378–1381.
- [BCE<sup>+</sup>20] Michael Brannan, Alexandru Chirvasitu, Kari Eifler, Samuel Harris, Vern Paulsen, Xiaoyu Su, and Mateusz Wasilewski, *Bigalois extensions and the graph isomorphism game*, Comm. Math. Phys. **375** (2020), no. 3, 1777–1809. MR 4091496
- [BCT99] Gilles Brassard, Richard Cleve, and Alain Tapp, *Cost of exactly simulating quantum entanglement with classical communication*, Phys. Rev. Lett. **83** (1999), 1874–1877.
- [BGH22] Michael Brannan, Priyanga Ganesan, and Samuel J. Harris, *The quantum-to-classical graph homomorphism game*, J. Math. Phys. **63** (2022), no. 11, Paper No. 112204, 34. MR 4505911
- [CM24] Ada Chan and William J. Martin, *Quantum isomorphism of graphs from association schemes*, Journal of Combinatorial Theory, Series B **164** (2024), 340–363.
- [CMN<sup>+</sup>07] Peter J. Cameron, Ashley Montanaro, Michael W. Newman, Simone Severini, and Andreas Winter, *On the quantum chromatic number of a graph*, Electron. J. Combin. **14** (2007), no. 1, Research Paper 81, 15. MR 2365980
- [DSW13] Runyao Duan, Simone Severini, and Andreas Winter, *Zero-error communication via quantum channels, noncommutative graphs, and a quantum Lovász number*, IEEE Trans. Inform. Theory **59** (2013), no. 2, 1164–1174. MR 3015725
- [EKS98] J. A. Erdos, A. Katavolos, and V. S. Shulman, *Rank one subspaces of bimodules over maximal abelian selfadjoint algebras*, J. Funct. Anal. **157** (1998), no. 2, 554–587. MR 1638277
- [EW19] Clive Elphick and Pawel Wocjan, *Spectral lower bounds for the quantum chromatic number of a graph*, Journal of Combinatorial Theory, Series A **168** (2019), 338–347.
- [FR87] Peter Frankl and Vojtěch Rödl, *Forbidden intersections*, Trans. Amer. Math. Soc. **300** (1987), no. 1, 259–286. MR 871675
- [Gan23] Priyanga Ganesan, *Spectral bounds for the quantum chromatic number of quantum graphs*, 2023, pp. 351–376. MR 4603835
- [GTW13] Viktor Galliard, Alain Tapp, and Stefan Wolf, *Deterministic quantum non-locality and graph colorings*, Theoretical Computer Science **486** (2013), 20–26, Theory of Quantum Communication Complexity and Non-locality.
- [Har24] Samuel Harris, *Universality of graph homomorphism games and the quantum coloring problem*, Annales Henri Poincaré (2024).

- [Ito84] Noboru Ito, *The spectrum of a conjugacy class graph of a finite group*, Math. J. Okayama Univ. **26** (1984), 1–10. MR 779767
- [Ito85a] ———, *Hadamard graphs. I*, Graphs Combin. **1** (1985), no. 1, 57–64. MR 796183
- [Ito85b] ———, *Hadamard graphs. II*, Graphs Combin. **1** (1985), no. 4, 331–337. MR 951024
- [Ji13] Zhengfeng Ji, *Binary constraint system games and locally commutative reductions*, ArXiv [abs/1310.3794](https://arxiv.org/abs/1310.3794) (2013).
- [JNV<sup>+</sup>22] Zhengfeng Ji, Anand Natarajan, Thomas Vidick, John Wright, and Henry Yuen, *Mip\*=re*, 2022.
- [MRV18] Benjamin Musto, David Reutter, and Dominic Verdon, *A compositional approach to quantum functions*, J. Math. Phys. **59** (2018), no. 8, 081706, 42. MR 3849575
- [OP16] Carlos M. Ortiz and Vern I. Paulsen, *Quantum graph homomorphisms via operator systems*, Linear Algebra Appl. **497** (2016), 23–43. MR 3466632
- [PSS<sup>+</sup>16] Vern I. Paulsen, Simone Severini, Daniel Stahlke, Ivan G. Todorov, and Andreas Winter, *Estimating quantum chromatic numbers*, J. Funct. Anal. **270** (2016), no. 6, 2188–2222. MR 3460238
- [Wea21] Nik Weaver, *Quantum graphs as quantum relations*, J. Geom. Anal. **31** (2021), no. 9, 9090–9112. MR 4302212
- [WED20] Pawel Wocjan, Clive Elphick, and Parisa Darbari, *Spectral lower bounds for the quantum chromatic number of a graph—Part II*, Electron. J. Combin. **27** (2020), no. 4, Paper No. 4.47, 11. MR 4245222

PERIMETER INSTITUTE FOR THEORETICAL PHYSICS, 51 CAROLINE STREET, WATERLOO, ONTARIO, CANADA

PURDUE UNIVERSITY, MATHEMATICAL SCIENCES BLDG, 150 N UNIVERSITY ST, WEST LAFAYETTE, IN 47907

*Email address:* [amcnamara@perimeterinstitute.ca](mailto:amcnamara@perimeterinstitute.ca)