# Improved Sample Complexity of Imitation Learning for Barrier Model Predictive Control\*

Daniel Pfrommer<sup>†</sup> Swati Padmanabhan<sup>‡</sup> Kwangjun Ahn<sup>§</sup> Jack Umenberger<sup>¶</sup>

Tobia Marcucci

Zakaria Mhammedi<sup>\*\*</sup> Al

Ali Jadbabaie<sup>††</sup>

October 2, 2024

#### Abstract

Recent work in imitation learning has shown that having an expert controller that is both suitably smooth and stable enables stronger guarantees on the performance of the learned controller. However, constructing such smoothed expert controllers for arbitrary systems remains challenging, especially in the presence of input and state constraints. As our primary contribution, we show how such a smoothed expert can be designed for a general class of systems using a log-barrier-based relaxation of a standard Model Predictive Control (MPC) optimization problem.

Improving upon our previous work, we show that barrier MPC achieves theoretically optimal error-to-smoothness tradeoff along some direction. At the core of this theoretical guarantee on smoothness is an improved lower bound we prove on the optimality gap of the analytic center associated with a convex Lipschitz function, which we believe could be of independent interest. We validate our theoretical findings via experiments, demonstrating the merits of our smoothing approach over randomized smoothing.

<sup>\*</sup>The first two authors contributed equally. This work extends our previous result in [38], which has been accepted for publication in CDC 2024. An earlier version of this manuscript was submitted as part of DP's Master's thesis [37]. <sup>†</sup>Massachusetts Institute of Technology. Email: dpfrom@mit.edu.

<sup>&</sup>lt;sup>†</sup>Massachusetts Institute of Technology. Email: pswt@mit.edu.

<sup>&</sup>lt;sup>§</sup>Microsoft Research. Email: kwangjunahn@microsoft.com.

<sup>&</sup>lt;sup>¶</sup>University of Oxford. Email: jack.umenberger@eng.ox.ac.uk.

Massachusetts Institute of Technology. Email: tobiam@mit.edu.

<sup>\*\*</sup>Massachusetts Institute of Technology. Email: mhammedi@mit.edu.

<sup>&</sup>lt;sup>††</sup>Massachusetts Insitute of Technology. Email: jadbabai@mit.edu.

# 1 Introduction

Imitation learning has emerged as a powerful tool in machine learning, enabling agents to learn complex behaviors by imitating expert demonstrations acquired either from a human demonstrator or a policy computed offline [3, 11, 12, 13]. Despite its significant success, imitation learning often suffers from a compounding error problem: Successive evaluations of the approximate policy could accumulate error, resulting in out-of-distribution failures [3]. Recent results in imitation learning [31, 32, 34] have identified *smoothness* (i.e., Lipschitzness of the derivative of the optimal controller with respect to the initial state) and *stability* of the expert as two key properties that circumvent this issue, thereby allowing for end-to-end performance guarantees for the final learned controller.

In this paper, our focus is on enabling such guarantees when the expert being imitated is a Model Predictive Controller (MPC), a powerful class of control algorithms based on solving an optimization problem over a receding prediction horizon [23]. In some cases, the solution to this multiparametric optimization problem, known as the explicit MPC representation [6], can be pre-computed. For instance, in our setup — linear systems with polytopic constraints — the optimal control input is a piecewise affine (and, hence, highly non-smooth) function of the state [6]. However, the number of these pieces may grow exponentially with the time horizon and the state and input dimension, which makes pre-computing and storing such a representation impractical in high dimensions.

While the approximation of a linear MPC controller has garnered significant attention [24, 27, 33], these prior works are primarily concerned with approximating the non-smooth explicit MPC using a neural network and then introducing schemes for enforcing the stability of the learned policy. In contrast, in our paper, we first construct a smoothed version of the expert and then apply theoretical results derived from the imitation of a smoothed expert.

In particular, we demonstrate — both theoretically and empirically — that a log-barrier formulation of the underlying MPC optimization yields smoothness properties similar to its randomizedsmoothing-based counterpart, while being faster to compute. Similar to prior works [9, 18, 19], our barrier MPC formulation replaces the constraints in the MPC optimization problem by a log-barrier in the objective (cf. Section 4). We show that, in conjunction with a black-box imitation learning algorithm, this provides end-to-end guarantees on the performance of the learned policy.

**Our Contributions.** It is known from classical optimization theory [15] that any smooth approximation of a nonsmooth function which is  $O(\epsilon)$  close everywhere must have a smoothness constant (Lipschitzness of the gradient) at least  $O(1/\epsilon)$ . The well-known randomized smoothing technique [16] (convolution with a smoothing kernel) is optimal in this sense; however, it does not preserve the stability properties of the underlying controller and hence is not well-suited for controls applications.

Our main result is that log-barrier-based MPC [9] is an optimal smoother along some direction and outperforms randomized smoothing for controls tasks. More formally, for a given MPC, letting  $u^*$  be the solution of the explicit MPC and  $u^{\eta}$  be the solution of the barrier-MPC formulation, with  $\eta$  being the weight on the barrier, our main contributions for barrier MPC are as follows.

We provide in Theorem 4.8 an upper bound of  $O(\frac{1}{\sqrt{\eta+d^2}-d})$  on the spectral norm of the Hessian of  $u^{\eta}$  with respect to  $x_0$ , where d is the distance of the unconstrained solution from the polytope boundary under the appropriate metric. Separately, we show that there exists a direction a (independent of  $\eta$ ) along which the error  $a^{\top}(u^{\eta}-u^*)$  is at most  $O(\sqrt{\eta+d^2}-d)$ . These two results together show that the controller smoothness and the error match along this direction a, from which we infer that barrier MPC is an optimal smoother along this direction. Along the way, we show (Lemma 4.6) that the Jacobian of the log-barrier solution can be written as a convex combination of the Jacobian of the explicit MPC. In particular, this shows that the rate of change

of  $u^{\eta}$  with respect to  $x_0$  is bounded independent of the weight  $\eta$  applied to the log barrier. We also show (Theorem 4.3) that overall, the distance of  $u^{\eta}$  from  $u^{\star}$  is bounded by  $O(\sqrt{\eta})$ . Finally, we demonstrate through numerical experiments that barrier MPC outperforms randomized smoothing, thus empirically affirming the merits of controls-aware smoothing techniques.

A crucial technical component in obtaining the aforementioned bound on the controller smoothness is a *lower* bound on the distance of  $u^{\eta}$  from the boundary of the polytope (equivalently, a lower bound on the optimality gap of the analytic center associated with a convex Lipschitz function). Intuitively, the nature of the self-concordant barrier already suggests that the solution to a problem with such a barrier in the objective cannot be too close to the boundary of the constraint set. However, obtaining the desired upper bound on the controller smoothness requires an *explicit quantification* of this distance. We provide (Theorem B.13) such a bound for general convex Lipschitz functions via a novel reduction to the setting of linear objectives and then invoking a result by Zong, Lee, and Yue [35]. Furthermore, our smoothing analysis demonstrates that our lower bound is tight up to constants. We believe this result could be broadly useful to the optimization community.

## 2 Problem Setup and Background

We first state our notation and setup that we use throughout. The notation  $\|\cdot\|$  refers to the  $\ell_2$  norm  $\|\cdot\|_2$  for vectors and, by extension, to the spectral norm (largest singular value) for square matrices. For a positive definite matrix H, we denote the local inner product  $\|x\|_H = \sqrt{x^\top H x}$ . Unless transposed, all vectors are column vectors. We use uppercase letters for matrices and lowercase letters for vectors. For a vector x, we use Diag(x) for the diagonal matrix with the entries of x along its diagonal. We use [n] for the set  $\{1, 2, \ldots, n\}$ . Given a matrix  $M \in \mathbb{R}^{n \times n}$  and  $\sigma \in \{0, 1\}^n$ , we denote by  $[M]_{\sigma}$  the principal submatrix of M corresponding to the rows and columns i for which  $\sigma_i = 1$ . We use  $M_{\sigma}^{-1}$  to denote the matrix obtained by first computing the inverse of the matrix  $[M]_{\sigma}$  and then appropriately padding it with zeroes so that the resulting matrix  $M_{\sigma}^{-1}$  has the same size as M. Similarly, we define  $\operatorname{adj}(M)_{\sigma}$  to be the matrix obtained by first computing the adjugate (the transpose of the cofactor matrix) of  $[M]_{\sigma}$  and then appropriately padding it with zeroes so that adj $(M)_{\sigma}$  has the same size as M. Lastly,  $\mathcal{O}(\cdot)$  denotes expressions where numerical constants have been suppressed.

We use the same setup as in our previous work [38] and consider constrained discrete-time linear dynamical systems of the form,

$$x_{t+1} = Ax_t + Bu_t, \quad x_t \in \mathcal{X}_t, u_t \in \mathcal{U}_t, \tag{2.1}$$

with state  $x_t \in \mathcal{X}_t \subseteq \mathbb{R}^{d_x}$  and control-input  $u_t \in \mathcal{U}_t \subseteq \mathbb{R}^{d_u}$  indexed by time step t, and state and input maps  $A \in \mathbb{R}^{d_x \times d_x}$ ,  $B \in \mathbb{R}^{d_x \times d_u}$ . The sets  $\mathcal{X}_t$  and  $\mathcal{U}_t$ , respectively, are the compact convex state and input constraint sets described by the polytopes

$$\mathcal{X}_t := \{ x \in \mathbb{R}^{d_x} \mid A_{x_t} x \le b_{x_t} \}, \quad \mathcal{U}_t := \{ u \in \mathbb{R}^{d_u} \mid A_{u_t} u \le b_{u_t} \},$$

where  $A_{x_t} \in \mathbb{R}^{k_x \times d_x}$ ,  $A_{u_t} \in \mathbb{R}^{k_u \times d_u}$ ,  $b_{x_t} \in \mathbb{R}^{k_x}$ , and  $b_{u_t} \in \mathbb{R}^{k_u}$ . We use  $A_x \in \mathbb{R}^{(T \cdot k_x) \times d_x}$ ,  $A_u \in \mathbb{R}^{(T \cdot k_u) \times d_u}$ ,  $b_x \in \mathbb{R}^{T \cdot k_x}$ , and  $b_u \in \mathbb{R}^{T \cdot k_u}$  to denote the vertically stacked constraints for the full sequences  $x_{1:T}$  and  $u_{0:T-1}$ . A constraint  $f(x) \leq 0$  is said to be "active" at y if f(y) = 0. Given a polytope  $Ax \leq b$ , we say that the quantity  $b_i - a_i^{\top} x$  is its "*i*<sup>th</sup> residual". For notational convenience, we overload  $\phi$  to compactly denote the vector of constraint residuals for a state x and input u as well as for the sequences  $x_{1:T}$  and  $u_{0:T-1}$ :

$$\phi_t(x_t, u_{t-1}) := \begin{bmatrix} b_{x_t} - A_{x_t} x_t \\ b_{u_{t-1}} - A_{u_{t-1}} u_{t-1} \end{bmatrix}, \quad \phi(x_0, u_{0:T-1}) := \begin{bmatrix} \phi_1(x_1, u_0) \\ \vdots \\ \phi_T(x_T, u_{T-1}) \end{bmatrix}.$$
(2.2)

We consider deterministic state-feedback control policies of the form  $\pi : \mathcal{X} \to \mathcal{U}$  and denote the closed-loop system under  $\pi$  by  $f_{cl}^{\pi}(x) := Ax + B\pi(x)$ . We use  $\pi^*$  to refer to the expert policy and  $\hat{\pi}$  to refer to its learned approximation.

In particular, our principal choice of  $\pi^*$  in this paper is an MPC with quadratic cost and linear constraints. The MPC policy is obtained by solving the following minimization problem over future actions  $u := u_{0:T-1}$  with quadratic cost in u and states  $x := x_{1:T}$ :

$$\begin{array}{ll} \text{minimize}_{u} & V(x_{0}, u) := \sum_{t=1}^{T} x_{t}^{\top} Q_{t} x_{t} + \sum_{t=0}^{T-1} u_{t}^{\top} R_{t} u_{t} \\ \text{subject to} & x_{t+1} := A x_{t} + B u_{t}, \\ & x_{T} \in \mathcal{X}, u_{0} \in \mathcal{U}, \\ & x_{t} \in \mathcal{X}, u_{t} \in \mathcal{U}, \ \forall t \in [T-1], \end{array}$$

$$(2.3)$$

where  $Q_t$  and  $R_{t-1}$  are positive definite for all  $t \in [T]$ . For a given state x, the corresponding input  $\pi_{mpc}$  of the MPC is:

$$\boldsymbol{\pi}_{\mathrm{mpc}}(x) := \arg\min_{u_0} \min_{u_{1:T-1}} V(x, u_{0:T-1}), \tag{2.4}$$

where the minimization is over the feasible set defined in Problem 2.3. For  $\pi_{mpc}$  to be well-defined, we assume that  $V(x_0, u)$  has a unique global minimum in u for all feasible  $x_0$ .

### 2.1 Explicit Solution to MPC

As first noted by Bemporad, Morari, Dua, and Pistikopoulos [6], explicit MPC rewrites Problem 2.4 as a multi-parametric quadratic program with linear inequality constraints and solves it for every possible combination of active constraints, building an analytical solution to the control problem. Following this known derivation (see [6, Section 4] and [23, Chapter 11]), we rewrite Problem 2.4 as the optimization problem, in variable  $u := u_{0:T-1} \in \mathbb{R}^{Td_u}$ , as described below:

$$\begin{array}{ll} \text{minimize}_u & \mathcal{V}(x_0, u) := \frac{1}{2} u^\top H u - x_0^\top F u \\ \text{subject to} & G u \leq w + P x_0, \end{array}$$

$$(2.5)$$

with cost matrices  $H \in \mathbb{R}^{T \cdot d_u \times T \cdot d_u}$  and  $F \in \mathbb{R}^{d_x \times T \cdot d_u}$  and m constraints captured via  $G \in \mathbb{R}^{m \times T \cdot d_u}$ , and  $P \in \mathbb{R}^{m \times d_x}$ , and vector  $w \in \mathbb{R}^m$ , all given by

$$H = R_{0:T-1} + \widehat{B}^{\top} Q_{1:T} \widehat{B}, \quad F = -2\widehat{A}^{\top} Q_{1:T} \widehat{B},$$
$$G = \begin{bmatrix} A_u \\ A_x \widehat{B} \end{bmatrix}, \quad P = \begin{bmatrix} 0 \\ -A_x \widehat{A} \end{bmatrix}, \quad w = \begin{bmatrix} b_u \\ b_x \end{bmatrix},$$

where  $Q_{1:T}$  and  $R_{0:T-1}$  are block diagonal with  $Q_1, \ldots, Q_T$  and  $R_0, \ldots, R_{T-1}$  on the diagonal, and  $\hat{B}$  and  $\hat{A}$  are,

$$\hat{A} = \begin{bmatrix} A \\ A^2 \\ \vdots \\ A^T \end{bmatrix}, \quad \hat{B} = \begin{bmatrix} B & 0 & \dots & 0 \\ AB & B & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ A^{T-1}B & A^{T-2}B & \dots & B \end{bmatrix}$$

so that  $x_{1:T} = \widehat{A}x_0 + \widehat{B}u$ .

**Assumption 2.1.** We assume that the constraint polytope in Problem 2.5 contains a ball of radius r and is contained inside an origin-centered ball of radius R.



Figure 1: The explicit MPC controller for  $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ ,  $B = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ , Q = I, R = 0.01, T = 10 with the constraints  $||x||_{\infty} \le 10$ ,  $|u| \le 1$ . For this simple 2-dimensional system there are 261  $K_{\sigma}$ . This figure appeared in our previous work [38].

We now state the solution to Problem 2.5 and later (in Lemma 4.6) show how it appears in the smoothness of the *barrier* MPC solution.

**Lemma 2.2** (Bemporad, Morari, Dua, and Pistikopoulos [6, Theorem 2]). Given a feasible initial state x, let  $\sigma(x) \in \{0,1\}^m$  denote the indicator of active constraints of the optimizer of Problem 2.5, with  $\sigma_i(x) = 1$  iff the ith constraint is active. For  $\sigma \in \{0,1\}^m$ , let  $P_{\sigma} = \{x | \sigma(x) = \sigma\}$  be the set of initial states x for which the solution has active constraints determined by  $\sigma$ . Then for  $x_0 \in P_{\sigma}$ , the solution u of Problem 2.5 is expressed as  $u = K_{\sigma}x_0 + k_{\sigma}$ , where  $K_{\sigma}$  and  $k_{\sigma}$  are defined as:

$$K_{\sigma} := H^{-1}[F^{\top} - G^{\top}(GH^{-1}G^{\top})_{\sigma}^{-1}(GH^{-1}F^{\top} - P)],$$
  

$$k_{\sigma} := H^{-1}G^{\top}(GH^{-1}G^{\top})_{\sigma}^{-1}w.$$
(2.6)

Using this result, one may pre-compute an efficient lookup structure mapping  $x \in P_{\sigma}$  to  $K_{\sigma}, k_{\sigma}$ . However, since every combination of active constraints may yield a potentially unique feedback law, the number of pieces to be computed may grow *exponentially* in the problem dimension or time horizon. For instance, even the simple two-dimensional toy system visualized in Figure 1 has 261 pieces. As a result, it might be computationally intractable to even merely enumerate or store all pieces of the explicit MPC in high dimensions or over long time horizons.

This observation motivates us to consider a learning-based approach. In the spirit of imitation learning discussed in Section 1, we approximate explicit MPC using a polynomial number of sample trajectories, collected offline. We introduce this framework in the next section.

# 3 Motivating Smoothness: Imitation Learning Frameworks

In this section, we instantiate the imitation learning framework to motivate our approach of barrier MPC. We use the Taylor series based imitation learning framework introduced by Pfrommer, Zhang, Tu, and Matni [31], which gives high-probability guarantees on the quality of an approximation.

### 3.1 Taylor Series Imitation Learning

We first introduce the setting for imitation learning. Suppose we are given an expert controller  $\pi^*$ , a policy class  $\Pi$ , a distribution of initial conditions  $\mathcal{D}$ , and N sample trajectories  $\{x_{0:K-1}^{(i)}\}_{i=1}^N$  of length K, with  $\{x_0^{(i)}\}_{i=1}^N$  sampled i.i.d from  $\mathcal{D}$ . As formalized in Fact 3.6, our goal is to find an approximate policy  $\widehat{\pi} \in \Pi$  such that given a suitably small accuracy parameter  $\epsilon$ , the closed-loop states  $\widehat{x}_t$  and  $x_t^*$  induced by  $\widehat{\pi}$  and  $\pi^*$ , respectively, satisfy, with high probability over  $x_0 \sim \mathcal{D}$ ,

$$\|\widehat{x}_t - x_t^\star\| \le \epsilon, \forall t > 0.$$

To understand the sufficient conditions for such a guarantee, we now introduce a few definitions. We first assume through Assumption 3.1 that  $\hat{\pi}$  has been chosen by a black-box supervised imitation learning algorithm which, given the input data, produces a  $\hat{\pi} \in \Pi$  such that, with high probability over the distribution induced by  $\mathcal{D}$ , the policy and its Jacobian are close to the expert.

**Assumption 3.1.** For some  $\delta \in (0,1), \epsilon_0 > 0, \epsilon_1 > 0$  and given N trajectories  $\{x_{0:K-1}^{(i)}\}_{i=1}^{(N)}$  of length K sampled i.i.d. from  $\mathcal{D}$  and rolled out under  $\pi^*$ , the approximating policy  $\hat{\pi}$  satisfies:

$$\mathbb{P}_{x_0 \sim \mathcal{D}} \bigg[ \sup_{k \ge 0} \|\widehat{\pi}(x_k) - \pi^{\star}(x_k)\| \le \epsilon_0 / N \quad \land \quad \sup_{k \ge 0} \left\| \frac{\partial \widehat{\pi}}{\partial x}(x_k) - \frac{\partial \pi^{\star}}{\partial x}(x_k) \right\| \le \epsilon_1 / N \bigg] \ge 1 - \delta.$$

As shown in [31], an example in which Assumption 3.1 holds is when  $\hat{\pi}$  is chosen as an empirical risk minimizer from a class of twice differentiable parametric functions with  $\ell_2$ -bounded parameters, e.g., dense neural networks with smooth activation functions and trained with  $\ell_2$  weight regularization. We refer the reader to [31, 32] for other valid examples of  $\Pi$ . Further, note that the above definition requires generalization on only the state distribution induced by the expert, rather than on the distribution induced by the learned policy, as is the case in [24, 33].

Next, we define a weaker variant of the standard *incremental input-to-state stability* ( $\delta$ ISS) [29] and assume, in Assumption 3.3, that this property holds for the expert policy.

**Definition 3.2** (Local Input-to-State Stability with Linear Gain, cf. [31]). For all initial conditions  $x_0 \in \mathcal{X}$  and bounded sequences of input perturbations  $\{\Delta_t\}_{t>0}$  that satisfy  $\|\Delta_t\| < \kappa$ , let  $\overline{x}_{t+1} = f_{cl}^{\pi}(\overline{x}_t, 0), \overline{x}_0 = x_0$  be the nominal trajectory, and let  $x_{t+1} = f_{cl}^{\pi}(x_t, \Delta_t)$  be the perturbed trajectory. We say that the closed-loop dynamics under  $\pi$  is  $(\kappa, \gamma)$ -locally-incrementally input-to-state stable for linear gain  $\gamma$  if  $\kappa, \gamma > 0$  and,

$$\|x_t - \overline{x}_t\| \le \gamma \cdot \max_{k < t} \|\Delta_k\|, \quad \forall t \ge 0.$$

**Assumption 3.3.** The expert policy  $\pi^*$  is  $(\kappa, \gamma)$ -locally incrementally input-to-state stable.

As noted in [31], local incremental input-to-state stability (local  $\delta ISS$ ) is a much weaker criterion than regular incremental input-to-state stability. We will later show in Lemma 4.10 that under mild assumptions even input-to-state stabilizing (ISS) policies (defined in (Definition 4.9)) are locally  $\delta ISS$ . There is considerable prior work (see, e.g., [14, 28]) demonstrating that ISS holds under mild conditions for both the explicit MPC and the barrier-based MPC under consideration in this paper. Putting these facts together then implies local  $\delta ISS$  of barrier MPC. Having established some preliminaries for stability, we now move on to the smoothness property we consider.

**Definition 3.4** ([31]). We say that an MPC policy  $\pi$  is  $(L_0, L_1)$ -smooth if for all  $x, y \in \mathcal{X}$ ,

$$\|\pi(x)-\pi(y)\| \leq L_0 \|x-y\|$$
 and  $\left\|rac{\partial \pi}{\partial x}(x)-rac{\partial \pi}{\partial x}(y)
ight\| \leq L_1 \|x-y\|$ 

**Assumption 3.5.** The expert policy  $\pi^*$  and the learned policy  $\hat{\pi}$  are both  $(L_0, L_1)$ -smooth.

At a high level, assuming smoothness of the expert and the learned policy helps implicitly ensure that the learned policy captures the stability of the expert in a neighborhood around the data distribution. If the expert or learned policy were to be only piecewise smooth (as is the case, e.g., with standard MPC-based solution of LQR), a transition from one piece to another in the expert not replicated by the learned policy could result in unstable closed-loop behavior.

Having stated all the necessary assumptions, we are now ready to state below the main export of this section, given by [31], guaranteeing closeness of the learned and expert policies.

**Fact 3.6** ([31], Corollary A.1). Provided  $\pi^*$ ,  $\hat{\pi}$  are  $(L_0, L_1)$ -smooth,  $(\kappa, \gamma)$ -locally incrementally stable, and  $\hat{\pi}$  satisfies Assumption 3.1 with  $\frac{\epsilon_0}{N} \leq \min\{\frac{1}{16\gamma^2 L_1}, \frac{1}{16\gamma}, \frac{\kappa}{8\gamma}\}$  and  $\frac{\epsilon_1}{N} \leq \frac{1}{4\gamma}$ ,  $\delta > 0$ , then with probability  $1 - \delta$  for  $x_0 \sim \mathcal{D}$ , we have

$$\|\widehat{x}_t - x_t^{\star}\| \le \frac{8\gamma\epsilon_0}{N} \quad \forall t \ge 0.$$

The upshot of this result is that provided the MPC policy  $\pi^*$  is  $(L_0, L_1)$ -smooth, to match the trajectory of  $\pi^*$  with high probability, we need to match the Jacobian and value of  $\pi^*$  on only NK pieces. This is in contrast to prior work such as [27, 26, 24] on approximating explicit MPC, which require sampling new control inputs during training (in a reinforcement learning-like fashion) or post-training verification of the stability properties of the network.

However, as we noted in Section 1, these strong guarantees crucially require a smooth expert controller. We investigate two approaches for smoothing  $\pi_{mpc}$ : randomized smoothing and barrier MPC. Before doing so, we first consider what constitutes an "optimal" smoothing approach in terms of the smallest possible Hessian norm for a given level of approximation error.

#### 3.2 An Overview of Optimal Smoothing

We begin by considering the properties of a general smoothing algorithm. For simplicity, in this section, we consider smoothing functions of the form  $f : \mathbb{R} \to \mathbb{R}$ , although we note that this analysis can easily be extended to  $f : \mathbb{R}^n \to \mathbb{R}^m$  by considering arbitrary paths  $\mathbb{R} \to \mathbb{R}^n$  and projection  $\mathbb{R}^m \to \mathbb{R}$ . This motivates the following definition of a smoothing algorithm.

**Definition 3.7** ( $\epsilon$ -Smoothing Algorithm). Let  $\epsilon > 0$ . An  $\epsilon$ -smoothing algorithm S for a function class  $\mathcal{F}$  is a map  $S : \mathcal{F} \to C^1$  where  $C^1$  is the class of functions  $\mathbb{R} \to \mathbb{R}$  with continuous derivatives. Furthermore S satisfies,

$$\sup_{x} \|\mathcal{S}(f(x)) - f(x)\| \le \epsilon \quad \forall f \in \mathcal{F}.$$

Analogously, we define a general smoothing algorithm which can smooth functions for arbitrary  $\epsilon$ .

**Definition 3.8** (Smoothing Algorithm). A general smoothing algorithm S for a function class  $\mathcal{F}$  is a map  $S : \mathbb{R} \times \mathcal{F} \to C^1$  where  $S(\epsilon, \cdot)$  is an  $\epsilon$ -smoothing algorithm.

It is known that for any  $\epsilon$ -smoothing algorithm S for the class of L-Lipschitz functions (which we denote by  $\mathcal{L}_L$ ), there exists  $f \in \mathcal{L}_L$  such that the derivative of g := S(f) has Lipschitz constant at least  $\mathcal{O}(\frac{1}{\epsilon})$ . For in-depth treatment of the subject under a more general setting, we direct the reader to Kornowski and Shamir [30] and Beck and Teboulle [15]. A simple example of such a function for which this bound holds is the scaled absolute value function  $x \to C|x|$ .

**Lemma 3.9** ([30], Lemma 30). Let S be any  $\epsilon$ -smoothing algorithm  $S : \mathcal{L}_L \to C^1$  for  $L, \epsilon > 0$  and let f(x) := L|x|, g(x) := S(f). Then there exists  $x, y \in \mathbb{R}$  such that,

$$|
abla g(x) - 
abla g(y)| \ge rac{L^2}{9\epsilon} |x - y|.$$

*Proof.* Consider the value of g(x) at  $x = -\frac{3\epsilon}{L}$ , x = 0, and  $x = \frac{3\epsilon}{L}$ . Since S is an  $\epsilon$ -smoothing algorithm and  $f(-\frac{3\epsilon}{L}) = f(\frac{3\epsilon}{L}) = 3\epsilon$  and f(0) = 0, we can conclude that  $g(-\frac{3\epsilon}{L}), g(\frac{3\epsilon}{L}) > 2\epsilon, g(0) < \epsilon$ . This implies that  $g(\frac{3\epsilon}{L}) - g(0) \ge \epsilon$  and  $g(0) - g(-\frac{3\epsilon}{L}) \le -\epsilon$ . By the mean value theorem, there exist points  $y \in [-\frac{3\epsilon}{L}, 0]$  and  $x \in [0, \frac{3\epsilon}{L}]$  such that,

$$abla g(y) < -rac{L}{3}, 
abla g(x) > rac{L}{3}.$$

Note that  $|x-y| \leq \frac{6\epsilon}{L}$  or that  $\frac{L^2}{9\epsilon}|x-y| \leq \frac{2L}{3}$ . Therefore, we may complete the proof by noting,

$$|\nabla g(x) - \nabla g(y)| \ge \frac{2L}{3} \ge \frac{L^2}{9\epsilon} |x - y|.$$

The above result suggests an inherent tradeoff between the approximation error  $\epsilon$  and the Lipschitzness of the derivative of the smoothed function. Using intuition from the above result, we now state a more general result for arbitrary piecewise twice differentiable functions where the derivatives at the boundaries of the pieces do not necessarily match.

**Lemma 3.10.** Let  $f : \mathbb{R} \to \mathbb{R}$  be a function with piecewise continuous derivatives. Let  $c \in \mathbb{R}$  be a point such that  $\lim_{x\to c^-} \nabla f(x) = a$  and  $\lim_{x\to c^+} \nabla f(x) = b$  where  $a \neq b$  (i.e. the derivative is discontinuous). Then for sufficiently small  $\epsilon$  and any  $\epsilon$ -smoothing algorithm S, we have that for g := S(f) there exist x, y such that,

$$|\nabla g(x) - \nabla g(y)| \ge \frac{|a-b|^2}{144\epsilon} |x-y|.$$

*Proof.* Without loss of generality, we can shift f so that c = 0, f(0) = 0. Similarly, we can also subtract off  $\frac{(a+b)}{2}x$  from both f and g as well as transform  $f(x) \to f(-x), g(x) \to g(-x)$  such that  $\lim_{x\to 0^-} \nabla f(x) = -\frac{|a-b|}{2}, \lim_{x\to 0^+} \nabla f(x) = \frac{|a-b|}{2}$ . Let  $d := \frac{|a-b|}{2}$ .

Since f has piecewise continuous derivative, by definition there exists some radius  $\delta > 0$  around 0 such that f is differentiable on  $(-\delta, 0)$  and  $(0, \delta)$  and that  $\nabla f(x) \leq -\frac{d}{2}$  for  $x \in (-\delta, 0)$  and  $\nabla f(x) \geq \frac{d}{2}$  for  $x \in (0, \delta)$ . We can therefore lower bound,

$$f(x) \ge -\frac{d}{2}x \quad \forall x \in (-\delta, 0), \qquad f(x) \ge \frac{d}{2}x \quad \forall x \in (0, \delta).$$

Similar to Lemma 3.9, we note that for  $\epsilon \leq \frac{d}{6}\delta$ ,  $f(\frac{6\epsilon}{d})$ ,  $f(\frac{6\epsilon}{d}) > 3\epsilon$ . Since f(0) = 0,  $g(0) \leq \epsilon$  and therefore  $g(0) - g(\frac{6\epsilon}{d}) < -\epsilon$ ,  $g(\frac{6\epsilon}{d}) - g(0) \geq \epsilon$ . Therefore for some  $x, y \in \mathbb{R}$  such that  $|x - y| \leq \frac{12\epsilon}{d}$ , we have that,

$$|\nabla g(x) - \nabla g(y)| \ge \frac{d}{3} \ge \frac{d^2}{36\epsilon} |x - y| = \frac{|a - b|^2}{144\epsilon} |x - y|.$$

The above result suggests that the derivative of an  $\epsilon$ -smoothed function has a Lipschitz constant lower bounded by the square of the "discontinuity" in the derivatives times the inverse of the largest approximation error. If the smoothed function is twice differentiable, this is equivalent to a lower bound on the Hessian.

Guided by the above results, we now state our definition for an "optimal smoothing" algorithm, which is a smoothing algorithm such that the above bound is tight, up to a constant. For simplicity, we define optimal smoothing only for L-Lipschitz functions.



Figure 2: Visualizations of the log-barrier MPC control policy and several trajectories for the same system as Figure 1 and different choices of  $\eta$ . This figure appeared in our previous work [38].

**Definition 3.11.** A smoothing algorithm  $S : \mathcal{R} \times \mathcal{F} \to C^1$  for a function class  $\mathcal{F}$  is worst-case optimal up to a constant if there exists C > 0 such that, for any sufficiently small  $\epsilon > 0$ , L > 0, and L-Lipschitz function  $f \in \mathcal{L}_L \subset \mathcal{F}$ , the following inequality holds with  $g := S(\epsilon, f)$ ,

$$\|\nabla g(x) - \nabla g(y)\| \le C \frac{L^2}{\epsilon} \|x - y\|.$$

Note that, by the above lemmas, an algorithm satisfying Definition 3.11 yields smoothed functions where the bound on the Hessian is at most a constant factor worse than the best possible bound for Lipschitz functions. Since the explicit MPC is always Lipschitz, for our purposes we will simply refer to smoothing algorithms satisfying Definition 3.11 as "optimal smoothers".

In the next two sections, we answer the question of whether an optimal smoothing algorithm can preserve the stability of an explicit MPC controller. We will show that while randomized smoothing is an optimal smoother, there exist systems for which randomized smoothing does not preserve the stability of the system. We will then introduce barrier MPC and prove that barrier MPC is an optimal smoother along a certain direction.

### 3.3 First Approach: Randomized Smoothing

We first consider randomized smoothing (see, e.g., [16]) as a baseline approach for smoothing the expert policy  $\pi^*$ . Here, the imitator is learned with a loss function that randomly samples with noise drawn from a chosen probability distribution in order to smooth the policy, effectively convolving the controller with a smoothing kernel. This approach corresponds to the following controller.

**Definition 3.12** (Randomized Smoothed MPC). Given a control policy  $\pi_{mpc}$  of the form Problem 2.4, a desired zero-mean noise distribution  $\mathcal{P}$ , and a smoothing parameter  $\sigma > 0$ , the randomizedsmoothing based MPC,  $\pi^{rs}$ , is defined as:

$$\boldsymbol{\pi}^{\mathrm{rs}}(x) := \mathbb{E}_{w \sim \mathcal{P}}[\boldsymbol{\pi}_{\mathrm{mpc}}(x + \sigma w)].$$

The distribution  $\mathcal{P}$  in Definition 3.12 is usually chosen such that the following guarantees on error and smoothness hold.

**Fact 3.13** (Duchi, Bartlett, and Wainwright [16, Appendix E]). For control policy  $\pi_{mpc} : \mathcal{X} \to \mathcal{U}$ ,  $\mathcal{X} \subset \mathbb{R}^{d_x}, \mathcal{U} \subset \mathbb{R}^{d_u}$  and  $\mathcal{P} \in \{\text{Unif}(B_{\ell_2}(1)), \text{Unif}(B_{\ell_\infty}(1)), \mathcal{N}(0, I)\}$ , there exist constants  $C_0, C_1 > 0$  that depend on  $d_x$ . Let L be the Lipschitz constant of the given control policy  $\pi_{mpc}$ . Then, for any

smoothing parameter  $\sigma > 0$ , the associated  $\pi^{rs}$  satisfies,

$$\begin{split} \|\boldsymbol{\pi}^{\mathrm{rs}}(x) - \boldsymbol{\pi}_{\mathrm{mpc}}(x)\| &\leq C_0 \sigma & \forall x \in \mathcal{X}, \\ \|\nabla \boldsymbol{\pi}^{\mathrm{rs}}(x) - \nabla \boldsymbol{\pi}^{\mathrm{rs}}(y)\| &\leq \frac{C_1 L^2}{\sigma} \|x - y\| & \forall x, y \in \mathcal{X}. \end{split}$$

This implies that randomized smoothing is an optimal smoother for the given choices of  $\mathcal{P}$ .

However, using randomized smoothing to obtain a smoothed policy has three key disadvantages. First,  $\mathbb{E}_{w \sim \mathcal{P}}[\pi_{mpc}(x + \epsilon w)]$  is evaluated via sampling, which means the expert policy must be continuously re-evaluated during training in order to guarantee convergence to the smoothed policy. Secondly, smoothing in this manner may cause  $\pi^{rs}$  to violate state constraints. Finally, simply smoothing the policy may not preserve the stability of  $\pi_{mpc}$ . The first two stated problems arise due to randomized smoothing *oversmoothing* the underlying controller. Consider the following example.

**Example 3.14.** Consider the system  $f(x_t, u_t) = 2x_t + u_t$  and controller  $\pi^*(x) = \min(\max(-2x, -1), 1)$ . We can see that as  $\sigma \to \infty$  (where  $\sigma$  is the smoothing parameter from Definition 3.12), we have  $\pi^{rs}(x) \to 0$  for all x.

The above example shows that  $\pi^{rs}$  is not stable for high  $\sigma$ . Ideally, we would like to aggressively smooth only the discontinuities that do not affect stability or constraint guarantees. This requires a smoothing technique that is aware of when more aggressive control inputs are being taken in order to more quickly stabilize versus preserve some constraint guarantees. As we shall show, barrier MPC is precisely one such method that also preserves state guarantees. We now define barrier MPC and bound the approximation error and smoothness of the resulting controller.

# 4 Our Approach to Smoothing: Barrier MPC

Having described the guarantees obtained via randomized smoothing, we now consider smoothing via self-concordant barrier functions, a notion introduced by Nesterov and Nemirovskii [4].

**Definition 4.1** (Nesterov and Nemirovskii [4]). A convex, thrice differentiable function  $\phi : \mathcal{Q} \to \mathbb{R}$ is a  $\nu$ -self-concordant barrier on an open convex set  $\mathcal{Q} \subseteq \mathbb{R}^n$  if the following conditions hold.

- (i) For all sequences  $x_i \in \mathcal{Q}$  converging to the boundary of  $\mathcal{Q}$ , we have  $\lim_{i \to \infty} \phi(x_i) \to \infty$ .
- (ii) For all  $x \in Q$  and  $h \in \mathbb{R}^n$ , we have the bound  $|\mathcal{D}^3 f(x)[h, h, h]| \leq 2(\mathcal{D}^2 f(x)[h, h])^{3/2}$ , where  $\mathcal{D}^k f(x)[h_1, \ldots, h_k]$  is the k-th derivative of f at x along directions  $h_1, \ldots, h_k$ ,
- (iii) For all  $x \in Q$ , we have  $\nabla f(x)^{\top} (\nabla^2 f(x))^{-1} \nabla f(x) \leq \nu$ . The parameter  $\nu$  satisfies  $\nu \geq 1$ .

The self-concordance property essentially says that locally, the Hessian does not change too fast — it has therefore proven extremely useful in interior-point methods to design fast algorithms for (constrained) convex programming [1, 2] and has also found use in model-predictive control [9, 18, 19, 20, 21] in order to ensure strict feasibility of the control inputs.

In this paper, we consider barrier MPC as a naturally smooth alternative to randomized smoothing of Problem 2.4. In barrier MPC, the inequality constraints occurring in the optimal control problem are eliminated by incorporating them into the cost function via suitably scaled barrier terms. We work only with the log-barrier, which turns a constraint  $f(x) \ge 0$  into the term  $-\eta \log(f(x))$  in the minimization objective and is the standard choice of barrier on polytopes [4]. Concretely, starting from Problem 2.5, the following is our barrier MPC.

**Problem 4.2** (Barrier MPC). Given an MPC as in Problem 2.5 and weight  $\eta > 0$ , the barrier MPC is defined by minimizing, over the input sequence  $u \in \mathbb{R}^{T \cdot d_u}$ , the cost

$$\mathcal{V}^{\eta}(x_0, u) := \frac{1}{2} u^{\top} H u - x_0^{\top} F u - \eta \left[ \mathbf{1}^{\top} \log(\phi(x_0, u)) - d^{\top} u \right],$$
(4.1)

where  $\phi(x_0, u) = Px_0 + w - Gu \in \mathbb{R}^m$  is the (vector) residual of constraints for  $x_0$  and u, and the vector d is set to  $d := \nabla_u \sum_{i=1}^m \log(\phi_i(0, u))|_{u=0}$ . We denote by  $u^{\eta}(x_0)$  the minimizer of (4.1) for a given  $x_0$  and by  $\pi^{\eta}_{mpc}(x) := \arg\min_{u_0} \min_{u_{1:T-1}} \mathcal{V}^{\eta}(x, u)$  the associated control policy.

Some remarks are in order. First, the choice of d in Problem 4.2 is made so as to ensure that  $\arg \min_{u^{\eta}} \mathcal{V}^{\eta}(0, u^{\eta}) = 0$ , i.e. that  $\pi^{\eta}_{\text{mpc}}$  satisfies  $\pi^{\eta}_{\text{mpc}}(0) = \pi_{\text{mpc}}(0) = 0$ , which is a necessary condition for the controller to be stabilizing at the origin. Further, note that  $||d||^2$  is a constant by construction, a fact that turns out to be useful in Theorem 4.8. Secondly, the technical assumptions about the constraint polytope in Problem 2.5 containing a full-dimensional ball of radius r and being contained inside an origin-centered ball of radius R are both inherited by Problem 4.2.

### 4.1 Error Bound for Barrier MPC

To kick off our analysis of the barrier MPC, we first give the following upper bound on the distance between the optimal solution of Problem 4.2 and that of explicit MPC in Problem 2.5. Our result is based on standard techniques to analyze the sub-optimality gap in interior-point methods and crucially uses the strong convexity of our quadratic cost in (4.1).

**Theorem 4.3.** Suppose that  $u^{\eta}$  and  $u^{\star}$  are, respectively, the optimizers of Problem 4.2 and Problem 2.5. Then we have the following bound in terms of the barrier parameter  $\eta$  in (4.1):

$$\|u^{\eta} - u^{\star}\| \le O(\sqrt{\eta}).$$

*Proof.* In this proof, we use  $\mathcal{K}$  for the constraint polytope of Problem 2.5. First, Lemma B.4 shows that the recentered log-barrier  $\phi_{\mathcal{K}}$  in Problem 4.2 is also a self-concordant barrier with some self-concordance parameter  $\nu$ . Since  $u^{\eta} = \arg \min_{u} q(u) + \eta \phi_{\mathcal{K}}(u)$ , where q is the quadratic cost function of Problem 4.2 and  $\phi_{\mathcal{K}}$  the recentered log-barrier on  $\mathcal{K}$ , we have by first-order optimality:

$$\nabla q(u^{\eta}) = -\eta \nabla \phi_{\mathcal{K}}(u^{\eta}). \tag{4.2}$$

Denote by  $\alpha$  the strong convexity parameter of the cost function in Problem 4.2 and by  $\nu$  the self-concordance parameter of the barrier  $\phi_{\mathcal{K}}$ . Then,

$$\{q(u^{\eta}) - q(u^{\star})\} + \frac{1}{2}\alpha \|u^{\eta} - u^{\star}\|^{2} \le \nabla q(u^{\eta})^{\top}(u^{\eta} - u^{\star}) = \eta \cdot \nabla \phi_{\mathcal{K}}(u^{\eta})^{\top}(u^{\star} - u^{\eta}) \le \eta \nu,$$

where the first step is by  $\alpha$ -strong convexity of q, the second step uses (4.2), and the final step applies Fact B.1 at the points  $u^{\eta}$  and  $u^{\star}$ . Since both  $q(u^{\eta}) - q(u^{\star})$  and  $\frac{1}{2}\alpha ||u^{\eta} - u^{\star}||^2$  are positive, we can bound the latter by  $\eta \nu$ . Finally, note that  $\nu \geq 1$  to finish the proof.

We note that the above bound of  $\mathcal{O}(\sqrt{\eta})$  is tight for arbitrary directions. However, provided that  $u^* \neq K_0 x_0$  (where  $K_0$  is the gain associated with the origin piece of the explicit MPC, i.e. the solution is not in the interior of the constraint set), we can show that there exists a direction (independent of  $\eta$  of the choice of barrier) along which the error scales with  $O(\eta/||u^* - K_0 x_0||)$  for small  $\eta$ . **Lemma 4.4.** Suppose that  $u^{\eta}$  and  $u^{\star}$  are, respectively, the optimizers of Problem 4.2 and Problem 2.5. Consider the case where  $u^{\star} \neq K_0 x_0$ , for  $K_0 = H^{-1} F^{\top}$ , i.e.  $K_0 x_0$  is the solution to the unconstrained problem. Let  $a = H(u^{\star} - K_0 x_0)/||H(u^{\star} - K_0 x_0)||$ . Then we have the following upper and lower bounds in terms of the barrier parameter  $\eta$  in Problem 4.2:

$$a^{\top}(u^{\eta} - u^{\star}) \leq \frac{1}{2\sqrt{\alpha_{1}}} \left( \sqrt{4m\eta + \|u^{\star} - K_{0}x_{0}\|_{H}^{2}} - \|u^{\star} - K_{0}x_{0}\|_{H} \right),$$
$$\frac{\overline{\alpha_{1}}}{\alpha_{2}} \cdot \frac{r}{R} \min\left\{ \frac{1}{\sqrt{m\alpha_{2}}} \left( \sqrt{\eta + \|u^{\star} - K_{0}x_{0}\|_{H}^{2}} - \|u^{\star} - K_{0}x_{0}\|_{H} \right), \sqrt{\frac{\alpha_{1}}{\alpha_{2}}} \cdot \frac{r}{2m + 4\sqrt{m}} \right\} \leq a^{\top}(u^{\eta} - u^{\star}),$$

where m is the number of constraints and  $\alpha_1 I \preceq H \preceq \alpha_2 I$ .

*Proof.* This is a direct application of Theorem B.13, which holds for general  $\nu$ -self-concordant barriers, using  $\nu = m$ , the self-concordant barrier parameter of our log barrier.

Since by Lemma 3.10, the spectral norm of the Hessian of  $u^{\eta}$  is lower-bounded by  $\mathcal{O}(1/\epsilon)$  for  $\mathcal{O}(\epsilon)$  error, the upper and lower error bounds of Lemma 4.4 suggest the tightest-possible upper bound on the Hessian that could be shown (and which would demonstrate barrier MPC is an optimal smoother along the *a* direction) is  $\mathcal{O}\left(1/\left[\sqrt{\eta + \|u^* - K_0 x_0\|^2} - \|u^* - K_0 x_0\|\right]\right)$ . We indeed establish this bound later in Theorem 4.8.

The above bound highlights that barrier MPC smooths in a manner which is shaped by problem constraints, unlike randomized smoothing, which generally smooths isotropically. Namely, note that the direction a is the direction of the gradient of the objective at  $u^*$ , which is a combination active constraint directions. This is indicative of barrier MPC smoothing less aggressively along directions associated with active constraints.

#### 4.2 First-Derivative Bound for the Barrier MPC

To prove our main result (Theorem 4.8) on the spectral norm of the Hessian, we first establish the following technical result bounding the first derivative of  $u^{\eta}$  with respect to  $x_0$ . This result may be of independent interest, since it formulates the Jacobian of the log-barrier smoothed solution as a convex combination of derivatives associated with sets of active constraints from the original MPC problem. Our proof starts with the first-order optimality condition for  $u^{\eta}$  and obtains the desired simplification by applying the Sherman-Morrison-Woodbury identity (Fact A.3).

**Lemma 4.5.** Consider Problem 4.2 with associated cost matrices H and F defined therein. Let  $\Phi :=$ Diag $(\phi(x_0, u^{\eta}(x_0)))$  be the diagonal matrix constructed using the (vector) residual  $\phi(x_0, u^{\eta}(x_0)) =$  $Px_0 + w - Gu \in \mathbb{R}^m$ . Then, the solution  $u^{\eta}$  to the barrier MPC in Problem 4.2 satisfies:

$$\frac{\partial u^{\eta}}{\partial x_0} = H^{-1} [F^{\top} - G^{\top} (GH^{-1}G^{\top} + \eta^{-1}\Phi^2)^{-1} (GH^{-1}F^{\top} - P)]$$

*Proof.* We first state the following first-order optimality condition associated with minimizing (4.1):

$$Hu^{\eta}(x_0) - F^{\top}x_0 + \eta \sum_{i=1}^m \left(\frac{g_i}{\phi_i(x_0, u^{\eta}(x_0))} + d_i\right) = 0$$

Differentiating with respect to  $x_0$  and rearranging yields

$$\frac{\partial u^{\eta}}{\partial x_0} = (H + \eta G^{\top} \Phi^{-2} G)^{-1} (F^{\top} + \eta G^{\top} \Phi^{-2} P).$$

$$\tag{4.3}$$

For the rest of the proof, we introduce the notation  $S = GH^{-1}G^{\top} + \eta^{-1}\Phi^2$ . Then, we have by applying the Sherman-Morrison-Woodbury identity (Fact A.3) to the inverse in (4.3) that

$$(H + \eta G^{\top} \Phi^{-2} G)^{-1} = H^{-1} - H^{-1} G^{\top} S^{-1} G H^{-1},$$

which simplifies our expression in (4.3) to

$$\frac{\partial u^{\eta}}{\partial x_{0}} = (H^{-1} - H^{-1}G^{\top}S^{-1}GH^{-1}) \cdot (F^{\top} + \eta G^{\top}\Phi^{-2}P) 
= H^{-1}F^{\top} - H^{-1}G^{\top}S^{-1}GH^{-1}F^{\top} + \underbrace{(H^{-1} - H^{-1}G^{\top}S^{-1}GH^{-1}) \cdot \eta G^{\top}\Phi^{-2}P}_{\text{Term 1}}.$$
(4.4)

We now show that "Term 1" may be simplified as follows.

$$(H^{-1} - H^{-1}G^{\top}S^{-1}GH^{-1}) \cdot \eta G^{\top}\Phi^{-2}P = H^{-1}G^{\top}S^{-1}P.$$
(4.5)

Once this is done, the claim is finished, since plugging the right-hand side from (4.5) into "Term 1" from (4.4) gives exactly the claimed expression in the statement of the lemma. Therefore, we now prove (4.5). To this end, we observe that by factoring out  $H^{-1}G^{\top}$  from the left and  $\eta \Phi^{-2}P$  from the right, we may re-write the left-hand side in (4.5) as

$$\begin{aligned} (H^{-1} - H^{-1}G^{\top}S^{-1}GH^{-1}) \cdot \eta G^{\top}\Phi^{-2}P &= H^{-1}G^{\top}(I - S^{-1}GH^{-1}G^{\top}) \cdot \eta \Phi^{-2}P \\ &= H^{-1}G^{\top}S^{-1} \cdot (S - GH^{-1}G^{\top}) \cdot \eta \Phi^{-2} \cdot P \\ &= H^{-1}G^{\top}S^{-1}P, \end{aligned}$$

where the last step is by using our definition of S and cancelling  $\eta^{-1}\Phi^2$  with  $\eta\Phi^{-2}$ .

Equipped with Lemma 4.5, we are now ready to state Lemma 4.6, where we connect the rate of change (with respect to the initial state  $x_0$ ) of the solution (2.6) of the constrained MPC and the barrier MPC solution (from Lemma 4.5). Put simply, Lemma 4.6 tells us that the barrier MPC solution implicitly interpolates between a potentially exponential number of affine pieces from the original explicit MPC problem. This important connection helps us get a handle on the smoothness of barrier MPC in Theorem 4.8. The starting point for our proof for this result is the expression for  $\frac{\partial u^{\eta}}{\partial x_0}$  from Lemma 4.5. To simplify this expression so it is  $\eta$ -independent, we crucially use our linear algebraic result (Lemma A.7) on products of the form  $L \cdot \operatorname{adj}(LL^{\top})_{\sigma}$  for which  $\operatorname{det}(LL^{\top})_{\sigma} = 0$ .

**Lemma 4.6.** Consider the setup in Problem 4.2 with associated cost matrices H and F, constraint matrices P and G, and barrier parameter  $\eta$ , all defined therein. We define the following quantities.

- (i) For any  $\boldsymbol{\sigma} \in \{0,1\}^m$ , define the matrix  $K_{\boldsymbol{\sigma}} = H^{-1}[F^{\top} G^{\top}(GH^{-1}G^{\top})_{\boldsymbol{\sigma}}^{-1}(GH^{-1}F^{\top} P)]$ , which, recall, in Lemma 2.2 describes the solution u to the constrained MPC.
- (ii) Recall from Problem 4.2 the residual  $\phi := \phi(x_0, u) = Px_0 + w Gu \in \mathbb{R}^m$ ; here, we denote it by  $\phi$ . For any  $\phi$  and  $\boldsymbol{\sigma} = \{0, 1\}^m$ , define the scaling factor  $h_{\boldsymbol{\sigma}} = \det([GH^{-1}G^{\top}]_{\boldsymbol{\sigma}}) \prod_{i=1}^m (\eta^{-1}\phi_i^2)^{1-\sigma_i}$ .
- (iii) We split the set  $\boldsymbol{\sigma} \in \{0,1\}^m$  into the following two sets:

$$S := \left\{ \boldsymbol{\sigma} \in \{0,1\}^m \mid \det([GH^{-1}G^{\top}]_{\boldsymbol{\sigma}}) > 0 \right\} \text{ and } S^{\complement} = \left\{ \boldsymbol{\sigma} \in \{0,1\}^m \mid \det([GH^{-1}G^{\top}]_{\boldsymbol{\sigma}}) = 0 \right\}$$

Then the rate of evolution, with respect to  $x_0$ , of the solution  $u^{\eta}$  to the barrier MPC (in Lemma 4.5) is connected to the solution of the constrained MPC (in (2.6)) as follows:

$$\frac{\partial u^{\eta}}{\partial x_0} = \frac{1}{\sum_{\sigma \in S} h_{\sigma}} \sum_{\sigma \in S} h_{\sigma} K_{\sigma}.$$

*Proof.* Let  $\Phi := \text{Diag}(\phi)$ . Then from Lemma 4.5, we have the following expression for  $\frac{\partial u^{\eta}}{\partial x_{0}}$ :

$$\frac{\partial u^{\eta}}{\partial x_0} = H^{-1} [F^{\top} - G^{\top} (GH^{-1}G^{\top} + \eta^{-1}\Phi^2)^{-1} (GH^{-1}F^{\top} - P)].$$

We now split  $G^{\top}(GH^{-1}G^{\top} + \eta^{-1}\Phi^2)$  above into the following two components via Lemma A.11.

$$G^{\top}(GH^{-1}G^{\top} + \eta^{-1}\Phi^2)^{-1} = \frac{1}{\sum_{\sigma \in S} h_{\sigma}} \left( \sum_{\sigma \in S} h_{\sigma} \cdot G^{\top}(GH^{-1}G^{\top})_{\sigma}^{-1} + \sum_{\sigma \in S^{\complement}} c_{\sigma} \cdot G^{\top} \operatorname{adj}(GH^{-1}G^{\top})_{\sigma} \right),$$

where  $c_{\sigma} := \prod_{i=1}^{m} (\eta^{-1} \phi_i^2)^{1-\sigma_i}$ . By definition of  $S^{\complement}$ , the second sum in the preceding equation comprises those terms for which  $\det(GH^{-1}G^{\top})_{\sigma} = 0$ . We now invoke Lemma A.10, which states that  $G^{\top} \operatorname{adj}(GH^{-1}G^{\top})_{\sigma} = \mathbf{0}$  for all  $\sigma \in S^{\complement}$ . Consequently, we may express  $\frac{\partial u^{\eta}}{\partial x_0}$  in terms of only the first set of terms in the preceding equation (zeroing out the second set of terms):

$$\frac{\partial u^{\eta}}{\partial x_0} = \frac{1}{\sum_{\sigma} h_{\sigma}} \sum_{\sigma \in S} h_{\sigma} H^{-1} [F^{\top} - G^{\top} (GH^{-1}G^{\top})_{\sigma}^{-1} (GH^{-1}F^{\top} - P)] = \frac{1}{\sum_{\sigma} h_{\sigma}} \sum_{\sigma \in S} h_{\sigma} K_{\sigma}.$$

where we plugged in (2.6) in the final step, thus concluding the proof.

The above theorem immediately implies that  $\left\|\frac{\partial u^{\eta}}{\partial x_{0}}\right\|$  is bounded from above as stated next in Corollary 4.7. We draw attention to the fact that Corollary 4.7 shows that the Lipschitz constant of  $u^{\eta}$  is independent of  $\eta$ , which demonstrates that the log-barrier does not worsen the Lipschitz constant of the controller, rather it changes only the interpolation between the different pieces.

**Corollary 4.7.** In the setting of Lemma 4.6, we have,

$$\left\|\frac{\partial u^{\eta}}{\partial x_0}\right\| \le L := \max_{\sigma \in S} \|K_{\sigma}\|.$$

*Proof.* From Lemma 4.6, we infer that  $\frac{\partial u^{\eta}}{\partial x_0}$  lies in the convex hull of  $\{K_{\sigma}\}_{\sigma \in S}$ , and note that  $|S| < \infty$ .

### 4.3 Main Result: Smoothness Bound for the Barrier MPC

We are now ready to state our main result, which effectively shows that  $u^{\eta}$  (and hence  $\pi_{mpc}^{\eta}$ ) satisfies the conditions of Assumption 3.5. Our proof of Theorem 4.8 starts with Lemma 4.5 and computes another derivative. To get an upper bound on the operator norm of the Hessian so obtained, our proof then crucially hinges on Lemma B.11 and Theorem B.13, which provide explicit lower bounds on residuals when minimizing a quadratic cost plus a self-concordant barrier over a polytope, a result we believe to be of independent interest to the optimization community.

**Theorem 4.8.** Consider the setting of Problem 4.2 with associated cost matrices H and F, constraint matrices P and G, barrier parameter  $\eta$ , number of constraints m, the recentering vector d, and the solution  $u^{\eta}$ , all defined therein. We define the following quantities.

- (i) Denote by L the Lipschitz constant of  $u^{\eta}$  from Corollary 4.7.
- (ii) We split the set  $\boldsymbol{\sigma} \in \{0,1\}^m$  into the following two sets:

$$S := \left\{ \boldsymbol{\sigma} \in \{0,1\}^m \mid \det([GH^{-1}G^{\top}]_{\boldsymbol{\sigma}}) > 0 \right\} \text{ and } S^{\complement} = \left\{ \boldsymbol{\sigma} \in \{0,1\}^m \mid \det([GH^{-1}G^{\top}]_{\boldsymbol{\sigma}}) = 0 \right\}$$

(iii) For  $\boldsymbol{\sigma} \in S$ , define the parameter  $C := \max_{\boldsymbol{\sigma} \in S} \|2H^{-1}G^{\top}(GH^{-1}G^{\top})_{\boldsymbol{\sigma}}^{\dagger}\|$ .

(iv) Denote by r and R the inner and outer radius, respectively, associated with Problem 4.2.

(v) Define the residual lower bound,

$$res_{\ell.b.} = \frac{\lambda_{\min}(H)}{\lambda_{\max}(H)} \cdot \frac{r}{R} \cdot \min\left\{\frac{1}{\sqrt{\nu\lambda_{\min}(H)}} \left(\sqrt{\eta + \|u^{\star} - K_0 x_0\|_H^2} - \|u^{\star} - K_0 x_0\|_H\right), \frac{r}{2\nu + 4\sqrt{\nu}}\right\},$$
(4.6)

with  $\nu = 20(m + R^2 ||d||^2)$  and  $K_0 x_0 := H^{-1} F^{\top} x_0$ , the solution to the unconstrained minimization of the quadratic objective. We denote  $||u||_H := \sqrt{u^{\top} H u}$ .

Then, the Hessian of  $u^{\eta}$  with respect to  $x_0$  is bounded by:

$$\left\|\frac{\partial^2 u^\eta}{\partial x_0^2}\right\| \leq \frac{C}{\operatorname{res}_{\ell.b.}} (\|P\| + \|G\|L)^2.$$

where  $\|\cdot\|$  denotes the spectral norm of the third-order tensor  $\frac{\partial^2 u^{\eta}}{\partial x_0^2}$ .

*Proof.* Recall from Lemma 4.5 the following expression for  $\frac{\partial u^{\eta}}{\partial x_0}$  evaluated at a particular  $x_0$ :

$$\frac{\partial u^{\eta}}{\partial x_0}(x_0) = H^{-1}[F^{\top} - G^{\top}(GH^{-1}G^{\top} + \eta^{-1}\Phi(x_0)^2)^{-1}(GH^{-1}F^{\top} - P)],$$

where  $\Phi(x_0) := \text{Diag}(Px_0 - w + Gu^{\eta}(x_0))$ . Let  $y \in \mathbb{R}^{d_x}$  be an arbitrary unit-norm vector, and define the univariate function

$$M(t) := GH^{-1}G^{\top} + \eta^{-1}\Phi(t)^2,$$

where we overload  $\Phi$  to mean  $\Phi(t) := \text{Diag}(P(x_0 + ty) - w + Gu^{\eta}(x_0 + ty))$ , the residual along the path  $t \mapsto x_0 + ty$ . We therefore have the following expression for  $\frac{\partial u^{\eta}}{\partial x_0}$  evaluated at  $x_0 + ty$ :

$$\frac{\partial u^{\eta}}{\partial x_0}(x_0 + ty) = H^{-1}[F^{\top} - G^{\top}M(t)^{-1}(GH^{-1}F^{\top} - P)].$$

Then by differentiating  $M(t)^{-1}$  and applying the chain rule, we get,

$$\begin{split} \frac{d}{dt} \left( \frac{\partial u^{\eta}}{\partial x_0} (x_0 + ty) \right) &= H^{-1} G^{\top} M(t)^{-1} \frac{dM(t)}{dt} M(t)^{-1} (GH^{-1} F^{\top} - P) \\ &= 2H^{-1} G^{\top} M(t)^{-1} \left( \frac{d\Phi(t)}{dt} \eta^{-1} \Phi(t) \right) M(t)^{-1} (GH^{-1} F^{\top} - P) \\ &= 2H^{-1} G^{\top} M(t)^{-1} \frac{d\Phi(t)}{dt} (\eta GH^{-1} G^{\top} \Phi^{-1}(t) + \Phi(t))^{-1} (GH^{-1} F^{\top} - P) \\ &= 2H^{-1} G^{\top} M(t)^{-1} \frac{d\Phi(t)}{dt} (\eta \Phi(t)^{-1} GH^{-1} G^{\top} \Phi(t)^{-1} + I)^{-1} \Phi(t)^{-1} (GH^{-1} F^{\top} - P), \end{split}$$

where the third and fourth steps factor out  $\Phi(t)$  from the right and left, respectively. We now bound groups of terms of the product on the right-hand side and then finish the bound by submultiplicativity of the spectral norm. First, since M(t) is a sum of a square matrix and a positive diagonal matrix, we may apply Lemma A.11 to express  $G^{\top}M(t)^{-1}$  as follows with appropriate  $h_{\sigma}$  and  $c_{\sigma}$ :

$$G^{\top}M(t)^{-1} = \frac{1}{\sum_{\sigma \in S} h_{\sigma}} \left( \sum_{\sigma \in S} h_{\sigma} G^{\top} (GH^{-1}G^{\top})_{\sigma}^{\dagger} + \sum_{\sigma \in S^{\complement}} c_{\sigma} G^{\top} \operatorname{adj}(GH^{-1}G^{\top})_{\sigma} \right).$$
(4.7)

Now note that for  $\boldsymbol{\sigma} \in S^{\complement}$ , we have  $\det(GH^{-1}G^{\top})_{\boldsymbol{\sigma}} = 0$ . We may then invoke Lemma A.10 to infer that for  $\boldsymbol{\sigma} \in S^{\complement}$ , we have  $G^{\top} \operatorname{adj}(GH^{-1}G^{\top})_{\boldsymbol{\sigma}} = \mathbf{0}$ . As a result, the second term on the right-hand side of (4.7) vanishes, thereby affording us the following simplification:

$$\begin{aligned} \|2H^{-1}G^{\top}M(t)^{-1}\| &= \left\|\frac{1}{\sum_{\sigma\in S}h_{\sigma}}\sum_{\sigma\in S}2h_{\sigma}H^{-1}G^{\top}(GH^{-1}G^{\top})_{\sigma}^{\dagger}\right\|\\ &\leq C:=\max_{\sigma\in S}\|2H^{-1}G^{\top}(GH^{-1}G^{\top})_{\sigma}^{\dagger}\|,\end{aligned}$$

where the second equality follows via Hölder's inequality. Next, from the definition of  $\Phi$ , we have that  $\frac{d\Phi}{dt} = P + G\left(\frac{\partial u^{\eta}}{\partial x_0}(x_0 + ty)y\right)$ . By the triangle inequality, the Lipschitzness L of  $u^{\eta}$  (from Corollary 4.7), and the fact that y is unit norm, we have

$$\left\|\frac{d\Phi}{dt}\right\| \le \|P\| + \|G\| \left\|\frac{\partial u^{\eta}}{\partial x_0}\right\| \le \|P\| + \|G\|L.$$

To bound  $\|(\eta \Phi^{-1}GH^{-1}G^{\top}\Phi^{-1}+I)^{-1}\Phi^{-1}\|$ , we first note that because  $\eta \Phi^{-1}GH^{-1}G^{\top}\Phi^{-1} \succeq \mathbf{0}$ , we have  $(\eta \Phi^{-1}GH^{-1}G^{\top}\Phi^{-1}+I)^{-1} \preceq I$ , which in turn implies that  $\|(\eta \Phi^{-1}GH^{-1}G^{\top}\Phi^{-1}+I)^{-1}\| \leq 1$ . Then, by submultiplicativity of the spectral norm, we have

$$\|(\eta \Phi^{-1} G H^{-1} G^{\top} \Phi^{-1} + I)^{-1} \Phi^{-1}\| \le \|\Phi^{-1}\| \le \frac{1}{\min_{i \in [m]} \phi_i}.$$

We may then plug in the lower bound on  $\min_{i \in [m]} \phi_i$  from Theorem B.13 that uses  $\nu = 20(m + R^2 ||d||^2)$ , the self-concordance parameter (computed via Lemma B.4) of the recentered log-barrier in Problem 4.2. Finally, recognizing  $H^{-1}F^{\top}$  as  $K_{\sigma}$  from (2.6) (with  $\sigma = \mathbf{0}^m$ ) yields

$$||GH^{-1}F^{\top} - P|| = ||GK_0 - P|| \le ||P|| + ||G||L.$$

Combining all the bounds obtained above, we may then finish the proof.

Thus, Theorem 4.8 establishes bounds analogous to those in Fact 3.13 for randomized smoothing, demonstrating that the Jacobian of the smoothed expert policy is sufficiently Lipschitz. Indeed, in this case our result is stronger, showing that the Jacobian is differentiable and the Hessian tensor is bounded. This theoretically validates the core proposition of our paper: the barrier MPC policy in Problem 4.2 is suitably smooth, and therefore the guarantees in Section 3 hold. We now briefly revisit our learning guarantees, applied to specifically to log-barrier MPC.

### 4.4 Learning Guarantees for Barrier MPC

We now revisit the learning guarantees discussed in Section 3, adapted specifically to a log-barrier MPC expert. We begin by considering the stability properties of barrier MPC. Since we are interested in establishing  $\|\hat{x}_t - x_t^*\| \leq \epsilon$ , where  $\hat{x}$  is the state under the learned policy and  $x^*$  is the state under the expert, and since we consider MPC controllers which stabilize to the origin, we can relax our local incremental input-to-state stability requirements to simply input-to-state stability (ISS) with minimal assumptions. Definition 4.9 introduces this weaker input-to-state stability property, and Lemma 4.10 shows that ISS policies are locally  $\delta$ ISS. We then observe that there is considerable prior work showing that ISS holds under minimal assumptions for barrier MPC, meaning Assumption 3.3 is satisfied for barrier MPC.

**Definition 4.9** (Input to State Stability [7]). A system  $x_{t+1} = f(x_t, u_t)$  is input-to-state stable (equivalently, a controller  $\pi$  is input-to-state stabilizing under f for  $x_{t+1} = f(x_t, \pi(x_t) + u_t)$ ) if there exists  $\beta \in \mathcal{KL}$  and  $\gamma \in \mathcal{K}$  (where  $f \in \mathcal{K}$  provided  $f : [0, a] \to \mathbb{R}_{\geq 0}$  for a > 0, f is strictly increasing, f(0) = 0, and  $g \in \mathcal{KL}$  provided  $g : [0, a] \times \mathbb{R}_{\geq 0} \to \mathbb{R}_{\geq 0}$  is class  $\mathcal{K}$  in the first argument for some a > 0 and monotonically decreasing in the second such that  $\lim_{t\to\infty} g(s, t) = 0 \forall s$ ). so that for all initial values  $x_0$  and all admissible inputs  $u_t$ , we have

$$||x_t|| \le \beta(||x_0||, t) + \gamma(||u_t||_{\infty}) \quad \forall t \ge 0.$$

**Lemma 4.10.** Let  $\pi$  be an L-Lipschitz controller which is input-to-state stabilizing for the dynamics  $x_{t+1} = Ax_t + Bu_t$  with gains  $\beta \in \mathcal{KL}, \gamma \in \mathcal{K}$ . Define  $\mathcal{B}^{-1}$  such that  $\beta(||x_0||, \mathcal{B}^{-1}(\epsilon)) \leq \epsilon$  for  $||x_0|| \leq B_x$ . As  $\beta \in \mathcal{KL}$ , we know that  $\mathcal{B}^{-1}$  exists and is monotonically decreasing. Define the gain,

$$v(\epsilon) := \min\left\{\gamma^{-1}(\epsilon/2), \epsilon \cdot (1 + ||A|| + (1 + L)||B||)^{-\mathcal{B}^{-1}(\epsilon/4)}\right\}.$$

Then,  $\pi$  is incrementally input-to-state stabilizing with gain  $\gamma' := v^{-1} \in \mathcal{K}$ , i.e., for  $||u||_{\infty} \leq v(\epsilon)$  we have that  $||x_t - \bar{x}_t|| \leq \epsilon$  where  $x_{t+1} = f(x_t, \pi(x_t) + u_t)$  and  $\bar{x}_{t+1} = f(\bar{x}_t, \pi(x_t))$  with  $\bar{x}_0 = x_0$ . Note that over a horizon of length K, we have

$$\gamma'(\|u\|_{\infty}) \le \max\{2\gamma(\|u\|_{\infty}), (1+\|A\|+(1+L)\|B\|)^{K}\|u\|_{\infty}\}.$$

*Proof.* We first show that  $\gamma' \in \mathcal{K}$ . Since  $\gamma \in \mathcal{K}, \gamma^{-1} \in \mathcal{K}$ . Furthermore, as  $\mathcal{B}^{-1}$  is monotonically decreasing,  $C^{-\mathcal{B}^{-1}(\epsilon)}$  is monotonically non-decreasing in  $\epsilon$  for  $C \geq 1$  and  $\epsilon \cdot C^{-\mathcal{B}^{-1}(\epsilon)} \in \mathcal{K}$ . Since v is the minimum of two class  $\mathcal{K}$  gains, it follows that  $v \in \mathcal{K}$  and therefore  $\gamma' := v^{-1} \in \mathcal{K}$ .

We now prove that  $\pi$  is incrementally input-to-state stabilizing with gain  $\gamma' := v^{-1}$ . Fix any  $\epsilon > 0$ . WTS that for  $||u||_{\infty} \leq v(\epsilon)$ ,  $||x - \bar{x}||_{\infty} \leq \epsilon$ . First, consider  $t \leq \beta^{-1}(\epsilon/4)$ . Note that,

$$||x_{t+1} - \bar{x}_{t+1}|| \le (||A|| + ||B||L)||x_t - \bar{x}_t|| + ||B|| \cdot ||u||_{\infty}.$$

By telescoping we can write,

$$\|x_t - \bar{x}_t\| \le (1 + \|A\| + (1 + L)\|B\|)^t \cdot \|u\|_{\infty} \quad .$$

$$(4.8)$$

We then use that  $t \leq \mathcal{B}^{-1}(\epsilon/4)$  and that  $||u||_{\infty} \leq v(\epsilon) \leq \epsilon(1+||A||+(1+L)||B||)^{-\mathcal{B}^{-1}(\epsilon/4)}$ . Combining with (4.8), we get  $||x_t - \bar{x}_t|| \leq \epsilon$ . We next consider the case  $t \geq \mathcal{B}^{-1}(\epsilon/4)$ . We now finish the proof by using the input-to-state stability of  $\pi$ :

$$\|x_t - \bar{x}_t\| \le \|x_t\| + \|\bar{x}_t\| \le 2\beta(\|x_0\|, t) + \gamma(\|u\|_{\infty}) \le 2\beta(\|x_0\|, \mathcal{B}^{-1}(\epsilon/4)) + \gamma(\gamma^{-1}(\epsilon/2)) \le \frac{\epsilon}{2} + \frac{\epsilon}{2} \le \epsilon$$

For stabilizable systems and proper choices of cost function and constraints, the barrier MPC is ISS [28, 19]. We therefore impose the following assumption on the parameters of the barrier MPC.

Assumption 4.11. The parameters of the barrier MPC controller  $\pi_{mpc}^{\eta}$  in Problem 4.2 are chosen such that the system is input-to-state stabilizing. Consequently, by Lemma 4.10 and Corollary 4.7, it is incrementally input-to-state stabilizing over  $t \leq K$  for some with linear gain function  $\gamma$ .

This shows that  $\pi_{\text{mpc}}^{\eta}$  satisfies the even weaker notion of locally  $\delta$ ISS as required in Assumption 3.3. We now state our end-to-end learning guarantee, an extension of Fact 3.6.



Figure 3: Left: The imitation error  $\max_t ||\hat{x} - x^*||$  for the trained MLP over 5 seeds, as a function of the expert smoothness for both randomized smoothing and log-barrier MPC. Center, Right: The  $L_0$  (gradient norm) and  $L_1$  (hessian norm) smoothness of  $\pi^*$  as a function of the smoothing parameter. This figure appeared in our previous work [38].

**Corollary 4.12.** Let  $\pi_{mpc}^{\eta}$  be a barrier MPC as in Problem 4.2 that satisfies Assumption 4.11 such that it is  $(\kappa, \gamma)$ -locally- $\delta ISS$  for linear gain  $\gamma$  over a horizon length K. Let L be as defined in Corollary 4.7 and overload  $\gamma$  to denote the constant associated with  $\gamma(\cdot)$ . Let m be the number of constraints and r, R be the radii associated with the constraint polytope in Problem 4.2.

Assume that  $||x_0|| \leq B_x$  and let  $\hat{\pi}$  be chosen such that, for some  $\epsilon_0, \epsilon_1 > 0$  and given N sample trajectories of length K under  $\pi_{mpc}^{\eta}$  from an initial condition distribution  $\mathcal{D}$ ,

$$\mathbb{P}_{x_0 \sim \mathcal{D}} \bigg[ \sup_{0 \le k \le K} \|\widehat{\pi}(x_k) - \pi_{\mathrm{mpc}}^{\eta}(x_k)\| \le \epsilon_0 / N \wedge \sup_{0 \le k \le K} \left\| \frac{\partial \widehat{\pi}}{\partial x}(x_k) - \frac{\partial \pi_{\mathrm{mpc}}^{\eta}}{\partial x}(x_k) \right\| \le \epsilon_1 / N \bigg] \ge 1 - \delta.$$

Let  $C, L, \operatorname{res}_{\ell.b.} > 0$  be defined as in Theorem 4.8 and P, G be the matrices associated with the original MPC problem, Problem 2.5. Then, provided that  $N \ge \mathcal{O}\left(\max\left\{\frac{C\gamma^2}{\operatorname{res}_{\ell.b.}}(\|P\| + \|G\|L), \gamma, \frac{\gamma}{\kappa}\right\}\epsilon_0\right)$ , and  $N \ge 4\gamma\epsilon_1$ , it follows that,

$$\|\hat{x}_t - x_t^{\eta}\| \le \frac{8\gamma\epsilon_0}{N} \quad \forall \, 0 \le t \le K.$$

Per Theorem 4.8, res<sub>l.b.</sub> scales with either  $\mathcal{O}(\eta)$  or  $\mathcal{O}(\sqrt{\eta})$ , depending on the direction of  $u^* - u^{\eta}$ .

*Proof.* This is an application of Fact 3.6, combined with our  $L_1$  smoothness bound on the Hessian (Theorem 4.8), where,  $L_1 \leq \frac{C}{\operatorname{res}_{\ell,\mathrm{b.}}}(\|P\| + \|G\|L)$ .

In conclusion, our analysis in this section shows that our key assumption of locally input-to-state stability (Assumption 3.3) of the barrier MPC controller  $\pi_{mpc}^{\eta}$  is satisfied. We also showed smoothness of  $\pi_{mpc}^{\eta}$ . Per the results of Pfrommer, Zhang, Tu, and Matni [31], these two properties together imply the learning guarantees in Corollary 4.12. This concludes our theoretical analysis of imitation learning using the barrier MPC. We now corroborate our theory with numerical experiments.

### 5 Experiments

The experiments presented below first appeared in our previous work [38]. We include these here for completeness.

We demonstrate the advantage of barrier MPC over randomized smoothing for the double integrator system visualized in Figure 1. The matrices describing the dynamics are  $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$  and  $B = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ , and the cost matrices are given by  $Q_t = I$ ,  $R_t = 0.01I$ , with horizon length T = 10. Our

constraints for Problem 4.2 are  $||x||_{\infty} \leq 10$  and  $||u||_{\infty} \leq 1$ . This is the same setup as in [33], which we note asymptotically stabilizes the system to the origin.

We sample  $N \in \{20, 50\}$  trajectories of length K = 20 using  $\pi_{\text{mpc}}^{\eta}$  and  $\pi^{\text{rs}}$  and smoothing parameters  $\eta$  (Problem 4.2) and  $\sigma$  (Definition 3.12) ranging from  $10^{-4}$  to  $10^3$  and  $10^{-4}$  to 20, respectively. We use  $\mathcal{P} = \mathcal{N}(0, I)$  for the randomized smoothing distribution. For each parameter set, we trained a 4-layer multi-layer perceptron (MLP) using GELU activations [22] to ensure smoothness of  $\Pi$ . We used AdamW [25] with a learning rate of  $3 \cdot 10^{-4}$  and weight decay of  $10^{-3}$  in order to ensure boundedness of the weights (see [31]).

We visualize the smoothness properties of the chosen expert  $\pi^*$  of each method (either  $\pi_{\text{mpc}}^{\eta}$ or  $\pi^{\text{rs}}$ ) across the choices of  $\eta, \sigma$  in Figure 3. For small Hessian norms (i.e. the large  $\eta, \sigma$  regime), barrier MPC has larger gradient norm  $\|\nabla \pi^*\|$  than randomized smoothing. This shows that  $\pi_{\text{mpc}}^{\eta}$ prevents oversmoothing in comparison to  $\pi^{\text{rs}}$ . While randomized smoothing reduces  $\|\nabla^2 \pi^*\|$  by essentially flattening the function,  $\pi_{\text{mpc}}^{\eta}$  achieves equally smooth functions while still maintaining control of the system. This effect is also seen in Figure 2, where we visualize the barrier MPC controller for different  $\eta$  and see that, even for large  $\eta$ , we successfully stabilize to the origin.

One interesting phenomenon is that the maximum gradient of  $\pi^{\eta}_{mpc}$  begins decreasing much earlier than  $\pi^{rs}$ . This is due to the fact that  $\pi^{rs}$  smooths only locally, meaning that if the smoothing radius is sufficiently small, the gradient will not be affected. Meanwhile,  $\pi^{\eta}_{mpc}$  always performs a global form of smoothing, so even for small  $\eta$ , the controller is smoothed everywhere.

In Figure 3, we also compare the trajectory error when imitating trajectories from  $\pi^{rs}$ ,  $\pi^{\eta}_{mpc}$  for equivalent levels of smoothness. We can see that for N = 20 and N = 50,  $\pi^{\eta}_{mpc}$  significantly outperforms  $\pi^{rs}$  across all smoothness levels. This effect is particularly pronounced in the very smooth regime, where imitating  $\pi^{rs}$  proves unstable due to the inherit instability of (A, B), leading to extremely large imitation errors. Meanwhile,  $\pi^{\eta}_{mpc}$  is strictly easier to imitate the more smoothing that is applied. Overall, these experiments confirm our hypothesis that not all smoothing techniques perform equally and that barrier MPC is an effective smoothing technique that outperforms randomized smoothing for the purposes of imitation learning.

### 6 Discussion

We consider two methods for smoothing MPC policies for constrained linear systems: randomized smoothing and barrier MPC. While the former is known to have the theoretically optimal ratio of approximation error to Hessian norm, it may not preserve the stability or constraint satisfaction properties of the underlying controller and hence is not always well-suited for controls applications. We show that the log-barrier-based MPC yields a smooth control with optimal error to smoothness ratio along some direction. Additionally, it better ensures constraint satisfaction while also retaining the stability properties of the original policy. We show how these properties enable theoretical guarantees when learning barrier MPC and demonstrate experimentally its better performance compared to a randomized smoothing baseline.

Our key technical contribution towards proving the smoothness of barrier MPC is a lower bound on the optimality gap of the analytic center associated with a convex Lipschitz function, which we hope could be of independent interest to the broader optimization community. Extending our results to smoothing nonlinear MPC policies would be a fruitful direction for future work.

# Acknowledgements

We gratefully acknowledge funding from ONR N00014-23-1-2299 and a Vannevar Bush Fellowship from the Office of Undersecretary of Defense for Research and Engineering (OUSDR&E).

# References

- I. I. Dikin. "Iterative solution of problems of linear and quadratic programming". English. In: Sov. Math., Dokl. (1967) (cited on page 10).
- [2] Narendra Karmarkar. "A new polynomial-time algorithm for linear programming". In: *Proceedings of the sixteenth annual ACM symposium on Theory of computing*. 1984 (cited on page 10).
- [3] Dean A Pomerleau. "Alvinn: An autonomous land vehicle in a neural network". In: Advances in neural information processing systems (1988) (cited on page 2).
- [4] Yurii Nesterov and Arkadii Nemirovskii. Interior-point polynomial algorithms in convex programming. SIAM, 1994 (cited on pages 10, 28).
- [5] James Renegar. A mathematical view of interior-point methods in convex optimization. SIAM, 2001 (cited on pages 33, 34).
- [6] Alberto Bemporad, Manfred Morari, Vivek Dua, and Efstratios N Pistikopoulos. "The explicit linear quadratic regulator for constrained systems". In: *Automatica* (2002) (cited on pages 2, 4, 5).
- [7] Hassan K Khalil. Control of nonlinear systems. Prentice Hall, New York, NY, 2002 (cited on page 17).
- [8] Alan J Laub. Matrix analysis for scientists and engineers. SIAM, 2004 (cited on page 25).
- [9] Adrian G Wills and William P Heath. "Barrier function based model predictive control". In: Automatica (2004) (cited on pages 2, 10).
- [10] Ilse CF Ipsen and Rizwana Rehman. "Perturbation bounds for determinants and characteristic polynomials". In: SIAM Journal on Matrix Analysis and Applications (2008) (cited on page 23).
- [11] Nathan D Ratliff, David Silver, and J Andrew Bagnell. "Learning to search: Functional gradient techniques for imitation learning". In: Autonomous Robots (2009) (cited on page 2).
- [12] Pieter Abbeel, Adam Coates, and Andrew Y Ng. "Autonomous helicopter aerobatics through apprenticeship learning". In: *The International Journal of Robotics Research* (2010) (cited on page 2).
- [13] Stéphane Ross, Geoffrey Gordon, and Drew Bagnell. "A reduction of imitation learning and structured prediction to no-regret online learning". In: *Proceedings of the fourteenth international conference on artificial intelligence and statistics*. JMLR Workshop and Conference Proceedings. 2011 (cited on page 2).
- [14] Majid Zamani and Rupak Majumdar. "A Lyapunov approach in incremental stability". In: 2011 50th IEEE conference on decision and control and European control conference. IEEE. 2011 (cited on page 6).
- [15] Amir Beck and Marc Teboulle. "Smoothing and first order methods: A unified framework". In: SIAM Journal on Optimization 22.2 (2012), pp. 557–580 (cited on pages 2, 7).

- [16] John C Duchi, Peter L Bartlett, and Martin J Wainwright. "Randomized smoothing for stochastic optimization". In: *SIAM Journal on Optimization* (2012) (cited on pages 2, 9).
- [17] Roger A Horn and Charles R Johnson. *Matrix analysis*. Cambridge university press, 2012 (cited on page 23).
- [18] Christian Feller and Christian Ebenbauer. "A barrier function based continuous-time algorithm for linear model predictive control". In: 2013 European Control Conference (ECC). IEEE. 2013, pp. 19–26 (cited on pages 2, 10).
- [19] Christian Feller and Christian Ebenbauer. "Barrier function based linear model predictive control with polytopic terminal sets". In: 53rd IEEE Conference on Decision and Control. IEEE. 2014, pp. 6683–6688 (cited on pages 2, 10, 17).
- [20] Christian Feller and Christian Ebenbauer. "Weight recentered barrier functions and smooth polytopic terminal set formulations for linear model predictive control". In: 2015 American Control Conference (ACC). IEEE. 2015, pp. 1647–1652 (cited on page 10).
- [21] Christian Feller and Christian Ebenbauer. "Relaxed logarithmic barrier function based model predictive control of linear systems". In: *IEEE Transactions on Automatic Control* (2016) (cited on page 10).
- [22] Dan Hendrycks and Kevin Gimpel. "Gaussian error linear units (gelus)". In: arXiv preprint arXiv:1606.08415 (2016) (cited on page 19).
- [23] Francesco Borrelli, Alberto Bemporad, and Manfred Morari. *Predictive control for linear and hybrid systems*. Cambridge University Press, 2017 (cited on pages 2, 4).
- [24] Steven Chen, Kelsey Saulnier, Nikolay Atanasov, Daniel D Lee, Vijay Kumar, George J Pappas, and Manfred Morari. "Approximating explicit model predictive control using constrained neural networks". In: 2018 Annual American control conference (ACC). IEEE. 2018 (cited on pages 2, 6, 7).
- [25] Ilya Loshchilov and Frank Hutter. "Decoupled Weight Decay Regularization". In: International Conference on Learning Representations. 2019. URL: https://openreview.net/forum?id= Bkg6RiCqY7 (cited on page 19).
- [26] Benjamin Karg and Sergio Lucia. "Efficient representation and approximation of model predictive control laws via deep learning". In: *IEEE Transactions on Cybernetics* (2020) (cited on page 7).
- [27] CG da S Moraes. "A neural network architecture to learn explicit MPC controllers from data". In: *IFAC-PapersOnLine* (2020) (cited on pages 2, 7).
- [28] Maxime Pouilly-Cathelain, Philippe Feyel, Gilles Duc, and Guillaume Sandou. "Stability of Barrier Model Predictive Control". In: 17th International Conference on Informatics in Control, Automation and Robotics (ICINCO). 2020 (cited on pages 6, 17).
- [29] Rick Voßwinkel and Klaus Röbenack. "Determining input-to-state and incremental input-tostate stability of nonpolynomial systems". In: International Journal of Robust and Nonlinear Control (2020) (cited on page 6).
- [30] Guy Kornowski and Ohad Shamir. "Oracle complexity in nonsmooth nonconvex optimization". In: Journal of Machine Learning Research 23.314 (2022), pp. 1–44 (cited on page 7).
- [31] Daniel Pfrommer, Thomas Zhang, Stephen Tu, and Nikolai Matni. "Tasil: Taylor series imitation learning". In: Advances in Neural Information Processing Systems (2022) (cited on pages 2, 5–7, 18, 19).

- [32] Stephen Tu, Alexander Robey, Tingnan Zhang, and Nikolai Matni. "On the sample complexity of stability constrained imitation learning". In: *Learning for Dynamics and Control Conference*. PMLR. 2022 (cited on pages 2, 6).
- [33] Kwangjun Ahn, Zakaria Mhammedi, Horia Mania, Zhang-Wei Hong, and Ali Jadbabaie.
   "Model Predictive Control via On-Policy Imitation Learning". In: Learning for Dynamics and Control Conference. PMLR. 2023 (cited on pages 2, 6, 19).
- [34] Adam Block, Ali Jadbabaie, Daniel Pfrommer, Max Simchowitz, and Russ Tedrake. "Provable Guarantees for Generative Behavior Cloning: Bridging Low-Level Stability and High-Level Behavior". In: *Thirty-seventh Conference on Neural Information Processing Systems*. 2023 (cited on page 2).
- [35] MR Zong, YT Lee, and MC Yue. "Short-step Methods Are Not Strongly Polynomial-Time". In: *Mathematical Programming* (2023) (cited on pages 3, 29).
- [36] Mehrdad Ghadiri, Yin Tat Lee, Swati Padmanabhan, William Swartworth, David P Woodruff, and Guanghao Ye. "Improving the Bit Complexity of Communication for Distributed Convex Optimization". In: Proceedings of the 56th Annual ACM Symposium on Theory of Computing. 2024, pp. 1130–1140 (cited on page 28).
- [37] Daniel Pfrommer. "On the Sample Complexity of Imitation Learning for Smoothed Model Predictive Control". PhD thesis. Massachusetts Institute of Technology, 2024 (cited on page 1).
- [38] Daniel Pfrommer, Swati Padmanabhan, Kwangjun Ahn, Jack Umenberger, Tobia Marcucci, Zakaria Mhammedi, and Ali Jadbabaie. On the Sample Complexity of Imitation Learning for Smoothed Model Predictive Control. 2024. arXiv: 2306.01914 [eess.SY]. URL: https: //arxiv.org/abs/2306.01914 (cited on pages 1, 3, 5, 9, 18).

# A Technical Results from Matrix Analysis

We use the notation introduced in Section 2. Additionally, we use  $e_i$  to denote the vector with one at the  $i^{\text{th}}$  coordinate and zeroes at the remaining coordinates. We first collect the following relevant facts from matrix analysis before proving our technical results.

**Fact A.1** ([17]). Given a square matrix  $A \in \mathbb{R}^{n \times n}$ , its  $(i, j)^{\text{th}}$  minor  $M_{i,j}$ , is defined as the determinant of the  $(n-1) \times (n-1)$  matrix resulting from deleting row *i* and column *j* of *A*. Next, the  $(i, j)^{\text{th}}$  cofactor is defined to be the  $(i, j)^{\text{th}}$  minor scaled by  $(-1)^{i+j}$ :

$$C_{ij} = (-1)^{i+j} M_{i,j}.$$
 (A.1)

We then define the cofactor matrix  $C \in \mathbb{R}^{n \times n}$  of A as the matrix of cofactors of all entries of A, *i.e.*,  $C = ((-1)^{i+j}M_{i,j})_{1 \le i,j \le n}$ . The adjugate of A is the transpose of the cofactor matrix C of A, and hence its  $(i, j)^{\text{th}}$  entry may be expressed as:

$$adj(A)_{ij} = (-1)^{i+j} M_{j,i}.$$
 (A.2)

In particular, if the matrix A is symmetric, then the  $(i, j)^{\text{th}}$  minor equals the  $(j, i)^{\text{th}}$  minor, implying

$$\operatorname{adj}(A)_{ij} = \operatorname{adj}(A)_{ji} \text{ for all } i, j \in [n].$$
 (A.3)

The minors of a matrix are also useful in computing its determinant. Specifically, the Laplace expansion of a matrix A along its column j is given as:

$$\det(A) = \sum_{i=1}^{n} (-1)^{i+j} a_{ij} M_{i,j}.$$
(A.4)

Finally, the adjugate adj(A) also satisfies the following important property:

$$\operatorname{adj}(A) \cdot A = A \cdot \operatorname{adj}(A) = \det(A) \cdot I.$$
 (A.5)

Fact A.2 (Matrix determinant lemma, [17]). For any M, the determinant for a unit-rank update may be expressed as:

$$\det(M + uv^{\top}) = \det(M) + v^{\top} \operatorname{adj}(M)u.$$

Fact A.3 (Sherman-Morrison-Woodbury identity). Given conformable matrices A, C, U, and V such that A and C are invertible, we have

$$(A + UCV)^{-1} = A^{-1} - A^{-1}U(C^{-1} + VA^{-1}U)^{-1}VA^{-1}.$$

We crucially use the following expansion for determinants of perturbed matrices.

**Fact A.4** (Ipsen and Rehman [10, Theorem 2.3]). Given  $A \in \mathbb{R}^{m \times m}$  as in Fact A.2, positive diagonal matrix  $\Lambda = \text{Diag}(\lambda) \in \mathbb{R}^{m \times m}$ , and  $A_{\sigma}$  denoting the principal submatrix formed by selecting A's rows and columns indexed by  $\sigma \in \{0,1\}^m$ , we have

$$\det(A + \Lambda) = \sum_{\sigma \in \{0,1\}^n} \left( \prod_{i=1}^m \lambda_i^{1-\sigma_i} \right) \det(A_{\sigma})$$

We now state and prove a technical result that we build upon to prove Lemma A.11, which we in turn use in the proof of Lemma 4.6.

**Lemma A.5.** Consider a matrix  $A = \begin{bmatrix} a & b^{\top} \\ b & D \end{bmatrix} \in \mathbb{R}^{n \times n}$ , where  $D \in \mathbb{R}^{(n-1) \times (n-1)}$  is a symmetric matrix. Then the adjugate  $\operatorname{adj}(A)$  may be expressed as follows:

$$\mathrm{adj}(A) = \begin{bmatrix} \det(D) & -b^{\top} \operatorname{adj}(D) \\ -\operatorname{adj}(D)b & a \cdot \operatorname{adj}(D) + K \end{bmatrix},$$

for some matrix K independent of a.

*Proof.* Let  $\tilde{D}_{ij}$  be the  $(n-2) \times (n-2)$  matrix obtained by deleting the  $i^{\text{th}}$  row and  $j^{\text{th}}$  column of D. Let  $\tilde{D}_j$  be the  $(n-1) \times (n-2)$  matrix formed by removing the  $j^{\text{th}}$  column of D, and let  $\tilde{b}_i \in \mathbb{R}^{(n-2)}$  be the vector obtained by deleting the  $i^{\text{th}}$  coordinate of b. With this notation in hand, we now compute some relevant cofactors.

First, observe that D is the matrix obtained by deleting the first row and column of A, and hence this fact along with (A.1) yields the (1,1)<sup>th</sup> cofactor:

$$C_{1,1} = \det(D).$$

Second, for some j > 0, observe that the matrix obtained by deleting the first row of A and the  $(1+j)^{\text{th}}$  column of A is exactly the horizontal concatenation of b and  $\tilde{D}_j$ . Applying this observation in (A.1) then gives the following expression for the  $(1, 1+j)^{\text{th}}$  cofactor:

$$C_{1,1+j} = (-1)^{j} \det\left(\begin{bmatrix} b & \widetilde{D}_{j}\end{bmatrix}\right) = (-1)^{j} \sum_{i=1}^{n-1} b_{i}(-1)^{i+1} \det(\widetilde{D}_{ij}) = -\sum_{i=1}^{n-1} b_{i} \operatorname{adj}(D)_{ij} = -\begin{bmatrix} b^{\top} \operatorname{adj}(D) \end{bmatrix}_{j},$$

where the second step is by using (A.4) to expand det  $\left(\begin{bmatrix} b & \tilde{D}_j \end{bmatrix}\right)$  along the column vector b, and the third step is by (A.2) and (A.3), which applies since D is assumed symmetric. Finally, to compute the  $(1 + i, 1 + j)^{\text{th}}$  cofactor, we first construct the matrix obtained by deleting the  $(1 + i)^{\text{th}}$  row and  $(1 + j)^{\text{th}}$  column of A. Based on the notation we introduced above, this may be expressed as  $\begin{bmatrix} a & \tilde{b}_j^{\mathsf{T}} \\ \tilde{b}_i & \tilde{D}_{ij} \end{bmatrix}$ , from which we have by (A.1):

$$C_{1+i,1+j} = (-1)^{i+j} \det \left( \begin{bmatrix} a & \widetilde{b}_j^\top \\ \widetilde{b}_i & \widetilde{D}_{ij} \end{bmatrix} \right),$$
(A.6)

which we now simplify. To this end, we observe that

$$\det\left(\begin{bmatrix}a & \widetilde{b}_{j}^{\top}\\ \widetilde{b}_{j} & \widetilde{D}_{ij}\end{bmatrix}\right) = \det\left(\begin{bmatrix}a & \widetilde{b}_{j}^{\top}\\ 0 & \widetilde{D}_{ij}\end{bmatrix} + \begin{bmatrix}0\\ \widetilde{b}_{j}\end{bmatrix} \cdot e_{1}^{\top}\right) = \det\left(\begin{bmatrix}a & \widetilde{b}_{j}^{\top}\\ 0 & \widetilde{D}_{ij}\end{bmatrix}\right) + e_{1}^{\top}\operatorname{adj}\left(\begin{bmatrix}a & \widetilde{b}_{j}^{\top}\\ 0 & \widetilde{D}_{ij}\end{bmatrix}\right) \begin{bmatrix}0\\ \widetilde{b}_{j}\end{bmatrix},$$
(A.7)

where we used Fact A.2 in the second step. The first term in the right-hand side of the preceding equation may be simplified to  $a \cdot \det(\tilde{D}_{ij})$ . To simplify the second term, we first observe that we wish to compute only the first row of  $\operatorname{adj} \begin{pmatrix} \begin{bmatrix} a & \tilde{b}_j^\top \\ 0 & \tilde{D}_{ij} \end{bmatrix} \end{pmatrix}$ ; to this end, we introduce the notation that  $X := \tilde{D}_{ij}$  and  $y = \tilde{b}_j$ ; we denote  $\tilde{X}_{\ell j}$  to be the matrix obtained by deleting the  $\ell^{\text{th}}$  row and  $j^{\text{th}}$ column of X; we use  $\tilde{X}_{\ell}$  for the matrix obtained by deleting the  $\ell^{\text{th}}$  row of X. Now observe that for  $\ell > 0$ , the  $(1 + \ell)^{\text{th}}$  entry of the desired first row may be computed as follows:

$$\operatorname{adj}\left(\begin{bmatrix}a & \widetilde{b}_{j}^{\top}\\ 0 & \widetilde{D}_{ij}\end{bmatrix}\right)_{1,1+\ell} = (-1)^{\ell} \operatorname{det}\begin{bmatrix}y^{\top}\\\widetilde{X}_{\ell}\end{bmatrix} = \sum_{j=1}^{n-1} y_{j} \cdot (-1)^{\ell+j} \operatorname{det}\widetilde{X}_{\ell j} = \sum_{j=1}^{n-1} y_{j} \cdot (\operatorname{adj}\widetilde{X})_{j\ell} = \widetilde{b}_{j}^{\top} \operatorname{adj}(\widetilde{D}_{ij})e_{\ell},$$

where the first step is by expressing the  $(1, 1 + \ell)^{\text{th}}$  entry of the adjugate in question in terms of its  $(1 + \ell, 1)^{\text{th}}$  minor (as per (A.2)), the second step is by the Laplace expansion of the determinant along its first row (analogous to (A.4)), the third step is by (A.2), and the final step plugs back the newly introduced notation. Hence, we have

$$e_1^{\top} \operatorname{adj} \left( \begin{bmatrix} a & \widetilde{b}_j^{\top} \\ 0 & \widetilde{D}_{ij} \end{bmatrix} \right) = \begin{bmatrix} a \cdot \det(\widetilde{D}_{ij}) & \widetilde{b}_j^{\top} \operatorname{adj}(\widetilde{D}_{ij}) \end{bmatrix}.$$
(A.8)

Multiplying the right-hand side of (A.8) by  $\begin{bmatrix} 0\\ \tilde{b}_j \end{bmatrix}$  and plugging the result back into (A.7) and eventually into (A.6) then gives

$$C_{1+i,1+j} = a \cdot \operatorname{adj}(D)_{ij} + (-1)^{i+j} \widetilde{b}_j^{\top} \operatorname{adj}(\widetilde{D}_{ij}) \widetilde{b}_j.$$

By mapping these cofactors back into the definition of the adjugate we want, one can then conclude the proof, with K collecting all the  $(-1)^{i+j}\tilde{b}_i^{\top} \operatorname{adj}(M_{ij})\tilde{b}_j$  terms.

**Corollary A.6.** Let 
$$A = \begin{bmatrix} a & b^{\top} \\ b & D \end{bmatrix}$$
 be a symmetric matrix. Then,  
 $\operatorname{adj}(A + \lambda e_1 e_1^{\top}) = \operatorname{adj}(A) + \lambda \begin{bmatrix} 0 & \mathbf{0}^{\top} \\ \mathbf{0} & \operatorname{adj}(D) \end{bmatrix}$ 

*Proof.* First, observe that by applying Lemma A.5, we have

$$\operatorname{adj}\left(\begin{bmatrix}a+\lambda & b^{\mathsf{T}}\\b & D\end{bmatrix}\right) = \begin{bmatrix}\operatorname{det}(D) & -b^{\mathsf{T}}\operatorname{adj}(D)\\-\operatorname{adj}(D)b & (a+\lambda)\cdot\operatorname{adj}(D)+K\end{bmatrix}.$$
(A.9)

Next, observe that based on the definition of A, the left-hand side of (A.9) is precisely  $\operatorname{adj}(A + \lambda e_1 e_1^{\top})$ ; based on the expression for  $\operatorname{adj}(A)$  from Lemma A.5, the right-hand side of (A.9) may be split into  $\operatorname{adj}(A) + \lambda \begin{bmatrix} 0 & 0 \\ 0 & \operatorname{adj}(D) \end{bmatrix}$ , as desired. This concludes the proof.

**Lemma A.7.** Consider a matrix  $L \in \mathbb{R}^{m \times n}$ , and define the matrix  $A = LL^{\top} \in \mathbb{R}^{m \times m}$ . Suppose that det(A) = 0. Then, the following equation holds:

$$\operatorname{adj}(A)L = \mathbf{0}.$$

To prove Lemma A.7, we use the following two technical results from matrix analysis.

**Fact A.8** (Theorem 4.18, [8]). Suppose  $A \in \mathbb{R}^{n \times p}$  and  $L \in \mathbb{R}^{n \times m}$ . Then  $AA^{\dagger}L = L$  if and only if the range spaces  $\mathcal{R}(L)$  and  $\mathcal{R}(A)$  satisfy the inclusion  $\mathcal{R}(L) \subseteq \mathcal{R}(A)$ .

**Fact A.9** (Theorem 3.21, [8]). Let  $L \in \mathbb{R}^{m \times n}$ . Then the range spaces  $\mathcal{R}(L)$  and  $\mathcal{R}(LL^{\top})$  satisfy the property  $\mathcal{R}(L) = \mathcal{R}(LL^{\top})$ .

*Proof of Lemma A.7.* We prove the claim by showing that

$$\operatorname{adj}(A)L = [\operatorname{adj}(A)A]A^{\dagger}L = \mathbf{0},$$

where the last equality follows from the property that  $\operatorname{adj}(A)A = \det(A)I = 0$ . All that remains is to prove that  $L = AA^{\dagger}L$ . By Fact A.8, this is true if and only if  $\mathcal{R}(L) \subseteq \mathcal{R}(A)$ . From Fact A.9, we know that this is true. This concludes the proof.

**Lemma A.10.** Given a binary vector  $\boldsymbol{\sigma} \in \{0,1\}^m$ , matrix  $G \in \mathbb{R}^{m \times n}$  and matrix  $H \in \mathbb{R}^{n \times n}$  with the properties  $H \succ 0$  and  $\det(GH^{-1}G^{\top})_{\boldsymbol{\sigma}} = 0$ , we have

$$G^{\top} \operatorname{adj}(GH^{-1}G^{\top})_{\sigma} = 0.$$

*Proof.* Without loss of generality, let  $G = \begin{bmatrix} G_1 \\ G_2 \end{bmatrix}$  where  $\sigma_i = 1$  for the rows and columns associated with  $G_2$ . Then we may express  $GH^{-1}G^{\top}$  in terms of  $G_1$  and  $G_2$  as follows:

$$GH^{-1}G^{\top} = \begin{bmatrix} G_1 H^{-1} G_1^{\top} & G_1 H^{-1} G_2^{\top} \\ G_2 H^{-1} G_1^{\top} & G_2 H^{-1} G_2^{\top} \end{bmatrix}.$$

Based on the expansion above, observe that  $(GH^{-1}G^{\top})_{\sigma} = G_2H^{-1}G_2^{\top}$ . As a result, we may express  $G^{\top} \operatorname{adj}(GH^{-1}G^{\top})_{\sigma}$ , our matrix product of interest, as follows:

$$G^{\top} \operatorname{adj}(GH^{-1}G^{\top})_{\boldsymbol{\sigma}} = \begin{bmatrix} G_1^{\top} & G_2^{\top} \end{bmatrix} \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \operatorname{adj}(G_2H^{-1}G_2^{\top}) \end{bmatrix} = \begin{bmatrix} \mathbf{0} & G_2^{\top} \operatorname{adj}(G_2H^{-1}G_2^{\top}) \end{bmatrix}.$$

All that remains is to show that  $G_2^{\top} \operatorname{adj}(G_2 H^{-1} G_2^{\top}) = \mathbf{0}$ . To this end, we note that

$$G_2^{ op} \operatorname{adj}(G_2 H^{-1} G_2^{ op}) = H^{1/2} \left[ H^{-1/2} G_2^{ op} \operatorname{adj}(G_2 H^{-1/2} H^{-1/2} G_2^{ op}) \right] = \mathbf{0}$$

where the last equality follows immeditely by applying Lemma A.7.

**Lemma A.11.** For a positive semi-definite matrix  $A \in \mathbb{R}^{m \times m}$ , a diagonal matrix  $\Lambda = \text{Diag}(\lambda)$ , and the indicator vector  $\boldsymbol{\sigma} \in \{0,1\}^m$ , define the following parameters:

- (i) Scaling factor  $h_{\sigma} = \det(A_{\sigma}) \prod_{i=1}^{m} \lambda_i^{1-\sigma_i}$ ,
- (ii) Normalizing factor  $h = \sum_{\sigma \in \{0,1\}^m} h_{\sigma}$ ,
- (iii) Adjugate scaling factor  $c_{\sigma} = \prod_{i=1}^{m} \lambda_i^{1-\sigma_i}$ .
- (iv) We split the set  $\boldsymbol{\sigma} \in \{0,1\}^m$  into the following two sets:

$$S := \left\{ \boldsymbol{\sigma} \in \{0,1\}^m \mid \det([GH^{-1}G^{\top}]_{\boldsymbol{\sigma}}) > 0 \right\} \text{ and } S^{\complement} = \left\{ \boldsymbol{\sigma} \in \{0,1\}^m \mid \det([GH^{-1}G^{\top}]_{\boldsymbol{\sigma}}) = 0 \right\}$$

Assume that  $A + \Lambda$  is invertible. Then we have the following decomposition of  $(A + \Lambda)^{-1}$  in terms of inverses and adjugates of  $A_{\sigma}$  (the principal submatrices of A), with the adjugate or inverse computed on the basis of whether or not det $(A_{\sigma}) = 0$ , as follows:

$$(A+\Lambda)^{-1} = \sum_{\sigma \in S} \frac{h_{\sigma}}{h} A_{\sigma}^{-1} + \sum_{\sigma \in S^{\complement}} \frac{c_{\sigma}}{h} \operatorname{adj}(A)_{\sigma}.$$
 (A.10)

*Proof.* We begin by proving the following simpler statement for  $c_{\sigma} = \prod_{i=1}^{m} \lambda_i^{1-\sigma_i}$ :

$$\operatorname{adj}(A + \Lambda) = \sum_{\sigma \in \{0,1\}^m} c_{\sigma} \operatorname{adj}(A)_{\sigma}.$$
(A.11)

Once this statement is proven, (A.10) is implied by the following argument: Per Fact A.4, we have that  $\det(A + \Lambda) = \sum_{\sigma \in \{0,1\}^m} h_{\sigma} = h$ , so dividing throughout by h yields  $(A + \Lambda)^{-1}$  on the left-hand side (by (A.5)); the term  $\sum_{\sigma \in \{0,1\}^m} \frac{c_{\sigma}}{h} \operatorname{adj}(A)_{\sigma}$  may be split into two sums of terms, one over those

vectors  $\boldsymbol{\sigma} \in \{0,1\}^m$  for which  $\det(A_{\boldsymbol{\sigma}}) = 0$  and the second over those choices of  $\boldsymbol{\sigma} \in \{0,1\}^m$  for which  $\det(A_{\boldsymbol{\sigma}}) \neq 0$ . For terms such that  $\det(A_{\boldsymbol{\sigma}}) \neq 0$ , we have

$$c_{\sigma} \operatorname{adj}(A)_{\sigma} = c_{\sigma} \operatorname{det}([A]_{\sigma}) \frac{1}{\operatorname{det}([A]_{\sigma})} \operatorname{adj}(A)_{\sigma} = h_{\sigma} \cdot (A)_{\sigma}^{-1}.$$

Hence, (A.11), when divided by h, gives (A.10), as desired. We now prove (A.11), proceeding via induction on nnz( $\Lambda$ ), the number of nonzero entries in  $\Lambda$ .

**Base case:** When the number of non-zero entries  $nnz(\Lambda) = 0$ , by definition,  $\Lambda = Diag(\mathbf{0})$ , which implies that the left-hand side of (A.11) is adj(A). Further, since by definition,  $c_{\sigma} = \prod_{i=1}^{m} \lambda_i^{1-\sigma_i}$ , for our choice of  $\Lambda = Diag(\mathbf{0})$ , this gives the following expression:

$$c_{\boldsymbol{\sigma}} = \begin{cases} 0 & \text{if } \boldsymbol{\sigma} \neq 1, \\ 1 & \text{if } \boldsymbol{\sigma} = 1. \end{cases}$$

With this choice of  $c_{\sigma}$ , the right-hand side of (A.11) reduces to  $\operatorname{adj}(A)$ , which matches the left-hand side of (A.11), thus implying that in this base case, (A.11) is true.

**Induction Step:** Suppose (A.11) holds for  $\operatorname{nnz}(\Lambda) = k$ . We now show that (A.11) holds for  $\operatorname{nnz}(\Lambda) = k + 1$  as well with some scaling factor  $c_{\sigma}$ . Without loss of generality, assume that  $\lambda_i \neq 0$  for  $i \in [k+1]$ . Let  $\widetilde{A}_{11}$  be the  $\mathbb{R}^{(m-1)\times(m-1)}$  matrix obtained by deleting the first row and first column of A. By expressing  $A + \Lambda$  as  $(A + \sum_{i=2}^{k+1} \lambda_i e_i e_i^{\top}) + \lambda_1 e_1 e_1^{\top}$ , we may use Corollary A.6 to expand  $\operatorname{adj}(A + \Lambda)$  as follows:

$$\operatorname{adj}(A+\Lambda) = \operatorname{adj}\left(A + \sum_{i=2}^{k+1} \lambda_i e_i e_i^{\mathsf{T}}\right) + \lambda_1 \begin{bmatrix} \mathbf{0} & \mathbf{0}^{\mathsf{T}} \\ \mathbf{0} & \operatorname{adj}\left(\widetilde{A}_{11} + \sum_{i=1}^k \lambda_{i+1} \widetilde{\mathbf{e}}_i \widetilde{\mathbf{e}}_i^{\mathsf{T}}\right) \end{bmatrix}, \qquad (A.12)$$

where note that the  $e_i \in \mathbb{R}^m$  and  $\tilde{e}_i \in \mathbb{R}^{m-1}$ . We observe that both the terms on the right-hand side have  $nnz(\Lambda) - 1 = k$  nonzero entries in their respective diagonal components. Hence, by our assumption, the induction hypothesis is applicable; therefore, suppose that by (A.11),

$$\operatorname{adj}(A + \sum_{i=2}^{\ell} \lambda_i e_i e_i^{\top}) = \sum_{\sigma \in \{0,1\}^m} \hat{c}_{\sigma} \operatorname{adj}(A)_{\sigma},$$
  
$$\operatorname{adj}(\widetilde{A}_{11} + \sum_{i=1}^{\ell-1} \lambda_{i+1} \widetilde{e}_i \widetilde{e}_i^{\top}) = \sum_{\sigma' \in \{0,1\}^{m-1}} \tilde{c}_{\sigma'} \operatorname{adj}(\widetilde{A}_{11})_{\sigma'},$$
  
(A.13)

where, based on the diagonal components in each of the terms on the left-hand side, the scaling factors on the respective right-hand sides are  $\hat{c}_{\sigma} = 0^{(1-\sigma_1)} \prod_{i=2}^{m} \lambda_i^{1-\sigma_i}$ , and  $\tilde{c}_{\sigma'} = \prod_{i=1}^{m-1} \lambda_{i+1}^{1-\sigma'_i}$ . As a consequence of these definitions, we can re-write the terms in (A.13) using  $c_{\sigma}$  as follows. First, observe that  $\hat{c}_{\sigma} = 0$  when  $\sigma_1 = 0$  and  $\hat{c}_{\sigma} = c_{\sigma}$  otherwise. This implies:

$$\sum_{\boldsymbol{\sigma} \in \{0,1\}^m} \tilde{c}_{\boldsymbol{\sigma}} \operatorname{adj}(A)_{\boldsymbol{\sigma}} = \sum_{\substack{\boldsymbol{\sigma} \in \{0,1\}^m, \\ \sigma_1 = 1}} c_{\boldsymbol{\sigma}} \operatorname{adj}(A)_{\boldsymbol{\sigma}}.$$
 (A.14)

Next, observe that for the vector  $\boldsymbol{\sigma} = [0; \boldsymbol{\sigma}']$  formed by concatenating zero with  $\boldsymbol{\sigma}'$ , we have  $c_{\boldsymbol{\sigma}} = \lambda_1 \tilde{c}_{\boldsymbol{\sigma}'}$ . This implies the following chain of equations:

$$\lambda_1 \begin{bmatrix} 0 & \mathbf{0}^\top \\ \mathbf{0} & \sum_{\sigma' \in \{0,1\}^{m-1}} \tilde{c}_{\sigma'} \operatorname{adj}(\tilde{A}_{11})_{\sigma'} \end{bmatrix} = \sum_{\sigma' \in \{0,1\}^{m-1}} \lambda_1 \tilde{c}_{\sigma'} \operatorname{adj}(A)_{\left[\begin{smallmatrix} 0\\ \sigma' \end{smallmatrix}\right]} = \sum_{\substack{\sigma \in \{0,1\}^m, \\ \sigma_0 = 0}} c_{\sigma} \operatorname{adj}(A)_{\sigma}, \quad (A.15)$$

where in the first step, we used the fact that  $A_{11}$  is, by definition, the principal submatrix of A obtained by deleting its first row and first column; the second step is by our prior observation connecting  $c_{\sigma}$  and  $\tilde{c}_{\sigma'}$ . Plugging the right-hand sides from (A.13) into that of (A.12) and then applying (A.14) and (A.15) gives

$$\begin{aligned} \operatorname{adj}\left(A+\Lambda\right) &= \sum_{\boldsymbol{\sigma} \in \{0,1\}^m} \hat{c}_{\boldsymbol{\sigma}} \operatorname{adj}(A)_{\boldsymbol{\sigma}} + \lambda_1 \begin{bmatrix} 0 & \mathbf{0}^\top \\ \mathbf{0} & \sum_{\boldsymbol{\sigma}' \in \{0,1\}^{m-1}} \tilde{c}_{\boldsymbol{\sigma}'} \operatorname{adj}(\tilde{A}_{11})_{\boldsymbol{\sigma}'} \end{bmatrix} \\ &= \sum_{\substack{\boldsymbol{\sigma} \in \{0,1\}^m, \\ \boldsymbol{\sigma}_0 = 1}} c_{\boldsymbol{\sigma}} \operatorname{adj}(A)_{\boldsymbol{\sigma}} + \sum_{\substack{\boldsymbol{\sigma} \in \{0,1\}^m, \\ \boldsymbol{\sigma}_0 = 0}} c_{\boldsymbol{\sigma}} \operatorname{adj}(A)_{\boldsymbol{\sigma}} \\ &= \sum_{\substack{\boldsymbol{\sigma} \in \{0,1\}^m}} c_{\boldsymbol{\sigma}} \operatorname{adj}(A)_{\boldsymbol{\sigma}}. \end{aligned}$$

Thus, we have shown (A.11) for  $nnz(\Lambda) = k + 1$ , thereby completing the induction and concluding the proof of (A.11) and, consequently, of the stated lemma.

# **B** Technical Results from Convex Analysis

**Fact B.1** ([4]). Let  $\Phi$  be a  $\nu$ -self-concordant barrier. Then for any  $x \in dom(\Phi)$  and  $y \in cl(dom)(\Phi)$ ,

$$abla \Phi(x)^+(y-x) \le 
u.$$

**Fact B.2** ([4, Proposition 2.3.1]). Let G be a closed convex domain in E, let F be a  $\nu$ -self-concordant barrier for G, and let  $x = \mathcal{A}(y)$  be an affine transformation from a space E' into E such that  $\mathcal{A}(E') \cap \operatorname{int}(G) \neq \emptyset$ . Let  $G' = \mathcal{A}^{-1}(G)$  and  $F'(y) = F(\mathcal{A}(y)) : \operatorname{int}(G') \to \mathbb{R}$ . Then, F' is a  $\nu$ -self-concordant barrier for G'.

To prove Theorem B.7, we use two simple properties of self-concordant barriers Lemma B.3 and Lemma B.4 that originally appeared in [36]. We provide their proofs here for completeness.

**Lemma B.3** ([36]). If f is a self-concordant barrier for a set  $\mathcal{K} \subseteq \mathcal{B}(0, R)$ , then for any  $x \in \mathcal{K}$ , we have

$$abla^2 f(x) \succeq rac{1}{9R^2} I.$$

*Proof.* For the sake of contradiction, suppose  $\nabla^2 f \succeq \frac{1}{9R^2}I$ . This is equivalent to, for some  $x \in \mathcal{K}$  and unit vector u,

$$(3Ru)^{\top} \nabla^2 f(x)(3Ru) < 1.$$
 (B.1)

Define the unit-radius Dikin ellipsoid around x as

$$\mathcal{E}_x(x,1) = \left\{ y : (y-x)^\top \nabla^2 f(x)(y-x) \le 1 \right\}.$$

Then, Inequality (B.1) is equivalent to the assertion that  $x + 3Ru \in \mathcal{E}_x(x, 1)$ . Because f is selfconcordant we have  $\mathcal{E}_x(x, 1) \subseteq \mathcal{K}$  (see, e.g., [4, Theorem 2.1.1]). This, combined with  $x + 3Ru \in \mathcal{E}_x(x, 1)$ , implies  $x + 3Ru \in \mathcal{K}$ . However, since  $\mathcal{K} \subseteq \mathcal{B}(0, R)$  and  $x \in \mathcal{K}$  by construction, the inclusion  $x + 3Ru \in \mathcal{K}$  cannot hold for any unit vector u, which implies that our initial assumption must be false, thus concluding the proof.

**Lemma B.4** ([36]). If f is a  $\nu$ -self-concordant barrier for a given convex set  $\mathcal{K}$  then  $g(x) = c^{\top}x + f(x)$  is a self-concordant barrier over  $\mathcal{K}$ . Further, if  $\mathcal{K} \subseteq \mathcal{B}(0, R)$ , then g has self-concordance parameter at most

$$20(
u + R^2 \|c\|^2).$$

*Proof.* Since  $\nabla^2 g = \nabla^2 f$ , we can conclude that g is also a self-concordant function. Since  $\mathcal{K} \subseteq \mathcal{B}(0, R)$ , Lemma B.3 applies, and we have  $\nabla^2 f(x) \succeq \frac{1}{9R^2}I$  for all  $x \in \mathcal{K}$ . Equivalently,

$$\nabla^2 f(x)^{-1} \preceq 9R^2 I \text{ for all } x \in \mathcal{K}.$$
 (B.2)

The self-concordance parameter (see Definition 4.1) of g is:

$$\|\nabla g(x)\|_{\nabla^2 g(x)^{-1}}^2 = \|c + \nabla f(x)\|_{\nabla^2 f(x)^{-1}}^2 \le 2\|c\|_{\nabla^2 f(x)^{-1}}^2 + 2\|\nabla f(x)\|_{\nabla^2 f(x)^{-1}}^2, \tag{B.3}$$

where the first step is by definition of self-concordance parameter of g. To finish the proof, we recall that  $||c||_{\nabla^2 f(x)^{-1}}^2 \leq 9R^2 ||c||_2^2$  by Inequality (B.2), and  $||\nabla f(x)||_{\nabla^2 f(x)^{-1}}^2 \leq \nu$  by the self-concordance parameter of f and put these bounds into Inequality (B.3).

Both our proofs of our main result lower bounding the residual crucially build upon the following result from [35], who proved the following highly non-trivial lower bound in Inequality (B.5) (the upper bound was known in the classical literature on interior-point methods).

**Fact B.5** ([35], Theorem 2). Fix a vector c, a polytope  $\mathcal{K}$ , and a point v. We assume that the polytope  $\mathcal{K}$  contains a full-dimensional ball of radius r. Let  $v^* = \arg \min_{u \in \mathcal{K}} c^{\top} u$ . We define, for c,

$$gap(v) = c^{\top}(v - v^{\star}). \tag{B.4}$$

Further, define  $v_{\eta} = \arg \min_{v} c^{\top} v + \eta \phi_{\mathcal{K}}(v)$ , where  $\phi_{\mathcal{K}}$  is a self-concordant barrier on  $\mathcal{K}$ . Then we have the following lower bound on this suboptimality gap evaluated at  $v_{\eta}$ :

$$\min\left\{\frac{\eta}{2}, \frac{r\|c\|}{2\nu + 4\sqrt{\nu}}\right\} \le \operatorname{gap}(v_{\eta}) = c^{\top}(v_{\eta} - v^{\star}) \le \eta\nu.$$
(B.5)

Finally, we need the following known claim about the minimizer of a convex combination of two convex functions. We provide its proof for the sake of completeness of this document.

**Claim B.6.** Let  $f : \mathbb{R} \to \mathbb{R}$  and  $g : \mathbb{R} \to \mathbb{R}$  be two convex, continuously differentiable functions with  $x_f$  and  $x_g$  satisfying  $f'(x_f) = 0$  and  $g'(x_g) = 0$ . Additionally, let g be strictly convex. Let  $x_f < x_g$ . Then, for any positive  $\alpha$  and  $\beta$ , we have that the point  $x_{\alpha f+\beta g}$  satisfying  $\alpha f'(x_{\alpha f+\beta g})+\beta g'(x_{\alpha f+\beta g})=0$  additionally satisfies  $x_{\alpha f+\beta g} \in [x_f, x_g]$ .

Proof. Since f and g are both convex, their gradients are monotone (and the gradients of g are strictly monotone). Since  $x_f < x_g$ , this then implies  $0 = f'(x_f) \le f'(x_g)$  and  $g'(x_f) < g'(x_g) = 0$ . Multiplying the first inequality by  $\alpha$  and the second by  $\beta$  and summing them gives  $\alpha f'(x_f) + \beta g'(x_f) < 0 < \alpha f'(x_g) + \beta g'(x_g)$ . By the mean value theorem, there must be a point  $x_{\alpha f + \beta g} \in [x_f, x_g]$  at which the derivative is zero. Since  $\alpha f + \beta g$  is strictly convex, the point where its derivative is zero must be unique. This concludes the proof.

#### B.1 A First Lower Bound on the Residual

Equipped with the tools from the previous sections, we now provide a preliminary lower bound on the residual, as desired to claim smoothness of our barrier MPC solution.

**Theorem B.7.** Let  $\mathcal{K} = \{x : Ax \ge b\}$  be a polytope such that each of m rows of A is normalized to be unit norm. Let  $\mathcal{K}$  contain a ball of radius r and be contained inside a ball of radius R centered at the origin. Let

$$u_{\eta} := \arg\min_{u} q(u) + \eta \phi_{\mathcal{K}}(u), \tag{B.6}$$

where q is a convex L-Lipschitz function and  $\phi_{\mathcal{K}}$  is a  $\nu$ -self-concordant barrier on  $\mathcal{K}$ . We show for  $res_i(u_n)$ , the *i*<sup>th</sup> residual at  $u_n$ , the following lower bound:

$$res_i(u_\eta) \ge \min\left\{\frac{\eta}{2}, \frac{r\eta^2}{150(\nu\eta^2 + R^2(L^2 + 1))}
ight\}$$

*Proof of Theorem B.7.* Applying the first-order optimality condition of  $u_n$  in (B.6) gives us that

$$\eta \nabla \phi_{\mathcal{K}}(u_{\eta}) + \nabla q(u_{\eta}) = 0. \tag{B.7}$$

From here on, we fix  $c = \nabla q(u_{\eta})$ , where  $u_{\eta}$  is as in (B.6). Then, we may conclude

$$u_{\eta} \in \arg\min_{u} c^{\top} u + \eta \phi_{\mathcal{K}}(u), \tag{B.8}$$

where we have replaced the cost q in (B.6) with a specific linear cost c; to see (B.8), observe that  $u_{\eta}$  satisfies the first-order optimality condition of (B.8) because of (B.7) and our choice of c.

We now define the function  $\tilde{\phi}_{\mathcal{K}}(x) = \eta^{-1} \cdot (c - a_i)^{\top} x + \phi_{\mathcal{K}}(x)$ . By Lemma B.4, we have that  $\tilde{\phi}_{\mathcal{K}}$  is a self-concordant-barrier on  $\mathcal{K}$  with self-concordance parameter

$$\widetilde{\nu} \le 20(\nu + R^2 \eta^{-2} (\|c\|^2 + \|a_i\|^2).$$
(B.9)

With this new self-concordant barrier in hand, we may now express  $u^{\eta}$  from (B.8) as the following optimizer:

$$u^{\eta} = \arg\min_{u} a_i^{\top} u + \eta \widetilde{\phi}_{\mathcal{K}}(u).$$
(B.10)

Further, let  $u^* \in \arg\min_{u \in \mathcal{K}} a_i^\top u$ . By applying Fact B.5 to  $u^\eta$  expressed as in (B.10), we have

$$\min\left\{\frac{\eta}{2}, \frac{r\|a_i\|}{2\tilde{\nu} + 4\sqrt{\tilde{\nu}}}\right\} \le a_i^{\top}(u_\eta - u^{\star}).$$
(B.11)

The lower bound in Inequality (B.11) may be expanded upon via (B.9), and chaining this with the observation  $a_i^{\top}(u_{\eta} - u^{\star}) = \operatorname{res}_i(u^{\eta}) - \operatorname{res}_i(u^{\star})$  gives:

$$\min\left\{\frac{\eta}{2}, \frac{r\|a_i\|}{150(\nu + R^2\eta^{-2}(\|c\|^2 + \|a_i\|^2))}\right\} \le \operatorname{res}_i(u^\eta) - \operatorname{res}_i(u^\star).$$

The definition of  $u^*$  implies  $\operatorname{res}_i(u^*) \ge 0$ , hence  $\operatorname{res}_i(u_\eta) \ge \min\left\{\frac{\eta}{2}, \frac{r}{150(\nu+R^2\eta^{-2}(L^2+1))}\right\}$ . Repeating this computation for each constraint of  $\mathcal{K}$  gives the claimed bound overall.

### B.2 An Improved Lower Bound on the Residual

In this section, we strengthen the bound from Theorem B.7 via a more careful analysis.

#### **B.2.1** Warmup: The One-Dimensional Case

We begin with a lemma on optimizing quadratics in one dimension to motivate our later results for arbitrary polytopes in higher dimensions.

**Lemma B.8.** Let  $\phi$  be a  $\nu$ -self-concordant barrier over (0,r) and q be a convex function such that  $\nabla q(v) = 0$  and  $0 < m \leq \nabla^2 q(x) \leq M$ . Define,

$$x^{\eta} := \arg\min_{x} q(x) + \eta \phi(x)$$

Then,

$$\min\left\{\frac{1}{2}\left(\sqrt{\frac{2\eta}{M} + v^2} + v\right), \frac{mr}{M(2\nu + 4\sqrt{\nu})}\right\} \le x^{\eta} \le \frac{1}{2}\left(\sqrt{\frac{4\eta\nu}{m} + v^2} + v\right).$$
(B.12)

*Proof.* Let  $c := \nabla q(x^{\eta})$ . Via the same trick as for (B.8), we can express  $x^{\eta}$  equivalently as  $x^{\eta} = \arg \min_{x} cx + \eta \phi(x)$ . Let  $x^{\star} := \arg \min_{x \in (0,r)} cx$  and  $\tilde{x} := \arg \min_{x} \phi(x)$ .

**Case 1**:  $v < \tilde{x}$ . Then by Claim B.6 applied to the functions q and  $\phi$ , we may deduce that  $v < x^{\eta} < \tilde{x}$ , meaning c > 0 and therefore  $x^* = 0$ .

Applying Fact B.5, we have:

$$\min\left\{\frac{\eta}{2}, \frac{rc}{2\nu + 4\sqrt{\nu}}\right\} \le cx^{\eta} \le \eta\nu.$$

Using that  $m(x^{\eta} - v) \le c \le M(x^{\eta} - v)$  we have,

$$\min\left\{\frac{\eta}{2M}, \frac{r(x^{\eta}-v)}{2\nu+4\sqrt{\nu}}\right\} \le (x^{\eta}-v)x^{\eta} \le \frac{\eta\nu}{m}.$$

Solving  $\frac{\eta}{2M} \leq (x^{\eta} - v)x^{\eta} \leq \frac{\nu\eta}{m}$  with the condition that  $x^{\eta} > v$ , we have,

$$\frac{1}{2}\left(\sqrt{\frac{2\eta}{M}+v^2}+v\right) \le x^{\eta} \le \frac{1}{2}\left(\sqrt{\frac{4\eta\nu}{m}+v^2}+v\right).$$

Combining with the minimum on the LHS bound, we arrive at

$$\min\left\{\frac{1}{2}\left(\sqrt{\frac{2\eta}{M}+v^2}+v\right),\frac{r}{2\nu+4\sqrt{\nu}}\right\} \le x^{\eta} \le \frac{1}{2}\left(\sqrt{\frac{4\eta\nu}{m}+v^2}+v\right).$$

**Case 2**:  $v \ge \tilde{x}$ . By the same reasoning as in the previous case, we have  $\tilde{x} \le x^{\eta} \le v$ . Note that by applying Fact B.5 with c = 1 and considering  $\eta \to \infty$ , we can deduce that  $\tilde{x} \ge \frac{r}{2\nu + 4\sqrt{\nu}}$ . We can see that (B.12) still holds as,

$$\min\left\{\frac{1}{2}\left(\sqrt{\frac{\eta}{M}+v^2}+v\right), \frac{r}{(2\nu+4\sqrt{\nu})}\right\} \le \frac{r}{2\nu+4\sqrt{\nu}} \le \tilde{x} \le x^{\eta} \le v \le \frac{1}{2}\left(\sqrt{\frac{4\eta\nu}{m}+v^2}+v\right).$$

The above result shows that if the minimizer of a strongly convex cost lies outside of the constraint set, we should expect to get a bound of the form  $O(\sqrt{\eta + v^2} - v)$ , where v is the distance to the constraint set.

#### **B.2.2** Upper Bounds on Approximation Error

**Lemma B.9.** Let  $\mathcal{K} \subset \mathbb{R}^n$  be a polytope and  $\phi$  be a  $\nu$ -self-concordant barrier on  $\mathcal{K}$ . Let  $x^{\eta} := \arg \min_{x \in \mathcal{K}} \frac{\alpha}{2} \|x - v\|^2 + \eta \phi(x)$  and  $x^* := \arg \min_{x \in \mathcal{K}} \frac{\alpha}{2} \|x - v\|^2$  for some  $v \in \mathbb{R}^n$ . Then,

$$\|x^{\eta} - x^{\star}\| \le \sqrt{\frac{\eta\nu}{\alpha}}.$$

*Proof.* We proceed similar to Theorem 4.3. Note that by  $\alpha$ -strong-convexity of  $q(x) := \frac{\alpha}{2} ||x - v||^2$ , we have that,

$$[\nabla q(x^{\eta}) - \nabla q(x^{\star})]^{\top}(x^{\eta} - x^{\star}) \ge \alpha \|x^{\eta} - x^{\star}\|^{2}.$$

Note that from the optimality condition  $\nabla q(x^{\eta}) + \eta \nabla \phi(x^{\eta}) = 0$ , and, by convexity of  $\mathcal{K}$ ,  $\nabla q(x^{\star})^{\top}[x^{\eta} - x^{\star}] \geq 0$ , and by Fact B.1, it follows,

$$\alpha \|x^{\eta} - x^{\star}\|^{2} \leq \eta \nabla \phi(x^{\eta})^{\top} [x^{\star} - x^{\eta}] - \nabla q(x^{\star})^{\top} [x^{\eta} - x^{\star}] \leq \nabla \phi(x^{\eta})^{\top} [x^{\star} - x^{\eta}] \leq \eta \nu.$$

Rearranging the terms then gives the claim.

Note that the above result can be generalized to  $\alpha$ -strongly-convex functions. In the next lemma, we show that we can make a similar bound along the gradient of the cost function at  $x^*$ .

**Lemma B.10.** Let  $\mathcal{K} \subset \mathbb{R}^n$  and  $\phi$  be a  $\nu$ -self-concordant barrier on  $\mathcal{K}$ . Let  $x^{\eta} := \arg \min_x \frac{\alpha}{2} ||x - v||^2 + \eta \phi(x)$  and  $x^* := \arg \min_{x \in \mathcal{K}} \frac{\alpha}{2} ||x - v||^2$  for some  $v \in \mathbb{R}^n$ . Assume that  $x^* \neq v$ , and let  $a = \frac{x^* - v}{||x^* - v||}$ . Then,

$$0 \le a^{\top}(x^{\eta} - x^{\star}) \le \frac{1}{2} \left( \sqrt{\frac{4\eta\nu}{\alpha} + \|x^{\star} - v\|^2} - \|x^{\star} - v\| \right).$$

(Note that for the case where  $x^* = v$ , Lemma B.9 can be chosen for any a)

*Proof.* Let  $q(x) := \frac{\alpha}{2} ||x - v||^2$ . Note that we can write:

$$abla q(x^\eta) = lpha \cdot a^ op (x^\eta - x^\star) a + lpha \cdot b^ op (x^\eta - x^\star) b + 
abla q(x^\star),$$

where ||a|| = ||b|| = 1 and  $b \perp a$ . Using (B.7) (valid because in this lemma we define  $x^{\eta} := \arg \min_x q(x) + \eta \phi(x)$  for a convex q) with Fact B.1 yields

$$\nabla q(x^{\eta})^{\top}(x^{\eta}-x^{\star}) \leq \eta \nu.$$

Then it follows that

$$\alpha \cdot [a^{\top}(x^{\eta} - x^{\star})]^{2} + \alpha [b^{\top}(x^{\eta} - x^{\star})]^{2} + \alpha ||x^{\star} - v|| \cdot a^{\top}(x^{\eta} - x^{\star}) \le \eta \nu.$$

We drop  $\alpha \cdot [b^{\top}(x^{\eta} - x^{\star})]^2$  and solve for  $a^{\top}(x^{\eta} - x^{\star})$  to prove our claimed upper bound on  $a^{\top}(x^{\eta} - x^{\star})$ . To derive the lower bound, we use that  $0 \leq \nabla q(x^{\star})^{\top}[x^{\eta} - x^{\star}] = c \cdot a^{\top}[x^{\eta} - x^{\star}]$  for some c > 0.  $\Box$ 

#### B.2.3 An Improved Lower Bound on the Residual

**Lemma B.11.** Fix a polytope  $\mathcal{K}$ , a convex function q, and a  $\nu$ -self-concordant barrier  $\phi$  over  $\mathcal{K}$ . Assume that the polytope  $\mathcal{K}$  contains a full-dimensional ball of radius r and is contained within a ball of radius R around some point  $\bar{x}$ , i.e.  $\mathcal{B}(\bar{x},r) \subseteq \mathcal{K} \subseteq \mathcal{B}(\bar{x},R)$ . Let  $x^{\eta} := \arg \min q(x) + \eta \phi(x)$  for arbitrary  $\eta > 0$ ,

$$\mathcal{B}\left(x^{\eta}, \frac{r}{R}\min\left\{\frac{\eta}{2\|\nabla q(x^{\eta})\|}, \frac{r}{2\nu + 4\sqrt{\nu}}\right\}\right) \subseteq \mathcal{K}.$$
(B.13)

*Proof.* Consider the line passing through  $\bar{x}$  and  $x^{\eta}$  given by  $S = \{\bar{x}t + x^{\eta}(1-t) : t\}$  and let  $x_1$  and  $x_2$  be the endpoints of  $\mathcal{K} \cap S$ . Equipped with these definitions, we will show

$$\min(\|x^{\eta} - x_1\|, \|x^{\eta} - x_2\|) \ge \min\left\{\frac{\eta}{2\|\nabla q(x^{\eta})\|}, \frac{r}{2\nu + 4\sqrt{\nu}}\right\}.$$
(B.14)

Before proving Inequality (B.14), we first show why it immediately gives the claimed result of (B.13). Pick  $\hat{x} \in \{x_1, x_2\}$  such that  $x^{\eta}$  lies on the line segment between  $\hat{x}$  and  $\bar{x}$ . Consider any direction c such

that ||c|| = 1. Consider the triangle formed by the points  $\hat{x}, \bar{x}$ , and  $\bar{x} + rc$ , and draw a line segment from  $x^{\eta}$ , parallel to c and intersecting the line segment from  $\hat{x}$  to  $\bar{x}$  at a point we label y. Then, by convexity,  $y \in \mathcal{K}$ . Then, we prove (B.13) by showing that  $||y - x^{\eta}|| \ge \frac{r}{R} \min\left\{\frac{\eta}{2||\nabla q(x^{\eta})||}, \frac{r}{2\nu + 4\sqrt{\nu}}\right\}$ . To see this inequality, we note that

$$\|y - x^{\eta}\| = \|\bar{x} - (\bar{x} + rc)\| \cdot \frac{\|x^{\eta} - \hat{x}\|}{\|\bar{x} - \hat{x}\|} = r \cdot \frac{\|x^{\eta} - \hat{x}\|}{\|\bar{x} - \hat{x}\|} \ge r \cdot \frac{1}{R} \cdot \min\left\{\frac{\eta}{2\|\nabla q(x^{\eta})\|}, \frac{r}{2\nu + 4\sqrt{\nu}}\right\},$$

where the first equation is by similarity of the triangles formed by  $\hat{x}$ ,  $\bar{x}$ , and  $\bar{x} + rc$  and by  $\hat{x}$ ,  $x^{\eta}$ , and y; the final step is by Inequality (B.14) and the assumed upper bound of R on the polytope diameter. We now proceed to prove Inequality (B.14). Without loss of generality, let  $x_1$  be such that  $\nabla q(x^{\eta})^{\top} \left(\frac{\bar{x}-x_1}{\|\bar{x}-x_1\|}\right) \geq 0$ . We use this characterization of  $x_1$  in both parts of our proof below.

**Lower bound on**  $||x^{\eta} - x_2||$ . Denote the restriction of the barrier  $\phi$  (defined on the polytope  $\mathcal{K}$ ) to the line  $\mathcal{S}$  by a univariate function  $\xi$ , so that

$$\xi(t) := \phi(tx_2 + (1-t)x^{\eta}), \text{ with } \xi'(t) = \nabla \phi(tx_2 + (1-t)x^{\eta})^{\top}(x_2 - x^{\eta}).$$
(B.15)

We note that  $\xi(0) = \phi(x^{\eta})$  and  $\xi(1) = \phi(x_2)$ . By definition of  $\phi$  as a barrier on  $\mathcal{K}$ , note that  $\xi$  is also a barrier defined only on  $\mathcal{K} \cap \mathcal{S}$  [5]. Define the following quantities associated with  $\xi$ :

$$t_{\rm ac} := \arg\min_t \xi(t), \text{ and } x_{\rm ac} := t_{\rm ac} x_2 + (1 - t_{\rm ac}) x^{\eta}.$$
 (B.16)

In other words,  $x_{ac}$  is the analytic center of the barrier  $\xi$  on S. We now apply Fact B.5 with  $c = x_{ac} - x_2$ ,  $\mathcal{K} = S$ , the barrier  $\xi$  on S,  $\eta \to \infty$ , and denoting  $\nu_{\xi}$  to be the self-concordance parameter of  $\xi$ . Then, combining the lower bound in Fact B.5 with the definition of  $x_{ac}$  and  $\nu_{\xi} \leq \nu$  [5] yields

$$||x_{\rm ac} - x_2|| \ge \frac{r}{2\nu_{\xi} + 4\sqrt{\nu_{\xi}}} \ge \frac{r}{2\nu + 4\sqrt{\nu}}.$$
 (B.17)

Next we have by the choice of  $x_1$  that  $\nabla q(x^{\eta})^{\top}[\bar{x} - x_1] \ge 0$ . By the first-order optimality condition of  $x^{\eta}$  (as in (B.7)), this is equivalent to  $\nabla \phi(x^{\eta})^{\top}[\bar{x} - x_1] \le 0$ . This final inequality implies

$$\nabla \phi(x^{\eta})^{\top}[x_2 - x^{\eta}] \le 0 \tag{B.18}$$

since  $x_1$  and  $x_2$  are the end points of  $\mathcal{K} \cap \mathcal{S}$  (the line segment whose interior contains  $\bar{x}$  and  $x^{\eta}$ ), and hence  $x_2 - x^{\eta}$  is a vector in the same direction as  $\bar{x} - x_1$ . Note that we have

$$\xi'(0) \le 0 \text{ and } \xi'(t_{\rm ac}) = 0,$$
 (B.19)

where the first inequality is by using (B.15) to equivalently rewrite Inequality (B.18), and the equality is by construction of  $t_{\rm ac}$  in (B.16) and by convexity of  $\xi$ . Since the univariate function  $\xi$  is strictly convex, its derivatives are strictly monotone; hence, Inequality (B.19) implies that  $t_{\rm ac} \geq 0$ . Recalling that by (B.15), t = 0 corresponds to  $x^{\eta}$  and t = 1 corresponds to  $x_2$ , and that  $x_{\rm ac} \in \operatorname{int}(\mathcal{S} \cap \mathcal{K})$ , we may deduce from  $t_{\rm ac} \geq 0$  that

$$x_{\rm ac} \in \{x^{\eta}t + (1-t)x_2 \mid t \in [0,1]\}.$$
(B.20)

Combining Inequalities (B.17) and (B.20), we have

$$||x^{\eta} - x_2|| \ge ||x_{\rm ac} - x_2|| \ge \frac{r}{2\nu + 4\sqrt{\nu}},$$
 (B.21)

which proves one part of Inequality (B.14).

**Lower bound on**  $||x^{\eta} - x_1||$ . We now parameterize S by  $\psi(t) = x_1 + t \frac{\bar{x} - x_1}{||\bar{x} - x_1||}$ . Define  $c := \nabla q(x^{\eta})^{\top} \left(\frac{\bar{x} - x_1}{||\bar{x} - x_1||}\right)$ . We then define the following two optimizers

$$t^{\star} := \arg\min_{t,\psi(t)\in\mathcal{K}} c \cdot t, \text{ and } t^{\eta} := \arg\min_{t} c \cdot t + \eta \phi(\psi(t)).$$

It follows from  $\psi([0, 2r]) \subseteq \mathcal{K}$  and  $c \geq 0$  (by our choice of  $x_1$ ) that  $t^* = 0$ . We then apply Fact B.5 with the above  $c, t^{\eta}, t^*$ , and barrier  $\phi(\psi(\cdot))$  (with its associated self-concordance parameter  $\nu_{\phi\circ\psi} \leq \nu$  [5]) and Cauchy-Schwarz inequality to conclude that,

$$\min\left\{\frac{\eta}{2\|\nabla q(x^{\eta})\|}, \frac{r}{2\nu + 4\sqrt{\nu}}\right\} \le \min\left\{\frac{\eta}{2c}, \frac{r}{2\nu + 4\sqrt{\nu}}\right\} \le t^{\eta}.$$
(B.22)

Finally, note that the optimality condition of  $x^{\eta}$  implies  $\nabla q(x^{\eta}) + \eta \nabla \phi(x^{\eta}) = 0$ , and specifically that  $[\nabla q(x^{\eta}) + \eta \nabla \phi(x^{\eta})]^{\top} \left(\frac{\bar{x} - x_1}{\|\bar{x} - x_1\|}\right) = 0$ . Since  $x^{\eta} \in S$ , we can write

$$\begin{split} 0 &= [\nabla q(x^{\eta}) + \eta \nabla \phi(x^{\eta})]^{\top} \left(\frac{\bar{x} - x_1}{\|\bar{x} - x_1\|}\right) \\ &= \nabla q(x^{\eta})^{\top} \left(\frac{\bar{x} - x_1}{\|\bar{x} - x_1\|}\right) + \eta \frac{d(\phi \circ \psi)}{dt}(\psi^{-1}(x^{\eta}))) \\ &= c + \eta \frac{d(\phi \circ \psi)}{dt}(\psi^{-1}(x^{\eta})). \end{split}$$

We can observe that  $c + \eta \frac{d}{dt} (\phi \circ \psi)(t)|_{t=t^{\eta}} = 0$  is the optimality condition of  $t^{\eta}$ . Since  $\phi$  and, by extension,  $\phi \circ \psi$  are strongly convex, we have that  $\psi^{-1}(x^{\eta}) = t^{\eta}$ . Since  $\psi$  is parameterized in terms of distance from  $x_1$ , we have that,

$$t^{\eta} = \|x^{\eta} - x_1\|. \tag{B.23}$$

Combining Inequalities (B.21) and (B.22) and (B.23) yields Inequality (B.14), which, as argued earlier, concludes the proof of the lemma.  $\Box$ 

Similar to Lemma B.10, we now adapt this to get a lower bound for isotropic quadratics.

**Lemma B.12.** Let  $\mathcal{K} \subset \mathbb{R}^n$  be a polytope and  $\phi$  be a  $\nu$ -self-concordant barrier function. Assume there exists  $\bar{x} \in \mathcal{K}$  such that  $\mathcal{B}(\bar{x}, r) \subseteq \mathcal{K} \subseteq \mathcal{B}(\bar{x}, R)$  for some r, R > 0. Let  $x^{\eta} := \arg \min_x \frac{\alpha}{2} ||x - v||^2 + \eta \phi(x)$ ,  $x^* := \arg \min_{x \in \mathcal{K}} \frac{\alpha}{2} ||x - v||^2$  for some  $v \in \mathbb{R}^n$ . Then we know the following ball centered around  $x^{\eta}$  is contained within  $\mathcal{K}$ .

$$\mathcal{B}\left(x^{\eta}, \frac{r}{R}\min\left\{\frac{1}{2\sqrt{\nu}}\left(\sqrt{\frac{\eta}{\alpha} + \|x^{\star} - v\|^{2}} - \|x^{\star} - v\|\right), \frac{r}{2\nu + 4\sqrt{\nu}}\right\}\right) \subseteq \mathcal{K}.$$

*Proof.* To prove this we use Lemma B.11 and techniques similar to Lemma B.10. By the triangle inequality and Lemma B.9, we have

$$\alpha \|x^{\eta} - v\| \le \alpha (\|x^{\eta} - x^{\star}\| + \|x^{\star} - v\|) \le \sqrt{\alpha} \sqrt{\eta \nu} + \alpha \|x^{\star} - v\|.$$

We next apply this upper bound to the result of Lemma B.11 with  $q(x) = \frac{\alpha}{2} ||x - v||^2$  and obtain:

$$\mathcal{B}\left(x^{\eta}, \frac{r}{R} \min\left\{\frac{\eta}{2(\sqrt{\alpha\eta}\sqrt{\nu} + \alpha \|x^{\star} - v\|)}, \frac{r}{2\nu + 4\sqrt{\nu}}\right\}\right) \subseteq \mathcal{K}.$$

With some rearranging, we may express the above bound as:

$$\mathcal{B}\left(x^{\eta}, \frac{r}{R} \min\left\{\frac{1}{2\sqrt{\nu}} \frac{\frac{\eta}{\alpha}}{\sqrt{\frac{\eta}{\alpha}} + \frac{1}{\sqrt{\nu}} \|x^{\star} - v\|}, \frac{r}{2\nu + 4\sqrt{\nu}}\right\}\right) \subseteq \mathcal{K}.$$

Observe that for any  $x > 0, y \in \mathbb{R}$ , we have that,

$$\sqrt{x+y^2} - y = \frac{x}{\sqrt{x+y^2} + y} \le \frac{x}{\sqrt{x} + y}.$$

Since  $\nu \ge 1$ , we have  $\frac{\eta/\alpha}{\sqrt{\eta/\alpha} + \frac{1}{\sqrt{\nu}} \|x^{\star} - v\|} \ge \frac{\eta/\alpha}{\sqrt{\eta/\alpha} + \|x^{\star} - v\|} \ge \sqrt{\frac{\eta}{\alpha} + \|x^{\star} - v\|^2} - \|x^{\star} - v\|$ . We can then simplify the last bound on the radius around  $x^{\eta}$  to match the claimed bound.

#### **B.2.4** Consolidated Upper and Lower Bounds

We now collect Lemma B.10 and Lemma B.12, performing a change of basis to provide bounds for arbitrary quadratic objective functions.

**Theorem B.13.** Let  $\mathcal{K} = \{x : Ax \ge b\}$  be a polytope for some  $A \in \mathbb{R}^{n_r \times n}, b \in \mathbb{R}^{n_r}$ . Let  $\phi$  be a  $\nu$ -self-concordant barrier over  $\mathcal{K}$ . Assume there exists  $\bar{x} \in \mathcal{K}$  such that  $\mathcal{B}(\bar{x}, r) \subseteq \mathcal{K} \subseteq \mathcal{B}(\bar{x}, R)$  for some r, R > 0. Let,

$$\begin{split} x^{\eta} &:= \operatorname*{arg\,min}_{x} \frac{1}{2} (x-v)^{\top} H(x-v) + \eta \phi(x), \\ x^{\star} &:= \operatorname*{arg\,min}_{x \in \mathcal{K}} \frac{1}{2} (x-v)^{\top} H(x-v), \end{split}$$

where  $mI \leq H \leq MI$ . Let  $a = \frac{H(x^*-v)}{\|H(x^*-v)\|}$  if  $\|x^*-v\| > 0$ . Then the following hold:

$$\begin{array}{l} (i) \ \|x^{\eta} - x^{\star}\| \leq \sqrt{\frac{\eta\nu}{m}}, \\ (ii) \ 0 \leq a^{\top}(x^{\eta} - x^{\star}) \leq \frac{1}{2\sqrt{m}} \left(\sqrt{4\eta\nu + \|x^{\star} - v\|_{H}^{2}} - \|x^{\star} - v\|_{H}\right), \\ (iii) \ \mathcal{B}\left(x^{\eta}, \sqrt{\frac{m}{M}} \cdot \frac{r}{R} \cdot \min\left\{\frac{1}{\sqrt{\nu M}} \left(\sqrt{\eta + \|x^{\star} - v\|_{H}^{2}} - \|x^{\star} - v\|_{H}\right), \sqrt{\frac{m}{M}} \cdot \frac{r}{2\nu + 4\sqrt{\nu}}\right\}\right) \subseteq \mathcal{K}. \end{array}$$

Note that this implies that, if a exists, then

$$\sqrt{\frac{m}{M}} \cdot \frac{r}{R} \cdot \min\left\{\frac{1}{\sqrt{\nu M}} \left(\sqrt{\eta + \|x^{\star} - v\|_{H}^{2}} - \|x^{\star} - v\|_{H}\right), \sqrt{\frac{m}{M}} \cdot \frac{r}{2\nu + 4\sqrt{\nu}}\right\} \le a^{\top}(x^{\eta} - x^{\star}).$$

Note that Theorem B.13(iii) implies a lower bound on the residual (distance from boundary of  $\mathcal{K}$ ) of  $u^{\eta}$ . This is the bound we directly use in our smoothness bound in Theorem 4.8.

Proof of Theorem B.13. For both the upper and lower bounds, we use the change of basis  $y = H^{1/2}x$ ,  $z = H^{1/2}v$ . We can then transform the assumed definitions of  $x^{\eta}$  and  $x^{\star}$  into the following optimization problem in y:

$$egin{aligned} &y^\eta := rgmin_y rac{1}{2} \|y-z\|^2 + \eta \cdot \phi(H^{-1/2}y), \ &y^\star := rgmin_{y \in H^{1/2} \cdot \mathcal{K}} rac{1}{2} \|y-z\|^2. \end{aligned}$$

By Fact B.2, we have that  $\phi \circ H^{-1/2}$  defined on the set  $H^{1/2} \cdot \mathcal{K}$  is still a self-concordant barrier with parameter  $\nu$ . Therefore, we have that,

$$||x^{\eta} - x^{\star}|| \le \frac{1}{\sqrt{m}} ||y^{\eta} - y^{\star}|| \le \sqrt{\frac{\eta\nu}{m}},$$

where the final step is by Lemma B.9. This completes the proof of Theorem B.13(i). Next, we define  $\tilde{a} := \frac{y^{\star}-z}{\|y^{\star}-z\|}$  and  $\hat{a} := \frac{1}{\sqrt{m}} \cdot H^{1/2}\tilde{a}$ . Then we have  $a = \hat{a}/\|\hat{a}\| = \frac{H(x^{\star}-v)}{\|H(x^{\star}-v)\|}$  and

$$0 \le a^{\top}(x^{\eta} - x^{\star}) = \frac{1}{\|\hat{a}\|} \hat{a}^{\top}(x^{\eta} - x^{\star}) = \frac{1}{\|\hat{a}\|\sqrt{m}} \tilde{a}^{\top}(y^{\eta} - y^{\star}) \le \frac{1}{2\sqrt{m}} \left(\sqrt{4\eta\nu + \|y^{\star} - z\|^2} - \|y^{\star} - z\|\right)$$

where the first and final inequalities are by Lemma B.10 applied to  $y^{\eta}$  and  $y^{\star}$ , and using that  $\|\hat{a}\| \ge 1$ . Finally, by again using this affine transformation between y and x, we obtain Theorem B.13(ii):

$$0 \le a^{\top}(x^{\eta} - x^{\star}) \le \frac{1}{2\sqrt{m}} \left( \sqrt{4\eta\nu + \|x^{\star} - v\|_{H}^{2}} - \|x^{\star} - v\|_{H} \right).$$

For the lower bound, we note that  $H^{1/2}\mathcal{K}$  contains a ball of radius at least  $r \cdot \sqrt{m}$  and is contained in a ball of radius at most  $R \cdot \sqrt{M}$ . Applying Lemma B.12, we have that,

$$\mathcal{B}\left(y^{\eta}, \sqrt{\frac{m}{M}} \cdot \frac{r}{R} \min\left\{\frac{1}{\sqrt{\nu}}\left(\sqrt{\eta + \|y^{\star} - z\|^2} - \|y^{\star} - z\|\right), \sqrt{m} \cdot \frac{r}{2\nu + 4\sqrt{\nu}}\right\}\right) \subseteq H^{1/2} \cdot \mathcal{K}.$$

Plugging in the definitions for y and z in the LHS, we have,

$$\mathcal{B}\left(y^{\eta}, \sqrt{\frac{m}{M}} \cdot \frac{r}{R} \min\left\{\frac{1}{\sqrt{\nu}}\left(\sqrt{\eta + \|x^{\star} - v\|_{H}^{2}} - \|x^{\star} - v\|_{H}\right), \sqrt{m} \cdot \frac{r}{2\nu + 4\sqrt{\nu}}\right\}\right) \subseteq H^{1/2} \cdot \mathcal{K}.$$

Since  $||H^{1/2}|| \leq \sqrt{M}$ , we have  $\sigma_{\min}(H^{-1/2}) \geq \frac{1}{\sqrt{M}}$ . This implies that the ball centered around  $x^{\eta}$  in  $\mathcal{K}$  is at least  $\frac{1}{\sqrt{M}}$  the radius of the ball centered around  $y^{\eta}$  in  $H^{1/2} \cdot \mathcal{K}$ . Thus, as claimed in Theorem B.13(iii), we have,

$$\mathcal{B}\left(x^{\eta}, \sqrt{\frac{m}{M}} \cdot \frac{r}{R} \min\left\{\frac{1}{\sqrt{\nu M}} \left(\sqrt{\eta + \|x^{\star} - v\|_{H}^{2}} - \|x^{\star} - v\|_{H}\right), \sqrt{\frac{m}{M}} \cdot \frac{r}{2\nu + 4\sqrt{\nu}}\right\}\right) \subseteq \mathcal{K}.$$

Finally, note that  $a := \frac{H(x^*-v)}{\|H(x^*-v)\|} = \frac{\nabla q(x^*)}{\|\nabla q(x^*)\|}$ , where  $q(x) := \frac{1}{2}(x-v)^{\top}H(x-v)$  is the objective for the hard-constrained problem. By the optimality of  $x^*$  for this problem, a is a combination of the active constraints at  $x^*$ , meaning that

$$\{x: a^{\top}(x - x^{\star}) \ge 0\} \supseteq \mathcal{K}.$$

Since, by Theorem B.13(iii), there exists a minimum-radius ball around  $x^{\eta}$  (which in turn implies that the minimum distance of  $x^{\eta}$  from the boundary of  $\mathcal{K}$  is at least this radius) and a is unit-norm, we can conclude that the distance of  $x^{\eta}$  from the hyperplane specified by a is at least this radius:

$$\sqrt{\frac{m}{M}} \cdot \frac{r}{R} \min\left\{\frac{1}{\sqrt{\nu M}} \left(\sqrt{\eta + \|x^{\star} - v\|_{H}^{2}} - \|x^{\star} - v\|_{H}\right), \sqrt{\frac{m}{M}} \cdot \frac{r}{2\nu + 4\sqrt{\nu}}\right\} \le a^{\top}(x^{\eta} - x^{\star}).$$