Mechanical Interactions Govern Self-Organized Ordering in Two-Dimensional Bacterial Colonies

Samaneh Rahbar, Ludger Santen, and Reza Shaebani *Department of Theoretical Physics and Center for Biophysics, Saarland University, 66123 Saarbr¨ucken, Germany*

We investigate the influence of mechanical interactions on the growth and self-organization of rodshaped bacteria confined within square and circular geometries. Our results reveal that mechanical interactions are crucial in constraining cell growth and shaping the degree of ordering within bacterial colonies. Colonies with larger aspect ratios exhibit higher levels of nematic ordering, while the geometry of confinements significantly impacts colony organization. Compression influences force magnitudes within colonies, with larger division lengths encountering greater forces. These findings enhance our understanding of bacterial self-organization in confined environments, with implications for microbial ecology and biotechnology.

PACS numbers: 87.15.-v, 87.10.-e, 87.10.Pq

I. Introduction

Bacteria can freely disperse in the environment as motile, self-propelled particles to probe the environment for nutrients, or they can form a biofilm as immobilized cells, that attach to any available surface. During cell attachment on the surface, the early stage of the microcolony formation comprises aggregation of the individual bacterial. As long as the surface provides moisture and nutrients, we are likely to find biofilm [\[1](#page-5-0)[–5\]](#page-6-0).

A freely growing colony of rod-shaped non-selfpropelled bacterial takes advantage of the force interaction resulting from the linear growth of the length in time to expand and proliferate. The expansion of the colony occurs over the interplay between adhesive interactions between bacteria and the substrate and the driven forces caused by the growth of the bacteria. Bacteria are attached to the surface due to the adhesive interaction; while force driven caused by the bacterial growth pushes the bacteria and leads to the proliferation and expansion of the colony. Due to mechanical interactions, locally aligned areas appeared, these alignments could be strengthened or weakened due to the torque generated. Such a bacterial colony is globally isotropic due to the active forces caused by the bacterial growth and imperfect alignment of the bacteria after division which perturb the orientational distribution. It has been known that the key factors on the geometrical properties of the aligned area are, the aspect ratio (length to width), growth rates ν_l , and bacterial division length l_d [\[6](#page-6-1)[–8\]](#page-6-2).

Several studies have been conducted on nonmotile bacteria both experimentally and computationally to explore the outcomes of confinement on the emergence of global ordering within the microbial community. In a specific study, cultured bacteria in a microfluidic channel exhibit a distinct nematic ordering as the channel becomes densely populated. The mechanical interaction between the proliferation of bacteria and the constraint of the channel causes the emergence of the ordered pattern.

Additionally, computational studies have revealed that anisotropic geometrical confinement, such as a straight channel, induces mechanical anisotropy, which significantly influences the collective arrangement of bacteria. The intricate interplay between the mechanical properties of the confinement and the responses of bacterial cells gives rise to phenomena where the bacteria, align themselves along the path of least stress. [\[6,](#page-6-1) [10\]](#page-6-3).

In this paper, we explore the intricate interplay between mechanical interactions and bacterial selforganization within confined environments. Through molecular dynamics simulations, we investigate how these interactions influence the growth dynamics, spatial organization, and degree of ordering within bacterial colonies. Specifically, we examine the effects of confinement geometry and bacterial division lengths on the emergence of aligned areas and the distribution of mechanical forces within the colony. Our findings shed light on the fundamental mechanisms driving bacterial selforganization, with implications for understanding microbial behavior in various natural and engineered environments.

Experimental studies have furthered our understanding of how external mechanical conditions impact bacterial colony growth. These investigations reveal that exposure to external pressure leads to a reduction in the rate of cell production within bacterial colonies. Interestingly, both lateral and longitudinal forces exert varying effects on bacterial growth by influencing DNA segregation, albeit with differing intensities [\[11](#page-6-4)[–13\]](#page-6-5). These findings underscore the significance of mechanical forces in shaping bacterial growth dynamics and highlight the importance of considering such forces in bacterial growth regulation. As such, we introduce a growth equation for bacterial cells that accounts for mechanical forces acting along the longitudinal axis, recognizing their pivotal role in bacterial growth regulation

FIG. 1: (a) Hertz contact force interaction between two rodshaped bacteria. (b) A bacterium with initial length L , the bacterium elongates to length l_d before undergoing division, giving rise to two daughter bacteria of specific lengths

II. Method

Each bacterium is modeled as a rod-shaped with a constant diameter d_0 , and length L , which grows as a function of time and force in a two-dimensional space. Bacteria mechanically interact according to the Hertz contact force,

$$
F_{ij}(h) = Ed_0^{\frac{1}{2}} (h_{ij})^{\frac{3}{2}}, \qquad (1)
$$

where E the Young's modulus of the bacteria, h_{ij} is the overlap distance between bacteria i and bacteria j . The resulting force exerts on the major axis of the bacteria and is normal to the bacterial surface. Bacterial over-damped dynamics including translational and orientational motion follow Newton's laws,

$$
\frac{d\vec{r}_i}{dt} = \frac{1}{\zeta L_i} \sum_j F_{ij} \cdot \frac{\vec{r}_{c_{ij}}}{|\vec{r}_{c_{ij}}|},\tag{2}
$$

$$
\frac{d\theta_i}{dt} = \frac{12}{\zeta L_i^3} \sum_j ((\vec{r}_{c_{ij}} - \vec{r}_i) \times \vec{F}_{ij}) \cdot \hat{z}
$$
(3)

where, r_i and θ_i are the position and orientation of the bacterial center of mass, and $\vec{r}_{c_{ij}}$, as it is shown in Fig1(a), is the contact vector between *i*th and *j*th bacteria. ζ represents the friction per unit length of bacteria due to the contribution of bacteria-substrate adhesion. In this model, ζ is an isotropic quantity as the force is proportional to the bacterium diameter and is independent of the cell orientation.

Mechanical interactions between bacteria significantly shape bacterial growth and impact various aspects of bacterial behavior. Confinement-induced effects alter cell division, motility, and spatial organization, impacting growth dynamics within confined environments. Based on this observation, we devised a growth equation

that captures the influence of mechanical forces on the colony's growth dynamics. Of particular significance is the effect of forces acting along the major axis of a cell on its growth. To address this, we formulated a growth equation that accounts for both time and force, integrating an average growth rate into the model ν_i ,

$$
\frac{\mathrm{d}L_i(t)}{\mathrm{d}t} = \begin{cases} \nu_{l_i} - \beta(|F_{p_i}|), & \nu_{l_i} > \beta(|F_i|), \\ 0, & \nu_{l_i} \le \beta(|F_i|), \end{cases} (4)
$$

where $|F_p|$ is the total value of the projected forces along the major axis of the *i*th bacteria and β characterizes the mechanosensitivity of the bacteria, representing the degree to which an individual bacterium responds to mechanical forces, either robustly or weakly.

The value of the $|F_p|$ on *i*th bacteria, has been computed with the following method. We have individual forces acting on each bacterium. It is crucial to note that all of these forces must be projected along the bacterium's major axis, represented by \hat{n} ($\hat{n} = \cos \phi_i \sin \theta_i \hat{x} +$ $\sin \theta_i \sin \phi_i \hat{y} + \cos \theta_i \hat{z})$).

The contribution of each force interaction projected along the major axis of the *i*th bacteria $(\hat{n_i})$ would be $|F_{p_i}| = \sum_{j=1}^{N_c} |\vec{F_{ij}} \cdot \hat{n_i}| = \sum_{j=1}^{N_c} |F_{x_{ij}} \cdot n_{x_i} + F_{y_{ij}} \cdot n_{y_i} + F_{z_{ij}}$ n_{z_i} , which the summation is conducted over the total number of contacts (N_c) for each bacterium. Following up on the second method, we consider a condition, which up to the deformation equal to half of the cell's diameter $\left(\frac{d_0}{2}\right)$, the cell stops growing, accordingly $\frac{dL}{dt} = 0$ [\[12,](#page-6-6) [13\]](#page-6-5).

The value of β could be estimated by the average value of the bacterial growth rate, and a threshold value for $|F_p|$ in a way that $\frac{dL}{dt} = 0$. The threshold value for $|F_p|$ is a force in which two bacteria are deformed around half of the cell's diameter which is approximately equal to $35 kPa(\mu m)^2$, so the corresponding β for average growth $\nu = \frac{4\mu m}{h}$ is $\beta = 0.11 \frac{1}{kPa(\mu m)h}$. A higher value of β corresponds to mechanical conditions where small forces have a greater impact on bacterial growth. Conversely, a smaller value of β indicates that higher forces have a greater effect on growth.

As the applied pressure increases, it is observed that the doubling time of bacteria lengthens, regardless of whether it arises from cell-cell or cell-confinement interactions [\[11](#page-6-4), [12\]](#page-6-6). Eventually, when bacteria reach a threshold length, denoted as l_d , cell division occurs $Fig1(b).$ $Fig1(b).$ $Fig1(b).$

Following division, the two daughter cells generally maintain a comparable orientation to that of the mother cell, albeit with minor random deviations between (0, 0.1745] as the alignment of the two daughter cells is not perfect. The growth rates of the daughter cells are randomly chosen from the uniform distribution of the average growth rate $\left[\frac{\nu_l}{2}, \frac{3\nu_l}{2}\right]$ [\[7](#page-6-7), [10](#page-6-3)]. The magnitude order of some parameters is similar to the other

FIG. 2: Visualizing Bacterial Colony Dynamics: Simulated configurations at concurrent time steps. (a) $l_d = 2\mu m$. $N_{bacteria} = 2853$ (b) $l_d = 3\mu m$, $N_{bacteria} = 1265$ and (c) $l_d = 4\mu m$, $N_{bacteria} = 505$.

literature [\[16\]](#page-6-8): $d_0 = 0.5 \ \mu m, \ \nu_t = 4 \frac{\mu m}{h}, \ \ l_d = [2 - 4] \mu m,$ $\zeta = 200$ Pah, $E = 400$ kPa.

The motions of bacteria Eqs. [\(2\)](#page-1-1)-[\(3\)](#page-1-2) have been computed numerically by solving the equation of motion using an Euler time-stepping algorithm with time step, $\Delta t = \frac{\zeta}{E}$. Δt defines the time required for a bacteria to move on a substrate over the physical adhesion contact, between the bacteria and bacteria-bacteria mechanical interaction. Those mentioned above Δt is valid in the growth equation as long as it doesn't lead to undamped or oscillatory behavior in the system. Each simulation is initiated with a randomly positioned and oriented cell in square and circular confinement in dimension $L_x \times L_y$ and diameter d_c accordingly. Mechanical interaction between bacteria and the confinement follows Hertz contact force according to Eq.1 in both x and y direction. We consider an impenetrable hard wall with an elastic modulus a few times greater than the bacterial elastic modulus $E_{wall} = 1000kPa.$

III. Results

We use a molecular dynamics simulation to examine the growth of the bacterial colony without and under constraint. Bacterial growth in confinement occurs inside square confinement in dimension $60 \times 60(\mu m)^2$, and circular confinement with diameter $d_c = 60 \mu m$, with a bacteria located at the center of the box. All the simulations have been done for different initial conditions, different division lengths l_d , and the projected forces' contribution to growth according to Eq [\(4\)](#page-1-3).

It is evident that following the initial growth phase, bacteria tend to align locally in a shared direction due to contact forces and mechanical interactions (Fi[g2\)](#page-2-0). These local alignments occur because the contact forces shared by the bacteria influence their orientation. However, at the colony scale, where bacteria grow freely, the orientational distribution does not exhibit any discernible preferential orientation [\[14](#page-6-9)].

The bacterial growth medium is supplied with nutrients, enabling bacteria to grow at a constant average

growth rate $\nu_i = 4 \frac{\mu m}{h}$. As Eq [\(4\)](#page-1-3) indicates, the primary factor regulating population growth is the mechanical interaction between adjacent bacteria.

We established square and circular confinements, which act as rigid confinement. The interaction between hard walls and the bacteria follows the Hertz contact force $Eq(1)$ $Eq(1)$. Simulations have been done for different division lengths l_d with one bacteria placed in the middle of a square and circle-like confinements. We allowed the simulations to evolve and monitored the magnitude of the nematic ordering across various division lengths l_d until the confinement reached a high density of bacteria Fig. [3.](#page-3-0)

Considering the scaler order parameter $S =$ $\langle 2\cos^2\theta_{ij} - 1\rangle_{ij}$, which θ_{ij} is the angle between *i*th and jth bacteria, allows us to evaluate the extent of bacterial alignment across different aspect ratios. Additionally, calculating the orientational correlation function in a colony in contrast to the orientation of the initial bacteria enables us to systematically analyze and quantify the level of alignment or ordering present among the bacterial cells concerning the initial bacteria [\[15\]](#page-6-10). This approach provides valuable insights into the organization of bacterial populations within confined environments, aiding in understanding emergent patterns and dynamics.

Accordingly, the magnitude of nematic ordering and orientational correlation function within the colony are directly influenced by a dynamic interplay of factors. These factors include the strength of mechanical interactions, the concentration of bacteria, the aspect ratios of individual bacteria, and the geometric intricacies of the confinement. All these factors play a significant role in shaping the patterns and organization observed within bacterial colonies.

As the simulation progresses, a consistent decrease is observed in both the scalar order parameter and the orientational correlation function at each time point. This decline is primarily driven by the anisotropic growth of the bacteria, which leads to an imbalance between contact forces and drag. This imbalance results in buckling instabilities in the bacterial configuration. Moreover, the inherent stochasticity and dynamic system characteristics, such as the increasing number of bacteria and their division, may intensify these instabilities.

The rate of decline in both the scalar order parameter and the orientational correlation function is slower for colonies with a larger aspect ratio, indicating that they evolve more gradually than those with smaller aspect ratios. This slower rate of decline can be attributed to the smaller population in these colonies, which undergoes fewer instabilities caused by cell division and mechanical forces. Consequently, colonies with larger aspect ratios tend to be stiffer and more resistant to deformation. Additionally, the random orientation generated by cell division also contributes to an anisotropic stress, further highlighting the stability of larger aspect ratio colonies

FIG. 3: Configuration of bacterial colonies for each division length under square and circular confinement. The figure depicts the time step at which the confinement is filled with bacteria. Panels (a-c) represent division lengths $l_d = [2, 3, 4] \mu m$ respectively in square-like confinement, while panels (d-f) represent the same division lengths in circle-like confinement. The color circle indicates the orientation of the bacteria.

(Fig. [4\(](#page-3-1)a,b)).

At a specific time point when the colony reaches the confinement's boundary, it stabilizes its scalar order parameter. This stabilization occurs approximately around 5.5 hours for $l_d = 2\mu m$, 7.5 hours for $l_d = 3\mu m$, and 10 hours for $l_d = 4\mu m$. The square confinement's distinct edges and corners offer clear alignment cues, resulting in higher degrees of nematic ordering. In contrast, the circular confinement, lacking such prominent features along its perimeter, exhibits comparatively lower degrees of alignment.

Taken together, colonies characterized by higher as-

FIG. 4: Sample-averaged orientational order parameter and orientational correlation function for both square and circlelike confinement. (a) Orientational order parameter illustrated for different values of division lengths $l_d = [2, 3, 4] \mu m$. (b) Orientational correlation function has shown for different lengths $l_d = [2, 3, 4] \mu m$.

FIG. 5: The distribution $P(|F|)$ of individual contact forces against the circle and square-like confinement for $l_d = 4$, $l_d =$ $3\mu m$, $l_d = 2\mu m$. (a) square-like, (b) circle-like. The Red dashed line shows $|F| = 13 kPa(\mu m)^2$.

pect ratios demonstrate elevated levels of nematic ordering and orientational correlation, primarily attributed to the augmented overlap between contact cells, a topic explored in detail later in the paper.

Bacterial colonies with larger division lengths tend to have more compact and organized structures than those with shorter division lengths, as depicted in Fig. [4.](#page-3-1) During growth and division, contact interactions between growing bacteria cause neighboring bacteria to align and reorient themselves, eventually resulting in lateral packing. As we mentioned earlier, anisotropic forces due to bacterial growth lead to buckling instabilities on microscopic scales, which in turn result in higher orientational entropy. To minimize orientational entropy and continued compressive forces, and to maximize translational entropy, bacteria become laterally packed, providing more space to grow. This behavior can be explained by Onsager's theory for the isotropic-nematic transition of rodlike particles and supported by experimental approaches

Continement $l_d(\mu m)$		
2	0.168	0.169
3	0.203	0.224
	0.286	0.303

FIG. 6: Comparison of probability values for individual contact forces exceeding $|\vec{F}| = 13 \; \kappa P a \mu m^2$ in square and circle confinements across different division lengths. This table presents the probability values associated with individual contact forces surpassing 13 for varying division lengths l_d under square and circle confinement conditions.

[\[14\]](#page-6-9). In this theory, the effective interaction potential is proportional to the square length of the bacteria, denoted by L^2 . Therefore, the effective interaction potential is larger for colonies with larger division lengths, resulting in a more ordered configuration. Fig. [4](#page-3-1) shows this behavior, in which the order parameter for the larger division length is higher than that of the shorter division length [\[14\]](#page-6-9).

The interaction between bacteria, both with each other and the confinement, leads to bacterial deformation and dense lateral packing. To study the alignment of force interactions among contact bacteria relative to the bacterial aspect ratio, we analyzed the distribution of individual contact forces of bacteria. In Fig. [5](#page-3-2) we present the resulting individual contact force distribution $P(|F|)$, under circular and square-like confinements when the boxes are filled with bacteria.

We observe that all curves corresponding to the three division lengths exhibit a Gaussian force distribution for individual force values around $|F| = 13 kPa\mu m^2$ on a log-linear scale. This intersection point suggests that the above $|F| = 13 kPa\mu m^2$, all curves demonstrate a Gaussian force distribution. Integration of the surface area above this point indicates that the probability of encountering larger forces is greater for the colonies with larger division lengths Fig. [6.](#page-4-0)

This result provides strong evidence that the interaction potential scales with the length of the bacteria. As a result, bacteria with larger aspect ratios are more likely to experience higher forces.

In Fig. [7,](#page-4-1) we selectively highlight force magnitudes greater than $|F| \geq 13$ $kPa(\mu m)^2$ and map the center of mass of the contacting bacteria to visualize the force chain. The high-range force chains are predominantly distributed laterally, spanning bacteria positioned side by side or at small relative angles. Forces below 13 $kPa(\mu m)^2$ have been excluded from this figure, as they mainly correspond to interactions at the bacterial

FIG. 7: Force network for individual contact forces exceeding 13 $kPa(\mu m)^2$ (black lines), overlaid on the bacterial colony configuration for each division length under square and circular confinement. Panels (a-c) represent division lengths $l_d = [2, 3, 4] \mu m$ respectively in square-like confinement, while panels (d-f) represent the same division lengths in circle-like confinement. The color circle indicates the orientation of the bacteria.

tips. This distribution reflects a characteristic pattern of bacterial growth along their major axis, which allows for some relaxation of force interactions at the tips.

IV. Discussion

The research utilized molecular dynamics simulations to investigate the growth and self-organization of overdamped rod-shaped bacteria within square and circlelike confinements. We found that mechanical interactions within the system heavily influence the self-organization and growth of bacteria. Specifically, we observed that higher levels of mechanical interaction corresponded to the emergence of more structured patterns within the

bacterial colonies, accompanied by slower growth rates. Interestingly, we noted that the mechanism underlying the formation of these structured patterns varied significantly depending on the bacterial species under study and the geometry of the confinement [\[9,](#page-6-11) [11,](#page-6-4) [19,](#page-6-12) [20](#page-6-13)].

Consistent with earlier predictions, the magnitude of nematic ordering and orientational correlation function within a colony of elastic bacteria can be influenced by various factors. One such factor is the intrinsic characteristics of the bacteria, such as their aspect ratio. Our findings corroborate this notion, as we observed that colonies composed of bacteria with a larger aspect ratio exhibited higher levels of nematic ordering and orientational correlation function.

Another influential factor is the geometry of the confinements. As the confinement becomes densely populated with bacteria at the final step, we observe slight variations in order. By its distinct edges and corners, square confinement promotes preferred directions, leading to higher ordering than circular confinement. Thus, the collective influence of bacteria's physical and geometrical properties and the confinement contributes to the overall degree of organization within the colony.

To delve into the fundamental differences between colonies with larger and smaller division lengths for both confinements, we examined a specific time step during which all colonies were freely growing. This allowed us to isolate and analyze the effects of division length on the emergent ordering within the colonies. We identify individual local islands based on a criterion that requires the relative angle between contact bacteria to be less than or equal to 10 degrees.

The probability distribution for local scalar order parameter indicates that the probability of higher nematic ordering is greater for colonies with larger division lengths. Certain characteristics, such as the number of bacteria in each island and the number of potential contacts for each division length, exhibit similar behavior.

Hence, the greater nematic ordering observed in our simulations stems from increased force interactions, leading to a higher degree of overlap between longer rods. This overlap is crucial for overcoming orientational entropy and reducing gaps, particularly between long bacteria, thereby facilitating sufficient space for bacterial growth.

We selected a specific threshold value for individual contact forces, ensuring all curves exhibit a Gaussian distribution. By integrating the probability values derived from these distributions, we arrived at a noteworthy finding. It became evident that as compression occurs, there is a higher probability of encountering greater forces in colonies with larger division lengths in contrast to colonies with shorter division lengths. This observation underscores the influence of compression on force distributions and sheds light on how division length impacts force magnitudes within the bacterial colony.

In addition to analyzing the probability distribution of individual contact forces within the colony, we conducted a thorough investigation into the spatial distribution of higher forces and their impact on the cellular configuration. This analysis aimed to ascertain whether higher forces predominantly originated from end-to-end interactions, end-to-lateral sides, or lateral-lateral sides between bacteria. Our findings revealed that the elevated values of individual contact forces primarily manifested between the lateral-lateral sides of the bacteria. This phenomenon can be attributed to the active growth of bacteria along their major axis. As elastic materials, bacteria tend to reduce forces along the major axis, resulting in smaller contact forces compared to those along the lateral sides.

Our work revealed that mechanical interactions due to bacterial growth play a crucial role in constraining cell growth within confined environments, influencing both self-organization and growth dynamics of bacterial colonies. We found that division length significantly impacts the degree of ordering within bacterial colonies, with larger division lengths correlating with higher levels of nematic ordering and correlation function. The geometry of confinements, particularly square confinement, was observed to foster preferred directions and result in higher ordering compared to circular confinement. Analysis of force distributions highlighted the influence of compression on force magnitudes within bacterial colonies, with colonies of larger division lengths exhibiting a higher probability of encountering greater forces. Our investigation into the spatial distribution of higher forces revealed that elevated contact forces primarily occur between the lateral-lateral sides of bacteria, influenced by the active growth of bacteria along their major axis. Additionally, previous experiments and simulations have explored the buckling instability in growing non-motile bacterial colonies, particularly focusing on Escherichia coli (E. coli). This instability, attributed to the anisotropy of the stress tensor within the colony, limits the orientational order in large bacterial colonies and leads to denser lateral packing of the bacteria [\[22](#page-6-14)].

By pursuing these directions, future studies can deepen our understanding of the complex interplay between mechanical interactions and bacterial behavior within confined environments, with potential implications for diverse fields ranging from microbial ecology to biomaterials science.

^[1] Jenna A Moore-OttSelena ChiuDaniel B AmchinTapomoy BhattacharjeeSujit S Datta (2022) A biophysical threshold for biofilm formation eLife 11:e76380.

^[2] Rosalind J Allen and Bartlomiej Waclaw 2019 *Rep. Prog. Phys.* 82 016601.

^[3] R. Phillips, J. Kondev, J. Theriot, H. Garcia, *Physical Biology of the Cell* (2nd edition, Garland Science, 2012).

- [4] Luís André de Almeida Campos, João Victor de Oliveira Santos, Iago Dillion Lima Cavalcanti, Azael Francisco Silva Neto, Mariane Cajubá de Britto Lira Nogueira, Nereide Stela Santos-Magalhães, Isabella Macário Ferro Cavalcanti, Natural products and combination therapy as strategies against bacterial biofilm, Studies in Natural Products Chemistry, Elsevier,77(2023).
- [5] Crouzet, M., Le Senechal, C., Brözel, V.S. et al. Exploring early steps in biofilm formation: set-up of an experimental system for molecular studies. BMC Microbiol 14, 253 (2014).
- [6] Isensee Jonas, Hupe Lukas, Golestanian Ramin and Bittihn Philip. 2022 Stress anisotropy in confined populations of growing rods. *J. R. Soc. Interface*. 19: 20220512.
- [7] You Z, Pearce DJG, Sengupta A, Giomi L. Geometry and Mechanics of Microdomains in Growing Bacterial Colonies. Physical Review X. 2018;8(3):31065.
- [8] DeCamp, S., Redner, G., Baskaran, A. et al. Orientational order of motile defects in active nematics. *Nature Mater* 14, 1110–1115 (2015).
- [9] D. Volfson, S. Cookson, J. Hasty, L. S. Tsimring, Biomechanical ordering of dense cell populations. *Proc. Natl. Acad. Sci. U.S.A.* 105, 15346–15351 (2008).
- [10] You Z, Pearce DJG, Giomi L. Confinement-induced selforganization in growing bacterial colonies. *Sci. Adv.* 7, eabc8685(2021).
- [11] P. Kumar, A. Libchar, *Pressure and temperature dependence of growth and morphology of Escherichia coli: Experiments and stochastic model,* Biophys J. 105, 783 (2013).
- [12] Si, F., Li, B., Margolin, W. et al. Bacterial growth and form under mechanical compression. *Sci Rep* 5, 11367

(2015).

- [13] Tuson, H.H., Auer, G.K., Renner, L.D., Hasebe, M., Tropini, C., Salick, M., Crone, W.C., Gopinathan, A., Huang, K.C. and Weibel, D.B. (2012), Measuring the stiffness of bacterial cells from growth rates in hydrogels of tunable elasticity. Molecular Microbiology, 84: 874- 891.
- [14] M. Doi, *Soft Matter Physics* (OUP Oxford, 2013).
- [15] Duvernoy, MC., Mora, T., Ardré, M. et al. Asymmetric adhesion of rod-shaped bacteria controls microcolony morphogenesis. *Nat Commun* 9, 1120(2018).
- [16] P. Ghosh, J. Mondal, E. Ben-Jacob, H.Levine. Mechanically-driven phase separation in a growing bacterial colony. *Proc. Natl. Acad. Sci. U.S.A.* 112(17), E2166- 732015 (2015).
- [17] H. A. Makse, D. L. Johnson, and Lawrence M. Schwartz, Phys. Rev. Lett. 84, 4160 (2000).
- [18] Daniel L. Blair, T. Neicu, and A. Kudrolli.*Phys. Rev. E* 67, 031303 (2003).
- [19] B. L. Shraiman, Mechanical feedback as a possible regulator of tissue growth. *Proc. Natl. Acad. Sci. U.S.A.* 102, 3318–3323 (2005).
- [20] Dell'Arciprete, D., Blow, M.L., Brown, A.T. et al. A growing bacterial colony in two dimensions as an active nematic. *Nat Commun* 9, 4190 (2018).
- [21] Cho H, Jönsson H, Campbell K, Melke P, Williams JW, et al. (2007) Self-Organization in High-Density Bacterial Colonies: Efficient Crowd Control. PLOS Biology 5(11).
- [22] D. Boyer, W. Mather, O. Mondragón-Palomino, S. OrozcoFuentes, T. Danino, J. Hasty and L. S. Tsimring, Buckling instability in ordered bacterial colonies, *Phys. Biol*., 2011, 8(2), 026008.