

Dissipative ground state preparation in *ab initio* electronic structure theory

Hao-En Li,¹ Yongtao Zhan,² and Lin Lin^{3,4,*}

¹*Department of Chemistry, Tsinghua University, Beijing 100084, China*

²*Institute of Quantum Information and Matter, California Institute of Technology, Pasadena, California 91125, USA*

³*Department of Mathematics, University of California, Berkeley, California 94720, USA*

⁴*Applied Mathematics and Computational Research Division,
Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

Dissipative engineering is a powerful tool for quantum state preparation, and has drawn significant attention in quantum algorithms and quantum many-body physics in recent years. In this work, we introduce a novel approach using the Lindblad dynamics to efficiently prepare the ground state for general *ab initio* electronic structure problems on quantum computers, without variational parameters. These problems often involve Hamiltonians that lack geometric locality or sparsity structures, which we address by proposing two generic types of jump operators for the Lindblad dynamics. Type-I jump operators break the particle number symmetry and should be simulated in the Fock space. Type-II jump operators preserve the particle number symmetry and can be simulated more efficiently in the full configuration interaction space. For both types of jump operators, we prove that in a simplified Hartree-Fock framework, the spectral gap of our Lindbladian is lower bounded by a *universal* constant. For physical observables such as energy and reduced density matrices, the convergence rate of our Lindblad dynamics with Type-I jump operators remains universal, while the convergence rate with Type-II jump operators only depends on coarse grained information such as the number of orbitals and the number of electrons. To validate our approach, we employ a Monte Carlo trajectory-based algorithm for simulating the Lindblad dynamics for full *ab initio* Hamiltonians, demonstrating its effectiveness on molecular systems amenable to exact wavefunction treatment.

I. INTRODUCTION

Quantum state preparation is a fundamental task in quantum simulation and quantum algorithm design [1–6]. While eigenstates of a Hamiltonian can, in principle, be prepared using quantum phase estimation (QPE) and its variants, these algorithms themselves often require an initial state that has a significant overlap with the target state [7–15]. Dissipative state engineering offers a very different perspective on this problem. Rather than treating dissipation as a source of decoherence due to system-environment coupling, properly designed dissipative dynamics, such as those governed by the Lindblad equation, can encode a wide variety of strongly correlated states as the steady states of a dynamical process. Dissipative techniques, and state preparation techniques using mid-circuit measurements in general, have been widely employed in preparing matrix product states, ground states of stabilizer codes, spin systems, and other states exhibiting long-range entanglement [2, 16–22]. There has been also growing recent interest in using Lindblad dynamics as an algorithmic tool for thermal state and ground state preparation [23–31]. However, many applications have focused on Hamiltonians with special structures, as the dissipative terms often need to be carefully engineered based on the special properties of the Hamiltonian. In contrast, in quantum chemistry and materials science, *ab initio* Hamiltonians lack specific geometric locality or sparsity, which significantly complicates the design of dis-

sipative terms.

In this work, we overcome this difficulty and present a novel method for using Lindblad dynamics to efficiently prepare the ground state for general *ab initio* electronic structure problems on quantum computers. Our approach builds upon recent developments in quantum ground state preparation [25], which has the advantage of being applicable to both commuting and non-commuting Hamiltonians on an equal basis. Unlike [25] which prepares the ground state using a single jump operator together with a coherent term, we propose two sets of Lindblad jump operators, termed Type-I and Type-II. Each set contains $\text{poly}(L)$ jump operators (L is the number of spatial orbitals), which are agnostic to chemical details and thus can readily be applied to *ab initio* Hamiltonians with unstructured and long range coefficients. The process does not involve variational parameters. Type-I Lindblad dynamics break the particle-number symmetry and must be simulated in the Fock space. In contrast, Type-II jump operators preserve the particle number, allowing for more efficient simulation (on both classical and quantum computers) in the full configuration interaction (FCI) space.

The efficiency of Lindblad dynamics for quantum state preparation is quantified by the mixing time, which is the time required to reach the target steady state to a certain precision from an *arbitrary* initial state [32, 33]. Theoretical analysis of the mixing time is in general a challenging task, and is often feasible only for specific systems, parameter regimes, or simplified settings [34–39]. Our strategy is to first theoretically analyze the spectral gap of the Lindbladian, as well as dynamics of observables, such as the energy and the reduced density

* linlin@math.berkeley.edu

matrix (RDM), within a simplified Hartree-Fock (HF) framework. In this setting, the combined action of the jump operators effectively implements a classical Markov chain Monte Carlo within the molecular orbital basis. We prove that the convergence can be provably agnostic to specific chemical details, and in some cases, the convergence rate can be universal. We then perform numerical simulations to examine the transferability of this behavior to the full *ab initio* Hamiltonian, using an approach based on unraveled Lindblad dynamics. This is applied to molecular systems such as BeH₂, H₂O, and Cl₂, which are amenable to exact wavefunction treatment within the FCI space. We also apply our method to investigate the stretched square H₄ system, which has nearly degenerate low-energy states and poses challenges for highly accurate quantum chemistry methods such as CCSD(T). In all cases, the Lindblad dynamics can prepare a quantum state with an energy that achieves chemical accuracy, even in the strongly correlated regime.

II. RESULTS

A. Dissipation engineering for ground state preparation

We consider Lindblad dynamics of the form:

$$\begin{aligned} \frac{d}{dt}\rho &= \mathcal{L}[\rho] = \mathcal{L}_H[\rho] + \mathcal{L}_K[\rho] \\ &= -i[\hat{H}, \rho] + \sum_k \hat{K}_k \rho \hat{K}_k^\dagger - \frac{1}{2} \{ \hat{K}_k^\dagger \hat{K}_k, \rho \}. \end{aligned} \quad (1)$$

For ground state preparation, the key object in Ref. [25] is the following the jump operator:

$$\hat{K}_k = \sum_{i,j} \hat{f}(\lambda_i - \lambda_j) \langle \psi_i | A_k | \psi_j \rangle | \psi_i \rangle \langle \psi_j |. \quad (2)$$

Here, $\{A_k\}$ are called (primitive) coupling operators, whose selection will be discussed in detail later. Each jump operator \hat{K}_k is derived by reweighting A_k in the eigenbasis $\{|\psi_i\rangle\}_{i=0}^{N-1}$ of the Hamiltonian \hat{H} by a filter function $\hat{f}(\omega)$ evaluated at the energy difference $\lambda_i - \lambda_j$. The filter function $\hat{f}(\omega)$ is only supported on the negative axis. In other words, $\hat{f}(\lambda_i - \lambda_j) = 0$ for any $i \geq j$ (assuming $\lambda_0 < \lambda_1 \leq \dots \leq \lambda_{N-1}$). As a result, the jump operator \hat{K}_k only allows transitions between the eigenvectors of \hat{H} that lower the energy. Since the Lindblad dynamics generates a completely positive, trace preserving (CPTP) map [40–42], $\langle \psi_i | \rho(t) | \psi_i \rangle$ is a normalized probability distribution at time any t . In the energy basis, the dynamics continuously “shovels” high-energy states towards low-energy ones, eventually reaching the ground state, as shown in Figure 1. Furthermore, we can easily verify $\hat{K}_k |\psi_0\rangle = 0$ since $\hat{f}(\lambda_j - \lambda_0) = 0$ for any $j \geq 0$. This immediately suggests that the ground state $|\psi_0\rangle \langle \psi_0|$ is a stationary point of the Lindblad dynamics.

This dynamics is ergodic if the ground state is the *only* stationary point, and this can be achieved by a carefully chosen set of coupling operators $\{A_k\}$.

The shoveling process shares some similarities with various forms of imaginary time evolution (ITE) [3, 43–49], though with notable differences. A direct implementation of ITE through the application of $e^{-\tau\hat{H}}$ does not lead to a completely positive trace-preserving (CPTP) map, and its quantum implementation again requires nontrivial overlaps between the initial and target states. The quantum imaginary time evolution (QITE) algorithm [3] addresses some of these challenges but introduces a process tomography procedure, whose analysis can be highly nontrivial.

At first glance, it may seem that constructing the jump operator in Eq. (2) requires diagonalizing \hat{H} , which would clearly defeat the purpose. However, we can reformulate the definition of \hat{K}_k in the time domain as

$$\hat{K}_k = \int_{\mathbb{R}} f(s) A_k(s) ds, \quad (3)$$

where $A_k(s) = e^{i\hat{H}s} A_k e^{-i\hat{H}s}$ is the Heisenberg evolution of A_k , and $f(s) = \frac{1}{2\pi} \int_{\mathbb{R}} \hat{f}(\omega) e^{-i\omega s} d\omega$ is the inverse Fourier transform of the filter function \hat{f} in the frequency domain.

In this form, the construction of the jump operator can be achieved using standard Trotter expansions for digitally simulating the Hamiltonian evolution. Additionally, the Trotter expansion can also be applied to simulate the Lindblad dynamics in Eq. (1). The choice of the filter function depends only on coarse-grained properties of the system, such as estimates of the spectral radius and the spectral gap of \hat{H} . A brief review of the selection of the filter function and the quantum simulation algorithm is provided in the Supplementary Information (SI).

B. Type-I and Type-II jump operators for *ab initio* calculations

Unlike lattice problems, the second quantized Hamiltonians in *ab initio* electronic structure calculations do not have clean forms such as nearest-neighbor interactions. Therefore it is important to choose simple and yet effective set of coupling operators $\{A_k\}$ that are easy to implement and allow the system to converge rapidly towards the ground state. In this work, we introduce two simple sets of primitive coupling operators, referred to as Type-I and Type-II, respectively. The corresponding jump operators can be constructed according to Eq. (3).

We choose the set of Type-I coupling operators to be $\mathcal{A}_I = \{a_i^\dagger \mid i = 1, 2, \dots, 2L\} \cup \{a_i \mid i = 1, 2, \dots, 2L\}$. This includes all of the $4L$ (counting spatial and spin degrees of freedom) fermionic creation and annihilation operators. Each operator can be expressed in the atomic orbital basis, molecular orbital basis, or some other basis sets. These different choices differ by a unitary matrix.




FIG. 1. A conceptual illustration of the “shoveling” process in ground state preparation via Lindbladians. The choice of jump operators ensures that the Lindbladian only allows transitions from high-energy eigenstates to low-energy eigenstates.

Given the linear relationship between the jump operator and the coupling operator, a unitary rotation of the coupling operators will correspondingly induce a unitary rotation of the jump operators. This unitary rotation can be viewed as a gauge degree of freedom. Ideally the numerical result should be independent of this gauge. We will verify that this is indeed the case in a simplified Hartree-Fock setting.

The set of Type-II coupling operators is $\mathcal{A}_{\text{II}} = \{a_i^\dagger a_j \mid i, j = 1, 2, \dots, 2L\}$ which includes every fermionic creation and annihilation pairs, and has $4L^2$ elements in total. Most Hamiltonians in *ab initio* electronic structure calculations are particle number preserving. Unlike Type-I coupling operators which break particle number symmetries and must be simulated in the Fock space, Type-II coupling operators (and the corresponding jump operators) preserve particle number symmetries. The corresponding Lindblad dynamics can be simulated in the full configuration interaction (FCI) space. Note that the dimension of density matrix in the Fock formulation is 4^{2L} , and that of the FCI space is $\binom{2L}{N_e}^2$, where N_e is the number of electrons. The difference becomes particularly significant when N_e is very small or large, such as simulating alkali metals and halogen elements in a small basis set. The particle number symmetry can be used also reduce the cost of quantum simulations, such as the Trotter error for Hamiltonian simulation $e^{-i\hat{H}t}$ [50, 51], or the block encoding subnormalization factors of the Hamiltonian [52, 53].

Both Type-I and Type-II sets are “bulk” coupling operators, meaning that dissipation is introduced on every (atomic or molecular) basis function. As will be seen below, this can be very effective in reducing the mixing time. On the other hand, this comes at the cost of introducing a large number of jump operators and add simulation cost, both on quantum computers [25] and in classical simulation. We will also discuss how to reduce the number using active space ideas.

C. Universal fast convergence with Type-I set in Hartree-Fock theory

We first consider the ground state preparation via Lindbladians at the HF level before moving on to the interacting regime. We refer readers to the SI for a brief review of the Hartree-Fock theory. Essentially, after self-consistency is reached, all the information of the Hartree-Fock theory is encoded in a non-interacting Hamiltonian $\hat{H} = \sum_{p,q=1}^{2L} F_{pq} a_p^\dagger a_q$, where the Hermitian matrix F is called the Fock matrix. Let Φ be the unitary matrix that diagonalizes F , then the new basis set, known as molecular orbitals, is obtained by transforming the atomic orbitals using Φ . For such Hamiltonians, the information contained in the many-body density operator ρ is entirely stored in the one-particle reduced density matrix (1-RDM), defined as

$$P_{ij} = \text{Tr}\left(\rho a_j^\dagger a_i\right), \quad 1 \leq i, j \leq 2L. \quad (4)$$

According to the Thouless theorem [54, 55],

$$e^{\hat{H}} a_i^\dagger = \sum_p a_p^\dagger (e^F)_{pi} e^{\hat{H}}, \quad e^{\hat{H}} a_i = \sum_q (e^{-F})_{iq} a_q e^{\hat{H}}. \quad (5)$$

Therefore, we have

$$\begin{aligned} \hat{K}_{p,+} &= \int_{\mathbb{R}} f(s) e^{i\hat{H}s} a_p^\dagger e^{-i\hat{H}s} ds \\ &= \int_{\mathbb{R}} f(s) \sum_{r=1}^{2L} a_r^\dagger (e^{iFs})_{r,p} e^{i\hat{H}s} e^{-i\hat{H}s} ds \\ &= \sum_{r=1}^{2L} a_r^\dagger (\hat{f}(F))_{r,p} \end{aligned} \quad (6)$$

and similarly

$$\begin{aligned} \hat{K}_{q,-} &= \int_{\mathbb{R}} f(s) e^{i\hat{H}s} a_q e^{-i\hat{H}s} ds \\ &= \int_{\mathbb{R}} f(s) \sum_{r=1}^{2L} (e^{-iFs})_{q,r} a_r e^{i\hat{H}s} e^{-i\hat{H}s} ds \\ &= \sum_{r=1}^{2L} a_r (\hat{f}(-F))_{q,r}. \end{aligned} \quad (7)$$

This implies that for the Type-I set, the jump operators are all linear in fermionic creation and annihilation operators. The corresponding Lindblad dynamics is quasi-free [56–59], and we can derive a closed form equation of motion for the 1-RDM

$$\partial_t P(t) = -i[F, P(t)] + B - \frac{1}{2}[P(t)(B+C) + (B+C)P(t)]. \quad (8)$$

Here

$$B = \hat{f}(F)\hat{f}(F)^\dagger = \hat{f}^2(F), \quad C = \hat{f}(-F)\hat{f}(-F)^\dagger = \hat{f}^2(-F), \quad (9)$$

and we use the fact that the Fock matrix F is Hermitian. A detailed derivation of Eq. (8) can be found in the SI. For any filter function \hat{f} satisfying $\hat{f}(\omega) = 1$ on $[-2\|\hat{H}\|_2, -\Delta]$ and $\hat{f}(\omega) = 0$ on $[0, +\infty)$, we have $B + C = \hat{f}^2(F) + \hat{f}^2(-F) = \mathbf{1}$, where $\mathbf{1}$ is the identity matrix. The equation of motion Eq. (8) then takes a very simple form

$$\partial_t P(t) = -i[F, P(t)] - P(t) + \hat{f}(F). \quad (10)$$

In particular, if we perform a unitary rotation of the fermionic creation and annihilation operators used to define the primitive coupling operators, this amounts to a gauge choice, and the final equation of motion (10) is gauge-invariant.

From Eq.(10) we can easily see that $P^* = \hat{f}(F)$ is the unique stationary point. In fact, let $P(t)$ and $P'(t)$ be the solution to solve Eq. (10) with different initial values $P(0)$ and $P'(0)$, then

$$\|P(t) - P'(t)\|_F = e^{-t}\|P(0) - P'(0)\|_F, \quad (11)$$

where $\|A\|_F := \sqrt{\text{Tr}(A^\dagger A)}$ denotes the Frobenius norm of matrices. The detailed derivation of Eq. (11) is provided in the SI.

We may get more insight by rewriting Eq. (10) in the energy basis. Specifically, we define $\tilde{P} = \Phi^\dagger P \Phi = \left(\text{Tr}(\rho c_j^\dagger c_i)\right)_{1 \leq i, j \leq 2L}$, where Φ is the coefficient matrix of the molecular orbitals. Then we have

$$\partial_t \tilde{P}(t) = -i[\Lambda, \tilde{P}(t)] - \tilde{P}(t) + \hat{f}(\Lambda). \quad (12)$$

Here we use $F\Lambda = \Lambda\Phi$ and $\Lambda = \text{diag}(\varepsilon_1, \dots, \varepsilon_{2L})$ with $\varepsilon_1 \leq \dots \leq \varepsilon_{N_e} \leq 0 < \varepsilon_{N_e+1} \leq \dots \leq \varepsilon_{2L}$. Therefore the stationary point \tilde{P}^* is given by $\tilde{P}^* = \hat{f}(\Lambda) = \text{diag}(\underbrace{1, \dots, 1}_{N_e}, \underbrace{0, \dots, 0}_{2L-N_e})$ which is consistent with the *aufbau principle*, and this is achieved without explicitly diagonalizing the Fock matrix. In particular, the diagonal elements of $\tilde{P}(t)$ evolves as follows:

$$\partial_t \langle n_i \rangle = \left(\tilde{P}(t)\right)_{ii} = \begin{cases} -\langle n_i \rangle + 1, & i = 1, \dots, N_e, \\ -\langle n_i \rangle, & i = N_e + 1, \dots, 2L. \end{cases} \quad (13)$$

Assume the initial occupation numbers of the molecular orbitals (i.e. the diagonal elements of the initial 1-RDM $\tilde{P}(0)$) is given by $n_i(0)$, then

$$\langle n_i \rangle = \begin{cases} 1 - (1 - n_i(0))e^{-t}, & i = 1, \dots, N_e, \\ n_i(0)e^{-t}, & i = N_e + 1, \dots, 2L. \end{cases} \quad (14)$$

Therefore in the energy basis, the Lindblad dynamics with Type-I jump operators can drive the occupation numbers of the lowest N_e molecular orbitals to approach 1 exponentially, while the occupation numbers of the remaining $2L - N_e$ high-energy molecular orbitals exponentially approach 0 (see Figure 2). The convergence rate is *universal* and is independent of any chemical details or initial starting point. A numerical validation of this statement is provided in Section III F.




FIG. 2. A conceptual illustration of the evolution of the diagonal elements of the 1-RDM under linear bulk dissipation for ground state preparation. The occupation numbers in each molecular orbital increase or decrease independently in an exponential rate.

D. Convergence with Type-II set oblivious to chemical details in Hartree-Fock theory

Let us now carry out the calculation for Type-II jump operators in the Hartree-Fock setting. Recall $\hat{f}(\omega) = 1$ on $[-2\|\hat{H}\|_2, -\Delta]$ and $\hat{f}(\omega) = 0$ on $[0, +\infty)$, using the Thouless theorem, the jump operators satisfy

$$\begin{aligned} \hat{K}_{ij} &= \int_{\mathbb{R}} f(s) e^{i\hat{H}s} a_i^\dagger a_j e^{-i\hat{H}s} ds \\ &= \sum_{p,q=1}^{2L} \hat{f}(\varepsilon_p - \varepsilon_q) c_p^\dagger c_q \Phi_{ip}^* \Phi_{jq} = \sum_{p < q} c_p^\dagger c_q \Phi_{ip}^* \Phi_{jq}. \end{aligned} \quad (15)$$

Note that \hat{K}_{ij} is a quadratic operator in the fermionic operators and not Hermitian. This means that the Lindblad dynamics is not quasi-free, and the 1-RDM cannot

satisfy a closed form equation of motion [59]. Despite this, we demonstrate below an explicit description of the dynamics in the energy basis. For the coherent part,

$$\mathcal{L}_H^\dagger(c_r^\dagger c_s) = i[\hat{H}, c_r^\dagger c_s] = i(\varepsilon_r - \varepsilon_s)c_r^\dagger c_s. \quad (16)$$

Here \mathcal{L}_H^\dagger denotes the adjoint of the superoperator \mathcal{L}_H with respect to the Hilbert-Schmidt inner product. Moreover, using that Φ consists of orthonormal columns, we find

$$\begin{aligned} \sum_{i,j=1}^{2L} \hat{K}_{ij}^\dagger \hat{K}_{ij} &= \sum_{ij} \sum_{p<q} \sum_{r<s} c_q^\dagger c_p c_r^\dagger c_s \Phi_{ip} \Phi_{jq}^* \Phi_{ir}^* \Phi_{js} \\ &= \sum_{p<q} c_q^\dagger c_p c_p^\dagger c_q = \sum_{p<q} (1 - n_p) n_q. \end{aligned} \quad (17)$$

Similarly, for $r \neq s$,

$$\begin{aligned} \mathcal{L}_K^\dagger(c_r^\dagger c_s) &= \sum_{ij} \hat{K}_{ij}^\dagger c_r^\dagger c_s \hat{K}_{ij} - \frac{1}{2} \{ \hat{K}_{ij}^\dagger \hat{K}_{ij}, c_r^\dagger c_s \} \\ &= -\frac{1}{2} (M_r + M_s + 1) c_r^\dagger c_s. \end{aligned} \quad (18)$$

Here

$$M_k = \sum_{p<k} c_p c_p^\dagger + \sum_{q>k} c_q^\dagger c_q = \sum_{p<k} (1 - n_p) + \sum_{q>k} n_q. \quad (19)$$

For $r = s$,

$$\mathcal{L}_K^\dagger(c_r^\dagger c_r) = \sum_{q>r} (1 - n_r) n_q - \sum_{p<r} (1 - n_p) n_r. \quad (20)$$

In all the expressions above, the operators occurring in the Lindblad dynamics are all invariant to the gauge choice in the primitive coupling operators, and can all be expressed using simple operators in molecular orbitals. For a detailed derivation of the above equations, interested readers can refer to the SI.

Consider the 1-RDM in the molecular orbital basis $\tilde{P} = (\text{Tr}(c_r^\dagger c_s \rho))_{1 \leq s, r \leq 2L} = \Phi^\dagger P \Phi$. Then the equation of motion of the entries of \tilde{P} depends on that of the 2-RDM. Specifically, for the off-diagonal elements,

$$\partial_t \tilde{P}_{sr} = \begin{cases} -i(\varepsilon_s - \varepsilon_r) \tilde{P}_{sr} - \frac{1}{2} \langle (M_r + M_s + 1) c_r^\dagger c_s \rangle & r < s, \\ -i(\varepsilon_s - \varepsilon_r) \tilde{P}_{sr} - \frac{1}{2} \langle c_r^\dagger c_s (M_r + M_s + 1) \rangle & r > s. \end{cases} \quad (21)$$

For the diagonal elements,

$$\begin{aligned} \partial_t \tilde{P}_{rr} &= -\sum_{p<r} \tilde{P}_{pp} + \sum_{q>r} \tilde{P}_{qq} + \sum_{p<r} \langle n_p n_r \rangle - \sum_{q>r} \langle n_r n_q \rangle \\ &= -\sum_{p<r} \langle (1 - n_p) n_r \rangle + \sum_{q>r} \langle (1 - n_r) n_q \rangle \\ &= -\langle M_r n_r \rangle + \sum_{q>r} \langle n_q \rangle. \end{aligned} \quad (22)$$

Further derivations of the equation of motion for the 2-RDM will lead to the 3-RDM, and so forth. It resembles the renowned Bogoliubov–Born–Green–Kirkwood–Yvon (BBGKY) hierarchy [60]. To make the system solvable, we can truncate the equations by neglecting the higher-order moment terms. At this point, if we consider these matrix elements as random processes, then the equations of motion describe a *classical* continuous-time Markov chain, with the stationary distribution 1-RDM approximately given by $\tilde{P}^* = \text{diag}(\underbrace{1, \dots, 1}_{N_e}, \underbrace{0, \dots, 0}_{2L - N_e})$ according to the *aufbau principle*. Nonetheless, from Eq. (21) and Eq. (22), it is evident that the evolution of the RDMs are oblivious to chemical details, and are solely determined by the number of orbitals L and the number of electrons N_e . Moreover, the dynamics of the diagonal entries is independent of that of the off-diagonal entries. For an intuitive understanding of this, it resembles a “mass transport” process from higher energy orbitals to lower energy orbitals. Therefore, the change in occupation number of each orbital is influenced by the electronic population of other orbitals, leading to the appearance of 2-RDM related terms in the equations (see Figure 3).




FIG. 3. A conceptual illustration of the evolution of the diagonal elements of the 1-RDM under quadratic bulk dissipation for ground state preparation, which is a “mass transport” process from higher energy orbitals to lower energy orbitals.

For the diagonal entries, applying the mean-field approximation on Eq. (22) gives:

$$\partial_t \tilde{P}_{rr} \approx -\langle M_r \rangle \tilde{P}_{rr} + \sum_{q>r} \tilde{P}_{qq}. \quad (23)$$

Note that $M_k \geq 1$ for all $r < N_e$ and $r > N_e + 1$, the convergence rates of the off-diagonal 1-RDM entries are exponential with the exponent of at least 1. The case $r \in \{N_e, N_e + 1\}$ is beyond the mean-field analysis. These convergence rate arguments can be verified numerically (see Section III F). A detailed explanation of the results above can be found in the SI.

E. Spectral gap of Lindbladian in Hartree-Fock theory

In this section, we provide a direct characterization of the spectral gap of the Lindbladian. Consider the following Hamiltonian, which corresponds to the dissipative part of the Lindblad dynamics.

$$\hat{H}_{\text{dp}} = \frac{1}{2} \sum_k \hat{K}_k^\dagger \hat{K}_k, \quad (24)$$

Since for every jump operator $\hat{K}_k |\psi_0\rangle = 0$ by construction, \hat{H}_{dp} can be viewed as a frustration-free parent Hamiltonian of the ground state $|\psi_0\rangle$. We have the following theorem (the proof is given in the SI).

Theorem 1. *If $[\hat{H}, \hat{H}_{\text{dp}}] = 0$, then the spectral gap of the Lindbladian \mathcal{L} is equal to the gap of \hat{H}_{dp} .*

In the Hartree-Fock setting, we have established that the equation of motion of the 1-RDM in the molecular orbital basis is independent of chemical details. Now we prove a stronger result, which states that the spectral gap of the Lindbladian is rigorously bounded from below using Type-I and Type-II jump operators in Hartree-Fock theory.

For Type-I jump operators, by calculating \hat{H}_{dp} in the molecular orbital basis (see the SI for details), we obtain

$$\hat{H}_{\text{dp}} = \frac{1}{2} \sum_{p \leq N_e} (1 - n_p) + \frac{1}{2} \sum_{q > N_e} n_q, \quad (25)$$

where n_p, n_q are number operators and commute with \hat{H} , which is diagonal in the molecular orbital basis. Applying Theorem 1, we find that the spectral gap of the Lindbladian is the same as the gap of \hat{H}_{dp} , which is equal to $\frac{1}{2}$.

For Type-II jump operators, from previous calculations,

$$\hat{H}_{\text{dp}} = \frac{1}{2} \sum_{p < q} (1 - n_p) n_q. \quad (26)$$

which again only consists of number operators. Applying Theorem 1, we find that the spectral gap of the Lindbladian is also equal to $\frac{1}{2}$.

For thermal state preparation, an estimate of the spectral gap provides a direct upper bound on the mixing time of the Lindblad dynamics in trace distance [32]. This relationship, however, assumes that the stationary state is invertible, which is not satisfied by the ground state density matrix. In practice, we observe that the convergence rate of observables aligns closely with the spectral gap analysis and does not exhibit dependence on system size.

F. Numerical verification in the Hartree-Fock setting

Given that the equations of motion of the entries of the 1-RDM are not closed, it is more challenging to analyze the convergence rate of the Type-II settings. Nonetheless, we may provide a qualitative estimation of the convergence rate using mean-field approximation. Let us first focus on the diagonal elements. For both $r > s$ and $r < s$, applying the mean-field approximation to Eq. (21) results in the following linear homogeneous equation:

$$\partial_t \tilde{P}_{sr} \approx \left[-\frac{1}{2}(1 + \langle M_r + M_s \rangle) + i(\varepsilon_r - \varepsilon_s) \right] \tilde{P}_{sr}. \quad (27)$$

So under the mean-field approximation, the off-diagonal entry \tilde{P}_{sr} converges exponentially to zero with an exponent of at least $\frac{1}{2}$, since M_k is positive semidefinite for any k .


In this section, we numerically verify the convergence rate of observables such as energy and 1-RDM at the HF level. Using the Type-I set in Section II C, Figure 4 (a) shows that the convergence of the energy towards the ground state energy follows a universal relation $\exp(-t)$, for any molecule in any basis set. This is because the effect of the collection action of the jump operators is to *independently* adjust the occupation number of each molecular orbital, until the *aufbau principle* is reached.

Figure 4 (b) shows the convergence rate of the energy using the Type-II set in Section II D. The test systems are F_2 (1.4 Å), LiH (1.546 Å), chain H_4 (2.0 Å) and square H_4 (2.0 Å) using STO-3G and STO-6G basis sets. For each molecule, STO-3G and STO-6G have the same number of orbitals and electrons. The two isomers of H_4 also share the same number of orbitals and electrons. We observe that the convergence of the systems with the same L and N_e are exactly identical up to renormalization, but it varies across those molecules with different numbers, indicating a nontrivial dependence on both L and N_e .

To further examine the convergence rate, we track the evolution of the diagonal and off-diagonal entries of the 1-RDM for the H_4 within STO-3G, as shown in Figure 5. From Figure 5 (a) and (b), we see that the convergence rates of the diagonal entries are faster than $\exp(-t)$ except for the highest occupied molecular orbital (HOMO) and the lowest unoccupied molecular orbital (LUMO). Additionally, the off-diagonal entries exhibit convergence exponents of at least $\frac{1}{2}$. All of the numerical results agree very well with the mean-field analysis discussed in Section II D.

G. Transferability to full *ab initio* calculations

Both Type-I and Type-II coupling operators can be readily applied in full *ab initio* calculations. However, the jump operators are no longer linear and quadratic fermionic operators. For small-sized systems up to 12



(a) Convergence of energy using the Type-I set for Hartree-Fock state preparation of the molecules H₄, LiH, H₂O, CH₄, HCN, C₂H₄, N₂, H₁₀ and SO₃. The y -axis is displayed on a logarithmic scale. We perform each simulation by propagating the 1-RDM according to Eq. (8) using the DOPRI5 solver. The convergence rate is universal.

(b) Convergence of energy using the Type-II set for F₂, LiH and H₄ molecules (square and chain) in STO-3G and STO-6G. In each simulation, we perform the simulation by directly propagating the many-body density operator using DOPRI5 solver in the number preserving sector with random initialization. The convergence rate only depends on the number of orbitals and the number of electrons.

FIG. 4. The numerical verification of the chemical-detail-independence in the Hartree-Fock setting under the two types of bulk dissipation. (a) presents the dynamics of the Hartree-Fock state preparation of various molecular systems with Type-I set. (b) demonstrates the energy convergence curves of molecular systems with various values of L and N_e with Type-II set.




FIG. 5. Numerical demonstration of the convergence rate of the diagonal and off-diagonal entries of the 1-RDM with Type-II set for the H₄ system in STO-3G. The y -axis is plotted on a logarithmic scale.

spin orbitals, we may choose to propagate a many-body density operator, or a stochastic wavefunction by “unraveling” Lindblad dynamics and performing stochastic averages (see the SI), in the Fock space or in the FCI space. For systems of larger sizes, the only feasible option for direct simulation is to simulate the stochastic wavefunction using Type-II jump operators in the FCI space. We quantify the convergence of the Lindblad dynamics based on the rate of energy convergence.

We note that both Type-I and Type-II sets involve a

large number of jump operators, which can lead to increased simulation costs. However, in practice, the number of jump operators can be significantly reduced with minimal impact on efficiency. This is because the primary challenges in simulating chemical systems often arise in the low-energy space, particularly near the Fermi surface when we start with the Hartree-Fock initial guess. As a result, we can apply “active space” techniques to reduce the number of jump operators, focusing only on the most relevant degrees of freedom. For instance, if

we start from the vacuum state, it is unnecessary to include all operators from the set $\mathcal{A}_I = \{a_p\}_{p=1}^{2L} \cup \{a_p^\dagger\}_{p=1}^{2L}$, as the Hartree-Fock state is confined to the low-energy sector. Therefore, we can instead select the subset $\mathcal{S}_I^r = \{c_{i,\uparrow}, c_{i,\uparrow}^\dagger, c_{i,\downarrow}, c_{i,\downarrow}^\dagger\}_{i=N_e-r+1}^{N_e+r}$, which includes only the $8r$ orbitals defined around the Fermi surface under the molecular orbital basis.

We perform the numerical tests for \mathcal{S}_I^r . In all of the four systems demonstrated in Figure 6, we choose the initial state to be the Hartree-Fock state. We observe that energy decreases to λ_0 with a fidelity within the chemical accuracy, which shows a good transferability of the active space reduction idea to full *ab initio* calculations for the Type-I setting.




FIG. 6. Discrete Monte-Carlo trajectory based simulations for hydrogen chain systems: H_2 at bondlength 0.7 \AA in STO-3G, H_2 at bondlength 0.7 \AA in 6-31G, chain-like H_4 at bondlength 0.7 \AA in STO-3G and chain-like H_6 at bondlength 0.7 \AA in STO-3G. We choose \mathcal{S}_I^r as the coupling matrices with $r = 1, 1, 1, 2$ respectively and initialize with $|\text{HF}\rangle$ in all of the four cases.

Similarly, for the Type-II set, we can start from the Hartree-Fock state or low excited Slater determinant and include only the jump operators defined on a small number of orbitals around the Fermi surface. In this case, we consider the following reduced set of particle-number preserving coupling operators

$$\mathcal{S}_{II}^r = \{c_{i,\sigma}^\dagger c_{j,\tau} | i, j \in \{N_e-r+1, \dots, N_e+r\}, \sigma, \tau \in \{\uparrow, \downarrow\}\}, \quad (28)$$

which has $16r^2$ spin orbitals in total. In practice, setting $r = 1$ or $r = 2$ is typically sufficient to achieve convergence of the system to its ground state. The corresponding numerical results are presented in Figure 7. To compare the convergence rates of the full *ab initio* and Hartree-Fock state preparation with the reduced set \mathcal{S}_{II}^r , we begin from the (triplet) excited Slater determi-

nant $c_{N_e+1,\uparrow}^\dagger c_{N_e,\downarrow} |\text{HF}\rangle$. Notably, the convergence rate of the full *ab initio* state preparation is observed to be not much slower than that of the Hartree-Fock method, in the sense of Lindblad simulation time required to reach chemical accuracy.

In fact, the set of Type-II orbitals can be further compressed. For instance, we may take the subset \mathcal{T}_{II}^r of \mathcal{S}_{II}^r that contains only the “nearest hopping” of molecular orbitals:

$$\mathcal{T}_{II}^r = \{c_{i,\sigma}^\dagger c_{j,\tau} | i, j \in \{N_e-r+1, \dots, N_e+r\}, |i-j|=1, \sigma, \tau \in \{\uparrow, \downarrow\}\} \quad (29)$$

The number of operators in \mathcal{T}_{II}^r increases only linearly with L for fixed r . The numerical convergence behaviors with \mathcal{T}_{II}^r are shown in Figure 7. We observe that the further compression of \mathcal{S}_{II}^r maintains the simulation efficiency, in the sense that the Lindblad simulation time required to achieve chemical accuracy with the \mathcal{T}_{II}^r set remains comparable to or only slightly longer than that needed with the \mathcal{S}_{II}^r set.

The ground state preparation via Lindbladians with the set \mathcal{T}_{II}^r can perform well even in strongly correlated systems where the Hartree-Fock state poorly approximates the true ground state. Examples like the stretched square H_4 highlight these challenges due to their nearly degenerate low energy states, which even highly accurate methods like CCSD(T), often referred to as the “gold standard” in molecular quantum chemistry, struggle with [47, 61, 62]. Our results show that Lindblad dynamics effectively captures the correlation energy. As shown in Figure 8 (a), as the bond length of the square H_4 system increases, the energy accuracy of the Hartree-Fock initial guess decreases. Concurrently, the initial overlap between the Hartree-Fock state and the true ground state diminishes, indicating a growing extent of strong correlation. However, as illustrated in Figure 8 (b), the convergence of Lindblad dynamics remains largely unaffected by the degree of strong correlation starting from the Hartree-Fock initial state at various bond lengths.

III. DISCUSSION

To our knowledge, this work is the first Lindblad-based ground state preparation algorithm for *ab initio* electronic structure calculations. The Lindblad dynamics is employed as an algorithmic tool for dissipative state engineering. Unlike imaginary time evolution or quantum phase estimation, this method can efficiently prepare the ground state starting from any initial configuration, provided even starting from a state with zero initial overlap. We prove that the Lindblad dynamics with Type-I and Type-II jump operators can converge rapidly in a simplified Hartree-Fock setting, and validate the transferability to full *ab initio* calculations even for systems exhibiting strong correlation behaviors.

In order to perform numerical simulation for larger systems with tens to hundreds of spin-orbitals, even prop-





FIG. 7. Discrete Monte-Carlo trajectory based simulations for full *ab initio* molecular systems within the particle-number preserving sector with S_{II}^r and $\mathcal{T}_{\text{II}}^r$. In all cases, we initialize with the (triplet) excited Hartree-Fock Slater determinant $C_{N_e+1,\uparrow}^\dagger C_{N_e,\downarrow} |\text{HF}\rangle$. Panels (a-f) display results for the H₂, F₂, LiH, Cl₂, H₂O, and BeH₂ molecules respectively. Panel (g) presents the Lindblad simulation time required to achieve chemical accuracy with full *ab initio* and simplified Hartree-Fock Hamiltonians.



(a) The accuracy of the HF, CCSD and CCSD(T) energy and the initial overlap between the HF state and the true ground state ρ_{g} under difference bondlengths for square H₄ system in STO-3G

(b) The energy error relative to FCI energy vs. Lindblad simulation time for $d = 1.2, 1.4, 1.6$ and 2.0 Å

FIG. 8. Discrete Monte-Carlo trajectory based simulations for the strong-correlated H₄ molecule in STO-3G at different bondlengths.

agating the state vector in the FCI space can be very costly, and advanced simulation methods such as quantum Monte Carlo methods or tensor network-based methods must be employed. Theoretically, we need to develop a more powerful framework for understanding convergence rates with a degenerate stationary state (such as the ground state preparation problem), which may be analogous to the role of log-Sobolev inequalities [33] in thermal state analysis. This work may also provide new perspectives of ground state preparation in other areas, such as nuclei physics, fermions with random coefficients (e.g., the SYK model), or optimization problems on unstructured graphs.

ACKNOWLEDGMENTS

This work was partially supported by the National Natural Science Foundation of China under Grant No.

223B1011 (H.-E. L.). The Tsinghua Xuetang Talents Program and High-Performance Computing Center of Tsinghua University were acknowledged for providing computational resources. This material is based upon work supported by the U.S. Department of Energy, Office of Science, Accelerated Research in Quantum Computing Centers, Quantum Utility through Advanced Computational Quantum Algorithms, grant no. DE-SC0025572 (L.L.). The Institute for Quantum Information and Matter is an NSF Physics Frontiers Center. L.L. is a Simons Investigator in Mathematics. The authors thank Zhiyan Ding, Zhen Huang, and Jakob Huhn for helpful discussions.

-
- [1] Aspuru-Guzik, A., Dutoi, A. D., Love, P. J. & Head-Gordon, M. Simulated quantum computation of molecular energies. *Science* **309**, 1704–1707 (2005).
 - [2] Verstraete, F., Wolf, M. M. & Cirac, I. Quantum computation and quantum-state engineering driven by dissipation. *Nature Physics* **5**, 633–636 (2009).
 - [3] Motta, M. *et al.* Determining eigenstates and thermal states on a quantum computer using quantum imaginary time evolution. *Nat. Phys.* **16**, 205–210 (2020).
 - [4] Albash, T. & Lidar, D. A. Adiabatic quantum computation. *Reviews of Modern Physics* **90**, 015002 (2018).
 - [5] Cerezo, M. *et al.* Variational quantum algorithms. *Nature Reviews Physics* **3**, 625–644 (2021).
 - [6] Zhang, X.-M., Li, T. & Yuan, X. Quantum state preparation with optimal circuit depth: Implementations and applications. *Phys. Rev. Lett.* **129**, 230504 (2022).
 - [7] O’Brien, T. E., Tarasinski, B. & Terhal, B. M. Quantum phase estimation of multiple eigenvalues for small-scale (noisy) experiments. *New J. Phys.* **21**, 023022 (2019).
 - [8] Ge, Y., Tura, J. & Cirac, J. I. Faster ground state preparation and high-precision ground energy estimation with fewer qubits. *J. Math. Phys.* **60**, 022202 (2019).
 - [9] Lin, L. & Tong, Y. Near-optimal ground state preparation. *Quantum* **4**, 372 (2020).
 - [10] Lin, L. & Tong, Y. Heisenberg-limited ground state energy estimation for early fault-tolerant quantum computers. *PRX Quantum* **3**, 010318 (2022).
 - [11] Wan, K., Berta, M. & Campbell, E. T. Randomized quantum algorithm for statistical phase estimation. *Phys. Rev. Lett.* **129**, 030503 (2022).
 - [12] Dutkiewicz, A., Terhal, B. M. & O’Brien, T. E. Heisenberg-limited quantum phase estimation of multiple eigenvalues with few control qubits. *Quantum* **6**, 830 (2022).
 - [13] Ding, Z. & Lin, L. Even shorter quantum circuit for phase estimation on early fault-tolerant quantum computers with applications to ground-state energy estimation. *PRX Quantum* **4**, 020331 (2023).
 - [14] Lee, S. *et al.* Evaluating the evidence for exponential quantum advantage in ground-state quantum chemistry. *Nature Comm.* **14**, 1952 (2023).
 - [15] Berry, D. W. *et al.* Rapid initial state preparation for the quantum simulation of strongly correlated molecules. *arXiv preprint arXiv:2409.11748* (2024).
 - [16] Kraus, B. *et al.* Preparation of entangled states by quantum markov processes. *Phys. Rev. A* **78**, 042307 (2008).
 - [17] Zhou, L., Choi, S. & Lukin, M. D. Symmetry-protected dissipative preparation of matrix product states. *Physical Review A* **104**, 032418 (2021).
 - [18] Cubitt, T. S. Dissipative ground state preparation and the dissipative quantum eigensolver. *arXiv preprint arXiv:2303.11962* (2023).
 - [19] Wang, Y., Snizhko, K., Romito, A., Gefen, Y. & Murch, K. Dissipative preparation and stabilization of many-body quantum states in a superconducting qutrit array. *Phys. Rev. A* **108**, 013712 (2023).
 - [20] Lu, T.-C., Lessa, L. A., Kim, I. H. & Hsieh, T. H. Measurement as a shortcut to long-range entangled quantum matter. *PRX Quantum* **3**, 040337 (2022).
 - [21] Foss-Feig, M. *et al.* Experimental demonstration of the advantage of adaptive quantum circuits. *arXiv preprint arXiv:2302.03029* (2023).
 - [22] Kalinowski, M., Maskara, N. & Lukin, M. D. Non-Abelian Floquet Spin Liquids in a Digital Rydberg Simulator. *Phys. Rev. X* **13**, 31008 (2023).
 - [23] Mozgunov, E. & Lidar, D. Completely positive master equation for arbitrary driving and small level spacing. *Quantum* **4**, 1–62 (2020).
 - [24] Chen, C.-F. & Brandão, F. G. S. L. Fast thermalization from the eigenstate thermalization hypothesis. *arXiv preprint arXiv:2112.07646* (2023).
 - [25] Ding, Z., Chen, C.-F. & Lin, L. Single-ancilla ground state preparation via Lindbladians. *Phys. Rev. Research* **6**, 033147 (2024).
 - [26] Rall, P., Wang, C. & Wocjan, P. Thermal State Preparation via Rounding Promises. *Quantum* **7**, 1132 (2023).

- [27] Chen, C.-F., Kastoryano, M. J., Brandão, F. G. & Gilyén, A. Quantum thermal state preparation. *arXiv preprint arXiv:2303.18224* (2023).
- [28] Chen, C.-F., Kastoryano, M. J. & Gilyén, A. An efficient and exact noncommutative quantum Gibbs sampler. *arXiv preprint arXiv:2311.09207* (2023).
- [29] Ding, Z., Li, B. & Lin, L. Efficient quantum Gibbs samplers with Kubo–Martin–Schwinger detailed balance condition. *arXiv preprint arXiv:2404.05998* (2024).
- [30] Gilyén, A., Chen, C.-F., Doriguello, J. F. & Kastoryano, M. J. Quantum generalizations of Glauber and Metropolis dynamics. *arXiv preprint arXiv:2405.20322* (2024).
- [31] Ding, Z., Li, B., Lin, L. & Zhang, R. Polynomial-time preparation of low-temperature Gibbs states for 2d toric code. *arXiv preprint arXiv:2410.01206* (2024).
- [32] Temme, K., Kastoryano, M. J., Ruskai, M. B., Wolf, M. M. & Verstraete, F. The χ^2 -divergence and mixing times of quantum markov processes. *Journal of Mathematical Physics* **51** (2010).
- [33] Kastoryano, M. J. & Temme, K. Quantum logarithmic sobolev inequalities and rapid mixing. *Journal of Mathematical Physics* **54** (2013).
- [34] Bardet, I. *et al.* Rapid thermalization of spin chain commuting Hamiltonians. *Phys. Rev. Lett.* **130**, 060401 (2023).
- [35] Kastoryano, M. J. & Brandao, F. G. Quantum Gibbs samplers: The commuting case. *Communications in Mathematical Physics* **344**, 915–957 (2016).
- [36] Temme, K., Pastawski, F. & Kastoryano, M. J. Hypercontractivity of quasi-free quantum semigroups. *Journal of Physics A: Mathematical and Theoretical* **47**, 405303 (2014).
- [37] Temme, K. Thermalization time bounds for Pauli stabilizer Hamiltonians. *Communications in Mathematical Physics* **350**, 603–637 (2017).
- [38] Bardet, I. *et al.* Entropy decay for davies semigroups of a one dimensional quantum lattice. *Communications in Mathematical Physics* **405**, 42 (2024).
- [39] Alicki, R., Fannes, M. & Horodecki, M. On thermalization in kitaev’s 2d model. *Journal of Physics A: Mathematical and Theoretical* **42**, 065303 (2009).
- [40] Davies, E. B. Markovian master equations. *Commun. Math. Phys.* **39**, 91–110 (1974).
- [41] Lindblad, G. On the generators of quantum dynamical semigroups. *Commun. Math. Phys.* **48**, 119–130 (1976).
- [42] Gorini, V., Kossakowski, A. & Sudarshan, E. C. G. Completely positive dynamical semigroups of n -level systems. *J. Math. Phys.* **17**, 821–825 (1976).
- [43] Zhang, S., Carlson, J. & Gubernatis, J. E. Constrained path quantum monte carlo method for fermion ground states. *Phys. Rev. Lett.* **74**, 3652–3655 (1995).
- [44] Zhang, S. & Krakauer, H. Quantum monte carlo method using phase-free random walks with slater determinants. *Phys. Rev. Lett.* **90**, 136401 (2003).
- [45] Booth, G. H., Thom, A. J. W. & Alavi, A. Fermion Monte Carlo without fixed nodes: A game of life, death, and annihilation in Slater determinant space. *J. Chem. Phys.* **131**, 054106 (2009).
- [46] McArdle, S. *et al.* Variational ansatz-based quantum simulation of imaginary time evolution. *npj Quantum Inf.* **5**, 75 (2019).
- [47] Huggins, W. J. *et al.* Unbiasing fermionic quantum monte carlo with a quantum computer. *Nature* **603**, 416–420 (2022).
- [48] Lee, J., Pham, H. Q. & Reichman, D. R. Twenty years of auxiliary-field quantum monte carlo in quantum chemistry: An overview and assessment on main group chemistry and bond-breaking. *J. Chem. Theory Comput.* **18**, 7024–7042 (2022).
- [49] Jiang, T. *et al.* Walking through hilbert space with quantum computers. *arXiv preprint arXiv:2407.11672* (2024).
- [50] Su, Y., Huang, H.-Y. & Campbell, E. T. Nearly tight Trotterization of interacting electrons. *Quantum* **5**, 495 (2021).
- [51] Low, G. H., Su, Y., Tong, Y. & Tran, M. C. Complexity of implementing trotter steps. *PRX Quantum* **4**, 020323 (2023).
- [52] Su, Y., Berry, D. W., Wiebe, N., Rubin, N. & Babbush, R. Fault-tolerant quantum simulations of chemistry in first quantization. *PRX Quantum* **2**, 040332 (2021).
- [53] Tong, Y., An, D., Wiebe, N. & Lin, L. Fast inversion, preconditioned quantum linear system solvers, fast green’s-function computation, and fast evaluation of matrix functions. *Phys. Rev. A* **104**, 032422 (2021).
- [54] Thouless, D. J. Stability conditions and nuclear rotations in the hartree-fock theory. *Nuclear Physics* **21**, 225–232 (1960).
- [55] Szabo, A. & Ostlund, N. S. *Modern quantum chemistry: introduction to advanced electronic structure theory* (Courier Corporation, 2012).
- [56] Prosen, T. Third quantization: a general method to solve master equations for quadratic open Fermi systems. *New J. Phys.* **10**, 043026 (2008).
- [57] Prosen, T. & Žunkovič, B. Exact solution of Markovian master equations for quadratic Fermi systems: thermal baths, open XY spin chains and non-equilibrium phase transition. *New J. Phys.* **12**, 025016 (2010).
- [58] Horstmann, B., Cirac, J. I. & Giedke, G. Noise-driven dynamics and phase transitions in fermionic systems. *Phys. Rev. A* **87**, 012108 (2013).
- [59] Barthel, T. & Zhang, Y. Solving quasi-free and quadratic lindblad master equations for open fermionic and bosonic systems. *J. Stat. Mech: Theory Exp.* **2022**, 113101 (2022).
- [60] Stefanucci, G. & Van Leeuwen, R. *Nonequilibrium many-body theory of quantum systems: a modern introduction* (Cambridge Univ. Pr., 2013).
- [61] Paldus, J., Piecuch, P., Pylypow, L. & Jeziorski, B. Application of hilbert-space coupled-cluster theory to simple $(h_2)_2$ model systems: Planar models. *Phys. Rev. A* **47**, 2738–2782 (1993).
- [62] Lee, J., Huggins, W. J., Head-Gordon, M. & Whaley, K. B. Generalized unitary coupled cluster wave functions for quantum computation. *J. Chem. Theory Comput.* **15**, 311–324 (2018).
- [63] Lidar, D. A. Lecture notes on the theory of open quantum systems. *arXiv preprint arXiv:1902.00967* (2019).
- [64] Dalibard, J., Castin, Y. & Mølmer, K. Wave-function approach to dissipative processes in quantum optics. *Phys. Rev. Lett.* **68**, 580 (1992).
- [65] Dum, R., Zoller, P. & Ritsch, H. Monte carlo simulation of the atomic master equation for spontaneous emission. *Phys. Rev. A* **45**, 4879–4887 (1992).
- [66] Christie, R., Eastman, J., Schubert, R. & Graefe, E.-M. Quantum-jump vs stochastic schrödinger dynamics for gaussian states with quadratic Hamiltonians and linear Lindbladians. *J. Phys. A: Math. Theor.* **55**, 455302 (2022).

- [67] Breuer, H.-P. & Petruccione, F. *The theory of open quantum systems* (OUP Oxford, 2002).
- [68] Moodley, M. & Petruccione, F. Stochastic wave-function unraveling of the generalized Lindblad master equation. *Phys. Rev. A* **79**, 042103 (2009).
- [69] Löwdin, P.-O. On the non-orthogonality problem connected with the use of atomic wave functions in the theory of molecules and crystals. *J. Chem. Phys.* **18**, 365–375 (1950).
- [70] Sun, Q. *et al.* PySCF: the Python-based simulations of chemistry framework. *Wiley Interdiscip. Rev. Comput. Mol. Sci.* **8**, e1340 (2018).

**SUPPLEMENTARY INFORMATION: DISSIPATIVE GROUND STATE PREPARATION IN AB INITIO
ELECTRONIC STRUCTURE THEORY**

CONTENTS

I.	Introduction	1
II.	Results	2
	A. Dissipation engineering for ground state preparation	2
	B. Type-I and Type-II jump operators for <i>ab initio</i> calculations	2
	C. Universal fast convergence with Type-I set in Hartree-Fock theory	3
	D. Convergence with Type-II set oblivious to chemical details in Hartree-Fock theory	4
	E. Spectral gap of Lindbladian in Hartree-Fock theory	6
	F. Numerical verification in the Hartree-Fock setting	6
	G. Transferability to full <i>ab initio</i> calculations	6
III.	Discussion	8
	Acknowledgments	10
	References	10
	Supplementary Information: Dissipative ground state preparation in <i>ab initio</i> electronic structure theory	13
A.	Choice of filter function and sketch of quantum simulation algorithm	13
B.	Review of Hartree-Fock theory in the second quantized representation	14
C.	Solving quasi-free Lindblad dynamics	15
D.	Equations of motion of 1-RDM with Type-II jump operators	16
E.	Convergence of 1-RDM in the simplified Hartree-Fock setting	18
F.	Proof of Theorem 1 for the spectral gap of the Lindbladian	19
G.	Monte-Carlo trajectory based method for unraveling Lindblad dynamics	20
H.	Monte-Carlo trajectory based simulation results for <i>ab initio</i> calculations	21
I.	Molecular geometries used for the numerical simulations	21

Appendix A: Choice of filter function and sketch of quantum simulation algorithm

In this section, we discuss the choice of filter function in Ref. [25] and briefly review quantum algorithms for simulating the Lindblad dynamics.

We begin by reviewing the definition of the jump operator \hat{K}_k . Since the eigenvectors of \hat{H} are typically not accessible, we express \hat{K}_k in the time domain as follows:

$$\hat{K}_k = \int_{\mathbb{R}} f(s) A_k(s) ds \quad (\text{A1})$$

where $A_k(s) = e^{i\hat{H}s} A_k e^{-i\hat{H}s}$ is the Heisenberg evolution of A_k and $f(s) = \frac{1}{2\pi} \int_{\mathbb{R}} \hat{f}(\omega) e^{-i\omega s} d\omega$ is the inverse Fourier transform of the filter \hat{f} in the frequency domain. One possible choice for the filter function in the frequency domain \hat{f} is given by the following form [25]

$$\hat{f}(\omega) := \frac{\operatorname{erf}\left(\frac{\omega+a}{\delta_a}\right) - \operatorname{erf}\left(\frac{\omega+b}{\delta_b}\right)}{2} \quad (\text{A2})$$

where $\text{erf}(\omega) := \int_0^\omega \frac{2}{\sqrt{\pi}} e^{-t^2} dt$ denotes the error function. Here a is chosen to be an energy cutoff satisfying $a > 2\|\hat{H}\|_2$ and b is chosen to be the spectral gap of the Hamiltonian $\Delta := \lambda_1 - \lambda_0$. The parameters δ_a and δ_b are chosen to be on the same order of a and b respectively. In this setup, \hat{f} is approximately supported on the interval $[-2\|\hat{H}\|_2, -\Delta]$ (Note that the largest eigenvalue difference is $|\lambda_i - \lambda_j| \leq 2\|\hat{H}\|_2$). The inverse Fourier transform f can also be computed analytically as

$$f(s) = \frac{1}{2\pi i s} (\exp(ias - \delta_a^2 s^2/4) - \exp(ibs - \delta_b^2 s^2/4)) \quad (\text{A3})$$

where $f(0) = \frac{a-b}{2\pi}$ is obtained by taking the limit $s \rightarrow 0$. $f(s)$ is a smooth complex-valued function with the modulus $|f(s)|$ exhibiting a rapid decay when $|s| \rightarrow \infty$. Specifically, $f(s)$ approximately vanishes when $|s| > S_s$ for some $S_s = \Theta(1/\Delta)$, which allows us to truncate the infinite integral and use the trapezoidal quadrature rule to approximate the jump operator \hat{K}_k :

$$\hat{K}_k \approx \int_{-S_s}^{S_s} f(s) A(s) ds \approx \sum_{l=-M_s}^{M_s} f(s_l) A(s_l) w_l \quad (\text{A4})$$

where M_s is the number of quadrature nodes on $[0, S_s]$ or $[-S_s, 0]$, $s_l = l\Delta s$ and $\Delta s = S_s/M_s$. The weights $\{w_l\}$ are chosen to be $\Delta s/2$ for $l = \pm M_s$ and Δs for $-M_s + 1 \leq l \leq M_s - 1$.

To simulate the Lindblad dynamics $\exp(\mathcal{L}t)$ on quantum devices, we can begin with a first order Trotter splitting $\exp(\mathcal{L}t) \approx (\exp(\mathcal{L}_H\tau) \cdot \exp(\mathcal{L}_K\tau))^{t/\tau}$. The coherent dynamics $\exp(\mathcal{L}_H\tau)$ is just the Hamiltonian simulation $\exp(-i\hat{H}\tau)$. For the nonunitary dissipative part $\exp(\mathcal{L}_K\tau)$, we can reduce this problem to a dilated Hamiltonian simulation up to a partial trace on the ancilla qubit, namely

$$\exp(\mathcal{L}_K\tau)[\rho] = \text{Tr}_a e^{-i\tilde{K}\sqrt{\tau}} (|0_a\rangle\langle 0_a| \otimes \rho) e^{i\tilde{K}\sqrt{\tau}} + O(\tau^2). \quad (\text{A5})$$

Here, Tr_a denotes the partial trace operation on the ancilla qubit. For simplicity we consider only one jump operator K , and \tilde{K} is defined by the Hermitian dilated matrix $\tilde{K} = \begin{pmatrix} 0 & K^\dagger \\ K & 0 \end{pmatrix}$. Then the dilated Hamiltonian simulation $e^{i\tilde{K}\sqrt{\tau}}$ can be efficiently performed on a quantum computer using a second order Trotter splitting according to the discretized time evolution of K shown in Eq. (A4) (see [25] for details).

Appendix B: Review of Hartree-Fock theory in the second quantized representation

The HF theory finds a self-consistent single-particle operator approximation to the many-body Hamiltonian taking the form

$$\hat{H}_{\text{HF}} = \sum_{p,q=1}^{2L} F_{pq} a_p^\dagger a_q, \quad F_{pq} = h_{pq} + V_{pq} - K_{pq}. \quad (\text{B1})$$

Here L is the number of spatial orbitals, and $2L$ is the number of spin orbitals. The operators a_p^\dagger and a_p are fermionic creation and annihilation operators with respect to the orthonormalized spin orbitals. Here h_{pq} is a fixed single-particle matrix. The direct Coulomb and Fock exchange terms, denoted by V_{pq} and K_{pq} , respectively, should be solved self-consistently with respect to the one-particle density matrix (1-RDM), defined as

$$D_{pq} = \sum_{i=1}^{N_e} \Phi_{pi} \Phi_{qi}^*, \quad p, q = 1, 2, \dots, 2L. \quad (\text{B2})$$

Here $\Phi \in \mathbb{C}^{2L \times 2L}$ is a unitary matrix called the molecular orbital coefficients, which are eigenfunctions of the Fock matrix F . We denote $F\Lambda = \Lambda\Phi$ with $\Lambda = \text{diag}(\varepsilon_1, \dots, \varepsilon_{2L})$.

Once D is fixed, the HF Hamiltonian in Eq. (B1) is a quadratic Hamiltonian. Its ground state has an explicit expression in the Slater-determinant form:

$$|\text{HF}\rangle = c_1^\dagger \cdots c_{N_e}^\dagger |\text{vac}\rangle, \quad (\text{B3})$$

where the new set of creation operators $\{c_i^\dagger\}$ are given by the unitary transform of $\{a_i^\dagger\}$ via

$$(c_1^\dagger, \dots, c_{2L}^\dagger) = (a_1^\dagger, \dots, a_{2L}^\dagger) \Phi. \quad (\text{B4})$$

Note that $|\text{HF}\rangle$ is the ground state of the converged Hamiltonian \hat{H}_{HF} with eigenvalue $E_0 = \sum_{k=1}^{N_e} \varepsilon_k$. This sum of eigenvalues can also be expressed as $\text{Tr}(FD)$, which differs from the Hartree-Fock energy by a nonlinear term that depends only on D .

Appendix C: Solving quasi-free Lindblad dynamics

We consider the solution of the Lindblad dynamics that is quasi-free, i.e. with Hamiltonians that are quadratic in the fermionic creation and annihilation operators and Lindblad jump operators that are linear in the fermionic creation and annihilation operators [56–59].

Specifically, we consider $\hat{H} = \sum_{p,q}^{2L} F_{pq} a_p^\dagger a_q$ with a Hermitian coefficient matrix $F = (F_{pq})_{1 \leq p,q \leq 2L}$, and the following Lindblad jump operators:

$$\hat{K}_{p,+} = \sum_{r=1}^{2L} a_r^\dagger K_{r,p}, \quad \hat{K}_{q,-} = \sum_{r=1}^{2L} a_r J_{q,r}, \quad K_{r,p}, J_{q,r} \in \mathbb{C}. \quad (\text{C1})$$

We will derive the equation of motion for the one-particle reduced density matrix (1-RDM) with respect to the many-body density operator ρ , defined as

$$P_{ij} = \text{Tr}(\rho a_j^\dagger a_i), \quad 1 \leq i, j \leq 2L. \quad (\text{C2})$$

From the Lindblad equation:

$$\partial_t \rho = -i[\hat{H}, \rho] + \sum_p (\hat{K}_{p,+} \rho \hat{K}_{p,+}^\dagger - \frac{1}{2} \{\hat{K}_{p,+}^\dagger \hat{K}_{p,+}, \rho\}) + \sum_q (\hat{K}_{q,-} \rho \hat{K}_{q,-}^\dagger - \frac{1}{2} \{\hat{K}_{q,-}^\dagger \hat{K}_{q,-}, \rho\}), \quad (\text{C3})$$

the equation of motion of P_{ij} is given by

$$\partial_t P_{ij}(t) = -i \text{Tr}([\hat{H}, \rho] a_j^\dagger a_i) + \sum_{p, \bullet \in \{+, -\}} \text{Tr}(\hat{K}_{p,\bullet}^\dagger a_j^\dagger a_i \hat{K}_{p,\bullet} \rho) - \frac{1}{2} \text{Tr}(\{\hat{K}_{p,\bullet}^\dagger \hat{K}_{p,\bullet}, a_j^\dagger a_i\} \rho). \quad (\text{C4})$$

According to the canonical anticommutation relations (CAR), each term in Eq. (C4) can be simplified as follows:

$$-i \text{Tr}([\hat{H}, \rho] a_j^\dagger a_i) = -i \sum_{p=1}^{2L} (F_{ip} P_{pj} - F_{pj} P_{ip}), \quad (\text{C5})$$

$$\sum_p \text{Tr}(\hat{K}_{p,+}^\dagger a_j^\dagger a_i \hat{K}_{p,+} \rho) - \frac{1}{2} \text{Tr}(\{\hat{K}_{p,+}^\dagger \hat{K}_{p,+}, a_j^\dagger a_i\} \rho) = \sum_p K_{jp}^* K_{ip} - \frac{1}{2} \sum_{p,r} (K_{jp}^* K_{rp} P_{ir} + K_{rp}^* K_{ip} P_{rj}), \quad (\text{C6})$$

and

$$\sum_p \text{Tr}(\hat{K}_{p,-}^\dagger a_j^\dagger a_i \hat{K}_{p,-} \rho) - \frac{1}{2} \text{Tr}(\{\hat{K}_{p,-}^\dagger \hat{K}_{p,-}, a_j^\dagger a_i\} \rho) = -\frac{1}{2} \sum_{p,r} (J_{jp}^* J_{rp} P_{ir} + J_{rp}^* J_{ip} P_{rj}). \quad (\text{C7})$$

Next we provide detailed derivations of Eq. (C5) to (C7).

For Eq. (C5),

$$\begin{aligned} \text{Tr}([\hat{H}, \rho] a_j^\dagger a_i) &= \sum_{p,q} \text{Tr}(F_{pq} a_p^\dagger a_q \rho a_j^\dagger a_i) - \text{Tr}(F_{pq} \rho a_p^\dagger a_q a_j^\dagger a_i) \\ &= \sum_{pq} F_{pq} \text{Tr}(\rho (a_j^\dagger a_i a_p^\dagger a_q - a_p^\dagger a_q a_j^\dagger a_i)) \\ &= \sum_{pq} F_{pq} (\delta_{ip} \text{Tr}(\rho a_j^\dagger a_q) - \delta_{qj} \text{Tr}(\rho a_p^\dagger a_i)) + \sum_{pq} F_{pq} \text{Tr}(\rho (a_p^\dagger a_j^\dagger a_q a_i - a_j^\dagger a_p^\dagger a_i a_q)) \\ &= \sum_q F_{iq} P_{qj} - \sum_p F_{pj} P_{ip} + \sum_{pq} F_{pq} \text{Tr}(\rho ((-a_j^\dagger a_p^\dagger)(-a_i a_q) - a_j^\dagger a_p^\dagger a_i a_q)) \\ &= \sum_q F_{iq} P_{qj} - \sum_p F_{pj} P_{ip}. \end{aligned} \quad (\text{C8})$$

Here we use the definition $P_{ij} = \text{Tr}(\rho a_j^\dagger a_i)$ and the canonical anticommutation relations.

For Eq. (C6), we first notice that

$$\mathrm{Tr}\left(\left(\hat{K}_{p,+}^\dagger a_j^\dagger a_i \hat{K}_{p,+} \rho\right) - \frac{1}{2}\{\hat{K}_{p,+}^\dagger \hat{K}_{p,+}, a_j^\dagger a_i\}\rho\right) = \frac{1}{2}\mathrm{Tr}\left([\hat{K}_{p,+}^\dagger, a_j^\dagger a_i]\hat{K}_{p,+} \rho\right) + \frac{1}{2}\mathrm{Tr}\left(\rho \hat{K}_{p,+}^\dagger [a_j^\dagger a_i, \hat{K}_{p,+}]\right). \quad (\text{C9})$$

For each term in Eq. (C9), we expand $\hat{K}_{p,+}$ and $\hat{K}_{p,+}^\dagger$, yielding

$$\begin{aligned} & \mathrm{Tr}\left(\left([\hat{K}_{p,+}^\dagger, a_j^\dagger a_i]\hat{K}_{p,+} + \hat{K}_{p,+}^\dagger [a_j^\dagger a_i, \hat{K}_{p,+}]\right)\rho\right) \\ &= \sum_{r,s=1}^{2L} \mathrm{Tr}\left(K_{rp}^* K_{sp} (a_r a_j^\dagger a_i a_s^\dagger - a_j^\dagger a_i a_r a_s^\dagger + a_r a_j^\dagger a_i a_s^\dagger - a_r a_s^\dagger a_j^\dagger a_i)\rho\right) \end{aligned} \quad (\text{C10})$$

Using the algebraic relations $[a_p, a_j^\dagger a_i] = \delta_{pj} a_i$, $[a_q^\dagger, a_j^\dagger a_i] = -\delta_{qi} a_j^\dagger$ (can be verified directly using CAR), we obtain:

$$\begin{aligned} \mathrm{Tr}\left(\left([\hat{K}_{p,+}^\dagger, a_j^\dagger a_i]\hat{K}_{p,+} + \hat{K}_{p,+}^\dagger [a_j^\dagger a_i, \hat{K}_{p,+}]\right)\rho\right) &= \sum_{r,s=1}^{2L} \mathrm{Tr}\left(K_{rp}^* K_{sp} ([a_r, a_j^\dagger a_i] a_s^\dagger - a_r [a_s^\dagger, a_j^\dagger a_i])\rho\right) \\ &= \sum_{r,s=1}^{2L} \mathrm{Tr}\left(K_{rp}^* K_{sp} (\delta_{rj} a_i a_s^\dagger + \delta_{si} a_r a_j^\dagger)\rho\right) \\ &= \sum_{s=1}^{2L} K_{j,p}^* K_{s,p} \mathrm{Tr}(a_i a_s^\dagger \rho) + \sum_{r=1}^{2L} K_{r,p}^* K_{i,p} \mathrm{Tr}(a_r a_j^\dagger \rho). \end{aligned} \quad (\text{C11})$$

Taking summation with respect to the index p on both sides, we get

$$\begin{aligned} & \sum_p \mathrm{Tr}\left(\left([\hat{K}_{p,+}^\dagger, a_j^\dagger a_i]\hat{K}_{p,+} + \hat{K}_{p,+}^\dagger [a_j^\dagger a_i, \hat{K}_{p,+}]\right)\rho\right) \\ &= \sum_p \left(\sum_{s=1}^{2L} K_{j,p}^* K_{s,p} [\delta_{is} - \mathrm{Tr}(a_s^\dagger a_i \rho)] + \sum_{r=1}^{2L} K_{r,p}^* K_{i,p} [\delta_{rj} - \mathrm{Tr}(a_j^\dagger a_r \rho)] \right) \\ &= 2 \sum_p K_{ip} K_{jp}^* - \sum_{p,r} (K_{jp}^* K_{rp} P_{ir} + K_{rp}^* K_{ip} P_{rj}), \end{aligned} \quad (\text{C12})$$

which proves Eq. (C6). Here we use $\mathrm{Tr} \rho = 1$. Eq. (C7) can be verified by following almost the same procedure.

Using Eq. (C5) to Eq. (C7), it follows readily that the equation of motion $P(t)$ has the following closed form:

$$\partial_t P(t) = -i[F, P(t)] + B - \frac{1}{2}[P(t)(B+C) + (B+C)P(t)]. \quad (\text{C13})$$

The positive semidefinite matrices B and C are defined as

$$B_{ij} = \sum_p K_{ip} K_{jp}^*, \quad C_{ij} = \sum_p J_{ip} J_{jp}^*. \quad (\text{C14})$$

Eq. (C13) is a linear autonomous system for $P(t)$ of size $2L \times 2L$. Therefore, we can solve Lindblad dynamics of the quasi-free fermionic systems of large sizes by studying the evolution of the 1-RDM. Once we have the equation of motion for $P(t)$, we can obtain the evolution of energy expectation value (or other quadratic observables) of $\rho(t)$ using the 1-RDM:

$$E(t) = \mathrm{Tr}(\rho(t)\hat{H}) = \sum_{p,q=1}^{2L} \mathrm{Tr}(\rho(t) a_p^\dagger a_q F_{pq}) = \mathrm{Tr}(P(t)F). \quad (\text{C15})$$

Appendix D: Equations of motion of 1-RDM with Type-II jump operators

In this section, we provide a detailed derivation of the equation of motion of the 1-RDM with the Type-II set. We consider the converged Hartree-Fock Hamiltonian $\hat{H} = \sum_{p,q=1}^{2L} F_{pq} a_p^\dagger a_q$ with a Hermitian coefficient matrix F and the following set of coupling operators:

$$\mathcal{A}_{\text{II}} = \{a_i^\dagger a_j \mid i, j = 1, 2, \dots, 2L\}. \quad (\text{D1})$$

Assume F can be diagonalized as $F\Lambda = \Lambda\Phi$, where Φ is a unitary matrix and the diagonal entries of $\Lambda = \text{diag}(\varepsilon_p)$ are ordered non-decreasingly. In the molecular orbital basis, the HF Hamiltonian can be further simplified as a weighted summation of the number operators on the molecular orbitals, namely $\hat{H} = \sum_{p=1}^{2L} \varepsilon_p c_p^\dagger c_p$.

Next we consider the equation of motion of the 1-RDM under the energy basis of the HF Hamiltonian. We first calculate

$$\begin{aligned} \mathcal{L}_H^\dagger(c_r^\dagger c_s) &= i[\hat{H}, c_r^\dagger c_s] = i \sum_p \varepsilon_p c_p^\dagger c_p c_r^\dagger c_s - \varepsilon_p c_r^\dagger c_s c_p^\dagger c_p \\ &= i \sum_p \delta_{rp} \varepsilon_p c_p^\dagger c_s - \varepsilon_p c_p^\dagger c_r^\dagger c_p c_s - \delta_{ps} \varepsilon_p c_r^\dagger c_p + \varepsilon_p c_r^\dagger c_p^\dagger c_s c_p \\ &= i \sum_p \delta_{rp} \varepsilon_p c_p^\dagger c_s - \varepsilon_p (-c_r^\dagger c_p^\dagger)(-c_s c_p) - \delta_{ps} \varepsilon_p c_r^\dagger c_p + \varepsilon_p c_r^\dagger c_p^\dagger c_s c_p \\ &= i(\varepsilon_r - \varepsilon_s) c_r^\dagger c_s. \end{aligned} \quad (\text{D2})$$

Next, we compute the jump operators using the Thouless theorem

$$\hat{K}_{ij} = \int_{\mathbb{R}} f(s) e^{i\hat{H}s} a_i^\dagger a_j e^{-i\hat{H}s} ds = \sum_{p,q=1}^{2L} \hat{f}(\varepsilon_p - \varepsilon_q) c_p^\dagger c_q \Phi_{ip}^* \Phi_{jq} = \sum_{p<q} c_p^\dagger c_q \Phi_{ip}^* \Phi_{jq}. \quad (\text{D3})$$

Here we use $\hat{f}|_{[-2\|\hat{H}\|_2, -\Delta]} = 1$ and $\hat{f}|_{[0, +\infty)} = 0$. Therefore, using the fact that Φ consists of orthonormal columns, we have

$$\sum_{i,j=1}^{2L} \hat{K}_{ij}^\dagger \hat{K}_{ij} = \sum_{ij} \sum_{p<q} \sum_{r<s} c_q^\dagger c_p c_r^\dagger c_s \Phi_{ip} \Phi_{jq}^* \Phi_{ir}^* \Phi_{js} = \sum_{p<q} c_q^\dagger c_p c_p^\dagger c_q = \sum_{p<q} (1 - n_p) n_q. \quad (\text{D4})$$

Similarly, we calculate

$$\mathcal{L}_K^\dagger(c_r^\dagger c_s) = \sum_{i,j} \hat{K}_{ij}^\dagger c_r^\dagger c_s \hat{K}_{ij} - \frac{1}{2} \{ \hat{K}_{ij}^\dagger \hat{K}_{ij}, c_r^\dagger c_s \} = \sum_{p<q} c_q^\dagger c_p c_r^\dagger c_s c_p^\dagger c_q - \frac{1}{2} \{ c_q^\dagger c_p c_p^\dagger c_q, c_r^\dagger c_s \}. \quad (\text{D5})$$

To simplify Eq. (D5), we begin with rewriting it as

$$\mathcal{L}_K^\dagger(c_r^\dagger c_s) = \frac{1}{2} \sum_{p<q} (c_q^\dagger c_p c_r^\dagger c_s c_p^\dagger c_q - c_q^\dagger c_p c_p^\dagger c_q c_r^\dagger c_s) + \frac{1}{2} \sum_{p<q} (c_q^\dagger c_p c_r^\dagger c_s c_p^\dagger c_q - c_r^\dagger c_s c_q^\dagger c_p c_p^\dagger c_q). \quad (\text{D6})$$

For the first term,

$$\begin{aligned} c_q^\dagger c_p c_r^\dagger c_s c_p^\dagger c_q - c_q^\dagger c_p c_p^\dagger c_q c_r^\dagger c_s &= \delta_{ps} c_q^\dagger c_p c_r^\dagger c_q - c_q^\dagger c_p c_r^\dagger c_p^\dagger c_s c_q - \delta_{rq} c_q^\dagger c_p c_p^\dagger c_s + c_q^\dagger c_p c_p^\dagger c_r^\dagger c_s c_q \\ &= \delta_{ps} c_q^\dagger c_p c_r^\dagger c_q - c_q^\dagger c_p (-c_p^\dagger c_r^\dagger)(-c_q c_s) - \delta_{rq} c_q^\dagger c_p c_p^\dagger c_s + c_q^\dagger c_p c_p^\dagger c_r^\dagger c_s c_q \\ &= \delta_{ps} c_q^\dagger c_p c_r^\dagger c_q - \delta_{rq} c_q^\dagger c_p c_p^\dagger c_s. \end{aligned} \quad (\text{D7})$$

Likewise, for the second term,

$$c_q^\dagger c_p c_r^\dagger c_s c_p^\dagger c_q - c_r^\dagger c_s c_q^\dagger c_p c_p^\dagger c_q = \delta_{pr} c_q^\dagger c_s c_p^\dagger c_q - \delta_{sq} c_r^\dagger c_p c_p^\dagger c_q. \quad (\text{D8})$$

So we have,

$$\mathcal{L}_K^\dagger(c_r^\dagger c_s) = \frac{1}{2} \left(\sum_{q>s} c_q^\dagger c_s c_r^\dagger c_q - \sum_{p<r} c_r^\dagger c_p c_p^\dagger c_s + \sum_{q>r} c_q^\dagger c_s c_r^\dagger c_q - \sum_{p<s} c_r^\dagger c_p c_p^\dagger c_s \right). \quad (\text{D9})$$

In particular, for $r \neq s$, we move c_q^\dagger to the right in the first term and move c_q to the left in the third term:

$$\begin{aligned} \mathcal{L}_K^\dagger(c_r^\dagger c_s) &= \frac{1}{2} \left(- \sum_{q>s} c_r^\dagger c_s c_q^\dagger c_q - \sum_{p<r} c_p c_p^\dagger c_r^\dagger c_s - \sum_{q>r} c_q^\dagger c_q c_r^\dagger c_s - \sum_{p<s} c_r^\dagger c_s c_p c_p^\dagger \right) \\ &= -\frac{1}{2} (M_r c_r^\dagger c_s + c_r^\dagger c_s M_s) \end{aligned} \quad (\text{D10})$$

Here, M_k is defined as follows:

$$M_k := \sum_{p < k} c_p c_p^\dagger + \sum_{q > k} c_q^\dagger c_q = \sum_{p < k} c_p c_p^\dagger + \sum_{q > k} c_q^\dagger c_q = \sum_{p < k} (1 - n_p) + \sum_{q > k} n_q. \quad (\text{D11})$$

Notice that for $r < s$:

$$\begin{aligned} [c_r^\dagger c_s, M_s] &= \sum_{p < s} c_r^\dagger c_s c_p c_p^\dagger + \sum_{q > s} c_r^\dagger c_s c_q^\dagger c_q - \sum_{p < s} c_p c_p^\dagger c_r^\dagger c_s - \sum_{q > s} c_q^\dagger c_q c_r^\dagger c_s \\ &= \sum_{p < s} \delta_{pr} c_p^\dagger c_s - \sum_{q > s} \delta_{qr} c_q^\dagger c_s = c_r^\dagger c_s. \end{aligned} \quad (\text{D12})$$

Similarly, for $r > s$:

$$\begin{aligned} [M_r, c_r^\dagger c_s] &= \sum_{p < r} c_p c_p^\dagger c_r^\dagger c_s + \sum_{q > r} c_q^\dagger c_q c_r^\dagger c_s - \sum_{p < r} c_r^\dagger c_s c_p c_p^\dagger - \sum_{q > r} c_r^\dagger c_s c_q^\dagger c_q \\ &= \sum_{p < r} \delta_{ps} c_r^\dagger c_p - \sum_{q > r} \delta_{qs} c_r^\dagger c_q = c_r^\dagger c_s. \end{aligned} \quad (\text{D13})$$

Using Eq. (D2) and Eq. (D10) to (D13), we obtain the equation of motion of the 1-RDM in the molecular orbital basis $\tilde{P} = \Phi^\dagger P \Phi = \left(\text{Tr}(\rho c_j^\dagger c_i) \right)_{1 \leq i, j \leq 2L}$. For the off-diagonal entries, we have

$$\partial_t \tilde{P}_{sr} = \begin{cases} -i(\varepsilon_s - \varepsilon_r) \tilde{P}_{sr} - \frac{1}{2} \langle (M_r + M_s + 1) c_r^\dagger c_s \rangle & r < s, \\ -i(\varepsilon_s - \varepsilon_r) \tilde{P}_{sr} - \frac{1}{2} \langle c_r^\dagger c_s (M_r + M_s + 1) \rangle & r > s. \end{cases} \quad (\text{D14})$$

For the diagonal entries, we have

$$\begin{aligned} \partial_t \tilde{P}_{rr} &= \sum_{q > r} \langle c_q^\dagger c_r c_r^\dagger c_q \rangle - \sum_{p < r} \langle c_r^\dagger c_p c_p^\dagger c_r \rangle = \sum_{q > r} (\tilde{P}_{qq} - \langle n_r n_q \rangle) + \sum_{p < r} (\tilde{P}_{pp} - \langle n_p n_r \rangle) \\ &= - \sum_{p < r} \langle (1 - n_p) n_r \rangle + \sum_{q > r} \langle (1 - n_r) n_q \rangle = - \langle M_r n_r \rangle + \sum_{q > r} \langle n_q \rangle. \end{aligned} \quad (\text{D15})$$

Appendix E: Convergence of 1-RDM in the simplified Hartree-Fock setting

For the Type-I setting, we will show the universal exponential convergence that is oblivious to specific chemical details. For the Type-II setting, we provide a qualitative analysis on the convergence rate of the entries of the 1-RDM via mean-field approximation.

As discussed in the main text, the dynamics with Type-I set will converge exponentially to the stationary point P^* and the mixing time is independent of the system size:

$$\|P(t) - P'(t)\|_F = e^{-t} \|P(0) - P'(0)\|_F. \quad (\text{E1})$$

We can easily see Eq. (E1) by vectorizing $P - P'$, i.e. let $X := \text{vec}(P - P')$, then the equation of motion for $X(t)$ takes the following homogeneous form:

$$\dot{X}(t) = -i\mathbf{F}X(t) - X(t) \quad \Rightarrow \quad X(t) = e^{-t} e^{-i\mathbf{F}t} X(0). \quad (\text{E2})$$

Since $\mathbf{F} = I \otimes F - F^T \otimes I$ is Hermitian, it follows readily that

$$\|P(t) - P'(t)\|_F = \|X(t)\|_2 = e^{-t} \|e^{-i\mathbf{F}t} X(0)\|_2 = e^{-t} \|X(0)\|_2 = e^{-t} \|P(0) - P'(0)\|_F. \quad (\text{E3})$$

If we define the mixing time of the quasi-free Lindblad dynamics as [25]

$$\|P(t_{\text{mix}}) - P'(t_{\text{mix}})\|_F \leq \frac{1}{2} \|P(0) - P'(0)\|_F \quad (\text{E4})$$

for any initial covariance matrices $P(0)$ and $P'(0)$, then the dynamics of HF state preparation has the minimum mixing time $\log 2$ by definition and is independent of system size.

For the Type-II setting, we first estimate the bound of the positive semidefinite matrices M_k , defined in Eq. (D11). Let us consider three cases:

1. When $k < N_e$, we have $\sum_{q>k} n_q \geq N_e - k - 1 \geq 1$, therefore $M_k \succeq 1$.
2. When $k > N_e + 1$, we have $\sum_{p<k} (1 - n_p) \geq k - N_e \geq 1$, therefore $M_k \succeq 1$.
3. For $k \in \{N_e, N_e + 1\}$, we do not have a nontrivial lower bound for M_k . Indeed, M_k is positive semidefinite but not positive definite in this case, since we can easily verify that the Hartree-Fock state $|\text{HF}\rangle \in \ker M_k$.

For any 2-RDM $\langle a_i^\dagger a_j a_k^\dagger a_\ell \rangle$, we adopt the following mean-field approximation

$$\langle a_i^\dagger a_j a_k^\dagger a_\ell \rangle \approx \langle a_i^\dagger a_j \rangle \langle a_k^\dagger a_\ell \rangle. \quad (\text{E5})$$

To qualitatively understand the convergence behavior, we consider the mean-field approximation of Eq. (D14) and Eq.(D15). We begin with the off-diagonal entries. For both $r > s$ and $r < s$, we have

$$\partial_t \tilde{P}_{sr} \approx \left[-\frac{1}{2}(1 + \langle M_r + M_s \rangle) + i(\varepsilon_r - \varepsilon_s) \right] \tilde{P}_{sr}. \quad (\text{E6})$$

Noticing from Eq. (D11) that each M_k is a diagonal matrix in the energy basis represented as the sum of two positive semidefinite matrices, we have $M_r + M_s \succeq 0$. It follows readily that under the mean-field approximation, the off-diagonal entry \tilde{P}_{sr} converges exponentially to zero with an exponent of at least $\frac{1}{2}$.

For the diagonal entries, we take the mean-field approximation on Eq. (D15), yielding

$$\partial_t \tilde{P}_{rr} \approx -\langle M_r \rangle \tilde{P}_{rr} + \sum_{q>r} \tilde{P}_{qq}. \quad (\text{E7})$$

These equations can be solved one-by-one. Specifically, we start from $r = 2L$:

$$\partial_t \tilde{P}_{2L,2L} \approx -\langle M_{2L} \rangle \tilde{P}_{2L,2L}. \quad (\text{E8})$$

This homogeneous equation gives that $\tilde{P}_{2L,2L}$ converges exponentially with an exponent being at least 1 since $M_{2L} \succeq 1$. Then we move on to $r = 2L - 1$ and this time $\tilde{P}_{2L,2L}$ becomes the inhomogeneous term, so the convergence exponent of $\tilde{P}_{2L-1,2L-1}$ is also greater than 1 and so forth. This argument holds for all $r > N_e + 1$. For $r < N_e$, we rewrite the mean-field equation of motion as follows:

$$\partial_t (1 - \tilde{P}_{rr}) \approx \langle M_r \rangle (1 - \tilde{P}_{rr}) + \sum_{p<r} (1 - \tilde{P}_{pp}). \quad (\text{E9})$$

We solve these equations from $r = 1$ and similarly we conclude that \tilde{P}_{rr} converges exponentially to 1 for all $r < N_e$.

Appendix F: Proof of Theorem 1 for the spectral gap of the Lindbladian

Proof. Since $[\hat{H}, \hat{H}_{\text{dp}}] = 0$, we can choose $\{|\psi_j\rangle\}$ to diagonalize \hat{H} and \hat{H}_{dp} simultaneously, and the eigenvalues for \hat{H} still satisfies $\lambda_0 < \lambda_1 \leq \dots$. We can also write $\hat{H}_{\text{dp}} |\psi_j\rangle = \xi_j |\psi_j\rangle$, but the eigenvalues ξ_j may not be ordered. Then for $\hat{J} = -i\hat{H} - \hat{H}_{\text{dp}}$ we have $\hat{J} |\psi_j\rangle = (-i\lambda_j - \xi_j) |\psi_j\rangle$.

We can then use $\{|\psi_i\rangle\langle\psi_j|\}$ as an ordered basis for expanding ρ , following a row-major order. Since \hat{K}_k is by construction an upper triangular matrix in the energy basis, the operator $\mathcal{K} : \rho \mapsto \sum_k \hat{K}_k \rho \hat{K}_k^\dagger$ is also upper triangular after vectorization. Meanwhile for $\rho = |\psi_i\rangle\langle\psi_j|$, we have

$$\hat{J}\rho + \rho\hat{J}^\dagger = (-i\lambda_i - \xi_i + i\lambda_j - \xi_j)\rho. \quad (\text{F1})$$

So the vectorized Lindbladian, denoted by \mathbf{L} , is an upper triangular matrix with eigenvalues $(-i\lambda_i - \xi_i + i\lambda_j - \xi_j)$. Since \hat{H}_{dp} has an eigenvalue 0, this immediately shows that the spectral gap of \mathbf{L} is equal to that of \hat{H}_{dp} . \square

We compute the jump operators with Type-I set under the molecular orbital basis. Recall that

$$\hat{K}_{p,+} = \sum_{r=1}^{2L} a_r^\dagger (\hat{f}(F))_{r,p}, \quad \hat{K}_{q,-} = \sum_{r=1}^{2L} a_r (\hat{f}(-F))_{q,r}. \quad (\text{F2})$$

Rewriting Eq. (F2) in the molecular orbital basis yields

$$\hat{K}_{p,+} = \sum_{r=1}^{2L} c_r^\dagger \hat{f}(\varepsilon_r) \Phi_{pr}^*, \quad \hat{K}_{q,-} = \sum_{r=1}^{2L} c_r \hat{f}(-\varepsilon_r) \Phi_{qr}. \quad (\text{F3})$$

Then

$$\hat{H}_{\text{dp}} = \frac{1}{2} \sum_p \hat{K}_{p,+}^\dagger \hat{K}_{p,+} + \frac{1}{2} \sum_q \hat{K}_{q,-}^\dagger \hat{K}_{q,-} = \frac{1}{2} \sum_{r=1}^{2L} \hat{f}^2(\varepsilon_r) c_r c_r^\dagger + \frac{1}{2} \sum_{r=1}^{2L} \hat{f}^2(-\varepsilon_r) c_r^\dagger c_r. \quad (\text{F4})$$

In other words,

$$\hat{H}_{\text{dp}} = \frac{1}{2} \sum_{p \leq N_e} (1 - n_p) + \frac{1}{2} \sum_{q > N_e} n_q. \quad (\text{F5})$$

Appendix G: Monte-Carlo trajectory based method for unraveling Lindblad dynamics

To solve the Lindblad dynamics for ground state preparation at the FCI level, we may directly propagate the many-body density operator $\rho(t) \in \mathbb{C}^{2^L \times 2^L}$ using a differential equation solver [63]. Another approach is to ‘‘unravel’’ the Lindblad dynamics for the many-body density operator [64, 65]. Broadly speaking, we employ a family of Monte-Carlo-type algorithms where we only propagate the state vector $|\psi(t)\rangle$ in some stochastic schemes, and the many-body density operator $\rho(t)$ at the time t can be retrieved by taking the average of the random matrix $|\psi(t)\rangle\langle\psi(t)|$, i.e. $\rho(t) = \mathbb{E}|\psi(t)\rangle\langle\psi(t)|$. The deterministic Lindblad equation for the dynamics of the many-body density operator is now expressed using stochastic pure-state trajectories, thus leading to a quadratic reduction in dimensionality, at the cost of incorporating statistical averaging across multiple runs.

The simplest setting is the discrete form of unraveling, or quantum jump method [66]. The quantum-jump pure-state dynamics is evolved under an effective non-Hermitian Hamiltonian $\hat{H} - i/2 \sum_k \hat{K}_k^\dagger \hat{K}_k$, with stochastic quantum jumps occurring intermittently throughout the evolution. Specifically, it can be described by the following stochastic differential equation (SDE):[66–68]

$$d\psi = \left(-i\hat{H} - \frac{1}{2} \sum_k (\hat{K}_k^\dagger \hat{K}_k - \langle \hat{K}_k^\dagger \hat{K}_k \rangle) \right) \psi dt + \sum_k \left(\frac{\hat{K}_k}{\sqrt{\langle \hat{K}_k^\dagger \hat{K}_k \rangle}} - 1 \right) \psi dN_t^k. \quad (\text{G1})$$

Here, N_t denotes a Poisson process with a splitting $N_t = \sum_k N_t^k$. For a sufficiently small time step Δt , the Poisson increment ΔN_t takes the values 0 (no jump) or 1 (jump) with expectation value $\mathbb{E}(\Delta N_t) = \sum_k \|\hat{K}_k \psi\|^2 \Delta t = \sum_k \langle \hat{K}_k^\dagger \hat{K}_k \rangle \Delta t$. The Poisson processes $\{N_t^k\}$ are mutually independent with intensities given by $\|\hat{K}_k \psi\|^2 = \langle \hat{K}_k^\dagger \hat{K}_k \rangle$. This implies that, in the event of a jump, we select the jump operator \hat{K}_k to apply to ψ with a probability proportional to $\langle \hat{K}_k^\dagger \hat{K}_k \rangle$ for $k = 1, 2, \dots, N$. It can be shown that the density operator, defined as $\rho(t) = \mathbb{E}|\psi(t)\rangle\langle\psi(t)|$ indeed solves the Lindblad dynamics, by calculating $\frac{d\psi\psi^\dagger}{dt}$ using Itô’s lemma for the Poisson process and then taking the expectation.

For a Monte-Carlo-type simulation of Eq. (G1), we first discretize the time interval by the time step Δt . Then at each step, we randomly pick up $k \in [N + 1]$ (assume we have N jump operators in total) with respect to the distribution

$$p_k = \begin{cases} \|\hat{K}_k \psi_n\|^2 \Delta t, & k \leq N \\ 1 - \sum_{\ell=1}^N p_\ell, & k = N + 1 \end{cases}. \quad \text{If } k = 1, \dots, N, \text{ we update the trajectory using } \psi_{n+1} = \hat{K}_k \psi_n / \langle \hat{K}_k^\dagger \hat{K}_k \rangle. \text{ If}$$

$k = N + 1$, we update using $\psi_{n+1} = \psi_n - (i\hat{H} + \frac{1}{2} \sum_k \hat{K}_k^\dagger \hat{K}_k) \psi_n \Delta t$. Then the many-body density operator $\rho(t_n)$ at time t_n can be approximated by taking the average of the pure states $|\psi(t_n)\rangle\langle\psi(t_n)|$ over the trajectories [63].

We can also consider a variant of the Monte Carlo-type algorithm that is slightly different but essentially equivalent to the one described above. Notice that [63]

$$\left\| \exp \left(-i \left(H - \frac{i}{2} \sum_k \hat{K}_k^\dagger \hat{K}_k \right) \Delta t \right) \psi(t) \right\|^2 = 1 - \sum_k \|\hat{K}_k \psi(t)\|^2 \Delta t + O((\Delta t)^2) \quad (\text{G2})$$

which implies that the decaying evolution governed by the non-Hermitian effective Hamiltonian primarily dictates the probability distribution. Consequently, we actually do not need to compute $\|\hat{K}_k \psi_n\|^2$ at every time step. Instead, we

can first calculate $\|\psi_n\|^2$ and sample a random number $R \sim \text{U}(0, 1)$ to decide whether jump or not. If $R < \|\psi_n\|^2$, we just propagate the trajectory using $\psi_{n+1} = \psi_n - \left(i\hat{H} + \frac{1}{2} \sum_k \hat{K}_k^\dagger \hat{K}_k\right) \psi_n \Delta t$. If $R \geq \|\psi_n\|^2$, we calculate each $\langle \hat{K}_k^\dagger \hat{K}_k \rangle$ and update $\psi_{n+1} = \hat{K}_k \psi_n / \langle \hat{K}_k^\dagger \hat{K}_k \rangle$ with a probability proportional to $\langle \hat{K}_k^\dagger \hat{K}_k \rangle$ for $k = 1, 2, \dots, N$. After this, we reset the random number $R \sim \text{U}(0, 1)$. Essentially, this corresponds to evolving the trajectory using $\hat{H} - i/2 \sum_k \hat{K}_k^\dagger \hat{K}_k$ deterministically causing the norm of ψ to decrease, until a random quantum jump occurs. At this point, the norm is restored to 1, and the process repeats [59]. In all of the steps in the unraveling algorithm, only matrix-vector multiplication is involved.

Appendix H: Monte-Carlo trajectory based simulation results for *ab initio* calculations

In this section, we implement the Monte-Carlo trajectory based method (as described in Section G) for interacting systems.

To construct the converged HF Hamiltonian in second quantized representation for the numerical validations of the convergence rate arguments of the simplified HF settings (with Type-I or Type-II set), we first apply the Löwdin's orthogonalization to obtain a set of orthonormal spin orbitals [55, 69]. Concretely, given that the overlap matrix S is Hermitian positive definite, we can diagonalize it as $S = UsU^\dagger$ where U is a unitary matrix and s is a diagonal matrix of positive eigenvalues. We define the transformation matrix $X = Us^{-1/2}$ and transform the Fock operator using $F \mapsto X^\dagger F X$. This transformation is equivalent to using the orthogonalized spin orbitals as the new basis set. The one-particle and two-particle integrals, as well as the overlap integrals within a given basis set are imported from the PySCF package [70].

We can then consider the electronic structure Hamiltonian in the FCI basis

$$\hat{H} = \sum_{p,q=1}^{2L} T_{pq} c_p^\dagger c_q + \frac{1}{2} \sum_{p,q,r,s=1}^{2L} S_{pqrs} c_p^\dagger c_q^\dagger c_r c_s, \quad (\text{H1})$$

where the T_{pq} and S_{pqrs} are the one-particle and two-particle integrals of molecular orbitals generated from the HF calculation.

For all of the simulations, we choose the parameters of the filter as follows: $a = \frac{5}{2} \|\hat{H}\|_2$, $\delta_a = a/5$, $b = \delta_b = \Delta$. For the integral truncation, we let $S_s = \frac{10}{\Delta}$ and for the quadrature nodes we set $M_s = \lceil S_s / (\pi/2a) \rceil$ and $\Delta s = S_s / M_s$ [25]. All units are in the atomic unit (a.u.).

We numerically verify that this method can indeed approximate Lindblad dynamics. We use the vacuum state as initial state and the reduced Type-I set \mathcal{S}_I as the jump operators to prepare the ground state of H_2 system. We apply the quantum jump unraveling method for these simulations with time step $\Delta t = 0.1$ and the stopping times are set to $T = 30$. In each simulation, we repeat the pure-state evolution on $N_{\text{traj}} = 10, 25, 50, 100$ and 500 trajectories and we approximate the many-body density operator at the time t_n as:

$$\rho(t_n) = \mathbb{E} |\psi(t_n)\rangle\langle\psi(t_n)| \approx \frac{1}{N_{\text{traj}}} \sum_{k=1}^{N_{\text{traj}}} |\psi_k(t_n)\rangle\langle\psi_k(t_n)|, \quad (\text{H2})$$


where $|\psi_k(t_n)\rangle$ denotes the state vector on the k -th trajectory at the time t_n . It can be observed in Figure 9 that as the number of trajectories increases, the results of the quantum-jump unraveling exhibit a trend of convergence to the exact Lindblad dynamics, which is obtained by propagating the density operator using the DOPRI5 solver.

Due to the quadrature error in the jump operators, the Monte Carlo trajectory evolution methods may exhibit numerical instability. In short, this instability arises because the quadrature error can occasionally lead to an over-estimation of the quantum jump probability, which can cause an increase in energy, particularly as the dynamics approach the ground state.


Appendix I: Molecular geometries used for the numerical simulations

For the numerical simulations of the ground state preparation at the Hartree-Fock (HF) level using the quasi-free approach, we conduct the simulation for these molecular systems: H_4 , LiH , H_2O , CH_4 , HCN , C_2H_4 , N_2 , H_{10} and SO_3 . The molecular geometries are listed below:

\mathbf{H}_4 in STO3G:
H 0.0 0.0 0.0



(a) Energy convergence curves



(b) Absolute error in the energy expectation value relative to the exact Lindblad dynamics vs. time

FIG. 9. The Lindblad dynamics of the H_2 system in the STO-3G basis set, obtained by exactly propagating the many-body density operator and using the quantum-jump unraveling method with 10, 25, 50, 100, and 500 trajectories, respectively. The (a) panel illustrates the energy convergence curves of each simulation. The (b) panel shows the average absolute error of the overlap $\langle \rho_g \rangle$ with the exact ground state relative to the exact Lindblad dynamics at each time point. The average error decreases approximately as $1/\sqrt{N_{\text{traj}}}$.

```
H 0.0 0.0 2.0
H 0.0 2.0 0.0
H 0.0 2.0 2.0
```

LiH in STO3G:

```
H 0.000 0.000 0.000
Li 0.000 0.000 1.546
```

H₂O in STO3G:

```
O 0.000000 0.000000 0.1271610
H 0.000000 0.7580820 -0.5086420
H 0.000000 -0.7580820 -0.5086420
```

HCN in STO3G:

```
C 0.000000 0.000000 -0.5000780
H 0.000000 0.000000 -1.5699900
N 0.000000 0.000000 0.6529230
```

C₂H₄ in STO3G:

```
C 0.000000 0.000000 0.6530360
C 0.000000 0.000000 -0.6530360
H 0.000000 0.9157470 1.2292130
H 0.000000 -0.9157470 1.2292130
H 0.000000 -0.9157470 -1.2292130
H 0.000000 0.9157470 -1.2292130
```

N₂ in cc-pVDZ:

```
N 0.000000 0.000000 0.5386530
N 0.000000 0.000000 -0.5386530
```

H₁₀ in cc-pVDZ:

H 0.0 0.0 0.0
H 0.0 0.0 0.7
H 0.0 0.0 1.4
H 0.0 0.0 2.1
H 0.0 0.0 2.8
H 0.0 0.0 3.5
H 0.0 0.0 4.2
H 0.0 0.0 4.9
H 0.0 0.0 5.6
H 0.0 0.0 6.3

SO₃ in cc-pVDZ:

S 0.0000000 0.0000000 0.0000000
O 0.0000000 1.4167620 0.0000000
O 1.2269520 -0.7083810 0.0000000
O -1.2269520 -0.7083810 0.0000000

molecule	basis set	bond length (Å)	number of qubits	dimension of \mathcal{F}
H ₂	STO3G	0.7	4	16
H ₂	6-31G	0.7	8	256
H ₄ (chain)	STO3G	0.7	8	256
H ₆ (chain)	STO3G	0.7	12	4096

TABLE I. The molecular geometries used in the simulations of interacting systems with Type-I

molecule	basis set	bond length (Å)	number of qubits	dimension of \mathcal{F}_{N_e}
H ₄ (square)	STO3G	-	8	70
H ₂	cc-pVDZ	0.7	20	190
F ₂	STO3G	1.4	20	190
LiH	STO3G	1.546	12	495
Cl ₂	STO3G	2.130	36	630
H ₆ (grid)	STO3G	2.0	12	924
H ₂ O	STO3G	-	14	1001
BeH ₂	STO3G	1.304	14	3003

TABLE II. The molecular geometries used in the simulations of interacting systems with Type-II

For the simulations of interacting systems using the Monte Carlo trajectory-based method, we employed the same geometry as in the quasi-free calculation for the H₂O molecule within the STO-3G basis set. The geometries for other molecular systems are listed in Table I and Table II. Additionally, we provide the dimension of the total Fock space \mathcal{F} and the FCI sector \mathcal{F}_{N_e} with N_e electrons, which corresponds to the size of the Hamiltonian matrix \hat{H} .