# Kinematic censorship and high energy particle collisions in the Schwarzschild background

A. V. Toporensky

Sternberg Astronomical Institute, Lomonosov Moscow State University, 119991 Moscow, Russia\*

O. B. Zaslavskii

Department of Physics and Technology, Kharkov V.N. Karazin National University, 4 Svoboda Square, Kharkov 61022, Ukraine<sup>†</sup>

We consider near-horizon collisions between two particles moving freely in the Schwarzschild metric in the region outside the horizon. One of them emerges from a white hole. We scrutiny when such a process can lead to the indefinitely large growth of the energy in the center of mass frame in the point of collision. We also trace how the kinematics of collision manifests itself in preserving the principle of kinematic censorship according to which the energy released in any event of collision cannot be literally infinite. According to this principle, the energy released in any event of collision, must remain finite although it can be made as large as one likes. Also, we find that particle decay near the singularity leads to unbounded release of energy independently of its initial value.

PACS numbers: 04.70.Bw, 97.60.Lf

### I. INTRODUCTION

High-energy particle collisions near black holes became a hot topic after an observation made in [1]. If two particles fall towards the rotating black hole and collide near its horizon, under certain conditions the energy  $E_{c.m.}$  in the center of mass frame can grow unbounded. This is called the BSW effect (after the names of its authors). It was found for extremal

<sup>\*</sup>Electronic address: atopor@rambler.ru

<sup>&</sup>lt;sup>†</sup>Electronic address: zaslav@ukr.net

rotating black holes and requires one of particles to be fine-tuned,  $E = \omega_H L$  [1], [2]. Here, E is the Killing energy, L being the angular momentum,  $\omega_H$  the angular speed of a black hole. For the Schwarzschild black hole this is impossible since  $\omega_H = 0$ . Meanwhile, there is another version (or analogue) of the BSW effect, valid even in the Schwarzschild background. It implies head-on collision between particles, when, say, particle 1 approaches the horizon while particle 2 moves away from it. In doing so, no fine-tuning is required,  $E_{c.m.}$  grows indefinitely for any individual energies. As a particle cannot emerge from the black hole horizon, this scenario implies the presence of a white hole. Collisions of such a type were considered in [3], [4]. Before, the fact that collision between particles with opposite radial velocity leads to unbounded  $E_{c.m.}$  was noticed in [5] - [7]. However, it was not pointed out there that any reasonable physical realization of such a scenario corresponds just to white (not black) hole horizons. A coherent scheme of possible high energy particle collisions involving particles that emerge from a white hole, was built in [8] depending on possible fine-tuning parameters of one of two particles.

Meanwhile, in the present article, we consider other aspects of head-on collisions. We take a simplest and physically relevant metric - the Schwarzschild one. Then, all particles are not fine-tuned, so we consider collisions between two usual particles. This corresponds to line 1 in Table 1 of Ref. [8]. Thus we restrict ourselves by one type of collision only but for this type scrutinize all possible types of trajectories in space-time and their kinematic properties. In this context, we make special emphasis on manifestation of the principle of kinematic censorship [9]. According to this principle, in any event of collision, only finite energy can be released. If some scenario admits literally infinite  $E_{c.m.}$ , it is based on some unphysical assumption or mistake and should be rejected. Sometimes, it is quite non-trivial task to trace how this principle reveals itself in concrete scenarios [10]. In the present work, we take a simplest case and consider collisions in our region (outside the horizon). Even in this case, corresponding explanation, how this principle reveals itself, is quite involved and relies heavily on some subtle details of particle kinematics.

One reservation is necessary. Although the fine-tuned trajectories in common sense do not exist in the Schwarzschild space-time, there exist very special scenarios in which a particle with small, although nonzero energy  $E \approx 0$  plays a crucial role [11] - [15]. We will not discuss this degenerate scenario in the present paper.

The notion of white holes was introduced in [16] where it was conjectured that they

can be realized as some kinds of remnants after expansion of Universe, see also [17]. They can be considered as some windows through which the energy flows in our universe [18], [19], [20], [21]. Such objects are probably unstable as was argued in [22] (see also Sec. 15. 2 in [23]). However, interest to them is still kept as they can lead to high energy collisions between matter emerging from them and matter in our universe Particle collisions under discussion can be a potential source of one more kind of instability of white holes. Independently of whether or not astrophysical white holes exist, consideration of physical processes in their vicinity is of big methodical interest. (For instance, see recent work [24] and references 43-65 therein.) General relativity is consistent with such objects and, moreover, the full geodesically complete structure of space-time includes both black and white hole regions that reveals itself also in potential physical processes in the geodesically complete background can model more realistic ones when collapsing matter collides with expanding one.

The paper is organized as follows. In Sec. II we give the expression for the metric, equations of motion of free particles and the main formulas for particle collisions. In Sec. III and IV we describe in detail main scenarios of particle collisions discussed in the present paper. In Sec. V we discuss the relations between the principle of kinematic censorship and properties of trajectories in question. In Sec. VI we discuss the properties of the Schwarzschild time for a particle moving near a white whole and relate them to our scenarios. In Sec. VII we describe these scenarios using the expanding Lemaître frame. In Sec. VIII we consider processes with particles near the singularity. In Sec. IX we give summary of the results obtained.

#### II. GENERAL SETUP

Let us consider the static spherically symmetric metric

$$ds^{2} = -dt^{2}f + f^{-1}dr^{2} + r^{2}d\omega^{2}.$$
(1)

If  $f = 1 - \frac{r_+}{r}$ , this corresponds to the Schwarzschild metric with  $r_+$  being the horizon radius. We restrict ourselves by pure radial motion. Choosing the plane in which a particle moves to be  $\theta = \frac{\pi}{2}$ , we have equations of motion

$$\dot{t} = \frac{\varepsilon}{f},\tag{2}$$

$$\dot{r} = \sigma P, P = \sqrt{\varepsilon^2 - f}.$$
 (3)

Here, dot denotes derivative with respect to the proper time  $\tau, \varepsilon = \frac{E}{m}$ , where E is the Killing energy, m being the particle's mass,  $\sigma = +1$  for motion in the outward direction and  $\sigma = -1$  for for motion in the inward one.

At the horizon, the metric coefficients in the coordinate frame (1) spoil. To repair this, there exists a number of other frames. We will stick to the Kruskal-Szekeres coordinates (see, e.g. the textbook [25], Chapter VIII. 31. 4 - 31.5. A reader should bear in mind that we use another notations). The concrete form of transformation from original coordinates to Kruskal-Szekeres ones depends on the region under consideration.

In what follows, we will be interested in the energy  $E_{c.m.}$  in the center of mass frame of two colliding particles. By definition,

$$E_{c.m.}^{2} = -(m_{1}u_{1\mu} + m_{2}u_{2\mu})(m_{1}u_{1}^{\mu} + m_{2}u_{2}^{\mu}) = m_{1}^{2} + m_{2}^{2} + 2m_{1}m_{2}\gamma,$$
(4)

where  $\gamma = -u_{1\mu}u_2^{\mu}$  is the Lorentz factor of relative motion. If particles 1 and 2 move in the same plane, it follows from (2), (3) that

$$\gamma = \frac{\varepsilon_1 \varepsilon_2 - \sigma_1 \sigma_2 P_1 P_2}{f}.$$
(5)

Hereafter, we will use, along with original Schwarzschild coordinates (1), also Kruskal-Szekeres ones since this enables us to avoid coordinate singularities on the horizon. The whole space-time contains four regions (see. e.g. Fig. 31.3 of [25]). However, for our purposes we need to consider only two of them. These are our region outside the horizon and the white hole one  $T^+$  under the horizon [23].

## III. SCENARIO 1. PARTICLE 1 IN OUR REGION, PARTICLE 2 FROM WHITE HOLE REGION. COLLISION NEAR THE PAST HORIZON

We consider a scenario in which particle 2 emerges from the white hole region and meets particle 1 in our region outside the horizon. See Fig. 1. We assume that collision occurs in the point  $r = r_c$  somewhere near the right branch of the past horizon, so in this point  $V_c \approx 0$  while  $U_c < 0$  is arbitrary.



Fig 1. Particle collision in Scenario 1

Particle 1 comes from remote past, particle 2 comes from a white hole crossing the past horizon in point P. Collision occurs near an arbitrary point C of the past horizon.

## A. Properties of the trajectory of particle 1

Let us consider movement of a particle in region  $r > r_+$ . Then, the relevant transformation from coordinates (1) to Kruskal-Szekeres ones is

$$U = -\exp(-\kappa u), \ u = t - r^*, \tag{6}$$

$$V = \exp(\kappa v), v = t + r^*, \tag{7}$$

$$r^* = \int_{r_1}^r \frac{dr'}{f(r')}.$$
 (8)

where  $r_1$  is some constant separated from  $r_+$ . Here,  $\kappa$  is the surface gravity. For the Schwarzschild black hole,  $\kappa = \frac{1}{2r_+}$ .

We can adjust a constant  $r_1$  in such a way that

$$r^* = r + r_+ \ln(\frac{r - r_+}{r_+}). \tag{9}$$

Eqs. (6), (7) can be rewritten in the form

$$U = -\sqrt{\frac{r}{r_{+}} - 1} \exp(\frac{r - t}{2r_{+}}),\tag{10}$$

$$V = \sqrt{\frac{r}{r_{+}} - 1} \exp(\frac{r+t}{2r_{+}}).$$
(11)

Let a particle move towards the horizon, so  $\sigma = -1$  in (3). Then, it follows from (2), (3) that along the trajectory,

$$\frac{dt}{dr} = -\frac{\varepsilon}{Pf},\tag{12}$$

whence

$$t = t_0 + \varepsilon \int_r^{r_0} \frac{dr'}{P(r')f(r')},\tag{13}$$

$$u = u_0 + I_+(r, r_0), \ I_+(r, r_0) = \int_r^{r_0} dr' \frac{\varepsilon + P}{Pf},$$
(14)

$$v = v_0 + I_-(r, r_0), \ I_-(r, r_0) = \int_r^{r_0} \frac{\varepsilon - P}{Pf} dr',$$
 (15)

where  $t_0$ ,  $r_0$ ,  $u_0$  and  $v_0$  are constants. Here,  $r_0$  corresponds to the point where particle 1 starts its motion at the initial moment  $t_0$ ,  $r \leq r_0$ , the constants are related to each other according to

$$u_0 = t_0 + \int_{r_0}^{r_1} \frac{dr'}{f(r')},\tag{16}$$

$$v_0 = t_0 - \int_{r_0}^{r_1} \frac{dr'}{f(r')},\tag{17}$$

where  $u_0 = t_0 = v_0$ , if  $r_0 = r_1$ .

It is also instructive to write down equations of motion in the form

$$\frac{dU}{dr} = \frac{\kappa U}{fP} (\varepsilon + P), \tag{18}$$

$$\frac{dV}{dr} = -\kappa V \frac{(\varepsilon - P)}{Pf}.$$
(19)

As  $U_c < 0$  is arbitrary according to assumption,  $u_c$  is finite (hereafter, subscript "c" means that the corresponding quantity refers to the point of collision). Then, according to (7)  $v_c$  should be large negative when  $r \to r_+$ . We assume that  $r_0$  is fixed and finite, so  $r_0^*$  is finite as well. Meanwhile,  $t_0$  is a parameter allowed to vary.

If  $t_0$  is finite,  $u_0$  is finite as well. As the integral in (14) diverges logarithmically when  $r \to r_+$ , it means that  $u \to +\infty, U \to 0$  and the point of collision is close to the future

horizon instead of being close to the past one, contrary to assumption. To repair this, we assume that  $u_0$  is big and negative to compensate the integral which is big positive. This is achieved by the choice of big negative  $t_0$ , so formally

$$t_0 \to -\infty. \tag{20}$$

For r close to the horizon,

$$f \approx 2\kappa (r - r_+). \tag{21}$$

Then, we choose

$$u_0 = \frac{2}{\kappa} \ln(r_c - r_+) + u_1 \tag{22}$$

where  $u_1$  is some other finite constant. As a result, in the limit  $r_c \to r_+$  we see that uremains finite, so  $U \to U_c < 0$  separated from zero. As far as  $v_c$  is concerned, the integral in (15) remains finite for  $r \to r_+$ . In doing so,  $v_c \approx v_0 + I_-(r_+, r_0)$ . Taking into account (20), we see that  $v_0 \to -\infty$ ,  $v_c \to -\infty$ ,  $V_c \to 0$ , while  $t_c$  is finite. Thus in the scenario under discussion  $U_c = O(1) < 0$ , and we are indeed near the right branch of the past horizon. In doing so,

$$V_c \approx \exp(\frac{2t_0}{r_+})V^*,\tag{23}$$

where the quantity  $V^*$  remains finite nonzero in the limit when  $r_c \to r_+$ . Thus  $V_c \approx 0$  due to (20).

If collision did not occur, particle 1 would have continued its motion and hit the future horizon U = 0 in the point with very small  $V \sim V_c$ . Thus it would pass very close to the bifurcation point U = V = 0.

#### B. Properties of the trajectory of particle 2

Particle 2 moves emerges from a white hole. In this region, the Kruskal-Szekeres transformation reads

$$U = -\exp(-\kappa u), u = t - r^*,$$
 (24)

$$V = -\exp(\kappa v), v = t + r^*.$$
<sup>(25)</sup>

Particle 2 should hit the right branch of the past horizon in the point with U < 0, V = 0. Therefore, we must have  $u_0, v_0$  for this particle finite. As it moves from smaller r to bigger r, we have  $\dot{r}_2 > 0$ . Meanwhile, as particle 1 moves towards the horizon,  $\dot{r}_1 < 0$ . Thus  $\sigma_1 \sigma_2 = -1$  and near the horizon we have from (5) $\sim$ 

$$\gamma \approx \frac{2\varepsilon_1 \varepsilon_2}{f} \tag{26}$$

that diverges in the limit  $r \to r_+, f \to 0$ .

It is instructive to rewrite this formula in the form that explicitly includes the parameter  $t_0$ . It follows from (6), (7) that

$$UV = -\exp(\frac{r}{r_{+}})\frac{(r-r_{+})}{r_{+}}$$
(27)

where we used (9) and the value  $\kappa = \frac{1}{2r_+}$  for the Schwarzschild metric. Near the horizon, (21) gives us

$$f \approx -UV.$$
 (28)

Eqs. (27) and (28) are general expressions valid independently of any scenario of collision.

Iw we consider collisions and assume Scenario 1, we must take into account (23) and the fact that  $U_c = O(1)$ . Then,

$$r_c - r_+ \backsim \exp(\frac{2t_0}{r_+}) \backsim V_c. \tag{29}$$

$$\gamma \sim V_c^{-1} \sim \exp(-\frac{2t_0}{r_+}) \sim \frac{1}{r_c - r_+},$$
(30)

Thus we see that the same parameter  $t_0$  (which is large negative) governs both the behavior of the Lorentz gamma factor (hence,  $E_{c.m.}$ ) and the proximity of the point of collision to the horizon.

## IV. SCENARIO 2. PARTICLE 1 IN OUR REGION, PARTICLE 2 FROM WHITE HOLE REGION. COLLISION NEAR THE FUTURE HORIZON

Now, we consider somewhat different scenario. Particle 2 comes from the white hole region and collides with particle 1 near the future horizon in region I. See Fig. 2. In our region we can use expressions (6), (7). In contrast to the scenario described in Sec. III, where in the point of collision  $U_c < 0$  is arbitrary and  $V_c \rightarrow 0$ , now  $U_c \rightarrow 0$  and  $V_c > 0$  is arbitrary,  $t_c \rightarrow +\infty$ .



Fig. 2. Particle collisions in Scenario 2

Particle 2 crosses the past horizon in point P near the bifurcation point. Collision occurs near an arbitrary point C of the future horizon.

### A. Properties of the trajectory of particle 1

For particle 1, this can be achieved if both  $u_0$  and  $v_0$  in (14), (15) are finite. In this sense, there are no special restrictions on this trajectory. In doing so,  $t_c \to +\infty$ .

### B. Properties of the trajectory of particle 2

Let us consider now particle 2. In the scenario in question it crosses the past horizon and further moves in the outward direction, so  $\sigma_2 = +1$ , and instead of (12), we have along the trajectory of particle 2

$$\frac{dt}{dr} = +\frac{\varepsilon}{Pf} \tag{31}$$

and

$$t = t_0^{(2)} - \varepsilon \int_r^{r_0} \frac{dr'}{P(r')f(r')},$$
(32)

$$u = u_0^{(2)} + \int_r^{r_0} dr' \frac{P - \varepsilon}{Pf},$$
(33)

$$v = v_0^{(2)} - \int_r^{r_0} \frac{\varepsilon + P}{Pf} dr',$$
 (34)

where

$$u_0^{(2)} = t_0^{(2)} + \int_{r_0}^{r_1} \frac{dr'}{f(r')},$$
(35)

$$v_0^{(2)} = t_0^{(2)} - \int_{r_0}^{r_1} \frac{dr'}{f(r')}.$$
(36)

In terms of the Kruskal-Szekeres coordinates, for particle 2 we have in region I

$$\frac{dU}{dr} = -\frac{\kappa U}{fP}(\varepsilon - P),\tag{37}$$

$$\frac{dV}{dr} = \frac{\kappa V}{fP} \left(\varepsilon + P\right). \tag{38}$$

We want to adjust the trajectory of particle 2 in such a way that near the future horizon  $V_c > 0$  is arbitrary,  $U_c \approx 0$ . It is seen from (33) that the integral remains finite in the limit  $r \to r_+$ ,  $f \to 0$  whereas the integral in (34) tends to  $+\infty$ . Therefore, the only way to achieve the aforementioned properties is to take  $u_0^{(2)}$  and  $v_0^{(2)}$  big positive. Then, when particle 2 crosses the past horizon, V = 0 and  $U = U_p < 0$  with  $|U_p| \ll 1$ . Meanwhile, when it approaches  $r_c$ , both terms in (34) compensate each other and  $V_c = O(1)$ . The fact that  $|U_p| \ll 1$  means that particle 2 crosses the past horizon near the bifurcation point U = 0 = V.

More precisely, it is sufficient to choose

$$v_0^{(2)} = \frac{2}{\kappa} \left| \ln \frac{r_c - r_+}{r_+} \right| + v_2, \tag{39}$$

where  $v_2$  is some unimportant constant. In doing so, particle 2 crosses the past horizon  $r = r_+$  in the point with

$$|U_p| \sim \exp(-\frac{2t_0^{(2)}}{r_+}) \sim r_c - r_+$$
 (40)

In the limit under discussion both  $t_0^{(2)}$ ,  $v_0^{(2)} \to +\infty$ , and  $t_c^{(2)} \to +\infty$  as well. In doing so,  $v_c$  remains finite. Thus collision occurs in infinite future (from the viewpoint of a remote observer in our world) as it should be.

As now in the point of collision  $\dot{r}_1 < 0$  and  $\dot{r}_2 > 0$ , the Lorentz factor (26) is unbounded. It can be rewritten in the form

$$\gamma \sim \frac{1}{r_c - r_+} \sim \frac{1}{|U_p|}.\tag{41}$$

The same parameter  $t_0^{(2)}$  governs the behavior of  $\gamma$  (hence  $E_{c.m.}$ ) in the point of collision  $r_c$  and the proximity of the trajectory of particle 2 to the bifurcation point when it crossed the past horizon before collision.

Geometrically, it is convenient to interpret the situation in terms of coordinates r = -T, t = y under the horizon [26],

$$ds^{2} = -\frac{dT^{2}}{g} + gdy^{2} + T^{2}d\omega^{2},$$
(42)

where  $g = -f \ge 0$ . The surface of a constant r looks like a hypercylinder. Then, the condition  $t_0^{(2)} \to +\infty$  is translated to  $y_0^{(2)} \to +\infty$  for particle 2 where  $y_0^{(2)}$  is an initial value of y. Thus particle 2 should start its motion very far on the leg of the aforementioned hypercylinder.

## V. DIVERGENT $E_{c.m.}$ , UNATTAINABILITY OF THE SPEED OF LIGHT AND KINEMATIC CENSORSHIP

We would like to stress some meaningful features of the scenarios under discussion that are connected with manifestation of kinematic censorship.

#### A. Scenario 1

For any finite  $t_0$ , the energy  $E_{c.m.}$  remains finite, although it can be made as big as one likes. In the limit  $t_0 \to -\infty$  it diverges. But, clearly, the moment  $t_0$  cannot be equal to infinity literally. To some extent, this example is complimentary to that for extremal black holes. For them,  $E_{c.m.}$  is infinite if collision happens precisely on the horizon and one of colliding particles is fine-tuned. But this requires infinite proper time  $\tau$  to reach the horizon [27]. We can take it not exactly fine-tuned, then  $\tau$  is finite. But  $E_{c.m.}$  is finite as well. In our example it is coordinate (not proper) time diverges and this restricts  $E_{c.m.}$ 

There is also another interesting property of this scenario. As is mentioned above, without collision particle 1 would continue its trajectory and hit the future horizon in the point very close to the bifurcation point. The proximity to it is governed by the same parameter  $t_0$ . Formally, infinite  $|t_0|$  would mean that particle 1 passes though the bifurcation point precisely. In turn, this would mean movement along the horizon. But a massive particle

cannot move along the light-like surface, its trajectory is time-like. Thus we can say that impossibility to reach infinite  $E_{c.m.}$  is connected (although indirectly) with impossibility for a massive particle to attain the speed of light.

## B. Scenario 2

In this scenario, for particle 2 the quantity  $t_0^{(2)} \to +\infty$  along with infinite  $E_{c.m.}$ . Under the horizon, as is said above, this implies infinite distance along the leg of cylinder that is impossible. Also, the same arguments as above, show that infinity  $t_0^{(2)}$  would mean that a massive particle moves along the light-like surface. The difference between scenarios 1 and 2 consists in this context in that in the first case the past horizon is relevant whereas in the second case it is the future horizon.

## VI. DISCUSSION: COLLISION NEAR WHITE HOLE AND BEHAVIOR OF THE SCHWARZSCHILD TIME

In the above consideration, we were faced with quite unusual behavior of the Schwarzschild time in Scenario 1: although collision happens near the horizon,  $t_c$  remains finite and does not diverge when  $r_c \rightarrow r_+$ . This is contrasted with Scenario 2, where a quite standard property well known from textbooks reveals itself: when a particle approaches a black hole, its time t measured by a remote observer, diverges. To elucidate corresponding subtleties, let us get back to the calculation of  $t_c$ . It is instructive to consider this for each of two particles separately.

#### A. Particle 1

For particle 1, eq. (13) with (16), (17), (22) taken into account, gives us that the big negative divergent  $t_0$  compensates the big positive integral in such a way that the net outcome is finite in the limit  $r_0 \rightarrow r_+$ . Thus  $t_c$  remains finite. We would like to stress that this conclusion has nothing to do with the existence of a white hole region under the past horizon and is valid independently of whether or not we consider collisions. However, it relies heavily on distinction between the future horizon and the past horizon. In the first case, when a particle approaches the future horizon, i.e. falls in a black hole,  $t \to \infty$ .

### B. Particle 2

It is instructive to look for the trajectory of particle 2 even in a more wide context. Let us trace how  $t^{(2)}(r)$  changes along the trajectory of particle 2. In the region outside the horizon  $t^{(2)}(r)$  is finite for any  $r > r_+$ . This allows us to arrange collision between particles 1 and 2 in some point  $r_c > r_+$ , so that  $t^{(2)}(r_c) = t^{(1)}(r_c)$ . Meanwhile, it follows from (31) that along the trajectory

$$t^{(2)}(r) = t_0^{(2)} - \int_r^{r_c} \frac{\varepsilon_2 dr}{P(r')f(r')},$$
(43)

where  $t_0^{(2)} = t^{(2)}(r_c)$ . When we trace behavior of  $t^{(2)}(r)$  back in time and  $r \to r_+$ ,  $t^{(2)}(r) \to -\infty$ . In other words, a particle crossed the past horizon in an infinite past.

The situation is opposite to that for ingoing particle 1 for which t(r) is finite for any  $r > r_+$  but  $t(r) \to +\infty$  when  $r \to r_+$ .

The contents of the present subsection is a counterpart of pages 114 - 115 of [28], where appearance of massless particles from a white hole was discussed. It was explained there that an observer in the outer world is able to see all interior of a white hole starting from the singularity. In somewhat different language, we argued above that a similar phenomenon is valid for massive particles. An observer residing at any  $r > r_+$  is able to catch particles coming from the interior and, as a consequence, to see all the interior of a white hole including the singularity.

Thus it is quite typical for outgoing particle 2 to have  $t_c$  finite. What is more unusual is that  $t_c$  remains finite in the limit  $r_c \to r_+$  for ingoing particle 1. It would seem that this is contrasted with the behavior known even from textbooks according to which  $t_c \to \infty$  in such circumstances. However, one should bear in mind two qualitative features here: (i) now the horizon is a past horizon, not a future one, (ii) collision occurs near a white (not a black) hole. Mathematically, finite  $t_c$  appears due to compensation of a big constant  $t_0^{(1)}$ and big (almost diverging) integral in (13) as is explained above in III.

## VII. HOW DOES THE PICTURE LOOK LIKE IN THE LEMAÎTRE COORDINATES?

Apart from the Kruskal-Szekeres coordinates, there exists one more popular frame that remains regular near the event horizon. This is the Lemaître frame. It is quite instructive to describe in this frame particle collisions under discussion. The connection between the Lemaître frame and the original curvature coordinate is well known and is described in textbooks (see, e.g. Sec. 102 in [29]). Here, one can distinguish between two different cases - the contracting Lemaître frame and the expanding one. The contracting Lemaître frame is composed by particles that fall freely from infinity with  $\varepsilon = 1$ . The expanding Lemaître frame is composed by particles with  $\varepsilon = 1$  that move from the singularity, cross the white hole region and enter our part of space. In our context, where the presence of a white hole is an essential ingredient, it is the expanding one that is preferable since it encompasses both the white hole region ( $T^+$  one according to the terminology of [23]) and the outer one (Rregion). Meanwhile, the contracting Lemaître frame is unable to describe the history of a particle in the  $T^+$  region.

In the expanding Lemaître frame we introduce new coordinates

$$\hat{t} = t - \int_{r_c}^r \frac{dr\sqrt{1-f}}{f},\tag{44}$$

where we choose the constant of integration such that  $\hat{t} = t$  at  $r = r_c$ ,

$$\rho = -t + \int_{r_c}^r \frac{dr}{f\sqrt{1-f}},\tag{45}$$

in which the metric reads

$$ds^{2} = -d\hat{t}^{2} + (1 - f)d\rho^{2} + r^{2}d\omega^{2}.$$
(46)

Along the trajectory of an outgoing particle 2, we have relations (31), (43), and

$$\hat{t} = t_0^{(2)} + \varepsilon \int_{r_c}^r \frac{dr'}{f(r')\sqrt{\varepsilon^2 - f}} - \int_{r_c}^r \frac{dr\sqrt{1 - f}}{f}.$$
(47)

Near the horizon when an initial point  $r \to r_+ + 0$ , the metric function  $f \to 0$  and both integrals compensate each other, so  $\hat{t}_c$  is finite for any finite  $t_0^{(2)}$ . This is quite natural as this frame is designed just for description of outgoing particles. In other words, divergent  $t_c(r)$  for  $r \to r_+$  is combined with the finiteness of  $\hat{t}(r)$ , as it should be. This agrees with the behavior of t(r) described in Sec. VI.

For an ingoing particle 1,

$$t = t_0^{(1)} - \varepsilon \int_{r_0}^r \frac{dr'}{f(r')P(r')}$$
(48)

and

$$\hat{t} = t_0^{(1)} - \varepsilon \int_{r_0}^r \frac{dr'}{f(r')\sqrt{\varepsilon^2 - f}} - \int_{r_0}^r \frac{dr\sqrt{1 - f}}{f}.$$
(49)

As particle 1 starts at  $r = r_0 > r_+$ , both integrals have the same sign and diverge when  $r \to r_+$ . More aspects of relation between different frames and particles moving in the background under discussion can be found in [31].

Let us consider the same scenarios as before.

#### A. Scenario 1

Particle 2 appears from the white hole region and moves in the outward direction where it meets particle 1 close to the past horizon. Particle 1 is ingoing. For it, (49) holds. Particle 2 is outgoing, so (47) holds. We want both particles to meet, so  $\hat{t}_1 = \hat{t}_2$ . To achieve this, we must adjust constants  $t_0^{(1)}$ ,  $t_0^{(2)}$ . Then, for collisions near the horizon, we take finite  $t_0^{(2)}$ and  $t_0^{(1)} \to -\infty$ . Thus we again come to the same conclusion obtained above in the analysis that uses Kruskal-Szekeres coordinates.

We see that there is full complementarity between pictures of particle motion and collision in both frames. The ingoing particle is "unnatural" with respect to the expanding Lemaître and has formally divergent Lemaître time when it starts its motion. Note, that collision itself occurs for a finite Lemaître time even if the collision energy is unbounded.

#### B. Scenario 2

Now, there are no special restriction imposed on particle 1,  $t_0^{(1)}$  is finite. As a result, for the point of collision  $\hat{t} \to +\infty$ . For particle 2, we choose  $t_0^{(2)} \to +\infty$  to achieve  $\hat{t}_1 = \hat{t}_2$ . Note that is this scenario the Lemaître time of the collision diverges for divergent energy.

#### VIII. PARTICLE DECAY NEAR SINGULARITY

Up to now, we mainly considered particle collisions near the horizon. Meanwhile, it is also of interest to consider an immediate vicinity of the singularity. In doing so, instead of collisions, we will discuss more simple case, when particle 0 decays to fragments 1 and 2. We will see that even in this simplified settings, results are quite interesting and have nontrivial consequences. In the present article we are concerned with processes outside the horizon. However, it makes sense to add the decay in question into a general scheme, provided we consider such a decay in the white hole region. Motivation comes from the fact that after decay new particles can appear in the outer region.

As we are interested in the region beyond the horizon, f = -g < 0, it is convenient to use the metric in the form (42).

As before, we assume that all angular momenta are zero. Then, equations of motion for a free particle read

$$m\dot{y} = -\frac{E}{g},\tag{50}$$

$$m\dot{T} = \sqrt{E^2 + gm^2},\tag{51}$$

where E is a constant. Now, the quantity E has the meaning of momentum (not energy) along the y-direction and  $m\dot{T}$  has the meaning of energy (up to the factor g). As now the metric is homogeneous and non-static, it is E which is conserved whereas the energy is not. However, we assume that in the point of decay conservation laws hold:

$$E_0 = E_1 + E_2, (52)$$

$$\sqrt{E_0^2 + gm_0^2} = \sqrt{E_1^2 + gm_1^2} + \sqrt{E_2^2 + gm_2^2}.$$
(53)

After some algebra, one obtains

$$E_1 = \frac{b_1}{2m_0^2} E_0 \pm \frac{\sqrt{d}\sqrt{E_0^2 + m_0^2 g}}{2m_0^2},\tag{54}$$

$$E_2 = \frac{b_2}{2m_0^2} E_0 \mp \frac{\sqrt{d}\sqrt{E_0^2 + m_0^2 g}}{2m_0^2},\tag{55}$$

where

$$b_{1,2} = m_0^2 + m_{1,2}^2 - m_{2,1}^2, (56)$$

$$d = b_1^2 - 4m_0^2 m_1^2 = b_2^2 - 4m_0^2 m_2^2.$$
(57)

These formulas are similar to eqs. 45, 46 and 51, 52 of [30] where decay outside the horizon was considered. However, now they hold for a non-static metric.

Near singularity  $g \to \infty$ . Then,

$$E_{1,2} \sim \pm \frac{\sqrt{d}}{2m_0} \sqrt{g} \tag{58}$$

independently of  $E_0$ .

The quantity E under the horizon has the meaning of momentum, not energy. It conserves due to homogeneity of the metric in this region. When a particle crosses the horizon, the value of this integral of motion remains the same but the meaning changes: momentum "converts" into energy. Therefore, if  $E_{1,2}$  is indefinitely big in the point of decay, it remains indefinitely big in the vicinity of the horizon from any side of it. Thus if decay occurs inside a white hole region near the singularity, particles with unbounded E can cross the horizon and enter the outer R region with the same unbounded E: particle with E > 0 enters the right R region and that with E < 0 enters the left R region. As a result, flux of such particles can cause strong backreaction and destroy the past horizon.

Although detailed consideration in the black hole region is beyond the scope of the present work, it is to the point to make short comparison. The aforementioned destruction does not happen for the black hole horizon (future horizon) since in the  $T^-$  region (we follow classification developed in [32]) decay near the singularity occurs when particles already crossed the horizon in the past. However, such strong backreaction can destroy, in principle, the black hole singularity. Thus the effect of decay in this case is very similar to that due to particle collisions near the singularity [33]. In this sense, there is some complementarity: decay near the white hole singularity seems to destroy the white hole horizon, decay near the black hole singularity destroys the singularity itself.

It is worth stressing that the results in question are obtained pure classically and do not require resort to quantum effects [34], [35].

## IX. SUMMARY AND CONCLUSIONS

High energy collisions in the Schwarzschild metric are possible due to participation of a particle coming from a white hole region that are absent in the standard BSW effect. The main properties of such collisions are summarized in Table I. We indicated in scenario 1 proximity of particle 1 to the bifurcation point. However, the reservation is in order here that this is "virtual" proximity for the trajectory which would continue without collision with particle 2. In line 2 we indicate the Schwarzschild time of collision  $t_c$ . We would like to stress that the signs  $-\infty$  or  $+\infty$  are pure conditional and used for the sake of brevity. They mean that in the horizon limit  $r_c \rightarrow r_+$  the corresponding quantity tends to plus or minus infinity, although for any  $r_c > r_+$  it is finite.

	Scenario 1	Scenario 2
$\hat{t}_c$	finite	$+\infty$
$t_c$	finite	$+\infty$
$t_{0}^{(1)}$	$-\infty$	finite
$t_0^{(2)}$	finite	$+\infty$
proximity to bifurcation point	particle 1 (virtually)	particle 2

Table I. Properties of constants  $t_0^{(1)}$ ,  $t_0^{(2)}$  and behavior of the Lemaître time in the point of collision with unbounded energy  $E_{c.m.}$ .

We would like to stress that for particle 2  $t_0^{(2)}$  cannot be identified as an initial moment. Under the horizon, where this particle merges from, the variable t has a spatial character, t = y, so its divergent value means that it starts its motion very far along the leg of a corresponding hypercilinder in the metric (42).

We traced in detail how the same conditions govern both the indefinite growth of  $E_{c.m.}$  of colliding particles, proximity of corresponding events to the horizon and unattainability of a speed of light for a massive particle. In this sense, we reveal physical meaning of kinematic censorship for scenarios under discussion.

We also considered process of particle decay near the singularity and found that it leads to indefinitely large energies of its debris. However, literally infinite energy requires a decay exactly at a singularity where known physics fails. This result is also obtained purely classically.

We described the processes under discussion in the Kruskal-Szekeres frames as well as in the expanding Lemaître one. Such a frame is rather rarely used in literature but now its using was motivated by a context since it is this version of the Lemaître one that is able to describe history of particles in the regions of interest.

It is reasonable to believe that unboundedness of the collision and decay energies from above (despite that literal infinity cannot be achieved) can be considered as a new argument against the existence of astrophysical white holes, using pure classical arguments and including processes both near the past horizon and near the white hole singularity. This issue needs further investigations.

## X. ACKNOWLEDGEMENT

We recall with gratitude A. A. Starobinsky with whom we had time to discuss this work before his sudden death.

- M. Bañados, J. Silk and S.M. West, Kerr black holes as particle accelerators to arbitrarily high energy, Phys. Rev. Lett. 103 (2009) 111102 [arXiv:0909.0169].
- [2] O.B. Zaslavskii, Acceleration of particles as universal property of rotating black holes, Phys. Rev. D 82 (2010) 083004 [arXiv:1007.3678].
- [3] A. Grib and Yu. V. Pavlov, Are black holes totally black? Grav. Cosmol. 21 (2015) 13, [arXiv:1410.5736].
- [4] O. B. Zaslavskii, On white holes as particle accelerators, Gravitation and Cosmology, 24 (2018) 92 [arXiv:1707.07864].
- [5] T. Piran, J. Katz, and J. Shaham, High efficiency of the Penrose mechanism for particle collision, Astrophys. J. 196 (1975) L107.
- [6] T. Piran and J. Shaham, Production of gamma-ray bursts near rapidly rotating accreting black holes, Astrophys. J. 214 (1977) 268.
- [7] T. Piran and J. Shanam, Upper bounds on collisional Penrose processes near rotating black hole horizons, Phys. Rev. D 16 (1977) 1615.
- [8] H.V.Ovcharenko, O.B.Zaslavskii, High energy head-on particle collisions near event horizons: classification of scenarios, Int. Journ. of Mod. Phys. D [arXiv:2404.03364].
- [9] Yu. V. Pavlov and O. B. Zaslavskii, Kinematic censorship as a constraint on allowed scenarios of high energy particle collisions, Grav. Cosmol. 25 (2019) 390, [arXiv:1805.07649].
- [10] H. V. Ovcharenko and O. B. Zaslavskii, Bañados-Silk-West effect with finite forces near different types of horizons: general classification of scenarios, Phys. Rev. D 108, 064029 (2023) [arXiv:2304.13087].

- [11] A. A. Grib, Yu. V. Pavlov, O. F. Piattella, High energy processes in the vicinity of the Kerr's black hole horizon Int. J. Mod. Phys. A 26 (2011) 3856 [arXiv:1105.1540].
- [12] E. Hackmann, H. Nandan, P. Sheoran, Particle collisions near static spherically symmetric black holes, Phys. Lett. B 810 (2020) 135850 [arXiv:2006.05045].
- [13] K. Ogasawara and N. Tsukamoto, Effect of the self-gravity of shells on a high energy collision in a rotating Bañados-Teitelboim-Zanelli spacetime, Phys. Rev. D 99 (2019) 024016 [arXiv:1810.03294].
- [14] O. B. Zaslavskii, On special case of the Banados-Silk-West effect. Phys. Rev. D 102 (2020) 044051. [arXiv:2006.11552].
- [15] O. B. Zaslavskii, The Banados-Silk-West effect with immovable particles near static black holes and its rotational counterpart, Grav. and Cosmology 29 (2023) 74, [arXiv:2207.03213].
- [16] I. D. Novikov, Delayed explosion of a part of the Fridman Universe, and quasars, Astr. Zhurnal
  41 (1964) 1075 (Soviet Astronomy AJ 8 (1965), 857).
- [17] Y. Ne'eman, Expansion as an energy source in quasi-stellar radio sources, Astrophys. J. 141 (1965), 1303.
- [18] J. V. Narlikar, K. M. V. Appa Rao and N. Dadhich, High energy radiation from white holes, Nature 251 (1974) 590.
- [19] Patil, M., Harada, T.: Extremal Kerr white holes as a source of ultra high energy particles.
   Phys. Rev. D 102 (2020) 024002 arXiv:2004.12874.
- [20] O. B. Zaslavskii. Super-Penrose process for extremal charged white holes. Modern Physics Letters A. 36 2150020 (2021), [arXiv:2005.11090].
- [21] O. B. Zaslavskii, Super-Penrose process for extremal rotating neutral white holes, Gen. Relat.
   Grav. 52 (2020) 104, [arXiv:2005.14241].
- [22] D. M. Eardley, Death of White Holes in the Early Universe, Phys. Rev. Lett. 33 (1974) 442.
- [23] V. P. Frolov and I. D. Novikov, Black Hole Physics: Basic Concepts and New Developments (Kluwer, 1998).
- [24] R. Gaur and M. Visser, Black holes, white holes, and near-horizon physics, J. High Energ. Phys. 2024, 172 (2024) [arXiv:2304.10692].
- [25] C.W. Misner, K. S. Thorne, and J. A. Wheeler, Gravitation (Freeman, San Francisco, 1973).
- [26] I.D. Novikov, Note on the space-time metric inside the Schwarzschild singular sphere, Sov.Astron. AJ5 (1961) 423 (Astron.Zh. 38, 564(1961)).

- [27] T. Jacobson, T.P. Sotiriou, Spinning Black Holes as Particle Accelerators, Phys. Rev. Lett. 104, 021101 (2010), [arXiv:0911.3363].
- [28] Ya. B. Zel'dovich and I. D.Novikov, Stars and relativity (University of Chicago Press, Chicago, Illinois, 1971).
- [29] L. D. Landau and E. M. Liefshitz, The Classical Theory of Fields (Pergamon Press, New York, 1971).
- [30] O.B. Zaslavskii, On general properties of the Penrose process with neutral particles in the equatorial plane, Phys. Rev. D 108, 084022 (2023), [arXiv:2307.06469]052].
- [31] A. Toporensky, O. Zaslavskii, General radially moving references frames in the black hole background, Eur. Phys. J. C 83 (2023) 225.
- [32] I.D. Novikov, R-and T-regions in space-time with spherically symmetric space, Soobshch.
   GAISh 132 (1964) 43 [Gen.Relativ.Gravit. 33 (2001) 2259].
- [33] A. Radosz, A. V. Toporensky, O. B. Zaslavskii, On particle dynamics near the singularity inside the Schwarzschild black hole and T-spheres, Eur. Phys. J. C 83 (2023) 650, [arXiv:2301.11651].
- [34] Ya. B. Zel'dovich, I. D. Novikov, and A. A. Starobinskii, Quantum effects in white holes, Zh. Eksp. Teor. Fiz. 66 1897 (1974) (Sov. Phys.-JETP 39 (1974), 933).
- [35] R. Wald and Sriram Ramaswamy, Particle production by white holes, Phys. Rev. D 21, 2736 (1980).