

The spectral density of astrophysical stochastic backgrounds

Enis Belgacem,^{1,2,*} Francesco Iacovelli,^{1,2,†} Michele Maggiore,^{1,2,‡} Michele Mancarella,^{3,§} and Niccolò Muttoni^{1,2,¶}

¹*Département de Physique Théorique, Université de Genève, 24 quai Ernest Ansermet, 1211 Genève 4, Switzerland*

²*Gravitational Wave Science Center (GWSC), Université de Genève, CH-1211 Geneva, Switzerland*

³*Aix-Marseille Université, Université de Toulon, CNRS, CPT, Marseille, France*

(Dated: November 7, 2024)

We provide a detailed derivation of the spectral density of the stochastic background generated by the superposition of coalescing compact binaries. We show how the expression often used in the literature emerges from an average over the extrinsic parameters of the binaries (times of arrival, polarization angles, arrival directions and orbit inclinations) and how the Stokes parameters related to circular and linear polarization are set to zero by such averaging procedure. We then consider the effect of shot noise, i.e. the fact that for the superposition of a finite number of sources these averages are only approximate, and we show how it generates circular and linear polarizations (even for isotropic backgrounds) as well as spatial anisotropies, and we compute them explicitly for a realistic population of binary black holes and binary neutron stars.

CONTENTS

I. Introduction	1
II. Statistical characterization of discrete stochastic backgrounds	2
A. Basic definitions	2
B. Averages over extrinsic and intrinsic parameters	3
III. Computation of the spectral density of the astrophysical background	5
A. Average over polarization angles	6
B. Average over arrival times	7
C. Average over arrival directions	8
D. Average over orbit inclinations	8
E. Summary	10
IV. Energy density of anisotropic and polarized astrophysical backgrounds	11
V. Effect of shot noise	12
A. Polarization from shot noise	13
B. Linear polarizations and spatial isotropy	16
C. Numerical results	17
D. Anisotropies from shot noise	17
VI. Conclusions	19

I. INTRODUCTION

Stochastic backgrounds of gravitational waves (GWs) are among the main targets of present and future GW

experiments. Such backgrounds can have an astrophysical origin, or being generated by cosmological processes in the early Universe (see Refs. [1–8] for reviews). Pulsar timing arrays have already provided evidence for a stochastic GW background which has a possible explanations as due to the superposition of signals from supermassive black hole binaries [9–12]. The astrophysical background due to the superposition of compact binaries coalescences (CBCs) might already be detectable in the most advanced stage of current (second generation) ground-based GW detectors [13–15]. At third-generation (3G) detectors, such as Einstein Telescope (ET) [16–19] and Cosmic Explorer (CE) [20–23], the astrophysical stochastic background from CBCs will be detectable with a very high signal-to-noise ratio [15, 24, 25]. Beside carrying important astrophysical information by itself, the astrophysical GW background can be a foreground to stochastic backgrounds of cosmological origin, and also from this point of view is therefore important to characterize it accurately, in view of subtracting it [26–40].

A first aim of this paper is to put on a firmer theoretical ground the derivation of a basic quantity used in the description of the astrophysical background from CBCs, namely its spectral density, presenting an accurate derivation of its expression. We will see in particular how it emerges from an average over the times of arrival, polarization angles, arrival directions and orbit inclinations of an ensemble of CBCs and how, in the process, the Stokes parameters related to circular and linear polarization are averaged to zero. We will then use our formalism to take into account the effect of shot noise, i.e. of the fact that any given realization of the astrophysical background, with a finite number of events, only corresponds to an imperfect average over these parameters, so it leaves a residual in quantities that otherwise would be averaged to zero [41, 42].

The paper is organized as follows. In Sec. II we recall standard definitions on discrete stochastic backgrounds and we define the averages over extrinsic and intrinsic parameters of an ensemble of CBCs. In Sec. III we compute the spectral density of the astrophysical background, by

* enis.belgacem@unige.ch

† francesco.iacovelli@unige.ch

‡ michele.maggiore@unige.ch

§ mancarella@cpt.univ-mrs.fr

¶ niccolo.muttoni@unige.ch

performing explicitly the averages over the extrinsic parameters. In Sec. IV we compute the energy density of a generic (anisotropic and polarized) stochastic background, showing that there is no energy density associated Stokes parameters describing linear and circular polarizations (a fact that sometimes gives rise to confusion in the literature). In Sec. V we use our formalism to describe the generation of polarization and anisotropies from shot noise. We will see, in particular, that shot noise can generate linear polarization even for an isotropic distribution of sources (contrary to statements in the literature). We will then estimate numerically these shot-noise effects on a realistic population of BBHs and BNSs. Section VI contains our conclusions.

II. STATISTICAL CHARACTERIZATION OF DISCRETE STOCHASTIC BACKGROUNDS

In this section we begin by introducing standard definitions for the astrophysical stochastic background, in order to fix the notation, and we will then discuss the averages that are involved in its characterization, distinguishing between the averages over extrinsic and intrinsic parameters.¹

A. Basic definitions

A superposition of GWs coming from all directions, with propagation directions labeled by a unit vector $\hat{\mathbf{n}}$, can be written as

$$h_{kl}(t, \mathbf{x}) = \int_{-\infty}^{\infty} df \int_{S^2} d^2 \hat{\mathbf{n}} \times \sum_{A=+, \times} \tilde{h}_A(f, \hat{\mathbf{n}}) e_{kl}^A(\hat{\mathbf{n}}) e^{-2\pi i f(t - \hat{\mathbf{n}} \cdot \mathbf{x}/c)}, \quad (1)$$

where $k, l = 1, 2, 3$ are spatial indices (which, in the TT gauge, reduce to indices taking two values, in the transverse plane), $A = +, \times$ labels the two polarizations, and e_{kl}^A are the polarization tensors, normalized as

$$e_{kl}^A(\hat{\mathbf{n}}) e_{kl}^{A'}(\hat{\mathbf{n}}) = 2\delta^{AA'}, \quad (2)$$

where the sum over the repeated spatial indices is understood (we follow the notation and conventions in Ref. [2]). A stochastic GW background can only be characterized statistically, through an ensemble average. Conceptually, this ensemble average is an average over many different realizations of the Universe. Of course, in practice we

only have a single realization of the Universe; cosmological backgrounds (such as, for instance, the background produced by the amplification of vacuum fluctuations in the early Universe), can be considered as due to the superposition of an effectively infinite number of signals, so they are fully characterized by such ensemble averages, through the corresponding correlators (or just the two-point correlator, for a Gaussian background). In contrast, the superposition of astrophysical signals will retain some dependence on the specific realization. For instance, a stochastic background due to the superposition of 10^4 BBHs (that correspond to the BBH coalescences taking place in the Universe in about 1 month, given our current knowledge of their rate [43]) will retain some dependence on the specific sample of sources that coalesced during that period; a subsequent data stretch of one more month will effectively correspond to a different realization of the same discrete stochastic process. In the following, we will use the notation $\langle \dots \rangle_U$ for the average over different realizations of the Universe. In Sec. II B we will discuss how to perform the ‘‘Universe’’ averages for an ensemble of CBCs, separating it into averages over extrinsic and intrinsic parameters.

If a stochastic background of GWs is stationary, isotropic and unpolarized, its two-point correlator can be written as

$$\langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_U = \delta(f - f') \frac{\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')}{4\pi} \delta_{AA'} \frac{1}{2} S_h(f), \quad (3)$$

where $\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')$ is a Dirac delta over the two-sphere,

$$\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}') = \delta(\phi - \phi') \delta(\cos \theta - \cos \theta'), \quad (4)$$

and (θ, ϕ) are the polar angles that define $\hat{\mathbf{n}}$. The function $S_h(f)$, defined by this expression, is the spectral density of the stochastic background.

Actually, Eq. (3) assumes an infinite observation time. Any realistic observation will only last for a finite observation time T , in which case the Dirac delta $\delta(f - f')$ must be replaced by a regularized Dirac delta, i.e.

$$\langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_U = \delta_T(f - f') \frac{\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')}{4\pi} \delta_{AA'} \frac{1}{2} S_h(f), \quad (5)$$

where

$$\delta_T(f - f') = \int_{-T/2}^{T/2} dt e^{-2\pi i(f - f')t}. \quad (6)$$

Note that, for $f = f'$, $\delta_T(f - f')$ takes the value

$$\delta_T(0) = T. \quad (7)$$

We will always be interested in situations where the frequencies f of interest are such that $fT \gg 1$. For ground-based detectors, even of third generation, that operates at $f \gtrsim (5 - 10)$ Hz, this is very well satisfied already for stretches of data of about 60 s (as typically used to estimate the PSD of a detector), and is extremely well

¹ This section has some overlap with the corresponding introductory section of the companion paper [40], where we use this formalism to study the subtraction of the astrophysical background in a two-detector correlation.

satisfied if we integrate the signal to observation periods of months or years. Therefore, the details of the regularization of the Dirac delta are irrelevant.² The main virtue of using a regularized expression is the possibility of using Eq. (7). Otherwise, we will use it as a normal Dirac delta, e.g. writing

$$\int_{-\infty}^{\infty} df' \delta_T(f - f') G(f') \simeq G(f), \quad (8)$$

and in fact, with an abuse of notation, we will in general replace here \simeq by an equal sign.

In general, we expect that the assumption of stationarity will be very well satisfied. The statistical properties of a population of astrophysical sources only change on cosmological timescales (say, millions of years), which are huge compared to the observation time T , and this is even more true for a background of cosmological origin; so, the underlying process is invariant under time translations (over a timescale of the order of the observation time, and in fact even much bigger than it) and the only deviation from a distribution of events uniform in time will be due to shot noise, i.e. to the specific finite realization. In contrast, a stochastic GW background need not be isotropic, nor unpolarized. The most general form of the two-point correlator is then [44] (see also Refs. [4, 45, 46] for reviews)

$$\begin{aligned} & \langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_U \\ &= \delta_T(f - f') \frac{\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')}{4\pi} \frac{1}{2} H_{AA'}(f, \hat{\mathbf{n}}). \end{aligned} \quad (9)$$

Note that we still assume that signals coming from different directions are not correlated, i.e. the correlator is still proportional to $\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')$; however, the possibility of an anisotropy is encoded in the fact that the function $H_{AA'}(f, \hat{\mathbf{n}})$ could depend on $\hat{\mathbf{n}}$; the possibility of a polarization of the background, instead, is encoded in the fact that $H_{AA'}(f, \hat{\mathbf{n}})$ is a 2×2 matrix in the polarization indices, not necessarily proportional to $\delta_{AA'}$. Taking the complex conjugate of Eq. (9) we see that $H_{AA'}(f, \hat{\mathbf{n}})$ is a Hermitian matrix, so it can be decomposed (with real coefficients) into the basis made by the identity matrix and the Pauli matrices,

$$\begin{aligned} H_{AA'}(f, \hat{\mathbf{n}}) &= I(f, \hat{\mathbf{n}}) \delta_{AA'} + U(f, \hat{\mathbf{n}}) \sigma_{AA'}^1 \\ &+ V(f, \hat{\mathbf{n}}) \sigma_{AA'}^2 + Q(f, \hat{\mathbf{n}}) \sigma_{AA'}^3, \end{aligned} \quad (10)$$

where the three Pauli matrices are given by

$$\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

In matrix form

$$H(f, \hat{\mathbf{n}}) = \begin{pmatrix} I(f, \hat{\mathbf{n}}) + Q(f, \hat{\mathbf{n}}) & U(f, \hat{\mathbf{n}}) - iV(f, \hat{\mathbf{n}}) \\ U(f, \hat{\mathbf{n}}) + iV(f, \hat{\mathbf{n}}) & I(f, \hat{\mathbf{n}}) - Q(f, \hat{\mathbf{n}}) \end{pmatrix}. \quad (11)$$

² Furthermore, we will show below that the average over the times of arrival gives precisely the expression (6).

The coefficients of this decompositions define the Stokes parameters of the GW stochastic background, and are real functions of f and $\hat{\mathbf{n}}$ describing intensity (I), linear polarization (U and Q) and circular polarization (V).

For the stochastic background generated by the superposition of astrophysical sources in an observation time T , we can write

$$\tilde{h}_A(f, \hat{\mathbf{n}}) = \sum_{i=1}^{N_{\text{ev}}} \tilde{h}_{A,i}(f) \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i), \quad (12)$$

where $\hat{\mathbf{n}}_i$ is the propagation direction of the i -th signal, and N_{ev} is the number of events reaching the detector in the observation time T . Then, Eq. (1) becomes

$$\begin{aligned} h_{kl}(t, \mathbf{x}) &= \sum_{i=1}^{N_{\text{ev}}} \int_{-\infty}^{\infty} df \\ &\times \sum_{A=+, \times} \tilde{h}_{A,i}(f) e_{kl}^A(\hat{\mathbf{n}}_i) e^{-2\pi i f(t - \hat{\mathbf{n}}_i \cdot \mathbf{x}/c)}. \end{aligned} \quad (13)$$

B. Averages over extrinsic and intrinsic parameters

We now discuss how to perform the average $\langle \dots \rangle_U$ for an astrophysical background. We will use a language appropriate to CBCs, although some aspects of the discussion below hold for more general astrophysical backgrounds, e.g. due to supernovae. An essential step is to separate the parameters of the waveform of a CBC into two groups, extrinsic and intrinsic parameters, and perform first the average over the extrinsic parameters.

In this work, we define the extrinsic parameters of a CBC (labeled by an index i), to be: its arrival time t_i (defined, e.g., as the time of arrival of the peak of the waveform, or the time of entry of the signal in the detector bandwidth); the polarization angle ψ_i ; the propagation direction $+\hat{\mathbf{n}}_i$ of the GW (so that the direction of the source in the sky is $-\hat{\mathbf{n}}_i$); and the inclination of the orbit, defined by the angle ι_i between the normal to the orbit and the line-of-sight (we will actually rather use $\cos \iota_i$).³ So, for a set N_{ev} of events, the extrinsic parameters are

$$\{t_i, \psi_i, \hat{\mathbf{n}}_i, \cos \iota_i\}_{i=1, \dots, N_{\text{ev}}}. \quad (14)$$

The intrinsic parameters are all the other parameters of the binary, such as the component masses, spins, luminosity distance, orbit eccentricity, tidal deformability (for neutron stars), etc.

A first rationale for this distinction is that events characterized by different extrinsic parameters (but the same intrinsic parameters) can be intuitively seen as different

³ In the case of precessing spins the orbital plane is not fixed, and the notion of ι becomes time-dependent. Anyway, all computations below will remain valid replacing ι by θ_{JN} , defined as the angle between the line-of-sight and the total angular momentum.

realizations of the same event; e.g., signals all corresponding to a BBH with given masses, spin, etc., but with different arrival times (within the time span of the observation), or arriving from different directions in the sky, can be considered as different realizations of the same event. In contrast, once we start sampling e.g. the mass distribution, there is no longer much sense in which we could say that an ensemble of signals always refers to the same BBH, just extracted with different component masses.

More technically, the averages over extrinsic parameters are made with a flat measure, that reflects some underlying symmetry principle. For instance, the average over the time of arrival t_i of a generic quantity (denoted by the dots) is just performed with a flat measure

$$\langle \dots \rangle_{t_i} = \frac{1}{T} \int_{-T/2}^{T/2} dt_i (\dots), \quad (15)$$

(where T is the observation time), reflecting the invariance under time translations of the probability of having a BBH or BNS merger during the observation time T , at least on the timescale T of the observation, which is typically at most months or years, and therefore is extremely small compared to the timescales over which the populations of BBHs and BNSs evolve.⁴ Similarly, the polarization angle parameterizes the freedom of the observer to choose the axes, in the plane transverse to the propagation direction of the i -th signal, with respect to which the plus and cross polarizations of the i -th signal are defined, so the average over the polarization angle ψ_i is naturally performed with the flat measure

$$\langle \dots \rangle_{\psi_i} = \int_0^{2\pi} \frac{d\psi_i}{2\pi} (\dots). \quad (16)$$

The inclination angle of the orbit with respect to the line-of-sight must also be distributed uniformly on the sphere, since the binary “knows nothing” about the location of the observer, and there is no reason why it should have preferential orientations with respect to the observer’s line of sight,⁵ so it is averaged as

$$\langle \dots \rangle_{\cos \iota_i} = \frac{1}{2} \int_{-1}^1 d \cos \iota_i (\dots). \quad (17)$$

In many situations it is also natural to assume that, at least in a first approximation, extragalactic sources such

as BBHs and BNSs are uniformly distributed in the sky corresponding, in the average over this extrinsic parameter, to

$$\langle \dots \rangle_{\hat{\mathbf{n}}_i} = \int_{S^2} \frac{d^2 \hat{\mathbf{n}}_i}{4\pi} (\dots), \quad (18)$$

where S^2 denotes the 2-sphere. However, the appropriateness of this assumption really depends on the context; in particular, if one is interested in the anisotropies of the astrophysical background and in its correlation with the galaxy field [41, 47–60], this average would select the “monopole” term of the distribution, while in this case one would be interested exactly in the small deviations from the uniform distributions, described by the higher multipoles; so, more generally, as long as we assume that the integrations over arrival directions of different signals factorize, for an anisotropic background we will write

$$\langle \dots \rangle_{\hat{\mathbf{n}}_i} = \int_{S^2} \frac{d^2 \hat{\mathbf{n}}_i}{4\pi} \mu(\hat{\mathbf{n}}_i) (\dots), \quad (19)$$

for some function $\mu(\hat{\mathbf{n}}_i)$, normalized as

$$\int_{S^2} \frac{d^2 \hat{\mathbf{n}}}{4\pi} \mu(\hat{\mathbf{n}}) = 1, \quad (20)$$

and that could reflect for instance a prior given by a galaxy catalog.

Therefore, assuming that the integration over all these parameters are independent, the average over the extrinsic parameters is defined as

$$\begin{aligned} \langle \dots \rangle_{\text{ext}} &= \prod_{i=1}^{N_{\text{ev}}} \int_{-T/2}^{T/2} \frac{dt_i}{T} \int_0^{2\pi} \frac{d\psi_i}{2\pi} \int_{-1}^1 \frac{d \cos \iota_i}{2} \\ &\times \int_{S^2} \frac{d^2 \hat{\mathbf{n}}_i}{4\pi} \mu(\hat{\mathbf{n}}_i) (\dots). \end{aligned} \quad (21)$$

As we will see below, the average over these extrinsic parameters allow us to extract, from the correlator $\langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_U$, Dirac delta functions or Kronecker’s deltas, or other simple mathematical structures.

In contrast, intrinsic parameters, such as the masses and spins of the component stars, the distance to the binary, etc., must be sampled according to non-trivial distributions that reflect specific astrophysical properties of the population, rather than uniform distributions reflecting symmetry principles.⁶

⁴ If the process were not stationary on the timescale T , we would rather have $\int dt_i \mu(t_i)$ for some measure $\mu(t_i)$. This happens on cosmological timescales, in which case the merger distribution has an important dependence on redshift.

⁵ Of course, here we are referring to the intrinsic distribution of the source parameters. The detection probability depends on $\cos \iota$, and in particular the amplitude of the waveform in the inspiral phase is maximized for $\cos^2 \iota = 1$, i.e. for face-on/face-off systems.

⁶ In general, in the literature, the distinction between extrinsic and intrinsic parameters is rather performed stating that the extrinsic parameters are those also have a dependence on the observer, while the intrinsic do not. For instance, the arrival time and the arrival direction depend on the observer’s position on Earth, the polarization angle on the axes in the transverse plane chosen by the observer to define the polarization, and $\cos \iota$ depends on the line of sight to the observer; in contrast, e.g.

Let us denote by $\langle \dots \rangle_{\text{ext}}$ the average over the extrinsic parameters, and by $\langle \dots \rangle_{\text{int}}$ that over the intrinsic ones. These operations define what one means as ‘‘average over different realization of the Universe’’ at the level of the parameters characterizing an individual CBC. On top of it, even the number N_{ev} of binaries coalescing in a given observation time T is a stochastic variable, and the average over different realizations of the ‘‘Universe’’ should include also an average over a Poisson distribution for the number of events, with mean \bar{N}_{ev} . Then, overall,

$$\langle \dots \rangle_U = \langle \langle \dots \rangle_{\text{ext}} \rangle_{\text{int}, N_{\text{ev}}}, \quad (22)$$

where we have denoted by $\langle \dots \rangle_{\text{int}, N_{\text{ev}}}$ the average over the intrinsic parameters as well as on N_{ev} .

III. COMPUTATION OF THE SPECTRAL DENSITY OF THE ASTROPHYSICAL BACKGROUND

We now compute the spectral density for an astrophysical stochastic background, by evaluating explicitly the correlator $\langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_U$. We compute first the average over the extrinsic parameters of all signals,

$$\langle \text{ext} \rangle = \{ \psi_k, t_k, \hat{\mathbf{n}}_k, \cos \iota_k \}_{k=1, \dots, N_{\text{ev}}}. \quad (23)$$

the (source frame) masses or the spins of the component stars are independent of the observer. We find that the distinction based on whether a parameter has a flat integration measure determined by symmetry principles, or a non-trivial integration measure due to astrophysical effects, is more pertinent. Indeed, the computation below will make it clear that the real reason to separate the averages into two groups is that the averages over extrinsic parameters produce Dirac or Kronecker delta's, or other simple structures. Note that, with our definition, also the luminosity distance d_L of the binary is included among the intrinsic parameters (despite the fact that, in principle, depends on the observer's location), because any change in the luminosity distance of the source that produces observable effects corresponds to placing the event at an appreciably different redshift, and the probability of having a coalescence at a given redshift depends on non-trivial astrophysics, and not just on symmetries. Within the same logic, not only the source-frame masses are obviously included among the intrinsic parameters, but also the detector-frame masses, despite the factor $(1+z)$ that connects them to the source-frame masses. Another way to appreciate the difference between time of arrival (which we include among the extrinsic parameters) and luminosity distance d_L (which instead we treat as an intrinsic parameter) is that the former enters in the phase of the waveform. Since phases are defined only modulo 2π , an absolute change Δt_i matters as long as $f\Delta t_i$ is non-negligible, even if Δt_i is of course extremely small compared to the travel time of the signal from the source to the observer. In contrast, d_L enters only in the amplitude, and changes in d_L are appreciable only if $\Delta d_L/d_L$ is non-negligible. For instance, changing the observer's position on Earth by, say, 1000 km, produces a Δt_i whose effect in the phase is significant, but the corresponding change Δd_L only affects the waveform through $\Delta d_L/d_L$, which is utterly negligible, and in this sense d_L does not depend on the observer's position, while t_i does.

Until now, when writing $\tilde{h}_A(f, \hat{\mathbf{n}})$ or $\tilde{h}_{A,i}(f)$, we have left implicit the dependence of the GW signal on the extrinsic parameters, as well as that on the intrinsic parameters. If we write explicitly at least the dependence on the extrinsic parameters, then

$$\tilde{h}_{A,i} = \tilde{h}_{A,i}(f; \psi_i, t_i, \cos \iota_i), \quad (24)$$

while $\hat{\mathbf{n}}_i$ enters through the Dirac delta in Eq. (12); correspondingly, from Eq. (12), $\tilde{h}_A(f, \hat{\mathbf{n}})$ should in principle be written as

$$\tilde{h}_A = \tilde{h}_A(f, \hat{\mathbf{n}}; \{ \hat{\mathbf{n}}_k \}, \{ \psi_k \}, \{ t_k \}, \{ \cos \iota_k \}), \quad (25)$$

where $\{ \hat{\mathbf{n}}_k \}$ denotes the collections of all $\hat{\mathbf{n}}_k$ with $k = 1, \dots, N_{\text{ev}}$, $\{ \psi_k \}$ denotes the collections of all ψ_k with $k = 1, \dots, N_{\text{ev}}$, and so on. However, in order not to burden too much the notation, we will often either still keep implicit the dependence on the extrinsic parameters, or else we will just write in the argument those relevant to the specific steps of the computation that we are performing. In any case, it is useful to keep in mind that the full dependencies are those given in Eqs. (24) and (25).

The averages over the extrinsic parameters are performed while keeping fixed the intrinsic parameters and the number of events N_{ev} . We use the notation $\langle \dots \rangle_{\{ \psi_k \}}$ to denote the average over all ψ_k with $k = 1, \dots, N_{\text{ev}}$, $\langle \dots \rangle_{\{ t_k \}}$ for the average over all t_k , $\langle \dots \rangle_{\{ \hat{\mathbf{n}}_k \}}$ for the average over all $\hat{\mathbf{n}}_k$, and $\langle \dots \rangle_{\{ \cos \iota_k \}}$ for the average over all $\cos \iota_k$. A label such as $\langle \dots \rangle_{\{ \psi_k, t_k, \hat{\mathbf{n}}_k \}}$ means that we average over all the corresponding parameters. We assume that the extrinsic parameters of different events are independent. Then, from Eq. (12),

$$\begin{aligned} \langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_{\{ \psi_k, t_k, \hat{\mathbf{n}}_k, \cos \iota_k \}} &= \sum_{i,j=1}^{N_{\text{ev}}} \\ &\times \langle \tilde{h}_{A,i}^*(f) \tilde{h}_{A',j}(f') \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \delta(\hat{\mathbf{n}}' - \hat{\mathbf{n}}_j) \rangle_{\{ \psi_k, t_k, \hat{\mathbf{n}}_k, \cos \iota_k \}} \\ &= \sum_{i,j=1}^{N_{\text{ev}}} \langle \tilde{h}_{A,i}^*(f) \tilde{h}_{A',j}(f') \rangle_{\{ \psi_k, t_k, \cos \iota_k \}} \\ &\times \langle \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \delta(\hat{\mathbf{n}}' - \hat{\mathbf{n}}_j) \rangle_{\{ \hat{\mathbf{n}}_k \}}, \end{aligned} \quad (26)$$

where in the last equality we used the fact that the directions of propagation $\hat{\mathbf{n}}_k$, with $k = 1 \dots, N_{\text{ev}}$, do not enter into $\tilde{h}_{A,i}(f)$ and $\tilde{h}_{A',j}(f')$, even when $k = i$ or $k = j$. Once again, in Eq. (26) the full dependencies of $\tilde{h}_{A,i}$ and \tilde{h}_A on the extrinsic parameters are given in Eqs. (24) and (25), but we will usually write explicitly only the arguments on which we are averaging; so, e.g., when performing the averages over ψ we write the correlator between $\tilde{h}_{A,i}$ and $\tilde{h}_{A',j}$ as

$$\langle \tilde{h}_{A,i}^*(f; \psi_i) \tilde{h}_{A',j}(f'; \psi_j) \rangle_{\{ \psi_k \}}. \quad (27)$$

A. Average over polarization angles

It is convenient to start from the average over polarization angles. Using Eq. (21),

$$\begin{aligned} & \langle \tilde{h}_{A,i}^*(f; \psi_i) \tilde{h}_{A',j}(f'; \psi_j) \rangle_{\{\psi_k\}} = \\ & = \prod_{k=1}^{N_{\text{ev}}} \int_0^{2\pi} \frac{d\psi_k}{2\pi} \tilde{h}_{A,i}^*(f; \psi_i) \tilde{h}_{A',j}(f'; \psi_j). \end{aligned} \quad (28)$$

Since $\tilde{h}_{A,i}(f)$ only depends on ψ_i and $\tilde{h}_{A',j}(f')$ only depends on ψ_j , in Eq. (28) the integrations over the parameters ψ_k with $k \neq i, j$ play no role as they trivially reduce to unity. Thus, if $i \neq j$,

$$\begin{aligned} & \langle \tilde{h}_{A,i}^*(f; \psi_i) \tilde{h}_{A',j}(f'; \psi_j) \rangle_{\{\psi_k\}} \\ & = \int_0^{2\pi} \frac{d\psi_i}{2\pi} \int_0^{2\pi} \frac{d\psi_j}{2\pi} \tilde{h}_{A,i}^*(f; \psi_i) \tilde{h}_{A',j}(f'; \psi_j), \end{aligned} \quad (29)$$

while, for $i = j$,

$$\begin{aligned} & \langle \tilde{h}_{A,i}^*(f; \psi_i) \tilde{h}_{A',i}(f'; \psi_i) \rangle_{\{\psi_k\}} \\ & = \int_0^{2\pi} \frac{d\psi_i}{2\pi} \tilde{h}_{A,i}^*(f; \psi_i) \tilde{h}_{A',i}(f'; \psi_i), \end{aligned} \quad (30)$$

(no sum over i). To compute the integrals in Eqs. (29) and (30) we need to write explicitly the dependence on the polarization angles. The polarization angle of the i -th event enters through a rotation by an angle $2\psi_i$ which (with a suitable choice of sign conventions) can be written as

$$\tilde{h}_{A,i}(f; \psi_i) = \sum_{B=+,\times} R_{AB}(2\psi_i) \tilde{h}_{B,i}(f; \psi_i = 0), \quad (31)$$

where $R_{AB}(2\psi_i)$ is the rotation matrix with entries

$$R_{AB}(2\psi_i) = \begin{pmatrix} \cos 2\psi_i & -\sin 2\psi_i \\ \sin 2\psi_i & \cos 2\psi_i \end{pmatrix}_{AB}, \quad (32)$$

and $\psi_i = 0$ is a fixed reference angle. It is convenient to decompose the rotation matrix as

$$R_{AB}(2\psi_i) = \delta_{AB} \cos 2\psi_i - i\sigma_{AB}^2 \sin 2\psi_i, \quad (33)$$

where σ^2 is the second Pauli matrix. Then, the integrations in Eqs. (29) and (30) are straightforward, and in particular the integral in Eq. (29), with $i \neq j$, vanishes. We then obtain

$$\begin{aligned} & \langle \tilde{h}_{A,i}^*(f; \psi_i) \tilde{h}_{A',j}(f'; \psi_j) \rangle_{\{\psi_k\}} \\ & = \frac{1}{2} \delta_{ij} \sum_{B,B'=+,\times} (\delta_{AA'} \delta_{BB'} - \sigma_{AA'}^2 \sigma_{BB'}^2) \\ & \quad \times \tilde{h}_{B,i}^*(f; \psi_i = 0) \tilde{h}_{B',i}(f'; \psi_i = 0), \\ & = \frac{1}{2} \delta_{ij} \delta_{AA'} \sum_{B=+,\times} \tilde{h}_{B,i}^*(f; \psi_i = 0) \tilde{h}_{B,i}(f'; \psi_i = 0) \end{aligned}$$

$$\begin{aligned} & -\frac{i}{2} \delta_{ij} \sigma_{AA'}^2 [\tilde{h}_{\times,i}^*(f; \psi_i = 0) \tilde{h}_{+,i}(f'; \psi_i = 0) \\ & \quad - \tilde{h}_{+,i}^*(f; \psi_i = 0) \tilde{h}_{\times,i}(f'; \psi_i = 0)], \end{aligned} \quad (34)$$

where, to get the second equality, we made use of the identity⁷

$$\delta_{AB} \delta_{A'B'} - \sigma_{AB}^2 \sigma_{A'B'}^2 = \delta_{AA'} \delta_{BB'} - \sigma_{AA'}^2 \sigma_{BB'}^2. \quad (40)$$

We observe that the average over the polarizations has produced a factor δ_{ij} , which means that, after this average, different events are uncorrelated. This will also simplify the computation of the subsequent averages. We also notice that it has produced only two structures in the (A, A') indices, i.e. $\delta_{AA'}$ and $\sigma_{AA'}^2$, out of the four possible combinations coming from the fact that the identity matrix and the three Pauli matrices are a complete basis for a 2×2 matrix; i.e. the terms proportional to $\sigma_{AA'}^1$ and $\sigma_{AA'}^3$, and therefore the Stokes parameters U and Q describing linear polarization, see Eq. (10), have been averaged to zero while, at this stage, we still have a non-vanishing circular polarization V .

In terms of $\tilde{h}_A(f, \hat{\mathbf{n}})$ given in Eq. (12), Eq. (34) gives

$$\begin{aligned} & \langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_{\{\psi_k\}} \\ & = \frac{1}{2} \delta_{AA'} \sum_{i=1}^{N_{\text{ev}}} \sum_{B=+,\times} \tilde{h}_{B,i}^*(f; \psi_i = 0) \tilde{h}_{B,i}(f'; \psi_i = 0) \\ & \quad \times \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \delta(\hat{\mathbf{n}}' - \hat{\mathbf{n}}_i) \\ & \quad - \frac{i}{2} \sigma_{AA'}^2 \sum_{i=1}^{N_{\text{ev}}} [\tilde{h}_{\times,i}^*(f; \psi_i = 0) \tilde{h}_{+,i}(f'; \psi_i = 0) \\ & \quad - \tilde{h}_{+,i}^*(f; \psi_i = 0) \tilde{h}_{\times,i}(f'; \psi_i = 0)] \\ & \quad \times \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \delta(\hat{\mathbf{n}}' - \hat{\mathbf{n}}_i), \end{aligned} \quad (41)$$

where the presence of δ_{ij} in Eq. (34) allowed us to reduce the double sum over i, j to a single sum. We further observe that

$$\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \delta(\hat{\mathbf{n}}' - \hat{\mathbf{n}}_i) = \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}') \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i). \quad (42)$$

⁷ This identity can be proven observing that the left-hand side is a 2×2 matrix (with complex matrix elements) in the indices (A, A') , and can therefore be expanded in the basis of $\delta_{AA'}$ and $\sigma_{AA'}^a$, with (complex) coefficients that carry the B, B' indices, i.e. it can be written in the form

$$\delta_{AB} \delta_{A'B'} - \sigma_{AB}^2 \sigma_{A'B'}^2 = \delta_{AA'} q_{BB'} + \sigma_{AA'}^a p_{BB'}^a, \quad (35)$$

for some coefficients $q_{BB'}$ and $p_{BB'}^a$. Contracting both sides with $\delta_{AA'}$ fixes

$$q_{BB'} = \delta_{BB'}, \quad (36)$$

while contracting both side with $\sigma_{AA'}^b$ gives

$$p_{BB'}^a = -\delta^{a2} \sigma_{BB'}^2. \quad (37)$$

Proceeding analogously, one also gets the identities

$$\delta_{AB} \delta_{A'B'} - \sigma_{AB}^1 \sigma_{A'B'}^1 = -\sigma_{AA'}^2 \sigma_{BB'}^2 + \sigma_{AA'}^3 \sigma_{BB'}^3, \quad (38)$$

$$\delta_{AB} \delta_{A'B'} - \sigma_{AB}^3 \sigma_{A'B'}^3 = \sigma_{AA'}^1 \sigma_{BB'}^1 - \sigma_{AA'}^2 \sigma_{BB'}^2. \quad (39)$$

Therefore, we can also extract a term $\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')$ from the sum over i , and we get

$$\begin{aligned}
& \langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_{\{\psi_k\}} \\
&= \frac{1}{2} \delta_{AA'} \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}') \\
&\times \sum_{i=1}^{N_{\text{ev}}} \sum_{B=+, \times} \tilde{h}_{B,i}^*(f; \psi_i = 0) \tilde{h}_{B,i}(f'; \psi_i = 0) \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \\
&\quad - \frac{i}{2} \sigma_{AA'}^2 \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}') \\
&\times \sum_{i=1}^{N_{\text{ev}}} [\tilde{h}_{\times,i}^*(f; \psi_i = 0) \tilde{h}_{+,i}(f'; \psi_i = 0) \\
&\quad - \tilde{h}_{+,i}^*(f; \psi_i = 0) \tilde{h}_{\times,i}(f'; \psi_i = 0)] \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i). \quad (43)
\end{aligned}$$

B. Average over arrival times

We next compute the average over the times of arrival of the GW events. Again, we need to make explicit the dependence on the times of arrival, that until now we kept implicit in $\tilde{h}_{A,i}(f)$. For the i -th event, t_i enters through a phase factor as

$$\tilde{h}_{A,i}(f; t_i) = e^{2\pi i f t_i} \tilde{h}_{A,i}(f, t_i = 0). \quad (44)$$

Since Eq. (34) is already proportional to δ_{ij} , we only need to compute the time average for $i = j$. This is given by

$$\begin{aligned}
& \langle \tilde{h}_{A,i}^*(f; \psi_i, t_i) \tilde{h}_{A',i}(f'; \psi_i, t_i) \rangle_{\{\psi_k, t_k\}} \\
&= \prod_{k=1}^{N_{\text{ev}}} \int_{-T/2}^{T/2} \frac{dt_k}{T} \langle \tilde{h}_{A,i}^*(f; \psi_i, t_i) \tilde{h}_{A',i}(f'; \psi_i, t_i) \rangle_{\{\psi_k\}} \\
&= \int_{-T/2}^{T/2} \frac{dt_i}{T} \langle e^{-2\pi i f t_i} \tilde{h}_{A,i}^*(f; \psi_i, t_i = 0) \\
&\quad \times e^{2\pi i f' t_i} \tilde{h}_{A',i}(f'; \psi_i, t_i = 0) \rangle_{\{\psi_k\}} \\
&= \frac{1}{T} \delta_T(f - f') \\
&\quad \times \langle \tilde{h}_{A,i}^*(f; \psi_i, t_i = 0) \tilde{h}_{A',i}(f'; \psi_i, t_i = 0) \rangle_{\{\psi_k\}}, \quad (45)
\end{aligned}$$

where $\delta_T(f - f')$ is given by Eq. (6). Inserting here $\langle \tilde{h}_{A,i}^*(f; \psi_i, t_i = 0) \tilde{h}_{A',i}(f'; \psi_i, t_i = 0) \rangle_{\{\psi_k\}}$ from Eq. (34), we get

$$\begin{aligned}
& \langle \tilde{h}_{A,i}^*(f; \psi_i, t_i) \tilde{h}_{A',j}(f', \psi_j, t_j) \rangle_{\{\psi_k, t_k\}} \\
&= \frac{1}{2} \delta_{ij} \delta_T(f - f') [\mathcal{I}_i(f) \delta_{AA'} + \mathcal{V}_i(f) \sigma_{AA'}^2], \quad (46)
\end{aligned}$$

where

$$\mathcal{I}_i(f) = \frac{|\tilde{h}_{+,i}(f)|^2 + |\tilde{h}_{\times,i}(f)|^2}{T}, \quad (47)$$

$$\mathcal{V}_i(f) = \frac{2 \text{Im} [\tilde{h}_{+,i}(f) \tilde{h}_{\times,i}^*(f)]}{T}. \quad (48)$$

Observe that the factors $1/T$ in these expressions have been inherited from Eq. (15). Note also that, if we set

$f' = f$ in Eq. (46) and use Eq. (7), they cancel with the factor $\delta_T(0) = T$.

Introducing the right-handed (R) and left-handed (L) GW fields

$$\tilde{h}_{R,i}(f; \psi_i, t_i) = \frac{\tilde{h}_{+,i}(f; \psi_i, t_i) + i \tilde{h}_{\times,i}(f; \psi_i, t_i)}{\sqrt{2}}, \quad (49)$$

$$\tilde{h}_{L,i}(f; \psi_i, t_i) = \frac{\tilde{h}_{+,i}(f; \psi_i, t_i) - i \tilde{h}_{\times,i}(f; \psi_i, t_i)}{\sqrt{2}}, \quad (50)$$

we see that Eqs. (47) and (48) can also be rewritten as

$$\mathcal{I}_i(f) = \frac{|\tilde{h}_{R,i}(f)|^2 + |\tilde{h}_{L,i}(f)|^2}{T}, \quad (51)$$

$$\mathcal{V}_i(f) = \frac{|\tilde{h}_{R,i}(f)|^2 - |\tilde{h}_{L,i}(f)|^2}{T}. \quad (52)$$

Note that $\mathcal{I}_i(f)$ and $\mathcal{V}_i(f)$ are invariant under rotations of the form (31), i.e.

$$\tilde{h}_{A,i}(f) \rightarrow \sum_{B=+, \times} R_{AB}(2\psi_i) \tilde{h}_{B,i}(f). \quad (53)$$

In terms of $\tilde{h}_{L,i}(f)$ and $\tilde{h}_{R,i}(f)$, Eq. (53) reads

$$\tilde{h}_{L,i}(f) \rightarrow e^{-2i\psi_i} \tilde{h}_{L,i}(f), \quad \tilde{h}_{R,i}(f) \rightarrow e^{+2i\psi_i} \tilde{h}_{R,i}(f), \quad (54)$$

which expresses the fact that $\tilde{h}_{L,i}(f)$ and $\tilde{h}_{R,i}(f)$ are eigenstates of helicity with eigenvalues -2 and $+2$, respectively.⁸ Therefore $\mathcal{I}_i(f)$ and $\mathcal{V}_i(f)$ lose any dependence on the reference values $\psi_i = 0$ introduced in Eq. (31). Similarly, $\mathcal{I}_i(f)$ and $\mathcal{V}_i(f)$ are also invariant under

$$\tilde{h}_{A,i}(f) \rightarrow e^{2\pi i f t_i} \tilde{h}_{A,i}(f), \quad (55)$$

and therefore $\mathcal{I}_i(f)$ and $\mathcal{V}_i(f)$ lose any dependence on the reference values $t_i = 0$ introduced in Eq. (44). For this reason, in Eqs. (47) and (48) we have omitted the arguments $\psi_i = 0$ and $t_i = 0$ from $h_A(f; \psi_i = 0, t_i = 0)$, and similarly in Eqs. (51) and (52).

We see that the average over polarizations and over arrival times have produced a Kronecker delta δ_{ij} that decorrelates different events, and a (finite-time) Dirac delta $\delta_T(f - f')$, that decorrelates different frequencies (in the limit $fT \gg 1$). These two averages did not involve any real physical assumption (such as the isotropy of the source distribution), since we just assumed that the result does not depend on the system of axes set by the observer to measure ψ , and that the astrophysical population does

⁸ Our sign convention on the definition of helicity is the same as in Eq. (2.197) of Ref. [2], that states that an eigenstate of helicity with eigenvalue h (not to be confused with a GW amplitude! both symbols are standard) transforms as $e^{ih\psi}$ under a rotation by an angle ψ . If one uses the opposite convention, so that an eigenstate of helicity with eigenvalue h transforms as $e^{-ih\psi}$, one must exchange the definitions of h_R and h_L .

not evolve appreciably over the observation time; the corresponding result is therefore very general. Plugging Eq. (46) into Eq. (26) we see that, at this stage, we can write

$$\begin{aligned} & \langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_{\{\psi_k, t_k, \hat{\mathbf{n}}_k, \cos \iota_k\}} \\ &= \frac{1}{2} \delta_T(f - f') \sum_{i=1}^{N_{\text{ev}}} \left[\langle \mathcal{I}_i(f) \delta_{AA'} + \mathcal{V}_i(f) \sigma_{AA'}^2 \rangle_{\cos \iota_i} \right. \\ & \quad \left. \times \langle \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \delta(\hat{\mathbf{n}}' - \hat{\mathbf{n}}_i) \rangle_{\hat{\mathbf{n}}_i} \right], \end{aligned} \quad (56)$$

where the presence of δ_{ij} in Eq. (46) allowed us to reduce Eq. (26) to a sum over a single index i ; correspondingly, for the i -th term inside the sum over i , the average over $\{\cos \iota_k\}_{k=1, \dots, N_{\text{ev}}}$ reduces to an average over $\cos \iota_i$ and, similarly, the average over the propagation directions reduces to an average over $\hat{\mathbf{n}}_i$. We can now further use

$$\langle \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \delta(\hat{\mathbf{n}}' - \hat{\mathbf{n}}_i) \rangle_{\hat{\mathbf{n}}_i} = \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}') \langle \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \rangle_{\hat{\mathbf{n}}_i}. \quad (57)$$

Then,

$$\begin{aligned} & \langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_{\{\psi_k, t_k, \hat{\mathbf{n}}_k, \cos \iota_k\}} \\ &= \frac{1}{2} \delta_T(f - f') \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}') \\ & \quad \times \sum_{i=1}^{N_{\text{ev}}} \left[\langle \mathcal{I}_i(f) \delta_{AA'} + \mathcal{V}_i(f) \sigma_{AA'}^2 \rangle_{\cos \iota_i} \langle \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \rangle_{\hat{\mathbf{n}}_i} \right]. \end{aligned} \quad (58)$$

We observe that, after averaging over polarization angles and times of arrival, the $\langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle$ correlator is proportional to both $\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')$ and $\delta_T(f - f')$, and linear polarizations have been averaged to zero.

C. Average over arrival directions

We next perform the average over the arrival directions $-\hat{\mathbf{n}}_i$ or, equivalently, over the propagation directions $\hat{\mathbf{n}}_i$. In a generic, non-isotropic case, using Eq. (18) we get

$$\begin{aligned} \langle \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \rangle_{\hat{\mathbf{n}}_i} &= \int_{S^2} \frac{d^2 \hat{\mathbf{n}}_i}{4\pi} \mu(\hat{\mathbf{n}}_i) \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \\ &= \frac{\mu(\hat{\mathbf{n}})}{4\pi}. \end{aligned} \quad (59)$$

Inserting this into Eq. (58), and including also the averages over intrinsic parameters and over N_{ev} as in Eq. (22), we then find

$$\begin{aligned} & \langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_U \\ &= \delta_T(f - f') \frac{\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')}{4\pi} \frac{1}{2} [S_h^{\text{astro}}(f; \hat{\mathbf{n}})]_{AA'}, \end{aligned} \quad (60)$$

where

$$[S_h^{\text{astro}}(f; \hat{\mathbf{n}})]_{AA'} = I(f; \hat{\mathbf{n}}) \delta_{AA'} + V(f; \hat{\mathbf{n}}) \sigma_{AA'}^2. \quad (61)$$

The functions $I(f; \hat{\mathbf{n}})$ and $V(f; \hat{\mathbf{n}})$ factorize into a function of f and a function of $\hat{\mathbf{n}}$,

$$I(f; \hat{\mathbf{n}}) = \mu(\hat{\mathbf{n}}) I(f), \quad (62)$$

$$V(f; \hat{\mathbf{n}}) = \mu(\hat{\mathbf{n}}) V(f), \quad (63)$$

where

$$\begin{aligned} I(f) &= \left\langle \sum_{i=1}^{N_{\text{ev}}} \langle \mathcal{I}_i(f) \rangle_{\cos \iota_i} \right\rangle_{\text{int}, N_{\text{ev}}} \\ &= \frac{1}{T} \left\langle \left[\sum_{i=1}^{N_{\text{ev}}} \langle (|\tilde{h}_{+,i}(f)|^2 + |\tilde{h}_{\times,i}(f)|^2) \rangle_{\cos \iota_i} \right] \right\rangle_{\text{int}, N_{\text{ev}}} \\ &= \frac{1}{T} \left\langle \left[\sum_{i=1}^{N_{\text{ev}}} \langle (|\tilde{h}_{R,i}(f)|^2 + |\tilde{h}_{L,i}(f)|^2) \rangle_{\cos \iota_i} \right] \right\rangle_{\text{int}, N_{\text{ev}}}, \end{aligned} \quad (64)$$

and

$$\begin{aligned} V(f) &= \left\langle \sum_{i=1}^{N_{\text{ev}}} \langle \mathcal{V}_i(f) \rangle_{\cos \iota_i} \right\rangle_{\text{int}, N_{\text{ev}}} \\ &= \frac{2}{T} \left\langle \left[\sum_{i=1}^{N_{\text{ev}}} \langle \text{Im}(\tilde{h}_{+,i}(f) \tilde{h}_{\times,i}^*(f)) \rangle_{\cos \iota_i} \right] \right\rangle_{\text{int}, N_{\text{ev}}} \\ &= \frac{1}{T} \left\langle \left[\sum_{i=1}^{N_{\text{ev}}} \langle (|\tilde{h}_{R,i}(f)|^2 - |\tilde{h}_{L,i}(f)|^2) \rangle_{\cos \iota_i} \right] \right\rangle_{\text{int}, N_{\text{ev}}}. \end{aligned} \quad (65)$$

Therefore, also $[S_h^{\text{astro}}(f; \hat{\mathbf{n}})]_{AA'}$ factorizes into a function of f times a function of $\hat{\mathbf{n}}$,

$$[S_h^{\text{astro}}(f; \hat{\mathbf{n}})]_{AA'} = \mu(\hat{\mathbf{n}}) [S_h^{\text{astro}}(f)]_{AA'}, \quad (66)$$

where

$$[S_h^{\text{astro}}(f)]_{AA'} = I(f) \delta_{AA'} + V(f) \sigma_{AA'}^2. \quad (67)$$

The isotropic case is recovered setting $\mu(\hat{\mathbf{n}}) = 1$.

Comparing Eqs. (60) and (61) with Eqs. (9) and (10) we see that $I(f, \hat{\mathbf{n}})$ is the Stokes parameter associated with intensity and $V(f, \hat{\mathbf{n}})$ is the Stokes parameter associated with circular polarization, while the terms associated with linear polarizations are absent. We also see that $I(f, \hat{\mathbf{n}})$ and $V(f, \hat{\mathbf{n}})$ factorize as in Eqs. (62) and (63).

D. Average over orbit inclinations

Let us now tackle the explicit computation of the average over $\cos \iota$. Similarly to the polarization angle and the time of arrival, the integration over $\cos \iota$ is also naturally carried out with a flat measure, Eq. (17), without the need of any non-trivial physical assumption: the binary knows nothing about the observer, and has no reason to be oriented preferentially in some direction with respect to the line of sight connecting it to the observer. However, the difference with the previous cases is that now the dependence on $\cos \iota$, for a full waveform including the effect of higher modes, is in general non-trivial.

In Ref. [42] the average over the inclination has been computed keeping only the quadrupolar mode, neglecting all higher modes in the amplitude. In that case, the dependence on $\cos \iota$ becomes very simple, and is given by (see Eqs. (4.34) and (4.35) of Ref. [2])

$$\tilde{h}_+(f; \cos \iota) = \mathcal{A}(f)e^{i\Psi(f)} \frac{1 + \cos^2 \iota}{2}, \quad (68)$$

$$\tilde{h}_\times(f; \cos \iota) = \mathcal{A}(f)ie^{i\Psi(f)} \cos \iota, \quad (69)$$

where $\mathcal{A}(f)$ and $\Psi(f)$ are an amplitude and a phase, whose explicit expression we do not need here. Since in Eqs. (64) and (65) the averages of $\mathcal{I}_i(f)$ and $\mathcal{V}_i(f)$ are just with respect to $\cos \iota_i$, with the same index i , in Eqs. (68) and (69), and in the following steps, for notational simplicity we just suppress the event index i . Note that the dependence on $\cos \iota$ in Eqs. (64) and (65) is a characteristic of the mode considered, independently of the post-Newtonian order at which the phase of the waveform is computed, and is valid even for a full inspiral-merger-ringdown waveform, again as far as only the lowest mode of the amplitude (i.e., the quadrupole) is retained. Then

$$|\tilde{h}_+(f)|^2 + |\tilde{h}_\times(f)|^2 = \mathcal{A}^2(f)Q_I(\cos \iota), \quad (70)$$

$$\text{Im} [\tilde{h}_+(f)\tilde{h}_\times^*(f)] = -\mathcal{A}^2(f)Q_V(\cos \iota), \quad (71)$$

where

$$Q_I(\cos \iota) = \left(\frac{1 + \cos^2 \iota}{2} \right)^2 + \cos^2 \iota, \quad (72)$$

$$Q_V(\cos \iota) = \frac{1 + \cos^2 \iota}{2} \cos \iota. \quad (73)$$

Averaging over $\cos \iota$ with the flat measure (17),

$$\langle Q_I(\cos \iota) \rangle_{\cos \iota} = \frac{4}{5}, \quad (74)$$

$$\langle Q_V(\cos \iota) \rangle_{\cos \iota} = 0. \quad (75)$$

Therefore, in the approximation in which one considers only the quadrupole mode in the waveform and neglects the higher modes, the circular polarization averages to zero, as already found in Ref. [42].

For the term related to intensity, using the fact that $Q_I(\cos \iota = 1) = 2$, so that $\langle Q_I(\cos \iota) \rangle_{\cos \iota} = (2/5)Q_I(\cos \iota = 1)$, we could write (reinstating explicitly the dependence on $\cos \iota_i$ in $\langle \mathcal{I}_i(f) \rangle_{\cos \iota_i}$),

$$\langle \mathcal{I}_i(f; \cos \iota_i) \rangle_{\cos \iota_i} = \frac{2}{5} \mathcal{I}_i(f; \cos \iota_i = 1). \quad (76)$$

The question of whether the circular polarization vanishes exactly or not is, however, quite important, because parity-violating mechanism in the early Universe could produce a cosmological stochastic background with a net circular polarization [61], and various techniques have been discussed to extract this effect from GW data [4, 44, 62–67]; such early-Universe signature might however be masked by

contamination from the astrophysical background, if even a small fraction of it should turn out to be polarized. It is therefore interesting to see if the vanishing of the circular polarization after averaging over $\cos \iota$ also holds when we include higher modes in the waveform. At first sight one might doubt it, given the complicated dependence on $\sin \iota$ and $\cos \iota$ of the higher modes. However, the vanishing of the circular polarization upon averaging over $\cos \iota$ is in fact a general feature, which simply follows from parity arguments. Let us denote by $\hat{\mathbf{n}}_{\text{orbit}}$ the normal to the orbit. In polar coordinates, choosing as polar axis the line-of-sight to the observer, $\hat{\mathbf{n}}_{\text{orbit}} = (\sin \iota \cos \varphi, \sin \iota \sin \varphi, \cos \iota)$, where φ is an angle in the plane transverse to the polar axis. If we perform a parity transformation, the normal to the orbit changes sign, $\hat{\mathbf{n}}_{\text{orbit}} \rightarrow -\hat{\mathbf{n}}_{\text{orbit}}$ which, recalling that $\iota \in [0, \pi]$, means $\{\iota, \varphi\} \rightarrow \{\pi - \iota, \varphi + \pi\}$, so in particular

$$\cos \iota \rightarrow -\cos \iota, \quad \sin \iota \rightarrow \sin \iota. \quad (77)$$

At the same time, under a parity transformation, h_+ transform as a scalar, $h_+ \rightarrow h_+$, while h_\times as a pseudoscalar, $h_\times \rightarrow -h_\times$.⁹ Therefore, the GWs produced by a CBC whose orbit has a normal $\hat{\mathbf{n}}_{\text{orbit}}$ are obtained from those produced by a CBC whose orbit has a normal $-\hat{\mathbf{n}}_{\text{orbit}}$, simply replacing $h_+ \rightarrow h_+$ and $h_\times \rightarrow -h_\times$, or, from Eqs. (49) and (50), exchanging h_L with h_R . Since this follows from a general symmetry argument, it holds for the full waveform, including higher modes.¹⁰ We see from Eqs. (51) and (52) that $\mathcal{I}_i(f)$ is even under $h_L \leftrightarrow h_R$, while $\mathcal{V}_i(f)$ is odd. As a result, for $I(f; \hat{\mathbf{n}})$ the integral over $\cos \iota$ between -1 and 1 is twice the integral from 0 to

⁹ This follows from the fact that the plus and cross polarization of a GW with propagation direction $\hat{\mathbf{n}}$ are defined introducing two unit vectors $\hat{\mathbf{u}}, \hat{\mathbf{v}}$ orthogonal to $\hat{\mathbf{n}}$ and among them, chosen so that $\{\hat{\mathbf{u}}, \hat{\mathbf{v}}, \hat{\mathbf{n}}\}$ form a right-handed oriented frame, i.e. $\hat{\mathbf{u}} \times \hat{\mathbf{v}} = \hat{\mathbf{n}}$; e.g., when $\hat{\mathbf{n}} = \hat{\mathbf{z}}$, one could choose $\hat{\mathbf{u}} = \hat{\mathbf{x}}$ and $\hat{\mathbf{v}} = \hat{\mathbf{y}}$. Then the polarization tensors associated with the plus and cross polarizations are, respectively,

$$e_{ij}^+(\hat{\mathbf{n}}) = \hat{\mathbf{u}}_i \hat{\mathbf{u}}_j - \hat{\mathbf{v}}_i \hat{\mathbf{v}}_j, \quad e_{ij}^\times(\hat{\mathbf{n}}) = \hat{\mathbf{u}}_i \hat{\mathbf{v}}_j + \hat{\mathbf{v}}_i \hat{\mathbf{u}}_j, \quad (78)$$

(see Eq. (1.54) of Ref. [2]). Under a parity transformation we have $\hat{\mathbf{n}} \rightarrow \hat{\mathbf{n}}' = -\hat{\mathbf{n}}$, $\hat{\mathbf{u}} \rightarrow \hat{\mathbf{u}}' = -\hat{\mathbf{u}}$ and $\hat{\mathbf{v}} \rightarrow \hat{\mathbf{v}}' = -\hat{\mathbf{v}}$. However, $\{\hat{\mathbf{u}}', \hat{\mathbf{v}}', \hat{\mathbf{n}}'\}$ no longer forms a right-handed frame, since $\hat{\mathbf{u}}' \times \hat{\mathbf{v}}' = -\hat{\mathbf{n}}'$. To recover a right-handed frame we must exchange the sign of either $\hat{\mathbf{u}}'$ or $\hat{\mathbf{v}}'$. As a result,

$$e_{ij}^+(-\hat{\mathbf{n}}) = e_{ij}^+(\hat{\mathbf{n}}), \quad e_{ij}^\times(-\hat{\mathbf{n}}) = -e_{ij}^\times(\hat{\mathbf{n}}). \quad (79)$$

¹⁰ This can be checked explicitly for the first five higher-mode amplitudes $H_{+,\times}^{(a)}$ (with $a = 1/2, 1, 3/2, 2, 5/2$), i.e., the amplitudes up to 2.5PN order, given explicitly in Refs. [68, 69] where we see that, indeed, under the transformation (77), the amplitudes $H_+^{(a)}$ are invariant while $H_\times^{(a)}$ change sign. It can also be verified by explicit numerical calculation (which we performed up to $l = 6$) that the $\cos \iota$ average of the combination $\text{Im}[\tilde{h}_+ \tilde{h}_\times^*]$ entering Eq. (65) vanishes exactly for all higher modes due to the structure of the spin-weighted spherical harmonics commonly used to decompose the signal.

1, while the integral of $V(f; \hat{\mathbf{n}})$ vanishes. Note that this holds already at the level of Eqs. (62) and (63), before making any assumption on the distribution of the arrival directions.

Therefore, after averaging over the inclination angle as in Eq. (17), the circular polarization vanishes, even for the exact waveform including higher modes. In contrast, the exact number in Eq. (74) is not protected by symmetries, and will be affected by higher modes. Then, for this term it can be better to leave the average over $\cos \iota$, rather than using Eq. (76). Depending on the accuracy required, one could perform the average numerically with the desired waveform including higher modes, as indeed we will do below.

To sum up, after averaging over $\cos \iota$ (and for a generic distribution $\mu(\hat{\mathbf{n}})$ of arrival directions), Eqs. (60)–(63) become

$$\begin{aligned} & \langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_U \\ &= \delta_T(f - f') \frac{\mu(\hat{\mathbf{n}}) \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')}{4\pi} \delta_{AA'} \frac{1}{2} S_h^{\text{astro}}(f), \end{aligned} \quad (80)$$

where $S_h^{\text{astro}}(f) \equiv I(f)$, and $I(f)$ is given by Eq. (64) so, writing explicitly also the dependence on $\cos \iota_i$ in $\tilde{h}_{A,i}(f; \cos \iota_i)$,

$$S_h^{\text{astro}}(f) = \frac{1}{T} \sum_{i=1}^{N_{\text{ev}}} \sum_{A=+, \times} \langle |\tilde{h}_{A,i}(f; \cos \iota_i)|^2 \rangle_{\cos \iota_i} \Big|_{\text{int}, N_{\text{ev}}}. \quad (81)$$

If we further assumes isotropy, then $\mu(\hat{\mathbf{n}}) = 1$.

E. Summary

To summarize this section, we have computed the spectral density due to an astrophysical background by performing explicitly the average of the intrinsic parameters. Going step-by-step, one can appreciate the effects of the different averages. The averages over the polarization angles decorrelate the different events, producing the factor δ_{ij} in Eq. (34) or, equivalently, decorrelate the arrival directions, producing a Dirac delta $\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')$ in Eq. (43). They also eliminate the linear polarizations, since in Eq. (34) the terms proportional to $\sigma_{AA'}^1$ and $\sigma_{AA'}^3$ have been averaged to zero. The averages over arrival times produce a (regularized) Dirac delta $\delta_T(f - f')$.¹¹

¹¹ One could have also started from the integration over the times of arrival. In that case, for $i \neq j$, using Eq. (44) the correlator $\langle \tilde{h}_{A,i}^*(f; t_i) \tilde{h}_{A',j}(f', t_j) \rangle_{\{t_k\}}$ becomes proportional to

$$\int_{-T/2}^{T/2} dt_i \int_{-T/2}^{T/2} dt_j e^{-2\pi i f t_i + 2\pi i f' t_j},$$

which does not vanish but is rather proportional to $\delta_T(f) \delta_T(f')$. This could however be set to zero considering that, for GWs, we are not interested in the static part of the field. In any case, performing first the average over the polarization angles produces already a factor δ_{ij} , so that we do not need this extra step.

The averages over the inclination angles remove the remaining Stokes parameter, associated with circular polarization. The result for $\langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle$ then becomes proportional also to $\delta_{AA'}$. The average over arrival directions can be performed, at no extra cost, without assuming isotropy and using a generic function $\mu(\hat{\mathbf{n}})$ in the average, Eq. (19), so eventually the two-point correlator takes the form given in Eq. (80), with a spectral density $S_h^{\text{astro}}(f; \hat{\mathbf{n}})$ given explicitly in Eq. (81).

We finally need to perform the remaining averages over $\cos \iota_i$ of the intensity Stokes parameter, as well as the averages over the intrinsic parameters of the binary, and on the number of events that arrive during the observation time T . When we have a large ensemble of CBC, whether coming from actual detections or generated synthetically according to a given distribution of parameters, the averages over these parameters will be automatically performed (modulo a dependence on the realization, i.e. shot noise effects, that we will discuss in Sec. V) by the sum over the events. For instance, when generating a CBC population, one will draw $\cos \iota_i$ from a uniform distribution and this, in the limit of an infinite number of events, will automatically produce an average over $\cos \iota_i$ (times the total number of events) with this flat distribution; similarly for the intrinsic parameters, e.g. a BBH population will be generated drawing the masses from the desired mass distribution. In this sense, we can remove the explicit reference to the average over $\cos \iota$ and over the intrinsic parameters in Eq. (81). Of course, this could have been done for all parameters, including times of arrival, polarization angles and arrival directions. However, in those cases we have been able to perform the averages analytically, and this allowed us to extract Dirac or Kronecker delta functions, that simplify considerably the structure of the result. Note that, for the average over $\cos \iota$, we are taking a mixed approach, where we use the information that the average sets to zero the circular polarization, while we compute the average of the intensity parameter by summing over the events; this is because, as we have seen, the vanishing of the circular polarization is an exact result related to a parity symmetry (again, modulo shot noise effects that we will discuss in Sec. V) while a numerical coefficient such as that in Eq. (76) is not protected by any symmetry argument, and will be affected by the inclusion of higher modes in the waveform.

With this understanding, for a generic distribution of arrival directions, we can rewrite Eq. (81) as

$$S_h^{\text{astro}}(f) = \frac{1}{T} \left\langle \sum_{i=1}^{N_{\text{ev}}} |\tilde{h}_{+,i}(f)|^2 + |\tilde{h}_{\times,i}(f)|^2 \right\rangle_{N_{\text{ev}}}, \quad (82)$$

keeping only the average over the Poisson distribution of the number of events detected in the observation time T . As a further simplification, we can eliminate even the latter average, and just replace N_{ev} with its average value

\bar{N}_{ev} , so in this approximation we can simply write

$$S_h^{\text{astro}}(f) = \frac{1}{T} \sum_{i=1}^{\bar{N}_{\text{ev}}} \left[|\tilde{h}_{+,i}(f)|^2 + |\tilde{h}_{\times,i}(f)|^2 \right], \quad (83)$$

which provides a simple expression for the spectral density generated by an astrophysical background in an observation time T .¹²

IV. ENERGY DENSITY OF ANISOTROPIC AND POLARIZED ASTROPHYSICAL BACKGROUNDS

Let us now recall that the energy associated with a single GW signal is given by (see Eq. (1.135) of Ref. [2])

$$\rho_{\text{gw}} = \frac{c^2}{32\pi G} \langle \dot{h}_{kl} \dot{h}_{kl} \rangle_t, \quad (84)$$

where the sum over the repeated indices k, l is understood, and $\langle \dots \rangle_t$ denotes a temporal average over a time interval much longer than the period of the GW. For a stochastic background, observed for a time T , we should further perform an average over Universe realizations. However, the latter also implicitly contains the time average $\langle \dots \rangle_t$ for each individual signal, through the average over arrival times so, for a stochastic background,

$$\rho_{\text{gw}} = \frac{c^2}{32\pi G} \langle \dot{h}_{kl} \dot{h}_{kl} \rangle_U. \quad (85)$$

For a stochastic background stationary, isotropic and unpolarized, whose two-point correlator is given by Eq. (3), a standard computation (see, e.g., Eq. (7.201) of Ref. [2]) then shows that

$$\frac{d\rho_{\text{gw}}}{d \log f} = \frac{\pi c^2}{2G} f^3 S_h(f). \quad (86)$$

¹² Despite the fact that Eq. (83), or in fact rather the equivalent expression (89) below for $\Omega_{\text{gw}}(f)$, has been often used in the literature, to our surprise we have not been able to find any real derivation of it; typically, either the result is just stated, or is attributed to references that, in reality, do not contain any derivation but just restate it, or a chain of citations eventually leads to [Phinney's](#) paper [70]; the latter is the closest reference for this result but, in fact, even this reference does not contain (and does not claim) a derivation of Eqs. (83) and (89); actually, Eq. (22) of Ref. [70] gives the energy flux, where indeed enters the quantity $f^3 \sum_i [|\tilde{h}_{+,i}(f)|^2 + |\tilde{h}_{\times,i}(f)|^2]$, which also appears at the numerator of Eq. (89), but it is not so obvious how to proceed from there to get the factor $1/T$ in Eqs. (83) and (89); sometimes this has been vaguely justified as a ‘‘normalization factor’’. The expressions (83) or (89) only appear in subsequent papers, with the factor $1/T$ added by hand, presumably on dimensional grounds, but without any real derivation. Our steps in this section provide such a derivation. As one can appreciate from these steps, this derivation involved several non-trivial conceptual and technical points, that we attempted to make as explicit as possible.

It is convenient to define the dimensionless quantity

$$\Omega_{\text{gw}}(f) \equiv \frac{1}{\rho_c} \frac{d\rho_{\text{gw}}}{d \log f}, \quad (87)$$

where $\rho_c = 3c^2 H_0^2 / (8\pi G)$ is the critical energy density for closing the Universe. In terms of $S_h(f)$, we therefore have

$$\Omega_{\text{gw}}(f) = \frac{4\pi^2}{3H_0^2} f^3 S_h(f). \quad (88)$$

When the background is due to the superposition of a discrete number of astrophysical events, the spectral density is given by Eq. (82) which, for large N_{ev} , can also be approximated by Eq. (83). Using for simplicity the latter expression, the energy density per logarithmic interval of frequency, normalized again to ρ_c , is then given by

$$\Omega_{\text{gw}}^{\text{astro}}(f) = \frac{4\pi^2}{3H_0^2} \frac{f^3}{T} \sum_{i=1}^{\bar{N}_{\text{ev}}} \left[|\tilde{h}_{+,i}(f)|^2 + |\tilde{h}_{\times,i}(f)|^2 \right]. \quad (89)$$

Observe that, in the limit of large observation time, this sum converges to a finite quantity. This can be better seen rewriting Eq. (89) as

$$\begin{aligned} \Omega_{\text{gw}}^{\text{astro}}(f) &= \frac{4\pi^2}{3H_0^2} f^3 \frac{\bar{N}_{\text{ev}}}{T} \\ &\times \frac{1}{\bar{N}_{\text{ev}}} \sum_{i=1}^{\bar{N}_{\text{ev}}} \left[|\tilde{h}_{+,i}(f)|^2 + |\tilde{h}_{\times,i}(f)|^2 \right]. \end{aligned} \quad (90)$$

In the large T limit, \bar{N}_{ev}/T converges to the merger rate, while the term in the second line is the average of $|\tilde{h}_{+,i}(f)|^2 + |\tilde{h}_{\times,i}(f)|^2$ over the population of events.

It is interesting to generalize this computation to a generic anisotropic and polarized background. This can be relevant both to cosmological backgrounds and to the astrophysical background, where anisotropies, at some level, will be present; furthermore, even if in this section we have shown that the average over extrinsic parameter sets to zero the polarization of the CBC background, we will see in Sec. V that some amount of polarization can be generated by shot noise.

The computation can be performed writing the correlator as in Eq. (9), with a generic function $H_{AA'}(f, \hat{\mathbf{n}})$ (without even assuming a factorized form). Inserting Eq. (1) into Eq. (85) we get

$$\begin{aligned} \rho_{\text{gw}} &= \frac{c^2}{32\pi G} \int_{-\infty}^{\infty} df df' \int_{S^2} d^2 \hat{\mathbf{n}} d^2 \hat{\mathbf{n}}' \sum_{A, A' = +, \times} \\ &\times (2\pi i f) (-2\pi i f') \langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_U \\ &\times e_{kl}^A(\hat{\mathbf{n}}) e_{kl}^{A'}(\hat{\mathbf{n}}') e^{2\pi i(f-f')t} e^{-2\pi i(f\hat{\mathbf{n}} - f'\hat{\mathbf{n}}') \cdot \mathbf{x}/c} \\ &= \frac{\pi c^2}{8G} \int_0^{\infty} df f^2 \int_{S^2} \frac{d^2 \hat{\mathbf{n}}}{4\pi} \\ &\times \sum_{A, A' = +, \times} H_{AA'}(f, \hat{\mathbf{n}}) e_{kl}^A(\hat{\mathbf{n}}) e_{kl}^{A'}(\hat{\mathbf{n}}), \end{aligned} \quad (91)$$

where we have written the result as an integral from $f = 0$ to $f = \infty$ using $\tilde{h}_A^*(f, \hat{\mathbf{n}}) = \tilde{h}_A(-f, \hat{\mathbf{n}})$ [which follows from the reality of $h_{kl}(t)$ and of the polarization tensors e_{kl}^+ , e_{kl}^\times , see Eq. (78)].

The crucial point now is that, thanks to the Dirac delta $\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')$ in Eq. (9), we recovered the structure $e_{kl}^A(\hat{\mathbf{n}})e_{kl}^{A'}(\hat{\mathbf{n}})$, in which the arguments in the two polarization tensors are the same. We can then use Eq. (2) (recall that the sum over the repeated spatial indices k, l is understood), and then

$$\begin{aligned} H_{AA'}(f, \hat{\mathbf{n}})e_{kl}^A(\hat{\mathbf{n}})e_{kl}^{A'}(\hat{\mathbf{n}}) &= 2\delta_{AA'}H_{AA'}(f, \hat{\mathbf{n}}) \\ &= 4I(f, \hat{\mathbf{n}}), \end{aligned} \quad (92)$$

where, in the last line, we wrote $H_{AA'}(f, \hat{\mathbf{n}})$ as in Eq. (10), and used the fact that the Pauli matrices are traceless.

Therefore the Stokes parameters associated with linear and with circular polarization do not contribute to the energy density of a stationary stochastic background (independently of whether it is isotropic or anisotropic), and

$$\rho_{\text{gw}} = \frac{\pi c^2}{2G} \int_0^\infty d \log f f^3 \int_{S^2} \frac{d^2 \hat{\mathbf{n}}}{4\pi} I(f, \hat{\mathbf{n}}), \quad (93)$$

so that

$$\frac{d\rho_{\text{gw}}}{d \log f d^2 \hat{\mathbf{n}}} = \frac{\pi c^2}{2G} \frac{f^3 I(f, \hat{\mathbf{n}})}{4\pi}. \quad (94)$$

We can also define

$$S_h(f) \equiv I(f) \equiv \int_{S^2} \frac{d^2 \hat{\mathbf{n}}}{4\pi} I(f, \hat{\mathbf{n}}), \quad (95)$$

so that we recover the standard expressions

$$\rho_{\text{gw}} = \frac{\pi c^2}{2G} \int_0^\infty d \log f f^3 S_h(f), \quad (96)$$

and

$$\begin{aligned} \frac{d\rho_{\text{gw}}}{d \log f} &= \frac{\pi c^2}{2G} f^3 S_h(f) \\ &= \frac{\pi c^2}{2G} f^3 \int_{S^2} \frac{d^2 \hat{\mathbf{n}}}{4\pi} I(f, \hat{\mathbf{n}}), \end{aligned} \quad (97)$$

Therefore, we can write the result in any of the equivalent forms

$$\begin{aligned} \Omega_{\text{gw}}(f) &= \frac{4\pi^2}{3H_0^2} f^3 S_h(f) \\ &= \frac{4\pi^2}{3H_0^2} f^3 I(f) \\ &= \frac{4\pi^2}{3H_0^2} f^3 \int_{S^2} \frac{d^2 \hat{\mathbf{n}}}{4\pi} I(f, \hat{\mathbf{n}}). \end{aligned} \quad (98)$$

In the literature, quantities analogous to $\Omega_{\text{gw}}(f)$ for the Stokes parameters associated with polarization are sometimes defined. For instance, Refs. [42, 66], together

with $\Omega^I = [(4\pi^2 f^3)/(3H_0^2)]I(f)$, also define a quantity $\Omega^V = [(4\pi^2 f^3)/(3H_0^2)]V(f)$ associated with circular polarization. One must be careful, however, with the interpretation of the latter quantity. As we showed here, there is no energy density associated with the V , U , and Q Stokes parameters. Only the spectral density associated with I contributes to the energy density. Of course, one can always multiply any spectral density, or any Stokes parameter, by $[(4\pi^2 f^3)/(3H_0^2)]$, in analogy to Eq. (88). This can sometimes be useful because, while any spectral density $S_h(f)$ has dimensions of $1/f$, and in the GW context typically has numerical values which are not very natural, such as 10^{-44} Hz^{-1} , the quantity $[(4\pi^2 f^3)/(3H_0^2)]S_h(f)$ is dimensionless and is naturally compared with the actual physical energy density per unit logarithmic frequency (normalized to ρ_c) of a GW background. The same can be done for the spectral density of the noise $S_n(f)$ (defined from the noise-noise correlator), from which it can be useful to form the dimensionless combination $[(4\pi^2 f^3)/(3H_0^2)]S_n(f)$. However, again, there is no physical energy density associated with this definition. One could simply work at the level of spectral densities, comparing e.g. the spectral density of the signal, $[S_h^{\text{astro}}(f)]_{AA'}$, with the spectral density of the noise, $S_n(f)$, or comparing among them the contributions to $[S_h^{\text{astro}}(f)]_{AA'}$ associated with the different Stokes parameters. Again, one can, of course, multiply all these spectral densities by $[(4\pi^2 f^3)/(3H_0^2)]$ to deal with dimensionless quantities, but one must then be aware that only the spectral density associated with the I Stokes parameter has an interpretation in terms of an energy density, and there is no energy density associated with the other Stokes parameters, or to noise.

V. EFFECT OF SHOT NOISE

In the previous section we have computed explicitly the averages over extrinsic parameters. However, when we deal with a finite ensemble of N_{ev} events collected in an observation time T , corresponding to a given realization of the underlying stochastic process, these formal averages will not represent the actually observed quantities. For instance, we will not have an infinite number of sources with polarization angle ψ distributed uniformly in $[0, 2\pi]$, but rather a specific sample of sources, with polarization angles $\psi_1, \dots, \psi_{N_{\text{ev}}}$; then, quantities that would vanish in a formal average over ψ computed as in Eq. (16), will in general be non-vanishing because the cancellations will only be partial, reflecting the discreteness of the underlying stochastic process, i.e. shot noise.

In this section, therefore, we no longer perform analytically the averages over the extrinsic parameters, and we instead work directly with the original sums over discrete events, with the aim of extracting, in the large N_{ev} limit, the difference between the discrete and continuous average. We will then eventually compute the results numerically, on a given specific realization of events. The formulas

below will be written for a generic number of events N_{ev} . Typically, we will be interested in the situation where N_{ev} is the number of events, in the given observation time T , that were below the detection threshold, since the actual astrophysical stochastic background is made by the superposition of the unresolved signals. However, the results below are general, and could also be applied to the sum of resolved and unresolved events.

For the astrophysical stochastic background, there are two situations where the effect of shot noise can be especially important. One is the computation of the polarization: we have seen that linear polarizations are set to zero by the average over the polarization angles ψ_i , while circular polarization survives this average but is set to zero by the average over $\cos \iota_i$. The second situation concerns the angular distribution of an astrophysical background. Even if the background is assumed to be exactly isotropic, anisotropies will be generated by shot noise. In both cases, these shot noise contributions could mask very interesting effects, such as a polarized cosmological background that might be generated by parity-violation mechanism in the early Universe [4, 44, 61–67], or actual anisotropies of the astrophysical background that provide information on the clustering of the matter distribution [41, 47–60]. It is therefore important to evaluate these shot-noise effects. The shot noise contribution to the polarization has recently been investigated in Ref. [42] (restricting, however, to circular polarization), while the anisotropies of the astrophysical background induced by shot noise, and their effect on GW observations, have been discussed in Ref. [41, 71].

The formalism that we have developed in Sec. III is well-suited to evaluate the shot noise contribution to the spectral density of the astrophysical background. In Sec. V A and V D we will then show how to evaluate these effects with our formalism. In this paper, we focus on the methodological aspects, and we will illustrate our results by summing over the contribution of all the sources that merge in a given time span (we will assume one year). However, at a specific detector network, some sources will be resolved and some will not; the resolved sources can then be subtracted, and the actual residual contribution will be due only to the unresolved sources, plus the accumulation of errors on the reconstruction of resolved sources. We defer a detailed study at specific 3G detector network to subsequent work.

Our starting point is given by Eq. (12), from which it follows that

$$\begin{aligned} & \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \\ &= \sum_{i,j=1}^{N_{\text{ev}}} \tilde{h}_{A,i}^*(f; \psi_i, t_i, \cos \iota_i) \tilde{h}_{A',j}(f'; \psi_j, t_j, \cos \iota_j) \\ & \quad \times \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \delta(\hat{\mathbf{n}}' - \hat{\mathbf{n}}_j), \end{aligned} \quad (99)$$

which is Eq. (26) before performing on it any average. Actually, the full dependence of $\tilde{h}_A(f, \hat{\mathbf{n}})$ and $\tilde{h}_{A'}(f', \hat{\mathbf{n}}')$ on the extrinsic parameters is as in Eq. (25), but we do not write it explicitly for notational simplicity.

We are not interested in the shot noise effect on the statistics of the times of arrival: it is difficult to imagine physical mechanism that, on the timescale T of the observation, would correlate the times of arrival of different CBCs, so the statistics of the times of arrival is just that of independent events, and does not carry specific astrophysical information. We therefore still perform the corresponding average as in Eq. (15). Then, we write

$$\begin{aligned} & \langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_{\{t_k\}} \\ &= \sum_{i,j=1}^{N_{\text{ev}}} \langle \tilde{h}_{A,i}^*(f; \psi_i, t_i, \cos \iota_i) \tilde{h}_{A',j}(f'; \psi_j, t_j, \cos \iota_j) \rangle_{\{t_k\}} \\ & \quad \times \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \delta(\hat{\mathbf{n}}' - \hat{\mathbf{n}}_j). \end{aligned} \quad (100)$$

As we have seen in Sec. III B, the average over arrival times produces a factor $(1/T)\delta_T(f - f')$, as well as a factor δ_{ij} (cf. footnote 11). Then, using also

$$\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \delta(\hat{\mathbf{n}}' - \hat{\mathbf{n}}_i) = \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}') \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i), \quad (101)$$

we get

$$\begin{aligned} & \langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_{\{t_k\}} = \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}') \delta_T(f - f') \quad (102) \\ & \quad \times \frac{1}{T} \sum_{i=1}^{N_{\text{ev}}} \tilde{h}_{A,i}^*(f; \psi_i, \cos \iota_i) \tilde{h}_{A',i}(f'; \psi_i, \cos \iota_i) \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i). \end{aligned}$$

This expression will be our starting point for evaluating the effect of shot noise, either on the polarization or on the anisotropies.

A. Polarization from shot noise

As we saw in Sec. III, linear polarizations are set to zero by the average over the polarization angles ψ_i , while circular polarization survives this average but is set to zero by the average over $\cos \iota_i$. In this section, we therefore do not perform explicitly these averages. In contrast, we are not interested here in the shot noise effect on the anisotropies. Therefore, we perform analytically the average over $\hat{\mathbf{n}}_k$, $k = 1, \dots, N_{\text{ev}}$ in Eq. (103) using Eq. (59), which gives

$$\begin{aligned} & \langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_{\{t_k, \hat{\mathbf{n}}_k\}} \\ &= \mu(\hat{\mathbf{n}}) \frac{\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')}{4\pi} \delta_T(f - f') \\ & \quad \times \frac{1}{T} \sum_{i=1}^{N_{\text{ev}}} \tilde{h}_{A,i}^*(f; \psi_i, \cos \iota_i) \tilde{h}_{A',i}(f'; \psi_i, \cos \iota_i). \end{aligned} \quad (103)$$

To compute this expression, we begin by extracting explicitly the dependence of $\tilde{h}_{A,i}(f; \psi_i, \cos \iota_i)$ on ψ_i . Using Eq. (31), we have

$$\begin{aligned} & \tilde{h}_{A,i}^*(f; \psi_i, \cos \iota_i) \tilde{h}_{A',i}(f'; \psi_i, \cos \iota_i) \quad (104) \\ &= M_{AA',BB'} \tilde{h}_{B,i}^*(f; \psi_i = 0, \cos \iota_i) \tilde{h}_{B',i}(f'; \psi_i = 0, \cos \iota_i), \end{aligned}$$

where

$$M_{AA',BB'} = R_{AB}(2\psi_i)R_{A'B'}(2\psi_i). \quad (105)$$

Treating $M_{AA',BB'}$ as a 2×2 complex matrix in the (A, A') indices, we can decompose it in the basis of the identity matrix $\delta_{AA'}$ and the Pauli matrices $\sigma_{AA'}^a$ ($a = 1, 2, 3$), with complex coefficients that are matrices in the (B, B') indices,

$$M_{AA',BB'} = \delta_{AA'}\alpha_{BB'} + \sum_{a=1}^3 \sigma_{AA'}^a \beta_{BB'}^a. \quad (106)$$

The matrix $\alpha_{BB'}$ can be obtained contracting Eq. (106) with $\delta_{AA'}$. This gives

$$\begin{aligned} 2\alpha_{BB'} &= \sum_{AA'} \delta_{AA'} M_{AA',BB'} \\ &= [R^T(2\psi_i)R(2\psi_i)]_{BB'}, \end{aligned} \quad (107)$$

where $R^T(2\psi_i)$ is the transpose matrix. Since R is a rotation matrix, $[R^T(2\psi_i)R(2\psi_i)]_{BB'} = \delta_{BB'}$ and we get

$$2\alpha_{BB'} = \delta_{BB'}. \quad (108)$$

Similarly, contracting both sides of Eq. (106) with $\sigma_{A'A}^b$ and using $\text{Tr}(\sigma^a \sigma^b) = 2\delta^{ab}$, we get

$$\begin{aligned} 2\beta_{BB'}^b &= \sum_{AA'} \sigma_{A'A}^b M_{AA',BB'} \\ &= [R^T(2\psi_i)(\sigma^b)^T R(2\psi_i)]_{BB'}. \end{aligned} \quad (109)$$

Writing $R(2\psi_i)$ as in Eq. (33) and recalling from Eq. (11) that $(\sigma^2)^T = -\sigma^2$ while $(\sigma^1)^T = \sigma^1$ and $(\sigma^3)^T = \sigma^3$, we get

$$2\beta_{BB'}^1 = \sigma_{BB'}^1 \cos 4\psi_i + \sigma_{BB'}^3 \sin 4\psi_i, \quad (110)$$

$$2\beta_{BB'}^2 = -\sigma_{BB'}^2, \quad (111)$$

$$2\beta_{BB'}^3 = -\sigma_{BB'}^1 \sin 4\psi_i + \sigma_{BB'}^3 \cos 4\psi_i. \quad (112)$$

Inserting this into Eq. (103),

$$\begin{aligned} &\langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_{\{t_k, \hat{\mathbf{n}}_k\}} \\ &= \mu(\hat{\mathbf{n}}) \frac{\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')}{4\pi} \delta_T(f - f') \frac{1}{2} \sum_{i=1}^{N_{\text{ev}}} \\ &\quad \times \left[\mathcal{I}_i(f; \cos \iota_i) \delta_{AA'} + \mathcal{U}_i(f; \cos \iota_i, \psi_i) \sigma_{AA'}^1 \right. \\ &\quad \left. + \mathcal{V}_i(f; \cos \iota_i) \sigma_{AA'}^2 + \mathcal{Q}_i(f; \cos \iota_i, \psi_i) \sigma_{AA'}^3 \right], \end{aligned} \quad (113)$$

where

$$\mathcal{I}_i(f; \cos \iota_i) = \frac{1}{T} \sum_{B=+, \times} |\tilde{h}_{B,i}(f; \psi_i = 0, \cos \iota_i)|^2, \quad (114)$$

$$\mathcal{V}_i(f; \cos \iota_i) = \frac{1}{T} \sum_{B, B'=+, \times} \quad (115)$$

$$\times [-\tilde{h}_{B,i}^*(f; \psi_i = 0, \cos \iota_i) \sigma_{BB'}^2 \tilde{h}_{B',i}(f; \psi_i = 0, \cos \iota_i)],$$

$$\mathcal{U}_i(f; \cos \iota_i, \psi_i) = \frac{1}{T} \sum_{B, B'=+, \times} \tilde{h}_{B,i}^*(f; \psi_i = 0, \cos \iota_i) \quad (116)$$

$$\times [\sigma_{BB'}^1 \cos 4\psi_i + \sigma_{BB'}^3 \sin 4\psi_i] \tilde{h}_{B',i}(f; \psi_i = 0, \cos \iota_i),$$

and

$$\mathcal{Q}_i(f; \cos \iota_i, \psi_i) = \frac{1}{T} \sum_{B, B'=+, \times} \tilde{h}_{B,i}^*(f; \psi_i = 0, \cos \iota_i) \quad (117)$$

$$\times [-\sigma_{BB'}^1 \sin 4\psi_i + \sigma_{BB'}^3 \cos 4\psi_i] \tilde{h}_{B',i}(f; \psi_i = 0, \cos \iota_i).$$

As we discussed below Eq. (54), \mathcal{I}_i and \mathcal{V}_i are independent of the reference value $\psi_i = 0$, so we can replace the argument $\psi_i = 0$ in Eqs. (114) and (115) by an arbitrary value ψ_i , i.e.

$$\mathcal{I}_i(f; \cos \iota_i) = \frac{1}{T} \sum_{B=+, \times} |\tilde{h}_{B,i}(f; \psi_i, \cos \iota_i)|^2, \quad (118)$$

$$\mathcal{V}_i(f; \cos \iota_i) = -\frac{1}{T} \sum_{B, B'=+, \times} \quad (119)$$

$$\times \tilde{h}_{B,i}^*(f; \psi_i, \cos \iota_i) \sigma_{BB'}^2 \tilde{h}_{B',i}(f; \psi_i, \cos \iota_i),$$

or simply suppress the argument ψ_i , as we will do below. In contrast, as we will see in a moment, \mathcal{Q}_i and \mathcal{U}_i depend on ψ_i . Using Eqs. (109) and (110) we see that

$$\sigma^1 \cos 4\psi_i + \sigma^3 \sin 4\psi_i = R^T(2\psi_i) \sigma^1 R(2\psi_i), \quad (120)$$

and therefore Eq. (116) can be rewritten as

$$\mathcal{U}_i(f; \cos \iota_i, \psi_i) = \frac{1}{T} \sum_{A, A'=+, \times} \quad (121)$$

$$\times \tilde{h}_{A,i}^*(f; \psi_i, \cos \iota_i) \sigma_{AA'}^1 \tilde{h}_{A',i}(f; \psi_i, \cos \iota_i),$$

and similarly

$$\mathcal{Q}_i(f; \cos \iota_i, \psi_i) = \frac{1}{T} \sum_{A, A'=+, \times} \quad (122)$$

$$\times \tilde{h}_{A,i}^*(f; \psi_i, \cos \iota_i) \sigma_{AA'}^3 \tilde{h}_{A',i}(f; \psi_i, \cos \iota_i).$$

In terms of $h_{L,i}$ and $h_{R,i}$, we have already found that

$$\mathcal{I}_i(f; \cos \iota_i) = \frac{1}{T} [|\tilde{h}_{R,i}(f; \cos \iota_i)|^2 + |\tilde{h}_{L,i}(f; \cos \iota_i)|^2], \quad (123)$$

and

$$\mathcal{V}_i(f; \cos \iota_i) = \frac{1}{T} [|\tilde{h}_{R,i}(f; \cos \iota_i)|^2 - |\tilde{h}_{L,i}(f; \cos \iota_i)|^2], \quad (124)$$

cf. Eqs. (64) and (65) (we are now also writing explicitly the argument $\cos \iota_i$). Similarly, we get

$$\mathcal{U}_i(f; \psi_i, \cos \iota_i) = \frac{i}{T} [\tilde{h}_{R,i}^*(f; \psi_i, \cos \iota_i) \tilde{h}_{L,i}(f; \psi_i, \cos \iota_i)$$

$$-\tilde{h}_{L,i}^*(f; \psi_i, \cos \iota_i) \tilde{h}_{R,i}(f; \psi_i, \cos \iota_i)], \quad (125)$$

and

$$\begin{aligned} \mathcal{Q}_i(f; \psi_i, \cos \iota_i) &= \frac{1}{T} [\tilde{h}_{R,i}^*(f; \psi_i, \cos \iota_i) \tilde{h}_{L,i}(f; \psi_i, \cos \iota_i) \\ &+ \tilde{h}_{L,i}^*(f; \psi_i, \cos \iota_i) \tilde{h}_{R,i}(f; \psi_i, \cos \iota_i)]. \quad (126) \end{aligned}$$

Using Eq. (54) we then see that, under $\psi_i \rightarrow \psi_i + \psi_0$, \mathcal{U}_i and \mathcal{Q}_i transform linearly among them, as

$$\mathcal{U}_i \rightarrow \cos(4\psi_0)\mathcal{U}_i + \sin(4\psi_0)\mathcal{Q}_i, \quad (127)$$

$$\mathcal{Q}_i \rightarrow -\sin(4\psi_0)\mathcal{U}_i + \cos(4\psi_0)\mathcal{Q}_i, \quad (128)$$

so that

$$\mathcal{Q}_i + i\mathcal{U}_i \rightarrow e^{i4\psi_0}(\mathcal{Q}_i + i\mathcal{U}_i), \quad (129)$$

$$\mathcal{Q}_i - i\mathcal{U}_i \rightarrow e^{-i4\psi_0}(\mathcal{Q}_i - i\mathcal{U}_i). \quad (130)$$

Therefore, while \mathcal{I}_i and \mathcal{V}_i transform as helicity-0 fields under rotations in the plane transverse to $\hat{\mathbf{n}}_i$, \mathcal{U}_i and \mathcal{Q}_i transform as helicity-4 field [4, 44, 72]. In particular, this implies that \mathcal{U}_i and \mathcal{Q}_i depend explicitly on the polarization angle ψ_i (and therefore on the choice of the reference value $\psi_i = 0$), which was not the case for \mathcal{I}_i and \mathcal{V}_i . Note, however, that the combination

$$\mathcal{P}_i = (\mathcal{U}_i^2 + \mathcal{Q}_i^2)^{1/2}, \quad (131)$$

is again a scalar under rotations of the polarization angle, and therefore does not depend on the choice of the reference angle $\psi_i = 0$.

So, in conclusion, Eq. (113) becomes

$$\begin{aligned} \langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_{\{t_k, \hat{\mathbf{n}}_k\}} \\ = \frac{\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')}{4\pi} \delta_T(f - f') \frac{1}{2} [S_h^{\text{astro}}(f; \hat{\mathbf{n}})]_{AA'}, \quad (132) \end{aligned}$$

where the spectral density $[S_h^{\text{astro}}(f; \hat{\mathbf{n}})]_{AA'}$ factorizes as

$$[S_h^{\text{astro}}(f; \hat{\mathbf{n}})]_{AA'} = \mu(\hat{\mathbf{n}}) [S_h^{\text{astro}}(f)]_{AA'}, \quad (133)$$

and

$$\begin{aligned} [S_h^{\text{astro}}(f)]_{AA'} &= I(f)\delta_{AA'} + V(f)\sigma_{AA'}^2 \\ &+ U(f)\sigma_{AA'}^1 + Q(f)\sigma_{AA'}^3. \quad (134) \end{aligned}$$

The explicit expressions for the Stokes parameters, reinstating explicitly all the dependencies as in Eqs. (24) and (25), are

$$I(f; \hat{\mathbf{n}}, \{\cos \iota_k\}) = \mu(\hat{\mathbf{n}})I(f; \{\cos \iota_k\}), \quad (135)$$

$$V(f; \hat{\mathbf{n}}, \{\cos \iota_k\}) = \mu(\hat{\mathbf{n}})V(f; \{\cos \iota_k\}), \quad (136)$$

$$U(f; \hat{\mathbf{n}}, \{\cos \iota_k\}, \{\psi_k\}) = \mu(\hat{\mathbf{n}})U(f; \{\cos \iota_k\}, \{\psi_k\}), \quad (137)$$

$$Q(f; \hat{\mathbf{n}}, \{\cos \iota_k\}, \{\psi_k\}) = \mu(\hat{\mathbf{n}})Q(f; \{\cos \iota_k\}, \{\psi_k\}), \quad (138)$$

where

$$I(f; \{\cos \iota_k\}) = \sum_{i=1}^{N_{\text{ev}}} \mathcal{I}_i(f; \cos \iota_i), \quad (139)$$

$$V(f; \{\cos \iota_k\}) = \sum_{i=1}^{N_{\text{ev}}} \mathcal{V}_i(f; \cos \iota_i), \quad (140)$$

$$U(f; \{\cos \iota_k\}, \{\psi_k\}) = \sum_{i=1}^{N_{\text{ev}}} \mathcal{U}_i(f; \cos \iota_i, \psi_i), \quad (141)$$

$$Q(f; \{\cos \iota_k\}, \{\psi_k\}) = \sum_{i=1}^{N_{\text{ev}}} \mathcal{Q}_i(f; \cos \iota_i, \psi_i). \quad (142)$$

and $\mathcal{I}_i(f; \cos \iota_i), \dots, \mathcal{Q}_i(f; \cos \iota_i)$ are given in Eqs. (114)–(117) or, equivalently, in Eqs. (118), (119), (121) and (122).

For an ensemble of events we also define

$$\begin{aligned} P(f; \{\cos \iota_k\}, \{\psi_k\}) \\ = [U^2(f; \{\cos \iota_k\}, \{\psi_k\}) + Q^2(f; \{\cos \iota_k\}, \{\psi_k\})]^{1/2}, \quad (143) \end{aligned}$$

or, if we do not want to write explicitly all arguments $\{\cos \iota_k\}, \{\psi_k\}$,

$$P(f) = [U^2(f) + Q^2(f)]^{1/2}. \quad (144)$$

Note that, from Eqs. (127) and (128) together with Eqs. (141) and (142), under a common transformation of all angles ψ_i as $\psi_i \rightarrow \psi_i + \psi_0$, we have (suppressing for notational simplicity all arguments)

$$U \rightarrow \cos(4\psi_0)U + \sin(4\psi_0)Q, \quad (145)$$

$$Q \rightarrow -\sin(4\psi_0)U + \cos(4\psi_0)Q, \quad (146)$$

and therefore P is invariant, i.e. P has helicity zero.¹³

The first two terms in Eq. (134), proportional to $\delta_{AA'}$ and to $\sigma_{AA'}^2$, are the same that we found performing the average over ψ_i by integrating over $d\psi_i/(2\pi)$. We see that they do not receive any correction from shot noise, cf. Eqs. (46)–(52). In contrast, we see that, because of shot noise, we get terms proportional to both $\sigma_{AA'}^1$ and $\sigma_{AA'}^3$, so the Stokes parameters U and Q associated with linear polarization are now non-vanishing, cf. Eq. (10). The results of Sec. III are recovered if we now superimpose an analytic average over polarization angles, since

$$\langle \cos 4\psi_i \rangle_{\psi_i} = \int_0^{2\pi} \frac{d\psi_i}{2\pi} \cos 4\psi_i = 0, \quad (148)$$

and similarly for the average of $\sin 4\psi_i$. Therefore, the averages over ψ_i of \mathcal{U}_i and of \mathcal{Q}_i vanish, as we see more easily from Eqs. (116) and (117).

¹³ Observe that, in terms of \mathcal{U}_i and \mathcal{Q}_i , we have defined

$$P = \left[\left(\sum_i \mathcal{U}_i \right)^2 + \left(\sum_i \mathcal{Q}_i \right)^2 \right]^{1/2} \quad (149)$$

rather than as $P = \sum_i \mathcal{P}_i$, where \mathcal{P}_i is defined by Eq. (131). The reason is that, in a stochastic background, the observable quantities are related to the correlator $\langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle$ and therefore to the Stokes parameters I, V, U and Q , see Eq. (9). A quantity such as $\sum_i \mathcal{P}_i$, in contrast, is not directly observable.

B. Linear polarizations and spatial isotropy

From Eqs. (137) and (138) we see that, once we include the effect of shot noise, linear polarization is in principle allowed even in the isotropic case, $\mu(\hat{\mathbf{n}}) = 1$. This is apparently at odds with the observation [4, 44] that, because of Eqs. (129) and (130), under a rotation $\psi_i \rightarrow \psi_i + \psi_0$ we have

$$Q(f; \hat{\mathbf{n}}) \pm iU(f; \hat{\mathbf{n}}) \rightarrow e^{\pm i4\psi_0} [Q(f; \hat{\mathbf{n}}) \pm iU(f; \hat{\mathbf{n}})]. \quad (149)$$

Therefore $Q(f; \hat{\mathbf{n}}) \pm iU(f; \hat{\mathbf{n}})$ have helicities ± 4 , and can be expanded in (± 4) spin-weighted spherical harmonics, ${}_{\pm 4}Y_{lm}(\hat{\mathbf{n}})$, as

$$Q(f; \hat{\mathbf{n}}) \pm iU(f; \hat{\mathbf{n}}) = \sum_{l=4}^{\infty} \sum_{m=-l}^l C_{lm}^{\pm}(f) {}_{\pm 4}Y_{lm}(\hat{\mathbf{n}}), \quad (150)$$

with the expansion starting from $l = 4$, since ${}_{\pm 4}Y_{lm}(\hat{\mathbf{n}})$ vanish for $l < 4$. The argument in Ref. [44], then, is that an isotropic distribution of sources, that only has the $l = 0$ multipole, cannot produce a non-vanishing value for $Q(f; \hat{\mathbf{n}}) \pm iU(f; \hat{\mathbf{n}})$, that both start from $l = 4$.

In contrast, from our explicit computation, we have found that both $U(f; \hat{\mathbf{n}})$ and $Q(f; \hat{\mathbf{n}})$ are proportional to $\mu(\hat{\mathbf{n}})$, which is a scalar function and can therefore be expanded in scalar spherical harmonics $Y_{lm}(\hat{\mathbf{n}})$, which starts from $l = 0$. In particular, for an isotropic distribution, $\mu(\hat{\mathbf{n}}) = 1$ and we find that $U(f; \hat{\mathbf{n}})$ and $Q(f; \hat{\mathbf{n}})$ can be non-vanishing.

The resolution of this apparent paradox is that the labels (l, m) in Eq. (150) are in fact an abuse of notation, which induces one to believe that these are the indices associated with the orbital angular momentum operator. This is not true, and these indices are associated with the total angular momentum operator, and should be more appropriately denoted as (j, j_z) , so Eq. (150) should rather be written as

$$Q(f; \hat{\mathbf{n}}) \pm iU(f; \hat{\mathbf{n}}) = \sum_{j=4}^{\infty} \sum_{j_z=-j}^j C_{jj_z}^{\pm}(f) {}_{\pm 4}Y_{jj_z}(\hat{\mathbf{n}}). \quad (151)$$

The issue was already discussed in Sec. 3.5.2 of Ref. [2] (see in particular Note 47 there). To understand this point, let us recall that the tensor spherical harmonics can be obtained by coupling the scalar spherical harmonics $Y_{l_z}(\hat{\mathbf{n}})$ to the spin functions χ_{ss_z} , with the appropriate Clebsch–Gordan coefficients which gives a state with total angular momentum $|j, j_z\rangle$,

$$Y_{jj_z}^{ls}(\theta, \phi) = \sum_{l_z=-l}^l \sum_{s_z=-s}^s \langle sls_z l_z | jj_z \rangle Y_{l_z}(\theta, \phi) \chi_{ss_z}. \quad (152)$$

By definition, the functions $Y_{jj_z}^{ls}(\theta, \phi)$ are eigenvectors of \mathbf{J}^2 , J_z^2 , \mathbf{L}^2 and \mathbf{S}^2 (where \mathbf{L} is the orbital angular momentum operator, \mathbf{S} the spin operator and $\mathbf{J} = \mathbf{L} + \mathbf{S}$), which commute among them, and can be diagonalized simultaneously (note that, in contrast, they are not eigenvectors

of L_z , which does not commute with \mathbf{J}^2 and cannot be diagonalized simultaneously). As an example, consider the vector spherical harmonics (we closely follow Sec. 3.5.2 of Ref. [2]). In this case the spin function χ is a vector, that we denote by $\boldsymbol{\xi}$. The eigenfunctions of S_z with eigenvalues $s_z = 0, \pm 1$ can be constructed from the unit vectors \mathbf{e}_x , \mathbf{e}_y and \mathbf{e}_z as

$$\boldsymbol{\xi}^{(\pm 1)} = \mp \frac{1}{\sqrt{2}} (\mathbf{e}_x \pm i\mathbf{e}_y), \quad \boldsymbol{\xi}^{(0)} = \mathbf{e}_z. \quad (153)$$

Then the vector spherical harmonics are

$$\mathbf{Y}_{jj_z}^l(\theta, \phi) = \sum_{l_z=-l}^l \sum_{s_z=0, \pm 1} \langle ll_s l_z | jj_z \rangle Y_{l_z}(\theta, \phi) \boldsymbol{\xi}^{(s_z)}, \quad (154)$$

where we have written $\mathbf{Y}_{jj_z}^{l1}$ simply as $\mathbf{Y}_{jj_z}^l$, since the fact that $s = 1$ is already implicit in the vector notation \mathbf{Y} . The vectors $\mathbf{Y}_{jj_z}^l(\theta, \phi)$ can have $j = l - 1, l, l + 1$ (if $l \neq 0$), or $j = 1$ if $l = 0$ (in a quantum language, the possible quantum combinations of spin $s = 1$ and orbital angular momentum l), but have no special property with respect to the radial unit vector $\hat{\mathbf{n}}$, i.e., they are neither purely transverse nor purely longitudinal. With them, we can however form the combinations

$$\mathbf{Y}_{jj_z}^E = (2j + 1)^{-1/2} [(j + 1)^{1/2} \mathbf{Y}_{jj_z}^{j-1} + j^{1/2} \mathbf{Y}_{jj_z}^{j+1}] \quad (155)$$

$$\mathbf{Y}_{jj_z}^B = i\mathbf{Y}_{jj_z}^j, \quad (156)$$

$$\mathbf{Y}_{jj_z}^R = (2j + 1)^{-1/2} [j^{1/2} \mathbf{Y}_{jj_z}^{j-1} - (j + 1)^{1/2} \mathbf{Y}_{jj_z}^{j+1}] \quad (157)$$

with $j \geq 1$, together with $\mathbf{Y}_{00}^R = Y_{00}\hat{\mathbf{n}}$; note that $\mathbf{Y}_{jj_z}^E$ and $\mathbf{Y}_{jj_z}^B$ are superposition of vector harmonics with different values of l , so they are no longer eigenfunctions of \mathbf{L}^2 . However, it can be shown that they are transverse to the propagation direction $\hat{\mathbf{n}}$. Since $\mathbf{Y}_{jj_z}^E$ and $\mathbf{Y}_{jj_z}^B$ start at $j = 1$, while $\mathbf{Y}_{jj_z}^R$ starts at $j = 0$, an arbitrary vector field $\mathbf{V}(t, r, \theta, \phi)$ can be expanded as

$$\mathbf{V}(t, r, \theta, \phi) = \sum_{j=0}^{\infty} \sum_{j_z=-j}^j R_{jj_z}(t, r) \mathbf{Y}_{jj_z}^R(\theta, \phi) + \sum_{j=1}^{\infty} \sum_{j_z=-j}^j [E_{jj_z}(t, r) \mathbf{Y}_{jj_z}^E(\theta, \phi) + B_{jj_z}(t, r) \mathbf{Y}_{jj_z}^B(\theta, \phi)]. \quad (158)$$

With a common abuse of notation, the indices (j, j_z) , that appear here, are typically written as (l, m) but it is clear, from the above derivation, that they label the eigenvalues of (\mathbf{J}^2, J_z) , not of (\mathbf{L}^2, L_z) . In several situations we are only interested in the highest helicity states, in this case $h \equiv s_z = \pm 1$. For instance, for a massless spin-1 particle such as the photon, only the states with $h = \pm 1$ are allowed, so the radial functions $R_{jj_z}(t, r) = 0$ vanish, and we only remain with the transverse states, whose multipole expansion starts from $j = 1$. Thus, it is impossible to put a photon in a state with total angular momentum $j = 0$. The vector spherical harmonics $\mathbf{Y}_{jj_z}^E(\theta, \phi)$ and $\mathbf{Y}_{jj_z}^B(\theta, \phi)$

are in correspondence with the ± 1 spin-weighted spherical harmonics ${}_{\pm 1}Y_{jj_z}(\theta, \phi)$. Once again, it is important to stress that the (j, j_z) indices here refer to total angular momentum, even if they are typically written as (l, m) .

The construction can be generalized to higher spin. For instance, for spin 2, one finds a set of five tensor spherical harmonics: $\mathbf{T}_{jj_z}^{S0}$, which has $j \geq 0$; $\mathbf{T}_{jj_z}^{E1}$ and $\mathbf{T}_{jj_z}^{B1}$, which have $j \geq 1$; and $\mathbf{T}_{jj_z}^{E2}$ and $\mathbf{T}_{jj_z}^{B2}$, which have $j \geq 2$. If one is studying a massive spin-2 field (such as a massive graviton), these five tensors field would describe the corresponding five degrees of freedom. However, if one is interested only in the states with helicity $h = \pm 2$, as for a GW (or a massless graviton), one must select the transverse-traceless tensor spherical harmonics $\mathbf{T}_{jj_z}^{E2}$ and $\mathbf{T}_{jj_z}^{B2}$, which can be mapped into the ± 2 spin-weighted spherical harmonics ${}_{\pm 2}Y_{jj_z}(\theta, \phi)$.¹⁴ The construction can be generalized to all higher spins.

Having understood that the indices in Eq. (151) refer to the total angular momentum, rather than to the orbital one, we see that there is no contradiction between the fact that the sum in Eq. (151) starts at $j = 4$, while an isotropic distribution of sources only has $l = 0$. Simply, we see from Eq. (149) that $Q \pm iU$ have $s = 4$, $s_z = \pm 4$. This operator combines with $\mu(\hat{\mathbf{n}})$, which in the isotropic case only has multipoles $l = 0, l_z = 0$, to build an object which has $j = 4, j_z = \pm 4$, and which can therefore appear as the lowest term of the expansion (151).

An equivalent, and in fact faster way of reaching the same conclusion is to observe that, with U and Q , we can form the combination P defined in Eq. (144), which is a scalar, and therefore can in principle receive contribution from the $l = m = 0$ multipole of the source distribution.

We conclude that (contrary to statements in the literature) an isotropic distribution of sources can in principle produce a GW background which also has a linear polarization.

C. Numerical results

The expressions given by Eqs. (139)–(142), together with Eqs. (114)–(117), are a natural starting point for a numerical evaluation of the Stokes parameters generated by a given finite ensemble of sources, including the effect of shot noise. While the above formalism holds for a generic set of N_{ev} signals, in practical applications the sums will only run over the unresolved sources, since these are those that contribute to the stochastic background. Similarly to what has been discussed in the literature for the energy density (see Refs. [26–40]), i.e. for the I Stokes parameter, one will also have to add to this the effect due to the accumulation of errors from the reconstruction of the resolved sources.

All these aspects, however, will be very dependent on the specific detector network considered, since this determines which sources will be resolved (in a given noise realization), and the accuracy of their reconstruction. We leave a full analysis for specific 3G detector networks for future work. In this paper we are rather concerned with the methodological aspects, and we illustrate here our results computing the Stokes parameters on an ensemble of BBHs and BNSs corresponding to one year of data in our population model, independently of whether, at a given network, these sources will be resolved or unresolved. In this sense, our results are really an upper bound on the actual contribution of the astrophysical background to the Stokes parameters or, equivalently, corresponds to the limit in which the threshold for CBC detection is set very high, so that all sources are deemed unresolved.

To compute the sums in Eqs. (139)–(142) we use the same catalogs of BBHs and BNSs described in Ref. [24], and from it we extract a random sample of events corresponding to one year of observation which, in our population model, corresponds to about 1×10^5 BBH and 7×10^5 BNS.¹⁵

The results are shown in Fig. 1, where we plot $I(f)$, $V(f)$, $Q(f)$, $U(f)$, as well as $P(f) = \sqrt{U^2(f) + Q^2(f)}$ for BBHs (left panel) and for BNSs (right panel). We see that the polarization parameters, even if suppressed with respect to I , are indeed present, and follow roughly the same frequency dependence. For BBHs the linear polarization parameter P is actually of the same order of magnitude as the circular polarization parameter V and in fact even slightly larger, and receives most of its contribution from Q , while U is smaller (so that the red curve describing $Q(f)$ is almost identical to the violet curve describing $P(f)$, and not visible because is covered by it, except at the highest frequencies). For the BNS case, the polarization terms are more suppressed with respect to I , which can be attributed to the larger sample of sources, and the polarization parameter V is dominant with respect to the other (suggesting a potentially different scaling with the number of events), while the hierarchy between Q and U is unchanged.

D. Anisotropies from shot noise

We next address the effect of shot noise on the anisotropies of the angular distribution. These will be a foreground to the (more physically interesting) anisotropies of the actual distribution of astrophysical sources, that can carry important information on the clustering of the underlying matter distribution. To this purpose we start from Eq. (58) and, rather than performing the average $\langle \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i) \rangle_{\hat{\mathbf{n}}_i}$ through Eq. (59), we leave it

¹⁴ See App. F and G of Ref. [4] for the explicit mapping between spin-weighted spherical harmonics and tensor spherical harmonics, for the spin-1 and spin-2 cases.

¹⁵ For the BNS case, we use the catalog featuring a Gaussian mass distribution of the sources, and adimensional tidal deformabilities computed assuming the APR equation of state [73].




FIG. 1. The quantities $I(f)$, $V(f)$, $Q(f)$, $U(f)$, as well as $P(f) = \sqrt{U^2(f) + Q^2(f)}$, for an ensemble of $\sim 10^5$ BBHs (left panel) and $\sim 7 \times 10^5$ BNSs (right panel), corresponding to 1 yr of data in our population model.




FIG. 2. The functions $c_l(f)$ for a selection of values $l = [1, 2, 3, 4, 5, 10, 50, 100]$, for an ensemble of $\sim 10^5$ BBHs (left panel) and $\sim 7 \times 10^5$ BNSs (right panel), corresponding to 1 yr of data in our population model.

in the form of a sum over the N_{ev} events. We also neglect the contribution from circular and linear polarizations that, as we have seen, are subdominant, and we leave the average of $\mathcal{I}_i(f)$ over $\cos \iota$ in the form of an explicit sum over the events, performing only the averages over polarization angles and times of arrival. We therefore write

$$\langle \tilde{h}_A^*(f, \hat{\mathbf{n}}) \tilde{h}_{A'}(f', \hat{\mathbf{n}}') \rangle_{\{\psi_k, t_k\}} = \delta_{AA'} \delta_T(f - f')$$

$$\times \frac{\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')}{4\pi} \frac{1}{2} I(f; \hat{\mathbf{n}}), \quad (159)$$

where

$$I(f; \hat{\mathbf{n}}) = 4\pi \sum_{i=1}^{N_{\text{ev}}} \mathcal{I}_i(f) \delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}_i), \quad (160)$$

and it is understood that $\mathcal{I}_i(f)$ is actually $\mathcal{I}_i(f; \cos \iota_i)$ and $I(f; \hat{\mathbf{n}})$ is actually $I(f; \hat{\mathbf{n}}, \{\cos \iota_k\})$. If we would now

average over $\hat{\mathbf{n}}_i$, using Eq. (19), we would get back Eq. (62) with $I(f)$ given by Eq. (64) (where we had also written explicitly the average over intrinsic parameters). However, here we are interested in computing the shot noise effect from the finite sum over events.

As with any scalar function of $\hat{\mathbf{n}}$, we can expand $I(f; \hat{\mathbf{n}})$ in spherical harmonics,

$$I(f; \hat{\mathbf{n}}) = \sum_{l=0}^{\infty} \sum_{m=-l}^l I_{lm}(f) Y_{lm}(\hat{\mathbf{n}}), \quad (161)$$

whose inversion is

$$\begin{aligned} I_{lm}(f) &= \int_{S^2} d^2\hat{\mathbf{n}} I(f; \hat{\mathbf{n}}) Y_{lm}^*(\hat{\mathbf{n}}) \\ &= 4\pi \sum_{i=1}^{N_{\text{ev}}} \mathcal{I}_i(f) Y_{lm}^*(\hat{\mathbf{n}}_i). \end{aligned} \quad (162)$$

The above expressions are valid for a generic underlying distribution $\mu(\hat{\mathbf{n}})$. We henceforth consider the isotropic case $\mu(\hat{\mathbf{n}}) = 1$, and study how shot noise generates anisotropies in the angular distribution. In this case we can follow the standard approach used for the CMB multipoles: for a given value of l , the coefficients I_{lm} with $m = -l, \dots, l$ are statistically equivalent and all information is contained in the quantities

$$C_l(f) = \frac{1}{2l+1} \sum_{m=-l}^l |I_{lm}(f)|^2. \quad (163)$$

We insert here Eq. (162) and use the addition theorem for spherical harmonics,

$$\sum_{m=-l}^l Y_{lm}(\hat{\mathbf{n}}_i) Y_{lm}^*(\hat{\mathbf{n}}_j) = \frac{2l+1}{4\pi} P_l(\hat{\mathbf{n}}_i \cdot \hat{\mathbf{n}}_j), \quad (164)$$

where P_l are the Legendre polynomials. Then, we get

$$C_l(f) = 4\pi \sum_{i,j=1}^{N_{\text{ev}}} \mathcal{I}_i(f) \mathcal{I}_j(f) P_l(\hat{\mathbf{n}}_i \cdot \hat{\mathbf{n}}_j). \quad (165)$$

Since $P_0(\cos\theta) = 1$, we have

$$C_0(f) = 4\pi I^2(f). \quad (166)$$

We are interested in computing the functions $C_l(f)$ for $l \neq 0$ which, compared to $C_0(f)$, will give a measure of the anisotropies of the astrophysical background generated by shot noise. It is therefore convenient to define

$$\begin{aligned} c_l(f) &= \frac{C_l(f)}{C_0(f)} \\ &= \frac{1}{I^2(f)} \sum_{i,j=1}^{N_{\text{ev}}} \mathcal{I}_i(f) \mathcal{I}_j(f) P_l(\hat{\mathbf{n}}_i \cdot \hat{\mathbf{n}}_j). \end{aligned} \quad (167)$$

We have computed numerically these quantities for our distribution of BBHs and of BNSs, using again one year

of data and taking all sources as unresolved. As already discussed in Sec. V C, this only has an illustrative purpose: at any given detector network, one will have to separate the resolved from unresolved sources, sum over the unresolved ones, and take into account the error in the reconstruction of the resolved sources. Figure 2 shows the functions $c_l(f)$ for a selection of values of l .

For BBHs we see that, for all multipoles shown, below about 100 Hz the anisotropies are at the percent level, and then they grow, reaching values of order one in the kHz region. This can be understood from the fact that most BBHs merge at lower frequencies; therefore, the effective number of BBHs contributing to the result in the kHz region is just a small fraction of the total, and the relative effect of shot noise increases when the effective number of sources decreases.

For BNS we observe that the anisotropies are considerably smaller as compared to the BBH case, with values of $\mathcal{O}(10^{-4})$ up to the 100 Hz region. This can be traced on the one hand to the larger amount of sources present in the BNS catalog, and on the other hand to the morphology of their signals in band. Indeed, most of the BNS systems merge at $\mathcal{O}(\text{kHz})$ frequencies, and sweep all through the frequency band of ground-based detectors. Also in this case, for $f \gtrsim 300$ Hz we observe a raise in the spectrum up to values of $\sim 10^{-1}$: this is again because part of the sources, especially at high redshifts, start to merge.

VI. CONCLUSIONS

In this paper we have shown how the spectral density of an ensemble of compact binary coalescences emerges from the average of the extrinsic parameters of the population: the times of arrival, the polarization angles, the arrival directions and the orbit inclinations. We have seen the role of each of these averages in arriving at the final result (83), which is often used in the literature although, to our knowledge, it lacked any convincing derivation. The step-by-step procedure that we have described allows us to understand the roles of the various averages: the averages over the polarization angles decorrelate the different events or, equivalently, different arrival directions, producing a Dirac delta $\delta(\hat{\mathbf{n}} - \hat{\mathbf{n}}')$ in Eq. (43), and furthermore eliminates the linear polarizations. The averages over arrival times produce a (regularized) Dirac delta $\delta_T(f - f')$ and, finally, the averages over the inclination angles remove the remaining Stokes parameter, associated with circular polarization.

For a finite sample of events, however, all these averages must be replaced with the sum over the actual events of the specific realizations, and the above cancellations are only approximate. Shot noise therefore generates a certain amount of polarization even in the CBC background, and a certain amount of anisotropy. In particular, we have seen that shot noise generates a linear polarization even when the underlying distribution of sources is isotropic (in

contrast with previous statements in the literature). We have computed numerically these shot noise contributions with a realistic catalog of BBHs and of BNSs (treating all sources as undetected). This formalism can be useful to disentangle possible polarized background produced by mechanisms in the early Universe, or anisotropies in the stochastic background that reflect the underlying matter distribution, from contamination effects due to shot noise of the astrophysical background.

ACKNOWLEDGMENTS

The research of F.I., M. Mag. and N.M. is supported by the Swiss National Science Foundation, grant

200020_191957, and by the SwissMap National Center for Competence in Research. E.B. and M. Mag. are supported by the SNSF grant CRSII5_213497. The work of M. Man. received support from the French government under the France 2030 investment plan, as part of the Initiative d'Excellence d'Aix-Marseille Université – A*MIDEX AMX-22-CEI-02. Computations made use of the Baobab cluster at the University of Geneva.

-
- [1] M. Maggiore, *Phys. Rept.* **331**, 283 (2000), [arXiv:gr-qc/9909001 \[gr-qc\]](#).
- [2] M. Maggiore, *Gravitational Waves. Vol. 1: Theory and Experiments* (Oxford University Press, 2007).
- [3] T. Regimbau, *Res. Astron. Astrophys.* **11**, 369 (2011), [arXiv:1101.2762 \[astro-ph.CO\]](#).
- [4] J. D. Romano and N. J. Cornish, *Living Rev. Rel.* **20**, 2 (2017), [arXiv:1608.06889 \[gr-qc\]](#).
- [5] M. Maggiore, *Gravitational Waves. Vol. 2: Astrophysics and Cosmology* (Oxford University Press, 2018).
- [6] C. Caprini and D. G. Figueroa, *Class. Quant. Grav.* **35**, 163001 (2018), [arXiv:1801.04268 \[astro-ph.CO\]](#).
- [7] N. Christensen, *Rept. Prog. Phys.* **82**, 016903 (2019), [arXiv:1811.08797 \[gr-qc\]](#).
- [8] J. Lawrence, K. Turbang, A. Matas, A. I. Renzini, N. van Remortel, and J. D. Romano, *Phys. Rev. D* **107**, 103026 (2023), [arXiv:2301.07675 \[gr-qc\]](#).
- [9] G. Agazie *et al.* (NANOGrav), *Astrophys. J. Lett.* **951**, L8 (2023), [arXiv:2306.16213 \[astro-ph.HE\]](#).
- [10] J. Antoniadis *et al.* (EPTA), *Astron. Astrophys.* **678**, A50 (2023), [arXiv:2306.16214 \[astro-ph.HE\]](#).
- [11] D. J. Reardon *et al.*, *Astrophys. J. Lett.* **951**, L6 (2023), [arXiv:2306.16215 \[astro-ph.HE\]](#).
- [12] H. Xu *et al.*, *Res. Astron. Astrophys.* **23**, 075024 (2023), [arXiv:2306.16216 \[astro-ph.HE\]](#).
- [13] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Phys. Rev. D* **100**, 061101 (2019), [arXiv:1903.02886 \[gr-qc\]](#).
- [14] C. Perigois, C. Belczynski, T. Bulik, and T. Regimbau, *Phys. Rev. D* **103**, 043002 (2021), [arXiv:2008.04890 \[astro-ph.CO\]](#).
- [15] T. Regimbau, *Symmetry* **14**, 270 (2022).
- [16] S. Hild, S. Chelkowski, and A. Freise, (2008), [arXiv:0810.0604 \[gr-qc\]](#).
- [17] M. Punturo *et al.*, *Proceedings, 14th Workshop on Gravitational wave data analysis (GWDAAW-14): Rome, Italy, January 26-29, 2010*, *Class. Quant. Grav.* **27**, 194002 (2010).
- [18] S. Hild *et al.*, *Class. Quant. Grav.* **28**, 094013 (2011), [arXiv:1012.0908 \[gr-qc\]](#).
- [19] M. Maggiore *et al.*, *JCAP* **2020** (03), 050, [arXiv:1912.02622 \[astro-ph.CO\]](#).
- [20] D. Reitze *et al.*, *Bull. Am. Astron. Soc.* **51**, 035 (2019), [arXiv:1907.04833 \[astro-ph.IM\]](#).
- [21] M. Evans *et al.*, (2021), [arXiv:2109.09882 \[astro-ph.IM\]](#).
- [22] M. Evans *et al.*, (2023), [arXiv:2306.13745 \[astro-ph.IM\]](#).
- [23] I. Gupta *et al.*, (2023), [arXiv:2307.10421 \[gr-qc\]](#).
- [24] M. Branchesi *et al.*, *JCAP* **2023** (07), 068, [arXiv:2303.15923 \[gr-qc\]](#).
- [25] F. Iacovelli, E. Belgacem, M. Maggiore, M. Mancarella, and N. Muttoni, *JCAP* **2024** (10), 085, [arXiv:2408.14946 \[gr-qc\]](#).
- [26] C. Cutler and J. Harms, *Phys. Rev. D* **73**, 042001 (2006), [arXiv:gr-qc/0511092](#).
- [27] J. Harms, C. Mahrtdt, M. Otto, and M. Priess, *Phys. Rev. D* **77**, 123010 (2008), [arXiv:0803.0226 \[gr-qc\]](#).
- [28] T. Regimbau, M. Evans, N. Christensen, E. Katsavounidis, B. Sathyaprakash, and S. Vitale, *Phys. Rev. Lett.* **118**, 151105 (2017), [arXiv:1611.08943 \[astro-ph.CO\]](#).
- [29] Z. Pan and H. Yang, *Class. Quant. Grav.* **37**, 195020 (2020), [arXiv:1910.09637 \[astro-ph.CO\]](#).
- [30] S. Sachdev, T. Regimbau, and B. S. Sathyaprakash, *Phys. Rev. D* **102**, 024051 (2020), [arXiv:2002.05365 \[gr-qc\]](#).
- [31] A. Sharma and J. Harms, *Phys. Rev. D* **102**, 063009 (2020), [arXiv:2006.16116 \[gr-qc\]](#).
- [32] M. Lewicki and V. Vaskonen, *Eur. Phys. J. C* **83**, 168 (2023), [arXiv:2111.05847 \[astro-ph.CO\]](#).
- [33] C. Perigois, F. Santoliquido, Y. Bouffanais, U. N. Di Carlo, N. Giacobbo, S. Rastello, M. Mapelli, and T. Regimbau, *Phys. Rev. D* **105**, 103032 (2022), [arXiv:2112.01119 \[astro-ph.CO\]](#).
- [34] B. Zhou, L. Reali, E. Berti, M. alıřkan, C. Creque-Sarbinowski, M. Kamionkowski, and B. S. Sathyaprakash, (2022), [arXiv:2209.01221 \[gr-qc\]](#).
- [35] B. Zhou, L. Reali, E. Berti, M. alıřkan, C. Creque-Sarbinowski, M. Kamionkowski, and B. S. Sathyaprakash, *Phys. Rev. D* **108**, 064040 (2023), [arXiv:2209.01310 \[gr-qc\]](#).
- [36] H. Zhong, R. Ormiston, and V. Mandic, *Phys. Rev. D* **107**, 064048 (2023), [arXiv:2209.11877 \[gr-qc\]](#).
- [37] Z. Pan and H. Yang, *Phys. Rev. D* **107**, 123036 (2023), [arXiv:2301.04529 \[gr-qc\]](#).
- [38] H. Zhong, B. Zhou, L. Reali, E. Berti, and V. Mandic, *Phys. Rev. D* **110**, 064047 (2024), [arXiv:2406.10757 \[gr-qc\]](#).
- [39] M. Li, J. Yu, and Z. Pan, (2024), [arXiv:2403.01846 \[gr-qc\]](#).

- [40] E. Belgacem, F. Iacovelli, M. Maggiore, M. Mancarella, and N. Muttoni, in prep. (2024).
- [41] A. C. Jenkins and M. Sakellariadou, *Phys. Rev. D* **100**, 063508 (2019), [arXiv:1902.07719 \[astro-ph.CO\]](#).
- [42] L. Valbusa Dall’Armi, A. Nishizawa, A. Ricciardone, and S. Matarrese, *Phys. Rev. Lett.* **131**, 041401 (2023), [arXiv:2301.08205 \[astro-ph.CO\]](#).
- [43] R. Abbott *et al.* (KAGRA, VIRGO, LIGO Scientific), *Phys. Rev. X* **13**, 011048 (2023), [arXiv:2111.03634 \[astro-ph.HE\]](#).
- [44] N. Seto and A. Taruya, *Phys. Rev. D* **77**, 103001 (2008), [arXiv:0801.4185 \[astro-ph\]](#).
- [45] A. Renzini, *Mapping the Gravitational-Wave Background*, Ph.D. thesis, Imperial Coll., London (2020).
- [46] A. I. Renzini, B. Goncharov, A. C. Jenkins, and P. M. Meyers, *Galaxies* **10**, 34 (2022), [arXiv:2202.00178 \[gr-qc\]](#).
- [47] C. R. Contaldi, *Phys. Lett. B* **771**, 9 (2017), [arXiv:1609.08168 \[astro-ph.CO\]](#).
- [48] G. Cusin, C. Pitrou, and J.-P. Uzan, *Phys. Rev.* **D96**, 103019 (2017), [arXiv:1704.06184 \[astro-ph.CO\]](#).
- [49] G. Cusin, C. Pitrou, and J.-P. Uzan, *Phys. Rev.* **D97**, 123527 (2018), [arXiv:1711.11345 \[astro-ph.CO\]](#).
- [50] A. C. Jenkins and M. Sakellariadou, *Phys. Rev.* **D98**, 063509 (2018), [arXiv:1802.06046 \[astro-ph.CO\]](#).
- [51] G. Cusin, I. Dvorkin, C. Pitrou, and J.-P. Uzan, *Phys. Rev. Lett.* **120**, 231101 (2018), [arXiv:1803.03236 \[astro-ph.CO\]](#).
- [52] A. C. Jenkins, M. Sakellariadou, T. Regimbau, and E. Slezak, *Phys. Rev.* **D98**, 063501 (2018), [arXiv:1806.01718 \[astro-ph.CO\]](#).
- [53] A. C. Jenkins, R. O’Shaughnessy, M. Sakellariadou, and D. Wysocki, *Phys. Rev. Lett.* **122**, 111101 (2019), [arXiv:1810.13435 \[astro-ph.CO\]](#).
- [54] G. Cusin, I. Dvorkin, C. Pitrou, and J.-P. Uzan, *Mon. Not. Roy. Astron. Soc.* **493**, L1 (2020), [arXiv:1904.07757 \[astro-ph.CO\]](#).
- [55] G. Cusin, I. Dvorkin, C. Pitrou, and J.-P. Uzan, *Phys. Rev. D* **100**, 063004 (2019), [arXiv:1904.07797 \[astro-ph.CO\]](#).
- [56] A. C. Jenkins, J. D. Romano, and M. Sakellariadou, *Phys. Rev. D* **100**, 083501 (2019), [arXiv:1907.06642 \[astro-ph.CO\]](#).
- [57] D. Bertacca, A. Ricciardone, N. Bellomo, A. C. Jenkins, S. Matarrese, A. Raccanelli, T. Regimbau, and M. Sakellariadou, *Phys. Rev. D* **101**, 103513 (2020), [arXiv:1909.11627 \[astro-ph.CO\]](#).
- [58] C. Pitrou, G. Cusin, and J.-P. Uzan, *Phys. Rev. D* **101**, 081301 (2020), [arXiv:1910.04645 \[astro-ph.CO\]](#).
- [59] N. Bellomo, D. Bertacca, A. C. Jenkins, S. Matarrese, A. Raccanelli, T. Regimbau, A. Ricciardone, and M. Sakellariadou, *JCAP* **06** (06), 030, [arXiv:2110.15059 \[gr-qc\]](#).
- [60] G. Capurri, A. Lapi, C. Baccigalupi, L. Boco, G. Scelfo, and T. Ronconi, *JCAP* **2021** (11), 032, [arXiv:2103.12037 \[gr-qc\]](#).
- [61] S. H.-S. Alexander, M. E. Peskin, and M. M. Sheikh-Jabbari, *Phys. Rev. Lett.* **96**, 081301 (2006), [arXiv:hep-th/0403069](#).
- [62] S. G. Crowder, R. Namba, V. Mandic, S. Mukohyama, and M. Peloso, *Phys. Lett. B* **726**, 66 (2013), [arXiv:1212.4165 \[astro-ph.CO\]](#).
- [63] K. Yagi and H. Yang, *Phys. Rev. D* **97**, 104018 (2018), [arXiv:1712.00682 \[gr-qc\]](#).
- [64] V. Domcke, J. Garcia-Bellido, M. Peloso, M. Pieroni, A. Ricciardone, L. Sorbo, and G. Tasinato, *JCAP* **2020** (05), 028, [arXiv:1910.08052 \[astro-ph.CO\]](#).
- [65] K. Martinovic, C. Badger, M. Sakellariadou, and V. Mandic, *Phys. Rev. D* **104**, L081101 (2021), [arXiv:2103.06718 \[gr-qc\]](#).
- [66] T. Callister, L. Jenks, D. Holz, and N. Yunes, (2023), [arXiv:2312.12532 \[gr-qc\]](#).
- [67] N. M. J. Cruz, A. Malhotra, G. Tasinato, and I. Zavala, (2024), [arXiv:2406.04957 \[astro-ph.CO\]](#).
- [68] L. Blanchet, B. R. Iyer, C. M. Will, and A. G. Wiseman, *Class. Quant. Grav.* **13**, 575 (1996), [arXiv:gr-qc/9602024](#).
- [69] K. G. Arun, L. Blanchet, B. R. Iyer, and M. S. S. Qu-sailah, *Class. Quant. Grav.* **21**, 3771 (2004), [Erratum: *Class. Quant. Grav.* **22**, 3115 (2005)], [arXiv:gr-qc/0404085](#).
- [70] E. S. Phinney, (2001), [arXiv:astro-ph/0108028](#).
- [71] N. Kouvatsos, A. C. Jenkins, A. I. Renzini, J. D. Romano, and M. Sakellariadou, *Phys. Rev. D* **109**, 103535 (2024), [arXiv:2312.09110 \[astro-ph.CO\]](#).
- [72] C. Conneely, A. H. Jaffe, and C. M. F. Mingarelli, *Mon. Not. Roy. Astron. Soc.* **487**, 562 (2019), [arXiv:1808.05920 \[astro-ph.CO\]](#).
- [73] A. Akmal, V. R. Pandharipande, and D. G. Ravenhall, *Phys. Rev. C* **58**, 1804 (1998), [arXiv:nucl-th/9804027](#).