

Quantum geometric tensor determines the i.i.d. conversion rate in the resource theory of asymmetry for any compact Lie group

Koji Yamaguchi,¹ Yosuke Mitsuhashi,² and Hiroyasu Tajima^{1,3}

¹*Department of Communication Engineering and Informatics,
University of Electro-Communications, 1-5-1 Chofugaoka, Chofu, Tokyo, 182-8585, Japan*

²*Department of Basic Science, University of Tokyo,
3-8-1 Komaba, Meguro-ku, Tokyo 153-8902, Japan*

³*JST, PRESTO, 4-1-8 Honcho, Kawaguchi, Saitama, 332-0012, Japan*

Symmetry is one of the most significant foundational principles underlying nature. The resource theory of asymmetry (RTA) is a resource-theoretic framework for investigating asymmetry as a resource to break constraints imposed by symmetries. It has recently undergone significant developments, resulting in applications in a variety of research areas since symmetry and its breaking are ubiquitous in physics. Nevertheless, the resource conversion theory at the core of RTA remains incomplete. In the independent and identically distributed (i.i.d.) setup, where identical copies of a state are converted to identical copies of another state, conversion theory among pure states has been completed only for $U(1)$ group and finite groups. Here, we establish an i.i.d. conversion theory among any pure states in RTA for any continuous symmetry described by a compact Lie group, which includes the cases where multiple conserved quantities are involved. We show that the quantum geometric tensor is an asymmetry monotone that determines the optimal approximate asymptotic conversion rate. Our formulation achieves a unified understanding of conversion rates in prior studies for different symmetries. As a corollary of the formula, we also affirmatively prove the Marvian-Spekkens conjecture on reversible asymptotic convertibility in RTA, which has remained unproven for a decade.

I. INTRODUCTION

Symmetry lies at the core of our fundamental understanding of nature. It governs conservation laws and interactions, influencing diverse areas of modern physics. A new perspective on symmetry has been recently developed [1–6] in the context of quantum resource theories [7]. Quantum resource theory is a versatile framework for investigating a situation where freely implementable operations and preparable states are restricted. These limitations can often be circumvented with the help of states that cannot be freely prepared, which are regarded as resources. Building on the pioneering development of entanglement theory [8], the framework of quantum resource theories expanded to broader concepts, such as coherence [9–11], athermality [12, 13], and asymmetry [1–6]. The resource theory of asymmetry (RTA) [1–6], which was also called the resource theory of quantum reference frame in earlier studies [1–3], is a branch of quantum resource theories where the restriction is imposed by a symmetry group, and asymmetry that breaks the symmetry is treated as a resource.

Symmetry and its breaking are ubiquitous in nature, playing crucial roles in numerous physical phenomena. Consequently, studies in RTA have highlighted its applications in broad research topics, such as super-selection rules [1–3], acceleration of quantum operations [14], no-broadcasting theorems for quantum coherence [15, 16], quantum thermodynamics [17–20], creation of accurate clocks [19, 21], quantum error correction [22–28], gate implementation in quantum computation [25, 27, 29, 30], quantum measurements [5, 27, 31–33], and black hole evaporation [25, 27].

Despite the significant progress in RTA, its resource manipulation theory, a central issue in any resource theory, is still incomplete. In the independent and identically distributed (i.i.d.) setup where identical copies of a state are converted to identical copies of another state, the exact and approximate settings are investigated in RTA, where exact means no error is allowed in conversion, while approximate implies asymptotically vanishing error is allowed in conversion. For a discrete symmetry described by a finite group, a recent study [34] established both exact and approximate i.i.d. conversion theory for pure states, which generalizes the result on \mathbb{Z}_2 in [2]. For a continuous symmetry described by a Lie group, however, little is known; an approximate conversion theory is completed only for $U(1)$ in [19, 21] generalizing the prior results [2, 35, 36], and partial results on specific examples on restricted sets of pure states are obtained for other groups [2, 37]. This means that a comprehensive theory is still lacking for interesting cases involving multiple conserved quantities where non-commutativity is essential. Even in a special case where the asymptotic conversion is reversible, a definitive understanding remains elusive. A necessary and sufficient condition for conversion to be asymptotically reversible was conjectured by Marvian and Spekkens in [4, 6]. However, this Marvian-Spekkens conjecture had remained unproven for a decade. It is worth mentioning that the asymptotic conversion rate in RTA cannot be derived from the approach in general resource theory for reversible conversion [38] using the generalized Stein’s lemma [39–42]. This is because this framework does not work well for RTA since the regularized relative entropy vanishes in RTA [3], as emphasized in [38, 41]. As such, a unified understanding

of asymptotic conversion theory has not yet been established.

Establishing an asymptotic resource conversion theory is essentially inseparable from identifying an important resource measure. Entanglement entropy in entanglement theory and the Helmholtz free energy in quantum thermodynamics are of distinguished significance as they characterize the asymptotic conversion rates of respective resources [12, 43]. In RTA, various asymmetry measures are proposed, such as G -asymmetry [44], the relative entropy of G -freeness [3], purity of coherence [19], the quantum Fisher information [21], the quantum Fisher information matrix [45, 46], and the metric adjusted skew information [47–49] and its extension [50]. Among others, the quantum Fisher information plays a significant role in RTA for $U(1)$ group since the asymptotic conversion rate among pure states is determined by the ratio of the quantum Fisher information [21]. However, it remains elusive how to extend this result to other continuous symmetries.

In this paper, we establish the i.i.d. approximate asymptotic conversion theory among pure states in RTA for an arbitrary compact Lie group. The asymptotic conversion rate is shown to be determined by the quantum geometric tensor (QGT), a local geometric characteristic of rays in a Hilbert space. The QGT is a complex Hermitian tensor whose real symmetric part corresponds to the quantum Fisher information matrix, while the imaginary anti-symmetric part gives the Berry curvature [51, 52]. For a Lie group G with dimension $\dim G$, the QGT is a $\dim G \times \dim G$ Hermitian matrix, which is equal to a non-symmetrized covariance matrix for generators of symmetry transformations. Our formula shows that the asymptotic conversion rate is determined by a finite number of matrix inequalities among the QGTs, where each inequality corresponds to a connected component of a compact Lie group.

The proof of our formula on conversion rate consists of the converse and direct parts, i.e., the impossibility of achieving the conversion rate beyond the formula and the achievability of the optimal conversion rate. The converse part is a consequence of the fact that QGT and its asymptotic rate are valid asymmetry measures for pure states, proven by using the properties of Petz’s monotone metrics [53] and a technique analyzing its asymptotic behavior [50]. In the proof of the direct part, we adopt results on quantum local asymptotic normality (QLAN) [54–57] to extract the asymptotic characteristic of i.i.d. copies of a state, and construct quantum channels that asymptotically transform original states into the desired states with exponentially small failure probability. Our result shows that the QGT plays a distinguished role in RTA for an arbitrary compact Lie group that determines the asymptotic conversion rate.

Our formula is valid for any continuous symmetry described by a compact Lie group. To illustrate its wide applicability, we provide four examples. First, since the asymptotic conversion theory among pure states has been completed in prior research exclusively for the case of

$U(1)$ [21], we confirm that this result immediately follows from our formula. Second, we provide a necessary and sufficient condition for conversion to be reversible. As a consequence, we affirmatively prove the Marvian-Spekkens conjecture, which had been unsolved for the past decade until today. Third, we analyze examples for $U(1)$ and $SU(2)$ in which the asymptotic conversion is irreversible. The conversion rate for the example in $SU(2)$ is shown to be consistent with the prior study on a specific class of pure states in [2]. Fourth, we calculate the conversion rate for finite groups. Although a finite group is not a Lie group, it is shown that our formula is applicable by appending a trivial $U(1)$ phase. We find a divergence in conversion rate, reproducing the prior result on the approximate conversion rate in RTA for finite groups in [34]. Based on our proof of the direct part, we also provide an intuitive explanation of the reason why the approximate conversion rate can diverge in RTA for finite groups.

The rest of this paper is organized as follows: We first review the framework of RTA and the definition of the QGT in Sec. II. The main result of this paper, i.e., the approximate asymptotic conversion rate among pure states in RTA for an arbitrary Lie group, is presented in Sec. III. In Sec. IV, we show that the QGT is asymmetry monotone for pure states in one-shot and asymptotic conversion. As a consequence, we prove an inequality providing a bound for the asymptotic conversion rate, i.e., the converse part of the main theorem. In Sec. V, we show the optimality of this bound, i.e., the direct part of the main theorem. In Sec. VI, we apply our formula for RTA for several setups, providing a unified understanding of prior studies on conversion rates in RTA for different groups [2, 4, 6, 21, 34, 37], and proves the Marvian-Spekkens conjecture [4, 6].

II. SETUP AND NOTATIONS

A. The resource theory of asymmetry (RTA)

In this paper, we follow the standard setup of RTA [1–6]. We study a symmetry described by a group G that is realized by a *projective unitary representation* U , which maps $g \in G$ to a unitary operator $U(g)$ on a quantum system. From the consistency of successive application of symmetry transformations, U must satisfy $U(g_1)U(g_2) = \omega(g_1, g_2)U(g_1g_2)$ for any $g_1, g_2 \in G$, where ω is a complex-valued function such that $|\omega(g_1, g_2)| = 1$. In a special case where $\omega(g_1, g_2) = 1$ for all $g_1, g_2 \in G$, U is called a *non-projective unitary representation* or a *unitary representation* for short.

Just like in other resource theories, including the entanglement theory, RTA is defined by specifying the *free states* and *free operations* that can be freely prepared and implemented. In RTA, free states are symmetric states, which are invariant under any symmetry transformations. That is, a state ρ is G -symmetric iff

$U(g)\rho U(g)^\dagger = \rho$ for all $g \in G$. Free operations are G -covariant channels. Here, a channel \mathcal{E} is G -covariant iff it satisfies $\mathcal{E} \circ \mathcal{U}_g = \mathcal{U}'_g \circ \mathcal{E}$ for all $g \in G$, where \mathcal{U}_g and \mathcal{U}'_g are defined by $\mathcal{U}_g(\cdot) := U(g)(\cdot)U(g)^\dagger$ and $\mathcal{U}'_g(\cdot) := U'(g)(\cdot)U'(g)^\dagger$ using projective unitary representations U, U' of G on the input and output systems. Throughout this paper, we use the prime symbol to represent a quantity related to the output system.

As a simple yet important characterization of convertibility among states, the symmetry subgroup is known, which is the set of group elements that leave the state invariant, i.e., $\text{Sym}_G(\rho) := \{g \in G \mid U(g)\rho U(g)^\dagger = \rho\}$ for the input state ρ , and similarly for the output state σ . It is shown in [4] that if a state ρ can be convertible to a state σ via some G -covariant operation without error, the symmetry subgroups must satisfy $\text{Sym}_G(\rho) \subset \text{Sym}_G(\sigma)$.

We analyze the i.i.d. setup in RTA where identical copies of a state are converted to identical copies of another state. In this setup, we use the tensor product of a representation, which corresponds to the conservation of global quantities such as total particle number in the $U(1)$ case and total spins in the $SU(2)$ case. Concretely, we say that a channel \mathcal{E} from N copies of input state to M copies of output state is G -covariant if \mathcal{E} satisfies $\mathcal{E} \circ \mathcal{U}_g^{\otimes N} = \mathcal{U}'_g^{\otimes M} \circ \mathcal{E}$ for all $g \in G$, where $\mathcal{U}_g(\cdot) := U(g)(\cdot)U(g)^\dagger$ and $\mathcal{U}'_g(\cdot) := U'(g)(\cdot)U'(g)^\dagger$.

We finally define the asymptotic conversion rate. We say that a state ρ is approximately asymptotically convertible to another state σ with a conversion rate r iff there exists a sequence $\{\mathcal{E}_N\}_N$ of G -covariant channels such that $\lim_{N \rightarrow \infty} T(\mathcal{E}_N(\rho^{\otimes N}), \sigma^{\otimes \lfloor rN \rfloor}) = 0$ with the trace distance T , which has operational significance in state distinguishability [58, 59]. We denote this conversion by

$$\{\rho^{\otimes N}\}_N \xrightarrow{G\text{-cov.}} \{\sigma^{\otimes \lfloor rN \rfloor}\}_N, \quad (1)$$

and we define the approximate asymptotic conversion rate $R(\rho \rightarrow \sigma)$ by the supremum of achievable conversion rates r . In the following, we simply call $R(\rho \rightarrow \sigma)$ conversion rate. In this paper, we establish a formula for calculating this conversion rate among pure states, which is applicable to any compact Lie group G .

B. Quantum geometric tensor (QGT)

Here, we introduce the QGT [51, 52]. For a given parametric family of pure states $|\xi(\boldsymbol{\lambda})\rangle$ with $\boldsymbol{\lambda} \in \mathbb{R}^m$, QGT at $|\psi\rangle := |\xi(\mathbf{0})\rangle$ is an $m \times m$ Hermitian matrix $\mathcal{Q}^{\psi, \xi}$ whose matrix elements are defined by

$$\mathcal{Q}_{\mu\nu}^{\psi, \xi} := \langle \partial_\mu \psi | (I - |\psi\rangle\langle\psi|) | \partial_\nu \psi \rangle, \quad (2)$$

where $|\partial_\mu \psi\rangle := \frac{\partial}{\partial \lambda^\mu} |\xi(\boldsymbol{\lambda})\rangle|_{\boldsymbol{\lambda}=\mathbf{0}}$. The real and symmetric part of QGT is the quantum Fisher information matrix, while the imaginary anti-symmetric part corresponds to the Berry curvature [51, 52, 60, 61].

In RTA for a Lie group, we can define QGT using a natural parametric family of a pure state defined with a projective unitary representation. Let G be a Lie group, and $\dim G$ denote its dimension as a smooth manifold. Elements in the neighborhood of the identity $e \in G$ can be parametrized as $g(\boldsymbol{\lambda}) = e^{i \sum_{\mu=1}^{\dim G} \lambda^\mu A_\mu}$ by using a basis $\{A_\mu\}_{\mu=1}^{\dim G}$ of the Lie algebra \mathfrak{g} , where the convention in physics for the definition of Lie algebra is adopted, which differs from that in mathematics by a factor of the imaginary unit. Given a projective unitary representation U , we define $|\xi(\boldsymbol{\lambda})\rangle := U(g(\boldsymbol{\lambda}))|\psi\rangle$ for a pure state $|\psi\rangle$. For simplicity, we assume that the map U is differentiable, which follows from continuity in the case of unitary representations. We treat the cases where U is continuous but not differentiable in Appendix A 1. For a given U , we introduce Hermitian operators

$$X_\mu := -i \frac{\partial}{\partial \lambda^\mu} U(g(\boldsymbol{\lambda})) \Big|_{\boldsymbol{\lambda}=\mathbf{0}} \quad (3)$$

for $\mu = 1, \dots, \dim G$, where this corresponds to $L(A_\mu)$ with the Lie algebra representation $L(A) := -i \frac{d}{dt} U(e^{itA})|_{t=0}$. Then the QGT for $|\xi(\boldsymbol{\lambda})\rangle$ at $|\psi\rangle$, which we simply denote by \mathcal{Q}^ψ , is equal to the non-symmetrized covariance matrix for $\mathbf{X} := \{X_\mu\}_{\mu=1}^{\dim G}$:

$$\begin{aligned} \mathcal{Q}_{\mu\nu}^\psi &= \langle \psi | X_\mu (I - |\psi\rangle\langle\psi|) X_\nu | \psi \rangle \\ &= \langle \psi | X_\mu X_\nu | \psi \rangle - \langle \psi | X_\mu | \psi \rangle \langle \psi | X_\nu | \psi \rangle. \end{aligned} \quad (4)$$

Note that in this case, the size of a QGT is determined only by the dimension of the Lie group, $\dim G$, independent of the dimensions of Hilbert spaces or the representations of the group. Since QGTs are Hermitian matrices, we can introduce a partial order $\mathcal{Q}^\psi \geq \mathcal{Q}^\phi$, which means that $\mathcal{Q}^\psi - \mathcal{Q}^\phi$ is positive semi-definite. We remark that this ordering is independent of the parametrization of the Lie group G since both \mathcal{Q}^ψ and \mathcal{Q}^ϕ transform as tensors under a coordinate transformation on the group.

III. MAIN RESULT

The main theorem of this paper is the following formula for the asymptotic conversion rate:

Theorem 1. *Let U and U' be projective unitary representations of a compact Lie group G on finite-dimensional Hilbert spaces \mathcal{H} and \mathcal{H}' . The conversion rate from a pure state $|\psi\rangle \in \mathcal{H}$ to another pure state $|\phi\rangle \in \mathcal{H}'$ is calculated as follows: (I) When $\text{Sym}_G(\psi) \not\subset \text{Sym}_G(\phi)$, $R(\psi \rightarrow \phi) = 0$. (II) When $\text{Sym}_G(\psi) \subset \text{Sym}_G(\phi)$,*

$$R(\psi \rightarrow \phi) = \sup\{r \geq 0 \mid \forall i, \mathcal{Q}^{U_{g_i}(\psi)} \geq r \mathcal{Q}^{U'_{g_i}(\phi)}\}, \quad (5)$$

where $g_i \in G_i$ is a representative element of each connected component G_i of G . In particular, if G is connected, then

$$R(\psi \rightarrow \phi) = \sup\{r \geq 0 \mid \mathcal{Q}^\psi \geq r \mathcal{Q}^\phi\}. \quad (6)$$

Since a compact Lie group has only a finite number of connected components (see, e.g., [62]), the conversion rate in Theorem 1 is expressed in terms of a finite number of matrix inequalities. In particular, Eq. (5) is inde-

pendent of the choice of the elements g_i , and thus can be easily evaluated. Although Theorem 1 is of practical significance, the following formula is simple and theoretically useful, where all group elements are treated equally:

$$R(\psi \rightarrow \phi) = \begin{cases} \sup\{r \geq 0 \mid \forall g \in G, \mathcal{Q}^{\mathcal{U}_g(\psi)} \geq r \mathcal{Q}^{\mathcal{U}'_g(\phi)}\} & (\text{if } \text{Sym}_G(\psi) \subset \text{Sym}_G(\phi)) \\ 0 & (\text{otherwise}) \end{cases} \quad (7)$$

The equivalence of Eq. (5) in Theorem 1 to Eq. (7) follows from the fact that if the matrix inequality holds at a point in a connected component of the group, then it also holds at any other point in the connected component. This is because the QGTs at different group elements in a connected component are interrelated by a congruence transformation independent of representation, as detailed in Appendix A 2. We note that the same argument for quantum Fisher information matrices instead of QGT can be found in [46].

In the following, we prove Eq. (7) under the assumption that U and U' are (non-projective) unitary representations of a compact Lie group G . The extension of the formula to any projective unitary representations is provided in Appendix A 3 by using the method in [34] that relates the conversion rate for projective unitary representations to (non-projective) unitary representations.

We remark that for any states ρ and σ , Proposition 5 in [4] shows that $\text{Sym}_G(\rho) \not\subset \text{Sym}_G(\sigma)$ implies that the conversion from ρ to σ without error is impossible via a G -covariant channel. As its generalization to approximate asymptotic conversion, we prove Eq. (8) in Appendix A 4.

IV. QGT AS ASYMMETRY MONOTONE

In this section, we show that QGT is an asymmetry monotone both in one-shot and asymptotic conversions. As a consequence, we find that QGT provides a bound on the conversion rate.

A. Monotonicity of QGT in one-shot conversion

We here prove the monotonicity of the QGT under conversion without error using a G -covariant channel. For this purpose, let us first relate the QGT to monotone metrics [53, 63], which are Riemannian metrics on state space contracting under information processing. The pioneering research on monotone metrics initiated by Morozova and Chentsov [63] was later completed by Petz [53], proving that monotone metrics are in a one-to-one correspondence with operator monotone functions. Here, a function $f : [0, \infty) \rightarrow [0, \infty)$ is called an operator mono-

tone iff $0 \leq A \leq B$ implies $f(A) \leq f(B)$. Petz's monotone metric associated with an operator monotone function f is defined by

$$\langle A, B \rangle_{f,\rho} := \text{Tr} \left(A^\dagger m_f^{-1}(\mathcal{R}_\rho, \mathcal{L}_\rho)(B) \right), \quad (9)$$

where $m_f(x, y) := yf(x/y)$, $\mathcal{R}_\rho(A) := A\rho$ and $\mathcal{L}_\rho(A) := \rho A$. Here, when ρ is not invertible, following the argument in [64], $m_f^{-1}(\mathcal{R}_\rho, \mathcal{L}_\rho)$ denotes the Moore-Penrose inverse, such that

$$m_f^{-1}(\mathcal{R}_\rho, \mathcal{L}_\rho)(B) := \sum_{k,l; m_f(p_k, p_l) > 0} \frac{\langle l|B|k \rangle}{m_f(p_k, p_l)} |l \rangle \langle k|, \quad (10)$$

where we used the eigenvalue decomposition of a state $\rho = \sum_k p_k |k \rangle \langle k|$. For a linear operator A , we denote $\|A\|_{f,\rho} := \sqrt{\langle A, A \rangle_{f,\rho}}$.

An additional symmetry condition $f(t) = tf(t^{-1})$ is often imposed to the monotone function, which is equivalent to $\langle A, B \rangle_{f,\rho} = \langle B^\dagger, A^\dagger \rangle_{f,\rho}$ for any linear operators A and B . A famous example is $f_{\text{SLD}}(x) := (1+x)/2$ associated with the symmetric logarithmic derivative (SLD) metric, which plays a central role in quantum estimation theory. However, such a symmetric monotone metric does not contain the information of the anti-symmetric part of the QGT. Therefore, we do not impose the symmetry condition.

In order to investigate the properties of QGTs, we consider a family of operator monotone functions given by

$$f_q(x) := (1-q) + qx, \quad q \in (0, 1). \quad (11)$$

Note that this family contains f_{SLD} as a special case for $q = 1/2$. Let us show that the QGT can be obtained as a limit of Petz's monotone metric for f_q . Since $m_{f_q}(p_k, p_l) = (1-q)p_l + qp_k$, for any pure state ψ and any linear operator O , it holds

$$\|i[\psi, O]\|_{f_q, \psi}^2 = \frac{1}{1-q} V(\psi, O) + \frac{1}{q} V(\psi, O)^*, \quad (12)$$

where we have defined a generalized variance V by

$$V(\psi, O) := \langle \psi | O(I - \psi) O^\dagger | \psi \rangle \quad (13)$$

for a pure state ψ and a linear operator O . Here, following a standard convention, we described a pure state and a unit vector by the same Greek letter, i.e., $\psi = |\psi\rangle\langle\psi|$. Note that when O is Hermitian, V is equivalent to the ordinary variance.

When O is given by $O := \gamma^\dagger \mathbf{X} := \sum_{\mu=1}^{\dim G} \gamma^{*\mu} X_\mu$ for $\gamma \in \mathbb{C}^{\dim G}$ and $\mathbf{X} = \{X_\mu\}_{\mu=1}^{\dim G}$ defined in Eq. (3), we get

$$\|i[\psi, O]\|_{f_q, \psi}^2 = \gamma^\dagger \left(\frac{1}{1-q} \mathcal{Q}^\psi + \frac{1}{q} (\mathcal{Q}^\psi)^* \right) \gamma. \quad (14)$$

Therefore, we find

$$\lim_{q \rightarrow 1^-} f_q(0) \|i[\psi, O]\|_{f_q, \psi}^2 = \gamma^\dagger \mathcal{Q}^\psi \gamma, \quad (15)$$

where we have used $f_q(0) = 1 - q > 0$.

We stress again that the operator monotone function satisfying the symmetry condition is not suitable for investigating the monotonicity of QGT. Indeed, for operator monotone functions $f^{(s)}$ satisfying $f^{(s)}(0) > 0$ and the symmetry condition $f^{(s)}(t) = t f^{(s)}(t^{-1})$, from the definition in Eq. (9), we find

$$\frac{f^{(s)}(0)}{2} \|i[\psi, O]\|_{f^{(s)}, \psi}^2 = \gamma^\dagger \left(\frac{1}{2} (\mathcal{Q}^\psi + (\mathcal{Q}^\psi)^*) \right) \gamma, \quad (16)$$

for $O := \gamma^\dagger \mathbf{X}$, meaning that it captures only the symmetric part of \mathcal{Q}^ψ .

Note that the quantity $\frac{f^{(s)}(0)}{2} \|i[\rho, H]\|_{f^{(s)}, \rho}^2$ for a Hermitian operator H is called the metric adjusted skew information [47]. It is known that the metric adjusted skew information for a pure state ψ is equal to the variance, i.e.,

$$\frac{f^{(s)}(0)}{2} \|i[\psi, H]\|_{f^{(s)}, \psi}^2 = V(\psi, H). \quad (17)$$

This equation can be viewed as a special case of Eq. (16), where γ is a real vector.

The metric $\langle \cdot, \cdot \rangle_{f_q, \rho}$ is monotonic under a quantum channel in the following sense:

Lemma 2. For $f_q(x) := (1-q) + qx$ with $q \in (0, 1)$, any state ρ , any linear operator O , and any quantum channel \mathcal{E} , it holds

$$\|i[\rho, O]\|_{f_q, \rho}^2 \geq \|\mathcal{E}(i[\rho, O])\|_{f_q, \mathcal{E}(\rho)}^2 \quad (18)$$

if there is an operator O' satisfying $\mathcal{E}(i[\rho, O]) = i[\mathcal{E}(\rho), O']$.

In literature, the proof of the monotonicity of monotone metric is sometimes provided under the assumption that ρ is invertible. Lemma 2 is a special case of Theorem 6.1 in [65], which is valid not only for invertible states but also non-invertible states. The details of the proof of Lemma 2 and its relation to Theorem 6.1 in [65] can be found in [66].

The monotonicity of the QGT in RTA is proven by using Lemma 2. Suppose that a state ρ is convertible without error to another state σ via a G -covariant channel \mathcal{E} , i.e., $\mathcal{E}(\rho) = \sigma$. Since the G -covariance of the channel implies $\mathcal{E}(\mathcal{U}_g(\rho)) = \mathcal{U}'_g(\sigma)$, its derivative with respect to the parameter λ for $g \in G$ yields $\mathcal{E}(i[\rho, X_\mu]) = i[\sigma, X'_\mu]$, where X_μ is given in Eq. (3), and X'_μ is defined similarly by differentiating the projective unitary representation U' of G on the output system instead of U .

By using the linearity of \mathcal{E} , we get $\mathcal{E}(i[\rho, O]) = i[\mathcal{E}(\rho), O']$ for $O := \gamma^\dagger \mathbf{X}$ and $O' := \gamma^\dagger \mathbf{X}'$ for any $\gamma \in \mathbb{C}^{\dim G}$. The monotonicity in Lemma 2 implies

$$\|i[\rho, O]\|_{f_q, \rho}^2 \geq \|i[\sigma, O']\|_{f_q, \sigma}^2. \quad (19)$$

Let us now apply this inequality to the case where ρ and σ are pure states, which we denote by ψ and ϕ , respectively. Multiplying Eq. (19) by $f_q(0) = 1 - q > 0$ and taking the limit of $q \rightarrow 1^-$, Eq. (15) implies, $\gamma^\dagger \mathcal{Q}^\psi \gamma \geq \gamma^\dagger \mathcal{Q}^\phi \gamma$ for any $\gamma \in \mathbb{C}^{\dim G}$, or equivalently,

$$\mathcal{Q}^\psi \geq \mathcal{Q}^\phi, \quad (20)$$

completing the proof of the monotonicity of QGT under exact conversion among pure states in RTA.

B. Asymptotic discontinuity of QGT

When the projective unitary representation of G on a single system is given by U , the representation of G on N i.i.d. copies of the system is given by $U^{\otimes N}$. Therefore, for an i.i.d. pure state $\psi^{\otimes N}$, the QGT is additive, i.e., $\mathcal{Q}^{\psi^{\otimes N}} = N \mathcal{Q}^\psi$ for any positive integer N . When $\{\psi^{\otimes N}\}_N \xrightarrow{G\text{-cov.}} \{\phi^{\otimes \lfloor rN \rfloor}\}_N$, combined with the monotonicity of QGT in Eq. (20), one would expect that the asymptotic rates of the QGTs for these sequences, i.e., $\lim_{N \rightarrow \infty} \frac{1}{N} \mathcal{Q}^{\psi^{\otimes N}}$ and $\lim_{N \rightarrow \infty} \frac{1}{N} \mathcal{Q}^{\phi^{\otimes \lfloor rN \rfloor}}$, are monotonically decreasing, implying that

$$\mathcal{Q}^\psi \geq r \mathcal{Q}^\phi. \quad (21)$$

Although this inequality is shown to be correct in the following subsection, its proof requires more careful analysis. This necessity arises from the asymptotic discontinuity [67, 68] of the QGT, i.e., the property that the QGT can drastically change due to a small error in conversion.

To gain a deeper understanding, let us consider the simplest case of $G = U(1)$, where its non-trivial representation is given by e^{iHt} with a Hermitian operator H . For $G = U(1)$, the QGT is equal to the variance of the Hamiltonian $V(\psi, H)$ and its asymptotic discontinuity was pointed out in [3] and further analyzed in [19, 21, 50]. To examine the discontinuity, let H_N be the Hamiltonian for N i.i.d. copies of the system. Since the operator norm of H_N grows linearly in N and the variance is quadratic in the Hamiltonian H_N , the variance can change $O(\epsilon_N N^2)$ when there is an error ϵ_N in states in the conversion. Therefore, the change in the variance

per copy is of the order of $O(\epsilon_N N)$, which may not vanish as $N \rightarrow \infty$ even when $\lim_{N \rightarrow \infty} \epsilon_N = 0$ [3, 19, 21]. Since the QGT for a generic Lie group has the same asymptotic discontinuity, a more detailed analysis is required for the proof of Eq. (21).

C. Monotonicity of QGT in asymptotic conversion

To show the monotonicity of the QGT in asymptotic conversion, we investigate the asymptotic behavior of Petz's monotone metrics near i.i.d. pure states. For this purpose, let us first review the results in [50]. For an arbitrary Hermitian operator H we denote its i.i.d. extension by H_N , defined as $H_N := \sum_{n=1}^N I^{\otimes n-1} \otimes H \otimes I^{\otimes N-n}$. For a sequence of states $\{\sigma_N\}_N$ such that $\limsup_{N \rightarrow \infty} T(\sigma_N, \phi^{\otimes N}) \leq \epsilon$ for $\epsilon > 0$, and an operator monotone function $f^{(s)}$ satisfying $f^{(s)}(0) > 0$ and the symmetry condition $f^{(s)}(t) = t f^{(s)}(t^{-1})$, Lemma 3 in [50] shows that under a certain regularity condition, if $\epsilon > 0$ is sufficiently small, it holds

$$\frac{f^{(s)}(0)}{2} \|\mathfrak{i}[\sigma_N, H_N]\|_{f^{(s)}, \sigma_N}^2 \geq NV(\phi, H) - Nh(\epsilon) + o(N) \quad (22)$$

for all sufficiently large N , where h is a real-valued function independent of N , satisfying $\lim_{\epsilon \rightarrow 0} h(\epsilon) = 0$. The central limit theorem provides an intuitive explanation of this lemma. In the i.i.d. setting, the probability distribution of $\phi^{\otimes N}$ with respect to the eigenbasis of H_N approaches to the normal distribution after properly redefining the random variable. It turns out that the left-hand side of Eq. (22) is approximated from below by the variance for a probability distribution modified due to the error between σ_N and $\phi^{\otimes N}$. In general, modifying the distribution at values far from the mean can result in a considerable change in the variance. However, since the normal distribution has an exponentially small tail, it is impossible to significantly reduce the variance. In other words, the metric adjusted skew information approximately takes a local minimum around the i.i.d. pure states [50], expressed by Eq. (22). Note that although the definition of convergence in the state conversion in [50] is slightly different from ours, both yield the equivalent convertibility condition, as shown in Appendix B 1.

To prove the monotonicity of QGT under approximate asymptotic conversion, we show a variation of Lemma 3 in [50]. Let us first introduce several notations. When $\{\psi^{\otimes N}\}_N \xrightarrow{G\text{-cov.}} \{\phi^{\otimes \lfloor rN \rfloor}\}_N$, there exists a sequence of quantum channels $\{\mathcal{E}_N\}_N$ such that $\lim_{N \rightarrow \infty} T(\mathcal{E}_N(\psi^{\otimes N}), \phi^{\otimes \lfloor rN \rfloor}) = 0$. For notational convenience, we define $M(N) := \lfloor rN \rfloor$, $\sigma_{M(N)} := \mathcal{E}_N(\psi^{\otimes N})$ and $\epsilon_{M(N)} := T(\sigma_{M(N)}, \phi^{\otimes M(N)})$. In the following, we omit the dependence of $M(N)$ on N and simply write M . Since the G -covariance of \mathcal{E}_N implies $\mathcal{E}_N(\mathcal{U}_g^{\otimes N}(\psi^{\otimes N})) = \mathcal{U}_g^{\otimes M}(\sigma_M)$, by repeating the argument presented above Eq. (19), we get $\mathcal{E}_N(\mathfrak{i}[\psi^{\otimes N}, O_N]) = \mathfrak{i}[\sigma_M, O'_M]$. Here, O_N

and O'_M are defined by $O_N := \sum_{n=1}^N I^{\otimes n-1} \otimes O \otimes I^{\otimes N-n}$ and $O'_M := \sum_{n=1}^M I^{\otimes n-1} \otimes O' \otimes I^{\otimes M-n}$ for $O := \gamma^\dagger \mathbf{X}$ and $O' := \gamma^\dagger \mathbf{X}'$. Applying Eq. (19) for $\psi^{\otimes N}$ and σ_M instead of ρ and σ , we find

$$\|\mathfrak{i}[\psi^{\otimes N}, O_N]\|_{f_q, \psi^{\otimes N}}^2 \geq \|\mathfrak{i}[\sigma_M, O'_M]\|_{f_q, \sigma_M}^2. \quad (23)$$

Note that the additivity of the monotone metric in the i.i.d. setting implies that the left-hand side is equal to $N \|\mathfrak{i}[\psi, O]\|_{f_q, \psi}^2$. To prove the monotonicity, we need to relate the right-hand side of this inequality to the QGT for $(\mathcal{U}'_g(\phi))^{\otimes M}$.

By modifying Eq. (22) to make it applicable to an arbitrary linear operator O instead of Hermitian operator H without assuming f is symmetric, we prove the following lemma, which yields a lower bound on the asymptotic behavior of the right-hand side of Eq. (23):

Lemma 3. *Let f be an arbitrary operator monotone function such that $f(0) > 0$ and $f(0) = \lim_{\epsilon \rightarrow 0^+} f(\epsilon)$. For a linear operator O on a finite-dimensional Hilbert space, we denote its i.i.d. extension by O_N . Let $\{\sigma_N\}_N$ be an arbitrary sequence of states such that $\lim_{N \rightarrow \infty} T(\sigma_N, \phi^{\otimes N}) = 0$ for a pure state ϕ . Then there exists a real-valued function h independent of N satisfying $\lim_{\epsilon \rightarrow 0} h(\epsilon) = 0$ such that for any sufficiently small parameter $\epsilon > 0$, it holds*

$$f(0) \|\mathfrak{i}[\sigma_N, O_N]\|_{f, \sigma_N}^2 \geq NV(\phi, O) - Nh(\epsilon) + o(N) \quad (24)$$

for all sufficiently large N .

Notice that the normalization of the left-hand sides of Eqs. (22) and (24) differs factor by 2. This difference also appeared in Eqs. (16) and (15), where the right-hand side of the former is symmetrized while that of the latter is not symmetrized.

Lemma 3 is proven by carefully modifying the proof of Lemma 3 in [50]. Since the proof is involved, the details are presented in Appendix B 2.

By using Lemma 3, Eq. (23) implies

$$f_q(0) \|\mathfrak{i}[\psi, O]\|_{f_q, \psi}^2 \geq \frac{M}{N} \gamma^\dagger \mathcal{Q}^\phi \gamma - \frac{M}{N} h(\epsilon) + \frac{1}{N} o(M), \quad (25)$$

where we have used $f_q(0) > 0$. Since $M = \lfloor rN \rfloor$, in the limit of $N \rightarrow \infty$, we get

$$f_q(0) \|\mathfrak{i}[\psi, O]\|_{f_q, \psi}^2 \geq r \gamma^\dagger \mathcal{Q}^\phi \gamma - r h(\epsilon). \quad (26)$$

Since this inequality holds for any sufficiently small $\epsilon > 0$, we get $f_q(0) \|\mathfrak{i}[\psi, O]\|_{f_q, \psi}^2 \geq r \gamma^\dagger \mathcal{Q}^\phi \gamma$. In the limit of $q \rightarrow 1^-$, from Eq. (15), we finally get $\gamma^\dagger \mathcal{Q}^\psi \gamma \geq r \gamma^\dagger \mathcal{Q}^\phi \gamma$ for any $\gamma \in \mathbb{C}^{\dim G}$, i.e., Eq. (21). Since $\{\psi^{\otimes N}\}_N \xrightarrow{G\text{-cov.}} \{\phi^{\otimes \lfloor rN \rfloor}\}_N$ is equivalent to $\{\mathcal{U}_g(\psi)^{\otimes N}\}_N \xrightarrow{G\text{-cov.}} \{\mathcal{U}'_g(\phi)^{\otimes \lfloor rN \rfloor}\}_N$, by repeating the argument so far for $\mathcal{U}_g(\psi)$ and $\mathcal{U}'_g(\phi)$ instead of ψ and ϕ , we get

$$\forall g \in G, \quad \mathcal{Q}^{\mathcal{U}_g(\psi)} \geq r \mathcal{Q}^{\mathcal{U}'_g(\phi)}, \quad (27)$$

which completes the proof of the monotonicity of QGT in asymptotic conversion.

In summary, we have proven that $\{\psi^{\otimes N}\}_N \xrightarrow{G\text{-cov.}} \{\phi^{\otimes \lfloor rN \rfloor}\}_N$ implies $\mathcal{Q}^{\mathcal{U}_g(\psi)} \geq r\mathcal{Q}^{\mathcal{U}'_g(\phi)}$ for all $g \in G$, which establishes the following upper bound using QGTs on the conversion rate:

Proposition 4 (Converse part). *Let U, U' be projective unitary representations of a Lie group G on finite-dimensional Hilbert spaces \mathcal{H} and \mathcal{H}' . For any pure states $|\psi\rangle \in \mathcal{H}$ and $|\phi\rangle \in \mathcal{H}'$, it holds*

$$R(\psi \rightarrow \phi) \leq \sup\{r \geq 0 \mid \forall g \in G, \mathcal{Q}^{\mathcal{U}_g(\psi)} \geq r\mathcal{Q}^{\mathcal{U}'_g(\phi)}\}. \quad (28)$$

Proposition 4 serves as the converse part of Eq. (7), i.e., the inequality showing the impossibility of achieving the rate beyond the right-hand side of Eq. (7). In the following section, we show that this rate is the optimal achievable rate.

We remark that Proposition 4 is valid even for non-compact Lie groups as long as the representation space is finite-dimensional because the proof only depends on the local properties of a state in the state space. Note also that Proposition 4 is applicable regardless of whether $\text{Sym}_G(\psi) \subset \text{Sym}_G(\phi)$ holds.

V. QGT DETERMINES ASYMPTOTIC CONVERSION RATE

In the previous section, we have proven an upper bound on the conversion rate, given in Eq. (28), which follows from the monotonicity of QGT in asymptotic conversion. In this section, we show that this bound is optimal by proving that if $r > 0$ satisfies $\mathcal{Q}^{\mathcal{U}_g(\psi)} \geq r\mathcal{Q}^{\mathcal{U}'_g(\phi)}$ for all $g \in G$, then $\{\psi^{\otimes N}\}_N \xrightarrow{G\text{-cov.}} \{\phi^{\otimes \lfloor (r-\delta)N \rfloor}\}_N$ holds for arbitrary $\delta \in (0, r)$, which establishes the opposite inequality of Eq. (28):

Proposition 5 (Direct part). *Let U, U' be (non-projective) unitary representations of a compact Lie group G on finite-dimensional Hilbert spaces \mathcal{H} and \mathcal{H}' . For any pure states $|\psi\rangle \in \mathcal{H}$ and $|\phi\rangle \in \mathcal{H}'$ such that $\text{Sym}_G(\psi) \subset \text{Sym}_G(\phi)$, it holds*

$$R(\psi \rightarrow \phi) \geq \sup\{r \geq 0 \mid \forall g \in G, \mathcal{Q}^{\mathcal{U}_g(\psi)} \geq r\mathcal{Q}^{\mathcal{U}'_g(\phi)}\}. \quad (29)$$

Combining Proposition 5 with Proposition 4, our main result in Eq. (7) is proven for (non-projective) unitary representations, meaning that QGT determines the asymptotic conversion rate. As mentioned in Sec. III, this result can be extended to projective unitary representations by using the method in [34], which is detailed in Appendix A3.

In the proof of Proposition 5, the results in QLAN on pure-state unitary models [56, 57] play a crucial role.

This can be intuitively understood through the following lemma, which provides an alternative yet equivalent characterization of the convertibility in RTA using quantum channels that are not assumed to be G -covariant:

Lemma 6. *Let U and U' be projective unitary representations of a compact Lie group G on finite-dimensional Hilbert spaces \mathcal{H} and \mathcal{H}' . Define $\mathcal{U}_g(\cdot) := U(g)(\cdot)U(g)^\dagger$ and $\mathcal{U}'_g(\cdot) := U'(g)(\cdot)U'(g)^\dagger$. Let ρ and σ be arbitrary states on \mathcal{H} and \mathcal{H}' , respectively. For any $\epsilon \geq 0$, the following two statements are equivalent:*

(i) *There exists a quantum channel \mathcal{E} such that $T(\mathcal{E}(\mathcal{U}_g(\rho)), \mathcal{U}'_g(\sigma)) \leq \epsilon$ for any $g \in G$.*

(ii) *There exists a G -covariant channel \mathcal{E}' such that $T(\mathcal{E}'(\rho), \sigma) \leq \epsilon$.*

For $\epsilon = 0$, i.e., for the conversion without error, the above is shown as Lemma 7 in [4]. Although its generalization to the case of $\epsilon > 0$ is straightforward, we provide the proof in Appendix C1 for completeness.

Lemma 6 implies that $\{\psi^{\otimes N}\}_N \xrightarrow{G\text{-cov.}} \{\phi^{\otimes \lfloor rN \rfloor}\}_N$ if and only if there exists a sequence of quantum channels $\{\mathcal{E}_N\}_N$ which asymptotically map $\mathcal{U}_g(\psi)^{\otimes N}$ to $\mathcal{U}'_g(\phi)^{\otimes \lfloor rN \rfloor}$ with vanishing error in $N \rightarrow \infty$ for any $g \in G$. Importantly, if we regard $g \in G$ as an unknown parameter, this sequence of quantum channels $\{\mathcal{E}_N\}_N$ achieve a conversion of a statistical model $\{\mathcal{U}_g(\psi)^{\otimes N}\}_{g \in G}$ to another statistical model $\{\mathcal{U}'_g(\phi)^{\otimes \lfloor rN \rfloor}\}_{g \in G}$ in the asymptotic limit. The asymptotic behavior of i.i.d. copies of parametric families of states has been studied in the context of QLAN [54–57], which aims to approximate i.i.d. statistical models by a limit model, called the Gaussian shift model. In the following, after briefly reviewing QLAN for i.i.d. pure-state models in [56, 57], we point out that QGT is a crucial characteristic of their asymptotics. Based on this observation, we construct quantum channels asymptotically converting a pure state to another pure state in RTA, proving Proposition 5.

A. Quantum local asymptotic normality

Here, let us briefly review studies in QLAN [54–57], which investigate the asymptotic properties of statistical models by associating them with their limit models.

We review particularly for QLAN on pure-state models [56, 57]. For a pure state ψ and a set of Hermitian operators $\mathbf{X} = \{X_i\}_{i=1}^m$, we define a pure-state statistical model $\mathcal{U}_\theta(\psi)$ with $\mathcal{U}_\theta(\cdot) = U_\theta(\cdot)U_\theta^\dagger$ and $U_\theta := e^{i\theta \cdot \mathbf{X}}$, where $\theta \cdot \mathbf{X} := \sum_{i=1}^m \theta^i X_i$ for $\theta \in \mathbb{R}^m$. To study its asymptotic property, we consider its i.i.d. copies of the state $(\mathcal{U}_{\mathbf{u}/\sqrt{N}}(\psi))^{\otimes N}$, where we introduced a local parameter $\mathbf{u} := \sqrt{N}\theta$. In the limit of $N \rightarrow \infty$, it is shown [56, 57] that this model is approximately interconvertible to the so-called Gaussian shift model with vanishing error.

For a more detailed explanation, let us introduce several notations in [56]. For a pure state $|\psi\rangle$ on a qudit system, we define a matrix $C : \mathbb{C}^m \rightarrow \mathbb{C}^{d-1}$ by $(C)_{ki} := -i \langle k|X_i|\psi\rangle$, where $\{|k\rangle\}_{k=1}^{d-1}$ denotes an orthonormal basis for the subspace orthogonal to $|\psi\rangle$. For $\mathbf{z} = (z_1, \dots, z_{d-1})^\top \in \mathbb{C}^{d-1}$, we denote a $(d-1)$ -mode coherent state with coherence amplitude \mathbf{z} by $|\mathbf{z}\rangle := \left(\bigotimes_{i=1}^{d-1} e^{z_i a_i^\dagger - z_i^* a_i} |0\rangle_i \right)$, where a_i^\dagger, a_i and $|0\rangle_i$ denote the creation and annihilation operator and the vacuum state for the i th mode. It is proven [57] that there exist sequences of quantum channels $\{\mathcal{T}_N\}_N$ and $\{\mathcal{S}_N\}_N$ such that

$$\begin{aligned} & \lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\mathcal{T}_N \left(\mathcal{U}_{\frac{\mathbf{u}}{\sqrt{N}}}(\psi)^{\otimes N} \right), |C\mathbf{u}\rangle \langle C\mathbf{u}| \right) \\ &= \lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\mathcal{U}_{\frac{\mathbf{u}}{\sqrt{N}}}(\psi)^{\otimes N}, \mathcal{S}_N(|C\mathbf{u}\rangle \langle C\mathbf{u}|) \right) \\ &= 0 \end{aligned} \quad (30)$$

for $\epsilon \in (0, 1/9)$.

B. QGT and asymptotic convertibility among i.i.d. statistical models

In the studies on QLAN, the main purpose is to approximate the i.i.d. statistical models using the Gaussian shift model in the asymptotic limit. Our focus is slightly different, and we aim to investigate the convertibility among i.i.d. statistical models to study the conversion rate in RTA. Our finding here is that the QGT plays a central role in characterizing asymptotic convertibility.

As shown in Eq. (30), $\left(\mathcal{U}_{\frac{\mathbf{u}}{\sqrt{N}}}(\psi) \right)^{\otimes N}$ is reversibly asymptotically convertible to the Gaussian-shift model $|C\mathbf{u}\rangle$, implying that the matrix C characterizes the asymptotic properties of the i.i.d. statistical model $\left(\mathcal{U}_{\frac{\mathbf{u}}{\sqrt{N}}}(\psi) \right)^{\otimes N}$. Such a matrix C is not uniquely determined since there remains the freedom in the choice of an orthonormal basis $\{|k\rangle\}_{k=1}^{d-1}$ for the subspace orthogonal to $|\psi\rangle$. Adopting another orthonormal basis, the matrix C is transformed to $C \mapsto \tilde{C} := VC$, where V denotes a $\mathbb{C}^{d-1} \times \mathbb{C}^{d-1}$ unitary matrix. By using the polar decomposition theorem, for matrices C and C' of the same size, we find that $\tilde{C} = VC$ holds for some unitary matrix V if and only if $C^\dagger C = \tilde{C}^\dagger \tilde{C}$. This fact implies that the essential characteristic of a statistical model is $C^\dagger C$, rather than the matrix C itself.

Importantly, the (i, j) element of $C^\dagger C$ is given by

$$\begin{aligned} (C^\dagger C)_{ij} &= \sum_{k=1}^{d-1} \langle \psi|X_i|k\rangle \langle k|X_j|\psi\rangle \\ &= \langle \psi|X_i(I - \psi)X_j|\psi\rangle, \end{aligned} \quad (31)$$

meaning that $C^\dagger C = \mathcal{Q}^\psi$, where \mathcal{Q}^ψ is the QGT for $\mathcal{U}_\theta(\psi)$. Therefore, if two pure-state statistical models

have the same QGT, they are asymptotically interconvertible to each other since both of them can be reversibly converted to the same Gaussian shift model, shown in Eq. (30).

As a generalization of the above observation, we prove the following:

Lemma 7. *For sets of Hermitian operators $\mathbf{X} := \{X_i\}_{i=1}^m$ and $\mathbf{X}' := \{X'_i\}_{i=1}^m$, we define $\mathcal{U}_\theta(\cdot) := e^{i\theta \cdot \mathbf{X}}(\cdot)e^{-i\theta \cdot \mathbf{X}}$ and $\mathcal{U}'_\theta(\cdot) := e^{i\theta \cdot \mathbf{X}'}(\cdot)e^{-i\theta \cdot \mathbf{X}'}$ for $\theta \in \mathbb{R}^m$. For given two pure states ψ and ϕ , we define pure-state statistical models by $\mathcal{U}_\theta(\psi)$ and $\mathcal{U}'_\theta(\phi)$ and denote their QGTs by \mathcal{Q}^ψ and \mathcal{Q}^ϕ , respectively. If $r > 0$ satisfies $\mathcal{Q}^\psi \geq r\mathcal{Q}^\phi$, then there exists a sequence of quantum channels $\{\mathcal{E}_N\}_N$ such that*

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\mathcal{E}_N \left(\mathcal{U}_{\frac{\mathbf{u}}{\sqrt{N}}}(\psi)^{\otimes N} \right), \mathcal{U}'_{\frac{\mathbf{u}}{\sqrt{N}}}(\phi)^{\otimes [rN]} \right) = 0 \quad (32)$$

for $\epsilon \in (0, 1/9)$.

In the proof of Lemma 7, the argument is divided into two parts: one focuses on the case where the QGT is conserved, while the other addresses the case where the QGT is reduced. We first prove that the growth rate of i.i.d. copies can be set to one by adjusting the scaling of the parameters. Concretely, $\mathcal{U}'_{\frac{\mathbf{u}}{\sqrt{N}}}(\phi)^{\otimes [rN]}$ is shown to be asymptotically interconvertible to $\mathcal{U}'_{\frac{\sqrt{r}\mathbf{u}}{\sqrt{N}}}(\phi)^{\otimes N}$. This asymptotic interconversion is achieved by combining the reversible conversion channels in QLAN, where the QGT is conserved. Then, we also prove that $\mathcal{U}_{\frac{\mathbf{u}}{\sqrt{N}}}(\psi)^{\otimes N}$ can be asymptotically converted into $\mathcal{U}'_{\frac{\sqrt{r}\mathbf{u}}{\sqrt{N}}}(\phi)^{\otimes N}$ if the QGT with respect to \mathbf{u} decreases in the sense of matrix inequality. The details of these lemmas and their proofs are provided in Appendix C 2.

C. Proof of achievability of the conversion rate

The process of transforming $\mathcal{U}_g(\psi)^{\otimes N}$ to $\mathcal{U}'_g(\phi)^{\otimes [rN]}$ with asymptotically vanishing error consists of two steps, which we may call the estimation and conversion steps. In the estimation step, we obtain a rough estimate \hat{g} of the true value g by consuming copies of $\mathcal{U}_g(\psi)$ sublinear in N . In the conversion step, based on the estimate \hat{g} , we convert the remaining copies of $\mathcal{U}_g(\psi)$ to i.i.d. copies of $\mathcal{U}'_g(\phi)$ by using the channel described in Lemma 7.

For a detailed explanation, let us first introduce notations. Let U and U' be (non-projective) unitary representations of a Lie group G on the input and output systems. Elements in the neighborhood of the identity $e \in G$ can be parametrized as $g(\boldsymbol{\lambda}) = e^{i \sum_{\mu=1}^{\dim G} \lambda^\mu A_\mu}$ by using a basis $\{A_\mu\}_{\mu=1}^{\dim G}$ of the Lie algebra \mathfrak{g} . Define operators $X_\mu := -i \frac{\partial}{\partial \lambda^\mu} U(g(\boldsymbol{\lambda}))|_{\boldsymbol{\lambda}=\mathbf{0}}$ and $X'_\mu := -i \frac{\partial}{\partial \lambda^\mu} U'(g(\boldsymbol{\lambda}))|_{\boldsymbol{\lambda}=\mathbf{0}}$, which correspond to the images of A_μ by the Lie algebra representations induced from

U and U' . At least locally, these operators satisfy $U(g(\boldsymbol{\lambda})) = e^{i \sum_{\mu=1}^{\dim G} \lambda^\mu X_\mu}$ and $U'(g(\boldsymbol{\lambda})) = e^{i \sum_{\mu=1}^{\dim G} \lambda^\mu X'_\mu}$. In this case, by slightly abusing our notation of \mathcal{U}_g and \mathcal{U}'_g , we also denote $\mathcal{U}_\theta(\cdot) := e^{i\theta \cdot \mathbf{X}}(\cdot)e^{-i\theta \cdot \mathbf{X}}$ and $\mathcal{U}'_\theta(\cdot) := e^{i\theta \cdot \mathbf{X}'}(\cdot)e^{-i\theta \cdot \mathbf{X}'}$ for $\theta \in \mathbb{R}^{\dim G}$, implicitly indicating $\|\theta\|$ is sufficiently small.

In the estimation step, we obtain a rough estimate $\hat{g} \in G$ of $g \in G$ by performing a measurement on n copies of the system in a state $\mathcal{U}_g(\psi)^{\otimes n}$, where $n := N^{1-\epsilon}$ for a fixed $\epsilon \in (0, 1/2)$. Let $p(\hat{g}|\mathcal{U}_g(\psi)^{\otimes n})$ denote the probability of obtaining an estimate \hat{g} when the system is in a state $\mathcal{U}_g(\psi)^{\otimes n}$. We say that the estimation is successful within an acceptable error of $\delta > 0$ if and only if the estimate $\hat{g} \in G$ is an element of the following set:

$$G_{\text{succ.}}(\delta) := \{\hat{g} \in G \mid \exists \theta, \mathcal{U}_\theta \circ \mathcal{U}_{\hat{g}}(\psi) = \mathcal{U}_g(\psi), \|\theta\| < \delta\}. \quad (33)$$

We define the success probability by

$$\begin{aligned} & \Pr_{\mathcal{U}_g(\psi)^{\otimes n}}(\|\theta\| < \delta) \\ & := \int_G d\mu_G(\hat{g}) p(\hat{g}|\mathcal{U}_g(\psi)^{\otimes n}) \chi_{G_{\text{succ.}}(\delta)}(\hat{g}), \end{aligned} \quad (34)$$

where μ_G is the Haar measure on the Lie group G , and χ_A denotes the indicator function of a set A . Note that the above success probability is well-defined since $G_{\text{succ.}}(\delta)$ is a measurable set, as shown in Appendix C3.

We show the following lemma, stating that the existence of an estimator that is successful within an acceptable error of $\delta = N^{-1/2+\epsilon}$ whose failure probability asymptotically vanishes:

Lemma 8. *Let G be a compact Lie group and ρ be an arbitrary state. Fix $\epsilon \in (0, 1/2)$. Then there exists an estimator of $g \in G$ by using consuming $\mathcal{U}_g(\rho)^{\otimes n}$ with $n = N^{1-\epsilon}$ such that its success probability satisfies*

$$\lim_{N \rightarrow \infty} \Pr_{\mathcal{U}_g(\rho)^{\otimes N^{1-\epsilon}}}(\|\theta\| < N^{-1/2+\epsilon}) = 1. \quad (35)$$

We remark that $\|\theta\| \sim N^{-1/2}$ for the error is the standard scaling in parameter estimation for i.i.d. models [69], including QLAN studies [54–57]. The proof is detailed in Appendix C4.

Proposition 5 is proven by combining Lemmas 7 and 8.

Proof of Proposition 5. From the assumption, $r > 0$ satisfies $\mathcal{Q}^{\mathcal{U}_g(\psi)} \geq r\mathcal{Q}^{\mathcal{U}'_g(\phi)}$ for all $g \in G$. Let $\hat{g} \in G$ be the estimator ensured to exist in Lemma 8 for $\epsilon \in (0, 1/9)$. Given the estimate \hat{g} , we apply Lemma 7 to pure states $(\mathcal{U}_{\hat{g}}(\psi))^{\otimes N'}$ and $(\mathcal{U}'_{\hat{g}}(\phi))^{\otimes \lfloor rN' \rfloor}$ for $N' := N - N^{1-\epsilon}$ and denote by $\{\mathcal{E}_{N'}^{\hat{g}}\}_{N'}$ the sequence of conversion channels. Applying $\mathcal{E}_{N'}^{\hat{g}}$ after the estimation step, the resulting state is given by

$$\begin{aligned} & \mathcal{E}_N(\mathcal{U}_g(\psi)^{\otimes N}) \\ & := \int d\mu_G(\hat{g}) p(\hat{g}|\mathcal{U}_g(\psi)^{\otimes N^{1-\epsilon}}) \mathcal{E}_{N'}^{\hat{g}}(\mathcal{U}_g(\psi)^{\otimes N'}). \end{aligned} \quad (36)$$

From the assumption $\text{Sym}_G(\psi) \subset \text{Sym}_G(\phi)$, if parameters θ satisfy $\mathcal{U}_\theta \circ \mathcal{U}_{\hat{g}}(\psi) = \mathcal{U}_g(\psi)$ for some $g, \hat{g} \in G$, then it also holds $\mathcal{U}'_\theta \circ \mathcal{U}'_{\hat{g}}(\phi) = \mathcal{U}'_g(\phi)$. Therefore, for $\hat{g} \in G_{\text{succ.}}(\delta)$ with $\delta > 0$, we have

$$\begin{aligned} & \text{Fid}\left(\mathcal{E}_{N'}^{\hat{g}}\left(\mathcal{U}_g(\psi)^{\otimes N'}\right), \mathcal{U}'_g(\phi)^{\otimes \lfloor rN' \rfloor}\right) \\ & = \text{Fid}\left(\mathcal{E}_{N'}^{\hat{g}}\left(\mathcal{U}_\theta(\psi_{\hat{g}})^{\otimes N'}\right), \mathcal{U}'_\theta(\phi_{\hat{g}})^{\otimes \lfloor rN' \rfloor}\right), \end{aligned} \quad (37)$$

where we have defined $\psi_{\hat{g}} := \mathcal{U}_{\hat{g}}(\psi)$ and $\phi_{\hat{g}} := \mathcal{U}'_{\hat{g}}(\phi)$, and Fid denotes the fidelity defined by $\text{Fid}(\rho, \sigma) := (\text{Tr}(\sqrt{\sqrt{\rho}\sigma\sqrt{\rho}}))^2$ for states ρ and σ . Since $N^{-1/2+\epsilon} < (N')^{-1/2+\epsilon}$, we have

$$\begin{aligned} f_N & := \inf_{\hat{g} \in G_{\text{succ.}}(N^{-1/2+\epsilon})} \text{Fid}\left(\mathcal{E}_{N'}^{\hat{g}}\left(\mathcal{U}_g(\psi)^{\otimes N'}\right), \mathcal{U}'_g(\phi)^{\otimes \lfloor rN' \rfloor}\right) \\ & \geq \inf_{\hat{g} \in G_{\text{succ.}}((N')^{-1/2+\epsilon})} \text{Fid}\left(\mathcal{E}_{N'}^{\hat{g}}\left(\mathcal{U}_g(\psi)^{\otimes N'}\right), \mathcal{U}'_g(\phi)^{\otimes \lfloor rN' \rfloor}\right) \\ & = \inf_{\|\mathbf{u}\| < (N')^\epsilon} \text{Fid}\left(\mathcal{E}_{N'}^{\hat{g}}\left(\mathcal{U}_{\frac{\mathbf{u}}{\sqrt{N'}}}(\psi_{\hat{g}})^{\otimes N'}\right), \mathcal{U}'_{\frac{\mathbf{u}}{\sqrt{N'}}}(\phi_{\hat{g}})^{\otimes \lfloor rN' \rfloor}\right) \\ & \xrightarrow{N \rightarrow \infty} 1, \end{aligned} \quad (38)$$

where in the last line, we have used Lemma 7 and the fact that the convergence in trace distance is equivalent to that in fidelity, shown by the Fuchs–van de Graaf inequalities.

From Eqs. (35), (36), and (38), we get

$$\begin{aligned} & \text{Fid}\left(\mathcal{E}_N(\mathcal{U}_g(\psi)^{\otimes N}), \mathcal{U}'_g(\phi)^{\otimes \lfloor rN' \rfloor}\right) \\ & = \int_G d\mu_G(\hat{g}) p(\hat{g}|\mathcal{U}_g(\psi)^{\otimes N^{1-\epsilon}}) \\ & \quad \times \text{Fid}\left(\mathcal{E}_{N'}^{\hat{g}}\left(\mathcal{U}_g(\psi)^{\otimes N'}\right), \mathcal{U}'_g(\phi)^{\otimes \lfloor rN' \rfloor}\right) \\ & \geq \int d\mu_G(\hat{g}) p(\hat{g}|\mathcal{U}_g(\psi)^{\otimes N^{1-\epsilon}}) \chi_{G_{\text{succ.}}(N^{-1/2+\epsilon})}(\hat{g}) \\ & \quad \times \text{Fid}\left(\mathcal{E}_{N'}^{\hat{g}}\left(\mathcal{U}_g(\psi)^{\otimes N'}\right), \mathcal{U}'_g(\phi)^{\otimes \lfloor rN' \rfloor}\right) \\ & \geq \Pr_{\mathcal{U}_g(\psi)^{\otimes N^{1-\epsilon}}}(\|\theta\| < N^{-1/2+\epsilon}) \times f_N \\ & \xrightarrow{N \rightarrow \infty} 1, \end{aligned} \quad (39)$$

implying that

$$\lim_{N \rightarrow \infty} T\left(\mathcal{E}_N(\mathcal{U}_g(\psi)^{\otimes N}), \mathcal{U}'_g(\phi)^{\otimes \lfloor rN' \rfloor}\right) = 0 \quad (40)$$

holds for any $g \in G$.

To complete the proof, we introduce $\delta \in (0, r)$. Then for all sufficiently large N , it holds $\lfloor rN' \rfloor > \lfloor (r-\delta)N \rfloor$. Denoting by $\{\Lambda_N\}_N$ the channel discarding $\lfloor rN' \rfloor - \lfloor (r-\delta)N \rfloor$ copies of the output system, we get

$$\lim_{N \rightarrow \infty} T\left(\Lambda_N \circ \mathcal{E}_N(\mathcal{U}_g(\psi)^{\otimes N}), \mathcal{U}'_g(\phi)^{\otimes \lfloor (r-\delta)N \rfloor}\right) = 0 \quad (41)$$

for any $g \in G$. From Lemma 6, this is equivalent to $\{\psi^{\otimes N}\}_N \xrightarrow{G\text{-cov.}} \{\phi^{\otimes \lfloor (r-\delta)N \rfloor}\}_N$, which complete the proof of Proposition 5. \square

VI. APPLICATIONS

Theorem 1 is valid for any compact Lie group. As a demonstration, we here apply our formula for several groups, which provides a unified understanding of prior studies and proves an unsolved conjecture on reversible transformations.

A. U(1) group

In a study [21] on RTA for time-translation symmetry, the approximate asymptotic conversion rate among states having the same finite period was calculated, which generalizes prior studies [2, 35, 36]. When the Hamiltonians of the input and output systems are H and H' , the time-translation unitary operators are given by e^{-iHt} and $e^{-iH't}$. For pure states ψ and ϕ , the periods are defined as $\tau := \inf\{t > 0 \mid e^{-iHt}\psi e^{iHt} = \psi\}$ and $\tau' := \inf\{t > 0 \mid e^{-iH't}\phi e^{iH't} = \phi\}$. In Theorem 1 in [21], it is proven that if $\tau = \tau'$, the conversion rate is given by the ratio of the variances, i.e.,

$$R(\psi \rightarrow \phi) = \frac{V(\psi, H)}{V(\phi, H')}. \quad (42)$$

We note that, when investigating the conversion among states with the same period, we can assume without loss of generality that the Hamiltonians have integer eigenvalues after appropriately redefining them [21, 70]. Therefore, this result corresponds to the case of $G = U(1)$.

In the notation of the present paper, we consider $G = U(1)$ and its representations $U(e^{i\theta}) = e^{iH\theta}$ and $U'(e^{i\theta}) = e^{iH'\theta}$ for $\theta \in [0, 2\pi)$, where H and H' are Hermitian operators whose eigenvalues are integers. Since $\dim G = 1$ for $G = U(1)$, the QGT is a scalar and given by $\mathcal{Q}^\psi = V(\psi, H)$ and $\mathcal{Q}^\phi = V(\phi, H')$. When the pure states ψ and ϕ has the same period, i.e., $\text{Sym}_G(\psi) = \text{Sym}_G(\phi)$, we find Eq. (42) immediately follows from Eq. (6).

B. Reversible asymptotic conversion: Proof of the Marvian-Spekkens conjecture

For pure states ψ and ϕ , we say that they are asymptotically reversibly convertible if and only if the conversion rates satisfy $R(\psi \rightarrow \phi)R(\phi \rightarrow \psi) = 1$. The conversion between pure states with equal periods is an example of reversible conversion, as Eq. (42) shows. Prior to the establishment of the conversion theory in RTA for $U(1)$ group [21], Marvian and Spekkens proposed a conjecture for a necessary and sufficient condition for reversible conversion in RTA for connected compact Lie groups [4, 6]. We prove a statement equivalent to the Marvian-Spekkens conjecture here, where the equivalence is shown later in this subsection.

Corollary 9. *For a connected compact Lie group G , pure states ψ and ϕ are asymptotically reversibly convertible if*

and only if both of the following conditions (A) and (B) are satisfied:

$$(A) \text{Sym}_G(\psi) = \text{Sym}_G(\phi).$$

$$(B) \exists! r > 0 \text{ such that } \mathcal{Q}^\psi = r\mathcal{Q}^\phi.$$

Note that the proportional constant r in condition (B) provides the conversion rate $R(\psi \rightarrow \phi)$.

Proof. Suppose that ψ and ϕ are asymptotically reversibly convertible. Then, $R(\psi \rightarrow \phi)$ and $R(\phi \rightarrow \psi)$ must be non-vanishing, which requires that the condition (A) must hold. In this case, Theorem 1 implies

$$R(\psi \rightarrow \phi) = \sup \{r \geq 0 \mid \mathcal{Q}^\psi \geq r\mathcal{Q}^\phi\}, \quad (43)$$

$$R(\phi \rightarrow \psi) = \sup \{r \geq 0 \mid \mathcal{Q}^\phi \geq r\mathcal{Q}^\psi\}. \quad (44)$$

These equations imply that $R(\psi \rightarrow \phi)R(\phi \rightarrow \psi) = 1$ holds only if the condition (B) holds. Conversely, if conditions (A) and (B) are satisfied, then Eqs. (43) and (44) imply $R(\psi \rightarrow \phi)R(\phi \rightarrow \psi) = 1$ holds. \square

Let us now review the statement of the Marvian-Spekkens conjecture and prove that it is equivalent to Corollary 9. Let $\mathbf{X} = \{X_k\}_k$ be the representation of a basis of the Lie algebra \mathfrak{g} . We define the *symmetrized* covariance matrix as

$$(C_{\mathfrak{g}}(\psi))_{kl} := \frac{1}{2} \langle \psi | (X_k X_l + X_l X_k) | \psi \rangle - \langle \psi | X_k | \psi \rangle \langle \psi | X_l | \psi \rangle. \quad (45)$$

The commutator subalgebra $i[\mathfrak{g}, \mathfrak{g}]$ is defined as the subalgebra spanned by $i[L_1, L_2]$ for all $L_1, L_2 \in \mathfrak{g}$. In general, the input and output Hilbert spaces \mathcal{H}_{in} and \mathcal{H}_{out} to which ψ and ϕ belong are different. However, as shown in Appendix B in [71], by considering a larger Hilbert space $\mathcal{H} := \mathcal{H}_{\text{in}} \oplus \mathcal{H}_{\text{out}}$ and a representation on it, it suffices to consider the case where the input and output Hilbert spaces are the same. The Marvian-Spekkens conjecture [4, 6] states that pure states ψ and ϕ are asymptotically reversibly convertible in RTA for a connected compact Lie group G if and only if all the following three conditions are satisfied:

$$(i) \text{Sym}_G(\psi) = \text{Sym}_G(\phi).$$

$$(ii) C_{\mathfrak{g}}(\psi) = R(\psi \rightarrow \phi)C_{\mathfrak{g}}(\phi).$$

$$(iii) \langle \psi | L | \psi \rangle = R(\psi \rightarrow \phi) \langle \phi | L | \phi \rangle \text{ for any element } L \text{ in the representation of the commutator subalgebra } i[\mathfrak{g}, \mathfrak{g}].$$

Condition (i) is the same as condition (A) in Corollary 9. Conditions (ii) and (iii) correspond respectively to the symmetric and anti-symmetric parts of condition (B) in Corollary 9.

We remark that other results on reversible conversion rates [2, 21, 37] also follow since the Marvian-Spekkens conjecture has now been proven.

C. Irreversibility in asymptotic conversion

The result in the previous subsection shows that conversion is asymptotically irreversible if and only if at least one of the conditions (A) and (B) in Corollary 9 is not satisfied.

Condition (A) can be violated even when $G = U(1)$. Indeed, if the periods τ and τ' of ψ and ϕ satisfy $\tau = k\tau'$ for some integer $k > 1$, then $\text{Sym}_G(\psi) \subsetneq \text{Sym}_G(\phi)$ holds, implying that $R(\psi \rightarrow \phi) = V(\psi, H)/V(\phi, H')$, while $R(\phi \rightarrow \psi) = 0$.

Condition (B) is easily violated when $\dim G > 1$. As a simple illustration, let us analyze $G = SU(2)$ and its unitary representation $e^{i\sum_{i=x,y,z}\theta^i J_i}$ with spin operators J_i . We adopt the z -axis as a quantization axis and denote simultaneous eigenstates of $J^2 := J_x^2 + J_y^2 + J_z^2$ and J_z by $|j, m\rangle$. To simplify the argument, we consider pure states $|\psi\rangle$ and $|\phi\rangle$ in a subspace spanned by the highest-weight states $\{|j, j\rangle\}_{j=0, \frac{1}{2}, 1, \frac{3}{2}, \dots, K}$, where K is an integer or a half-integer introduced to make the Hilbert space finite-dimensional. Following the notation in [2], we define

$$\mathcal{J} := \sum_{j=0, \frac{1}{2}, 1, \frac{3}{2}, \dots, K} j |j, j\rangle \langle j, j|, \quad (46)$$

$$\mathcal{M}(\psi) := 2 \langle \psi | \mathcal{J} | \psi \rangle, \quad (47)$$

$$\mathcal{V}(\psi) := 4(\langle \psi | \mathcal{J}^2 | \psi \rangle - \langle \psi | \mathcal{J} | \psi \rangle^2). \quad (48)$$

Since the QGT is calculated for ψ as

$$\mathcal{Q}^\psi = \frac{1}{4} \begin{pmatrix} \mathcal{M}(\psi) & \frac{1}{i}\mathcal{M}(\psi) & 0 \\ -\frac{1}{i}\mathcal{M}(\psi) & \mathcal{M}(\psi) & 0 \\ 0 & 0 & \mathcal{V}(\psi) \end{pmatrix}, \quad (49)$$

and similarly for ϕ , condition (B) holds only if $\mathcal{M}(\psi)/\mathcal{M}(\phi) = \mathcal{V}(\psi)/\mathcal{V}(\phi)$. This can also be explicitly confirmed from the conversion rate $R(\psi \rightarrow \phi) = \min \left\{ \frac{\mathcal{M}(\psi)}{\mathcal{M}(\phi)}, \frac{\mathcal{V}(\psi)}{\mathcal{V}(\phi)} \right\}$ calculated from Eq. (6) for pure states satisfying $\text{Sym}_G(\psi) \subset \text{Sym}_G(\phi)$. We remark that this conversion rate is consistent with the prior result on $SU(2)$ in Theorem 24 in [2], where the conversion rate is studied for a restricted set of pure states.

D. Finite groups

Here, we analyze the approximate asymptotic conversion theory in RTA for a finite group. As proven in [34], the conversion rate diverges for a finite group. We show that this fact can also be derived from our formula in Eq. (5) by lifting a finite group to a compact Lie group by appending a trivial phase, though a finite group itself is not a Lie group.

Let G be a finite group given by $G = \{g_i \mid i = 0, \dots, k\}$. Let U and U' be projective unitary representations of G on the input and output Hilbert spaces. Let us introduce unitary representations \tilde{U} and \tilde{U}' of $\tilde{G} := G \times U(1)$ such that $\tilde{U}(g, e^{i\theta}) := U(g)e^{i\theta}$ and $\tilde{U}'(g, e^{i\theta}) := U'(g)e^{i\theta}$ for $g \in G$ and $\theta \in [0, 2\pi)$. Since

the phase θ does not affect the state, a channel is G -covariant if and only if \tilde{G} -covariant. Therefore, appending the trivial phase leaves the conversion rate invariant. Since \tilde{G} is a compact Lie group, we can apply Eq. (5). The QGT vanishes for any pure state for \tilde{U} and \tilde{U}' , implying that $\mathcal{Q}^\psi \geq r\mathcal{Q}^\phi$ holds for any r . In addition, $\text{Sym}_G(\psi) \subset \text{Sym}_G(\phi)$ holds iff $\text{Sym}_{\tilde{G}}(\psi) \subset \text{Sym}_{\tilde{G}}(\phi)$. Therefore, we get

$$R(\psi \rightarrow \phi) = \begin{cases} \infty & (\text{if } \text{Sym}_G(\psi) \subset \text{Sym}_G(\phi)) \\ 0 & (\text{otherwise}) \end{cases}, \quad (50)$$

reproducing the result on the approximate asymptotic conversion rate in [34].

The proof of the direct part provides a clear understanding of the reason why the approximate asymptotic conversion rate diverges for finite groups. Since there is only a finite number of different elements in $\{\mathcal{U}_g(\psi) \mid g \in G\}$, it is possible to identify the state $\mathcal{U}_g(\psi)$ with an exponentially small failure probability using a state tomography protocol on $\mathcal{U}_g(\psi)^{\otimes N}$. Let $\hat{g} \in G$ denote the estimate of the true value g obtained by this tomography protocol. Depending on the estimate \hat{g} , we can prepare an arbitrary number of copies of $\mathcal{U}'_{\hat{g}}(\phi)$. The condition $\text{Sym}_G(\psi) \subset \text{Sym}_G(\phi)$ implies that if $\mathcal{U}_{\hat{g}}(\psi) = \mathcal{U}_g(\psi)$ is satisfied, then $\mathcal{U}'_{\hat{g}}(\phi) = \mathcal{U}'_g(\phi)$ also holds. Therefore, the approximate asymptotic conversion rate diverges since it is possible to create an arbitrary number of copies of $\mathcal{U}'_{\hat{g}}(\phi)$ from $\mathcal{U}_g(\psi)^{\otimes N}$ with asymptotically vanishing failure probability. We emphasize that the exact asymptotic conversion rate in [34] is *not* covered by this analysis because there always remains an error in the tomography process.

VII. CONCLUSION

In this paper, we established the approximate asymptotic conversion theory for i.i.d. pure states in RTA for a compact Lie group. Our result provides a formula for the asymptotic conversion rate in RTA for any compact Lie group by using QGTs, implying that QGT fulfills the same function in RTA as entanglement entropy does in entanglement theory. In other words, the present study unveils the distinguished role of QGT as an asymmetry measure governing the conversion rate in resource manipulation in RTA. As a demonstration, we applied our formula to several different setups. Prior to the present research, the i.i.d. asymptotic conversion theory among pure states in RTA for a Lie group is established only for $U(1)$ group [21], and a few examples of restricted sets of pure states have been studied for other Lie groups [2, 37]. We first proved that the formula for the conversion rate in RTA for $U(1)$ in [21] is reproduced from our formula. We also affirmatively proved the Marvian-Spekkens conjecture [6, 72] on reversible conversion in RTA as a corollary of our main theorem, which has remained unproven for a decade. We also studied the examples where the

asymptotic conversion is irreversible for $U(1)$ and $SU(2)$ groups. The conversion rate calculated for this example of $SU(2)$ is consistent with the result in [2], where the conversion rate is studied on a restricted set of pure states. Furthermore, we demonstrated that RTA for finite groups can also be analyzed in our framework by appending a trivial phase, providing an alternative proof of the divergence in the approximate asymptotic conversion rate for any finite group [34].

ACKNOWLEDGMENTS

The authors would like to thank Tomohiro Shitara for providing his expert knowledge on monotone metrics, Yui Kuramochi for valuable comments on an early draft of this manuscript, and also Iman Marvian for the valuable discussion. K.Y. acknowledges support from JSPS KAKENHI Grant No. JP24KJ0085. Y.M. is supported by JSPS KAKENHI Grant No. JP23KJ0421. H.T. was supported by MEXT KAKENHI Grant-in-Aid for Transformative Research Areas B “Quantum Energy Innovation” Grant Numbers 24H00830 and 24H00831, and JST PRESTO No. JPMJPR2014, JST MOONSHOT No. JPMJMS2061.

-
- [1] S. D. Bartlett, T. Rudolph, and R. W. Spekkens, *Reviews of Modern Physics* **79**, 555 (2007).
- [2] G. Gour and R. W. Spekkens, *New Journal of Physics* **10**, 033023 (2008).
- [3] G. Gour, I. Marvian, and R. W. Spekkens, *Physical Review A* **80**, 012307 (2009).
- [4] I. Marvian Mashhad, *Symmetry, Asymmetry and Quantum Information*, Ph.D. thesis, University of Waterloo (2012).
- [5] K. Korzekwa, *Resource theory of asymmetry*, Master’s thesis, Imperial College London (2013).
- [6] I. Marvian and R. W. Spekkens, *Physical Review A* **90**, 014102 (2014).
- [7] E. Chitambar and G. Gour, *Reviews of Modern Physics* **91**, 025001 (2019).
- [8] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, *Reviews of Modern Physics* **81**, 865 (2009).
- [9] J. Aberg, [arXiv:quant-ph/0612146](https://arxiv.org/abs/quant-ph/0612146) (2006).
- [10] T. Baumgratz, M. Cramer, and M. Plenio, *Physical Review Letters* **113**, 140401 (2014).
- [11] A. Streltsov, G. Adesso, and M. B. Plenio, *Reviews of Modern Physics* **89**, 041003 (2017).
- [12] F. G. S. L. Brandão, M. Horodecki, J. Oppenheim, J. M. Renes, and R. W. Spekkens, *Physical Review Letters* **111**, 250404 (2013).
- [13] F. Brandão, M. Horodecki, N. Ng, J. Oppenheim, and S. Wehner, *Proceedings of the National Academy of Sciences* **112**, 3275 (2015).
- [14] I. Marvian, R. W. Spekkens, and P. Zanardi, *Physical Review A* **93**, 052331 (2016).
- [15] I. Marvian and R. W. Spekkens, *Physical Review Letters* **123**, 020404 (2019).
- [16] M. Lostaglio and M. P. Müller, *Physical Review Letters* **123**, 020403 (2019).
- [17] M. Lostaglio, K. Korzekwa, D. Jennings, and T. Rudolph, *Physical Review X* **5**, 021001 (2015), publisher: American Physical Society.
- [18] P. Faist, J. Oppenheim, and R. Renner, *New J. Phys.* **17**, 043003 (2015).
- [19] I. Marvian, *Nature Communications* **11**, 25 (2020).
- [20] H. Tajima and R. Takagi, “Gibbs-preserving operations requiring infinite amount of quantum coherence,” (2024), [arXiv:2404.03479](https://arxiv.org/abs/2404.03479) [quant-ph].
- [21] I. Marvian, *Physical Review Letters* **129**, 190502 (2022).
- [22] A. Kubica and R. Demkowicz-Dobrzański, *Physical Review Letters* **126**, 150503 (2021).
- [23] S. Zhou, Z.-W. Liu, and L. Jiang, *Quantum* **5**, 521 (2021).
- [24] Y. Yang, Y. Mo, J. M. Renes, G. Chiribella, and M. P. Woods, *Physical Review Research* **4**, 023107 (2022).
- [25] H. Tajima and K. Saito, [arXiv:2103.01876](https://arxiv.org/abs/2103.01876) [cond-mat, physics:quant-ph] (2021), 10.48550/arXiv.2103.01876.
- [26] Z.-W. Liu and S. Zhou, [arXiv:2111.06360](https://arxiv.org/abs/2111.06360) [quant-ph] (2022), <https://doi.org/10.48550/arXiv.2111.06360>.
- [27] H. Tajima, R. Takagi, and Y. Kuramochi, “Universal trade-off structure between symmetry, irreversibility, and quantum coherence in quantum processes,” (2022), [arXiv:2206.11086](https://arxiv.org/abs/2206.11086) [cond-mat, physics:hep-th, physics:quant-ph].
- [28] Z.-W. Liu and S. Zhou, *npj Quantum Information* **9**, 1 (2023).
- [29] H. Tajima, N. Shiraishi, and K. Saito, *Physical Review Letters* **121**, 110403 (2018).
- [30] H. Tajima, N. Shiraishi, and K. Saito, *Physical Review Research* **2**, 043374 (2020).
- [31] M. Ahmadi, D. Jennings, and T. Rudolph, *New J. Phys.* **15**, 013057 (2013).
- [32] I. Marvian and R. W. Spekkens, (2012), [arXiv:1212.3378](https://arxiv.org/abs/1212.3378).
- [33] H. Tajima and H. Nagaoka, [arXiv:1909.02904](https://arxiv.org/abs/1909.02904) [cond-mat, physics:quant-ph] (2019), 10.48550/arXiv.1909.02904.
- [34] T. Shitara, Y. Mitsuhashi, and H. Tajima, “The i.i.d. state convertibility in the resource theory of asymmetry for finite groups,” (2024).
- [35] N. Schuch, F. Verstraete, and J. I. Cirac, *Physical Review Letters* **92**, 087904 (2004).
- [36] N. Schuch, F. Verstraete, and J. I. Cirac, *Physical Review A* **70**, 042310 (2004).
- [37] Y. Yang, G. Chiribella, and Q. Hu, *New Journal of Physics* **19**, 123003 (2017).
- [38] F. G. Brandão and G. Gour, *Physical Review Letters* **115**, 070503 (2015).
- [39] F. G. S. L. Brandão and M. B. Plenio, *Communications in Mathematical Physics* **295**, 791 (2010).
- [40] M. Berta, F. G. S. L. Brandão, G. Gour, L. Lami, M. B. Plenio, B. Regula, and M. Tomamichel, *Quantum* **7**, 1103 (2023).

- [41] M. Hayashi and H. Yamasaki, “Generalized Quantum Stein’s Lemma and Second Law of Quantum Resource Theories,” (2024).
- [42] L. Lami, “A solution of the generalised quantum Stein’s lemma,” (2024).
- [43] C. H. Bennett, H. J. Bernstein, S. Popescu, and B. Schumacher, *Physical Review A* **53**, 2046 (1996).
- [44] J. A. Vaccaro, F. Anselmi, H. M. Wiseman, and K. Jacobs, *Physical Review A* **77**, 032114 (2008).
- [45] D. Kudo and H. Tajima, *Physical Review A* **107**, 062418 (2023).
- [46] L. Gao, H. Li, I. Marvian, and C. Rouzé, *Communications in Mathematical Physics* **405**, 180 (2024).
- [47] F. Hansen, *Proceedings of the National Academy of Sciences* **105**, 9909 (2008).
- [48] C. Zhang, B. Yadin, Z.-B. Hou, H. Cao, B.-H. Liu, Y.-F. Huang, R. Maity, V. Vedral, C.-F. Li, G.-C. Guo, and D. Girolami, *Physical Review A* **96**, 042327 (2017).
- [49] R. Takagi, *Scientific Reports* **9**, 14562 (2019).
- [50] K. Yamaguchi and H. Tajima, *Quantum* **7**, 1012 (2023).
- [51] J. P. Provost and G. Vallee, *Communications in Mathematical Physics* **76**, 289 (1980).
- [52] M. V. Berry, in *Geometric Phases in Physics*, Vol. 5 (World Scientific Singapore, 1989) pp. 3–28.
- [53] D. Petz, *Linear Algebra and its Applications* **244**, 81 (1996).
- [54] M. Guță and J. Kahn, *Physical Review A* **73**, 052108 (2006).
- [55] J. Kahn and M. Guță, *Communications in Mathematical Physics* **289**, 597 (2009).
- [56] F. Girotti, A. Godley, and M. Guță, *Journal of Physics A: Mathematical and Theoretical* **57**, 245304 (2024).
- [57] S. Lahiry and M. Nussbaum, *Bernoulli* **30**, 610 (2024).
- [58] C. W. Helstrom, *Journal of Statistical Physics* **1**, 231 (1969).
- [59] A. S. Holevo, *Journal of Multivariate Analysis* **3**, 337 (1973).
- [60] R. Cheng, “Quantum Geometric Tensor (Fubini-Study Metric) in Simple Quantum System: A pedagogical Introduction,” (2013), arXiv:1012.1337 [math-ph, physics:quant-ph].
- [61] J. Liu, H. Yuan, X.-M. Lu, and X. Wang, *Journal of Physics A: Mathematical and Theoretical* **53**, 023001 (2019).
- [62] B. C. Hall, *Lie Groups, Lie Algebras, and Representations: An Elementary Introduction*, Graduate Texts in Mathematics, Vol. 222 (Springer International Publishing, Cham, 2015).
- [63] E. A. Morozova and N. N. Chentsov, *Itogi Nauki i Tekhniki* **36**, 69 (1989).
- [64] D. Petz and C. Ghinea, in *Quantum Probability and Related Topics*, QP-PQ: Quantum Probability and White Noise Analysis, Vol. Volume 27 (WORLD SCIENTIFIC, 2011) pp. 261–281.
- [65] M. Hayashi, *Quantum Information Theory: Mathematical Foundation*, Graduate Texts in Physics (Springer Berlin Heidelberg, Berlin, Heidelberg, 2017).
- [66] T. Shitara, K. Yamaguchi, and H. Tajima, (2024), in preparation.
- [67] M. J. Donald, M. Horodecki, and O. Rudolph, *Journal of Mathematical Physics* **43**, 4252 (2002).
- [68] M. Plenio and S. Virmani, *Quantum Information and Computation* **7**, 1 (2007).
- [69] R. D. Gill and S. Massar, *Physical Review A* **61**, 042312 (2000).
- [70] K. Yamaguchi and H. Tajima, *Physical Review Letters* **131**, 200203 (2023).
- [71] I. Marvian and R. W. Spekkens, *New Journal of Physics* **15**, 033001 (2013).
- [72] I. Marvian and R. W. Spekkens, arXiv:1212.3378 [quant-ph] (2012).
- [73] A. W. Knap, *Lie Groups Beyond an Introduction* (Springer Science & Business Media, 2002).
- [74] T. Ogawa and H. Nagaoka, *IEEE Transactions on Information Theory* **46**, 2428 (2000).
- [75] D. Pérez-García, M. M. Wolf, D. Petz, and M. B. Ruskai, *Journal of Mathematical Physics* **47**, 083506 (2006).
- [76] M. Guță, J. Kahn, R. Kueng, and J. A. Tropp, *Journal of Physics A: Mathematical and Theoretical* **53**, 204001 (2020).

Appendix A: Proof for the main theorem for any projective unitary representations

In the main text, we proved Eq. (7) for (non-projective) unitary representations of a compact Lie group. In Appendix A 1, we first review the result in [34] that relates the conversion rates for projective unitary representations to that for (non-projective) unitary representations and explain a formula for the conversion rate for projective unitary representations that are continuous but not differentiable. In Appendix A 2, we first show that the QGTs at different points of a connected component of a Lie group are interrelated by a congruence transformation, which plays an essential role in proving Theorem 1 from Eq. (7). In Appendix A 3, we complete the proof for Eq. (5) in Theorem 1 and Eq. (7) for any projective unitary representations that are differentiable. Furthermore, Eq. (8) is proven Sec. A 4.

1. Differentiability of representation

A (non-projective) unitary representation is differentiable as long as it is continuous [62], whereas this is not always the case for a projective unitary representation. However, by adopting the following result in [34], the conversion rate can be calculated even when projective unitary representation is not differentiable.

Lemma 10. *Let U, U' be projective unitary representations of G on finite-dimensional Hilbert spaces \mathcal{H} and \mathcal{H}' . Define maps \tilde{U} and \tilde{U}' by*

$$\tilde{U}(g) := \frac{U(g)^{\otimes d}}{\det(U(g))}, \quad \tilde{U}'(g) := \frac{U'(g)^{\otimes d'}}{\det(U'(g))}, \quad (\text{A1})$$

where $d := \dim \mathcal{H}$ and $d' := \dim \mathcal{H}'$. Then, \tilde{U} and \tilde{U}' are a (non-projective) unitary representations of G on $\mathcal{H}^{\otimes d}$ and $\mathcal{H}'^{\otimes d'}$, respectively. Furthermore, it holds

$$R(\rho \rightarrow \sigma) = \frac{d'}{d} R(\rho^{\otimes d} \rightarrow \sigma^{\otimes d'}), \quad (\text{A2})$$

where on the right-hand side, the convertibility is defined with respect to the unitary representations \tilde{U} and \tilde{U}' .

The proof can be found in [34]. Note that this lemma is valid for any groups, including both finite groups and Lie groups, as long as the dimensions of the representation spaces are finite.

For projective unitary representations U and U' that are continuous, (non-projective) unitary representations \tilde{U} and \tilde{U}' in Eq. (A1) are continuous and hence differentiable. Therefore, the conversion rate among pure state is obtained from Eq. (A2) since its right-hand side can be calculated from Theorem 1.

Except for this subsection, we only consider representations that are differentiable. In the following subsections, by using Eq. (A2), we show that the formula for conversion rate described by QGTs in Theorem 1 is valid for projective unitary representations U and U' that are differentiable.

2. Relation among matrix inequalities between QGTs at different group elements for unitary representations

Let us prove the relation of the QGT at different points of a connected compact Lie group.

Lemma 11. *Let U be a (non-projective) unitary representation of a connected compact Lie group G . For each $g \in G$, there exists an invertible real matrix $V(g)$ independent of the representation U such that $\mathcal{Q}^{U_g(\psi)} = V(g)^\top \mathcal{Q}^\psi V(g)$.*

We remark the same argument for the congruence transformation of quantum Fisher information matrices can be found in [46].

Proof of Lemma 11. Let us first review several useful facts on a compact Lie group G and its Lie algebra \mathfrak{g} . Let $GL(n, \mathbb{C})$ denote the general linear group. A closed subgroup of $GL(n, \mathbb{C})$ is referred to as a closed linear group. Any compact Lie group G is isomorphic to a closed linear group (see, e.g., Corollary 4.22 in [73]). Through this identification, we can regard the exponential map from the Lie algebra \mathfrak{g} to a compact Lie group G as the matrix exponential. Now, let us fix an arbitrary basis $\{B_\mu\}_{\mu=1}^{\dim G}$ of \mathfrak{g} . For any $g \in G$, the set $\{g^{-1}B_\mu g\}_{\mu=1}^{\dim G}$ is also a basis of \mathfrak{g} . Therefore, we can introduce an invertible real matrix $V(g)$ by

$$g^{-1}B_\mu g =: \sum_{\nu=1}^{\dim G} V(g)_{\nu\mu} B_\nu. \quad (\text{A3})$$

If we further assume that G is a connected and compact Lie group, then the exponential map is surjective (see e.g., [73]), implying that there is $A_g \in \mathfrak{g}$ such that $g = \exp(iA_g)$. Therefore,

$$g^{-1}B_\mu g = \exp(-iA_g)B_\mu \exp(iA_g) = \sum_{n=0}^{\infty} \frac{i^n}{n!} [\cdots [B_\mu, \underbrace{A_g, A_g}_{n}, \cdots], A_g]. \quad (\text{A4})$$

This means that the matrix $V(g)$ is determined by the Lie algebraic structure and is independent of the representation if G is a connected compact Lie group.

Let us fix a local coordinate of G in a neighborhood of the identity $e \in G$, which parametrizes the elements as $g(\boldsymbol{\lambda})$ such that $g(\mathbf{0}) = e$. In this coordinate system, the QGT for ψ is given by

$$\mathcal{Q}_{\mu\nu}^\psi = \langle \psi | X_\mu X_\nu | \psi \rangle - \langle \psi | X_\mu | \psi \rangle \langle \psi | X_\nu | \psi \rangle, \quad (\text{A5})$$

where $X_\mu := -i \frac{\partial}{\partial \lambda^\mu} U(g(\boldsymbol{\lambda})) \big|_{\boldsymbol{\lambda}=\mathbf{0}}$. Similarly, the QGT for $\mathcal{U}_g(\psi) = |\psi(g)\rangle \langle \psi(g)|$, where $|\psi(g)\rangle := U(g) |\psi\rangle$, is given by

$$\mathcal{Q}_{\mu\nu}^{\mathcal{U}_g(\psi)} = \langle \psi(g) | X_\mu (I - |\psi(g)\rangle \langle \psi(g)|) X_\nu | \psi(g) \rangle \quad (\text{A6})$$

$$= \langle \psi | X_\mu(g) X_\nu(g) | \psi \rangle - \langle \psi | X_\mu(g) | \psi \rangle \langle \psi | X_\nu(g) | \psi \rangle, \quad (\text{A7})$$

where $X_\mu(g) := U(g)^\dagger X_\mu U(g)$. By using the matrix defined in Eq. (A3), we get

$$U(g)^\dagger X_\mu U(g) = \sum_{\nu=1}^{\dim G} V(g)_{\nu\mu} X_\nu. \quad (\text{A8})$$

Therefore, $\mathcal{Q}^{\mathcal{U}_g(\psi)} = V(g)^\top \mathcal{Q}^\psi V(g)$. □

As an immediate corollary, we find that a matrix inequality between QGTs holds for any point of a connected compact Lie group G if it holds at a point in G .

Corollary 12. *Let U and U' be a (non-projective) unitary representation of a connected compact Lie group G . If a matrix inequality $\mathcal{Q}^{\mathcal{U}_g(\psi)} \geq r \mathcal{Q}^{\mathcal{U}'_g(\phi)}$ holds for a point $g \in G$, then it also holds for any $g \in G$.*

Proof. Since the reversible matrix $V(g)$ in Lemma 11 is independent of unitary representation, we immediately get

$$\begin{aligned} \forall g \in G, \quad \mathcal{Q}^{\mathcal{U}_g(\psi)} &\geq r \mathcal{Q}^{\mathcal{U}'_g(\phi)} \\ \iff \exists g \in G, \quad \mathcal{Q}^{\mathcal{U}_g(\psi)} &\geq r \mathcal{Q}^{\mathcal{U}'_g(\phi)}. \end{aligned} \quad (\text{A9})$$

□

We can generalize the above corollary to any compact Lie group G . Let G_0 be the identity component of G . Since G_0 is a normal subgroup of G , we can introduce a quotient group G/G_0 , which characterizes the disconnected group elements. In other words, the component of G connected to $g \in G$, i.e., the set of group elements that are connected to $g \in G$ can be written as $G_0 g$. Note that a compact Lie group G has only a finite number of components (see, e.g., [62]). We denote the decomposition into the connected components by $G = \bigsqcup_{i=0}^k G_i$. We arbitrarily fix $g_i \in G_i$ for $i = 0, \dots, k$. Applying the argument in Corollary 12 for each connected component, we get

$$\begin{aligned} \forall g \in G, \quad \mathcal{Q}^{\mathcal{U}_g(\psi)} &\geq r \mathcal{Q}^{\mathcal{U}'_{g_i}(\phi)} \\ \iff \forall i = 0, \dots, k, \quad \forall g \in G_i, \quad \mathcal{Q}^{\mathcal{U}_g(\psi)} &\geq r \mathcal{Q}^{\mathcal{U}'_{g_i}(\phi)} \\ \iff \forall i = 0, \dots, k, \quad \mathcal{Q}^{\mathcal{U}_{g_i}(\psi)} &\geq r \mathcal{Q}^{\mathcal{U}'_{g_i}(\phi)}. \end{aligned} \quad (\text{A10})$$

3. Conversion rates for projective unitary representations and (non-projective) unitary representations

Using a projective unitary representation that is differentiable, the QGT is obtained as Eq. (4). This QGT is proportional to the QGT defined with the unitary representation in Eq. (A1):

Lemma 13. *Let U be a projective unitary representation of a Lie group G on a d -dimensional Hilbert space \mathcal{H} . Define a (non-projective) unitary representation \tilde{U} by Eq. (A1). Then it holds*

$$\forall g \in G, \quad \mathcal{Q}^{\mathcal{U}_g(\psi)} = \frac{1}{d} \mathcal{Q}^{\tilde{\mathcal{U}}_g(\psi)^{\otimes d}} \quad (\text{A11})$$

Proof. Let us fix a local coordinate $g(\boldsymbol{\lambda})$ in the neighborhood of the identity $e \in G$ such that $g(\mathbf{0}) = e$. From the definition of \tilde{U} , we get

$$\partial_\mu \tilde{U}(g(\boldsymbol{\lambda})) \Big|_{\boldsymbol{\lambda}=\mathbf{0}} \quad (\text{A12})$$

$$= \frac{1}{\det(U(g(\boldsymbol{\lambda})))} \sum_{i=1}^d U(g(\boldsymbol{\lambda}))^{\otimes i-1} \otimes \partial_\mu U(g(\boldsymbol{\lambda})) \otimes U(g(\boldsymbol{\lambda}))^{\otimes d-i} \Big|_{\boldsymbol{\lambda}=\mathbf{0}} + U(g(\boldsymbol{\lambda}))^{\otimes d} \partial_\mu \frac{1}{\det(U(g(\boldsymbol{\lambda})))} \Big|_{\boldsymbol{\lambda}=\mathbf{0}} \quad (\text{A13})$$

$$= e^{-i\theta} \sum_{i=1}^d I^{\otimes i-1} \otimes \partial_\mu U(g(\boldsymbol{\lambda})) \Big|_{\boldsymbol{\lambda}=\mathbf{0}} \otimes I^{\otimes d-i} + c_\mu e^{i\theta d} I^{\otimes d} \quad \left(U(e) =: e^{i\theta} I, c_\mu := \partial_\mu \frac{1}{\det(U(g(\boldsymbol{\lambda})))} \Big|_{\boldsymbol{\lambda}=\mathbf{0}} \right). \quad (\text{A14})$$

Note that the second term, which is proportional to the identity matrix, does not contribute to the QGT. Thus, we get

$$\begin{aligned} \frac{1}{d} \mathcal{Q}_{\mu\nu}^{\psi^{\otimes d}} &= \frac{1}{d} \sum_{i=1}^d (\langle \partial_\mu U(g(\boldsymbol{\lambda}))\psi | \partial_\nu U(g(\boldsymbol{\lambda}))\psi \rangle - \langle \partial_\mu U(g(\boldsymbol{\lambda}))\psi | U(g(\boldsymbol{\lambda}))\psi \rangle \langle U(g(\boldsymbol{\lambda}))\psi | \partial_\nu U(g(\boldsymbol{\lambda}))\psi \rangle) \Big|_{\boldsymbol{\lambda}=\mathbf{0}} \\ &= \mathcal{Q}_{\mu\nu}^\psi. \end{aligned} \quad (\text{A15})$$

Since this equality holds for an arbitrary pure state ψ , we get

$$\forall g \in G, \quad \mathcal{Q}^{\mathcal{U}_g(\psi)} = \frac{1}{d} \mathcal{Q}^{\tilde{\mathcal{U}}_g(\psi^{\otimes d})}. \quad (\text{A16})$$

□

Now we are ready to prove Eq. (7) for any projective unitary representations.

Proof of Eq. (7). We consider the case where $\text{Sym}_G(\psi) \subset \text{Sym}_G(\phi)$. Note that the case where $\text{Sym}_G(\psi) \not\subset \text{Sym}_G(\phi)$ is separately treated in Appendix A4.

From Proposition 4 and Proposition 5, we have proven Eq. (7) for any (non-projective) unitary representations. For any projective unitary representations U and U' on \mathcal{H} and \mathcal{H}' , we define unitary representations \tilde{U} and \tilde{U}' of G on $\mathcal{H}^{\otimes d}$ and $\mathcal{H}'^{\otimes d'}$ by Eq. (A1). From Eq. (A2), we have

$$R(\psi \rightarrow \phi) = \frac{d'}{d} R(\psi^{\otimes d} \rightarrow \phi^{\otimes d'}). \quad (\text{A17})$$

Since Eq. (7) is already proven for any (non-projective) unitary representations, we get

$$R(\psi^{\otimes d} \rightarrow \phi^{\otimes d'}) = \sup\{r \geq 0 \mid \forall g \in G, \mathcal{Q}^{\tilde{\mathcal{U}}_g(\psi^{\otimes d})} \geq r \mathcal{Q}^{\tilde{\mathcal{U}}'_g(\phi^{\otimes d'})}\}. \quad (\text{A18})$$

By using Eq. (A11), we get

$$\sup\{r \geq 0 \mid \forall g \in G, \mathcal{Q}^{\tilde{\mathcal{U}}_g(\psi^{\otimes d})} \geq r \mathcal{Q}^{\tilde{\mathcal{U}}'_g(\phi^{\otimes d'})}\} = \frac{d}{d'} \sup\{r \geq 0 \mid \forall g \in G, \mathcal{Q}^{\mathcal{U}_g(\psi)} \geq r \mathcal{Q}^{\mathcal{U}'_g(\phi)}\}. \quad (\text{A19})$$

Therefore,

$$R(\psi \rightarrow \phi) = \sup\{r \geq 0 \mid \forall g \in G, \mathcal{Q}^{\mathcal{U}_g(\psi)} \geq r \mathcal{Q}^{\mathcal{U}'_g(\phi)}\}. \quad (\text{A20})$$

□

By using the result in Appendix A2, we prove Eq. (5) in Theorem 1.

Proof of Eq. (5) in Theorem 1. For given projective unitary representations U and U' of G , we introduce unitary representations by Eq. (A1). Then it holds,

$$\sup\{r \geq 0 \mid \forall g \in G, \mathcal{Q}^{\mathcal{U}_g(\psi)} \geq r \mathcal{Q}^{\mathcal{U}_g(\phi)}\} \stackrel{\text{Eq. (A11)}}{=} \frac{d'}{d} \sup\{r \geq 0 \mid \forall g \in G, \mathcal{Q}^{\tilde{\mathcal{U}}_g(\psi^{\otimes d})} \geq r \mathcal{Q}^{\tilde{\mathcal{U}}'_g(\phi^{\otimes d'})}\} \quad (\text{A21})$$

$$\stackrel{\text{Eq. (A10)}}{=} \frac{d'}{d} \sup\{r \geq 0 \mid \forall i = 0, \dots, k, \mathcal{Q}^{\tilde{\mathcal{U}}_{g_i}(\psi^{\otimes d})} \geq r \mathcal{Q}^{\tilde{\mathcal{U}}'_{g_i}(\phi^{\otimes d'})}\} \quad (\text{A22})$$

$$\stackrel{\text{Eq. (A11)}}{=} \sup\{r \geq 0 \mid \forall i = 0, \dots, k, \mathcal{Q}^{\mathcal{U}_{g_i}(\psi)} \geq r \mathcal{Q}^{\mathcal{U}'_{g_i}(\phi)}\}. \quad (\text{A23})$$

Applying Eq. (7), we complete the proof of Eq. (5) in Theorem 1. □

4. Symmetry subgroup and the conversion rate

We here prove that $R(\psi \rightarrow \phi) = 0$ unless $\text{Sym}_G(\psi) \subset \text{Sym}_G(\phi)$. In fact, we can prove this fact not only for pure states but also for generic states:

Proposition 14. *If $\text{Sym}_G(\rho) \not\subset \text{Sym}_G(\sigma)$, then $R(\rho \rightarrow \sigma) = 0$.*

Proof. Suppose that $\text{Sym}_G(\rho) \not\subset \text{Sym}_G(\sigma)$. Then there exists an element $g_* \in G$ such that $g_* \in \text{Sym}_G(\rho)$ and $g_* \notin \text{Sym}_G(\sigma)$. Define $\Delta := \text{Fid}(\mathcal{U}_{g_*}(\sigma), \sigma)$. Since $g_* \notin \text{Sym}_G(\sigma)$, we have $\text{Fid}(\mathcal{U}_{g_*}(\sigma), \sigma) < 1$ and hence $\text{Fid}(\mathcal{U}_{g_*}(\sigma)^{\otimes N}, \sigma^{\otimes N}) = \text{Fid}(\mathcal{U}_{g_*}(\sigma), \sigma)^N = \Delta^N \leq \Delta$, implying that $T(\mathcal{U}_{g_*}(\sigma)^{\otimes N}, \sigma^{\otimes N}) \geq 1 - \sqrt{\text{Fid}(\mathcal{U}_{g_*}(\sigma), \sigma)^N} \geq 1 - \sqrt{\Delta}$ for any positive integer N .

Assume that $\{\rho^{\otimes N}\}_N \xrightarrow{G\text{-cov.}} \{\sigma^{\otimes \lfloor rN \rfloor}\}_N$. Then for any $\epsilon > 0$, there exists a sequence of G -covariant channels $\{\mathcal{E}_N\}_N$ such that

$$T\left(\mathcal{E}_N(\rho^{\otimes N}), \sigma^{\otimes \lfloor rN \rfloor}\right) \leq \epsilon \quad (\text{A24})$$

for all sufficiently large N . Note that this also implies

$$\forall g \in G, \quad T\left(\mathcal{E}_N(\mathcal{U}_g(\rho)^{\otimes N}), \mathcal{U}'_g(\sigma)^{\otimes \lfloor rN \rfloor}\right) \leq \epsilon \quad (\text{A25})$$

since \mathcal{E}_N is a G -covariant channel. If $r > 0$, we have $\lfloor rN \rfloor \geq 1$ for $N \geq 1/r$. However, from the triangle inequality of trace distance,

$$T\left(\mathcal{U}_{g_*}(\sigma)^{\otimes \lfloor rN \rfloor}, \sigma^{\otimes \lfloor rN \rfloor}\right) \quad (\text{A26})$$

$$\leq T\left(\mathcal{U}_{g_*}(\sigma)^{\otimes \lfloor rN \rfloor}, \mathcal{E}_N(\mathcal{U}_{g_*}(\rho)^{\otimes N})\right) + T\left(\mathcal{E}_N(\mathcal{U}_{g_*}(\rho)^{\otimes N}), \mathcal{E}_N(\rho^{\otimes N})\right) + T\left(\mathcal{E}_N(\rho^{\otimes N}), \sigma^{\otimes \lfloor rN \rfloor}\right) \quad (\text{A27})$$

$$= T\left(\mathcal{U}_{g_*}(\sigma)^{\otimes \lfloor rN \rfloor}, \mathcal{E}_N(\mathcal{U}_{g_*}(\rho)^{\otimes N})\right) + T\left(\mathcal{E}_N(\rho^{\otimes N}), \sigma^{\otimes \lfloor rN \rfloor}\right) \quad (\text{A28})$$

$$\leq 2\epsilon \quad (\text{A29})$$

holds for all sufficiently large N , which contradicts $T(\mathcal{U}_{g_*}(\sigma)^{\otimes N}, \sigma^{\otimes N}) \geq 1 - \sqrt{\Delta}$. Therefore, $r = 0$. \square

Appendix B: Proofs of lemmas in the converse part

1. Proof of the equivalence between two alternative definitions of asymptotic convertibility

Proposition 15. *For any sequences of states $\{\rho_N\}_N$ and $\{\sigma_N\}_N$, the following two conditions are equivalent:*

(i) *For any $\epsilon > 0$, there exists N_0 such that for any positive integer $N \geq N_0$, there exists a G -covariant channel \mathcal{E}_N satisfying $T(\mathcal{E}_N(\rho_N), \sigma_N) \leq \epsilon$.*

(ii) *There exists a sequence of G -covariant channels $\{\mathcal{E}_N\}_N$ such that $\lim_{N \rightarrow \infty} T(\mathcal{E}_N(\rho_N), \sigma_N) = 0$.*

Proof. (ii) \implies (i): Any sequence of G -covariant channels $\{\mathcal{E}_N\}_N$ such that $\lim_{N \rightarrow \infty} T(\mathcal{E}_N(\rho_N), \sigma_N) = 0$ provides an example of the channels satisfying condition (i).

(i) \implies (ii): Fix a sequence of positive numbers $\{\epsilon_k\}_k$ such that $\lim_{k \rightarrow \infty} \epsilon_k = 0$. Assume the condition (i) is true. Then for each $\epsilon_k > 0$, $\exists N_0(\epsilon_k) > 0$ such that $\forall N > N_0(\epsilon_k)$, $\exists \mathcal{E}_{N, \epsilon_k} : G\text{-cov. s.t. } T(\mathcal{E}_{N, \epsilon_k}(\rho_N), \sigma_N) \leq \epsilon_k$. Without loss of generality, we can assume $N_0(\epsilon_{k+1}) > N_0(\epsilon_k)$. Let us define G -covariant channels $\{\mathcal{E}_N\}_N$ by

$$\mathcal{E}_N := \mathcal{E}_{N, \epsilon_k} \text{ if } N_0(\epsilon_k) \leq N < N_0(\epsilon_{k+1}) \quad (\text{B1})$$

for $N \geq N_0(\epsilon_1)$. For $N < N_0(\epsilon_1)$, we arbitrarily fix a G -covariant channel \mathcal{E}_N . Then, for any $N \geq N_0(\epsilon_k)$, we have $T(\mathcal{E}_N(\rho_N), \sigma_N) \leq \epsilon_k$. Thus, we get $\lim_{N \rightarrow \infty} T(\mathcal{E}_N(\rho_N), \sigma_N) \leq \epsilon_k$. In the limit of $k \rightarrow \infty$, we find $\lim_{N \rightarrow \infty} T(\mathcal{E}_N(\rho_N), \sigma_N) = 0$ since $\{\epsilon_k\}_k$ is assumed to satisfy $\lim_{k \rightarrow \infty} \epsilon_k = 0$. \square

For example, in studies on convertibility in RTA, condition (i) is adopted in [50, 70], while condition (ii) is employed in [19, 21, 34] and the present paper. Note that such an equivalence is not specific to the convertibility via G -covariant channels, and a similar argument can be found, e.g., in [74].

2. Proof of Lemma 3

Let us first prove a lemma relating an inner product on a (mixed) state to a covariance matrix of its eigenstate with the largest eigenvalue when the state is close to a pure state.

Lemma 16 (Modification of Corollary 1 in [19] and Lemma 9 in [50]). *Let σ be an arbitrary state and ϕ be a pure state. Suppose that their infidelity by $\delta := 1 - \langle \phi | \sigma | \phi \rangle$ satisfies $\delta < 1/2$. For the eigenvector Φ with the largest eigenvalue of σ , it holds*

$$|\langle \phi | \Phi \rangle|^2 \geq 1 - 2\delta. \quad (\text{B2})$$

Let f be an operator monotone function such that $f(0) > 0$. Then for any linear operator O , it holds

$$\langle i[\sigma, O], i[\sigma, O] \rangle_{f, \sigma} \geq \frac{1}{f\left(\frac{\delta}{1-\delta}\right)} (1 - 2\delta)^2 \langle \Phi | O(I - \Phi)O^\dagger | \Phi \rangle. \quad (\text{B3})$$

The first half is proven in Corollary 1 in [19]. The second half is a modification of Corollary 1 in [19] and Lemma 9 in [50]. In Corollary 1 in [19] and Lemma 9 in [50], a similar inequality was proven for Hermitian operators O and monotone functions f satisfying the symmetry condition $f(t) = tf(t^{-1})$. Equation (B3) differs by a factor of 2 from those in [19, 50], and it is applicable to monotone functions without assuming the symmetry condition.

Proof of Lemma 16. Let p be the largest eigenvalue of σ . The eigenvalue decomposition of σ is written as

$$\sigma = \sum_{k=0}^{d-1} \lambda_k |k\rangle \langle k| = p |\Phi\rangle \langle \Phi| + \sum_{k=1}^{d-1} \lambda_k |k\rangle \langle k|, \quad (\text{B4})$$

where $\lambda_0 := p$ and $|0\rangle := |\Phi\rangle$. Since p is assumed to be the largest eigenvalue, we have

$$1 - \delta = p |\langle \phi | \Phi \rangle|^2 + \sum_{k=1}^{d-1} \lambda_k |\langle \phi | k \rangle|^2 \leq p \sum_{k=0}^{d-1} |\langle \phi | k \rangle|^2 = p. \quad (\text{B5})$$

Therefore, we get

$$1 - \delta = \langle \phi | \sigma | \phi \rangle \quad (\text{B6})$$

$$= p |\langle \phi | \Phi \rangle|^2 + \sum_{k=1}^{d-1} \lambda_k |\langle \phi | k \rangle|^2 \quad (\text{B7})$$

$$\leq p |\langle \phi | \Phi \rangle|^2 + (1-p) \sum_{k=1}^{d-1} |\langle \phi | k \rangle|^2 \quad (\text{B8})$$

$$\leq p |\langle \phi | \Phi \rangle|^2 + \delta(1 - |\langle \phi | \Phi \rangle|^2), \quad (\text{B9})$$

meaning that

$$|\langle \phi | \Phi \rangle|^2 \geq \frac{1 - 2\delta}{p - \delta} \geq 1 - 2\delta, \quad (\text{B10})$$

where we have used $0 < 1 - 2\delta \leq p - \delta \leq 1$.

The second half of this lemma is a modification of Corollary 1 in [19] and Lemma 9 in [50]. For any operator monotone function f , we have

$$\langle i[\sigma, O], i[\sigma, O] \rangle_{f, \sigma} = \sum_{k, l; \lambda_l f(\lambda_k/\lambda_l) > 0} \frac{1}{\lambda_l f(\lambda_k/\lambda_l)} \langle k | [\sigma, O^\dagger] | l \rangle \langle l | [O, \sigma] | k \rangle \quad (\text{B11})$$

$$= \sum_{k, l; \lambda_l f(\lambda_k/\lambda_l) > 0} \frac{(\lambda_l - \lambda_k)^2}{\lambda_l f(\lambda_k/\lambda_l)} |\langle l | O | k \rangle|^2 + \sum_{k, l; \lambda_l f(\lambda_k/\lambda_l) > 0, l \neq 0} \frac{(\lambda_l - \lambda_k)^2}{\lambda_l f(\lambda_k/\lambda_l)} |\langle l | O | k \rangle|^2 \quad (\text{B12})$$

$$\geq \sum_{k; pf(\lambda_k/p) > 0} \frac{(p - \lambda_k)^2}{pf(\lambda_k/p)} |\langle \Phi | O | k \rangle|^2 \quad (\text{B13})$$

$$= \sum_{k=1}^{d-1} \frac{(p - \lambda_k)^2}{pf(\lambda_k/p)} |\langle \Phi | O | k \rangle|^2, \quad (\text{B14})$$

where we have used $pf(\lambda_k/p) \geq pf(0) > 0$ for all $k = 0, \dots, d-1$ in the last equality. Since $p - \lambda_k \geq 1 - 2\delta$, we have $(p - \lambda_k)^2 \geq (1 - 2\delta)^2$ if $\delta \leq 1/2$. Moreover, since $\lambda_k/p \leq \delta/(1 - \delta)$ and $p \leq 1$, we get $pf(\lambda_k/p) \leq f(\lambda_k/p) \leq f(\delta/(1 - \delta))$ for any monotonic function f . Therefore,

$$\langle i[\sigma, O], i[\sigma, O] \rangle_{f, \sigma} \geq \sum_{k=1}^{d-1} \frac{(p - \lambda_k)^2}{pf(\lambda_k/p)} |\langle \Phi | O | k \rangle|^2 \quad (\text{B15})$$

$$\geq \frac{1}{f\left(\frac{\delta}{1-\delta}\right)} (1 - 2\delta)^2 \sum_{k=1}^{d-1} |\langle \Phi | O | k \rangle|^2 \quad (\text{B16})$$

$$= \frac{1}{f\left(\frac{\delta}{1-\delta}\right)} (1 - 2\delta)^2 \langle \Phi | O(I - \Phi)O^\dagger | \Phi \rangle. \quad (\text{B17})$$

□

In the asymptotic conversion scenario, the initial i.i.d. pure state is transformed into a state close to i.i.d. copies of the target pure state. In order to further bound the right hand side of Eq. (B3), let us consider any linear operator O on \mathcal{H} and its i.i.d. extension O_N defined by

$$O_N := \sum_{n=1}^N O^{(n)}, \quad O^{(n)} := I^{\otimes n-1} \otimes O \otimes I^{\otimes N-n}. \quad (\text{B18})$$

By decomposing O into Hermitian and anti-Hermitian parts as

$$O = A + iB, \quad A := \frac{O + O^\dagger}{2}, \quad B := \frac{O - O^\dagger}{2i}, \quad (\text{B19})$$

we get

$$O_N = A_N + iB_N, \quad (\text{B20})$$

$$A_N := \sum_{n=1}^N A^{(n)}, \quad A^{(n)} := I^{\otimes n-1} \otimes A \otimes I^{\otimes N-n}, \quad (\text{B21})$$

$$B_N := \sum_{n=1}^N B^{(n)}, \quad B^{(n)} := I^{\otimes n-1} \otimes B \otimes I^{\otimes N-n}. \quad (\text{B22})$$

For any pure state Φ_N , we find

$$\left\langle \Phi_N \left| O_N (I - \Phi_N) O_N^\dagger \right| \Phi_N \right\rangle = V(\Phi_N, A_N) + V(\Phi_N, B_N) - i \langle \Phi_N | [A_N, B_N] | \Phi_N \rangle, \quad (\text{B23})$$

where V denotes the variance $V(\Phi_N, A_N) := \langle \Phi_N | A_N^2 | \Phi_N \rangle - \langle \Phi_N | A_N | \Phi_N \rangle^2$. We derive a lower bound for each contribution in Eq. (B23) under the assumption that Φ_N is sufficiently close to an i.i.d. copies of a pure state $\phi^{\otimes N}$.

A bound for the last term in Eq. (B23) can be easily obtained as follows:

Lemma 17. *Let A, B be arbitrary Hermitian operators. We denote their i.i.d. extensions by A_N and B_N defined in Eqs. (B21) and (B22). For arbitrary state ρ_N and σ_N , it holds*

$$|\text{Tr}(i[A_N, B_N]\rho_N) - \text{Tr}(i[A_N, B_N]\sigma_N)| \leq N \|i[A, B]\|_\infty \|\rho_N - \sigma_N\|_1, \quad (\text{B24})$$

where $\|\cdot\|_\infty$ denotes the operator norm.

Proof. Since

$$i[A_N, B_N] = i \sum_{n, n'=1}^N [A^{(n)}, B^{(n')}] = i \sum_{n=1}^N [A^{(n)}, B^{(n)}] \quad (\text{B25})$$

we get

$$\|i[A_N, B_N]\|_\infty \leq \sum_{n=1}^N \|i[A^{(n)}, B^{(n)}]\|_\infty = N \|i[A, B]\|_\infty. \quad (\text{B26})$$

Therefore,

$$|\text{Tr}(i[A_N, B_N]\rho_N) - \text{Tr}(i[A_N, B_N]\sigma_N)| = |\text{Tr}(i[A_N, B_N](\rho_N - \sigma_N))| \quad (\text{B27})$$

$$\leq \|i[A_N, B_N]\|_\infty \|\rho_N - \sigma_N\|_1 \quad (\text{B28})$$

$$\leq N \|i[A, B]\|_\infty \|\rho_N - \sigma_N\|_1. \quad (\text{B29})$$

□

Since $i[A_N, B_N]$ is Hermitian, its expectation value is a real number. Applying this lemma for pure states Φ_N and $\phi^{\otimes N}$, we get

$$-i \langle \Phi_N | [A_N, B_N] | \Phi_N \rangle \geq -i \langle \phi^{\otimes N} | [A_N, B_N] | \phi^{\otimes N} \rangle - N \|i[A, B]\|_\infty \|\Phi_N - \phi^{\otimes N}\|_1. \quad (\text{B30})$$

On the other hand, the variances can change more drastically in general. Nevertheless, for a pure state close to an i.i.d. pure state, we can prove the following bound:

Lemma 18 (Extension of Lemma 10 in [50]). *Let A be an arbitrary Hermitian operator on a finite-dimensional Hilbert space \mathcal{H} . We denote their i.i.d. extensions by A_N defined in Eq. (B21). Fix a pure state $|\phi\rangle \in \mathcal{H}$ and a sufficiently small parameter ϵ . There exists a function $h_1(\epsilon)$ independent of N satisfying $\lim_{\epsilon \rightarrow 0} h_1(\epsilon) = 0$ such that for any sequence of pure states $\{\Phi_N\}_N$ satisfying $\limsup_{N \rightarrow \infty} T(\Phi_N, \phi^{\otimes N}) < \epsilon$, it holds*

$$V(\Phi_N, A_N) \geq (1 - h_1(\epsilon)) V(\phi^{\otimes N}, A_N) \quad (\text{B31})$$

for all sufficiently large N .

As we have mentioned in Sec. IV B, the variance per copy can generally change of the order of $O(N\epsilon_N)$ for a state Φ_N that is ϵ_N -close to $\phi^{\otimes N}$ in trace distance. Even when $\lim_{N \rightarrow \infty} \epsilon_N = 0$, the variance per copy may change significantly in the limit of $N \rightarrow \infty$, and therefore, the variance has an asymptotic discontinuity. Nevertheless, Lemma 18 shows that when the state deviates slightly from $\phi^{\otimes N}$, the variance per copy can decrease at most $h_1(\epsilon)V(\phi, A)$, which is independent of N . The central limit theorem provides insight into this mechanism. The probability distribution of $\phi^{\otimes N}$ with respect to the observable A_N is approximated by the normal distribution as $N \rightarrow \infty$ after properly redefining the random variable. When this probability distribution is modified due to the change in state, the variance cannot be reduced radically because the tail of the normal distribution is exponentially small.

Proof of Lemma 18. When all the eigenvalues of A are degenerate, the statement trivially holds since $V(\Phi_N, A_N) = V(\phi^{\otimes N}, A_N) = 0$. Below, we assume the difference between the largest and smallest eigenvalues of A is non-vanishing, which we denote $\Delta a > 0$. Similarly, if the variance $\sigma^2 := \langle \phi | A^2 | \phi \rangle - \langle \phi | A | \phi \rangle^2$ vanishes for $|\phi\rangle$, then the inequality is trivial since $V(\phi^{\otimes N}, A_N) = N\sigma^2 = 0$. Therefore, we also assume that $\sigma > 0$.

Define

$$\Delta\mu_N := \langle \phi^{\otimes N} | A_N | \phi^{\otimes N} \rangle - \langle \Phi_N | A_N | \Phi_N \rangle, \quad (\text{B32})$$

$$Q_N := (A_N - \langle \Phi_N | A_N | \Phi_N \rangle I)^2 = (A_N - \langle \phi^{\otimes N} | A_N | \phi^{\otimes N} \rangle I + \Delta\mu_N I)^2. \quad (\text{B33})$$

Since

$$\langle \phi^{\otimes N} | (A_N - \langle \Phi_N | A_N | \Phi_N \rangle I)^2 | \phi^{\otimes N} \rangle = \langle \phi^{\otimes N} | (A_N - \langle \phi^{\otimes N} | A_N | \phi^{\otimes N} \rangle I)^2 | \phi^{\otimes N} \rangle + |\Delta\mu_N|^2, \quad (\text{B34})$$

we have

$$V(\Phi_N, A_N) - V(\phi^{\otimes N}, A_N) = |\Delta\mu_N|^2 - \text{Tr}(Q_N(\phi^{\otimes N} - \Phi_N)). \quad (\text{B35})$$

Here, we derive an upper bound of $\text{Tr}(Q_N(\phi^{\otimes N} - \Phi_N))$. Let us denote the eigenvalue decomposition of A by $A = \sum_{j=1}^d a_j |j\rangle \langle j|$. Its i.i.d. extension is written as

$$A_N = \sum_{j_1=1}^{d-1} \cdots \sum_{j_N=1}^{d-1} s(\mathbf{j}) |j\rangle \langle j|, \quad |\mathbf{j}\rangle := |j_1 \otimes j_2 \otimes \cdots \otimes j_N\rangle, \quad s(\mathbf{j}) := \sum_{n=1}^N a_{j_n}, \quad (\text{B36})$$

where $\mathbf{j} := (j_1, \dots, j_N)$. In the following, we abbreviate $\sum_{j_1=1}^{d-1} \cdots \sum_{j_N=1}^{d-1}$ as $\sum_{\mathbf{j}}$.

We decompose labels for eigenvectors of A_N into two disjoint sets:

$$\mathcal{A}^{(\text{core})}(x) := \left\{ \mathbf{j} \left| \left| \frac{\sqrt{N}}{\sigma} \left(\frac{s(\mathbf{j})}{N} - \mu \right) \right| \leq x \right. \right\}, \quad \mathcal{A}^{(\text{tail})}(x) := \left\{ \mathbf{j} \left| \left| \frac{\sqrt{N}}{\sigma} \left(\frac{s(\mathbf{j})}{N} - \mu \right) \right| > x \right. \right\}, \quad (\text{B37})$$

where $\mu := \langle \phi | A | \phi \rangle$, $\sigma^2 := \langle \phi | A^2 | \phi \rangle - \langle \phi | A | \phi \rangle^2$, and $x > 0$ is a positive parameter that will be fixed later.

Since Q_N is diagonal in $\{|\mathbf{j}\rangle\}$ basis, we can also decompose Q_N as follows:

$$Q_N = Q_N^{(\text{core})}(x) + Q_N^{(\text{tail})}(x), \quad (\text{B38})$$

$$Q_N^{(\text{core})}(x) := \sum_{\mathbf{j} \in \mathcal{A}^{(\text{core})}(x)} (s(\mathbf{j}) - N\mu + \Delta\mu_N)^2 |\mathbf{j}\rangle \langle \mathbf{j}|, \quad (\text{B39})$$

$$Q_N^{(\text{tail})}(x) := \sum_{\mathbf{j} \in \mathcal{A}^{(\text{tail})}(x)} (s(\mathbf{j}) - N\mu + \Delta\mu_N)^2 |\mathbf{j}\rangle \langle \mathbf{j}|. \quad (\text{B40})$$

Thus, we get

$$\text{Tr}(Q_N(\phi^{\otimes N} - \Phi_N)) = \text{Tr}(Q_N^{(\text{core})}(x)(\phi^{\otimes N} - \Phi_N)) + \text{Tr}(Q_N^{(\text{tail})}(x)(\phi^{\otimes N} - \Phi_N)). \quad (\text{B41})$$

Since the largest eigenvalue of $Q_N^{(\text{core})}(x)$ is less than $(x\sqrt{N}\sigma + |\Delta\mu_N|)^2$, the first term is bounded as

$$\text{Tr}(Q_N^{(\text{core})}(x)(\phi^{\otimes N} - \Phi_N)) \leq \left| \text{Tr}(Q_N^{(\text{core})}(x)(\phi^{\otimes N} - \Phi_N)) \right| \quad (\text{B42})$$

$$\leq (x\sqrt{N}\sigma + |\Delta\mu_N|)^2 \|\phi^{\otimes N} - \Phi_N\|_1 \quad (\text{B43})$$

$$\leq 2(x\sqrt{N}\sigma^2 + |\Delta\mu_N|)^2 \epsilon \quad (\text{B44})$$

for all sufficiently large N .

Since $Q_N^{(\text{tail})}(x)$ is a positive operator, we have

$$\text{Tr} \left(Q_N^{(\text{tail})}(x) (\phi^{\otimes N} - \Phi_N) \right) \leq \text{Tr} \left(Q_N^{(\text{tail})}(x) \phi^{\otimes N} \right). \quad (\text{B45})$$

Let p denote a probability distribution defined by $p(j) := |\langle j | \phi \rangle|^2$. We denote its i.i.d. extension by

$$p^{\otimes N}(\mathbf{j}) := p(j_1)p(j_2) \cdots p(j_N). \quad (\text{B46})$$

Then we have

$$\text{Tr} \left(Q_N^{(\text{tail})}(x) \phi^{\otimes N} \right) = \sum_{\mathbf{j} \in \mathcal{A}^{(\text{tail})}(x)} (s(\mathbf{j}) - N\mu + \Delta\mu_N)^2 p^{\otimes N}(\mathbf{j}) \quad (\text{B47})$$

$$= \sum_{\mathbf{j} \in \mathcal{A}^{(\text{tail})}(x)} \left((s(\mathbf{j}) - N\mu)^2 + 2\Delta\mu_N(s(\mathbf{j}) - N\mu) + |\Delta\mu_N|^2 \right) p^{\otimes N}(\mathbf{j}) \quad (\text{B48})$$

$$\leq \sum_{\mathbf{j} \in \mathcal{A}^{(\text{tail})}(x)} \left((s(\mathbf{j}) - N\mu)^2 + 2|\Delta\mu_N| |s(\mathbf{j}) - N\mu| + |\Delta\mu_N|^2 \right) p^{\otimes N}(\mathbf{j}). \quad (\text{B49})$$

From the Cauchy-Schwarz inequality, the second term is bounded as

$$\sum_{\mathbf{j} \in \mathcal{A}^{(\text{tail})}(x)} |s(\mathbf{j}) - N\mu| p^{\otimes N}(\mathbf{j}) \leq \sqrt{\sum_{\mathbf{j} \in \mathcal{A}^{(\text{tail})}(x)} |s(\mathbf{j}) - N\mu|^2 p^{\otimes N}(\mathbf{j})} \sqrt{\sum_{\mathbf{j} \in \mathcal{A}^{(\text{tail})}(x)} p^{\otimes N}(\mathbf{j})}. \quad (\text{B50})$$

Therefore, we get

$$\text{Tr} \left(Q_N^{(\text{tail})}(x) \phi^{\otimes N} \right) \leq \left(\sqrt{\sum_{\mathbf{j} \in \mathcal{A}^{(\text{tail})}(x)} (s(\mathbf{j}) - N\mu)^2 p^{\otimes N}(\mathbf{j})} + |\Delta\mu_N| \sqrt{\sum_{\mathbf{j} \in \mathcal{A}^{(\text{tail})}(x)} p^{\otimes N}(\mathbf{j})} \right)^2. \quad (\text{B51})$$

For i.i.d. random variables $Y_1, \dots, Y_N \sim p$, we denote

$$Z_N := \frac{\sqrt{N}}{\sigma} \left(\frac{1}{N} \sum_{i=1}^N Y_i - \mu \right) = \frac{1}{\sqrt{N}\sigma} \left(\sum_{i=1}^N Y_i - N\mu \right). \quad (\text{B52})$$

The tail contribution can be recast into

$$\sum_{\mathbf{j} \in \mathcal{A}^{(\text{tail})}(x)} (s(\mathbf{j}) - N\mu)^2 p^{\otimes N}(\mathbf{j}) = N\sigma^2 - \sum_{\mathbf{j} \in \mathcal{A}^{(\text{core})}(x)} (s(\mathbf{j}) - N\mu)^2 p^{\otimes N}(\mathbf{j}) \quad (\text{B53})$$

$$= N\sigma^2 (1 - \mathbb{E}_N[f(Z_N)]), \quad (\text{B54})$$

where we defined

$$f(z) := \begin{cases} z^2 & (\text{if } -x \leq z \leq x) \\ 0 & (\text{otherwise}) \end{cases}. \quad (\text{B55})$$

Since the Hilbert space is assumed to be finite-dimensional, the variance σ is finite. Therefore, we can apply the central limit theorem to Z_N and find that Z_N converges in distribution to the standard normal $\mathcal{N}(0, 1)$ as $N \rightarrow \infty$. Since $f(z)$ is bounded and its points of discontinuity $z = \pm x$ have measure zero in $\mathcal{N}(0, 1)$, we have

$$\lim_{N \rightarrow \infty} \mathbb{E}[f(Z_N)] = \int dz \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} f(z) = \int_{-x}^x dz \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} z^2. \quad (\text{B56})$$

Therefore, for any $\epsilon > 0$, $\mathbb{E}[f(Z_N)] \geq (1 - \epsilon) \int_{-x}^x dz \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} z^2$ holds for all sufficiently large N . In this case, it holds

$$\sum_{\mathbf{j} \in \mathcal{A}^{(\text{tail})}(x)} (s(\mathbf{j}) - N\mu)^2 p^{\otimes N}(\mathbf{j}) \leq N\sigma^2 \left(1 - (1 - \epsilon) \int_{-x}^x dz \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} z^2 \right) =: N\sigma^2 \tilde{g}_2(\epsilon, x). \quad (\text{B57})$$

On the other hand, from the Hoeffding inequality we have

$$\sum_{j \in \mathcal{A}^{(\text{tail})}(x)} p^{\otimes N}(j) \leq 2 \exp\left(-\frac{2\sigma^2 x^2}{(\Delta a)^2}\right) =: \tilde{g}_0(x), \quad (\text{B58})$$

where $\Delta a > 0$ denotes the difference between the largest and smallest eigenvalue of A . Therefore, we get

$$\text{Tr}\left(Q_N^{(\text{tail})}(x)\phi^{\otimes N}\right) \leq \left(\sqrt{N\sigma^2\tilde{g}_2(\epsilon, x)} + |\Delta\mu_N|\sqrt{\tilde{g}_0(x)}\right)^2. \quad (\text{B59})$$

From Eqs. (B44) and (B59), we have proven

$$V(\Phi_N, A_N) - V(\phi^{\otimes N}, A_N) = |\Delta\mu_N|^2 - \text{Tr}\left(Q_N(\phi^{\otimes N} - \Phi_N)\right) \quad (\text{B60})$$

$$\geq |\Delta\mu_N|^2 - \left(2(x\sqrt{N\sigma^2} + |\Delta\mu_N|)^2\epsilon + \left(\sqrt{N\sigma^2\tilde{g}_2(\epsilon, x)} + |\Delta\mu_N|\sqrt{\tilde{g}_0(x)}\right)^2\right) \quad (\text{B61})$$

holds for all sufficiently large N .

Now, let us take the parameter to be $x := \epsilon^{-1/3}$ and define

$$g_2(\epsilon) := \tilde{g}_2(\epsilon, \epsilon^{-1/3}), \quad g_0(\epsilon) := \tilde{g}_0(\epsilon^{-1/3}), \quad (\text{B62})$$

which satisfy $\lim_{\epsilon \rightarrow 0} g_2(\epsilon) = \lim_{\epsilon \rightarrow 0} g_0(\epsilon) = 0$.

For sufficiently small $\epsilon > 0$ satisfying $1 - 2\epsilon - g_0(\epsilon) > 0$, we have

$$|\Delta\mu_N|^2 - \left(2(\epsilon^{-1/3}\sqrt{N\sigma^2} + |\Delta\mu_N|)^2\epsilon + \left(\sqrt{N\sigma^2\tilde{g}_2(\epsilon, \epsilon^{-1/3})} + |\Delta\mu_N|\sqrt{\tilde{g}_0(\epsilon^{-1/3})}\right)^2\right) \quad (\text{B63})$$

$$= (1 - 2\epsilon - g_0(\epsilon)) \left(|\Delta\mu_N| - \frac{2\epsilon^{2/3}\sqrt{N\sigma^2} + \sqrt{N\sigma^2 g_0(\epsilon) g_2(\epsilon)}}{1 - 2\epsilon - g_0(\epsilon)} \right)^2 \quad (\text{B64})$$

$$- \left(\frac{(2\epsilon^{2/3}\sqrt{N\sigma^2} + \sqrt{N\sigma^2 g_0(\epsilon) g_2(\epsilon)})^2}{1 - 2\epsilon - g_0(\epsilon)} + 2\epsilon^{1/3}N\sigma^2 + N\sigma^2 g_2(\epsilon) \right). \quad (\text{B65})$$

Therefore, the right-hand side of Eq. (B61) is bounded as

$$|\Delta\mu_N|^2 - \left(2(\epsilon^{-1/3}\sqrt{N\sigma^2} + |\Delta\mu_N|)^2\epsilon + \left(\sqrt{N\sigma^2\tilde{g}_2(\epsilon, \epsilon^{-1/3})} + |\Delta\mu_N|\sqrt{\tilde{g}_0(\epsilon^{-1/3})}\right)^2\right) \geq -N\sigma^2 h_1(\epsilon), \quad (\text{B66})$$

where

$$h_1(\epsilon) := \frac{(2\epsilon^{2/3} + \sqrt{g_0(\epsilon)g_2(\epsilon)})^2}{1 - 2\epsilon - g_0(\epsilon)} + 2\epsilon^{1/3} + g_2(\epsilon). \quad (\text{B67})$$

We remark that this function satisfies $\lim_{\epsilon \rightarrow 0} h_1(\epsilon) = 0$ because $\lim_{\epsilon \rightarrow 0} g_0(\epsilon) = \lim_{\epsilon \rightarrow 0} g_2(\epsilon) = 0$. Since $N\sigma^2 = V(\phi^{\otimes N}, A_N)$, we find

$$V(\Phi_N, A_N) \geq (1 - h_1(\epsilon))V(\phi^{\otimes N}, A_N) \quad (\text{B68})$$

holds for all sufficiently large N , as long as $\epsilon > 0$ is sufficiently small so that $1 - 2\epsilon - g_0(\epsilon) > 0$. \square

From Lemmas 17 and 18, we obtain the following corollary:

Lemma 19. *Let O be an arbitrary linear operator on a finite-dimensional Hilbert space \mathcal{H} . We denote by O_N its i.i.d. extension defined in Eq. (B18). Fix a pure state $|\phi\rangle \in \mathcal{H}$ and a sufficiently small parameter ϵ . Then there exists a function $h_2(\epsilon)$ independent of N satisfying $\lim_{\epsilon \rightarrow 0} h_2(\epsilon) = 0$ such that for any sequence of pure states $\{\Phi_N\}_N$ satisfying $\limsup_{N \rightarrow \infty} T(\Phi_N, \phi^{\otimes N}) < \epsilon$, it holds*

$$\left\langle \Phi_N \left| O_N(I - \Phi_N)O_N^\dagger \right| \Phi_N \right\rangle \geq \langle \phi^{\otimes N} | O_N(I - \phi^{\otimes N})O_N^\dagger | \phi^{\otimes N} \rangle - Nh_2(\epsilon) \quad (\text{B69})$$

for all sufficiently large N .

Proof. From Lemma 18, we get

$$V(\Phi_N, A_N) + V(\Phi_N, B_N) \geq (1 - h_{1,A}(\epsilon))V(\phi^{\otimes N}, A_N) + (1 - h_{1,B}(\epsilon))V(\phi^{\otimes N}, B_N), \quad (\text{B70})$$

where $h_{1,A}$ and $h_{1,B}$ are real functions ensured to exist in Lemma 18, each of which in general depends on the difference between the largest and smallest eigenvalues and the variance of A and B , respectively. Combining Eqs. (B30) and (B70), we obtain the following bound:

$$\left\langle \Phi_N \left| O_N(I - \Phi_N)O_N^\dagger \right| \Phi_N \right\rangle \geq \langle \phi^{\otimes N} | O_N(I - \phi^{\otimes N})O_N^\dagger | \phi^{\otimes N} \rangle - Nh_2(\epsilon) \quad (\text{B71})$$

for all sufficiently large N , where the function

$$h_2(\epsilon) := h_{1,A}(\epsilon)V(\phi, A) + h_{1,B}(\epsilon)V(\phi, B) + 2\|i[A, B]\|_\infty \epsilon \quad (\text{B72})$$

is independent of N and satisfies $\lim_{\epsilon \rightarrow 0} h_2(\epsilon) = 0$. \square

From Lemma 16 and Lemma 19, we now prove Lemma 3.

Lemma (Restatement of Lemma 3). *Let f be an arbitrary operator monotone function such that $f(0) > 0$ and $f(0) = \lim_{\epsilon \rightarrow 0^+} f(\epsilon)$. For a linear operator O on a finite-dimensional Hilbert space, we denote its i.i.d. extension by O_N . Let $\{\sigma_N\}_N$ be an arbitrary sequence of states such that $\lim_{N \rightarrow \infty} T(\sigma_N, \phi^{\otimes N}) = 0$ for a pure state ϕ . Then there exists a real-valued function h independent of N satisfying $\lim_{\epsilon \rightarrow 0} h(\epsilon) = 0$ such that for any sufficiently small parameter $\epsilon > 0$, it holds*

$$f(0)\|i[\sigma_N, O_N]\|_{f, \sigma_N}^2 \geq NV(\phi, O) - Nh(\epsilon) + o(N) \quad (\text{B73})$$

for all sufficiently large N .

Proof. From Lemma 16, we get

$$\langle i[\sigma_N, O], i[\sigma_N, O] \rangle_{f, \sigma_N} \geq \frac{1}{f\left(\frac{\delta_N}{1-\delta_N}\right)} (1 - 2\delta_N)^2 \left\langle \Phi_N \left| O_N(I - \Phi)O_N^\dagger \right| \Phi_N \right\rangle, \quad (\text{B74})$$

where $\delta_N := 1 - \langle \phi^{\otimes N} | \sigma_N | \phi^{\otimes N} \rangle$ denotes the infidelity. For a sufficiently small $\epsilon > 0$, from Lemma 19, we further find

$$\langle i[\sigma_N, O], i[\sigma_N, O] \rangle_{f, \sigma_N} \geq \frac{1}{f\left(\frac{\delta_N}{1-\delta_N}\right)} (1 - 2\delta_N)^2 (N \langle \phi | O(I - \phi)O^\dagger | \phi \rangle - Nh_2(\epsilon)) \quad (\text{B75})$$

Note that from the Fuchs–van de Graaf inequalities, the infidelity satisfies $1 - \sqrt{1 - \delta_N} \leq \epsilon_N$, where $\epsilon_N := T(\sigma_N, \phi^{\otimes N})$. Since $\lim_{N \rightarrow \infty} \epsilon_N = 0$, $\delta_N \leq 1 - (1 - \epsilon)^2$ for all sufficiently large N . Since $c(\delta) := \frac{1}{f\left(\frac{\delta}{1-\delta}\right)} (1 - 2\delta)^2$ is a monotonically decreasing function of δ if $\delta \in [0, 1/2]$, we get

$$f(0) \langle i[\sigma_N, O], i[\sigma_N, O] \rangle_{f, \sigma_N} \geq \frac{f(0)}{f\left(\frac{\delta_N}{1-\delta_N}\right)} (1 - 2\delta_N)^2 N \langle \phi | O(I - \phi)O^\dagger | \phi \rangle - Nh(\epsilon), \quad (\text{B76})$$

where

$$h(\epsilon) := f(0)c(1 - (1 - \epsilon)^2)h_2(\epsilon) \quad (\text{B77})$$

is a real-valued function satisfying $\lim_{\epsilon \rightarrow 0} h(\epsilon) = 0$. Therefore, we get

$$f(0) \langle i[\sigma_N, O], i[\sigma_N, O] \rangle_{f, \sigma_N} \geq N \langle \phi | O(I - \phi)O^\dagger | \phi \rangle - Nh(\epsilon) + N \langle \phi | O(I - \phi)O^\dagger | \phi \rangle \left(\frac{f(0)}{f\left(\frac{\delta_N}{1-\delta_N}\right)} (1 - 2\delta_N)^2 - 1 \right). \quad (\text{B78})$$

The last term is $o(N)$ since

$$\lim_{N \rightarrow \infty} \frac{f(0)}{f\left(\frac{\delta_N}{1-\delta_N}\right)} (1 - 2\delta_N)^2 - 1 = \frac{f(0)}{f(0)} - 1 = 0, \quad (\text{B79})$$

where we have used $\lim_{N \rightarrow \infty} \delta_N = 0$ and $f(0) = \lim_{\epsilon \rightarrow 0^+} f(\epsilon)$. \square

Appendix C: Proofs of lemmas in the direct part

1. Proof of Lemma 6

Proof of Lemma 6. (ii) \implies (i): This is a direct consequence of the definition of the G -covariance of a channel and the fact that the trace distance is invariant under unitary transformations.

(i) \implies (ii): Define $\mathcal{E}' := \int_{g \in G} d\mu_G(g) \mathcal{U}'_{g^{-1}} \circ \mathcal{E} \circ \mathcal{U}_g$, where μ_G denotes the normalized right-invariant Haar measure on G . This map \mathcal{E}' is a G -covariant channel since

$$\forall h \in G, \quad \mathcal{E}' \circ \mathcal{U}_h = \int_{g \in G} d\mu_G(g) \mathcal{U}'_g \circ \mathcal{E} \circ \mathcal{U}_{g^{-1}h} \quad (\text{C1})$$

$$= \mathcal{U}'_h \circ \int_{g \in G} d\mu_G(g) \mathcal{U}'_{h^{-1}g} \circ \mathcal{E} \circ \mathcal{U}_{(h^{-1}g)^{-1}} \quad (\text{C2})$$

$$= \mathcal{U}'_h \circ \mathcal{E}' \quad (\text{C3})$$

holds. From the convexity of the trace distance, we get

$$T(\mathcal{E}'(\rho), \sigma) \leq \int_{g \in G} d\mu_G(g) T(\mathcal{U}'_{g^{-1}} \circ \mathcal{E} \circ \mathcal{U}_g(\rho), \sigma) \quad (\text{C4})$$

$$= \int_{g \in G} d\mu_G(g) T(\mathcal{E} \circ \mathcal{U}_g(\rho), \mathcal{U}'_g(\sigma)) \leq \epsilon. \quad (\text{C5})$$

□

2. Proof of Lemma 7

To prove Lemma 7, we divide the argument into two lemmas, Lemma 24 and Lemma 25, which are detailed later. See Fig. 1 for the relations among these lemmas. On the one hand, Lemma 24 shows that the change in the asymptotic rate r is effectively equivalent to the change in the scale of the parameters, which is proven by using results on QLAN for pure-state models [56, 57]. The asymptotics of QGTs are conserved in the interconversion among models in Lemma 24. On the other hand, Lemma 25 shows that the asymptotic conversion between two pure-state statistical models with rate one is possible when QGTs satisfy a matrix inequality. As a direct consequence of these lemmas, we complete the proof of Lemma 7 at the end of this subsection.




FIG. 1. A schematic figure of the relations among Lemma 7, Lemma 24 and Lemma 25. The direction of the arrow indicates the convertibility with vanishing error. In Lemma 24, we find the growth rate of i.i.d. copies can be set to 1 by adjusting the scaling of the parameters \mathbf{u} when analyzing the asymptotic convertibility. Lemma 25 shows that the asymptotic conversion between two pure-state models are possible if QGT is reduced during the conversion in the sense of matrix inequality. Combining these lemmas, Lemma 7 is proven.

To proceed further, we here prove several lemmas. In the analysis below, we expand unitary operators to extract the asymptotic behaviors of pure-state models. The following two lemmas serve as a foundation for our analysis.

Lemma 20. Let \mathcal{H} be a finite-dimensional Hilbert space and f be a function from \mathbb{R} to linear operators on \mathcal{H} and of class C^3 . Then,

$$\left\| f(1) - \left(f(0) + f'(0) + \frac{1}{2}f''(0) \right) \right\|_1 \leq \frac{1}{6} \max_{\theta \in [0,1]} \{ \|f^{(3)}(\theta)\|_1 \}. \quad (\text{C6})$$

Proof. By Taylor's theorem, we have

$$f(1) - \left(f(0) + f'(0) + \frac{1}{2}f''(0) \right) = \int_0^1 \frac{(1-\theta)^2}{2} f^{(3)}(\theta) d\theta. \quad (\text{C7})$$

By the triangle inequality, we get

$$\left\| f(1) - \left(f(0) + f'(0) + \frac{1}{2}f''(0) \right) \right\|_1 \leq \int_0^1 \left\| \frac{(1-\theta)^2}{2} f^{(3)}(\theta) \right\|_1 d\theta \leq \int_0^1 \frac{(1-\theta)^2}{2} M d\theta = \frac{M}{6}, \quad (\text{C8})$$

where $M := \max_{\theta \in [0,1]} \{ \|f^{(3)}(\theta)\|_1 \}$. □

Lemma 21. Let $l \in \mathbb{N}$, \mathcal{H} and \mathcal{H}' be finite-dimensional Hilbert spaces, ρ be a linear operator on \mathcal{H} , O be a Hermitian operator on \mathcal{H} , O' be a Hermitian operator on \mathcal{H}' , \mathcal{E} be a linear map from linear operators on \mathcal{H} to those on \mathcal{H}' , and $F_{\rho, \mathcal{E}}$ be a function from \mathbb{R} to linear operators on \mathcal{H}' defined by

$$F_{\rho, \mathcal{E}}(\theta) := e^{-i\theta O'} \mathcal{E}(e^{i\theta O} \rho e^{-i\theta O}) e^{i\theta O'} \quad \forall \theta \in \mathbb{R}. \quad (\text{C9})$$

Then, for any $\theta \in \mathbb{R}$,

$$F_{\rho, \mathcal{E}}^{(l)}(\theta) = \sum_{j=0}^l \binom{l}{j} e^{-i\theta O'} \underbrace{[-iO', \dots, [-iO', \mathcal{E}(e^{i\theta O} \underbrace{[iO, \dots, [iO, \rho] \dots] e^{-i\theta O}}_{l-j})] \dots]}_j e^{i\theta O'}, \quad (\text{C10})$$

and

$$\left\| F_{\rho, \mathcal{E}}^{(l)}(\theta) \right\|_1 \leq \|\rho\|_1 \|\mathcal{E}\| \cdot 2^l (\|O\|_\infty + \|O'\|_\infty)^l, \quad (\text{C11})$$

where $\|\mathcal{E}\|$ is defined by

$$\|\mathcal{E}\| := \max_{\|L\|_1=1} \{ \|\mathcal{E}(L)\|_1 \}. \quad (\text{C12})$$

Proof. We note that $\frac{d}{d\theta} e^{i\theta A} B e^{-i\theta A} = e^{i\theta A} [iA, B] e^{-i\theta A}$ for all linear operators A and B . By using the Leibniz rule, we get the expression for $F_{\rho, \mathcal{E}}^{(l)}(\theta)$. By applying the triangle inequality to the expression, we get

$$\left\| F_{\rho, \mathcal{E}}^{(l)}(\theta) \right\|_1 \leq \sum_{j=0}^l \binom{l}{j} \left\| e^{-i\theta O'} \underbrace{[-iO', \dots, [-iO', \mathcal{E}(e^{i\theta O} \underbrace{[iO, \dots, [iO, \rho] \dots] e^{-i\theta O}}_{l-j})] \dots]}_j e^{i\theta O'} \right\|_1. \quad (\text{C13})$$

We note that by the triangle inequality and Hölder's inequality, we have

$$\|[A, B]\|_1 = \|AB - BA\|_1 \leq \|AB\|_1 + \|BA\|_1 \leq \|A\|_\infty \|B\|_1 + \|B\|_1 \|A\|_\infty = 2\|A\|_\infty \|B\|_1 \quad (\text{C14})$$

for all linear operators A and B . By using this relation, the definition of $\|\mathcal{E}\|$, and the unitary invariance of the trace

norm, we get

$$\left\| F_{\rho, \mathcal{E}}^{(l)}(\theta) \right\|_1 \leq \sum_{j=0}^l \binom{l}{j} \left\| \underbrace{[-iO', \dots, [-iO', \mathcal{E}(e^{i\theta O} \underbrace{[iO, \dots, [iO, \rho] \dots]}_{l-j}) \dots]}_j e^{-i\theta O} \right\|_1 \quad (\text{C15})$$

$$\leq \sum_{j=0}^l \binom{l}{j} (2\|O'\|_\infty)^j \left\| \mathcal{E}(e^{i\theta O} \underbrace{[iO, \dots, [iO, \rho] \dots]}_{l-j}) \right\|_1 \quad (\text{C16})$$

$$\leq \sum_{j=0}^l \binom{l}{j} (2\|O'\|_\infty)^j \left\| \mathcal{E} \left\| e^{i\theta O} \underbrace{[iO, \dots, [iO, \rho] \dots]}_{l-j} e^{-i\theta O} \right\|_1 \right\|_1 \quad (\text{C17})$$

$$= \sum_{j=0}^l \binom{l}{j} (2\|O'\|_\infty)^j \left\| \mathcal{E} \left\| \underbrace{[iO, \dots, [iO, \rho] \dots]}_{l-j} \right\|_1 \right\|_1 \quad (\text{C18})$$

$$\leq \sum_{j=0}^l \binom{l}{j} (2\|O'\|_\infty)^j \left\| \mathcal{E} \right\| (2\|O\|_\infty)^{l-j} \|\rho\|_1 \quad (\text{C19})$$

$$= \|\rho\|_1 \|\mathcal{E}\| \cdot 2^l (\|O\|_\infty + \|O'\|_\infty)^l. \quad (\text{C20})$$

□

Using Lemma 20 and Lemma 21, we prove a lemma that relates the setup in QLAN and ours:

Lemma 22. *Let ϕ be an arbitrary pure state in a finite-dimensional Hilbert space. Fix a set of Hermitian operators $\mathbf{X} = \{X_i\}_{i=1}^m$. We define another set of Hermitian operator $\tilde{\mathbf{X}} = \{\tilde{X}_i\}_{i=1}^m$ by $\tilde{X}_i := \Pi_\phi^\perp X_i \Pi_\phi + \Pi_\phi X_i \Pi_\phi^\perp$, where $\Pi_\phi := \phi$ and $\Pi_\phi^\perp := I - \phi$ are projectors onto a subspace spanned by ϕ and its orthogonal space, respectively. For $\epsilon \in (0, 1/6)$, it holds*

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T\left(\phi_{\mathbf{u}/\sqrt{N}}^{\otimes N}, \tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes N}\right) = 0, \quad (\text{C21})$$

where $\phi_\theta := e^{i\theta \cdot \mathbf{X}} \phi e^{-i\theta \cdot \mathbf{X}}$ and $\tilde{\phi}_\theta := e^{i\theta \cdot \tilde{\mathbf{X}}} \phi e^{-i\theta \cdot \tilde{\mathbf{X}}}$ with $\theta \cdot \mathbf{X} = \sum_{i=1}^m \theta^i X_i$ and $\theta \cdot \tilde{\mathbf{X}} = \sum_{i=1}^m \theta^i \tilde{X}_i$ for $\theta \in \mathbb{R}^m$.

Proof. Let us define $O := \frac{1}{\sqrt{N}} \sum_{i=1}^m u^i X_i$ and $O' := \frac{1}{\sqrt{N}} \sum_{i=1}^m u^i \tilde{X}_i = \Pi_\phi^\perp O \Pi_\phi + \Pi_\phi O \Pi_\phi^\perp$ for $\mathbf{u} \in \mathbb{R}^m$ such that $\|\mathbf{u}\| < N^\epsilon$. Introducing a finite non-negative number L by

$$L := \max \left\{ \max_{\|\theta\|=1} \|\theta \cdot \mathbf{X}\|_\infty, \max_{\|\theta\|=1} \|\theta \cdot \tilde{\mathbf{X}}\|_\infty \right\}, \quad (\text{C22})$$

we have $\|O\|_\infty \leq L \frac{\|\mathbf{u}\|}{\sqrt{N}}$ and $\|O'\|_\infty \leq L \frac{\|\mathbf{u}\|}{\sqrt{N}}$. Applying Lemma 20 and Lemma 21 for $f(\theta) := e^{-i\theta O'} e^{i\theta O} \phi e^{-i\theta O} e^{i\theta O'}$, we find

$$\left| \left\langle \phi \left| f(1) - \left(f(0) + f'(0) + \frac{1}{2} f''(0) \right) \right| \phi \right\rangle \right| \leq \left\| f(1) - \left(f(0) + f'(0) + \frac{1}{2} f''(0) \right) \right\|_1 \quad (\text{C23})$$

$$\leq \frac{2^3}{6} (\|O\|_\infty + \|O'\|_\infty)^3 \leq \frac{2^6}{6} \left(L \frac{\|\mathbf{u}\|}{\sqrt{N}} \right)^3. \quad (\text{C24})$$

Since

$$\langle \phi | O'^n O^m | \phi \rangle = \begin{cases} 1 & (n, m) = (0, 0) \\ \langle \phi | O | \phi \rangle & (n, m) = (0, 1), \\ 0 & (n, m) = (1, 0) \end{cases}, \quad (\text{C25})$$

we get

$$\langle \phi | f(0) | \phi \rangle = 1, \quad (\text{C26})$$

$$\langle \phi | f'(0) | \phi \rangle = i \langle \phi | O | \phi \rangle - i \langle \phi | O' | \phi \rangle = 0. \quad (\text{C27})$$

Similarly, since

$$\langle \phi | O^m O^m | \phi \rangle = \begin{cases} \langle \phi | O^2 | \phi \rangle & (n, m) = (0, 2) \\ \langle \phi | O(I - \phi) O | \phi \rangle & (n, m) = (1, 1), (2, 0) \end{cases} \quad (\text{C28})$$

and hence

$$\sum_{\substack{n, m=0,1,2 \\ n+m=2}} \frac{(-i)^n i^m}{n! m!} \langle \phi | O^m O^m | \phi \rangle = -\frac{1}{2} \langle \phi | O^2 | \phi \rangle + \frac{1}{2} \langle \phi | O(I - \phi) O | \phi \rangle = -\frac{1}{2} \langle \phi | O | \phi \rangle^2 \quad (\text{C29})$$

we get

$$\langle \phi | f''(0) | \phi \rangle = \sum_{\substack{n, m, n', m'=0,1,2 \\ n+m+n'+m'=2}} \frac{(-i)^n i^m}{n! m!} \frac{(-i)^{m'} i^{n'}}{m'! n'!} \langle \phi | O'^n O^m | \phi \rangle \langle \phi | O^{m'} O^{n'} | \phi \rangle \quad (\text{C30})$$

$$= -\frac{1}{2} \langle \phi | O | \phi \rangle^2 + \langle \phi | O | \phi \rangle^2 - \frac{1}{2} \langle \phi | O | \phi \rangle^2 = 0. \quad (\text{C31})$$

Therefore, by using $\langle \phi | f(1) | \phi \rangle = \text{Fid}(\phi_{\mathbf{u}/\sqrt{N}}, \tilde{\phi}_{\mathbf{u}/\sqrt{N}})$, Eq. (C24) implies

$$\left| \text{Fid}(\phi_{\mathbf{u}/\sqrt{N}}, \tilde{\phi}_{\mathbf{u}/\sqrt{N}}) - 1 \right| \leq \frac{2^3}{6} (\|O\|_\infty + \|O'\|_\infty)^3 \leq \frac{2^6}{6} \left(L \frac{\|\mathbf{u}\|}{\sqrt{N}} \right)^3 \leq \frac{2^6}{6} (LN^{-1/2+\epsilon})^3 \quad (\text{C32})$$

for any \mathbf{u} such that $\|\mathbf{u}\| < N^\epsilon$.

Since $N^{-1/2+\epsilon} = o(1/N)$ if $\epsilon \in (0, 1/6)$, we get

$$\inf_{\|\mathbf{u}\| < N^\epsilon} \text{Fid}(\phi_{\mathbf{u}/\sqrt{N}}^{\otimes N}, \tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes N}) = \inf_{\|\mathbf{u}\| < N^\epsilon} \left(\text{Fid}(\phi_{\mathbf{u}/\sqrt{N}}, \tilde{\phi}_{\mathbf{u}/\sqrt{N}}) \right)^N \geq \left(1 + o\left(\frac{1}{N}\right) \right)^N, \quad (\text{C33})$$

implying that

$$\lim_{N \rightarrow \infty} \inf_{\|\mathbf{u}\| < N^\epsilon} \text{Fid}(\phi_{\mathbf{u}/\sqrt{N}}^{\otimes N}, \tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes N}) = \lim_{N \rightarrow \infty} \left(1 + o\left(\frac{1}{N}\right) \right)^N = 1. \quad (\text{C34})$$

From Fuchs-van de Graaf's inequalities, we also get

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T(\phi_{\mathbf{u}/\sqrt{N}}^{\otimes N}, \tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes N}) = 0. \quad (\text{C35})$$

□

Therefore, as long as the asymptotic properties of an i.i.d. copies of $\phi_{\mathbf{u}/\sqrt{N}} = e^{i\mathbf{u} \cdot \mathbf{X}/\sqrt{N}} \phi e^{-i\mathbf{u} \cdot \mathbf{X}/\sqrt{N}}$ is concerned, it suffice to analyze $\tilde{\phi}_{\mathbf{u}/\sqrt{N}} := e^{i\mathbf{u} \cdot \tilde{\mathbf{X}}/\sqrt{N}} \phi e^{i\mathbf{u} \cdot \tilde{\mathbf{X}}/\sqrt{N}}$.

The asymptotic behavior of $\tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes N}$ can be understood by studies on QLAN. The result in [57] is complicated since it aims to prove a general statement applicable to a wide class of statistical models. In this paper, we only focus on pure-state models. For pure-state unitary models, the result in [57] can be stated as follows:

Lemma 23 (Theorem 3.3 in [57], particularized to pure-state model. See also [56]). *Let $|\phi\rangle$ be an arbitrary state in a d -dimensional Hilbert space. For any fixed set of Hermitian operators $\{X_i\}_{i=1}^m$, we define a pure-state qudit model $\tilde{\phi}_\theta$ by*

$$|\tilde{\phi}_\theta\rangle := \exp\left(-i \sum_{i=1}^m \theta^i \tilde{X}_i\right) |\phi\rangle, \quad \tilde{X}_i := (I - \phi) X_i \phi + \phi X_i (I - \phi). \quad (\text{C36})$$

We introduce a $(d-1) \times m$ matrix defined by

$$iC_{ki} := \langle k | \tilde{X}_i | 0 \rangle, \quad (\text{C37})$$

where $\{|k\rangle\}_{k=1}^{d-1}$ denotes an orthonormal basis of the subspace orthogonal to $|\phi\rangle$.

Let \mathcal{H}_{d-1} denote the $(d-1)$ -mode Fock space. For $\mathbf{z} = (z_1, \dots, z_{d-1})^\top \in \mathbb{C}^{d-1}$, we denote the coherent state by

$$|\mathbf{z}\rangle := \left(\bigotimes_{i=1}^{d-1} e^{z_i a_i^\dagger - z_i^* a_i} |0\rangle_i \right) \in \mathcal{H}_{d-1}, \quad (\text{C38})$$

where a_i^\dagger, a_i and $|0\rangle_i$ denote the creation and annihilation operator and the vacuum state for the i th mode.

For $\epsilon \in (0, 1/9)$, there exists a sequence of quantum channels $\{\mathcal{T}_N\}_N$ and $\{\mathcal{S}_N\}_N$ such that

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T\left(\mathcal{T}_N(\tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes N}), |C\mathbf{u}\rangle \langle C\mathbf{u}|\right) = 0, \quad (\text{C39})$$

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T\left(\tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes N}, \mathcal{S}_N(|C\mathbf{u}\rangle \langle C\mathbf{u}|)\right) = 0. \quad (\text{C40})$$

By using Lemma 22 and Lemma 23, we prove the relation between the asymptotic rate and the scale in the parameters:

Lemma 24. Let $|\phi\rangle$ be an arbitrary pure state in a finite-dimensional Hilbert space \mathcal{H} . Fix any sets of Hermitian operators $\mathbf{X} = \{X_i\}_{i=1}^m$ and introduce a pure-state statistical model $\phi_\theta := e^{i\theta \cdot \mathbf{X}} \phi e^{-i\theta \cdot \mathbf{X}}$ for $\theta \in \mathbb{R}^m$. For any $r > 0$, there exist sequences of quantum channels $\{\mathcal{E}_N\}_N$ and $\{\mathcal{E}'_N\}_N$ such that

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T\left(\mathcal{E}_N\left(\phi_{\frac{\mathbf{u}}{\sqrt{rN}}}^{\otimes N}\right), \phi_{\frac{\mathbf{u}}{\sqrt{rN}}}^{\otimes [rN]}\right) = 0, \quad (\text{C41})$$

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T\left(\phi_{\frac{\mathbf{u}}{\sqrt{rN}}}^{\otimes N}, \mathcal{E}'_N\left(\phi_{\frac{\mathbf{u}}{\sqrt{rN}}}^{\otimes [rN]}\right)\right) = 0 \quad (\text{C42})$$

for $\epsilon \in (0, 1/9)$.

Proof. For $\tilde{X}_i := (I - \phi)X_i\phi + \phi X_i(I - \phi)$, consider a sequence of statistical models $\{(\tilde{\phi}_\theta)^{\otimes [rN]}\}_N$, where $r > 0$ and $\tilde{\phi}_\theta := e^{i\theta \cdot \tilde{\mathbf{X}}} \phi e^{-i\theta \cdot \tilde{\mathbf{X}}}$. From Lemma 23, there exist sequences of quantum channels $\{\mathcal{T}_N\}_N$ and $\{\mathcal{S}_N\}_N$ such that

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < [rN]^\epsilon} T\left(\mathcal{T}_{[rN]}\left(\tilde{\phi}_{\frac{\mathbf{u}}{\sqrt{[rN]}}}^{\otimes [rN]}\right), |C\mathbf{u}\rangle \langle C\mathbf{u}|\right) = 0, \quad (\text{C43})$$

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < [rN]^\epsilon} T\left(\tilde{\phi}_{\frac{\mathbf{u}}{\sqrt{[rN]}}}^{\otimes [rN]}, \mathcal{S}_{[rN]}(|C\mathbf{u}\rangle \langle C\mathbf{u}|)\right) = 0. \quad (\text{C44})$$

for $\epsilon \in (0, 1/9)$. Introducing a new parameter $\mathbf{u}'/\sqrt{N} := \mathbf{u}/\sqrt{[rN]}$, we get

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}'\| < [rN]^\epsilon \sqrt{\frac{[rN]}{N}}} T\left(\mathcal{T}_{[rN]}\left(\tilde{\phi}_{\frac{\mathbf{u}'}{\sqrt{N}}}^{\otimes [rN]}\right), \left|\sqrt{\frac{[rN]}{N}} C\mathbf{u}'\right\rangle \left\langle \sqrt{\frac{[rN]}{N}} C\mathbf{u}'\right|\right) = 0, \quad (\text{C45})$$

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}'\| < [rN]^\epsilon \sqrt{\frac{[rN]}{N}}} T\left(\tilde{\phi}_{\frac{\mathbf{u}'}{\sqrt{N}}}^{\otimes [rN]}, \mathcal{S}_{[rN]}\left(\left|\sqrt{\frac{[rN]}{N}} C\mathbf{u}'\right\rangle \left\langle \sqrt{\frac{[rN]}{N}} C\mathbf{u}'\right|\right)\right) = 0. \quad (\text{C46})$$

For any $\epsilon' \in (0, \epsilon)$,

$$\lim_{N \rightarrow \infty} \frac{N^{\epsilon'}}{[rN]^\epsilon \sqrt{\frac{[rN]}{N}}} = 0, \quad (\text{C47})$$

meaning that $[rN]^\epsilon \sqrt{\frac{[rN]}{N}} \geq N^{\epsilon'}$ holds for all sufficiently large N . In addition, it holds

$$\lim_{N \rightarrow \infty} \left| \left\langle \sqrt{\frac{[rN]}{N}} C\mathbf{u}' \middle| \sqrt{r} C\mathbf{u}' \right\rangle \right| = \lim_{N \rightarrow \infty} e^{-\|C\mathbf{u}'\|^2 \left(\sqrt{r} - \sqrt{\frac{[rN]}{N}}\right)^2} = 1 \quad (\text{C48})$$

for any $\mathbf{u}' \in \mathbb{R}^m$. Therefore, we get

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^{\epsilon'}} T \left(\mathcal{T}_{\lfloor rN \rfloor} \left(\tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes \lfloor rN \rfloor} \right), |\sqrt{r}C\mathbf{u}\rangle \langle \sqrt{r}C\mathbf{u}| \right) = 0, \quad (\text{C49})$$

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^{\epsilon'}} T \left(\tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes \lfloor rN \rfloor}, \mathcal{S}_{\lfloor rN \rfloor} (|\sqrt{r}C\mathbf{u}\rangle \langle \sqrt{r}C\mathbf{u}|) \right) = 0. \quad (\text{C50})$$

Note that we can take an arbitrary value in $(0, 1/9)$ as ϵ' by appropriately taking $\epsilon \in (0, 1/9)$.

On the other hand, from Lemma 23, we have

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\mathcal{T}_N \left(\tilde{\phi}_{\sqrt{r}\mathbf{u}/\sqrt{N}}^{\otimes N} \right), |\sqrt{r}C\mathbf{u}\rangle \langle \sqrt{r}C\mathbf{u}| \right) = 0, \quad (\text{C51})$$

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\tilde{\phi}_{\sqrt{r}\mathbf{u}/\sqrt{N}}^{\otimes N}, \mathcal{S}_N (|\sqrt{r}C\mathbf{u}\rangle \langle \sqrt{r}C\mathbf{u}|) \right) = 0. \quad (\text{C52})$$

Therefore, we get

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\mathcal{S}_{\lfloor rN \rfloor} \circ \mathcal{T}_N \left(\tilde{\phi}_{\sqrt{r}\mathbf{u}/\sqrt{N}}^{\otimes N} \right), \tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes \lfloor rN \rfloor} \right) = 0, \quad (\text{C53})$$

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\tilde{\phi}_{\sqrt{r}\mathbf{u}/\sqrt{N}}^{\otimes N}, \mathcal{S}_N \circ \mathcal{T}_{\lfloor rN \rfloor} \left(\tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes \lfloor rN \rfloor} \right) \right) = 0 \quad (\text{C54})$$

for $\epsilon \in (0, 1/9)$. From Lemma 22, we also get

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\mathcal{S}_{\lfloor rN \rfloor} \circ \mathcal{T}_N \left(\phi_{\sqrt{r}\mathbf{u}/\sqrt{N}}^{\otimes N} \right), \phi_{\mathbf{u}/\sqrt{N}}^{\otimes \lfloor rN \rfloor} \right) = 0, \quad (\text{C55})$$

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\phi_{\sqrt{r}\mathbf{u}/\sqrt{N}}^{\otimes N}, \mathcal{S}_N \circ \mathcal{T}_{\lfloor rN \rfloor} \left(\phi_{\mathbf{u}/\sqrt{N}}^{\otimes \lfloor rN \rfloor} \right) \right) = 0 \quad (\text{C56})$$

for $\epsilon \in (0, 1/9)$. □

Importantly, for both $\phi_{\sqrt{r}\mathbf{u}/\sqrt{N}}^{\otimes N}$ and $\phi_{\mathbf{u}/\sqrt{N}}^{\otimes \lfloor rN \rfloor}$, the QGT with respect to the parameters \mathbf{u} is equal to $r \text{Cov}(\phi, \mathbf{X})$ in the limit of $N \rightarrow \infty$, where the elements of covariance matrix are defined by $(\text{Cov}(\phi, \mathbf{X}))_{ij} := \langle \phi | X_i (I - \phi) X_j | \phi \rangle$. In other words, the QGTs are preserved under reversible asymptotic conversion between $\{\phi_{\sqrt{r}\mathbf{u}/\sqrt{N}}^{\otimes N}\}_N$ and $\{\phi_{\mathbf{u}/\sqrt{N}}^{\otimes \lfloor rN \rfloor}\}_N$.

From the above lemma states, the asymptotic rate can be effectively set to $r = 1$ by adjusting the scale of the parameters. The following lemma claims that the asymptotic conversion with rate $r = 1$ is possible only if the covariance matrix of the original state is larger than that of the target state in the sense of matrix inequality:

Lemma 25. *Let $|\psi\rangle \in \mathcal{H}$ and $|\phi\rangle \in \mathcal{H}'$ be arbitrary pure states in finite-dimensional Hilbert spaces $\mathcal{H}, \mathcal{H}'$. Fix any sets of Hermitian operators $\mathbf{X} = \{X_i\}_{i=1}^m$ and $\mathbf{X}' = \{X'_i\}_{i=1}^m$. If $\text{Cov}(\psi, \mathbf{X}) \geq \text{Cov}(\phi, \mathbf{X}')$ holds, then there exists a quantum channel \mathcal{E} such that*

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\mathcal{E}^{\otimes N} \left(\psi_{\mathbf{u}/\sqrt{N}}^{\otimes N} \right), \phi_{\mathbf{u}/\sqrt{N}}^{\otimes N} \right) = 0 \quad (\text{C57})$$

for $\epsilon \in (0, 1/6)$, where $\psi_{\boldsymbol{\theta}} := e^{i\boldsymbol{\theta} \cdot \mathbf{X}} \psi e^{-i\boldsymbol{\theta} \cdot \mathbf{X}}$ and $\phi_{\boldsymbol{\theta}} := e^{i\boldsymbol{\theta} \cdot \mathbf{X}'} \phi e^{-i\boldsymbol{\theta} \cdot \mathbf{X}'}$ for $\boldsymbol{\theta} \in \mathbb{R}^m$.

Proof. Following Lemma 22, we introduce $\tilde{X}_i := (I - \psi)X_i\psi + \psi X_i(I - \psi)$ and $\tilde{X}'_i := (I - \phi)X'_i\phi + \phi X'_i(I - \phi)$ and denote

$$\tilde{\psi}_{\mathbf{u}/\sqrt{N}} := e^{i\frac{1}{\sqrt{N}}\mathbf{u} \cdot \tilde{\mathbf{X}}} \psi e^{-i\frac{1}{\sqrt{N}}\mathbf{u} \cdot \tilde{\mathbf{X}}}, \quad \tilde{\phi}_{\mathbf{u}/\sqrt{N}} := e^{i\frac{1}{\sqrt{N}}\mathbf{u} \cdot \tilde{\mathbf{X}'}} \phi e^{-i\frac{1}{\sqrt{N}}\mathbf{u} \cdot \tilde{\mathbf{X}'}}. \quad (\text{C58})$$

Let us fix orthonormal bases $\{|k\rangle\}_{k=1}^{d-1}$ and $\{|k'\rangle\}_{k'=1}^{d'-1}$ for the orthogonal complements to $|\psi\rangle$ and $|\phi\rangle$, respectively, where $d := \dim \mathcal{H}$ and $d' := \dim \mathcal{H}'$. For matrices $C : \mathbb{C}^m \rightarrow \mathbb{C}^{d-1}$, $C' : \mathbb{C}^m \rightarrow \mathbb{C}^{d'-1}$ whose elements are given by

$$(C)_{ki} := -i \langle k | X_i | \psi \rangle, \quad (C')_{\nu i} := -i \langle \nu | X'_i | \phi \rangle, \quad (\text{C59})$$

we define matrices $Z : \mathbb{C}^{d-1} \rightarrow \mathbb{C}^{d-1}$ and $\Gamma := \mathbb{C}^{d-1} \rightarrow \mathbb{C}^{d-1}$ by

$$Z := C' C^+, \quad (\text{C60})$$

and

$$\Gamma := I - Z^\dagger Z, \quad (\text{C61})$$

where C^+ denotes the Moore–Penrose inverse of C . Note that the assumption that $\text{Cov}(\psi, \mathbf{X}) \geq \text{Cov}(\phi, \mathbf{X}')$ is equivalent to

$$C^\dagger C \geq C'^\dagger C'. \quad (\text{C62})$$

Let P be the projector to the range of C , given by $P = CC^+$. By using Eq. (C62), it holds

$$\Gamma = I - Z^\dagger Z \geq P - Z^\dagger Z = P^2 - Z^\dagger Z = (C^+)^{\dagger} (C^\dagger C - C'^\dagger C') C^+ \geq (C^+)^{\dagger} C^+ \geq 0, \quad (\text{C63})$$

i.e., Γ is positive-semidefinite. By using the square root $\sqrt{\Gamma}$ of Γ , the operators $\{K_k\}_{k=0}^{d-1}$ defined by

$$K_0 := |\phi\rangle \langle \psi| + \sum_{k'=1}^{d'-1} \sum_{k=1}^{d-1} z_{k'k} |k'\rangle \langle k|, \quad (\text{C64})$$

$$K_k := \sum_{l=1}^{d-1} \left(\sqrt{\Gamma} \right)_{kl} |\phi\rangle \langle l| \quad (k = 1, 2, \dots, d-1), \quad (\text{C65})$$

satisfy the normalization condition:

$$\sum_{k=0}^{d-1} K_k^\dagger K_k = |\psi\rangle \langle \psi| + \sum_{k,l=1}^{d-1} \sum_{k',l'=1}^{d'-1} z_{k'l}^* z_{kl} \delta_{k'l'} |k\rangle \langle l| + \sum_{k,l=1}^{d-1} \sum_{m=1}^{d-1} \left(\sqrt{\Gamma} \right)_{mk}^* \left(\sqrt{\Gamma} \right)_{ml} |k\rangle \langle l| \quad (\text{C66})$$

$$= |\psi\rangle \langle \psi| + \sum_{k,l=1}^{d-1} |k\rangle \langle l| (Z^\dagger Z + \Gamma)_{kl} = |\psi\rangle \langle \psi| + \sum_{k=1}^{d-1} |k\rangle \langle k| = I. \quad (\text{C67})$$

Therefore, the linear map $\mathcal{E}(\cdot) := \sum_{k=0}^{d-1} K_k(\cdot) K_k^\dagger$ is a completely positive trace-preserving map, i.e., a quantum channel. In the following, we prove that this quantum channel satisfies Eq. (C57).

Let us define Hermitian operators O, O' by $O := \mathbf{u} \cdot \tilde{\mathbf{X}} / \sqrt{N}$ and $O' := \mathbf{u}' \cdot \tilde{\mathbf{X}}' / \sqrt{N}$. Introducing a finite non-negative number L by

$$L := \max \left\{ \max_{\|\boldsymbol{\theta}\|=1} \|\boldsymbol{\theta} \cdot \tilde{\mathbf{X}}\|_\infty, \max_{\|\boldsymbol{\theta}\|=1} \|\boldsymbol{\theta} \cdot \tilde{\mathbf{X}}'\|_\infty \right\}, \quad (\text{C68})$$

we have $\|O\|_\infty \leq L \frac{\|\mathbf{u}\|}{\sqrt{N}}$ and $\|O'\|_\infty \leq L \frac{\|\mathbf{u}'\|}{\sqrt{N}}$. Applying Lemma 20 and Lemma 21 for $f(\theta) := e^{-i\theta O'} \mathcal{E}(e^{i\theta O} \psi e^{-i\theta O}) e^{i\theta O'}$, we get

$$\left| \left\langle \phi \left| f(1) - \left(f(0) + f'(0) + \frac{1}{2} f''(0) \right) \right| \phi \right\rangle \right| \leq \left\| f(1) - \left(f(0) + f'(0) + \frac{1}{2} f''(0) \right) \right\|_1 \quad (\text{C69})$$

$$\leq \frac{2^3}{6} (\|O\|_\infty + \|O'\|_\infty)^3 \leq \frac{2^6}{6} \left(L \frac{\|\mathbf{u}\|}{\sqrt{N}} \right)^3, \quad (\text{C70})$$

where we have used the fact that $\|\mathcal{E}\| \leq 1$ holds for any positive and trace-preserving map \mathcal{E} [75].

Let us evaluate each term of $\langle \phi | (f(0) + f'(0) + \frac{1}{2} f''(0)) | \phi \rangle$. From the definition of the channel \mathcal{E} , we get $\mathcal{E}(\psi) = \phi$, implying that

$$\langle \phi | f(0) | \phi \rangle = 1. \quad (\text{C71})$$

Since $\langle \phi | O' | \phi \rangle = 0$, we have $\langle \phi | O' \mathcal{E}(\psi) | \phi \rangle = 0$ and $\langle \phi | \mathcal{E}(\psi) O' | \phi \rangle = 0$. In addition, since

$$\mathcal{E}(O\psi) = \frac{-i}{\sqrt{N}} \sum_{k'=1}^{d'-1} (ZC\mathbf{u})_{k'} |k'\rangle \langle \phi|, \quad (\text{C72})$$

we also get $\langle \phi | \mathcal{E}(O\psi) | \phi \rangle = 0$ and $\langle \phi | \mathcal{E}(\psi O) | \phi \rangle = 0$. Therefore,

$$\langle \phi | f'(0) | \phi \rangle = 0. \quad (\text{C73})$$

From $\mathcal{E}(\psi) = \phi$ and $\langle \phi | O' | \phi \rangle = 0$, we get

$$\frac{1}{2} \langle \phi | (O'^2 \mathcal{E}(\psi) + \mathcal{E}(\psi) O'^2) | \phi \rangle = \langle \phi | O'^2 | \phi \rangle = \frac{1}{N} \sum_{i,j=1}^m u^i u^j \langle \phi | \tilde{X}'_i \tilde{X}'_j | \phi \rangle = \frac{1}{N} (C'^\dagger \mathbf{u})^\dagger C' \mathbf{u}, \quad (\text{C74})$$

$$\langle \phi | O' \mathcal{E}(\psi) O' | \phi \rangle = 0. \quad (\text{C75})$$

From Eq. (C72), we find $\langle \phi | \mathcal{E}(O\psi) = 0$ and $\mathcal{E}(\psi O) | \phi \rangle = 0$. Therefore,

$$\langle \phi | \mathcal{E}(O\psi) O' | \phi \rangle = 0, \quad \langle \phi | O' \mathcal{E}(\psi O) | \phi \rangle = 0. \quad (\text{C76})$$

From Eq. (C72), we also get

$$O' \mathcal{E}(O\psi) = \frac{1}{N} (C' \mathbf{u})^\dagger Z C \mathbf{u} | \phi \rangle \langle \phi |, \quad \mathcal{E}(\psi O) O' = \frac{1}{N} (Z C \mathbf{u})^\dagger C' \mathbf{u} | \phi \rangle \langle \phi |, \quad (\text{C77})$$

implying that

$$\langle \phi | O' \mathcal{E}(O\psi) | \phi \rangle = \frac{1}{N} (C' \mathbf{u})^\dagger Z C \mathbf{u}, \quad \langle \phi | \mathcal{E}(\psi O) O' | \phi \rangle = \frac{1}{N} (Z C \mathbf{u})^\dagger C' \mathbf{u}. \quad (\text{C78})$$

From the definition of the quantum channel \mathcal{E} , we get

$$\mathcal{E}(O^2 \psi) = \frac{1}{N} \mathcal{E}((C \mathbf{u})^\dagger (C \mathbf{u}) | \psi \rangle \langle \psi |) = \frac{1}{N} (C \mathbf{u})^\dagger (C \mathbf{u}) | \phi \rangle \langle \phi |, \quad (\text{C79})$$

$$\mathcal{E}(O\psi O) = \frac{1}{N} \sum_{k,l=1}^{d-1} (C \mathbf{u})_k (C \mathbf{u})_l^* \mathcal{E}(|k\rangle \langle l|) = \frac{1}{N} \left(\sum_{k',l'=1}^{d'-1} (Z C \mathbf{u})_{k'} (Z C \mathbf{u})_{l'}^* |k'\rangle \langle l'| + (C \mathbf{u})^\dagger \Gamma C \mathbf{u} | \phi \rangle \langle \phi | \right), \quad (\text{C80})$$

and hence

$$\frac{1}{2} \langle \phi | \mathcal{E}(O^2 \psi + \psi O^2) | \phi \rangle = \frac{1}{N} (C \mathbf{u})^\dagger C \mathbf{u}, \quad \langle \phi | \mathcal{E}(O\psi O) | \phi \rangle = \frac{1}{N} (C \mathbf{u})^\dagger \Gamma (C \mathbf{u}). \quad (\text{C81})$$

By combining these results, we get

$$\begin{aligned} \left\langle \phi \left| \frac{1}{2} f'(0) \right| \phi \right\rangle &= -\frac{1}{2} \langle \phi | (O'^2 \mathcal{E}(\psi) - 2O' \mathcal{E}(\psi) O' + \mathcal{E}(\psi) O'^2) | \phi \rangle + \langle \phi | (O' \mathcal{E}(O\psi - \psi O) - \mathcal{E}(O\psi - \psi O) O') | \phi \rangle \\ &\quad - \frac{1}{2} \langle \phi | \mathcal{E}(O^2 \psi - 2O\psi O + \psi O^2) | \phi \rangle \end{aligned} \quad (\text{C82})$$

$$= \frac{1}{N} \left(-(C'^\dagger \mathbf{u})^\dagger C' \mathbf{u} + (C' \mathbf{u})^\dagger Z C \mathbf{u} + (Z C \mathbf{u})^\dagger C' \mathbf{u} - (C \mathbf{u})^\dagger C \mathbf{u} + (C \mathbf{u})^\dagger \Gamma (C \mathbf{u}) \right) \quad (\text{C83})$$

$$= -\frac{1}{N} \mathbf{u}^\dagger (C' - ZC)^\dagger (C' - ZC) \mathbf{u}. \quad (\text{C84})$$

Let Q be the projector Q to the support of C , given by $Q = C^+ C$. From $C^\dagger C \geq C'^\dagger C'$, we get

$$(C' - ZC)^\dagger (C' - ZC) = (I - Q)(C'^\dagger C')(I - Q) \leq (I - Q)C^\dagger C(I - Q) = (I - Q)C^\dagger (C - CC^+ C) = 0. \quad (\text{C85})$$

We thus have $C' - ZC = 0$, which implies

$$\left\langle \phi \left| \frac{1}{2} f'(0) \right| \phi \right\rangle = 0. \quad (\text{C86})$$

Since $\text{Fid}(\mathcal{E}(\tilde{\psi}_{\mathbf{u}/\sqrt{N}}), \tilde{\phi}_{\mathbf{u}/\sqrt{N}}) = \langle \phi | f(1) | \phi \rangle$, from Eqs. (C70), (C71), (C73), and (C86), we have proven

$$\left| \text{Fid}(\mathcal{E}(\tilde{\psi}_{\mathbf{u}/\sqrt{N}}), \tilde{\phi}_{\mathbf{u}/\sqrt{N}}) - 1 \right| \leq \frac{2^6}{6} \left(L \frac{\|\mathbf{u}\|}{\sqrt{N}} \right)^3 \leq \frac{2^6}{6} (LN^{-1/2+\epsilon})^3 \quad (\text{C87})$$

for any \mathbf{u} such that $\|\mathbf{u}\| < N^\epsilon$.

Since $N^{-1/2+\epsilon} = o(1/N)$ if $\epsilon \in (0, 1/6)$, we get

$$\inf_{\|\mathbf{u}\| < N^\epsilon} \text{Fid} \left(\mathcal{E}^{\otimes N}(\tilde{\psi}_{\mathbf{u}/\sqrt{N}}^{\otimes N}), \tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes N} \right) = \inf_{\|\mathbf{u}\| < N^\epsilon} \text{Fid} \left(\mathcal{E}(\tilde{\psi}_{\mathbf{u}/\sqrt{N}}), \tilde{\phi}_{\mathbf{u}/\sqrt{N}} \right)^N \geq \left(1 + o\left(\frac{1}{N}\right) \right)^N, \quad (\text{C88})$$

and hence

$$\lim_{N \rightarrow \infty} \inf_{\|\mathbf{u}\| < N^\epsilon} \text{Fid} \left(\mathcal{E}^{\otimes N}(\tilde{\psi}_{\mathbf{u}/\sqrt{N}}^{\otimes N}), \tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes N} \right) = 1 \quad (\text{C89})$$

for $\epsilon \in (0, 1/6)$. From the Fuchs-van de Graaf inequalities, we also have

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\mathcal{E}^{\otimes N}(\tilde{\psi}_{\mathbf{u}/\sqrt{N}}^{\otimes N}), \tilde{\phi}_{\mathbf{u}/\sqrt{N}}^{\otimes N} \right) = 0. \quad (\text{C90})$$

Therefore, by using Lemma 22, we get

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\mathcal{E}^{\otimes N}(\psi_{\mathbf{u}/\sqrt{N}}^{\otimes N}), \phi_{\mathbf{u}/\sqrt{N}}^{\otimes N} \right) = 0 \quad (\text{C91})$$

for $\epsilon \in (0, 1/6)$. □

By using Lemma 24 and Lemma 25, we complete the proof of Lemma 7.

Lemma (Restatement of Lemma 7). *For sets of Hermitian operators $\mathbf{X} := \{X_i\}_{i=1}^m$ and $\mathbf{X}' := \{X'_i\}_{i=1}^m$, we define $\mathcal{U}_\theta(\cdot) := e^{i\theta \cdot \mathbf{X}}(\cdot)e^{-i\theta \cdot \mathbf{X}}$ and $\mathcal{U}'_\theta(\cdot) := e^{i\theta \cdot \mathbf{X}'}(\cdot)e^{-i\theta \cdot \mathbf{X}'}$ for $\theta \in \mathbb{R}^m$. For given two pure states ψ and ϕ , we define pure-state statistical models by $\mathcal{U}_\theta(\psi)$ and $\mathcal{U}'_\theta(\phi)$ and denote their QGTs by \mathcal{Q}^ψ and \mathcal{Q}^ϕ , respectively. If $r > 0$ satisfies $\mathcal{Q}^\psi \geq r\mathcal{Q}^\phi$, then there exists a sequence of quantum channels $\{\mathcal{E}_N\}_N$ such that*

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\mathcal{E}_N \left(\mathcal{U}_{\frac{\mathbf{u}}{\sqrt{N}}}(\psi)^{\otimes N} \right), \mathcal{U}'_{\frac{\mathbf{u}}{\sqrt{N}}}(\phi)^{\otimes \lfloor rN \rfloor} \right) = 0 \quad (\text{C92})$$

for $\epsilon \in (0, 1/9)$.

Proof. The assumption $\mathcal{Q}^\psi \geq r\mathcal{Q}^\phi$ is equivalent to $\text{Cov}(\psi, \mathbf{X}) \geq r\text{Cov}(\phi, \mathbf{X}')$. Lemma 25 implies that there is a channel \mathcal{E} such that

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\left(\mathcal{E} \left(e^{i\frac{\mathbf{u} \cdot \mathbf{X}}{\sqrt{N}}} \psi e^{-i\frac{\mathbf{u} \cdot \mathbf{X}}{\sqrt{N}}} \right) \right)^{\otimes N}, \left(e^{i\frac{\mathbf{u} \cdot \sqrt{r}\mathbf{X}'}{\sqrt{N}}} \phi e^{-i\frac{\mathbf{u} \cdot \sqrt{r}\mathbf{X}'}{\sqrt{N}}} \right)^{\otimes N} \right) = 0 \quad (\text{C93})$$

for $\epsilon \in (0, 1/9)$. On the other hand, Lemma 24 implies there is a sequence of channels $\{\mathcal{E}'_N\}_N$ such that

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\mathcal{E}'_N \left(e^{i\frac{\mathbf{u} \cdot \sqrt{r}\mathbf{X}'}{\sqrt{N}}} \phi e^{-i\frac{\mathbf{u} \cdot \sqrt{r}\mathbf{X}'}{\sqrt{N}}} \right)^{\otimes N}, \left(e^{i\frac{\mathbf{u} \cdot \mathbf{X}'}{\sqrt{N}}} \phi e^{-i\frac{\mathbf{u} \cdot \mathbf{X}'}{\sqrt{N}}} \right)^{\otimes \lfloor rN \rfloor} \right) = 0. \quad (\text{C94})$$

Therefore, for $\mathcal{E}'_N \circ \mathcal{E}^{\otimes N}$, we have

$$\lim_{N \rightarrow \infty} \sup_{\|\mathbf{u}\| < N^\epsilon} T \left(\mathcal{E}'_N \circ \mathcal{E}^{\otimes N} \left(e^{i\frac{\mathbf{u} \cdot \mathbf{X}}{\sqrt{N}}} \psi e^{-i\frac{\mathbf{u} \cdot \mathbf{X}}{\sqrt{N}}} \right)^{\otimes N}, \left(e^{i\frac{\mathbf{u} \cdot \mathbf{X}'}{\sqrt{N}}} \phi e^{-i\frac{\mathbf{u} \cdot \mathbf{X}'}{\sqrt{N}}} \right)^{\otimes \lfloor rN \rfloor} \right) = 0. \quad (\text{C95})$$

□

3. Proof of measurability of the set defined in Eq. (33)

For $\delta > 0$, let us define

$$H(\delta) := \{\hat{g} \in G \mid \exists \theta, \mathcal{U}_\theta \circ \mathcal{U}_{\hat{g}}(\psi) = \mathcal{U}_{\hat{g}}(\psi), \|\theta\| \leq \delta\} \quad (\text{C96})$$

and prove that $H(\delta)$ is a closed set. Let $\{h_n\}_{n \in \mathbb{N}}$ be a convergent sequence in $H(\delta)$ such that $h_n \rightarrow h$ as $n \rightarrow \infty$. From the definition of $H(\delta)$, there exists a sequence $\{\theta_n\}_{n \in \mathbb{N}}$ such that $\mathcal{U}_{\theta_n} \circ \mathcal{U}_{h_n}(\psi) = \mathcal{U}_g(\psi)$ and $\|\theta_n\| \leq \delta$. Due to the compactness of the closed ball of radius δ , we can extract a convergent subsequence $\{\theta_{n_k}\}_{k \in \mathbb{N}}$ such that $\theta_{n_k} \rightarrow \theta_0$ for some θ_0 as $k \rightarrow \infty$. Then, we have

$$\mathcal{U}_g(\psi) = \mathcal{U}_{\theta_{n_k}} \circ \mathcal{U}_{h_{n_k}}(\psi) \rightarrow \mathcal{U}_{\theta_0} \circ \mathcal{U}_h(\psi), \quad \delta \geq \|\theta_{n_k}\| \rightarrow \|\theta_0\| \quad (\text{C97})$$

as $k \rightarrow \infty$. Therefore, $h \in H(\delta)$, implying that $H(\delta)$ is a closed set.

From the definition of $G_{\text{succ.}}(\delta)$, we have

$$G_{\text{succ.}}(\delta) = \bigcup_{n \in \mathbb{N}} H\left(\delta - \frac{1}{n}\right). \quad (\text{C98})$$

Therefore, $G_{\text{succ.}}(\delta)$ is a countable union of closed sets and hence is a measurable set.

4. Proof of Lemma 8

In [76], a bound on the error in a state tomography process is proven:

Lemma 26 (Theorem 1 in [76]). *Fix any qudit state ρ . There exists an estimator $\hat{\rho}_n$ consuming n samples of states, $\rho^{\otimes n}$, such that*

$$\Pr(\|\hat{\rho}_n - \rho\|_1 \geq \delta) \leq de^{-\frac{n\delta^2}{43g(d)r^2}}, \quad \delta \in [0, 1], \quad (\text{C99})$$

where $r := \min\{\text{rank}(\rho), \text{rank}(\hat{\rho}_n)\}$ and $g(d)$ is a constant depending only on the dimension of the Hilbert space d .

To prove Lemma 8, we relate the magnitude of error in trace distance to that in the parameter θ . For this purpose, we begin by proving several lemmas. Since any compact Lie group G is isomorphic to a closed linear group, we assume G is a compact linear Lie group and adopt its Schatten norm in the following arguments.

Lemma 27. *Let U be a unitary representation of a compact linear Lie group G . Then,*

$$\forall \epsilon > 0 \exists \delta > 0 \forall g, g' \in G \left(\|U(g')\rho U(g')^\dagger - U(g)\rho U(g)^\dagger\|_1 < \delta \rightarrow \min_{h \in \text{Sym}_G(\rho)} \|g' - gh\|_\infty < \epsilon \right). \quad (\text{C100})$$

Proof. Since

$$\|U(g')\rho U(g')^\dagger - U(g)\rho U(g)^\dagger\|_1 = \|U(g^{-1}g')\rho U(g^{-1}g')^\dagger - \rho\|_1, \quad (\text{C101})$$

$$\|g' - gh\|_\infty = \|g^{-1}g' - h\|_\infty, \quad (\text{C102})$$

it is sufficient to prove that

$$\forall \epsilon > 0 \exists \delta > 0 \forall g \in G \left(\|U(g)\rho U(g)^\dagger - \rho\|_1 < \delta \rightarrow \min_{h \in \text{Sym}_G(\rho)} \|g - h\|_\infty < \epsilon \right). \quad (\text{C103})$$

We suppose that this statement does not hold. Then, we can take $\epsilon > 0$ and a sequence $(g_n)_{n \in \mathbb{N}}$ in G such that

$$\|U(g_n)\rho U(g_n)^\dagger - \rho\|_1 < \frac{1}{n}, \quad (\text{C104})$$

$$\min_{h \in \text{Sym}_G(\rho)} \|g_n - h\|_\infty \geq \epsilon \quad (\text{C105})$$

for all $n \in \mathbb{N}$. Since G is sequentially compact, we can take some subsequence $(g_{n(j)})_{j \in \mathbb{N}}$ that converges to $k \in G$. This subsequence satisfies

$$\|U(g_{n(j)})\rho U(g_{n(j)})^\dagger - \rho\|_1 < \frac{1}{n(j)} \leq \frac{1}{j}, \quad (\text{C106})$$

$$\min_{h \in \text{Sym}_G(\rho)} \|g_{n(j)} - h\|_\infty \geq \epsilon \quad (\text{C107})$$

for all $j \in \mathbb{N}$. By taking the limit of $j \rightarrow \infty$ in Eq. (C106), we get $\|U(k)\rho U(k)^\dagger - \rho\|_1 = 0$, which implies $k \in \text{Sym}_G(\rho)$. Thus Eq. (C107) implies that

$$\|g_{n(j)} - k\|_\infty \geq \min_{h \in \text{Sym}_G(\rho)} \|g_{n(j)} - h\|_\infty \geq \epsilon \quad (\text{C108})$$

for all $j \in \mathbb{N}$. This contradicts with the fact that the subsequence $(g_{n(j)})_{j \in \mathbb{N}}$ converges to k . \square

Lemma 28. *Let ψ be a linear operator and A be a Hermitian operator. Then,*

$$\|e^{iA}\psi e^{-iA} - \psi\|_1 \geq (1 - \|A\|_\infty) \|[A, \psi]\|_1. \quad (\text{C109})$$

Proof. We define $F(u) := e^{iuA}\psi e^{-iuA}$ for $u \in \mathbb{R}$. Then, we have

$$e^{iA}\psi e^{-iA} - \psi - i[A, \psi] = F(1) - F(0) - F'(0) = \int_0^1 du \int_0^u dv F''(v). \quad (\text{C110})$$

By the triangle inequality and Hölder's inequality, we get

$$\|e^{iA}\psi e^{-iA} - \psi - i[A, \psi]\|_1 \leq \int_0^1 du \int_0^u dv \|F''(v)\|_1 = \frac{1}{2} \|[A, [A, \psi]]\|_1 \leq \|A\|_\infty \|[A, \psi]\|_1. \quad (\text{C111})$$

By using the triangle inequality again, we have

$$\|e^{iA}\psi e^{-iA} - \psi - i[A, \psi]\|_1 \geq \|[A, \psi]\|_1 - \|e^{iA}\psi e^{-iA} - \psi\|_1. \quad (\text{C112})$$

By Eqs. (C111) and (C112), we get Eq. (C109). \square

We show an upper bound of $\|e^{i(A+B)}e^{-iA} - I\|_\infty$ for general Hermitian operators A and B .

Lemma 29. *Let A and B be Hermitian operators and satisfy $\|A + B\|_\infty + 2\|B\|_\infty < 1$. Then,*

$$\|e^{i(A+B)}e^{-iA} - I\|_\infty \leq \frac{\|B\|_\infty}{1 - \|A + B\|_\infty - 2\|B\|_\infty}. \quad (\text{C113})$$

Proof. By using the triangle inequality, we have

$$\begin{aligned} \|e^{i(A+B)}e^{-iA} - I\|_\infty &= \|e^{i(A+B)} - e^{iA}\|_\infty \\ &= \left\| \sum_{k=0}^{\infty} \frac{i^k}{k!} [(A+B)^k - A^k] \right\|_\infty \\ &\leq \sum_{k=0}^{\infty} \frac{1}{k!} \|[A+B]^k - A^k\|_\infty. \end{aligned} \quad (\text{C114})$$

Now let us prove $\|[A+B]^k - A^k\|_\infty \leq (\|A\|_\infty + \|B\|_\infty)^k - \|A\|_\infty^k$ by the mathematical induction. This inequality is trivial for $k=0$. Assume that the inequality holds for some k . Then we get

$$\|(A+B)^{k+1} - A^{k+1}\|_\infty = \|(A+B)((A+B)^k - A^k) - BA^k\|_\infty \quad (\text{C115})$$

$$\leq \|(A+B)\|_\infty \|(A+B)^k - A^k\|_\infty + \|BA^k\|_\infty \quad (\text{C116})$$

$$\leq (\|A\|_\infty + \|B\|_\infty)((\|A\|_\infty + \|B\|_\infty)^k - \|A\|_\infty^k) + \|B\|_\infty \|A\|_\infty^k \quad (\text{C117})$$

$$= (\|A\|_\infty + \|B\|_\infty)^{k+1} - \|A\|_\infty^{k+1}. \quad (\text{C118})$$

Thus, the inequality also holds for $k+1$.

Therefore, we get

$$\begin{aligned} \|e^{i(A+B)}e^{-iA} - I\|_\infty &\leq \sum_{k=0}^{\infty} \frac{1}{k!} [(\|A\|_\infty + \|B\|_\infty)^k - \|A\|_\infty^k] \\ &= e^{\|A\|_\infty + \|B\|_\infty} - e^{\|A\|_\infty} \\ &\leq \|B\|_\infty e^{\|A\|_\infty + \|B\|_\infty} \\ &\leq \|B\|_\infty e^{\|A+B\|_\infty + 2\|B\|_\infty} \\ &\leq \frac{\|B\|_\infty}{1 - \|A+B\|_\infty - 2\|B\|_\infty}, \end{aligned} \quad (\text{C119})$$

where we used $e^x - 1 \leq xe^x$ for all $x \in \mathbb{R}$ in the second inequality, the triangle inequality in the third inequality, and $e^x \leq 1/(1-x)$ for all $x \in [0, 1)$ in the fourth inequality. \square

When $\|A + B\|_\infty + 3\|B\|_\infty < 1$, we show an upper bound of $\|\log(e^{i(A+B)}e^{-iA})\|_\infty$. Note that $\log(e^{i(A+B)}e^{-iA})$ is guaranteed to be defined by Lemma 29.

Lemma 30. *Let A and B be Hermitian operators and satisfy $\|A + B\|_\infty + 3\|B\|_\infty < 1$. Then,*

$$\left\| \log \left(e^{i(A+B)} e^{-iA} \right) \right\|_\infty \leq \frac{\|B\|_\infty}{1 - \|A + B\|_\infty - 3\|B\|_\infty}. \quad (\text{C120})$$

Proof. By the definition of the logarithm, we have

$$\log \left(e^{i(A+B)} e^{-iA} \right) = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \left(e^{i(A+B)} e^{-iA} - I \right)^k. \quad (\text{C121})$$

By the triangle inequality, we have

$$\left\| \log \left(e^{i(A+B)} e^{-iA} \right) \right\|_\infty \leq \sum_{k=1}^{\infty} \frac{1}{k} \left\| e^{i(A+B)} e^{-iA} - I \right\|_\infty^k. \quad (\text{C122})$$

By applying Lemma 29 to the right-hand side of this inequality, we get

$$\begin{aligned} \left\| \log \left(e^{i(A+B)} e^{-iA} \right) \right\|_\infty &\leq \sum_{k=1}^{\infty} \frac{1}{k} \left(\frac{\|B\|_\infty}{1 - \|A + B\|_\infty - 2\|B\|_\infty} \right)^k \\ &\leq \sum_{k=1}^{\infty} \left(\frac{\|B\|_\infty}{1 - \|A + B\|_\infty - 2\|B\|_\infty} \right)^k \\ &= \frac{\|B\|_\infty}{1 - \|A + B\|_\infty - 3\|B\|_\infty}. \end{aligned} \quad (\text{C123})$$

□

By using these lemmas, we prove the following:

Lemma 31. *Let ρ be an operator, and U be a unitary representation of a compact Lie group G . Then there exist positive constants $\delta_*, c > 0$ such that for any $g, \hat{g} \in G$,*

$$\|\mathcal{U}_g(\rho) - \mathcal{U}_{\hat{g}}(\rho)\|_1 < \delta_* \implies \exists \varphi \text{ s.t. } \begin{cases} \mathcal{U}_\varphi \circ \mathcal{U}_{\hat{g}}(\rho) = \mathcal{U}_g(\rho) \\ \|\varphi\| \leq c \|\mathcal{U}_g(\rho) - \mathcal{U}_{\hat{g}}(\rho)\|_1 \end{cases}, \quad (\text{C124})$$

where $\mathcal{U}_\varphi(\cdot) := e^{i\varphi \cdot \mathbf{Y}}(\cdot)e^{-i\varphi \cdot \mathbf{Y}}$ with $\varphi \in \mathbb{R}^n$ and $\mathbf{Y} = \{Y_i\}_{i=1}^n$ is a basis of the linear span of $\{X_\mu\}_{\mu=1}^{\dim G}$, where X_μ is defined in Eq. (3).

Proof. We denote by \mathcal{A} the linear space spanned by $\{Y_i\}_{i=1}^n$, and define its linear subspace \mathcal{A}_0 by

$$\mathcal{A}_0 := \{A \in \mathcal{A} \mid [A, \rho] = 0\} \quad (\text{C125})$$

and its complementary subspace \mathcal{A}_1 .

Let us first consider an exceptional case where $\mathcal{A}_1 = \emptyset$, i.e., $\mathcal{A} = \mathcal{A}_0$. Let $G = \bigsqcup_{i=0}^k G_i$ be the decomposition into the connected components. Since X_μ commutes with ρ for all $\mu = 1, \dots, \dim G$, we find $\mathcal{U}_g(\rho) = \mathcal{U}_{g'}(\rho)$ if g and g' are in the same component. Note that the number of connected components is finite since G is assumed to be a compact Lie group. Therefore, there are only a finite number of distinct elements in the set $\{\mathcal{U}_g(\rho) \mid g \in G\}$. By setting

$$\delta_* := \min_{g, g' \in G; \mathcal{U}_g(\rho) \neq \mathcal{U}_{g'}(\rho)} \|\mathcal{U}_g(\rho) - \mathcal{U}_{g'}(\rho)\|_1 \quad (\text{C126})$$

and c to be arbitrary positive number, Eq. (C145) holds for $\varphi = \mathbf{0}$. We remark that this essentially includes the case where G is a finite group.

From now on, we consider the case where $\mathcal{A}_1 \neq \emptyset$. In this case, let us define

$$m := \min_{\|D\|_\infty=1, D \in \mathcal{A}_1} \|[D, \rho]\|_1 > 0. \quad (\text{C127})$$

By Lemma 27 and the continuity of U , we can take $\delta > 0$ such that

$$\forall g, g' \in G \quad (\|\mathcal{U}_g(\rho) - \mathcal{U}_{g'}(\rho)\|_1 < \delta \rightarrow \min_{h \in \text{Sym}_G(\rho)} \|U(g^{-1}\hat{g}h) - I\|_\infty < 1 - e^{-\frac{1}{8}}). \quad (\text{C128})$$

We define

$$l := \min_{\|\gamma\|=1} \left\| \sum_k \gamma^k Y_k \right\|_\infty. \quad (\text{C129})$$

Note that $l > 0$ since $\{Y_k\}$ are assumed to be linearly independent. We set $\delta_* := \min\{\delta, m/16\}$ and $c := 1/(lm)$ and prove that they satisfy Eq. (C145).

Fix arbitrary $g, \hat{g} \in G$ satisfying $\|\mathcal{U}_g(\rho) - \mathcal{U}_{\hat{g}}(\rho)\|_1 < \delta$, and find appropriate φ satisfying Eq. (C145). By the definition of δ , we can take $h \in \text{Sym}_G(\rho)$ such that $\|U(g^{-1}\hat{g}h) - I\|_\infty \leq 1 - e^{-1/8} < 1$. Thus we can define

$$A := -i \log(U(g^{-1}\hat{g}h)). \quad (\text{C130})$$

By the definition of the logarithm, we have

$$\begin{aligned} \|A\|_\infty &= \left\| \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} (U(g^{-1}\hat{g}h) - I)^k \right\|_\infty \\ &\leq \sum_{k=1}^{\infty} \frac{1}{k} \|U(g^{-1}\hat{g}h) - I\|_\infty^k \\ &= -\log(1 - \|U(g^{-1}\hat{g}h) - I\|_\infty) \\ &\leq \frac{1}{8}, \end{aligned} \quad (\text{C131})$$

where we used the triangle inequality in the first inequality and the second inequality follows from the choice of δ . By the definition of A , we have

$$\begin{aligned} \|U(\hat{g})\rho U(\hat{g})^\dagger - U(g)\rho U(g)^\dagger\|_\infty &= \|U(g^{-1}\hat{g}h)\rho U(g^{-1}\hat{g}h)^\dagger - \rho\|_\infty \\ &= \|e^{iA}\rho e^{-iA} - \rho\|_\infty \\ &\geq (1 - \|A\|_\infty) \|[A, \rho]\|_\infty \\ &\geq \frac{1}{2} \|[A, \rho]\|_\infty, \end{aligned} \quad (\text{C132})$$

where we used Lemma 28 in the first inequality and Eq. (C131) in the second inequality. Since \mathcal{A}_0 and \mathcal{A}_1 are complementary subspaces of \mathcal{A} , A can be decomposed into $A = A_0 + A_1$ with some $A_0 \in \mathcal{A}_0$ and $A_1 \in \mathcal{A}_1$. We can take some $D \in \mathcal{A}_1$ such that $A_1 = \|A_1\|_\infty D$ and $\|D\|_\infty = 1$. Then, we have

$$\|[A, \rho]\|_\infty = \|[A_1, \rho]\|_\infty = \|A_1\|_\infty \|[D, \rho]\|_1 \geq m \|A_1\|_\infty. \quad (\text{C133})$$

By using Eqs. (C133) and (C132), we have

$$\|A_1\| \leq \frac{1}{m} \|[A, \rho]\|_\infty \leq \frac{2}{m} \|U(\hat{g})\rho U(\hat{g})^\dagger - U(g)\rho U(g)^\dagger\|_\infty. \quad (\text{C134})$$

Since $\|U(\hat{g})\rho U(\hat{g})^\dagger - U(g)\rho U(g)^\dagger\|_\infty$ is upper bounded by δ and $\delta \leq m/16$, we get

$$\|A_1\| \leq \frac{2\delta}{m} \leq \frac{1}{8}. \quad (\text{C135})$$

By Eqs. (C131) and (C135) and Lemma 29, we get $\|e^{iA}e^{-iA_0} - I\|_\infty \leq 1/5 < 1$, which enables us to define

$$B := -i \log(e^{iA}e^{-iA_0}). \quad (\text{C136})$$

We define φ by

$$\sum_k \varphi^k Y_k = U(g)BU(g)^\dagger \quad (\text{C137})$$

and show that

$$\|\boldsymbol{\theta}\| \leq \frac{1}{lm} \|U(\hat{g})\rho U(\hat{g})^\dagger - U(g)\rho U(g)^\dagger\|_1, \quad (\text{C138})$$

$$e^{-i\sum_k \varphi^k Y_k} U(\hat{g})\rho U(\hat{g})^\dagger e^{i\sum_k \varphi^k Y_k} = U(g)\rho U(g)^\dagger. \quad (\text{C139})$$

As for the proof of Eq. (C139), by using Eqs. (C137), (C136), and (C130) we have

$$\begin{aligned} U(\hat{g})^\dagger e^{i\sum_k \varphi^k Y_k} &= U(\hat{g}^{-1})U(g)e^{iB}U(g)^\dagger \\ &= U(\hat{g}^{-1}g)e^{iA}e^{-iA_0}U(g)^\dagger \\ &= U(\hat{g}^{-1}g)U(g^{-1}\hat{g}h)e^{-iA_0}U(g)^\dagger \\ &= U(h)e^{-iA_0}U(g)^\dagger, \end{aligned} \quad (\text{C140})$$

which implies that

$$e^{-i\sum_k \varphi^k Y_k} U(\hat{g})\rho U(\hat{g})^\dagger e^{i\sum_k \varphi^k Y_k} = U(g)e^{iA_0}U(h)^\dagger \rho U(h)e^{-iA_0}U(g)^\dagger = U(g)\rho U(g)^\dagger. \quad (\text{C141})$$

Finally, we prove Eq. (C138). By Eqs. (C131) and (C135) and Lemma 30, we get

$$\|B\|_\infty \leq \frac{\|A_1\|_\infty}{1 - \|A\|_\infty - 3\|A_1\|_\infty} \leq \frac{\|A_1\|_\infty}{2}. \quad (\text{C142})$$

Since we can take γ satisfying $\boldsymbol{\varphi} = \|\boldsymbol{\varphi}\|\boldsymbol{\gamma}$ and $\|\boldsymbol{\gamma}\| = 1$, we have

$$\|B\|_\infty = \left\| \sum_k \varphi^k Y_k \right\|_\infty = \|\boldsymbol{\varphi}\| \left\| \sum_k \varphi^k Y_k \right\|_\infty \geq l\|\boldsymbol{\varphi}\|. \quad (\text{C143})$$

By using Eqs. (C143),(C142), and (C134), we get

$$\|\boldsymbol{\varphi}\| \leq \frac{\|B\|_\infty}{l} \leq \frac{\|A_1\|_\infty}{2l} \leq \frac{1}{lm} \|U(\hat{g})\psi U(\hat{g})^\dagger - U(g)\psi U(g)^\dagger\|_\infty \leq \frac{1}{lm} \|U(\hat{g})\psi U(\hat{g})^\dagger - U(g)\psi U(g)^\dagger\|_1. \quad (\text{C144})$$

□

As a generalization of this lemma, we prove the following:

Lemma 32. *Let ρ be an operator, and U be a unitary representation of a compact Lie group G . Then there exist positive constants $\delta_*, c > 0$ such that for any $g, \hat{g} \in G$,*

$$\|\mathcal{U}_g(\rho) - \mathcal{U}_{\hat{g}}(\rho)\|_1 < \delta_* \implies \exists \boldsymbol{\theta} \text{ s.t. } \begin{cases} \mathcal{U}_{\boldsymbol{\theta}} \circ \mathcal{U}_{\hat{g}}(\rho) = \mathcal{U}_g(\rho) \\ \|\boldsymbol{\theta}\| \leq c\|\mathcal{U}_g(\rho) - \mathcal{U}_{\hat{g}}(\rho)\|_1 \end{cases}, \quad (\text{C145})$$

where $\mathcal{U}_{\boldsymbol{\theta}}(\cdot) := e^{i\boldsymbol{\theta} \cdot \mathbf{X}}(\cdot)e^{-i\boldsymbol{\theta} \cdot \mathbf{X}}$ for $\boldsymbol{\theta} \in \mathbb{R}^{\dim G}$, where $\mathbf{X} = \{X_\mu\}_{\mu=1}^{\dim G}$ is defined in Eq. (3).

Proof. Since $\mathbf{Y} = \{Y_i\}_{i=1}^n$ in Lemma 31 is a basis of the linear span of $\{X_\mu\}_{\mu=1}^{\dim G}$, there is an $n \times \dim G$ matrix L such that $L\mathbf{X} = \mathbf{Y}$. By using the transpose L^\top of L , we define $\boldsymbol{\theta} \in \mathbb{R}^{\dim G}$ by $\boldsymbol{\theta} := L^\top \boldsymbol{\varphi}$, which satisfies $\boldsymbol{\theta} \cdot \mathbf{X} = \boldsymbol{\varphi} \cdot \mathbf{Y}$. Note that $\|\boldsymbol{\theta}\| \leq \|L^\top\|_2 \|\boldsymbol{\varphi}\|$ holds, where $\|L^\top\|_2$ denotes the matrix norm induced from the 2-norm for vectors, which is finite. Therefore, Lemma 32 holds by replacing the constant c in Lemma 31 to $c\|L^\top\|_2$. □

Combining Lemma 26 and Lemma 32, we finally prove Lemma 8:

Lemma (Restatement of Lemma 8). *Let G be a compact Lie group and ρ be an arbitrary state. Fix $\epsilon \in (0, 1/2)$. Then there exists an estimator of $g \in G$ by using consuming $\mathcal{U}_g(\rho)^{\otimes n}$ with $n = N^{1-\epsilon}$ such that its success probability satisfies*

$$\lim_{N \rightarrow \infty} \Pr_{\mathcal{U}_g(\psi)^{\otimes N^{1-\epsilon}}} (\|\boldsymbol{\theta}\| < N^{-1/2+\epsilon}) = 1. \quad (\text{C146})$$

Proof. Fix $\epsilon \in (0, 1/2)$. Let $\delta_*, c > 0$ be constants assured to exist by Lemma 32. Applying Lemma 26 to a given state $\mathcal{U}_g(\rho)$ with $\delta := N^{-1/2+\epsilon}/(2c)$, we obtain a bound for the error probability in state tomography given by

$$\begin{aligned} & \Pr\left(\|\hat{\rho}_n - \mathcal{U}_g(\rho)\|_1 < N^{-1/2+\epsilon}/(2c)\right) \\ & \geq 1 - de^{-\frac{n(N^{-1/2+\epsilon}/(2c))^2}{43g(d)r^2}} \\ & = 1 - de^{-\frac{N^\epsilon}{172g(d)r^2c^2}} \end{aligned} \quad (\text{C147})$$

by performing measurements on $n = N^{1-\epsilon}$ copies of the system in a state $\mathcal{U}_g(\rho)^{\otimes n}$. From the estimated state $\hat{\rho}_n$, we pick up any $\hat{g} \in G$ satisfying $\|\mathcal{U}_{\hat{g}}(\rho) - \hat{\rho}_n\|_1 < N^{-1/2+\epsilon}/(2c)$ as an estimate of the true value $g \in G$. If there is no \hat{g} satisfying $\|\mathcal{U}_{\hat{g}}(\rho) - \hat{\rho}_n\|_1 < N^{-1/2+\epsilon}/(2c)$, which may happen with an exponentially small probability, we arbitrarily select \hat{g} as an element in G , such as $e \in G$. Whenever $\|\hat{\rho}_n - \mathcal{U}_g(\rho)\|_1 < N^{-1/2+\epsilon}/(2c)$ holds, the estimated value $\hat{g} \in G$ satisfies $\|\mathcal{U}_{\hat{g}}(\rho) - \mathcal{U}_g(\rho)\|_1 < N^{-1/2+\epsilon}/c$ due to the triangle inequality. In this case, for all sufficiently large N such that $N^{-1/2+\epsilon}/c < \delta_*$, we get $\|\boldsymbol{\theta}\| < N^{-1/2+\epsilon}$. Therefore, the success probability is bounded as

$$\Pr_{\mathcal{U}_g(\psi)^{\otimes N^{1-\epsilon}}}\left(\|\boldsymbol{\theta}\| < N^{-1/2+\epsilon}\right) \geq 1 - c_1 e^{-c_2 N^\epsilon}, \quad (\text{C148})$$

where $c_1 := d$ and $c_2 := \frac{1}{172g(d)r^2c^2}$. In other words, the failure probability is exponentially small in N . In particular, it holds

$$\lim_{N \rightarrow \infty} \Pr_{\mathcal{U}_g(\psi)^{\otimes N^{1-\epsilon}}}\left(\|\boldsymbol{\theta}\| < N^{-1/2+\epsilon}\right) = 1. \quad (\text{C149})$$

□